Divergence of *germ cell-less* roles in germ line development across insect species

Authors: Jonchee A. Kao[1], Ben Ewen-Campen[2,3], Cassandra G. Extavour[1,2,4]

[1]Department of Molecular and Cellular Biology, Harvard University, 16 Divinity Ave, Cambridge, MA 02138, USA

[2]Department of Organismic and Evolutionary Biology, Harvard University, 16 Divinity Ave, Cambridge, MA 02138, USA

[3]Current address: Department of Genetics, Blavatnik Institute, Harvard Medical School, Harvard University, Boston, MA 02115, USA

[4]Howard Hughes Medical Institute, Chevy Chase, MD, USA

Corresponding author: Cassandra G. Extavour

Email: extavour@oeb.harvard.edu

Phone: (617) 496-1935

ORCIDs: JAK 0000-0003-1701-3265; BE-C 0000-0002-4326-3033, CGE 0000-0003-2922-5855

Keywords: *Oncopeltus fasciatus*; *Gryllus bimaculatus*; germ cell specification; *torso*; pole cells

Abstract

During development, sexually reproducing animals must specify and maintain the germ line, the

lineage of cells that gives rise to the next generation of animals. In the fruit fly *Drosophila*

melanogaster, *germ cell-less* (*gcl*) is required for the formation of primordial germ cells in the

form of cells that cellularize at the posterior pole of the embryo, called pole cells. Forming pole

cells is a mechanism of germ cell formation unique to a subset of insects. Even though most

animals do not form pole cells as primordial germ cells, *gcl* is conserved across Metazoa, raising

the question of how this conserved gene acquired its central role in the evolutionarily derived

process of pole cell formation. Here, we examine the functions of *gcl* in two other insects with

different modes of germ cell specification: the milkweed bug *Oncopeltus fasciatus* and the

cricket *Gryllus bimaculatus*. We found that *gcl* is involved in germ cell development, but not

strictly required for germ cell specification, in *O. fasciatus*, although it appears to function

through a mechanism different from that in *D. melanogaster*. In contrast, we could not detect any

impact on the embryonic germ line upon *gcl* knockdown in *G. bimaculatus*. This work serves as

a case study into how the roles of genes in the process of germ line development can change over

evolutionary time across animals.

Keywords: Germ cells, pole cells, *Oncopeltus fasciatus*, *Gryllus bimaculatus*, Hemimetabola,
milkweed bug, cricket, Torso

Highlights

•   *germ cell-less* knockout reduces germ cell number in *Oncopeltus fasciatus*.
•   *germ cell-less* knockdown does not affect germ cell number in *Gryllus bimaculatus*.
•   The genetic mechanisms through which *gcl* affects germ cell number have diverged among insects.

Graphical abstract

Introduction

      One of the most crucial processes in animal development is the specification and maintenance of the germ line, the lineage of cells uniquely capable of giving rise to the next generation of animals (Extavour and Akam, 2003). Despite the importance of this process, mechanisms of germ line development vary widely even within closely related taxa, including across insects (Extavour and Akam, 2003). For example, in the fruit fly *Drosophila melanogaster*, germ cells first appear at the posterior pole of the embryo very early in embryogenesis, before cellularization occurs in the rest of the embryo (Huettner, 1923). The formation of germ cells in embryos of this fly depends on the presence of the germ plasm, a specialized region of the cytoplasm at the posterior pole containing maternally synthesized RNAs and proteins that collectively direct the formation and specification of germ cells (Illmensee and Mahowald, 1974). In contrast, germ cells in the cricket *Gryllus bimaculatus* are first detectable comparatively later in embryogenesis, after segmentation of the abdomen has begun (Ewen-Campen et al., 2013a). There is no evidence of germ plasm in cricket embryos, and germ cells are thought to be specified instead by inductive signals from neighboring cells, including BMP signaling (Donoughe et al., 2014; Ewen-Campen et al., 2013a). In yet another mechanism of insect germ cell specification, germ cells of the milkweed bug *Oncopeltus fasciatus* appear at a stage intermediate between those of *D. melanogaster* and *G. bimaculatus*, after blastoderm cellularization but before abdominal segmentation (Butt, 1949; Ewen-Campen et al., 2013b). There is also no evidence of germ plasm in embryos of this bug (Ewen-Campen et al., 2013b), but specific germ cell specification signals have not been identified.

      Despite the variation in modes of germ line development across species, many genes involved in this process are conserved. These include *vasa* and *piwi*, which are commonly used

as germ cell markers in many different animals (Ewen-Campen et al., 2010). *germ cell-less* (*gcl*)

is also sometimes used as a germ cell marker, but its role in germ line development is less well

understood (Jongens et al., 1992). Embryos laid by *D. melanogaster* females homozygous for a

loss of function mutation in *gcl* have significantly fewer germ cells than controls, if any (Jongens

et al., 1994, 1992). The gene is thought to have a role in suppressing somatic cell fate by

targeting the receptor tyrosine kinase Torso for degradation (Colonnetta et al., 2021; Pae et al.,

2017). *gcl* is also required for germ cell transcriptional quiescence in flies (Leatherman et al.,

2002). However, *gcl* is not required for germ cell specification in the mouse *Mus musculus* nor in

the flour beetle *Tribolium castaneum*, but it is required for proper mouse spermatogenesis

(Ansari et al., 2018; Kimura et al., 2003). Outside the germ line, *gcl* plays a role in establishing

the anterior–posterior axis in *T. castaneum* (Ansari et al., 2018). Despite the wide divergence in

their physiological roles, *gcl* homologs are highly conserved in amino acid sequence, with 45%

similarity between homologs from *D. melanogaster* and *M. musculus*. In addition, these

homologs appear to share at least some biochemical functions, since the mouse *gcl* homolog can

partially restore germ cell formation in a *gcl*-null *D. melanogaster* embryo, though not to wild

type numbers (Leatherman et al., 2000). *gcl* expression has also been observed in the embryonic

germ line in other arthropod species (Ewen-Campen et al., 2013a; Gupta and Extavour, 2013).

However, to our knowledge, functional tests of whether *gcl* is required for insect germ line

development or function remain limited to *D. melanogaster* and *T. castaneum*.

Here, we examine the function of *germ cell-less* in the embryonic and adult germ lines of

O. fasciatus and *G. bimaculatus*. Our data suggest that *gcl* is not required for germ cell

specification nor for adult gametogenesis in either species. However, *gcl* mutant *O. fasciatus*

embryos have significantly fewer germ cells than wild type. Knockdown of *Of-torso* does not

affect germ cell number, suggesting that *gcl* in this species affects germ cell number through a

different mechanism from that in *D. melanogaster*. Overall, these studies into *gcl* function across

species inform our understanding of the evolution of gene function in germ line development.

Materials and Methods

Gene identification

germ cell-less. *Of-germ cell-less* is annotated as OFAS015300 on Scaffold 2113 of the current *O.*

fasciatus genome assembly (Panfilio et al., 2019). *Gb-germ cell-less* is annotated as GBIM03474

on Scaffold 14 of the current *G. bimaculatus* genome assembly (Ylla et al., 2021). We amplified

both genes from mixed stage embryonic cDNA of the respective species to verify the full-length

coding sequences.

Of-torso. A tBLASTn search was performed on the *O. fasciatus* genome (Panfilio et al., 2019)

using *D. melanogaster* Torso as a query, and a reciprocal BLASTp was performed on the top

results with e-value < 1e–40. Of the initial results, OFAS005359 on Scaffold 336 returned *torso*

orthologs from other insects. A reciprocal BLASTp search using the adjacent upstream annotated

gene, OFAS005360, also returned *torso* orthologs from other insects. PCR was performed from

mixed stage embryonic cDNA to confirm that OFAS005359 and OFAS005360 correspond to the

same transcript. To obtain full-length sequence, RACE was performed using the SMARTer

RACE 5'/3' Kit (Takara Bio 634859) following manufacturer's instructions.

The initial tBLASTn search using *D. melanogaster* Torso as a query also returned

OFAS025175 as a result. A reciprocal BLASTp search using the predicted peptide sequence of

this hit returned *fgf receptor* orthologs from other insects. Since the FGF receptor is a family of

receptor tyrosine kinases closely related to Torso (Duncan et al., 2013), we wanted to include

OFAS025175 for phylogenetic analysis. Therefore, we performed RACE using the SMARTer

RACE 5'/3' Kit to obtain the full-length sequence of this gene. Full length sequences for both

Of-torso and *Of-fgfr* have been deposited at GenBank with accession numbers PV361543 and

PV361544, respectively.

Phylogenetic analysis was performed using MEGA 11 (Tamura et al., 2021). Insect

orthologs of Torso were aligned using MUSCLE (Edgar, 2004) along with orthologs of Ret

oncogene and FGF receptor, which are the closest related families of receptor tyrosine kinases

(Duncan et al., 2013). The multiple sequence alignment was used to generate a gene tree using

the Maximum Likelihood method with the JTT substitution model (Jones et al., 1992), a discrete

Gamma-distributed rate model and 1000 bootstrap replicates.

Molecular biology

<u>RNA extraction and cDNA synthesis</u>. Embryos or tissues of interest were homogenized in TRIzol

(Invitrogen 15596026), and RNA was extracted with phenol–chloroform extraction and

isopropanol precipitation. cDNA was synthesized with SuperScript III First-Strand Synthesis

SuperMix for qRT-PCR (Invitrogen 11752250) following manufacturer's instructions.

<u>Cloning and in vitro transcription</u>. Fragments of genes of interest were PCR-amplified from

embryonic cDNA and each cloned into the pGEM-T Easy vector (Promega A137A) using

manufacturer's instructions. A fragment of *gfp* was subcloned from an existing plasmid

containing the *gfp* sequence. Insertion was confirmed by Sanger sequencing. Primers used are

listed in Supplemental Table 1.

For *in situ* probe synthesis, gene fragments were PCR-amplified from plasmids using T7

and Sp6 primers. *In vitro* transcription was performed using T7 RNA polymerase (Promega

P2075) or Sp6 RNA polymerase (Roche 10810274001) with DIG RNA labeling mix (Roche

11277073910). 100 ng of template was used for a 20-µl reaction run at 37°C overnight. RNA

was purified with LiCl/ethanol precipitation, then resuspended in 20 µl DEPC-treated water.

Concentration was measured using a NanoDrop Spectrophotometer (ThermoFisher), and the

sample was diluted to 100 ng/µl with hybridization buffer (50% formamide, 1x phosphate-

buffered saline (PBS), 5x saline–sodium citrate (SSC), 0.1% Triton X-100, 0.1% CHAPS, 5 mM

EDTA, 50 µg/mL heparin, 1 µg/mL yeast tRNA). Probes were stored at −20°C.

For double-stranded RNA (dsRNA) synthesis, gene fragments were PCR-amplified from

plasmids using a forward T7 primer and a reverse primer containing both T7 and Sp6 promoter

sequences to generate a template for *in vitro* transcription with the T7 promoter sequence on both

ends. *In vitro* transcription was performed using the MEGAscript T7 Transcription Kit

(Invitrogen AM13345). 100 ng of template was used for an 80-µl reaction run at 37°C overnight.

RNA was purified with LiCl/ethanol precipitation, then resuspended in 100 µl DEPC-treated

water. To anneal the two strands, this solution was heated to 95°C on a heat block for 5 minutes,

then allowed to cool slowly by leaving the tube on the heat block with the power switched off. A

1:10 dilution of annealed RNA was analyzed using a NanoDrop Spectrophotometer

(ThermoFisher) to measure its concentration and purity, then run on a 1% agarose gel to confirm

its size and integrity. For embryonic RNAi, dsRNA was diluted in DEPC-treated water to 2 µg/µl

(*O. fasciatus*) or 5 µg/µl (*G. bimaculatus*), with 0.05% phenol red. For *G. bimaculatus* adult

RNAi, dsRNA was diluted to 7.5 µg/µl in DEPC-treated water.

<u>*CRISPR/Cas9-mediated mutagenesis*</u>. Alt-R S.p. Cas9 Nuclease V3 (cat. #1081059) and Alt-R

CRISPR-Cas9 tracrRNA (cat. #1072533) were ordered from IDT, along with gene-specific Alt-R

CRISPR-Cas9 crRNA. tracrRNA and crRNA were resuspended to 100 µM in Nuclease-Free

Duplex Buffer (IDT cat. #11-01-03-01). 5 µl of each RNA were combined in a PCR tube, heated

to 95°C for 5 minutes, and allowed to cool to room temperature to form 50 µM gRNA duplex. 3

µl gRNA and 2 µl Cas9 protein (10 µg/µl or ~62 µM) were combined in a PCR tube and

incubated at room temperature for 10 minutes to form Cas9/gRNA ribonucleoprotein complex

(RNP) at roughly 25 µM. Finally, RNP was diluted tenfold in 1x PBS with 0.05% phenol red.

Of-Vasa antibody generation

A rabbit polyclonal antibody was raised against a recombinant *Of*-Vasa protein fragment

(McGill Biology CIAN facility, Canada). The sequence encoding the 565 C-terminal amino acids

of *Of*-Vasa, the stop codon and the first 96 base pairs of 3' UTR (GenBank accession KC261571

(Ewen-Campen et al., 2013b)) was cloned into the pET151/dTOPO expression vector (Invitrogen

K151-01), thus introducing an N-terminal 6x-polyhistidine tag. Protein expression was induced

in *E. coli* BL21 (DE3) by addition of IPTG to a final concentration of 0.25 mM and shaking at

30°C for four hours. The overexpressed proteins were highly insoluble, so the cell pellets were

disrupted by sonication in non-denaturing buffer, and the soluble fraction was separated by

centrifugation and discarded. The pellet was resuspended in 8 M urea (crude prep). Crude preps

were further purified by SDS gel-purification. Protein bands were excised from the gel after

reverse staining with zinc chloride and the proteins were collected by electroelution. Acetone

precipitation was performed, and the precipitated protein was washed and resuspended in 8 M

urea. Protein concentration was adjusted to 1–2 mg/mL for injection. Rabbits were injected using

a standard 80-day protocol with three boosts after the initial injection; the final bleed was

performed after 87 days. The serum was processed by addition of sodium azide to 0.02% (w/v).

The antibody was affinity-purified against bacterially expressed *Of*-Vasa protein.

Embryo microinjection

Needles for injection were made by pulling glass capillaries (World Precision Instruments

1B100-4) with a Sutter P-97 Flaming/Brown micropipette puller. Needle tips were manually

opened with fine forceps just prior to injection. Eggs were injected with a Narishige IM300

microinjector.

O. fasciatus were allowed to lay eggs in dry cotton for four hours, after which eggs were

picked out of the cotton for injection. Two glass microscope slides were taped together with

double-sided tape to create a ledge with exposed double-sided tape against which embryos were

lined up. Eggs were then submerged in tap water for at least 20 minutes to soften the eggshell

before injection. Slides with injected eggs were kept at 28°C to continue development until the

desired age.

G. bimaculatus were allowed to lay eggs in wet cotton for four hours, after which eggs

were washed out of the cotton for injection. Eggs were aligned in grooves (570 μm wide × 2930

μm long × 650 μm deep) made in 2% agarose in Milli-Q water using a custom mold as

previously described (Donoughe et al., 2018). Eggs were injected while submerged in 1x PBS.

After injection, the solution was replaced with fresh 1x PBS. Injected eggs were kept submerged

in 1x PBS at 28°C overnight. The following day, dead embryos, if any, were removed, and the

remaining eggs were transferred onto damp Kimwipes in a covered Petri dish kept at 28°C to

continue development. Three days after injection, eggs were submerged in tap water again and

monitored every day until they reached the desired stage for dissection.

Embryo dissection and fixation

O. fasciatus eggs were stuck to the bottom of a 60-mm Petri dish with double-sided tape

and a slit was created in the eggshell with a microscalpel. 1x PBS with 4% paraformaldehyde

(PFA) was then poured into the dish to fix the embryos for 30 minutes at room temperature,

followed by at least three washes with PBT (1x PBS with 0.1% Triton X-100). Embryos were

then dissected free of their yolk in PBT with fine forceps. Embryos were washed at least three

times with PBT at room temperature before continuing with staining or dehydration in methanol.

G. bimaculatus embryos were dissected free of their yolk in 1x PBS with fine forceps and

fixed overnight in 1x PBS with 4% PFA at 4°C. The following morning, they were washed at

least three times with PBT at room temperature before continuing with staining or dehydration in

methanol.

G. bimaculatus adult RNA interference (RNAi)

2 μl dsRNA was injected into the abdomen of an adult cricket with a Hamilton syringe

through the base of a metathoracic leg. Gonads were dissected seven days after injection for

analysis.

Colorimetric in situ hybridization

Dissected embryos were stored in methanol at −20°C overnight or longer before *in situ* hybridization. After rehydration in PBT and washing in hybridization buffer, embryos were hybridized overnight in a 70°C water bath with 1 ng/µl probe in hybridization buffer. The following day, the following series of 20-minute washes were performed at 70°C: once in hybridization buffer, three times in wash 1 (2x SSC, 0.1% CHAPS) and three times in wash 2 (0.2x SSC, 0.1% CHAPS). Embryos were then washed three times for ten minutes each at room temperature in KTBT (1x Tris-buffered saline with KCl and 0.1% Triton X-100) followed by an hour-long incubation at room temperature in *in situ* blocking solution (1x PBS, 0.1% Tween-20, 5% NGS, 2 mg/mL bovine serum albumin, 1% dimethyl sulfoxide). Embryos were incubated for two hours at room temperature with sheep anti-digoxigenin–AP (Roche 11093274910), diluted 1:2000 in *in situ* blocking solution, then washed at least four times over two hours with KTBT and incubated in KTBT overnight at 4°C. The following day, embryos were washed with NTMT (100 mM Tris pH 9.5, 100 mM NaCl, 50 mM $MgCl_2$, 0.1% Tween-20), then developed in the dark at 37°C with NBT/BCIP in NTMT with 5% polyvinyl alcohol. Development was monitored at least every hour under a dissection microscope. To stop development of the enzymatic reaction, embryos were washed in PBT, incubated with 5 ng/µl Hoechst 33342 in PBT at room temperature for an hour, post-fixed in 4% PFA in PBS, then incubated in 70% glycerol before mounting in 100% glycerol.

Immunohistochemistry

Embryos were incubated in blocking solution (PBT with 5% normal goat serum (Jackson ImmunoResearch 005-000-121)) for one hour at room temperature, then incubated overnight at 4°C with primary antibodies diluted in blocking solution. The following day, embryos were washed for at least four times over four hours with PBT, incubated for an hour in blocking solution, then incubated overnight at 4°C with secondary antibodies diluted in blocking solution. Finally, embryos were washed in PBT, post-fixed in 4% PFA in PBS, then incubated in 70% glycerol before mounting in glycerol. Antibody staining was also performed with a no-primary control.

Primary antibodies used were: rabbit anti-*Gb*-Piwi (Ewen-Campen et al., 2013a) at a final concentration of 1:300; rabbit anti-*Of*-Vasa at 1:100; mouse anti-α-tubulin (DM1A; Sigma T9026) at 1:100; rabbit anti-β-catenin (Sigma C2206) at 1:500; and mouse anti-RNA polymerase II (H5; Covance MMS-129R) (Bregman et al., 1995; Warren et al., 1992) at 1:100. Secondary antibodies used were goat anti-rabbit conjugated to Alexa Fluor 647 (Invitrogen A-21246), goat anti-rabbit Alexa Fluor 488 (Invitrogen A-11008), goat anti-rabbit Alexa Fluor 568 (Invitrogen A-11011), donkey anti-mouse Alexa Fluor 488 (Invitrogen A-21202), and goat anti-mouse Alexa Fluor 647 (Invitrogen A-21236). All secondary antibodies were used at a final concentration of 1:400 along with Hoechst 33342 (Sigma B2261) at 5 ng/µl.

Embryos were imaged on a Nikon CSU-W1 spinning disk inverted confocal microscope with 10x air or 20x air objectives or a Zeiss AxioObserver inverted compound microscope with 10x air or 20x air objectives. Images were analyzed in Fiji.

Quantitative RT-PCR

Quantitative RT-PCR was performed on an AriaMx Real-time PCR system (Agilent)

using PerfeCTa SYBR Green SuperMix, Low ROX (Quantabio 95056). Fold-change differences

were calculated using the $\Delta\Delta C_t$ method for RNAi experiments. Expression levels were

normalized against *Of-gapdh* or *Gb-β-tubulin*. All primers used are listed in Supplemental Table

1.

Results.

germ cell-less expression across insect species.

During oogenesis in *D. melanogaster*, Gcl protein is detectable at the nuclear membrane

of nurse cells and oocytes (Moore et al., 2009). In addition, *gcl* transcripts are transcribed by

nurse cells, transported into the oocyte and deposited at the oocyte posterior, where they remain

as part of the germ plasm into early embryogenesis (Jongens et al., 1992). These transcripts are

translated into protein starting at the cleavage stages of embryogenesis (Jongens et al., 1992).

Gcl protein is localized to the nuclear membranes of germ cells after they form and remains

detectable in germ cells until mid-gastrulation, when germ cells enter the primordial midgut

(Jongens et al., 1992). We asked whether *gcl* is also expressed in other insect embryos at the time

and place of germ cell specification.

We assessed *gcl* transcript levels using quantitative RT-PCR on *O. fasciatus* embryos

collected in 12-hour windows spanning the first four days of embryogenesis. *Of-gcl* transcripts

were detected at the highest levels 24–36 hours after egg-lay at 28°C (Figure 1A). This window

corresponds to the late blastoderm stage, when germ cells are first distinguishable (Ewen-

Campen et al., 2013b), and the beginning of germ band elongation. We speculate that this peak in

expression is likely due to zygotic rather than maternal expression, since it is higher than

expression in freshly laid eggs. However, we were unable to detect *Of-gcl* transcripts by *in situ*

hybridization above background levels in embryos at any stage of development, nor in adult

ovaries or testes (Supplemental Figure 1A–B).

We similarly assessed *gcl* transcript levels using qRT-PCR on *G. bimaculatus* embryos

collected in 8-hour windows spanning the first two days of embryogenesis and 24-hour windows

spanning the following eight days of embryogenesis. *Gb-gcl* transcripts were detectable at the

highest levels in the first eight hours after egg-lay (Figure 1B), suggesting maternal deposition of

Gb-gcl transcript. Consistent with this hypothesis, *Gb-gcl* transcript was detectable by *in situ*

hybridization in ovaries but not during embryogenesis (Supplemental Figure 1C–D).

Although we were unable to detect localized *gcl* expression in either *O. fasciatus* or *G.*

bimaculatus embryos, we still sought to determine whether the gene's documented roles in germ

cell specification (*D. melanogaster*) or adult spermatogenesis (*M. musculus*) might be conserved

(Jongens et al., 1992; Kimura et al., 2003). To this end, we analyzed *gcl* loss of function

conditions in embryos and adult gonads of both species. *gcl* is also required for transcriptional

quiescence in germ cells in *D. melanogaster* (Leatherman et al., 2002). However, we could not

detect any reduction in transcriptional activity in germ cells relative to somatic cells of either *O.*

fasciatus or *G. bimaculatus* (Supplemental Figure 2), so we did not assess this phenotype in our

functional studies of *gcl*.

Of-gcl mutation decreases germ cell number in milkweed bug embryos.

To determine whether *gcl* is required for germ cell specification in *O. fasciatus*, we generated an *Of-gcl* putative protein-null allele with CRISPR/Cas9-mediated mutagenesis (Figure 2A–B). This allele contains a 2-bp insertion at amino acid residue 14. The resulting frameshift leads to a premature stop codon, resulting in a predicted peptide of 30 amino acid residues, compared with the 478-residue wild type protein. Surprisingly, individuals homozygous for this mutant allele were viable and fertile, and we have maintained a stable mutant line for at least twelve generations at the time of writing. There were no detectable morphological defects in adult ovaries or testes from this mutant line, nor any evidence of abnormal gametogenesis in either males or females (Supplemental Figures 3–4).

In *O. fasciatus*, germ cells are first detectable in stage 3 on the interior of the cellular blastoderm, forming part of an invaginating structure called the posterior pit (Ewen-Campen et al., 2013b). The germ cells remain at the posterior end of the germ band as it forms by invaginating into the egg (Ewen-Campen et al., 2013b) (Figure 2B; Supplemental Figure 5). At stage 3, *gcl* mutant embryos had on average 35% fewer germ cells than wild type embryos ($p <$ 0.001; $n =$ 14 wild type, 8 *gcl* embryos; Figure 2C–D; Supplemental Table 2). Germ cell number remained significantly lower in *gcl* mutant embryos relative to wild type embryos throughout all embryonic stages examined (Figure 2D). We detected a marginal decrease in germ cell proliferation in *gcl* mutant embryos relative to wild type only at stages 6 and 9, as measured by the proportion of germ cells with detectable phospho-histone 3 (Figure 2E; Supplemental Table 2). However, there was no detectable difference in germ cell proliferation between the two genotypes at any other embryonic stage examined (Figure 2E; Supplemental Table 2). We were

also unable to detect cleaved caspase 3-dependent cell death in stages 3–4 of either genotype ($n =$

27 wild type, 23 *gcl* embryos). Given that the discrepancy in germ cell number did not appear to

be the result of differences in cell proliferation or cell death, these data suggest that *Of-gcl* is

involved in, but not strictly necessary for, one or more of germ cell specification, formation or

maintenance.

In *D. melanogaster*, Gcl promotes germ cell fate by targeting the receptor Torso for

degradation, and *torso* RNAi in *gcl* mutant flies restores germ cell numbers to wild type levels

(Pae et al., 2017). To determine whether this mechanism of germ cell number determination is

conserved in *O. fasciatus*, we asked whether *torso* loss of function also affects germ cell number

in this species. Despite previous reports that the *O. fasciatus* genome lacks a *torso* ortholog

(Skelly et al., 2019; Weisbrod et al., 2013), we identified a high-confidence homolog in the

published genome assembly (Panfilio et al., 2019) (Figure 3A). Knockdown of *Of-torso* by

embryonic RNAi did not affect germ cell number in stages 3–4 in either wild type or *gcl* mutant

backgrounds (Figure 3B–C). This result suggests that the mechanism through which *gcl* controls

germ cell number is not conserved between *O. fasciatus* and *D. melanogaster*.

Gb-gcl knockdown does not affect germ cell number in cricket embryos.

To determine whether *gcl* is required for germ cell specification in *G. bimaculatus*, we

knocked down *gcl* expression using embryonic RNAi. We found no significant difference in total

germ cell number in stage 8 embryos under *gcl* knockdown relative to *gfp* control, nor did we

detect any significant difference in germ cell number in any segment (Figure 4). qPCR analysis

of injected embryos confirmed that RNAi was effective at reducing *gcl* transcript levels to 34%

relative to controls. It is possible that knockdown efficiency is not high enough with RNAi to

affect germ cell specification. However, we note that we have previously detected significant

changes in germ cell number with comparable or lower knockdown efficiencies (Barnett et al.,

2019; Donoughe et al., 2014; Nakamura and Extavour, 2016). Therefore, we speculate that the

role of *Gb-gcl* in germ cell specification or maintenance, if any, must be subtle.

Next, we asked whether *gcl* has any role in adult gametogenesis in *G. bimaculatus* by

performing RNAi in adult crickets. We could not detect any morphological or gametogenesis

defects in either ovaries or testes (Supplemental Figures 1–2). These data suggest that *gcl* is not

required for adult gametogenesis in this cricket.

Discussion.

Our analysis of *gcl* loss of function in embryos of two different hemimetabolous insects

suggests that *gcl* is not required for germ cell specification in these species, in contrast to the

gene's importance in germ cell formation in *D. melanogaster*. However, *Of-gcl* knockout

embryos have significantly fewer germ cells than wild type embryos, suggesting that the gene

may be involved in germ cell specification or maintenance in this bug, even if it is not strictly

required.

Gcl protein suppresses the somatic cell fate in *D. melanogaster* by targeting the receptor

tyrosine kinase Torso for degradation (Colonnetta et al., 2021; Pae et al., 2017). Germ cell

number can thus be modulated in flies by up- or downregulating Torso activity (Pae et al., 2017).

However, RNAi knockdown of *torso* in *O. fasciatus* does not affect germ cell number in either

wild type or *gcl* mutant embryos (Figure 3). Therefore, we hypothesize that germ cell

specification in this species does not require downregulation of Torso by Gcl, and that *Of-gcl*

regulates germ cell specification through some other unknown pathway. Torso's role in germ cell

specification may have been co-opted from its ancestral functions in neuroendocrine signaling,

just as it is thought to have been co-opted into its role in terminal patterning in flies (Duncan et

al., 2013; Schoppmeier and Schröder, 2005).

In *D. melanogaster*, the interaction between and Gcl and Torso proteins depends on

sequence motifs present on both proteins (Pae et al., 2017). In Gcl protein, this motif is the GCL

domain, which is conserved across insect homologs (Supplemental Figure 6) (Pae et al., 2017).

Torso interacts with Gcl through a motif in the linker domain between its split kinase domains.

This linker region is significantly longer in *Dm*-Torso compared with other insect homologs

(Supplemental Figure 7), and the specific motif identified by Pae and colleagues (Pae et al.,

2017) is not present in other homologs. The conservation of the GCL domain despite the lack of

conservation in the Torso interaction motif may suggest that Gcl has some other unknown

interaction partner, or that the Gcl–Torso interaction may not be as sequence-specific as

originally thought. AlphaFold3 (Abramson et al., 2024) predicts potential interaction between

Gcl and Torso homologs in both *O. fasciatus* and *G. bimaculatus* through the split kinase linker

domain of Torso (Supplemental Figure 8). Further experiments will be needed to test whether

these proteins interact *in vivo*, and to identify additional conserved interaction partners of Gcl.

In mice, *gcl* knockout results in abnormal nuclear morphology in the liver, pancreas and

testis, though the only functional defect identified was impaired spermatogenesis (Kimura et al.,

2003). Similarly, *gcl* is expressed at low levels in somatic tissues in flies including the gut,

muscle and nervous system, but no defects have been reported in those tissues in *gcl*-null flies

(Jongens et al., 1992; Robertson et al., 1999). We were also unable to detect any defects in

spermatogenesis under either *Of-gcl* knockout or *Gb-gcl* knockdown conditions (Supplementary

Figure 1). Given that the mouse *gcl* homolog can rescue germ cell formation in *gcl*-null flies

(Leatherman et al., 2000), its biochemical functions are likely to be conserved across these

species. We speculate that *gcl* has a ubiquitous but subtle cell biological role in maintaining

nuclear morphology that only yields a phenotype in certain tissue contexts, such as in mouse

spermatogenesis. Under this hypothesis, *gcl* then gained an additional role in germ cell

specification in certain lineages of insects. In addition, the mechanism underlying this germ cell

specification has diverged between insect lineages, perhaps through the evolution of new

interaction partners such as Torso. This model reflects the complexity and variation present in the

mechanisms of germ cell specification and their evolution, including the co-option of new genes

with the shifting mechanisms of this process across animals.

Author contributions

JAK: Conceptualization, Methodology, Formal Analysis, Investigation, Writing – Original Draft,

Writing – Review & Editing, Visualization; CGE: Conceptualization, Writing – Review &

Editing, Supervision, Funding Acquisition

Acknowledgments

The authors thank members of the Extavour lab for valuable discussions. This research was

supported by National Science Foundation (NSF) Awards IOS-2220747 and IOS-0817678 to

CGE, and an NSF Graduate Research Fellowship to BE-C.

Figure 1. *gcl* expression is detectable by qPCR in *O. fasciatus* (A) and *G. bimaculatus* (B) embryos. C_q values were normalized against *Of-gapdh* or *Gb-tubulin* expression and averaged over four biological replicates for each time point. Error bars represent standard deviations. The width of each bar represents the length of time for each embryo collection.

A

Of-Gcl | 73 | 173 181 | 259 | 348 387 | 478
BTB | BACK | GCL

mutant ——— 30

B

allele	DNA sequence	translation
wild type	...CTGCCC<u>AG--A GCGTGTTGCTGCAACTCTC</u>...	...YCPERVAATL...
mutant	**...CTGCCCAG****CC****A GCGTGTTGCTGCAACTCTC...**	**...YCP****ASVLLQL****...**

C

Of-Vasa

wild type | gcl

D — total germ cells vs stage

stage	3	4	5	6	7	8	9	10	11	12	13	14
significance	***	***	***	***	***	***	**	***	***	*	***	***
n =	14 8	22 20	54 36	52 20	59 26	28 20	65 14	21 13	31 25	26 17	39 35	33 30

E — PH3+ germ cells vs stage

stage	3	4	5	6	7	8	9	10	11	12	13	14
significance				*			*					
n =	14 18	22 20	49 36	47 20	46 26	22 20	43 14	17 13	30 25	23 17	26 35	33 30

● wild type ● *gcl* mutant

Figure 2 (previous page). *Of-gcl* mutation decreases germ cell number in embryos.

(**A**) Predicted domain architecture of *Of*-Gcl and predicted mutant protein. Wild type *Of*-Gcl has
two predicted folded domains (BTB and BACK domains) as well as a conserved motif called the
GCL domain. The predicted mutant protein contains none of these domains and is truncated after
residue 30.

(**B**) *Of-gcl* mutant allele contains a two base pair insertion, creating a frameshift and leading to a
predicted truncated protein. Guide sequence is underlined; arrowhead marks predicted Cas9 cut
site. Inserted bases and predicted mutated residues are indicated in magenta.

(**C**) Representative antibody stains against *Of*-Vasa to mark germ cells in stage 3 wild type and
gcl mutant embryos. Mutant embryos still have germ cells at the expected location, indicated by
arrows. Scale bar = 100 µm.

(**D**) Germ cell counts across embryogenesis in wild type and *gcl* mutant embryos. Mutant
embryos have significantly fewer germ cells per embryo compared to wild type.

(**E**) Proportion of PH3$^+$ germ cells across embryogenesis in wild type and *gcl* mutant embryos.
There is no significant difference in the proportion of proliferating germ cells between wild type
and *gcl* mutant embryos.

All statistical tests are non-parametric bootstrap comparisons. Embryonic stages are defined in
Supplemental Figure 5 and Supplemental Table 3. * $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$

Figure 3 (previous page). *Of-torso* knockdown does not affect embryonic germ cell number.

(**A**) Phylogenetic analysis of *torso* and related receptor tyrosine kinases (*Ret oncogene* and *FGF*
receptor) across insects. Each clade is individually highlighted for clarity. The putative *torso*
homolog from *O. fasciatus* (OFAS005359/OFAS0005360) falls within the *torso* clade.

(**B**) Germ cell counts in stage 3–4 *Of-torso* RNAi embryos, performed in wild type and *gcl*
mutant backgrounds. *torso* knockdown does not affect germ cell number in either background.

(**C**) Proportion of PH3$^+$ germ cells in stage 3–4 *Of-torso* RNAi embryos, performed in wild type
and *gcl* mutant backgrounds. *torso* knockdown marginally reduces germ cell proliferation in *gcl*
mutant embryos but not in wild type embryos.

All statistical tests are non-parametric bootstrap comparisons.
* $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$

A

Gb-Gcl ———— 75 ———— 175 183 ———— 261 ————— 349 388 ———— 498
 BTB   BACK   GCL

dsRNA

B

| control | *Gb-gcl* RNAi |

Gb-Piwi

C

n.s.

germ cells per embryo

ctrl   *gcl*

n =   46   68

D

n.s.   *n.s.*   *n.s.*

germ cells per hemisegment

A2   A3   A4

n =   92   136   92   136   92   136

Figure 4 (previous page). ***Gb-gcl* knockdown does not affect embryonic germ cell number.**

(**A**) Predicted domain architecture of *Gb*-Gcl. *Gb*-Gcl has two predicted folded domains (BTB
and BACK domains) and a conserved motif called the GCL domain. The fragment region used
for dsRNA synthesis is marked.

(**B**) Representative antibody stains against *Gb*-Piwi to mark germ cells in stage 8 control and *gcl*
knockdown embryos. Embryos in which *Gb-gcl* has been knocked down still have germ cells in
the expected positions. Scale bar = 100 μm.

(**C**) Germ cell counts in stage 8 *Gb-gcl* RNAi embryos. Germ cell numbers are not significantly
different in *Gb-gcl* knockdown embryos relative to control. Each point represents one embryo.

(**D**) Germ cell counts in stage 8 *Gb-gcl* RNAi embryos, divided into hemisegments (*i.e.*, the left
or right half of one segment). *Gb-gcl* knockdown does not affect the number of germ cells in any
individual segment. Each point represents a hemisegment.

All statistical tests are non-parametric bootstrap comparisons.

Supplemental Figure 1. *gcl* is not detectable with *in situ* hybridization during embryogenesis in *O. fasciatus* or *G. bimaculatus*.

(**A–B**) *In situ* hybridization against *Of-gcl* in embryos (C) and ovaries (D). Numbers indicate embryonic stages. Scale bars = 500 μm (embryos) or 200 μm (ovaries).

(**C–D**) *In situ* hybridization against *Gb-gcl* in embryos (E) and ovaries (F). Numbers indicate embryonic stages. Scale bars = 500 μm (embryos) or 200 μm (ovaries).

Supplemental Figure 2. Germ cells in *O. fasciatus* and *G. bimaculatus* are not transcriptionally quiescent.

(**A**) Antibody stains in stage 3 *O. fasciatus* embryos against *Of*-Vasa to visualize germ cells and against hyperphosphorylated RNA polymerase II to visualize transcriptional activity. Germ cells are transcriptionally active. Scale bar = 25 µm.

(**B**) Antibody stains in stage 7 *G. bimaculatus* embryos against *Gb*-Piwi to visualize germ cells and against hyperphosphorylated RNA polymerase II to visualize transcriptional activity. Germ cells are transcriptionally active. Scale bar = 25 µm.

Supplemental Figure 3. Lack of *gcl* activity does not affect the progression of adult spermatogenesis in either *O. fasciatus* or *G. bimaculatus*. Representative images of Hoechst-stained testes showing different stages of spermatogenesis in both species under both wild type and *gcl* loss of function conditions. Scale bar = 50 μm.

Supplemental Figure 4. Lack of *gcl* activity does not affect adult oogenesis in either *O. fasciatus* or *G. bimaculatus*.
Representative images of single 1 μm optical sections of ovarioles under both wild type ((A) *O. fasciatus*; (C) *G. bimaculatus*) and (B)
gcl knockout (*O. fasciatus*) or (D) knockdown (*G. bimaculatus*) conditions. Scale bars = 100 μm in (A) (applies also to (B)) and (C)
(applies also to (D)).

Supplemental Figure 5. Wild type germ line development in *O. fasciatus*.

(**A**) Schematics showing the position of the *O. fasciatus* embryo within the egg, roughly
corresponding to the stages shown below in (**B**). Germ cells are indicated in magenta.

(**B**) Representative images of *Of*-Vasa antibody stains in stage 3–11 *O. fasciatus* embryos. Top
row shows whole mount embryos, while the bottom row shows a higher magnification view of
the germ cells. Scale bar = 500 μm in top row, 100 μm in bottom row.

(**C**) *Of*-Vasa antibody stains in stages 10–14 showing coalescence of germ cells into the
embryonic gonad. Right gonad shown in all images. Scale bar = 50 μm.

(**D**) Timeline of embryonic stages for wild type development at 28°C, 60% humidity.

Supplemental Figure 6. MUSCLE alignment of select *gcl* homologs. Darker shading represents greater degree of conservation at each position. The conserved GCL domain identified by Pae and colleagues (Pae et al., 2017) is marked in magenta. In that study, the role of the GCL domain was directly tested by mutating the residue marked by the arrowhead (R377).

511
512
513
514

Supplemental Figure 7. MUSCLE alignment of the intracellular domain of select insect *torso* **homologs.** Darker shading represents greater degree of conservation at each position. The Gcl-interacting motif identified by Pae and colleagues (Pae et al., 2017) is marked in magenta. Note that this motif is not conserved in other insects; the surrounding domain is much longer in *D. melanogaster* compared with other homologs.

Supplemental Figure 8. AlphaFold3 predicts potential interactions between Gcl and Torso homologs. Predicted alignment error plots are shown for AlphaFold3 structure predictions of Gcl and Torso homologs from *D. melanogaster* (**A**), *O. fasciatus* (**B**) and *G. bimaculatus* (**C**). Darker color represents higher confidence (lower predicted error) in the relative positions of the scored and aligned residues. Overall confidence in the structure of the Gcl–Torso complex is low for all species. However, in all species, there is a region of Torso protein (marked in magenta) for which its predicted alignment error with Gcl is lower, suggesting a potential interaction between the two proteins through this region of Torso.

Supplemental Table 1. Primers used.

gene	direction	primer sequence	expected length
full-length sequence			
Of-gcl	forward	ATGGGTAAAGTATTCAGTGCAGTTGCT	1437 bp
	reverse	TTAGTCTTCTTTTAATTCTCTCTCTGTAAATAG	
Gb-gcl	forward	CGTCGATGAGCTAGTTGAGAGCTGA	1804 bp
	reverse	AGATGTTCAAGCATAGCCGCCCAAG	
in situ probe and dsRNA			
Of-gcl	forward	GCAGTTGCTTATTACTGCCCAGAGCGT	441 bp
	reverse	TGCAGCAGCCAGCACACCAACAACA	
Of-torso	forward	TCGAGGTCGCCGTCAAATCCCT	665 bp
	reverse	GGCAACAGACGCTGTCGGAACA	
Gb-gcl	forward	TGTCCGCTCGTGGGTCGGCAGAAAA	413 bp
	reverse	ACAGAGAGGCTGCAGCAAGCACTGA	
qPCR			
Of-gcl	forward	TCCAGAAGGTTGGTTGCTCGATGCCT	257 bp
	reverse	TGCTGCTGTCAAACTGTGTACGCTGCC	
Of-torso	forward	TGAGCATGGTGGGCGATTGCTG	288 bp
	reverse	AGGGATTTGACGGCGACCTCGA	
Gb-gcl	forward	GCACCACCATTTCGTGCACTTCGCC	258 bp
	reverse	ACCTGTCCATCGCCACACATGTTCT	
Of-gapdh	forward	TGGCAAACTAACTGGCATGGCA	171 bp
	reverse	ACAACCTGGTCATCAGTGTACGCCA	
Gb-β-tubulin	forward	TGGACTCCGTCCGGTCAGGC	166 bp
	reverse	TCGCAGCTCTCGGCCTCCTT	
RACE			
Of-torso	forward 1 (3' RACE)	ACTGGAGGAGGCTCCCAACTCGAGGTCGCC	
	forward 2 (3' RACE)	ACAGGAGGGCCACGGCAGAGGACTTGGCTT	
	reverse 1 (5' RACE)	TGGAGAAAGGCGGCAACAGACGCTGTCGGA	
	reverse 2 (5' RACE)	CGCTGGGAGGTAGAGAAGCCGACCAGGGAG	
Of-fgfr	forward 1 (3' RACE)	AACAAGGGCATACAGATGCAGAAATGACTG	
	forward 2 (3' RACE)	GCATCGGCCATCATCTGGTTATGAACCTGC	
	reverse 1 (5' RACE)	TATTCCTTGCTGCCAGGTCTCGATGTAAAC	
	reverse 2 (5' RACE)	TGACAGGTATTCCATGCCTCTAGCAACTTG	

Supplemental Table 2. Quantification of germ cell development in wild type and *gcl* mutant *O. fasciatus* embryos. All *p*-values were calculated using a non-parametric bootstrap test.

| stage | total germ cells per embryo | | | | | | proportion of PH3+ germ cells per embryo | | | | |
| | wild type | | *gcl* | | | | wild type | | *gcl* | | |
	mean	*n*	mean	*n*	ratio	*p*-value	mean	*n*	mean	*n*	*p*-value
3	102.00	14	68.13	8	0.668	<0.001	0.0126	14	0.0110	8	0.4221
4	108.00	22	56.55	20	0.524	<0.001	0.0181	22	0.0124	20	0.0953
5	115.69	54	77.58	36	0.671	<0.001	0.0221	49	0.0279	36	0.3386
6	124.71	52	65.70	20	0.527	<0.001	0.0250	47	0.0236	20	0.0382
7	133.54	59	86.19	26	0.645	<0.001	0.0137	46	0.0143	26	0.2572
8	118.57	28	75.70	20	0.638	<0.001	0.0108	22	0.0132	20	0.0745
9	122.19	65	96.36	14	0.789	0.002	0.0081	43	0.0067	14	0.0164
10	130.24	21	79.77	13	0.612	<0.001	0.0040	17	0.0058	13	0.4213
11	113.36	31	80.52	25	0.710	0.001	0.0040	30	0.0045	25	0.4566
12	91.04	26	70.29	17	0.772	0.017	0.0030	23	0.0042	17	0.2247
13	89.95	39	68.63	35	0.763	<0.001	0.0014	26	0.0012	35	0.1769
14	101.67	33	69.03	30	0.679	<0.001	0.0015	33	0.0005	30	0.0605

Supplemental Table 3. Staging table for wild type *O. fasciatus* embryogenesis at 28°C.

stage	hours ael (28°C)	stage description	germ cell development
1		Freshly laid egg.	
2		Uniform blastoderm.	
3	26–28	An indentation appears at the posterior end of the egg, called the posterior pit.	Primordial germ cells are first observed just inside the posterior pit, but we cannot currently perform antibody stains at this stage.
4	28–32	The germ band has invaginated into the posterior pit and extends toward the anterior. The orientation of the germ band is reversed with respect to the egg, *i.e.*, the posterior end of the germ band is closer to the anterior of the egg.	Primordial germ cells remain in a cluster at the posterior tip of the germ band. The Vasa antibody stain is inconsistent at this stage.
5	32–36	The thorax region is wider than the abdomen region (mediolaterally), but limb buds are not yet distinct. Segments have been specified up to A5–A7 but are not yet distinguishable without staining for a segmentation marker.	Germ cells are located in the bulbous part of the germ band, primarily in A5–A6, but with some in A4 and A7.
6	36–44	Limb buds are now distinct semicircles. Segments have been specified up to A7–A9 but are not yet distinguishable without staining for a segmentation marker.	Germ cells are located in a single cluster primarily in A4–A6, with some in A7. Segments in which germ cells are located are narrower in the medial–lateral axis compared with the posterior tip.
7	42–50	Segments start to become visibly distinct in the anterior abdomen around the time A9 is specified. Posterior tip is still round. Neural groove has formed at most up to A3.	Germ cells separate into segmental clusters. When there are many germ cells, they may still appear as one large medial cluster, especially as one spanning both segments A4 and A5.
8	48–54	Legs and gnathal appendages are triangular rather than semicircular lobes. A11 starts to thicken but the posterior tip is still mostly round.	Germ cells split into bilateral clusters as the neural groove forms in each segment. In this stage, germ cells are located right at the edges of the neural groove. When there are many germ cells in a particular segment, they may still appear to be one large medial cluster in that segment.
9	52–64	Neural groove has formed up to A8–A9. "Coelomic cavities" are visible as pits in all abdominal segments up to A9, though they may sometimes be obscured by germ cells, especially in A4–A6. Posterior end has folded over to form the proctodeum. Lateral	Germ cells have moved laterally away from the neural groove and clusters usually appear rounder.

		margins of the germ band are thickening in the gnathal and thoracic segments.	
10	60–64	Lateral margins of the germ band are thickening in the abdomen. Ganglia appear as two parallel rectangles in each segment, separated by the median cord along the midline. "Coelomic cavities" are no longer distinct.	Germ cell clusters lengthen along the AP axis and narrow along the medial–lateral axis. Adjacent clusters may fuse, but a segmental grouping is still visible.
11	64–68	Median cord clusters appear at segmental boundaries, and the ganglionic masses are now slightly crooked.	Germ cells are lined up in the genital ridges on each side of the embryo, parallel to the AP axis.
12	68–72	Abdominal cross commissures have formed. With a nuclear stain, ganglia appear as pairs of half circles facing away from each other.	Germ cells remain in the genital ridges.
13	72–76	The lateral cords move dorsolaterally and become parallel with the median cord again. Gonadal follicle primordia become visible in the nuclear channel.	Germ cells remain in the genital ridges.
14	76–	The embryo undergoes katatrepsis and begins dorsal closure. This stage will likely be subdivided further.	Germ cells start to migrate laterally into gonadal follicle primordia.

References

Abramson, J., Adler, J., Dunger, J., Evans, R., Green, T., Pritzel, A., Ronneberger, O., Willmore, L., Ballard, A.J., Bambrick, J., Bodenstein, S.W., Evans, D.A., Hung, C.-C., O'Neill, M., Reiman, D., Tunyasuvunakool, K., Wu, Z., Žemgulytė, A., Arvaniti, E., Beattie, C., Bertolli, O., Bridgland, A., Cherepanov, A., Congreve, M., Cowen-Rivers, A.I., Cowie, A., Figurnov, M., Fuchs, F.B., Gladman, H., Jain, R., Khan, Y.A., Low, C.M.R., Perlin, K., Potapenko, A., Savy, P., Singh, S., Stecula, A., Thillaisundaram, A., Tong, C., Yakneen, S., Zhong, E.D., Zielinski, M., Bapst, V., Kohli, P., Jaderberg, M., Hassabis, D., Jumper, J.M., 2024. Accurate structure prediction of biomolecular interactions with AlphaFold 3. Nature 630, 493–500. https://doi.org/10.1038/s41586-024-07487-w

Ansari, S., Troelenberg, N., Dao, V.A., Richter, T., Bucher, G., Klingler, M., 2018. Double abdomen in a short-germ insect: Zygotic control of axis formation revealed in the beetle,Tribolium castaneum. Proc. Natl. Acad. Sci. 115, 1819–1824. https://doi.org/10.1073/pnas.1716512115

Barnett, A.A., Nakamura, T., Extavour, C.G., 2019. Hox genes limit germ cell formation in the short germ insect,Gryllus bimaculatus. Proc. Natl. Acad. Sci. 116, 16430–16435. https://doi.org/10.1073/pnas.1816024116

Bregman, D.B., Du, L., Zee, S. van der, Warren, S.L., 1995. Transcription-dependent redistribution of the large subunit of RNA polymerase II to discrete nuclear domains. J. cell Biol. 129, 287–298. https://doi.org/10.1083/jcb.129.2.287

Butt, F.H., 1949. Embryology of the Milkweed Bug, Oncopeltus fasciatus (Hemiptera). Cornell Experiment Station Memoir 3–43.

Colonnetta, M.M., Lym, L.R., Wilkins, L., Kappes, G., Castro, E.A., Ryder, P.V., Schedl, P., Lerit, D.A., Deshpande, G., 2021. Antagonism between,germ cell-less,and Torso receptor regulates transcriptional quiescence underlying germline/soma distinction. eLife 10, e54346. https://doi.org/10.7554/elife.54346

Donoughe, S., Kim, C., Extavour, C.G., 2018. High-throughput live-imaging of embryos in microwell arrays using a modular specimen mounting system. Biol. Open 7, bio031260. https://doi.org/10.1242/bio.031260

Donoughe, S., Nakamura, T., Ewen-Campen, B., Green, D.A., Henderson, L., Extavour, C.G., 2014. BMP signaling is required for the generation of primordial germ cells in an insect. Proc. Natl. Acad. Sci. 111, 4133–4138. https://doi.org/10.1073/pnas.1400525111

Duncan, E.J., Benton, M.A., Dearden, P.K., 2013. Canonical terminal patterning is an evolutionary novelty. Dev. Biol. 377, 245–261. https://doi.org/10.1016/j.ydbio.2013.02.010

Edgar, R.C., 2004. MUSCLE: multiple sequence alignment with high accuracy and high throughput. Nucleic Acids Res. 32, 1792–1797. https://doi.org/10.1093/nar/gkh340

Ewen-Campen, B., Donoughe, S., Clarke, D.N., Extavour, C.G., 2013a. Germ cell specification requires zygotic mechanisms rather than germ plasm in a basally branching insect. Curr. Biol. 23, 835–842. https://doi.org/10.1016/j.cub.2013.03.063

Ewen-Campen, B., Jones, T.E.M., Extavour, C.G., 2013b. Evidence against a germ plasm in the milkweed bug,Oncopeltus fasciatus, a hemimetabolous insect. Biol. Open 2, 556–568. https://doi.org/10.1242/bio.20134390

Ewen-Campen, B., Schwager, E.E., Extavour, C.G., 2010. The molecular machinery of germ line specification. Mol. Reprod. Dev. 77, 3–18. https://doi.org/10.1002/mrd.21091

Extavour, C.G., Akam, M., 2003. Mechanisms of germ cell specification across the metazoans: epigenesis and preformation. Development 130, 5869–5884. https://doi.org/10.1242/dev.00804

Gupta, T., Extavour, C.G., 2013. Identification of a putative germ plasm in the amphipod,Parhyale hawaiensis. EvoDevo 4, 34. https://doi.org/10.1186/2041-9139-4-34

Huettner, A.F., 1923. The origin of the germ cells in,Drosophila melanogaster. J. Morphol. 37, 385–423. https://doi.org/10.1002/jmor.1050370204

Illmensee, K., Mahowald, A.P., 1974. Transplantation of posterior polar plasm in,Drosophila. Induction of germ cells at the anterior pole of the egg. Proc. Natl. Acad. Sci. 71, 1016–1020. https://doi.org/10.1073/pnas.71.4.1016

Jones, D.T., Taylor, W.R., Thornton, J.M., 1992. The rapid generation of mutation data matrices from protein sequences. Bioinformatics 8, 275–282. https://doi.org/10.1093/bioinformatics/8.3.275

Jongens, T.A., Ackerman, L.D., Swedlow, J.R., Jan, L.Y., Jan, Y.N., 1994. Germ cell-less encodes a cell type-specific nuclear pore-associated protein and functions early in the germ-cell specification pathway of Drosophila. Genes Dev. 8, 2123–2136. https://doi.org/10.1101/gad.8.18.2123

Jongens, T.A., Hay, B., Jan, L.Y., Jan, Y.N., 1992. The germ cell-less,gene product: A posteriorly localized component necessary for germ cell development in,Drosophila. Cell 70, 569–584. https://doi.org/10.1016/0092-8674(92)90427-e

Kimura, T., Ito, C., Watanabe, S., Takahashi, T., Ikawa, M., Yomogida, K., Fujita, Y., Ikeuchi, M., Asada, N., Matsumiya, K., Okuyama, A., Okabe, M., Toshimori, K., Nakano, T., 2003. Mouse Germ cell-less as an essential component for nuclear integrity. Mol. Cell. Biol. 23, 1304–1315. https://doi.org/10.1128/mcb.23.4.1304-1315.2003

Leatherman, J.L., Kaestner, K.H., Jongens, T.A., 2000. Identification of a mouse,germ cell-less,homologue with conserved activity in,Drosophila. Mech. Dev. 92, 145–153. https://doi.org/10.1016/s0925-4773(99)00335-4

Leatherman, J.L., Levin, L., Boero, J., Jongens, T.A., 2002. germ cell-less,acts to repress transcription during the establishment of the Drosophila germ cell lineage. Curr. Biol. 12, 1681–1685. https://doi.org/10.1016/s0960-9822(02)01182-x

Moore, J., Han, H., Lasko, P., 2009. Bruno negatively regulates,germ cell-less,expression in a BRE-independent manner. Mech. Dev. 126, 503–516. https://doi.org/10.1016/j.mod.2009.04.002

Nakamura, T., Extavour, C.G., 2016. The transcriptional repressor Blimp-1 acts downstream of BMP signaling to generate primordial germ cells in the cricket,Gryllus bimaculatus. Development 143, 255–263. https://doi.org/10.1242/dev.127563

Pae, J., Cinalli, R.M., Marzio, A., Pagano, M., Lehmann, R., 2017. GCL and CUL3 Control the Switch between Cell Lineages by Mediating Localized Degradation of an RTK. Dev. Cell 42, 130-142.e7. https://doi.org/10.1016/j.devcel.2017.06.022

Panfilio, K.A., Jentzsch, I.M.V., Benoit, J.B., Erezyilmaz, D., Suzuki, Y., Colella, S., Robertson, H.M., Poelchau, M.F., Waterhouse, R.M., Ioannidis, P., Weirauch, M.T., Hughes, D.S.T., Murali, S.C., Werren, J.H., Jacobs, C.G.C., Duncan, E.J., Armisén, D., Vreede, B.M.I., Baa-Puyoulet, P., Berger, C.S., Chang, C., Chao, H., Chen, M.-J.M., Chen, Y.-T., Childers, C.P., Chipman, A.D., Cridge, A.G., Crumière, A.J.J., Dearden, P.K., Didion, E.M., Dinh, H., Doddapaneni, H.V., Dolan, A., Dugan, S., Extavour, C.G., Febvay, G., Friedrich, M., Ginzburg, N., Han, Y., Heger, P., Holmes, C.J., Horn, T., Hsiao, Y., Jennings, E.C., Johnston, J.S., Jones, T.E., Jones, J.W., Khila, A., Koelzer, S., Kovacova, V., Leask, M., Lee, S.L., Lee, C.-Y., Lovegrove, M.R., Lu, H., Lu, Y., Moore, P.J., Munoz-Torres, M.C., Muzny, D.M., Palli, S.R., Parisot, N., Pick, L., Porter, M.L., Qu, J., Refki, P.N., Richter, R., Rivera-Pomar, R., Rosendale, A.J., Roth, S., Sachs, L., Santos, M.E., Seibert, J., Sghaier, E., Shukla, J.N., Stancliffe, R.J., Tidswell, O., Traverso, L., Zee, M. van der, Viala, S., Worley, K.C., Zdobnov, E.M., Gibbs, R.A., Richards, S., 2019. Molecular evolutionary trends and feeding ecology diversification in the Hemiptera, anchored by the milkweed bug genome. Genome Biol. 20, 64. https://doi.org/10.1186/s13059-019-1660-0

Robertson, S.E., Dockendorff, T.C., Leatherman, J.L., Faulkner, D.L., Jongens, T.A., 1999. germ cell-less,is required only during the establishment of the germ cell lineage of,Drosophila,and has activities which are dependent and independent of its localization to the nuclar envelope. Dev. Biol. 215, 288–297. https://doi.org/10.1006/dbio.1999.9453

Schoppmeier, M., Schröder, R., 2005. Maternal Torso Signaling Controls Body Axis Elongation in a Short Germ Insect. Curr. Biol. 15, 2131–2136. https://doi.org/10.1016/j.cub.2005.10.036

Skelly, J., Pushparajan, C., Duncan, E.J., Dearden, P.K., 2019. Evolution of the Torso activation cassette, a pathway required for terminal patterning and moulting. Insect Mol. Biol. 28, 392–408. https://doi.org/10.1111/imb.12560

Tamura, K., Stecher, G., Kumar, S., 2021. MEGA11: Molecular Evolutionary Genetics Analysis Version 11. Mol. Biol. Evol. 38, 3022–3027. https://doi.org/10.1093/molbev/msab120

Warren, S.L., Landolfi, A.S., Curtis, C., Morrow, J.S., 1992. Cytostellin: a novel, highly conserved protein that undergoes continuous redistribution during the cell cycle. J. Cell Sci. 103, 381–388. https://doi.org/10.1242/jcs.103.2.381

Weisbrod, A., Cohen, M., Chipman, A.D., 2013. Evolution of the insect terminal patterning system—Insights from the milkweed bug,Oncopeltus fasciatus. Dev. Biol. 380, 125–131. https://doi.org/10.1016/j.ydbio.2013.04.030

Ylla, G., Nakamura, T., Itoh, T., Kajitani, R., Toyoda, A., Tomonari, S., Bando, T., Ishimaru, Y., Watanabe, T., Fuketa, M., Matsuoka, Y., Barnett, A.A., Noji, S., Mito, T., Extavour, C.G., 2021. Insights into the genomic evolution of insects from cricket genomes. Commun. Biol. 4, 733. https://doi.org/10.1038/s42003-021-02197-9