Archaeological Bone Assemblages Trace Changes in Turtle and Tortoise Diversity in Continental Southeast Asia

Corentin Bochaton[1*], Wilailuck Naksri[3], Sirikanya Chantasri[1,2], Melada Maneechote[4], Supalak Mithong[1], Sophady Heng[5], Hubert Forestier[6], Prasit Auetrakulvit[4], Valéry Zeitoun[7], Jutinach Bowonsachoti[1,2], Julien Claude[1,8]

[1]ISEM, Univ Montpellier, CNRS, IRD, Montpellier, France.
[2]12th Regional Office of Fine Art Department, Nakhon Si Thammarat, Thailand,
[3]Northeastern Research Institute of Petrified Wood and Mineral Resources, Nakhon Ratchasima Rajabhat University, Nakhon Ratchasima, Thailand.
[4]Department of Archaeology, Faculty of Archaeology, Silpakorn University, Bangkok, Thailand.
[5]Ministry of Culture and Fine Arts of Cambodia, Phnom Penh, Cambodia.
[6]Muséum national d'Histoire naturelle, Musée de l'Homme, UMR 7194 HNHP, Paris, France.
[7]UMR 7207. CR2P-CNRS-MNHN-SU, Sorbonne Université, campus Jussieu, T. 46-56, 5ème étage, case 104, 4, place Jussieu, 75 252 Paris Cedex 05, France.
[8]Faculty of Science, University of Chulalongkorn, Bangkok, Thailand.

*Corresponding author
Correspondence: Corentin.bochaton@cnrs.fr

ABSTRACT

Among tropical regions, continental Southeast Asia is a major hotspot of biodiversity. However, due to rapid urban expansion and severe deforestation, this area is recognized as one of the top global conservation priorities. As in other tropical regions, the prevailing idea is that the decline of local biodiversity in this region is a direct result of the overwhelming environmental changes that have occurred over the last 80 years. However, paleoecological data indicate that tropical forests have been exploited by human populations since at least the Late Pleistocene, and forest clearance for farming began several thousand years ago. To gain a complete overview of the long-term evolution of vertebrate fauna in continental Southeast Asia, it is necessary to investigate the past. This task is currently out of reach, mainly due to the lack of paleontological and historical studies regarding small animals, including emblematic taxa like turtles. Indeed, although some evidence suggests that the biodiversity of turtles in continental Southeast Asia may have changed over the last millennia, the paucity of historical records and archaeological assemblages studies currently precludes further assessment of the potential effect of past environmental changes, including potential human disturbances. The present study consists of a paleontological analysis of turtle bone assemblages collected from four prehistoric sites in Thailand and Cambodia. The aim is to test the hypothesis of changes in turtle biodiversity over the last 10,000 years. Our results indicate that turtle assemblages in most areas have experienced only limited changes across the Holocene, except for

Laang Spean in Cambodia. However, a few rare taxa may represent extirpated or
extinct species of Testudinidae in continental Southeast Asia at an unknown time
during the Holocene. We also observed significant changes in the geographic
distribution of some genera, such as *Batagur*, *Cyclemys*, and *Amyda*. The fact that
turtle fauna has been rather resilient and unchanged for several millennia shows that
past human impact in most investigated areas had limited effects on these taxa. The
strong modifications observed in Cambodia could, however, serve as a warning about
the potential fate of turtle biodiversity in this region if further loss of natural
environments occurs.

Keywords: Quaternary; Paleontology; Human Impact; Testudines; Southeast Asia

Introduction

In our modern world, the environmental degradations caused by human activities have resulted
in a current rate of species extinction that is estimated to be 100 to 1,000 times greater than the
"normal rate" observed throughout geological time and a genus extinction rate 35 greater (Cowie
et al., 2022; Ceballos & Ehrlich, 2023). Researchers estimate that, since 1500, between 0.8% and
1.5% of vertebrate species on Earth have gone extinct (Ceballos et al., 2015; Johnson et al., 2017;
Ceballos & Ehrlich, 2018). In this context, a deep understanding of the dynamics of these
extinctions is critical to better evaluate the human impact on the environment. However,
paradoxically, these are still not fully understood. This is particularly true for the tropics, known to
be among the most threatened ecosystems. One of the main gaps in our knowledge concerns the
poor understanding the scientific community has of the deep history of human impact on
ecosystems and biodiversity in these regions. Notably, modern tropical rainforests are often seen
as wild areas, never having been impacted by preindustrial societies and removed from human
exploitation (Denevan, 1992; Willis et al., 2004; Barlow et al., 2012; Roberts, 2019). However,
increasing numbers of paleoenvironmental and archaeological studies are demonstrating that this
view is highly romanticized and misleading: these environments have been extensively exploited
and modified by human groups since prehistory (Denham et al., 2003; Willis et al., 2004; Roberts,
2019; Stephens et al., 2019). An evaluation of the evolution of the biodiversity in these regions
across the Quaternary is thus needed to clarify whether or not human populations had a major
impact on it.

Among tropical areas, Southeast Asia is a major hotspot of biodiversity, hosting at least 6,220
non-fish vertebrate species (Myers et al., 2000). Due to rapid urban expansion and strong
deforestation, this geographic area is recognized as one of the worldwide top conservation priorities
(Sodhi et al., 2010; Duckworth et al., 2012). As a result of these anthropogenic disturbances,
Southeast Asia has the highest proportion of its vascular plant, reptile, bird, and mammal species
categorized as globally threatened on the IUCN Red List (Sodhi et al., 2010). Within Southeast
Asia, Thailand, and Cambodia are informative areas to study past ecological phenomena in
continental Southeast Asia. These countries are also among the places with the highest biodiversity
and greatest human-induced threats of extinction. Both countries are located in one of the 25
identified Earth biodiversity hotspots: the Indo-Burma area, with Thailand also part of the
Sundaland area (south of its peninsula) (Myers et al., 2000). Combined, these two hotspots host
3,985 non-fish vertebrate species, representing 14% of non-fish vertebrate total biodiversity on our
planet. Thailand and Cambodia respectively host 1,923 and 1,028 non-fish vertebrate species,
among which 9.6% and 10.4% are considered currently threatened by the IUCN
(www.iucnredlist.org). However, although the animal biodiversity of Southeast Asia is currently
declining and could decline even more by 2050 (Trisurat et al., 2010; Paradis, 2018), the overall

environmental situation presents strong variations between the different countries, and this
tendency could be reversed should the appropriate measures be taken (Pomoim et al., 2022;
Botterill-James et al., 2024). As in other tropical areas, the prevailing idea is that the decline of
biodiversity in Thailand is a direct effect of the modern economic revolution, which started around
1950 (Falkus, 1991), with the beginning of broad-scale agricultural exploitation (especially for rice
and teak wood). These recent ecological impacts led to the loss of most of Thailand's forest cover
in less than forty years (Ramitanondh, 1989; FAO, 2020a). The situation appears similar in
Cambodia, even if this phenomenon started later, around 1970 (Tsujino et al., 2019). Overwhelming
environmental changes have obviously occurred in the last 80 years in continental Southeast Asia,
and these changes have had a strong impact on the native vertebrate biodiversity (Baker &
Phongpaichit, 2014). However, paleoecological data indicate that tropical forests have been
exploited by human populations since the Late Pleistocene at least (Conrad, 2015). Likewise, the
first traces of mass forest clearance for farming started around 2000 B.C. (White et al., 2004).
Investigating the deeper past might improve our understanding of biodiversity loss dynamic in
continental Southeast Asia.

Moreover, little is known about the past large mammal diversity of Southeast Asia (e.g., Zeitoun
et al., 2010; Conrad, 2015), and almost nothing is known about the past diversity of the small
animals accounting for 97% of the current terrestrial vertebrate biodiversity of this region
(Chaimanee, 1998; Bochaton et al., 2024). Among these small vertebrate species are turtles and
tortoises, animals that are, for half of them, considered strongly exposed to human disturbance and
at risk of extinction on a global scale (Böhm et al., 2013). These animals present significant diversity
in Thailand and Cambodia, with 27 and 14 species of terrestrial and freshwater turtles respectively
(http://www.iucnredlist.org). However, nearly no data exist about the evolution of their diversity over
the last 10,000 years, although preliminary data have begun to emerge, suggesting that part of that
diversity might have been impacted by human-driven past environmental changes. Indeed, a study
from the central plain of Thailand has shown that turtles and tortoises were more diverse during
historical times than they are today (Claude et al., 2019). In that pioneering study, eleven taxa were
identified in several archaeological assemblages dated between 4,000 and 1,000 BP. These taxa
include *Batagur affinis*, *Manouria emys*, and *Geochelone* sp., three genera that are no longer
present in that area nowadays. Although not figured and represented by a single individual,
Siebenrockiella crassicolis may also have been identified in the northernmost part of Thailand in
the Late Pleistocene/Early Holocene site of Spirit Cave (Conrad et al., 2016), a place where it does
not seem to occur anymore. These few pieces of evidence suggest that the biodiversity of turtles
in continental Southeast Asia has potentially changed over the last millennia, but the lack of
occurrence data collected from archaeological assemblages currently precludes further
assessment of this potentially effect of past environmental changes, including potential past human
disturbances.

The present study aims to investigate this question further by conducting an in-depth
paleontological study of the turtle bone assemblage collected from four prehistoric sites in Thailand
and Cambodia to test the hypothesis of changes in turtle faunas over the last 10,000 years. This
material was first examined in a zooarchaeological study focused on the species representing most
of the assemblages, *Indotestudo elongata* (Bochaton et al., 2023). In the present work, we will
focus on the investigation of the other species found in the assemblages and will discuss their
identification in greater detail than is possible in a classical zooarchaeological study.

Material and Methods

Taxonomic identification of the bone remains

The archaeological assemblages of bones studied here correspond to all the non-*Indotestudo*
turtle remains identified in the framework of a general zooarchaeological investigation of various

bone samples (Bochaton et al., 2023). During this study, key bones presenting morphological
characteristics compatible with Trionychidae, Geoemydidae, and non-*Indotestudo* Testudinidae
species were photographed for further comparisons and subjected to more detailed morphological
and taxonomic analysis. The identification of these bones was performed based on direct
comparison with pictures and 3D models of most of the turtle species currently occurring in
Southeast Asia (see specimen list in Appendix 1). This process was complicated by the fact that,
as with all zooarchaeological assemblages, most of the studied material consisted of isolated bones
and plates that were more or less altered by post-mortem manipulations by archaeological human
populations or natural post-depositional processes. These factors made the identification of this
material more challenging than that of samples corresponding to at least partially complete
specimens. Consequently, precise identification at the species level was possible for only a minor
proportion of the complete bone assemblage. Thus, with few exceptions, the taxa identified in the
assemblages are represented by a very limited number of remains bearing potentially discriminant
osteological criteria of taxonomic value. Such criteria were difficult to define because only a few
studies have been conducted on the identification of isolated plates of Southeast Asian turtle
species (Pritchard et al., 2009; Claude et al., 2019). We therefore had to define new identification
criteria based on a sample of modern specimens gathered from several museum institutions (see
specimen list in Appendix 1). Unless explicitly stated in the text, all statements regarding the
morphological specificity of each taxon are made within the framework of a comparison with the
species present in our reference sample only. This means that if we state a character is unique to
a particular species, it is unique within the sample of modern species we studied, which includes
nearly all Southeast Asian turtle species but not all turtles globally. These osteological characters
were defined following the nomenclatures of Zangerl (1969) for the carapace and Szczygielski and
Piechowski (2023) for the postcranial elements. We also created drawings of the carapaces and
plastrons of as many Southeast Asian Testudinidae and Geoemydidae taxa as we could find in
museum collections and the literature. These drawings are included in Appendix 2 and although
we were not able to figure all the taxa, we hope that this initiative will facilitate future studies in the
region. The reader should however keep in mind that these figures do not represent the full
variability of the specimens included in our comparison dataset (see Appendix. 1).

The studied archaeological assemblages

The studied deposits were described in detail in a previous publication (Bochaton et al., 2023).
They include the Doi Pha Kan rock shelter located in northern Thailand (E 99° 46' 37.2''; N 18° 26'
57.0''), where three Hoabinhian burials dated between 11,170 ± 40 and 12,930 ± 50 BP were found
(Imdirakphol et al., 2017; Zeitoun et al., 2019). This site also provided a rich archaeological
assemblage corresponding to a Hoabinhian occupation older than the burials. The turtle sample
studied includes 4,762 bone remains, with identified turtle taxa corresponding to *Indotestudo*
elongata and rare other Testudinidae (71%), Geoemydidae (28%), and Trionychidae (0.6%).

The second site is Moh Khiew Cave, an archaeological rock shelter located in southern
Thailand, in the Krabi Province (E 98° 55' 49.27''; N 08° 09' 36.32''). It was first excavated by S.
Pookajorn between 1991 and 1994 (Pookajorn, 2001) and again in 2008 by P. Auetrakulvit
(Auetrakulvit et al., 2012). The stratigraphy of the site comprises several archaeological layers
dated from the Holocene to the Late Pleistocene, most of which correspond to Hoabinhian
occupations. The material studied in the present work corresponds to that collected in 2008. The
turtle sample from this site includes 5,740 bone remains, with identified turtle taxa corresponding
to *Indotestudo elongata* (67%), Geoemydidae (32%), and Trionychidae (1%). These bones mainly
come from two layers. The first layer provided two ^{14}C dates: 7,520 ± 420 BP and 8,660 ± 480 BP
(Auetrakulvit et al., 2012). The second layer provided two additional dates: 8,730 ± 480 BP and
9,270 ± 510 BP(Auetrakulvit et al., 2012).

The third site is Khao Ta Phlai Cave, located in southern Thailand, in the province of Chumphon
(E 99° 5' 49.08''; N 10° 36' 12.39''). It has been excavated by the 12[th] Regional Office of the Fine

Arts Department, Nakhon Si Thammarat, since 2014. The turtle sample from this site includes
5,236 bone remains, with identified turtle taxa corresponding to *Indotestudo elongata* (63%),
Geoemydidae (34%), and Trionychidae (1.9%). This bone sample is divided into two chronological
assemblages corresponding to the Metal Ages and Neolithic periods. These layers have been
dated based on the typology of the archaeological artifacts they have provided, such as lithic tools,
ceramic shards, and metal objects. However, no direct dating is available yet, as the detailed study
of the site is still ongoing.

The last site is Laang Spean Cave, a large cave of more than 1,000 m² located in northwest
Cambodia, in the Battambang Province (E 102° 51' 00.0''; N 12° 51' 00.0''). The site was first
excavated between 1965 and 1968 by R. Mourer and C. Mourer-Chauviré (Mourer-Chauviré et al.,
1970; Mourer-Chauviré & Mourer, 1970). Following this initial exploration, the site underwent a new
detailed archaeological excavation by H. Forestier between 2009 and 2019 (Forestier et al., 2015;
Sophady et al., 2016). These excavations led to the discovery of an important undisturbed
Hoabinhian layer dated between 4,785 ± 25 cal. BP and 11,000 ± 100 cal. BP as well as Neolithic
sepulture and a Neolithic occupation date between 3,310 ± 29 BP and 3,370 ± 25 BP (Zeitoun et
al., 2024). The turtle sample we studied from this site includes 8,749 bone remains, with identified
turtle taxa corresponding to *Indotestudo elongata* and rare other Testudinidae (85%),
Geoemydidae (13%), and Trionychidae (0.4%).

Results

Paleontological study

Geoemydidae Theobald, 1868

Batagur sp. Gray, 1856 – Figure. 01

Material: Two fused peripheral plates from Laang Spean. (**Table. 01**)

Descriptions and comments:

The two identified plates (**Fig. 01**) correspond to the right first and second peripheral plates of
large size that are fused to the point where their separation line is no longer visible. In dorsal view,
the overall aspect of the plates is smooth; they are relatively elongated, and their anterolateral
margin is rounded and smooth. The sulci of the 1st pleural, and of the 1st and 2nd marginal scutes
are thin and well-visible. The first vertebral scute sulcus cannot be seen and was certainly included
in the nuchal plate. In ventral view, a ventrodorsal depression is found after the scute sulcus. This
element has been attributed to *Batagur* sp. because it is large, the dorsal margin of the marginal
scute is situated well below the costo-peripheral suture, the fusion between plates (first and second
peripheral plates and probably the first costal plate, of which a small fragment is visible on the
dorsal margin of the fragment), and the two fused plates form a strongly arched anterolateral
margin. The attribution to *Batagur* and not to *Orlitia* is based on the reduced extension of the
marginal scutes, which are far from the costo-peripheral suture, and on the fact that the first
vertebral scute does not cover the first peripheral plate and is restricted to the nuchal bone. These
characters are valid for adult specimens that we did not figure in Appendix 2 because the suture
between the plates completely disappeared at some point during the maturation of the specimen.

Figure 01 - Two fused peripheral plates of *Batagur* sp. in dorsal (left) and ventral (right) views (Laang
Spean). <u>Abbreviations</u>: **c1**: 1st costal plate; **c-p s**: costo-peripheral suture; **v.d.**: ventrodorsal depression;
M1&2: 1st and 2nd marginal scute; **P1**: 1st pleural scute. Scalebars=1cm.

Cuora Gray, 1856

Cuora couro (Leschenault de la Tour, 1812) – Figure.02 - A, D-H

Material: 277 remains distributed in the four sites (**Table. 01**): five nuchal plates, nine neural
plates, height suprapygal plates, height pygal plates, 10 costal plates, 173 peripheral plates, 22
epiplastra, two entoplastra, five hyoplastra, 12 fragmented hyoplastra or hypoplastra, seven
hypoplastra, six xiphiplastra, one scapula, and nine ilia bones.

Description of key elements and comments:

The plates attributed to this species present a smooth external surface with sutural contacts
between the plates that are composed of many densely packed, very small pits, spikes, and crests
of limited elevation and depth (**Fig. 02: B**). The plates are also thick relative to their size, and the
limits between the scutes are marked by thin sulci. The significant thickness of the plates relative
to their small size, the ligamentous contact between the plastron and the carapace, and the sutural
contacts are characteristics we observed in *Cuora couro* and in the *C. galbinifrons* complex, and
that were absent in *C. mouhotii* (**Fig. S2 and S3**).

The nuchal plate is thick and vaulted. In dorsal view, the bone is slightly wider than long with a
smooth and slightly curved anterior margin. The imprint of the cervical scute is thin and triangular
with concave lateral and posterior borders. The sulcus of the anterior margin of the first pleural
scute is visible on a small portion of the most lateral extremity of the plate. In ventral view, the
cervical scute is broader than in dorsal view. The combination of these criteria was observed only
in *C. couro* (**Fig. S2**).

The second suprapygal plate (**Fig. 02: D**) is pentagonal due to the medial contact of the two 7th
costal plates and sometimes hexagonal (figured specimen) in posterior view. These plates are
slightly wider than high. If present, the contact with the 11th peripheral plates is the same length as
that of the pygal plate. The posterior sulcus of the 5th vertebral scute is visible on the posterior part
of the bone and is slightly convex in posterior view. The sulcus is straigth at the level of the contact
with the 12th marginal scutes. The second suprapygal plate also presents the above general
characteristics (smooth surface, thin sulci, thickness). Within geoemydids and among our reference
samples, the combination of these characteristics was found only in *C. couro* (**Fig. S3**), but it may
also occur in species absent from continental Southeast Asia, namely *C. amboinensis* and *C.
philippinensis*. This plate presents a morphology similar to that of *Malayemys macrocephala* and

Siebenrockiella crassicollis (**Fig. S13, S26**), but in these species, the plates are thinner and less
thick relative to their size by comparison to those of *C. amboinensis* and the *C. galbinifrons* complex
(*C. galbinifrons*, *C. bourreti*, and *C. picturata*).

The peripheral plates of the bridge present a nearly smooth contact area with the plastron,
indicating that the attachment of the plastron and carapace is ligamentous, with no sutural contact
between the peripherals and the hypoplastron and hyoplastron. Additionally, a strong pointed
extension is present on the ventromedial surface of the 5th peripheral plate (**Fig. 02: A**). This
process is followed by the ligamentous contact between the hyoplastron and hypoplastron. Among
our sample of modern species, these criteria have only been observed in *C. couro* and the *C.
galbinifrons* complex (**Fig. S3**) but are absent in our specimen of *C. mouhotii*. This character could
also be lacking in other non-Southeast Asian *Cuora* species, such as *C. aurocapitata*, or be partly
absent in others, such as in *C. pani*.

The pygal plate, as the peripheral plates (**Fig. 02: E**), is small and thick. It is square in dorsal
view, and its ventral margin does not exhibit any indentation. It is intersected by the medial
separation of the 12th marginal scutes and exhibits the general characteristics of *C. couro* (smooth
surface, thin sulci, thickness, and morphology of the contact areas). In our modern sample, this
morphology is only encountered in *C. couro*, while the specimens of *C. galbinifrons* we observed
have a small ventral indentation (**Fig. S2**). No other Southeast Asian taxa present a thick, small-
sized, square pygal plate.

The epiplastron (**Fig. 02: F**) is short, flat, and thin. Its lateral margin is smooth, regular, and
rounded. In visceral view, the gular and humeral scutes cover at least half of its surface, with the
area covered by each scute being nearly equal. The area occupied by these scutes presents a
slightly rugged surface. In ventral view, the humeral scute covers two-thirds of the surface of the
bone. The combination of these morphological criteria, along with the small size of the elements,
is unique to *C. couro* and *C. galbinifrons* (**Fig. S2**). It is absent in *C. mouhotii* but might be rather
general in other *Cuora* species.

The hyoplastron and hypoplastron (hypoplastron in **Fig. 02: H**) present a contact area with the
carapace that is completely smooth, similar to the one described for the peripheral plates of the
bridge. The inguinal and axillary buttresses are weakly developed. The contact area between the
hyoplastron and hypoplastron is smooth, indicating a ligamentous connection. Among living and
subfossil Geoemydidae, this morphology is, in our sample, observed only in *Cuora*. In our sample,
it is unique to *C. couro* and *C. galbinifrons*, but it is absent in *C. mouhotii* (**Fig. S2, and S3**) although
this might reflect a bias of our comparative sample. Indeed, this character is subject to some degree
of ontogenetic variations. Ligamentous articulations between the hyo and hypoplastra have been
observed in other taxa (*Cyclemys*, *Notochelys*) but the articulations are not as smooth as in *Cuora*.
In addition, the femoral scute present a very short midline length, a character only observed in *C.
couro*, *C. flavomarginata*, and *C. picturata* and absent in *C. galbinifrons* (Naksri et al., 2013).

The xiphiplastron (**Fig. 02: H**) is thin, flat, and wider than it is long in ventral view. In this view,
its lateral margin is very convex and presents only a very small indentation at the level of the
boundary between the femoral and anal scutes. It bears no trace of an anal notch, but the inter-
anal sulcus is present. This morphology is unique to most species of *Cuora*, and in our sample, to
C. couro and the *C. galbinifrons* complex (**Fig. S2**). The anal notch is also absent in *C.
flavomarginata* and *C. bourreti* (Naksri et al., 2013).

The only scapula (**Fig. 02: C**) attributed to this taxon is a fragment from Laang Spean
representing the dorsal process. In lateral view, this process is quite short and broad. Its orientation
changes close to the glenoid cavity, as the dorsal three-fourths of its length is slightly inclined
medially. Because of this, its lateral margin forms an obtuse angle. The dorsal extremity of the
dorsal process is also slightly enlarged. The medial inclination of the dorsal process of the scapula

is found in all the observed members of *Cuora*, but in most taxa, this process is uniformly arched,
and its lateral margin forms an obtuse angle only in our specimens of *C. couro*. An enlargement of
the dorsal part of the dorsal process has been seen in all the *Cuora* in our sample, but the strong
build of the described fossil scapula is only observed in *C. couro*, *C. trifasciata*, and *C. galbinifrons*,
while it is more elongated in the other species.

The ilium (**Fig. 02: G**) is moderately elongated. In lateral view, its ilium neck is thin and rounded,
and its posterior process is enlarged and sub-quadrangular in shape. The dorsal and posterior
edges of this process are rugged. This general shape of the ilium was observed in *C. galbinifrons*
and *C. couro*, but in *C. galbinifrons*, the posterior process is shorter, and its posterior edge is more
rounded, making the overall shape of the whole process more hook-like (see also Yasukawa,
Hirayama, and Hikida, 2001). In contrast, this process is longer with more angular margins in *C.*
couro.

Figure 02 - A) Fused 5th and 6th left peripheral plates of *C. couro* in lateral (top-left), medial (top-right), and ventral (bottom) views (Khao Ta Phlai); **B)** Detailed view of the contact areas between peripheral plates in a modern specimen of *C. couro*; **C)** Partial right scapula of *C. couro* in lateral view

(Laang Spean); **D**) 2nd suprapygal plate of *C. couro* in posterior view (Moh Khiew); **E**) Left 11th peripheral
plate of *C. couro* in posterior (left) and anterior (right) view (Laang Spean); **F**) Right epiplastron of *C.*
couro in dorsal (left) and ventral (right) views (Moh Khiew); **G**) Right ilium bone of *C. couro* in lateral
view (Doi Pha Kan); **H**) Left hypoplastron and xiphiplastron of *C. couro* cemented together by calcite
(Laang Spean). <u>Abbreviations</u>: **AN.**: annal scute; **c. p.**: contact area with the plastron; **c. py.**: contact
area with the pygal plate; **c. p11**: contact area with the 11th peripheral plate; **d. p.**: dorsal process; **FE.**:
femoral scute; **g. c.**: glenoid cavity; **g. s.**: gular scute; **h. s.**: humeral scute; **i.**: indentation; **i. b.**: inguinal
buttress; **i.n.**: ilium neck; **hyp.**: hypoplastron; **M12**: 12th marginal scute; **p. e.**: pointed extension; **p. p.**:
posterior process; **V5**: 5th vertebral scute; **x.**: xiphiplastron. Scalebars=1cm.

Figure 03 - Nuchal plate of *Cuora* sp. in dorsal (left) and ventral (right) views (Moh Khiew).
<u>Abbreviations</u>: **C.**: cervical scute; **V1**: first vertebral scute. Scalebar=1cm.

Cuora sp. – Figure. 03

Material: One partial nuchal plate from Moh Khiew. (**Table. 01**)

Description and comments:

This nuchal plate (**Fig. 03**) is only represented by its anterior portion; its posterior part is well
projected in the dorsal direction. In dorsal view, the lateral sulcus of the first vertebral scute extends
beyond the limits of the plate. In the same view, the cervical scute is elongated and very thin. In
ventral view, the scute is triangular and approximately 2.5 times longer than wide. In our reference
sample, a similar morphology of the cervical scute was only observed in *C. mouhotii* (**Fig. S3**), but
in this species, although very thin, the scute is usually shorter and the anterior margin of the first
vertebral scute is more convex. Additionally, the anterolateral margins of the same scute do not
extend onto the adjacent plates. Outside of our references, the cervical scute morphology
described here was observed as a rare morphology in several species (*Cuora couro*, *Cuora*
galbinifrons complex *Cuora philippinensis*, and *C. mouhotii*) (J. C. Pers. Obs.). It thus seems likely
that this plate is just an abnormal specimen of *C. couro*.

Cyclemys cf. *oldhamii* Gray, 1863 – Figure. 04

Material: 489 remains distributed in all the sites (**Table. 01**): 20 nuchal plates, 32 costal plates,
30 neural plates, 292 peripheral plates, three 1st suprapygal plates, four 2nd suprapygal plates, four
pygal plates, 27 epiplastra, three entoplastra, 20 hyoplastra, 18 hypoplastra, 24 fragments of hyo
or hypoplastra, 12 xiphiplastra.

Descriptions and comments:

The nuchal plates (**Fig. 04: A, C**) exhibit a very typical morphology observed in *Cyclemys*
oldhamii (**Fig. S5**). The bone is nearly as long as it is wide, with imprints of the 1st pleural scutes

occupying a significant part of its lateral sides in dorsal view. In the dorsal view, the cervical scute
presents a slender and elongated triangular shape, laterally depressed in its anterior half and
concave posteriorly. The imprint of the cervical scute is not visible in the ventral view, but this
condition is variable (J. C. Pers. Obs.). In the two specimens of *Cyclemys atripons* we observed
(**Fig. S4**), the cervical scute is shorter, and the bone is thickened and slightly arched below the
scute, making this area very prominent. This characteristic was only observed in this species
among all the Southeast Asian taxa we examined. In our six specimens of *Cyclemys dentata*, the
cervical scute is shorter and a bit thicker than the morphology observed in *C. oldhamii*. We did not
observe any specimens of *C. enigmatica*, *C. fusca*, and *C. pulchristriata* as these three species are
currently absent from the studied area although some recent observations indicate possible
occurrences of *C. enigmatica* in the Kra isthmus. Considering that some species are missing from
our sample and that we had access to a limited number of comparison specimens, we propose an
identification of *Cyclemys* cf. *oldhamii* while keeping in mind that the species of *Cyclemys* in the
area of Moh Khiew and Khao Ta Phlai could be *C. enigmatica*.

The peripheral plates (**Fig. 04: A, B**) are thin and small in size, with the dorsal third of the height
of their lateral side occupied by the pleural scutes. However, the peripheral plates of the bridge
present an articulation morphology with the plastron that stands out from most other taxa. The 3th
to 7th peripheral plates have a ventral part that projects medially with a thick contact area with the
plastron, visible only in medial and ventromedial views. This contact area is marked by deep ridges
mainly located in the dorsal and ventral parts of the contact area, while its center is ornamented
with weakly marked ridges. The 3rd to 4th and 7th peripherals exhibit sub-circular imprints of the
ligaments attaching the carapace to the buttresses on their medial sides. The imprint of the axillary
buttress can be either fully located on the third peripheral or at the intersection of the 3rd and 4th. It
can also be in contact with the lateral articular surface of the plastron or separated from it by a
ridge. We have not been able to determine if this variability is of taxonomic significance at the
interspecific level within the genus. The imprint of the inguinal buttress is located on the 7th
peripheral and might slightly overlap with the 8th. This attachment type is intermediate between the
strong joint observed in nearly all investigated taxa, where the articular areas of the bones are
strongly interlocked (as seen in the other carapace plates of *Cyclemys* species) (**Fig. S5**), and the
completely smooth contact surface in some species of *Cuora* (*C. couro* and *C. galbinifrons*) (**Fig.
S2**). The intermediate condition observed in *Cyclemys* only occurs in *Cuora mouhotii* (**Fig. S3**),
where the attachment for the axillary buttresses is reduced and triangular.

The neural plates are thin and of trapezoid shape in dorsal view. Only a few of them, mostly
from posterior positions, present traces of a keel. These bones do not exhibit characteristics unique
to *Cyclemys* but have been attributed to this taxon due to its prevalence in the investigated samples
and because their morphology does not differ from that of our comparative specimens of *Cyclemys
oldhamii* (**Fig. S5**).

The 1st suprapygal (**Fig. 04: D**) is fused with the 8th neural plate. It has an elongated shape with
a broad anterior part and an elongated posterior part in dorsal view. The two parts are separated
by a transverse anterior sulcus between the 4th and 5th vertebral scutes, where the bone is slightly
constricted. This morphology is common in the *Cyclemys* genus but is rare in other living
geoemydid genera.

The epiplastron (**Fig. 04: E**) is thin and moderately elongated, with a circular antero-lateral
margin and a well-marked acute indentation at the level of the sulcus between the humeral and the
gular scutes in ventral view. The epiplastral lip is weakly developed in visceral view. Still in visceral
view, the delimitation of the humeral and gular scutes is visible. This morphology is more similar to
C. oldhamii (**Fig. S5**). Other taxa of comparable size and morphology lack a similar indentation,
which can be either absent (*Cuora couro*, *Cuora mouhotii*, *Cuora trifasciata*) (**Fig. S2, and S3**) or
more obtuse (*Cuora galbinifrons*, *Cyclemys atripons*, *Heosemys spinosa*) (**Fig. S4, and S9**). Some
other taxa, like *Geoemyda* (**Fig. S6**), *Indotestudo*, *Melanochelys* (**Fig. S13**), *Morenia*, *Manouria*

(**Fig. S11, and S12**), and *Notochelys* (**Fig. S14**), present a well-marked indentation similar to the
one observed in *Cyclemys oldhamii*, but in these taxa, the anterior part of the bone forms an
epiplastral projection or quadrangular shape that is absent in *C. oldhamii* (**Fig. S5**). In *Notochelys*
(**Fig. S14**), the strong development of the epiplastral lips forms an indentation, absent in *Cyclemys*,
at the level of the inter-gular sulcus.

The entoplastron is diamond-shaped, with its posterior part slightly more developed than the
anterior one beyond the posterior humeral sulcus. Its posterior margin is subcircular in ventral view,
and is crossed by the posterior sulci of the humeral scutes around its mid-length and by the
posterior sulci of the gular scutes near its anterior tip. Entoplastra with similar morphologies are
present in *Cuora couro* (**Fig. S2**), *Cuora galbinifrons*, and *Siebenrockiella crassicollis* (**Fig. S18**),
but in these taxa, there is a more developed posterior entoplastral process in visceral view. In
Melanochelys trijuga (**Fig. S13**) and *Geoemyda spengleri* (**Fig. S6**), the posterior sulcus of the
humeral scute is closer to the posterior margin of the bone. In *Mauremys annamensis*, the
entoplastron presents both of these differences compared to *Cyclemys*. In *Indotestudo elongata*,
although the morphology is similar to *Cyclemys*, the entire bone is clearly thicker.

The hyoplastron (**Fig. 04: G**) is moderately elongated. In ventral view, the humero-pectoral
sulcus is only visible near the anterior tip of the bone and intersects the suture line with the
posterolateral part of the entoplastron. The posterior sulcus of the pectoral scute is located near
the posterior margin of the bone. The articular surface with the plastron presents a morphology
similar to the one described on the peripheral plates of *Cyclemys*: it is composed of a medial furrow
located between two crests bearing small spikes that fit the grooves present on the peripheral
plates of the bridge. This type of articular surface, also observed on the hypoplastron, is unique to
Cyclemys among the considered modern species. In *Cuora*, although the contact between these
bones is also ligamentous, the contact area in *Cyclemys* is never completely smooth. Additionally,
in *Cyclemys*, the abdominal scute partly covers the hyoplastron, which is not the case in *Cuora*.

The xiphiplastron (**Fig. 04: F**) is 1.3 times longer than it is wide. Its lateral margin is smooth and
circular, with a moderately developed anal notch. The femoro-anal sulcus is positioned posteriorly
but close to the mid-length of the bone and runs parallel to the suture line with the hypoplastron.
The combination of these characteristics is unique to *Cyclemys*. A nearly similar morphology exists
in *Melanochelys trijuga* (**Fig. S13**), but the anal notch is more developed in this latter species, and
the lateral lips of the anal scutes are more elevated.

Figure 04 - A) Fragments of a carapace of *Cyclemys* cf. *oldhamii* in dorsal view (Khao Ta Phlai); **B**)
The plastral articular portion of the carapace illustrated in A; **C)** Nuchal plate of *Cyclemys* cf. *oldhamii* in
dorsal view (Moh Khiew); **D**) 1st suprapygal plate of *Cyclemys* cf. *oldhamii* (Doi Pha Kan); **E**) Epiplastron

of *Cyclemys* cf. *oldhamii* in ventral (left) and dorsal (right) views (Doi Pha Kan); **F)** Xiphiplastron of
Cyclemys cf. *oldhamii* in ventral view (Doi Pha Kan); **G)** Hyoplastron of *Cyclemys* cf. *oldhamii* in ventral
view (Khao Ta Phlai). <u>Abbreviations</u>: **A.**: abdominal scute; **AN.**: annal scute; **a. n.**: annal notch; **C.**:
cervical scute; **c. a.**: contact area; **F.**: femoral scute; **G.**: gular scute; **H.**: humeral scute; **i.**: indentation;
i. a. b.: imprint of the axillary buttress; **M1**: 1^{st} marginal scute; **M2**: 2^{nd} marginal scute; **P.**: pleural scute;
PE.: pectoral scute; **P1**: 1^{st} pleural scute; **p3**: 3^{rd} peripheral plate; **p5**: 5^{th} peripheral plate; **t. a. s.**:
transverse anterior sulcus; **V1**: 1^{st} vertebral scute. Scalebars=1cm.

Heosemys Stejneger, 1902

Heosemys annandalii (Boulenger, 1903) – Figure. 05 - A

Material: Three nuchal plates from Khao Ta Phlai and Laang Spean (**Table. 01**).

Descriptions and comments:

These nuchal plates (**Fig. 05: A**) are large, but their complete shape cannot be observed as
only the anterior portions of the plates are preserved. The imprints of the 1^{st} pleural scutes are well
visible and expanded on the postero-lateral areas of the bone. In dorsal view, the anterior margin
of the bone presents a weakly marked central depression. The imprint of the cervical scute is large
and heart-shaped. The attribution of these plates to *Heosemys* is based on their size, the large
expansion of the 1^{st} pleural scutes on the posterolateral part of the nuchal plate, and the shape of
the cervical scute. Indeed, in other large-sized taxa present in the area (*Manouria*, *Batagur*, and
Orlitia) (**Fig. S1, S11, S12, S15, and S16**), the 1^{st} pleural scutes are either not in contact or poorly
expanded on the nuchal plate, and the cervical scute is not heart-shaped. The only difference we
observed between *Heosemys annandalii* (**Fig. S7**) and *H. grandis* (**Fig. S8**) in dorsal view concerns
the shape of the imprint of the cervical scute on the nuchal plate. In *H. annandalii*, it is large and
heart-shaped, whereas, in *H. grandis*, the scute is reduced, with a shape ranging from nearly
square to a somewhat heart shape presenting a very depressed posterior margin. On their ventral
surface, the nuchal plate of *H. annandalii* presents a very well-marked bulge with a strong medial
depression while the same bulge in *H. grandis* is less developed and lacks the medial depression.

Heosemys grandis (Gray, 1860) – Figure. 05 - B and C

Material: Three nuchal plates and two peripheral plates from Laang Spean (**Table. 01**).

Descriptions and comments:

These nuchal plates (**Fig. 05: B**) present the same characteristics as those attributed to *H.*
annandalii. The only difference from the previously described morphotype concerns the shape of
the cervical scute, which is reduced in the three identified specimens and nearly square in the
figured one (see previous paragraph).

Posterior peripheral plates (**Fig. 05: C**) with significant serration can be attributed to this species
based on their large size and morphology. Although several taxa present limited serrations on their
posterior peripheral plates (9^{th} to 11^{th}) (namely *Cuora mouhotii* (**Fig. S3**), *Cyclemys atripons* (**Fig.**
S4), *Notochelys platynota* (**Fig. S14**), and *Heosemys annandalii* (**Fig. S7**)), strong serrations are
only present in *Heosemys grandis* (**Fig. S8**), *Heosemys spinosa* (**Fig. S9**), and *Geoemyda*
spengleri (**Fig. S6**). These plates were thus attributed to *H. grandis* in Laang Spean because of
their large size.

Heosemys cf. *spinosa* (Gray, 1830) – Figure. 05 - D

Material: 15 peripheral plates and two pygal plates from Khao Ta Phlai, and Moh Khiew (**Table.**
01).

Descriptions and comments:

Peripheral plates have been tentatively attributed to *H. spinosa* based on their significant serration and small size (see previous comments about the peripheral plates of *H. grandis*). It is possible that these plates could correspond to young *H. grandis* specimens, but no other plates pertaining to this species were recorded at Khao Ta Phlai, and Moh Khiew.

The two pygal plates (**Fig. 05: D**) found in Khao Ta Phlai and Moh Kiew present a nearly unique morphology among the observed modern species. The plate is wider than it is high, tectiform, and has a thin, deep indentation of its ventral margin. In *H. grandis* (**Fig. S8**) and *H. annandalii* (**Fig. S7**), the posterior margin of the plate is smooth with a wide and moderately deep indentation, and the plate is not tectiform. Somehow similar morphologies were observed in the genus *Cyclemys* (**Fig. S4, and S5**), but in *H. spinosa* (**Fig. S9**) the plate is consistently wider than it is high which is not the case for the latter. Additionally, we found one of the two pygal plates in association with a series of peripheral plates from a juvenile individual that were too thick to be associated with *Cyclemys* but also lack the strong serrations observed in the other peripheral plates we attributed to *H. spinosa*. Given the limited number of elements attributed to *H. spinosa* and the potential ambiguity of the morphological characters considered in these attributions in our sample, we consider the identification of *H. spinosa* in our material as tentative.

Heosemys cf. *grandis/annandalii* – Figure. 05 - E

Material: 179 remains from Moh Khiew, Khao Ta Phlai, and Laang Spean (**Table. 01**): four nuchal plates, 21 neural plates, four 2^{nd} suprapygal plates, one pygal plate, four costal plates, 61 peripheral plates, five epiplastra, six entoplastra, six hyoplastra, seven hypoplastra, 30 fragmented hyoplastra or hypoplastra, nine xiphiplastra, nine humeri, six ilia bones, one ischium bone, one pubis bone, and three femora.

Descriptions and comments:

These bones were attributed to a large species of *Heosemys* based on their large size and the following morphological characteristics that distinguish them from the other large species considered in our comparative sample (*Batagur baska*, *Batagur borneoensis*, *Orlitia borneensis*, *Manouria impressa*, and *Manouria emys*). Regarding the general characteristics of the subfossil elements: despite their relatively large size, none of the carapace and plastron bones identified were fused to any neighboring elements. This precludes attributing these elements to *Batagur* and *Orlitia*, in which these elements tend to fuse once the animal has reached its adult size. Additionally, the subfossil elements considered here are relatively thick, which precludes an attribution of the neural and costal plates to *Manouria*, as these bones remain thin even in large individuals of this genus.

More specific observations were made on the following bones:

The 2^{nd} suprapygal plates are large, thick, hexagonal in shape, not fused to adjacent plates, and crossed by the dorsal 12^{th} marginal sulcus. The peripheral plates are thick, not fused to the adjacent plates, and high. The pleuro-marginal sulci are visible and located below the costoperipheral sutures of all the peripheral plates. A moderate serration is present exclusively along the margins of the 8^{th} to the 11^{th} peripheral plates. The epiplastron is thick, moderately elongated, and presents a moderately expanded anterior projection with a transverse anterior margin, giving the anterolateral margin of the whole bone a slightly angular shape in ventral view. Its articular surfaces with the other epiplastron and the hyoplastron are of similar lengths. The entoplastron has its ventral face crossed by the posterior gular sulcus and the posterior humeral sulcus; this characteristic is unique to *Heosemys* among the large Southeast Asian turtles. The hypoplastron is moderately elongated and presents nearly parallel anterior and posterior margins.

The xiphiplastron is crossed around its mid-length by the femoro-anal sulcus, which is parallel to
the straight anterior margin of the bone, and a strongly marked anal notch is present. The humerus
presents the characteristics of a large Geoemydidae, but we did not identify characters specific to
a given genus within this family. The shaft is moderately curved, a character shared with
Trionychidae, differing from Testudinidae in which the shaft is more strongly curved. The medial
and lateral processes are moderately divergent, and the deep trochanteric fossa separating them
forms an angle of around 80-90°. This angle is lower in *Manouria* (around 60°) and higher in
Trionychidae (around 100-110°). Also, in Trionychidae, the medial process is more individualized
from the humeral head than in Geoemydidae and *Manouria*, in which the two structures are not
separated. The medial process is also more extended posteriorly in Geoemydidae and
Trionychidae than in *Manouria*. The ilium is antero-laterally flared (Yasukawa, Hirayama, and
Hikida, 2001) as in all Geoemydidae, and its anterior part is triangular in lateral view. In *Manouria*,
the anterior part of the bone is also triangular but is not antero-laterally flared. Additionally, the
lateral ilial process (sensu Szczygielski and Piechowski, 2023) that is projected in a dorsolateral
direction in large Geoemydidae is projected dorsally and slightly incurved in the medial direction in
Manouria. In Trionychidae, the anterior part of the ilium is cylindrical along its entire length.
Regarding the femur, in *Heosemys* and other large Geoemydidae, the trochanter major is higher
than the trochanter minor and the two are coalesced, but the crest linking them is depressed in
ventral view. In *Manouria*, the trochanter minor is higher than the trochanter major in ventral view,
and the crest linking them is less depressed. The distal part of the bone is curved in *Manouria*,
while the whole diaphysis is nearly straight in Geoemydidae. The morphology of Trionychidae is
completely different, as the two trochanters are not coalesced and are separated by a deep
intertrochanteric fossa, similar to the one observed on the humerus. In Trionychidae, the trochanter
major is also much more developed in the dorsal direction and is triangular with a blunt apex.

Figure 05 - A) Nuchal plate of *Heosemys annandalii* in dorsal view (Khao Ta Phlai); **B**) Nuchal plate of *Heosemys grandis* in dorsal view (Laang Spean); **C**) Peripheral plate of *Heosemys grandis* in lateral view (Laang Spean); **D**) Pygal plate of *Heosemys* cf. *spinosa* in posterior view (Khao Ta Phlai); **E**) Right

7th to 11th peripherals, pygal and suprapygal plates attributed to *Heosemys* cf. *annandalii/grandis* (Laang
Spean). <u>Abbreviations</u>: **C.**: cervical scute; **i.**: indentation; **M1**: 1st marginal scute; **M12**: 12th marginal
scute; **P1**: 1st pleural scute; **p11**: 11th peripheral plate; **s.-p. 2**: 2nd suprapygal plate; **V1**: 1st vertebral
scute. Scalebars=1cm.

Malayemys sp. Lindholm, 1931 – Figure. 06

Material: Three costal plates, one fragment of hyo or hypoplastron, and two fused dentaries
from Khao Ta Phlai, Moh Khiew, and Doi Pha Kan. (**Table. 01**)

Descriptions and comments:

Three costal plates (5th and 6th) (**Fig. 06: A**) have been attributed to *Malayemys* sp. because
their medial surface presents an impression of the inguinal buttress in the ventral part of the contact
area of the two plates, A faint lateral keel in the dorsal area of the plates, and a relatively straight
and longitudinal vertebro-pleural sulcus. This condition is found only in *Malayemys* (**Fig. S10**)
among Southeast Asian turtles. In Southeast Asia, Geoemyda has also keels, but they are located
more distally, and the inguinal buttress only reaches the 5th costal plate. An imprint of the inguinal
buttress is also present on the 5th and 6th costal plates in *Batagur, Hardella,* and *Pangshura*.
However, in these taxa, the buttress imprint is much stronger, covering half or more than half the
length of the plates. *Geoclemys*, *Orlitia*, and *Siebenrockiella* present inguinal buttress imprints
similar to *Malayemys*, but *Orlitia* and *Siebenrockiella* have mushroom-shaped vertebral scutes,
resulting in non-continuous and non-longitudinal vertebro-pleural sulci. These two species also
either lack (*Orlitia*) or bear lateral keels weaker than in *Malayemys* (*Siebenrockiella)*. In *Geoclemys,*
the lateral keels are conversely more strongly marked than in *Malayemys*. The plates that are
figured present a complex vermiculation on their lateral surface that could be either related to an
antemortem phenomenon (parasitosis/senescence) or to a postmortem taphonomic alteration.
Such alterations are often observed on *Malayemys* skeletal specimens.

The dentaries (**Fig. 06: B**) present no visible suture line. In dorsal view, the dentaries are short
with a very wide and flat triturating surface. They present large imprints of the coronoid bones on
their posterior areas, and long rami of the mandible. Long and flat lower triturating surface is known
in *Malayemys*, some species of *Mauremys (M. nigricans and M. megalocephala)*, and *Geoclemys*.
However, in *Geocle*mys the symphysis is shorter than in the three other species. While in
Mauremys specimens presenting a long symphysis, the rami of the mandibles are smaller than in
Malayemys.

A fragment of hyo- or hypoplastron (**Fig. 06: C**) has also been attributed to this genus because
it presented very small anastomosed vermiculations organized in polygones on its ventral surface.
Such anastomosis pattern can also be seen in *Morenia,* but in the latter, the furrows are deeper
and slightly larger, while they are very thin and not deep in *Malayemys*.

Figure 06 - A) Left 5th and 6th costal plates of *Malayemys* sp. in dorsal (left) and ventral (right) views
(Moh Khiew); **B**) Dentaries of *Malayemys* sp. in dorsal (top) and ventral (bottom) views (Moh Khiew); **C**)
Hyo- or hypoplastron of *Malayemys* sp. in ventral view (Doi Pha Kan). <u>Abbreviations</u>: **a. p.**: anterior
portion; **i. c.**: imprint of the coronoid bone; **i. i. b.**: impression of the inguinal buttress; **v.**: vermiculation.
Scalebars=1cm.

Geoemydidae indet. Theobald, 1868

Material: 767 remains distributed in the four sites and representing nearly all of the skeletal
parts: nine nuchal plates, 131 neural plates, 49 costal plates, 229 Peripheral plates, 11 suprapygal
plates, four pygal plates, 13 epiplastra, nine entoplastra, 16 hyoplastra, 24 hypoplastra, 29
xiphiplastra, 8 undetermined fragments of plastral plates, one skull fragment, 37 scapula, 31
coracoids, 65 humeri, height radius, three ulna, 26 ilia, five ischia, 14 pubis, 38 femora, 5 tibia, 1
metapodial, and 1 phalanx. **(Table. 01)**

Descriptions and comments:

These bones are either fragmented or lack distinctive morphological features necessary for
attribution to a specific genus. They have been classified under Geoemydidae based on their
differences from elements attributed to *Indotestudo elongata* and Trionychidae. Most of these
classifications are not founded on distinctive characteristics of the Geoemydidae family but are
included in this study to offer a rough estimate of the family's significance within the complete
assemblage.

Testudinidae Batsch, 1788

Geochelone sp. Fitzinger, 1835– Figure. 07

Material: One ilium from Laang Spean. (**Table. 01**)

The ilium bone (**Fig. 07**) is partially complete and covered by a layer of calcite, with the dorsal part damaged. In lateral view, the bone is elongated, with its dorsal and ventral extremities enlarged. Its anterior and posterior edges are slightly incurved but remain nearly straight and parallel to each other in the central part of the bone. A shallow dorsal fossa is present in the dorsal area of the lateral face of the bone, though its full dorsal extent is unknown. This fossa is bordered by a thick postero-dorsal crest and a sharp antero-dorsal crest marking the anterior edge. In posterior view, the bone is slightly incurved medially, and the thick postero-dorsal crest marking the posterior delimitation of the dorsal fossa is visible. In medial view, the medial surface is smooth, and the dorsal extremity of the bone presents an angulation where the postero-dorsal crest projects laterally.

The fact that the ilium presents enlargement of its dorsal and ventral extremities, a nearly straight anterior edge, and a distal fossa are characteristics of testudinoid turtles (Zug, 1971). Among all the Southeast Asian Geoemydidae we observed, the dorsal portion of the ilium presents a broad range of morphologies (also figured in Yasukawa, Hirayama, and Hikida, 2001), but its anterior and posterior edges are never close to being parallel. Among the specimens we saw, such morphology is only encountered in *Geochelone elegans* (we did not have access to a *G. platynota* specimen) and *I. elongata*. However, in the latter species, the dorsal fossa is less deep, and its limits are well-defined on the lateral surface. Additionally, *I. elongata* does not exhibit dorsal enlargement, and this seems to be the case in our fossil, even if the morphology of its dorsal part cannot be fully assessed. We thus attribute our ilium to an undefined member of the genus *Geochelone*, as we were not able to assess the inter-specific variability of the ilium within this genus. There is a possibility that this remain could be associated with the Testudinidae indet. nuchal plate described below, but since the remains were not found at the same site, this cannot be demonstrated so far.

Figure 07 - Ilium of *Geochelone* sp. in medial (right) and lateral (left) views (Laang Spean). Abbreviations: **a.**: angulation; **a. c.**: antero-dorsal crest; **d. f.**: dorsal fossa; **p. c.**: postero-dorsal crest. Scalebar=1cm.

Manouria impressa (Günther, 1882) – Figure. 08 - A, B, C

Material: Three neural plates, five costal plates, and one pygal plate from Doi Pha Kan and Laang Spean. (**Table. 01**)

Description and comments:

The neural plates and costal plates (**Fig. 08: A and B**) attributed to *Manouria* exhibit a distinct morphology with prominent elevations along the sulcus. In the costal plates, the anterolateral, posterolateral, and medial parts of the costal bone, separated by well-defined elevations along the sulcus, are strongly depressed. Additionally, the morphology of these plates retains a juvenile appearance as they do not contact each other along the full length of their lateral margins. The ribs are also visible on half of the length of the costal plates that do not bear the transverse sulcus of the pleural scutes. This morphology has only been observed in *Manouria impressa* (**Fig. S12**).

The pygal plate (**Fig. 08: C**) is wider than it is high, with a rectangular shape in posterior view and a slightly depressed dorsal margin. It is crossed by the posterior sulcus of the 5th vertebral scute, which does not overlap with the dorsal margin of the bone. Additionally, the sulcus between the 12th marginal scutes is present. This morphology is specific to *M. impressa* (**Fig. S12**); on the pygal plate of *Manouria emys* (**Fig. S11**), the posterior sulcus of the 5th vertebral scute overlaps with the dorsal margin of the bone.

Manouria sp. Gray, 1854 – Figure. 08 – D, E, and F

Material: One nuchal plate, two peripheral plates, one pygal plate, one entoplastron, and one humerus from Doi Pha Kan and Laang Spean. (**Table. 01**)

Description and comments:

The nuchal plate (**Fig. 08: D**) is partly broken but was likely longer than wide in dorsal view. In lateral view, the dorsal margin of the bone forms a 110° angle. In dorsal view, the plate presents long anterolateral margins crossed at their mid-length by the anterior sulcus of the 1st vertebral scute. The lateral sulcus of the 1st vertebral scute is not visible. The cervical scute is quadrangular and marked by a well-defined indentation of the anterior margin of the bone. In ventral view, the cervical scute is nearly square. This nuchal plate has been attributed to a juvenile individual of *Manouria* based on the combination of the following characteristics only encountered in this genus: nuchal plate longer than wide, and occurrence of a small quadrangular cervical scute (**Fig. S11, and S12**).

The two peripheral plates attributed to *Manouria* sp. (**Fig. 08: E**) present a very unusual morphology. They both have a sinusoidal ventral margin with a medial bulge. Additionally, they bear a peculiar pathological condition; their dorsal surface exhibits a scaly appearance formed of deep irregular furrows that mask the sulci of the marginal scutes. These two pieces have been associated with *Manouria* based on the observation of similar elements on a modern specimen of *Manouria* sp. However, this modern specimen did not present the pathological condition observed in these fossils.

The pygal plate (**Fig. 08: F**) is relatively thin and has a trapezoidal shape in posterior view. The posterior sulcus of the 5th vertebral overlaps with the dorsal limit of the plate, the 12th marginal inter-sulcus is present, and a small notch is visible on the central part of the ventral margin of the bone. In our comparative sample, the closest correspondence we found is *Manouria emys*. However, the pygal plate of *M. emys* is always wider than high in juvenile and adult specimens, and the shape of the bone is less triangular than our fossil. As we lack appropriate references for *M. impressa* to rule out the possibility that this plate might be attributed to a modern species of *Manouria*, we tentatively refer this plate to an undefined species of *Manouria*.

The ventral face of the only subfossil entoplastron attributed to *Manouria* sp. is crossed by the medial sulcus of the humeral and pectoral scutes but does not exhibit any other sulci. This condition has only been observed in *Manouria* (**Fig. S11, and S12**). This plate tends to be as wide as long in adult individuals but is more elongated in juvenile individuals, a condition observed in our subfossil element.

A humeral diaphysis can be associated with *Manouria* due to its large size and its morphological similarity to this taxon. It exhibits limited torsion and a well-developed medial process (*sensu* Hermanson et al., 2024) in a vertical position parallel to the main proximo-distal axis of the diaphysis.

Figure 08 - A) Costal plate of *Manouria impressa* in dorsal view (Laang Spean); **B**) Costal plate of *Manouria impressa* in lateral view (Doi Pha Kan); **C**) Pygal plate of *Manouria impressa* in anterior (right) and posterior view (left) (Doi Pha Kan); **D**) Nuchal plate of *Manouria* sp. in dorsal view (Doi Pha Kan); **E**) Peripheral plates of *Manouria* sp. in lateral view (Laang Spean); **F**) Pygal plate of *Manouria* sp. in anterior (right) and posterior (left) views (Laang Spean). <u>Abbreviations</u>: **b.**: bulge; **C.**: cervicale scute; **e.**: elevation; **M1**: 1st marginal scute; **M12**: 12th marginal scute; **n.**: notch; **V1**: 1st vertebral scute; **V5**: 5th vertebral scute. Scalebar=1cm.

Testudinidae indet. Batsch, 1788– Figure. 09

Material: One nuchal plate from Doi Pha Kan. (**Table. 01**)

Descriptions and comments:

The nuchal plate (**Fig. 09**) is pentagonal in dorsal view. In lateral view, the ventral margin of the
bone forms a 110° angle as its posterior part projects dorsally. The cervical scute is absent in both
dorsal and ventral views. In dorsal view, the anteromedial margins of the 1^{st} pleural scutes are
visible in the posterior half of the bone. The anterior margin of the plate is notched at the level of
the sulcus between the 1^{st} marginal scutes. On the unbroken right side of the plate, the 1^{st} marginal
scute extends well posteriorly and nearly reaches the suture between the 1^{st} peripheral and 1^{st}
costal plates. In ventral view, sulcus between the 1^{st} marginal scutes extends over more than half
the length of the plate. The contact with the first peripheral plate is well-extended, and longer than
the contact with the costal plate.

Several characters support the attribution of this fossil to the family Testudinidae: the absence
of a cervical scute, the dorsal posterior extension of the marginal that nearly reaches the suture
between the 1^{st} peripheral and 1^{st} costal plates, and the visceral extension of the marginal scute.
The first character is found in several species of Testudinidae, while the second and third are
present in all Testudinidae. On the other hand, in most Testudinidae and all its Asian forms, the
pleural scutes rarely cover the nuchal to a large extent, which is a character more frequently
observed in Geoemydidae. However, the absence of a cervical scute is known to occur very rarely
and only in a few taxa of Geoemydidae (*Heosemys*, *Cuora*) (J. C. pers. obs.). The absence of a
cervical scute is also found in Podocnemididae Cope 1869, a family currently present only in
Madagascar and South America and in the tertiary fossil record of Eurasia (Rhodin et al., 2021).
Some species of this group may have been present until the Late Miocene in continental Southeast
Asia (Hirayama et al., 2023), but the nuchal plate is usually flat in visceral view for this group while
it shows a concavity for Cryptodira. Assuming this plate belongs to a Testudinidae, the comparison
with the two modern Southeast Asian species lacking a cervical scute, namely *Geochelone*
platynota and *Indotestudo travancorica*, is rather inconclusive. In *I. travancorica*, the nuchal plate
is flatter, its anterior margin is not strongly indented, and the posterior margin presents a well-
marked concavity for the 1^{st} neural plate. In *G. platynota*, the medial contact between the 1^{st}
marginals is much shorter than in our fossil, and the pleural scute does not or not strongly overlap
the nuchal plate. Therefore, we attribute our nuchal plate to an unknown Testudinidae, pending for
new material in the Doi Pha Kan locality.

Regarding previously published Quaternary fossils, Claude et al. (2019) mentioned the
presence of another testudinid in the central plain of Thailand, different from *Manouria* and
Indotestudo, tentatively assigning this material to *Geochelone platynota* based on geographic
arguments. It is possible that the material we attribute to *Geochelone* sp. and Testudinidae indet.
could be related to the same taxon. This or these extinct Testudinidae might have been present in
Thailand and Cambodia during the Holocene and possibly the terminal Pleistocene.

Figure 09 - Nuchal plate of *Testudinidae* indet. in dorsal (left) and ventral (rigth) views (Doi Pha
Kan). <u>Abbreviations</u>: **M1**: 1st marginal scute; **P1**: 1st pleural scute; **V1**: 1st vertebral scute. Scalebar=1cm.

Trionychidae Gray, 1825

Amyda sp. Geoffroy Saint-Hilaire, 1809 – Figure. 10

Material: 15 remains from Doi Pha Kan, Khao Ta Phlai, Laang Spean, and Moh Khiew: one
neural plate, 11 costal or unidentified carapace plates, one hyoplastron, one xiphiplastron, and one
scapula. (**Table. 01**)

Descriptions and comments:

One neural plate (**Fig. 10: A**) has been tentatively attributed to *Amyda* sp. The plate is complete,
measuring 2.5 cm in length, and its dorsal surface is covered with irregular pits. The maximum
number of pits across the width of the plate is six, which is consistent with what is observed in other
large trionychids. The pits appear shallower than those in the genus *Chitra* and do not become
more elongated near the anterior or posterior margins of the bone.

Six fragments of costal or unidentified plates (**Fig. 10: B and C**) have been attributed to *Amyda*
based on the morphology of their dermal ornamentation. This ornamentation consists of
vermiculated sculptures and sharp ridges, both forming well-defined, irregular polygonal
depressions across all external surfaces of the bones. Some elements also exhibit longitudinal
ridges of higher terrain (sensu Pritchard et al. 2009). The ornamentation pattern remains consistent
across all surfaces, showing little variation near the edges of the bones, except at the lateral
margins of the costal plates, where the polygons become less defined and shallower. The presence
of longitudinal ridges is characteristic of *Amyda* among the genera currently living in Thailand but
is also observed in the southeast asian genera *Palea* and *Pelodiscus*. In contrast, the
ornamentation patterns of other Thai taxa differ: in *Pelochelys* and *Chitra*, the ornamentation is less
random and changes near the anterior and posterior sutures of the costal plates. Additionally, in
Pelochelys, the ornamentation pattern remains consistent near the lateral margins of the costal
plates (Pritchard et al., 2009).

A fragment of the medial side of a hyoplastron (**Fig. 10: D**) can be attributed to *Amyda* sp. The
fragment is 4 cm wide, but the total width is estimated to be approximately 9 to 10 cm. The bone is
narrow in its axillary region. The ventral surface is covered by a callosity ornamented with small
ridges and punctuations. The medial area features three or four well-developed spikes that are

closely packed and project in the same anteromedial direction. Among the taxa currently present in Thailand, this fragment can only be referred to *Amyda*. In *Pelochelys*, the axillary region of the hyoplastron is broad, while the medial spikes are either absent or poorly developed in *Dogania* and oriented in varying directions, from anteromedial to medial, in *Chitra*.

One left xiphiplastron (**Fig. 10: E**), missing a small portion of its anterior side, has been attributed to *Amyda* sp. The bone is triangular in shape, approximately 8 cm in length, with a strongly concave lateral side. On the medial side, spikes and sockets are present for articulation with the right xiphiplastron. The ventral surface features a callosity that covers most of the area and is ornamented with small, vermiculated ridges running parallel to the lateral side. This ornamentation becomes less prominent and fades near the bone's margins. The triangular shape of the bone, the presence of medial spikes and sockets, the lateral concavity, and the morphology of the ornamentation are diagnostic of the genus *Amyda*.

One scapula (**Fig. 10: F**), with an angle of approximately 60° between its ventral and dorsal branches, has been tentatively attributed to *Amyda* due to the absence of distinguishing features separating it from this species. However, a detailed comparison with other Trionychidae taxa currently living in Thailand was not conducted.

Figure 10 - A) Neural plate of *Amyda* sp. in dorsal view (Laang Spean); **B**) Coastal plate of *Amyda* sp. in dorsal view (Doi Pha Kan); **C**) Coastal plate of *Amyda* sp. in lateral view (Laang Spean); **D**) hyoplastron of *Amyda* sp. in ventral view (Moh Khiew); **E**) Xiphiplastron of *Amyda* sp. in ventral view (Moh Khiew); **F**) Scapula of *Amyda* sp. in anterior view (Moh Khiew); **G**) Coastal plate of *Dogania* sp. in

lateral view (Moh Khiew). <u>Abbreviations</u>: **a. r.**: axillary region; **d. b.**: dorsal branch; **s.**: spikes; **l. c.**: lateral
convexity; **v. b.**: ventral branch. Scalebars=1cm.

Dogania sp. Gray, 1844 – Figure. 10

Material: 17 remains from Moh Khiew: one neural plate, 13 coastal plates, one hyoplastron,
and two unidentified carapace plates. (**Table. 01**)

All the remains attributed to *Dogania* exhibit a very distinctive dermal ornamentation pattern
(**Fig. 10: G**), which is much finer than that of *Amyda* and other Southeast Asian taxa. This pattern
consists of very small, deep depressions of irregular shape, along with small tubercles located in
the distal area of the costal plates (Pritchard et al., 2009).

Trionychidae indet. Gray, 1825

Material: 57 remains distributed in the four sites: 33 costal plates, one epiplastron, two
unidentified carapace plates, one dentary, one coracoid, three scapula, six humeri, one radius, two
ilia, one pubis, four femora, and two tibias. **(Table. 01)**

Descriptions and comments:

These bones correspond to either fragmented elements or anatomical parts whose morphology
was not distinctive enough to be attributed to a specific genus, or for which appropriate comparative
references were not available. However, they were attributed to Trionychidae based on direct
comparisons with several members of this family (postcranial and cranial elements) and on the
presence of ornamentation that, although not well-preserved, was insufficient to permit a subfamily-
level attribution (carapace elements).

Distribution of the species in the investigated assemblages (**Table. 01**)

Regarding the Geoemydidae: the genus *Batagur* has been identified only at the site of Laang
Spean in a Neolithic context. The genera *Cuora* (*Cuora couro*) and *Cyclemys* have been identified
in all four assemblages. *Heosemys* have been identified in all sites except in northern Thailand (Doi
Pha Kan), with two or three species represented (*H. spinosa*, *H. grandis*, and H. *annandalii*). The
genus *Malayemys* was identified in Moh Khiew, Doi Pha Kan and Khao Ta Phlai. For the family
Testudinidae, *Indotestudo elongata* was present in all the assemblages (Bochaton et al., 2023).
Additional Testudinidae species were identified in the present study in Doi Pha Kan and Laang
Spean, with *Manouria impressa*, *Manouria* sp., and at least one undetermined species attributed
to Testudinidae sp. in Doi Pha Kan and to *Geochelone* sp. in Laang Spean. Regarding Laang
Spean we observed no chronological tendency in the distribution of the identified species. For the
Trionychidae: the genus *Amyda* was identified in Laang Spean, Khao Ta Phlai, Doi Pha Kan and
Moh Khiew, and the genus *Dogania* only in Moh Khiew. Unidentified Trionychidae were also
identified in the four sites.

Species	Doi Pha Kan	Khao Ta Phlai	Laang Spean	Moh Khiew
Geoemydidae				
Batagur sp.			1	
Cuora couro	23	30	74	150
Cuora sp.				1
Cyclemys cf. *oldhamii*	229	76	115	69
Heosemys annandalii		1	2	
Heosemys grandis			5	
Heosemys spinosa		1		16
Heosemys cf. *grandis/annandalii*		87	54	38
Malayemys sp.	1	1		3
Geoemydidae indet.	190	136	149	292
Testudinidae				
Geochelone sp.			1	
Manouria impressa	5		4	
Manouria sp.	1		5	
Testudinidae indet.	1			
Trionychidae				
Amyda sp.	3	5	2	7
Dogania sp.				17
Trionychidae indet.	10	14	13	15

Table 1 - Distribution of the identified species in the different assemblages in terms of Number of Remains. The complete account of the turtle bone remains found at the sites is published in Bochaton et al. (2023). The small discrepancy in the bone numbers corresponds to bones ultimately moved to the category of non-identified turtle bone remains between the two studies.

Discussion

Meaning of the archaeological accumulation in terms of past biodiversity

Before discussing the significance of the data provided by paleontological analyses in terms of past biodiversity, it is necessary to address the question of the significance of these data and the mode of accumulation of the studied archaeological bone assemblages. Bone samples do not represent a frozen image of the surrounding biodiversity; they provide distorted images that reflect more on how the bone assemblage was created than the biodiversity of the surrounding animals at the site. To understand and interpret the biodiversity data provided by a bone sample, it is first necessary to understand how it was formed, in an attempt to identify, if not correct, its biases. To achieve this, taphonomic analyses and, in the case of archaeological sites, zooarchaeological analyses are needed. Such analyses were not conducted in the present study but have been performed in previous studies (Auetrakulvit, 2004; Forestier et al., 2015; Frère et al., 2018; Bochaton et al., 2019, 2023). Regarding the turtle remains, these studies unambiguously demonstrate that they correspond to animals gathered by prehistoric hunters for consumption and that they were therefore transported over some distance. The potential contribution of additional accumulation agents (i.e., birds or carnivores) is insignificant. This has major implications, as all the specimens present at the site are of relatively large size, a compositional bias that could be amplified by the bone recovery protocol in the field (see Bochaton et al., 2023). In the four sites, the hunters focused their gathering efforts on a single species, *Indotestudo elongata*. The methods

that might have been used to trap these tortoises could include trapping (Santos et al., 2020;
Bochaton et al., 2023), similar to the methods possibly used for other small animals like monitor
lizards (Bochaton et al., 2019). The same traps could have also caught *Cuora* individuals, which
are often active outside of water or can be found in shallow water. However, this cannot account
for the more aquatic species, including *Cyclemys*, which is well-represented in the assemblages.
This probably indicates the use of fishing techniques, corroborated by the presence of strictly
aquatic fauna (mollusks and fish) in all the sites (Auetrakulvit, 2004; Forestier et al., 2015; Frère et
al., 2018; Bochaton et al., 2019, 2023). Nevertheless, the exploitation of freshwater fauna was
clearly secondary in all the assemblages. This could be related to the distance to freshwater
sources. In addition to a likely lower representation of aquatic species, the studied bone samples
clearly suffer from a near absence of large individuals. As far as turtles are concerned, this
phenomenon is probably explained by the weight of these animals relative to their meat yield.
Indeed, transporting these animals was probably not easy, but they are also simpler to keep captive
and alive than mammals or birds and could thus be kept in the settlement for future consumption.
Large animals that were impossible to move might have been excluded from the archaeological
assemblages because their carcasses could have been left at the killing site, with only their meat
carried to the settlement. Conversely, small animals might not have been collected because they
did not represent an abundant enough source of meat to justify their transport and culinary
preparation. Of course, hunting strategies could also involve less straightforward and practical
considerations, such as ritual hunting (Balée, 1985), which are impossible to address with the
archaeological record. Overall, it is clear that the archaeological assemblages are not a faithful
representation of past turtle biodiversity, as aquatic animals are poorly represented and large and
small animals are excluded. Despite these biases, it is interesting to observe that the coverage of
freshwater species in the different sites seems fairly good. Indeed, the species in the sites reveal
species associations similar to those observed today in most freshwater environments in
continental Southeast Asia (*Amyda ornata*, *Cuora couro*, *Cyclemys oldhamii*, *Heosemys*
annandalii, *Heosemys grandis*, and *Siebenrockiella crassicollis*) in Cambodia (Platt et al., 2008),
Vietnam (Le, 2007), and Laos (Brakels, 2018).

Biodiversity Changes Information Provided by the Studied Archaeological Samples

The taxa identified at the investigated sites provide information about the local changes of turtle
fauna across the Holocene. These can be categorized into species that were: I) not or weakly
impacted, II) species whose distribution shrank, and III) species that went extinct.

I) Weakly Impacted Species (N=7): *Manouria impressa* is still present today in the two areas
where it has been identified (northern Thailand and southwest Cambodia) (Rhodin et al., 2021)
(Figure 11-A). This turtle is a terrestrial species living in mountainous forests, mostly active during
the hot and rainy season, and primarily consumes mushrooms (Wanchai et al., 2013; Cota et al.,
2021). Regarding its occurrence in Laang Spean, it is interesting to note that the site is not in an
elevated area, which could indicate a past occurrence of *M. impressa* in low-elevation forests that
no longer exist nowadays, but the data are currently too scarce to investigate this question. The
same remarks could be made about *Manouria emys* (Claude et al., 2019). The species of
Heosemys identified in the samples are all present in the vicinity of the sites today (Figure 11-B).
Heosemys grandis and *H. annandalii* are freshwater species commonly found in swamps, while *H.*
spinosa is a lowland forest species (Sharma & Tisen, 2000; Das, 2010). Some other taxa might
have been impacted but only in part of their range. For instance, *Cuora couro* (Figure 11-C) and
Malayemys sp. (Figure 11-D) are both still present today in the areas of the sites where they have
been identified but not in Laang Spean. However, both of these taxa are still present a few dozen
kilometers away from this last site, which indicates that if their distribution area has effectively
shrunk, this might be a very recent phenomenon. *Cuora couro* is a semi-aquatic species inhabiting
many types of lowland freshwater habitats. It is mostly active at night and can feed both in water
and on land (Schoppe & Das, 2015). *Malayemys* is a genus of aquatic species that estivate on land

in forests and occupy a variety of freshwater habitats (Dawson et al., 2018, 2020). Regarding
Trionychidae, we identified *Dogania* inside its modern range in Southern Thailand (Fig. 12-B).

II) Species Whose Distribution Area Shrunk (N=4): The distribution of *Cyclemys* cf. *oldhamii*
may have undergone significant changes, as it is now absent in the vicinity of all the sites, although
it has been recorded a few dozen kilometers away from Laang Spean, Khao Ta Phlai, and Moh
Khiew (Figure 11-E). This could reflect either shrink of distribution, lack of survey of modern
species, or transportation by prehistoric populations. However, in Doi Pha Kan, the distance to the
modern distribution area is around 100 km, making the hypothesis of a reduction in the distribution
area of this taxon more likely although it might still be related to a lack of biodiversity survey in that
area. A similar observation can be made for our finding of *Batagur* in Laang Spean (Figure 11-F).
Batagur affinis (Cantor 1847) is the only species of the *Batagur* genus currently present in
Cambodia. It inhabits the floodplain of the Tonle Sap, which is only 50 km away from Laang Spean
(Moll et al., 2015). It is an aquatic species living in estuaries and tidal regions of large rivers (Moll
et al., 2015; Thomson, 2021). The distribution of these two species has undoubtedly shrunk since
the early/middle Holocene. *Indotestudo elongata*, previously mentioned in Laang Spean (Bochaton
et al., 2023) but not described in this paper, should also be added to this list, as its closest modern
occurrence is around 50 km away from the site. Regarding the softshell turtle *Amyda* (Fig. 12-A),
we identified its remains in Doi Pha Kan in Northern Thailand, over 150 km from its closest modern
range near the Myanmar border (Rhodin et al., 2021) but, although this genus might be rare and
absent in several areas of northern Thailand, its complete absence could reflect a lack of
documentation in several areas (J. C. pers. obs). *Amyda* was also identified in Khao Ta Phlai
between the modern distributions of *A. ornata* and *A. cartilaginea* (Fig. 12) in an area where the
genus is absent nowadays. Bones of this genus have also been found at the Laang Spean site,
although *Amyda* is still present just a few kilometers from this site today. Based on its modern
range, it is likely that *Amyda* has experienced a significant reduction in its distribution area in
Northern Thailand.

III) Species That Went Extinct (N=1 or 2): The only clear extinction, or at least a massive
reduction in distribution, is associated with the one or two species of Testudinidae (*Geochelone* sp.
and Testudinidae indet.) identified in the archaeological samples. Whether the material is attributed
to *Geochelone* or another Testudinidae taxon, it indicates that at least one species of this family
was present in Thailand and Cambodia in the past and that it completely disappeared over the
course of the Holocene. Such event is unlikely to reflect an overexploitation by human groups as
the remains of these turtles are very scarce in the archaeological assemblages. There is currently
not a single extinction of turtle species documented in Southeast Asia during the Late Pleistocene
and the Holocene (Rhodin et al., 2015).

Figure. 11 - Modern distribution maps of the Testudinidae and Geoemydidae species identified in
the four archaeological assemblages, along with the locations of the archaeological sites that provided
remains of the concerned taxa. The distribution data are from Rhodin et al. (2021) and Chaianunporn
et al., (2025) for *Malayemys.*

Figure. 12 - Modern distribution maps of the Trionychidae species identified in the four
archaeological assemblages, along with the locations of the archaeological sites that provided remains
of the concerned taxa. The distribution data are from Rhodin et al. (2021).

Has Southeast Asian Turtle Biodiversity Changed in the Last 10,000 Years?

The scarcity of available data regarding past turtle biodiversity in Southeast Asia currently
precludes detailed analyses, but some interesting observations can be made. Our results clearly
indicate that the different investigated areas had varying trajectories concerning the evolution of
their turtle biodiversity, as the impacted or non-impacted species were not randomly distributed
across the different sites. The two sites in southern Thailand show a turtle fauna that was very
close to the modern one, with only one species that may have suffered a limited reduction in its
distribution range (*Cyclemys* sp.). The situation is similar in northern Thailand (Doi Pha Kan), even
though Trionychidae turtles seem to have disappeared from that area, and an unknown extinct
testudinid species is present in this deposit. It is currently impossible to link this extinction or change
of distribution to a specific phenomenon, as it is presently impossible to estimate when this species
disappeared. Aside from this observation, the turtle fauna of Thailand appears to exhibit remarkable
stability, which stands in stark contrast to the observations made in the central plain of Thailand
(Claude et al., 2019). In this latter region, at least two forest taxa and one large freshwater turtle
may have been extirpated following environmental changes and deforestation in the second half of
the Holocene. This suggests that not all areas of Thailand were impacted in the same way.
Regarding the sites we studied, the only locality that truly stands out in our sample is Laang Spean,
where seven of the eleven species we identified (Table. 01 – *Batagur* sp., *Cuora couro*, *Cyclemys*
sp., *Geochelone* sp., *Indotestudo elongata*, and *Amyda* sp.) are absent from the surrounding area
(Figs. 11 and 12).

We believe that the biodiversity modifications observed at the sites reflect recent human impact
rather than deep historical alterations of the environments, and this is for two reasons: 1) the
distribution changes observed are mostly minor, typically only a few dozen kilometers in most
cases, and 2) the fact that the Cambodian site shows the most significant changes in its turtle fauna
compared to the sites in Thailand can be easily explained by the magnitude of the very recent
environmental modifications these two countries have experienced. Indeed, compared to Thailand,
the impact of deforestation has been much more severe in Cambodia. Thailand's forest cover was
estimated to be about 70% of the country's surface in 1930 (35.7 million ha), shrinking to around
29.6 million ha in 1952 and about 17.8 million ha in 1978 (Ramitanondh, 1989). This corresponds
to an annual deforestation rate of r=1.4%. However, the deforestation rate has significantly
decreased since then, being reduced by more than tenfold between 1990 and 2016, and
maintaining an annual rate of r=0.028% until at least 2020 (Puyravaud, 2003; FAO, 2020a; b). In
Cambodia, the situation is very different. Forest cover represented 73.3% of the country's area in
the 1960s (13.3 million ha) but decreased to 10.937 million ha by 1990 (Tsujino et al., 2019; FAO,
2020c), corresponding to an annual deforestation rate of r=0.6%. The subsequently documented
annual deforestation rate was r=0.23% between 1990 and 2010, but this rate dramatically
increased to r=4.5% between 2010 and 2015, and r=2.1% between 2015 and 2020 (Puyravaud,
2003; FAO, 2020b; c). This situation is already well known, and the future of Cambodia's
biodiversity is now the most concerning among continental Southeast Asian countries (Botterill-
James et al., 2024). Laang Spean, being located outside of a protected area, has suffered massive
and very fast environmental changes in recent decades. This observation, if not related to an overall
paucity of biodiversity inventory in Cambodia, could be a striking demonstration of the quick effect
of massive environmental alterations that could wipe out taxa proven to be strongly resilient even
in disturbed areas of Thailand.

Overall, our data point to the resilience of turtle assemblages outside the areas that have been
heavily anthropized, such as the central plain of Thailand and Cambodia. In these two regions, the
extent of environmental modifications has exceeded the adaptation capacity of the taxa, leading to
a significant reduction in local biodiversity. That said, other factors might also be at play, such as
climatic events and human-induced changes in the distant past. These phenomena are
characterized by significant reductions in geographic distribution and even the extinction of some
rare taxa, but these events cannot be accurately dated and thus characterized at this time.
However, these impacts appear to be limited, as most local species assemblages have persisted
since the Late Pleistocene/Early Holocene without major modifications. This could also be the case
in other Southeast Asian area. For instance, in the Late Pleistocene archaeological layers of Niah
Cave in Sarawak, the three species identified in the second phase and the three species identified
in the first phase are still extant in the vicinity of the site today (Pritchard et al., 2009). Only the
study of additional bone assemblages and well-dated occurrences of critically important taxa will
enhance our understanding of the causes behind these modifications.

1026

Conclusion

Our observations indicate that the turtle assemblages in northern and southern Thailand, and
southwest Cambodia, have experienced only limited changes across the Holocene, with the
exception of Laang Spean in Cambodia. We also raise questions about the existence of one or
several potentially extinct species of Testudinidae in continental Southeast Asia at an unknown
time during the Holocene. Additionally, we observed significant reductions in geographic
distribution affecting some species, such as *Batagur*, *Cyclemys*, and *Amyda*. Unfortunately, it is
currently difficult to determine the timing and causes of the biodiversity changes we observed, and
we have had to rely on probabilistic hypotheses. To address these uncertainties, it is now
necessary to improve the quality of the available paleontological and zooarchaeological records,

particularly for more recent periods (historical and modern). Radiometric dating of the remains of critically important taxa will be essential to clarify their periods of occurrence in different areas. This should be done in parallel with the enhancement of modern reference collections and possibly molecular approaches to identify past taxonomic diversity in Southeast Asia as reliably as possible. In conclusion, although the results described here show that it is not too late to preserve the native biodiversity of several areas in continental Southeast Asia, our observations serve as a warning of the potential fate of turtle biodiversity in this region if further loss of natural environments, such as what is currently happening in Cambodia, is not prevented. We also emphasize the importance of biodiversity surveys to accurately track the evolutionary dynamics of species facing environmental disturbances.

Funding

This work has been done thanks to the support of several funding agencies: the FYSSEN foundation, the DIM-MAP funded by the Region île-de-France, the French Ministry of Europe and Foreign Affairs, and the IRN PalBioDivASE 0846 funded by the CNRS. It was also supported by the Mission Préhistorique FrancoCambodgienne, the Mission Préhistorique Franco-Thaïe, and the Mission Paléolithique Franco-Thaïe of the French Ministry of Europe and Foreign Affairs (MEAE, Paris).

Conflict of interest disclosure

The authors declare that they comply with the PCI rule of having no financial conflicts of interest in relation to the content of the article.

Author contributions

Conceptualization: Corentin Bochaton ; **Data curation**: Corentin Bochaton ; **Formal analysis** : Corentin Bochaton, Julien Claude ; **Funding acquisition** : Corentin Bochaton, Valéry Zeitoun, Hubert Forestier ; **Investigation** : Corentin Bochaton, Wilailuck Naksri, Sirikanya Chantasri, Melada Maneechote, Supalak Mithong, Julien Claude ; **Methodology** : Corentin Bochaton ; **Resources** : Sophady Heng, Hubert Forestier, Prasit Auetrakulvit, Valéry Zeitoun, Jutinach Bowonsachoti ; **Supervision** : Corentin Bochaton ; **Visualization**: Corentin Bochaton; **Writing – original draft**: Corentin Bochaton, Julien Claude; **Writing – review & editing**: Corentin Bochaton, Julien Claude.

References

Ascarrunz E, Claude J, Joyce WG (2021) The phylogenetic relationships of geoemydid turtles from the Eocene Messel Pit Quarry: a first assessment using methods for continuous and discrete characters. *PeerJ*, **9**, e11805. https://doi.org/10.7717/peerj.11805

Auetrakulvit P (2004) Faunes du pléistocène final à l'holocène de Thaïlande : approche archéozoologique. PhD thesis Thesis. Aix Marseille 1, Marseille.

Auetrakulvit P, Forestier H, Khaokhiew C, Zeitoun V (2012) New excavation at MohKhiew site (Southern Thailand). In: *Crossing Borders in Southeast Asian Archaeology* (eds Bonatz D, Reinecke A, Tjoa-Bonatz ML), pp. 62–74. Singapore.

Baker C, Phongpaichit P (2014) *A History of Thailand*. Cambridge University Press, Cambridge.

Balée W (1985) Ka'apor ritual hunting. *Human Ecology*, **13**, 485–510.

Barlow J, Gardner TA, Lees AC, Parry L, Peres CA (2012) How pristine are tropical forests? An ecological perspective on the pre-Columbian human footprint in Amazonia and implications for contemporary conservation. *Biological Conservation*, **151**, 45–49. https://doi.org/10.1016/j.biocon.2011.10.013

Bochaton C, Chantasri S, Maneechote M, Claude J, Griggo C, Naksri W, Forestier H, Sophady H, Auertrakulvit P, Bowonsachoti J, Zeitoun V (2023) Zooarchaeological investigation of the Hoabinhian exploitation of reptiles and amphibians in Thailand and Cambodia with a focus on the Yellow-headed Tortoise (*Indotestudo elongata* (Blyth, 1854)). *Peer Community Journal*, **3**, 1–32. https://doi.org/10.24072/pcjournal.328

Bochaton C, Claude J, Auetrakulvit P, Zeitoun V (2024) The past vertebrate biodiversity of Southeast Asia and its relevance to ongoing environmental issues: current limitations and perspectives. In: *On the edge of sixth extinction in biodiversity hotspots: Facts, needs, solutions and opportunities in Thailand and adjacent countries* (eds Claude J, Kitana N), pp. 255–280. CU Press.

Bochaton C, Hanot P, Frère S, Claude J, Naksri W, Auetrakulvit P, Zeitoun V (2019) Size and weight estimations of subfossil monitor lizards (Varanus sp. Merrem 1820) with an application to the Hoabinhian assemblage of Doi Pha Kan (Late Pleistocene, Lampang province, Thailand). *Annales de Paléontologie*, **4**, 295–304. https://doi.org/10.1016/j.annpal.2019.05.003

Böhm M, Collen B, Baillie JEM, Bowles P, Chanson J, Cox N, Hammerson G, Hoffmann M, Livingstone SR, Ram M, Rhodin AGJ, Stuart SN, van Dijk PP, Young BE, Afuang LE, Aghasyan A, García A, Aguilar C, Ajtic R, Akarsu F, Alencar LRV, Allison A, Ananjeva N, Anderson S, Andrén C, Ariano-Sánchez D, Arredondo JC, Auliya M, Austin CC, Avci A, Baker PJ, Barreto-Lima AF, Barrio-Amorós CL, Basu D, Bates MF, Batistella A, Bauer A, Bennett D, Böhme W, Broadley D, Brown R, Burgess J, Captain A, Carreira S, Castañeda M del R, Castro F, Catenazzi A, Cedeño-Vázquez JR, Chapple DG, Cheylan M, Cisneros-Heredia DF, Cogalniceanu D, Cogger H, Corti C, Costa GC, Couper PJ, Courtney T, Crnobrnja-Isailovic J, Crochet P-A, Crother B, Cruz F, Daltry JC, Daniels RJR, Das I, de Silva A, Diesmos AC, Dirksen L, Doan TM, Dodd Jr. CK, Doody JS, Dorcas ME, Duarte de Barros Filho J, Egan VT, El Mouden EH, Embert D, Espinoza RE, Fallabrino A, Feng X, Feng Z-J, Fitzgerald L, Flores-Villela O, França FGR, Frost D, Gadsden H, Gamble T, Ganesh SR, Garcia MA, García-Pérez JE, Gatus J, Gaulke M, Geniez P, Georges A, Gerlach J, Goldberg S, Gonzalez J-CT, Gower DJ, Grant T, Greenbaum E, Grieco C, Guo P, Hamilton AM, Hare K, Hedges SB, Heideman N, Hilton-Taylor C, Hitchmough R, Hollingsworth B, Hutchinson M, Ineich I, Iverson J, Jaksic FM, Jenkins R, Joger U, Jose R, Kaska Y, Kaya U, Keogh JS, Köhler G, Kuchling G, Kumlutaş Y, Kwet A, La Marca E, Lamar W, Lane A, Lardner B, Latta C, Latta G, Lau M, Lavin P, Lawson D, LeBreton M, Lehr E, Limpus D, Lipczynski N, Lobo AS, López-Luna MA, Luiselli L, Lukoschek V, Lundberg M, Lymberakis P, Macey R, Magnusson WE, Mahler DL, Malhotra A, Mariaux J, Maritz B, Marques OAV, Márquez R, Martins M, Masterson G, Mateo JA, Mathew R, Mathews N, Mayer G, McCranie JR, Measey GJ, Mendoza-Quijano F, Menegon M, Métrailler S, Milton DA, Montgomery C, Morato SAA, Mott T, Muñoz-Alonso A, Murphy J, Nguyen TQ, Nilson G, Nogueira C, Núñez H, Orlov N, Ota H, Ottenwalder J, Papenfuss T, Pasachnik S, Passos P, Pauwels OSG, Pérez-Buitrago N, Pérez-Mellado V, Pianka ER, Pleguezuelos J, Pollock C, Ponce-Campos P, Powell R, Pupin F, Quintero Díaz GE, Radder R, Ramer J, Rasmussen AR, Raxworthy C, Reynolds R, Richman N, Rico EL, Riservato E, Rivas G, da Rocha PLB, Rödel M-O, Rodríguez Schettino L, Roosenburg WM, Ross JP, Sadek R, Sanders K, Santos-Barrera G, Schleich HH, Schmidt BR, Schmitz A, Sharifi M, Shea G, Shi H-T, Shine R, Sindaco R, Slimani T, Somaweera R, Spawls S, Stafford P, Stuebing R, Sweet S, Sy E, Temple HJ, Tognelli MF, Tolley K, Tolson PJ, Tuniyev B, Tuniyev S, Üzüm N, van Buurt G, Van Sluys M, Velasco A, Vences M, Veselý M, Vinke S, Vinke T, Vogel G, Vogrin M, Vogt RC, Wearn OR, Werner YL, Whiting MJ, Wiewandt T, Wilkinson J, Wilson B, Wren S, Zamin T, Zhou K, Zug G (2013) The conservation status of the world's reptiles. *Biological Conservation*, **157**, 372–385. https://doi.org/10.1016/j.biocon.2012.07.015

Botterill-James T, Yates LA, Buettel JC, Brook BW (2024) The future of Southeast Asia's biodiversity: A crisis with a hopeful alternative. *Biological Conservation*, **296**, 110641. https://doi.org/10.1016/j.biocon.2024.110641

Bourret R (1941) *Les tortues de l'Indochine*. Institut Océanographique de l'Indochine, Vietnam.

Brakels P (2018) *Assessment of the Distribution, Status and Conservation Priorities of Freshwater Turtles from Xe Champhone and Beung Kiat Ngong Ramsar Sites, Lao PDR*. IUCN, Vientiane, Laos.

Ceballos G, Ehrlich PR (2018) The misunderstood sixth mass extinction. *Science*, **360**, 1080–1081. https://doi.org/10.1126/science.aau0191

Ceballos G, Ehrlich PR (2023) Mutilation of the tree of life via mass extinction of animal genera. *Proceedings of the National Academy of Sciences*, **120**, e2306987120. https://doi.org/10.1073/pnas.2306987120

Ceballos G, Ehrlich PR, Barnosky AD, García A, Pringle RM, Palmer TM (2015) Accelerated modern human–induced species losses: Entering the sixth mass extinction. *Science Advances*, **1**, e1400253. https://doi.org/10.1126/sciadv.1400253

Chaianunporn T, Mokarat M, Ruangsakunporn A, Kutanan W, Lomthaisong K, Chaianunporn K, Keithmaleesatti S (2025) Distribution Ranges of Snail-Eating Turtle Species (*Malayemys* spp.) in Northeastern and Eastern Thailand. *Chelonian Conservation and Biology*, **23**, 216–227. https://doi.org/10.2744/CCB-1515.1

Chaimanee Y (1998) *Plio-pleistocene rodents of Thailand*. National Center for Genetic Engineering and Biotechnology, Bangkok, Thailand.

Claude J, Auetrakulvit P, Naksri W, Bochaton C, Zeitoun V, Tong H (2019) The recent fossil turtle record of the central plain of Thailand reveals local extinctions. *Annales de Paléontologie*, **4**, 305–315. https://doi.org/10.1016/j.annpal.2019.04.005

Conrad C (2015) Archaeozoology in Mainland Southeast Asia: Changing methodology and Pleistocene to Holocene forager subsistence patterns in Thailand and peninsular Malaysia. *Open Quaternary*, **1**. https://doi.org/10.5334/oq.af

Conrad C, Higham C, Eda M, Marwick B (2016) Palaeoecology and Forager Subsistence Strategies during the Pleistocene – Holocene Transition: A Reinvestigation of the Zooarchaeological Assemblage from Spirit Cave, Mae Hong Son Province, Thailand. *Asian Perspectives*, **55**, 2–27. https://doi.org/10.1353/asi.2016.0013

Cota M, Li P, Rao D-Q, Stanford C, Wang L, Wanchai P (2021) *Manouria impressa. he IUCN Red List of Threatened Species*, **2021**, e.T12775A508518. https://www.researchgate.net/deref/https%3A%2F%2Fdx.doi.org%2F10.2305%2FIUCN.UK.2021-2.RLTS.T12775A508518.en?_tp=eyJjb250ZXh0Ijp7ImZpcnN0UGFnZSI6InB1YmxpY2F0aW9uIiwicGFnZSI6InB1YmxpY2F0aW9uIiwicG9zaXRpb24iOiJwYWdlSGVhZGVyIn19

Cowie RH, Bouchet P, Fontaine B (2022) The Sixth Mass Extinction: fact, fiction or speculation? *Biological Reviews*, **97**, 640–663. https://doi.org/10.1111/brv.12816

Das I (2010) *A field guide to the reptiles of South-East Asia*. Bloomsbury Publishing, London, New Delhi, New York, Sydney.

Dawson JE, Ihlow F, Ettmar S, Van Dijk PP, Thirakhupt K (2018) *Malayemys macrocephala* (Gray 1859) – Malayan Snail-Eating Turtle, Rice-Field Terrapin. In: *Conservation Biology of Freshwater Turtles and Tortoises* Chelonian Research Monographs. (eds Rhodin AGJ, Iverson JB, Van Dijk PP, Stanford CB, Goode EV, Buhlmann KA, Pritchard PCH, Mittermeier RA), pp. 1–16. Chelonian Research Foundation and Turtle Conservancy. https://doi.org/10.3854/crm.5

Dawson JE, Ihlow F, Platt SG (2020) *Malayemys subtrijuga* (Schlegel and Müller 1845) Mekong Snail-Eating Turtle. In: *Conservation Biology of Freshwater Turtles and Tortoises* Chelonian Research Monographs. (eds Rhodin AGJ, Iverson JB, Van Dijk PP, Stanford CB, Goode EV, Buhlmann KA, Mittermeier RA), pp. 1–24. Chelonian Research Foundation and Turtle Conservancy. https://doi.org/10.3854/crm.5

Denevan WM (1992) The Pristine Myth: The Landscape of the Americas in 1492. *Annals of the Association of American Geographers*, **82**, 369–385. https://doi.org/10.1111/j.1467-8306.1992.tb01965.x

Denham TP, Haberle SG, Lentfer C, Fullagar R, Field J, Therin M, Porch N, Winsborough B (2003) Origins of Agriculture at Kuk Swamp in the Highlands of New Guinea. *Science*, **301**, 189–193. https://doi.org/10.1126/science.1085255

Duckworth JW, Batters G, Belant JL, Bennett EL, Brunner J, Burton J, Challender DWS, Cowling V, Duplaix N, Harris JD, Hedges S, Long B, Mahood SP, McGowan PJK, McShea WJ, Oliver WLR, Perkin S, Rawson BM, Shepherd CR, Stuart SN, Talukdar BK, Dijk PP van, Vié J-C, Walston JL, Whitten T, Wirth R (2012) Why South-east Asia should be the world's priority for averting imminent species extinctions, and a call to join a developing cross-institutional programme to tackle this urgent issue. *S.A.P.I.EN.S. Surveys and Perspectives Integrating Environment and Society*.

Falkus M (1991) The economic history of Thailand. *Australian Economic History Review*, **31**, 53–71. https://doi.org/10.1111/aehr.311004

FAO (2020a) *Global Forest Resources Assessment 2020: Thailand Report*. FAO, Rome. https://doi.org/10.4060/ca9825en

FAO (2020b) *Global Forest Resources Assessment 2020: Main Report*. FAO, Rome. https://doi.org/10.4060/ca9825en

FAO (2020c) *Global Forest Resources Assessment 2020: Cambodia Report*. FAO, Rome. https://doi.org/10.4060/ca9825en

Forestier H, Sophady H, Puaud S, Celiberti V, Frère S, Zeitoun V, Mourer-Chauviré C, Mourer R, Than H, Billault L (2015) The Hoabinhian from Laang Spean Cave in its stratigraphic, chronological, typo-technological and environmental context (Cambodia, Battambang province). *Journal of Archaeological Science: Reports*, **3**, 194–206. https://doi.org/10.1016/j.jasrep.2015.06.008

Frère S, Auetrakulvit P, Zeitoun V, Sophady H, Forestier H (2018) Doi Pha Kan (Thailand), Ban Tha Si (Thailand) and Laang Spean (Cambodia) Late Paleolithic animal bone assemblages. A new perception of meat supply strategies for Early Holocene Mainland South-east Asia? In: *Advancing Southeast Asian Archaeology* (ed Tan NH), pp. 100–108, 320–323. Bangkok.

Hermanson G, Arnal FAM, Szczygielski T, Evers SW (2024) A systematic comparative description of extant turtle humeri, with comments on humerus disparity and evolution based on fossil comparisons. *The Anatomical Record*, **307**, 3437–3505. https://doi.org/10.1002/ar.25450

Hirayama R, Sonoda T, Takai M, Prefectural F (2023) A new skull of the side-necked turtle (Order Testudines: Infraorder Pleurodira; Family Podocnemididae) from the Neogene of Myanmar. *Waseda Global Forum*, **20**, 1–10.

Imdirakphol S, Zazzo A, Auetrakulvit P, Tiamtinkrit C, Pierret A, Forestier H, Zeitoun V (2017) The perforated stones of the Doi Pha Kan burials (Northern Thailand): A Mesolithic singularity? *Comptes Rendus Palevol*, **16**, 351–361. https://doi.org/10.1016/j.crpv.2016.12.003

Johnson CN, Balmford A, Brook BW, Buettel JC, Galetti M, Guangchun L, Wilmshurst JM (2017) Biodiversity losses and conservation responses in the Anthropocene. *Science*, **356**, 270–275. https://doi.org/10.1126/science.aam9317

Le M (2007) Conservation of turtles in Vietnam: A survey of Cat Tien National Park. *Oryx*, **41**, 544–547. https://doi.org/10.1017/S0030605307012148

Moll EO, Platt SG, Chan EH, Horne BD, Platt K, Praschag P, Chen PN, van Dijk PP (2015) *Batagur affinis* (Cantor 1847) – Southern River Terrapin, Tuntong. In: *Conservation Biology of Freshwater Turtles and Tortoises: A Compilation Project of the IUCN/SSC Tortoise and Freshwater Turtle Specialist Group* Chelonian Research Monographs. (eds Rhodin A, Pritchard P, Van Dijk PP, Saumure R, Buhlmann K, Iverson J, Mittermeier R), pp. 1–17. Chelonian Research Foundation. https://doi.org/10.3854/crm.5

Mourer-Chauviré C, Mourer R (1970) The Prehistoric Industry of Laang Spean, Province of Battambang, Cambodia. *Archaeology and Physical Anthropology in Oceania*, **5**, 128–146. https://doi.org/10.1002/j.1834-4453.1970.tb00110.x

Mourer-Chauviré C, Mourer R, Thommeret Y (1970) Premières datations absolues de l'habitat préhistorique de la grotte de Laang Spean, province of Battambang (Cambodge). *C. R. Acad. Sci. Paris, Ser. D*, **270**, 471–473.

Myers N, Mittermeier RA, Mittermeier CG, Fonseca GAB da, Kent J (2000) Biodiversity hotspots for conservation priorities. *Nature*, **403**, 853. https://doi.org/10.1038/35002501

Naksri W, Tong H, Lauprasert K, Suteethorn V, Claude J (2013) A new species of *Cuora* (Testudines: Geoemydidae) from the Miocene of Thailand and its evolutionary significance. *Geological Magazine*, **150**, 908–922. https://doi.org/10.1017/S0016756812001082

Paradis E (2018) Nonlinear relationship between biodiversity and human population density: evidence from Southeast Asia. *Biodiversity and Conservation*, **27**, 2699–2712. https://doi.org/10.1007/s10531-018-1563-5

Platt S, Sovannara H, Kheng L, Holloway R, Stuart B, Rainwater T (2008) Biodiversity, Exploitation, and Conservation of Turtles in the Tonle Sap Biosphere Reserve, Cambodia, with Notes on Reproductive Ecology of Malayemys subtrijuga. *Chelonian Conservation and Biology*, **7**, 195–204. https://doi.org/10.2744/CCB-0703.1

Pomoim N, Trisurat Y, Hughes AC, Corlett RT (2022) Can Thailand Protect 30% of Its Land Area for Biodiversity, and Will This Be Enough? *Diversity*, **14**, 344. https://doi.org/10.3390/d14050344

Pookajorn S (2001) New perspectives for Palaeolithic research in Thailand. In: *Origine des peuplements et chronologie des cultures paléolithiques dans le sud-est asiatique*, pp. 167–188. Semenanjung, Paris, France.

Pritchard PCH, Rabett RJ, Piper PJ (2009) Distinguishing species of Geoemydid and Trionychid turtles from shell fragments: evidence from the Pleistocene at Niah Caves, Sarawak. *International Journal of Osteoarchaeology*, **19**, 531–550. https://doi.org/10.1002/oa.1038

Puyravaud J-P (2003) Standardizing the calculation of the annual rate of deforestation. *Forest Ecology and Management*, **177**, 593–596. https://doi.org/10.1016/S0378-1127(02)00335-3

Ramitanondh S (1989) Forests and deforestation in Thailand: a pandisciplinary approach. In: *Culture and Environment in Thailand: A Symposium of the Siam Society*, pp. 23–50. The Siam Societey, Bangkok, Thailand.

Rhodin AGJ, Iverson J, Bour R, Fritz U, Georges A, Shaffer BH, van Dijk PP (2021) *Turtles of the World: Annotated Checklist and Atlas of Taxonomy, Synonymy, Distribution, and Conservation Status (9th Ed.)*. Chelonian Research Foundation and Turtle Conservancy, United States of America.

Rhodin AGJ, Thomson S, Georgalis GL, Karl HV, Danilov IG, Takahashi A, De La Fuente MS, Bourque JR, Delfino M, Bour R, Iverson JB, Shaffer BH, van Dijk PP (2015) Turtles and tortoises of the world during the rise and global spread of humanity: first checklist and review of extinct Pleistocene and Holocene chelonians. *Chelonian Research Monographs*, **5**, 1–66.

Roberts P (2019) *Tropical forests in prehistory, history, and modernity*. Oxford, United Kingdom.

Santos A, Mayor P, Loureiro L, Gilmore M, Perez Peña P, Bowler M, Lemos L, Svensson M, Nekaris KA, Nijman V, Valsecchi J, Morcatty T (2020) Widespread Use of Traditional Techniques by Local People for Hunting the Yellow-Footed Tortoise (Chelonoidis denticulatus) Across the Amazon. *Journal of Ethnobiology*, **40**, 268. https://doi.org/10.2993/0278-0771-40.2.268

Schoppe S, Das I (2015) *Cuora amboinensis* (Riche in Daudin 1801) Southeast Asian Box Turtle. In: *Conservation Biology of Freshwater Turtles and Tortoises: A Compilation Project of the IUCN/SSC Tortoise and Freshwater Turtle Specialist Group* Chelonian Research Monographs. (eds Rhodin A, Pritchard P, Van Dijk PP, Saumure R, Buhlmann K, Iverson J, Mittermeier R), pp. 1–53. Chelonian Research Foundation. https://doi.org/10.3854/crm.5

Sharma DSK, Tisen OB (2000) Freshwater Turtle and Tortoise Utilization and Conservation Status in Malaysia. In: *Asian Turtle Trade: Proceedings ofa Woiksliop on Consen'ation and Trade ofFre1zirarer Turtles and Tortoises in Asia* Chelonian Research Monographs. (eds Van Dijk PP, Stuart BL, Rhodin AGJ), pp. 120–128.

Sodhi NS, Posa MRC, Lee TM, Bickford D, Koh LP, Brook BW (2010) The state and conservation of Southeast Asian biodiversity. *Biodiversity and Conservation*, **19**, 317–328. https://doi.org/10.1007/s10531-009-9607-5

Sophady H, Forestier H, Zeitoun V, Puaud S, Frère S, Celiberti V, Westaway K, Mourer R, Mourer-Chauviré C, Than H, Billault L, Tech S (2016) Laang Spean cave (Battambang province): A tale of occupation in Cambodia from the Late Upper Pleistocene to Holocene. *Quaternary International*, **416**, 162–176. https://doi.org/10.1016/j.quaint.2015.07.049

Stephens L, Fuller D, Boivin N, Rick T, Gauthier N, Kay A, Marwick B, Armstrong CG, Barton CM, Denham T, Douglass K, Driver J, Janz L, Roberts P, Rogers JD, Thakar H, Altaweel M, Johnson AL, Vattuone MMS, Aldenderfer M, Archila S, Artioli G, Bale MT, Beach T, Borrell F, Braje T, Buckland PI, Cano NGJ, Capriles JM, Castillo AD, Çilingiroğlu Ç, Cleary MN, Conolly J, Coutros PR, Covey RA, Cremaschi M, Crowther A, Der L, Lernia S di, Doershuk JF, Doolittle WE, Edwards KJ, Erlandson JM, Evans D, Fairbairn A, Faulkner P, Feinman G, Fernandes R, Fitzpatrick SM, Fyfe R, Garcea E, Goldstein S, Goodman RC, Guedes JD, Herrmann J, Hiscock P, Hommel P, Horsburgh KA, Hritz C, Ives JW, Junno A, Kahn JG, Kaufman B, Kearns C, Kidder TR, Lanoë F, Lawrence D, Lee G-A, Levin MJ, Lindskoug HB, López-Sáez JA, Macrae S, Marchant R, Marston JM, McClure S, McCoy MD, Miller AV, Morrison M, Matuzeviciute GM, Müller J, Nayak A, Noerwidi S, Peres TM, Peterson CE, Proctor L, Randall AR, Renette S, Schug GR, Ryzewski K, Saini R, Scheinsohn V, Schmidt P, Sebillaud P, Seitsonen O, Simpson IA, Sołtysiak A, Speakman RJ, Spengler RN, Steffen ML, Storozum MJ, Strickland KM, Thompson J, Thurston TL, Ulm S, Ustunkaya MC, Welker MH, West C, Williams PR, Wright DK, Wright N, Zahir M, Zerboni A, Beaudoin E, Garcia SM, Powell J, Thornton A, Kaplan JO, Gaillard M-J, Goldewijk KK, Ellis E (2019) Archaeological assessment reveals Earth's early transformation through land use. *Science*, **365**, 897–902. https://doi.org/10.1126/science.aax1192

Szczygielski T, Piechowski R (2023) Limb anatomy of the Triassic turtles: appendicular osteology of *Proterochersis* (Testudinata, Proterochersidae). *Zoological Journal of the Linnean Society*, **199**, 771–831. https://doi.org/10.1093/zoolinnean/zlad057

Thomson SA (2021) Turtles of the World: Annotated Checklist and Atlas of Taxonomy, Synonymy, Distribution, and Conservation Status. *Phyllomedusa: Journal of Herpetology*, **20**, 225–228. https://doi.org/10.11606/issn.2316-9079.v20i2p225-228

Trisurat Y, Alkemade R, Verburg PH (2010) Projecting Land-Use Change and Its Consequences for Biodiversity in Northern Thailand. *Environmental Management*, **45**, 626–639. https://doi.org/10.1007/s00267-010-9438-x

Tsujino R, Kajisa T, Yumoto T (2019) Causes and history of forest loss in Cambodia. *International Forestry Review*, **21**, 372–384. https://doi.org/10.1505/146554819827293178

Wanchai P, Stanford CB, Tharapoom K, Thirakhupt K (2013) Activity Budget of the Impressed Tortoise, *Manouria impressa* (Günther, 1882), in Phu Luang Wildlife Sanctuary, Thailand. *Tropical Natural History*, **13**, 39–48.

White JC, Penny D, Kealhofer L, Maloney B (2004) Vegetation changes from the late Pleistocene through the Holocene from three areas of archaeological significance in Thailand. *Quaternary International*, **113**, 111–132. https://doi.org/10.1016/j.quaint.2003.09.001

Willis KJ, Gillson L, Brncic TM (2004) How "Virgin" Is Virgin Rainforest? *Science*, **304**, 402–403. https://doi.org/10.1126/science.1093991

Yasukawa Y, Hirayama R, Hikida T (2001) Phylogenetic relationships of Geoemydine turtles (Reptilia: Bataguridae). *Current Herpetology*, **20**, 105–133.

Zangerl R (1969) The turtle shell. In: *Biology of the Reptilia, Morphology A, Volume 1* (eds Gans C, Bellairs AA, Parsons TS), pp. 311–339. Academic Press, London and New York.

1340   Zeitoun V, Auetrakulvit P, Zazzo A, Pierret A, Frère S, Forestier H (2019) Discovery of an
1341   outstanding Hoabinhian site from the Late Pleistocene at Doi Pha Kan (Lampang province,
1342   northern   Thailand).   *Archaeological   Research   in   Asia*,   **18**,   1–16.
1343   https://doi.org/10.1016/j.ara.2019.01.002
1344   Zeitoun V, Lenoble A, Laudet F, Thompson J, Rink WJ, Mallye J-B, Chinnawut W (2010) The Cave
1345   of the Monk (Ban Fa Suai, Chiang Dao wildlife sanctuary, northern Thailand). *Quaternary*
1346   *International*, **220**, 160–173. https://doi.org/10.1016/j.quaint.2009.11.022
1347   Zeitoun V, Pérez-Balarezo A, Sophady H, Zhou Y, Puaud S, Zazzo A, Tombret O, Détroit F, Gollette
1348   L, Griggo C, Bochaton C, Pois V, Kosal N, Forestier H (2024) An integrative study of new
1349   environmental and cultural data from the Hoabinhian of Laang Spean Cave (Cambodia)
1350   including modern human remains. *Quaternary Environments and Humans*, 100020.
1351   https://doi.org/10.1016/j.qeh.2024.100020
1352   Zug GR (1971) Buoyancy, locomotion, morphology of the pelvic girdle and hindlimb, and
1353   systematics of cryptodiran turtles. *Miscellaneous Publications of the Museum of Zoology,*
1354   *University of Michigan*, **142**, 1–98.

Appendix 1. List of the specimens used in the study. Abbreviations: **CUMZ-R**: Chulalongkorn
University Museum of Zoology (Bangkok, Thailand) reptile collection; **MNHN-RA**: Muséum national
d'Histoire naturelle (Paris, France) reptiles and amphibians collection; **MNHN-RA**: Muséum
national d'Histoire naturelle (Paris, France) comparative anatomy collection; **MRHNB**: Musée
Royal d'Histoire naturelle de Belgique; **MVZ**: Museum of Vertebrate Zoology (Berkeley, USA); **UF**:
University of Florida Museum of Natural History (Gainesville, USA).

Geoemydidae: *Batagur baska* (MNHN-RA-0.9094; CUMZ-R-TG1105; CUMZ-R-TG1108;
CUMZ-R-TG151697; CUMZ-R-TG155191); *Cuora couro* (MNHN-RA-1889-182; CUMZ-R-TG364;
CUMZ-R-TG342 ; UF 134824; UF 134833); *Cuora galbinifrons* (UF 81775; UF 154221); *Cuora*
mouhotii (UF 53668; UF 61986; UF 81980) ; *Cuora trifasciata* (UF 103395); *Cyclemys oldhamii*
(CUMZ-R-TG969); *Cyclemys atripons* (UF 105993; UF 119584, UF 116449); *Cyclemys dentata*
(CUMZ-R-TG954 ; UF 135096 ; UF 139828 ; UF 154136 ; UF 154138 ; UF 155192); *Cyclemys* sp.
(CUMZ-R-TG959; CUMZ-R-TG960; CUMZ-R-TG1432; CUMZ-R-TG1053); *Heosemys annandalii*
(CUMZ-R-TG606; CUMZ-R-TG608; CUMZ-R-TG609; CUMZ-R-TG613; CUMZ-R-TG614; CUMZ-
R-TG615; CUMZ-R-TG618; CUMZ-R-TG619; CUMZ-R-TG622; CUMZ-R-TG624; CUMZ-R-
TG633; CUMZ-R-TG647; CUMZ-R-TG656; CUMZ-R-TG762; UF 154245; UF 155196); *Heosemys*
grandis (CUMZ-R-TG863; UF 154202; UF 154244); *Heosemys spinosa* (CUMZ-R-TG1005 ; UF
32853 ; UF 154230 ; UF 154227); *Leucocephalon yuwonoi* (UF 97335 ; UF 109835); *Malayemys*
subtrijuga (CUMZ-R-TG42 ; CUMZ-R-TG51 ; CUMZ-R-TG58 ; CUMZ-R-TG147 ; MNHN-RA-
1874-78 ; UF 62085 ; UF 67527 ; UF 67638 ; UF 69380 ; UF 150281); *Mauremys annamensis* (UF
88811 ; UF 151054 ; UF 154218) ; *Mauremys mutica* (UF 107174 ; UF 88806); *Melanochelys*
trijuga (CUMZ-R-TG1151 ; MNHN-RA-1900-184; UF 61987); *Morenia ocellata* (MNHN-RA-
0.9167); *Notochelys platynota* (UF 123862 ; UF 151154 ; UF 154260); *Orlitia borneensis* (MRHNB
1941.97; UF 152476 ; UF 154020 ; UF 154234); *Siebenrockiella crassicollis* (CUMZ-R-TG551;
MNHN-ZM-AC 1875-88). **Testudinidae**: *Geochelone elegans* (MVZ-Herp-23519); *Geochelone*
platynota (AMNH-R-57451); *Manouria emys* (CUMZ-R-TT11; CUMZ-R-TT12; CUMZ-R-TT15; UF
52644; UF 27470; UF 67618; UF 134830; UF 155203). **Platysternidae**: *Platysternon*
megacephalum (UF 56669; UF 56950; UF 57111; UF 57608). **Trionychidae**: *Amyda cartilaginea*
(UF 57728); *Dogania subplana* (UF 61810; UF 69259); *Lissemys punctata* (UF 56017); *Pelodiscus*
sinensis (UF 175673; UF 175674; UF 175673; UF 175674).

Appendix 2. Supplementary figures: Drawings of the carapace and plastron of the modern
Southeast Asian turtles (Testudinidae and Geoemydidae). The species are ordered alphabetically
and *Indotestudo elongata* is not included as already figured in Bochaton et al. (2023).

1389

1390   **Fig. S1** - Carapace and plastron of a subadult specimen of *Batagur borneoensis* (Schlegel & Müller,
1391   1845) (CUMZ-R-TG1108) in dorsal view (upper left), ventral view (lower left), posterior view (upper
1392   right), and anterior view (lower left). Scalebars=2cm.

1393

1394   **Fig. S2** - Carapace (left) and plastron (right) of *Cuora couro* (Leschenault de la Tour in Schweigger,
1395   1812) (CUMZ-R-TG342). From top to bottom for carapace: dorsal, ventral, lateral, and posterior views,
1396   and for plastron: ventral, dorsal, and lateral views. Scalebar=2cm.

1397

1398   **Fig. S3** - Carapace and plastron of *Cuora mouhotii* (Gray, 1862) (UF 81980). Carapace in dorsal
1399   (upper left), posterior (upper right), and lateral (medium left) views. Plastron in dorsal (bottom left) and
1400   ventral (bottom right) views. A detailed view of the attachment surface of the carapace with the plastron
1401   is also figured (medium right). Scalebars=2cm.

1402

1403   **Fig. S4** - Carapace (left) and plastron (right) of *Cyclemys atripons* Iverson & McCord, 1997 (UF
1404   116449). No scalebar.

1405

Fig. S5 - Carapace and plastron of *Cyclemys oldhamii* Gray, 1863 (CUMZ-R-TG969). Carapace in dorsal (upper), and posterior (medium left) views. Plastron in dorsal (bottom left) and ventral (bottom right) views. A detailed view of the attachment surface of the carapace with the plastron is also figured (medium right). Scalebars=2cm.

1410

1411   **Fig. S6** - Carapace (left, dorsal view) and plastron (right, ventral) of *Leucocephalon spengleri*
1412   (Gmelin, 1789) (PCHP 12207). Modified from Ascarrunz, Claude, and Joyce (2021). No scalebar.

1413

1414

1415   **Fig. S7** - Carapace of *Heosemys annandalii* (Boulenger, 1903) (CUMZ-R-TG647) in ventral (upper
1416 left), dorsal (bottom left), lateral (upper right), and posterior (bottom right) views. Scalebars=2cm.

1417

1418   **Fig. S8** - Carapace of *Heosemys grandis* (Gray, 1860) (Chulalongkorn Museum – no number) in
1419   ventral (upper left), dorsal (bottom left), and posterior (right) views. Scalebar=4cm.

1420

1421   **Fig. S9** - Carapace of *Heosemys spinosa* (Gray, 1830) (UF 32853) in dorsal (upper left), ventral
1422 (upper right), lateral (bottom left), and posterior (bottom right) views. Scalebar=3cm.

1423

1424   **Fig. S10** - Carapace of *Malayemys macrocephala* (Gray, 1859) (CUMZ-R-TG42) in ventral (upper
1425   left), dorsal (medium left), lateral (bottom left), and posterior (right) views. Scalebar=2cm.

1426

1427   **Fig. S11** - Carapace of *Manouria emys* (Schlegel & Müller, 1840) (UF 134830) in dorsal (upper left),
1428   ventral (upper right), lateral (bottom left), and posterior (bottom right) views. Scalebar=5cm.

1429

1430   **Fig. S12** - Carapace of *Manouria impressa* (Günther, 1882) modified from Bourret (1941) in dorsal
1431   (left) and ventral (right) views. Scalebar=5cm.

1432

1433   **Fig. S13** - Carapace of *Melanochelys trijuga* (Schweigger, 1812) (CUMZ-R-TG51) in dorsal (upper
1434   left), ventral (lower left), lateral (upper right), and posterior (bottom right) views. Scalebars=3cm.

1435

1436   **Fig. S14** - Carapace and plastron of *Notochelys platynota* (Gray, 1834) (UF 123862). Carapace in
1437   dorsal (upper left), lateral (upper right), and posterior (medium right) views. Plastron in dorsal (bottom
1438   left) and ventral (bottom right) view. Scalebars=2cm.

1439

1440   **Fig. S15** - Carapace of a juvenile specimen of *Orlitia borneensis* Gray, 1873 (MRHNB 1941.97) in
1441   dorsal (upper left), ventral (bottom left), lateral (upper right), and posterior (bottom right) views.
1442   Scalebars=2cm.

1443

1444   **Fig. S16** - Carapace of an adult specimen of *Orlitia borneensis* Gray, 1873 (UF 152476) in dorsal
1445   (upper), and ventral (bottom) views. Scalebars=5cm.

1446

1447   **Fig. S17** - Carapace (UF 57111) and plastron (UF 56669) of *Platysternon megacephalum* Gray,
1448   1831. Carapace in dorsal view (left) and plastron in dorsal (upper right) and ventral (bottom right) views.
1449   Scalebars=2cm.

1450

1451   **Fig. S18** - Carapace of *Siebenrockiella crassicollis* (Gray, 1830) (CUMZ-R-TG551) in ventral (upper
1452   left), lateral (bottom left), dorsal (upper right), and posterior (bottom right) views. Scalebars=2cm.

1453