Title: Illuminating the mystery of thylacine extinction: a role for relaxed selection and gene

loss

Buddhabhushan Girish Salve, Nagarjun Vijay

Computational Evolutionary Genomics Lab, Department of Biological Sciences, IISER

Bhopal, Bhauri, Madhya Pradesh, India

Author for Correspondence: Nagarjun Vijay, nagarjun@iiserb.ac.in

Abstract

Gene loss shapes lineage-specific traits but is often overlooked in species survival. In this

study, we investigate the role of ancestral gene loss using the extinction icon—thylacine

(*Thylacinus cynocephalus*). While studies of neutral genetic variation indicate a population

decline before extinction, the impact of thylacine-specific ancestral gene losses remains

unexplored. The availability of a chromosomal-level genome of the extinct thylacine offers a

unique opportunity for such comparative studies. Here, we leverage palaeogenomic data to

compare gene presence/absence patterns between the Tasmanian devil and thylacine. We

discovered ancestral (between 13-1 Ma) loss of *SAMD9L*, *HSD17B13*, *CUZD1* and *VWA7*

due to multiple gene-inactivating mutations, corroborated by short-read sequencing. The loss

of *SAMD9* correlates with a carnivorous diet, with ancestral gene loss timing mirroring the

thylacine's shift to hypercarnivory. Our genome-wide analysis reveals olfactory receptor loss

and relaxed selection, aligning with reduced olfactory lobes in the thylacine, indicating

olfaction is not its primary hunting sense. By integrating palaeogenomic data with

comparative genomics, our study reveals ancestral gene losses and their impact on species

fitness while providing a foundation for selecting candidate species and genes for thylacine

de-extinction. Our approach can be extended to other extinct/endangered species,

highlighting genetic factors for de-extinction/conservation efforts.

Keywords

thylacine, extinct species, gene loss, RELAX selection, comparative genomics, de-extinction, hypercarnivorous diet

1) Introduction

Traditional genetics-based approaches to species conservation have focused mainly on measures of genetic diversity, the prevalence of mutational load and recent demographic histories [1–4]. However, there has been little focus on ancestral gene loss and its potential role in species survival and extinction. This aspect remains underexplored in conservation genetics despite its potential impact on long-term species survival and resilience to rapid environmental changes due to anthropogenic activities. Several studies have shown that gene loss plays a key role in shaping phenotypic evolution [5], such as adaptations for blood-feeding in bats [6], modifications of the epidermis for aquatic environments in cetaceans [7], dietary variation and retention/pseudogenisation of *ADH7* [4] and loss of the *PLAAT1* gene in vertebrates for adaptation to low-light environments [8]. Nonetheless, ancestral gene losses have demonstrated implications for species survival. For instance, (i) the ancient loss of the *PON1* gene in marine mammals eliminates their primary defence against neurotoxicity from anthropogenic organophosphorus compounds, posing potential risks in modern environments [9], (ii) in odontocetes, the loss of *MX1* and *MX2* genes leads to decreased resistance to viral infections and long-term survival [10], and (iii) the loss of the *CNR2* gene makes parrots more susceptible to neuroinflammation [11], highlighting the consequences of gene loss in these species in terms of long-term survival and disease susceptibility. In previous studies, we highlighted that the loss of ciliary and immune cell genes contributes to differential susceptibility to influenza in bird species [12,13]. These findings emphasise that species-specific ancestral gene losses, which may have conferred advantages in adapting to unique

environments in the distant past, can now have adverse effects in the present day because of

anthropogenic activities [7] and underscore a critical link between gene loss and increased

extinction risk.

As a promising approach, knowledge of gene losses in extinct species could provide valuable

insights into the causes of species extinction, refine de-extinction strategies, and offer clues

about evolutionary processes, such as trait evolvability. For example, the loss of the *FOXQ1*

gene among extinct woolly mammoths has been linked to a reduced ability to adapt to cold

climates, ultimately leading to lower fitness [14]. The advent of machine learning-based

comparative genomics tools such as TOGA [15] and the increase in publicly available

genomic data from both extant and extinct species make it possible to conduct genome-wide

screening for ancestral gene loss in extinct species [15–17]. Gene loss can result from genetic

drift, positive or relaxed selection, leading to exon deletions or the accumulation of mutations

that disrupt the reading frame, such as premature stop codons, frameshifts, dysfunctional

splice sites, or repeat insertions or chromosomal rearrangements [6,8,12,13]. The scarcity of

multiple samples from extinct species hampers the robust estimation of genetic diversity and

the prevalence of mutational load, making it challenging to fully understand their genetic

dynamics and evolutionary history [18]. However, single-sample genomic data are sufficient

for ancestral gene loss inference. Genome-wide screening for ancestral gene loss becomes

vital when the only surviving species within a clade becomes extinct. For example, the

thylacine (*Thylacinus cynocephalus*; also known as the Tasmanian tiger or Tasmanian wolf)

was the largest modern-day carnivorous marsupial and the sole surviving member of the

genus *Thylacinus* and the family *Thylacinidae* until modern times [19]. The *Thylacinidae*

were small-bodied, generalist faunivores until the middle Miocene climatic transition

(MMCT; ~15–13 Ma). Following this period, they increased in size and specialised in a

hypercarnivorous diet [20]. This raises a critical question in conservation genetics: which

genetic changes are unique to a species, and what are their functional impacts, particularly in terms of the species' long-term survivability, resilience to environmental changes, and vulnerability to novel diseases?

Thylacines were once widespread across continental Australia and New Guinea, but their disappearance from the mainland during the late Holocene was likely driven by climate change, human intensification, and competition with dingoes [21–23]. In Tasmania, however, thylacine numbers had already declined significantly by the early 1800s, following European settlement, the introduction of farming, a government hunting bounty, and disease, leading to eventual extinction, with the last recorded thylacine dying in 1936 [16,17,24]. However, mitochondrial and genomic DNA sequences suggest that limited genetic diversity in thylacine precedes climatic transition-mediated extinction [17,25]. Previous research has focused primarily on thylacine's canid-like craniofacial convergence, estimation of extinction date, phylogenetic relationships, demographic history, annotation of immune genes, genetic diversity, biology and anatomy [16,17,19,22,25–34].

In this study, we leverage paleogenomic data of thylacine to examine gene presence and absence patterns compared with the Tasmanian devil (*Sarcophilus harrisii*). We began with a genome-wide screen for gene losses in the thylacine, corroborating these loss events with short-read Illumina sequencing data. Additionally, we reconstructed the gene loss history using genomes from other marsupial species and assessed relaxed selection signatures. Our results offer novel insights into the ancestral loss of key genes in the thylacine— *SAMD9L*, *HSD17B13*, *CUZD1* and *VWA7* potentially compromising its health by affecting antiviral defence, metabolic processes, lactation, pancreatitis and tumour susceptibility, which may have influenced its extinction and environmental adaptations, providing a basis for refining genomics-informed genetic-engineering based de-extinction strategies. Although the present study focuses on the thylacine, our approach can easily be extended to other species (extinct

99 and extant) and improve our understanding of species-specific biology, thereby refining de-

100 extinction strategies and extending conservation efforts to include genetic-engineering-based

101 enhancements.

2) Methods

(a) Genome-wide screen for thylacine-specific gene loss

We identified gene loss in the thylacine (*Thylacinus cynocephalus*) by comparing its genome

to the Tasmanian devil (*Sarcophilus harrisii*). Firstly, to compute pairwise genome

alignments between the Tasmanian devil (GCF_902635505.1; chromosome-wise; as target)

and the thylacine (GCA_007646695.3; whole genome; as query), we used the

make_lastz_chains v2.0.8 pipeline (https://github.com/hillerlab/make_lastz_chains), which

employs LASTZ v1.04.15 with the following parameters: K = 2400, L = 3000, Y = 9400, H =

2000. Next, we applied the machine learning-based tool TOGA v1.1.7 [15] to determine the

status of genes in the thylacine genome, using chain alignment and NCBI RefSeq annotation

of the Tasmanian devil (52,692 transcripts) (see electronic supplementary material, figure

S1).

(b) Reconstruction of gene loss history and its validation

We assessed the gene orthology of the gene loss candidates on the basis of their syntenic

position within the genomes of human (*Homo sapiens*), opossum (*Monodelphis domestica*),

yellow-footed antechinus (*Antechinus flavipes*) and Tasmanian devil (*Sarcophilus harrisii*)

(see electronic supplementary material, table S1-S2). The gene-inactivating events inferred

by TOGA for "clearly lost" genes were corroborated with the CDS of Tasmanian devil as

query using the BLASTn v2.13.0 (parameters: -evalue 0.01 -max_target_seqs 5000 -outfmt

'17 SQ') search in ~223 Gb short-read data of thylacine (SRR5055303, SRR5055304,

SRR5055305, and SRR5055306) [35]. The alignments were then used to generate

visualizations with IGV-reports v1.12.0, and the results were inspected to confirm the

presence of in-frame stop codons and frame-disrupting events. Additionally, we used bam-

readcount v1.0.1 [36] to determine the number of reads supporting the reference versus the

alternative (disruptive) base or indel at the positions of predicted events by TOGA v1.1.7,

providing further support for the inferred gene loss. Additionally, cross-species mapping of

thylacine short-read data to Tasmanian devil was done using BWA-mem v0.7.17 [37]. We

reassessed the whole-genome sequencing short-read data for the presence of in-frame stop

codons using HybPiper v2.3.1 and Patchwork v0.5.2. To assess the integrity of the

HSD17B13 locus in the chromosomal-level assembly of the opossum (*Monodelphis*

domestica), we first aligned long-read data of PacBio HiFi to the opossum reference genome

using BWA-MEM v0.7.17 [37] (parameters: bwa mem -x pacbio). The resulting BAM file

was then used as the input for the scan_alignment and alignment_plot subprograms in the

klumpy v1.0.11 [38]. However, similar analyses could not be performed for the numbat

(*Myrmecobius fasciatus*), coppery ringtail possum (*Pseudochirops cupreus*), plush-coated

ringtail possum (*Pseudochirops corinnae*), and thylacine due to the unavailability of long-

read sequencing data for these species.

To reconstruct the history of gene loss, we downloaded available chromosomal-level

marsupial genomes from NCBI. In addition, we included scaffold-level assemblies of

phylogenetically closely related species, numbat (*Myrmecobius fasciatus*) (n=21; see

electronic supplementary material, table S3). We started with a BLASTn v2.13.0 search of

marsupial genomes using the Tasmanian devil gene sequences as queries. The resulting hits

were processed using mergeBed -d 10000 to merge close intervals and slopBed -b 30000 to

expand the regions on both sides. These regions were then extracted using getFastaFromBed

of BEDTools v2.27.1 [39]. The resulting FASTA sequences were used to generate pairwise

alignments between Tasmanian devil (as target genomes) and twenty marsupial genomes (as

query) using make_lastz_chains pipeline, as mentioned in the previous section. We ran

TOGA with parameters --o2o to process only the genes with a single orthologous chain, using

NCBI RefSeq annotation of the Tasmanian devil.

(c) Transcriptional status of lost genes

We investigated the RNA expression of the genes lost in thylacine using the publicly

available Tasmanian devil and dunnart transcriptome sequencing data from 27 tissues

(allantois, amnion, axillary nerve, bone marrow, brachial plexus nerve, brain stem,

cerebellum, cerebrum, distal yolk sac, endometrium, eye, heart, liver, lung, optic nerve, ovary,

oviduct, pancreas, prostate gland, proximal yolk sac, sciatic nerve, skin, spleen, testis,

thyroid, trigeminal nerve and uterus) of both males and females. (BioProject ID:

PRJEB34650 [40] and PRJNA1028148 [41]; see electronic supplementary material, table

S4). We obtained transcripts per million (TPM) values by performing pseudo-alignments of

the reads to the longest isoforms of the Tasmanian devil and dunnart transcripts using kallisto

v0.51.0 [42].

We screened the expression of the lost gene in the thylacine miRNA-seq short-read database

using BLASTn v2.13.0. To validate the presence of protein-coding gene expression in this

dataset, we also queried (as a positive control) the sequences of *LOC100913894* (*ACTA*) and

LOC100925998 (*MYH7*), based on a previous report [43]. Additionally, we assessed the

transcriptional status of the *VWA7* gene using RNA-Seq data from the tongue, lung, and liver

of the numbat (*Myrmecobius fasciatus*) [44] and *SAMD9/9L* using transcriptome data from

testes, spleen, lung, liver, kidney and heart of tammar wallaby (*Macropus eugenii*).

(d) Evaluating the strength of selection, limitations with GC-content and estimation of

gene loss timing

To test the relaxation of selection in gene loss candidates, we obtained pairwise codon

alignments between the Tasmanian devil and other marsupials, including the thylacine from

TOGA [15]. Codons affected by frameshifting insertions or deletions and premature stop

codons were replaced by "NNN" to maintain a reading frame. A multiple sequence alignment

(MSA) of the protein-coding sequence was done using PRANK v.170427 in GUIDANCE2

v2.01. The sequence saturation in MSA was checked in the DAMBE v7.3.32 program as

proposed by Xia et al. [45]. We considered MSA for molecular evolutionary analysis when

Iss < Iss.c (Iss.c is the value at which the sequences will begin to fail to recover the true tree)

and p-value < 0.05. Furthermore, the alignments and phylogenetic tree obtained from the

Timetree website (https://timetree.org/) were used to investigate whether the gene evolves

under relaxed selection specifying one branch as the foreground and all other branches as

background at a time in the RELAX model of the HyPhy v2.5.48 program [46] and codeml of

PAML v4.9f program [47]. We also implemented aBSREL and BUSTED, which are branch

and gene-wide tests, respectively, to test for signatures of selection on branches of the

marsupial phylogeny for the lost genes. Additionally, site-specific models such as FEL and

MEME were used to identify sites evolving under positive and negative selection.

The gene loss timing was estimated using the method proposed by Meredith et al. [49], which

is based on two assumptions: (i) $dS_f = dS_p$ (the synonymous substitution rate in the mixed

branch   (dS_f)   does   not   change   after   the   gene   is   pseudogenised

(dS_p), i.e. the 1ds method), and (ii) $dS_f = 0.7\ dS_p$ (the synonymous substitution rate of a

functional copy of the gene (dS_f) is 0.7 times the substitution rate after pseudogenisation

(dS_p), i.e. the 2ds method). We used 38.5 Ma as the divergence time for the thylacine from its

most recent common ancestor (MRCA) and dN/dS (ω) estimated for functional (ω_f), mixed

(ω_m), and pseudogene (ω_p) using a branch-free model in the codeml program from PAML

using unrooted trees.

(e) *SAMD9* and *SAMD9L* gene phylogeny

To confirm that the *SAMD9* and *SAMD9L* sequences used are 1-to-1 orthologs, we evaluate the maximum likelihood tree generated from marsupial species' *SAMD9* and *SAMD9L* orthologs. The MSA of *SAMD9/9L* was used to generate a gene tree using IQ-TREE v2.3.6 (parameters: *-m MFP --alrt 1000 -B 1000 --boot-trees*) [50]. We also calculated a concordance factor for each branch of the gene tree, comparing the gene tree computed from the complete MSA against the fraction of gene trees of the 1 kb window sequence.

(f) Identifying *SAMD9* loss and phylogenetic logistic regression

We investigated the status of *SAMD9* and *SAMD9L* genes across mammals using TOGA. To examine whether *SAMD9* gene loss is associated with the percentage of endothermic vertebrates in the diet, we performed phylogenetic logistic regression analyses using the R package "phylolm" [51]. Specifically, we tested whether the proportion of endothermic vertebrates in the diet of each species correlated with the intactness (coded as 1) or loss (coded as 0) of *SAMD9*. We were able to collate the proportion of endothermic vertebrates for a total of 72 mammal species whose *SAMD9* gene status is also available from Jiao, Hengwu et al. [52] and Kapsetaki, Stefania E et al. [53] (see electronic supplementary material, table S5).

For detailed methods and parameter settings, see the electronic supplementary data.

3) Results

In brief, our comparative genomic analysis identified ancestral gene losses in the extinct thylacine, attributed to frame-disrupting changes and/or in-frame stop codons in synteny-confirmed orthologs of *SAMD9L*, *HSD17B13*, *CUZD1*, and *VWA7* (figure 1). Signatures of relaxed selection were detected in the thylacine lineage for *SAMD9/9L*, *CUZD1*, and *VWA7* genes using RELAX and/or codeml (see electronic supplementary material, table S6). The

estimated timing of gene loss for these genes ranges from approximately 13-6 Ma, and the loss of the *SAMD9* gene is negatively correlated with the percentage of endothermic vertebrates in the diet (figure 2; see electronic supplementary material, table S5-S7). Additionally, we observed rampant gene losses in olfactory receptor genes and other genes prone to pseudogenisation accompanied by signatures of relaxed selection (figure 3; see electronic supplementary material, table S1 and S6). The GC content of the four lost genes ranges from 37.65% to 51.71% and is not susceptible to lower coverage due to GC bias. The intermediate levels of GC among these genes suggest these loss events are not a consequence of intragenomic mutational heterogeneity [54].

(a) Candidate Gene Losses: Identification and Validation

The thylacine genome was assembled using Illumina sequencing (HiSeq and NextSeq), which exhibits reduced coverage in regions with high GC content [16,17,55]. This limitation can lead to false positives in gene loss identification [56]. In our analysis, many genes categorised by TOGA as missing, partially intact, partially missing, uncertain loss, or clearly lost were associated with high GC content, low or no coverage, missing exons, or deleted exons (figure 3a; see electronic supplementary material, table S1 and figure S1-S2). We filtered out gene loss candidates with high GC content, prevalence of G/C stretches, and insufficient coverage to minimise false positives, ensuring robust identification of true gene losses. Similarly, paralogs can affect the assembly, ultimately leading to ambiguity in validating gene-disrupting events for specific genes (see electronic supplementary material, figure S2-S5 and data). Additionally, genes annotated as "Low-quality protein" in the Tasmanian devil genome were excluded, as they are potentially lost in the Tasmanian devil itself.

Furthermore, we observed widespread gene loss and signatures of relaxed selection in

olfactory genes (figure 3b; see electronic supplementary material, table S1 and S6). These

loci experience reduced purifying selection and evolve rapidly in many mammals, exhibiting

high rates of gene gain and loss [57]. Therefore, we did not pursue these findings further in

our study. Also, we observe the loss of genes prone to pseudogenisation (as per NCBI

annotation), such as *CASPASE-16*, *PTCHD3*, and *PRSS45* (see electronic supplementary

material, table S1 for supporting references).

(b) Loss of *SAMD9L* and signatures of relaxed selection in *SAMD9/9L*

Our genome-wide analysis identified that *SAMD9L* is lost in the thylacine due to multiple

gene-inactivating mutations. These mutations include the deletion of one base (A) at codon-

287, an in-frame stop codon (CGA→TGA) at codon-511, a deletion of four bases (GGTG)

between 1932-1937th position, an insertion of one base (C) at codon-667, and in-frame stop

codons   at   codon-680   (CGA→TGA),   codon-1207   (CAG→TAG),   and   codon-1463

(CAA→TAA) (figure 1). All seven events are supported by short-read sequencing data (see

electronic supplementary material, figure S6-S8 and data). Further, HybPiper found non-

terminal stop codons for *SAMD9L*, ruling out potential paralog-to-contig assignment errors.

Among the screened marsupial genomes, *SAMD9L* loss was unique to the thylacine (figure

1), with an estimated timing of gene loss at approximately 11–8 Ma (figure 2a; see electronic

supplementary material, table S7). While other marsupial species such as the opossum,

yellow-footed antechinus, and Tasmanian devil have intact *SAMD9L* genes with *TMPPE*,

CRTAP, and *SUSD5* on the left flank, and *SAMD9*, *CDK6*, *FAM133B*, and *PEX1* on the right

flank. In human *CALCR*, *VPS50* and *HEPACAM2* genes are on the left flank of *SAMD9L*, and

SAMD9, *CDK6*, *FAM133B,* and *PEX1* genes are on the right flank (see electronic

supplementary material, table S2). In addition to the synteny-based identification of *SAMD9L*

orthologs, gene sequence-based phylogenetic trees also delineated the true orthologs with

good bootstrap support (see electronic supplementary material, figure S9-S10). In the

Tasmanian devil, *SAMD9L* is annotated as *LOC100924681* (XM_031940019.1), with a GC

content of 37.65%. The *SAMD9L* is expressed in the Tasmanian devil and dunnart lung,

ovary, and thyroid transcriptomes (figure 3c; see electronic supplementary material, table S4).

Notably, we found signatures of relaxed selection using RELAX in *SAMD9* of the thylacine

(p-value = 0.004; k-value = 0.16; figure 3b); in contrast, relaxed selection signatures are not

found in closely related species of the *Dasyuridae* family. Additionally, we observed relaxed

selection signatures for *SAMD9L* using codeml in PAML (see electronic supplementary

material, table S6) and episodic diversifying positive selection using aBSREL (p-value =

0.00019) and BUSTED (p-value = 0.0061) models of the Hyphy program in thylacine. Some

marsupial species of the *Macropodidae* family also exhibit signatures of relaxed selection

(see electronic supplementary material, figure S11 and table S6). However, we found

evidence of robust expression in the tammar wallaby (*Macropus eugenii*) (see electronic

supplementary material, figure S12).

The signatures of relaxed selection in *SAMD9/9L* and *SAMD9L* gene loss due to multiple

gene-inactivating mutations indicate that thylacine may have become vulnerable to the

critical function provided by *SAMD9* and *SAMD9L* genes. Interestingly, we found that the

SAMD9 gene is annotated as "Low-quality protein" in 22 out of 130 annotated species on

NCBI, of which 7/7 are carnivore species. Although most mammals retain an intact *SAMD9*,

widespread losses of *SAMD9* were detected in the carnivore lineage. We detected inactivating

mutations leading to loss of function, including frameshifting indels and inframe stop codons,

in mammalian orders such as Carnivora and Pholidota (see electronic supplementary

material, figure S13 and table S5). However, shared loss of *SAMD9* is rare in the

Perissodactyla and Artiodactyla orders. Widespread loss of *SAMD9* in carnivores, in contrast

to conservation in other vertebrates, suggested a tentative link between diet and dispensability

of this gene. We found a negative correlation between the percentage of endothermic

vertebrates in the diet and the *SAMD9* gene loss (slope = -0.01, p-value = 0.3426 for

logistic_MPLE method (Insignificant); slope = -0.03, p-value = 0.0061 for

logistic_IG10 method (Significant), (figure 2b, see electronic supplementary material, table

S5)) strengthening the relationship between *SAMD9* dispensability and carnivorous diet.

(c) Loss of *HSD17B13*

In the thylacine, *HSD17B13* was inactivated through a splice-site mutation (gt →ga) at

Exon1, and the deletion of one base (G) in Exon5 (figure 1c; see electronic supplementary

material, figure S6). The estimated timing of this gene loss is approximately 2–1 Ma (figure

2a; see electronic supplementary material, table S7). The Tasmanian devil's *HSD17B13*

(XM_003774508.4) has a GC content of 39.98% and is expressed in the liver and pancreas

(figure 3c; see electronic supplementary material, table S4). The micro-synteny of the human

HSD17B13 gene is flanked by *IBSP*, *DMP1*, *DSPP*, *SPARCL1*, *SCPPPQ1*, *NUDT9*, and

HSD17B11 on the left, while the right flank consists of *KLHL8*, *AFF1*, *C6H4orf36*, and

SLC10A6 genes. In the yellow-footed antechinus and Tasmanian devil, the left flank includes

IBSP, *DMP1*, *DSPP*, *NUDT9*, and *HSD17B11*, while the right flank remains consistent with

humans, containing *KLHL8*, *AFF1*, *C6H4orf36*, and *SLC10A6,* confirming 1-to-1 orthology

(see electronic supplementary material, table S2).

Notably, *HSD17B13* annotation is missing from the annotated genome of the opossum in the

NCBI Genome Data Viewer (see electronic supplementary material, figure S14). We found

that *HSD17B13* is independently lost in opossum due to an in-frame stop codon (TGG →

TGA) in Exon1 (BLASTn search of short-read data revealed polymorphic hits for this event)

and deletion of Exons 3-to-7 (figure 1b, see electronic supplementary material, figure S14).

We identified an assembly gap at the syntenic locus (see electronic supplementary material,

figure S15); however, a BLASTn search in the long-read database returned spurious hits for

Exons 3-to-7. Additionally, signatures of relaxed selection were found in the opossum using

RELAX (p-value = 0.0008; k-value = 0.00) and PAML's codeml (see electronic

supplementary material, table S6). At the same time, episodic diversifying positive selection

was detected in the opossum's *HSD17B13* using aBSREL (p-value = 0.00) and BUSTED (p-

value = 0.00) models (see electronic supplementary material, table S6).

(d) Loss of *CUZD1*

We identified multiple gene-inactivating mutations in *CUZD1* that contributed to its loss in

thylacine (figure 1c). These include the deletion of one base (G) at codons-67 and 95, an in-

frame stop codon (TAC → TAG) at codon-119, and another premature stop codon (TGC →

TGA) at codon-283 (see electronic supplementary material, figure S6). Furthermore,

BLASTn analysis recovered only partial sequences for Exon6, whereas no BLASTn hits were

obtained for Exons 7-8, indicating its likely deletion. The estimated timing of gene loss is

approximately 13-9 Ma (figure 2a). The *CUZD1* gene, annotated as *LOC100927202*

(XM_003755291.4) in the Tasmanian devil, has a GC content of 39.35% and is expressed in

the pancreas (figure 3c; see electronic supplementary material, table S4). Notably, we found

signatures of relaxed selection by the PAML codeml. At the same time, we found evidence

for intensification by RELAX model (p-value = 0.00; k-value = 3.72) and signatures of

episodic diversifying positive selection are detected in thylacine *CUZD1* by aBSREL (p-

value = 0.00) and BUSTED (p-value = 0.00) models of Hyphy (see electronic supplementary

material, table S6). Additionally, *CUZD1* independently lost in the golden ringtail possum

(*Pseudochirops corinnae*) and the coppery ringtail possum *(Pseudochirops cupreus)*; both

species belong to the marsupial family *Pseudocheiridae*. In golden ringtail possum, no

BLASTn hits were found for Exons 1-to-4 in the genomic short-read database, while Exon5

exhibited two in-frame stop codons (CAA → TAA and TCA → TGA) and one base (A)

deletion. In coppery ringtail possum, an in-frame stop codon (CAA → TAA) and one base

(A) deletion led to the inactivation of *CUZD1* (figure 1b; see electronic supplementary

material, figure S16). Two gene-inactivating mutations in Exon5 of these two species are

shared. At the same time, this gene remains intact in the western ringtail possum

(*Pseudocheirus occidentalis*), suggesting that *CUZD1* was lost independently within the

Pseudochirops genus at approximately 25-20 Ma (figure 1b).

Our micro-syntenic analysis found that in humans, *KZF5*, *PSTK*, *LOC124902519*,

C10orf88B, and *FAM24B* genes occur on the left flank of *CUZD1* and *LOC124900290*,

DMBT1L1, and *SPADHI* on the right flank. However, among marsupials (opossum, yellow-

footed antechinus and Tasmanian devil), the right flank has *C2H10orf88*, *PSTK,* and *IKZF5,*

whereas, on the left flank, *HTRA1* and *DMBT1* were annotated in the NCBI genome data

viewer and located approximately 350 kb away supporting 1-to-1 orthology (see electronic

supplementary material, table S2).

(e) Loss of *G7c* (*VWA7*)

This gene has acquired multiple gene-inactivation mutations in thylacine (figure 1c).

Insertion of one base (C) in codon-37, two base deletion of CA in codon-171, insertion of one

base (T) at codon-403, TGG → TAG premature stop codon at codon-575 and insertion of two

bases (TA) leading to an in-frame stop codon (GCC → TAG) at codon-706 (see electronic

supplementary material, figure S6). The estimated timing of gene loss is approximately 8-6

Ma (figure 2a). The GC content of Tasmanian devil *VWA7* (XM_003768930.3) is 51.71 %

and expressed in the dunnart oviduct (figure 3; see electronic supplementary material, table

S4). The micro-synteny of *VWA7* in humans includes *HSPA1A*, *HSPA1L*, *LSM2*, and *VARS1*

on the left flank, whereas *SAPCD1*, *MSH5*, *CLIC1*, and *DAH2* are on the right flank. A

similar orthology relationship was found in marsupials such as the opossum, yellow-footed

antechinus, and Tasmanian devil (see electronic supplementary material, table S2). We found that *VWA7* is lost in numbat due to an in-frame stop codon in Exon5 and Exon14 and the deletion of one base in Exon14 (figure 1, also see electronic supplementary material, figure S17). We found signatures of relaxed selection by RELAX (p-value = 0.01; k-value = 0.40) as well as by codeml in thylacine, and lack of expression at its syntenic locus, suggesting *VWA7* gene loss in numbat (see electronic supplementary material, table S6, and figure S18).

4) Discussion

In this study, we used thylacine, an extinction icon, as a case study to investigate ancestral gene losses through a comparative genomic approach. Our genome-wide analysis identified the ancestral loss of *SAMD9L*, *HSD17B13*, *CUZD1*, *VWA7* genes and several olfactory receptors in thylacine. Short-read sequencing confirmed all gene-inactivating mutations, validating gene loss. Our investigation into the timing of gene loss events revealed a staggered pattern, occurring over a span of approximately 13-1 Ma. This temporal distribution indicates that the gene losses were not confined to a single evolutionary event but occurred progressively after the MMCT. Interestingly, the loss of *SAMD9* among mammals is negatively correlated with a carnivorous diet. The genes lost in the thylacine are also lost in about 10% of screened marsupials and show limited evidence of relaxed selection. Relaxation signals detected by RELAX and/or codeml, along with lower dN/dS values on functional branches compared to mixed branches, suggest these genes evolved under gradually relaxed constraints (see electronic supplementary material, table S6-S8) [49,58]. This indicates that their loss was not widespread in marsupials and likely reflects lineage-specific pressures or stochastic events in the thylacine. The lost genes are expressed in similar tissues in the Tasmanian devil and dunnart, implying conserved functions [59] and involved in antiviral defence, metabolism, lactation signalling, pancreatitis, and tumour susceptibility, and their loss may have reduced thylacine fitness and adaptability, potentially contributing to

extinction. More broadly, identifying such gene losses informs conservation genomics and de-extinction, guiding genome editing strategies to restore extinct traits. Identifying and targeting these loci can facilitate the selection of appropriate candidate species and genes for genome editing, thereby improving the feasibility and success of de-extinction efforts and genetic engineering in conservation.

(a) Impact of ancestral gene losses on thylacine fitness and adaptation

The thylacine genome exhibits the loss of four genes– *SAMD9L*, *HSD17B13*, *CUZD1*, and *VWA7* –each with important biological functions and implications for species fitness. The *SAMD9L* (Sterile Alpha Motif Domain Containing 9 Like) is a paralog of *SAMD9* [60]. The *SAMD9/9L* genes are critical antiviral factors and tumour suppressors, playing crucial roles in the innate immune defence [61,62]. The *SAMD9* gene is lost (deleted) in the mouse due to evolutionary genomic rearrangements [60], and mouse *SAMD9L* is not a functional paralog of human *SAMD9* [63]. Notably, species-specific differences in *SAMD9L* among rodents act as a natural barrier to cross-species orthopoxvirus infections, underscoring the adaptive significance of these genes in shaping immune defences across different lineages [64]. Furthermore, *SAMD9L* is the only gene that experienced convergent amino acid substitutions and an accelerated evolutionary shift among bats and birds, which are natural hosts of several zoonotic pathogens, highlighting its crucial role in conferring resistance to viruses [65]. Our study found that *SAMD9* and *SAMD9L* are highly conserved across marsupials, except for thylacine. However, *SAMD9* is lost among mammals like carnivores and pangolins while intact in closely related groups such as odd-toed ungulates, indicating a possible association between gene loss and dietary diversification. The phylogenetic correlation between *SAMD9* gene loss and the fraction of endothermic vertebrates in the diet suggests that a shift towards carnivory might have favoured the loss of *SAMD9/9L* [20]. Hence, the loss of *SAMD9L* in thylacine may have conferred a benefit following a shift to hypercarnivory in the past. At the

same time, in the modern environment, it may have increased vulnerability to zoonotic viral infections and compromised immune responses. Another lost gene, *HSD17B13* (Hydroxysteroid 17-beta-dehydrogenase 13), encodes a hepatic lipid droplet-associated enzyme [66]. Loss-of-function variants in *HSD17B13* are associated with protection against the progression of steatosis to more severe liver conditions [67], suggesting that *HSD17B13* gene loss could have functional adaptive significance. The CUB and zona pellucida-like domain-containing protein 1 (*CUZD1*) is also lost in the thylacine. *CUZD1* interacts with *JAK1*/*JAK2* and *STAT5*, which are mediators of prolactin signalling critical for mammary gland development. Its absence abolishes *STAT5* phosphorylation during amelogenesis, leading to defective lactation [68]. Furthermore, *CUZD1* is abundantly expressed in pancreas acinar cells of human and mouse [69,70]. We also found that *CUZD1* is expressed in the Tasmanian devil's pancreas. *CUZD1* deficient mice demonstrated increased severity of experimentally induced acute pancreatitis, suggesting *CUZD1* is a strong candidate as a pancreatitis susceptibility gene [70,71]. Finally, *VWA7* (Von Willebrand Factor A Domain Containing 7), also known as *G7c*, is located in the mouse and human MHC class III region [72]. The *VWA7* gene has been implicated in controlling lung tumour susceptibility and host immune response to viral infections, suggesting its potential role in disease susceptibility [73,74].

Interestingly, the timing of gene losses for *SAMD9L*, *CUZD1*, and *VWA7* is estimated to be approximately 13–6 Ma, a period that follows the middle Miocene climatic transition (MMCT; ~15–13 Ma). This period marked significant ecological and climatic changes, during which *Thylacinidae*, previously small-bodied, unspecialised faunivores, began to undergo notable shifts. After MMCT, they increased in size and developed adaptations to a hypercarnivorous diet, likely driven by the aridification of the Australian environment [20]. Besides this, the thylacine and grey wolf exemplify remarkable convergent evolution,

displaying strikingly similar craniofacial morphologies despite ~160 million years of

evolutionary divergence [16,17]. Notably, these lost genes are located near a candidate

craniofacial cis-regulatory element, indicating a possible connection with phenotypic

convergence (see supplementary table of [79]). This pattern underscores the complexity of

gene loss events and their potential role in shaping convergent phenotypic traits, which

remains to be verified experimentally [80]. In addition, our analysis revealed a drastic loss of

olfactory receptors in the thylacine, paired with relaxed selection, aligning with earlier

findings of its reduced olfactory lobes [20,77]. Unlike the Tasmanian Devil, a scavenger, the

taller thylacine likely depended on vision and sound for hunting, aided by its ability to see

over vegetation. This shift in sensory reliance is further evidenced by the pronounced ridges

on its neocortex, suggesting heightened cognitive abilities to support its hunting strategies

[78]. Notably, gene loss has also been observed during major evolutionary transitions, such as

from unicellular to multicellular life and from land to water [75,76].

Whatever the cause for gene loss, the loss of *SAMD9L*, *HSD17B13*, *CUZD1*, and *VWA7*

genes in the thylacine possibly had negative pleiotropic effects, potentially compromising its

health by affecting antiviral defence, metabolic processes, lactation, pancreatitis and tumour

susceptibility. Although model-based population viability analyses suggest that the disease

played only a minor role in thylacine extinction [81], other studies suggest that a 'canine-

distemper-like' disease played a role in exacerbating its extinction [82–84]. Therefore,

adaptive ancestral gene loss may have provided specific advantages to the thylacine in

response to environmental pressures during the MMCT, while being deleterious in the recent

past, similar to the example of *PON1* gene loss in marine mammals [9].

(b) Studying gene loss in extinct species: Opportunities, constraints, and conservation

implications

Comparative studies of extinct species offer valuable insights into host-specific immune responses and phenotypic evolution [5,12]. Such studies can play a pivotal role in refining de-extinction strategies by assisting in selecting candidate species and genetic loci for genome editing, thereby informing decisions about diet and habitat preferences [85–87]. Advances in comparative genomics and ancient DNA sequencing enable researchers to link gene loss with phenotypic diversity and evolutionary mechanisms, though such analyses depend heavily on genome assembly quality [6,7,9–11,14]. For extinct species, reliance on Illumina sequencing—despite its limitations in GC-rich regions—can lead to false positives in gene loss detection [16,56]. The fragmentary nature of ancient DNA in such species further restricts the use of long-read sequencing technologies [88]. This lack of long-read sequencing, DNA damage and the absence of transcriptomic data make it even more challenging to differentiate between assembly artefacts and true gene losses [14,16].

Specifically for thylacine, the genome assembly (GCA_007646695.3) was assembled using Illumina HiSeq and NextSeq platforms. In line with previous research, our study finds that genes with high GC content tend to have lower sequencing coverage [56] and, therefore, are classified as missing, partially intact, or partially lost by TOGA. As a result, this might lead to incomplete or less reliable assemblies in GC-rich genomic regions, affecting the accuracy of downstream analyses and may lead to false positives for gene loss [56]. Although miRNA-Seq data is available [43], it does not provide insights into protein-coding gene expression. The gene losses we identified involve gene-inactivating mutations (e.g., splice-site changes, stop codons, insertions, and deletions) supported by forward and reverse short reads, indicating they are genuine rather than artefacts of ancient DNA damage. Assessment of gene loss in large multi-gene families with complex orthology relationships remains challenging and must be interpreted cautiously [6], especially in reference-guided assemblies [89]. Future improvements to the thylacine genome assembly will strengthen the assertion of gene losses.

Although habitat loss, hunting, introduced species, inbreeding depression, and loss of genetic

diversity are widely recognised as key contributors to species extinction [2,90], these factors

have primarily dominated conservation genetics studies focused on mitigating risks to

threatened species [3,91–93]. In contrast, the role of ancestral gene loss in species extinction

remains largely unexplored and raises the question whether genetically altering a gene or

gene combination will help the species survive in the modern world, and how the genetic

background (the allelic states of genes, prevalence of regulatory elements, gene content, etc,

in the rest of the genome) will affect the fitness [9–11]. Insights into ancestral lineage-

specific gene losses can deepen our understanding of extinct species evolution and refine de-

extinction approaches. Our study also offers a framework to test hypotheses on why specialist

phylogenetic lineages struggle to persist and diversify [7,94] and provides a novel perspective

on the interplay between species diversity and genetic diversity.

5) Conclusion

Our study offers an immediate framework that includes practical approaches in comparative

genomics and insights into de-extinction strategies. Our findings from this case study of an

extinction icon- the thylacine—establishes the ancestral gene loss of *SAMD9L*, *HSD17B13*,

CUZD1, and *VWA7*, contributing unique insights into the genetic factors influencing

extinction. Additionally, we observe gene loss in olfactory receptors.

In line with temporal estimations of gene losses, our study provides an empirical case where

we observe the coincidence of gene loss timing (approximately 13–1 Ma, following the

middle Miocene climatic transition ~15–13 Ma). This period marked significant ecological

and climatic changes, during which *Thylacinidae*, previously small-bodied, unspecialised

faunivores, underwent notable shifts such as an increase in size and developing adaptations to

a hypercarnivorous diet, likely driven by the aridification of the Australian environment [20].

We find evidence that *SAMD9* gene loss is associated with a switch to hypercarnivory. As the thylacine was the only surviving member of the genus *Thylacinus* and the family *Thylacinidae* until modern times, our study also offers a hypothesis on why specialist phylogenetic lineages struggle to persist and diversify over macroevolutionary timescales [7,19,94]. Furthermore, our analysis reveals a pervasive loss of olfactory receptors in the thylacine, reflecting relaxed selection and correlating with its reduced olfactory lobes. This shift in sensory reliance, possibly driven by changes in traits such as height and vision, contrasts with that of the Tasmanian devil.

In addition, our analysis provides critical information for identifying which species are better candidates for cloning and which genetic loci are suitable for genome editing, contributing to future de-extinction efforts for thylacine and guiding decisions about diet and habitat preferences [85–87]. Our study provides a framework for identifying ancestral gene losses that may improve the probability of success of de-extinction programs.

Ethics

This work did not require ethical approval from a human subject or animal welfare committee.

Data accessibility

This study's data and relevant code are available on Github: https://github.com/CEGLAB-Buddhabhushan/Thylacine_Genome-wide_GENELOSS.git. The reviewer URL: https://github.com/CEGLAB-Buddhabhushan/Thylacine_Genome-wide_GENELOSS.git

Declaration of AI use

We have not used AI-assisted technologies to create this article.

Conflict of interest declaration

We declare that we have no competing interests.

Authors' contributions

B.G.S.: conceptualisation, data curation, formal analysis, investigation, methodology,

validation, visualisation, writing—original draft, writing—review and editing; N.V.:

conceptualisation, funding acquisition, supervision, investigation, methodology, validation,

writing—original draft, writing — review, and editing.

Both authors gave final approval for publication and agreed to be held accountable for the

work performed therein.

Funding

This article was funded by the Department of Biotechnology, Ministry of Science and

Technology, India (grant no. BT/11/IYBA/2018/03) and Science and Engineering Research

Board (grant no. ECR/2017/001430) provided funds for procuring computational resources

(i.e. Har Gobind Khorana Computational Biology cluster) used.

Acknowledgements

We thank the Council of Scientific & Industrial Research for a fellowship to B.G.S. We used

BioRender (https://biorender.com) to arrange figures and images of species. We thank

Sandhya Sharma for the animal paintings in Figure 1.

References

1.   Bouzat JL. 2010 Conservation genetics of population bottlenecks: the role of chance,

selection, and history. *Conserv Genet* **11**, 463–478. (doi:10.1007/s10592-010-0049-0)

2.   Frankham R. 2005 Genetics and extinction. *Biol Conserv* **126**, 131–140.

(doi:10.1016/j.biocon.2005.05.002)

3.   Khan A *et al.* 2021 Genomic evidence for inbreeding depression and purging of

deleterious genetic variation in Indian tigers. *Proc. Natl. Acad. Sci.* **118**, e2023018118.

(doi:10.1073/pnas.2023018118)

4.   Pinto SL *et al.* 2023 Diet and the evolution of ADH7 across seven orders of mammals.

R. Soc. Open Sci. **10**. (doi:10.1098/RSOS.230451)

5.   Albalat R, Cañestro C. 2016 Evolution by gene loss. *Nat. Rev. Genet. 2016 177* **17**,

379–391. (doi:10.1038/nrg.2016.39)

6.   Blumer M *et al.* 2022 Gene losses in the common vampire bat illuminate molecular

adaptations to blood feeding. *Sci. Adv.* **8**, 6494. (doi:10.1126/sciadv.abm6494)

7.   Sharma V, Hecker N, Roscito JG, Foerster L, Langer BE, Hiller M. 2018 A genomics

approach reveals insights into the importance of gene losses for mammalian

adaptations. *Nat. Commun. 2018 91* **9**, 1–9. (doi:10.1038/s41467-018-03667-1)

8.   Drabeck DH *et al.* 2024 Gene loss and relaxed selection of plaat1 in vertebrates

adapted to low-light environments. *Proc. R. Soc. B* **291**.

(doi:10.1098/RSPB.2023.2847)

9.   Meyer WK *et al.* 2018 Ancient convergent losses of Paraoxonase 1 yield potential

risks for modern marine mammals. *Science (80-.).* **361**, 591–594.

(doi:10.1126/science.aap7714)

10.   Braun BA, Marcovitz A, Camp JG, Jia R, Bejerano G. 2015 Mx1 and Mx2 key

antiviral proteins are surprisingly lost in toothed whales. *Proc. Natl. Acad. Sci.* **112**,

8036–8040. (doi:10.1073/pnas.1501844112)

11.   Divín D *et al.* 2022 Cannabinoid receptor 2 evolutionary gene loss makes parrots more

susceptible to neuroinflammation. *Proc. R. Soc. B* **289**. (doi:10.1098/RSPB.2022.1941)

12.   Salve BG, Kurian AM, Vijay N. 2023 Concurrent loss of ciliary genes WDR93 and

583   CFAP46 in phylogenetically distant birds. (doi:10.1098/rsos.230801)

584   13.   Salve BG, Sharma S, Vijay N. 2024 Evolutionary diversity of CXCL16-CXCR6:

585   Convergent substitutions and recurrent gene loss in sauropsids. *Immunogenet. 2024*

586   *765* **76**, 397–415. (doi:10.1007/S00251-024-01357-5)

587   14.   Rogers RL, Slatkin M. 2017 Excess of genomic defects in a woolly mammoth on

588   Wrangel island. *PLoS Genet.* **13**, e1006601. (doi:10.1371/journal.pgen.1006601)

589   15.   Kirilenko BM *et al.* 2023 Integrating gene annotation with orthology inference at

590   scale. *Science (80-.).* **380**. (doi:10.1126/science.abn3107)

591   16.   Feigin CY *et al.* 2018 Genome of the Tasmanian tiger provides insights into the

592   evolution and demography of an extinct marsupial carnivore. *Nat. Ecol. Evol.* **2**, 182–

593   192. (doi:10.1038/s41559-017-0417-y)

594   17.   Feigin C, Frankenberg S, Pask A. 2022 A Chromosome-Scale Hybrid Genome

595   Assembly of the Extinct Tasmanian Tiger (Thylacinus cynocephalus). *Genome Biol.*

596   *Evol.* **14**. (doi:10.1093/GBE/EVAC048)

597   18.   Dutoit L, Vijay N, Mugal CF, Bossu CM, Burri R, Wolf J, Ellegren H. 2017

598   Covariation in levels of nucleotide diversity in homologous regions of the avian

599   genome long after completion of lineage sorting. *Proc. R. Soc. B Biol. Sci.* **284**.

600   (doi:10.1098/RSPB.2016.2756)

601   19.   Krajewski C, Driskell AC, Baverstock PR, Braun MJ. 1992 Phylogenetic relationships

602   of the thylacine (Mammalia: Thylacinidae) among dasyuroid marsupials: Evidence

603   from cytochrome b DNA sequences. *Proc. R. Soc. B Biol. Sci.* **250**, 19–27.

604   (doi:10.1098/rspb.1992.0125)

605   20.   Rovinsky DS, Evans AR, Adams JW. 2019 The pre-Pleistocene fossil thylacinids

606   (Dasyuromorphia: Thylacinidae) and the evolutionary context of the modern thylacine.

607   *PeerJ* **2019**, 1–41. (doi:10.7717/peerj.7457)

608   21.   Letnic M, Fillios M, Crowther MS. 2012 Could direct killing by larger dingoes have

609   caused the extinction of the thylacine from mainland Australia? *PLoS One* **7**, 19–23.

610   (doi:10.1371/journal.pone.0034877)

611   22.   White LC, Mitchell KJ, Austin JJ. 2018 Ancient mitochondrial genomes reveal the

612   demographic history and phylogeography of the extinct, enigmatic thylacine

613   (Thylacinus cynocephalus). *J. Biogeogr.* **45**, 1–13. (doi:10.1111/jbi.13101)

614   23.   White LC, Saltré F, Bradshaw CJA, Austin JJ. 2018 High-quality fossil dates support a

615   synchronous, Late Holocene extinction of devils and thylacines in mainland Australia.

616   *Biol. Lett.* **14**, 10–13. (doi:10.1098/rsbl.2017.0642)

617   24.   Gough N. 2001 Robert Paddle 2000, The Last Tasmanian Tiger: The History and

618   Extinction of the Thylacine. Cambridge University Press, Oakleigh, Victoria. *Aust. J.*

619   *Environ. Educ.* **17**, 162–163. (doi:10.1017/S0814062600003670)

620   25.   Menzies BR, Renfree MB, Heider T, Mayer F, Hildebrandt TB, Pask AJ. 2012 Limited

621   genetic diversity preceded extinction of the Tasmanian tiger. *PLoS One* **7**.

622   (doi:10.1371/journal.pone.0035433)

623   26.   Wroe S, Clausen P, McHenry C, Moreno K, Cunningham E. 2007 Computer

624   simulation of feeding behaviour in the thylacine and dingo as a novel test for

625   convergence and niche overlap. *Proc. R. Soc. B Biol. Sci.* **274**, 2819–2828.

626   (doi:10.1098/rspb.2007.0906)

27.   Pask AJ, Behringer RR, Renfree MB. 2008 Resurrection of DNA function in vivo

from an extinct genome. *PLoS One* **3**, 3–7. (doi:10.1371/journal.pone.0002240)

28.   Peel E, Frankenberg S, Hogg CJ, Pask A, Belov K. 2021 Annotation of immune genes

in the extinct thylacine (Thylacinus cynocephalus). *Immunogenetics* **73**, 263–275.

(doi:10.1007/s00251-020-01197-z)

29.   Rovinsky DS, Evans AR, Martin DG, Adams JW. 2020 Did the thylacine violate the

costs of carnivory? Body mass and sexual dimorphism of an iconic Australian

marsupial. *Proc. R. Soc. B Biol. Sci.* **287**. (doi:10.1098/rspb.2020.1537)

30.   Haines E, Bailey E, Nelson J, Fenlon LR, Suárez R. 2023 Clade-specific forebrain

cytoarchitectures of the extinct Tasmanian tiger. *Proc. Natl. Acad. Sci. U. S. A.* **120**.

(doi:10.1073/PNAS.2306516120)

31.   Janis CM, Figueirido B. 2014 Forelimb anatomy and the discrimination of the

predatory behavior of carnivorous mammals: The thylacine as a case study. *J.*

Morphol. **275**, 1321–1338. (doi:10.1002/jmor.20303)

32.   Old JM. 2015 Immunological insights into the life and times of the extinct tasmanian

tiger (Thylacinus cynocephalus). *PLoS One* **10**, 1–17.

(doi:10.1371/journal.pone.0144091)

33.   Newton AH, Weisbecker V, Pask AJ, Hipsley CA. 2021 Ontogenetic origins of cranial

convergence between the extinct marsupial thylacine and placental gray wolf.

Commun. Biol. **4**, 1–9. (doi:10.1038/s42003-020-01569-x)

34.   Miller W *et al.* 2009 The mitochondrial genome sequence of the Tasmanian tiger

[Thylacinus cynocephalus]. *Genome Res.* **19**, 213–220. (doi:10.1101/gr.082628.108)

35.   Thorvaldsdottir H, Robinson JT, Mesirov JP. 2013 Integrative Genomics Viewer

(IGV): high-performance genomics data visualization and exploration. *Brief.*

651  *Bioinform.* **14**, 178–192. (doi:10.1093/bib/bbs017)

652 36. Khanna A *et al.* 2022 Bam-readcount - rapid generation of basepair-resolution

653   sequence metrics. *ArXiv* **7**, 3722. (doi:10.21105/joss.03722)

654 37. Li H. 2013 Aligning sequence reads, clone sequences and assembly contigs with

655   BWA-MEM. *arxiv*

656 38. Madrigal G, Minhas BF, Catchen J. 2024 Klumpy: A tool to evaluate the integrity of

657   long-read genome assemblies and illusive sequence motifs. *Mol. Ecol. Resour.* ,

658   e13982. (doi:10.1111/1755-0998.13982)

659 39. Quinlan AR, Hall IM. 2010 BEDTools: a flexible suite of utilities for comparing

660   genomic features. *Bioinformatics* **26**, 841–2.

661 40. Stammnitz MR *et al.* 2023 The evolution of two transmissible cancers in Tasmanian

662   devils. *Science (80-.).* **380**, 283–293. (doi:10.1126/science.abq6453)

663 41. Ibeh N, Feigin CY, Frankenberg SR, McCarthy DJ, Pask AJ, Romero IG. 2024 De

664   novo transcriptome assembly and genome annotation of the fat-tailed dunnart

665   (Sminthopsis crassicaudata). *GigaByte* **2024**, 35–38. (doi:10.46471/gigabyte.118)

666 42. Bray NL, Pimentel H, Melsted P, Pachter L. 2016 Near-optimal probabilistic RNA-seq

667   quantification. *Nat. Biotechnol. 2016 345* **34**, 525–527. (doi:10.1038/nbt.3519)

668 43. Mármol-Sánchez E *et al.* 2023 Historical RNA expression profiles from the extinct

669   Tasmanian tiger. *Genome Res.* **33**, 1299–1316. (doi:10.1101/gr.277663.123)

670 44. Peel E *et al.* 2022 Genome assembly of the numbat (Myrmecobius fasciatus), the only

671   termitivorous marsupial. *Gigabyte* **2022**, 1–17. (doi:10.46471/GIGABYTE.47)

672 45. Xia X, Xie Z, Salemi M, Chen L, Wang Y. 2003 An index of substitution saturation

and its application. *Mol. Phylogenet. Evol.* (doi:10.1016/S1055-7903(02)00326-3)

46.   Wertheim JO, Murrell B, Smith MD, Kosakovsky Pond SL, Scheffler K. 2015

RELAX: Detecting Relaxed Selection in a Phylogenetic Framework. *Mol. Biol. Evol.*

32, 820–832.

47.   Yang Z. 2007 PAML 4: Phylogenetic Analysis by Maximum Likelihood. *Mol. Biol.*

Evol. **24**, 1586–1591. (doi:10.1093/MOLBEV/MSM088)

48.   Wickham H. 2016 ggplot2. (doi:10.1007/978-3-319-24277-4)

49.   Meredith RW, Gatesy J, Murphy WJ, Ryder OA, Springer MS. 2009 Molecular decay

of the tooth gene enamelin (ENAM) mirrors the loss of enamel in the fossil record of

placental mammals. *PLoS Genet.* **5**. (doi:10.1371/journal.pgen.1000634)

50.   Minh BQ, Schmidt HA, Chernomor O, Schrempf D, Woodhams MD, Von Haeseler A,

Lanfear R, Teeling E. 2020 IQ-TREE 2: New Models and Efficient Methods for

Phylogenetic Inference in the Genomic Era. *Mol. Biol. Evol.* **37**, 1530–1534.

(doi:10.1093/MOLBEV/MSAA015)

51.   Ives AR, Garland T. 2010 Phylogenetic Logistic Regression for Binary Dependent

Variables. *Syst. Biol.* **59**, 9–26. (doi:10.1093/SYSBIO/SYP074)

52.   Jiao H, Zhang L, Xie H, Simmons N, Liu H, Zhao H. 2019 Trehalase Gene as a

Molecular Signature of Dietary Diversification in Mammals. *Mol. Biol. Evol.* **36**,

2171–2183. (doi:10.1093/MOLBEV/MSZ127)

53.   Kapsetaki SE, Basile AJ, Compton ZT, Rupp SM, Duke EG, Boddy AM, Harrison

TM, Sweazea KL, Maley CC. 2025 The relationship between diet, plasma glucose, and

cancer prevalence across vertebrates. *Nat. Commun.* **16**, 2271. (doi:10.1038/S41467-

025-57344-1)

54.   Hara Y, Kuraku S. 2025 Intragenomic mutational heterogeneity: structural and

functional insights from gene evolution. *Trends Genet.*

(doi:10.1016/J.TIG.2025.03.007,)

55.   Benjamini Y, Speed TP. 2012 Summarizing and correcting the GC content bias in

high-throughput sequencing. *Nucleic Acids Res.* **40**, e72. (doi:10.1093/NAR/GKS001)

56.   Kim J *et al.* 2022 False gene and chromosome losses in genome assemblies caused by

GC content variation and repeats. *Genome Biol.* **23**, 1–29. (doi:10.1186/s13059-022-

02765-0)

57.   Nei M, Niimura Y, Nozawa M. 2008 The evolution of animal chemosensory receptor

gene repertoires: roles of chance and necessity. *Nat. Rev. Genet.* **9**, 951–963.

(doi:10.1038/NRG2480)

58.   Li WH, Gojobori T, Nei M. 1981 Pseudogenes as a paradigm of neutral evolution. *Nat.*

1981 2925820 **292**, 237–239. (doi:10.1038/292237a0)

59.   Hara Y, Kuraku S. 2023 The impact of local genomic properties on the evolutionary

fate of genes. *Elife* **12**. (doi:10.7554/eLife.82290)

60.   Lemos de Matos A, Liu J, McFadden G, Esteves PJ. 2013 Evolution and divergence of

the mammalian SAMD9/SAMD9L gene family. *BMC Evol. Biol.* **13**, 121.

(doi:10.1186/1471-2148-13-121)

61.   Meng X, Zhang F, Yan B, Si C, Honda H, Nagamachi A, Sun LZ, Xiang Y. 2018 A

paralogous pair of mammalian host restriction factors form a critical host barrier

against poxvirus infection. *PLoS Pathog.* (doi:10.1371/journal.ppat.1006884)

62.   Legrand A *et al.* 2024 SAMD9L acts as an antiviral factor against HIV-1 and primate

lentiviruses by restricting viral and cellular translation. *PLOS Biol.* **22**, e3002696.

(doi:10.1371/JOURNAL.PBIO.3002696)

63. Li Q, Guo H, Matsui H, Honda H, Inaba T, Sundberg JP, Sprecher E, Uitto J. 2012

Mouse Samd9l is not a functional paralogue of the human SAMD9, the gene mutated

in normophosphataemic familial tumoral calcinosis. *Exp. Dermatol.* **21**, 554–556.

(doi:10.1111/J.1600-0625.2012.01524.X)

64. Zhang F, Meng X, Townsend MB, Satheshkumar PS, Xiang Y. 2019 Identification of

CP77 as the Third Orthopoxvirus SAMD9 and SAMD9L Inhibitor with Unique

Specificity for a Rodent SAMD9L. *J. Virol.* **93**. (doi:10.1128/JVI.00225-19)

65. Matsuda Y, Makino T. 2024 Comparative genomics reveals convergent signals

associated with the high metabolism and longevity in birds and bats. *Proc. R. Soc. B*

291. (doi:10.1098/RSPB.2024.1068)

66. Su W, Mao Z, Liu Y, Zhang X, Zhang W, Gustafsson JA, Guan Y. 2019 Role of

HSD17B13 in the liver physiology and pathophysiology. *Mol. Cell. Endocrinol.* **489**,

119–125. (doi:10.1016/J.MCE.2018.10.014)

67. Abul-Husn NS *et al.* 2018 A Protein-Truncating HSD17B13 Variant and Protection

from Chronic Liver Disease. *N. Engl. J. Med.* **378**, 1096–1106.

(doi:10.1056/NEJMOA1712191)

68. Mapes J, Li Q, Kannan A, Anandan L, Laws M, Lydon JP, Bagchi IC, Bagchi MK.

2017 CUZD1 is a critical mediator of the JAK/STAT5 signaling pathway that controls

mammary gland development during pregnancy. *PLoS Genet.* **13**.

(doi:10.1371/JOURNAL.PGEN.1006654)

69. Fagerberg L *et al.* 2014 Analysis of the human tissue-specific expression by genome-

wide integration of transcriptomics and antibody-based proteomics. *Mol. Cell.*

Proteomics **13**, 397–406. (doi:10.1074/MCP.M113.035600)

70.   Rygiel AM *et al.* 2022 Variants in the pancreatic CUB and zona pellucida-like
domains 1 (CUZD1) gene in early-onset chronic pancreatitis - A possible new
susceptibility gene. *Pancreatology* **22**, 564–571. (doi:10.1016/J.PAN.2022.04.015)

71.   Imamura T, Asada M, Vogt SK, Rudnick DA, Lowe ME, Muglia LJ. 2002 Protection
from Pancreatitis by the Zymogen Granule Membrane Protein Integral Membrane-
associated Protein-1. *J. Biol. Chem.* **277**, 50725. (doi:10.1074/JBC.M204159200)

72.   Van Kooij M, De Groot K, Van Vugt H, Aten J, Snoek M. 2001 Genotype versus
phenotype: conflicting results in mapping a lung tumor susceptibility locus to the G7c
recombination interval in the mouse MHC class III region. *Immunogenetics* **53**, 656–
661. (doi:10.1007/S00251-001-0381-0)

73.   Kumánovics A, Fischer Lindahl K. 2001 G7c in the lung tumor susceptibility (Lts)
region of the Mhc class III region encodes a von Willebrand factor type A domain
protein. *Immunogenetics* **53**, 64–68. (doi:10.1007/S002510100297)

74.   Snoek M, Albertella MR, Van Kooij M, Wixon J, Van Vugt H, De Groot K, Campbell
RD. 2000 G7c, a novel gene in the mouse and human major histocompatibility
complex class III region, possibly controlling lung tumor susceptibility.
Immunogenetics **51**, 383–386. (doi:10.1007/S002510050634)

75.   Jiménez-Marín B, Rakijas JB, Tyagi A, Pandey A, Hanschen ER, Anderson J, Heffel
MG, Platt TG, Olson BJSC. 2023 Gene loss during a transition to multicellularity. *Sci.
Rep.* **13**, 5268. (doi:10.1038/S41598-023-29742-2)

76.   Huelsmann M, Hecker N, Springer MS, Gatesy J, Sharma V, Hiller M. 2019 Genes
lost during the transition from land to water in cetaceans highlight genomic changes

associated with aquatic adaptations. *Sci. Adv.* **5**, 6671–6696.

(doi:10.1126/sciadv.aaw6671)

77. Moeller H. 1970 Vergleichende Untersuchungen zum Evolutionsgrad der Gehirne

großer Raubbeutler (Thylacinus, Sarcophilus und Dasyurus). *J. Zool. Syst. Evol. Res.*

8, 69–80. (doi:10.1111/J.1439-0469.1970.TB00866.X)

78. Dixon JM. 1989 Chapter 20 - Thylacinidae. In *Fauna of Australia* (eds DW Walton,

BJ Richardson), pp. 1–36. Canberra: Australian Government Publishing Service.

79. Feigin CY, Newton AH, Pask AJ. 2019 Widespread cis-regulatory convergence

between the extinct Tasmanian tiger and gray Wolf. *Genome Res.* **29**, 1648–1658.

(doi:10.1101/GR.244251.118/-/DC1)

80. Stern DL. 2013 The genetic causes of convergent evolution. *Nat. Rev. Genet. 2013*

1411 **14**, 751–764. (doi:10.1038/nrg3483)

81. Prowse TAA, Johnson CN, Lacy RC, Bradshaw CJA, Pollak JP, Watts MJ, Brook

BW. 2013 No need for disease: Testing extinction hypotheses for the thylacine using

multi-species metamodels. *J. Anim. Ecol.* **82**, 355–364. (doi:10.1111/1365-

2656.12029)

82. De Castro F, Bolker B. 2005 Mechanisms of disease-induced extinction. *Ecol. Lett.* **8**,

117–126. (doi:10.1111/J.1461-0248.2004.00693.X)

83. Paddle R. 2012 The thylacine's last straw: epidemic disease in a recent mammalian

extinction. *Aust. Zool.* **36**, 75–92. (doi:10.7882/AZ.2012.008)

84. Carlson CJ, Bond AL, Burgio KR. 2018 Estimating the extinction date of the thylacine

with mixed certainty data. *Conserv. Biol.* **32**, 477–483. (doi:10.1111/cobi.13037)

85. Seddon PJ, Moehrenschlager A, Ewen J. 2014 Reintroducing resurrected species:

selecting DeExtinction candidates. *Trends Ecol. Evol.* **29**, 140–147.

(doi:10.1016/J.TREE.2014.01.007)

86. Peers MJL, Thornton DH, Majchrzak YN, Bastille-Rousseau G, Murray DL. 2016 De-

extinction potential under climate change: Extensive mismatch between historic and

future habitat suitability for three candidate birds. *Biol. Conserv.* **197**, 164–170.

(doi:10.1016/J.BIOCON.2016.03.003)

87. Simons M. 2024 Why I work to revive the Tasmanian tiger. *Nature* **631**, 474.

(doi:10.1038/D41586-024-02228-5)

88. Paabo S. 1989 Ancient DNA: extraction, characterization, molecular cloning, and

enzymatic amplification. *Proc. Natl. Acad. Sci.* **86**, 1939–1943.

(doi:10.1073/PNAS.86.6.1939)

89. Lin J *et al.* 2022 Probing the genomic limits of de-extinction in the Christmas Island

rat. *Curr. Biol.* **32**, 1650-1656.e3. (doi:10.1016/j.cub.2022.02.027)

90. Spielman D, Brook BW, Frankham R. 2004 Most species are not driven to extinction

before genetic factors impact them. *Proc. Natl. Acad. Sci.* **101**, 15261–15264.

(doi:10.1073/pnas.0403809101)

91. Khan A, Sil M, Thekaekara T, Garg KM, Sinha I, Khurana R, Sukumar R,

Ramakrishnan U. 2024 Divergence and serial colonization shape genetic variation and

define conservation units in Asian elephants. *Curr. Biol.* **34**, 4692-4703.e5.

(doi:10.1016/J.CUB.2024.08.062)

92. Pinto A V., Hansson B, Patramanis I, Morales HE, van Oosterhout C. 2024 The impact

of habitat loss and population fragmentation on genomic erosion. *Conserv. Genet.* **25**,

49–57. (doi:10.1007/s10592-023-01548-9)

93.   Willi Y, Kristensen TN, Sgro CM, Weeks AR, Ørsted M, Hoffmann AA. 2022

Conservation genetics as a management tool: the five best-supported paradigms to

assist the management of threatened species. *PNAS* **119**, e2105076119.

(doi:10.1073/pnas.2105076119)

94.   Day EH, Hua X, Bromham L. 2016 Is specialization an evolutionary dead end?

Testing for differences in speciation, extinction and trait transition rates across diverse

phylogenies of specialists and generalists. *J. Evol. Biol.* **29**, 1257–1267.

(doi:10.1111/JEB.12867)

Figure legends

Figure 1. Gene losses in the thylacine (*Thylacinus cynocephalus*). (a) Organs where the lost

genes play an important role in the thylacine. We used dog anatomy to illustrate the

corresponding organs in the thylacine. (b) Status of the lost genes across 21 marsupial

species. A red box indicates gene loss in a given species, while a turquoise box represents an

intact gene. The phylogenetic tree used for the analysis was obtained from the TimeTree

website (https://timetree.org/) and annotated using iTOL (https://itol.embl.de/). The timescale

is in Ma. (c) Frame-disrupting events leading to gene loss. The blue box represents intact

exons, while the events of exons affected by frame-disrupting changes are shown with yellow

highlights. Exons with no BLASTn hits are shown in red, and those with partial BLASTn hits

are depicted in orange. Short-read support is shown for four gene-inactivating events of the

SAMD9L gene (for all events, see electronic supplementary material, figure S6).

Figure 2. Timing of gene loss in the thylacine mirrors the shift to hypercarnivory. (a) The

species tree illustrates the gene loss timing in the thylacine lineage, inferred using the 1ds

method under the F1x4 codon frequency model in PAML (see electronic supplementary

material, table S7). The mixed branch is coloured in blue, while the functional branches are in

black. The yellow-filled rectangle highlights the Middle Miocene Climatic Transition

(approximately 15–13 Ma). The timescale is in Ma. (b) The phylogenetic logistic regression

finds a negative correlation between the proportion of endothermic vertebrates in the diet and

SAMD9 gene loss in mammals (n = 72). We ran two logistic regression models, the

logistic_MPLE (red) and the logistic_IG10 (blue). Diet and genotypes of the species included

are shown in black, with the size of the dot indicating sample size.

Figure 3. The GC content, relaxed selection signatures, and expression analysis of lost genes.

(a) Genes classified by TOGA as missing (M), partially missing (PM), partially intact (PI),

unclear loss (UL), and clearly lost (L) exhibit high GC content with the prevalence of long

stretches of G/C in the thylacine. Vertical and horizontal dotted lines indicate the third

quartile values for a particular classification. The lost genes are marked with red-filled

triangles and labelled accordingly. (b) Relaxed selection signatures in the thylacine. The x-

axis represents the k-value, and the y-axis represents the -log10(p-value) obtained using the

RELAX model of HYPHY (see electronic supplementary material, table S6). (c) Heatmap

depicting transcript per million (TPM) values of lost genes across tissues. The tissue names

and their corresponding SRR IDs are shown on the x-axis. The colour gradient is from yellow

(lowest) to blue (highest). All genes were expressed in at least one tissue above a significant

threshold (>1 TPM; see electronic supplementary material, table S4).

(a)

CUZD1 — Pancreas
HSD17B13 — Liver
VWA7 — Lung
SAMD9L — Spleen

(b)

	SAMD9L	HSD17B13	CUZD1	VWA7
Gracilinanus agilis				
Monodelphis domestica		■		
Dromiciops gliroides				
Macrotis lagotis				
Thylacinus cynocephalus	■	■	■	■
Myrmecobius fasciatus				■
Sminthopsis crassicaudata				
Antechinus flavipes				
Dasyurus viverrinus				
Sarcophilus harrisii				
Vombatus ursinus				
Pseudocheirus occidentalis				
Pseudochirops corinnae			■	
Pseudochirops cupreus			■	
Trichosurus vulpecula				
Phalanger gymnotis				
Potorous gilbertii				
Lagorchestes hirsutus				
Notamacropus eugenii				
Macropus giganteus				
Macropus fuliginosus				

(c)

SAMD9L

Markers: -1 TGA -4 +1 TGA TAG TAA

```
XM_031940019.1     GGAACCTAGGTTTGTGGAAGTCTTACTGCAGGATAA
VAHE01001033.1     GGAACCCAGGTTTGTGGANGTCTTATTGCAGGATAA
CM040578.1         GGAACCCAGGTTTGTGGA-GTCTTATTGCAGGATAA
SRR5055306.222563758  GGAACCCAGGTTTGTGGA-GTCTTATTGCAGGATAA
SRR5055305.27772038   GGAACCCAGGTTTGTGGA-GTCTTATTGCAGGATAA
SRR5055306.86522301   GGAACCCAGGTTTGTGGA-GTCTTATTGCAGGATAA
SRR5055305.68588415   GGAACCCAGGTTTGTGGA-GTCTTATTGCAGGATAA
SRR5055306.212285426  GGAACCCAGGTTTGTGGA-GTCTTATTGCAGGATAA
```

```
XM_031940019.1     TCCTTCAGGTCACCAATACAAATATATGAGTC
VAHE01001033.1     TCCTTCAGGTCATAATACAAATATATGAGTC
CM040578.1         TCCTTCAGGTCATAATACAAATATATGAGTC
SRR5055306.134518679  TCCTTCAGGTCATAATACAAATATATGAGTC
SRR5055306.229106607  TCCTTCAGGTCATAATACAAATATATGAGTC
SRR5055305.75301732   TCCTTCAGGTCATAATACAAATATATGAGTC
SRR5055306.176096220  TCCTTCAGGTCATAATACAAATATATGAGTC
SRR5055305.251291565  TCCTTCAGGTCATAATACAAATATATGAGTC
```

```
XM_031940019.1     AGAAGATTTCATGGGTGCTCTAGAAATTCTC
VAHE01001033.1     AGAAGATTCCATGNNTGCTCTAGAAATTCTC
CM040578.1         AGAAGATTCCATG----CTCTAGAAATTCTC
SRR5055306.205021789  AGAAGATTCCATG----CTCTAGAAATTCTC
SRR5055306.82759021   AGAAGATTCCATG----CTCTAGAAATTCTC
SRR5055306.78732274   AGAAGATTCCATG----CTCTAGAAATTCTC
SRR5055306.206588657  AGAAGATTCCATG----CTCTAGAAATTCTC
SRR5055305.112418896  AGAAGATTCCATG----CTCTAGAAATTCTC
```

```
XM_031940019.1     AGAGCATTTTTATCGAGGTGGCAAAGTTTCT
VAHE01001033.1     AGAGCATTTTTATTGAGGTGGCAAAGTTTCT
CM040578.1         AGAGCATTTTTATTGAGGTGGCAAAGTTTCT
SRR5055304.16935107   AGAGCATTTTTATTGAGGTGGCAAAGTTTCT
SRR5055304.60391055   AGAGCATTTTTATTGAGGTGGCAAAGTTTCT
SRR5055303.851173     AGAGCATTTTTATTGAGGTGGCAAAGTTTCT
SRR5055305.104896029  TGAGCATTTTTTTTGAGGTGGCATTGTTTCT
SRR5055304.35542997   AGAGCATTTTTATTGAGGTGGCAAAGTTTCT
```

HSD17B13

Markers: ga -1

CUZD1

Markers: -1 -1 TGA TGA

VWA7

Markers: +1 -2 +1 TAG +2 TAG

(a)

Macrotis lagotis

SAMD9L
(10.93 Ma)

CUZD1
(13.14 Ma)

VWA7
(8.03 Ma)

Thylacinus cynocephalus

HSD17B13
(1.91 Ma)

Myrmecobius fasciatus

Sminthopsis crassicaudata

Antechinus flavipes

Dasyurus viverrinus

Sarcophilus harrisii

Middle Miocene Climate Transition
(MMCT; ~ 15-13 Ma)

(b)

Intact
1

SAMD9 gene status

logistic_MPLE
Slope   = -0.01
p   = 0.3426

logistic_IG10
Slope   = -0.03
p   = 0.0061

Dot size   Number of species

Loss
0

0   20   40   60   80   100
Endothermic vertebrates in diet (%)