Genome reannotation and effector candidate identification in *Meloidogyne chitwoodi* through gland-specific transcriptome analysis

Marcella Teixeira[1][¶], Itsuhiro Ko[1][¶], Sapinder Bali[1][¶#], Paulo Vieira[2], Thomas R. Maier[3], Thomas J. Baum[3] and Cynthia Gleason[1*]

[1]Department of Plant Pathology, Washington State University, Pullman, WA, United States of America

[2]Mycology and Nematology Genetic Diversity and Biology Laboratory, USDA Agricultural Research Service, Beltsville, MD, United States of America

[3]Department of Plant Pathology, Entomology and Microbiology, Iowa State University, Ames, IA, United States of America

#Current Address: Simplot Plant Sciences, J. R. Simplot company, Boise, ID United States of America

¶Equal contribution
*Corresponding author cynthia.gleason@wsu.edu

Short title: Gland-specific effector identification in *Meloidogyne chitwoodi*

Authors summary

The root-knot nematode *Meloidogyne chitwoodi* is a major problem for potato farmers in the western U.S., reducing crop yield and quality. These nematodes produce special proteins, called effectors, in their esophageal glands, which help them infect plants. Since these glands are the main source of effectors, we isolated them from juvenile nematodes and analyzed their gene expression. This helped us improve the nematode's genome map and identify genes specific to the glands. From this study, we found 111 potential effector genes. One of them, *Mc15g003960*, was highly active before the nematode started feeding. When we introduced this gene into *Arabidopsis* plants, the nematodes caused more damage, but it didn't seem to weaken the plant's basic defense system. This suggests *Mc15g003960* is not suppressing plant defenses and has a different role in helping the nematode with successful infection. Overall, our approach helped us identify key effectors more efficiently, which could lead to better ways to manage nematode infestations in the future.

Abstract

The root-knot nematode (RKN) *Meloidogyne chitwoodi* is a threat for potato production in the western United States, negatively impacting potato yield and product value*. Meloidogyne chitwoodi* produce proteins, called effectors, in their esophageal glands that are secreted during parasitism and play integral roles in plant-nematode interactions. Because the esophageal glands are the main effector secretory organ, we performed juvenile gland isolation and gland transcriptome analysis. The data allowed us to improve the *M. chitwoodi* genome annotation. Additionally, the gland-specific transcriptome data gave us an enrichment of gland-localized genes, which was validated by *in situ* hybridization. The gland transcriptome analysis led to the identification of 111 effector candidates. One of the effectors, Mc15g003960, which was highly expressed in the pre-parasitic J2 gland tissue, was further characterized. Expression of Mc15g003960 in Arabidopsis resulted in increased galling by *M. chitwoodi*. However, the ectopic expression of Mc15g003960 in planta did not suppress defense-related callose deposition, suggesting that this effector might be involved in processes other than interfering with plant basal defense responses. Our data shows that using the gland transcriptome, good quality genome annotation and stringent criteria, we can increase the efficiency of effector identification, which can be used to develop more sustainable management tools.

Introduction

Root-knot nematodes (RKN, *Meloidogyne* spp.) cause substantial damage to crop plants, amounting to millions of dollars in agricultural losses each year [1]. *Meloidogyne chitwoodi* is a RKN that is endemic to the potato-producing regions of Oregon, Idaho, and Washington, where over half of the USA potatoes are grown [2, 3]. This species is problematic for potato growers because the nematodes can infect both the roots and tubers. The tuber infections result in potatoes with rough, bumpy appearance and brown spots under the skin, and there is little industry tolerance for such potato blemishes. Moreover, *M. chitwoodi* is considered a quarantine pest in certain export markets for potatoes grown in the USA, and their presence in a shipment bound for such a market could mean that the entire shipment is rejected [4]. There is currently no genetic resistance available in commercially available potato cultivars [5]. To develop new strategies of *M. chitwoodi* management, we need information about how the nematodes are successful at parasitism, which calls for investigation into nematode parasitism genes, also known as effector genes.

Nematode effector genes encode proteins that facilitate successful infections, either by suppressing the plant immune responses and/or by altering the plant cell metabolism to form the nematode feeding site [6]. The expression of effector genes may occur at any life stage. Following embryogenesis and initial development in the egg stage, the RKN will hatch as second-stage juvenile (J2). This is the motile infective life stage that must enter the plant root and migrate to the vasculature to establish its feeding site. As a result, the J2 releases effector proteins, which include cell wall-degrading enzymes, such as cellulases and polygalacturonases. Once inside the root, the J2 induces root

galling and will choose between 6-10 plant cells to convert into its feeding sites called

giant-cells [7]. The plant cells that form into giant-cells undergo synchronous waves of

mitotic activity uncoupled from cytokinesis, and they are transcriptionally active, resulting

in enlarged, multinucleate giant-cells that the nematodes rely on for nutrition. After initially

feeding, the J2 will quickly molt 3 more times into the adult stage. The adult female will

feed on the giant-cells and produce eggs in a gelatinous matrix on the surface of the root.

Because nematode parasitic success requires root penetration and giant-cell

formation, the effectors secreted during early stages of parasitism, in the pre-parasitic

nematode J2 and during early feeding site formation (parasitic J2), are critical [8]. The

pre-parasitic J2 faces the challenge of penetrating the roots and migrating towards the

vascular cylinder without triggering insurmountable plant defense responses. In addition,

the effectors must suppress any defenses, such as those inadvertently triggered by the

pathogen associated molecular patterns (PAMPs), such as ascaroside #18 [9]. These

basal plant defenses, known as PAMP-triggered immunity (PTI), are counteracted by

nematode effectors that specifically inhibit PTI responses [8, 10].

The identification of J2 effectors is not only interesting from the viewpoint of

understanding nematode parasitism, but these effectors could also be targets for gene

silencing as a possible future nematode control strategy. As a result, we have focused

our current efforts on the pre-parasitic J2 transcriptomes. Previously, we studied the

motile *M. chitwoodi* J2 and sedentary parasitic J2 life-stages and found 2,248 genes up-

regulated in expression in parasitic nematodes in potato roots at 8 days post-inoculation

(dpi) compared to the motile, pre-parasitic J2 life stage [11]. Of these 2,248 genes, 269

contained a signal peptide (SP) sequence and no transmembrane (TM) domain(s). These

two characteristics are key criteria used in effector discovery pipelines because they

suggest the genes encode secreted proteins. While bioinformatic predictions based on

the presence of a signal peptide sequence and absence of transmembrane domains can

help identify potential effectors, additional confirmation is typically needed. This is usually

done through in situ hybridization (ISH) assays. The ISH assays are performed for each

gene of interest to determine whether the transcript is localized to the secretory organ.

Most nematode effectors are found to be made in the esophageal glands and then

secreted out of the nematode stylet into the plant [12]. Thus, the presence of an ISH probe

signal localized in the glands of the nematode is indirect confirmation of secretion.

Unfortunately, such assays are laborious, and in previous experiments, only about 20%

of the transcriptionally upregulated genes in whole J2 transcriptome analysis that contain

signal peptide sequence and no transmembrane domain could also be confirmed to be

gland localized by in situ hybridization assays [13]. As a result, whole gland

transcriptomes may offer a more gland-targeted approach to effector discovery. Whole

gland transcriptomes have been optimized from isolated glands for pinewood, cyst, root-

knot and root-lesion nematodes, and have been useful in identifying nematode

effectoromes [14-19].

Here we have undertaken a gland transcriptome analysis of *M. chitwoodi* pre-

parasitic J2s. This data has allowed us to improve *M. chitwoodi* genome annotation and

identify effector candidates that would be gland localized and relevant for early stages of

parasitism. The gland transcriptome has shown that pre-parasitic J2s express known and

putative novel effectors. We selected a few effectors (both known and novel) and

confirmed their gland localization by ISH, indicating that the gland library transcriptomes

can provide sets of highly confident effector candidates. By identifying and characterizing

novel effectors through gland transcriptome analysis, we better understand nematode

parasitism, which will ultimately facilitate developing effective management strategies for

this economically damaging nematode species.

Results

Evidence-based genome annotation improved the existing annotation of the *M. chitwoodi* genome

To better predict *M. chitwoodi* protein-coding genes from transcriptome analyses,

the first step was to improve the *M. chitwoodi* genome annotation (S1 Fig). To reannotate

the genome, the *M. chitwoodi* J2 RNA-seq data [11] was combined with the newly

generated *M. chitwoodi* pre-parasitic J2 gland-specific RNA-seq data (this paper). In

addition, to reannotate gene structure, the sequences of proteins from Invertebrate

OrthoDB [20] were used with the sequences of proteins identified from: 1) three RKN

species (*M. javanica, M. arenaria, M. incognita*), 2) more distantly related PPN,

Heterodera schachtii (cyst nematode), 3) animal-parasitic nematodes *Trichinella spiralis,*

Loa loa, Brugia mallayi, Necator americanus, and 4) free-living nematodes

Caenorhabditis remanei, C. elegans, and C. briggsae. Using this approach, a total of

12,205 *M. chitwoodi* genes were predicted. This number is similar to the 12,295 coding

genes annotated for the publicly available *M. chitwoodi* genome assembly (Table 1).

However, a high annotation BUSCO score of *M. chitwoodi* reannotated genome was

achieved, meaning a better completeness of gene content compared to that of the publicly

available genome (Table 1).

To further assess the quality of our new *M. chitwoodi* annotation, a species shared

and specific gene family (orthogroup) analysis was performed using Orthofinder [21]. We

compared our newly reannotated *M. chitwoodi* genome to the publicly available *M.*

chitwoodi and *M. arenaria* genome annotations. *M. arenaria* was chosen because it has

the best publicly available RKN genome annotation, as shown by its BUSCO score (Table

1). There were 1,823 orthogroups shared between the two versions of *M. chitwoodi*

annotations. However, the analysis indicated that there were 2,153 shared orthogroups

between our *M. chitwoodi* genome annotation and *M. arenaria* genome annotation that

were missing in the publicly available *M. chitwoodi* genome annotation (Fig 1). This

indicates that our new annotation has a more complete gene family profile compared to

the current available *M. chitwoodi* genome annotation.

Table 1. Genome annotation features for root-knot nematodes (RKNs). The table shows a comparison

between previous gene annotations for RKN and the current Meloidogyne chitwoodi annotation. BUSCO

scores were calculated based on eukaryota_odb10 dataset.

SPECIES	SOURCE	COMPLETE	SINGLE	DUPLICATED	FRAGMENTED	MISSING
M. chitwoodi	This paper	70.8	68.1	2.7	3.8	25.4
M. chitwoodi	PRJNA666745	49.3	48	1.3	6.8	43.9
M. arenaria	PRJEB8714	74.3	8.6	65.7	4	21.7
M. javanica	PRJEB8714	73.2	14.2	59	4.6	22.2
M. incognita	PRJEB8714	73.2	12.1	61.1	4.2	22.6

Fig 1. The newly re-annotated *M. chitwoodi* genome shares more orthogroups with *M. arenaria* than with the previous *M. chitwoodi* genome annotation. Venn diagram showing the shared gene families (orthogroups) between the newly re-annotated *M. chitwoodi,* current version *M. chitwoodi*, and *M. arenaria* genomes annotations.

We noticed that several putative effectors were not properly predicted by the current annotation, such as *M. chitwoodi* homologue of MiPG1, a polygalacturonase with a signal peptide and fibronectin and glycoside hydrolase 28 domains [22]. A BLASTp search for homologues using the Wormbase ParaSite resulted in the gene Mc1_00280, which is predicted to have a kinase domain (Fig 2A). Alignment of Mc1_00280 and homologues from other RKNs show that only *M. chitwoodi*'s genome predicts a kinase domain (Fig 2A). Further alignment of the gland transcriptome reads with the publicly available genome annotation shows that there are no reads present for the region comprising the predicted kinase domain (Fig 2B). The new genome annotation predicts

Fig 2. Genome reannotation improved the *M. chitwoodi* gene structure prediction.
A. Protein alignments of polygalacturonases from *M. chitwoodi* (Mc1_00280), *M. arenaria* (MA_PG1), *M. enterolobii* (ME_PG1), *M. incognita* (MI_PG1) and *M. javanica* (MJ_PG1), showing that only *M. chitwoodi*'s current genome annotation predicts a protein with a kinase domain. **B.** Alignment of gland transcriptome reads, current genome annotation (in blue) and new annotation (in orange), validating the gene structure predicted by the new annotation. Domains are highlighted in dashed boxes and domains that differ from the others in *Meloidogyne* spp. are in red and blue. Protein alignment was performed using Genious software and transcriptome alignment to the genome was performed using IGV. Created in BioRender. Teixeira, M. (2025) https://BioRender.com/o29t552

that the region has two distinct transcripts: a polygalacturonase with its expected canonical domains (Mc1g003100.t1) and a casein kinase (Mc1g003100.t2). This shows that the polygalacturonase expressed in the gland transcriptome indeed does not include a kinase domain and that the second gene that was further annotated, the casein kinase (Mc1g003100.t2), is not expressed in the glands of pre-parasitic J2s and, for that reason, we cannot observe any reads in our gland transcriptome.

228

M. chitwoodi gland transcriptome analysis revealed putative effectors

231

Pre-parasitic J2s esophageal gland transcriptomes generated between 18.3 to

43.3 million high quality raw reads in each of three bio-replicated cDNA libraries. Around

0.7 to 6.8 million reads were mapped to the M. chitwoodi genome (S1 Table). The reads

mapped to 5,213 transcripts present in all three biological replicates. Although the number

of mapped reads appears relatively low, it is along the same percentage of mapped reads

found in the Pratylenchus gland library [14] and may reflect: a large proportion of read

mapping ribosomal genes and/or the stringency of the gland transcript analysis pipeline

in mapping to the genome. Most of the plant-parasitic nematode (PPN) effectors are

secreted through the canonical ER-Golgi secretory pathway and therefore, characterized

by the presence of a predicted signal peptide and lack of a predicted transmembrane

domain [12]. Using these criteria, 348 putative effectors were identified and represented

in this first gland library generated for M. chitwoodi (S2 Table)

Using our new M. chitwoodi genome annotation, we revisited the previously

published J2 transcriptome data [11]. The new analysis resulted in 10,001 transcripts

identified from the whole pre-parasitic J2 transcriptome, compared to the 5,213 transcripts

identified from the gland cells transcriptome. Using the criteria that effectors must have

signal peptide sequence and no transmembrane domain(s), 783 possible effectors were

retrieved from the J2 transcriptome (Fig 3). The comparison between the gland and the

250

Fig 3. Transcriptome analysis summary. A. Comparison between the number of transcripts retrieved from whole J2s and their glands. **B.** Venn diagram showing number of overlapping putative effectors identified in pre-parasitic J2 and their glands. Created in BioRender. Teixeira, M. (2025) https://BioRender.com/f42i870

256
257

whole J2 transcriptomes revealed 337 genes in common between the gland the J2

transcriptome libraries (Fig 3, S3 Table). The 337 genes were subjected to a BLAST

homology search and there were 18 genes identified that were homologous to previously

described RKN effectors (S4 Table). The normalized read counts (transcripts per million,

TPM) of all 18 known effectors were assessed, and Mc14g000910, a putative homolog

of the known effector MiXyl1 [23], had the lowest read count from the gland transcriptome.

The expression of this gene was used as the threshold of minimum effector gene

expression. From the initial 337 putative effectors, 111 effector candidates (S5 Table)

passed the following criteria: 1) gene expression is above the minimum expression level

set by Mc14g000910; 2) encode proteins with predicted secretion signal sequences; 3)

encode proteins with no predicted TM domains. Among these 111 candidate effectors,

72 contained known protein domain sequences, including sequences for calreticulin,

glycoside hydrolase, c-type lectin-like, peptidase, which are protein domains found in

plant-parasitic nematode effectors (S2 Fig). Thirty-nine out of 111 candidate effectors

have no identified domain(s) based on Interpro or CDD (NCBI Conserved Domain

Database) searches (S6 Table). Twenty-six had protein sequences with more than 50%

identity to proteins from *M. graminicola* or *M. enterolobii*, or both. The remaining 13 had

no significant similarity to any proteins from other organisms, indicating they may be *M.*

chitwoodi specific (S6 Table). The 13 *M. chitwoodi* specific putative effectors were further

analysed by comparing number of reads for each in the pre-parasitic J2s and the parasitic

J2 [11]. From these 13 putative effectors, 7 were more highly expressed in the pre-

parasitic J2 stage compared to the parasitic life stage (8 dpi in potato roots) (Table 2)

To confirm our *in-silico* analysis of the gland library, the expression of 6 genes was

quantified by qRT-PCR in three life stages (eggs, pre-parasitic J2s and parasitic

nematodes at 4 dpi) (Table 2). First, we looked at gene expression of three known

effectors identified in our gland library: Mc5g006680 and Mc1g003100, which are

glycoside hydrolases with high similarity to MiENG1 [24] and MiPG1 [22], respectively,

and Mc5g002180, which has similarity to the described venom allergen protein Mi-msp-

1 [25]. Next, we looked at the expression of Mc11229, a proteinase-inhibitor like, which

was previously characterized as a gland localized effector from *M. chitwoodi* [26]. Lastly,

we measured gene expression for two putative effectors from the gland library analysis.

One was an effector called Mc4g000590 that had putative homologs in *M. graminicola*

and *M. enterolobii*. The second was Mc15g003960, which was *M. chitwoodi* specific

based on BLAST analysis and shows one of the highest fold change differences in the

comparison between the pre-parasitic J2s and the parasitic stage (8 dpi in potato roots)

(Table 2).

Table 2. Transcripts of putative *Meloidogyne chitwoodi* specific effectors, with unknown domains. Table
shows transcripts per million (TPM) from pre-parasitic J2s and for parasitic nematodes at 8 dpi. Fold
change was calculated relative to pre-parasitic J2 TPMs.

*Transcript Mc20.g000500 was not expressed at 8dpi.

Gene id	J2	8 dpi	Fold change
Mc15.g003960	328	38	0.1
Mc20.g000500	266	0	-*
Mc17.g000180	209	107	0.5
Mc17.g001050	198	96	0.5
Mc15.g000880	153	24	0.2
Mc4.g003190	69	6	0.1
Mc11.g005210	62	1825	29.3
Mc1.g006580	27	37	1.4
Mc14.g000410	33	105	3.2
Mc19.g000680	49	641	13.1
Mc17.g000220	1441	1300	0.9
Mc19.g000260	101	187	1.9
Mc7.g004450	354	1037	2.9

All 6 genes whose expression was investigated by qRT-PCR were highly

expressed in pre-parasitic J2 stage compared to the egg stage (Fig 4). The three known

effector genes (the two glycoside hydrolases [22, 24] and a Mi-msp-1 [25] exhibited the

highest level of expression in parasitic nematodes (4 dpi in potato) compared to eggs and

pre-parasitic J2s. Transcript abundance of two novel effectors, Mc4g000590 and

Mc15g003960, and the previously identified effector Mc11229 were significantly higher in

pre-parasitic J2s relative to the eggs and parasitic nematodes (4 dpi).

Fig 4. Relative expression of selected effector candidate genes at three
nematode life stages (egg, pre-parasitic J2 and parasitic J2 at 4 dpi). Data is
presented as the mean of fold change of two biological replicates +/- standard deviation
(SD).

To further validate the gland library, ISH were performed on these six genes (Table

3, Fig 5). The anti-sense probes for all six genes localized to the esophageal gland region

of the J2, making them "candidate effectors". To test the stringency of our effector

discovery method, we tested two potential effectors (presence of SP, no TM domain) that

did not pass our threshold of expression in gland tissue, Mc6g004130 and Mc8g001490.

The ISH showed these transcripts were not localized to the glands, i.e., Mc6g004130

localized to the hypodermis and Mc8g001490 localized to two punctate structures

posteriorly to the nerve ring (Fig 5). Thus, only the putative effectors that pass our gland

specific criteria were proven to be expressed in the glands, validating the gland

transcriptome data and our effector discovery method.

Table 3. Transcripts chosen for further characterization. Domains were predicted using Interpro and NCBI

Conseved Domain Database. Putative orthologs were obtained performing BLASTn searches on NCBI and

WormBase parasite.

Gene id	Predicted domain	Putative orthologs
Mc4.g000590	Unknown	-
Mc15.g003960	Unknown	-
Mc16.g000220	Protease inhibitor-like	Mc11229 [26]
Mc5.g006680	Glycoside hydrolase family 5	MiENG1[24]
Mc1.g003100	Glycoside hydrolase family 28	MiPG1[22]
Mc5.g002180	VAP domain	Mi-msp1[25]

329
330

Fig 5. Representative photos of *in situ* hybridizations of a digoxigenin-labeled antisense probes for transcripts expressed in the J2s A-F, gland localized effectors. **A.** scaffold4.g000590 (novel *M. chitwoodi* effector, with putative homologs in *M. enterolobii* and *M. graminicola*), **B.** scaffold15.g003960 (*M. chitwoodi* specific), **C.** scaffold5.g006680 (putative glycosyl hydrolase with high similarity to MiENG1), **D.** scaffold1.g003100 (putative glycosyl hydrolase with high similarity to MiPG1), **E.** scaffold17.g005060 (effector with a cysteine-rich secretory proteins, antigen 5, and pathogenesis-related 1 proteins domain, with similarity to MiVAP), **F.** scaffold1.g000780 (described by Roze et al. 2018). **G.** scaffold6.g004130 and **H.** scaffold8.g001490, transcripts that didn't pass our effector criteria for gland expression.

The novel effector, Mc15g003960, facilitated *M. chitwoodi* parasitism.

Mc15g003960 was further investigated because it was discovered as a *M. chitwoodi* specific effector in the gland transcriptome analysis and confirmed to be gland localized by ISH. Moreover, the gene is more highly expressed in the pre-parasitic J2 stage compared to other life stages, as found in our whole J2 transcriptome (Table 2) and in the qRT-PCR expression analyses (Fig 4).

Mc15g003960 is a 438 bp coding sequence that encodes a 145 amino acid protein. The signal peptide sequence cleavage site was predicted to be between the amino acids 26 and 27. To characterize its role in *M. chitwoodi* parasitism, two independent Arabidopsis transgenic lines were generated that expressed *Mc15g003960* without its signal peptide *(Mc15g003960Δsp),* driven by the CaMV35S promoter. The transformation events in the two transgenic lines were first confirmed with genomic PCR and the expression of the *Mc15g003960* in the two lines was measured by qRT-PCR (S1 Fig). The two independent Arabidopsis lines expressing *Mc15g003960Δsp* were infected with *M. chitwoodi* and assessed at 15 dpi for root galling. The fresh shoot and root weight of the two transgenic lines showed no significant difference as compared to wildtype plants (S3 Fig), suggesting that the expression of *Mc15g003960Δsp* has no effects on plant growth or development. Remarkably, both transgenic lines exhibited more galling compared to the wildtype plants at 15 dpi (Fig 6).

We hypothesized that the transgenic lines expressing *Mc15g003960Δsp* showed enhanced susceptibility to *M. chitwoodi* infection because the effector might be suppressing PTI. We had previously identified a RKN effector that can block defense

Fig 6. Expression of Mc15g003960 in Arabidopsis makes the plant more susceptible to *M. chitwoodi* infection. Two independent lines of Arabidopsis expressing Mc15g003960 (1.3 and 1.12) and the wildtype control plants (Col-0) were inoculated with *M. chitwoodi* Race 1 J2s. At 14 dpi, galls per plant were counted at 14 dpi. Data represents the mean number of galls per plant +/- SD (Mann-Whitney test, error bars represent SD, ns: not significant p-value>0.05, ***p-value ≤0.001, ****p-value ≤0.0001). N= 15. Experiment was repeated three times with similar results.

related callose deposition [27]. Callose deposits in leaves can be quickly triggered by the bacterial

defense elicitor flg22 [28]. To determine if *Mc15g003960* would affect flg22-induced callose

deposition in leaves, flg22 was infiltrated into the leaves of wildtype and *Mc15g003960Δsp*

transgenic lines. Subsequently, flg22-induced callose deposits were detected using aniline blue

staining. The leaves from both *Mc15g003960Δsp* transgenic lines showed similar levels of callose

deposits when compared with the wildtype after flg22 treatment. These results suggest that flg22-

triggered callose response in *Arabidopsis* leaves was not affected by the constitutive expression

of *Mc15g003960* (Fig 7).

Fig 7. Mc15g003960 does not suppress callose deposition triggered by flg22. A. Quantification of aniline blue-stained callose deposits in the leaf tissue of the *Arabidopsis* plants expressing Mc15g003960 and wild-type plants after flg22 infiltration, calculated using ImageJ software. **B.** Aniline blue staining shows that callose deposition was equivalent in the leaves of transgenic *Arabidopsis* plants expressing Mc15g003960 compared with wild-type plants after flg22 treatment. (n>25).

Discussion

Proper prediction of protein-coding genes in nematode genomes is essential for

effector identification and subsequent characterization. Thus, generating new

transcriptome data provides ample opportunity for enhancing the genome annotation of

plant-parasitic nematodes. Among RKNs, *M. arenaria* (ASM90000398v1) currently has

the best annotated genome for a published genome based on a BUSCO score of 74.3%,

while *M. chitwoodi* (ASM1518303v1) presents the lowest BUSCO score of 49.3% (Table

1). Improving the genome annotation results in adequate gene structure prediction. A search for effectors using the current annotation indicated incorrect prediction for a few well characterized effectors from other RKNs. The genome re-annotation can benefit from both transcriptome and proteome data, which confer more robustness to annotation prediction [29]. Therefore, to improve *M. chitwoodi* genome annotation, we combined our previously published J2 transcriptome and the current gland transcriptome datasets, with protein datasets from other PPNs, animal-parasitic nematodes, and free-living nematodes. Although the number of predicted genes, 12,205, remains very similar to the current annotation (12,295), we achieved a better prediction of gene structure, as shown by the new BUSCO score of 70.8%. The quality of this updated *M. chitwoodi* annotation was further assessed by comparing the othogroups of two annotation versions with those of the high-quality *M. arenaria* genome. The newly annotated genome identified 2,153 orthogroups between *M. chitwoodi* and *M. arenaria* that were absent in the previous version, indicating a more complete *M. chitwoodi* genetic makeup.

The whole pre-parasitic J2 transcriptome analysis indicated that there were 783 possible effectors, following the criteria of coding for proteins with a signal peptide sequence and no transmembrane domains. However, it is important to note that the majority of known effectors are mainly produced in the nematode esophageal glands and secreted via the stylet [12]. Therefore, focusing on genes expressed within the nematode glands would be a more efficient way to identify putative effectors relevant for successful parasitism. For instance, gland transcriptomes generated for other nematode species, such as *Globodera rostochiensis*, *Pratylenchus penetrans, Radopholus similis, Bursaphelenchus xylophilus*; *M. incognita*, *H. schachtii*, and *H. glycines* [14, 15, 17, 18,

30-32], reveal large sets of highly confident effector candidates. In the case of *M. chitwoodi*, 337 of the 783 transcripts coding for secreted proteins were found in both the whole J2 and the gland transcriptomes. This suggests that, akin to other PPNs, the transcriptome of the glands library offers a more refined and targeted pool of potential effector candidates. By focusing on this specific subset, it allows for a more precise identification of key molecular players, providing a higher level of specificity compared to broader approaches. It should be noted that not being gland specific does not preclude genes from being effectors as they may be secreted from nematode organs other than the stylet.

A more conservative criterion was applied by considering only genes with expression levels exceeding that of the known effector with the lowest expression level, leading to a final set of 11 putative genes. These represent around 2% of the 5,213 total transcripts initially retrieved from our gland transcriptome. Other publications found that putative effectors represent 2.42% [30] and 7.5% [31] in *M. incognita* glands, and 2% in *P. penetrans* gland transcriptomes [14]. These variations might reflect the different bioinformatic strategies deployed for effector discovery. Our approach used known effectors in RKN to define a minimum level of effector gene expression. This stringent filtering step may exclude true effectors that have expression lower than this cutoff but allowed us to narrow our list of effectors to a more manageable number of candidates.

Among our final candidates, the majority (65%) encode proteins with known domains, many of which have been found in other RKN species. The remaining transcripts (35%) are novel proteins with unknown domains, from which 12% encode *M. chitwoodi* specific proteins and 23%, proteins with similarity to *M. enterolobii* or *M.*

graminicola, or both. Thus, our results show a small proportion of candidate genes

encoding pioneer sequences, which suggests that most of the effectors were present

before diverging from the last common ancestor, unlike *Pratylenchus*, which contains a

large number (57%) of unique effectors [14].

To characterize novel effectors from *M. chitwoodi* involved in the early stages of

parasitism, we analyzed their expression using our previously published transcriptome

data from whole pre-parasitic and parasitic J2s [11]. Among the thirteen *M. chitwoodi*-

specific effectors, six were found to be highly expressed in pre-parasitic J2s compared to

parasitic nematodes. Of these, Mc15g003960 was selected for further study. When

expressed in *Arabidopsis*, Mc15g003960 increased the plant's susceptibility to *M.*

chitwoodi. While many root-knot nematode effectors are known to suppress plant defense

responses, Mc15g003960 did not inhibit flg22-induced callose deposition, suggesting it

may target other aspects of PTI or perform additional roles in facilitating nematode-plant

interactions. Future studies will further explore Mc15g003960's role in nematode

parasitism and identify its interacting plant proteins.

Materials and Methods

Reannotation of *M. chitwoodi* genome

M. chitwoodi softmasked genome (ASM1518303v1) was obtained from Wormbase

ParaSite database under Bioproject ID: PRJNA666745 [5]. Genome quality was

assessed using eukaryote_odb10 dataset [32]. The genome annotation was conducted

at the high-performance computer cluster "Kamiak", at Washington State University and

a pre-built genome structural and functional annotation workflow [33] was adapted in this

study. In brief, GEMmaker (v2.1.0) [34] was used to map *M. chitwoodi* gland

transcriptome generated from this study and previously generated *M. chitwoodi* J2 RNA-

seq data [11] to the softmasked genome. The generated bam files were used as hint files

for BRAKER1 [35] an RNA-seq based genome structural annotator. As suggested by

Gabriel *et al.* 2021 [36], the annotation accuracy can be enhanced by running the protein

homology-based genome annotator BRAKER2 [37]. Therefore, Invertebrate OrthoDB

[20], *Heterodera schachtii, M. arenaria, M. javanica, M. incognita, Trichinella spiralis, Loa*

loa, Brugia mallayi, Necator americanus, *Caenorhabditis remanei, C. elegans, and C.*

briggsae protein datasets were used as hints for BRAKER2 to annotate *M. chitwoodi*

genome. Later, TSEBRA [36] was used to combine the BRAKER1 and BRAKER2 results.

To further improve the annotation, EviAnn was used to annotate genes purely based on

the same set of protein and rna-seq data used previously. Bedtool2 intersect [38] was

used to combine the BRAKER, EviAnn, and previous annotation [5]. The final structurally

annotated *M. chitwoodi* genome assembly was fed into EnTAPnf [39] for gene functional

annotation with gene ontology assignment. The completeness of functionally annotated

genome assembly was assessed by BUSCO using eukyrote_odb10 dataset [32]. The

enrichment of gene expression in the gland transcriptome was visualized using

Integrative Genomics Viewer (IGV) software (v. 2.10.3) [40]. The species-specific gene

family (orthogroup) analysis was adapted from Zhang et al. (2022) [21]. Orthofinder [41]

was used to assign each gene to a protein family.

Transcriptomic analysis

Five to 10 esophageal glands were extracted from pre-parasitic second stage

juveniles of *M. chitwoodi* race 1 and pooled to form one sample [42]. Libraries were

prepared using Takara Bio's SMART-seq mRNA LP (634768) kit, which includes RNA

isolation and library preparation. Three libraries were sequenced by Novogene, using

Illumina Novaseq 6000 (PE 150bp with 250M read depth).

Raw sequencing data were process using the gene expression analysis workflow

GEMmaker (v2.1.0) [34]. STAR aligner implemented in GEMmaker was used to map the

transcriptomes to the *M. chitwoodi* race 1 draft genome [5]. The resulting sequence

mapping quality report and normalized gene count matrix were used in this study.

511   Prediction of putative effectors

For possible effector gene identification from all transcripts, signal Peptides were

predicted using SignalP6.0 [43, 44] and transmembrane domains were predicted using

DeepTMHMM1.0 [45], both using default settings. Domain prediction was performed

using Interpro [46] and NCBI's conserved domain database [47], using amino acid

sequences. To identify homolog of candidate effector gene, BLASTp and BLASTn search

was carried out at the NCBI online servers using non-redundant protein dataset and the

expressed sequence tags (ESTs) database respectively using the default parameter.

Manual curation of putative effector sequences was performed aligning the gland

transcriptome data to the genome annotation using Integrative Genomics Viewer (IGV)

software v. 2.10.3) [42].

523   Cloning and transgenic lines generation

Mc15g003960 coding sequence, without its predicted signal peptide sequence

(*Mc15g003960Δsp*), was amplified from *M. chitwoodi* race 1 cDNA by PCR (S7 table)

and inserted into the Gateway donor vector pDONR207. It further was recombined into

Gateway destination vector pB2GW7 to produce 35S::*Mc15g003960Δsp*.After

sequencing, the plasmid was used to transform *Agrobacterium tumefaciens* GV3101,

which was used to transform Arabidopsis Col-0 following the floral dip protocol of Clough

and Bent (1998) [48]. Homozygous T3 seeds from two independent transgenic lines were

collected from T2 lines after segregation analysis on BASTA-containing medium.

Nematode gene expression in the transgenic plants was confirmed by qPCR (S7 Table).

Nematode cultures

535   *M. chitwoodi* race 1 (initially provided by Dr Charles Brown from USDA-ARS,

536   Prosser, WA) was multiplied on susceptible tomato *Solanum lycopersicum* cv. Rutgers

537   under greenhouse conditions. Egg mass collection and J2 hatching was performed as

538   previously described [11].

Nematode infection assays

Arabidopsis seeds were surface-sterilized and sown on Murashige & Skoog

(Duchefa) agar plates (½MS salts, 1% sucrose, 0.8% agar, pH 6.4). Three week old

seedlings were transferred to 500 ml sand-filled cone-tainers kept in growth chamber

(14 h:10 h light:dark cycle, 23 °C). One week after transplanting to sand, the plants were

inoculated with 750 *M. chitwoodi* eggs per plant. At 15 days post-inoculation, the plants

were gently removed from the cone-tainers. Each root system was weighed and the

number of galls per plant was assessed. Nematode infection assays were repeated twice

with similar results.

RNA extraction and Real time PCR validation

Susceptible Russet Burbank plants were propagated in tissue culture for three

weeks and then transferred to 500 ml cone-tainers filled with sand and subsequently

maintained in growth chambers at 14 h:10 h light:dark cycle, 23 °C. Fourteen days after

transplanting to sand, the plants were inoculated with 500 freshly hatched *M. chitw*oodi

J2s. Samples were collected 4 days post inoculation (dpi) and immediately frozen for

RNA extraction.

For gene expression validation, two biological samples of eggs, pre parasitic J2s

and galled potato root tissue at 4 dpi were collected. Total RNA was extracted using the

Trizol method. Briefly, trizol was added to ground sample and RNA was precipitated

overnight using isopropanol. RNA was then purified using ethanol and resuspended in

DEPC water. 1ug of RNA was treated with DNA-freeTM DNA Removal Kit

(ThermoFisher) to remove residual DNA and ProtoScript II First Strand cDNA Synthesis

Kit (New England Biolabs, USA) was used to prepare cDNA. Quantitative PCR was

performed using SsoAdvancedTM Universal SYBR Green Supermix on a CFX96 Real-

Time PCR Detection System (Bio-Rad, USA) and the primers listed in S7 Table. The qRT-

PCR conditions were: 95C for 3 min, 40 cycles for 95C for 15 sec, 53C for 15sec and 72C

for 20 sec, followed by a melting curve analysis from 65C to 95C with 0.5C increments at

5sec. Each experiment consisted of three technical replicates per sample.

Expression levels of genes of interest in nematode were normalized to the

expression of Internal transcribed spacer 2 (ITS2) rRNA, a housekeeping gene from *M.*

chitwoodi [50]. The relative expression levels of each target gene from pre parasitic J2s

and 4 dpi tissue were calculated by comparing with those in eggs using the 2$-\Delta\Delta$Ct

method [49].

Callose deposition assay

Leaves from 4-week-old Arabidopsis plants were infiltrated with 1uM flg22. After

24h, leaves were punched using a cork borer and fixed in 95% ethanol overnight.

Samples were then rehydrated with a serial incubation with 50% ethanol for 30 minutes,

followed by two 30 min incubations with a phosphate-buffered saline, stained with 0.8%

aniline blue for 1h and visualized under an AxioObserver A1 inverted microscope using

a DAPI filter. Callose deposition was quantified with ImageJ [50]. At least 25 leaf punches

were collected from four plants per genotype using a 4 mm cork borer, all images were

taken at 10X magnification and the experiment was repeated twice.

In situ hybridization

Whole mount in situ hybridizations were performed for *M. chitwoodi* J2 following

the protocol of de Boer et al. [51]. Specific PCR primers were used to amplify the genes

of interest from *M. chitwoodi* J2 cDNA (S7 Table). The resulting PCR products were then

used as a template for generation of sense and antisense DIG-labeled probes using a

DIG-nucleotide labeling kit (Roche). Hybridized probes within the nematode tissues were

detected using an anti-DIG antibody conjugated to alkaline phosphatase and its

substrate. Nematode segments were observed using a Nikon Eclipse 5*i* light microscope.

Statistics

595 Statistical tests were performed using GraphPad Prism version 8.00 for Windows

596 (GraphPad Software, La Jolla California, USA).

Acknowledgements

599 All computation works were performed using Dr. Stephen Ficklin's "Computational

600 Biology Lab" node resource and "Kamiak" HPC at Washington State University, US.

Competing interests

603 This study declares no competing interests.

Data Availability

606 The *M. chitwoodi* RNA-seq data have been deposited at the NCBI database under

607 BioProject ID PRJNA1190951 and the M. chitwoodi genome re-annotation is found

608 at **10.5281/zenodo.**14542289.

References

1.   Nicol, JM et al. Current nematode threats to world agriculture. In: Jones, J., Gheysen, G. and Fenoll, C., Editors, Genomics and Molecular Genetics of Plant-Nematode Interactions. Dordrecht: Springer; 2011. pp 21-43. https://doi.org/10.1007/978-94-007-0434-3_2

2.   Santo GS, O'bannon JH, Finley A, Golden AM. Occurrence and host range of a new root-knot nematode (*Meloidogyne chitwoodi*) in the Pacific northwest. Plant Disease. 1980;64:951-2.

3.   Zasada IA, Kitner M, Wram C, Wade N, Ingham RE, Hafez S, et al. Trends in occurrence, distribution, and population densities of plant-parasitic nematodes in the Pacific Northwest of the United States from 2012 to 2016. Plant Health Progress. 2019;20(1):20-8. doi: 10.1094/php-11-18-0077-rs.

4.   Ingham RE, Hamm PB, Baune M, David NL, Wade NM. Control of *Meloidogyne chitwoodi* in potato with shank-injected metam sodium and other nematicides. J Nematol. 2007;39(2):161-8. PubMed PMID: 19259485.

5.   Bali S, Hu S, Vining K, Brown C, Mojtahedi H, Zhang L, et al. Nematode genome announcement: draft genome of *Meloidogyne chitwoodi,* an economically important pest of potato in the Pacific Northwest. Mol Plant Microbe Interact. 2021;34(8):981-6. doi: 10.1094/mpmi-12-20-0337-a.

6.   Vieira P, Gleason C. Plant-parasitic nematode effectors - insights into their diversity and new tools for their identification. Curr Opin Plant Biol. 2019;50:37-43. doi: 10.1016/j.pbi.2019.02.007.

7.   Escobar C, Barcala M, Cabrera J, Fenoll C. Chapter One - Overview of root-knot Nematodes and giant cells. In: Escobar C, Fenoll C, editors. Advances in Botanical Research. 73: Academic Press; 2015. p. 1-32.

8.   Bali S, Gleason C. Unveiling the diversity: plant parasitic nematode effectors and their plant interaction partners. Mol Plant Microbe Interact. 2024;37(3):179-89. doi: 10.1094/mpmi-09-23-0124-fi. PubMed PMID: 37870371.

9.   Manosalva P, Manohar M, von Reuss SH, Chen S, Koch A, Kaplan F, et al. Conserved nematode signalling molecules elicit plant defenses and pathogen resistance. Nat Commun. 2015;6:7795. doi: 10.1038/ncomms8795.

10.   Khan M, Khan AU. Plant parasitic nematodes effectors and their crosstalk with defense response of host plants: A battle underground. Rhizosphere. 2021;17:100288. doi: https://doi.org/10.1016/j.rhisph.2020.100288.

11.   Zhang L, Gleason C. Transcriptome Analyses of pre-parasitic and parasitic *Meloidogyne Chitwoodi* Race 1 to identify putative effector genes. J Nematol. 2021;53. doi: 10.21307/jofnem-2021-084.

12.   Mitchum MG, Hussey RS, Baum TJ, Wang X, Elling AA, Wubben M, et al. Nematode effector proteins: an emerging paradigm of parasitism. New Phytologist. 2013;199(4):879-94. doi: https://doi.org/10.1111/nph.12323.

13.   Vieira P, Maier TR, Eves-van den Akker S, Howe DK, Zasada I, Baum TJ, et al. Identification of candidate effector genes of *Pratylenchus penetrans*. Mol Plant Pathol. 2018;19(8):1887-907.

14.   Vieira P, Shao J, Vijayapalani P, Maier TR, Pellegrin C, Eves-van den Akker S, et al. A new esophageal gland transcriptome reveals signatures of large scale de novo effector birth in the root lesion nematode *Pratylenchus penetrans*. BMC Genomics. 2020;21(1):738. doi: 10.1186/s12864-020-07146-0.

15.   Maier TR, Masonbrink RE, Vijayapalani P, Gardner M, Howland AD, Mitchum MG, et al. Esophageal gland RNA-Seq resource of a virulent and avirulent population of the soybean cyst nematode *Heterodera glycines*. Mol Plant Microbe Interact. 2021;34(9):1084-7. doi: 10.1094/mpmi-03-21-0051-a.

16.   Molloy B, Shin DS, Long J, Pellegrin C, Senatori B, Vieira P, et al. The origin, deployment, and evolution of a plant-parasitic nematode effectorome. PLoS Pathog. 2024;20(7):e1012395. doi: 10.1371/journal.ppat.1012395.

17.   Elashry AM, Habash SS, Vijayapalani P, Brocke-Ahmadinejad N, Blümel R, Seetharam A, et al. Transcriptome and parasitome analysis of beet cyst nematode *Heterodera schachtii*. Sci Rep. 2020;10(1):3315. doi: 10.1038/s41598-020-60186-0.

18.   Rocha RO, Hussey RS, Pepi LE, Azadi P, Mitchum Melissa G. Discovery of novel effector protein candidates produced in the dorsal gland of adult female root-knot nematodes. Mol Plant Microbe Interact. 2023;36(6):372-80. doi: 10.1094/mpmi-11-22-0232-r.

19. Espada M, Eves-van den Akker S, Maier T, Vijayapalani P, Baum T, Mota M, et al. STATAWAARS: a promoter motif associated with spatial expression in the major effector-producing tissues of the plant-parasitic nematode *Bursaphelenchus xylophilus*. BMC Genomics. 2018;19(1):553. doi: 10.1186/s12864-018-4908-2.

20.   Kuznetsov D, Tegenfeldt F, Manni M, Seppey M, Berkeley M, Kriventseva EV, et al. OrthoDB v11: annotation of orthologs in the widest sampling of organismal diversity. Nucleic Acids Res. 2023;51(D1):D445-d51. doi: 10.1093/nar/gkac998.

21.   Zhang H, Wafula EK, Eilers J, Harkess Alex E, Ralph PE, Timilsena PR, et al. Building a foundation for gene family analysis in Rosaceae genomes with a novel workflow: A case study in *Pyrus* architecture genes. Front Plant Sci. 2022;13. doi: 10.3389/fpls.2022.975942.

24.   Rosso MN, Favery B, Piotte C, Arthaud L, De Boer JM, Hussey RS, et al. Isolation of a cDNA encoding a beta-1,4-endoglucanase in the root-knot nematode *Meloidogyne incognita* and expression analysis during plant parasitism. Mol Plant Microbe Interact. 1999;12(7):585-91. doi: 10.1094/mpmi.1999.12.7.585.

22.   Jaubert S, Laffaire JB, Abad P, Rosso MN. A polygalacturonase of animal origin isolated from the root-knot nematode Meloidogyne incognita. FEBS Lett. 2002;522(1-3):109-12. doi: 10.1016/s0014-5793(02)02906-x.

25.   Ding X, Shields J, Allen R, Hussey RS. Molecular cloning and characterisation of a venom allergen AG5-like cDNA from *Meloidogyne incognita*. Int J Parasitol. 2000;30(1):77-81. doi: 10.1016/s0020-7519(99)00165-4.

26.   Roze E, Hanse B, Mitreva M, Vanholme B, Bakker J, Smant G. Mining the secretome of the root-knot nematode *Meloidogyne chitwoodi* for candidate parasitism genes. Mol Plant Pathol. 2008;9(1):1-10. doi: 10.1111/j.1364-3703.2007.00435.x.

23.   Mitreva-Dautova M, Roze E, Overmars H, de Graaff L, Schots A, Helder J, et al. A symbiont-independent endo-1,4-beta-xylanase from the plant-parasitic nematode *Meloidogyne incognita*. Mol Plant Microbe Interact. 2006;19(5):521-9. doi: 10.1094/mpmi-19-0521.

27.   Gleason C, Polzin F, Habash SS, Zhang L, Utermark J, Grundler FM, et al. Identification of
two *Meloidogyne hapla* genes and an investigation of their roles in the plant-nematode
interaction. Mol Plant Microbe Interact. 2017;30(2):101-12. doi: 10.1094/mpmi-06-16-0107-r.
28.   Ellinger D, Voigt C. Callose biosynthesis in Arabidopsis with a focus on pathogen response:
what we have learned within the last decade. Ann Bot. 2014;114. doi: 10.1093/aob/mcu120.
29.   Gabriel L, Brůna T, Hoff KJ, Ebel M, Lomsadze A, Borodovsky M, et al. BRAKER3: Fully
automated genome annotation using RNA-seq and protein evidence with GeneMark-ETP,
AUGUSTUS, and TSEBRA. Genome Res. 2024;34(5):769-77. doi: 10.1101/gr.278090.123.
30.   Rutter WB, Hewezi T, Maier TR, Mitchum MG, Davis EL, Hussey RS, et al. Members of the
Meloidogyne avirulence protein family contain multiple plant ligand-like motifs. Phytopathol.
2014;104(8):879-85.
31.   Huang G, Gao B, Maier T, Allen R, Davis EL, Baum TJ, et al. A profile of putative parasitism
genes expressed in the esophageal gland cells of the root-knot nematode Meloidogyne incognita.
Mol Plant Microbe Interact. 2003;16(5):376-81.. doi: 10.1094/mpmi.2003.16.5.376.
32.   Manni M, Berkeley MR, Seppey M, Zdobnov EM. BUSCO: assessing genomic data quality
and beyond. Curr Protoc. 2021;1(12):e323. doi: https://doi.org/10.1002/cpz1.323.
33.   Zhang H, Ko I, Eaker A, Haney S, Khuu N, Ryan K, et al. A haplotype-resolved, chromosome-
scale genome for Malus domestica Borkh. 'WA 38'. G3 Bethesda. 2024;14(12). doi:
10.1093/g3journal/jkae222.
34.   Hadish JA, Biggs TD, Shealy BT, Bender MR, McKnight CB, Wytko C, et al. GEMmaker:
process massive RNA-seq datasets on heterogeneous computational infrastructure. BMC
Bioinform. 2022;23(1):156. doi: 10.1186/s12859-022-04629-7.
35.   Hoff KJ, Lange S, Lomsadze A, Borodovsky M, Stanke M. BRAKER1: Unsupervised RNA-
Seq-Based Genome Annotation with GeneMark-ET and AUGUSTUS. Bioinformatics.
2015;32(5):767-9. doi: 10.1093/bioinformatics/btv661.
36.   Gabriel L, Hoff KJ, Brůna T, Borodovsky M, Stanke M. TSEBRA: transcript selector for
BRAKER. BMC Bioinform. 2021;22(1):566. doi: 10.1186/s12859-021-04482-0.
37.   Brůna T, Hoff KJ, Lomsadze A, Stanke M, Borodovsky M. BRAKER2: automatic eukaryotic
genome annotation with GeneMark-EP+ and AUGUSTUS supported by a protein database. NAR
Genom Bioinform. 2021;3(1). doi: 10.1093/nargab/lqaa108.
38.   Quinlan AR, Hall IM. BEDTools: a flexible suite of utilities for comparing genomic features.
Bioinformatics. 2010;26(6):841-2. doi: 10.1093/bioinformatics/btq033.
~~39~~.   Hart AJ, Ginzburg S, Xu MS, Fisher CR, Rahmatpour N, Mitton JB, et al. EnTAP: Bringing
faster and smarter functional annotation to non-model eukaryotic transcriptomes. Mol Ecol
Resour. 2020;20(2):591-604. doi: 10.1111/1755-0998.13106.
40.   Robinson JT, Thorvaldsdóttir H, Winckler W, Guttman M, Lander ES, Getz G, et al.
Integrative genomics viewer. Nat Biotech. 2011;29(1):24-6. doi: 10.1038/nbt.1754.
41.   Emms DM, Kelly S. OrthoFinder: phylogenetic orthology inference for comparative
genomics. Genome Biol. 2019;20(1):238. doi: 10.1186/s13059-019-1832-y.
42.   Maier TR, Hewezi T, Peng J, Baum TJ. Isolation of whole esophageal gland cells from plant-
parasitic nematodes for transcriptome analyses and effector identification. Mol Plant-Microbe
Interact. 2013;26(1):31-5. doi: 10.1094/mpmi-05-12-0121-fi.

43.   Almagro Armenteros JJ, Tsirigos KD, Sønderby CK, Petersen TN, Winther O, Brunak S, et
al. SignalP 5.0 improves signal peptide predictions using deep neural networks. Nat Biotechnol.
2019;37(4):420-3. doi: 10.1038/s41587-019-0036-z.
44.   Nielsen H, Teufel F, Brunak S, von Heijne G. SignalP: the evolution of a web server. In:
Lisacek F, editor. Protein Bioinformatics. New York, NY: Springer US; 2024. p. 331-67.
45.   Hallgren J, Tsirigos KD, Pedersen MD, Almagro Armenteros JJ, Marcatili P, Nielsen H, et al.
DeepTMHMM predicts alpha and beta transmembrane proteins using deep neural networks.
bioRxiv. 2022:2022.04.08.487609. doi: 10.1101/2022.04.08.487609.
46.   Paysan-Lafosse T, Blum M, Chuguransky S, Grego T, Pinto BL, Salazar Gustavo A, et al.
InterPro in 2022. Nucleic Acids Res. 2022;51(D1):D418-D27. doi: 10.1093/nar/gkac993.
47.   Wang J, Chitsaz F, Derbyshire MK, Gonzales NR, Gwadz M, Lu S, et al. The conserved
domain database in 2023. Nucleic Acids Res. 2022;51(D1):D384-D8. doi: 10.1093/nar/gkac1096.
48.   Clough SJ, Bent AF. Floral dip: a simplified method for Agrobacterium-mediated
transformation   of   *Arabidopsis   thaliana*.   Plant   J.   1998;16(6):735-43.   doi:
https://doi.org/10.1046/j.1365-313x.1998.00343.x.
49.   Livak KJ, Schmittgen TD. Analysis of relative gene expression data using real-time
quantitative PCR and the 2(-Delta Delta C(T)) Method. Methods. 2001;25(4):402-8. doi:
10.1006/meth.2001.1262.
50.   Schneider CA, Rasband WS, Eliceiri KW. NIH Image to ImageJ: 25 years of image analysis.
Nat Methods. 2012;9(7):671-5. doi: 10.1038/nmeth.2089.
51.   de Boer, J. M., Y. Yan, G. Smant, E. L. Davis, and T. J. Baum. In-situ hybridization to
messenger RNA in *Heterodera glycines.* J Nematol. 1998; 30: 309-12.

Supporting Information

S1 Fig. A schematic workflow of the genome annotation pipelines used in this study. Teixeira, M. (2025) https://BioRender.com/f18v715

S2 Fig. Domain prediction of proteins encoded by putative effectors. Prediction performed using Interpro and NCBI Conserved Domain Database.

S3 Fig. Evaluation of two independent lines expressing Mc15g003960 (1.3 and 1.12). A. transgene expression from each line. **B**. Root weight of the two lines and Col-0 at 14 dpi. Data show mean weight of the roots +/-SD. (Mann Whitney test, error bars represent SD, ns: not significant). N=15. Experiment was repeated three times with similar results.

S1 Table. Summary of Gland transcriptome sequencing and alignment data.

S2 Table. Gland transcripts with SP and no TM domain

S3 Table. Common transcripts between glands and J2

S4 Table. Putative homologs of known effectors

S5 Table. Putative Mc1 effectors after using homologs as cut-off criteria.

S6 Table. Transcripts for which no domain was predicted using Interpro and NCBI Conserved Domain Database.

S7 Table. Primers used in this paper