The development of visual acuity and crowding: Finding the balance between fine detail and 'gist' processing

John A. Greenwood[1], Marilia Kyprianou[1], & Tessa M. Dekker[1,2]

[1]Experimental Psychology, University College London, London, UK
[2]Institute of Ophthalmology, University College London, London, UK

Keywords: Perception, development, vision, crowding, foveal vision.

Corresponding author: John A. Greenwood
Email: john.greenwood@ucl.ac.uk
Website: http://eccentricvision.com

Abstract

The adult visual system is characterised by high-resolution foveal vision and a peripheral field limited by crowding, the disruption to object recognition in clutter that gives a summary 'gist' in place of fine detail. In children, crowding is elevated in the fovea, with estimates of the age where foveal crowding drops to adult-like levels varying widely from 5 to 12+ years. As crowding restricts key processes like reading, better characterisation of this developmental trajectory is critical. We developed methods optimised to measure crowding in children, whereby adults and typically developing children aged 3-13 years (n=119) judged the orientation of a foveal 'VacMan' target either in isolation or surrounded by 'ghost' flankers. Stimulus sizes and separation were scaled adaptively. In the isolated condition, acuity (measured as gap-size thresholds) dropped rapidly to adult-like levels at 5-6 years. Thresholds rose when flankers were added, demonstrating foveal crowding at all ages. These elevations were highest at 3-4 years and persisted at 5-6 years, dropping to adult-like levels at 7-8 years. A meta-analysis of our thresholds and those from 11 prior studies reveals the same overall developmental trajectory, despite widely varying stimuli and procedures, with some divergent estimates likely driven by measurement confounds such as underestimated adult crowding levels. We further demonstrate that developmental foveal crowding shows the same selectivity for target-flanker similarity as peripheral crowding, consistent with common mechanisms. This prolonged development of crowding reveals a shifting balance in the visual system between the processing of fine detail vs. the 'gist' of the scene.

Introduction

The human visual system is *foveated* – our ability to recognise objects is greater in central (foveal) vision than in the periphery. Although the decline in peripheral vision is in part related to reduced acuity (Anstis, 1974), a greater disruption arises through *crowding*, a process whereby surrounding clutter interferes with the recognition of an otherwise visible target object (Strasburger et al., 2011; Whitney & Levi, 2011). Crowding provides a summary or 'gist' of the peripheral field, at the expense of fine detail (Rosenholtz, 2016). In the fovea, where our sensitivity to fine detail is greatest, crowding is typically minimal (Coates et al., 2018). This balance between fine detail and 'gist' differs in children, however, with pronounced elevations in crowding in their foveal vision (Atkinson & Braddick, 1983). These elevations would limit key abilities like reading (Kwon et al., 2007) and visual search (Gheri et al., 2007). This prolonged development is also likely to make the visual system vulnerable to disruption, as seen with the further elevations in crowding in developmental disorders ranging from amblyopia (Levi & Klein, 1985) to dyslexia (Atkinson, 1991). Estimates of the age at which crowding becomes adult-like vary widely, however. Though acuity is adult-like at 5-6 years (Birch et al., 1983; Ellemberg et al., 1999; Leat et al., 2009), estimates of the age of maturity for crowding vary from 5 through to 12 years or later (Atkinson et al., 1987; Jeon et al., 2012). Our aim was to better characterise the developmental trajectory of crowding using procedures optimised to measure crowding in a child-friendly format, and to compare these measures of crowding with those from prior studies to gain a broader view of this developmental process.

In peripheral vision, crowding occurs when flanker objects fall within an interference zone around a target (Bouma, 1970). Interference zones increase in size with eccentricity (Toet & Levi, 1992; Pelli et al., 2004) such that the centre-to-centre separation of objects needs to be beyond around half the target eccentricity to avoid crowding, regardless of object size (Levi, Hariharan, et al., 2002; Pelli et al., 2004). Crowding disrupts the recognition of visual modalities ranging from orientation (Wilkinson et al., 1997) to colour and motion (Bex & Dakin, 2005; Greenwood & Parsons, 2020), and complex elements such as faces (Louie et al., 2007; Kalpadakis-Smith et al., 2018). The mechanisms underlying crowding are contested, with high-level accounts proposing that crowding derives from attentional resolution (He et al., 1996) or Gestalt processes of grouping (Herzog & Manassi, 2015). These high-level accounts cannot explain the systematic nature of crowded errors however, whereby errors made regarding a given target are not random but rather follow the appearance of the

surrounding flankers (Greenwood et al., 2009). Pooling models explain these errors as the unwanted combination of target and flanker elements (Rosenholtz et al., 2019), with population pooling models (Harrison & Bex, 2015) in particular able to predict the errors made regarding foveal stimuli presented to children (Kalpadakis-Smith et al., 2022). It is this pooling that provides the 'gist' of the visual scene, which may be an efficient way to represent information-rich scenes with limited resources (Freeman & Simoncelli, 2011; Rosenholtz, 2016). In children, the elevations in foveal crowding suggest that this drive for pooling and the computation of the 'gist' of the scene occurs more extensively than in adults.

Numerous studies have demonstrated that foveal crowding is elevated in children relative to adults (Atkinson et al., 1987; Semenov et al., 2000; Bondarko & Semenov, 2005; Jeon et al., 2010; Norgett & Siderov, 2011, 2014; Doron et al., 2015). Estimates for the age at which these elevations recede to adult-like levels vary widely, however, as shown in Table 1. The earliest of these estimates suggest that crowding reaches adult-like levels around the age of 5-7 years (Atkinson et al., 1987; Atkinson et al., 1988), obtained using target letters surrounded by flanker letters. Others using letters report a slightly later estimate of 7-9 years (Norgett & Siderov, 2014), aligning with letter-chart based measurements suggesting maturity around 7 years (Doron et al., 2015). Later estimates include measurements with Landolt-C and tumbling-E elements that show maturity at 9 years (Semenov et al., 2000; Bondarko & Semenov, 2005), while others find crowding that is still immature at 11 years, suggesting maturity at 12 or beyond (Jeon et al., 2010). In addition to the above-noted stimulus differences, prior studies vary widely in their stimulus arrangement, with variations in the separation between elements particularly likely to impact the measures. Several studies varied the spacing between target and flankers directly, and quantified performance via edge-to-edge separation (Semenov et al., 2000; Bondarko & Semenov, 2005; Jeon et al., 2010), which is suboptimal given that crowding is driven by centre-to-centre separation (Levi, Hariharan, et al., 2002; Pelli et al., 2004). Others have scaled the size and separation of elements (Atkinson et al., 1987; Norgett & Siderov, 2011, 2014; Doron et al., 2015), though with widely-spaced elements that are less effective at estimating the strength of crowding than arrays with closely separated elements (Song et al., 2014; Facchin et al., 2019; Waugh & Fronius, 2024). To resolve these conflicting conclusions about the development of crowding, we need to re-examine this developmental trajectory using approaches optimised for the measurement of crowding in both children and adults.

Table 1. A comparison of estimates of the developmental trajectory of crowding, including (spoiler alert) the current study. We include only studies where childhood crowding is compared directly against adult or late-adolescent crowding. Where multiple conditions were tested, we include those with the most 'standard' configuration (e.g. a target and 4 adjacent flankers). Red blocks show ages at which crowding levels were significantly elevated above those of adult participants; grey blocks show non-significant differences ('adult-like' performance). Block widths indicate the grouping of participants by age. Example stimuli are depicted for each study, with their centre-to-centre separation in multiples of element diameter, except where this property was directly manipulated (marked as 'variable'). Studies are sorted by the age at which crowding becomes adult-like, excluding the current study.

With this aim, we developed child-friendly procedures adapted from those previously shown to optimally measure crowding in adult peripheral vision. Participants judged the orientation of a videogame character (Visual Acuity Man, or 'VacMan') viewed foveally, either in isolation or surrounded by 'ghost' flankers (Figure 1), as used previously in both typically developing children and those with amblyopia (Greenwood et al., 2012; Kalpadakis-Smith et al., 2022). The advantage of this paradigm is its use of target and flanker elements that are similar enough to induce crowding, but with distinct identities introduced through the videogame-like context in order to enhance task comprehension and minimise source confusion. Given the optimality of measuring crowding as the centre-to-centre separation between closely spaced elements (Song et al., 2014), we measured crowding using scaled elements with a centre-to-centre separation between the target and its flankers of 1.1x the stimulus size. We assessed acuity and crowding using independent measurements, with sizes varied using QUEST and catch trials included to ensure task engagement. A further advantage of this approach is that threshold estimates can be

converted to reveal the spatial extent of crowding, as well as other metrics used to assess crowding previously, which we used to perform a meta-analysis of prior studies.

As well as characterising the developmental trajectory, these methods further allow us to address the debate on whether foveal and peripheral crowding derive from the same mechanism (Levi, Klein, et al., 2002). Although the scale of foveal crowding has been noted to make it difficult to distinguish from masking effects in the adult fovea, the elevated crowding levels in children allow us to more clearly address this question. A key property of crowding in this regard is its selectivity for target-flanker similarity (Kooi et al., 1994) – crowding is reduced when target and flanker elements differ in properties such as contrast polarity, colour and depth. For instance, a black target amongst white flankers gives less crowding than an all-black array (Kooi et al., 1994; Chakravarthi & Cavanagh, 2007; Chung & Mansfield, 2009). In typical adults, these effects have similarly been observed in the fovea (Hess, Dakin, & Kapoor, 2000; Hess, Dakin, Kapoor, et al., 2000), though it is unclear whether the same differences reduce crowding in children. We thus examined the selectivity of crowding for target-flanker similarity in a subset of children, and the developmental trajectory of this selectivity.

We thus sought to examine both the age at which foveal crowding recedes to adult-like levels, and whether foveal crowding shows the same selectivity for target-flanker similarity as observed in peripheral vision. To this end, crowding and its selectivity was measured in children between the ages of 3-13, with comparison to adults conducting the same procedures.

Methods

Design

Children and adults judged the orientation of a black target element, known as Visual Acuity Man ('VacMan'), either in isolation or flanked by four black 'ghost' elements (Figure 1). All participants completed these two conditions, with a subset also completing a third condition where the black target was surrounded by white flankers. Thresholds were measured in each condition to find the smallest gap size in the target element that allowed accurate recognition. When present, flankers were scaled in both size and separation, as described below.

Figure 1. Example stimuli. **A.** Schematic depiction of the monitor display during unflanked trials, with Visual Acuity Man (VacMan) at the centre and four coloured reference ghosts moving around the monitor edge and presented on external cards to the side. **B.** An example target stimulus during unflanked trials. **C.** Example stimuli from the flanked-same condition, in which four same-polarity (black) flankers surrounded the target at random orientations. **D.** An example trial from the different-polarity condition, where flankers were white.

Participants

A total of 125 participants were recruited, including 95 children and 30 adults. All were required to have normal or corrected-to-normal vision, with normal colour-naming abilities (checked using the stimuli prior to participation), and no history of neurological abnormalities. Participants were required to complete at least one block of trials to be included in the study. 3 children failed to meet these criteria, and a further 3 began testing but failed to complete the first condition. All 6 were thus excluded.

The final sample comprised 89 children (49 female) with ages between 3.2-13.1 years (mean: 8.2 years), and 30 adults (20 female, and including the 3 authors), with ages between 18.3-37.6 (mean: 23.1 years). For analysis, children were divided into 5 groups by age, with 10 3-4 year-olds (mean age 3.8 years, range 3.2-4.4), 22 5-6 year-olds (mean age 6.2, range 5.1-6.9), 24 7-8 year-olds (mean age 7.8, range 7.0-8.9), 14 9-10 year-olds (mean age 9.9, range 9.0-10.9 years), and 19 11-12 year-olds (mean age 12.2, range 11.1-13.1 years). Adults were paid for their participation and children rewarded with toys. All participants gave informed assent/consent, with ethical approval from the UCL Experimental Psychology Research Ethics Committee.

Apparatus

Stimuli and experimental procedures were programmed using MATLAB (The MathWorks, Ltd., Cambridge, UK) on a Dell PC (Dell, Round Rock, TX) running PsychToolbox (Brainard, 1997; Pelli, 1997). Stimuli were presented on a 27" ASUS VG278HE LCD monitor, with 1920 × 1080 resolution and 120 Hz refresh rate. The monitor was calibrated using a Minolta photometer (Konica Minolta Sensing Americas, Ramsey, NJ) and linearised in software, to give a maximum luminance of 222 cd/m². Stimuli were viewed from 3 m, with the experimenter recording participants' verbal responses via keypad.

Stimuli and procedures

The experiment involved five video-game characters adapted from a previous study (Greenwood et al., 2012). The VacMan target was presented at the centre of the screen as a circle with a horizontal gap on one side, similar to a 'filled-in' Landolt-C (Figure 1B). The gap portrayed its 'mouth' and had a stroke width equivalent to one-fifth the stimulus diameter, as with Sloan letters (Sloan, 1959). During trials, the VacMan target was presented dark against the mid-gray background at a Weber contrast of 75%. Four achromatic ghost characters were also presented, with a semi-circular head and square 'legs' with gaps that were also one-fifth the stimulus diameter. When achromatic, the ghosts served as flanker stimuli in the flanked conditions (Figure 1C-D). In the unflanked condition, distinctly coloured animations of the ghosts (green above, red to the right, orange below, and blue to the left) moved along the screen boundaries (Figure 1A). Coloured ghosts were also shown on reference cards adhered to the edges of the monitor to serve as response options for the identification of the VacMan target orientation.

All participants began with the unflanked condition, where on each trial they reported which of the four ghosts VacMan was facing in order to help VacMan catch the ghost that he wanted to eat (a single-interval stimulus with a 4 Alternative Forced Choice response). Each ghost had a distinct colour and moved slowly along the monitor edges with a centre-to-centre separation approximately 5.5° from the target horizontally and 3.0° vertically (Figure 1A), making the chance of any crowding with the target unlikely. Responses based on ghost colour were used to avoid left–right response confusions (Corballis, 1974), though children could indicate the spatial location of the ghost if they wished. The use of clearly individuated characters also helped children with task comprehension, particularly in the flanked conditions – referring to the VacMan character helped make clear which element

they were required to respond to, minimising the possibility of errors arising from source confusion (Harrison & Bex, 2017), particularly if children were confused about which element to report. Prior to testing, subjects were familiarised with the characters via printed cards illustrating VacMan, the ghosts and an example of a trial, through which normal colour-naming abilities were also checked. Feedback was given after each trial via brief animations whereby a colour-rendered VacMan either smiled or frowned. After every third correct response, a longer animation occurred where VacMan ran towards and "ate" the correct ghost. This feedback animation acted both as a reward and as a reminder of the task.

In the two flanked conditions, Vac-Man and the four ghost flankers were achromatic and presented without eyes to ensure that the target and its flankers were sufficiently similar to induce crowding. On each trial of these flanked conditions, the achromatic ghost-flankers were presented at a random orientation (with one at each cardinal orientation). Children were instructed that the ghosts became colourless to 'hide' from VacMan, but it was emphasised that they maintained the same colour/position mapping as before (e.g., as in the unflanked condition, the green ghost was above VacMan, the red to the right, etc.), which was reinforced through the presence of the reference cards. As before, participants reported the ghost that VacMan was facing by referring either to its colour (via the four reference cards at the monitor boundaries) or location. All children completed one flanked condition where target and flanker elements had the same contrast polarity ('flanked same polarity'), with all elements presented dark against the mid-gray background at 75% Weber contrast (Figure 1C). A subset of children were also tested with a second flanked condition where a dark target (as in the other conditions) was flanked by light ('white') ghost-flankers, again at 75% Weber contrast (Figure 1D). 40 of the children and 10 adults completed this third condition.

In all three conditions, the size of the ghosts and VacMan (including the visibility of its "mouth") was varied using QUEST (Watson & Pelli, 1983), set to converge on a gap size (for VacMan's mouth gap) that gave 62.5% correct performance. Each block of trials consisted of a single QUEST staircase run on one of the conditions for 35 trials (not including practice trials). The target size was always scaled such that the diameter of VacMan was 5 times the mouth size. When flanked, the size of all elements was matched, with a centre-to-centre distance between the target and flankers set to 1.1 times the stimulus diameter. This separation is recommended by prior research as optimal for the robust

measurement of crowding (Song et al., 2014), and has been used to measure crowding in both adults and children previously (Tailor et al., 2021; Kalpadakis-Smith et al., 2022).

The QUEST procedure was modified in four ways to increase its suitability for testing children. First, five practise trials were presented to begin each block of trials, where the gap size was 5.3′, well above any thresholds observed during pilot testing. These trials were not included in final analyses. If performance on these trials was poor, the task was explained again and the test re-started. Second, easier "catch" trials were presented periodically (on trials 5, 15, 25 and 35) where stimulus size was scaled to triple the current QUEST estimate of threshold. These catch trials were included both to minimise the frustration and loss of motivation that can occur when placing continuous trials near threshold, and to measure inattentiveness during the block, as recommended by prior work (Manning et al., 2018). Third, to further reduce the effect of inattentiveness on threshold estimates, thresholds were determined by fitting psychometric functions to the data (also recommended by Manning et al., 2018). Catch trials were included in the final data used in this fitting procedure. The fourth modification was related to this later fitting – because QUEST ordinarily places stimulus sizes close to threshold, the resulting range of values can be insufficient for curve fitting. Stimulus size on each trial was thus jittered using Gaussian noise added to the current QUEST threshold estimate (with a standard deviation of half the threshold estimate). This jitter increased the range of stimulus sizes, in turn improving the subsequent fitting of psychometric functions to the data.

We aimed for each condition to be completed two times by children and three for adults, though testing was stopped after at least one block in each condition if cooperation waned or the participant did not wish to continue. As above, all children completed the unflanked and flanked-same polarity conditions, with a subset also completing the third flanked-opposite polarity condition. In each block, participants would earn points after correct responses to each of the four catch trials. When children achieved high scores in each condition, they received stickers that could be exchanged for a toy at the end of the experiment. These prizes were used to maintain children's motivation throughout the study. Rest breaks were permitted between the conditions. The full duration of the experiment was 30-60 minutes per person, followed by a debrief.

Results

Screening and analyses

The total number of trials completed per condition ranged from 35-105, corresponding to 1-3 blocks of trials. When split into the 5 age groups, this gave a mean number of trials of 45.5 per condition for 3-4 year olds, 59.8 for 5-6 year olds, 51.1 for 7-8 year olds, 56.6 for 9-10 year olds, 51.5 for 11-12 year olds and 65.8 for the adult group.

Performance on catch trials with large stimulus sizes was uniformly high, as shown by age in Figure 2A. Each age group had some individuals who failed on at least one of the catch trials, which was slightly higher in the youngest 3-4 year old group. Mean percent-correct performance for both the unflanked and flanked same-polarity conditions was nonetheless above 98% in all groups and conditions save for the flanked condition in the 3-4 year olds, which dropped slightly to average 93.8% correct. Given that catch trials were correctly reported on the majority of trials in all cases, all of these children were included in the final analyses. Supplementary analyses were also run after removing children who scored below 100% correct on catch trials, which did not differ from the main analyses (Figure S1).

To determine the threshold gap size required for performance in each condition, repeated blocks of trials were first combined for each participant and stimulus condition, with the corresponding proportion correct scores collated for each stimulus size (including catch trials). Psychometric functions were then fit to the data using a cumulative Gaussian function with two free parameters (midpoint and slope). Given the variable number of trials at each gap size (driven by the variability in the QUEST algorithm, as above), these fits were determined by weighting the least-squared error by the number of trials at each point and seeking to minimise this value.

Example proportion correct values and best-fitting psychometric functions for one observer (aged 6) are shown in Figure 2B. The number of trials at each gap size is depicted via the size of each point, with the weighted psychometric function fit emphasising those with the most trials. Performance in the unflanked condition (dark blue) begins at chance (25% correct) for the smallest gaps and rises rapidly to ceiling as stimulus size increases. A slower progression from chance to ceiling is evident in the flanked-same condition (light blue) where the black target was surrounded by black flankers. For each condition, gap-size

thresholds were taken from these psychometric functions at the size where performance reached 62.5% correct (midway between chance and ceiling).

Figure 2. The development of acuity and crowding. **A.** Percent-correct performance on catch trials (large stimulus sizes) as a function of age. Individuals are shown as small points and the mean within each age range (shown via grey/white bands) as large points, separately for the unflanked (dark blue) and flanked-same (light blue) conditions. **B.** Example percent-correct data and best-fitting psychometric functions for one observer (age 6), plotted as a function of stimulus size separated for the unflanked (dark blue) and flanked-same (light blue) conditions. Dot size indicates the number of trials at each point. **C.** Gap-size thresholds in minutes of arc, plotted against age. Individuals are shown as small points and the mean within each age range (shown via grey/white bands) as large points, separately for the unflanked (dark blue) and flanked-same (light blue) conditions. Adult data has been collapsed to fall near 18 years. Lines show the best fitting logistic function to each dataset.

The developmental trajectory of acuity and crowding

Figure 2C shows the final gap-size thresholds in minutes of arc, plotted as a function of age. These values represent the smallest gap that can be identified in the VacMan target (at 62.5% correct) when unflanked (dark blue) or surrounded by flankers with the same polarity (light blue). A two-way mixed effects ANOVA was run to examine the overall pattern, with a between-subjects factor for age and a within-subjects factor for flanker condition. This analysis revealed a significant main effect of age ($F_{5,113} = 20.85$, $p < .0001$), reflecting the drop in thresholds with age, a significant main effect of flanker condition ($F_{1,113} = 193.50$, $p < .0001$), reflecting the rise in thresholds when flankers were added, and a significant interaction between the two ($F_{5,113} = 9.71$, $p < .0001$).

To examine the effect of age in each condition, a series of planned contrasts were performed. Unflanked thresholds, a measure of visual acuity, were highest in the youngest children, but rapidly dropped to adult-like levels. This pattern is supported by independent samples t-tests between children in each age group and adults, which differed significantly at 3-4 years ($t_{38} = 5.148$, $p < 0.0001$), but not at 5-6 years ($t_{50} = 1.556$, $p = 0.126$) or beyond at 7-8 years ($t_{52} = 1.643$, $p = 0.106$), 9-10 years ($t_{42} = -0.755$, $p = 0.455$) or 11-12 years ($t_{47} = 1.307$, $p = 0.198$). Acuity is thus elevated in young children aged 3-4 years and converges to adult levels at the age of 5-6 years.

Flanked thresholds were higher than unflanked thresholds for all age groups, following the main effect of crowding condition. To examine the development of crowding, flanked thresholds in each age group for children were compared with those of adults. Flanked thresholds were considerably elevated in 3-4 year olds relative to adults ($t_{38} = 8.610$, $p < 0.0001$), with continued elevation at 5-6 years ($t_{50} = 3.592$, $p = 0.001$). Thresholds dropped to become indistinguishable from adults at 7-8 years ($t_{52} = 0.452$, $p = 0.653$) and remained flat thereafter with non-significant differences at 9-10 ($t_{42} = 1.007$, $p = 0.320$) and 11-12 years ($t_{47} = 0.812$, $p = 0.421$). Flanked performance thus indicates that crowding levels are initially strongly elevated in young children between the ages of 3-6, becoming indistinguishable from adult levels at the age of 7-8 years, later than the point at which acuity becomes adult-like.

As outlined in the introduction and Table 1, prior studies of developmental crowding show a striking variability in the age at which crowding has been argued to become adult-like, ranging from 5 to 12+ years. Our estimate of 7-8 years for the point of maturity falls

between these extremes. We next sought to investigate the cause of these discrepancies, both to build consensus around the developmental trajectory and to trace these differences back to their most likely methodological explanation (in turn considering the best-practice approach to measure crowding in children). With this aim, we first examined whether these discrepancies can be explained by different approaches to the measurement of crowding (specifically by varying flanker target-separation directly, or by scaling element size and separation), and then conducted a meta-analysis to gain a broader view of the developmental trajectory of crowding.

A comparison of separation vs. scaling approaches to measure crowding

As outlined in the introduction, crowding has been measured using a variety of stimuli, procedures, and measurement approaches. One apparent division in Table 1 is between studies where elements were scaled in both size and inter-element separation (as in the current study), which show earlier ages of maturation than those that varied separation directly using fixed-size elements (Semenov et al., 2000; Bondarko & Semenov, 2005; Jeon et al., 2010). Because the VacMan stimuli of the current study have been used to measure crowding previously using both the scaling approach (in the current study and Kalpadakis-Smith et al., 2022) and with direct manipulations of inter-element separation (Greenwood et al., 2012), we can directly test whether thresholds differ under these two approaches. To do so, thresholds in all three studies were analysed using the same procedures (see Supporting Information) and converted to give estimates of the centre-to-centre separation between elements at threshold.

Thresholds for the centre-to-centre separation between elements in the flanked-same condition in these three studies are plotted in Figure 3. Values from the current study again show a clear developmental trajectory, with thresholds from Greenwood et al. (2012) and Kalpadakis-Smith et al. (2022) largely interspersed amongst these values to follow this trajectory, aside from a small number of outliers. The lack of overall difference between these three studies was confirmed by taking values across a common age range for the three studies (4-8 years): Independent samples t-tests reveal that thresholds did not differ significantly between Greenwood et al. (2012) and the current study ($t_{54} = 1.401$, $p = 0.167$), between Kalpadakis-Smith et al. (2022) and the current study ($t_{55} = 0.104$, $p = 0.918$), nor between the two prior studies ($t_{33} = 0.973$, $p = 0.338$). We conclude that thresholds obtained using a scaled approach (as in the current study and Kalpadakis-Smith et al, 2022) are

comparable to those that vary inter-element separation directly (Greenwood et al., 2012). If anything, the direct manipulation of inter-element separation would appear to produce more variable data, with two children from Greenwood et al (2012) in particular showing very large critical spacing estimates for their age. We suspect that this approach could lead to problems if acuity thresholds were over-estimated, which would make it difficult for children to reach ceiling performance in the flanked conditions (since stimuli may be too small to reach a high level of recognition, even at the largest separations). As the scaling approach measures flanked and unflanked performance independently, there are no such issues with the cross-dependency of thresholds.

Figure 3. A comparison of different approaches to measure the critical separation of crowding, expressed as the centre-to-centre separation between elements at threshold, in minutes of arc. Thresholds from the flanked-same condition of the current study are shown in light blue, along with the best-fitting logistic function for these values. Thresholds are shown in equivalent units of centre-to-centre separation from Greenwood et al. (2012) in red (which measured inter-element separation directly) and Kalpadakis-Smith et al. (2022) in purple (which used the same scaling approach as the current study). Each point represents an individual child.

A meta-analysis of the developmental trajectory of crowding

We next sought a more direct comparison of our data against that from prior examinations of the developmental trajectory of crowding, both to obtain an overall consensus trajectory and to provide a quantitative basis for assessing the methodological characteristics that can bias the estimates of individual studies away from the central tendency. A major complication that arises in comparing prior estimates is the wide range of metrics used to measure crowding. However, a great advantage of the scaling approach in the present study (measuring flanked and unflanked thresholds by scaling element size and inter-

element separation proportionally) is that it allows for conversion into the same units as these previously-used metrics. We thus undertook a series of comparisons between prior datasets and our threshold values after conversion to the same metric, building towards a meta-analysis of the developmental trajectory of crowding.

Methods for data extraction and conversion for each study are described in the Supporting Information, along with individual comparisons between our values and those from 11 prior studies in their respective units of measurement (Figure S2). This includes both the studies listed in Table 1 as well as additional ones that lacked direct comparison to adult performance levels. Taking these analyses a step further, we then normalised each of these 11 datasets to span the same range, using our dataset range as a reference. The resulting estimates of normalised crowding across development are plotted in Figure 4.

Normalised crowding levels show some variation across studies, but an overall developmental trajectory is nonetheless clearly apparent. To examine this trajectory statistically, values were pooled in similar age ranges to those of the current dataset. Crowding levels were highest in the 3-4 year age range, which differed significantly from adult levels (t_{16}= 5.148, $p < 0.0001$). These levels remained significantly elevated at 5-6 years ($t_{15} = 3.131$, $p= 0.007$), but dropped to levels that did not differ significantly from adults at 7-8 years ($t_{16} = 1.224$, $p= 0.239$), and remained equivalent at 9-10 years ($t_7 = 0.635$, $p= 0.545$), 11-12 years ($t_7 = -0.451$, $p= 0.666$) and 13-17 years ($t_6 = -0.882$, $p= 0.412$). This overall pattern thus follows the same trajectory as that of the current dataset, despite variations in stimulus type (e.g. letters vs. symbols), inter-element separation (from intermediate to narrow spacings, and studies that manipulate this property directly), and measurement approach (including the use of scaling vs. separation manipulations, the number of response options, use of single targets vs. letter-chart configurations, and so on).

Figure 4. A meta-analysis of studies measuring the developmental trajectory of crowding. Crowding is plotted in normalised units for direct comparison (see Supporting Information), as a function of age on the x-axis. Small symbols show estimates from individual studies (see legend), while the large black points show the mean within each age range (indicated via shaded regions). The black line is the best-fitting logistic function fit to all studies. Some adult thresholds have been displaced on the x-axis for visibility.

Some divergence from this developmental trend is however apparent in individual studies, which can be traced to methodological elements. Although their estimated age of maturity agrees with ours, the estimates of Doron et al. (2015) run higher than others in the youngest children. It is likely that this elevation reflects the use of high numbers of elements shown in a letter chart arrangement, causing uncertainty regarding the target response in the youngest children. Interestingly, these values then drop to very low levels in the oldest children (producing the slight dip in mean normalised crowding levels in the oldest groups, due to the small number of studies with data in this range). This too may reflect the letter-chart arrangement – the wide spacing would mean that each element is only flanked closely by up to 2 elements (depending on the letter position in each line), which would underestimate the low crowding levels at older ages in particular. Divergences are also apparent in the estimates of Jeon et al. (2010), who directly varied the edge-to-edge separation between target and flanker elements of a fixed size. These estimates are high and unvarying in children, but drop considerably in adults, where estimates are much lower

than those of the current study and others reporting adult values. It may be that the adults tested by Jeon et al. (2010) were more experienced with psychological testing than other samples, or that differences were present in testing procedures between children and adults that could underestimate crowding in the latter. We also show in the comparison to individual studies (Figure S2) that the use of relative measures (which combine estimates of acuity and crowding to express crowding as a multiple of acuity levels) lead to noisier estimates compared to studies where acuity and crowding are measured independently (as with the scaling approach of the current study). Nonetheless, Figure 4 shows that the age of maturity estimate from the combination of these prior studies is consistent with that found in the present study, despite the apparent divergence in estimated ages reported by individual analyses.

The release from crowding

Given suggestions that foveal crowding may involve distinct processes in children and adults, we next sought to examine a key feature of crowding in peripheral vision – the selectivity of foveal crowding for target-flanker similarity. Differences in the contrast polarity of target and flanker elements have reliably been found to reduce the strength of crowding in peripheral vision (Kooi et al., 1994; Chakravarthi & Cavanagh, 2007; Chung & Mansfield, 2009). Here we compare the foveal vision of children and adults to test whether this same selectivity is present, using a subset of children and adults who completed an additional condition where the black target was flanked by white flankers ('flanked different').

Gap-size thresholds from 40 children and 10 adults are presented in Figure 5. Thresholds in the unflanked condition follow the same developmental trajectory as shown in Figure 2, with elevations in the youngest children that quickly recede to adult-like levels. Performance in the flanked-same condition is also similar to that of Figure 2, with elevations evident at all ages, but largest in the youngest children. Thresholds in the flanked-different condition were reduced relative to the flanked-same condition, though did not reach the levels of unflanked performance. Given the reduced sample size, statistical tests were run by taking a median split of the children by age, giving one group of 3-6 year olds and another of 7-12 year olds. Thresholds were significantly reduced in the flanked-different condition, relative to flanked-same performance, for children in both the 3-6 year old ($t_{19} = 2.648$, $p = 0.016$) and 7-12 year old groups ($t_{19} = 2.987$, $p = 0.008$), as well as

in adults ($t_9 = 3.263$, $p = 0.010$). If we consider this reduction as a proportion of the elevation in crowding (i.e. the percentage reduction of the difference between unflanked and flanked-same conditions), then performance was improved by approximately 50% in both children and adults. Both children and adults thus show reduced crowding when flankers are reversed in their contrast polarity, demonstrating a common selectivity between the developing fovea, adult fovea, and adult periphery.

Figure 5. The release from crowding with age. Gap-size thresholds are in minutes of arc, plotted against age. Individuals are shown as small points and the mean within each age range (shown via grey/white bands) as large points, separately for the unflanked (dark blue), flanked-same (light blue), and flanked-different (green) conditions. Adult data has been collapsed to fall near 18 years. Lines show the best fitting logistic function to each condition.

Discussion

Our results confirm that foveal vision is disrupted by clutter in young children. Elevations in the spatial extent of crowding were most prominent in the youngest children – for 3-4 year old children, thresholds for the recognition of our VacMan stimuli were almost doubled in the presence of surrounding clutter. These elevations recede with age, but do not reach adult-like levels until the age of 7-8 years when clutter increases thresholds by a factor of about 1.4. This development is slower than that of acuity, which reaches adult-like levels by around 5-6 years, though not as slow as some of the estimates of developmental crowding reported in prior studies. Despite the discrepancy in these previous reports, our meta-analysis reveals good overlap in the combined developmental trajectory of 12 studies (including our own) examining developmental crowding. This direct comparison was achieved after converting our thresholds to the metrics used by these studies, and gave an

age of maturity that again fell at 7-8 years, matching that of our dataset. Finally, we show that developmental elevations in foveal crowding also share a key property of peripheral crowding, with changes in target-flanker similarity (differences in contrast polarity) reducing crowding by 50% in all age groups. We conclude that developmental crowding derives from the same process as the disruptive effect evident in the adult visual system.

Our finding that foveal crowding matures at 7-8 years is similar to estimates from several prior studies (Norgett & Siderov, 2014; Doron et al., 2015), with our meta-analysis of 12 studies providing further evidence that crowding becomes adult-like at 7-8 years. Some discrepancies between individual estimates in this meta-analysis were nonetheless apparent, most notably in the study reporting the most prolonged developmental trajectory (Jeon et al., 2010). The comparison of these measures across studies suggests that this discrepancy derives from an apparent underestimation of adult crowding levels, which could reflect differences in the procedures used to test adults vs. children, the exclusive use of psychophysically-trained adults as a comparison, or the use of relative measures to calculate the magnitude of crowding (whereby flanked and unflanked performance are combined to yield multiple or difference values). This latter issue was further evident in the conversion of our thresholds to the relative measures used in other studies (e.g. Atkinson et al., 1987), which led to more variable estimates than when acuity and crowding were measured independently (Figure S2). Divergences were also apparent in studies using widely spaced letter-chart arrangements, which tended to underestimate crowding levels in older children in particular (Doron et al., 2015), consistent with prior observations that wider separations underestimate foveal crowding (Song et al., 2014; Facchin et al., 2019; Waugh & Fronius, 2024). Finally, though our measurements derive from stimuli with a target-flanker separation that was scaled relative to element size, we show with comparison to prior studies using the same stimuli (Greenwood et al., 2012; Kalpadakis-Smith et al., 2022) that this scaling approach yields estimates of crowding that are indistinguishable from those that directly manipulate target-flanker separation (though separation measures do show a tendency for increased variability, perhaps due to the dependence on fixed element sizes set relative to acuity levels). From this we recommend that developmental crowding is best measured against naïve adult observers, using tightly spaced stimuli, and with independent estimates of unflanked and flanked performance.

The studies compared in Table 1 and Figure 4 used a variety of stimuli and measurement procedures. Where these prior studies used multiple conditions to assess crowding, we

selected those with the most standard configuration (typically one target and four surrounding flankers). When these configurations are directly compared however, arrays with larger numbers of elements have been found to show a more pronounced elevation in children aged 7-9 than arrays with fewer elements (Norgett & Siderov, 2014). However, rather than reflecting an immaturity in crowding itself, the increased complexity of these stimulus arrangements may increase uncertainty regarding the location of target stimuli (Goldberg et al., 2001; Dye & Bavelier, 2010). This uncertainty may conflate the development of crowding with other processes such as spatial attention, which certainly interact with crowding (Mareschal et al., 2010), but are nonetheless clearly dissociable processes (Dakin et al., 2009). Other studies have reported slower developmental trajectories with children required to judge whether square-wave grating elements were horizontal or vertical (Bondarko & Semenov, 2005). Task comprehension with these more abstract stimuli may be an issue with children, and we suggest that the use of more gamified tasks is a useful means to clarify task requirements and ensure that they respond to the correct element.

Although it is clear that children make errors when judging the appearance of a fixated target in cluttered displays, it is not a given that these errors arise for the same reason as those found in adult peripheral vision. For instance, elevations in crowding could arise from changes in properties such as fixational stability (Aring et al., 2007) smearing elements across the retina, a factor that contributes to adult foveal crowding levels (Clark et al., 2024) and which becomes magnified in cases of infantile nystagmus (Chung & Bedell, 1995; Pascal & Abadi, 1995; Tailor et al., 2021). Contrary to this possibility, here we observe that crowding is reduced in children when elements are dissimilar in their contrast polarity. This effect of target-flanker similarity matches prior observations of polarity-based reductions in the adult fovea (Hess, Dakin, & Kapoor, 2000; Hess, Dakin, Kapoor, et al., 2000; Sayim et al., 2008) and adult periphery (Kooi et al., 1994; Chakravarthi & Cavanagh, 2007; Chung & Mansfield, 2009). This is also consistent with prior observations that children make the same kinds of errors in crowded arrays as adults in peripheral vision (Kalpadakis-Smith et al., 2022), which can be accounted for using population pooling models of crowding (Harrison & Bex, 2015; Greenwood & Parsons, 2020). In peripheral vision, this pooling process generates the 'gist' of the visual scene at the expense of fine detail (Freeman & Simoncelli, 2011; Rosenholtz et al., 2019). We suggest it is the same pooling process that is being refined over this period in the fovea as the visual system seeks

to balance the neural resources allocated to the recognition of fine detail vs. the extraction of summary statistics over a wider region to gain a simpler 'gist' of the scene.

Alternative accounts of crowding posit that these effects derive from grouping processes (Herzog & Manassi, 2015), including variations in contrast polarity (Sayim et al., 2008). It is unclear what aspect of the visual system would be maturing by these accounts however, and given that these mechanisms are usually depicted as heavily top down (Herzog & Manassi, 2015), one may expect a later maturation than that observed in the current study. Accordingly, processes that are often linked with grouping, such as the perception of illusory contours (Hadad et al., 2010) and the recognition of ambiguous two-tone images (Milne et al., 2024), do not mature until around 10-12 years, suggesting that the grouping processes that modulate crowding should show a similarly late maturation. Attentional processes also appear to develop late, with abilities such as multiple object tracking (Dye & Bavelier, 2010), attentional cueing (Goldberg et al., 2001), and the visual search for feature conjunctions (Trick & Enns, 1998) improving well into adolescence. This late maturation similarly seems inconsistent with an attentional basis for crowding, as some have proposed (He et al., 1996). These higher-level processes can nonetheless clearly interact with the magnitude of crowding, as above, and their role on developmental vision is important to ascertain.

Our results are consistent with a developmental cascade of maturation within the visual hierarchy (Hensch, 2004; Chomiak & Hu, 2017), whereby visual abilities that rely on higher levels of the cortical hierarchy tend to develop later (Lewis & Maurer, 2005; Braddick & Atkinson, 2011). Similar to prior estimates (Birch et al., 1983; Ellemberg et al., 1999; Leat et al., 2009), we find that acuity converges on adult-like levels at around the age of 5-6 years, earlier than the development of crowding. The developmental trajectory of acuity may then reflect the earlier maturation of retinal ganglion cells and primary visual cortex, given the link with acuity and these areas (Curcio et al., 1990; Duncan & Boynton, 2003). In contrast, the later maturation of crowding is consistent with observations that crowding relies on higher cortical levels than acuity, including candidate areas from V2-V4 (Motter & Simoni, 2007; Anderson et al., 2012; He et al., 2019; Henry & Kohn, 2022). The exact properties that might be developing in these areas is less clear. Though crowding is often linked with receptive field size (Motter & Simoni, 2007; He et al., 2019), both pRF sizes and surround inhibition components in areas V1-V4 have been found to be adult-like by 6 years of age (Dekker et al., 2019; Himmelberg et al., 2023). There are nonetheless some indications that

cortical magnification around the fovea may develop later in childhood within areas V2 and V3 (Himmelberg et al., 2023). The later development of these areas may then create the above interplay between fine detail and 'gist' processing during development.

The slow maturation of crowding may make this process more susceptible to disruptions during development. Similar proposals have been raised for motion perception, which develops late in childhood, with associated deficits in motion perception apparent in a range of developmental disorders (Braddick et al., 2003; Braddick & Atkinson, 2011). Crowding is similarly elevated in a range of developmental disorders of vision including amblyopia (Levi & Klein, 1985; Greenwood et al., 2012), nystagmus (Chung & Bedell, 1995; Pascal & Abadi, 1995; Tailor et al., 2021), and cerebral visual impairment (Huurneman, Boonstra, Cox, et al., 2012), which all have their onset in the age ranges where crowding is still maturing. Reports of elevated crowding in individuals with dyslexia (Atkinson, 1991; Martelli et al., 2009) and dyscalculia (Castaldi et al., 2020) could similarly be the result of this late maturation, given their typical onset within this age range. Disruptions to the spatial and/or featural selectivity of the visual system would seem to create an increased drive for 'gist' processing in spatial vision. This in turn would limit processes that rely on the fine detail of the fovea, such as reading (Kwon et al., 2007). Indeed, increased inter-letter spacing has been found to reduce reading errors in both dyslexic (Zorzi et al., 2012; Marinus et al., 2016) and typically developing children (Hakvoort et al., 2017).

Altogether, our results suggest that the optimisation of spatial vision takes time – while acuity becomes adult-like around the age of 5-6, crowding continues to develop until 7-8 years of age. These developmental elevations in crowding share core patterns of selectivity with adult peripheral vision, suggesting that foveal vision faces similar trade-offs between the representation of fine vision and 'gist' during development. The slow development of this balance renders these abilities particularly susceptible to disruption and presents challenges for everyday tasks such as reading. Measurement of these processes is also a continuing challenge, in both typical and clinical cohorts. Here we suggest that crowding be measured in a child-friendly format with equivalent comparisons to adult data, using independent measures of stimulus conditions (e.g. unflanked and flanked performance), and with tightly spaced stimuli. The careful measurement of these effects will be important for future work attempting to examine whether other properties of crowding, such as its modulation by higher-level processes like grouping, develop at a similar rate.

Acknowledgements

This work was funded by a UK MRC Career Development Award (MR/K024817/1) to JG. Our thanks to Alexandra Kalpadakis-Smith for help with code.

References

Anderson, E. J., Dakin, S. C., Schwarzkopf, D. S., Rees, G., & Greenwood, J. A. (2012). The neural correlates of crowding-induced changes in appearance. *Current Biology*, *22*(13), 1199-1206.

Anstis, S. M. (1974). A chart demonstrating variations in acuity with retinal position. *Vision Research*, *14*(7), 589-592.

Aring, E., Grönlund, M. A., Hellström, A., & Ygge, J. (2007). Visual fixation development in children. *Graefe's Archive for Clinical and Experimental Ophthalmology*, *245*, 1659-1665.

Atkinson, J. (1991). Review of human visual development: Crowding and dyslexia. In J. F. Stein (Ed.), *Vision and visual dysfunction* (Vol. 13, pp. 44-57). Macmillan Press.

Atkinson, J., Anker, S., Evans, C., Hall, R., & Pimm-Smith, E. (1988). Visual acuity testing of young children with the Cambridge Crowding Cards at 3 and 6 m. *Acta Ophthalmologica*, *66*, 505-508.

Atkinson, J., Anker, S., Evans, C., & McIntyre, A. (1987). The Cambridge Crowding Cards for preschool visual acuity testing. In M. Lenk--Schaefer (Ed.), *Transactions of the Sixth International Orthoptic Congress* (pp. 482-486). British Orthoptic Society.

Atkinson, J., & Braddick, O. J. (1983). Assessment of visual acuity in infancy and early childhood. *Acta Ophthalmologica*, *157 Suppl*, 18-26.

Bex, P. J., & Dakin, S. C. (2005). Spatial interference among moving targets. *Vision Research*, *45*, 1385-1398.

Birch, E. E., Gwiazda, J., Bauer Jr, J. A., Naegele, J., & Held, R. (1983). Visual acuity and its meridional variations in children aged 7–60 months. *Vision Research*, *23*(10), 1019-1024.

Bondarko, V. M., & Semenov, L. A. (2005). Visual acuity and the crowding effect in 8-to 17-year-old schoolchildren. *Human Physiology*, *31*, 532-538.

Bouma, H. (1970). Interaction effects in parafoveal letter recognition. *Nature*, *226*, 177-178.

Braddick, O. J., & Atkinson, J. (2011). Development of human visual function. *Vision Research*, *51*(13), 1588-1609.

Braddick, O. J., Atkinson, J., & Wattam-Bell, J. (2003). Normal and anomalous development of visual motion processing: motion coherence and 'dorsal-stream vulnerability'. *Neuropsychologia*, *41*(13), 1769-1784.

Brainard, D. H. (1997). The Psychophysics Toolbox. *Spatial Vision, 10*(4), 433-436.

Castaldi, E., Turi, M., Gassama, S., Piazza, M., & Eger, E. (2020). Excessive visual crowding effects in developmental dyscalculia. *Journal of Vision, 20*(8), 7-7.

Chakravarthi, R., & Cavanagh, P. (2007). Temporal properties of the polarity advantage effect in crowding. *Journal of Vision, 7*(2), 1-13.

Chomiak, T., & Hu, B. (2017). Mechanisms of hierarchical cortical maturation. *Frontiers in Cellular Neuroscience, 11*, 272.

Chung, S. T. L., & Bedell, H. E. (1995). Effect of retinal image motion on visual acuity and contour interaction in congenital nystagmus. *Vision Research, 35*(21), 3071-3082.

Chung, S. T. L., & Mansfield, J. S. (2009). Contrast polarity differences reduce crowding but do not benefit reading performance in peripheral vision. *Vision Research, 49*(23), 2782-2789.

Clark, A. M., Huynh, A., & Poletti, M. (2024). Oculomotor contributions to foveal crowding. *Journal of Neuroscience, 44*(48), e0594242024.

Coates, D. R., Levi, D. M., Touch, P., & Sabesan, R. (2018). Foveal crowding resolved. *Scientific Reports, 8*(1), 1-12.

Corballis, M. C. (1974). The left-right problem in psychology. *Canadian Psychologist, 15*(1), 16-33.

Curcio, C. A., Sloan, K. R., Kalina, R. E., & Hendrickson, A. E. (1990). Human photoreceptor topography. *Journal of Comparative Neurology, 292*(4), 497-523.

Dakin, S. C., Bex, P. J., Cass, J. R., & Watt, R. J. (2009). Dissociable effects of attention and crowding on orientation averaging. *Journal of Vision, 9(11):28*, 1-16.

Dekker, T. M., Farahbakhsh, M., Atkinson, J., Braddick, O. J., & Jones, P. R. (2020). Development of the spatial contrast sensitivity function (CSF) during childhood: Analysis of previous findings and new psychophysical data. *Journal of Vision, 20*(13), 4-4.

Dekker, T. M., Schwarzkopf, D. S., de Haas, B., Nardini, M., & Sereno, M. I. (2019). Population receptive field tuning properties of visual cortex during childhood. *Developmental Cognitive Neuroscience, 37*, 100614.

Doron, R., Spierer, A., & Polat, U. (2015). How crowding, masking, and contour interactions are related: A developmental approach. *Journal of Vision, 15*(8), 5-5.

Duncan, R. O., & Boynton, G. M. (2003). Cortical magnification within human primary visual cortex correlates with acuity thresholds. *Neuron, 38*, 659-671.

Dye, M. W. G., & Bavelier, D. (2010). Differential development of visual attention skills in school-age children. *Vision Research, 50*(4), 452-459.

Ellemberg, D., Lewis, T. L., Hong Liu, C., & Maurer, D. (1999). Development of spatial and temporal vision during childhood. *Vision Research, 39*(14), 2325-2333.

Facchin, A., Maffioletti, S., Martelli, M., & Daini, R. (2019). Different trajectories in the development of visual acuity with different levels of crowding: The Milan Eye Chart (MEC). *Vision Research*, *156*, 10-16.

Freeman, J., & Simoncelli, E. P. (2011). Metamers of the ventral stream. *Nature Neuroscience*, *14*(9), 1195-1201.

Gheri, C., Morgan, M. J., & Solomon, J. A. (2007). The relationship between search efficiency and crowding [Perception]. *Perception*, *36*(12), 1779-1787.

Goldberg, M. C., Maurer, D., & Lewis, T. L. (2001). Developmental changes in attention: The effects of endogenous cueing and of distractors. *Developmental Science*, *4*(2), 209-219.

Greenwood, J. A., Bex, P. J., & Dakin, S. C. (2009). Positional averaging explains crowding with letter-like stimuli. *Proceedings of the National Academy of Sciences of the United States of America*, *106*(31), 13130-13135.

Greenwood, J. A., & Parsons, M. J. (2020). Dissociable effects of visual crowding on the perception of color and motion. *Proceedings of the National Academy of Sciences of the United States of America*, *117*(14), 8196-8202.

Greenwood, J. A., Tailor, V. K., Sloper, J. J., Simmers, A. J., Bex, P. J., & Dakin, S. C. (2012). Visual acuity, crowding and stereo-vision are linked in children with and without amblyopia. *Investigative Ophthalmology & Visual Science*, *53*(12), 7655-7665.

Hadad, B.-S., Maurer, D., & Lewis, T. L. (2010). The development of contour interpolation: Evidence from subjective contours. *Journal of Experimental Child Psychology*, *106*(2-3), 163-176.

Hakvoort, B., van den Boer, M., Leenaars, T., Bos, P., & Tijms, J. (2017). Improvements in reading accuracy as a result of increased interletter spacing are not specific to children with dyslexia. *Journal of Experimental Child Psychology*, *164*, 101-116.

Harrison, W. J., & Bex, P. J. (2015). A Unifying Model of Orientation Crowding in Peripheral Vision. *Current Biology*, *25*(24), 3213-3219.

Harrison, W. J., & Bex, P. J. (2017). Visual crowding is a combination of an increase of positional uncertainty, source confusion, and featural averaging [Article]. *Scientific Reports*, *7*, 45551.

He, D., Wang, Y., & Fang, F. (2019). The critical role of V2 population receptive fields in visual orientation crowding. *Current Biology*, *29*(13), 2229-2236. e2223.

He, S., Cavanagh, P., & Intriligator, J. (1996). Attentional resolution and the locus of visual awareness. *Nature*, *383*, 334-337.

Henry, C. A., & Kohn, A. (2022). Feature representation under crowding in macaque V1 and V4 neuronal populations. *Current Biology*, *32*(23), 5126-5137. e5123.

Hensch, T. K. (2004). Critical period regulation. *Annual Review of Neuroscience*, *27*(1), 549-579.

Herzog, M. H., & Manassi, M. (2015). Uncorking the bottleneck of crowding: a fresh look at object recognition. *Current Opinion in Behavioral Sciences*, *1*, 86-93.

Hess, R. F., Dakin, S. C., & Kapoor, N. (2000). The foveal 'crowding' effect: Physics or physiology? *Vision Research*, *40*, 365-370.

Hess, R. F., Dakin, S. C., Kapoor, N., & Tewfik, M. (2000). Contour interaction in fovea and periphery. *Journal of the Optical Society of America. A, Optics, Image Science, and Vision*, *17*(9), 1516-1524.

Himmelberg, M. M., Tünçok, E., Gomez, J., Grill-Spector, K., Carrasco, M., & Winawer, J. (2023). Comparing retinotopic maps of children and adults reveals a late-stage change in how V1 samples the visual field. *Nature Communications*, *14*(1), 1561.

Huurneman, B., Boonstra, F. N., Cillessen, A. H. N., van Rens, G., & Cox, R. F. A. (2012). Crowding in central vision in normally sighted and visually impared children aged 4 to 8 years: the influence of age and test design. *Strabismus*, *20*(2), 55-62.

Huurneman, B., Boonstra, F. N., Cox, R. F. A., Cillessen, A. H. N., & Van Rens, G. (2012). A systematic review on 'Foveal Crowding' in visually impaired children and perceptual learning as a method to reduce crowding. *BMC Ophthalmology*, *12*, 1-14.

Jeon, S. T., Hamid, J., Maurer, D., & Lewis, T. L. (2010). Developmental changes during childhood in single-letter acuity and its crowding by surrounding contours. *Journal of Experimental Child Psychology*, *107*(4), 423-437.

Jeon, S. T., Maurer, D., & Lewis, T. L. (2012). The effect of video game training on the vision of adults with bilateral deprivation amblyopia. *Seeing and Perceiving*, *25*(5), 493-520.

Kalpadakis-Smith, A. V., Goffaux, V., & Greenwood, J. A. (2018). Crowding for faces is determined by visual (not holistic) similarity: Evidence from judgements of eye position. *Scientific Reports*, *8*(12556), 1-14.

Kalpadakis-Smith, A. V., Tailor, V. K., Dahlmann-Noor, A. H., & Greenwood, J. A. (2022). Crowding changes appearance systematically in peripheral, amblyopic, and developing vision. *Journal of Vision, 22(6):3*, 1-32.

Kooi, F. L., Toet, A., Tripathy, S. P., & Levi, D. M. (1994). The effect of similarity and duration on spatial interaction in peripheral vision. *Spatial Vision*, *8*(2), 255-279.

Kwon, M., Legge, G. E., & Dubbels, B. R. (2007). Developmental changes in the visual span for reading. *Vision Research*, *47*(22), 2889-2900.

Leat, S. J., Yadav, N. K., & Irving, E. L. (2009). Development of visual acuity and contrast sensitivity in children. *Journal of Optometry*, *2*(1), 19-26.

Levi, D. M., Hariharan, S., & Klein, S. A. (2002). Suppressive and facilitatory spatial interactions in peripheral vision: Peripheral crowding is neither size invariant nor simple contrast masking. *Journal of Vision*, *2*, 167-177.

Levi, D. M., & Klein, S. A. (1985). Vernier acuity, crowding and amblyopia. *Vision Research*, *25*(7), 979-991.

Levi, D. M., Klein, S. A., & Hariharan, S. (2002). Suppressive and facilitatory spatial interactions in foveal vision: Foveal crowding is simple contrast masking. *Journal of Vision*, *2*(2), 140-166.

Lewis, T. L., & Maurer, D. (2005). Multiple sensitive periods in human visual development: Evidence from visually deprived children. *Developmental Psychobiology*, *46*(3), 163-183.

Louie, E. G., Bressler, D. W., & Whitney, D. (2007). Holistic crowding: Selective interference between configural representations of faces in crowded scenes. *Journal of Vision*, *7(2):24*, 1-11.

Manning, C., Jones, P. R., Dekker, T. M., & Pellicano, E. (2018). Psychophysics with children: Investigating the effects of attentional lapses on threshold estimates. *Attention, Perception, & Psychophysics*, 1-14.

Mareschal, I., Morgan, M. J., & Solomon, J. A. (2010). Attentional modulation of crowding. *Vision Research*, *50*(8), 805-809.

Marinus, E., Mostard, M., Segers, E., Schubert, T. M., Madelaine, A., & Wheldall, K. (2016). A special font for people with dyslexia: Does it work and, if so, why? *Dyslexia*, *22*(3), 233-244.

Martelli, M., Di Filippo, G., Spinelli, D., & Zoccolotti, P. (2009). Crowding, reading, and developmental dyslexia. *Journal of Vision*, *9*(4), 1-18.

Milne, G. A., Lisi, M., McLean, A., Zheng, R., Groen, I. I. A., & Dekker, T. M. (2024). Perceptual reorganization from prior knowledge emerges late in childhood. *iScience*, *27*(2).

Motter, B. C., & Simoni, D. A. (2007). The roles of cortical image separation and size in active visual search performance. *Journal of Vision*, *7*(2(6)), 1-15.

Norgett, Y., & Siderov, J. (2011). Crowding in children's visual acuity tests—Effect of test design and age. *Optometry and Vision Science*, *88*(8), 920-927.

Norgett, Y., & Siderov, J. (2014). Foveal crowding differs in children and adults. *Journal of Vision*, *14*(12), 23-23.

Pascal, E., & Abadi, R. V. (1995). Contour interaction in the presence of congenital nystagmus. *Vision Research*, *35*(12), 1785-1789.

Pelli, D. G. (1997). The VideoToolBox software for visual psychophysics: Transforming numbers into movies. *Spatial Vision*, *10*(4), 437-442.

Pelli, D. G., Palomares, M., & Majaj, N. J. (2004). Crowding is unlike ordinary masking: Distinguishing feature integration from detection. *Journal of Vision*, *4*(12), 1136-1169.

Põder, E. (2008). Crowding with detection and coarse discrimination of simple visual features. *Journal of Vision*, *8(4):24*, 1-6.

Rosenholtz, R. (2016). Capabilities and limitations of peripheral vision. *Annual Review of Vision Science*, *2*, 437-457.

Rosenholtz, R., Yu, D., & Keshvari, S. (2019). Challenges to pooling models of crowding: Implications for visual mechanisms. *Journal of Vision*, *19*(7), 1-25.

Sayim, B., Westheimer, G., & Herzog, M. H. (2008). Contrast polarity, chromaticity, and stereoscopic depth modulate contextual interactions in vernier acuity. *Journal of Vision*, *8(8):12*, 1-9.

Semenov, L. A., Chernova, N. D., & Bondarko, V. M. (2000). Measurement of visual acuity and crowding effect in 3–9-year-old children. *Human Physiology*, *26*, 16-20.

Sloan, L. L. (1959). New test charts for the measurement of visual acuity at far and near distances. *American Journal of Ophthalmology*, *48*(6), 807-813.

Song, S., Levi, D. M., & Pelli, D. G. (2014). A double dissociation of the acuity and crowding limits to letter identification, and the promise of improved visual screening. *Journal of Vision*, *14*(5), 1-37.

Strasburger, H., Rentschler, I., & Jüttner, M. (2011). Peripheral vision and pattern recognition: A review. *Journal of Vision*, *11*(5:13), 1-82.

Tailor, V. K., Theodorou, M., Dahlmann-Noor, A. H., Dekker, T. M., & Greenwood, J. A. (2021). Eye movements elevate crowding in idiopathic infantile nystagmus syndrome. *Journal of Vision*, *21*(13:9), 1-23.

Toet, A., & Levi, D. M. (1992). The two-dimensional shape of spatial interaction zones in the parafovea. *Vision Research*, *32*(7), 1349-1357.

Trick, L. M., & Enns, J. T. (1998). Lifespan changes in attention: The visual search task. *Cognitive Development*, *13*(3), 369-386.

Watson, A. B., & Pelli, D. G. (1983). QUEST: A Bayesian adaptive psychometric method. *Perception & Psychophysics*, *33*(2), 113-120.

Waugh, S. J., & Fronius, M. (2024). Landolt C-Tests With "Fixed" Arcmin Separations Detect Amblyopia But Underestimate Crowding in Moderate-to-Severe Amblyopic Children and Adults. *Investigative Ophthalmology & Visual Science*, *65*(10), 33-33.

Whitney, D., & Levi, D. M. (2011). Visual crowding: A fundamental limit on conscious perception and object recognition. *Trends in Cognitive Sciences*, *15*(4), 160-168.

Wilkinson, F., Wilson, H. R., & Ellemberg, D. (1997). Lateral interactions in peripherally viewed texture arrays. *Journal of the Optical Society of America. A, Optics, Image Science, and Vision*, *14*(9), 2057-2068.

Zorzi, M., Barbiero, C., Facoetti, A., Lonciari, I., Carrozzi, M., Montico, M., Bravar, L., George, F., Pech-Georgel, C., & Ziegler, J. C. (2012). Extra-large letter spacing improves reading in dyslexia. *Proceedings of the National Academy of Sciences of the United States of America*, *109*(28), 11455-11459.

Supporting information

The influence of attentional lapses

Given suggestions that the frequency of attentional lapses can hinder the detection of visual abilities (Dekker et al., 2020), we examined the influence of these lapses on our data. We can quantify attentional lapses through the catch trials, whereby stimuli were presented at sizes three times larger than the QUEST estimate of threshold for that trial. Here we removed participants achieving less than 100% correct on these trials, independently for each of the unflanked and flanked-same conditions. There were not a great number of children who failed to meet this criteria in either case, though lapses were slightly higher in the flanked-same trials (as shown in Figure 2A). Amongst the 3-4 year olds, none were removed from the unflanked condition, with 2 removed from the flanked-same to leave 8. One 5-6 year old was removed from both conditions, leaving 21. One 7-8 year old was removed from the unflanked and two from the flanked-same, leaving 23 and 22, respectively. Of the 9-10 year olds, 3 were removed from the unflanked and 1 from the flanked-same, leaving 11 and 13, respectively. No 11-12 year olds were removed from the unflanked condition, and just one from the flanked-same to give 19 and 18, respectively. Six adults were removed from the unflanked condition to leave 24, and two from the flanked-same condition to give 28.

Thresholds from the remaining participants are shown in Figure S1. Mean values did not vary appreciably from those presented in the main analysis in Figure 2C. As before, isolated performance was significantly elevated relative to adult thresholds in the youngest children at 3-4 years ($t_{32} = 4.853$, $p < 0.0001$), with no significant difference for any age groups beyond this (5-6 years: $t_{43} = 1.495$, $p = 0.142$; 7-8 years: $t_{45} = 1.575$, $p = 0.122$; 9-10 years: $t_{33} = -0.872$, $p = 0.390$; 11-12 years: $t_{41} = 1.239$, $p = 0.222$). The addition of flankers similarly produced clear elevations in threshold, which were significantly larger than adults at 3-4 years ($t_{34} = 7.783$, $p < 0.0001$) and at 5-6 years ($t_{47} = 3.141$, $p = 0.003$), but not at

7-8 years ($t_{48} = 0.500$, p= 0.619) or beyond (9-10 years: $t_{39} = 0.728$, p = 0.471; 11-12 years: $t_{44} = 0.548$, p= 0.586). Given that the removal of participants with catch-trial performance below 100% does not alter the pattern of results, we conclude that differences in attentional lapse rates with age are unlikely to have played a major role in the pattern of results.

Figure S1. Gap-size thresholds plotted by age after filtering to remove individuals with performance on catch trials that was below 100% correct. As in Figure 2C, thresholds are plotted in minutes of arc, with individuals shown as small points and the mean in each age range (shown via grey/white bands) as large points, separately for the unflanked (dark blue) and flanked-same (light blue) conditions. Adult data has been collapsed to fall near 18 years. Lines show the best fitting logistic function to each dataset.

Comparisons to prior studies using scaling vs. spacing approaches

Prior reports of the age at which crowding reaches adult levels, summarised in Table 1, show an apparent divergence between studies where elements were scaled (as in the current study) and those using fixed stimulus sizes where inter-element separation was varied directly using elements of a fixed size (typically set as a multiple of unflanked acuity, using values measured prior to the measurement of crowding). Because two of our prior studies have taken these varied approaches, we sought to compare the resulting estimates of crowding to consider the influence of these measurement approaches.

The direct manipulation of inter-element spacing was previously used by Greenwood et al. (2012) using the same VacMan stimuli to examine crowding in children aged 4-8, where typical development data served as a control for studying disorders. Using stimuli of a fixed size (2.5 times above acuity thresholds), Greenwood et al. (2012) varied the centre-to-centre separation between elements to find the separation where performance reached 62.5% correct. Trial-by-trial data was extracted from this study and re-fit with psychometric functions for comparison to the current study. To consider the reliability of our estimates, we can further compare our values with another recent study measuring crowding in typical children (Kalpadakis-Smith et al., 2022), where a scaling method similar to the current study was used with children aged 3-8 years. Because Kalpadakis-Smith et al. (2022) used a higher threshold of 80% correct to measure the full extent of crowding, their trial-by-trial data was extracted and re-fit to obtain thresholds at 62.5% correct. Values from these two studies can then be compared with those of the current study by calculating the centre-to-centre separation of the elements in the current study at threshold. We can do so by multiplying our gap-size thresholds by 5 (to give stimulus diameter) and again by 1.1 (the scaled centre-to-centre separation) to give thresholds for the centre-to-centre separation between elements in minutes of arc. The resulting individual values are plotted against those of the current study in Figure 3.

Meta-analysis of the developmental trajectory of crowding

To examine the correspondence between our estimates of the developmental trajectory of crowding and those of prior studies, we sought to compare these estimates directly. As well as the studies reported in Table 1 (which only includes studies with comparison to adult data), here we also included studies without comparison to adult values, allowing a broader view of the literature. Values from prior studies were extracted from the figures of each publication using the GRABIT tool for MATLAB. Mean values were taken in all but one case where individual values were plotted (as below). For comparison to our dataset, values were plotted either at the reported age values or at the middle of range values, where used. For each comparison, individual values from the present study were converted to the same units of measurement and re-fit with a logistic function. To compare values against those of the current study, we derived error terms for the logistic fit to our dataset (separately after conversion to each measurement unit) by taking the 95% range of values obtained by fitting the logistic function to 1000 bootstrapped samples of the data.

A common metric used in prior studies is to report crowding as multiples of acuity thresholds, measuring the elevation in crowding relative to baseline acuity levels. This approach was first taken by Atkinson et al. (1987) using the Cambridge Crowding Cards. Figure S2A shows that these values tend to be slightly lower than ours, consistent with the broader spacing of their elements (1.5× vs. 1.1× element size in the present study), which has been shown to underestimate the amount of crowding (Song et al., 2014). Thresholds nonetheless follow roughly the same developmental trend as the current study and fall within the 95% range of our fitted values for both sets of children. Similar measures were taken by Huurneman, Boonstra, Cillessen, et al. (2012) using acuity charts. Their most-closely spaced condition (1.0× element size) is also plotted in Figure S2A, which again follows largely the same developmental trend. The slight undershoot in their values at age 7 relative to ours is likely due to their use of 'linear' stimulus arrangements, where each element is flanked by a maximum of 2 elements (depending on the position of the letter in each line), which would give less crowding than our 4 flankers given that crowding increases with flanker number (Põder, 2008). These estimates of crowded elevation are not therefore widely divergent from those of the present study. Of note however is the individual variability in these estimates – whereas the individual values plotted separately for unflanked and flanked-same conditions clearly follow the developmental trajectory in Figure 2, their greater dispersal here makes the trajectory harder to discern. We suggest that it is better to take independent measures of performance and to compare them across conditions than to use estimates that combine multiple sources of error.

We can also compare our thresholds to prior work measuring crowding as the log difference between flanked and unflanked thresholds. Our data can be converted to this format by converting gap-size thresholds in minutes of arc to log units and then subtracting flanked thresholds from unflanked. Values from Norgett and Siderov (2014) obtained with a standard target-flanker arrangement (a letter target with 4 flankers, as in the current study) are plotted in Figure S2B. These values lie close to ours, with the youngest children falling within our 95% range, though both older children and adults sit slightly below ours, likely driven by the wider spacing of 1.5× element size underestimating crowding levels, as above. Similar estimates can be seen in an earlier study using the same approach (Norgett & Siderov, 2011), also plotted in Figure S2B. Also included in this subplot is data from Facchin et al. (2019), who measured logMAR acuity using letter-chart arrangements at a range of inter-element spacings. For comparison, log thresholds were taken from the widest (2.0× element size) and narrowest (1.125×) centre-to-centre spacing conditions, which were

then subtracted to give log difference values. These estimates again agree closely with our own.

Log difference values were also computed by Doron et al. (2015), measured with letter-chart arrays of Tumbling-E stimuli. Because individual values were plotted in this study, we recorded these and calculated mean crowding levels within each of the age ranges used in the current study, as plotted in Figure S2C. Although this approach gives an estimate for the age of maturity that is very similar to that of the present study, the estimates of crowding are higher than ours for the youngest children, likely due to the use of high numbers of elements causing uncertainty in the youngest children. An undershoot in crowding levels is also visible in the oldest children, likely driven by the letter-chart arrangement whereby each element is flanked by a maximum of 2 elements (depending on the position of the letter in each line), as with Huurneman et al (2012). Thus, although the absolute values diverge from ours, the developmental trajectory remains largely the same.

The study with the oldest age of maturity for crowding comes from Jeon et al. (2010), who took measures of the edge-to-edge separation between target and flanker elements, with elements presented at a size 1.2× acuity (measured initially) and varied in their separation. We can convert our values for comparison to theirs with some assumptions. The scaling of our elements meant that the edge-to-edge separation was always 0.1× the element size. We can however estimate what our values might have been had we varied spacing using fixed-size elements by taking the centre-to-centre separation between elements at threshold and subtracting the diameter of elements at the acuity threshold multiplied by 1.2, before dividing by the gap-size threshold to give units of stroke width. The resultant values are plotted against those of Jeon et al. (2010) in Figure S2D. Only 1 of 3 estimates of childhood crowding coincide with our 95% range, though the largest discrepancy occurs in the adult data, where the estimates from Jeon et al. (2010) are considerably lower than those of the adults measured with our approach. One possibility is that the adult sample collected by Jeon et al. (2010) were more experienced with psychological testing than ours. Differences in the nature of the testing procedures used in children and adults could also give rise to this divergence. In comparison, our sample included predominantly adults who were inexperienced with psychological testing, with the same testing conditions used for both adults and children, and independent measures of the two abilities. The relative nature of these estimates (with flanked thresholds divided by unflanked) also makes it difficult to assess whether the difference reflects changes in crowding, acuity, or both.

Figure S2. Comparison to prior datasets, plotted as a function of age on the x-axis. **A.** Crowding expressed in multiples of unflanked thresholds, with individuals shown as small blue dots and means for each age range (as in Figure 2) shown as large blue points. Comparison data is shown from Atkinson et al. (1987) in red and Huurneman, Boonstra, Cillessen, et al. (2012) in purple. Lines show the best-fitting logistic function with shaded region indicating the 95% range of bootstrapped values. **B.** Crowding expressed as the difference between flanked and unflanked thresholds in log units, plotted against values from Norgett and Siderov (2011), Norgett and Siderov (2014), and Facchin et al (2019). **C.** As in panel B, plotted against both individual (small points) and mean (large) values from Doron et al. (2015). **D.** Crowding expressed as the estimated edge-to-edge separation between elements, in multiples of stroke width, with comparison to data from Jeon et al. (2010) in red. **E.** Crowding as edge-to-edge separation in multiples of element size, plotted against data from Semenov et al. (2000). **F.** Crowding as edge-to-edge separation in minutes of arc, plotted against data from Bondarko and Semenov (2005).

Similar calculations allow comparison with the data of Semenov et al. (2000), who varied the edge-to-edge separation between target and flanker elements presented at a size equal to acuity thresholds. Values were reported as edge-to-edge separation in multiples of unflanked whole-element size, which we can again calculate for our thresholds by taking the centre-to-centre separation between elements at threshold and subtracting the diameter of elements at the acuity threshold, before dividing by element diameter to give multiples of unflanked size. Semenov et al. (2000) used a subjectively determined high level of performance to take threshold, which we attempted to match by taking thresholds at 90% correct (instead of 62.5% correct elsewhere). As shown in Figure S2E, the estimates of Semenov et al. (2000) are elevated relative to ours in both children and adults, and show a period of continued elevation. Given the derivation of thresholds at high performance levels however, it is possible that some of this prolongation is due to attentional lapses, which are a primary cause of disrupted performance at the upper end of the psychometric function in particular (Manning et al., 2018; Dekker et al., 2020), which would also make the subjective estimation of asymptotic performance difficult.

Finally, Bondarko and Semenov (2005) conducted similar procedures to Semenov et al., measuring the edge-to-edge separation between target and flanker elements, with elements presented at a size equal to acuity thresholds and varied in their separation. Here, values were reported as edge-to-edge separation in minutes of arc, which are more easily compared to ours (with the above assumptions regarding stimulus size), as plotted in Figure S2F. Here although their youngest 8-year-old participants show a slight elevation relative to ours, the bulk of the measurements align well with ours. The age-range in our data nonetheless reveals that far greater changes in the developmental trajectory are evident before the age of 8 years. As with the estimates of Semenov et al. (2000), the use of subjective fitting procedures to asymptotic data ranges may have led to the slight elevation in performance seen here with the 8 year old participants. Estimating a lower performance criterion may alter these estimates.

To combine these disparate metrics of crowding for meta-analysis, we used our dataset in each case as a reference. Each of the above 11 datasets were thus normalised by subtracting the minimum value from the mean values across age in our dataset (after conversion into the relevant metric), before dividing by the maximum mean value of our dataset. This set our dataset to span the range 0-1, with other datasets varying around this. The resultant values could then be combined in the same space. Mean values were taken within similar

age ranges to those of the current study, which gave 13 estimates for 3-4 year-olds, 12 for 5-6 year-olds, 13 for 7-8 year-olds, 4 for 9-10 year-olds, 4 for 11-12 year-olds, 3 for 13-17 year-olds and 5 adult comparisons. Data for this meta-analysis are plotted in Figure 5.