Structural Insights into Kainate Receptor Desensitization

Changping Zhou[1,2], Guadalupe Segura-Covarrubias[1,2], and Nami Tajima[✓1]

[1] Department of Physiology and Biophysics, Case Western Reserve University School of Medicine, Ohio, 44106, USA.

[2] Equal contribution

[✓] Correspondence to nxt193@case.edu (N.T.)

Highlights

1. We present cryo-EM structures of GluK2 kainate receptors with engineered cysteine crosslinks at the inter-dimer interface, which restrict subunit lateral rotation and attenuate receptor desensitization.

2. The structure of GluK2 double cysteine mutant in complex with the allosteric potentiator BPAM344 and glutamate represents a non-desensitized state, highlighting the critical conformational changes required for ion channel gating.

3. The glutamate-bound GluK2 mutant adopts multiple conformations, representing both shallow- and deep-desensitized states. Electrophysiological recordings indicate that the GluK2 kainate receptor mutant recovers from desensitization more rapidly, resembling AMPA receptors. Our structural and functional data suggest that shallow-desensitized KARs remain conductive, implying that the large lateral LBD rotation during KAR desensitization is essential for complete channel closure, distinguishing KARs from other iGluRs.

Keywords

Kainate-type ionotropic glutamate receptors; structure; desensitization mechanism.

Abstract

Kainate receptors (KARs), along with AMPA and NMDA receptors, belong to the ionotropic glutamate receptor (iGluR) family and play critical roles in mediating excitatory neurotransmission throughout the central nervous system. KARs also regulate neurotransmitter release and modulate neuronal excitability and plasticity. Receptor desensitization plays a critical role in modulating the strength of synaptic transmission and synaptic plasticity. While KARs share overall structural similarity with AMPA receptors, the desensitized state of KARs differs strikingly from that of other iGluRs. Despite extensive studies on KARs, a fundamental question remains unsolved: why do KARs require large conformational changes upon desensitization, unlike other iGluRs? To address this, we present cryo-electron microscopy structures of GluK2 with double cysteine mutations in non-desensitized, shallow-desensitized and deep-desensitized conformations. In the shallow-desensitized conformation, two cysteine crosslinks stabilize the receptors in a conformation that resembles the desensitized state of AMPA receptors. However, unlike the tightly closed pore observed in the deep-desensitized KAR and desensitized AMPAR conformations, the channel pore in the shallow-desensitized state remains incompletely closed. Patch-clamp recordings and fluctuation analysis suggest that this state remains ion-permeable, indicating that the lateral rotational movement of KAR ligand-binding domains (LBDs) is critical for complete channel closure and stabilization of the receptor in desensitization states. Together with the multiple conformations representing different degree of desensitization, our results define the unique mechanism and conformational dynamics of KAR desensitization.

Main

Kainate receptors (KARs) belongs to the ionotropic glutamate receptor (iGluR) superfamily together with α-amino-3-hydroxy-5-methyl-4-isoxazolepropionic acid (AMPA), *N*-methyl-d-aspartate (NMDA) and delta receptors. They are cation selective ligand-gated ion channels activated by an excitatory neurotransmitter glutamate and mediate fast excitatory synaptic transmission in mammalian brains[1,2]. In addition to their role in mediating the synaptic transmission at post-synapses, KARs also function pre- and extra-synaptically, and regulate excitatory and inhibitory neurotransmitter release and modulate synaptic excitability[3-7]. Abnormal expression and dysfunction of KARs are associated with various neurological diseases and disorders, including schizophrenia, mood disorders, epilepsy, and Huntington's disease[8-17]. Therefore, KARs are considered potential therapeutic drug targets due to their physiological and pathophysiological roles[18-21].

Kainate receptors (KARs) assemble as homo- or heterotetramers composed of five subunits: GluK1–5[22-25]. Extensive studies revealed that KARs adopt a Y-shaped architecture with four distinct layers: the amino-terminal domain (ATD), which is critical for receptor assembly; the ligand-binding domain (LBD), which contains ligand-binding sites and regulates channel gating; the transmembrane domain (TMD), which forms the ion channel within the plasma membrane; and a short disordered C-terminal domain (CTD), which interacts with intracellular proteins[1,26]. Recently determined structures of intact KARs in various liganded states have elucidated the structure-function correlations underlying the mechanisms of activation/desensitization[25,27-31], allosteric modulation[32], inhibition[33], and channel block[34]. A characteristic structural feature of KARs lies in their unique conformation during desensitization. Structures of intact GluK2, first reported in 2014 by the Mayer/Subramaniam lab, have indicated that KAR LBD dimer assemblies are completely dissociated upon desensitization, with a large horizontal-plane rigid-body rotation of the LBDs in the BD subunits, resulting in an approximately four-fold symmetric structures, while

other iGluRs typically retain two-fold symmetric conformations viewed from the top when they are

desensitized[26,28]. In addition, a key functional characteristic of KARs is their significantly slower

recovery from desensitization — over 50 fold slower compared to AMPARs[35,36]— which limits KAR

mediated rapid and high-frequency transmission in neurons[37]. Due to the rapid onset of

desensitization induced by the neurotransmitter glutamate, which closes the ion channel within a

few milliseconds, previous KAR structures in complex with glutamate have been stabilized in fully

or deep-desensitized states, with the ion channel fully closed[25,27-31,33,38-40]. Despite their unique

structural and functional features, why only KARs exhibit such a strikingly different conformation,

which is not conserved in other iGluRs, and how these drastic conformational changes control

receptor gating remains elusive.

To address this outstanding question, we present the cryo-electron microscopy (cryo-EM)

structure of the GluK2 KAR in multiple functional states, including non-desensitized, shallow-

desensitized, intermediate (between shallow and deep desensitized), and deep-desensitized

conformations, using cysteine cross-linking in the presence of glutamate and the positive

allosteric modulator BPAM344. The structural comparison revealed that the large in-plane rotation

of LBDs in both BD subunits and the resulting twisted motion of the LBD-TM3 linkers are essential

for tight ion channel closure and stabilization of receptors in the deep-desensitized state—a

feature not observed in AMPARs and NMDARs. Functionally, the crosslinked GluK2 mutant

exhibits large fractional steady-state currents, as revealed by patch-clamp recording. The

stationary noise analysis results indicate that the GluK2 mutant remains conductive when

desensitized, resembling the behavior of the conductive GluA2(R) AMPAR under similar

conditions[41]. These structural and functional results show that robust in-plane rotations of the

LBDs are essential for proper ion channel closure. Overall, our study provides new insights into

the unique desensitization mechanism of the KAR subfamily.

Results

Trapping GluK2 KARs in a non-desensitizing state

Previous studies have shown that LBD dimer reorientations switch the gating of KARs[27-29]. To gain insight into the detailed conformational transitions from active to desensitized states, we first generated a homology model of the intact open GluK2 receptor, comprising the ATD, LBD, and TMD layers based on the structure of the activated GluA2 AMPA receptor in the presence of glutamate, potentiator, and auxiliary subunit in its higher conductive open state (Protein Data Bank (PDB) code 5WEO)[42-44] using SWISS-MODEL[45-49] (Fig. S1A, B). In the model, the G/E helices of subunit A and the K helix of subunit B come into close proximity (Fig. S1C, D), resembling the recently determined structure of GluK2 bound to glutamate, an allosteric positive modulator BPAM344, and Concanavalin A (ConA) in an open state, using time-resolved cryo-EM (PDB code 9B36)[27]. Specifically, the distances between the Cα atoms of K676 in subunit A and N802 in subunit B are 6.2 Å and 6.7 Å in the homology model and the active/open GluK2 structure, respectively. By contrast, these distances are much greater in the BAPM344-bound (no agonist supplemented) inactive state (PDB code 8FWS)[32], as well as in the antagonized (PDB code 5KUH)[28], and desensitized (PDB code 9B38, 5KUF)[27,28] states (Fig. S1D). Cysteine crosslinking approaches have been used to analyze the dynamics and the channel function of iGluRs[50-54]. Additionally, a recent report showed that the conformational changes of KARs upon activation resembles those of AMPARs[27]. Based on these reports, we hypothesized that an engineered disulfide bond between residues K676 and N802 would restrict the mobility of the LBD and trap a dimer-of-dimers LBD conformation, potentially altering desensitization kinetics. To test this hypothesis, we introduced a pair of cysteine residues designed to form disulfide bonds between the LBD dimer pairs (Fig. S1 E, F) and assessed the channel activity of the full-length rat GluK2 K676C/N802C mutant using whole-cell patch-clamp recordings. Note that the distance between the Cα atoms of two cysteines required to form a disulfide bond is reported to be less than 7 Å[55].

GluK2 wild type (WT) demonstrated the rapid desensitization, as observed previously[35].

By contrast, the GluK2 K676C/N802C mutant, activated by 1 mM glutamate, showed earlier rapid

desensitization and a steady-state current (Fig. 1A). When we assessed the GluK2 K676C and

GluK2 N802C single cysteine mutants, both exhibited nearly complete desensitization, similar to

GluK2 WT, without displaying the steady-state current observed when the GluK2 K676C/N802C

double cysteine mutant was activated by glutamate (Fig. S2A). These results confirm that disulfide

bond formation between K676C and N802C are responsible for generating the steady-state

current.

An earlier study showed that BPAM344 significantly slows the desensitization kinetics of

GluK2 WT and enhances the glutamate-evoked current by 140-fold and 21-fold, respectively[56].

To assess how BPAM344 modulates the activation and desensitization of the GluK2

K676C/N802C mutant, we next analyzed its effect. Indeed, the percent desensitization of GluK2

K676C/N802C elicited by 1 mM glutamate is dramatically decreased in the presence of 0.5 mM

BPAM344, with only 25% desensitization observed after 20 seconds (Fig. 1 A, B). While the peak

current amplitude (I_{peak}) was increased 1.5-fold (1.5 ± 0.3, n=10), similar to the WT in the presence

of BPAM344, the steady-state current (I_{ss}) amplitudes were also significantly increased, by

approximately 4.7-fold (4.7 ± 0.6, n=10) (Fig. 1C). Thus, we sought to capture the conformation

of receptors in this non-desensitizing state using cryo-EM.

GluK2 K676C/N802C adopts multiple conformations

We expressed the intact homotetrameric GluK2 KAR incorporating K676C/N802C double

cysteine mutations in HEK293S GnTI⁻ cells and purified the proteins in the detergent, n-Dodecyl-

B-D-Maltoside (DDM). The final protein sample (Fig. S2B, C) was pre-incubated with 1 mM

glutamate, with or without 0.5 mM BPAM344, prior to grid freezing. We initially performed 3-

dimensional (3D) refinement without imposing symmetry (Table S1). Our initial global

reconstruction of the glutamate/BPAM344-supplemented dataset generated six main classes (Fig.

S3a). One of the classes has a two-fold rotational symmetrical conformation (Fig. 1D), with the

extracellular domains resembling the open-state conformation (PDB code 9B36)[27]. To improve

the density, we further performed signal subtraction on the ATD and LBD-TMD layers separately.

The local refinements with focused masks yielded 3.7 Å and 3.9 Å reconstruction of the ATD and

LBD-TMD, respectively, enabling reliable model building (Fig. S3a, S4 A–C). The reconstruction

of the LBD-TMD showed clear density for BPAM344 at the LBD dimer interface consistent with

previous reports[27,32,57] (Fig. 1E–G). All four LBD bi-lobes are closed in the presence of glutamate,

compared to its open conformation in an apo GluK2 (PBD code 9CAZ) structure (Fig. 1H, I). While

the extracellular domain adopts an active-like conformation, the ion channel pore remains closed.

Therefore, we refer to this class as a non-desensitized conformation.

In contrast, the remaining five classes showed no BPAM344 molecules at the LBD D1-D1

interface, leading to D1-D1 rupture. In all of these five classes, the LBD bi-lobes of all four subunits

are tightly closed in the presence of glutamate, with closure degrees ranging from 23° to 27°,

indicating glutamate binding to the receptors. The overall structures of these five conformations

are stabilized in two- or four-fold symmetrical, or asymmetrical arrangements. In order to analyze

conformations in the absence of allosteric potentiators, we also collected cryo-EM data of GluK2

K676C/N802C supplemented with glutamate. These data revealed four main three-dimensional

(3D) classes (Fig. S3b), each are superimposable with the glutamate-bound, BPAM344-unbound

classes from the BPAM344-supplemented sample (Fig. S3a). Thus, we combined these two data

sets and further performed processing and refinement to improve the density quality. We defined

these classes as shallow-desensitized, intermediate, and deep-desensitized states, based on the

observations described below. Ultimately, the final resolutions of desensitized classes range from

3.7 Å to 4.0 Å (Fig. S4).

GluK2 K676C/N802C mutant structure bound to glutamate and BPAM344 represents a non-desensitized conformation

While all six 3D reconstructions showed three domains—ATD, LBD, and TMD—as previously reported[25,28,30,31,33], each class showed distinct conformations (Fig. S3). In the non-desensitized structure, two molecules of BPAM344 per LBD dimer bind and stabilize the LBD D1-D1 interfaces, with each BPAM344 forming hydrogen bonds with the side chain of P532, as recently reported[27,32,57] (Fig. 1G). In addition to BPAM344, clear densities corresponding to disulfide bonds between K676C and N802C were observed in this conformation (Fig. 1J). These two disulfide bonds trapped the LBDs in a dimer-of-dimers conformation in the presence of the agonist, as designed (Fig. 1D, F). Previously, the structure of open GluK2 was determined in the presence of two positive allosteric modulators, concanavalin A (ConA), BPAM344, as well as the agonist glutamate (PDB code 9B36)[27]. In the structure, the ~50 kDa ConA dimer, along with BPAM344 and glutamate, binds at the ATD-LBD interface, elevating the ATD by 16 Å and rotating the LBD relative to the ATD layer by 28° (Fig. S5). The same tendency was observed in the presence of NETO auxiliary proteins, which also bind to the ATD and LBD layers and modulate KAR kinetics[30]. Interestingly, a comparison between structures without an orthosteric ligand (PDB codes 8FWQ, 9CAZ)[32] and our non-desensitized structure revealed that the distance between the center of mass (COM) of the ATD and LBD was also increased by 7.5 Å in the non-desensitized conformation, even in the absence of ConA and NETO auxiliary proteins. By contrast, in the desensitized state, this distance between the ATD and LBD decreased by 2–4 Å compared to the apo state. While, the ATDs in all structures are superimposable and do not undergo any conformational changes (Fig. S5B), the ATD in the non-desensitized class also displayed a rigid-body rotation of 45° counterclockwise (Fig. S5C), following the same movement tendency observed in the open state of KARs[27] and AMPARs[58].

Focusing on the LBD layer, the LBDs are stabilized in a dimer-of-dimers arrangement that

resembles the open conformation (Fig. 2A), with the D1–D1 association stabilized by BPAM344

and the D2–D2 separation maintained within each LBD dimer in the presence of glutamate (Fig.

2B). Analysis of the center-of-mass (COM) distances between D1-D1 and D2-D2 in the two

structures further indicates their similar conformational arrangement. Specifically, the distance

between the Cα atoms of S670, located at the bottom of the D2 lobes in the AD and BC subunit

pairs, is equally separated as seen in the open-state conformation. Key features of the open-state

conformation of iGluRs include increased tension in LBD-TM3 linkers and the expansion of the

ion channel pore held by the linkers, particularly at the gate region[1,59]. Thus, we next compared

the LBD-TM3 linker arrangement, TM3 helix orientation, and channel openings in the BPAM344-

bound (without an orthosteric ligand), non-desensitized, and open GluK2 conformations. The

densities of all TM3 helices are well resolved (Fig. S6). In the non-desensitized conformation, the

TM3 helices are unwound at A656 in the BD subunits and at E662 in the AC subunits (position

(3) and (2'), respectively), and therefore, T660 (BD subunits) and M664 (AC subunits) are pulled

away from the ion channel central axis, while BPAM344-bound GluK2 (PDB code 8FWS)[32]

displays an additional half-turn of the helix at the pore entrance, and the TM3 helices unwind at

T660 in the AC subunits and at M664 in the BD subunits (position (2) and (1), respectively) (Fig.

2C). Consequently, the top of the gating region, which contains the conserved SYTANLAAF

motif[60,61], is more dilated in the non-desensitized conformation compared to the more tightly

closed BPAM344-bound GluK2 conformation. By contrast, the TM3 helices in the open GluK2

conformation (PDB code 9B36)[27] display kinking at a deeper position, specifically at L655

(position (4)) in all four subunits (Fig. 2D). Additionally, the open GluK2 conformation exhibits the

rotational movements of TM3, which put away T652 from the central axis of the pore, thus opening

the channel. However, these rotational movements are not observed in the non-desensitized

conformation. Overall, the ion channel in the non-desensitized conformation remains closed (Fig.

2E, F).

Disulfide bonds between K676C-N802C stabilized GluK2 KARs in a desensitized AMPA

receptor like conformation

The glutamate-bound, potentiator-unbound GluK2 K676C/N802C generated five classes

(Fig. 3A). The LBD layer conformations in these classes appear desensitized; however, they

represent different levels of desensitization (Fig. 3B). In one of these desensitized classes, the

absence of BPAM344 results in D1–D1 dissociation, a hallmark structural feature of iGluR

desensitization, while two disulfide bonds between K676C and N802C (Fig. 3C) stabilize LBD

dimers forming a dimer-of-dimers conformation. We refer to this as the 'shallow-desensitized'

conformation to distinguish it from the typical deep-desensitized conformations previously

observed[28]. Another class, termed the intermediate state, contains an LBD dimer on one side and

a disrupted dimer on the other. In this class, one LBD dimer pair maintains its dimer formation

with D1–D1 dissociation, resembling the shallow-desensitized state, while the disrupted dimer

adopts a conformation characteristic of deep-desensitized classes. Therefore, we propose that

this structure represents an intermediate state between the shallow- and deep-desensitized

conformations in the presence of glutamate. In contrast, the other three structures lacked inter-

subunit disulfide bonds between K676C and N802C, resulting in the LBD layers stabilizing an

approximately four-fold symmetric conformation, though with varying in-plane rotation angles of

the LBDs (Fig. 3B). We observed the formation of a 'desensitization ring' (Fig. 3D, E) in all three

deep-desensitized conformations, consistent with previous reports[28,29,62]. This ring formation plays

a key role in regulating desensitization and recovery from desensitization[28,29]. Accordingly, we

classified these classes as 'deep-desensitized' states."

Comparison of the non-desensitized and shallow-desensitized conformations reveals

multiple structural rearrangements in the context of the full-length receptor structure. The distance

between the ATD and LBD decreases in the shallow-desensitized state, measuring 2–3 Å shorter

than in the apo and BPAM-bound GluK2 structures, making it more similar to the deep-

desensitized state than to the open-state conformation (Fig. S5A). Focusing on the LBD layer,

the LBDs in the AC subunits are rotated by 5° parallel to the membrane when viewed from the

top, compared to the non-desensitized conformation (Fig. 4A). Viewed from the side, the entire

LBDs appear rolled down (Fig. 4B), which is associated with the rearrangement of the dimer-of-

dimers formation and the rupture of the D1-D1 interface. These conformational changes are also

observed during AMPAR desensitization[44,63-66]. Therefore, we next compared the shallow-

desensitized conformation of GluK2 K676C/N802C with the desensitized conformation of GluA2

AMPAR (PDB code 7RYZ)[64]. Viewed from the top, the distance between the Cα atoms of S700

(S662 in GluA2 AMPAR) in both structures are reduced compared to their respective open states,

although the arrangements in these two structures differ due to the restricted mobility of GluK2

caused by cysteine crosslinks (Fig. 4C). The D1 lobes of GluK2 roll down approximately 25°, in

contrast to the 33° rolling-down motion observed in the D1 lobes of GluA2 (Fig. 4D). The D2

distances in the shallow-desensitized GluK2 and desensitized GluA2 are comparable, measuring

20 Å and 24 Å, respectively. Overall, the shallow-desensitized conformation of the GluK2 mutant

appears to resemble the desensitized conformation of AMPA receptors. However, this

conformation in KAR is energetically unstable and has not been observed in the absence of

crosslinks. Comparison of the TM3 in the non-desensitized and shallow-desensitized

conformations revealed that the pore in the shallow-desensitized state is occluded at T660, and

M664 in the BD and AC subunits, respectively (position (2) and (1)) (Fig. 4E), while the ion channel

pore in the deep-desensitized conformations is sealed at M664 (position (1)) in all four subunits

(Fig. 4F).

A large in-plane rotation of the LBD is required for proper ion channel closure during

desensitization in GluK2 KAR.

We next assessed the ion channel pore in these desensitized conformations. Although the

extracellular domain in the shallow-desensitized GluK2 KAR and the desensitized GluA2 AMPAR

adopt similar conformations, we observed distinct differences in their ion channel pore structures

(Fig. 5A–C). The shallow-desensitized conformation has restriction points around T652, A656,

and T660, which correspond to T617, A621, and T625 in the desensitized GluA2 (PDB code

7RYZ)[64]. Notably, the pore diameter at T660 and T652 in the shallow desensitized GluK2 KAR is

wider compared to that in the desensitized GluA2 AMPAR. The ion channel pore in the transition

state is also similar to that in the shallow-desensitized state, except for the presence of a narrower

restriction point (<1 Å) around T652 (Fig. 5A). By contrast, all three deep-desensitized

conformations exhibit multiple tight constrictions, particularly at the top of the gate region (Fig. 5A,

D).

As previously reported[28,29], the 'desensitization ring' at the bottom of the LBD layer in these

deep-desensitized conformations is stabilized by numerous polar interactions, supporting the

four-fold symmetrical deep-desensitized conformations, whereas the shallow-desensitized and

intermediate conformations exhibit disrupted ring formation, resulting in wider gate regions.

Moreover, the large in-plane rotation of the LBDs in the BD subunits during desensitization twists

the LBD-TM3 linker, creating an additional constriction at M664 at the top of the channel pore

(Fig. 4F), which is not seen in AMPARs. We observe that the formation of the desensitization ring

at the bottom of the LBD layer tightly closes the ion channel in the deep-desensitized

conformations, whereas the shallow-desensitized state lacks this secure channel pore lock at

T660 and M664 (Fig. 5A).

Based on these functional and structural observations, we hypothesized that the GluK2

KAR K676C/N802C mutant, which forms cysteine crosslinks, undergoes shallow rather than deep

desensitization. This may allow the receptors to remain ion-permeable, exhibit a higher open

probability, or recover quickly from desensitization, similar to AMPARs, thereby enabling more

efficient activation of the receptors.

302 K676C-N802C cysteine crosslinks partially blocks receptors from undergoing deep

303 desensitization

To test our hypothesis, we first assessed the kinetics of the GluK2 K676C/N802C mutant

using whole-cell patch-clamp electrophysiology. Remarkably, we observed steady-state currents

of the cross-link mutant even at the highest concentration of glutamate (50 mM), which were not

observed in the wild type GluK2 receptors (Fig. 6A). There were no significant differences neither

in the current density at peak current between GluK2 WT and the mutant (GluK2 WT: 830 ± 50

pA/pF, n=6; GluK2 K676C/N802C: 700 ± 150 pA/pF, n=9), nor in the EC_{50} for glutamate (GluK2

WT: 110 ± 30 µM, n=7; GluK2 K676C/N802C: 150 ± 50 µM, n=5), indicating that the cysteine

crosslinks do not significantly affect agonist binding affinity or receptor activation (Fig. 6A-D).

GluK2 K676C/N802C partially desensitized even by 0.01 mM glutamate, a concentration below

the EC_{50} (approximately 0.17 mM). This suggests that even partial occupancy is sufficient to

initiate the conformational changes leading to desensitization, consistent with earlier findings[67,68].

While GluK2 WT desensitizes almost completely with a time constant of 8 ± 1 ms (n=14), the

GluK2 K676C/N802C receptors in the presence of 0.01–50 mM glutamate exhibited

approximately 60% desensitization, except when activated by 0.001 mM glutamate where only

the steady-state current was observed with a decreased peak amplitude (Fig. 6E). Therefore, the

degree of desensitization is not glutamate concentration dependent.

To assess the desensitization state of the crosslinked mutant, we next analyzed the rate

of recovery from desensitization. As previously reported, recovery from desensitization in the

GluK2 WT is extremely slow, requiring 1,150 ± 90 ms (n = 9) for full recovery. By contrast, the

GluK2 K676C/N802C mutant exhibited a faster recovery from desensitization, characterized by

two distinct components: τ_1 (100 ms, 60%) and τ_2 (1,000 ms, 40%) (n = 9). The τ mean of the

mutant was 440 ± 40 ms, approximately three times faster than that of GluK2 WT (Fig. 6F–H).

This shift brings its kinetics closer to those of AMPARs, which recover in the order of tens to

hundreds of milliseconds (e.g. GluA1 and GluA2 wild-type recovery constants are approximately

200 ms and 9-50 ms, respectively)[69-71]. This result highlights the impact of cysteine cross-linking

in accelerating desensitization recovery, potentially by stabilizing the dimer-of-dimers

conformation.

GluK2 KAR stabilized in a shallow-desensitized conformation remains ion-permeable

To assess whether the non-desensitized current and faster recovery kinetics result from

crosslinking between the K676C and N802C, we performed whole-cell recordings in the presence

of 5 mM dithiothreitol (DTT), applied for 7 minutes prior to the second glutamate activation. The

peak intensity of both GluK2 WT and GluK2 K676C/N802C, in the presence and absence of DTT,

showed no significant changes, confirming our earlier analysis that the crosslinks do not affect

activation kinetics (Fig. 7A). By contrast, the application of 5 mM DTT for 7 minutes led to a >90%

reduction in the steady-state current of GluK2 K676C/N802C (steady-state current prior to and

after DTT application: 0.4 ± 0.1 nA and DTT: 0.03 ± 0.03 nA, respectively, n=8) (Fig. 7B).

Additionally, recovery from desensitization of the DTT treated cells expressing GluK2

K676C/N802C was 1,040 ms ± 50 (n=5), similar to the WT (Fig. 7C, D). This experiment, together

with the assessment of the single cysteine mutants, GluK2 K676C and GluK2 N802C (Fig. S2A),

confirms that the observed non-desensitizing efficacy is specifically due to disulfide bond

formation, which stabilizes the two-fold symmetrical non-desensitized or shallow-desensitized

conformations as originally intended.

Next, we aimed to determine whether crosslinked receptors remain ion-conductive when

exhibiting steady-state current. A key feature of KARs is their exceptionally small conductance

and low open-channel probability, which makes it technically challenging to directly observe

channel activity at the single-channel level, unlike other channels, including AMPARs[72,73].

Therefore, we conducted noise analysis to assess its conductivity. We first performed

nonstationary fluctuation analysis (NSFA) on the desensitizing phase of macroscopic currents

from GluK2 WT and GluK2 K676C/N802C, activated by glutamate pulses (10 mM, 100 ms). The

variance of fluctuations around the expected current was analyzed. The weighted mean single-

channel conductance of GluK2 WT was calculated as 23 ± 7.8 pS (n = 5), consistent with previous

report[72]. While the current-variance relationship of the wild type was well-fitted by a parabolic

function, the analysis of GluK2 K676C/N802C revealed an anomalous current-variance

relationship, preventing the calculation of its conductance (Fig. 7E). However, the mutant

generates a unique large fractional steady-state current. Thus, we next analyzed stationary

fluctuation on the steady-state current of the desensitized receptors in the presence and absence

of BPAM344.

To assess unitary conductance and the number of active channels, we analyzed the

microscopic currents of GluK2 K676C/N802C activated by 10 mM glutamate for 30 seconds under

desensitizing conditions (Fig. 7F). When activated by 10 mM glutamate, the mutant exhibited a

unitary conductance of 10 pS—less than half of the wild-type conductance calculated by NSFA—

and was similar to the lowest conductive level (O1) previously determined for KARs[72]. Variance-

current analysis across individual cells showed that BPAM344 shifted receptor behavior toward

higher variance and mean current values (Fig. 7G), suggesting an increase in the number of open

receptors and their open probability. Interestingly, the conductance of the mutant activated by 10

mM glutamate was similar in the presence and absence of BPAM344 (GluK2 K676C/N802C, 10

mM glutamate: 10 ± 3 pS, n = 6; GluK2 K676C/N802C, 10 mM glutamate + 0.5 mM BPAM344:

10 ± 2 pS, n = 6) (Fig. 7H). Whereas the number of active channels in the presence of BPAM344

was approximately four-fold higher than in its absence (GluK2 K676C/N802C, 10 mM glutamate:

1,040 ± 600, n = 6; GluK2 K676C/N802C, 10 mM glutamate + 0.5 mM BPAM344: 4,080 ± 460, n

= 6) (Fig. 7I). In summary, these experiments indicate that desensitized GluK2 K676C/N802C

KARs remain conductive at the single channel level, although with significantly lower conductance

compared to wild-type channels activated by glutamate.

Discussion

Receptor desensitization is a fundamental property of ligand-gated ion channels which

depends on agonist, receptor subtypes, and auxiliary proteins. Recent advances in iGluR

research have revealed significant differences in the kinetics and conformational dynamics of

KARs during desensitization compared to other iGluRs[1,25,28,29,31,33]. The desensitized conformation

of KARs is characterized by the quasi-four-fold symmetrical LBDs, by contrast to typical

desensitized iGluRs, which remain in two-fold symmetrical conformations. In addition, functionally,

the majority of KARs enter deep-desensitized states, with recovery from desensitization occurring

5- to 130-fold slower than AMPARs. By contrast, most AMPARs typically adopt shallow-

desensitized states, exhibiting deep desensitization only under certain conditions[74,75]. In this study,

we employed a classical cysteine crosslinking approach to answer the fundamental question why

KARs require such dramatical conformational changes for proper channel closure and complete

desensitization, which are not conserved in other iGluRs.

In the previously determined structure of BPAM344- and ConA-bound GluK2 in a highly

ion-conductive open state, ConA binds both the ATD and LBD, occupying the space between

these two layers[27]. As a result, the organization of the ATD-LBD layers upon activation remains

unclear. In this study, we analyzed the extracellular domain organization in the absence of ConA.

The assembly of the extracellular domains of our glutamate- and BPAM344-bound GluK2

K676C/N802C in the non-desensitized state resembles that of GluK2 in its open state.

Interestingly, in the absence of ConA, the ATD-LBD distance in the non-desensitized state of

GluK2 K676C/N802C is wider. Thus, upon activation, the ATD layer is elevated and undergoes a

counterclockwise in-plane rotation. By contrast, the apo and desensitized receptors adopt more

compact extracellular domain conformations, with ATD and LBD layers positioned closer. Since

ATD-deleted KARs remain functional as ion channels[27,76], further investigation is needed to

determine how these conformational changes in the ATD layer contribute to receptor activation.

In addition, we also observed the TM3, which form the gating region of the ion channel pore, kinks

at the position between the kinking observed in the apo and the open structure (Fig. 8), along with

rotational movement of TM3. This observation suggests that TM3 kinking at a higher position

(A656/E662) within the gating region is insufficient for channel opening—particularly full opening

with high conductivity—which instead requires TM3 rotation and kinking at a lower position (L655)

in all four subunits (Fig. 8).

Conversely, the glutamate-bound GluK2 K676C/N802C structures represent shallow- and

deep- desensitized conformations as well as an intermediate state between these two

desensitized states. The shallow-desensitized conformation demonstrates that two inter-dimer

crosslinks stabilize receptors in a two-fold symmetrical conformation with D1-D1 ruptured LBDs,

similar to the desensitized state of AMPARs[42,44,63-65]. While most previously determined AMPAR

structures in desensitized states are stabilized in the classical two-fold symmetrical conformation,

the partial agonist, fluorowilliardiine-bound GluA2 AMPAR, in a desensitized state exhibited an

asymmetrical conformation with the LBD of subunit D undergoing a large conformational

rearrangement similar to our GluK2 in the intermediate-state conformation[77]. Additionally, recent

studies have shown that extracellular protons influence extracellular domains and alter LBD

conformation, causing minor in-plane rotation of LBDs within the submit B and D, although these

changes are not as pronounced as those observed in the deep-desensitized KAR

conformations[70,71,78]. Nevertheless, adopting the four-fold symmetrical deep-desensitized

conformation is not energetically favorable for AMPARs during desensitization[74], whereas KARs

exhibit greater stability in this four-fold conformation. Previous mutagenesis studies demonstrated

that various mutations in the D2 lobe of KARs alter desensitization and recovery from

desensitization, but no mutation completely blocked desensitization unlike AMPARs[62]. This

suggests that multiple non-conserved residues contribute to the stability of the four-fold

desensitized conformation of KARs, despite the overall architectural similarity between AMPAR

and KAR LBDs. Our electrophysiological experiments demonstrate that cysteine crosslinking

prevents the receptors from undergoing complete desensitization, resulting in significantly

stronger and faster recovery—resembling the kinetics of AMPARs rather than those of GluK2 WT.

Upon treatment with 5 mM DTT, which breaks the disulfide bonds in the GluK2 K676C/N802C

mutant, recovery from desensitization resembled that of the WT, indicating a correlation between

LBD conformation and desensitization recovery in GluK2 KAR.

Furthermore, we aimed to understand why KARs have to form the four-fold symmetrical

arrangement upon desensitization. Previous studies have shown that in the "desensitized ring", a

network of polar interactions stabilizes the deep-desensitized conformation of the LBD layer. This,

in turn, helps to stabilize the conformation of the TMD layer, thereby reinforcing receptor

desensitization and influencing the kinetics of desensitization recovery[28,29]. In the shallow-

desensitized conformation, lacking desensitization ring formation with the two-fold symmetric

LBDs, we observed that the ion-channel is dilated by contrast to channels observed in

desensitized AMPARs and fully/deep desensitized KARs, which show much narrower channel

pores throughout the gating region with the channel pore tightly sealed on the top of the gating

region. Previously, a conducting desensitized state of AMPAR and acetylcholine receptors has

been reported[41,79]. In addition, it was microscopically shown that the partially desensitized KARs

are conductive, as defined by the number of desensitized subunits per KAR tetramer[80]. The

structural observation together with the characteristic steady-state current of the GluK2

K676C/N802C indicates that the receptors in the shallow-desensitized state could form weakly

conductive channels. Indeed, our noise analysis demonstrates that the desensitized GluK2

K676C/N802C receptors, exhibiting steady-state current, remains conductive at the single-

channel level; however, its conductance is approximately 30% weaker than the GluK2 WT,

regardless of the presence of the positive allosteric modulator BPAM344. In addition, the

crosslinked receptors exhibit up to ten-fold faster recovery from desensitization, whereas reducing

the disulfide bonds with DTT restores currents similar to those of the GluK2 WT.

Taken together, this study answers the fundamental question by demonstrating that KARs

absolutely require large conformational rearrangements of the LBD layer and stabilization in a

four-fold symmetrical conformation to achieve proper channel closure and full desensitization into

a non-conductive state (Fig. 8). This structural feature distinguishes KARs from other iGluR

subfamilies, in which such symmetry and conformational changes are not observed.

Materials and Methods

Plasmid construction

For Cryo-EM studies, the construct used for the structural studies was full-length rat GluK2 KAR

(GenBank 54257, Uniprot code P42260) with two mutations of N802C and K656C. The gene was

cloned into the pFW vector (gift from Dr. Hiro Furukawa at Cold Spring Harbor Laboratory) and

fused to a C-terminal human rhinovirus 3C protease recognition site and a 1D4 affinity tag. The

construct was referred to as GluK2 N802C/K656C.

Cell culture

For structural studies, HEK293S GnTI$^-$ cells were grown to a density of 3.2×10^6 cells / ml

in FreeStyle 293 medium at 37°C and 8% CO_2 supplemented with 2 % fetal bovine serum. The

rGluK2 N802C/K676C mutant bacmid and baculovirus were generated as previously described.

The P1 and P2 viruses were produced in Sf9 cells. Cells were infected with the baculovirus

harboring GluK2 EM and incubated at 37°C for 12 hrs. Cells were supplemented with 10mM

sodium butyrate and 20 μM DNQX and temperature was shifted to 30°C at 12hrs post-infection.

The cell culture was incubated additional 60 hrs and harvested by low-speed centrifugation at

5,000 g for 20min and stored at -80°C until use.

For electrophysiology experiments, human embryonic kidney 293T (HEK293T) cells were

cultured in Dulbecco's modified Eagle medium (DMEM, Corning) supplemented with 10% FBS at

37°C in a 95% O_2–5% CO_2 atmosphere. Wild-type rat GluK2 and rat GluK2 K676C/N802C DNAs

were cloned into the pCAG-IRES-EGFP vector (Addgene plasmid #119739). HEK293T cells were

transfected with 1 μg/μl cDNA using the TransIT2020 transfection reagent (Mirus) according to

the manufacturer's instructions. Cells expressing GluK2 wild-type were transfected for 24 hours,

while those expressing K676C/N802C were transfected for 18 hours. After transfection, cells were

dissociated using Accutase (Innovative Cell Technologies, Inc.), resuspended, and plated onto

35 mm poly-D-lysine-coated dishes (Neuvitro). Electrophysiological recordings were performed

2–4 hours later[81].

Electrophysiological recordings

All whole-cell recordings were performed using HEKA EPC10 amplifiers (HEKA Elektronik,

Lambrecht, Germany) with thin-wall borosilicate glass pipettes (2–5 MΩ) coated with dental wax

to reduce electrical noise. Currents were recorded at a holding potential of −70 mV, with a

sampling frequency of 10 kHz, and filtered at 2.6 kHz. The external solution contained (in mM):

145 NaCl, 2.5 KCl, 1.8 $CaCl_2$, 1 $MgCl_2$, 5 glucose, and 5 HEPES. The internal pipette solution

contained (in mM): 105 NaCl, 20 NaF, 5 Na_4BAPTA, 0.5 $CaCl_2$, 10 Na_2ATP, and 5 HEPES. The

pH and osmotic pressure of the external and internal solutions were adjusted to 7.4 and 300–290

mOsm/kg, respectively. L-glutamate was applied using theta glass tubing mounted on a

piezoelectric stack (MXPZT-300 series, Siskiyou) driven by a HEKA EPC10 amplifier. Typical 10–

90% rise times were 250–300 μs, measured from junction potentials at the open tip of the patch

pipette after recordings. Data acquisition was performed using PULSE software (HEKA Elektronik, Lambrecht, Germany).

Data analysis

The macroscopic rate of desensitization (τ_desensitization) was measured by fitting an exponential function to the decay of current from ~80% of its peak amplitude (I_peak) to baseline in recordings where glutamate was applied for 1 second. For the double mutant N802C/K676C, only the first exponential component, which exhibited desensitization kinetics, was considered, while the steady-state current was omitted. Desensitization kinetics were fitted using a single-exponential, one-term Levenberg-Marquardt fitting approach.

To evaluate recovery from desensitization, two protocols were employed: (1) a two-pulse application of 10 mM glutamate for 100 ms per pulse, with interpulse intervals increasing in 100 ms steps from 50 ms to 2 s; and (2) a second protocol with interpulse intervals increasing in 2 s steps, ranging from 50 ms to 20 s. In all cases, the first peak was considered the control (100% activation), while the second peak represented recovery at a given time point. Data from both protocols were combined, and a recovery curve was generated, plotting the percentage of recovery against time. The experimental data were fitted to one-term exponential model, from which the recovery time constant (τ) was derived.

To analyze the EC_{50} for glutamate in wild-type (WT) and mutant receptors, varying concentrations of L-glutamate were applied to a single cell using a theta glass tubing mounted on a piezoelectric stack (MXPZT-300 series, Siskiyou), driven by a HEKA EPC10 amplifier, for 1 second. One input of the theta glass tubing was connected to the control solution, while the other was connected to a six-inlet manifold for different L-glutamate concentrations. A 2-minute wash was used between concentration changes to ensure complete exchange before applying the next solution. The L-glutamate concentrations tested were 50, 10, 1, 0.1, 0.01, and 0.001 mM, with 50 mM serving as the normalization reference. The normalized data are presented as mean ± S.E.M.

in a dose-response curve, with response on the y-axis and concentration on the x-axis.

Experimental data from independent biological measurements (each defined as data from one

cell with at least five concentration responses) were fitted to the Hill equation. The resulting EC_{50}

values were then averaged and presented as mean ± S.E.M.

Non-stationary fluctuation analysis (NSFA)

NSFA was performed on the decay phase of currents evoked by 200 ms applications of

10 mM glutamate (60–150 successive applications). Currents were sampled at 50 kHz and filtered

at 5 kHz. The variance and mean current were calculated for each set of currents and then binned

into 15 bins. The single-channel current and total number of channels (N) were determined by

plotting the binned variance against the binned mean current and fitting the data with a parabolic

function:

$$\sigma^2 = i\bar{I} - \frac{\bar{I}^2}{N} + \sigma_B^2$$

Where σ_B^2 represents the background variance, which was calculated from the baseline before

glutamate application. The σ_B^2 value was used to correct the variance, while the mean I_B was used

to correct the mean current.

Stationary fluctuation analysis (SFA)

HEK293T cells transfected with GluK2 K676C/N802C were activated with 10 mM

glutamate for 30 seconds, followed by a 2-minute incubation with 500 µM BPAM344. The cells

were then reactivated with 10 mM glutamate + 500 µM BPAM344. Currents were sampled at 10

kHz and filtered at 2 kHz. For each condition (glutamate alone and glutamate + BPAM344), the

last 10 seconds of the steady-state current were used to calculate the variance and mean current.

Since the stationary current represented approximately 35% of the peak current, we assumed

that the open probability (P_0) in the steady-state current was $P_0 < 1$. The unitary current (i) was

then estimated using the following equation:

$$i = \frac{\sigma^2}{\bar{I}}$$

The unitary current was then used to estimate the unitary conductance (γ) as follows:

$$\gamma = \frac{i}{Vm - Vr}$$

where V_m is the holding potential (−70 mV) and the reversal potential (V_r) is 0 mV.

The number of channels (N) was estimated using the equation:

$$N = \frac{\bar{I}^2}{\sigma^2}$$

Statistical analyses were performed using OriginLab 2025. Statistical significance was calculated

using a two-sided two-sample t-test, with significance assumed if $P < 0.05$.

Protein expression and purification

Cell pellets were resuspended in ice-cold lysis buffer containing 20 mM Tris-Cl (pH 8.0),

150 mM NaCl, 0.2 mM PMSF, 20 µg/mL DNase I, and 20 µM DNQX. Cells were lysed by

sonication using a Qsonica sonicator with an amplitude of 40% for 10 minutes total (10 s on, 10

s off). The lysate was centrifuged at 7,000 g for 20 min to remove cell debris and unbroken cells,

and the supernatant was subjected to ultracentrifugation at 200,000 g for 30 min to pellet the cell

membranes. The membrane pellet was homogenized and solubilized by gentle nutation for 2

hours at 4 °C in buffer consisting of 20 mM Tris-Cl (pH 8.0), 150 mM NaCl, 1% Dodecyl-β-D-

Maltosid (w/v, DDM), and 20 µM DNQX, followed by centrifugation at 200,000 g for 30 min to

remove insoluble material. The supernatant was incubated with Rho 1D4 affinity resin overnight

at 4 °C. The protein-bound resin was extensively washed with 100 mL of buffer consisting of 20

mM Tris-Cl (pH 8.0), 500 mM NaCl, 1 mM DNQX, and 0.02% DDM at room temperature, and

then slowly washed with 200 mL of buffer consisting of 20 mM Tris-Cl (pH 8.0), 300 mM NaCl,

and 0.02% DDM at room temperature at a rate of approximately 0.3 ml/min. The protein was

eluted in 50 mL of buffer with 20 mM Tris-Cl (pH 8.0), 300 mM NaCl, 0.02% DDM, and 0.25 mM

1D4 peptide. The eluted protein was concentrated and injected into a Superose 6 10/300 GL size-

exclusion column equilibrated with buffer containing 20 mM Tris-Cl (pH 8.0), 150 mM NaCl, and

0.02% DDM. The peak fractions corresponding to tetrameric GluK2 were pooled and

concentrated to 3.4 mg/mL for grid freezing.

Grid preparation and cryo-EM data

The cryo-EM grids were prepared as follows. UltrAuFoil R1.2/1.3 300 mesh gold grids

were glow-discharged for 50 seconds before sample application. A final concentration of 1 mM

glutamate was added to the sample, and then a droplet of 3 µL of protein sample was rapidly

applied onto the grids. For the BPAM344/glutamate sample, proteins were pre-incubated with 500

µM BPAM344 for 30 minutes on ice, followed by the addition of 1 mM glutamate to the sample.

3.5 µl of the protein sample was applied to the grids, and the grids are frozen within 10 seconds.

An FEI Vitrobot Mark IV (Thermo Fisher Scientific) was used to plunge-freeze the grids into liquid

ethane after sample application at 4°C and 100% humidity, with a blot time of 3 seconds and a

blot force of −5.

All cryo-EM data were collected on a Titan Krios 300 kV microscope (Thermo Fisher

Scientific) equipped with a GIF-Quantum energy filter (Gatan) with the slit set to 20 eV and a

Gatan K3 Summit direct electron detection camera (Gatan) at Case Western Reserve University.

Raw movies were collected using EPU in super-resolution mode, with a physical image pixel size

of 1.06 Å and a defocus range of −0.8 to −1.6 µm. The total dose of around 50 e⁻/Å² was used

for each movie, distributed across 50 frames for a total exposure time of approximately 3.2

seconds.

Cryo-EM image processing

For dataset **a** (GluK2 K676C/N802C supplemented with 1 mM glutamate and 500 µM BPAM344), the same motion correction procedure was applied using Patch Motion Correction in CryoSPARC4. CTF calculations were also carried out on the motion-corrected images using Patch CTF Estimation in CryoSPARC4. Micrographs unsuitable for further analysis were manually removed. Initial particle selection from 1000 micrographs was done with Blob Picker, followed by 2D classification to generate good templates. Template Picker was used to pick 903,654 particles, which were extracted with a pixel size of 2.12 Å. After several rounds of 2D classification, 612,540 particles were selected for initial 3D model generation via ab initio reconstruction. A total of 352,269 particles were selected and re-extracted with a pixel size of 1.06 Å following multiple rounds of heterogeneous refinement. NU-refinement for each class resulted in maps with resolutions of 6.16 Å, 5.80 Å, 4.14 Å, 4.00 Å, 4.00 Å, and 3.83 Å, respectively.

For dataset **b** (GluK2 K676C/N802C supplemented with 1 mM glutamate), the raw movie stacks were motion-corrected using Patch Motion Correction in CryoSPARC4. The Contrast Transfer Function (CTF) was calculated from the motion-corrected images using the Patch CTF Estimation in CryoSPARC4. Micrographs deemed unsuitable for further analysis were removed through manual inspection. Initial particle selection was performed using Blob Picker from 2,000 micrographs, followed by 2D classification to generate good templates. Subsequently, 1,005,623 particles were picked using Template Picker and extracted with pixel size of 2.12 Å. After several rounds of 2D classification, 765,893 particles were selected to generate the initial 3D model through ab initio reconstruction. A total of 277,369 particles were selected and re-extracted with a pixel size of 1.06 Å after multiple rounds of heterogeneous refinement. NU-refinement for each class yielded maps with resolutions of 3.92 Å, 4.35 Å, 3.85 Å, and 3.82 Å, respectively.

To improve the structure resolution for classes present in both datasets, particles from each class were combined and subjected to several rounds of heterogeneous refinement to remove junk

particles. Separate NU-refinements for each class generated maps with resolutions of 3.67 Å,

4.02 Å, 3.81 Å, and 3.74 Å, respectively. For Class 2, focused masks around the ATD and LBD-

TMD regions were created separately and used for local refinements of the corresponding regions.

As a result, 3.42 Å reconstruction of the ATD region and 3.86 Å reconstruction of the LBD-TMD

region were obtained through local refinement with focused masks. For the non-desensitized state,

particle subtraction was performed for both ATD-focused and LBD-TMD-focused heterogeneous

refinement. After several rounds of heterogeneous refinement, 80,325 particles were used for

subsequent NU-refinement, resulting in a 3.70 Å resolution of the ATD 3D reconstruction. A total

of 117,714 particles were subjected to NU-refinement for the LBD-TMD region, generating 3.93

Å maps. The Gold-standard Fourier Shell Correlation (FSC) 0.143 criteria were used to estimate

the map resolution for each class. For the shallow-desensitized state, focused masks around the

ATD and LBD-TMD regions were created separately and used for local refinements of the

corresponding regions. As a result, 3.42 Å reconstruction of the ATD region and 3.86 Å

reconstruction of the LBD-TMD region were obtained through local refinement with focused

masks.

Model building and refinement

The model building for the non-desensitized class was initiated by individually fitting rigid-

body components of the LBD extracted from the BPAM344-, and ConA-bound GluK2 structure

(PDB code 9B36), and the TMD and ATD from the BPAM344 bound GluK2 structure (PDB code

8FWQ) into the cryo-EM map using Chimera. Subsequent manual adjustments, including fitting

the backbone and side chains of the ATD-LBD and LBD-TM3 linkers into the density, as well as

deleting unresolved side chains and making subtle local adjustments to minimize clashes, were

performed in Coot.

For the shallow-desensitized class, the single LBD, entire ATD, and TMD were separately

extracted from the BPAM344 bound GluK2 structure (PDB code 8FWQ) and manually docked

into the corresponding cryo-EM map using Chimera. Further manual adjustments of the backbone

and side chains were completed in Coot.

For intermediate and deep-desensitized classes, the single LBD, entire ATD, and TMD

were extracted from the deep-desensitized GluK2 structure (PDB: 5KUF) and individually docked

into the corresponding cryo-EM maps for each class using Chimera. Additional manual

adjustments to the backbone and side chains were completed in Coot.

All four initial models were then refined in real space using Phenix against the

corresponding cryo-EM maps. The models were validated through comprehensive validation

procedures in Phenix. Glutamate ligands and BPAM344 compounds were extracted from the

cryo-EM structure (PDB code 9B36) and manually fitted into the corresponding densities in Coot.

Key source table

Reagent type (species) or resource	Designation	Source or reference	Identifiers	Additional information
Gene (*Rattus norvegicus*)	GRIK2_RAT	Genscript (synthesized)	P42260	
Cell line (*Homo sapiens*)	HEK293S GnTI-	ATCC	Cat. No. CRL-3022	
Cell line (*Homo sapiens*)	HEK293T	ATCC	Cat. No. CRL-3216	
Cell line (insect)	Sf9	Novagen	Cat. No. 71104	

Chemical compound, drug	L-glutamate	Sigma Aldrich	Cat. No. 49621	
Software, algorithm	Pulse	HEKA	https://scicrunch.org/resolver/SCR_018399	https://www.heka.com/
Software, algorithm	OriginPro 2020	OriginLab	RRID:SCR_014212	https://www.originlab.com/2020
Software, algorithm	cryoSPARC 4.	doi:10.1038/nmeth.4169.	RRID:SCR_016501	https://cryosparc.com/
Software, algorithm	UCSF Chimera	doi:10.1002/jcc.20084	RRID:SCR_004097	http://plato.cgl.ucsf.edu/chimera/
Software, algorithm	UCSF ChimeraX	doi:10.1002/pro.3943	RRID:SCR_015872	https://www.cgl.ucsf.edu/chimerax/
Software, algorithm	HOLE	doi:10.1016/s0263-7855(97)00009-x		http://www.holeprogram.org
Software, algorithm	COOT 0.9x	doi:10.1107/S0907444910007493	RRID:SCR_014222	http://www2.mrc-lmb.cam.ac.uk/personal/pemsley/coot/
Software, algorithm	Phenix 1.20	doi:10.1107/S2059798319011471	RRID:SCR_014224	https://www.phenix-online.org/

Software, algorithm	MolProbity	doi:10.1107/S0907444909042073	RRID:SCR_014226	http://molprobity.biochem.duke.edu
Software, algorithm	deepEMhancer	https://doi.org/10.1038/s42003-021-02399-1		https://github.com/rsanchezgarc/deepEMhancer

Acknowledgment

We thank Dr. Matthias Buck (CWRU) for discussion and critical feedback on the manuscript. We thank Drs. Kunpeng Li and Kyle Whiddon at Case Western Reserve University (CWRU) for their cryo-EM technical support, and the High-Performance Computing Resource CWRU Core facility for their computational support. We are grateful to Drs. Corey Smith and Shyue-An Chan at CWRU for their technical assistance with patch-clamp electrophysiology. We thank Ashlee Hoffman (CWRU) for proofreading the manuscript. This work was funded by the National Institute of Health (1R35GM147266-01 to N.T.), Whitehall Foundation (N.T.), and American Heart Association https://doi.org/10.58275/AHA.23POST1019193.pc.gr.174253 (S.G.).

Author contributions

C.Z., S.G., and N.T. conceived the project and designed the experimental procedures. C.Z. designed the constructs, conduced the protein expression and purification, and performed the EM data collection and data analysis. S.G. conducted electrophysiological recordings. C.Z., S.G., and N.T. wrote the manuscript. All authors reviewed the final draft.

Competing interests

The authors declare no competing interests.

683 **Data and materials availability**

684 Cryo-EM density maps and atomic coordinates for GluK2-apo and GluK2-domoate were

685 deposited in the electron microscopy data bank under the accession codes listed below.

Name	Accession code
GluK2 ATD in the non-desensitized state	PDB 9MZQ, EMD-48765
GluK2 LBD-TMD bound to BPAM344 and glutamate in the non-desensitized state	PDB 9MZR, EMD-48766
GluK2 bound to BPAM344 and glutamate in the non-desensitized state, composite map	PDB 9MZS, EMD-48767
GluK2 ATD in the shallow-desensitized state	PDB 9MZN, EMD-48762
GluK2 LBD-TMD in the shallow-desensitized state	PDB 9MZO, EMD-48763
GluK2 in the shallow-desensitized state, consensus map	PDB 9MZM, EMD-48761
GluK2 in the shallow-desensitized state, composite map	PDB 9MZP, EMD-48764
GluK2 in the intermediate state	PDB 9MZL, EMD-48760
GluK2 in the deep-desensitized state 1	PDB 9MZK, EMD-48759
GluK2 in the deep-desensitized state 2	PDB 9MZJ, EMD-48758
GluK2 in the deep-desensitized state 3	PDB 9MZI, EMD-48757

Figure legends

Figure 1. Functional and structural characterization of GluK2 K676C/N802C.

(A) Representative whole-cell patch-clamp trace showing response of GluK2 WT or GluK2

K676C/N802C to a 20-second application of 1 mM glutamate in the absence and presence of

BPAM344 (BPAM). Arrows indicate the maximum peak current (I_{peak}) and the steady-state current

(I$_{ss}$). **(B)** Quantification of percent desensitization for currents evoked by 1 mM glutamate. **(C)**

Potentiation of the peak current amplitudes (I$_{peak}$) of GluK2 WT (left), the peak current amplitudes

(I$_{peak}$) of K676C/N802C (middle), and the steady-state current amplitude (I$_{ss}$) of GluK2

K676C/N802C measured in the presence of BPAM344 compared to currents in the absence of

BPAM344. **(D)** Cryo-EM map and model of GluK2 K676C/N802C in complex with glutamate and

BPAM344 in a non-desensitized/non-active state. **(E)** Chemical structure of BPAM344. **(F)**

Structure of the LBD layer viewed parallel to the membrane, with BPAM344 and glutamate shown

as space-filling models, colored pink and cyan, respectively. **(G)** Close-up view of the BPAM344

binding site at the AD subunit interface, indicating by a black dotted box in panel F. **(H)**

Comparison of the apo GluK2 LBD (PDB code 9CAZ) and the glutamate-bound GluK2 LBD in the

non-desensitized LBD conformation. The LBD is a bi-lobed structure, consisting of D1 and D2

lobes, with glutamate binding in the cleft between them. The D1 lobe is superimposed, and the

rotation angles to align the D2 residues were calculated. The glutamate-bound bi-lobe is closed

20° than the apo state LBD conformation. **(I)** Cryo-EM density for bound glutamate and the

surrounding residues in the agonist binding pocket. **(J)** The cryo-EM densities for disulfide bonds

between K676C and N802C in AB and CD subunits. The crosslinks in the LBD layers are indicated

by the red dotted box in the panel F.

Figure 2. Structural mechanism of GluK2 upon activation and desensitization.

(A) Comparison of Cryo-EM reconstructions and models of BPAM344- and glutamate-bound

GluK2 K676C/N802C in a non-desensitized conformation, and BPAM344-, glutamate-, and

Concanavalin A-bound GluK2 in an open (active) state (PDB code 9B36). The distances between

S700 in the AC subunits and E788 in the BD subunits at the LBD dimer-dimer interface are colored

red and blue, respectively. **(B)** Structural comparison of LBD dimers in the non-desensitized and

open GluK2, viewed perpendicular to the membrane. The locations of the centers of mass (COM)

of D1 and D2 are indicated by blue and green spheres, respectively, with the COM distances

between D1 and D2 of subunit AD LBDs (shown as circles) displayed below. Arrows indicate LBD

bi-lobe closure, with the D2 lobe moving closer to D1. Distances between the Cα atoms of P773

(D1-D1 distance) and S670 (D2-D2 distance) in the AD subunits are also indicated. **(C)**

Comparison of LBD-TM3 linkers and TM3 formation in BPAM344-bound, agonist-unbound GluK2

(PDB code 8FWS)[32] and non-desensitized GluK2, showing the subunit BD pair (left) and the

subunit AC pair (right). S670 is shown as spheres in corresponding colors for the non-

desensitized state and in gray for BPAM344-bound GluK2. The cross-dimer distances between

the Cα atoms of S670 (in the BD and AC subunit pairs) are indicated above. Residues forming

the ion channel pore are displayed as sticks. The locations of the TM3 gating hinge at A656

(subunit BD) and E662 (subunit AC) in the non-desensitized conformation are highlighted in red.

The dotted line indicates the positions of pore-lining residues: (1) M664, (2) T660, (3) A656, and

(4) T652. (2') E662, which serves as the gating hinge, is also highlighted in this panel. Helices

surrounding T664 and M664 in subunit BD, as well as helices around M664 in subunit AC, are

unwound in the non-desensitized conformation. As a result, the ion channel pore above the gating

region in the non-desensitized conformation is wider compared to that in the BPAM344-bound

GluK2 structure, with black allows indicating the conformational changes. **(D)** Comparison of LBD-

TM3 linkers and TM3 formation in non-desensitized GluK2 and BPAM344-, ConA- and glutamate

bound open GluK2 (PDB code 9B36)[27], showing the subunit BD pair (left) and subunit AC pair

(right). The cross-dimer distances between the Cα atoms of S670 are indicated above. Residues

forming the ion channel pore are displayed as sticks. The locations of the gating hinge at L655 in

A-D subunits of the open GluK2 (PDB 9B36) structure are highlighted in gray, for comparison with

the gating hinge in the non-desensitized conformation shown in panel C. The dotted line indicates

the positions of pore-lining residues: (1) M664, (2) T660, (3) A656, and (4) T652, as well as (4')

L655, which forms the gating hinge. Yellow, green, blue, and pink arrows indicate the

rearrangement of TM3 in each subunit during the transition from the non-desensitized to the open

conformation. **(E)** Pore profile in the non-desensitized and open (PBD 9B36) structures. Pore-

delineating dots are colored according to pore radius: red for regions with a radius less than 1.1

Å and gray for regions with a radius greater than 1.1 Å, which would allow passage of a

dehydrated ion (1.1 Å for calcium). **(F)** The pore radius for the non-desensitized (red) and open

(PDB 9B36, gray) conformations, calculated using HOLE.

Figure 3. Structures of GluK2 in desensitized states.

(A) Three-dimensional (3D) cryo-EM reconstructions of glutamate-bound GluK2 K676C/N802C

in desensitized conformations. **(B)** Top views of glutamate-bound GluK2 K676C/N802C, focusing

on the LBD layers. The in-plane rotation angles of the LBDs are indicated. Disulfide bonds formed

between the two LBD dimers are highlighted in red dotted boxes. **(C)** Close-up view of the disulfide

bonds between K676C and N802C at the CD subunit interface (left) and the AC subunit interface

(right) in the shallow-desensitized conformation. **(D)** Top view of the LBD layer in the deep-

desensitized conformation, highlighting helices E and G, which form the desensitization ring. The

view is parallel to the membrane, with coloring consistent with panel A. **(E)** The desensitization

ring formed in three deep-desensitized conformations, viewed perpendicular to the membrane.

The in-plane rotation of the LBD shown in panel B shifts the LBD-TM3 linker, positioning it above

the ion channel pore formed by the TM3 helices.

Figure 4. GluK2 stabilized in a shallow-desensitized-like conformations resembbles the

desensitized conformation of AMPARs.

(A, B) Structural comparison between the non-desensitized and shallow-desensitized

conformations, viewed parallel (A) and perpendicular (B) to the membrane. Black arrows indicate

the degree of LBD domain rotation. **(C, D)** Structural comparison of LBD layers in GluK2

K676C/N802C in the shallow-desensitized state and GluA2 AMPAR in the desensitized state

(PDB code 7RYZ)[64], viewed parallel (C) and perpendicular (D) to the membrane. Distances

between S700 in the AC subunits and E788 in the BD subunits at the LBD dimer-dimer interface

are shown in red and blue, respectively. The locations of the centers of mass (COM) of D1 and

D2 are indicated by blue and green spheres, respectively, with the COM distances between D1

and D2 of subunit AD LBDs (shown as circles) displayed below. Arrows indicate D1 lobe rotations

compared to the non-desensitized (KAR) or apo (AMPAR) states. Distances between the Cα

atoms of P773 (D1-D1 distance) and S670 (D2-D2 distance) in the AD subunits are also indicated.

(E) Comparison of LBD-TM3 linkers and TM3 helices in non-desensitized and shallow-

desensitized conformations, showing the subunit BD pair (left) and subunit AC pair (right). S670

is shown as spheres in corresponding colors for the shallow-desensitized conformation and in

gray for the non-desensitized conformation. The cross-dimer distances between the Cα atoms of

S670 are indicated. Residues forming the ion channel pore are displayed as sticks. The locations

of the TM3 gating hinge at T660 (subunit BD) and M664 (subunit AC) in the shallow-desensitized

conformation are highlighted in red. The dotted line indicates the positions of pore-lining residues:

(1) M664, (2) T660, (3) A656, and (4) T652. The shallow-desensitized conformation contains an

additional half-turn of helices at T660 compared to the non-desensitized conformation. **(F)**

Comparison of LBD-TM3 linkers and TM3 formation in shallow- and deep-desensitized GluK2,

showing the subunit BD pair (left) and the subunit AC pair (right). S670 is represented as spheres

in corresponding colors for the deep-desensitized state and in gray for the shallow-desensitized

GluK2. In the deep-desensitized conformation, the ion channel is sealed at M664, completely

closing the ion channel pore, in contrast to the shallow-desensitized conformation, which has a

kink at T660 in the BD subunits.

Figure 5. Ion channel pore of desensitized GluK2 KAR.

(A) Permeation pathway dictating pore diameter of GluK2 K676C/N802C in desensitized states.

Pore-delineating dots are colored according to pore radius: red for regions with a radius less than

1.1 Å and gray for regions with a radius greater than 1.1 Å. **(B)** Permeation pathway determining

the pore diameter of the GluA2 AMPAR in a desensitized state (PDB code 7RYZ)[64], using the

same color code as (A). **(C–D)** Pore profile traces comparing GluK2 K676C/N802C KAR.

Comparison between the shallow-desensitized conformation and the desensitized GluA2 AMPAR

(PDB: 7RYZ) (C); and comparison among three deep-desensitized states (D).

Figure 6. Functional characterization of GluK2 K676C/N802C receptor.

(A) Representative whole-cell patch-clamp recordings of GluK2 WT (left) and GluK2

K676C/N802C (right) in response to varying concentrations of glutamate (0.001–50 mM). The

black bar at the top of the recording represents the glutamate application. The cells were held at

-70 mV. **(B)** Quantification of current density (pA/pF) at the peak current for GluK2 WT (gray) and

K676C/N802C mutant (blue) activated by 10 mM glutamate. No significant differences were

observed (NS, $p > 0.05$). **(C)** Glutamate dose-response curves for GluK2 WT (n=12) and

K676C/N802C (n=8), normalized to the maximal response at 50 mM glutamate. Glutamate was

applied for 1 s, and the peak current was measured at each concentration. All response were

recorded at each concentrations using a single cell. **(D)** EC_{50} values for glutamate activation of

GluK2 WT (gray) and K676C/N802C (blue). No significant differences were detected (NS, $p >$

0.05). **(E)** Ratio of steady-state to peak current ($I_{steady-state}$ / I_{peak}) at various glutamate

concentrations. The values of the steady-state current were taken at the end of the activation. **(F)**

Representative current traces of GluK2 WT (black) and GluK2 K676C/N802C (blue) during a two-

pulse protocol, showing the interval between glutamate exposures ranging from 50 ms to 20 s,

highlighting altered desensitization kinetics in the mutant. **(G)** Recovery from desensitization at

increasing inter-sweep intervals, fitted with one-term exponential functions for GluK2 WT

(R²=0.99) and two-term exponential functions for GluK2 K676C/N802C (R²=0.99). GluK2

K676C/N802C exhibits a faster recovery than GluK2 WT. **(H)** Time constant (τ) of recovery from

desensitization, showing a significant reduction in τ for the K676C/N802C mutant ($p < 0.0001$,

two-sided two-sample t-test). The data for the WT was obtained from the fit in panel G, and for

K676C/N802C, a mean τ was calculated using $\tau_{mean} = ((A1\ \tau1 + A2\ \tau2) / (A1 + A2))$. Biological

independent measurements: n=9 for GluK2 WT and n=9 for GluK2 K676C/N802C.

Error bars represent Standard Deviation (SD). Statistical significance is indicated as NS (not

significant) or with p-values where appropriate.

Figure 7. Cysteine crosslinking modulates the kinetics of GluK2 K676C/N802C

(A) Representative whole-cell patch-clamp recordings of GluK2 WT (left) and GluK2

K676C/N802C (right) in the presence of 10 mM glutamate before (black) and after treatment with

5 mM DTT for 7 minutes (red). **(B)** Quantification of the I_{ss}/I_{peak} ratio in GluK2 K676C/N802C before

(blue) and after (red) DTT treatment, showing a significant reduction in steady-state current post-

treatment ($p < 0.0001$, two-sided two-sample t-test). **(C)** Recovery from desensitization at

increasing inter-sweep intervals after treatment with 5 mM DTT. The GluK2 WT recovery curve

from Figure 6, panel C is shown in black for comparison. The recovery kinetics of GluK2

K676C/N802C post-DTT treatment were fitted with a one-term exponential function ($R^2 = 0.98$),

showing a recovery profile similar to WT. **(D)** Time constant (τ) of recovery from desensitization,

showing that DTT treatment restores the recovery kinetics of GluK2 K676C/N802C to a level

comparable to WT (NS, $p > 0.05$). **(E)** Representative whole-cell responses to 10 mM glutamate

(200 ms, -70 mV; black bar) from HEK293T cells expressing GluK2 WT (left, average current in

black, 60 individual responses in gray) and GluK2 K676C/N802C (right, average current in blue,

100 individual responses in gray). Inset: Current-variance relationship, with WT data fitting a

parabolic function (dotted line), whereas the K676C/N802C mutant does not follow this trend. **(F)**

Whole-cell recordings of GluK2 K676C/N802C in response to 10 mM glutamate in the absence

(blue) and presence (green) of 500 µM BPAM344. Right panel: Representative noise traces at

the end of glutamate application (~10 s) under both conditions, indicated by blue and green boxes.

Background noise (baseline) was taken from the current recording before glutamate application,

as indicated by the gray box. **(G)** Variance-mean current relationship for individual cells in the

presence of glutamate alone or with BPAM344, indicating an enhanced channel variance-current

relationship upon BPAM344 application. **(H)** Unitary conductance determined by variance-mean

analysis, showing no significant difference between two conditions with and without BPAM344

(NS, $p > 0.05$). (I) Estimated number of open channels, showing a significant increase upon

BPAM344 application ($p = 0.0025$, two-sided paired t-test). Biologically independent

measurements: $n = 6$. Data for glutamate and BPAM344 conditions were obtained from the same

cell.

Error bars represent standard deviation (SD). Statistical significance is indicated as NS (not

significant) or with p-values where appropriate.

Figure 8. Schematic representation of conformational rearrangements during activation

and desensitization. The proposed model illustrates how conformational changes in the LBD

layers during desensitization regulate ion channel permeation. Wild-type GluK2 typically

transitions into a deeply desensitized state, adopting a stabilized, classically four-fold symmetrical

conformation. By contrast, receptors that stabilize in a two-fold symmetrical, desensitized

AMPAR-like conformation, maintained by engineered disulfide bonds, remain conductive,

although their conductivity is lower than that of receptors in the open state. The position of the

gating kink differs among the non-desensitized, open, and shallow-desensitized conformations.

Additionally, the ATD layers elevate and rotate in both the non-desensitized (non-active) and

open-state GluK2 conformations.

Figure S1. Design of the double cysteine crosslink mutant of GluK2.

(A) Schematic illustration of the KAR subunit topology. Each KAR subunit is composed of four domains: ATD, LBD,TMD, and CTD. The LBD contains two segments termed D1 (upper lobe) and D2 (lower lobe). Agonists bind to the interface between the D1 and D2 lobes of the LBD. **(B)** Homology model of the active GluK2 KAR, generated based on the active GluA2 AMPAR (PDB code 5WEO) as a template. **(C)** The inter-protomer interface between subunit A and subunit B in the homology model. **(D)** The distance between residues K676 and N802 in the BPAM344-, concanavalin A-, and glutamate-bound open/active GluK2 (PDB code 9B36)[27], BPAM344-bound (without orthosteric ligands) inactive GluK2 (PDB code 8FWS)[32], and glutamate-bound deep desensitized GluK2 (PDB code 9B38)[27] structures. **(E, F)** Two designed disulfide crosslinks at the inter-protomer interface, shown in the homology model and the schematic figure.

Figure S2. Channel properties of the single cysteine mutant and protein preparation.

(A) Representative electrophysiological traces from HEK293T cells expressing GluK2 WT, GluK2 K676C, and GluK2 N802C, activated by 10 mM glutamate. **(B)** Size exclusion chromatography profile of the GluK2 N802C/K676C mutant eluted from the Superose 6 column chromatography. The fractions used for the cryo-EM studies are indicated by the black dotted line. **(C)** SDS-PAGE analysis of the purified GluK2 N802C/K676C mutant, with the black line indicating the full-length GluK2 protein band.

Figure S3. Cryo-EM data processing workflow for structures of the GluK2 K676C/N802C mutant in the presence of glutamate and BPAM344 (a), and glutamate alone (b)

Figure S4. Cryo-EM analysis of GluK2 K676C/N802C.

(A, D) Local resolution maps, **(B, E)** Fourier shell correlation (FSC) curves. The threshold was set

at 0.143 for unmasked and masked curves, and a 0.5 threshold was used for the map-to-model

curve. **(C, F)** Angular distribution of particles calculated using the 3D refinement reconstruction

algorithm in cryoSPARC.

Figure S5. Comparison of ATD-LBD layers.

(A) GluK2 structures, aligned parallel to the membrane. The distances between the center-of-

mass (COM, black circles) of the ATD and LBD layers are indicated. **(B)** Superimposition of the

ATD layer in the non-desensitized, intermediate, and desensitized GluK2 structures. The color

codes match those in panel A. **(C)** ATD-LBD rotation. The LBD layers in the BPAM344- and

glutamate-bound non-desensitized, and the BPAM344-, ConA-, and glutamate-bound open

GluK2 conformations, are rotated counterclockwise when the ATD layers are superimposed.

Figure S6. Models and cryo-EM densities of the transmembrane domain.

TM3 helices in non-desensitized, shallow-desensitized, intermediate, and deep-desensitized

structures, with the protein structural model shown as ribbons and sticks. Cryo-EM densities are

displayed in transparent gray.

915 **References**
916

1   Hansen, K. B. *et al.* Structure, Function, and Pharmacology of Glutamate Receptor Ion
Channels.   *Pharmacol   Rev   **73**,   298-487   (2021).
https://doi.org:10.1124/pharmrev.120.000131

2   Lerma, J. Roles and rules of kainate receptors in synaptic transmission. *Nat Rev Neurosci*
4, 481-495 (2003). https://doi.org:10.1038/nrn1118

3   Chittajallu, R. *et al.* Regulation of glutamate release by presynaptic kainate receptors in
the hippocampus. *Nature* **379**, 78-81 (1996). https://doi.org:10.1038/379078a0

4   Rodríguez-Moreno, A., Herreras, O. & Lerma, J. Kainate receptors presynaptically
downregulate GABAergic inhibition in the rat hippocampus. *Neuron* **19**, 893-901 (1997).
https://doi.org:10.1016/s0896-6273(00)80970-8

5   Cossart, R. *et al.* Presynaptic kainate receptors that enhance the release of GABA on CA1
hippocampal interneurons. *Neuron* **29**, 497-508 (2001). https://doi.org:10.1016/s0896-
6273(01)00221-5

6   Polenghi, A. *et al.* Kainate Receptor Activation Shapes Short-Term Synaptic Plasticity by
Controlling Receptor Lateral Mobility at Glutamatergic Synapses. *Cell Rep* **31**, 107735
(2020). https://doi.org:10.1016/j.celrep.2020.107735

7   Ramos, C. *et al.* Activation of Extrasynaptic Kainate Receptors Drives Hilar Mossy Cell
Activity. *J Neurosci* **42**, 2872-2884 (2022). https://doi.org:10.1523/jneurosci.0922-21.2022

8   Lerma, J. & Marques, J. M. Kainate receptors in health and disease. *Neuron* **80**, 292-311
(2013). https://doi.org:10.1016/j.neuron.2013.09.045

9   Aller, M. I., Pecoraro, V., Paternain, A. V., Canals, S. & Lerma, J. Increased Dosage of
High-Affinity Kainate Receptor Gene grik4 Alters Synaptic Transmission and Reproduces
Autism   Spectrum   Disorders   Features.   *J   Neurosci*   **35**,   13619-13628   (2015).
https://doi.org:10.1523/jneurosci.2217-15.2015

10   Begni, S. *et al.* Association between the ionotropic glutamate receptor kainate 3 (GRIK3)
ser310ala   polymorphism   and   schizophrenia.   *Mol   Psychiatry*   **7**,   416-418   (2002).
https://doi.org:10.1038/sj.mp.4000987

11   Shaltiel, G. *et al.* Evidence for the involvement of the kainate receptor subunit GluR6
(GRIK2) in mediating behavioral displays related to behavioral symptoms of mania. *Mol*
Psychiatry **13**, 858-872 (2008). https://doi.org:10.1038/mp.2008.20

12   Shibata, H. *et al.* Association study of polymorphisms in the GluR7, KA1 and KA2 kainate
receptor genes (GRIK3, GRIK4, GRIK5) with schizophrenia. *Psychiatry Res* **141**, 39-51
(2006). https://doi.org:10.1016/j.psychres.2005.07.015

13   Diguet, E. *et al.* Experimental basis for the putative role of GluR6/kainate glutamate
receptor subunit in Huntington's disease natural history. *Neurobiol Dis* **15**, 667-675 (2004).
https://doi.org:10.1016/j.nbd.2003.12.010

14   Schiffer, H. H. & Heinemann, S. F. Association of the human kainate receptor GluR7 gene
(GRIK3) with recurrent major depressive disorder. *Am J Med Genet B Neuropsychiatr*
Genet **144b**, 20-26 (2007). https://doi.org:10.1002/ajmg.b.30374

15   Pickard, B. S. *et al.* A common variant in the 3'UTR of the GRIK4 glutamate receptor gene
affects transcript abundance and protects against bipolar disorder. *Proc Natl Acad Sci U*
S A **105**, 14940-14945 (2008). https://doi.org:10.1073/pnas.0800643105

16   Catches, J. S., Xu, J. & Contractor, A. Genetic ablation of the GluK4 kainate receptor
subunit causes anxiolytic and antidepressant-like behavior in mice. *Behav Brain Res* **228**,
406-414 (2012). https://doi.org:10.1016/j.bbr.2011.12.026

17   Lowry, E. R., Kruyer, A., Norris, E. H., Cederroth, C. R. & Strickland, S. The GluK4 kainate
receptor subunit regulates memory, mood, and excitotoxic neurodegeneration.
Neuroscience **235**, 215-225 (2013). https://doi.org:10.1016/j.neuroscience.2013.01.029

18   Matute, C. Therapeutic potential of kainate receptors. *CNS Neurosci Ther* **17**, 661-669
(2011). https://doi.org:10.1111/j.1755-5949.2010.00204.x

19   Jane, D. E., Lodge, D. & Collingridge, G. L. Kainate receptors: pharmacology, function
and   therapeutic   potential.   *Neuropharmacology*   **56**,   90-113   (2009).
https://doi.org:10.1016/j.neuropharm.2008.08.023

20   Negrete-Díaz, J. V., Falcón-Moya, R. & Rodríguez-Moreno, A. Kainate receptors: from
synaptic   activity   to   disease.   *Febs   j*   **289**,   5074-5088   (2022).
https://doi.org:10.1111/febs.16081

21   Swanson, G. T. Targeting AMPA and kainate receptors in neurological disease: therapies
on   the   horizon?   *Neuropsychopharmacology*   **34**,   249-250   (2009).
https://doi.org:10.1038/npp.2008.158

22   Cui, C. & Mayer, M. L. Heteromeric kainate receptors formed by the coassembly of GluR5,
GluR6,   and   GluR7.   *J   Neurosci*   **19**,   8281-8291   (1999).
https://doi.org:10.1523/jneurosci.19-19-08281.1999

23   Mulle, C. *et al.* Subunit composition of kainate receptors in hippocampal interneurons.
Neuron **28**, 475-484 (2000). https://doi.org:10.1016/s0896-6273(00)00126-4

24   Reiner, A., Arant, R. J. & Isacoff, E. Y. Assembly stoichiometry of the GluK2/GluK5 kainate
receptor   complex.   *Cell   Rep*   **1**,   234-240   (2012).
https://doi.org:10.1016/j.celrep.2012.01.003

25   Selvakumar, P. *et al.* Structural and compositional diversity in the kainate receptor family.
Cell Rep **37**, 109891 (2021). https://doi.org:10.1016/j.celrep.2021.109891

26   Mayer, M. L. Structural biology of kainate receptors. *Neuropharmacology* **190**, 108511
(2021). https://doi.org:10.1016/j.neuropharm.2021.108511

27   Gangwar, S. P. *et al.* Kainate receptor channel opening and gating mechanism. *Nature*
(2024). https://doi.org:10.1038/s41586-024-07475-0

28   Meyerson, J. R. *et al.* Structural basis of kainate subtype glutamate receptor
desensitization. *Nature* **537**, 567-571 (2016). https://doi.org:10.1038/nature19352

29   Khanra, N., Brown, P. M., Perozzo, A. M., Bowie, D. & Meyerson, J. R. Architecture and
structural dynamics of the heteromeric GluK2/K5 kainate receptor. *Elife* **10** (2021).
https://doi.org:10.7554/eLife.66097

30   He, L. *et al.* Kainate receptor modulation by NETO2. *Nature* (2021).
https://doi.org:10.1038/s41586-021-03936-y

31   Kumari, J., Vinnakota, R. & Kumar, J. Structural and Functional Insights into GluK3-
kainate Receptor Desensitization and Recovery. *Sci Rep* **9**, 10254 (2019).
https://doi.org:10.1038/s41598-019-46770-z

32   Gangwar, S. P., Yen, L. Y., Yelshanskaya, M. V. & Sobolevsky, A. I. Positive and negative
allosteric modulation of GluK2 kainate receptors by BPAM344 and antiepileptic
perampanel. *Cell Rep* **42**, 112124 (2023). https://doi.org:10.1016/j.celrep.2023.112124

33   Kumari, J. *et al.* Structural dynamics of the GluK3-kainate receptor neurotransmitter
binding domains revealed by cryo-EM. *Int J Biol Macromol* **149**, 1051-1058 (2020).
https://doi.org:10.1016/j.ijbiomac.2020.01.282

34   Gangwar, S. P. *et al.* Trapping of spermine, Kukoamine A, and polyamine toxin blockers
in GluK2 kainate receptor channels. *Nat Commun* **15**, 10257 (2024).
https://doi.org:10.1038/s41467-024-54538-x

35   Paternain, A. V., Rodríguez-Moreno, A., Villarroel, A. & Lerma, J. Activation and
desensitization properties of native and recombinant kainate receptors.
Neuropharmacology **37**, 1249-1259 (1998). https://doi.org:10.1016/s0028-
3908(98)00098-7

36   Weston, M. C., Schuck, P., Ghosal, A., Rosenmund, C. & Mayer, M. L. Conformational
restriction blocks glutamate receptor desensitization. *Nat Struct Mol Biol* **13**, 1120-1127
(2006). https://doi.org:10.1038/nsmb1178

37   Jones, M. V. & Westbrook, G. L. The impact of receptor desensitization on fast synaptic
transmission. *Trends Neurosci* **19**, 96-101 (1996). https://doi.org:10.1016/s0166-
2236(96)80037-3

38   Meyerson, J. R. *et al.* Structural mechanism of glutamate receptor activation and
desensitization. *Nature* **514**, 328-334 (2014). https://doi.org:10.1038/nature13603

39   Dhingra, S., Chopade, P. M., Vinnakota, R. & Kumar, J. Functional implications of the
exon 9 splice insert in GluK1 kainate receptors. *Elife* **12** (2024).
https://doi.org:10.7554/eLife.89755

40   Schauder, D. M. *et al.* Glutamate receptor desensitization is mediated by changes in
quaternary structure of the ligand binding domain. *Proc Natl Acad Sci U S A* **110**, 5921-
5926 (2013). https://doi.org:10.1073/pnas.1217549110

41 Coombs, I. D. *et al.* Homomeric GluA2(R) AMPA receptors can conduct when
desensitized. *Nat Commun* **10**, 4312 (2019). https://doi.org:10.1038/s41467-019-12280-9

42 Twomey, E. C., Yelshanskaya, M. V., Grassucci, R. A., Frank, J. & Sobolevsky, A. I.
Channel opening and gating mechanism in AMPA-subtype glutamate receptors. *Nature*
549, 60-65 (2017). https://doi.org:10.1038/nature23479

43 Yelshanskaya, M. V., Patel, D. S., Kottke, C. M., Kurnikova, M. G. & Sobolevsky, A. I.
Opening of glutamate receptor channel to subconductance levels. *Nature* **605**, 172-178
(2022). https://doi.org:10.1038/s41586-022-04637-w

44 Chen, S. *et al.* Activation and Desensitization Mechanism of AMPA Receptor-TARP
Complex by Cryo-EM. *Cell* **170**, 1234-1246.e1214 (2017).
https://doi.org:10.1016/j.cell.2017.07.045

45 Waterhouse, A. *et al.* SWISS-MODEL: homology modelling of protein structures and
complexes. *Nucleic Acids Res* **46**, W296-w303 (2018). https://doi.org:10.1093/nar/gky427

46 Bienert, S. *et al.* The SWISS-MODEL Repository-new features and functionality. *Nucleic*
Acids Res **45**, D313-d319 (2017). https://doi.org:10.1093/nar/gkw1132

47 Guex, N., Peitsch, M. C. & Schwede, T. Automated comparative protein structure
modeling with SWISS-MODEL and Swiss-PdbViewer: a historical perspective.
Electrophoresis **30 Suppl 1**, S162-173 (2009). https://doi.org:10.1002/elps.200900140

48 Studer, G. *et al.* QMEANDisCo-distance constraints applied on model quality estimation.
Bioinformatics **36**, 1765-1771 (2020). https://doi.org:10.1093/bioinformatics/btz828

49 Bertoni, M., Kiefer, F., Biasini, M., Bordoli, L. & Schwede, T. Modeling protein quaternary
structure of homo- and hetero-oligomers beyond binary interactions by homology. *Sci Rep*
7, 10480 (2017). https://doi.org:10.1038/s41598-017-09654-8

50 Esmenjaud, J. B. *et al.* An inter-dimer allosteric switch controls NMDA receptor activity.
Embo j **38** (2019). https://doi.org:10.15252/embj.201899894

51 Salazar, H., Eibl, C., Chebli, M. & Plested, A. Mechanism of partial agonism in AMPA-type
glutamate receptors. *Nat Commun* **8**, 14327 (2017).
https://doi.org:10.1038/ncomms14327

52 Baranovic, J. & Plested, A. J. Auxiliary subunits keep AMPA receptors compact during
activation and desensitization. *Elife* **7** (2018). https://doi.org:10.7554/eLife.40548

53 Plested, A. J. & Mayer, M. L. AMPA receptor ligand binding domain mobility revealed by
functional cross linking. *J Neurosci* **29**, 11912-11923 (2009).
https://doi.org:10.1523/jneurosci.2971-09.2009

54 Daniels, B. A., Andrews, E. D., Aurousseau, M. R., Accardi, M. V. & Bowie, D. Crosslinking
the ligand-binding domain dimer interface locks kainate receptors out of the main open
state. *J Physiol* **591**, 3873-3885 (2013). https://doi.org:10.1113/jphysiol.2013.253666

55 Careaga, C. L. & Falke, J. J. Thermal motions of surface alpha-helices in the D-galactose
chemosensory receptor. Detection by disulfide trapping. *J Mol Biol* **226**, 1219-1235 (1992).
https://doi.org:10.1016/0022-2836(92)91063-u

56 Larsen, A. P. *et al.* Identification and Structure-Function Study of Positive Allosteric
Modulators of Kainate Receptors. *Mol Pharmacol* **91**, 576-585 (2017).
https://doi.org:10.1124/mol.116.107599

57 Bay, Y. *et al.* The positive allosteric modulator BPAM344 and L-glutamate introduce an
active-like structure of the ligand-binding domain of GluK2. *FEBS Lett* **598**, 743-757 (2024).
https://doi.org:10.1002/1873-3468.14824

58 Twomey, E. C. & Sobolevsky, A. I. Structural Mechanisms of Gating in Ionotropic
Glutamate Receptors. *Biochemistry* **57**, 267-276 (2018).
https://doi.org:10.1021/acs.biochem.7b00891

59 Zhou, C. & Tajima, N. Structural insights into NMDA receptor pharmacology. *Biochem Soc*
Trans **51**, 1713-1731 (2023). https://doi.org:10.1042/bst20230122

60 Klein, R. M. & Howe, J. R. Effects of the lurcher mutation on GluR1 desensitization and
activation kinetics. *J Neurosci* **24**, 4941-4951 (2004).
https://doi.org:10.1523/jneurosci.0660-04.2004

61 Wilding, T. J. & Huettner, J. E. Cadmium opens GluK2 kainate receptors with cysteine
substitutions at the M3 helix bundle crossing. *J Gen Physiol* **151**, 435-451 (2019).
https://doi.org:10.1085/jgp.201812234

62 Fleck, M. W., Cornell, E. & Mah, S. J. Amino-acid residues involved in glutamate receptor
6 kainate receptor gating and desensitization. *J Neurosci* **23**, 1219-1227 (2003).
https://doi.org:10.1523/jneurosci.23-04-01219.2003

63 Twomey, E. C., Yelshanskaya, M. V., Grassucci, R. A., Frank, J. & Sobolevsky, A. I.
Structural Bases of Desensitization in AMPA Receptor-Auxiliary Subunit Complexes.
Neuron **94**, 569-580.e565 (2017). https://doi.org:10.1016/j.neuron.2017.04.025

64 Klykov, O., Gangwar, S. P., Yelshanskaya, M. V., Yen, L. & Sobolevsky, A. I. Structure
and desensitization of AMPA receptor complexes with type II TARP γ5 and GSG1L. *Mol*
Cell **81**, 4771-4783.e4777 (2021). https://doi.org:10.1016/j.molcel.2021.09.030

65 Herguedas, B. *et al.* Mechanisms underlying TARP modulation of the GluA1/2-γ8 AMPA
receptor. *Nat Commun* **13**, 734 (2022). https://doi.org:10.1038/s41467-022-28404-7

66 Gangwar, S. P. *et al.* Modulation of GluA2-γ5 synaptic complex desensitization, polyamine
block and antiepileptic perampanel inhibition by auxiliary subunit cornichon-2. *Nat Struct*
Mol Biol **30**, 1481-1494 (2023). https://doi.org:10.1038/s41594-023-01080-x

67 Reiner, A. & Isacoff, E. Y. Tethered ligands reveal glutamate receptor desensitization
depends on subunit occupancy. *Nat Chem Biol* **10**, 273-280 (2014).
https://doi.org:10.1038/nchembio.1458

68 Fisher, M. T. & Fisher, J. L. Contributions of different kainate receptor subunits to the properties of recombinant homomeric and heteromeric receptors. *Neuroscience* **278**, 70-80 (2014). https://doi.org:10.1016/j.neuroscience.2014.08.009

69 Salazar, H., Mischke, S. & Plested, A. J. R. Measurements of the Timescale and Conformational Space of AMPA Receptor Desensitization. *Biophys J* **119**, 206-218 (2020). https://doi.org:10.1016/j.bpj.2020.05.029

70 Zhang, D. *et al.* Structural mobility tunes signalling of the GluA1 AMPA glutamate receptor. *Nature* **621**, 877-882 (2023). https://doi.org:10.1038/s41586-023-06528-0

71 Larsen, A. H., Perozzo, A. M., Biggin, P. C., Bowie, D. & Kastrup, J. S. Recovery from desensitization in GluA2 AMPA receptors is affected by a single mutation in the N-terminal domain interface. *J Biol Chem* **300**, 105717 (2024). https://doi.org:10.1016/j.jbc.2024.105717

72 Zhang, W. *et al.* A transmembrane accessory subunit that modulates kainate-type glutamate receptors. *Neuron* **61**, 385-396 (2009). https://doi.org:10.1016/j.neuron.2008.12.014

73 Swanson, G. T., Feldmeyer, D., Kaneda, M. & Cull-Candy, S. G. Effect of RNA editing and subunit co-assembly single-channel properties of recombinant kainate receptors. *J Physiol* **492 (Pt 1)**, 129-142 (1996). https://doi.org:10.1113/jphysiol.1996.sp021295

74 Aittoniemi, J., Jensen, M., Pan, A. C. & Shaw, D. E. Desensitization dynamics of the AMPA receptor. *Structure* **31**, 724-734.e723 (2023). https://doi.org:10.1016/j.str.2023.03.013

75 Carbone, A. L. & Plested, A. J. Coupled control of desensitization and gating by the ligand binding domain of glutamate receptors. *Neuron* **74**, 845-857 (2012). https://doi.org:10.1016/j.neuron.2012.04.020

76 Plested, A. J. & Mayer, M. L. Structure and mechanism of kainate receptor modulation by anions. *Neuron* **53**, 829-841 (2007). https://doi.org:10.1016/j.neuron.2007.02.025

77 Dürr, K. L. *et al.* Structure and dynamics of AMPA receptor GluA2 in resting, pre-open, and desensitized states. *Cell* **158**, 778-792 (2014). https://doi.org:10.1016/j.cell.2014.07.023

78 Ivica, J. *et al.* Proton-triggered rearrangement of the AMPA receptor N-terminal domains impacts receptor kinetics and synaptic localization. *Nat Struct Mol Biol* **31**, 1601-1613 (2024). https://doi.org:10.1038/s41594-024-01369-5

79 Revah, F. *et al.* Mutations in the channel domain alter desensitization of a neuronal nicotinic receptor. *Nature* **353**, 846-849 (1991). https://doi.org:10.1038/353846a0

80 Bowie, D. & Lange, G. D. Functional stoichiometry of glutamate receptor desensitization. *J Neurosci* **22**, 3392-3403 (2002). https://doi.org:10.1523/jneurosci.22-09-03392.2002

81 Dawe, G. B. *et al.* Defining the structural relationship between kainate-receptor deactivation and desensitization. *Nat Struct Mol Biol* **20**, 1054-1061 (2013). https://doi.org:10.1038/nsmb.2654

1138

1139

1140

1141

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7

Figure 8