TITLE: PCNA encircling primer/template junctions is eliminated by exchange of RPA for Rad51: Implications for the interplay between human DNA damage tolerance pathways

Jessica L. Norris[1], Lindsey O. Rogers[1], Grace Young[1], Kara G. Pytko[1], Rachel L. Dannenberg[1], Samuel Perreault[1], Vikas Kaushik[2], Edwin Antony[2], and Mark Hedglin[1,*]

From the [1]Department of Chemistry, The Pennsylvania State University, University Park, PA 16802

From the [2]The Saint Louis University School of Medicine, Department of Biochemistry and Molecular Biology, St. Louis MO, 63104

Running Title:

[*]To whom correspondence should be addressed: Mark Hedglin: Department of Chemistry, The Pennsylvania State University, University Park, PA 16802; muh218@psu.edu; Tel. (814) 863-1080

Keywords:

ABSTRACT

The DNA genome is constantly exposed to agents, such as ultraviolet radiation (UVR), that can alter or eliminate its coding properties through covalent modifications of the template bases. Many of these damaging modifications (i.e., lesions) persist into S-phase of the cell cycle where they may stall the canonical DNA replication machinery. In humans, these stalling events are circumvented by one of at least three interconnected DNA damage tolerance (DDT) pathways; translesion DNA synthesis (TLS), Template Switching (TS), and Homology-dependent Recombination (HDR). Currently, the functional interplay between these pathways is unclear, leaving wide gaps in our fundamental understanding of human DDT. To gain insights, we focus on the activation mechanisms of the DDT pathways. PCNA sliding clamps encircling primer/template (P/T) junctions of stalled replication sites are central to the activation of both TLS and TS whereas exchange of RPA for Rad51 filaments on the single strand DNA (ssDNA) sequences of stalled replication sites is central to HDR activation. Utilizing direct, ensemble FRET approaches developed by our lab, we independently monitor and directly compare PCNA occupancy and RPA/Rad51 exchange at P/T junctions under a variety of conditions that mimic *in vivo* scenarios. Collectively, the results reveal that assembly of stable Rad51 filaments at P/T junctions via RPA/Rad51 exchange causes complete and irreversible unloading of the resident PCNA, both in the presence and absence of an abundant PCNA-binding protein complex. Further investigations decipher the mechanism of RPA/Rad51 exchange-dependent unloading of PCNA. Collectively, these studies provide critical insights into the interplay between human DDT pathways and direction for future studies.

Introduction

Proper genetic inheritance requires that the double strand DNA (dsDNA) genome of each cell is accurately replicated for transfer to its progeny. The bulk of DNA replication is carried out by a canonical pathway that relies on the high-fidelity DNA polymerases (pols), ε and δ, and essential accessory factors such as the processivity sliding clamp ring, proliferating cell nuclear antigen (PCNA), the PCNA clamp loader, replication factor C (RFC), and the major single

strand DNA (ssDNA) binding protein complex, replication protein A (RPA) (**Figure 1A**). RFC
utilizes ATP to load PCNA onto primer/template (P/T) junctions such that the "front face" of
loaded PCNA is oriented towards the 3′ terminus of the primer strand from where DNA
synthesis initiates from (i.e., in the direction of DNA synthesis)[1-4]. Pols ε and δ are responsible
for the replication of leading and lagging strand ssDNA templates, respectively, and like many
DNA processing enzymes, anchor to the front face of loaded PCNA forming holoenzymes with
maximal efficiency. RPA directly engages P/T junctions and coats ssDNA templates as they are
exposed during replication, forming directional filaments that protect the underlying ssDNA
from being degraded or forming secondary structures that limit its accessibility[5-12]. The affinity
of human RPA for dsDNA is nearly 3 orders of magnitude weaker than its affinity for ssDNA [13-
15]. Hence, conversion of template ssDNA to dsDNA by DNA pols indirectly and rapidly releases
RPA into solution such that RPA filaments are short and transient under non-stressed
conditions[15-18].
The DNA genome is constantly exposed to agents, such as ultraviolet radiation (UVR), that
can alter or eliminate its coding properties through covalent modifications of the template bases.
Many of these damaging modifications (i.e., DNA lesions), such as the major photoproducts
generated from UVR exposure, cannot be accommodated by pols ε and δ and, consequently, stall
canonical replication, generating long, persistent RPA filaments that directly engage the afflicted
P/T junctions and the unreplicated ssDNA templates downstream. The resident PCNA are left
behind on the dsDNA regions upstream of the afflicted P/T junctions [19,20]. In humans, these
stalling events are circumvented by one of at least three non-canonical DNA replication

Figure 1. DNA damage and human DNA replication. Nascent DNA (primer) and parental DNA (template) strands
are blue and black, respectively. RPA subunits are color-coded and depicted to illustrate the OB-folds (A – E). The
"front" and "back" faces of PCNA are green and grey, respectively. For clarity, RFC is not shown. **A**) Canonical
DNA replication. Shown as an example is replication of lagging strand DNA template by pol δ. The ssDNA
template to be replicated is extended by an RPA filament in which each RPA engages the ssDNA in an orientation-
specific manner (as depicted). **B**) DNA damage tolerance. All pathways emanate from a site of stalled canonical
DNA replication (highlighted in yellow). The offending DNA lesion is indicated by an "X." Activation of a given
pathway is indicated by ➡.

pathways that bypass (i.e., "tolerate") the offending lesion, permitting canonical DNA replication
to resume downstream[21-23]. This process is referred to as DNA damage tolerance (DDT) and is
imperative for cell survival following mild exposure to DNA damaging agents. Previous studies
have identified the fundamental components of each DDT pathway and deciphered their basic
mechanisms and interdependencies (**Figure 1B**)[21-25].
Translesion DNA synthesis (TLS) relies on the attachment of single ubiquitin moieties to
K164 residue(s) on the "back face" of PCNA encircling sites of stalled canonical DNA
replication (i.e., stalled replication site). This posttranslational modification (PTM), referred to as
PCNA monoubiquitination, is catalyzed by the $Rad6(Rad18)_2$ complex and activates TLS by
recruiting essential TLS factors to stalled replication sites. TLS culminates with one or more
specialized DNA pols (i.e., TLS pols) utilizing monoubiquitinated PCNA to extend the aborted
primer strand directly opposite and beyond the offending lesion. TLS pols all contain expanded
active sites that accommodate an array of DNA lesions, albeit with lowered fidelities[23,24,26,27]. In
template switching (TS), the single ubiquitin moieties attached to the "back face" of PCNA by
the $Rad6(Rad18)_2$ complex are extended to a K63-linked polyubiquitin chain by the MMS2-
Ubc13 complex in conjunction with either HLTF or SHPRH. This PTM, referred to as PCNA
polyubiquitination, activates TS by recruiting the AH2/ZRANB3 DNA translocase, which
subsequently catalyzes replication fork reversal, displacing RPA. The resultant Holliday
junction-like structure permits an unidentified DNA pol to utilize the nascent DNA strand from
the undamaged, homologous sister chromatid as a template to extend the aborted primer[28-31]. In
homology-dependent recombination (HDR), RPA is exchanged for ATP-bound Rad51 (i.e.,
Rad51•ATP) on the unreplicated ssDNA templates of stalled replication sites. Rad51 is an ATP-
dependent DNA recombinase and activates HDR by catalyzing strand invasion in which the
aborted primer strand invades the undamaged, homologous sister chromatid at a complimentary
sequence. In this context, Rad51-catalyzed strand invasion results in a multi-branched, H-shaped
structure. An unidentified DNA pol then utilizes the nascent DNA strand from the undamaged,
homologous sister chromatid as a template to extend the aborted primer. Unlike TLS and TS,
HDR is independent of PCNA monoubiquitination and a recent study suggests PCNA does not
serve a functional role in Rad51-catalyzed strand invasion during HDR[19,21,22,32]. In both TS and
HDR pathways, the aborted primer strands are extended to a degree such that when they are
subsequently re-annealed to the damaged template strand (via reversal of the higher-order DNA
structures) their 3′ termini are placed downstream of the offending lesion.
All DDT pathways emanate from stalled replication sites, contribute to cell survival
following exposure to common DNA damaging agents such as UVR, and, if successfully
executed, yield the same results. Namely, faithful replication of a DNA lesion and the contiguous
DNA sequence and the resumption of canonical DNA replication downstream of the offending
lesion. Currently, the functional interplay between DDT pathways at stalled replication sites is
unclear, leaving wide gaps in our fundamental understanding of human DDT. To gain insights
we focus on the basic activation mechanisms of DDT pathways. PCNA encircling P/T junctions
of stalled replication sites is central to the activation of both TLS and TS, as described above and
depicted in **Figure 1B**. Multiple mechanisms collectively maintain loaded PCNA at/near these
sites. For example, "protein roadblocks" restrict diffusion of loaded PCNA away from stalled
replication sites[15,33]. Large DNA-binding protein complexes that rapidly and tightly engage
nascent DNA as it is generated during DNA replication restrict diffusion of the resident PCNA
along the dsDNA upstream of stalled replication sites[34-36]. RPA•ssDNA complexes immediately
adjacent to P/T junctions of stalled replication sites prevent diffusion of loaded PCNA onto and

along the unreplicated ssDNA templates[4,37]. Furthermore, these RPA•ssDNA complexes prohibit
RFC-catalyzed unloading of PCNA and permit RFC-catalyzed re-loading of PCNA back onto
stalled replication sites that are devoid of the sliding clamp[4,15,24]. Exchange of RPA for
Rad51•ATP filaments on the unreplicated ssDNA templates of stalled replication sites is central
to HDR activation (**Figure 1B**). In the present study, we utilize direct, ensemble FRET
approaches developed by our lab to investigate the interplay between RPA/Rad51 exchange and
PCNA occupancy at P/T junctions under a variety of conditions that mimic *in vivo* scenarios[15,38].
Collectively, the results from the present study provide critical insights into the functional
interplay between human DDT pathways and direction for future studies.

Materials and Methods

Oligonucleotides

Oligonucleotides were synthesized by Integrated DNA Technologies (Coralville, IA) and
purified via denaturing polyacrylamide gel electrophoresis. The concentrations of unlabeled
DNAs were determined from the absorbance at 260 nm using the provided extinction
coefficients. The concentrations of Cy5-labeled DNAs were determined from the extinction
coefficient at 650 nm for Cy5 ($\varepsilon_{650} = 250,000$ $M^{-1}cm^{-1}$). Concentrations of Cy3-labeled DNAs
were determined from the extinction coefficient at 550 nm for Cy3 ($\varepsilon_{550} = 136,000$ $M^{-1}cm^{-1}$).
For annealing two ssDNAs (as depicted in **Figure S1**), the primer and corresponding
complementary template strands were mixed in equimolar amounts in 1X Annealing Buffer (10
mM TrisHCl, pH 8.0, 100 mM NaCl, 1 mM EDTA), heated to 95 $^\circ$C for 5 minutes, and allowed
to slowly cool to room temperature.

Recombinant Human Proteins

Human RPA, Cy5-PCNA, exonuclease-deficient Pol δ (referred to herein as simply Pol δ)
and RFC were obtained as previously described[2,39-41]. The plasmid (pCH1/RAD51o) for the
expression of wild-type human Rad51 was a gift from Maria Spies (Addgene plasmid #102562;
http://n2t.net/addgene:102562; RRID:Addgene_102562) and human Rad51 was obtained as
previously described[41]. Human RPA containing a Cy5 label at residue 215 of the oligonucleotide
binding (OB) fold A of the RPA1 subunit (RPA-OBA-Cy5) was obtained as described
previously[15,42]. The concentrations of active RPA, RPA-OBA-Cy5, and Rad51 were determined
via a FRET-based activity assays as described previously[15,41,43].

FRET Measurements

All FRET experiments were performed at room temperature (23 ± 2 °C) in 1X Mg^{2+}/Ca^{2+}
buffer (20 mM HEPES, pH 7.5, 150 mM KCl, 5 mM $MgCl_2$, 5 mM $CaCl_2$) supplemented with 1
mM DTT, 1 mM ATP, and the ionic strength was adjusted to physiological (200 mM) by the
addition of appropriate amounts of KCl. Ca^{2+} is included to stabilize active Rad51 nucleoprotein
filaments on ssDNA via restriction of ATP hydrolysis (but not ATP binding) by Rad51[41]. Ca^{2+}
promotes more regular and stable Rad51 nucleoprotein filaments on ssDNA[41,44] but does not
affect the amount of RPA that binds to ssDNA nor the amount of PCNA loaded onto DNA by
RFC via a reaction mechanism that requires ATP binding and hydrolysis[15]. All experiments were
performed in a 16.100F-Q-10/Z15 sub-micro fluorometer cell (Starna Cells) and monitored in a
Horiba Scientific Duetta-Bio fluorescence/absorbance spectrometer as described previously[15]. In
short, reaction solutions are excited at 514 nm and the fluorescence emission intensities (*I*) are
monitored essentially simultaneously (Δt = 0.118 ms) at the peak emission wavelengths for Cy3
(563 nm, I_{563}) and Cy5 (665 nm, I_{665}) over time, recording *I* every 0.17 s. For all experiments,
excitation and emission slit widths are 10 nm. All recorded fluorescence emission intensities are

corrected by a respective dilution factor and all time courses are adjusted for the time between
the addition of each component and the fluorescence emission intensity recordings (i.e., dead
time $\leq$ 10 s). For any recording of the fluorescence emission intensities (I_{665} and I_{563}), the
approximate FRET efficiency is estimated from the equation $E_{FRET} = \frac{I_{665}}{I_{665} + I_{563}}$. For each
experiment below, the final concentrations of all reaction components are indicated. The
concentrations of ATP in all experimental solutions described below is 1.0 mM and, hence, this
concentration is maintained upon mixing.

Pre-steady state FRET experiments to monitor RPA/Rad51 exchange

For experiments performed in the absence of PCNA, RFC, and pol δ, a solution containing
ddP/5′TCy3 DNA (20 nM, **Figure S1**), NeutrAvidin (80 nM homotetramer), and ATP is pre-
incubated with an RPA (25 nM active heterotrimer, RPA or RPA-OBA-Cy5), the resultant
solution is transferred to a fluorometer cell, and the cell is placed in the instrument. I_{665} and I_{563}
are monitored over time until both stabilize for at least 1 min. Within this stable region, E_{FRET}
values are calculated from the observed I_{665} and I_{563} values and averaged to obtain the E_{FRET}
value observed prior to addition of Rad51•ATP. Then, pre-formed Rad51•ATP (500 nM active
Rad51 monomer, ATP) is added, the resultant solution is mixed by pipetting, and I_{665} and I_{563} are
monitored beginning $\leq$ 10 s after the addition of Rad51•ATP.
Experiments in the presence of PCNA and RFC are performed as described above except that
PCNA is first pre-loaded onto DNA, as previously described[15]. In short, an RPA (25 nM active
heterotrimer, RPA or RPA-OBA-Cy5) is added to a solution containing ddP/5′TCy3 DNA (20
nM, **Figure S1**), NeutrAvidin (80 nM homotetramer), and ATP. PCNA (20 nM homotrimer,
PCNA or Cy5-PCNA) and ATP-bound RFC (i.e., RFC•ATP, 20 nM RFC heteropentamer, ATP)
are then added in succession. RPA-OBA-Cy5 fully supports RFC-catalyzed loading of PCNA
onto P/T junctions[15]. The resultant solution is transferred to a fluorometer cell, and the cell is
placed in the instrument. 5 min after the addition of RFC•ATP (i.e., PCNA loading completed),
I_{665} and I_{563} are monitored over time until both stabilize for at least 1 min. Within this stable
region, E_{FRET} values are calculated from the observed I_{665} and I_{563} values and averaged to obtain
the E_{FRET} value observed prior to addition of Rad51•ATP. Finally, pre-formed Rad51•ATP (500
nM active Rad51 monomer, ATP) is added and the experiment is continued as described above.
Experiments in the presence of PCNA, RFC, pol δ, +/- dGTP experiments are performed as
described above except pol δ holoenzymes are first pre-assembled on P/T DNA, as previously
described[15]. In short, a PCNA (20 nM homotrimer, PCNA or Cy5-PCNA) is loaded by
RFC•ATP (20 nM RFC heteropentamer, ATP) onto ddP/5′TCy3 DNA (20 nM, **Figure S1**) that
is pre-engaged by NeutrAvidin (80 nM homotetramer) and an RPA (25 nM active heterotrimer,
RPA or RPA-OBA-Cy5) exactly as described above. 5 min after the addition of RFC•ATP (i.e.,
PCNA loading complete), dGTP (0 or 1.0 mM) and pol δ (40 nM heterotetramer) are added in
immediate succession, the resultant solution is transferred to a fluorometer cell, and the cell is
placed in the instrument. 5 min after the addition of pol δ +/- dGTP (i.e., formation of pol δ
holoenzymes or adoption of initiation states of pol δ holoenzymes for DNA synthesis is
completed), I_{665} and I_{563} are monitored over time until both stabilize for at least 1 min. Within
this stable region, E_{FRET} values are calculated from the observed I_{665} and I_{563} values and averaged
to obtain the E_{FRET} value observed prior to addition of Rad51•ATP. Finally, pre-formed
Rad51•ATP (500 nM active Rad51 monomer, ATP) are added and the experiment is continued
as described above.

To determine the predicted E_{FRET} trace for RPA-OBA-Cy5 remaining completely disengaged
from the ddP/5′TCy3 DNA substrate in the aforementioned experiments, I_{665} and I_{563} for RPA-
OBA-Cy5 alone (I_{665}^{RPA} and I_{563}^{RPA}) and a corresponding reaction mixture lacking only RPA-OBA-
Cy5 (I_{665}^{DNA} and I_{563}^{DNA}) are each monitored, and E_{FRET} is calculated as follows:

$$\frac{(I_{665}^{RPA} + I_{665}^{DNA})}{(I_{665}^{RPA} + I_{665}^{DNA}) + (I_{563}^{RPA} + I_{563}^{DNA})}$$

Pre-steady FRET Experiments to monitor PCNA Loading

Experiments in the presence of RPA are carried out exactly as described in a previous
report[15]. In short, RPA (25 nM active heterotrimer) is added to a solution containing 5′ddPCy3/T
DNA (20 nM, **Figure S1**), NeutrAvidin (80 nM homotetramer), and ATP. Cy5-PCNA (20 nM
homotrimer) is added, the resultant solution is transferred to a fluorometer cell, and the cell is
placed in the instrument. I_{665} and I_{563} are monitored over time until both stabilize for at least 1
min. Within this stable region, E_{FRET} values are calculated from the observed I_{665} and I_{563} values
and averaged to obtain the E_{FRET} value observed prior to addition of RFC•ATP. Finally,
RFC•ATP (20 nM RFC heteropentamer, ATP) is added, the resultant solution is mixed by
pipetting, and I_{665} and I_{563} are monitored beginning $\leq$ 10 s after the addition of RFC•ATP.
Experiments in the presence of Rad51•ATP are carried out exactly as described above except
that RPA is replaced with pre-formed Rad51•ATP (200 nM active Rad51 monomer, ATP).

To determine the predicted E_{FRET} trace for Cy5-PCNA remaining completely disengaged from
5′ddPCy3/T nucleoprotein complexes/filaments (i.e., Rad51 and RPA present) in the
aforementioned experiments, I_{665} and I_{563} for Cy5-PCNA alone (I_{665}^{PCNA} and I_{563}^{PCNA}) and a
corresponding reaction mixture lacking only Cy5-PCNA (I_{665}^{DNA} and I_{563}^{DNA}) are each monitored,
and E_{FRET} is calculated as follows:

$$\frac{(I_{665}^{PCNA} + I_{665}^{DNA})}{(I_{665}^{PCNA} + I_{665}^{DNA}) + (I_{563}^{PCNA} + I_{563}^{DNA})}$$

Results

PCNA encircling P/T junctions engaged by RPA are irreversibly unloaded through RPA/Rad51
exchange

In a previous study, we developed ensemble FRET assays that directly and continuously
monitor interactions of RPA and PCNA with P/T junctions during pol δ holoenzyme assembly
and the initiation of DNA synthesis[15,38]. In the present study, we adapt these assays to monitor
interactions of RPA and PCNA with P/T junctions during RPA/Rad51 exchange under a variety
of conditions that mimic *in vivo* scenarios. A high-fidelity DNA polymerase, such as pol δ,
dissociates into solution upon encountering DNA lesions it cannot accommodate, leaving loaded
PCNA behind on the dsDNA regions upstream of the vacated P/T junctions. Thus, we initially
focus on RPA/Rad51 exchange at P/T junctions encircled by PCNA (**Figure 2A**). Here, a Cy3-
labeled P/T DNA substrate (**Figure S1**, ddP/5′TCy3 or 5′ddPCy3/T) is first pre-bound with
NeutrAvidin and the respective RPA (RPA or RPA-OBA-Cy5). NeutrAvidin avidly binds biotin
attached to the 5′ termini of the primer strands (K_D ~ pM) and the resultant complex prohibits
diffusion of loaded PCNA off the blunt duplex ends of the P/T DNA substrates, emulating the
aforementioned protein roadblocks upstream of stalled replication sites[2,4,34-37].The lengths of the
template ssDNA regions (33 nucleotides, nts) of the P/T DNA substrates accommodate 1 RPA
heterotrimer in the 30 nucleotide binding mode ($K_D \leq$ 100 pM) at the RPA:DNA ratios utilized[5-

[7,15,45]. As indicated above, RPA promotes PCNA occupancy of P/T junctions through multiple
means. Together, the biotin•NeutrAvidin and RPA•DNA complexes maintain loaded PCNA on
the dsDNA regions of the P/T DNA substrates[4,15,24]. Next, a PCNA is added (PCNA or Cy5-
PCNA) followed by RFC•ATP and the resultant solution is incubated until it reaches equilibrium
where a PCNA (PCNA or Cy5-PCNA) is loaded and maintained on all P/T DNA•RPA
complexes[4,15]. Finally, Rad51•ATP is added. All assays in the current study are performed at
physiological ionic strength (200 mM I) in the absence of ATP hydrolysis by Rad51 (see
Materials and Methods). Under these conditions, Rad51 has an exceptionally strong, intrinsic
preference for forming filaments on ssDNA (**Figure S2**) such that dsDNA binding is
dramatically reduced if observed at all[46-48]. The ssDNA regions of the Cy3-labeled P/T DNA

Figure 2. All PCNA encircling P/T junctions engaged by RPA are irreversibly unloaded through RPA/Rad51 exchange (**A**) Schematic representations of the FRET pairs and experiments to monitor the interactions of RPA (*Left*) and PCNA (*Right*) with P/T junctions during RPA/Rad51 exchange (**B - C**) Data. Each trace is the mean of at least three independent traces with the S.E.M. shown in grey. The time trajectories of I_{563} (magenta) and I_{665} (cyan) are displayed in the top panels and the corresponding E_{FRET} (yellow) is displayed in the bottom panels. The times at which Rad51•ATP are added is indicated by red arrows. Changes in I_{563} and I_{665} are indicated in the top panels by magenta and cyan arrows, respectively. For observation, the I_{563} and I_{665}, and the corresponding E_{FRET} values observed prior to the addition of Rad51•ATP are each fit to dashed flat lines that are extrapolated to the axes limits. The predicted E_{FRET} traces for no interaction between the respective Cyanine-labeled species are displayed in pink in each bottom panel, offset to the time after Rad51•ATP is added and fit to a dashed flat line that is extrapolated to the axis limits. Data for interactions of RPA and PCNA with P/T junctions during RPA/Rad51 exchange are shown in panels **B** and **C,** respectively. The rate constant(s) for the E_{FRET} decrease observed after the addition of Rad51•ATP in each bottom panel is reported in **Table 1**.

substrates (**Figure S1**, ddP/5′TCy3 or 5′ddPCy3/T) accommodate Rad51•ATP filaments
comprised of 11 Rad51•ATP monomersa.
For experiments monitoring interactions of RPA with P/T junctions (**Figure 2A**, *Left*), I_{563}
and I_{665} values are stabilized (**Figure 2B**, *Top*) and a significant, constant E_{FRET} is observed prior
to the addition of Rad51•ATP (**Figure 2B**, *Bottom*) due to the stable interaction of RPA-OBA-
Cy5 with the ddP/5′TCy3 DNA[15]. Upon addition of Rad51•ATP, I_{563} increases simultaneously
with decreases in I_{665} (**Figure 2B**, *Top*) . These synchronized, anti-correlated changes in I_{563} and
I_{665} are indicative of a direct FRET decrease due to the increase in distance between the Cy3
FRET donor on the ddP/5′TCy3 DNA substrate and the Cy5 FRET acceptor on RPA-OBD-Cy5.
The E_{FRET} decreases in a monophasic manner down to the E_{FRET} trace predicted for no interaction
between ddP/5′TCy3 DNA and RPA-OBA-Cy5 (**Figure 2B**, *Bottom*). This indicates that
Rad51•ATP filaments irreversibly exchange with RPA on all ddP/5′TCy3 DNA, displacing all
RPA into solution. The E_{FRET} decrease observed for RPA/Rad51 exchange is best fit to a one-
step, irreversible kinetic model (i.e., single exponential decline) with a rate constant of $k_{obs\ dec}$ =
8.59 ± 0.05 (x 10^{-3}) s^{-1} (**Table 1**). Interestingly, RPA/Rad51 exchange observed in **Figure 2B** is
~2.2-fold slower compared to that observed for the reference condition where the P/T
DNA•RPA complexes are devoid of PCNA ($k_{obs\ dec}$ = 1.87 ± 0.01 (x 10^{-2}) s^{-1}, **Figure S3**, **Table**
1). Hence, the pre-occupancy of P/T DNA•RPA complexes by loaded PCNA slows but does not
prohibit RPA/Rad51exchange.
For experiments monitoring interactions of PCNA with P/T junctions (**Figure 2A**, *Right*),
I_{563} and I_{665} values are stabile over time (**Figure 2D**, *Top*) and a significant, constant E_{FRET} is
observed prior to the addition of Rad51•ATP (**Figure 2D**, *Bottom*) due to the loading and
maintenance of Cy5-PCNA on all 5′ddPCy3/T DNA•RPA complexes[15]. Upon addition of
Rad51•ATP, I_{563} increases simultaneously with decreases in I_{665} (**Figure 2D**, *Top*) . These
synchronized, anti-correlated changes in I_{563} and I_{665} are indicative of a direct FRET decrease
due to the increase in distance between the Cy3 FRET donor on the 5′ddPCy3/T DNA and the
Cy5 FRET acceptor on Cy5-PCNA. The E_{FRET} decreases in a biphasic manner down to the E_{FRET}
trace predicted for no interaction between 5′ddPCy3/T DNA and Cy5-PCNA (**Figure 2D**,
Bottom). This indicates that all PCNA encircling 5′ddPCy3/T DNA irreversibly unload. The
observed E_{FRET} decrease is best fit to a two-step, irreversible kinetic model with rate constants of
$k_{obs\ dec\ 1}$ = 5.47 ± 0.03 (x 10^{-3}) s^{-1} and $k_{obs\ dec\ 2}$ = 9.64 ± 0.02 (x 10^{-4}) s^{-1} where $k_{obs\ dec\ 2}$ reports on
PCNA unloading (**Table 1**). Direct comparisons of the data presented in **Figures 2B** and **2C**
indicate that upon addition of Rad51•ATP both RPA/Rad51 exchange and PCNA unloading go
to completion and are irreversible. Next, we sought to identify correlations between RPA/Rad51
exchange and the occupancy of P/T junctions by PCNA.
The E_{FRET} decreases observed in **Figures 2B** and **Figure 2C** were each normalized to their
respective ranges and the results plotted as function of time (after addition of Rad51•ATP). As
observed in **Figure S4**, PCNA begins unloading from P/T junctions immediately upon addition
of Rad51•ATP and becomes significant (i.e., $\geq$ 10%) when RPA/Rad51 exchange $\geq$ 90%
completed. ~40% of PCNA unloads as RPA/Rad51 exchange is occurring on P/T junctions; the
remaining ~60 % unloads after RPA/Rad51 exchange has been completed on all P/T junctions
(i.e., RPA has been replaced by Rad51•ATP filaments on all P/T junctions). Altogether, this
reveals that unloading of PCNA from P/T DNA•RPA complexes requires RPA/Rad51 exchange
but not vice versa. Rather, the occupancy of P/T DNA•RPA complexes by loaded PCNA slows
but does not prohibit RPA/Rad51exchange (**Figures 2B**, **S3**, and **Table 1**).

An abundant PCNA-binding protein complex does not stabilize loaded PCNA on P/T junctions during RPA/Rad51 exchange.

Pol δ is maintained at stalled replication sites induced by UVR exposure and may re-engage the resident PCNA, forming holoenzymes that re-iteratively associate and dissociate from the P/T junctions. However, pol δ holoenzymes cannot support stable insertion of a dNTP opposite the offending lesion. Furthermore, hundreds of additional PCNA-binding proteins/protein complexes are maintained or enriched on PCNA encircling P/T junctions of stalled replication sites induced by UVR exposure[15,24,49-54]. To emulate RPA/Rad51 exchange in this scenario where PCNA-binding proteins/protein complexes are locally abundant, we repeated the experiments described above (in **Figure 2**) exactly as indicated except that PCNA encircling P/T DNA•RPA complexes is preincubated with a 2-fold excess of pol δ prior to the addition of Rad51•ATP (**Figure 3A**). Under these conditions, pol δ binding to loaded PCNA is stoichiometric and unloading of PCNA from P/T DNA•RPA complexes is not observed up to a 4-fold excess of pol δ[15]. Excess pol δ is utilized to ensure all PCNA encircling P/T DNA•RPA complexes are engaged in pol δ holoenzymes prior to the addition Rad51•ATP[15]. However, in the absence of dNTPs, pol δ holoenzymes have dramatically low affinity, if any, for native (i.e., undamaged) P/T junctions. Thus, pol δ holoenzymes do not stably engage P/T junctions in these experiments, emulating that observed at P/T junctions of stalled replication sites[4,15,50-53].

For experiments monitoring interactions of RPA with P/T junctions (**Figure 3A**, *Left*), the time-dependent traces for I_{563} and I_{665} (**Figure 3B**, *Top*) and E_{FRET} (**Figure 3B**, *Bottom*) are similar to those observed in **Figure 2B** where pol δ was absent. This indicates that Rad51•ATP filaments irreversibly exchange with RPA on all ddP/5′TCy3 DNA under these conditions, displacing all RPA into solution. The E_{FRET} decrease observed for RPA/Rad51 exchange is best fit to a one-step, irreversible kinetic model (i.e., single exponential decline) with a rate constant [$k_{obs\ dec}$ = 1.23 ± 0.01 (x 10^{-2}) s^{-1}] that is ~1.5-fold slower compared to that observed for the reference condition (**Figure S3**, **Table 1**). Together with the observations described above, this indicates that occupancy of P/T DNA•RPA complexes by either PCNA alone (**Figure 2B**) or PCNA engaged in pol δ holoenzymes (**Figure 3B**) slows but does not prohibit RPA/Rad51exchange.

For experiments monitoring interactions of PCNA with P/T junctions (**Figure 3A**, *Right*), the behaviors of the time-dependent traces for I_{563} and I_{665} (**Figure 3C**, *Top*) and E_{FRET} (**Figure 3C**, *Bottom*) are similar to those observed in **Figure 2C** where pol δ was absent. This indicates that, despite being pre-engaged by in pol δ holoenzymes, all PCNA encircling ddP/5'TCy3 DNA irreversibly unload. Interestingly, the E_{FRET} decrease observed for PCNA unloading under these conditions (**Figure 3C**, *Bottom*) is best fit to a triple exponential decline with a fast rate constant ($k_{obs\ dec\ 1}$ = 4.19 ± 0.3 (x 10^{-2}) s^{-1}) that accounts for 19.5% ± 0.8% of the decline and is significantly greater than the slower rate constants ($k_{obs\ dec\ 2}$ = 9.16 ± 0.2 (x 10^{-3}) s^{-1}, $k_{obs\ dec\ 3}$ = 1.16 ± 0.05 (x 10^{-3}) s^{-1}) (**Table 1**). The fast rate constant is not observed when pol δ is omitted (**Figure 2C**, *Bottom*). The slower rate constants ($k_{obs\ dec\ 2}$ and $k_{obs\ dec\ 3}$) observed in the presence of pol δ are very similar to the rate constants ($k_{obs\ dec\ 1,}$ $k_{obs\ dec\ 2}$) observed in the absence of pol δ (**Table 1**). Consequently, upon addition of Rad51•ATP, subsequent unloading of PCNA from P/T DNA is achieved relatively faster when pre-loaded PCNA is engaged in pol δ holoenzymes compared to when it is not (**Figure S5**). Thus, pol δ holoenzymes do not impart any stability to loaded PCNA during RPA/Rad51 exchange. This suggests that tight interactions of PCNA-binding proteins/protein complexes with loaded PCNA do not stabilize sliding clamps on P/T

junctions during RPA/Rad51 exchange. Direct comparisons of the data presented in **Figures 3B**
and **3C** indicate that upon addition of Rad51•ATP both RPA/Rad51 exchange and PCNA
unloading go to completion and are irreversible under these conditions. Next, we compared the
timescales of RPA/Rad51 exchange and PCNA unloading under the conditions depicted in
Figure 3A to identify correlations between these two processes.
As observed in **Figure S6**, PCNA begins unloading from P/T junctions immediately upon
addition of Rad51•ATP and becomes significant (i.e., $\geq 10\%$) when RPA/Rad51 exchange is
only ~ 10 % completed. ~75% of PCNA unloads as RPA/Rad51 exchange is occurring on the
P/T DNA and the remaining ~25 % unloads after RPA has been replaced by Rad51•ATP
filaments on all P/T DNA. This contrasts that observed in the absence of pol δ where the
majority (~60%) of PCNA unloads after RPA/Rad51 exchange has been completed on all P/T

Figure 3. Pol δ does not stabilize loaded PCNA on P/T junctions during RPA/Rad51 exchange. (**A**) Schematic representations of the FRET pairs and experiments to monitor the interactions of RPA (*Left*) and PCNA (*Right*) with P/T junctions during RPA/Rad51 exchange in the presence of pol δ (**B - C**) Data. Each trace is the mean of at least three independent traces with the S.E.M. shown in grey. The time trajectories of I_{563} (magenta) and I_{665} (cyan) are displayed in the top panels and the corresponding E_{FRET} (yellow) is displayed in the bottom panels. The times at which Rad51•ATP are added is indicated by red arrows. Changes in I_{563} and I_{665} are indicated in the top panels by magenta and cyan arrows, respectively. For observation, the I_{563} and I_{665}, and the corresponding E_{FRET} values observed prior to the addition of Rad51•ATP are each fit to dashed flat lines that are extrapolated to the axes limits. The predicted E_{FRET} traces for no interaction between the respective Cyanine-labeled species are displayed in pink in each bottom panel, offset to the time after Rad51•ATP is added and fit to a dashed flat line that is extrapolated to the axis limits. Data for interactions of RPA and PCNA with P/T junctions during RPA/Rad51 exchange in the presence of pol δ are shown in panels **B** and **C,** respectively. The rate constant(s) for the E_{FRET} decrease observed after the addition of Rad51•ATP in each bottom panel is reported in **Table 1**.

DNA (**Figure S4**). Regardless, this confirms that unloading of PCNA from P/T DNA•RPA
complexes requires RPA/Rad51 exchange but not vice versa. Next, we investigate the
mechanism(s) of RPA/Rad51 exchange-dependent PCNA unloading.

RPA/Rad51 exchange on P/T DNA eliminates the resident PCNA by inhibiting RFC
In the experiments described in **Figures 2** and **3** above, P/T junctions are initially engaged by
RPA and ATP-bound RFC (i.e., RFC•ATP) pre-loads PCNA onto all P/T junctions. After
dissociation of the RFC•ADP complex into solution, loaded PCNA is left behind on the dsDNA
region upstream of the P/T junctions[15,33]. The biotin-neutravidin complexes prohibit diffusion of
loaded PCNA off the blunt duplex ends of the P/T DNA. RPA•ssDNA complexes prohibit
diffusion of loaded PCNA off the ssDNA ends of the P/T DNA as well as RFC-catalyzed
unloading of PCNA[4,37]. Thus, prior to the addition of Rad51•ATP, the only pathway for
unloading of PCNA from DNA is through spontaneous opening of the PCNA ring. This non-
catalytic process is dramatically slow and, in the event that it occurs, RPA permits RFC•ATP to
instantly "re-load" free PCNA back onto P/T junctions such that unloading is not
observed[2,4,15,24]. This is confirmed for the experimental system in the current study (**Figure S7**).
Hence, the stable (i.e., equilibrium) E_{FRET} traces observed prior to the addition of Rad51•ATP in
Figures 2C and **3C** represent RFC•ATP maintaining loaded Cy5-PCNA on all P/T DNA•RPA
complexes.
In the present study, all rate constants observed for RPA/Rad51 exchange-dependent PCNA
unloading are $\geq$ ~13-fold slower than the rate constant for RFC-catalyzed unloading of PCNA
($k_{obs} = 0.54 \pm 0.09$ s^{-1})[2]. This indicates that RPA/Rad51 exchange on P/T junctions does not
promote RFC-catalyzed unloading of the resident PCNA. Furthermore, ATP hydrolysis by
Rad51 is restricted in the present study, Rad51 lacks any known PCNA-binding motif(s), and, to
the best of our knowledge, direct interactions between Rad51 and PCNA have yet to be reported.
This indicates that Rad51 does not catalyze PCNA unloading. Altogether, this suggests that
RPA/Rad51 exchange-dependent PCNA unloading is non-catalytic. Two general populations
are observed for this process. The "slow" population is $\leq$ 2.0-fold slower than spontaneous
opening of the PCNA ring ($k_{dec,obs} = 1.93 \pm 0.01$ (x 10^{-3}) s^{-1}, **Figure S7**) and accounts for 100%
and 44.3 ± 1.1 % of the unloading observed in **Figure 2C** ($k_{obs\ dec\ 2}$) and **Figure 3C** ($k_{obs\ dec\ 3}$),
respectively. The "fast" population is $\geq$ ~ 4.74-fold faster than spontaneous opening of the
PCNA ring, accounts for 55.7 ± 1.1 % of the unloading observed in **Figure 3C** ($k_{obs\ dec\ 1}$, $k_{obs\ dec}$
$_2$), and only appears when Rad51•ATP is added to loaded PCNA pre-assembled in pol δ
holoenzymes. Altogether, this suggests that RPA/Rad51 exchange-dependent PCNA unloading
is non-catalytic and occurs through spontaneous opening of the PCNA ring or a mechanism
where Rad51•ATP and pol δ in conjunction somehow destabilize the PCNA ring.
RPA/Rad51-dependent PCNA unloading observed in **Figures 2B** and **2**C generates "free"
PCNA in solution, goes to completion, and is irreversible, despite the continuous presence of
RFC•ATP. This is only possible if assembly of stable Rad51•ATP filaments on P/T junctions
via RPA/Rad51 exchange also prohibits RFC•ATP from loading free PCNA onto P/T junctions
that are devoid of the sliding clamp. To confirm this, we monitor and directly compare RFC-
catalyzed loading of "free" PCNA on P/T junctions engaged by either RPA or Rad51•ATP
filaments (**Figure 4**)[15]. 5'ddPCy3/T DNA substrates are first pre-bound with NeutrAvidin and
either RPA or Rad51•ATP filaments. Next, Cy5-PCNA is added and the resultant solution is
equilibrated. Finally, RFC•ATP is added. For experiments on P/T junctions engaged by RPA
(**Figure 4A**, *Left*), I_{563} and I_{665} values are stable over time (**Figure 4B**, *Top*) and a low, constant

E_{FRET} is observed prior to the addition of RFC•ATP (**Figure 4B**, *Bottom*) due to the presence of

both the Cy3 donor and the Cy5 acceptor. E_{FRET} values observed during this period are a true

experimental baseline representing the absence of any interactions between Cy5-PCNA and the

5′ddPCy3/T DNA•RPA complexes[15]. Upon addition of RFC•ATP, I_{665} increases concomitantly

with a decrease in I_{563} after which both fluorescence emission intensities stabilize and persist

(**Figure 4B**, *Top*). These synchronized, anti-correlated changes in I_{563} and I_{665} are direct

indicators of the appearance and increase in FRET (**Figure 4B**, *Bottom*) due to the decrease in

distance between the Cy3 FRET donor on the 5′ddPCy3/T DNA substrate and the Cy5 FRET

acceptor on Cy5-PCNA. The observed E_{FRET} increase (**Figure 4B**, *Bottom*) is biphasic with k_{obs}

$_{inc\ 1}$ = 4.46 $\pm$ 0.2 (x 10^{-2}) s^{-1} and $k_{obs\ inc\ 2}$ = 1.70 $\pm$ 0.12 (x 10^{-2}) s^{-1}. These rate constants are in

excellent agreement with values obtained in identical experiments from a recent study[15]. At the

plateau of the E_{FRET} increase, Cy5-PCNA is loaded and maintained on all 5′ddPCy3/T•RPA

complexes[15].

Figure 4. Rad51•ATP filaments prohibit RFC-catalyzed loading of PCNA onto P/T junctions. (**A**) Schematic representations of the FRET pair and experiments to monitor the RFC-catalyzed loading of PCNA onto P/T junctions engaged by RPA (*Left*) or Rad51•ATP filaments (*Right*) (**B - C**) Data. Each trace is the mean of at least three independent traces with the S.E.M. shown in grey. The time trajectories of I_{563} (magenta) and I_{665} (cyan) are displayed in the top panels and the corresponding E_{FRET} (mustard) is displayed in the bottom panels. The time at which RFC•ATP is added is indicated by a red arrow. Changes in I_{563} and I_{665} are indicated in the top panel by magenta and cyan arrows, respectively. For observation, the I_{563} and I_{665}, and the corresponding E_{FRET} values observed prior to the addition RFC•ATP are each fit to dashed flat lines that are extrapolated to the axis limits. (**B**) Data for RFC-catalyzed loading of PCNA onto P/T junctions engaged by RPA. (**C**) Data for RFC-catalyzed loading of PCNA onto P/T junctions engaged by Rad51•ATP filaments.

For experiments on P/T junctions engaged by Rad51•ATP filaments (**Figure 4A**, *Right*), I_{563}
and I_{665} values are stabile over time (**Figure 4C**, *Top*) and a low, constant E_{FRET} is observed prior
to the addition of RFC•ATP (**Figure 4C**, *Bottom*) due to the presence of both the Cy3 donor and
the Cy5 acceptor. Again, E_{FRET} values observed during this period are a true experimental
baseline representing the absence of any interactions between Cy5-PCNA and the 5′ddPCy3/T
DNA•Rad51•ATP complexes. Upon addition of RFC•ATP, both I_{665} and I_{563} slightly but
reproducibly decrease within the dead time after which both I values remain flat (**Figure 4B**,
Top). These synchronized, correlated changes in I_{563} and I_{665} indicate that any observed changes
in FRET are indirect[15,55]. Nevertheless, a change in E_{FRET} is not observed upon addition of
RFC•ATP due to I_{563} and I_{665} each decreasing by the same relative amount (I_{665} by 10.6 ± 3.1 %
and I_{563} by 10.3 ± 2.0 %). This confirms that assembly of stable Rad51•ATP filaments on P/T
junctions via RPA/Rad51 exchange prohibit RFC•ATP from loading "free" PCNA on P/T
junctions that are devoid of the sliding clamp.
424
Rad51•ATP does not require access to 3′ primer termini of P/T junctions and the ssDNA
immediately downstream for RPA/Rad51 exchange
Formation of Rad51•ATP filaments occurs in a stepwise in a manner. First, 1 - 3 Rad51
monomers, each engaged with ATP, serve as a nucleus to initiate filament assembly (i.e.,
nucleate) on a template ssDNA. Next, the filament nuclei extend (i.e., grow/elongate) in either
direction by the addition of Rad51 monomers, each engaged with ATP[46-48]. Previous single
molecule studies on a "gapped" DNA substrate comprised of dsDNA regions (i.e., handles)
connected by an RPA-coated ssDNA gap revealed that nucleation of Rad51 is intrinsically
enriched at/near the 3′ dsDNA/ssDNA junction compared to the 5′ dsDNA/ssDNA junction[47]. A
3′ dsDNA/ssDNA junction contains a recessed 3′ primer terminus (from where DNA synthesis
can initiate from) and a 5′ template ssDNA overhang. These junctions are amenable to those
utilized in the present study (i.e., P/T junctions). Thus, we next sought to address whether
Rad51•ATP requires access to the recessed 3′ primer termini of P/T junctions for nucleation and,
hence, RPA/Rad51 exchange. To do so, we repeated the experiments described above in **Figure**
3 above except that loaded PCNA is preincubated with a 2-fold excess of pol δ and dGTP prior
to the addition of Rad51•ATP (**Figure 5A**). Under these conditions, pol δ engages the "front
face" of loaded PCNA, forming a holoenzyme, engages the P/T junction, and aligns an incoming
dGTP at the 3′ terminus of the primer strand in a complimentary bp with the template nucleotide
(a Cytosine) immediately 5′ of the P/T junction. However, both extension of the primer (via pol
δ DNA polymerase activity) and degradation of the primer (via pol δ exonuclease activity) are
prohibited due to the 3′ dideoxy-terminated primer strand and the use of exonuclease-deficient
pol δ. Hence, the pol δ holoenzymes are trapped in initiation states for DNA synthesis. In this
state (depicted in **Figure 5A**), RPA has vacated the 3′ primer termini of P/T junctions and ~ 6 nts
of the template ssDNA immediately downstream and these sites are sequestered within the DNA
polymerase active site of pol δ[4,15,52,56].
For experiments monitoring interactions of RPA with P/T junctions (**Figure 5A**, *Left*), the
time-dependent traces for I_{563} and I_{665} (**Figure 5B**, *Top*) and E_{FRET} (**Figure 5B**, *Bottom*) are
similar to those observed in **Figures 2B** and **3B** where either dGTP was absent (**Figure 3B**) or
both pol δ and dGTP were absent (**Figure 2B**). This indicates that Rad51•ATP filaments
irreversibly exchange with RPA on all ddP/5′TCy3 DNA under these conditions, displacing all
RPA into solution. Again, the E_{FRET} decrease observed during RPA/Rad51 exchange is best fit to

a one-step, irreversible kinetic model with a rate constant of $k_{obs\ dec} = 2.05 \pm 0.03$ (x 10^{-3}) s^{-1},

which is faster than all rate constants observed for RPA/Rad51 exchange in the current study

(**Table 1**). For experiments monitoring interactions of PCNA with P/T junctions (**Figure 5A**,

Right), I_{563} and I_{665} values are stabile over time (**Figure 5C**, *Top*) and a significant, constant

E$_{FRET}$ is observed prior to the addition of Rad51•ATP (**Figure CD**, *Bottom*) due to the initiation

states of pol δ holoenzymes[15]. Upon addition of Rad51•ATP, the I_{563} and I_{665} traces are

maintained (**Figure 5C**, *Top*) and a change in E$_{FRET}$ is not observed for over 15 mins (**Figure**

5C, *Bottom*). This confirms that the initiation state of pol δ holoenzymes is achieved in these

experiments prior to the addition of Rad51•ATP and reveals that this state stabilizes loaded

Figure 5. Rad51•ATP does not require access to 3′ primer termini of P/T junctions and the ssDNA immediately downstream for RPA/Rad51 exchange. (**A**) Schematic representations of the FRET pairs and experiments to monitor the interactions of RPA (*Left*) and PCNA (*Right*) with P/T junctions during RPA/Rad51 exchange in the presence of pol δ and dGTP (**B - C**) Data. Each trace is the mean of at least three independent traces with the S.E.M. shown in grey. The time trajectories of I_{563} (magenta) and I_{665} (cyan) are displayed in the top panels and the corresponding E$_{FRET}$ (mustard) are displayed in the bottom panels. The times at which Rad51•ATP is added are indicated by red arrows. Changes in I_{563} and I_{665}, if they occur, are indicated in the top panels by magenta and cyan arrows, respectively. For observation, the I_{563} and I_{665}, and the corresponding E$_{FRET}$ values observed prior to the addition Rad51•ATP are each fit to dashed flat lines that are extrapolated to the axis limits. The predicted E$_{FRET}$ traces for no interaction between the respective Cyanine-labeled species are displayed in pink in each bottom panel, offset to the time after Rad51•ATP is added and fit to a dashed flat line that is extrapolated to the axis limits. Data for interactions of RPA and PCNA with P/T junctions during RPA/Rad51 exchange in the presence of pol δ and dGTP are shown in panels **B** and **C,** respectively. The rate constant for the E$_{FRET}$ decrease observed after the addition of Rad51•ATP in panel **B** is reported in **Table 1**.

PCNA on P/T junctions during RPA/Rad51 exchange (for ≥ 1000 s). Direct comparisons of the data presented in **Figures 5B** and **5C** indicate that Rad51•ATP nucleates and completely exchanges with RPA (**Figure 5B**) while the 3′ primer termini of the P/T junctions and the template ssDNA immediately downstream are sequestered by pol δ holoenzymes in the initiation state (**Figure 5C**). Thus, Rad51•ATP does not require access to the recessed 3′ primer termini of P/T junctions for nucleation and, hence, RPA/Rad51 exchange.

Table 1. Kinetic rate constants observed in the present study for RPA/Rad51 exchange and RPA/Rad51 exchange-dependent PCNA unloading. N/A = Not Applicable

	Figure	S3B	2B	2C	3B	3C	5B	5C
Reaction Components	RPA	+ Cy5-RPA	+ Cy5-RPA	RPA	+ Cy5-RPA	RPA	+ Cy5-RPA	RPA
	RFC, PCNA	-	+ RFC, PCNA	+ RFC, Cy5-PCNA	+ RFC, PCNA	+ RFC, Cy5-PCNA	+ RFC, PCNA	+ RFC, Cy5-PCNA
	Pol δ	-	-	-	+	+	+	+
	dGTP	-	-	-	-	-	+	+
	Rad51	+	+	+	+	+	+	+
Rate Constant(s)	RPA/Rad51 Exchange	1.87 + 0.01 (x 10^{-2}) s^{-1}	8.59 + 0.05 (x 10^{-3}) s^{-1}	N/A	1.23 + 0.01 (x 10^{-2}) s^{-1}	N/A	2.05 + 0.03 (x 10^{-2}) s^{-1}	N/A
	RPA/Rad51 Exchange-dependent PCNA Unloading	N/A	N/A	5.47 + 0.03 (x 10^{-3}) s^{-1} 9.64 + 0.02 (x 10^{-4}) s^{-1}	N/A	4.19 + 0.3 (x 10^{-2}) s^{-1} 9.16 + 0.2 (x 10^{-3}) s^{-1} 1.16 + 0.05 (x 10^{-3}) s^{-1}	N/A	Not Observed

Discussion

Upon the stalling of canonical DNA replication at DNA lesions, such as UVR-induced photoproducts, the resident PCNA are left behind on the dsDNA regions upstream of the afflicted P/T junctions, which are directly engaged by RPA (**Figure 6**, *Top Left*). RPA also coats the unreplicated ssDNA templates downstream[19,20,57]. In humans, these stalling events are circumvented by one of at least three interconnected pathways; TLS, TS, HDR. PCNA encircling P/T junctions of stalled replication sites are central to the activation of both TLS and TS (**Figure 1B**). As described above, "protein roadblocks" restrict diffusion of loaded PCNA on DNA, collectively localizing the sliding clamps at/near P/T junctions of stalled replication sites[2,4,15,23,24,33-37,58,59]. Nevertheless, PCNA can unload from these sites and this process can be catalyzed by enzyme complexes, such as ATAD5, or occur via spontaneous opening of the

Figure 6. Model for interplay between RPA/Rad51 exchange and PCNA occupancy at P/T junctions of stalled replication sites. (*Left*) RFC maintains PCNA on P/T junctions engaged by RPA, promoting TLS and/or TS. (*Right*) Assembly of stable Rad51·ATP filaments on P/T junctions of stalled replication sites via RPA/Rad51 exchange ultimately leads to the complete elimination of the resident PCNA, prohibiting TLS and TS and promoting HDR.

PCNA ring (**Figure S7**)[2,60-65]. For loading PCNA onto P/T junctions, assembled
RFC•ATP•PCNA complexes, i.e., loading complexes, require direct interactions with the 3′
termini of primer strands and the template ssDNA immediately downstream[52,66]. RPA directly
engages these sites but permits access to loading complexes through microscopic dissociation
events within the RPA complex. Consequently, RFC loads and maintains PCNA on P/T
junctions that are directly engaged by RPA (**Figure S7**)[4,15]. This suggests that at stalled
replication sites where the P/T junctions are directly engaged by RPA, RFC maintains loaded
PCNA at these sites, promoting TLS and/or TS (**Figure 6**, *Left*).

Exchange of RPA for Rad51•ATP along the unreplicated ssDNA template sequences of
stalled replication sites is central to the activation of HDR (**Figure 1B**)[21,22,32,67,68]. The results
from the present study reveal that PCNA encircling P/T junctions engaged by RPA unload
slowly via non-catalytic mechanisms during and after (the complete) exchange of RPA for
Rad51•ATP on P/T junctions (**Figures 2-3**, **S4**, **S6**, **S7**). Furthermore, assembly of stable
Rad51•ATP filaments on P/T junctions via complete RPA/Rad51 exchange prohibit RFC•ATP
from loading "free" PCNA back onto P/T junctions that are devoid of the sliding clamp (**Figure 4**). Together, this leads to complete elimination of loaded PCNA from P/T junctions. This
phenomenon is observed both in the absence and presence of saturating pol δ, an abundant, high-
affinity PCNA-binding protein complex that is maintained at stalled replication sites induced by
UVR exposure (**Figures 2-3**, **S4-S6**)[49,54]. Multiple studies from many independent groups
revealed that stable Rad51•ATP filaments directly and statically engage the 3′ primer termini of
P/T junctions and the template ssDNA immediately downstream and these interactions
effectively shelter (i.e., "protect") the 3′ primer termini from processing by DNA metabolic
enzymes, such as 3′→5′ exonucleases[69]. Thus, stable Rad51•ATP filaments on P/T junctions
prohibit PCNA loading (as observed in **Figure 4B**) by restricting access of loading complexes to
3′ primer termini of P/T junctions and template ssDNA immediately downstream. Altogether,
this suggests that assembly of stable Rad51•ATP filaments at P/T junctions of stalled replication
sites via RPA/Rad51 exchange ultimately leads to the complete elimination of the resident
PCNA, prohibiting TLS and TS and promoting HDR (**Figure 6**, *Right*).

According to the model proposed in **Figure 6**, upon stalling of canonical replication at UVR-
induced lesions, the occupancy of the afflicted P/T junctions by PCNA, RPA, and Rad51•ATP
are anti-correlated temporally, leading to a transition from TLS/TS to HDR. This agrees with
results from recent iPOND studies of human cells treated with UVR. Here, as stalling prolonged
at UVR-induced lesions, the levels of PCNA, RPA, and pol η at P/T junctions of stalled
replications sites decreased over time as Rad51 levels increased at these sites[19]. Pol η is the TLS
pol primarily responsible for the bypass of UVR-induced lesions via TLS. These and other
observations led the authors to conclude a chronology of DDT pathways at UVR-stalled
replication sites where TLS is the favored, early DDT pathway elicited. The results from the
present study support this model.

The unreplicated ssDNA template sequences of UVR-stalled replication sites can reach 200 –
450 nts in length in human cells, which equates to RPA•ssDNA filaments comprised of ~6 - 15
RPA heterotrimers each in the 30 nucleotide binding mode[5-7,19,20,57,70]. Human Rad51 has an
intrinsic preference for nucleating at/near P/T junctions in the presence of RPA but does not
require access to these sites for nucleation and subsequent RPA/Rad51 exchange (**Figure 5**)[47 47].
Nevertheless, Rad51•ATP filaments must ultimately engage the P/T junctions via RPA/Rad51
exchange to eliminate the resident PCNA (**Figure 6**, *Right*). *In vivo*, other cellular factors likely

influence the temporal localization of RPA, PCNA, and Rad51 at the P/T junctions of stalled replication sites. For example, BRCA2, a molecular chaperone for Rad51 that accelerates filament nucleation, enhances Rad51•ATP deposition at dsDNA/ssDNA junctions in the presence of RPA[47,71]. Such activity at UVR-stalled replication sites may promote elimination of PCNA from the afflicted P/T junctions via inhibition of RFC (**Figure 4**). Furthermore, a recent study of human cells treated with hydroxyurea (HU) discovered that the PCNA unloader ATAD5 promotes interactions of Rad51 with P/T junctions of HU-stalled replication sites by catalytically unloading PCNA from these sites and subsequently recruiting Rad51 via direct protein•protein interactions. Hydroxyurea stalls canonical replication by depleting cellular dNTP pools[72]. At UVR-stalled replication sites, these ATAD5 activities may collectively eliminate PCNA from the afflicted P/T junctions by catalytically unloading PCNA from these sites and promoting Rad51-dependent inhibition of RFC (**Figure 4**). Also, recent studies have identified RadX, a negative regulator of Rad51 that promotes filament disassembly by dual mechanisms. RadX binds to and oligomerizes on the ends of growing Rad51•ATP filaments, blocking filament elongation. Also, RadX directly and selectively interacts with Rad51•ATP filaments, stimulating ATP hydrolysis and dismantling filaments[73,74]. At UVR-stalled replication sites, these RadX activities may stall and/or reverse RPA/Rad51 exchange at the afflicted P/T junctions, promoting occupancy of these sites by loaded PCNA (**Figure 4**). These and other hypotheses are currently under investigation.

SUPPLEMENTARY INFORMATION

Supplemental Information includes Supplementary Results, Supplementary Methods, and Supplementary Figures S1 – S7.

ACKNOWLEDGEMENTS. We would like to thank all members of the Hedglin and Antony labs for their efforts in reviewing/proofreading the current manuscript. This work was supported by funding from the National Institutes of Health to S.K. (F99CA274696), E.A. (R01 GM130756, R01 GM133967, R35 GM149320, and S10 OD030343) and M.H. (R35 GM147238-02),

CONFLICTS OF INTEREST

The authors declare that they have no conflicts of interest with the contents of this article.[1]

AUTHOR CONTRIBUTIONS

J.L.N., G.Y., K.G.P., R.L.D., S.P., and V.K. expressed, purified, and characterized all proteins. J.L.N., L.O.R., and G.Y. performed the experiments. M.H., J.L.N., and G.Y. designed the experiments. J.L.N., G.Y., M.H. and E.A. analyzed the data and wrote the paper.

REFERENCES

1   Hedglin, M., Kumar, R. & Benkovic, S. J. Replication clamps and clamp loaders. *Cold Spring Harb Perspect Biol* **5**, a010165 - a010183, doi:10.1101/cshperspect.a010165 (2013).

2   Hedglin, M., Perumal, S. K., Hu, Z. & Benkovic, S. Stepwise assembly of the human replicative polymerase holoenzyme. *Elife* **2**, e00278, 00271 - 00220, doi:10.7554/eLife.00278 (2013).

3   Yuzhakov, A., Kelman, Z., Hurwitz, J. & O'Donnell, M. Multiple competition reactions for
RPA order the assembly of the DNA polymerase delta holoenzyme. *EMBO J* **18**, 6189-6199,
doi:10.1093/emboj/18.21.6189 (1999).

4   Hedglin, M. & Benkovic, S. J. Replication Protein A Prohibits Diffusion of the PCNA
Sliding Clamp along Single-Stranded DNA. *Biochemistry* **56**, 1824-1835,
doi:10.1021/acs.biochem.6b01213 (2017).

5   Kim, C., Paulus, B. F. & Wold, M. S. Interactions of human replication protein A with
oligonucleotides. *Biochemistry* **33**, 14197-14206 (1994).

6   Kim, C., Snyder, R. O. & Wold, M. S. Binding properties of replication protein A from
human and yeast cells. *Mol Cell Biol* **12**, 3050-3059 (1992).

7   Kim, C. & Wold, M. S. Recombinant human replication protein A binds to polynucleotides
with low cooperativity. *Biochemistry* **34**, 2058-2064 (1995).

8   Kolpashchikov, D. M. *et al.* Polarity of human replication protein A binding to DNA.
Nucleic Acids Res **29**, 373-379, doi:10.1093/nar/29.2.373 (2001).

9   Pestryakov, P. E., Khlimankov, D. Y., Bochkareva, E., Bochkarev, A. & Lavrik, O. I. Human
replication protein A (RPA) binds a primer-template junction in the absence of its major
ssDNA-binding domains. *Nucleic Acids Res* **32**, 1894-1903, doi:10.1093/nar/gkh346 (2004).

10   Pestryakov, P. E. *et al.* Human replication protein A. The C-terminal RPA70 and the central
RPA32 domains are involved in the interactions with the 3'-end of a primer-template DNA. *J
Biol Chem* **278**, 17515-17524, doi:10.1074/jbc.M301265200 (2003).

11   Kolpashchikov, D. M. *et al.* Interaction of the p70 subunit of RPA with a DNA template
directs p32 to the 3'-end of nascent DNA. *FEBS Lett* **450**, 131-134, doi:10.1016/s0014-
5793(99)00484-6 (1999).

12   Kolinjivadi, A. M. *et al.* Smarcal1-Mediated Fork Reversal Triggers Mre11-Dependent
Degradation of Nascent DNA in the Absence of Brca2 and Stable Rad51 Nucleofilaments.
Mol Cell **67**, 867-881 e867, doi:10.1016/j.molcel.2017.07.001 (2017).

13   Lao, Y., Lee, C. G. & Wold, M. S. Replication protein A interactions with DNA. 2.
Characterization of double-stranded DNA-binding/helix-destabilization activities and the role
of the zinc-finger domain in DNA interactions. *Biochemistry* **38**, 3974-3984,
doi:10.1021/bi982371m (1999).

14   Binz, S. K., Dickson, A. M., Haring, S. J. & Wold, M. S. Functional assays for replication
protein A (RPA). *Methods Enzymol* **409**, 11-38, doi:10.1016/S0076-6879(05)09002-6
(2006).

15   Norris, J. L. *et al.* Replication protein A dynamically re-organizes on primer/template
junctions to permit DNA polymerase delta holoenzyme assembly and initiation of DNA
synthesis. *Nucleic Acids Res*, doi:10.1093/nar/gkae475 (2024).

16 Suarez, S. C., Toffton, S. M. & McCulloch, S. D. Biochemical analysis of DNA polymerase
eta fidelity in the presence of replication protein A. *PLoS One* **9**, e97382,
doi:10.1371/journal.pone.0097382 (2014).

17 Belan, O. *et al.* POLQ seals post-replicative ssDNA gaps to maintain genome stability in
BRCA-deficient cancer cells. *Mol Cell* **82**, 4664-4680 e4669,
doi:10.1016/j.molcel.2022.11.008 (2022).

18 Solomon, D. A., Cardoso, M. C. & Knudsen, E. S. Dynamic targeting of the replication
machinery to sites of DNA damage. *J Cell Biol* **166**, 455-463, doi:10.1083/jcb.200312048
(2004).

19 Benureau, Y. *et al.* Changes in the architecture and abundance of replication intermediates
delineate the chronology of DNA damage tolerance pathways at UV-stalled replication forks
in human cells. *Nucleic Acids Res* **50**, 9909-9929, doi:10.1093/nar/gkac746 (2022).

20 Despras, E., Daboussi, F., Hyrien, O., Marheineke, K. & Kannouche, P. L. ATR/Chk1
pathway is essential for resumption of DNA synthesis and cell survival in UV-irradiated XP
variant cells. *Hum Mol Genet* **19**, 1690-1701, doi:10.1093/hmg/ddq046 (2010).

21 Adar, S., Izhar, L., Hendel, A., Geacintov, N. & Livneh, Z. Repair of gaps opposite lesions
by homologous recombination in mammalian cells. *Nucleic Acids Res* **37**, 5737-5748,
doi:10.1093/nar/gkp632 (2009).

22 Izhar, L., Ziv, O., Cohen, I. S., Geacintov, N. E. & Livneh, Z. Genomic assay reveals
tolerance of DNA damage by both translesion DNA synthesis and homology-dependent
repair in mammalian cells. *Proc Natl Acad Sci U S A* **110**, E1462-1469,
doi:10.1073/pnas.1216894110 (2013).

23 Hedglin, M. & Benkovic, S. J. Regulation of Rad6/Rad18 Activity During DNA Damage
Tolerance. *Annu Rev Biophys* **44**, 207-228, doi:10.1146/annurev-biophys-060414-033841
(2015).

24 Hedglin, M. & Benkovic, S. J. Eukaryotic Translesion DNA Synthesis on the Leading and
Lagging Strands: Unique Detours around the Same Obstacle. *Chem Rev* **117**, 7857-7877,
doi:10.1021/acs.chemrev.7b00046 (2017).

25 Brun, J., Chiu, R. K., Wouters, B. G. & Gray, D. A. Regulation of PCNA polyubiquitination
in human cells. *BMC Res Notes* **3**, 85, doi:10.1186/1756-0500-3-85 (2010).

26 Ripley, B. M., Gildenberg, M. S. & Washington, M. T. Control of DNA Damage Bypass by
Ubiquitylation of PCNA. *Genes (Basel)* **11**, doi:10.3390/genes11020138 (2020).

27 Cipolla, L., Maffia, A., Bertoletti, F. & Sabbioneda, S. The Regulation of DNA Damage
Tolerance by Ubiquitin and Ubiquitin-Like Modifiers. *Front Genet* **7**, 105,
doi:10.3389/fgene.2016.00105 (2016).

28 Vujanovic, M. *et al.* Replication Fork Slowing and Reversal upon DNA Damage Require
PCNA Polyubiquitination and ZRANB3 DNA Translocase Activity. *Mol Cell* **67**, 882-890
e885, doi:10.1016/j.molcel.2017.08.010 (2017).

29 Masuda, Y. & Masutani, C. Spatiotemporal regulation of PCNA ubiquitination in damage
tolerance pathways. *Crit Rev Biochem Mol Biol* **54**, 418-442,
doi:10.1080/10409238.2019.1687420 (2019).

30 Ciccia, A. *et al.* Polyubiquitinated PCNA recruits the ZRANB3 translocase to maintain
genomic integrity after replication stress. *Mol Cell* **47**, 396-409,
doi:10.1016/j.molcel.2012.05.024 (2012).

31 Halder, S., Ranjha, L., Taglialatela, A., Ciccia, A. & Cejka, P. Strand annealing and motor
driven activities of SMARCAL1 and ZRANB3 are stimulated by RAD51 and the paralog
complex. *Nucleic Acids Res* **50**, 8008-8022, doi:10.1093/nar/gkac583 (2022).

32 Livneh, Z. *et al.* High-resolution genomic assays provide insight into the division of labor
between TLS and HDR in mammalian replication of damaged DNA. *DNA Repair (Amst)* **44**,
59-67, doi:10.1016/j.dnarep.2016.05.007 (2016).

33 Kochaniak, A. B. *et al.* Proliferating cell nuclear antigen uses two distinct modes to move
along DNA. *J Biol Chem* **284**, 17700-17710, doi:10.1074/jbc.M109.008706 (2009).

34 Shibahara, K. & Stillman, B. Replication-dependent marking of DNA by PCNA facilitates
CAF-1-coupled inheritance of chromatin. *Cell* **96**, 575-585 (1999).

35 Fennessy, R. T. & Owen-Hughes, T. Establishment of a promoter-based chromatin
architecture on recently replicated DNA can accommodate variable inter-nucleosome
spacing. *Nucleic Acids Res* **44**, 7189-7203, doi:10.1093/nar/gkw331 (2016).

36 Smith, D. J. & Whitehouse, I. Intrinsic coupling of lagging-strand synthesis to chromatin
assembly. *Nature* **483**, 434-438, doi:10.1038/nature10895 (2012).

37 Hedglin, M., Aitha, M. & Benkovic, S. J. Monitoring the Retention of Human Proliferating
Cell Nuclear Antigen at Primer/Template Junctions by Proteins That Bind Single-Stranded
DNA. *Biochemistry* **56**, 3415-3421, doi:10.1021/acs.biochem.7b00386 (2017).

38 Norris, J. L. & Hedglin, M. Direct, ensemble FRET approaches to monitor transient state
kinetics of human DNA polymerase delta holoenzyme assembly and initiation of DNA
synthesis. *Methods Enzymol* **705**, 271-309, doi:10.1016/bs.mie.2024.08.002 (2024).

39 Henricksen, L. A., Umbricht, C. B. & Wold, M. S. Recombinant replication protein A:
expression, complex formation, and functional characterization. *J Biol Chem* **269**, 11121-
11132 (1994).

40 Walsh, E., Wang, X., Lee, M. Y. & Eckert, K. A. Mechanism of replicative DNA polymerase
delta pausing and a potential role for DNA polymerase kappa in common fragile site
replication. *J Mol Biol* **425**, 232-243, doi:10.1016/j.jmb.2012.11.016 (2013).

41 Subramanyam, S. & Spies, M. Expression, Purification, and Biochemical Evaluation of
Human RAD51 Protein. *Methods Enzymol* **600**, 157-178, doi:10.1016/bs.mie.2017.11.011
(2018).

42 Yates, L. A. *et al.* A structural and dynamic model for the assembly of Replication Protein A
on single-stranded DNA. *Nat Commun* **9**, 5447, doi:10.1038/s41467-018-07883-7 (2018).

43 Li, M., Sengupta, B., Benkovic, S. J., Lee, T. H. & Hedglin, M. PCNA Monoubiquitination
Is Regulated by Diffusion of Rad6/Rad18 Complexes along RPA Filaments. *Biochemistry*,
doi:10.1021/acs.biochem.0c00849 (2020).

44 Ristic, D. *et al.* Human Rad51 filaments on double- and single-stranded DNA: correlating
regular and irregular forms with recombination function. *Nucleic Acids Res* **33**, 3292-3302,
doi:10.1093/nar/gki640 (2005).

45 Hoitsma, N. M. *et al.* Mechanistic insight into AP-endonuclease 1 cleavage of abasic sites at
stalled replication fork mimics. *Nucleic Acids Res*, doi:10.1093/nar/gkad481 (2023).

46 Candelli, A. *et al.* Visualization and quantification of nascent RAD51 filament formation at
single-monomer resolution. *Proc Natl Acad Sci U S A* **111**, 15090-15095,
doi:10.1073/pnas.1307824111 (2014).

47 Bell, J. C., Dombrowski, C. C., Plank, J. L., Jensen, R. B. & Kowalczykowski, S. C. BRCA2
chaperones RAD51 to single molecules of RPA-coated ssDNA. *Proc Natl Acad Sci U S A*
120, e2221971120, doi:10.1073/pnas.2221971120 (2023).

48 Hilario, J., Amitani, I., Baskin, R. J. & Kowalczykowski, S. C. Direct imaging of human
Rad51 nucleoprotein dynamics on individual DNA molecules. *Proc Natl Acad Sci U S A* **106**,
361-368, doi:10.1073/pnas.0811965106 (2009).

49 Srivastava, M. *et al.* Replisome Dynamics and Their Functional Relevance upon DNA
Damage through the PCNA Interactome. *Cell Rep* **25**, 3869-3883 e3864,
doi:10.1016/j.celrep.2018.11.099 (2018).

50 Hedglin, M., Pandey, B. & Benkovic, S. J. Stability of the human polymerase delta
holoenzyme and its implications in lagging strand DNA synthesis. *Proc Natl Acad Sci U S A*
113, E1777-1786, doi:10.1073/pnas.1523653113 (2016).

51 Hedglin, M., Pandey, B. & Benkovic, S. J. Characterization of human translesion DNA
synthesis across a UV-induced DNA lesion. *Elife* **5**, e19788, 19781 - 19718,
doi:10.7554/eLife.19788 (2016).

52 Tsurimoto, T. & Stillman, B. Replication factors required for SV40 DNA replication in vitro.
I. DNA structure-specific recognition of a primer-template junction by eukaryotic DNA
polymerases and their accessory proteins. *J Biol Chem* **266**, 1950-1960 (1991).

53 Zhou, Y., Meng, X., Zhang, S., Lee, E. Y. & Lee, M. Y. Characterization of human DNA
polymerase delta and its subassemblies reconstituted by expression in the MultiBac system.
PLoS One **7**, e39156, doi:10.1371/journal.pone.0039156 (2012).

54 Chea, J. *et al.* Spatiotemporal recruitment of human DNA polymerase delta to sites of UV
damage. *Cell Cycle* **11**, 2885-2895, doi:10.4161/cc.21280 (2012).

55 Tims, H. S. & Widom, J. Stopped-flow fluorescence resonance energy transfer for analysis of
nucleosome dynamics. *Methods* **41**, 296-303, doi:10.1016/j.ymeth.2007.01.001 (2007).

56 Lancey, C. *et al.* Structure of the processive human Pol delta holoenzyme. *Nat Commun* **11**,
1109, doi:10.1038/s41467-020-14898-6 (2020).

57 Zellweger, R. *et al.* Rad51-mediated replication fork reversal is a global response to
genotoxic treatments in human cells. *J Cell Biol* **208**, 563-579, doi:10.1083/jcb.201406099
(2015).

58 Li, M., Sengupta, B., Benkovic, S. J., Lee, T. H. & Hedglin, M. PCNA Monoubiquitination
Is Regulated by Diffusion of Rad6/Rad18 Complexes along RPA Filaments. *Biochemistry*
59, 4694-4702, doi:10.1021/acs.biochem.0c00849 (2020).

59 Hedglin, M., Aitha, M., Pedley, A. & Benkovic, S. J. Replication protein A dynamically
regulates monoubiquitination of proliferating cell nuclear antigen. *J Biol Chem* **294**, 5157-
5168, doi:10.1074/jbc.RA118.005297 (2019).

60 Kang, M. S. *et al.* Regulation of PCNA cycling on replicating DNA by RFC and RFC-like
complexes. *Nat Commun* **10**, 2420, doi:10.1038/s41467-019-10376-w (2019).

61 Kubota, T., Myung, K. & Donaldson, A. D. Is PCNA unloading the central function of the
Elg1/ATAD5 replication factor C-like complex? *Cell Cycle* **12**, 2570-2579,
doi:10.4161/cc.25626 (2013).

62 Lee, K. Y., Fu, H., Aladjem, M. I. & Myung, K. ATAD5 regulates the lifespan of DNA
replication factories by modulating PCNA level on the chromatin. *J Cell Biol* **200**, 31-44,
doi:10.1083/jcb.201206084 (2013).

63 Ryu, E. *et al.* Distinct Motifs in ATAD5 C-Terminal Domain Modulate PCNA Unloading
Process. *Cells* **11**, doi:10.3390/cells11111832 (2022).

64 Wang, F., He, Q., Yao, N. Y., O'Donnell, M. E. & Li, H. The human ATAD5 has evolved
unique structural elements to function exclusively as a PCNA unloader. *Nat Struct Mol Biol*
31, 1680-1691, doi:10.1038/s41594-024-01332-4 (2024).

65 Perumal, S. K., Xu, X., Yan, C., Ivanov, I. & Benkovic, S. J. Recognition of a Key Anchor
Residue by a Conserved Hydrophobic Pocket Ensures Subunit Interface Integrity in DNA
Clamps. *J Mol Biol* **431**, 2493-2510, doi:10.1016/j.jmb.2019.04.035 (2019).

66 Li, H., O'Donnell, M. & Kelch, B. Unexpected new insights into DNA clamp loaders:
Eukaryotic clamp loaders contain a second DNA site for recessed 5' ends that facilitates
repair and signals DNA damage: Eukaryotic clamp loaders contain a second DNA site for
recessed 5' ends that facilitates repair and signals DNA damage. *Bioessays* **44**, e2200154,
doi:10.1002/bies.202200154 (2022).

67 Piberger, A. L. *et al.* PrimPol-dependent single-stranded gap formation mediates homologous
recombination at bulky DNA adducts. *Nat Commun* **11**, 5863, doi:10.1038/s41467-020-
19570-7 (2020).

68 Mason, J. M., Chan, Y. L., Weichselbaum, R. W. & Bishop, D. K. Non-enzymatic roles of
human RAD51 at stalled replication forks. *Nat Commun* **10**, 4410, doi:10.1038/s41467-019-
12297-0 (2019).

69 Bhat, K. P. & Cortez, D. RPA and RAD51: fork reversal, fork protection, and genome
stability. *Nat Struct Mol Biol* **25**, 446-453, doi:10.1038/s41594-018-0075-z (2018).

70 Kang, Y. *et al.* Alteration of replication protein A binding mode on single-stranded DNA by
NSMF potentiates RPA phosphorylation by ATR kinase. *Nucleic Acids Res* **51**, 7936-7950,
doi:10.1093/nar/gkad543 (2023).

71 Belan, O. *et al.* Visualization of direct and diffusion-assisted RAD51 nucleation by full-
length human BRCA2 protein. *Mol Cell* **83**, 2925-2940 e2928,
doi:10.1016/j.molcel.2023.06.031 (2023).

72 Park, S. H. *et al.* ATAD5 promotes replication restart by regulating RAD51 and PCNA in
response to replication stress. *Nat Commun* **10**, 5718, doi:10.1038/s41467-019-13667-4
(2019).

73 Adolph, M. B. *et al.* RADX controls RAD51 filament dynamics to regulate replication fork
stability. *Mol Cell* **81**, 1074-1083 e1075, doi:10.1016/j.molcel.2020.12.036 (2021).

74 Balakrishnan, S. *et al.* Structure of RADX and mechanism for regulation of RAD51
nucleofilaments. *Proc Natl Acad Sci U S A* **121**, e2316491121,
doi:10.1073/pnas.2316491121 (2024).

Supplemental Information

TITLE: PCNA encircling primer/template junctions is eliminated by exchange of RPA for
Rad51: Implications for the interplay between human DNA Damage Tolerance pathways

Jessica L. Norris[1], Lindsey O. Rogers[1], Grace Young[1], Kara G. Pytko[1], Rachel L. Dannenberg[1],
Samuel Perreault[1], Vikas Kaushik[2], Edwin Antony[2], and Mark Hedglin[1,*]

[*]Corresponding author, to whom correspondence may be addressed: Email: muh218@psu.edu

Affiliations:

[1]The Pennsylvania State University, Department of Chemistry, University Park PA, 16802

[2]The Saint Louis University School of Medicine, Department of Biochemistry and Molecular
Biology, St. Louis MO, 63104

Supplementary Results

Rad51 has a strong, intrinsic preference for forming filaments on ssDNA at physiological ionic
strength. Human Rad51 binds both ssDNA and dsDNA to form right-handed helical
nucleoprotein filaments where each Rad51 monomer binds either 3 bp (for dsDNA) or 3 nt (for
ssDNA) [1-3]. However, in the absence of ATP hydrolysis, Rad51•ATP has an exceptionally
strong, intrinsic preference for forming filaments on ssDNA at physiological ionic strength (200
mM I) such that dsDNA binding is dramatically reduced, if observed at all. Binding of
Rad51•ATP to dsDNA is only significant at low ionic strength[4-6]. To confirm that the specificity
of Rad51•ATP for ssDNA at physiological ionic strength is achieved in the present studies we
carried out Rad51 titrations on a P/T DNA substrate (FRET P/T, **Figure S1**) in which the ssDNA
sequence is labeled with a Cy3 FRET donor and a Cy5 FRET acceptor. The former is located 3
nt downstream of the P/T junction and the latter is located 3 nt from the 5′ terminus of the
ssDNA. The FRET P/T DNA substrate is comprised of a 15 bp duplex region that can
accommodate 5 Rad51•ATP monomers and a 30 nt ssDNA region that can accommodate 10
Rad51•ATP monomers. In isolation, the ssDNA template of the FRET P/T DNA substrate
forms a collapsed, flexible structure, bringing the two cyanine fluorophores close together and
yielding a high E_{FRET} (**Figure S2A**). Stable interactions of Rad51•ATP with the ssDNA extends
the engaged sequence ~1.6-fold relative to the B-form, thereby increasing the Cy3–Cy5 distance
and reducing E_{FRET}[7]. Thus, only saturation of the ssDNA sequence at increased concentrations of
Rad51•ATP minimizes E_{FRET}. The titration data is plotted as a function of the ratio of the amount
Rad51 to the Total Amount Rad51-binding sites (i.e., dsDNA-binding sites + ssDNA-binding
sites). Under the conditions of the assay, E_{FRET} decreases until the ssDNA sequence is saturated
with Rad51•ATP, after which E_{FRET} remains constant at a minimal value. The fraction of Rad51
binding sites that are ssDNA is 0.66 (Rad51 ssDNA-binding sites/Total Rad51 binding sites =
10/15 = 0.667). Thus, if Rad51•ATP binds exclusively to ssDNA, the ssDNA sequence will be
saturated with Rad51•ATP and the E_{FRET} will minimize at an equivalence point of Rad51:Total
Rad51•ATP Binding Sites ~ 0.667. On the contrary, if Rad51•ATP engages both dsDNA- and
ssDNA-binding sites indiscriminately, the ssDNA sequence will be saturated with Rad51•ATP

and the E_{FRET} will minimize only when all Rad51 binding sites are saturated, i.e., at an

equivalence point of Rad51:Total Rad51•ATP Binding Sites ~ 1.0. At physiological ionic

strength (200 mM I), E_{FRET} decreases linearly and is minimized at an equivalence point of

Rad51:Total Rad51-binding sites = 0.673 $\pm$ 0.014 (**Figure S2B**). This indicates that binding of

Rad51•ATP to the ssDNA sequence of the FRET P/T DNA substrate is stoichiometric at 200

mM I, despite the presence of 15 bp of dsDNA. However, at low ionic strength (88 mM I), E_{FRET}

decreases linearly and is minimized at an equivalence point of essentially Rad51:ssDNA Rad51-

binding sites = 1.0 (0.981 $\pm$ 0.096, **Figure S2C**). This indicates that Rad51•ATP binds

indiscriminately to dsDNA and ssDNA at low ionic strength. Altogether, this agrees with the

results from previous studies and confirms that Rad51•ATP has an exceptionally strong, intrinsic

preference for forming filaments on ssDNA over dsDNA under the experimental conditions of

the current study (i.e., physiological ionic strength).

RFC•ATP complexes maintain loaded PCNA on all P/T DNA engaged by RPA. RPA engaged

with P/T junctions permit RFC•ATP complexes to instantly "re-load" non-catalytically unloaded

PCNA back onto P/T DNA such that the loss of PCNA is not observed over time[8-11]. Thus,

RFC•ATP complexes continuously maintain loaded PCNA on all P/T junctions engaged by

RPA. To confirm that this is established prior to the addition of Rad51•ATP complexes in the

experiments described in **Figures 2C** and **3C** in the main text, we continuously monitor

interactions of PCNA with P/T junctions engaged by RPA as reaction conditions progressively

evolve (**Figure S7A**). First, 5′ddPCy3/T DNA is pre-bound with Neutravidin and native RPA.

Next, Cy5-PCNA is added followed by RFC•ATP complexes and the fluorescence emission

intensities are monitored over time until PCNA loading reaches equilibrium where PCNA is

loaded and stabilized on all P/T junctions[8,9]. Under the conditions of the assay, RFC•ATP

engages the "front face" of a free PCNA (in solution), opens the clamp, and the resultant

complex, referred to as the loading complex, engages a P/T junction such that the "front face" of

PCNA is oriented towards the 3′ terminus of the primer [12]. Upon engaging a P/T junction, the

loading complex adopts an activated conformation in which ATP hydrolysis by RFC is

optimized. ATP hydrolysis by RFC simultaneously closes PCNA around the P/T junction and the

closed (i.e., loaded) PCNA is subsequently released onto the dsDNA region of the P/T junction.

The resultant RFC•ADP complex then releases into solution via dissociation from the resident

RPA and subsequently exchanges ADP for ATP, re-forming RFC•ATP [8,11-13]. In this scenario,

PCNA loading is biphasic. First, all steps up to and including release of loaded PCNA onto the

P/T junction are rate-limited by a kinetic step ($k_{obs\ inc\ 1}$) along the pathway that occurs prior to

and much slower than binding of loading complexes to P/T junctions. Second, loaded PCNA

repositions relatively slowly ($k_{obs\ inc\ 2}$) on the dsDNT region of the P/T junctions concurrent with

release of RFC•ADP complexes into solution[8,9]. Thus, $k_{obs\ inc\ 1}$ directly reports on the kinetics of

loading "free" PCNA onto P/T junctions engaged by RPA. Once PCNA loading has reached

equilibrium where PCNA is loaded and stabilized on all P/T junctions, unlabeled, native PCNA

is added in excess and the fluorescence emission intensities are monitored over time. Under these

conditions, the only pathway for unloading of Cy5-PCNA from P/T junctions is through

spontaneous opening of the Cy5-PCNA ring. Once Cy5-PCNA is unloaded from the Cy3-labeled

P/T DNA, re-loading of "free" Cy5-PCNA is prohibited due to the 70-fold excess of unlabeled,

native PCNA. Consequently, the observed FRET decreases. Here, the disappearance of FRET is

rate-limited by and directly reports on spontaneous opening of the PCNA ring.

Upon addition of RFC•ATP, I_{665} increases concomitantly with decreases in I_{563} after which
both fluorescence emission intensities stabilize and persist (**Figure S7B**, *Top*). These
synchronized, anti-correlated changes in I_{563} and I_{665} are indicative of the appearance and
increase in FRET (**Figure S7B**, *Bottom*). As expected, the increase in E_{FRET} observed upon
addition of RFC•ATP (**Figure S7B**, *Bottom*) is biphasic with $k_{obs\ inc\ 1} = 4.46 \pm 0.2$ (x 10^{-2}) s^{-1} and
$k_{obs\ inc\ 2} = 1.70 \pm 0.12$ (x 10^{-2}) s^{-1}. These rate constants are in excellent agreement with values
obtained in identical experiments from a recent study[9]. The increase in E_{FRET} observed upon
addition of RFC•ATP plateaus at values significantly above the E_{FRET} traces observed for no
interaction between Cy5-PCNA and the 5′ddPCy3/T•RPA complexes. At this point, Cy5-PCNA
is loaded and stabilized on all 5′ddPCy3/T•RPA complexes and RFC•ADP complexes have
released into solution and exchanged ADP for ATP. Upon addition of excess, unlabeled PCNA,
I_{665} decreases concomitantly with an increase in $I_{563,}$ after which both fluorescence emission
intensities stabilize and persist over time (**Figure S7B**, *Top*). These synchronized, anti-correlated
changes in I_{563} and I_{665} are indicative of a decrease in FRET (**Figure S7B**, *Bottom*). The E_{FRET}
decrease observed upon addition of excess, unlabeled PCNA bottoms out at E_{FRET} values
observed for no interaction between Cy5-PCNA and the 5′ddPCy3/T•RPA complexes (i.e., all
Cy5-PCNA remaining "free" in solution) indicating that all loaded Cy5-PCNA has non-
catalytically unloaded from all 5′ddPCy3/T•RPA complexes (**Figure S7B**, *Bottom*).
Furthermore, the observed decrease E_{FRET} is comprised of a single phase (i.e., monophasic) with
an observed rate constant [$k_{dec,obs} = 1.93 \pm 0.01$ (x 10^{-3}) s^{-1}] that is in good agreement with the
rate constant for spontaneous opening of the PCNA ring [$k_{open} = 1.25 \pm 0.32$ (x 10^{-3}) s^{-1}][14]. This
confirms that dissociation of PCNA from P/T junctions engaged by RPA is governed entirely by
spontaneous opening of the PCNA ring. Furthermore, the observed rate constant for RFC-
catalyzed loading of "free" PCNA on P/T junctions engaged by RPA ($k_{obs\ inc\ 1} = 4.46 \pm 0.2$ (x 10^{-2}) s^{-1})
is ~23-fold faster than the observed rate constant for non-catalytic unloading of PCNA
from P/T DNA ($k_{dec,obs} = 1.93 \pm 0.01$ (x 10^{-3}) s^{-1}). Hence, upon non-catalytic unloading of Cy5-
PCNA from P/T junctions engaged by RPA, RFC•ATP complexes instantly reload Cy5-PCNA
back onto the P/T junctions such that the loss of loaded Cy5-PCNA from 5′ddPCy3/T•RPA
complexes is not observed[8,11]. In other words, for the experiments described in **Figures 2** and **3**
in the main text, RFC•ATP complexes maintain loaded Cy5-PCNA on all 5′ddPCy3/T•RPA
complexes prior to the addition of Rad51•ATP complexes.
118
Supplementary Methods

Steady state FRET assays to monitor DNA-binding preference of Rad51. A solution containing
FRET P/T DNA (200 nM, **Figure S1**) and ATP was titrated with increasing concentrations of
Rad51•ATP and E_{FRET} at each concentration was calculated. Specifically, at each Rad51•ATP
concentration, I_{665} and I_{563} are monitored over time until both stabilize for at least 1 min. Within
this stable region, E_{FRET} values are calculated from the observed I_{665} and I_{563} values and averaged
to obtain the E_{FRET} value observed for a given concentration of Rad51•ATP.

Pre-steady state FRET assays to monitor PCNA loading and unloading. A solution containing
5′ddPCy3/T DNA (20 nM, **Figure S1**), neutravidin (80 nM homotetramer), and ATP is pre-
incubated with RPA (25 nM heterotrimer). Then, Cy5-PCNA (20 nM homotrimer) is added, the
resultant solution is transferred to a fluorometer cell, and the cell is placed in the instrument. I_{665}
and I_{563} are monitored over time until both stabilize for at least 1 min. Within this stable region,

E_{FRET} values are calculated from the observed I_{665} and I_{563} values and averaged to obtain the
E_{FRET} value observed prior to addition of RFC•ATP complexes. Next, pre-formed RFC•ATP (20
nM RFC heteropentamer, 1 mM ATP) is added, the resultant solution is mixed via pipetting, and
I_{665} and I_{563} are monitored beginning $\leq$ 10 s after the addition of RFC•ATP (dead time $\leq$ 10 s)
and continue to be until both stabilize for at least 1 min. Within this stable region, E_{FRET} values
are calculated from the observed I_{665} and I_{563} values and averaged to obtain the E_{FRET} value
observed prior to addition of unlabeled PCNA. Finally, unlabeled PCNA (1.4 µM homotrimer) is
added, the resultant solution is mixed by pipetting, and I_{665} and I_{563} are monitored beginning $\leq$ 10
s after the addition of unlabeled PCNA (dead time $\leq$ 10 s).

Supplemental Figures

Figure S1. P/T DNA substrates utilized in this study. The sequences and lengths of the dsDNA
and ssDNA regions are indicated. For the ddP/5′TCy3 and 5′ddPCy3/T DNA substrates, the size
of the dsDNA region (29 bp) is in agreement with the requirements for assembly of a PCNA ring
onto DNA by RFC[8,11,15]. The ssDNA regions of these substrates accommodate 1 RPA
heterotrimer or 11 Rad51 monomers[1,2,16-18]. RPA prevents loaded PCNA from sliding off the
ssDNA end of these substrates[8]. When pre-bound to neutravidin, the biotin attached to the 5′-end
of the primer strands of these substrates prevents loaded PCNA from sliding off the dsDNA
ends. Primers terminated at the 3′ end with a dideoxy C nucleotide cannot be extended by pol δ.
For the FRET P/T DNA substrate, the dsDNA (15 bp) and ssDNA (30 nt) regions accommodate
5 and 10 Rad51•ATP, respectively[1,2].

Figure S2. Rad51 has a strong, intrinsic preference for forming filaments on ssDNA over
dsDNA at physiological ionic strength (**A**) Schematic representation of the FRET experiment
utilizing a FRET P/T DNA substrate (**Figure S1**) and Rad51•ATP. The 30 nt ssDNA template
(**Figure S1**) is labeled with an internal 3′ Cy3 (FRET donor) 3 nt from the P/T junction and a
Cy5 (FRET acceptor) 3 nt from the 5′ terminus. In the absence of Rad51, the ssDNA template
forms a collapsed, flexible structure, bringing the two cyanine fluorophores close together and
yielding a high E_{FRET}. Saturation of the ssDNA template with Rad51•ATP extends the engaged
sequence ~1.6-fold relative to the B-form[7], thereby increasing the Cy3–Cy5 distance and
reducing E_{FRET}. (**B** - **C**) FRET data. 200 nM FRET P/T DNA is titrated with Rad51•ATP at
either physiological (200 mM I) or low ionic strength (88 mM I) and E_{FRET} values are monitored.
The observed E_{FRET} values are plotted (on the y-axes) as functions of the ratios of active Rad51
to Total Rad51-binding sites (on the x-axes). Each data point represents the mean ± S.E.M. of
three independent measurements. Data for each titration is fit to two segments lines; a linear
regression with a negative slope and a flat line. The equivalence point (indicated with standard
errors of the calculation) for each titration is calculated from the intersection of the two segment
lines. Titrations carried out at physiological (200 mM I) and low ionic strength (88 mM) are
displayed in panels **B** and **C**, respectively.

Figure S3. RPA/Rad51 exchange on P/T junctions devoid of PCNA. (**A**) Schematic
representation of the FRET pair and experiment to monitor RPA/Rad51 exchange on P/T
junctions devoid of PCNA. Experiments are performed exactly as described in the left panel of
Figure 2A in the main text except RFC and PCNA were omitted. (**B**) Data. Each trace is the
mean of at least three independent traces with the S.E.M. shown in grey. The time trajectories of
I_{563} (magenta) and I_{665} (cyan) are displayed in the top panel and the corresponding E_{FRET} (yellow)
is displayed in the bottom panel. The time at which Rad51•ATP is added is indicated by a red
arrow. Changes in I_{563} and I_{665} are indicated in the top panel by magenta and cyan arrows,
respectively. For observation, the I_{563} and I_{665}, and the corresponding E_{FRET} values observed prior
to the addition of Rad51•ATP complexes are each fit to dashed flat lines that are extrapolated to
the axis limits. The predicted E_{FRET} trace (pink) for no interaction between RPA-OBA-Cy5 and
the ddP/5′TCy3 DNA is displayed in the bottom panel, offset to the time after Rad51•ATP is
added, and fit to a dashed flat line that is extrapolated to the axis limits. The rate constant for the
E_{FRET} decrease ($k_{obs\ dec}$) observed after the addition of Rad51•ATP complexes is reported in
Table 1 in the main text.

Figure S4. PCNA unloads from P/T junctions during RPA/Rad51 exchange and after
RPA/Rad51 exchange has been completed. E_{FRET} traces from **Figures 2B** (RPA/Rad51
exchange) and **2C** (PCNA unloading) in the main text were each normalized to their respective
ranges and the results plotted as a function of time (after Rad51•ATP is added) with the S.E.M.
shown in grey. The normalized trace for RPA/Rad51 exchange (shown in brown) is fit to one-
step, irreversible kinetic model that is extrapolated to the axis limits. The normalized trace for
PCNA unloading (shown in orange), i.e. "% P/T Junctions Encircled by PCNA," is fit to a two-
step, irreversible kinetic model that is extrapolated to the axis limits. An extended time course is
shown to highlight the extrapolated kinetics fits and lifetimes for complete unloading of PCNA
from P/T junctions and complete RPA/Rad51 exchange on P/T junctions. For observation, the
time at which RPA/Rad51 exchange is completed is indicated by a vertical dashed line.

Figure S5. Pol δ holoenzymes do not stabilize loaded PCNA on P/T junctions during
RPA/Rad51 exchange. E_{FRET} traces for unloading of PCNA from P/T junctions in the absence
(**Figure 2C**) and presence of pol δ (**Figure 3C**) in the main text were each normalized to their
respective ranges and the results plotted as a function of time (after Rad51●ATP complexes are
added) with the S.E.M. shown in grey. The normalized traces for PCNA unloading, i.e., "% P/T
Junctions Encircled by PCNA," in the absence (- pol δ) and presence of pol δ (+ pol δ) are fit to
a two-step, irreversible kinetic model and a triple exponential, irreversible kinetic model,
respectively. Each kinetic model is extrapolated to the axis limits. A dashed flat line denotes 0%
P/T junctions encircled by PCNA.

Figure S6. PCNA pre-engaged by pol δ in holoenzymes unloads from P/T junctions during
RPA/Rad51 exchange and after RPA/Rad51 exchange has been completed. E_{FRET} traces from
Figures 3B (RPA/Rad51 exchange) and **3C** (PCNA Unloading) in the main text were each
normalized to their respective ranges and the results plotted as function of time (after
Rad51•ATP complexes are added) with the S.E.M. shown in grey. The normalized trace for
RPA/Rad51 exchange (shown in brown) is fit to a one-step, irreversible kinetic model that is
extrapolated to the axis limits. The normalized trace for PCNA unloading (shown in orange), i.e.,
"% P/T Junctions Encircled by PCNA," is fit to a triple exponential decline that is extrapolated
to the axis limits. An extended time course is shown to highlight the extrapolated kinetics fits
and lifetimes for complete unloading of PCNA from P/T junctions and complete RPA/Rad51
exchange on P/T junctions. For observation, the time at which RPA/Rad51 exchange is
completed is indicated by a vertical dashed line.

Figure S7. Dynamics of PCNA encircling P/T junctions engaged by RPA. (**A**) Schematic
representation of the FRET pair and experiment to monitor the dynamics of PCNA encircling
P/T junctions that are engaged by RPA. (**B**) Data. Each trace is the mean of at least three
independent traces with the S.E.M. shown in grey. The time trajectories of I_{563} (magenta) and I_{665}
(cyan) are displayed in the top panel and the corresponding E_{FRET} (yellow) is displayed in the
bottom panel. The times at which RFC•ATP and unlabeled PCNA are added are indicated by red
arrows. Changes in I_{563} and I_{665} are indicated in the top panel by magenta and cyan arrows,
respectively. For observation, the I_{563} and I_{665}, and the corresponding E_{FRET} values observed prior
to the addition RFC•ATP are each fit to dashed flat lines that are extrapolated to the axis limits.
Also, dashed flat lines are drawn to highlight the plateau I values observed after the addition of
RFC•ATP. The increase in E_{FRET} observed upon the addition of RFC•ATP complexes is fit to a
double exponential rise, yielding rate constants $k_{obs\ inc\ 1} = 4.46 \pm 0.2$ (x 10^{-2}) s^{-1} and $k_{obs\ inc\ 2} =$
1.70 ± 0.12 (x 10^{-2}) s^{-1}. The decrease in E_{FRET} observed upon the addition of unlabeled PCNA is
fit to a single exponential decline, yielding the rate constant $k_{dec,obs} = 1.93 \pm 0.01$ (x 10^{-3}) s^{-1}. The
data for the PCNA loading portion of the plots is identical to that shown in **Figure 4A** in the
main text.

References

334
1   Erratum for the Report "Base triplet stepping by the Rad51/RecA family of recombinases" by
J. Y. Lee, T. Terakawa, Z. Qi, J. B. Steinfeld, S. Redding, Y. Kwon, W. A. Gaines, W. Zhao,
P. Sung, E. C. Greene. *Science* **350**, aad6940, doi:10.1126/science.aad6940 (2015).
2   Lee, J. Y. *et al.* DNA RECOMBINATION. Base triplet stepping by the Rad51/RecA family
of recombinases. *Science* **349**, 977-981, doi:10.1126/science.aab2666 (2015).
3   Bonilla, B., Hengel, S. R., Grundy, M. K. & Bernstein, K. A. RAD51 Gene Family Structure
and Function. *Annu Rev Genet* **54**, 25-46, doi:10.1146/annurev-genet-021920-092410 (2020).
4   Candelli, A. *et al.* Visualization and quantification of nascent RAD51 filament formation at
single-monomer resolution. *Proc Natl Acad Sci U S A* **111**, 15090-15095,
doi:10.1073/pnas.1307824111 (2014).
5   Bell, J. C., Dombrowski, C. C., Plank, J. L., Jensen, R. B. & Kowalczykowski, S. C. BRCA2
chaperones RAD51 to single molecules of RPA-coated ssDNA. *Proc Natl Acad Sci U S A*
120, e2221971120, doi:10.1073/pnas.2221971120 (2023).
6   Hilario, J., Amitani, I., Baskin, R. J. & Kowalczykowski, S. C. Direct imaging of human
Rad51 nucleoprotein dynamics on individual DNA molecules. *Proc Natl Acad Sci U S A* **106**,
361-368, doi:10.1073/pnas.0811965106 (2009).
7   Subramanyam, S. & Spies, M. Expression, Purification, and Biochemical Evaluation of
Human RAD51 Protein. *Methods Enzymol* **600**, 157-178, doi:10.1016/bs.mie.2017.11.011
(2018).
8   Hedglin, M. & Benkovic, S. J. Replication Protein A Prohibits Diffusion of the PCNA
Sliding Clamp along Single-Stranded DNA. *Biochemistry* **56**, 1824-1835,
doi:10.1021/acs.biochem.6b01213 (2017).
9   Norris, J. L. *et al.* Replication protein A dynamically re-organizes on primer/template
junctions to permit DNA polymerase delta holoenzyme assembly and initiation of DNA
synthesis. *Nucleic Acids Res*, doi:10.1093/nar/gkae475 (2024).
10   Hedglin, M. & Benkovic, S. J. Eukaryotic Translesion DNA Synthesis on the Leading and
Lagging Strands: Unique Detours around the Same Obstacle. *Chem Rev* **117**, 7857-7877,
doi:10.1021/acs.chemrev.7b00046 (2017).
11   Hedglin, M., Perumal, S. K., Hu, Z. & Benkovic, S. Stepwise assembly of the human
replicative polymerase holoenzyme. *Elife* **2**, e00278, 00271 - 00220,
doi:10.7554/eLife.00278 (2013).
12   Hedglin, M., Kumar, R. & Benkovic, S. J. Replication clamps and clamp loaders. *Cold*
Spring Harb Perspect Biol **5**, a010165 - a010183, doi:10.1101/cshperspect.a010165 (2013).
13   Yuzhakov, A., Kelman, Z., Hurwitz, J. & O'Donnell, M. Multiple competition reactions for
RPA order the assembly of the DNA polymerase delta holoenzyme. *EMBO J* **18**, 6189-6199,
doi:10.1093/emboj/18.21.6189 (1999).
14   Perumal, S. K., Xu, X., Yan, C., Ivanov, I. & Benkovic, S. J. Recognition of a Key Anchor
Residue by a Conserved Hydrophobic Pocket Ensures Subunit Interface Integrity in DNA
Clamps. *J Mol Biol* **431**, 2493-2510, doi:10.1016/j.jmb.2019.04.035 (2019).
15   Hedglin, M., Aitha, M. & Benkovic, S. J. Monitoring the Retention of Human Proliferating
Cell Nuclear Antigen at Primer/Template Junctions by Proteins That Bind Single-Stranded
DNA. *Biochemistry* **56**, 3415-3421, doi:10.1021/acs.biochem.7b00386 (2017).
16   Kim, C., Paulus, B. F. & Wold, M. S. Interactions of human replication protein A with
oligonucleotides. *Biochemistry* **33**, 14197-14206 (1994).

17 Kim, C., Snyder, R. O. & Wold, M. S. Binding properties of replication protein A from
human and yeast cells. *Mol Cell Biol* **12**, 3050-3059 (1992).
18 Kim, C. & Wold, M. S. Recombinant human replication protein A binds to polynucleotides
with low cooperativity. *Biochemistry* **34**, 2058-2064 (1995).