Genome-wide dynamic nascent transcript profiles reveal that most paused RNA polymerases terminate

Rudradeep Mukherjee[a] **and Michael J. Guertin**[a,b] ✉

[a]Center for Cell Analysis and Modeling, University of Connecticut Health Center, Farmington, CT, United States of America
[b]Department of Genetics and Genome Sciences, University of Connecticut Health Center, Farmington, CT, United States of America

We present a simple model for analyzing and interpreting data from kinetic experiments that measure engaged RNA polymerase occupancy. The framework represents the densities of nascent transcripts within the pause region and the gene body as steady-state values determined by four key transcriptional processes: initiation, pause release, premature termination, and elongation. We validate the model's predictions using data from experiments that rapidly inhibit initiation and pause release. The model successfully classified factors based on the steps in early transcription that they regulate, confirming TBP and ZNF143 as initiation factors and HSF and GR as pause release factors. We found that most paused polymerases terminate and paused polymerases are short-lived with half lives less than a minute. We make this model available as software to serve as a quantitative tool for determining the kinetic mechanisms of transcriptional regulation.

Introduction

Gene expression regulation plays a crucial role in determining cell fate during development and influences how cells respond to environmental signals. Early events in the transcription cycle are major determinants of a gene's transcriptional output. The earliest events of transcription include the formation of a preinitiation complex, promoter opening, rounds of abortive initiation, promoter escape, and ultimately pausing of RNA polymerases 50–100 bases downstream from the transcription start site (1–3). We refer to these early events collectively as *initiation* throughout this study because most molecular measurements in cells cannot distinguish between these transient and unstable states. After initiation, the paused RNA polymerases may either proceed into productive elongation or prematurely terminate and disassociate from the DNA (4–6). Regulation of these steps is crucial for the precise control of transcription.

Previous studies have used imaging or high-throughput genomics assays to investigate and model the kinetics of transcription. Imaging studies track the generation of transcripts linked to fluorescent reporters and use models to interpret the time series of fluorescence signal (7, 8). Others use fluorescence recovery after photobleaching or photoactivatable molecules to determine residence times of RNA polymerases in different stages of transcription (9, 10). Genome-wide nuclear run-on assays, which precisely measure the position and orientation of transcriptionally engaged RNA polymerases, have been combined with inhibitors of initiation and pause release to dissect contribu-

tions of each regulated step (11). A recent study employed generative probabilistic modeling of nascent transcription datasets to estimate rates of initiation and pause release (12), highlighting the rate-limiting nature of RNA polymerase escape from the pause region. Despite the numerous studies and diverse methodologies employed, there is no consensus on the absolute and relative rates of these key regulatory steps.

Transcription factors (TFs) interact with various cofactors to coordinate the recruitment and activity of RNA polymerases (13–25). TFs bind to sequence motifs and assemble complexes that facilitate the loading and progression of RNA polymerases (26–29). Understanding the molecular functions of transcription factors, specifically how they contribute to the regulation of RNA polymerase activity at these various steps in early elongation, is crucial to understanding the complex mechanisms that drive gene expression (30, 31). Many foundational studies have leveraged rapidly inducible transcription factors to reveal their regulatory roles by tracking changes in RNA polymerase profiles at target genes (31–36). New chemical and physical methods can more broadly perturb TF activity or degrade TFs, significantly expanding the range of factors whose molecular functions can be studied (37–42). Determining the specific role of each transcription factor and their combinatorial effects on transcription regulation requires methods to acutely activate or perturb TF activity, quantify changes in RNA polymerase activity, and a modeling framework to interpret the data.

We present a Compartment Model framework, which represents RNA polymerase densities in the pause region and gene body as steady-state values determined by four key transcriptional processes: initiation, pause release, premature termination, and elongation (Fig. 1A). Unless stated otherwise, we define *initiation* as the sequence of events spanning from the incorporation of the first RNA base to the transition into the pause site. For each gene, the model estimates the magnitudes of pause release and initiation rates, as well as changes in these rates for all genes after experimental perturbation (Fig. 1B-C). We describe the framework and validate the model's predictions through analysis of publicly available experimental data that inhibited either initiation or pause release. Analysis of kinetic triptolide experimental datasets, which inhibit initiation, reveals that premature termination occurs at a faster rate than pause release. We further demonstrate that pause release serves as a critical regulatory step that significantly influences transcription output, but only when the premature termination rate of

Fig. 1. Kinetic modeling of transcription data determines the effect of treatments and perturbations. A) A gene has two compartments in the model–the pause window (p) and the gene body (b). RNA polymerase enters the pause region with an initiation rate constant k_{init}, prematurely terminates with a rate constant k_{pre}, escapes from pause region with a release rate k_{rel}, and proceeds through the gene body based on the elongation rate k_{elong}. Herein, we assume that k_{pre} and k_{elong} remain constant for a gene between conditions. B) The ideal input data for the model is generated from genome-wide nascent transcriptome (PRO-seq) experiments upon acute treatment with a perturbation or stimulus. Standard methods identify differentially expressed genes and the normalized data provides densities of reads in pause and gene body compartments. The model uses these densities to infer initiation and pause release rates for input genes. C) The regulatory effects on pause release and initiation, induced by general perturbations or the acute activation/inhibition of specific transcription factors, are determined by the rate changes explained by the model.

paused polymerases exceeds the release rate of paused polymerases into the gene body. We analyze publicly available transcription factor activation and perturbation data to confirm that ZNF143 and TBP predominantly regulate initiation, while the glucocorticoid receptor (GR) and *Drosophila* heat shock factor (HSF) regulate pause release. The model is available as a command-line tool that processes relative pause and gene body RNA polymerase densities from two experimental conditions as inputs and provides estimates of initiation rates, pause release rates, premature termination rates, and absolute RNA polymerase densities in pause regions and gene bodies.

Materials & Methods

Processing of nascent transcript datasets.
We processed six datasets spanning across a broad range of treatments - triptolide and flavopiridol in mouse v6.5 cells (11), degradation of TBP in HAP1 cells (43) and ZNF143 in HEK293T cells (44), dexamethasone treatment in A549 cells (45) and heat shocked *Drosophila* S2 cells (33). We downloaded the datasets from Gene Expres-

sion Omnibus (Table S1) and analyzed the raw data using a general pipeline (46, 47). We normalized signal within datasets using `sizeFactors` from DESeq2 (48). Since triptolide (12.5 min) and flavopiridol (25 min) treatments cause universal repression, we normalized these datasets by using read counts from gene regions > 50 kb for triptolide (> 100 kb for flavopiridol) as *controlGenes* in DESeq2 (48). We chose these gene regions because an elongation rate of 4kb/min is beyond the higher end of contemporary estimates of elongation rates (49). The body length of genes within these data sets were constrained to 25 kb and 50 kb to capture the region where RNA polymerase clears out upon treatment. We provide complete processing steps in the accompanying analysis vignette. We defined Heat Shock Factor (HSF) activated genes in S2 cells (33) by first aligning all data to an *Hsp70* consensus sequence. We mapped the remaining unmapped reads to the `dm6` genome, counted reads within genes, and normalized the replicates using DESeq2 `sizeFactors` (48). We only considered the 249 activated genes selected from Table S2 in Duarte et. al., (33) because these genes were carefully curated for

read-through transcription. We considered 65 heat shock activated genes (FDR < 0.001) that are dependent upon the presence of HSF.

Hsp70 consensus sequence.

We generated the *Hsp70* consensus gene sequence by combining the *Hsp70* sequences from the BDGP6.46 (release 112) reference. We used the start and end coordinates of six Hsp70 genes (*Hsp70Aa/Hsp70Ab*, *Hsp70Ba*, *Hsp70Bb*, *Hsp70Bbb*, *Hsp70Bc*) and included an additional 50 bases upstream of the reference start site. Using these coordinates, we extracted the six Hsp70 sequences from the reference genome with `bedtools getfasta`. We then created an alignment file for these sequences using `clustalw` (50). The consensus sequence was derived by using Biopython's `dumb_consensus()` function (51) to the alignment file with a threshold of 0.6.

Estimation of Pol II densities at pause region and gene body region.

To define the pause region of a gene, we searched for a 50 bp window with the maximum summed PRO-seq signal within a 200 bp region (or 300 bp for GRO-seq) downstream from the most prominent transcription start site (TSS) as defined by `primaryTranscriptAnnotation` (52). This 50 bp window is designated the *pause region*. The pause density or pause sum is the summed signal across all base pairs in this window. We sum the reads in pause region to accommodate signals from RNA polymerase pausing at any arbitrary position within the pause window. We define the gene body as the region 500 bp from end of the pause window to the inferred transcription termination site (52) or the distance an initiated polymerase can travel during the time of treatment if RNA polymerase elongates at a rate of 2kb/min, whichever is shorter. We define the body density as the mean signal in this window.

The Compartment Model.

We describe the time evolution of pause and body densities in the following manner:

$$\frac{dp}{dt} = k_{init} - (k_{pre} + k_{rel}) \cdot p \tag{1}$$

$$\frac{db}{dt} = k_{rel} \cdot p - k_{elong} \cdot b \tag{2}$$

In the above equations, t is time, p is the density in pause region, b is mean density in gene body and k_{init}, k_{rel}, k_{pre}, k_{elong} are rates of initiation, pause release, premature termination, and elongation, respectively. The measurements of pause sum and body densities from sequencing data are assumed to be steady-state values. The steady state assumption is reasonable because the measured rates of these various processes are much faster than the experimental treatment times. At steady-state, we have:

$$\frac{dp}{dt} = 0 \implies p = \frac{k_{init}}{k_{pre} + k_{rel}} \tag{3}$$

$$\frac{dp}{dt} = 0 \implies b = \frac{k_{rel}}{k_{elong}} \cdot p \tag{4}$$

For each gene, we consider that initiation (k_{init}) and pause release (k_{rel}) rates are regulated between treatments, while elongation (k_{elong}) and premature termination (k_{pre}) rates remain constant. Although the range of elongation rates for genes tend to be 1.5-4 kb/min (11, 34, 53–55), we cannot accurately estimate average elongation rates for all genes for each condition. However, we make this assumption of constant elongation rate because RNA polymerase signal within gene bodies correlate well with accumulated RNA-seq signal (56). If elongation rate increased during gene activation, then RNA polymerase density would decrease as mRNA output increases.

Considering two conditions under study as "control" and "treatment", we have the following relations for change in initiation and pause release rates:

$$
\begin{aligned}
fold\ change(k_{init}) &= \frac{k_{init}^{treatment}}{k_{init}^{control}} \\
&= \frac{p^{treatment}}{p^{control}} \cdot \frac{k_{pre} + k_{rel}^{treatment}}{k_{pre} + k_{rel}^{control}}
\end{aligned} \tag{5}
$$

$$
\begin{aligned}
fold\ change(k_{rel}) &= \frac{k_{rel}^{treatment}}{k_{rel}^{control}} \\
&= \frac{b^{treatment}}{b^{control}} \cdot \frac{p^{control}}{p^{treatment}} \\
&= \frac{fold\ change(b)}{fold\ change(p)}
\end{aligned} \tag{6}
$$

In Eq. 5, the fraction $\frac{k_{pre} + k_{rel}^{treatment}}{k_{pre} + k_{rel}^{control}}$ varies between two values depending upon relative values of premature termination rate (k_{pre}) and pause release rate (k_{rel}):

$$k_{pre} >> k_{rel} \implies \frac{k_{pre} + k_{rel}^{treatment}}{k_{pre} + k_{rel}^{control}} \to 1 \tag{7}$$

$$k_{pre} << k_{rel} \implies \frac{k_{pre} + k_{rel}^{treatment}}{k_{pre} + k_{rel}^{control}} \to \frac{k_{rel}^{treatment}}{k_{rel}^{control}} \tag{8}$$

From Eq. 5-8, fold change in initiation rate (k_{init}) for a gene can vary between fold change in pause density and the fold change in body density:

$$k_{pre} >> k_{rel} \implies fold\ change(k_{init}) \to \frac{p^{treatment}}{p^{control}} \tag{9}$$

$$k_{pre} << k_{rel} \implies fold\ change(k_{init})$$
$$\rightarrow \frac{k_{rel}^{treatment}}{k_{rel}^{control}} \cdot fold\ change(p) \quad \textbf{(10)}$$
$$= fold\ change(b)$$

Further, from Eq. 6, the fold change in pause release is an inverse ratio of change in "pausing index" (the ratio of pause density to body density.) As accurate measures of genic elongation rate become available, we can calculate the change in pause release rate by dividing the change in elongation rate by the change in pausing index (Eq. 4). Note that the normalization or scaling of read signals to estimate steady-state p and b values does not affect the fold change in pause release and bounds on changes in initiation rate. Consequently, the inference of the regulatory effects of a perturbation on initiation and pause release does not depend on the scaling of pause and body densities. For absolute values of rates we derive the following from Eq. 3 & 4:

$$k_{rel} = k_{elong} \cdot \frac{b}{p} \quad \textbf{(11)}$$

$$k_{init} = p \cdot (k_{pre} + k_{rel})$$
$$= p \cdot (k_{pre} + k_{elong} \cdot \frac{b}{p}) \quad \textbf{(12)}$$
$$= p \cdot k_{pre} + k_{elong} \cdot b$$

From Eq. (11), the absolute value of pause release rate (k_{rel}) for a gene depends upon elongation rate (usually between 30-60 bp/sec) and the inverse of the pause to body density ratio. Therefore, the value of pause release rate constant (k_{rel}) is also not dependent upon normalization or scaling criteria. From Eq. (12), the initiation rate (k_{init}) depends upon the absolute values of pause and body densities (i.e. the number of RNA polymerase in each compartment). Also, the amount of polymerases releasing into gene body, the effective release rate ($k_{rel} \times p$), provides a way to calculate spacing between polymerases on the gene body in base pairs:

$$Spacing = \frac{k_{elong}}{k_{rel} \times p} \quad \textbf{(13)}$$
$$= \frac{1}{b}$$

where, elongation rate (k_{elong}) is in units of $\frac{base\ pairs}{time}$ and pause release (k_{rel}) has the same time unit. From Eq. 11, the magnitude of occupancy is also equal to reciprocal of body density (b).

Further, a relation between premature termination rate (k_{pre}) and pause release rate ($k_{rel}^{control}$) can be derived from Eq. 5:

$$\frac{k_{pre}}{k_{rel}^{control}} = \frac{fold\ change(k_{rel}) - X}{X - 1} \quad \textbf{(14)}$$

where $X = \frac{fold\ change(k_{init})}{fold\ change(p)}$ with p representing the pause density for a gene. Since values and changes in pause release rate can be estimated using elongation rate, pause

sums and gene body densities (Eq. 6 & Eq. 11), the rates of premature termination can be assigned to each gene promoter using the relation in Eq. 14. Further, the ratio is not dependent on the scaling criteria used on pause and body densities.

The ratio in Eq. 14 becomes zero or undefined when numerator or denominator becomes zero:

$$X = fold\ change(k_{rel})$$
$$=> \frac{fold\ change(k_{init})}{fold\ change(p)} = fold\ change(k_{rel})$$
$$=> fold\ change(k_{init}) \quad \textbf{(15)}$$
$$= fold\ change(k_{rel}) \cdot fold\ change(p)$$
$$=> fold\ change(k_{init}) = fold\ change(b)$$

While Eq. 15 provides the condition where $\frac{k_{pre}}{k_{rel}^{control}}$ becomes zero, the ratio becomes undefined when denominator in Eq. 14 becomes zero:

$$X = 1 => fold\ change(k_{init}) = fold\ change(p) \quad \textbf{(16)}$$

The values in Eqs. 15-16, which render $\frac{k_{pre}}{k_{rel}^{control}}$ as zero or undefined, represent the extreme bounds on changes in initiation rate as described in Eqs. 9-10. However, transcription is expected to operate within a regime where the rates of premature termination (k_{pre}) and pause release (k_{rel}) yield a non-zero ratio for $\frac{k_{pre}}{k_{rel}^{control}}$. Consequently, the extreme cases for the relative values of k_{pre} and k_{rel}, as described in Eqs. 9-10, are not considered.

Pause and body densities as occupancy values.

To infer rates of initiation at different gene promoters we first transformed pause and gene body densities into absolute occupancy values. Occupancy refers to the number of RNA polymerases in the pause region or gene body per gene length. We assume that a fully occupied pause region will contain only one RNA Polymerase and that some genes would approach full occupancy of the pause region. In an ideal dataset without aneuploidy, gene duplications, PCR biases, and sequence mapping ambiguities, the ranked pause sums would saturate to a maximum occupancy. However, over all the datasets examined, the observed ranked pause sums exhibit a sudden uptick at the highest pause densities (Fig. S1A-F). Manual inspection of genes with high pause signals revealed multicopy genes located within non-uniquely mappable regions and genes characterized by highly variable read densities in the gene body. We decided to estimate a fully occupied pause region by fitting ranked pause densities of all expressed genes to a hyperbolic tangent function ($y = A \cdot tanh(B \cdot x) + C$) where density of a fully occupied pause region is assumed at $saturation = A + C$. We fitted the hyperbolic tangent function to log-transformed ranked pause sums using Scipy's `optimize.curve_fit()` function (57).

We reasoned that a saturation value should be robust to

amount of input data used to fit the curve, and be minimally influenced by the observed exponential increase in pause sums. For datasets with Unique Molecular Identifiers (UMIs) from which PCR duplicated reads could be confidently removed, the saturation values from incremental fits to pause densities (70th to 100th percentile) showed a modest increase by 1.5- to 4-fold (Fig. S1D-F & Fig. S2A). In contrast, the saturation values from incremental fits for datasets without UMIs varied from 23-fold to 600-fold (Fig. S1A-C & Fig. S2A). The saturation fits for datasets with UMIs tended to stabilize and reach 90th percentile of the ranked paused sums (Fig. S1D-F). Since the saturation values for datasets with UMIs were robust, we used the saturation values obtained from fits to all genes as the maximum occupancy signal. To remain consistent and mitigate the influence of the exponential increase in saturation values, we considered the pause sum at 90th percentile as maximum occupancy signal in datasets without UMIs. Using the maximum occupancy value, we scaled all pause and gene body densities from 0 to 1 to estimate initiation rates (Eq. 12). The assumption that the pause sum saturates at the 90th percentile of ranked pause sums in datasets without UMIs may not be accurate, as the pause sum at the 90th percentile is lower than the pause signals observed for heat shock activated genes (such as *Hsp70*) in *Drosophila* (Fig. S2B&C). Although our initiation rate calculations are based on these assumptions, more accurate estimates can be obtained by calibrating saturation values using datasets with known RNA polymerase densities in specific genomic regions.

Vignette and Model Availability.
A Python package is currently installable from PyPi: `https://pypi.org/project/compartmentModel` using the command: `python3 -m pip install compartmentModel`.

The details of installation, running the model and processing of sequencing data is available this vignette: `https://github.com/guertinlab/compartment_model/tree/master/compModel_vignette/compModel_vignette`.

The source files for building the package, figures, and UCSC Genome Browser track hubs are available: `https://github.com/guertinlab/compartment_model`. The analysis vignette is available here: `https://guertinlab.github.io/compartment_model/compModel_vignette/compModel_vignette/compartmentModel_vignette.html`

Results

Direct inhibition of pause release and initiation verifies the model predictions.
The compartment model we developed interprets the mechanistic steps of RNA polymerase initiation, pause release, premature termination, and elongation as rate constants that explain observed changes in nascent transcription after acute

perturbation and treatments. To validate the model predictions for changes in pause release rates (Eq. 6), we analyzed GRO-seq data following the inhibition of pause release with flavopiridol (11). Flavopiridol is a kinase inhibitor that targets CDK9 (58, 59), a key component of the positive transcription elongation factor b (P-TEFb) complex. P-TEFb promotes the release of paused RNA polymerase II into productive elongation, partially through the phosphorylation of the RPB1 C-terminal domain (4, 58, 60). Treatment with flavopiridol (0.3 μM) for 25 minutes results in a decrease of the pause release rate for ~82% of repressed genes, with a median rate decrease of ~57% (Fig. 2A, Fig. S3, & Fig. S4A). Additionally, we observed a modest reduction in the initiation rate, with a median decrease ranging from 0% to 52% for 50-100% of repressed genes, depending on the relative rates of premature termination and pause release (Fig. 2B).

The model links changes in a gene's initiation rate to changes in body and pause densities. The change in initiation rate for a gene is constrained by the fold change in either pause density or body density, depending on the relative rates of premature termination and pause release (Eq. 9 and Eq. 10). To validate the model predictions, we analyzed genome-wide nascent transcriptome data after acute treatments with an inhibitor of initiation, triptolide (TRP)(11). Triptolide blocks initiation by inhibiting XPB helicase (63), a component of TFIIH (64, 65). Treatment with 0.5 μM triptolide for 12.5 mins (11) causes the expected decrease in initiation rate for more than 98% of repressed genes, with a median decrease of ~77% when premature termination is faster than pause release ($k_{pre} >> k_{rel}$, Fig. 2B, Fig. S4B). When pause release is comparable to or faster than premature termination ($k_{rel} = k_{pre}$ or $k_{rel} >> k_{pre}$), triptolide treatment leads to a median decrease in initiation rate of approximately 54% or 32%, respectively (Fig. 2B). Since initiation rate changes are constrained within a range and are dependent on relative rates of premature termination and pause release, we examined the results more closely to better understand these relative rates.

Inhibition by triptolide indicates that premature termination is faster than pause release.
Since changes in initiation rate are defined by relative values of premature termination and pause release (Eqs. 9-10), we sought to determine which rate is likely to be faster. To quantify the relationship between k_{pre} and k_{rel}, we considered changes in the initiation rate (Eq. 5) for repressed genes after triptolide treatment (Fig. 2B). Specifically, we reasoned that the fold change in initiation rate for all repressed genes will be close to 0.25, based on the results that initiation is decreased by ~75% after 10 minutes of 1μM triptolide treatment (61). Approximately 55% (3367 of 6142) of repressed genes had fold-change in initiation rate bounds spanning 0.25; 45% (2775 of 6142) of the genes had bounds that didn't include 0.25 (Fig. S5). For genes with bounds of changes in initiation rate spanning 0.25, we assumed their change in initiation rate after triptolide treatment to be 0.25. For genes with bounds on initiation rate

Fig. 2. Inhibition of initiation and pause release validate the model. A) Flavopiridol treatment decreases pause release for ~82% of repressed genes, consistent with its known inhibitory effect on CDK9. Inspection of genes with increased pause release reveal a decrease in pause density for all genes (Fig. S3A-C). This unexpected reduction in pause density is attributed to spiky signals in the pause region or insufficient reads to accurately define the pause window (Fig. S3D-I). Triptolide increases pause release at most genes. Although this was not expected, this phenomenon could be a result of a general compensatory mechanism to increase transcription when initiation is inhibited. B) The bounds on changes in initiation rate are estimated with distinct relative values of pause release (k_{rel}) and premature termination rate (k_{pre}) (Materials & Methods). As expected, nearly all repressed genes (>99%) decrease their initiation rate upon triptolide treatment. Flavopiridol does not show any clear effect on initiation if $k_{pre} >> k_{rel}$, but appears inhibitory if $k_{pre} << k_{rel}$ or $k_{pre} = k_{rel}$.

changes not spanning 0.25, we set their changes in initiation rate to the bound closest to 0.25, with a 5% offset (i.e; $fold\ change(k_{init}) = (1 \pm 0.05) \times closestBound$). The 5% buffer prevents $\frac{k_{pre}}{k_{rel}^{control}}$ values of zero or undefined as described in Eqs. 15-16. We then calculated $\frac{k_{pre}}{k_{rel}^{control}}$ (Eq. 14) which resulted in a median value of 6.7, an inter-decile range of 1.2 - 30, and $k_{pre} > k_{rel}$ for 90% of genes (5509 of 6142) (Fig. 3A). This trend is consistent among genes that are classified according to their expression level (Fig. S6). This suggests that the rate of premature termination is much faster than pause release at the majority of gene promoters, and on average only one out of seven paused polymerases proceeds into productive elongation. These results are consistent with previous reports that suggest premature termi-

nation rates are 4-12 times faster than pause release rates (9, 66) (Table S2).

The result of flavopiridol inhibition on pause release also suggests that $k_{pre} >> k_{rel}$. The pause inhibition data was generated with 300 nM flavopiridol (FP) for 25 minutes, which should cause a decrease in pause release without preferentially stimulating or inhibiting initiation. However, our analysis indicates that if $k_{pre} << k_{rel}$, then nearly every gene would have to decrease initiation after FP treatment to explain the observed changes in pause and gene body densities (Fig. 2B) (11). While flavopiridol may interact with other kinases, such as CDK7 (59), at this concentration flavopiridol is unlikely to cause a decrease in initiation at nearly all genes. Moreover, the observed increase in RNA polymerase density at pause sites following flavopiridol

Fig. 3. Experimental data and the model equations indicate that premature termination is faster than pause release. A) We set all TRP-repressed genes to have fold change (FC) in initiation rate (k_{init}) close to 0.25 (61). We calculated ratio of premature termination and pause release rates (Eq. 14). The median value of $\frac{k_{pre}}{k_{rel}^{control}}$ was 6.7 with an inter-decile range of 1.2 - 30. B) Changes in pause release affect changes in body density (transcription output) when premature termination k_{pre} is faster ($k_{pre} >> k_{rel}$). When k_{pre} is relatively slower than k_{rel}, change in body density is closely dependent upon changes in k_{init} and is minimally affected by changes in k_{rel}. As termination k_{pre} becomes faster than pause release k_{rel}, the body density becomes dependent on changes in k_{rel} and approaches the product of changes in initiation k_{init} and pause release k_{rel} rates. Pausing is regulated to control expression output in metazoans (62), so these results support premature termination being faster than pause release at most genes.

treatment (11) contrasts with the decrease in pause density reported for CDK7 inhibition (67). Both the triptolide and flavopiridol analyses support the conclusion that premature termination occurs at a rate substantially faster than pause release.

Change in pause release affects transcription output when premature termination is faster than pause release.

To better understand how the rates of initiation, pause release, and premature termination collectively influence gene expression, we examined how changes in these rates determine transcription output. We use RNA polymerase density in the gene body as a proxy for change in transcription output because transcript levels are determined by the product of three quantities: (a) the fold change in initiation rate (k_{init}), (b) the fold change in pause release rate (k_{rel}) and (c) the inverse of fold change in "net exit rate" from the pause region, $\frac{k_{pre}+k_{rel}^{control}}{k_{pre}+k_{rel}^{treatment}}$. Depending on the relative values of pause release (k_{rel}) and termination rate (k_{pre}), the inverse of fold change in exit rate ($\frac{k_{pre}+k_{rel}^{control}}{k_{pre}+k_{rel}^{treatment}}$) is either 1 ($k_{pre} >> k_{rel}$) or the inverse of fold change in

pause release ($\frac{k_{rel}^{control}}{k_{rel}^{treatment}}$ when $k_{pre} << k_{rel}$) (Eq. 7-8).

Thus, when pause release is much faster compared to termination ($k_{rel} >> k_{pre}$), the change in transcription output closely mirrors the change in initiation rate and variations in pause release (k_{rel}) have minimal impact on transcription output (Fig. 3B left panel). As k_{pre} becomes faster than k_{rel}, changes in pause release (k_{rel}) become more influential, and transcription output approaches the product of change in initiation and change in pause release rate (Fig. 3B right panel). These results demonstrate that regulating pause release has a bigger effect on gene expression output when premature termination occurs at a relatively faster rate.

These analyses, taken together with the analysis of pause-inhibition and initiation-inhibition data, as well as previous work (Table S2), suggests that premature termination is faster than pause release at most genes. Therefore, we maintain this assumption throughout the main figures and text, meaning that changes in initiation rate are interpreted as fold change in pause densities as measured by PRO-seq (Eq. 9).

Fig. 4. Modeling kinetic PRO-seq data determines the regulatory roles of activated and degraded transcription factors. A) Degradation of ZNF143 (44) and TBP (43) reduces initiation for approximately 90% of repressed genes in each dataset. Dexamethasone-activated genes (45) and Heat Shock Factor (HSF)-activated genes in S2 cells (33) do not show a clear direction of change in initiation rate, suggesting that *Drosophila* HSF and human glucocorticoid receptor (GR) may not regulate initiation. Fig. S8 contains the estimated fold change in initiation rates if premature termination is less than pause release. B) Dexamethasone (45) enhances pause release in approximately 89% of activated genes, indicating that GR's role is to regulate pause release. In *Drosophila* S2 cells, HSF activates target genes by facilitating pause release. Degradation of TBP and ZNF143 tends to increase pause release for repressed genes, similar to the effect seen in repressed genes after triptolide treatment (Fig. 2.)

Mechanistic functions of transcription factors.

To determine the regulatory roles of transcription factors, we analyzed existing PRO-seq datasets that rapidly degraded TBP and ZNF143 (43, 44). The degradation of TBP reduces the initiation rates for more than ~86% of repressed genes (Fig. 4A, Fig. S7A), highlighting TBP's established role in initiation (68). Similarly, the degradation of ZNF143 led to a decrease in initiation rates for over ~87% of repressed genes (Fig. 4A, Fig. S7B), indicating that ZNF143 stimulates initiation. Previous work proposed the role of ZNF143 as an initiation factor by demonstrating that ZNF143 influences transcription start site (TSS) selection, with alternative TSSs being utilized in repressed genes following ZNF143 degradation (44).

Next, we investigated the function of glucocorticoid receptor (GR) by analyzing PRO-seq data after acute dexamethasone (Dex) treatment (45). Upon binding dexamethasone, GR translocates to the nucleus, binds DNA and directly activates transcription (69–71). After 45 mins of dexamethasone treatment in A549 cells (45), pause release rate increased for 89% of activated genes (Fig. 4B, Fig. S7C) with no clear change in initiation rate for Dex-activated genes (Fig. 4A), indicating that GR predominantly activates its target genes by stimulating productive release of paused RNA polymerases.

Promoter-proximal pausing was first characterized at *Drosophila* heat shock genes (2) and the *Drosophila* heat shock factor (HSF) is known to stimulate pause release (33, 72, 73). Our analyses confirms that nearly all (~97%) HSF-activated genes in *Drosophila* S2 cells show an increase in pause release upon heat shock (Fig. 4B, Fig. S7D). Taken together, these analyses of existing kinetic PRO-seq data demonstrates the utility of this approach when assigning molecular functions to transcription factors.

Inferring absolute and effective pause release rates.

Our model estimates the absolute values of pause release rates as the product of elongation rates and the inverse of a gene's *pausing index* (Eq. 11). Since these estimates are derived from the ratio of body to pause densities, they are independent of the scaling or normalization criteria applied to genomic data. We considered 2000 bp/min a representative elongation rate and we kept this constant for all genes because we cannot estimate accurate gene-specific elongation rates. We report the pause release rates before and after Dex treatment because GR regulates pause release (Fig. 4B) (45). The median pause release rate constants increase from 0.86 events/min to 1.35 events/min (Fig. 5A), with an inter-decile range of 0.27 - 4.1 events/min in untreated conditions and 0.42 - 7 events/min after dexamethasone treatment.

We also estimated an effective pause release rate, $k_{rel} \times P$, representing the number of polymerases entering productive elongation at steady state. Since calculating effective pause release requires an estimation of pause densities as absolute occupancy values, we used a saturation fit to scale pause densities (Fig. S1E) (Materials & Methods). Upon dexamethasone treatment, the median effective pause release rate of Dex-activated genes increased

Fig. 5. Rates of initiation and pause release can be estimated using steady-state densities of pause and gene body regions. A) Assuming an elongation rate of k_{elong} = 2000 bp/min, the median pause release rates of Dex-activated genes increase from 0.86 events/min (control) to 1.35 events/min. 89% of the Dex-activated genes (537 out of 605) increase their pause release rate. B) The effective pause release rate represents the release of RNA polymerase into the gene body and is calculated as the product of the pause release rate constant and the scaled pause density. The pause and body densities was transformed into occupancy levels of RNA polymerase (Fig. S1E). The median effective pause release rate for Dex-activated genes increases 1.6-fold, from 0.21 RNAP/min to 0.34 RNAP/min under treatment. All Dex-activated genes increase their effective pause release rate. C) The density in gene body increased after dexamethasone treatment with median density increasing from 1 RNAP every 9.6kb to 1 RNAP every 5.8kb. The inter-decile ranges were 1 RNAP every 3.1 kb - 36.3 kb in untreated conditions, increasing to 1 RNAP every 1.8 kb - 20.3 kb after dexamethasone treatment. D) Degradation of TBP in HAP1 cells and ZNF143 in HEK293T cells decreases the median initiation rate from 0.5 RNAP/min to 0.27 RNAP/min and 0.71 RNAP/min to 0.51 RNAP/min respectively.

1.6-fold from 0.21 RNA polymerase (RNAP) /min to 0.34 RNAP/min (Fig. 5B) with an inter-decile range of 0.06 - 0.63 RNAP/min under untreated conditions and 0.09 - 1 RNAP/min after dexamethasone treatment. Effective pause release rates increase for all Dex-activated genes (Fig. 5B). Ninety percent of Dex-activated genes have $\leq$ 1 RNAP en-

tering the gene body per minute (Fig. 5B).

Next, we calculated RNA polymerase occupancy levels in the gene body of Dex-activated genes in both conditions (Eq. 13). Based on the effective pause release rates, the median estimated occupancy in untreated cells was 1 RNA polymerase per 9.6 kilobases (kb), with an interdecile range of 1 RNA polymerase per 3.2 kb - 33.3 kb (Fig. 5C). The density of RNA polymerase in the gene body increased after dexamethasone treatment to median of 1 RNA polymerase every 5.8 kb and inter-decile range of 1 RNA polymerase every 1.8 kb - 20.3 kb (Fig. 5C). These results align with previous imaging studies showing that two-thirds of nascent RNA transcripts are spaced more than 12 kb apart from another RNA polymerase, while the remaining 33% of transcripts exhibit densities ranging from 1 RNA polymerase per kb to 1 RNA polymerase per 8 kb (74, 75). Previously reported "initiation" rates of 0.16–0.4 events/min in single-molecule imaging studies (76) can be interpreted as effective pause release rates, corresponding to a body density of 1 RNA polymerase every 5–12.5 kb. Another study estimated effective pause release rates of 0.19–0.29 events/min, yielding a body density of 1 RNA polymerase every 6.9–10.5 kb (77). Our calculated pause release rates and genomic estimates of RNA polymerase occupancy align with independent imaging and single-molecule studies, underscoring the validity of our approach in converting unit-less genomic RNA polymerase profiling values to absolute RNA polymerase occupancy estimates.

Inferring absolute initiation rates.

The initiation rate estimate incorporates premature termination rates, elongation rates, with pause and body densities as occupancy values (Eq. 12) (Materials & Methods). We concluded faster premature termination rates (k_{pre}) than pause release rates (k_{rel}) at gene promoters using experimental data from triptolide-treated cells (Fig. 3A). For the rapid degradation of TBP and ZNF143 datasets, we assumed that premature termination is 6.7x pause release (Fig. 3A). Upon TBP degradation in HAP1 cells, the median initiation rate decreased from 0.5 events/min to 0.27 RNAP/min with inter-decile ranges of 0.07 - 2.7 RNAP/min under untreated conditions and 0.04 - 2 RNAP/min upon TBP degradation (Fig. 5D). While, upon ZNF143 degradation in HEK293T cells, the median initiation rate decreased from 0.71 RNAP/min to 0.51 RNAP/min with inter-decile ranges of 0.3 - 1.9 RNAP/min under untreated conditions and 0.17 - 1.4 RNAP/min upon ZNF143 degradation (Fig. 5D). These calculations indicate that initiation rates are, on average, half of pause release rate constants, but the distributions of these rates largely fall within comparable ranges.

Most paused polymerases turn over in less than a minute.

The residency time of a paused RNA polymerase is determined by the rates of premature termination and pause release. Based on Fig. 3A, we considered $k_{pre} = 6.7 \times k_{rel}$ at each gene promoter to calculate the residency time of paused RNA polmerase. The median half-lives of paused RNA polymerases, calculated as $\log_2 / (k_{pre} + k_{rel})$, across all treatments was approximately 8.2 seconds (Fig. 6), with

Fig. 6. Most paused polymerases turn over rapidly. Assuming premature termination rates at each gene promoter as 6.7 times faster than pause release rates (Fig. 3A), the median half-life of paused polymerases across all datasets and treatments is 8.2 seconds, with an inter-decile range of 1.6 to 36.2 seconds. For repressed genes under triptolide treatment, the median half-life decreased from 7.9 seconds to 6.1 seconds. For HSF-activated genes, the median half-life decreased from 17.4 seconds to 12 seconds. The half-lives in other datasets showed only minor changes after treatment.

an inter-decile range of 1.6 to 36.2 seconds, indicating that most paused polymerases turn over within a minute. Premature termination rates primarily determine half-lives rather than pause release rates. For instance, at $k_{pre} = 10 \times k_{rel}$ (Fig. S9), the turnover time and interdecile range decrease by an approximate factor of 1.5 (i.e., $10/6.7$). Although several studies are consistent with our estimates of short lived paused polymerases (9, 61, 78), others report longer half-lives (10, 11, 66, 79). In the discussion we provide possible explanations for the disparate estimates in an attempt to reconcile seemingly inconsistent findings.

Parameter sensitivity analysis supports the robustness of our conclusions.

The three values that we base many of these calculations and our conclusions upon are: 1) the average elongation rate of a gene; 2) the efficacy of triptolide inhibition; and 3) the absolute density of RNA polymerases in the pause region. We varied these values over a wide range to determine how changing these parameters influence rates, ratios of rates, and our general conclusions.

In the previous sections we assumed that the average elongation rate is 2kb/min. However, elongation rates can vary between genes and within a gene (34, 49, 80, 81). Our analysis ignores the fact that elongation rate varies within a gene because we average the gene body density, which effectively averages the elongation rate in our model. We varied the average elongation rate from 1-5kb/min to determine how elongation speed influences other rates and our conclusions. In short, changing the elongation rate causes linear and proportional changes in all other rates (Fig. S10), since pause release is linearly dependent on elongation (Eq. 11) and premature termination is related to pause release (Eq. 14). For example, if we increase elongation rate by 2 fold, then all other rates will change by two-fold. The values of elongation rate have no effect on k_{pre}/k_{rel} (Eq. 14), fold changes in initiation (Eq. 9), or fold change in pause release (Eq. 6) between experimental conditions. However, faster elongation rates would correspond to shorter residency times of paused RNA polymerase due to faster termination and pause release (Fig. S10A&B).

We assumed that 12.5 minutes of 0.5 μM triptolide caused an average fold-change in the initiation rate of 0.25 for triptolide-repressed genes in mESC cells (11). This estimate is based on an *in vitro* estimate of genome-wide decrease in initiation after a time course (1, 3, 10 and 30 mins) of 1μM triptolide treatment (61). Uncertainty in this parameter will affect our estimates of absolute premature termination rates, absolute initiation rates, and the premature termination to pause release ratio (Fig. S11). If triptolide treatment is less effective than we assume (i.e. fold-change in initiation is >0.25), then the premature termination rate decreases (Fig. S11A, Eq. 14). The pause release rate is only dependent upon elongation rate, so this remains unchanged. If triptolide causes a modest 0.45 fold-change in initiation, then approximately half of genes have a $k_{pre} > k_{rel}$ (i.e. termination and pause release rates are comparable) (Fig. S11A). The reciprocal is also true: if we are underestimating trip-

tolide efficacy, then premature termination of a paused RNA polymerase is more likely than we report. Our estimate of triptolide efficacy is based on experimental data and 75% inhibition spans the empirically determined bounds of initiation rate fold change (Fig. S5), but other studies assume that this same triptolide treatment is nearly complete and more rapid (11). Although we only focus on triptolide-repressed genes herein, all genes are likely differentially affected by triptolide treatment, some completely inhibited and others insensitive. In the future, gene-specific measurements or estimates are needed to quantify the relationship between termination and pause release at individual genes.

We used a saturation function to estimate "full" occupancy of the pause region, which allowed us to calculate the absolute density of RNA polymerase in the gene body (Fig. S1A-F). Since the triptolide dataset (11) doesn't have UMIs, we had considered 90th percentile as saturation to be consistent with saturation values observed in datasets with UMIs (Fig. S1D-F). To investigate how saturation values may affect occupancy, we varied them by 5-fold in each direction (Fig. S12). Based on pause signal of *x* arbitrary units at the 90th percentile, we considered pause sums at 48, 70, 90, 95, 99.8 percentiles of ranked pause sums (P = 0.2*x*, 0.4*x*, *x*, 1.8*x* and 5*x*). The ratio of premature termination rate to pause release rate remains stable over this wide range of absolute RNAP density estimates (Fig. S12A), with only modest <2-fold differences in both rates over the 25-fold range (0.2*x* to 5*x*) (Fig. S12B&C). Initiation rate is more sensitive to the RNAP density, decreasing proportionally as RNA polymerase density estimates increase (Fig. S12D).

Scaling absolute pause occupancy proportionally affects gene body density calculations. By inferring the RNAP density in the pause region, we can calculate the median values of RNA Polymerase density within genes. These median gene body densities equate to 1 RNAP every 1.5 kb (48 percentile; 0.2*x*), 1 RNAP every 5.9 kb (90 percentile; *x*), and 1 RNAP every 28.3 kb (99.8 percentile; 5*x*) (Fig. S12E). The 90th percentile saturation values (Fig. 5C, Fig. S12E) equate to gene body densities that are supported by multiple imaging studies: 1 RNAP every 8-12 kb (74), 1 RNAP every 5-12.5 kb (76) and 1 RNAP every 6.9-10.5 kb (77). Although, absolute RNAP occupancy estimates influence all calculations except the k_{pre}/k_{rel} ratio, defining "full" occupancy using the 90th percentile of pause density–guided by saturation curve estimates (Fig. S1)–yields gene body densities that align best with independent RNA polymerase occupancy measurements.

Discussion

The results of this study provide insights into the kinetic mechanisms of transcriptional regulation, with a focus on the fate of paused RNA polymerases. Our modeling framework, validated with kinetic PRO-seq datasets, reveals that most paused polymerases terminate prematurely rather than progressing into productive elongation (Fig. 7). We found that the rate of premature termination (k_{pre}) is signif-

icantly faster than the rate of pause release (k_{rel}) for most genes. By leveraging triptolide inhibition data, we estimated the ratio of premature termination to pause release rates (k_{pre}/k_{rel}) and found that premature termination occurs approximately seven times more frequently than productive elongation. This aligns with imaging-based studies reporting rapid turnover of paused polymerases within seconds (9). Moreover, the dependency of transcriptional output on pause release rate changes further supports the predominance of premature termination. The relative rates of termination versus pause release has significant implications for understanding transcriptional control. Transcription factors that enhance pause release, such as HSF and GR, have greater influence on gene expression in contexts with fast termination rates. The ability of our model to quantify these rates provides a framework for determining how different factors modulate transcription.

Our framework was validated using datasets in which initiation and pause release were directly inhibited. The effects of triptolide on initiation and flavopiridol on pause release were consistent with known molecular mechanisms, demonstrating that our model accurately captures kinetic changes in transcription. Furthermore, the model successfully classified transcription factors based on their regulatory roles, confirming TBP and ZNF143 as initiation factors and HSF and GR as regulators of pause release.

Model limitations.

Sensitivity analyses further support the robustness of our conclusions. While absolute values of rates depend on estimates of elongation speed and triptolide efficacy, the key finding that termination is faster than pause release remains consistent across a broad range of estimates. Moreover, our calculation of gene body RNA polymerase densities from the modeling align with previous imaging studies, reinforc-

ing the validity of our RNA polymerase occupancy estimates.

One limitation of our current framework is that it does not explicitly take transcriptional bursting into account. Transcription occurs in stochastic bursts, where genes switch between active and inactive states, leading to fluctuations in RNA polymerase occupancy (82). Although our model assumes steady-state transcription rates, it can be adapted to incorporate bursting kinetics if the burst frequency and duration are known for a given gene. For example, if 20% of a gene's alleles are active, then the saturation curve would be re-calibrated so that complete occupancy would be considered 20% of the estimated full occupancy. Future work could integrate live cell imaging data to measure these values and refine the model to more fully capture dynamic transcriptional regulation for each gene.

Reconciling disparate estimates of rates and pause residency times from previous studies.

We found that premature termination is much faster than pause release (Fig. 3A) and paused RNA polymerases turn over rapidly, on the order of tens of seconds (Fig. 6). Some previous studies report slower pause release and premature termination rate constants (Table S2) (66), with estimated half-lives in the range of several minutes (66, 79). A key distinction between these studies and our approach is that they exclusively measured capped short nascent RNA (66, 79), while we quantify all engaged RNA polymerases that are competent to elongate (83, 84). We speculate that paused RNA polymerases with capped RNA are less likely to prematurely terminate than polymerases with uncapped RNA, potentially accounting for the observed differences. If nearly all initiated polymerases terminate prematurely (9) (Fig. S13), the slow turnover rates observed in (66, 79) may primarily reflect the termination dynamics of the minority

Fig. 7. A kinetic model of initiation, termination, and early elongation. The cartoon represents the conclusions from our modeling of kinetic nascent transcriptome data. The initiation rate of 0.5 RNA polymerases per minute assumes a non-limiting amount of RNA polymerase available to initiate. The translucent RNA polymerase represents pre-initiation or abortive initiation complexes; so the term *initiation* rate within our model refers to the series of events that have to occur for an RNA polymerase to successfully proceed into the pause region. The average pause release rate constant is 0.25/minute and the average termination rate of pause RNA polymerases is approximate 7 time faster at 1.7/minute. These rates are derived from well established rates of elongation (~2 kilobases/minute) averaged over a gene body, experimental triptolide efficacy data, and empirical estimates of fully occupied pause sites.

population of paused RNA polymerases with capped RNA, with minimal contribution of pause release to measurements in both untreated and flavopiridol conditions.

Another study that reported longer half-lives in minutes had used 0.5µM triptolide treatment over several time points (11). Since the initial publication of this study, comparable times and concentrations of triptolide were shown to be insufficient to fully inhibit initiation (61). Our analysis also provides an upper bound for initiation inhibition, which supports these *in vitro* measurements of triptolide efficacy (61). Anything less than immediate and complete inhibition of initiation would cause an inflation of pause residency time because all newly initiated RNA polymerases will be considered stably paused.

A live cell imaging study used a photoactivatable GFP-RNA Pol II to measure pause stability at an uninduced and heat shock-induced *Hsp70* transgene (10). This work was performed in polytene chromosomes, averaging the RNAPII signal over a thousand copies of *Hsp70*. This analysis assumes that paGFP-RNAPII will dissociate from the locus upon termination, but if terminated RNAPII reinitiates at a proximal *Hsp70* gene, it will not contribute to the decay of GFP signal from the locus. This will result in an inflated half-life of paused polymerases because any local reinitiation will be considered stably paused. They also used a biochemical fractionation experiment upon inhibition of initiation with 10µM triptolide to quantify decay of nascent paused RNA (10). Again, subsequent studies found that even 10µM triptolide is not sufficient to immediately and completely inhibit initiation (61). This leads to a deflated decay rate because newly initiated RNA polymerases are indistinguishable from polymerases that are stably paused.

Future Directions.

Although our model is based on known mechanisms of initiation and early elongation and provides valuable insights into transcription dynamics, future work could refine these estimates using direct measurements of elongation rates at individual genes. Single-molecule imaging approaches could complement our genomic estimates by providing real-time observations of polymerase behavior. Expanding this model to incorporate gene-specific variations in elongation and bursting kinetics would enhance its predictive power.

By quantifying initiation, pause release, and termination rates, our framework and corresponding `compartmentModel` software are powerful tools for dissecting the kinetic mechanisms of transcription and determining the molecular functions of transcription factors.

ACKNOWLEDGEMENTS
This work was funded by R35-GM128635 to MJG. We thank Pedro Mendes for modeling discussions and providing parameter estimations techniques.

AUTHOR CONTRIBUTIONS
RM and MJG conceptualized and developed the project. RM and MJG developed the model. RM analyzed the data. RM developed the software. RM and MJG wrote the manuscript.

References

1. Saunders A, Core LJ, Lis JT. Breaking barriers to transcription elongation. Nature reviews Molecular cell biology. 2006;7(8):557–567.

2. Rougvie AE, Lis JT. The RNA polymerase II molecule at the 5 end of the uninduced hsp70 gene of D. melanogaster is transcriptionally engaged. Cell. 1988;54(6):795–804.

3. Core LJ, Lis JT. Transcription regulation through promoter-proximal pausing of RNA polymerase II. Science. 2008;319(5871):1791–1792.

4. Price DH. P-TEFb, a cyclin-dependent kinase controlling elongation by RNA polymerase II. Molecular and cellular biology. 2000;.

5. Stein CB, Field AR, Mimoso CA, Zhao C, Huang KL, Wagner EJ, et al. Integrator endonuclease drives promoter-proximal termination at all RNA polymerase II-transcribed loci. Molecular Cell. 2022;82(22):4232–4245.e11. doi:10.1016/j.molcel.2022.10.004.

6. Tatomer DC, Elrod ND, Liang D, Xiao MS, Jiang JZ, Jonathan M, et al. The Integrator complex cleaves nascent mRNAs to attenuate transcription. Genes & development. 2019;33(21-22):1525–1538.

7. Suter DM, Molina N, Gatfield D, Schneider K, Schibler U, Naef F. Mammalian Genes Are Transcribed with Widely Different Bursting Kinetics. Science. 2011;332(6028):472–474. doi:10.1126/science.1198817.

8. Zenklusen D, Larson DR, Singer RH. Single-RNA counting reveals alternative modes of gene expression in yeast. Nature Structural & Molecular Biology. 2008;15:1263–1271. doi:10.1038/nsmb.1514.

9. Steurer B, Janssens RC, Geverts B, Geijer ME, Wienholz F, Theil AF, et al. Live-cell analysis of endogenous GFP-RPB1 uncovers rapid turnover of initiating and promoter-paused RNA Polymerase II. Proceedings of the National Academy of Sciences of the United States of America. 2018;115(19):E4368–E4376. doi:10.1073/pnas.1717920115.

10. Buckley MS, Kwak H, Zipfel WR, Lis JT. Kinetics of promoter Pol II on Hsp70 reveal stable pausing and key insights into its regulation. Genes & development. 2014;28(1):14–19.

11. Jonkers I, Kwak H, Lis JT. Genome-wide dynamics of Pol II elongation and its interplay with promoter proximal pausing, chromatin, and exons. Elife. 2014;3:e02407. doi:10.7554/eLife.02407.

12. Zhao Y, Liu L, Hassett R, Siepel A. Model-based characterization of the equilibrium dynamics of transcription initiation and promoter-proximal pausing in human cells. Nucleic Acids Research. 2023;51(21):e106. doi:10.1093/nar/gkad843.

13. Matsui T, Segall J, Weil P, Roeder R. Multiple factors required for accurate initiation of transcription by purified RNA polymerase II. Journal of Biological Chemistry. 1980;255(24):11992–11996.

14. Dynan WS, Tjian R. Isolation of transcription factors that discriminate between different promoters recognized by RNA polymerase II. Cell. 1983;32(3):669–680.

15. Parker CS, Topol J. A Drosophila RNA polymerase II transcription factor contains a promoter-region-specific DNA-binding activity. Cell. 1984;36(2):357–369.

16. Sopta M, Carthew RW, Greenblatt J. Isolation of three proteins that bind to mammalian RNA polymerase II. Journal of Biological Chemistry. 1985;260(18):10353–10360.

17. Feaver WJ, Gileadi O, Li Y, Kornberg RD. CTD kinase associated with yeast RNA polymerase II initiation factor b. Cell. 1991;67(6):1223–1230.

18. Goodrich JA, Tjian R. Transcription factors IIE and IIH and ATP hydrolysis direct promoter clearance by RNA polymerase II. Cell. 1994;77(1):145–156.

19. Cortes P, Flores O, Reinberg D. Factors involved in specific transcription by mammalian RNA polymerase II: purification and analysis of transcription factor ILA and identification of transcription factor IIJ. Molecular and cellular biology. 1992;12(1):413–421.

20. Zhou Q, Lieberman PM, Boyer TG, Berk AJ. Holo-TFIID supports transcriptional stimulation by diverse activators and from a TATA-less promoter. Genes & Development. 1992;6(10):1964–1974.

21. Killeen MT, Greenblatt JF. The general transcription factor RAP30 binds to RNA polymerase II and prevents it from binding nonspecifically to DNA. Molecular and cellular biology. 1992;12(1):30–37.

22. Akoulitchev S, Mäkelä TP, Weinberg RA, Reinberg D. Requirement for TFIIH kinase activity in transcription by RNA polymerase II. Nature. 1995;377(6549):557–560.

23. Klages N, Strubin M. Stimulation of RNA polymerase II transcription initiation by recruitment of TBP in vivo. Nature. 1995;374(6525):822–823.

24. Dvir A, Garrett KP, Chalut C, Egly JM, Conaway JW, Conaway RC. A Role for ATP and TFIIH in Activation of the RNA Polymerase II Preinitiation Complex Prior to Transcription Initiation (). Journal of Biological Chemistry. 1996;271(13):7245–7248.

25. Holstege F, Van der Vliet P, Timmers H. Opening of an RNA polymerase II promoter occurs in two distinct steps and requires the basal transcription factors IIE and IIH. The EMBO journal. 1996;15(7):1666–1677.

26. Zawel L, Kumar KP, Reinberg D. Recycling of the general transcription factors during RNA polymerase II transcription. Genes & development. 1995;9(12):1479–1490.

27. Horikoshi M, Hai T, Lin YS, Green MR, Roeder RG. Transcription factor ATF interacts with the TATA factor to facilitate establishment of a preinitiation complex. Cell. 1988;54(7):1033–1042.

28. Buratowski S, Hahn S, Guarente L, Sharp PA. Five intermediate complexes in transcription initiation by RNA polymerase II. Cell. 1989;56(4):549–561.

29. Giniger E, Ptashne M. Cooperative DNA binding of the yeast transcriptional activator GAL4. Proceedings of the National Academy of Sciences. 1988;85(2):382–386.

30. Dong J, Guertin MJ. Classifying the molecular functions of transcription factors beyond activation and repression. Genes & Development. 2024;38(21-24):931–932.

31. Rogers JM, Mimoso CA, Martin BJ, Martin AP, Aster JC, Adelman K, et al. Notch induces transcription by stimulating release of paused RNA polymerase II. Genes & Development. 2024;38(21-24):965–978.

32. Hah N, Danko CG, Core L, Waterfall JJ, Siepel A, Lis JT, et al. A rapid, extensive, and transient transcriptional response to estrogen signaling in breast cancer cells. Cell. 2011;145(4):622–634.

33. Duarte FM, Fuda NJ, Mahat DB, Core LJ, Guertin MJ, Lis JT. Transcription factors GAF and HSF act at distinct regulatory steps to modulate stress-induced gene activation. Genes & Development. 2016;30(15):1731–1746. doi:10.1101/gad.284430.116.

34. Danko CG, Hah N, Luo X, Martins AL, Core L, Lis JT, et al. Signaling pathways differentially affect RNA polymerase II initiation, pausing, and elongation rate in cells. Molecular cell. 2013;50(2):212–222.

35. Kwak H, Fuda NJ, Core LJ, Lis JT. Precise Maps of RNA Polymerase Reveal

How Promoters Direct Initiation and Pausing. Science. 2013;339(6122):950–953. doi:10.1126/science.1229386.

36. Core LJ, Waterfall JJ, Lis JT. Nascent RNA sequencing reveals widespread pausing and divergent initiation at human promoters. Science. 2008;322(5909):1845–1848. doi:10.1126/science.1162228.

37. Clopper KC, Taatjes DJ. Chemical inhibitors of transcription-associated kinases. Current Opinion in Chemical Biology. 2022;70:102186. doi:10.1016/j.cbpa.2022.102186.

38. Henley MJ, Koehler AN. Advances in targeting 'undruggable' transcription factors with small molecules. Nature Reviews Drug Discovery. 2021;20:669–688. doi:10.1038/s41573-021-00199-0.

39. Phuc Le P, Friedman JR, Schug J, et al. Glucocorticoid receptor-dependent gene regulatory networks. PLoS Genetics. 2005;1(2):e16. doi:10.1371/journal.pgen.0010016.

40. Nabet B, Roberts JM, Buckley DL, et al. The dTAG system for immediate and target-specific protein degradation. Nature Chemical Biology. 2018;14:431–441. doi:10.1038/s41589-018-0021-8.

41. Nishimura K, Fukagawa T, Takisawa H, Kakimoto T, Kanemaki M. An auxin-based degron system for the rapid depletion of proteins in nonplant cells. Nature methods. 2009;6(12):917–922.

42. Sathyan KM, McKenna BD, Anderson WD, Duarte FM, Core L, Guertin MJ. An improved auxin-inducible degron system preserves native protein levels and enables rapid and specific protein depletion. Genes & Development. 2019;33(19-20):1441–1455. doi:10.1101/gad.328237.119.

43. Santana JF, Collins GS, Parida M, Luse DS, Price DH. Differential dependencies of human RNA polymerase II promoters on TBP, TAF1, TFIIB and XPB. Nucleic Acids Research. 2022;50(16):9127–9148. doi:10.1093/nar/gkac678.

44. Dong J, Sathyan KM, Scott TG, Mukherjee R, Guertin MJ. ZNF143 binds DNA and stimulates transcription initiation to activate and repress direct target genes. Nucleic Acids Research. 2024; p. gkae1182. doi:10.1093/nar/gkae1182.

45. Wissink EM, Martinez DM, Ehmsen KT, Yamamoto KR, Lis JT. Glucocorticoid receptor collaborates with pioneer factors and AP-1 to execute genome-wide regulation. bioRxiv. 2021;doi:10.1101/2021.06.01.444518.

46. Smith JP, Dutta AB, Sathyan KM, et al. PEPPRO: quality control and processing of nascent RNA profiling data. Genome Biology. 2021;22(1):155. doi:10.1186/s13059-021-02349-4.

47. Scott TG, Martins AL, Guertin MJ. Processing and evaluating the quality of genome-wide nascent transcription profiling libraries. bioRxiv. 2022;doi:10.1101/2022.12.14.520463.

48. Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. Genome Biology. 2014;15(12):550. doi:10.1186/s13059-014-0550-8.

49. Jonkers I, Lis J. Getting up to speed with transcription elongation by RNA polymerase II. Nature Reviews Molecular Cell Biology. 2015;16:167–177. doi:10.1038/nrm3953.

50. Thompson JD, Higgins DG, Gibson TJ. CLUSTAL W: improving the sensitivity of progressive multiple sequence alignment through sequence weighting, position-specific gap penalties and weight matrix choice. Nucleic acids research. 1994;22(22):4673–4680.

51. Cock PJ, Antao T, Chang JT, Chapman BA, Cox CJ, Dalke A, et al. Biopython: freely available Python tools for computational molecular biology and bioinformatics. Bioinformatics. 2009;25(11):1422.

52. Anderson WD, Duarte FM, Civelek M, Guertin MJ. Defining data-driven primary transcript annotations with primaryTranscriptAnnotation in R. Bioinformatics. 2020;36(9):2926–2928. doi:10.1093/bioinformatics/btaa011.

53. O'Brien T, Lis JT. Rapid changes in Drosophila transcription after an instantaneous heat shock. Molecular and cellular biology. 1993;13(6).

54. Thummel CS, Burtis KC, Hogness DS. Spatial and temporal patterns of E74 transcription during Drosophila development. Cell. 1990;61(1):101–111.

55. Singh J, Padgett RA. Rates of in situ transcription and splicing in large human genes. Nature structural & molecular biology. 2009;16(11):1128–1133.

56. Core LJ, Waterfall JJ, Gilchrist DA, Fargo DC, Kwak H, Adelman K, et al. Defining the status of RNA polymerase at promoters. Cell reports. 2012;2(4):1025–1035.

57. Virtanen P, Gommers R, Oliphant TE, Haberland M, Reddy T, Cournapeau D, et al. SciPy 1.0: fundamental algorithms for scientific computing in Python. Nature methods. 2020;17(3):261–272.

58. Chao SH, Price DH. Flavopiridol inactivates P-TEFb and blocks most RNA polymerase II transcription in vivo. The Journal of Biological Chemistry. 2001;276:31793–31799. doi:10.1074/jbc.M102306200.

59. Karaman MW, Herrgard S, Treiber DK, Gallant P, Atteridge CE, Campbell BT, et al. A quantitative analysis of kinase inhibitor selectivity. Nature biotechnology. 2008;26(1):127–132.

60. Marshall NF, Peng J, Xie Z, Price DH. Control of RNA polymerase II elongation potential by a novel carboxyl-terminal domain kinase. Journal of Biological Chemistry. 1996;271(43):27176–27183.

61. Nilson KA, Lawson CK, Mullen NJ, Ball CB, Spector BM, Meier JL, et al. Oxidative stress rapidly stabilizes promoter-proximal paused Pol II across the human genome. Nucleic acids research. 2017;45(19):11088–11105.

62. Adelman K, Lis JT. Promoter-proximal pausing of RNA polymerase II: emerging roles in metazoans. Nature Reviews Genetics. 2012;13(10):720–731.

63. Titov DV, Gilman B, He QL, Bhat S, Low WK, Dang Y, et al. XPB, a subunit of TFIIH, is a target of the natural product triptolide. Nature Chemical Biology. 2011;7:182–188. doi:10.1038/nchembio.522.

64. Rimel JK, Taatjes DJ. The essential and multifunctional TFIIH complex. Protein Science. 2018;27(6):1018–1037.

65. Roeder RG. The role of general initiation factors in transcription by RNA polymerase II. Trends in biochemical sciences. 1996;21(9):327–335.

66. Zimmer JT, Rosa-Mercado NA, Canzio D, Steitz JA, Simon MD. STL-seq reveals pause-release and termination kinetics for promoter-proximal paused RNA polymerase II transcripts. Molecular cell. 2021;81(21):4398–4412.

67. Velychko T, Mohammad E, Ferrer-Vicens I, Parfentev I, Werner M, Studniarek C, et al.

68. CDK7 kinase activity promotes RNA polymerase II promoter escape by facilitating initiation factor release. Molecular Cell. 2024;.

68. Hernandez N. TBP, a universal eukaryotic transcription factor? Genes dev. 1993;7(7B):1291–308.

69. Tsurufuji S, Sugio K, Takemasa F. The role of glucocorticoid receptor and gene expression in the anti-inflammatory action of dexamethasone. Nature. 1979;280(5721):408–410.

70. Reddy TE, Pauli F, Sprouse RO, Neff NF, Newberry KM, Garabedian MJ, et al. Genomic determination of the glucocorticoid response reveals unexpected mechanisms of gene regulation. Genome research. 2009;19(12):2163–2171.

71. Vockley CM, D'Ippolito AM, McDowell IC, Majoros WH, Safi A, Song L, et al. Direct GR binding sites potentiate clusters of TF binding across the human genome. Cell. 2016;166(5):1269–1281.

72. Lis JT, Mason P, Peng J, Price DH, Werner J. P-TEFb kinase recruitment and function at heat shock loci. Genes & development. 2000;14(7):792–803.

73. Ni Z, Saunders A, Fuda NJ, Yao J, Suarez JR, Webb WW, et al. P-TEFb is critical for the maturation of RNA polymerase II into productive elongation in vivo. Molecular and cellular biology. 2008;28(3):1161–1170.

74. Jackson DA, Iborra FJ, Manders EM, Cook PR. Numbers and organization of RNA polymerases, nascent transcripts, and transcription units in HeLa nuclei. Molecular biology of the cell. 1998;9(6):1523–1536.

75. Beyer AL, Bouton AH, Hodge LD, Miller Jr OL. Visualization of the major late R strand transcription unit of adenovirus serotype 2. Journal of Molecular Biology. 1981;147(2):269–295.

76. Larson DR, et al. Real-Time Observation of Transcription Initiation and Elongation on an Endogenous Yeast Gene. Science. 2011;332:475–478. doi:10.1126/science.1202142.

77. Darzacq X, Shav-Tal Y, De Turris V, Brody Y, Shenoy SM, Phair RD, et al. In vivo dynamics of RNA polymerase II transcription. Nature structural & molecular biology. 2007;14(9):796–806.

78. Price DH. Transient pausing by RNA polymerase II. Proceedings of the National Academy of Sciences. 2018;115(19):4810–4812.

79. Henriques T, Scruggs BS, Inouye MO, et al. Widespread transcriptional pausing and elongation control at enhancers. Genes & Development. 2018;32(1):26–41. doi:10.1101/gad.309351.117.

80. Liu L, Zhao Y, Siepel A. DNA-sequence and epigenomic determinants of local rates of transcription elongation. bioRxiv. 2023;.

81. Ardehali MB, Lis JT. Tracking rates of transcription and splicing in vivo. Nature structural & molecular biology. 2009;16(11):1123–1124.

82. Rodriguez J, Ren G, Day CR, Zhao K, Chow CC, Larson DR. Intrinsic dynamics of a human gene reveal the basis of expression heterogeneity. Cell. 2019;176(1):213–226.

83. Rasmussen EB, Lis JT. In vivo transcriptional pausing and cap formation on three Drosophila heat shock genes. Proceedings of the National Academy of Sciences. 1993;90(17):7923–7927.

84. O'Brien T, Hardin S, Greenleaf A, Lis JT. Phosphorylation of RNA polymerase II C-terminal domain and transcriptional elongation. Nature. 1994;370(6484):75–77.

85. O'Brien T, Lis JT. RNA polymerase II pauses at the 5 end of the transcriptionally induced Drosophila hsp70 gene. Molecular and cellular biology. 1991;11(10):5285–5290.

Fig. S1. Saturation curves for datasets with Unique Molecular Identifiers (UMIs) provide estimates of RNA polymerase occupancy. The plots show the saturation curves and values from fits to the hyperbolic tangent function for datasets lacking Unique Molecular Identifiers (UMIs) (A-C) or containing UMIs (D-F). Datasets lacking UMIs do not have stable saturation values when ranges of pause sum percentiles are used to fit the saturation curve.

Fig. S2. Saturation values from datasets lacking Unique Molecular Identifiers (UMIs) are unreliable as maximum occupancy signal of RNA polymerase. A) The ratio of saturation values derived from fits to 100% of the data compared to 70% of the data (ignoring the top 30% of pause sum genes) is substantially higher for datasets without UMIs, ranging from 23-fold to 600-fold. In contrast, datasets with UMIs show much lower variability, with ratios ranging from 1.6-fold to 4-fold. B) These plots show the normalized read signal for the *Hsp70* consensus sequence from the dataset (33) used in this study. C) The pause sum of *Hsp70* (P = 992) is five times higher than the pause sum of the gene at the 90th percentile (P = 182). The violin plot represents the pause sums for all HSF-activated genes, most of which are higher than the 90th percentile value. Considering that *Hsp70* has a fully occupied pause region during heat shock (85), the assumption that the 90th percentile pause sum represents the maximum occupancy signal is not supported in this dataset lacking UMIs.

Fig. S3. Genes with increased pause release after flavopiridol treatment have atypical read distributions. A) Approximately 82% of flavopiridol-repressed genes decrease their pause release rate (k_{rel}). B) All genes with increased pause release rate also have a decrease in pause density. C) Sixty-one percent of repressed genes with decreased pause release have an expected increase in pause density after flavopiridol treatment. D-I) The UCSC Genome Browser snapshots show representative genes with increased pause release (k_{rel}) and decreased pause density. High pause signals under untreated conditions in these genes may result from the lack of UMIs in the libraries (11).

Fig. S4. Flavopiridol and Triptolide induce genome-wide repression and changes in RNA polymerase distribution. A) Flavopiridol (FP) treatment for 25 mins (11) resulted in 12994 *Down* and 310 *Up* genes. Composite profiles of the *Down* genes show an increase in RNA polymerase density in pause window upon FP treatment. B) Triptolide treatment for 12.5 mins (11) resulted in 9632 *Down* and 104 *Up* genes. The *Down* genes have a decrease in RNA polymerase density in pause window and gene body upon triptolide treatment.

Fig. S5. The model restrains the fold change initiation rates after triptolide inhibition for each genes within a calculated range. We consider repressed genes from a dataset that used 0.5 µM triptolide for 12.5 mins (11). A previous study reported a 0.25 fold change (dashed vertical line) in K_{init} upon treatment with 1µM triptolide at approximately 12 mins (61). We found 3367 (55% of 6142) TRP-repressed genes had bounds on changes in initiation rate spanning 0.25 (A). 2775 (45% of 6142) TRP-repressed genes had bounds on changes in initiation rate not spanning 0.25 (B).

Fig. S6. Premature termination is faster than pause release regardless of gene expression levels. We assumed that the repressed genes decrease initiation by ~75% and calculated ratio of premature termination and pause release rates for the indicated gene expression quantiles.

Fig. S7. Acute treatments induce genomic changes in gene expression and RNA polymerase distribution. A) Degradation of TBP (43) led to 493 *Down* and 86 *Up* genes. The *Down* genes have reduced RNA polymerase density in pause region upon TBP degradation. B) Rapid degradation of ZNF143 (44) led to 358 *Down* genes and 181 *Up* genes. The *Down* genes decrease pause density. C) Dexamethasone treatment for 45 mins (45) resulted in 862 *Up* and 738 *Down* genes. The *Up* genes have increased RNA polymerase density in pause window and gene body upon dexamethasone treatment. D) Heat shock (HS) in *Drosophila S2* cells resulted in 3922 *Up* genes, although many of these genes are false positives due to read-through transcription (33). The genes that are confidently HSF-activated (Materials & Methods) increase RNA polymerase density in both the pause region and gene body.

Fig. S8. Changes in initiation rate for datasets used in this study vary substantially depending on the $k_{rel} >> k_{pre}$ assumption. This figure reproduces the data from 4 and includes changes in initiation rates for faster pause release as well ($k_{rel} >> k_{pre}$).

Unique K_{pre} at each gene promoter

Fig. S9. Turnover times of paused polymerases can become extremely rapid, reaching sub-second timescales at high premature termination rates. Violin plots illustrate the distribution of paused RNA polymerase half lives if premature termination rates at each gene promoter are ten times faster than pause release rates (i.e., $k_{pre} = 10 \times k_{rel}$). The median half-life of paused RNA polymerases across all datasets and treatments is 5.83 seconds, with an inter-decile range of 1.2 to 26 seconds.

Fig. S10. The rates of pause release, premature termination, and initiation are proportional to elongation rates. A) We varied elongation rates from 1–5 kb/min. Under untreated conditions for repressed genes under TRP inhibition (11), pause release rates (k_{rel}) had interdecile ranges of 0.24–4.1 events/min, 0.6–10.8 events/min, 0.82–14.1 events/min, and 1.2–20.7 events/min for elongation rates of 1, 2.6, 3.4, and 5 kb/min, respectively. Median values are indicated in the boxplots. B) We calculated premature termination rates (k_{pre}) based on pause release using Eq. 14. The fold change in initiation rate (k_{init}) was set to 0.25. The inter-decile ranges of premature termination rates were 0.7-43.5 events/min, 1.8-113.1 events/min, 2.3-148.0 events/min and 3.4-217.5 events/min for elongation rates of 1, 2.6, 3.4, and 5 kb/min, respectively. Median values are indicated. C) We derived initiation rates (k_{init}) using values of premature termination and elongation rates (Eq. 12). The inter-decile ranges of initiation rates were 0.3-9.5 RNAP/min, 0.72-24.8 RNAP/min, 0.94-32.4 RNAP/min and 1.4-47.7 RNAP/min for 1, 2.6, 3.4, and 5 kb/min, respectively. Median values are indicated.

Fig. S11. Parameter sensitivity analysis indicates that the premature termination to pause release rate ratio is >1 if triptolide reduces initiation rate to less than 0.4 the control initiation rate. A) We varied the global decrease in k_{init} from 95% to 15%, corresponding to a fold change (FC) in k_{init} from 0.05 to 0.85 for repressed genes under TRP-inhibition (11). We set the elongation rate at a representative value of 2000 bp/min. We then calculated ratio of premature termination to pause release using Eq. 14. Median ratios are indicated. The inter-decile range at FC (k_{init}) = 0.05 is 4.0 to 108, decreasing to inter-decile range of 0.06-0.6 at FC (k_{init}) = 0.65 and 0.06-0.3 at FC (k_{init}) = 0.85. B) We estimated premature termination (k_{pre}) based on the ratio in (A). Considering elongation rate of 2000 bp/min, the pause release rate (Eq. 11) has a median of 3.9 events/min with an inter-decile range of 0.5-8.3 events/min. The inter-decile range of k_{pre} is 5.5-338 events/min at FC (k_{init}) = 0.05, decreasing to 0.05 - 2.7 events/min at FC (k_{init}) = 0.65 and 0.04-1.7 events/min at FC (k_{init}) = 0.85. C) We calculated initiation rate as indicated in Eq. 12. The inter-decile range changes from 1.5-62 RNAP/min at FC (k_{init}) = 0.05 which change to 0.16 - 1.5 RNAP/min at FC (k_{init}) = 0.65 and 0.15-1.4 RNAP/min at FC (k_{init}) = 0.85.

Fig. S12. Increasing signal from a fully occupied pause region decreases rates of pause release, initiation and premature termination. We selected saturation values at different percentiles of ranked pause sums for repressed genes under TRP-inhibition (11). To achieve 25-fold dynamic range around the saturation value considered at 90th percentile of ranked pause sums (P = 169, 90th percentile), we varied saturation values from 34 (at 48th percentile) to 844 (at 99.8th percentile). A) We calculated ratios of premature termination and pause release using Eq. 14. We set the foldchange (FC) in initiation rate to 0.25 and elongation rate to 2000 bp/min. The inter-decile ranges of ratio increases from 0.7-27.0 at 48th percentile to 1.44-31.5 at 99.8th percentile. B) The pause release rates (Eq. 11) have an inter-decile range of 0.9-11.6 events/min at 48th percentile to 0.3-7.24 events/min at 99.8th percentile. Median values are indicated on the boxplots. C) The premature termination rates (Eq. 14) have an inter-decile range of 1.4-133.9 events/min at 48th percentile to 1.2-74.7 events/min at 99.8th percentile. D) The initiation rates (Eq. 12) have an inter-decile range of 1.3-52.0 polymerase/min at 48th percentile and 0.1-3.2 polymerase/min at 99.8th percentile. E) We calculated the density of RNA Polymerase on gene body using Eq. 13. The RNA polymerases were less densely spaced on gene body with increasing signal of a fully occupied pause region. The inter-decile ranges were 1 polymerase every 585 bp - 5.4 kb at 48th percentile which decreased to 1 polymerase every 2.4 kb - 19.7 kb at 90th percentile and to 1 polymerase every 11.6 kb - 93.2 kb at 99.8th percentile.

Fig. S13. Most initiated RNA polymerases prematurely terminate. We categorized genes based on percentiles of paused polymerase density (A) and gene expression (B). We then estimated the ratio of initiation rate to effective premature termination ($k_{pre} \times P$) under untreated conditions. The median values ranged from 1.1 to 1.2, with a highly skewed distribution centered around 1. This suggests that the effective premature termination rate ($k_{pre} \times P$) is generally equivalent to k_{init}, regardless of paused polymerase density or gene expression levels.

Table S1

Treatment	Cell line	Species	Time	Citation	GEO Identifier
Triptolide	v6.5	Mouse	12.5 mins	Jonkers et al., 2014	GSE48895
Flavopiridol	v6.5	Mouse	25 mins	Jonkers et al., 2014	GSE48895
Dexamethasone	A549	Human	45 mins	Wissink et al., 2021	GSE168767
ZNF143 dTAG	HEK293T	Human	30 mins	Dong et al., 2024	GSE266491
TBP dTAG	HAP1	Human	2 hrs	Santana et al., 2022	GSE194153
Heat Shock	S2	Drosophila	20 mins	Duarte et al., 2016	GSE77607

Table S1. Gene Expression Omnibus (GEO) accession numbers for datasets used in this study.

Table S2

Study	Premature Termination	Pause Release
Zimmer et al., 2021	median 0.11 min^{-1}; inner 90% range 0.027–0.23 min^{-1}	median 0.027 min^{-1}; inner 90% range 0.0015–0.31 min^{-1}
Steurer et al., 2018	Frequency: 12 out of 13 Pol2 terminate at promoter proximal pause region (~92.4%, Fig. 3B)	Frequency: 1 out of 13 Pol2 proceed into elongation (7.6%, Fig. 3B)

Table S2. Multiple independent studies report faster premature termination rate compared to pause release.