SMC5/6-Mediated Plasmid Silencing is Directed by SIMC1-SLF2 and

Antagonized by LT

Martina Oravcová[1], Minghua Nie[1], Takanori Otomo[2,3], and Michael N. Boddy[1,3]

[1]Department of Molecular and Cellular Biology, The Scripps Research Institute, La Jolla,

CA 92037, USA.

[2]San Diego Biomedical Research Institute, San Diego, CA 92121, USA.

[3]Co-corresponding Authors: nboddy@scripps.edu and totomo@sdbri.org

Running Head: SIMC1 antagonizes plasmid gene expression.

Keywords: SIMC1, C5orf25, Nse5, SLF2, NSMCE5, SMC5, SMC6.

Abstract

SMC5/6 is unique amongst the Structural Maintenance of Chromosomes (SMC) complexes in its ability to repress transcription from extrachromosomal circular DNA (ecDNA), including viral genomes and plasmids. Previously, we showed that human SMC5/6 is regulated by two mutually exclusive subcomplexes—SIMC1-SLF2 and SLF1/2—the counterparts of yeast Nse5/6 (Oravcová, eLife, 2022). Notably, only SIMC1-SLF2 recruits SMC5/6 to SV40 Large T antigen (LT) foci in PML nuclear bodies (PML NBs), suggesting that these regulatory subcomplexes direct distinct roles of SMC5/6 on chromosomal versus ecDNA. However, their roles in plasmid repression remain unclear. Here, we demonstrate that SMC5/6-mediated repression of plasmid transcription depends exclusively on SIMC1-SLF2, whereas SLF1/2 is dispensable. Reinforcing its specialized role in ecDNA suppression, SIMC1-SLF2 does not participate in SMC5/6 recruitment to chromosomal DNA lesions. We further show that plasmid silencing requires a conserved interaction between SIMC1-SLF2 and SMC6, mirroring the functional relationship observed between yeast Nse5/6 and Smc6. As for viral silencing, plasmid repression depends on the SUMO pathway; however, unlike viral silencing, it does not require PML NBs. Additionally, we find that LT interacts with SMC5/6 and increases plasmid transcription to levels observed in SIMC1-SLF2-deficient cells–echoing the antagonistic roles of HBx (HBV) and Vpr (HIV-1) in viral genome repression. These findings expand the paradigm of viral antagonism against SMC5/6-mediated silencing, positioning LT as a novel player in this evolutionary tug-of-war.

Introduction

The Structural Maintenance of Chromosomes (SMC) complexes, present in both prokaryotes and eukaryotes, are core modulators of chromosome architecture and organization, supporting the development and homeostasis of all living organisms (1,2). The SMC1/3 (cohesin) and SMC2/4 (condensin) complexes ensure accurate genome propagation by folding chromosomes into distinct structures. Meanwhile, SMC5/6 plays a key role in maintaining genome stability and integrity, contributing to processes such as homologous recombination (HR), alternative lengthening of telomeres (ALT), and ribosomal DNA (rDNA) replication (3-7). Recently, SMC5/6 has also been recognized as a viral restriction factor, highlighting its broader role in managing extrachromosomal DNAs (ecDNAs, (8-10)).

Beyond viral restriction, the repression of transcription from simple plasmid DNA has also been added to the activities of the human SMC5/6 "Swiss Army knife" of DNA manipulation (8-12). SMC5/6 selectively represses the transcription of circular but not linear plasmid DNA (12), which is likely related to their topological states. That is, DNA that accumulates superhelical stress during transcription is the preferred binding substrate for SMC5/6, whereas linear DNA that dissipates such tension is not stably bound by the complex (12-15). Once DNA bound, SMC5/6 may further compact and thereby restrict transcriptional activity on the ecDNA.

All SMC complexes share a fundamental structure consisting of a SMC heterodimer connected by a kleisin protein. These complexes form a ring-like structure that traps DNA. The SMC proteins have a bent-arm shape and can open to encircle DNA when their ATPase head domains bind ATP, aiding DNA movement through the ring.

Kleisin plays a critical role for connecting and locking the SMC heads together (6,16-22).

The SMC5/6 complex is particularly adept at recognizing and compacting special DNA

structures, such as HR intermediates and supercoiled DNA (13,15,23,24).

The SMC5/6 complex is composed of a highly conserved "core" hexamer: the

SMC5/6 heterodimer and non-SMC-elements (NSE) 1-4 (9). A Kleisin (Nse4 in yeast;

NSMCE4 in human) and two interacting partners (Nse1 and Nse3 in yeast; NSMCE1 and

NSMCE3 in human) form a trimer that joins the SMC5/6 head domains. The SUMO ligase

Nse2 (NSMCE2 in human) attaches to the SMC5 arm. The complex's functional

specificity is further defined by non-core subunits, the Nse5/6 heterodimer (9,18,25).

Yeast Nse5/6 plays key roles in regulating Smc5/6's genome stability functions,

facilitating its chromatin loading, inhibiting its ATPase activity, and enhancing Nse2

SUMO ligase activity (19,26-32). Despite its critical regulatory roles, the identification of

a human Nse5/6-like complex lagged, largely due to the lack of cross-species protein

sequence conservation. Nevertheless, identified in proteomic approaches, the human

SLF1 (SMC5-SMC6 Complex Localization Factor 1)-SLF2 heterodimer (SLF1/2) was

shown to fulfill some genome stability roles performed by Nse5/6 in yeast (7). Interestingly,

however, with the possible exception of HIV-1 proviral DNA, the repression of viral DNA

transcription by SMC5/6 depends on SLF2 but not SLF1 (8,9,11). Therefore, SLF2 either

functions as a monomer in viral repression, or there is a cofactor that replaces SLF1 in

this role. Towards resolving this issue, we recently discovered SIMC1 as a missing link in

SMC5/6 regulation and demonstrated that the SIMC1-SLF2 complex is a human

counterpart of yeast Nse5/6 (33).

Our structural and mutational analyses revealed that SIMC1 and SLF2 contain

Nse5 and Nse6-like domains, respectively, which interact through a conserved interface.

SIMC1-SLF2 interacts with and directs SMC5/6 to subnuclear compartments at PML

nuclear bodies (PML NBs) induced by the SV40 polyomavirus large T antigen (LT).

Further, we showed that SIMC1-SLF2 and SLF1/2 are mutually exclusive subcomplexes

of SMC5/6 that likely support the distinct antiviral and DNA repair activities of the

holocomplex. Thus, SIMC1-SLF2 and SLF1/2 act as Nse5/6-like regulatory elements that

modulate the broad activities of the SMC5/6 complex (33).

Despite the foregoing advances, the specific roles of the SLF2-based SIMC1 and

SLF1 subcomplexes in SMC5/6 regulation remained unclear. Although we previously

proposed distinct functions for SIMC1-SLF2 and SLF1/2 in ecDNA silencing and DNA

repair, respectively, these mechanisms had yet to be demonstrated. Additionally,

SUMOylation, PML NBs, and SV40 LT were implicated in the regulation of SMC5/6-based

ecDNA silencing, but their contributions were not fully understood. Here, we address

these gaps directly. In line with our prior model, we show that SIMC1-SLF2, but not

SLF1/2, mediates plasmid silencing through a direct and evolutionarily conserved

interaction with SMC6. Conversely, SLF1/2 specifically recruits SMC5/6 to DNA lesions.

Furthermore, we demonstrate that plasmid silencing is facilitated by the SUMO pathway

but, unlike the SMC5/6-mediated silencing of HBV (34), does not require PML NBs. Lastly,

we reveal that SV40 LT binds SMC5/6 and antagonizes plasmid silencing, thereby adding

polyomaviruses to the growing list of pathogenic viruses that counteract SMC5/6-based

transcriptional repression.

Results

SIMC1-SLF2 associates with SMC6 through a conserved patch

Our previous structural analysis revealed that the Nse5-like domain of SIMC1 (SIMC1^{Nse5})

and the Nse6-like domain of SLF2 (SLF2^{Nse6}) both adopt α-solenoid-like structures (33).

These domains interact in a head-to-tail orientation, forming an ellipsoid-shaped dimer

(**Figure 1A**). This dimerization clusters conserved residues from the N-terminus of

SIMC1^{Nse5} and the C-terminus of SLF2^{Nse6} at one edge of the ellipsoid (**Figure 1B**). We

previously used a SIMC1 mutant harboring four substitutions in α1 of SIMC1 (SIMC1^{combo},

Figure 1B) to demonstrate that this conserved patch of SIMC1 is essential for binding to

SMC6 and for the localization of the SMC5/6 complex at LT-containing PML NBs (33).

In this study, we extended our analysis to SLF2 by introducing two sets of

mutations into the Nse6-like region of SLF2 (SLF2^{CTD}; residues 635-1173): **mut 1**

(E1121A/K1122A/K1125D/C1126R/E1130K in α16) and **mut 2**

(T1138R/K1141E/D1142R/A1145R/G1149E in the subsequent loop and α17). These

mutations were designed to disrupt charge interactions, hydrogen bonds, and van der

Waals contacts, following the same strategy used with SIMC1^{combo} to disrupt binding (33).

Using HEK293 cells co-expressing FLAG-tagged SLF2^{CTD} mutants, GFP-tagged SIMC1

and Myc-tagged SMC6, we performed GFP pull-downs followed by Western blot analysis

(**Figure 1C**). Wild-type SLF2^{CTD} interacted with both SIMC1 and SMC6 (Lane 1), whereas

both **mut 1** and **mut 2** disrupted SMC6 binding while preserving the interaction with

SIMC1 (Lanes 2, 3).

We also examined several control mutants. To confirm the specificity of the

SIMC1-SMC6 interaction, we introduced a new SIMC1 control mutant containing four

Figure 1: SIMC1-SLF2 composite surface patch is required for SMC6 association.
(A) The cryo-EM structure of the SIMC1-SLF2 complex (PDB ID: 7T5P) (33). Residues
mutated for interaction analysis are shown with side chains. **(B)** Conservation mapping
on the surface of SIMC1-SLF2 complex, showing the N-terminus of SIMC1 and the C-
terminus of SLF2. Conservation scores obtained from the Consurf server (35) are shown
by the color graduation as indicated. Amino acids mutated in SIMC1^{combo}
(R473D/N477A/E480K/E481K in α1 of SIMC1), SLF2 mut1
(E1121A/K1122A/K1125D/C1126R/E1130K in α16 of SLF2) and SLF2 mut2
(T1138R/K1141E/D1142R/A1145R/G1149E in α17 of SLF2) are labeled. **(C)** Western
blot of GFP-Trap immunoprecipitation from HEK293 cells transfected with plasmids
expressing GFP-SIMC1, Myc-SMC6, and FLAG-tagged C-terminal domain (CTD) of
SLF2, either WT or containing mutations (mut1, mut2). Input and GFP PD were detected
with anti-GFP, FLAG, or SMC6 antibody.

substitutions (Q842A/H846A/K849E/D857R) in α17, located at the opposite end of the

ellipsoid (**Figure 1A**). This mutant retained binding to both SLF2 and SMC6, indicating

that the residues altered in SIMC1[combo] are specifically required for SMC6 interaction

(**Figure 1—figure supplement 1A**). We additionally evaluated the most C-terminal

residues of SLF2 (1161–1173), which although conserved, are unstructured; however,

deleting these residues (SLF2[635-1160]) did not affect SMC6 binding (**Figure 1—figure**

supplement 1B). Collectively, these results confirm that the conserved surface patch

mediating the SIMC1-SLF2 interaction is essential for binding to SMC6.

SIMC1-SLF2 subcomplex contacts the neck region of SMC6

We used AlphaFold-Multimer (36) to predict the structure of the SIMC1-SLF2-SMC6

complex (**Figure 2A, Figure 2—figure supplement 1A**). The resulting high-confidence

model, as assessed by the predicted aligned error (PAE) plot (**Figure 2—figure**

supplement 1B), aligns well with mutational data and highlights the conserved surface

patch of SIMC1-SLF2 as the interface with SMC6 (**Figure 2A**). SMC6 is a long, hairpin-

like polypeptide in which its N- and C-termini fold together to form the ATPase "head"

domain at one end. The "hinge" domain at the bending end mediates heterodimerization

with SMC5 (**Figure 2—figure supplement 1A**). Connecting the head and hinge is a

flexible coiled-coil; the portion adjacent to the head is referred to as the "neck" (**Figure**

2A).

In the model, the conserved SIMC1-SLF2 surface patch interacts with C-terminal

residues of SMC6, centered on the neck and opposite the ATPase active site where it

dimerizes with SMC5's ATPase domain (21,30). This interface shows excellent shape

168
Figure 2: SIMC1-SLF2 subcomplex contacts the neck region of SMC6.
(A) The AlphaFold-Multimer model of the SIMC1^{Nse5}-SLF2^{Nse6}-SMC6 complex. The
disordered regions in the N- and C-termini, the hinge domain, and the majority of the
coilded-coil that do not contact SIMC1-SLF2 are omitted for clarity. The entire model and
the Predicted Aligned Error (PAE) plot of the prediction are shown in **Figure 2 - figure**
supplement 1. Residues mutated for interaction analysis are shown with side chains. **(B)**
A close-up view of the interface between SMC6's neck and the composite patch of
SIMC1-SLF2. **(C)** Western blot of GFP-Trap immunoprecipitation from HEK293 cells
transfected with plasmids expressing 2Myc-SIMC1, FLAG-SLF2^{CTD}, and WT or mutant
GFP-SMC6 (SIMC1 only: K942A, D946A, K957A; SLF2 only: R940A, Y944A, N947A;
Mixed: L939A, R940A, K942A, L943A). Input and GFP PD were detected with anti-GFP,
FLAG, or Myc antibody. **(D)** Schematic of yeast (left) and human (right) SMC5/6 complex
illustrating positions of its subunits within the complex and Nse5/Nse6 and SIMC1/SLF2
cofactors, respectively, interacting with the neck region of SMC6.

complementarity—SMC6's convex neck helix fits snugly into the concave surface formed

by α1 and α2 of SIMC1 and α17 of SLF2 (**Figure 2A**). Additional interactions arise from

SLF2's α16 and the α16-α17 loop with SMC6's head residues, spanning a total buried

surface area of 1516 Å².

This model explains the roles of the amino acids altered in our binding studies

(**Figure 2B; Figure 2—figure supplement 1C, D**). The four residues modified in

SIMC1^{combo} all make direct contact with SMC6's neck helix, including three salt bridges:

R473^{SIMC1}-D946^{SMC6}, E480^{SIMC1}-K957^{SMC6}/K942^{SMC6}, and E481^{SIMC1}-K942^{SMC6}. The SLF2

mutations also engage SMC6's neck region, forming salt bridges (E1130^{SLF2}-R940^{SMC6}

and K1141^{SLF2}-E1008^{SMC6}), a cation-π interaction (K1125^{SLF2}-Y944^{SMC6}), and multiple

van der Waals contacts (**Figure 2A; Figure 2—figure supplement 1C, D**). Thus, the

model and mutational data are consistent.

Next, we validated these SMC6's interface contacts using alanine substitutions.

We introduced three sets of mutations: 1) **SIMC1-only** (K942A/D946A/K957A), whose

wild-type residues form salt bridges with SIMC1; 2) **SLF2-only** (R940A/Y944A/N947A),

where R940 and Y944 interact with SLF2 while N947 forms salt bridges with SIMC1; 3)

Mixed (L939A/R940A/K942A/L943A), which alters two leucines that insert into a

hydrophobic cleft between α1 of SIMC1 and α17 of SLF2, along with two other residues

involved in salt bridges (**Figure 2B**). When these SMC6 mutants were co-expressed with

SIMC1 and SLF2, pull-down assays showed significantly reduced or abolished binding

(**Figure 2C, lane 2-4**). Thus, our mutational data strongly support the AlphaFold model.

We conclude that SIMC1-SLF2 associates with the neck region of SMC6, opposite

the ATPase active site used for heterodimerization with SMC5 (**Figure 2D; Figure 2—**

figure supplement 1A). A recent cryo-EM study on the *S. cerevisiae* Smc5/6 complex

similarly demonstrated that Nse5/6 engages with the neck region of Smc6 (30),

reinforcing the idea that this mode of interaction is evolutionarily conserved (**Figure 2D**).

This has implications for the function of SIMC1-SLF2 in the regulation of SMC5/6 activity,

based on the findings for yeast Nse5/6 (see discussion).

SMC5/6 mediated plasmid silencing requires the Nse5/6-like SIMC1-SLF2 complex

To examine the role of SIMC1 and SLF2 in SMC5/6-mediated episomal DNA silencing,

we used CRISPR-Cas9 to create U2OS SIMC1$^{-/-}$ or SLF2$^{-/-}$ cells. We compared the

expression of a transfected GFP reporter plasmid between wild-type (WT) and null cells.

GFP transcripts, quantified by qPCR, showed a strong increase over WT in both SIMC1$^{-/-}$

and SLF2$^{-/-}$ cells (**Figure 3A**). We also used fluorescence activated cell sorting (FACS)

to monitor GFP expression (37) and found that both SIMC1$^{-/-}$ and SLF2$^{-/-}$ cells expressed

significantly more GFP than WT (**Figure 3B**). In addition to U2OS cells, SIMC1$^{-/-}$ HEK293

cells also showed increased GFP reporter expression (**Figure 3—figure supplement 1**).

Importantly, expressing WT SIMC1 or SLF2 in the corresponding null cells reduced

reporter expression to levels observed in WT cells (**Figure 3B**), demonstrating the

involvement of SIMC1-SLF2 in plasmid silencing. Notably, however, SLF2^{mut1} that

abolishes SIMC1-SLF2 interaction with SMC6 (**Figure 1C**) did not reduce reporter

expression in SLF2$^{-/-}$ cells (**Figure 3B**). These findings demonstrate that SIMC1-SLF2,

and their interaction with SMC6, are required for plasmid repression.

228
229

Figure 3: SIMC1/SLF2-SMC6 interaction is critical for plasmid silencing.
(A) Expression of a GFP reporter transiently transfected into U2OS WT and SIMC1[-/-] or SLF2[-/-] cell lines was measured by quantitative PCR 72 hours after transfection. GFP expression was normalized to the expression of beta-actin (mean ± s.d. from n=4 independent experiments, two-tailed unpaired t-test; (**) $p<0.005$). **(B)** U2OS WT, SIMC1[-/-] or SLF2[-/-] cells with integrated empty vector or vector expressing SIMC1 or SLF2 variants (WT, mut1), respectively, were transiently transfected with GFP reporter. After 72 hours, GFP intensity was measured by FACS and is displayed relative to GFP intensity measured in WT cells. Data are the average ± s.d. from n=3 independent experiments, two-tailed unpaired t-test; (**) $p<0.005$). No significant difference ($p>0.05$) was found between WT and SIMC1[-/-] + pSIMC1 (left panel), or between WT and SLF2[-/-] + pSLF2 (right panel).

SIMC1 and SLF1 direct SMC5/6 to plasmid silencing and DNA repair, respectively

We previously showed that SLF1/2 and SIMC1-SLF2 form mutually exclusive complexes with SMC5/6 and recruit it to distinct cellular locations (33). Specifically, epitope-tagged SLF1 colocalized with gamma-H2AX at laser stripes, whereas SIMC1 did not. Here we more directly examined the roles of SLF1 and SIMC1 in recruiting SMC5/6 to DNA damage sites. To this end, we induced localized DNA lesions using laser microirradiation in WT, SLF1$^{-/-}$, and SIMC1$^{-/-}$ U2OS cells. In WT and SIMC1$^{-/-}$ cells SMC5/6 accumulated at laser-induced lesions marked by gamma-H2AX, whereas no such accumulation was seen in SLF1$^{-/-}$ cells (**Figure 4A**). Thus, SLF1 but not SIMC1 plays a critical role in directing SMC5/6 to DNA damage sites. Conversely, when comparing GFP reporter expression in WT and SLF1 null cells, we found no change in GFP levels in the absence of SLF1 (**Figure 4B**). Thus, unlike SIMC1, SLF1 does not play a role in SMC5/6-mediated plasmid silencing, which is consistent with what has been observed in SLF1$^{-/-}$ HepG2 cells (10).

260
Figure 4: SLF1 is involved in SMC5/6 DNA damage repair, not plasmid silencing.
(A) Representative immunofluorescence images of U2OS WT, SIMC1$^{-/-}$, SLF1$^{-/-}$ cells
exposed to laser microirradiation upon treatment with 50 uM angelicin and 1 ug/ml
Hoechst 33342 for 30 min. One hour after microirradiation, cells were pre-extracted, fixed
and stained with gamma-H2A.X (phosphorylation of histone H2A.X Ser139; green),
SMC6 (red) antibodies and DAPI (blue). Scale bar 10 µm. **(B)** GFP intensity was
measured by flow cytometry 72 hr after reporter transient transfection in U2OS WT cells
or SLF1$^{-/-}$ cells with integrated empty vector or stably expressing SLF1. Data are the
average ± s.d. from n=3 independent experiments, two-tailed unpaired t-test (p-
value >0.05).
271

SUMOylation supports plasmid repression

Given that SIMC1 contains SIMs, and the SMC5/6 complex is a SUMO ligase that

contains other SUMO-interacting interfaces, we also tested for a potential involvement of

the SUMO pathway in plasmid silencing (33,38,39). Cells were treated with the SUMO

pathway inhibitor (SUMOi), TAK981 (40). We observed a strong dose-dependent

increase in reporter gene expression in both U2OS and RPE cells (**Figure 5—figure**

supplement 2), supporting an involvement of the SUMO pathway in plasmid silencing.

This is consistent with the broadly observed transcriptionally repressive activity of

SUMOylation (41,42). Interestingly, SUMOi treatment does not significantly increase the

level of GFP reporter expression in SIMC1$^{-/-}$ and SLF2$^{-/-}$ over its effect on WT, despite the

pre-existing defect in plasmid silencing in SIMC1 and SLF2 null cell lines (**Figure 5A**).

The lack of additivity between SUMOi and SIMC1$^{-/-}$ or SLF2$^{-/-}$ suggests that SMC5/6-

mediated plasmid silencing operates in large part under the broad "umbrella" of

SUMOylation-based transcriptional repression.

288
Figure 5: Plasmid silencing depends on the SUMO pathway and LT, not PML NBs.
(A) U2OS WT, SIMC1$^{-/-}$ or SLF2$^{-/-}$ cells were transiently transfected with GFP reporter
and treated with 100 nM TAK-981(SUMOi) or DMSO at the same time. After 72 hours,
GFP intensity was determined by FACS. Data are normalized to WT cells treated with
DMSO and represent the average ± s.d. from n=4 independent experiments, two-tailed
unpaired t-test; (**) p<0.005. No significant difference was found between WT and
SIMC1$^{-/-}$ or SLF2$^{-/-}$ when treated with SUMOi (p>0.05). **(B)** U2OS WT or PML$^{-/-}$ cells with
integrated empty vector or vector expressing PML, respectively, were transiently
transfected with GFP reporter. Intensity of GFP was measured by FACS 72 hours after
transfection. Data are the average ± s.d. from n=3 independent experiments, two-tailed
unpaired t-test; (*) p<0.05). No significant difference (p>0.05) was found between WT and
PML$^{-/-}$ + pPML. **(C)** Western blot of GFP-trap immunoprecipitation from HEK293 cells
transiently transfected with either GFP-SMC5 or GFP alone in combination with SV40
vector expressing LT and Myc-SMC6. Signals were visualized using GFP, LT and SMC6
antibodies. **(D)** GFP intensity measured by FACS in U2OS WT, SIMC1$^{-/-}$ or SLF2$^{-/-}$ cells
with integrated empty vector, vector expressing large T antigen (LT) or LT-HR684 variant,
respectively that were transiently transfected with GFP reporter for 72 hours. Data are the
average ± s.d. from n=4 independent experiments. (∗) p<0.05; (∗∗∗) p<0.0005; (n.s.)
p>0.05 (two-tailed unpaired t-test).

PML NBs are not required for plasmid silencing

We previously showed that SIMC1-SLF2 directs SMC5/6 to LT antigen-containing SV40

replication centers at PML NBs (33). Moreover, SMC5/6-mediated restriction of HBV

occurs at PML NBs (34,43) and, the localization of SMC5/6 at PML NBs is SLF2-

dependent in HepG2 and PHH cells (10,43). We also now find that SLF2 is required for

the localization of SMC5/6 to PML NBs in U2OS cells (**Figure 5—figure supplement 1**).

Based on the foregoing, PML NBs have been proposed, but not directly shown, to support

simple plasmid repression (10). Therefore, we tested the impact of PML NBs on plasmid

silencing by measuring GFP reporter expression in WT and PML$^{-/-}$ U2OS cells (44).

Compared to SIMC1 and SLF2 null cells, we observed only a minor increase in GFP

expression in PML$^{-/-}$ versus WT cells (**Figure 5B**). Moreover, the stable re-expression of

PML in PML$^{-/-}$ cells did not significantly reduce the level of GFP expression, as compared

to vector control, suggesting the slight increase observed is due to clonal variation from

CRISPR editing (**Figure 5B**). Thus, we conclude that PML NBs are not needed for the

SMC5/6-mediated silencing of plasmids.

SV40 Large T antigen (LT) overcomes plasmid silencing

Considering that HBx and Vpr antagonize SMC5/6 to induce HBV and HIV-1 transcription,

respectively (45-47), we tested if SV40 LT similarly impacts plasmid silencing. In this

regard, we previously found that SMC5/6 colocalizes with LT-induced nuclear foci at PML

NBs in a SIMC1-SLF2-dependent manner (33). Here, we show that ectopically expressed

LT coimmunoprecipitates with GFP-SMC5/6 but not a GFP control (**Figure 5C**). Therefore,

LT, like HBx and Vpr, may contact SMC5/6 to antagonize its ecDNA repression activity.

Indeed, the stable expression of LT in WT cells promoted GFP reporter plasmid

expression, as compared to WT cells with an empty vector control (**Figure 5D; Figure 5**–

figure supplement 3). The host range defective mutant LT-HR684 had a similar effect

as WT LT, indicating that the C-terminus of LT, which inhibits the SV40 host range

restriction factor FAM111A (48), is not involved in overcoming plasmid repression (**Figure**

5D). Notably, the enhancement of plasmid DNA transcription by LT is not additive with

that already present in SIMC1$^{-/-}$ and SLF2$^{-/-}$ cells (**Figure 5D**). The epistatic relationship

between the positive effects of SIMC1$^{-/-}$, SLF2$^{-/-}$, and LT expression on plasmid

transcription indicates that LT overcomes SMC5/6-mediated repression.

Discussion

Our structural analyses indicate that SIMC1-SLF2 engages the SMC5/6 core at the neck

region of SMC6, situated opposite the ATPase active site. These findings, together with

recent cryo-EM analyses of the *S. cerevisiae* Smc5/6 complex (20,21,31), highlight a

conserved mode of interaction in which Nse5/6 or Nse5/6-like subunits bind to the

backside of SMC6's head domain. This binding site is ideal for allosteric regulation of the

ATPase and DNA-binding functions of the SMC5/6 family. Indeed, yeast Nse5/6 reduces

Smc5/6 ATPase activity and disrupts the Nse4 (kleisin)-Smc6 interface, which may open

the gate to enable topological DNA binding by Smc5/6 (**Figure 6**, (20,30,31)). The clear

overlap between our validated SIMC1-SLF2-SMC6 structure and that of yeast Nse5/6-

Smc6 suggests that SIMC1-SLF2, and its orthologue SLF1/2, may similarly modulate

SMC5/6 ATPase and DNA binding activities, albeit at distinct DNA targets.

354
Figure 6: SIMC1-SLF2 and SLF1/2 subcomplexes direct SMC5/6 to separate
pathways of plasmid silencing and DNA lesion repair. SIMC1-SLF2 and SMC5/6-
mediated plasmid silencing is facilitated by the SUMO pathway and antagonized by SV40
LT. PML NBs are not involved in plasmid silencing. S stands for SUMO; U for ubiquitin.
Both SIMC1-SLF2 and SLF1/2 likely contact SMC6 directly through their conserved
Nse5/6-like domains, but the SIMC1 SIMs and SLF1 BRCT/ARD domains contact distinct
posttranslational modifiers to direct the complex to its separate roles.
362

The distinct targeting functions of the SIMC1 and SLF1-based SLF2 complexes is

underscored by our current analysis of microirradiated cells that lack each complex

(**Figure 4**). While the deletion of SLF1 abrogates SMC5/6 localization at DNA lesions, the

loss of SIMC1 has no effect (**Figure 4A**). On the other hand, SIMC1 is required for the

recruitment of SMC5/6 to LT-containing PML NBs (33). Lastly, we found that SLF1 does

not impact plasmid silencing (**Figure 4B**), consistent with several analyses of HIV-1, rAAV,

and HBV silencing (10,43,45,49) but contrasting with another study (50). It remains

unclear why this latter discrepancy exists. While there may still be overlapping functions

of SIMC1 and SLF1 to uncover at the organismal level, it is clear they support distinct

cellular responses to ecDNAs and DNA lesions (**Figure 6**).

In keeping with the SUMO-binding ability of SIMC1 (**Figure 6**, (39)), we found that

the inhibition of SUMOylation by TAK-981 strongly enhanced plasmid transcription

(**Figure 5**–**figure supplement 2**). This is likely a broad effect that includes the loss of

repressive modifications of transcription factors (42); as well as the lifting of SMC5/6-

mediated repression to enhance ecDNA transcription. A similar effect of TAK-981 was

seen on HIV-1 proviral DNA gene expression, which was elevated by SUMO inhibition to

a similar degree as that observed in SMC5Δ cells (50). The authors concluded that the

effect of TAK-981 was entirely through the inhibition of SMC5/6-mediated SUMOylation

(NSE2 subunit). This is different to our results, which show that SUMOylation inhibition

has a larger effect than the loss of the SIMC1-SLF2 complex on plasmid gene expression.

The distinct natures of the ecDNAs under study, plasmid versus HIV-1 proviral DNA, and

the genetic backgrounds may contribute to this difference. Nevertheless, our data support

a key role of the SUMO pathway in plasmid repression, and that SMC5/6 works within it.

Despite the fact that SIMC1-SLF2 localizes SMC5/6 to PML NBs that contain SV40

LT (31), and HBV is restricted at PML NBs (34), they are dispensable for plasmid silencing.

While the reason for the different impact of PML NBs in HBV versus simple plasmid

silencing remain unknown, it may be related to the natural localization of each ecDNA.

That is, HBV localizes to PML NBs as part of the viral life cycle, in a manner dependent

on the SUMOylation of its core protein (51). In contrast, the direct visualization of plasmids

revealed they are pan-nuclear and apparently excluded from PML NBs (52). It is also

noteworthy that a recent genome-scale CRISPR screen for repressors of rAAV

transcription identified all components of the SMC5/6 complex, including SIMC1 and

SLF2, but not SLF1 or PML (49). This raises interesting questions about the mechanism

of SMC5/6-mediated transcriptional silencing, since it can occur outside PML NBs. In this

regard, SUMOylation was recently shown to support the recombination-based alternative

lengthening of telomeres (ALT) pathway in the absence of PML NBs (53), which are

normally key components of the fully active pathway (44). Therefore, the SUMOylation of

plasmid associated proteins, and or SMC5/6, may support transcriptional repression of

plasmids in the absence of PML NBs, consistent with our findings using TAK-981.

Interestingly, the expression of LT enhances plasmid transcription in a non-additive

manner with the deletion of SIMC1 or SLF2, suggesting an epistatic relationship. While

LT is known to be a promiscuous transcriptional activator (54,55) that does not rule out a

co-existing role in antagonizing SMC5/6. Indeed, these findings are reminiscent of HBx

from HBV and Vpr of HIV-1, both of which are known promiscuous transcriptional

activators that also directly antagonize SMC5/6 to relieve transcriptional repression

(12,45-47,56-58). Unlike HBx and Vpr, the interaction of LT with SMC5/6 does not appear

to induce the degradation of its subunits (33). Nevertheless, LT binds and sterically

inhibits several cellular proteins instead of inducing their degradation, including Rb and

p53 (59). Further analysis will be needed to define LT-SMC5/6 contacts, and its potential

mode of inhibition. This will have important ramifications for the lifecycle of the pathogenic

polyomaviruses since their LT proteins share functions with SV40 LT (60).

In conclusion, our structure-function analyses delineate a specific function for the

SIMC1-SLF2 complex in SMC5/6-mediated plasmid repression. This resolves a

discrepancy in the field, wherein SLF2 was thought to act alone in ecDNA repression,

since SLF1 is not required. While further studies are needed, SIMC1-SLF2 is most likely

responsible for supporting SMC5/6-mediated ecDNA silencing, including that of multiple

pathogenic viruses. Although HBV silencing requires PML NBs, we find that plasmid

silencing does not. This represents a departure from current dogma, suggesting that the

SMC5/6-mediated repression mechanism does not always rely on PML NB-resident

factors. In some cases, the SUMOylation of ecDNA-associated proteins may be sufficient

to support SMC5/6-based repression. Finally, because the SV40 LT antigen interacts with

SMC5/6 and antagonizes its repression of plasmids, the orthologous polyomaviral LT

antigens are likely to block SMC5/6-mediated restriction of pathogenic polyomaviruses.

Material and Methods

Construction of recombinant plasmids

Construction of plasmid DNA was described in (33). Mutations in SIMC1 (combo ctrl), SLF2 (mut1, mut2) and SMC6 (mixed, SIMC-only, SLF2-only) were either introduced in primers or a gBlock sequence containing desired mutations was purchased (IDT). All plasmids created in this study have been verified by sequencing service provided by Plasmidsaurus. Plasmid used in this study are listed in Appendix 1. Additional details of plasmid construction are available upon request.

Cell culture, transfection, stable line generation

All cell lines and their derivatives (see Appendix 1) were cultured in DMEM (Gibco) supplemented with 10% (v/v) fetal bovine serum (Omega Scientific), 1% (v/v) antibiotic-antimycotic (Gibco) and maintained at 37 °C in humidified air with 5% CO_2.

Transient plasmid transfections and generation of stable cell lines was detailed in (33). A complete list of stable mammalian cell lines generated, and reagents used in this study is provided (Appendix 1).

Cell line generation by CRISPR/Cas9

SIMC1, SLF1 and SLF2 knockout clones were generated via reverse transfection of CRISPR/Cas9 Ribonucleoprotein. A mixture of three sgRNAs specific for each gene (Synthego; 24 nM) was combined with 6 nM Cas9-NLS (Synthego) in reduced-serum medium OptiMEM and incubated for 10 min followed by a 15 min incubation with TransIT-X2 (Mirus). 6×10^4 cells were added and plated into 12-well plates. After 72 hours, cells

were diluted and plated to isolate single cell clones that were genotyped to confirm

successful editing. sgRNA sequences and diagnostic primers used in this study are listed

in Appendix 1.

Co-immunoprecipitation and Western blotting

GFP-labeled proteins were immunoprecipitated using the Nano-Trap magnetic agarose

(Chromotek) following the manufacturer's instructions. The protocol as well as whole cell

lysate preparation for SDS-PAGE and western blot are described in detail in (33).

Antibodies used for immunoblotting are listed in Appendix 1.

RNA extraction and quantitative PCR

For monitoring GFP expression by qPCR, cells were harvested for RNA isolation 72 hours

after GFP vector transient transfection. Total RNA precipitated using RNeasy Plus Mini

kit (Quiagen) was treated with DNaseI kit (Invitrogen) before cDNA synthesis by

SuperScript III First-Strand Synthesis System for RT-PCR (Invitrogen). SensiFAST SYBR

No-ROX kit (Meridian Bioscience) was used for qPCR and gene expression was

calculated as $2^{-\Delta Ct}$ using beta-actin as a reference. Oligonucleotides are listed in

Appendix 1.

Flow cytometry

GFP fluorescence measurements were based on (37). GFP reporter vector was

transiently transfected into cells using TransIT-LT1 (Mirus). After 72 hours, cells were

washed in PBS, dissociated using 0.25% trypsin and resuspended in DMEM. For SUMOi

experiments, cells were treated with 100 nM TAK-981 (Cayman Chemical) or 0.1%

DMSO in control plates at the time of GFP vector transfection. Median GFP intensity was

measured on Cytoflex S (Beckman Coulter) using the CytExpert software. A minimum of

1100 GFP-positive cells were analyzed.

DNA damage by laser microirradiation

Protocol adapted from (61). U2OS cells (1.5×10^5) were plated in a 35 mm µ-Grid dish

(Ibidi) a day before microirradiation. Cells were treated with 50 uM angelicin

(ThermoFisher) and 1 µg/ml Hoechst 33342 (Invitrogen) for 30 min before transferring to

the stage incubator of LSM880 Airyscan confocal laser scanning microscope (Zeiss).

Using the bleaching mode in ZEN software, DNA damage was induced by irradiation of

a 5-pixel wide region with a 405 nm diode laser (15% power, 1 iteration, zoom 1,

averaging 1, pixel dwell time 0.27 µs = speed 7). One hour after the irradiation, cells were

treated with extraction buffer (0.5% Triton X-100, 20 mM Hepes, 50 mM NaCl, 3 mM

$MgCl_2$, 300 mM sucrose) for 2 min, washed twice in PBS, fixed in 4% formaldehyde in

PBS for 15 min and stained with anti-phospho-Histone H2A.X (Ser139) and anti-SMC6

antibodies following the immunofluorescence protocol.

Immunofluorescence and microscopy

Immunofluorescence and confocal microscopy were done as described (33).

For optimal detection of SMC6, we pre-extracted the chromatin with extraction buffer (0.5%

Triton X-100, 20 mM Hepes, 50 mM NaCl, 3 mM $MgCl_2$, 300 mM sucrose) for 2 min on

ice. Antibodies used in this study are listed in Appendix 1.

Luciferase and MTT assays

U2OS cells were seeded at 4 x 10^3 cells per well of a 96-well plate and transfected with

20 ng of luciferase reporter plasmid (expresses secreted Gaussian luciferase under

EF1alpha promoter) using TransIT-LT1 (Mirus) by reverse transfection. TAK-981 in

indicated concentrations or DMSO were added at the time of transfection and Luciferase

activities were measured 3 days later. Ten microliters of culture media were mixed with

50 ul Gaussia Luciferase Glow Assay buffer that contains freshly diluted coelenterazine

substrate (Pierce^{TM} Gaussia Luciferase Glow Assay Kit). Luminescence was measured

after ten minutes on a Tecan Infinite M200 plate reader (Tecan Life Sciences). The

number of viable cells were estimated using MTT assay kit (Sigma) following

manufacturer's instructions.

AlphaFold-Multimer Structural modeling

510   AlphaFold-Multimer modeling (36) was performed using default settings on the following

511   protein sequences: SIMC1^{Nse5} (425-872), SLF2^{Nse6} (733-1173), and SMC6 (1-1091) .

Material Availability

514   All materials produced during this work are available upon written request, in keeping with

515   the requirements of the journal, funding agencies, and The Scripps Research Institute.

References

1.   Hirano T. At the heart of the chromosome: SMC proteins in action. Nat Rev Mol Cell Biol. 2006;7(5):311-22.

2.   Nasmyth K, Haering CH. The structure and function of SMC and kleisin complexes. Annu Rev Biochem. 2005;74:595-648.

3.   Peng XP, Lim S, Li S, Marjavaara L, Chabes A, Zhao X. Acute Smc5/6 depletion reveals its primary role in rDNA replication by restraining recombination at fork pausing sites. PLoS Genet. 2018;14(1):e1007129. PMCID: PMC5779651

4.   Potts PR, Yu H. The SMC5/6 complex maintains telomere length in ALT cancer cells through SUMOylation of telomere-binding proteins. Nat Struct Mol Biol. 2007;14(7):581-90.

5.   Wehrkamp-Richter S, Hyppa RW, Prudden J, Smith GR, Boddy MN. Meiotic DNA joint molecule resolution depends on Nse5-Nse6 of the Smc5-Smc6 holocomplex. Nucleic Acids Res. 2012;40(19):9633-46. PMCID: PMC3479181

6.   Aragon L. The Smc5/6 Complex: New and Old Functions of the Enigmatic Long-Distance Relative. Annu Rev Genet. 2018;52:89-107.

7.   Raschle M, Smeenk G, Hansen RK, Temu T, Oka Y, Hein MY, Nagaraj N, Long DT, Walter JC, Hofmann K, Storchova Z, Cox J, Bekker-Jensen S, Mailand N, Mann M. DNA repair. Proteomics reveals dynamic assembly of repair complexes during bypass of DNA cross-links. Science. 2015;348(6234):1253671.

8.   Irwan ID, Cullen BR. The SMC5/6 complex: An emerging antiviral restriction factor that can silence episomal DNA. PLoS Pathog. 2023;19(3):e1011180. PMCID: PMC9980727

9.   Peng XP, Zhao X. The multi-functional Smc5/6 complex in genome protection and disease. Nat Struct Mol Biol. 2023;30(6):724-34. PMCID: PMC10372777

10.   Abdul F, Diman A, Baechler B, Ramakrishnan D, Kornyeyev D, Beran RK, Fletcher SP,

Strubin M. Smc5/6 silences episomal transcription by a three-step function. Nat Struct Mol Biol.

2022;29(9):922-31.

11.   Liu HW, Roisne-Hamelin F, Gruber S. SMC-based immunity against extrachromosomal

DNA elements. Biochem Soc Trans. 2023;51(4):1571-83. PMCID: PMC10586767

12.   Diman A, Panis G, Castrogiovanni C, Prados J, Baechler B, Strubin M. Human Smc5/6

recognises transcription-generated positive DNA supercoils. Nat Commun. 2024;15(1):7805.

PMCID: PMC11379904

13.   Serrano D, Cordero G, Kawamura R, Sverzhinsky A, Sarker M, Roy S, Malo C, Pascal

JM, Marko JF, D'Amours D. The Smc5/6 Core Complex Is a Structure-Specific DNA Binding

and Compacting Machine. Mol Cell. 2020;80(6):1025-38 e5. PMCID: PMC8224240

14.   Jeppsson K, Pradhan B, Sutani T, Sakata T, Umeda Igarashi M, Berta DG, Kanno T,

Nakato R, Shirahige K, Kim E, Bjorkegren C. Loop-extruding Smc5/6 organizes transcription-

induced positive DNA supercoils. Mol Cell. 2024;84(5):867-82 e5.

15.   Gutierrez-Escribano P, Hormeno S, Madariaga-Marcos J, Sole-Soler R, O'Reilly FJ,

Morris K, Aicart-Ramos C, Aramayo R, Montoya A, Kramer H, Rappsilber J, Torres-Rosell J,

Moreno-Herrero F, Aragon L. Purified Smc5/6 Complex Exhibits DNA Substrate Recognition

and Compaction. Mol Cell. 2020;80(6):1039-54 e6. PMCID: PMC7758880

16.   Kim E, Barth R, Dekker C. Looping the Genome with SMC Complexes. Annu Rev

Biochem. 2023;92:15-41.

17.   Uhlmann F. SMC complexes: from DNA to chromosomes. Nat Rev Mol Cell Biol. 2016.

18.   Adamus M, Lelkes E, Potesil D, Ganji SR, Kolesar P, Zabrady K, Zdrahal Z, Palecek

JJ. Molecular Insights into the Architecture of the Human SMC5/6 Complex. J Mol Biol.

2020;432(13):3820-37.

565 19. Yu Y, Li S, Ser Z, Sanyal T, Choi K, Wan B, Kuang H, Sali A, Kentsis A, Patel DJ, Zhao

566 X. Integrative analysis reveals unique structural and functional features of the Smc5/6 complex.

567 Proc Natl Acad Sci U S A. 2021;118(19). PMCID: PMC8126833

568 20. Taschner M, Basquin J, Steigenberger B, Schafer IB, Soh YM, Basquin C, Lorentzen

569 E, Raschle M, Scheltema RA, Gruber S. Nse5/6 inhibits the Smc5/6 ATPase and modulates

570 DNA substrate binding. EMBO J. 2021;40(15):e107807. PMCID: PMC8327961

571 21. Li Q, Zhang J, Haluska C, Zhang X, Wang L, Liu G, Wang Z, Jin D, Cheng T, Wang H,

572 Tian Y, Wang X, Sun L, Zhao X, Chen Z, Wang L. Cryo-EM structures of Smc5/6 in multiple

573 states reveal its assembly and functional mechanisms. Nat Struct Mol Biol. 2024;31(10):1532-

574 42. PMCID: PMC11479838

575 22. Pradhan B, Kanno T, Umeda Igarashi M, Loke MS, Baaske MD, Wong JSK, Jeppsson

576 K, Bjorkegren C, Kim E. The Smc5/6 complex is a DNA loop-extruding motor. Nature.

577 2023;616(7958):843-8. PMCID: PMC10132971

578 23. Palecek JJ. SMC5/6: Multifunctional Player in Replication. Genes (Basel). 2018;10(1).

579 PMCID: PMC6356406

580 24. Ryu T, Spatola B, Delabaere L, Bowlin K, Hopp H, Kunitake R, Karpen GH, Chiolo I.

581 Heterochromatic breaks move to the nuclear periphery to continue recombinational repair. Nat

582 Cell Biol. 2015;17(11):1401-11. PMCID: PMC4628585

583 25. Palecek J, Vidot S, Feng M, Doherty AJ, Lehmann AR. The Smc5-Smc6 DNA repair

584 complex. bridging of the Smc5-Smc6 heads by the KLEISIN, Nse4, and non-Kleisin subunits. J

585 Biol Chem. 2006;281(48):36952-9.

586 26. Bustard DE, Ball LG, Cobb JA. Non-Smc element 5 (Nse5) of the Smc5/6 complex

587 interacts with SUMO pathway components. Biol Open. 2016;5(6):777-85. PMCID: PMC4920195

27.   Oravcova M, Gadaleta MC, Nie M, Reubens MC, Limbo O, Russell P, Boddy MN. Brc1

Promotes the Focal Accumulation and SUMO Ligase Activity of Smc5-Smc6 during Replication

Stress. Mol Cell Biol. 2019;39(2). PMCID: PMC6321882

28.   Pebernard S, Wohlschlegel J, McDonald WH, Yates JR, 3rd, Boddy MN. The Nse5-

Nse6 dimer mediates DNA repair roles of the Smc5-Smc6 complex. Mol Cell Biol.

2006;26(5):1617-30.

29.   Wan B, Wu J, Meng X, Lei M, Zhao X. Molecular Basis for Control of Diverse Genome

Stability Factors by the Multi-BRCT Scaffold Rtt107. Mol Cell. 2019;75(2):238-51 e5. PMCID:

PMC6745058

30.   Li S, Yu Y, Zheng J, Miller-Browne V, Ser Z, Kuang H, Patel DJ, Zhao X. Molecular

basis for Nse5-6 mediated regulation of Smc5/6 functions. Proc Natl Acad Sci U S A.

2023;120(45):e2310924120. PMCID: PMC10636319

31.   Hallett ST, Schellenberger P, Zhou L, Beuron F, Morris E, Murray JM, Oliver AW.

Nse5/6 is a negative regulator of the ATPase activity of the Smc5/6 complex. Nucleic Acids

Res. 2021;49(8):4534-49. PMCID: PMC8096239

32.   Etheridge TJ, Villahermosa D, Campillo-Funollet E, Herbert AD, Irmisch A, Watson AT,

Dang HQ, Osborne MA, Oliver AW, Carr AM, Murray JM. Live-cell single-molecule tracking

highlights requirements for stable Smc5/6 chromatin association in vivo. Elife. 2021;10. PMCID:

PMC8075580

33.   Oravcova M, Nie M, Zilio N, Maeda S, Jami-Alahmadi Y, Lazzerini-Denchi E,

Wohlschlegel JA, Ulrich HD, Otomo T, Boddy MN. The Nse5/6-like SIMC1-SLF2 complex

localizes SMC5/6 to viral replication centers. Elife. 2022;11. PMCID: PMC9708086

34.   Niu C, Livingston CM, Li L, Beran RK, Daffis S, Ramakrishnan D, Burdette D, Peiser L,

Salas E, Ramos H, Yu M, Cheng G, Strubin M, Delaney Iv WE, Fletcher SP. The Smc5/6

Complex Restricts HBV when Localized to ND10 without Inducing an Innate Immune Response

and Is Counteracted by the HBV X Protein Shortly after Infection. PLoS One.

2017;12(1):e0169648. PMCID: PMC5240991 does not alter our adherence to PLOS ONE

policies on sharing data and materials.

35. Ashkenazy H, Abadi S, Martz E, Chay O, Mayrose I, Pupko T, Ben-Tal N. ConSurf

2016: an improved methodology to estimate and visualize evolutionary conservation in

macromolecules. Nucleic Acids Res. 2016;44(W1):W344-50. PMCID: PMC4987940

36. Evans R, O'Neill M, Pritzel A, Antropova N, Senior A, Green T, Žídek A, Bates R,

Blackwell S, Yim J, Ronneberger O, Bodenstein S, Zielinski M, Bridgland A, Potapenko A,

Cowie A, Tunyasuvunakool K, Jain R, Clancy E, Kohli P, Jumper J, Hassabis D. Protein

complex prediction with AlphaFold-Multimer. bioRxiv. 2021:2021.10.04.463034.

37. Soboleski MR, Oaks J, Halford WP. Green fluorescent protein is a quantitative reporter

of gene expression in individual eukaryotic cells. FASEB J. 2005;19(3):440-2. PMCID:

PMC1242169

38. Cho T, Hoeg L, Setiaputra D, Durocher D. NFATC2IP is a mediator of SUMO-

dependent genome integrity. Genes Dev. 2024;38(5-6):233-52. PMCID: PMC11065178

39. Sun H, Hunter T. Poly-small ubiquitin-like modifier (PolySUMO)-binding proteins

identified through a string search. J Biol Chem. 2012;287(50):42071-83. PMCID: PMC3516753

40. Langston SP, Grossman S, England D, Afroze R, Bence N, Bowman D, Bump N, Chau

R, Chuang BC, Claiborne C, Cohen L, Connolly K, Duffey M, Durvasula N, Freeze S, Gallery M,

Galvin K, Gaulin J, Gershman R, Greenspan P, Grieves J, Guo J, Gulavita N, Hailu S, He X,

Hoar K, Hu Y, Hu Z, Ito M, Kim MS, Lane SW, Lok D, Lublinsky A, Mallender W, McIntyre C,

Minissale J, Mizutani H, Mizutani M, Molchinova N, Ono K, Patil A, Qian M, Riceberg J, Shindi

V, Sintchak MD, Song K, Soucy T, Wang Y, Xu H, Yang X, Zawadzka A, Zhang J, Pulukuri SM.

Discovery of TAK-981, a First-in-Class Inhibitor of SUMO-Activating Enzyme for the Treatment

of Cancer. J Med Chem. 2021;64(5):2501-20.

41.   Boggio R, Colombo R, Hay RT, Draetta GF, Chiocca S. A mechanism for inhibiting the

SUMO pathway. Mol Cell. 2004;16(4):549-61.

42.   Boulanger M, Chakraborty M, Tempe D, Piechaczyk M, Bossis G. SUMO and

Transcriptional Regulation: The Lessons of Large-Scale Proteomic, Modifomic and Genomic

Studies. Molecules. 2021;26(4). PMCID: PMC7915335

43.   Yao Q, Peng B, Li C, Li X, Chen M, Zhou Z, Tang D, He J, Wu Y, Sun Y, Li W. SLF2

Interacts with the SMC5/6 Complex to Direct Hepatitis B Virus Episomal DNA to Promyelocytic

Leukemia Bodies for Transcriptional Repression. J Virol. 2023;97(7):e0032823. PMCID:

PMC10373549

44.   Loe TK, Li JSZ, Zhang Y, Azeroglu B, Boddy MN, Denchi EL. Telomere length

heterogeneity in ALT cells is maintained by PML-dependent localization of the BTR complex to

telomeres. Genes Dev. 2020;34(9-10):650-62. PMCID: PMC7197349

45.   Dupont L, Bloor S, Williamson JC, Cuesta SM, Shah R, Teixeira-Silva A, Naamati A,

Greenwood EJD, Sarafianos SG, Matheson NJ, Lehner PJ. The SMC5/6 complex compacts

and silences unintegrated HIV-1 DNA and is antagonized by Vpr. Cell Host Microbe.

2021;29(5):792-805 e6. PMCID: PMC8118623

46.   Decorsiere A, Mueller H, van Breugel PC, Abdul F, Gerossier L, Beran RK, Livingston

CM, Niu C, Fletcher SP, Hantz O, Strubin M. Hepatitis B virus X protein identifies the Smc5/6

complex as a host restriction factor. Nature. 2016;531(7594):386-9.

47.   Murphy CM, Xu Y, Li F, Nio K, Reszka-Blanco N, Li X, Wu Y, Yu Y, Xiong Y, Su L.

Hepatitis B Virus X Protein Promotes Degradation of SMC5/6 to Enhance HBV Replication. Cell

Rep. 2016;16(11):2846-54. PMCID: PMC5078993

48.   Fine DA, Rozenblatt-Rosen O, Padi M, Korkhin A, James RL, Adelmant G, Yoon R,

Guo L, Berrios C, Zhang Y, Calderwood MA, Velmurgan S, Cheng J, Marto JA, Hill DE, Cusick

ME, Vidal M, Florens L, Washburn MP, Litovchick L, DeCaprio JA. Identification of FAM111A as

an SV40 host range restriction and adenovirus helper factor. PLoS Pathog.

2012;8(10):e1002949. PMCID: PMC3475652

49.   Ngo AM, Puschnik AS. Genome-Scale Analysis of Cellular Restriction Factors That

Inhibit Transgene Expression from Adeno-Associated Virus Vectors. J Virol.

2023;97(4):e0194822. PMCID: PMC10134838

50.   Irwan ID, Bogerd HP, Cullen BR. Epigenetic silencing by the SMC5/6 complex

mediates HIV-1 latency. Nat Microbiol. 2022;7(12):2101-13. PMCID: PMC9712108

51.   Hofmann S, Plank V, Groitl P, Skvorc N, Hofmann K, Luther J, Ko C, Zimmerman P,

Bruss V, Stadler D, Carpentier A, Rezk S, Nassal M, Protzer U, Schreiner S. SUMO

Modification of Hepatitis B Virus Core Mediates Nuclear Entry, Promyelocytic Leukemia Nuclear

Body Association, and Efficient Formation of Covalently Closed Circular DNA. Microbiol Spectr.

2023;11(3):e0044623. PMCID: PMC10269885

52.   Mearini G, Nielsen PE, Fackelmayer FO. Localization and dynamics of small circular

DNA in live mammalian nuclei. Nucleic Acids Res. 2004;32(8):2642-51. PMCID: PMC419472

53.   Zhao R, Xu M, Yu X, Wondisford AR, Lackner RM, Salsman J, Dellaire G, Chenoweth

DM, O'Sullivan RJ, Zhao X, Zhang H. SUMO promotes DNA repair protein collaboration to

support alternative telomere lengthening in the absence of PML. Genes Dev. 2024;38(13-

14):614-30. PMCID: PMC11368244

54.   Sullivan CS, Pipas JM. T antigens of simian virus 40: molecular chaperones for viral

replication and tumorigenesis. Microbiol Mol Biol Rev. 2002;66(2):179-202. PMCID:

PMC120785

55.   Gilinger G, Alwine JC. Transcriptional activation by simian virus 40 large T antigen:

requirements for simple promoter structures containing either TATA or initiator elements with

variable upstream factor binding sites. J Virol. 1993;67(11):6682-8. PMCID: PMC238107

56.   Qadri I, Conaway JW, Conaway RC, Schaack J, Siddiqui A. Hepatitis B virus

transactivator protein, HBx, associates with the components of TFIIH and stimulates the DNA

helicase activity of TFIIH. Proc Natl Acad Sci U S A. 1996;93(20):10578-83. PMCID: PMC38195

57.   Aufiero B, Schneider RJ. The hepatitis B virus X-gene product trans-activates both

RNA polymerase II and III promoters. EMBO J. 1990;9(2):497-504. PMCID: PMC551692

58.   Felzien LK, Woffendin C, Hottiger MO, Subbramanian RA, Cohen EA, Nabel GJ. HIV

transcriptional activation by the accessory protein, VPR, is mediated by the p300 co-activator.

Proc Natl Acad Sci U S A. 1998;95(9):5281-6. PMCID: PMC20252

59.   Ahuja D, Saenz-Robles MT, Pipas JM. SV40 large T antigen targets multiple cellular

pathways to elicit cellular transformation. Oncogene. 2005;24(52):7729-45.

60.   Googins MR, An P, Gauthier CH, Pipas JM. Polyomavirus large T antigens: Unraveling

a complex interactome. Tumour Virus Res. 2024;19:200306. PMCID: PMC11720896

61.   Tampere M, Mortusewicz O. DNA Damage Induction by Laser Microirradiation. Bio-

protocol 2016;6(23):e2039.

Acknowledgements

This work was supported by NIH grants R35 GM136273 to M.N.B. and GM092740 to T.O.

Author information

Affiliations

Department of Molecular and Cellular Biology, The Scripps Research Institute, La Jolla, CA 92037, USA.

Martina Oravcová, Minghua Nie & Michael N. Boddy

San Diego Biomedical Research Institute, San Diego, CA 92121, USA.

Takanori Otomo

Contributions

Conceptualization, M.N.B., T.O. Design of Methodology, M.O., M.N., M.N.B., T.O. Writing – Original Draft, M.N.B., M.N., T.O. Writing – Review & Editing, M.O., M.N., T.O., M.N.B. Formal analysis/Investigation, M.O., M.N., M.N.B., T.O. Validation, M.O., M.N. Resources, M.O., M.N. Visualization, M.O., M.N., T.O., M.N.B. Funding acquisition, M.N.B., T.O. Supervision, M.N.B., T.O. Project administration, M.N.B.

Corresponding authors:

Correspondence to Michael N. Boddy and Takanori Otomo.

727 **Ethics declarations**

728 Competing interests

729 The authors declare no competing interests.

Supplementary figures:

732
733
Figure 1 - figure supplement 1.
(A) Western blot of GFP-trap immunoprecipitation from HEK293 cells transiently
transfected respectively with GFP-SIMC1, GFP-SIMC1 combo mutant or GFP-SIMC1
combo control mutant in combination with FLAG-SLF2^{CTD} and Myc-SMC6. Signals were
detected using FLAG, SMC6 and GFP antibodies. **(B)** Western blot of GFP-trap
immunoprecipitation from HEK293 cells transiently transfected with either FLAG-SLF2^{CTD}
or FLAG-SLF2$^{635-1160}$ co-transfected with GFP-SIMC1 and Myc-SMC6. Signals were
visualized using FLAG, Myc and GFP antibodies.
742

743
744
Figure 2 - figure supplement 1
(A) The AlphaFold-Multimer model of the SIMC1[Nse5] (425-872)-SLF2[Nse6] (733-1173)-
SMC6 (1-1091) complex. The entire model is shown, colored according to pLDDT values
(left) or in the same manner as in **Figure 2A**. **(B)** The Predicted Aligned Error (PAE) plot
of the AlphaFold-Multimer prediction. **(C, D)** Close-up views of the SMC6 neck-SIMC1 (C)
and SMC6 neck-SLF2 (D) interfaces.

751
752 **Figure 3 - figure supplement 1**
753 HEK293 WT or SIMC1$^{-/-}$ cells were transiently transfected with GFP reporter and GFP
754 intensity was assessed 72 hours later by FACS. Data are the average ± s.d. from n=6
755 independent experiments, two-tailed unpaired t-test; (***) p<0.0005.

756
Figure 5 - figure supplement 1
Representative immunofluorescence images of U2OS WT or SLF2$^{-/-}$ cells fixed and
stained with SMC6 (green) and PML (red) antibodies along with DAPI (blue). Scale bar
10 µm.

Figure 5 - figure supplement 2

Arbitrary luminescence units of U2OS (**A**) or RPE (**B**) cells expressing transfected pLuc reporter. Cells were treated with TAK-981 or DMSO at the time of transfection. Means and error bars (s.d.) were derived from a minimum of n=3 independent transfections representing biological replicates. Two-tailed unpaired t-test was performed ((∗∗) p<0.005; (∗∗∗) p<0.0005).

768
Figure 5 - figure supplement 3
Immunoblot from U2OS WT, SIMC1$^{-/-}$ or SLF2$^{-/-}$ cells stably expressing large T antigen
(LT), LT-HR684 variant or have an empty vector integrated, respectively. PSTAIR serves
as a loading control.

List of Supplementary files

Figure 1—figure supplement 1. A, B

Figure 2—figure supplement 1. A-D

Figure 3—figure supplement 1

Figure 5—figure supplement 1.

Figure 5—figure supplement 2. A, B.

Figure 5—figure supplement 3.

List of source data

Figure 1 – source data 1. Full and unedited blots corresponding to panel C.

Figure 2 – source data 1. Full and unedited blots corresponding to panel C.

Figure 3 – source data 1. GFP transcripts qPCR corresponding to panel A.

Figure 3 – source data 2. GFP FACS corresponding to panel B.

Figure 4 – source data 1. GFP FACS corresponding to panel B.

Figure 5 – source data 1. GFP FACS corresponding to panel A.

Figure 5 – source data 2. GFP FACS corresponding to panel B.

Figure 5 – source data 3. Full and unedited blots corresponding to panel C.

Figure 5 – source data 4. GFP FACS corresponding to panel D.

Figure 1 – figure supplement 1 – source data 1. Full and unedited blots corresponding

to panel A.

Figure 1 – figure supplement 1 – source data 2. Full and unedited blots corresponding

to panel B.

Figure 3 – figure supplement 1 - source data 1. GFP FACS.

Figure 5 – figure supplement 2 - source data 1. Luciferase measurement corresponding

to panel A.

Figure 5 – figure supplement 2 - source data 2. Luciferase measurement corresponding

to panel B.

Figure 5 – figure supplement 3 - source data 1. Full and unedited blots.

802
803