Alectinib enhances response to RBM39 degradation via SRPK1 inhibition

Authors: Yurui Ma[1], Evon Poon[2], Chenchen Jin[1], Barbara Martins da Costa[2], Yuewei Xu[1,3],

Sadiya Quazi[1], Nikolaos Zourdoumis[1], Chiharu Wickremesinghe[1], Louis Chesler[2], Hector C.

Keun[1]*, Anke Nijhuis[1]*¥

Affiliations:

[1]Department of Surgery & Cancer, Imperial College London; London, UK.

[2]Division of Clinical Studies, The Institute of Cancer Research, London, UK

[3]Wisdom Lake Academy of Pharmacy, Xi'an Jiaotong-Liverpool University, Suzhou 215123,

China (current affiliation)

*Co-corresponding author

¥Lead contact: Dr Anke Nijhuis

Address: Division of Cancer, Imperial College London, London, UK, W12 0NN.

Email: a.nijhuis@imperial.ac.uk   **Phone**: +44 (0) 20 7594 3161

One Sentence Summary: Dual RNA splicing interference by indisulam and alectinib enhances

sensitivity in MYCN-driven malignancies such as high-risk neuroblastoma, increasing DNA

damage and R-loop accumulation.

Abstract: RNA splicing is a critical process in gene expression regulation and RNA splicing

factors are increased in expression in many cancers. We have previously demonstrated that the

molecular glue indisulam degrades RNA splicing factor RBM39 producing aberrant splicing

transcripts that is very effective in high-risk neuroblastoma and can potentiate the sensitivity to DNA damaging agents in ovarian high-grade serous carcinoma. RNA splicing inhibitors have the ability to generate vulnerabilities that can be exploited for combinatorial therapeutic strategies. In this study, we employed combination drug screens to identify potential synergistic drug-drug interactions between indisulam and other FDA approved clinical compounds. We demonstrate that alectinib, a clinically approved ALK inhibitor, has strong potency against serine/arginine protein kinase 1 (SRPK1), a key modulator of RNA splicing. Combined treatment with alectinib and indisulam leads to synergistic cell growth inhibition, increased cell apoptosis and cell cycle arrest in multiple cancer cell lines including high-risk neuroblastoma. Co-treatment exacerbated RNA splicing defects in genes involved in DNA damage response and R-loop resolution, leading to R-loop accumulation and increased DNA damage. In the Th-MYCN/ALK^{F1174L} neuroblastoma mouse model, combination therapy induced complete tumour regression and significantly improved survival rates compared with monotherapies.

Thus, the combination of indisulam and alectinib is a promising approach to treat aggressive malignancies such MYCN-driven high-risk neuroblastoma, exploiting the hitherto untapped pharmacology of alectinib as a clinical RNA splicing inhibitor. Our work demonstrates the concept that co-targeting of functionally interdependent RNA splicing factors is a strategy that can yield new and highly synergistic anti-cancer combination therapies effective against otherwise hard-to-treat cancers.

Main Text:

INTRODUCTION

The splicing of precursor messenger RNAs (pre-mRNAs) is a critical regulatory step in gene expression, during which non-coding introns are excised to produce functional mRNA(*1*). This process is tightly controlled by splicing factors, governing the recognition and selection of splice sites, thereby generating mRNAs with varying exon combinations and consequently diverse protein isoforms(*1*). Dysregulated alternative splicing can promote tumorigenesis and chemoresistance. Aryl sulfonamides such as indisulam have emerged as promising anticancer agents by targeting a pre-mRNA splicing factor RBM39 (RNA binding motif 39)(*2, 3*). These molecular glues facilitate the association of RBM39 with the CUL4-DCAF15 ubiquitin ligase, resulting in RBM39 polyubiquitination and subsequent proteasomal degradation(*2, 4*). Suppressing RBM39 efficiently inhibits tumour growth and advancement through modulating the transcription of various genes associated with tumorigenesis (reviewed in (*3*)).

One cancer type particularly vulnerable to splicing perturbation is neuroblastoma (*5-8*), a common and aggressive extracranial solid tumour in young children, which represents a significant contributor to mortality in children with cancer(*9*). High-risk neuroblastoma, comprising approximately 15% of all neuroblastoma cases, represents a distinct subtype characterized by aggressive tumour behaviour and an unfavourable prognosis(*10*). One pivotal factor contributing to the progression of high-risk neuroblastoma is the amplification of the MYC family transcription factor, MYCN, which can block the differentiation of neural progenitors, the key event in neuroblastoma initiation(*11*), and plays a crucial role in facilitating the transcriptional repression of tumour suppressor genes(*9, 10*). Furthermore, oncogenic gain-of-function mutations in the *ALK* (Anaplastic Lymphoma Kinase) occur in 10–15% of primary tumours, making ALK the most common single-gene alteration in high-risk neuroblastoma for which targeted clinical therapies have been developed(*9, 12*) Alectinib, an FDA-approved ATP-competitive ALK tyrosine kinase

inhibitor (TKI),was initially developed for non-small-cell lung cancer and has shown considerable

promise in addressing neuroblastoma in extensive preclinical studies(*13-15*). However, tumour

heterogeneity and resistance to treatment with ALK inhibitors (and molecularly-targeted drugs in

general) limits their use as single agents (*16*). Thus, novel strategies, particularly combination

therapies, remain important as a means to reduce the toxicity of any single drug and overcome

treatment resistance more efficiently. This approach is particularly advantageous in childhood

cancer, where long-term toxicity is a major lifelong issue (*17*).

Our previous study(*6*) and others(*8*) have recognized the exquisite sensitivity of high-risk

neuroblastoma to indisulam, with complete responses observed in multiple xenografts, patient-

derived xenografts (PDX) and transgenic mouse models. Notably, our study reported that cells

with MYCN amplification displayed heightened sensitivity to aryl sulfonamides in comparison to

those without MYCN amplification. This could be attributed to the increased reliance of

neuroblastoma cells on RNA processing due to elevated N-Myc/c-Myc activity, rendering them

more susceptible to RNA splicing interference (*18*). Through the induction of splicing errors in

genes within key oncogenic pathways, indisulam can render cancer cells susceptible to additional

disruptions caused by other anticancer drugs. For example, we have previously shown that several

key DNA damage repair (DDR) genes are mis-spliced after RBM39 depletion (such as *BRCA1*,

RAD51D and *ATM*), and therefore that indisulam treatment can increase sensitivity to PARP

inhibitors in ovarian high-grade serous carcinoma (*19*).

In this study, we aimed to identify additional vulnerabilities through drug-drug library screens

combining indisulam with FDA approved anti-cancer agents. Through this, we demonstrate a

novel synergistic interaction between indisulam and ALK-inhibitor alectinib, likely due to activity

against a recently discovered secondary target of alectinib, RNA Serine-Arginine Protein Kinase

1 (SRPK1) (*20*). SRPK1 promotes the nuclear translocation and activation of a family of SR

splicing factors (SRSFs). The combination of indisulam and alectinib simultaneously depletes

RBM39 and inhibits SRPK1-mediated SRSF phosphorylation, resulting in exacerbated RNA

splicing defects. The subsequent failure to resolve accumulated DNA damage and R loops results

in cell cycle arrest, cell growth inhibition and apoptosis of neuroblastoma cells *in-vitro* and tumour

regression *in vivo*. Our study provides important proof-of-concept evidence for the co-targeting of

splicing factors as a viable strategy to treat aggressive malignancies such as MYCN-driven high-

risk neuroblastoma.

RESULTS

Combination drug screens revealed synergy between indisulam and alectinib.

We previously demonstrated that indisulam-mediated RBM39 degradation causes wide-spread

RNA splicing defects, and an enrichment in aberrant expression of proteins that mediate DNA

damage repair and cell cycle regulation, across various cancer types(*6, 19*). This led us to speculate

that indisulam is likely to generate drug-induced-vulnerabilities that potentiate the efficacy of

existing, clinically approved anti-cancer compounds, such as PARP inhibitors(*19*). RBM39

degraders, such as indisulam and E7820, have been trialed in 47 clinical studies(*3*), highlighting

potential therapeutic combinations ready for clinical evaluation.

To explore this concept, we conducted two drug-library screens combining indisulam with 147

FDA-approved anticancer drugs (NCI plate number #4891, #4892) in an indisulam-sensitive cell

line (HCT116, colorectal cancer) and an indisulam-resistant line (KURAMOCHI, ovarian cancer).

Anticancer compounds were tested at 5 doses ranging from 16nM to 10μM based on the average

GI50 value, and combined with DMSO or indisulam at a fixed dose (GI30=50nM, for HCT116,

10μM for KURAMOCHI) (**Fig. 1A, Fig. S1A**). Bliss scores(*21*) were calculated to identify

synergistic interactions with indisulam. Depletion of RBM39 by indisulam is dependent on

DCAF15 expression, thus we generated CRISPR-Cas9 knockouts of *DCAF15* to identify RBM39-

specific drug-drug interactions (HCT116 *DCAF15* wild-type (*DCAF15*^{WT}) clone #1 and #2 and

DCAF15 knock-out clones (*DCAF15*^{KO}) clone #1 and #2). Drugs with Bliss score>0 in both

HCT116-*DCAF15*^{WT} clones were selected as hits which were further validated in triplicate

experiments with selected doses in both HCT116-*DCAF15*^{WT} and *DCAF15*^{KO} clones #1 and #2

and ranked by mean Bliss scores of both cell lines. Most of the synergistic drug-drug interactions

were rescued in HCT116-*DCAF15*^{KO} cells suggesting that the synergy is dependent on *DCAF15*

and RBM39 degradation (**Fig. 1B left**). In KURAMOCHI, the top 20 hits were ranked by mean

Bliss scores across all doses (**Fig. 1B right**). The clinical ALK inhibitor alectinib demonstrated

strong synergy in both HCT116 (top hit) and KURAMOCHI (second hit) and was therefore

selected for further validation using a wider range of drug concentrations. Strong synergistic

growth inhibition was observed across a range of doses of indisulam and alectinib in HCT116 (**Fig.**

1C) and KURAMOCHI (**Fig. 1D**), and the synergy was abrogated in HCT116-*DCAF15*^{KO} cells.

These data confirm that the combination of alectinib with RBM39 degraders exert strong

synergistic inhibition of cell growth in HCT116 and KURAMOCHI cells and this highlights an

opportunity for the development of a novel combination therapy across genomically diverse cancer

types.

Figure 1. Combination drug screens revealed a synergistic growth inhibition of indisulam and alectinib. (A) Workflow of combination drug screens in HCT116 and KURAMOCHI using indisulam and 147 FDA-approved anticancer compounds. **(B)** Top 20 hits from drug screens in HCT116 (left) and KURAMOCHI (right). Two HCT116-*DCAF15*^{WT} or HCT116-*DCAF15*^{KO} cell lines treated with selected hits with or without indisulam. Bliss scores were calculated based on cell growth measured by SRB assay. Hits are ranked by mean bliss scores of both *DCAF15*^{WT} clones. Bliss score > 0 (red) indicates synergism, n=3 replicates. KURAMOCHI cells were treated with indicated doses of drugs, with or without indisulam; Top 20 hits are ranked by mean bliss score. Bliss score > 0 (red) indicates synergism, n=3 replicates. **(C-F)** Cell lines used for synergy screen **(C, D)** and indicated HRNB cell lines **(E, F)** were treated with alectinib combining indisulam for 72 hours and analysed for cell growth by SRB. Synergy scores were calculated by Synergyfinder (https://synergyfinder.fimm.fi/) using the Loewe method.

Indisulam and alectinib exhibit synergistic anticancer efficacy in high-risk neuroblastoma

Considering the potency demonstrated by both indisulam and alectinib as monotherapies in high-risk neuroblastoma(*6, 14*), we further investigated the combination of indisulam and alectinib in four high-risk neuroblastoma cell lines (KELLY, IMR32, SKNBE2C, IMR5) (**Fig. 1E-F**). We demonstrate that combining alectinib with indisulam significantly inhibits cell proliferation compared to monotherapies and display great synergy across a wide-range doses of both compounds on neuroblastoma cell lines, but only additive or antagonistic effects were found in KELLY-*DCAF15*^{KO} cell line confirming the necessity of RBM39 degradation for this pharmacological interaction (**Fig. 1E right**). Similar synergy profiles were observed between alectinib and E7820, a second aryl sulfonamide RBM39 degrader (*2*)(**Fig. S1B-C**).

In addition, alectinib treatment alone did not induce cell apoptosis, but significantly increased the percentage of apoptotic cells when combined with indisulam in KELLY and IMR32 (**Fig. 2A-B, Fig. S2D**). Correspondingly, a significant increase of cleaved caspase 3 was detected in the

combination treatment group in KELLY cells (**Fig. 2C**). To evaluate cell cycle progression, we

used flow cytometry analysis and detected a notable accumulation of cells in the G2/M phases

following 48-hour treatment of indisulam which is consistent with a previous study(*8*). Co-

treatment with indisulam and alectinib increased the percentage of cells in the G1-phase, implying

enhanced cell cycle arrest in both G2/M and G1 phase, with a substantial decrease in the percentage

of cells in S-phase (**Fig. 2D-E**). These data confirm that indisulam and alectinib act synergistically

to suppress the proliferation of neuroblastoma cells, disrupt cell cycle and induce cell apoptosis.

Figure 2. Indisulam and alectinib induce apoptosis and cell cycle disruption in high-risk neuroblastoma. (A-B) Apoptosis analysis via flow cytometry in KELLY cells following exposure to 1.25μM indisulam, 4μM alectinib, or their combination for 48 hours. Cells were labelled by Annexin V and propidium iodide (PI). The significance of difference of total apoptotic cells was calculated by one-way ANOVA. (**C**) Western blot of KELLY cells treated with indicated doses of indisulam and alectinib for 48 hours. (n=3 independent experiments) significance of difference was calculated by one-way ANOVA. (**D-E**) Profiling of cell cycle distribution in KELLY cells was performed using flow cytometry after a 48-hour treatment with indisulam, Alectinib, or the combination. Cellular DNA content was labelled with PI, while replicating cells were labelled with nucleotide analogues 5-Iodo-2-deoxyuridine (IdU). Quantitative analysis of cell cycle distribution (n=3 independent experiments). Two-way ANOVA were employed to determine the statistical significance among the groups. n.s. not significant, * $p < 0.05$, **$p < 0.01$, ***$p < 0.001$, ****$p < 0.0001$.

ALK inhibition by alectinib is not required for synergistic interaction with indisulam.

According to data from the Human Protein Atlas (proteinatlas.org) (*22*) ALK and RET, both

primary targets of alectinib, are expressed in NB cells but not in HCT116 and KURAMOCHI (**Fig.**

3A, Table S1) in which synergy was initially discovered. Western blotting confirmed an absence

of ALK expression in these cells (**Fig. 3B**). To further test whether the synergy is driven by ALK

inhibition, we transfected KELLY, HCT116, and KURAMOCHI cells with an siRNA targeting

ALK (siALK) prior to indisulam dosing (**Fig. 3B**). We found no significant difference in cell

growth inhibition between siALK and non-targeting control (siNTC) transfected cells (**Fig. 3C**),

indicating that ALK inhibition alone does not increase sensitivity to indisulam. We also tested a

second ALK inhibitor with a distinct chemical scaffold and binding site, lorlatinib, and found that

this compound did not enhance sensitivity to indisulam, except in KELLY cells, which express a

mutant, oncogenic form of ALK (ALK^{F1174L}) (**Fig. 3D and Fig. S2A**). However, in cell lines with

wild type ALK (IMR32) or lacking ALK expression (HCT116 and KURAMOCHI) there was no

sensitization (**Fig. 3D**), consistent with other studies (*23*), indicating that the efficacy of lorlatinib

is specific to ALK mutated neuroblastoma. Further experimental validation is needed to clarify the

interaction between inhibition of mutant ALK-driven tumours and RBM39 depletion by indisulam.

Nonetheless, these findings suggest that ALK inhibition is not a significant contributor to the

synergy we observe between alectinib and the RBM39 degraders indisulam and E7820. This

suggests that synergy with RBM39 degradation relates primarily to inhibition of a secondary target

of alectinib.

Figure 3. ALK inhibition by alectinib is not an essential determinant of indisulam synergy. (**A**) RNA expression

data as normalized transcript per million (nTPM) values of cancer cell lines were obtained from proteinatlas.org. (**B**)

Western blot confirming the knock-down of ALK gene expression in cells post transfection with siNTC or siALK. (**C**)

Indicated cell lines were transfected with siALK or siNTC for 16 hours, followed by exposure to indisulam for 72

hours. Cell growth was measured by SRB assay. (**D**) Indicated cell lines were treated with lorlatinib and indisulam

concentration matrix for 72 hours. Loewe synergy scores are plotted. n=3 replicates.

Alectinib exerts potent inhibition of SRPK1.

Comprehensive kinase inhibitor screens indicate that alectinib inhibits an array of additional kinases other than ALK (*20*). The observation that alectinib is a potent inhibition of Serine/arginine (SR) Protein Kinase 1/2 (SRPK1/2) (*20, 24*) is particularly relevant to our research of aberrant RNA splicing. Both SRPK1 and 2 play a central role in regulation of alternative splicing via phosphorylation and activation and nuclear translocation of the Serine/Arginine-rich Splicing Factors (SRSFs) (*25, 26*). In MYCN-amplified neuroblastoma, SRPK1 expression is elevated and associated with reduced survival rates (**Fig. S2B-C**). In contrast, lower survival rates are not associated with increased expression of SRPK2 nor SRPK3, indicating an increased dependency on SRPK1 to drive tumour growth in MYCN-amplified tumours. Furthermore, a genome-wide CRISPR study showed that loss of SRPK1 sensitizes to indisulam treatment in various solid tumour cell lines (*27*). Thus, we further explored the treatment combination of SRPK1i and RBM39 degraders. To this end, we assessed the synergy between indisulam and SPHINX31, a known SRPK1 inhibitor(*28*). Co-treatment of SPHINX31 and indisulam exhibited synergistic growth inhibition cancer across all three cell lines (**Fig. 4A**). Time-lapse cell confluency monitoring in KELLY-*DCAF15*^WT and *DCAF15*^KO showed that co-treatment significantly suppressed the growth of KELLY-*DCAF15*^WT, compared to individual drug treatments. Conversely, in the absence of *DCAF15*, the combination therapy displayed similar potency to SPHINX31 monotherapy (**Fig. 4B**), suggesting this interaction is DCAF15-dependent and thus RBM39-dependent.

The inhibition of SRPK1 by alectinib was confirmed by enzyme activity assays which reported a half-maximal inhibitory concentration (IC50) of 26nM. This inhibition is more potent than that of

SPHINX31 (IC50 = 375nM) with lorlatinib showing very little inhibitory action (**Fig. 4C**).

Notably, alectinib inhibited SRPK2 with an IC50 of 700nM while SPHINX31 showed no

inhibitory effect (**Fig. S2D**). Comparison of the crystal structures of the SRPK1-Alectinib and

SRPK1-SPHINX31 complexes confirmed that alectinib and SPHINX31 interact with the ATP-

binding groove of SRPK1, with reasonable affinity and occupying similar positions within the

binding site (**Fig. 4D** and **Fig. S3**).

SRPK1 phosphorylates SRSFs, triggering their nuclear translocation and activity in modulating

RNA splicing (**Fig. 4E**). Both alectinib and SPHINX31 reduced phosphorylated SRSF (p-SRSF)

in KELLY, HCT116 and KURAMOCHI cells (**Fig. 4F)** and decreased the nuclear-to-cytoplasmic

ratio of p-SRSF intensity within 4 hours in KELLY cells, while lorlatinib lacked this effect (**Fig.**

4G). SRSFs are known to promote selection of the proximal splice site in *VEGF* pre-mRNA and

inhibiting SRPK1 alters *VEGF* splicing to generate the *VEGF165b* isoform(*20, 29*). Treatment of

both alectinib and SPHINX31 resulted in a switch and increased *VEGF165b* mRNA in high-risk

neuroblastoma cells (**Fig. 4H**).

In conclusion, our findings reveal that alectinib is a potent inhibitor of SRPK1, consequently

modulating downstream function and RNA splicing through a reduction in the phosphorylation of

SRSF splicing factors. Notably, alectinib demonstrates more robust inhibition of SRPK1 as

compared to the selective inhibitor SPHINX31, an intriguing observation that could potentially

expedite the clinical repurposing of alectinib as an SRPK1 inhibitor.

Figure 4. SRPK1 inhibition is synergistic with indisulam in high-risk neuroblastoma. (**A**) Indicated cell lines

were treated with indisulam and SPHINX31. Loewe synergy scores as indicated. (**B**) KELLY-*DCAF15*^{WT} and

DCAF15^{KO} cells were treated with indisulam and SPHINX31 individually or in combination for 72 hours. Cell

confluence was monitored every 6 hours using the Incucyte live cell imaging system. One-way ANOVA was used to

determine significance. (**C**) SRPK1 enzyme activity measured with ADP-Glo™ Kinase Assay with indicated doses of

compounds. (**D**) Alectinib (pink) docked into SRPK1 complexed with SPHINX31(orange). (**E**) Diagram depicting

Alectinib inhibition against SRPK1. Made with Biorender. (**F**) Western blotting of p-SRSF in KELLY, HCT116 and

KURAMOCHI post-treatment. (**G**) Immunofluorescent staining of p-SRSF in KELLY post-treatment. Nuclear

intensity and cytoplasmic staining intensity are calculated using DAPI and Phalloidin (F-actin) segmentation. Scale

bar = 20μm. (**H**) qRT-PCR of *VEGF165b* and total *VEGF165* in KELLY and IMR32 cells normalised to GAPDH. *

$p < 0.05$, **$p < 0.01$, ***$p < 0.001$. n = 3 independent experiments.

Alectinib and indisulam exacerbates RNA splicing defects, increase R-loop formation and

DNA damage.

Proteomic data from our previous study (*6*) showed that indisulam exposure leads to an increase

in protein expression of most SRSF family members (**Fig. S4A**). Here, we observed enhanced

nuclear translocation of p-SRSF post-indisulam treatment, which was rescued in *DCAF15*^{KO} cells

(**Fig. S4B-C**). These observations suggest that induction of the SRPK1/SRSF-pathway regulates

alternative splicing and is an adaptive response to indisulam treatment in cells compensating for

the loss of RBM39. RBM39 has previously been identified as a potential target of SRPK1 [25] and

we observed a direct interaction between RBM39 and SRPK1 in KELLY cells via Proximity

Ligation Assay (PLA) (**Fig. S4D**). We therefore hypothesize that the synergism observed between

alectinib and indisulam arises as a result of enhanced disruption of RNA splicing, and dual

targeting of key spliceosomal components with functional interdependence, thereby reducing the

opportunity for adaptive resistance.

We have previously demonstrated that degradation of RBM39 with indisulam leads to RNA splicing defects preferentially targeting genes implicated in DNA damage and HR repair (*BRCA1* and *RAD51D*), Fanconi Anaemia (*FANCI, FANCD2*) and Non-Homologous End Joining repair pathways (*TP53BP1*, and *RIF1*)(*6, 19*). To understand the consequences of dual inhibition of RBM39 and SRPK1 on these RBM39 targets, we tested the abundance of RNA splicing alterations in these genes through custom PCR primer design of affected exon regions (**Fig. 5A**). We found that RBM39 targets were increasingly affected when combined with SRPK1 inhibition either with alectinib or SPHINX31 in KELLY and IMR32 (**Fig. 5A-B, Fig. S5A**). This enhancement in mis-splicing was not found when indisulam was combined with siALK transfection or lorlatinib (**Fig. S5B-D**). In some cases, for example *CHK1*, the expression of the non-spliced canonical isoform was almost completely lost with co-treatment of indisulam with either alectinib or SPHINX31. Moreover, CHK1 protein was depleted following indisulam and alectinib combination treatment (**Fig. S5E**), These data confirm a complementary function of SRPK1 and RBM39 in RNA splicing. It also suggests that the synergy observed between alectinib and indisulam may be attributed to the compound reduction of canonical isoform expression and enhanced functional loss in effected pathways, such as DNA damage repair.

To assess whether the exacerbated impact on aberrant RNA splicing of DNA damage repair related genes led to increased DNA damage, we tested the expression of γH2AX as a marker for DNA double stranded breaks(*30*). γH2AX foci detected by immunofluorescence (**Fig. 5C-D**) and western blot analysis (**Fig. S6A**) demonstrated a significant increase following co-treatment of indisulam and alectinib compared to monotherapies. Furthermore, γH2AX expression levels did

not increase when indisulam was combined with lorlatinib, compared to indisulam alone (**Fig.**

S6B-C), again, demonstrating that increased DNA damage following treatment with alectinib and

indisulam is due to the dual targeting of RNA splicing factors, rather than ALK inhibition.

R-loops are DNA:RNA hybrids that are a product of intermediate transcription and must be

resolved to permit progression of replication forks (*31, 32*). R-loop accumulation results in

transcription-replication collisions, which generate replication stress and double-strand breaks

(*33*). Importantly, DNA damage repair-related factors play pivotal roles in resolving R-loops. For

example, BRCA1 interacts with the putative RNA helicase Senataxin (SETX), forming a complex

that localizes to R-loops(*34*). Several Fanconi Anemia genes recruit RNA processing enzymes to

resolve R-loops generated by moderate replication stress(*35*), while RIF1 facilitates replication

fork protection and efficient restart to maintain genome stability(*36*). Notably, SR proteins

regulated by SRPK1 can impede R-loop formation, ensuring faithful transcriptional elongation

and enhancing genomic stability(*31*). Dysregulation of these functions lead to accumulation of

unresolved R-loops, replication fork breakage, and genome instability (*31*).

We propose that aberrant splicing of DNA damage repair genes and inhibition of SRPK1 result in

R-loop accumulation, subsequent DNA damage and ultimately cell death. In KELLY cells, we

therefore quantitated R-loops using S9.6 antibody(*37*). Treatment with either alectinib or

SPHINX31 led to increased nuclear S9.6 staining indicating the accumulation of R-loops. The

combination treatment significantly increased R-loop formation as compared to monotherapies

(**Fig. 5E-F, Fig. S6D-F**). Since unresolved R loops impede the progression of replication forks,

causing transcription replication conflicts, and subsequent replication stress(*31, 38*), we quantified

the intensity of phosphorylated RPA32 serine 33 (pRPA), a marker of single-stranded DNA (ssDNA) at stalled replication forks. We observed an increase following exposure to the combination treatment relative as compared to monotherapy (**Fig. 5G, Fig. S6E**). To directly examine altered fork dynamics, we performed DNA fibre analysis using BrdU and IdU labelling (**Fig. S6G-H**). This uncovered a reduction in track length and the significantly decreased ratio of IdU to BrdU when treating cells with indisulam and alectinib, indicating increased fork stalling and replication stress. Collectively, these findings suggest that the combination of indisulam and alectinib or SRPK1 inhibitor leads to the accumulation of unresolved R-loops, promoting replication stress, and contributing to subsequent DNA damage in high-risk neuroblastoma.

Figure 5. Combination of indisulam and alectinib induce DNA damage, R-loop accumulation and replication stress. (**A**) PCR validation of exon skipping events in DDR-related genes (representative image of n=3). *GAPDH* was included as a control. (**B**) Densitometry analysis of alternative splicing products, Two-way ANOVA was used to detect the significance. (**C-D**) Immunofluorescence staining of γH2AX in KELLY cells treated with indisulam in combination of alectinib. Representative images of KELLY treated with 4μM alectinib and 1.25μM indisulam are shown. Scale bars = 20μm. Nuclear foci >5 indicates positive cells. n=3 independent experiments. (**E**) Representative images for nuclear S9.6 and RPA32 (S33) staining in KELLY cells, n = 3 independent experiments, Scale bars = 2.5μm. (**F-G**) Quantification of nuclear S9.6 intensity (F) and pRPA32 (G) intensity in KELLY cells. Cells were treated with 4μM alectinib monotherapy or with 1.25μM indisulam; nuclear signal intensity were measured by merging to DAPI; n>300 cells from 3 independent experiments were quantified. Significance was analysed among the median values.

Combination of indisulam and alectinib improves survival in a Th-MYCN/ALK^{F1174L} neuroblastoma model

Gain-of-function mutations in ALK account for 10-15% of high-risk neuroblastoma (*9*). ALK inhibitors have been shown to be safe and effective in treating ALK-driven neuroblastoma in a Phase I trial(*39, 40*), and are currently being investigated in several Phase III trials(*39, 41, 42*), highlighting their potential for clinical use in ALK-positive neuroblastoma. Considering the clinic relevance and benefit, we used an allograft mouse model implanted with Th-MYCN/ALK^{F1174L} tumour cells derived from genetically engineered mouse models (GEMMs) (**Fig.6A**).

Tumour-bearing mice were treated (indisulam 10mg/kg, alectinib 40mg/kg or combination) for 2-weeks with a 45-day follow-up. Both monotherapies significantly enhanced survival however all alectinib treated mice relapsed on treatment and 5/8 mice in the indisulam monotherapy group relapsed after withdrawal of treatment. Complete tumour regression without subsequent relapse

resulting in a 100% survival rate were reported in the combination group (**Fig. 6B-C**) consistent

with the increased efficacy in-vitro.

Tumour samples were collected for pharmacodynamic analysis after 3 days of treatment. Western

blotting showed a clear depletion of RBM39 following indisulam and combination therapy,

confirming the on-target effect of indisulam (**Fig. 6D**). Furthermore, treatment with 10 mg/kg

indisulam reduced MYCN expression, as previously reported by Singh et al (*43*), suggesting a

potential link between RBM39 inhibition and suppression of MYCN (**Fig. 6D**). Additionally, a

decrease in p-ALK **(Fig. S7A)**, and a reduction in p-SRSF (**Fig. 6F**) confirmed the on-target effects

of alectinib on SRPK1 and ALK. Double strand breaks measured by accumulation of γH2AX foci,

were increased in combination therapies (**Fig. 6G**), in line with our observations *in-vitro.* However,

indisulam alone did not significantly induce γH2AX accumulation, possibly due to the transient

nature of γH2AX focus formation during the DNA damage response. However, clear nuclear

fragmentation was observed in mice treated with indisulam and combination therapy (**Fig. 6G**).

Importantly, no significant weight loss was observed during the trial, indicating no significant

toxicity (**Fig. S7B)**, and we observed no significant RBM39 loss in other organs after 3-day

treatment (**Fig. S7C**).

Taken together, these data confirm that in neuroblastoma the therapeutic benefit of RBM39-

depleting agents can be enhanced through combination with SRPK1 inhibitors, such as alectinib.

This study provides proof-of concept for the co-targeting of functionally interdependent splicing

factors using repurposed clinical agents to more effectively target aggressive childhood cancers

such as neuroblastoma. The strategy could apply more widely to other oncogene-driven cancers

that exhibit a high dependency on faithful splicing of mRNA.

Figure.6. Combination of indisulam and alectinib suppress neuroblastoma tumour growth. (**A**) Th-ALK^F1174L/Th-MYCN GEMM tumour cells were injected into a cohort of mice subcutaneously, intravenous (i.v.) administration of 10 mg/kg of indisulam and oral gavage (p.o.) 40 mg/kg of alectinib for 2 weeks with 5 days on and 2 days off. (**B**) Tumour volume measured 3 times a week until the humane end-point was reached. Cyan shade indicates the treatment period. n = 8 for each group. (**C**) Survival plot of mice bearing Th-ALK^F1174L/Th-MYCN allografts treated with vehicle or alectinib or indisulam or combination therapy. (**D**) Western blotting in tumour tissues after administration of vehicle or alectinib or indisulam or combination therapy for 3 days (n = 4 per group). (**E-H**) Immunofluorescence staining was performed on tumour sections, with DAPI used for nuclear staining. Nuclear p-

396 SRSF intensity (E-F) was measured by merging to DAPI. Nuclear γH2AX (G-H) foci >5 indicates positive cells. Scale

397 bar = 20 μm. n.s. not significant, * $p < 0.05$, **$p < 0.01$, ***$p < 0.001$, ****$p < 0.0001$.

DISCUSSION

Indisulam has been recently identified as a molecular glue capable of regulating RNA alternative

splicing(*2*). By inducing splicing errors of genes in key oncogenic pathways, indisulam and other

spliceosomal inhibitors may render cancer cells susceptible to additional disruptions caused by

other anticancer drugs. Thus, it holds potential for implementation as combination therapy. We

performed extensive drug screens with FDA-approved anticancer agents which revealed several

potential drug-drug interactions with indisulam. Notably, the synergy between indisulam and

PARP inhibitors, also observed in our screens, has been corroborated by other studies in high-

grade ovarian carcinoma(*19*), has also been verified independently in colorectal cancer, and breast

cancer(*44*), giving confidence in our screening approach. Our study also highlights the fact that

such screens can reveal beneficial pharmacological interactions based on off-target effects (targets

other than the primary focus of clinical development), presenting novel rationales for drug

repurposing.

High-risk neuroblastoma, which constitutes about 15% of all neuroblastoma cases, is characterized

by aggressive behaviour and poor prognosis(*10*). Gain-of-function mutations in ALK account for

10-15% of primary tumours (*9*), hence a number of trials have investigated the use of alectinib and

other ALK inhibitors in this clinical setting. While the more selective ALK inhibitor loratinib has

shown greater traction for the treatment of mutant ALK-driven neuroblastoma(*39*), treatment

resistance remains a significant clinical challenge(*45*). Our study demonstrates that the activity of

alectinib against SRPK1 can be exploited therapeutically in combination with other spliceosomal

inhibitors. This presents an interesting strategy to treat patients with emerging resistance to ALK

inhibitor therapy, including in lung cancer where alectinib has demonstrated much greater clinical

use(*41*) and beyond into other cohorts irrespective of ALK status. While other inhibitors of SRPKs

and related SRSF regulators such as Cirtuvivint(*46*) and TG-003(*47*) are currently in early

development, the fact that an established therapeutic tool can be repurposed in this way could

accelerate the clinical testing of SRPK1/2 inhibition and co-targeting of splicing factors as an anti-

cancer treatment strategy, particularly in tumours where an ALK inhibitor might normally be

indicated.

While we have successfully demonstrated the principle of co-targeting functionally interdependent

splicing as an anticancer strategy, several resulting questions arise. Our investigation into the

functional consequences of combined RBM39 degradation and SRPK1 inhibition underscores

their interdependency in regulating alternative splicing, but it is not yet clear if RBM39 is a direct

target of SPRK1 as has been suggested by global interaction profiling (*48*) and our co-localisation

experiments. More broadly, we need further investigation to understand the interactions and co-

regulation of different splicing factors that are also cancer targets such as RBM39 and SPRK1,

and to develop more detailed mechanistic models of how these regulate alternative splicing

together.

Interestingly, when alectinib/SRPK1 inhibition was used alone, there was no detectable induction

of DDR genes mis-splicing observed in neuroblastoma. This is in contrast to interference with

RBM39 function(*19*) and what we might expect if the effect of alectinib was acting purely through

direct regulation of RBM39 activity and the mis-splicing of a specific gene family. As with the

targeting of spliceosomal components in general, it is unclear if the sequelae of accumulating R

loops, cell cycle arrest, DNA damage and eventual cell death that we observe in combined

SRPK1/RBM39 interference are lineage-dependent, result from mis-splicing and loss of function

of a select set of genes, or the accumulative impact of hundreds or even thousands of splicing

events within a functional pathway. Further work to resolve such mechanistic questions will be

important to understand what the most relevant pharmacodynamic readouts will be for future

discovery experiments and which predictive biomarkers can best identify patients most likely to

benefit from such combinations (i.e. for stratification).

In summary, the present study elucidates a novel therapeutic approach utilizing the combination

of indisulam and alectinib in high-risk neuroblastoma, and our results support a model in which

co-treatment indisulam and alectinib synergistically impacts RNA splicing through simultaneous

RBM39 degradation and SRPK1 inhibition. This dual RNA splicing interference exacerbates

splicing errors in key genes involved in DNA damage repair and in R-loop resolution, leading to

R-loop accumulation and increased DNA damage. These events eventually result in cellular

apoptosis and growth inhibition in high-risk neuroblastoma *in vitro*, as well as suppression of

tumour growth *in vivo*. These insights shed new light on translating research on spliceosome

inhibition into clinical applications, ultimately aiming to improve the prognosis for patients with

high-risk neuroblastoma.

MATERIALS AND METHODS

Study design

The objectives of this study were to identify the synergistic mechanism of action and therapeutic potential of indisulam combined with alectinib in high-risk neuroblastoma. We conducted drug screens in cancer cell lines and confirmed synergy in neuroblastoma models. Genetic and pharmacological experiments determined whether dual interference with RNA splicing—via RBM39 degradation and SRPK1 inhibition—could exacerbate RNA splicing defects in DNA damage repair genes, leading to enhanced DNA damage, R-loop accumulation, apoptosis, and tumor suppression. Animals were treated with indisulam and alectinib for two weeks. Sample sizes were based on prior experience with model growth profiles, providing sufficient statistical power to detect differences. Tumour volumes were measured every 2 to 3 days, and animals were euthanized upon reaching humane endpoints or impaired health criteria.

Cell lines and cell culture

Human neuroblastoma cell lines were cultured in were cultured in MEM (Gibco™) supplemented with 10% fetal bovine serum (FBS), 1% penicillin/streptomycin (P/S), 1mM sodium pyruvate and 2 mM L-Glutamine. All cell lines were maintained at 37 °C with 5% CO_2 in a controlled environment.

Sulforhodamine B (SRB) assay

Cells at the experiment's endpoint were fixed with 10% (w/v) cold trichloroacetic acid (TCA) (Sigma) and stained with 0.4% protein-bound SRB dye for 30 minutes. Excess dye was removed by washing the cells three times with 1% (v/v) acetic acid. The bound SRB dye was then solubilized in 10 mM Tris-Base solution, and the optical density (OD) was measured at 565 nm

using a ClarioStar microplate reader (BMG Labtech). The results were normalized to the vehicle

control values, with the control set as 100%.

Synergy analysis

Cell growth inhibition was assessed by SRB assay after 72 hours of drug exposure. The inhibition

was calculated by normalizing the mean OD values of each treatment group to vehicle control.

The Loewe additivity model was used to describe the interaction between two agents[32]. Cells

were treated with specific concentration matrices of the drugs, and cell growth data were analysed

using SynergyFinder 2.0 (https://synergyfinder.fimm.fi) to compute synergy scores. A synergy

score greater than 10 indicates synergism, a score between -10 and 10 represents additivity, and a

score below -10 reflects antagonism.

Western blotting|

Whole-cell lysates were prepared using RIPA buffer (Sigma Aldrich) with 1% HALT protease and

phosphatase inhibitor cocktail (Thermo Scientific). Equal amounts of protein from each sample

were separated on a 4-20% Bis-Tris SurePAGE gel (GenScript) and then transferred onto PVDF

membranes. The membranes were blocked with 5% milk for 1 hour at room temperature, followed

by an overnight incubation at 4°C with CHK1 (#2360, Cell Signaling Technology), p-CHK1

(#2348, Cell Signaling Technology), γH2AX (#05–636, Sigma-Aldrich). After washing, the

membranes were incubated with horseradish peroxidase (HRP)-conjugated secondary antibody for

1 hour at room temperature. Chemiluminescent detection was performed using SuperSignal West

Pico PLUS Chemiluminescent Substrate (Thermo Scientific).

End-point and quantification PCR reactions

1500ng RNA was utilized for reverse transcription, employing the High-Capacity RNA-to-cDNA™ Kit (Applied Biosystems) in accordance with the provided instructions. The resulting cDNA templates were subsequently employed for qPCR using thePowerTrack SYBR Green Master Mix (Applied Biosystems) or End-point PCR utilizing the Q5® High-Fidelity PCR Kit (New England Biolab). Primers were specifically designed to investigate alternative splicing events of the targeted genes. The PCR products were separated by electrophoresis using 2% agarose gels with SYBRsafe DNA gel stain, and the images were captured for densitometry analysis by Fiji V2.0.0.

Target genes	Primer (5' to 3')	Tm
BRCA1	AACTCTGAGGACAAAGCAGC CTTCTCAGTGGTGTTCAAATC	62
CHK1	GTGGAAGACTGGGACTTGGT ACCCCTGCCATGAGTTGATG	68
FANCD2	CGGTTACAGGCTTCCCAAGT GCTGTTCTTGAATGCAGCCT	67
FANCI	GAGTCAGGCCGAGAAGGTTC CAACGGCAGCAGGTTTCTC	68
RAD51D	TGGCTCAGTTCTCGGCTTT AGATGTCAAATGCATGCACCA	66
TP53BP1	GCTGTTGCTGAGTCTGTTGC TGCGTACTTCCCGGATTGTT	68
RIF1	AGGTGGGCAAAACTGGTCAT TGGGTGCTCCACTACGAAAT	67
VEGF165b	AAGAAAATCCCTGTGGGCCTT	60

	TGGTGAGAGATCTGCAAGTACGTT	
VEGF165	AGAGCAAGACAAGAAAATCC	53
	TACAAACAAATGCTTTCTCC	
GAPDH	GGCTGCTTTTAACTCTGG	62
	GGAGGGATCTCGCTCC	

Immunofluorescence staining

KELLY cells were fixed using 4% paraformaldehyde (PFA) for 15 minutes, followed by permeabilization with 0.1% Triton-X100 for 20 minutes. The cells were then blocked with 5% BSA in PBS-T for 30mins at room temperature. Subsequently, they were incubated overnight at 4°C with anti-phosphoepitope SR proteins antibody (1:100, #MABE50, Sigma-Aldrich) or γH2AX (1:500, #05–636, Sigma-Aldrich) in blocking buffer. Following this, cells were washed before staining with a mix of Alexa Fluor 488-conjugated goat anti-mouse antibody (1:500, AF488, Thermo Scientific), DAPI (1μg/ml, #40043, Biotium,), and Phalloidin-647 (1:50, #8940S, Cell Signaling Technology) for 1 hour at room temperature. The cells were washed again three times with PBS-T before imaging with an INCell Analyzer 1000 microscope (GE Healthcare) at 20x magnification.

S9.6 (1:250, MABE1095, Sigma-Aldrich) and phospho RPA32-S33 (1:250, #NB100-544, Novus) were detected in KELLY cells fixed with cold methanol at -20°C for 10 minutes and slides were imaged using the Aurox laser-free confocal microscope. To confirm the specificity of the S9.6 antibody, RNase H (Ribonuclease H, 1:100, #M0297S, NEB) was applied to the fixed cells before antibody incubation.

Image analysis was performed using Cell Profiler V4.2.6, with nuclear regions masked by DAPI staining and cytoplasmic regions masked by Phalloidin staining, excluding areas overlapping with DAPI.

SRPK1/2 kinase activity assay

The SRPK1 Kinase Enzyme System (#VA7558, Promega) and SRPK2 Kinase Enzyme System (#VA7561, Promega) were utilised with the ADP-Glo™ Kinase Assay (#V6930, Promega), following the manufacturer's protocol. Briefly, 1 µl of inhibitor, diluted in 1X kinase reaction buffer, was added to the wells of a 384-well low-volume plate and combined with 8 ng of SRPK1 enzyme. Following this, 2 µl of a substrate/ATP mixture (100 µM ATP, 0.1 µg/µl MBP) was added. The reaction was incubated at room temperature for 1 hour. After incubation, 5 µl of ADP-Glo™ Reagent was added to each well, and the plate was incubated at room temperature for 40 minutes. Subsequently, 10 µl of Kinase Detection Reagent was added, and the plate was incubated for an additional 30 minutes. Luminescence was recorded.

Proximity Ligation Assay (PLA)

The Duolink® Proximity Ligation Assay (PLA) was used to detect protein-protein interactions in KELLY cells. Samples were fixed with 4% paraformaldehyde, permeabilised with 0.1% Triton X-100, and blocked with the Duolink® blocking solution. Primary antibodies against RBM39 (1:500, 67420, Proteintech) and SRPK1 (1:250, ab90527, Abcam) were applied overnight at 4°C. Species-specific PLA probes (Sigma-Aldrich) were added, followed by ligation and rolling-circle amplification according to the manufacturer's instructions.

DNA fibre analysis

Cells were sequentially pulse-labeled with 10 μM BrdU and 200 mM IdU for 30 minutes each.

KELLY cells were spotted on SuperFrost slides, air-dried for 5 minutes, and lysed with a buffer

(200 mM Tris-HCl, pH 7.5, 50 mM EDTA, 0.5% SDS) to release DNA. After air-drying for 5

minutes, the slides were tilted to allow DNA to spread, then dried, fixed in 75% methanol/25%

acetic acid, and treated with 2.5M HCl for 1 hour to denature the DNA. After PBS washes, slides

were blocked with 1% BSA in PBS-T for 1 hour. BrdU was detected by incubating slides with rat

anti-BrdU antibody (1:50, ab6326, Abcam) followed by AF647-conjugated secondary antibody

(1:100, A21247, Invitrogen). IdU was labeled with mouse anti-BrdU antibody (1:50, #BD347580)

and AF488-conjugated secondary antibody (1:100, #A11001, Invitrogen). Slides were mounted

and imaged using a laser-free confocal microscope, with DNA track lengths measured in Fiji V2.0.

In vivo allograft model of neuroblastoma

Syngeneic tumours derived from TH-MYCN/ALK^{F1174L} genetically engineered mouse models

(GEMMs), were implanted into immune competent 129×1/SvJ mice. Treatment began when the

implanted tumours reached a volume of over 200 mm³. The dosing schedule continued for 2 weeks,

with daily administration for 5 days followed by 2 days off. Alectinib was dissolved in saline

containing 40% PEG300, 10% DMSO, and 5% Tween, and was administered orally at a dose of

10 μL/kg. Indisulam was dissolved in saline with 3.5% DMSO and 6.5% Tween80 and was

administered intravenously at a dose of 5 μL/kg. Tumour tissues were excised two hours after the

final dose of each compound. Half of the tumour was fixed in 4% PFA, while the other half was

snap-frozen for subsequent analysis.

Immunoassays on tissue samples

Snap frozen tumours were lysed in 5% CHAPS buffer and western blotting was carried out for RBM39 (#67420, Proteintech) ERK1/2 (#4695 Cell Signaling Technology), pERK1/2 (#9101, Cell Signaling Technology), phosphoepitope SR proteins (#MABE50, Sigma-Aldrich) and GAPDH (#2118, Cell Signaling). ALK Meso Scale Discovery® immunoassays were performed as previously described(*49*)

Immunofluorescence on tissue sections

Tumours were processed using an ASP300S tissue processor (Leica) following the manufacturer's guidelines. Sections were deparaffinized and rehydrated using Xylene and a graded series of alcohol, followed by a 5-minute rinse in water. They were then boiled in 1% citric-acid buffer for 20 minutes and allowed to cool to room temperature. The sections were blocked for 1 hour in TBS with 0.01% Triton (TBST) and 5% BSA. Afterwards, γH2AX antibody (#9718, Cell Signaling Technology) was applied at a 1:100 dilution in blocking buffer and incubated overnight at 4°C. After 3 TBST washes, a secondary Alexa Fluor 568 goat anti-rabbit antibody (1:1000) was incubated at room temperature for 1 hour. Images were captured using a confocal microscope (LSM700, LSM T-PMT) and processed with ZEN2009 software (Black version, Zeiss).

Statistical Analysis

Statistical analyses were conducted using GraphPad Prism 10. For normally distributed data, a Student's t-test was used for two-group comparisons, while one-way ANOVA with multiple comparisons was applied to three or more groups. Two-way ANOVA with multiple comparisons

603   was used for multi-group analyses involving two factors. Specific statistical details are provided

604   in the figure legends.

List of Supplementary Materials

607   Fig S1 to S7 for multiple supplementary figures

608   Table S1 for supplementary table

References and Notes

1.   A. A. Patel, J. A. Steitz, Splicing double: insights from the second spliceosome. *Nat Rev Mol Cell Biol* **4**,

960-970 (2003).

2.   T. Han, M. Goralski, N. Gaskill, E. Capota, J. Kim, T. C. Ting, Y. Xie, N. S. Williams, D. Nijhawan,

Anticancer sulfonamides target splicing by inducing RBM39 degradation via recruitment to DCAF15.

Science **356**, (2017).

3.   C. Xu, X. Chen, X. Zhang, D. Zhao, Z. Dou, X. Xie, H. Li, H. Yang, Q. Li, H. Zhang, C. Di, RNA-binding

protein 39: a promising therapeutic target for cancer. *Cell Death Discov* **7**, 214 (2021).

4.   T. Uehara, Y. Minoshima, K. Sagane, N. H. Sugi, K. O. Mitsuhashi, N. Yamamoto, H. Kamiyama, K.

Takahashi, Y. Kotake, M. Uesugi, A. Yokoi, A. Inoue, T. Yoshida, M. Mabuchi, A. Tanaka, T. Owa,

Selective degradation of splicing factor CAPERalpha by anticancer sulfonamides. *Nat Chem Biol* **13**, 675-

680 (2017).

5.   J. Bojko, M. Kollareddy, M. Szemes, J. Bellamy, E. Poon, A. Moukachar, D. Legge, E. E. Vincent, N.

Jones, S. Malik, A. Greenhough, A. Paterson, J. H. Park, K. Gallacher, L. Chesler, K. Malik, Spliceosomal

vulnerability of MYCN-amplified neuroblastoma is contingent on PRMT5-mediated regulation of

epitranscriptomic and metabolomic pathways. *Cancer Lett* **604**, 217263 (2024).

6.   A. Nijhuis, A. Sikka, O. Yogev, L. Herendi, C. Balcells, Y. Ma, E. Poon, C. Eckold, G. N. Valbuena, Y.

Xu, Y. Liu, B. M. da Costa, M. Gruet, C. Wickremesinghe, A. Benito, H. Kramer, A. Montoya, D. Carling,

E. J. Want, Y. Jamin, L. Chesler, H. C. Keun, Indisulam targets RNA splicing and metabolism to serve as a

therapeutic strategy for high-risk neuroblastoma. *Nat Commun* **13**, 1380 (2022).

7.   A. Salib, N. Jayatilleke, J. A. Seneviratne, C. Mayoh, K. De Preter, F. Speleman, B. B. Cheung, D. R.

Carter, G. M. Marshall, MYCN and SNRPD3 cooperate to maintain a balance of alternative splicing events

that drives neuroblastoma progression. *Oncogene* **43**, 363-377 (2024).

8.   S. Singh, W. Quarni, M. Goralski, S. Wan, H. Jin, L. A. Van de Velde, J. Fang, Q. Wu, A. Abu-Zaid, T.

Wang, R. Singh, D. Craft, Y. Fan, T. Confer, M. Johnson, W. J. Akers, R. Wang, P. J. Murray, P. G.

Thomas, D. Nijhawan, A. M. Davidoff, J. Yang, Targeting the spliceosome through RBM39 degradation

results in exceptional responses in high-risk neuroblastoma models. *Sci Adv* **7**, eabj5405 (2021).

9.   K. K. Matthay, J. M. Maris, G. Schleiermacher, A. Nakagawara, C. L. Mackall, L. Diller, W. A. Weiss,

Neuroblastoma. *Nat Rev Dis Primers* **2**, 16078 (2016).

10.   N. Bechmann, F. Westermann, G. Eisenhofer, HIF and MYC signaling in adrenal neoplasms of the neural

crest: implications for pediatrics. *Front Endocrinol (Lausanne)* **14**, 1022192 (2023).

11.   J. Otte, C. Dyberg, A. Pepich, J. I. Johnsen, MYCN Function in Neuroblastoma Development. *Front Oncol*

10, 624079 (2020).

12.   Y. Chen, J. Takita, Y. L. Choi, M. Kato, M. Ohira, M. Sanada, L. Wang, M. Soda, A. Kikuchi, T. Igarashi,

A. Nakagawara, Y. Hayashi, H. Mano, S. Ogawa, Oncogenic mutations of ALK kinase in neuroblastoma.

Nature **455**, 971-974 (2008).

13.   A. T. Shaw, L. Gandhi, S. Gadgeel, G. J. Riely, J. Cetnar, H. West, D. R. Camidge, M. A. Socinski, A.

Chiappori, T. Mekhail, B. H. Chao, H. Borghaei, K. A. Gold, A. Zeaiter, W. Bordogna, B. Balas, O. Puig,

V. Henschel, S. I. Ou, i. study, Alectinib in ALK-positive, crizotinib-resistant, non-small-cell lung cancer: a

single-group, multicentre, phase 2 trial. *Lancet Oncol* **17**, 234-242 (2016).

14.   M. W. Alam, M. Borenas, D. E. Lind, D. Cervantes-Madrid, G. Umapathy, R. H. Palmer, B. Hallberg,

Alectinib, an Anaplastic Lymphoma Kinase Inhibitor, Abolishes ALK Activity and Growth in ALK-

Positive Neuroblastoma Cells. *Front Oncol* **9**, 579 (2019).

15.   J. Lu, S. Guan, Y. Zhao, Y. Yu, S. E. Woodfield, H. Zhang, K. L. Yang, S. Bieerkehazhi, L. Qi, X. Li, J.

Gu, X. Xu, J. Jin, J. A. Muscal, T. Yang, G. T. Xu, J. Yang, The second-generation ALK inhibitor alectinib

effectively induces apoptosis in human neuroblastoma cells and inhibits tumor growth in a TH-MYCN

transgenic neuroblastoma mouse model. *Cancer Lett* **400**, 61-68 (2017).

16.   A. Zafar, W. Wang, G. Liu, X. Wang, W. Xian, F. McKeon, J. Foster, J. Zhou, R. Zhang, Molecular

657   targeting therapies for neuroblastoma: Progress and challenges. *Med Res Rev* **41**, 961-1021 (2021).

658   17.   P. J. Houghton, R. T. Kurmasheva, Challenges and Opportunities for Childhood Cancer Drug

659   Development. *Pharmacol Rev* **71**, 671-697 (2019).

660   18.   O. Anczukow, A. R. Krainer, The spliceosome, a potential Achilles heel of MYC-driven tumors. *Genome*

661   *Med* **7**, 107 (2015).

662   19.   Y. Xu, S. Spear, Y. Ma, M. P. Lorentzen, M. Gruet, F. McKinney, Y. Xu, C. Wickremesinghe, M. R.

663   Shepherd, I. McNeish, H. C. Keun, A. Nijhuis, Pharmacological depletion of RNA splicing factor RBM39

664   by indisulam synergizes with PARP inhibitors in high-grade serous ovarian carcinoma. *bioRxiv*,

665   2023.2001.2018.524417 (2023).

666   20.   J. M. Hatcher, G. Wu, C. Zeng, J. Zhu, F. Meng, S. Patel, W. Wang, S. B. Ficarro, A. L. Leggett, C. E.

667   Powell, J. A. Marto, K. Zhang, J. C. Ki Ngo, X. D. Fu, T. Zhang, N. S. Gray, SRPKIN-1: A Covalent

668   SRPK1/2 Inhibitor that Potently Converts VEGF from Pro-angiogenic to Anti-angiogenic Isoform. *Cell*

669   *Chem Biol* **25**, 460-470 e466 (2018).

670   21.   C. I. BLISS, THE TOXICITY OF POISONS APPLIED JOINTLY1. *Annals of Applied Biology* **26**, 585-

671   615 (1939).

672   22.   M. Uhlen, E. Bjorling, C. Agaton, C. A. Szigyarto, B. Amini, E. Andersen, A. C. Andersson, P. Angelidou,

673   A. Asplund, C. Asplund, L. Berglund, K. Bergstrom, H. Brumer, D. Cerjan, M. Ekstrom, A. Elobeid, C.

674   Eriksson, L. Fagerberg, R. Falk, J. Fall, M. Forsberg, M. G. Bjorklund, K. Gumbel, A. Halimi, I. Hallin, C.

675   Hamsten, M. Hansson, M. Hedhammar, G. Hercules, C. Kampf, K. Larsson, M. Lindskog, W. Lodewyckx,

676   J. Lund, J. Lundeberg, K. Magnusson, E. Malm, P. Nilsson, J. Odling, P. Oksvold, I. Olsson, E. Oster, J.

677   Ottosson, L. Paavilainen, A. Persson, R. Rimini, J. Rockberg, M. Runeson, A. Sivertsson, A. Skollermo, J.

678   Steen, M. Stenvall, F. Sterky, S. Stromberg, M. Sundberg, H. Tegel, S. Tourle, E. Wahlund, A. Walden, J.

679   Wan, H. Wernerus, J. Westberg, K. Wester, U. Wrethagen, L. L. Xu, S. Hober, F. Ponten, A human protein

680   atlas for normal and cancer tissues based on antibody proteomics. *Mol Cell Proteomics* **4**, 1920-1932

681   (2005).

23.   N. R. Infarinato, J. H. Park, K. Krytska, H. T. Ryles, R. Sano, K. M. Szigety, Y. Li, H. Y. Zou, N. V. Lee,

T. Smeal, M. A. Lemmon, Y. P. Mosse, The ALK/ROS1 Inhibitor PF-06463922 Overcomes Primary

Resistance to Crizotinib in ALK-Driven Neuroblastoma. *Cancer Discov* **6**, 96-107 (2016).

24.   H. Sakamoto, T. Tsukaguchi, S. Hiroshima, T. Kodama, T. Kobayashi, T. A. Fukami, N. Oikawa, T.

Tsukuda, N. Ishii, Y. Aoki, CH5424802, a selective ALK inhibitor capable of blocking the resistant

gatekeeper mutant. *Cancer Cell* **19**, 679-690 (2011).

25.   M. Tufail, C. Wu, SRPKs: a promising therapeutic target in cancer. *Clin Exp Med*, (2023).

26.   I. P. Nikas, S. C. Themistocleous, S. A. Paschou, K. I. Tsamis, H. S. Ryu, Serine-Arginine Protein Kinase 1

(SRPK1) as a Prognostic Factor and Potential Therapeutic Target in Cancer: Current Evidence and Future

Perspectives. *Cells* **9**, (2019).

27.   Z. Pogacar, K. Groot, F. Jochems, M. Dos Santos Dias, A. Mulero-Sanchez, B. Morris, M. Roosen, L.

Wardak, G. De Conti, A. Velds, C. Lieftink, B. Thijssen, R. L. Beijersbergen, R. Bernards, R. Leite de

Oliveira, Genetic and compound screens uncover factors modulating cancer cell response to indisulam. *Life*

Sci Alliance **5**, (2022).

28.   J. Batson, H. D. Toop, C. Redondo, R. Babaei-Jadidi, A. Chaikuad, S. F. Wearmouth, B. Gibbons, C.

Allen, C. Tallant, J. Zhang, C. Du, J. C. Hancox, T. Hawtrey, J. Da Rocha, R. Griffith, S. Knapp, D. O.

Bates, J. C. Morris, Development of Potent, Selective SRPK1 Inhibitors as Potential Topical Therapeutics

for Neovascular Eye Disease. *ACS Chem Biol* **12**, 825-832 (2017).

29.   E. M. Amin, S. Oltean, J. Hua, M. V. R. Gammons, M. Hamdollah-Zadeh, G. I. Welsh, M. K. Cheung, L.

Ni, S. Kase, E. S. Renne, K. E. Symonds, D. G. Nowak, B. Royer-Pokora, M. A. Saleem, M. Hagiwara, V.

A. Schumacher, S. J. Harper, D. R. Hinton, D. O. Bates, M. R. Ladomery, WT1 Mutants Reveal SRPK1 to

Be a Downstream Angiogenesis Target by Altering VEGF Splicing. *Cancer Cell* **20**, 768-780 (2011).

30.   S. Burma, B. P. Chen, M. Murphy, A. Kurimasa, D. J. Chen, ATM phosphorylates histone H2AX in

response to DNA double-strand breaks. *J Biol Chem* **276**, 42462-42467 (2001).

31.   J. M. Santos-Pereira, A. Aguilera, R loops: new modulators of genome dynamics and function. *Nat Rev*

Genet **16**, 583-597 (2015).

32.   S. Hamperl, K. A. Cimprich, Conflict Resolution in the Genome: How Transcription and Replication Make

It Work. *Cell* **167**, 1455-1467 (2016).

710 33. S. C. Sridhara, S. Carvalho, A. R. Grosso, L. M. Gallego-Paez, M. Carmo-Fonseca, S. F. de Almeida,

711   Transcription Dynamics Prevent RNA-Mediated Genomic Instability through SRPK2-Dependent DDX23

712   Phosphorylation. *Cell Rep* **18**, 334-343 (2017).

713 34. E. Hatchi, K. Skourti-Stathaki, S. Ventz, L. Pinello, A. Yen, K. Kamieniarz-Gdula, S. Dimitrov, S.

714   Pathania, K. M. McKinney, M. L. Eaton, M. Kellis, S. J. Hill, G. Parmigiani, N. J. Proudfoot, D. M.

715   Livingston, BRCA1 recruitment to transcriptional pause sites is required for R-loop-driven DNA damage

716   repair. *Mol Cell* **57**, 636-647 (2015).

717 35. Y. Okamoto, M. Abe, A. Itaya, J. Tomida, M. Ishiai, A. Takaori-Kondo, M. Taoka, T. Isobe, M. Takata,

718   FANCD2 protects genome stability by recruiting RNA processing enzymes to resolve R-loops during mild

719   replication stress. *FEBS J* **286**, 139-150 (2019).

720 36. C. Mukherjee, V. Tripathi, E. M. Manolika, A. M. Heijink, G. Ricci, S. Merzouk, H. R. de Boer, J.

721   Demmers, M. van Vugt, A. Ray Chaudhuri, RIF1 promotes replication fork protection and efficient restart

722   to maintain genome stability. *Nat Commun* **10**, 3287 (2019).

723 37. D. D. Phillips, D. N. Garboczi, K. Singh, Z. Hu, S. H. Leppla, C. E. Leysath, The sub-nanomolar binding of

724   DNA-RNA hybrids by the single-chain Fv fragment of antibody S9.6. *J Mol Recognit* **26**, 376-381 (2013).

725 38. A. Promonet, I. Padioleau, Y. Liu, L. Sanz, A. Biernacka, A. L. Schmitz, M. Skrzypczak, A. Sarrazin, C.

726   Mettling, M. Rowicka, K. Ginalski, F. Chedin, C. L. Chen, Y. L. Lin, P. Pasero, Topoisomerase 1 prevents

727   replication stress at R-loop-enriched transcription termination sites. *Nat Commun* **11**, 3940 (2020).

728 39. K. C. Goldsmith, J. R. Park, K. Kayser, J. Malvar, Y. Y. Chi, S. G. Groshen, J. G. Villablanca, K. Krytska,

729   L. M. Lai, P. T. Acharya, F. Goodarzian, B. Pawel, H. Shimada, S. Ghazarian, L. States, L. Marshall, L.

730   Chesler, M. Granger, A. V. Desai, R. Mody, D. A. Morgenstern, S. Shusterman, M. E. Macy, N. Pinto, G.

731   Schleiermacher, K. Vo, H. C. Thurm, J. Chen, M. Liyanage, G. Peltz, K. K. Matthay, E. R. Berko, J. M.

732   Maris, A. Marachelian, Y. P. Mosse, Lorlatinib with or without chemotherapy in ALK-driven

733   refractory/relapsed neuroblastoma: phase 1 trial results. *Nat Med* **29**, 1092-1102 (2023).

734 40. Y. Suk, S. K. Singh, Safety and efficacy of lorlatinib against ALK-driven refractory or relapsed

735   neuroblastoma. *Cell Rep Med* **4**, 101071 (2023).

736 41. S. Peters, D. R. Camidge, A. T. Shaw, S. Gadgeel, J. S. Ahn, D. W. Kim, S. I. Ou, M. Perol, R.

737   Dziadziuszko, R. Rosell, A. Zeaiter, E. Mitry, S. Golding, B. Balas, J. Noe, P. N. Morcos, T. Mok, A. T.

Investigators, Alectinib versus Crizotinib in Untreated ALK-Positive Non-Small-Cell Lung Cancer. *N Engl*

J Med **377**, 829-838 (2017).

42.   B. J. Solomon, G. Liu, E. Felip, T. S. K. Mok, R. A. Soo, J. Mazieres, A. T. Shaw, F. de Marinis, Y. Goto,

Y. L. Wu, D. W. Kim, J. F. Martini, R. Messina, J. Paolini, A. Polli, D. Thomaidou, F. Toffalorio, T. M.

Bauer, Lorlatinib Versus Crizotinib in Patients With Advanced ALK-Positive Non-Small Cell Lung

Cancer: 5-Year Outcomes From the Phase III CROWN Study. *J Clin Oncol* **42**, 3400-3409 (2024).

43.   S. Singh, W. Quarni, M. Goralski, S. B. Wan, H. J. Jin, L. A. Van de Velde, J. Fang, Q. Wu, A. Abu-Zaid,

T. T. Wang, R. Singh, D. Craft, Y. P. Fan, T. Confer, M. Johnson, W. J. Akers, R. N. Wang, P. J. Murray,

P. G. Thomas, D. Nijhawan, A. M. Davidoff, J. Yang, Targeting the spliceosome through RBM39

degradation results in exceptional responses in high-risk neuroblastoma models. *Science Advances* **7**,

(2021).

44.   S. Kohsaka, S. Yagishita, Y. Shirai, Y. Matsuno, T. Ueno, S. Kojima, H. Ikeuchi, M. Ikegami, R. Kitada,

K. I. Yoshioka, K. Toshimitsu, K. Tabata, A. Yokoi, T. Doi, N. Yamamoto, T. Owa, A. Hamada, H. Mano,

A molecular glue RBM39-degrader induces synthetic lethality in cancer cells with homologous

recombination repair deficiency. *NPJ Precis Oncol* **8**, 117 (2024).

45.   E. R. Tucker, I. Jimenez, L. Chen, A. Bellini, C. Gorrini, E. Calton, Q. Gao, H. Che, E. Poon, Y. Jamin, B.

Martins Da Costa, K. Barker, S. Shrestha, J. C. Hutchinson, S. Dhariwal, A. Goodman, E. Del Nery, P.

Gestraud, J. Bhalshankar, Y. Iddir, E. Saberi-Ansari, A. Saint-Charles, B. Geoerger, M. E. Marques Da

Costa, C. Pierre-Eugene, I. Janoueix-Lerosey, D. Decaudin, F. Nemati, A. M. Carcaboso, D. Surdez, O.

Delattre, S. L. George, L. Chesler, D. A. Tweddle, G. Schleiermacher, Combination Therapies Targeting

ALK-aberrant Neuroblastoma in Preclinical Models. *Clin Cancer Res* **29**, 1317-1331 (2023).

46.   B. Y. Tam, K. Chiu, H. Chung, C. Bossard, J. D. Nguyen, E. Creger, B. W. Eastman, C. C. Mak, M.

Ibanez, A. Ghias, J. Cahiwat, L. Do, S. Cho, J. Nguyen, V. Deshmukh, J. Stewart, C. W. Chen, C. Barroga,

L. Dellamary, S. K. Kc, T. J. Phalen, J. Hood, S. Cha, Y. Yazici, The CLK inhibitor SM08502 induces anti-

tumor activity and reduces Wnt pathway gene expression in gastrointestinal cancer models. *Cancer Lett*

473, 186-197 (2020).

47.   M. Sakuma, K. Iida, M. Hagiwara, Deciphering targeting rules of splicing modulator compounds: case of

TG003. *Bmc Mol Biol* **16**, (2015).

48.   M. Buljan, R. Ciuffa, A. van Drogen, A. Vichalkovski, M. Mehnert, G. Rosenberger, S. Lee, M. Varjosalo,

767   L. E. Pernas, V. Spegg, B. Snijder, R. Aebersold, M. Gstaiger, Kinase Interaction Network Expands

768   Functional and Disease Roles of Human Kinases. *Mol Cell* **79**, 504-520 e509 (2020).

769   49.   E. R. Tucker, J. R. Tall, L. S. Danielson, S. Gowan, Y. Jamin, S. P. Robinson, U. Banerji, L. Chesler,

770   Immunoassays for the quantification of ALK and phosphorylated ALK support the evaluation of on-target

771   ALK inhibitors in neuroblastoma. *Mol Oncol* **11**, 996-1006 (2017).

Acknowledgments: We are very grateful to the technicians from the Institute of Cancer

Research BSU animal center, Barbara Martins da Costa and Kevin Greenslade for supporting

animal work. We thank Prof. Matthew J Fuchter from Imperial College London and Dr Lizzie

Tucker from Institute of Cancer Research for providing helpful comments.

Funding: Imperial College London and China Scholarship Council joint PhD studentship

202208310101 (YM), 201808060050 (YX). AstraZeneca/NIHR Imperial BRC Imperial College

Research Fellowship (AN). Cancer Research UK Fellowship RCCCDF-Nov23/100003 (AN).

Cancer Research UK Discovery Programme Grant (A28278) (EP, BMC, LC); HEFCE/RAE

(LC). Ovarian Cancer Action (CW, AN). Cancer Research UK convergence Science Centre PhD

studentships SEBCATP-2024/100009 (NZ), CANCTA-2022\100007 (SQ).

Author contributions:

Conceptualization: YM, AN, HCK

Methodology: YM, EP, YX, SQ, AN

Investigation: YM, EP, BMC, YX, CJ, SQ, NZ, LC, HCK, AN

Visualization: YM, YX, CJ, SQ, AN

Funding acquisition: LC, HCK, AN

Project administration: HCK, AN

Supervision: AN, HCK, LC,

Writing – original draft: YX, LC, HCK, AN

Writing – review & editing: all authors contributed to the review and editing of the paper.

Competing interests: Authors declare that they have no competing interests.

Data and materials availability: All data supporting the conclusions of this paper are available

within the main text and/or Supplementary Materials.

801