Genomics and reproductive biology of *Leptopilina malgretoutensis* Buffington, Lue, Davis &

Tracey sp. nov. (Hymenoptera: Figitidae): An asexual parasitoid of Caribbean *Drosophila*

Amelia RI Lindsey*[1], Chia-Hua Lue[2], Jeremy S. Davis[3], Lydia J. Borjon[4,5], Stephanie E. Mauthner[5], Laura

C Fricke[1], Lauren Eads[4], Molly Murphy[4], Melissa K Drown[6,7], Christopher Faulk[6], Matthew L Buffington*[8],

W Daniel Tracey Jr*[4,5]

[1]Department of Entomology, University of Minnesota, Saint Paul, Minnesota, USA

[2]Department of Biology, Hood College, Frederick, Maryland, USA

[3]Department of Molecular Biosciences, Northwestern University, Evanston, Illinois, USA

[4]Gill Institute for Neuroscience, Indiana University, Bloomington, Indiana, USA

[5]Department of Biology, Indiana University, Bloomington, Indiana, USA

[6]Department of Animal Science, University of Minnesota, Saint Paul, Minnesota, USA

[7]Current Affiliation: Oxford Nanopore Technologies, New York, New York, USA

[8]Systematic Entomology Laboratory, ARS/USDA c/o Smithsonian Institution, National Museum of Natural

History, Washington, DC, USA

*To whom correspondence should be addressed (ARIL: alindsey@umn.edu; MLB:

matt.buffington@usda.gov; WDT: dtracey@iu.edu)

ORCIDs:
ARIL 0000-0003-3354-7419
CHL 0000-0002-5245-603X
JSD 0000-0002-5214-161X
LJB 0000-0002-3680-4145
SEM 0000-0002-4159-6571
CF 0000-0002-9749-9658
MLB 0000-0003-1900-3861
WDT 0000-0003-4666-8199

ABSTRACT

Drosophila and parasitic wasps in the genus *Leptopilina* have long been a model for understanding host-

parasite interactions. Indeed, parasitic wasps are important drivers of ecological and evolutionary

processes broadly, but we are generally lacking information about the diversity, natural history, and

evolution of these relationships. We collected insects from the Caribbean Island of Saint Lucia, home to

the eastern Caribbean '*dunni*' subgroup of *Drosophila*: a clade long appreciated for its recent patterns of

speciation and adaptation. Here we present an integrative approach that incorporates natural history,

taxonomy, physiology, and genomics to describe *Leptopilina malgretoutensis* Buffington, Lue, Davis &

Tracey sp. nov. (Hymenoptera: Figitidae), a virulent parasitoid of *dunni* group flies, especially *Drosophila*

antillea. *Leptopilina malgretoutensis* is nested within an early-branching clade of *Leptopilina*, offering

insights into the evolution of this important genus of *Drosophila* parasitoids. We present a high-quality

assembly for this wasp's 1Gbp genome, and for its bacterial endosymbiont: *Wolbachia* strain "*w*Lmal".

Furthermore, we show that *w*Lmal induces parthenogenesis in the wasp, and that these wasps are reliant

upon their *Wolbachia* infections to produce female offspring. Finally, comparison to historical museum

specimens indicate that *Leptopilina malgretoutensis* had been collected approximately 40 years prior from

the nearby island of Guadeloupe and were also asexually reproducing. This work represents one of only

a handful of studies in which field biology, taxonomy, systematics, genomics, and experimental biology

are integrated into a species description: showcasing the possibilities for biodiversity research in the

genomic era.

KEYWORDS

dunni, *cardini*, *Wolbachia*, parthenogenesis, Saint Lucia, Antilles, host-parasite

INTRODUCTION

Interactions between insects and their parasitoids are ubiquitous across terrestrial ecosystems, and have been postulated as critical, but underappreciated, drivers of ecological speciation in insect groups [1, 2]. Yet compared to other ecological interactions, considerably less attention is paid to parasitism as a driver of insect speciation, in part because of the challenges of identifying often cryptic parasitoid species. Molecular-based identification methods have rapidly revolutionized species identification, but issues persist for many parasitoid lineages [3]. Most critically, while parasitoids are commonly acknowledged as one of the most speciose groups of organisms, they are persistently under-described and under-represented in insect conservation efforts [4-6]. As such, more effort to find and describe the ecology and unique biology of new parasitoid species is needed to better understand the pattern and scope of biodiversity on earth.

Island species groups have been attractive systems for the study of speciation events and adaptive radiations since as far back as Darwin (1859), who recognized the immense number of endemic species on archipelago chains. Island archipelagos offer unique opportunities to study peripatric speciation following discrete patterns of colonization between single islands in an archipelago chain [7]. *Drosophila* species groups have been of particular interest for studying patterns of island colonization and speciation. Indeed, the Hawaiian *Drosophila* group represents one of the most well studied examples of island radiations – home to at least 500 species and research spanning decades [8-10]. While Hawaiian *Drosophila* are the most intensely studied group, several other island clades of *Drosophila* have received research attention (e.g., *sechellia-simulans* [11, 12]). Another such clade is the *D. dunni* group – a subgroup of the larger *D. cardini* group that is endemic to the Caribbean islands, particularly the Lesser Antilles [13, 14]. This subgroup is characterized by species or subspecies with distinct abdominal pigmentation endemic to just one or two major islands in this archipelago – frequently just 25-50 km apart. As such, this group represents an excellent model for the study of peripatric speciation and morphological evolution across archipelagos [12, 14]. Despite years of appreciation for the *dunni* group as a model, little is known about the parasitoid complex of these Caribbean flies.

To better understand the ecology of *dunni* group *Drosophila,* and to investigate their potential parasitoids,

we (JSD, SEM, WDT) conducted an expedition to the islands of the Lesser Antilles February through May

of 2019. We traveled among the islands aboard the sailing vessel Sea Salt (a 42-foot Fountaine Pajot

Catamaran) which also served as a mobile laboratory. We describe here results with collected flies and

associated parasitoids from the island of Saint Lucia. Other results of our expedition will be described

elsewhere. We describe a new species of *Drosophila* parasitoid: *Leptopilina malgretoutensis* Buffington,

Lue, Davis & Tracey sp. nov. (Hymenoptera: Figitidae). *Leptopilina malgretoutensis* appears to be

endemic to the Lesser Antilles and we find that it has high virulence against *dunni* group flies, particularly

Drosophila antillea also collected on Saint Lucia. To our knowledge, this is the first host-parasitoid

association recorded for *cardini* and *dunni* group *Drosophila.* We find that the wasps reproduce via

thelytokous parthenogenesis (the asexual reproduction of females) and that this is caused by infection

with a bacterium: a parthenogenesis-inducing *Wolbachia* symbiont. We report *de novo* genome

assemblies for *Leptopilina malgretoutensis* and its *Wolbachia* strain, hereinafter "*w*Lmal". *Leptopilina*

malgretoutensis has the largest *Leptopilina* genome reported to-date (~1Gbp) and will serve as a useful

resource for understanding the evolution of *Drosophila-Leptopilina* relationships in the Caribbean islands,

island speciation events, and parasitoid genome evolution more broadly.

METHODS

Collection: Field sites and field rearing

After sailing from Wallliabou Bay of Saint Vincent and the Grenadines, we moored the research vessel

and operated out of Soufrière and Marigot Bays of Saint Lucia. Verbal permission for collection of

Drosophila and their parasitoids was obtained in a February 2019 meeting with Dr. Hannah Romain of the

Saint Lucia Ministry of Agriculture. Sampling was conducted in 2019 on Saint Lucia and leveraged baited

trapping with either banana or calabasa, a locally available cucurbit fruit (likely *Cucurbita moschata*).

Calabasa and banana traps contained chopped fruit mixed with water and baker's yeast that had been

kept in a closed bucket for one or more days to ferment. The Calabasa bait was chosen based on

communication with Hope Hollecher who found it to be particularly attractive to *dunni* group *Drosophila.*

Traps were set at two separate sites from which flies and wasps in this study derive. The first site, known

as STL3 was placed on 24 February 2019 with a total of 2 calabasa traps on a short trail leading off the

main road near Ladera resort and Hotel Chocolat (GPS: lat 13.83, lon -61.05, elevation 289.6m, Figure

1). The second site, known as STL6 was placed on 26 February 2019 with a total of eight traps (5

calabasa, 3 banana) along Barre d'Isle trail (GPS: lat 13.924, lon -60.959, elevation 308.7m, Figure 1).

Each trapping site was checked daily for three days and insects were collected off the fruit each time.

Flies were collected using either cardboard mailing tubes for vertical trapping or mouth aspiration, and

wasps were collected by mouth aspiration while they searched for larvae in the bait.

Figure 1. Collections on Saint Lucia: field notebook entry from JSD on 1 March 2019: "*...a 40-minute brisk hike, in which I found 0 wasps in the traps - [bummer]. Until I remembered I had set out one more trap off the path a little that I had not checked yet. And [nevertheless], my lucky [telson] finds the wasp we needed sitting in this cup. I managed to nab it and was pretty excited leaving here.*" **(A)** Map of Saint Lucia (GoogleMaps 2023), annotated with (1) locations of the STL3 and STL6 trap placements, (2) the route and mooring locations (Soufrière in the south, and Marigot in the north) of the Sea Salt, and (3) Malgretoute: the location of the La Soufrière Volcano. **(B)** The STL6 calabasa trap. **(C)** The inside of a calabasa trap.

Collection: Colony establishment and maintenance

After collection, flies and wasps were transferred to individual rearing vials with instant fly media (Formula

4-24® Instant Drosophila Medium, Blue; Carolina Biological Supply Company). Isofemale lines of

Drosophila antillea and *Drosophila willistoni*, were initiated using flies collected at STL3 (see results). A

single wasp was collected from STL6 and was left in a vial with developing STL3 *Drosophila willistoni*

larvae. This vial was then shipped back to Indiana University Bloomington under USDA APHIS

Importation Permit Number P526P-19-01355. Since 2019, wasps have been maintained continuously on

host cultures of *D. willistoni* on Bloomington fly food medium and further reared in IU USDA regulated

containment facility number 5035.

Taxonomy: Imaging

Voucher specimens were sent to two of us (CHL and MB) for identification. These specimens were

compared with specimens deposited in the USNM as well as those on loan from Goran Nordlander

(Uppsala Agricultural University, retired). Additionally, notes from Yves Carton were shared with MB, and

his work with G. Nordlander in the mid 1980's. These notes revealed an undescribed *Leptopilina* from

Guadeloupe that matched the species described herein. We are therefore including these older

specimens in the description. Images were acquired using a Hitachi 3000 desktop Scanning Electron

Microscope (SEM), after a 120nm coating of Au-Pd from a Cressington sputter-coater. Light micrographs

were shot using a Macropod 3D Pro camera setup. All type specimens collected in this study are housed

in the USNM (registration numbers below); additional types, examined by G. Nordlander, reside in the

Uppsala Agricultural University collection overseen by G. Nordander. Terminology for these descriptions

follows Lue et al. (2016) [15].

Taxonomy: Systematics of *Leptopilina*

Lue et al. (2016) [15] provided a detailed description on the species limits of *Leptopilina*. Species in this

genus are common throughout the Holarctic Region and have recently been investigated for their

biological control potential of pestiferous Drosophilidae [16, 17]. Some 29 species are currently

recognized worldwide [15]. While most described species are endemic to the Holarctic Region, many

undescribed species remain in collections, awaiting description. A relatively common trait among

Neotropical *Leptopilina* are polychromatic antennal segments, specifically black/brown, cream, and

orange-colored segments. These traits are either rare, or wholly missing, from the Paleotropics and

Australia (Buffington, Lue, pers. obsv.). This is mentioned here as the new species, herein described, has

rather unique color patterns among the antennal segments. This character makes this species relatively

straightforward to diagnose from other *Leptopilina*.

Biology: Virulence assessment

To test the virulence of *L. malgretoutensis* on *Drosophila* host species, infection vials were established at

a wasp-to-larvae ratio of 1:10 (typically 5 female wasps and 50 drosophila larvae). Two isofemale lines

from two *Drosophila* species that were collected from the island of Saint Lucia were assessed: *Drosophila*

willistoni (line STL3 iso3) and *Drosophila antillea* (line STL3 iso1). To collect the age-matched larvae, 20-

30 male and female adult flies were placed in embryo collection cages. The embryo collection plates (60

x 15mm) contained 5mL of *Drosophila* banana food medium (Archon Scientific) and a small amount of

yeast paste (active dry yeast mixed with water). The embryo collection cages were maintained in an

incubator set at 25°C and 75% humidity on a 12h/12h light/dark cycle. Embryos from a restricted 12-hour

egg-laying window were allowed to develop in the incubator for 3 days until they reached the size of 2^{nd}

instar larvae. Larvae were carefully gently collected using extreme caution so as to not damage them.

Fifty larvae were placed into individual vials containing *Drosophila* banana food and a small cotton dental

plug that was inserted into the center of the food. The cotton plug was necessary to provide a place for

pupariation for the *D. antillea* species (which avoids pupariation on the side of the culture vials). Five

naïve *L. malgretoutensis* female wasps were placed into each infection vial and were allowed to infect the

larvae for 24 hours. During the infection period, the vials were closed with Flugs (Genesee Scientific

cat.no.49-102) dampened with honey and water (1:1 mixture) to ensure that the wasps had access to

food. After the infection period, the wasps were removed, and the vials were incubated as described

above. Uninfected mock control vials were treated in the same way but were not exposed to wasps. The

vials were assessed daily for the emergence of adult insects. Adult flies eclosed after 10-14 days, and

adult wasps eclosed after 25-30 days. Any newly eclosed adult flies and wasps were counted and

removed from the vials. The difference between the total number of adults and the initial number of larvae

in each vial accounts for larval mortality.

Biology: Western blots for _Wolbachia_ detection

Active _Wolbachia_ infection was assessed with western blot following published protocols [18]. Positive

and negative controls included individual _Drosophila melanogaster_ from previously validated _Wolbachia_-

infected and -uninfected stocks (RRID: BDSC_145) [19]. Individual insects were homogenized with a

pestle in 1.7 ml microcentrifuge tubes containing lysis buffer (150mM NaCl, 1% Triton X-100, 50mM

TrisHCl (pH8) containing HALT protease inhibitor cocktail (Thermo Scientific) and 5 mM EDTA).

Homogenates were pelleted via centrifugation (8,000Xg for 1 minute) and the supernatant was incubated

for 5 minutes at 95°C in Laemmli sample buffer with 5% β-mercaptoethanol (Bio-Rad). Proteins were

separated via SDS-PAGE on 4–20% Tris-Glycine gels (NuSep) in 1X Tris/Glycine/SDS running buffer

(Bio-Rad) alongside a PageRuler prestained protein ladder (Thermo Scientific), transferred to a PVDF

membrane in Tris-Glycine transfer buffer with 15% methanol for 3 hours at 40v, and blocked for 5 minutes

in Starting Block T20 (Thermo Scientific). The membrane was co-incubated in primary antibodies

overnight at 4°C. After washing, the blot was incubated with the secondary antibody anti-mouse:HRP

(Invitrogen) at 1:5000. Finally, HRP was detected with SuperSignal West Pico Chemiluminescent

Substrate (Thermo Scientific). Primary antibodies included (1) mouse monoclonal Anti-Wolbachia Surface

Protein (WSP), NR-31029 from BEI Resources, NIAID, NIH at 1:10,000, and (2) mouse anti-actin

monoclonal at 1:20,000 (Seven Hills Bioreagents) as a loading control.

Biology: Determining reproductive mode

To test whether the observed parthenogenesis was induced by infection with a bacterial symbiont, _L._

malgretoutensis female wasps were fed with the antibiotic rifampicin mixed into honey (100 mg of

rifampicin powder (Sigma-Aldrich cat.no. R3501-250MG) with 1 mL of honey) and reproductive biology

was assessed following standard protocols [20]. To feed the wasps, a Flug was dipped into water to

dampen one side, onto which a small amount of the rifampicin-honey mixture was spread (approx. 10-20

μL). A fresh Flug with rifampicin-honey was replaced once a day for six days. To prepare vials of fly

larvae, adult *Drosophila* were allowed to seed vials for 5 days, after which the adults were removed. After

the six days of rifampicin treatment, the treated wasps were placed into the vials with fly larvae and

allowed to infect as many larvae as possible. Wasp offspring eclosed after 25-30 days and all female and

male offspring were collected and counted. Rifampicin treatment trials were performed with larval hosts

from fly species that were collected on the island of Saint Lucia, either *D. melanogaster* or *D. willistoni*.

Regardless of the host species, the sex ratio of the wasp offspring in each generation was always almost

all female or almost all male (see results, Figure 5), and therefore the data from the trials with different

host species were combined. The treatment conditions of each generation were as follows: G_1 unmated

females received 6 days of rifampicin treatment. The G_2 offspring were almost all female and also

received 6 days of rifampicin treatment, after which they produced F_0 offspring. In order to cross

Wolbachia depleted males and females together, a separate generation of G_1 females was treated with

rifampicin to produce G_2 females that would eclose at the same time as the F_0 males. These G_2 females

then received 6 days of rifampicin treatment and were then mated to F_0 males. Subsequent generations

(F_1 - F_5) were sexed and propagated by attempting sibling mating (when possible, see results) without

further rifampicin treatments.

Genomics: Oxford Nanopore library preparation and sequencing

DNA was extracted from a pool of ~100 adult female wasps using a Gentra PureGene Tissue Kit

(Qiagen). 3 ug of DNA was end prepped using NEBNext FFPE Repair Mix (M6630), NEBNext Ultra II End

repair/dA-tailing Module (E7546), and NEBNext Quick Ligation Module (E6056) and subsequently library

prepped using the ONT supplied SQK- LSK110 kit. Sequencing took place on an Oxford Nanopore

Minion over 3 days with nuclease wash followed by reloading after 24 hours. The sequencing computer

was built with an AMD Ryzen 3900x processor with 12 threads, 24 cores, 64 Gb RAM and a 1 Tb SSD,

and a GeForce 2080Ti with 4352 CUDA cores running Ubuntu Linux 20.04. We monitored the CUDA

cores with nvtop (https://github.com/Syllo/nvtop). Fastq files were generated from the Oxford Nanopore

Minion instrument and called with Guppy v6.1.5. We used the GPU enabled version of guppy in concert

to enable live basecalling with the fast base calling model and for post-hoc base calling with the super

(sup) accuracy model called via Minknow v22.05.5. All resulting fastq.gz files were concatenated and

used for further processing.

<u>Genomics: Illumina sequencing</u>

Total DNA was extracted from a pool of ten adult wasps using the DNeasy blood and tissue kit (Qiagen).

Sequencing library preparation was performed with the NEBNext Ultra II DNA library preparation kit (New

England Biolabs) according to the manufacturer's protocols including 10 cycles of PCR amplification. The

resulting library was subjected to 150-bp paired-end sequencing on the Illumina NextSeq 500 platform at

the Indiana University Center for Genomics and Bioinformatics (Bloomington, IN) to generate

181,561,108 read pairs. To aid in gene annotation we also sequenced wasp transcriptomes. Total RNA

was extracted from a flash-frozen, mixed stage pool of 10 wasps (adults, pupae, larvae) using the

Monarch® Total RNA Miniprep Kit (New England Biolabs) including the DNase treatment step. An RNA-

Seq library was prepared with the Illumina TruSeq Stranded mRNA Library Prep Kit, including rRNA

removal with oligo-dT beads, and sequenced on a NovaSeq with S4 chemistry at the University of

Minnesota Genomics Center to generate 64,677,726 150-bp paired-end reads.

<u>Genomics: Nuclear genome assembly, curation, and quality control</u>

All reads greater than 5kb were filtered with SeqKit v2.5.1 [21], assembled with Flye v2.9 [22], and the

assembly was polished with Medaka v1.5.0 (https://github.com/nanoporetech/medaka). Haplotigs were

purged with PurgeHaplotigs v1.1.2 [23]. Pilon v.1.22 [23] was used for polishing: first WGS Illumina reads

were mapped to the long-read assembly with bwa v0.7.17 [24], alignment files were converted and sorted

with samtools v1.10 [25], and finally pilon was called for polishing to generate an updated assembly. Four

rounds of pilon polishing were performed, each time mapping the short reads to the most recent pilon

output prior to pilon correction. Manual curation of the assembly and identification of cytoplasmic

genomes was informed by results derived from running blobtools [26]. We discarded contigs that were

deemed anomalous for the following reasons: (1) less than 5,000 bp in length with no BUSCOs (because

>5k reads were used during assembly, these are very likely to be chimeric and/or haplotigs), or, (2)

having blast hits to non-arthropod taxa across the length of the contig, unusual sequencing

coverage(<15X; >800X), and unusual GC content (outside of 23-33%). Next, we checked the previously

purged contigs and recovered those that appeared to encode for *Leptopilina* rRNAs. Assemblies were

assessed with BUSCO v.5.3.2 [27] against the hymenoptera_odb10 database in genome mode ('-m

genome'). Finally, NCBI's Foreign Contamination Screen (FCS) was used to detect and remove other

contaminants [28]. Specifically, a small number of adaptor sequences were found and trimmed from the

ends of contigs. Three contigs had adaptor sequence in the middle (>5,000 bp from the ends): for these,

the adaptor sequence was removed and the contig split into two separate contigs, as per NCBI/FCS

recommendations [28].

<u>Genomics: Methylation</u>

Nanopore runs were base called with guppy v6.3.8 using model

"dna_r9.4.1_450bps_modbases_5mc_cg_sup" to call 5-methylcytosines in a CpG context only. Aligned

bam reads containing modified base information were converted into bed format with modbam2bed, with

strict parameters on canonical and modified base probabilities where bases are only called as canonical

or modified if they have a greater than 95% probability estimate. Parameters: modbam2bed -m 5mC -t 32

-e -p total.modmapped --aggregate --cpg dtwasp-final-curated.fa total.modmapped.bam -a 0.05 -b 0.95 >

total.modmapped.bed". Global methylation was calculated by an awk script.

<u>Genomics: Mitogenome</u>

Mitogenome annotation with MITOS2 [29] and manual rearrangement to start the circular mitogenome at

Cox1 (as is convention) was done as in [30]. MITOS2 parameters included the RefSeq 63 Metazoa

reference and the invertebrate mitochondrial translation code. Manual curation was facilitated by

comparison to published mitochondrial genomes from *Leptopilina boulardi* (KU665622.1) and *Leptopilina*

syphax (MT649407.1). A phylogeny of *Leptopilina* was constructed based on COX1 ("COI") sequences

available from the DROP database [3]. A subset of the available *Leptopilina* sequences were selected

based on (1) sampling across the genus, (2) limiting the total number of sequences for any given species,

(3) length of the barcoding region sequenced, and (4) checking for open reading frame. In total 55

sequences were used for phylogenetic reconstruction. These include an outgroup (*Ganaspis kimorum*),

previously sequenced barcodes for the *Leptopilina malgretoutensis* STL6 colony [3], and the COI

sequence derived from the mitogenome assembly. Nucleotide sequences were aligned with MAFFT

v7.511 [31] using the L-INS-i alignment algorithm and a five point gap open penalty (-op 5). Comparison

to the translated amino acid alignment verified that codon relationships were maintained and we did not

need to back-translate an amino acid alignment. Ends of the nucleotide alignment were trimmed such that

all sequences were represented at all positions. Phylogenetic reconstruction was then performed with

IQtree v1.6.11 [32] using a pre-analysis model optimization step to select the final model (here,

TIM3+F+I+G4) plus 1000 "ultrafast" bootstraps.

Genomics: Repeat identification

Repetitive DNA content was estimated using RepeatModeler2 [33] using the curated Dfam database

release 3.5 July 2021 [34], combined with the Repbase RepeatMasker database downloaded on October

26, 2018. RepeatMasker was run with the newly identified library to annotate the assembly with the

following parameters: RepeatMasker v4.1.2-pl -pa 24 -s [sensitive mode] [using rmblast 2.11.0+] -lib

ultimatefamilies.fa ultimateGenome.fasta. Each repeat library was also mapped against all other species

genomes with the following parameters: RepeatMasker v4.1.2-pl -pa 24 -s [sensitive mode] [using rmblast

2.11.0+] -lib families.fa genome.fasta.

Genomics: Gene annotation and ortholog clustering

Gene models were predicted with GeMoMa v1.9 [35] and leveraged three lines of evidence: (1) RefSeq

annotations for *Leptopilina heterotoma* (GCF_015476425.1) (2) RefSeq annotations for *Leptopilina*

boulardi (GCF_019393585.1) and (3) RNA-seq reads derived from the mixed pool of wasps. RNA-seq

reads were mapped to a hard masked version of the reference genome using STAR v2.5.3a [36].

GeMoMa was called with the following additional parameters: tblastn=TRUE, AnnotationFinalizer.n=false,

AnnotationFinalizer.r=SIMPLE, AnnotationFinalizer.t=true, AnnotationFinalizer.p=LMAL, pc=true, o=true.

To generate annotations for *Leptopilina clavipes* and *Ganaspis kimorum*, we ran the same annotation

pipeline on hard-masked assemblies, only without the RNA-seq data. Protein sequences from each

genome were clustered into orthologous groups using OrthoFinder v.2.5.4 [37]. For genes represented by

multiple transcript variants, only the longest isoform was used in clustering.

Genomics: *Wolbachia* genome

A single contig was identified to be a *Wolbachia* genome, based on size, circularity, and Blobtools results.

Genome completeness was assessed with BUSCO v5.3.2 [27] against the rickettsiales_odb10 database

in genome mode ('-m genome'). Genome annotation and phylogenetics were performed by running the

Wolbachia Phylogeny Pipeline (WHOP; https://github.com/gerthmicha/WHOP [38]). In brief, the pipeline

takes a provided database of bacterial genomes (*Wolbachia* strains and Rickettsiales outgroups) plus

additional genomes provided by the user, annotates them with Prokka v1.14.6 [39], clusters protein

sequences into orthogroups with OrthoFinder v2.5.4 [37], aligns the single-copy orthologs with MAFFT L-

INS-i v7.487 [31], removes recombining genes with PhiPack v1.1 [40], and then concatenates the

alignments prior to phylogenetic reconstruction with IQtree v2.2.3 [32]. IQtree was first run with a pre-

analysis model optimization step (-m LG+I+G4 -pre -redo -fast), followed by final reconstruction with the

selected model and 1000 bootstrap replicates. The *w*Lmal genome was re-oriented to start at *dnaA* and

proksee (https://proksee.ca/ [41]) was used for visualization. Prophage regions were identified with

VirSorter2 v.2.2.4 [42], and Phigaro v.2.3.0 [43], and other mobile elements were identified with

mobileOG-db v.1.0.1 [44], all implemented in proksee with default parameters. To aid in manual

annotation of *w*Lmal we leveraged the orthogroup relationships between *w*Lmal and *w*Mel generated by

WHOP to check for specific proteins. Additionally, we used blastp and tblastn with default parameters to

query against the protein sequences or genome, respectively, for a select number of functionally relevant

Wolbachia proteins not found in *w*Mel. All manual annotation queries and matches are in Supplemental

Tables S5-6.

Data visualization

Data visualization was carried out in R version 4.1.3 [45] and leveraged Paul Tol's "Medium-contrast

qualitative colour scheme", (R: khroma [46]). Newick files from phylogenetic reconstructions were

visualized in FigTree v1.4.4 (https://github.com/rambaut/figtree/) and annotated in Inkscape. Proksee

(https://proksee.ca/ [41]) was used for genome visualization, and the r package 'gggenes' v.0.5.1 was

used to plot gene models.

RESULTS

<u>**Collections on Saint Lucia**</u>

Banana and calabasa fruit traps were used to collect *Drosophila* and their associates on Saint Lucia in

2019. The calabasa traps at STL3 yielded six female *Drosophila antillea* and one female *Drosophila*

willistoni. Isofemale lines of *D. antillea* and *D. willistoni* were established from single females that

originated from the STL3 collection site. At site STL6, banana traps contained a total of 12 *melanogaster*-

like flies and 1 *repleta* group fly. Calabasa traps contained 4 *melanogaster*-like flies, 4 *repleta* group flies

and 20 *dunni*-group flies. Wasp "STL6" was collected from a calabasa trap on the morning of 1 March

2019: a clear day with a high temperature of 27C (Figure 1). The STL6 wasp was placed in a vial with

developing *Drosophila willistoni* (from the STL3 site, see above) to initiate a culture, and then shipped

back to Indiana University where they have been maintained since (see methods).

<u>***Leptopilina malgretoutensis* Buffington, Lue, Davis & Tracey sp. nov.**</u>

Species Diagnosis. *Leptopilina malgretoutensis* is distinguishable from other *Leptopilina* species by a

number of characters, including: flagellomere colors, in which F1-F4 are honey-brown, F5-F10 are dark

brown, and F11 is yellow (Figure 2A); the typical flagellomere color in *Leptopilina* do not match the unique

combination here. Latero-posterior margin of pronotal plate is with elongate, golden setae that reach the

anterior margin of the mesoscutum (Figure 2B (LM) and Figure 3A-B (SEM)). No other known *Leptopilina*

have setae so long off the pronotal plate (Figure 2B); other species have hairs here, but not so long.

Members of the *clavipes* group *sensu* [15] tend to also have sparse, scattered setae over the

mesoscutum, whereas *L. malgretoutensis* is entirely glabrous (Figure 3A-B).

Morphological Description

Coloration with head, mesosoma, metasoma black to dark brown, legs light brown. Sculpture on vertex,

lateral surface of pronotum and mesoscutum absent, surface smooth.

Antenna. Female antenna composed of scape, pecidel, 11 flagellomeres (F1-F11). F1-F4 sub-

cylinderical, slightly expanded distally, smooth, non-claval segments. F5-F11 claval segments, abruptly

broader than non-claval segments, distinctly barrel-shaped, with slightly more erect, dense setae; claval

segments with placoidal sensillae distributed evenly around each segment. Single campaniform sensillae

present ventro-apically on F6-F10 (arrow, Figure 3C; compare to Figure 3D). Articulation between

flagellomeres in antenna moniliform, segments distinctly separated by narrow neck-like articulation. Male

antenna composed of 13 flagellomeres, F1 as long as F2, slightly curved and excavated distally;

campaniform sensillae on all flagellomeres.

Head. In anterior view, broadly triangular. Pubescence on head sparse, single setae scattered over face.

Sculpture along lateral margin of occiput absent. Gena (measured from compound eye to posterolateral

margin of head) short, ratio of length of gena to length of compound eye in dorsal view <0.3. Sculpture of

gena absent, smooth. Lateral margin of occiput evenly rounded, not well defined. Occiput (except

extreme lateral margin) smooth. Carina issuing from lateral margin of postocciput absent. Ocelli small,

ratio of maximum diameter of a lateral ocellus to shortest distance between lateral ocelli 0.2-0.4. Anterior

ocellus far from posterior ocelli, clearly separate anterior ocellus to anterior margins of posterior ocelli.

Relative position of antennal sockets intermediate, ratio of vertical distance between inner margin of

antennal foramen and ventral margin of clypeus to vertical distance between anterior ocellus and

antennal rim 2.0-4.0. Median keel absent. Vertical carina adjacent to ventral margin of antennal socket

absent; facial sculpture absent, surface smooth. Facial impression absent, face flat. Antennal scrobe

absent. Anterior tentorial pits small. Vertical delineations on lower face absent. Ventral clypeal margin

laterally, close to anterior mandibular articulation, straight. Ventral clypeal margin medially straight, flat,

with 4-6 setae overhanging base of mandibles. Clypeus smooth. Malar space adjacent to anterior

articulation of mandible evenly rounded, smooth. Malar sulcus present, composed of single groove. Eye

removed from lateral ocelli, ratio of distance between compound eye and posterior mandibular articulation

to distance between posterior ocellus and compound eye <1.2. Compound eyes, in dorsal view,

moderately protruding from the surface of the head. Pubescence on compound eyes absent. Orbital

furrows absent. Lateral frontal carina of face absent. Dorsal, posterior aspect of vertex smooth. Hair

punctures on lateral aspect of vertex absent. Posterior surface of head deeply impressed around

postocciput.

Mouthparts. Apical segment of maxillary palp with pubescence, consisting of one long erect setae. Apical

seta on apical segment of maxillary palp longer than twice length of second longest apical seta. Maxillary

palp composed of four segments. Last two segments of maxillary palp (in normal repose) straight. Apical

segment of maxillary palp about 1.5 times as long as preceding segment. Left mandible with three teeth;

scattered setae dorsally.

Mesosoma. Long auburn-golden setae present, originating from anterior inflection of pronotum, extending

to anterior margin of mesoscutum; rest of pronotum glabrous (Figure 3A-B). Anteroventral inflection of

pronotum narrow. Pubescence on lateral surface of pronotum absent. No ridges present posterior to

pronotal plate. Anterior flange of pronotal plate distinctly protruding anteriorly, completely smooth. Ridges

extending posteriorly from lateral margin of pronotal plate absent. Lateral pronotal carina absent. Crest of

pronotal plate absent. Dorsal margin of pronotal plate (in anterior view) spatulate. Submedian pronotal

depressions open laterally, deep. Lateral margin of pronotal plate defined all the way to the dorsal margin

of the pronotum. Width of pronotal plate narrow, not nearly as wide as head.

Mesoscutal surface convex, evenly curved. Sculpture on mesoscutum absent, entire surface smooth,

shiny, with sparse long hairs. Notauli absent. Median mesoscutal carina absent. Anterior admedial lines

absent. Median mesoscutal impression absent. Parascutal carina nearly straight.

Mesopleuron entirely smooth. Subpleuron entirely smooth, glabrous. Lower pleuron entirely smooth,

glabrous. Epicnemial carina absent. Lateroventral mesopleural carina present, marking abrupt change of

slope of mesopectus. Mesopleural triangle absent. Subalar pit not visible. Speculum absent. Mesopleural

carina present, complete, composed of one complete, uninterrupted carina. Anterior end of mesopleural

carina inserting above notch in anterior margin of mesopleuron; posterior end terminates at setal pit of

lower metapleuron.

Dorsal surface of scutellum foveate-areolet (Figure 3E). Circumscutellar carina present, complete,

delimiting dorsal and ventral halves of scutellum. Posterior margin of axillula marked by distinct ledge,

axillula distinctly impressed adjacent to ledge. Latero-ventral margin of scutellum, posterior to axillula,

weakly rugulose. Dorsoposterior part of scutellum rounded. Transverse median carina on scutellar plate

absent. Dorsal part of scutellum entirely areolate. Scutellar plate, in dorsal view, medium sized, exposing

about half of scutellum, gently foveate with three to four single setae; glandular release pit at posterior

end, oval. Scutellar fovea present, two, distinctly margined posteriorly, smooth. Longitudinal scutellar

carinae absent. Single longitudinal carina separating scutellar foveae present, elongage, ending past

posterior margin of foveae, gently expanding into teardrop-shaped scutellar plate. Postero-lateral margin

of scutellum rounded. Lateral bar smooth, narrow.

Posterior impression of metepimeron absent. Metapectal cavity anterodorsal to metacoxal base present,

well-defined (Figure 3F, arrow A). Anterior margin of metapectal-propodeal complex meeting

mesopleuron at same level at point corresponding to anterior end of metapleural carina. Posteroventral

corner of metapleuron (in lateral view) not extended posteriorly. Anterior impression of metepimeron

absent. Posterior margin of metepimeron distinct, separating metepimeron from propodeum, with distinct

ridge (Figure 3F, arrow B). Anteroventral setal pit of lower metapleuron present, with foamy setae.

Subalar area broadened anteriorly, narrowed posteriorly. Prespiracular process present, blunt, lobe-like,

polished, at end of polished prespiracular groove; posteroventral corner of prespiracular groove extended

to distinct lobe. Dorsellum absent. Anterior impression of metepisternum, immediately beneath anterior

end of metapleural carina, present. Pubescence consisting of few hairs on posterior part of

metepisternum, few sparse hairs on propodeum, but with distinct posterior swatch of foamy setae at

midline of metapleuron.

Pubescence posterolaterally on metacoxae sparse, with a single postero-dorsal button of dense

pubescence. Microsculpture on hind coxa absent. Longitudinal ridge on the posterior surface of metatibia

absent. Metafemoral tooth present, elongate, with adjacent serrate ridge posteriorly. Ratio of first

metatarsal segment to remaining 4 segments less than 10; metatarsus 5 distinctly elongate relative to 2

through 4.

Wing vein M absent. Pubescence of fore wing present, long, moderately dense on most of surface. Apical

margin of female fore wing rounded. Rs+M of forewing defined but nebulous at point of origin from basal

vein at posterior third. Mesal end of Rs+M vein situated closer to anterior margin of wing, directed

towards middle of basalis. Vein R1 forming marginal cell completely. Basal abscissa of R1 (the abscissa

between 2r and the wing margin) of fore wing as broad as adjacent wing veins. Coloration of wing absent,

entire wing hyaline. Marginal cell of fore wing membranous, similar to other wing cells. Areolet absent.

Hair fringe along apical margin of fore wing present, long or very long (Figure 2C).

Propodeal spurs absent. Lateral propodeal carinae present, not reaching scutellum. Ventral end of lateral

propodeal carina reaching nucha, carinae separated from each other, elbowed at midpoint, not parallel.

Inter propodeal carinae space lightly setose, underlying surface appearing smooth. Petiolar rim of uniform

width along entire circumference. Petiolar foramen removed from metacoxae, directed posteriorly.

Horizontal carina running anteriorly from lateral propodeal carina absent, moderately setose. Calyptra, in

lateral view, rounded. Propodeum not strongly drawn out posteriorly. Calyptra, in posterior view,

dorsoventrally elongate or rounded.

Metasoma. Petiole about as long as wide. Surface of petiole longitudinally costate, ventral keel absent.

Posterior part of female petiole abruptly widened. Ventral and lateral parts of petiolar rim broad.

Setal band (hairy ring) at base of tergum 3 present, interrupted dorsally, ventrally. Tergum 3 indistinct,

fused with syntergum. Posterior margin of tergum 3 indistinct, fused with tergum 4 in syntergum. Posterior

margin of tergum 4 evenly rounded. Sternum 3 encompassed by syntergum. Sculpture on metasomal

terga absent. Syntergum present with terga 3 to 5 fused, ventral margin rounded. Peglike setae on T6-T7

absent. Postero-ventral cavities of female metasoma T7 present, glabrous save for few, long setae.

Female postero-ventral margin of T6-T7 straight, parallel. Terebrum and hypopygium (in lateral view)

curved, pointing upward. Ovipositor clip present.

Etymology. The species epithet refers to locations on Saint Lucia: Malgretoute, a town near the island's

active volcano, and the near-by Malgretoute Beach, from which the iconic Saint Lucia Pitons are visible.

Material examined. Holotype: Female. Saint Lucia, Barr d'Isle trail, central. lat 13.924, long -60.959,

elevation 308.7m. 17.II.2019, Jeremy S. Davis, collected from calabasa bait trap. USNMENT01025555.

[1] Actual specimen taken from Indiana University colony derived from the above collection locality.

Paratypes: 9 females, 3 males. Same collection data as holotype. USNMENT01557161,

USNMENT01025556, USNMENT01025557, USNMENT01025554, USNMENT01557162,

USNMENT01557158, USNMENT01448469, USNMENT01448467, USNMENT01932975-

USNMENT01932978.

Other material examined. 8 females. INRA laboratory, Petit Bourg, Guadeloupe, January 1982. Carton,

coll. USNMENT01025775-USNMENT01025782

504
505 **Figure 2. *Leptopilina malgretoutensis.* (A)** Lateral view, **(B)** lateral view of mesosoma, and **(C)** wing
506 vein pattern.

507
508 **Figure 3. SEM images of *Leptopilina malgretoutensis.* (A)** Top view of mesoscutum, **(B)** Lateral-
509 posterior view of mesosoma, **(C)** campaniform sensillae, white arrow, compared to **(D)** no campaniform
510 sensillae, **(E)** scutellum plate, and **(F)** metepimeron.
511

Leptopilina malgretoutensis is highly virulent on _D. antillea_ flies

As noted extensively above, our laboratory cultures of _L. malgretoutensis_ were collected on Saint Lucia.

This island is also the only reported location for the _Drosophila antillea_ species of the dunni subgroup. To

our knowledge, the susceptibility of dunni group _Drosophila_ to parasitoid wasps has not been previously

investigated. Thus, we wondered if the _D. antillea_ flies would be susceptible, or if they were possibly

resistant, to infection by a wasp species that is also found on Saint Lucia. We allowed female wasps to

oviposit for 24 hours on larval cultures that were derived from isofemale lines of either _D. antillea_ or _D._

willistoni (that both originated from the STL3 collection site) and then determined the proportion of wasps

and flies that emerged from the infection vials. The results clearly indicated a high degree of virulence of

L. malgretoutensis on _D. antillea_ (76.5% ± 5.6% wasp eclosion) relative to _D. willistoni_ (20% ± 3.8% wasp

eclosion) as host species (Figure 4)_.

**Figure 4. _Leptopilina malgretoutensis_ shows high virulence on a host fly species that is exclusive
to Saint Lucia.** Percentage of adult flies and wasps that eclosed from vials infected by _L._
malgretoutensis, and from uninfected control vials. Mortality represents the percentage of larvae that did
not develop to adulthood. _L. mal._ is clearly more virulent on _Drosophila antillea_ ("D. ant") hosts than on a
Drosophila willistoni ("D. wil") host strain that was also collected on Saint Lucia. Data are plotted as mean
± SEM; n = 8 infection vials, n = 3 uninfected control vials.

Wolbachia* mediates obligate thelytokous parthenogenesis in *Leptopilina malgretoutensis

While maintaining *L. malgretoutensis* wasp lines we noticed that the cultures were primarily female (>95%). And, in one generation of passaging, the cultures became entirely female. Fearing that we would lose the culture, we infected flies with these female wasps in an attempt to obtain haploid male offspring that could be mated back to their mothers to obtain more diploid females. However, the cultures generated by the virgin female wasps were again entirely female. This indicated that this species of wasp was capable of asexually producing females (*i.e.*, thelytokous parthenogenesis). Furthermore, the observation that males were rarely obtained in the cultures was unlikely to be strictly a consequence of the wasps having especially high rates of fertilization (which gives rise to female offspring in arrhenotokous, sexual, haplodiploids). In many parasitic wasp species, including a suite of other *Leptopilina sp.*, thelytokous parthenogenesis is mediated by vertically transmitted bacterial symbionts such as *Wolbachia* (α-proteobacteria: Rickettsiales) [47-49]. We tested for the presence of *Wolbachia* infection by Western blot, targeting a *Wolbachia* specific protein and found that indeed the wasps were infected with this microbe (Supplemental Figure S1).

To determine if a microbe was responsible for the asexual production of females, we used antibiotic treatment with rifampicin to remove the infection and assess changes in the offspring sex ratio derived from unmated females (Figure 5A). After treating unmated adult females (G_1) with rifampicin, they were allowed to parasitize host larvae. Their broods (G_2) were almost entirely female (98.6% ± 2.6% female). Critically, *Leptopilina* are largely proovigenic: all their eggs develop and mature during the pupal period [50]. This means that symbiont-mediated effects are likely established much earlier, and treating adults is "too late" (as has been seen in other wasps [51]). Thus, we performed a second generation of rifampicin treatment: G_2 rifampicin-treated females produced offspring (F_0) that were almost all male (99.7% ± 0.3% male), a clear indicator of symbiont-mediated parthenogenesis.

We attempted to generate a symbiont-free sexually reproducing line of wasps (Figure 5B) but were unsuccessful. Specifically, symbiont depleted males and treated G_2 females were crossed to assess sexual reproduction, but offspring continued to be almost all male (F_1-F_2, 91.2% ± 1.5% and 98.5% ±

561   0.6% male, respectively), indicating no sexual reproduction (Figure 5C). After antibiotic treatment was

562   discontinued, rare female progeny that were allowed to reproduce produced sex ratios that gradually

563   returned to being almost entirely female (F_3-F_4, 80% and 100% female, respectively), and unmated

564   females were again able to produce female offspring by parthenogenesis (F_5-F_6, 98.5% and 100%

565   female, respectively).

566
Figure 5. *Leptopilina malgretoutensis* reproduces by symbiont-mediated thelytokous
parthenogenesis. (A) Schematic depicting the production of male progeny after two generations of
antibiotic treatment ("+rif"), which supports the hypothesis of symbiont mediated parthenogenesis. **(B)**
Schematic of antibiotic treatment and crossing scheme to test for sexual reproduction after symbiont
depletion, and an attempt to generate a sexual, symbiont free strain. **(C)** Percentage of female and male
offspring in each generation. Data is plotted as mean ± SEM. For generations G_2 to F_2, n = 4-8 infection
vials; for generations F_3 to F_6, n = 1 infection vial.
574

***Leptopilina malgretoutensis* genome sequencing and assembly**

We generated a genome assembly for *Leptopilina malgretoutensis* based on nanopore long read

sequencing (Table 1, Supplemental Tables S1-S3). The draft assembly with Flye leveraged all nanopore

reads longer than 5,000 bp (Supplemental Table S1) and generated 1,666 contigs totaling just over 1Gbp

in total length (Table 1; Supplemental Figure S2). The assembly was then polished with illumina short

reads: short-read mapping rates were very high (98.9% of reads mapped). After polishing, separating

cytoplasmic genomes, removing haplotigs, purging contaminants, and removing otherwise spurious

contigs, the ultimate nuclear genome assembly was 1,411 contigs and ~995Mbp in size with an N50 of

>1.8Mbp (Table 1). The longest contig in the assembly was close to 13Mbp in length. Assembly quality

was assessed with BUSCO. Just under 90% of the hymenopteran BUSCO proteins were complete,

similar to other assemblies in the genus (Table 2). We recovered two cytoplasmic genomes:

mitochondrion and *Wolbachia*. Additional low-coverage bacterial sequences that were purged from the

assembly belonged to typical associates of the fly food and microbiome (Supplemental Results in File S1

and Supplemental Table S3). No other symbionts were identified in the contigs (e.g., *Cardinium*,

Spiroplasma, etc [52-54]). Thus, we determine that *Wolbachia* is likely the cause of thelytokous

parthenogenesis.

Table 1. Assembly statistics for *Leptopilina malgretoutensis*

Metric	Draft	Ultimate
Contigs	1,666	1,411
Length (bp)	1,004,830,435	994,936,849
Min length	499	5,452
Avg length	603,134	705,129
Max length	12,918,385	12,924,377
N50	1,831,405	1,832,545
%GC	29.85	29.75
BUSCO* (Hymenoptera)	C:89.3%[S:88.8%,D:0.5%], F:2.9%,M:7.8%,n:5991	C:89.7%[S:89.2%,D:0.5%],F :2.7%,M:7.6%,n:5991

593 *Standard BUSCO annotation: Complete BUSCOs (C) [Complete and single-copy BUSCOs (S),
594 Complete and duplicated BUSCOs (D)], Fragmented BUSCOs (F), Missing BUSCOs (M), Total BUSCO
595 groups searched (n)
596

604 **Table 2. Additional genomes used in comparative analyses.**

Species	Assembly Accession	Size (bp)	Protein Coding Genes	BUSCO* (Hymenoptera)
Leptopilina boulardi	ZJU_Lbou_2.1	354,804,354	13,647	C:91.4%[S:90.9%,D:0.5%], F:1.7%,M:6.9%,n:5991
Leptopilina clavipes	ASM185565v1	255,343,917	13,920	C:77.8%[S:77.3%,D:0.5%], F:7.3%,M:14.9%,n:5991
Leptopilina heterotoma	ASM1547642v1	484,694,486	14,450	C:91.1%[S:90.6%,D:0.5%], F:1.9%,M:7.0%,n:5991
Ganaspis kimorum[#]	GCA_037103525.1	1,025,447,244	20,567	C:89.4%[S:88.8%,D:0.6%], F:2.7%,M:7.9%,n:5991

605 * Standard BUSCO annotation: Complete BUSCOs (C) [Complete and single-copy BUSCOs (S),
606   Complete and duplicated BUSCOs (D)], Fragmented BUSCOs (F), Missing BUSCOs (M), Total
607   BUSCO groups searched (n). Hymenoptera dataset used.
608 [#] Previously '*Ganaspis* near *brasiliensis'* (including in genome report [55]), but recently re-assigned [6].
609

Methylation

We used a methyl-sensitive base-calling model to predict 5-methylcytosines in a CpG context. Across the

nuclear, mitochondrial, and *Wolbachia* genomes, fractional methylation was less than 1% (Genome:

0.600%, mitochondrion: 0.547%, *Wolbachia*: 0.598%). These estimates likely do not represent low levels

of true genomic methylation, rather, they are probably background noise due to error rates inherent to the

sequencing and base calling.

Mitogenome

We recovered a circular mitochondrial genome (14,490 bp; 20.6% GC; 3,562X sequencing coverage,

Figure 6, Supplemental Table S3). Using the barcoding sequence region from the mitochondrial genome,

plus representative sequences across the genus, we reconstructed a phylogeny of *Leptopilina sp.* (Figure

6A). *Leptopilina malgretoutensis* sequences were all identical, and formed a monophyletic group more

than 10% divergent from the next most closely related species in the dataset, further supporting its

designation as a new species. Importantly, *Leptopilina malgretoutensis* is sister to a clade within the

Leptopilina genus that contains all other species for which genome sequences are available (*Leptopilina*

clavipes, Leptopilina boulardi, and *Leptopilina heterotoma*). Annotation of the mitochondrial genome

revealed that gene order was largely conserved as compared to other *Leptopilina* mitochondria.

Additionally, we annotated the expected rRNAs and protein coding genes. However, insect mitochondrial

tRNAs are often difficult to annotate [56], and in *Leptopilina malgretoutensis* we were unable to find

tRNAs for asparagine (N), alanine (A), aspartic acid (D), pheylalanine (F), serine anticodon-1 (S1), and

arginine (R). Related to that, the region between the 3' end of the 16S rRNA (rrnL) and the 5' end of

nad2, was quite divergent as compared to *Leptopilina boulardi,* in which there is a cluster of tRNAs

(including D, F, S1, N, A, plus glutamic acid, glycine, and histidine). It is likely that divergence and/or

noncanonical tRNA structures, as is often the case in insect mitochondria, may be precluding a complete

annotation.

635
Figure 6. Mitochondrial phylogeny and genome of *Leptopilina malgretoutensis*. (A) Phylogenetic
reconstruction of *Leptopilina,* based on 480 nucleotides of Cox1, aligned to maintain codon position
relationships. Node colors indicate bootstrap support as indicated in legend. The three *Leptopilina*
malgretoutensis sequences are identical. The two other *L. malgretoutensis* sequences are from

previously barcoded STL6 specimens [3]. Sequences assigned to provisional species are indicated by
'sp' followed by a number, as per DROP database [3]. Accession numbers are in gray at the end of each
leaf label. **(B)** Map of the assembled mitochondrial genome and annotations.
643

Gene Annotation

Annotation of the repeat-masked *Leptopilina malgretoutensis* genome revealed 16,402 genes (Table 3),

which is in-line with annotations of closely related species (Table 2). To better understand genome

evolution, we clustered protein sequences (n=78,896 in total) of the four *Leptopilina* species plus

outgroup species *Ganaspis kimorum* [6, 55, 57-59]. As expected with a small set of closely related

species, we identified orthologs for most genes: 97.1% of the genes were clustered into 13,659

orthogroups, each of which contained 5.6 genes on average. 7,266 orthogroups contained genes from all

five species, and of those, 6,389 orthogroups were strictly single-copy. 1,077 orthogroups were species-

specific and only contained in-paralogs.

Table 3. Annotation metrics for the *Leptopilina malgretoutensis* genome.

Metric	Value*
Number of genes	16,402
Number of mRNAs	18,165
Number of exons	97,163
Number of introns	78,998
Mean mRNAs per gene	1.1
Mean exons per mRNA	5.3
Total gene length	158,352,048 bp
Longest gene	235,629 bp
Mean gene length	9,654 bp
Longest CDS	76,353 bp
Mean CDS length	1,541 bp
Longest exon	16,278 bp
Mean exon length	288 bp

655 *Base pairs = bp
656

Repetitive elements

Given the larger than expected size of the *Leptopilina malgretoutensis* genome relative to other

Leptopilina species, we analyzed the repeat content of the assembly using a *de novo* repeat identification

pipeline. 66.81% of the nearly 1Gbp genome assembly was identified as repetitive (Figure 7,

Supplementary Table S4). Retroelements and DNA transposons only constituted 9.57% and 5.09% of the

length of the assembly, respectively. The majority of the repeat content (48.97% of the assembly) was

categorized as unclassified repetitive elements. To determine if this large repetitive fraction was unique to

Leptopilina malgretoutensis we ran the same *de novo* repeat identification pipeline on the three other

Leptopilina species with genome assemblies available, plus an outgroup from a closely related genus,

Ganaspis kimorum (Supplemental Table S4). Indeed, the other *Leptopilina* species were less repeat

dense with 35.40%, 48.01%, and 57.27% repeat content for *L. clavipes, boulardi, and heterotoma,*

respectively. Similar to *Leptopilina malgretoutensis*, the majority of these repetitive elements were

unclassified. In fact, the overall size and makeup of the *Leptopilina malgretoutensis* genome was much

more similar to the outgroup *Ganaspis kimorum* (Figure 7A,B)*.* However, the *Ganaspis kimorum* and

Leptopilina malgretoutensis genomes contained approximately double the amount of non-repetitive

genome as compared to the other *Leptopilina* (black bars in Figure 7B). Thus, while *Leptopilina*

malgretoutensis had the largest repetitive content in the genus (both by total length and percentage),

repeat content alone does not completely explain the overall difference in genome size relative to the

other congenerics.

To understand the large differences in repeat content across the *Leptopilina* genomes, we performed an

all-versus-all analysis in which we mapped each species' repeat library to all other species' genomes

(Figure 7C). Then we determined how many of the unique repeat families were present to infer if there

was simply a difference in the number of repeat occurrences (e.g., family size), or, if entire repeat families

had been gained or lost across the clade. The most repeat dense genomes, *Ganaspis kimorum* and

Leptopilina malgretoutensis, contained 3,563 and 4,002 repeat families respectively, likely indicating a

gain of repeats in the *Leptopilina* common ancestor. Accordingly, while 84.1% of the *Ganaspis kimorum*

repeat families occur in the *Leptopilina malgretoutensis* genome, only 82.5% of the *Leptopilina*

malgretoutensis repeat families occur in *Ganaspis kimorum*. There appears to have been a large loss of

repeat families in the ancestor of the clade containing *L. boulardi*, *L. heterotoma*, and *L. clavipes*, and

additional loss in the lineage leading to *L. clavipes*. These genomes contained 2,727, 2,742, and 1,718

families respectively, and between 88-91% of these families were found in *L. malgretoutensis*.

Furthermore, while only 78-80% of the *L. malgretoutensis* repeats were identified in these other

Leptopilina genomes, an even lower percentage of the *Ganaspis kimorum* repeats were present (73.3-

77.5%). Finally, the comparisons within the clade of *L. boulardi*, *L. heterotoma*, and *L. clavipes*, indicate

high levels of shared repeats (85.8-93.9%). All together, these data indicate that (1) the ancestor of the

four *Leptopilina* species here acquired repetitive elements that are not present in *Ganaspis kimorum*, (2)

many of the repeats have since been lost in *L. boulardi*, *L. heterotoma*, and especially in *L. clavipes*.

696
Figure 7. Evolution of *Leptopilina* genomes. (A) Species tree inferred by OrthoFinder, rooted on
outgroup, *Ganaspis kimorum*. Phylogeny is annotated with (1) the number of species-specific genes (gray
boxes), which include both singletons, and species-specific orthogroups, and (2) inferred gains and
losses of orthogroups. "Gained" orthogroups are those that are unique to and present in all taxa within the
clade, and "lost" orthogroups are present in all the outgroups, but none of the ingroups. **(B)** Genome

content of each representative assembly. **(C)** All-verus-all comparison of repetitive elements. Libraries of
the *de novo* identified repeats from each species were mapped to all other genomes, and the number of
unique repeat families recovered in each comparison is indicated within each box in the heat map
(number of families identified in "genome"/number of unique families in "library").
706

***Wolbachia* Strain *w*Lmal**

We identified a circular *Wolbachia* genome in our assembly, hereinafter strain "*w*Lmal" (Table 4). A single

contig was sequenced at 344X coverage and blast results from blobtools revealed similarity to *Wolbachia*

strain *w*Ha from *Drosophila simulans* across most of the contig. BUSCO genome completeness metrics

were very high, with 99.2% of Rickettsiales BUSCOs present in single copy (Table 4). Phylogenetic

reconstruction based on core single-copy orthologs revealed that *w*Lmal is closely related to *Drosophila*

infecting *Wolbachia* strains (e.g., *w*Mel, *w*Ha*, w*Ri, *w*Au) in *Wolbachia* Supergroup A (Figure 8A). This is

notable not only due to the host-parasitoid relationship of *Leptopilina* and *Drosophila*, but also because

the parthenogenesis-inducing "*w*Lcla" strain from *Leptopilina clavipes* is quite divergent, and nested

within a different major *Wolbachia* clade: Supergroup B [60, 61]. The size (~1.2Mbp) and coding content

of the *w*Lmal genome is quite typical of an insect-infecting *Wolbachia* strain [62, 63] (Figure 8B, Table 4).

Two putative prophage regions were identified. The first (1,00,3471 bp - 1,051,238 bp) is partial and is

likely remnants of a *Wolbachia* "WO" prophage. This prophage region contains some phage structural

elements (primarily proteins for the tail) and what looks to be part of a "Eukaryotic Association Module"

(EAM), a region associated with WO phages which often contains proteins for manipulating host

reproduction and cell biology [64]. The second predicted prophage region (824,983 bp - 864,973 bp)

appears much more degenerate and only has a few remaining open reading frames encoding for phage

proteins.

We searched for the two recently identified putative parthenogenesis-inducing effector proteins, *pifA* and

pifB, which are found in the parthenogenesis inducing strains *w*Lcla and *w*Tpre [60]. *pifA* is also present

in the PI-*Wolbachia* infecting *Encarisa formosa* (designated *"piff"* in that strain) [65]*.* We identified two

identical copies of *pifB*, but did not find evidence for *pifA* in *w*Lmal (Supplemental File S1). Closer

inspection revealed that the *pifB* copies are in opposite orientations, and one copy of *pifB* is surrounded

by insertion elements and is just downstream of a putative *hin* DNA invertase, likely from the WO phage.

We also queried the *w*Lmal genome for a suite of other notable *Wolbachia* proteins and identified other

putative or confirmed effector proteins. Surprisingly, these included the genes *cifA* and *cifB* (here,

specifically "type I" or *cid*-type [66] which cause an entirely different host reproductive phenotype: the

conditional sperm-egg incompatibility known as cytoplasmic incompatibility (CI).

Table 4. *Wolbachia* strain *w*Lmal genome assembly and annotation.

Metric	*w*Lmal
Contigs	1
Length (bp)	1,198,858
%GC	35.1
BUSCO* (Rickettsiales)	C:99.2%[S:99.2%,D:0.0%], F:0.3%,M:0.5%,n:364
CDS	1,356
rRNAs	3
tRNAs	34

738   *Standard BUSCO annotation: Complete BUSCOs (C) [Complete and single-copy BUSCOs (S),
739   Complete and duplicated BUSCOs (D)], Fragmented BUSCOs (F), Missing BUSCOs (M), Total BUSCO
740   groups searched (n)

Figure 8. *Wolbachia* strain *w*Lmal. (A) Phylogenetic relationships of *Wolbachia* strains and Rickettsiales relatives based on 127 core single-copy orthologs. **(B)** Map of the complete *w*Lmal genome. Each concentric circle is annotated with a different set of genomic information. The innermost circle is annotated with a basepair marker, the second circle indicates GC content, the third circle indicates prophage regions identified by VirSorter, and the outermost circle shows the coding regions of the genome. Abbreviations: Type IV Secretion system (T4SS), noncoding RNA (ncRNA), coding sequences

(CDS). Putative effector proteins are indicated with the gene name where possible, otherwise the wLmal
CDS number is indicated and the *w*Mel homologs can be found in Supplemental Table S5. **(C)** Gene
models in the prophage region containing remnant WO phage structural proteins and a suite of effector
proteins homologs. Notably the two *pifB* coding sequences are present in an inverted repeat. Open
reading frames labeled with numbers (e.g., 1167) are gene annotation IDs, provided to benchmark the
region.
754

Historical specimens of *Leptopilina malgretoutensis*

As noted above, our Saint Lucian specimens matched previously undescribed *Leptopilina* collected from

the Caribbean in the 1980's (specimens USNMENT01025775-USNMENT01025782). Specifically, those

Leptopilina (n=16, all female) were collected in 1982 and 1983 from Grande Terre, Guadeloupe (~270 km

from our collection site on Saint Lucia), some of which were used to initiate a now extinct colony, "PB10-

5". The PB10-5 wasps were previously characterized as an undescribed species nested within an early-

branching *Leptopilina* lineage, in the same clade as *Leptopilina longipes* [67, 68]. Our COI phylogenetic

reconstruction of *Leptopilina* recapitulated these relationships (Figure 6A). In addition to all the collected

Guadeloupe wasps being female, Yves Carton's notes indicated "*[the PB10-5 colony] emerged only*

females: we can conclude that this species is parthenogenetic." In summary, our data indicate that *L.*

malgretoutensis was previously collected ~40 years ago on the neighboring island of Guadeloupe and

also reproduced via thelytokous parthenogenesis.

DISCUSSION

Our study represents a comprehensive description of *Leptopilina malgretoutensis*, the first recorded

parasitoid of *dunni* group *Drosophila* from their native range. By integrating classic taxonomy and

systematics approaches with genomics and experimental biology, we were able to generate a more

comprehensive view of the biology and evolution of this new species. Furthermore, historical records and

museum specimens enabled the discovery that asexually reproducing *Leptopilina malgretoutensis* have

also been found on another island in the Antilles. This work represents one of a limited number of studies

in which description of a new metazoan species is coupled with genome sequencing [69-73]. Indeed, this

is the fifth such example in insects [74-77], wherein body size often precludes whole genome sequencing

of holotypes.

Despite the interest in speciation patterns of the *dunni* group *Drosophila*, the proximal ecological or

evolutionary mechanisms driving morphological and species diversity in this group remain unresolved.

One study found elevated nonsynonymous substitution rates in the pigmentation gene *yellow* in two

members of this group [78]. This is consistent with selection for pigmentation diversity, but no studies

have yet identified the selective agent responsible for this pattern. Importantly, melanization is a process

in many insect lineages that is critical in both the development of cuticle pigmentation as well as in

immune responses that occur during encapsulation of foreign pathogens [79]. In *Drosophila*, melanization

is a well-studied response to infection by eukaryotic parasites (including *Leptopilina*) and is linked to the

same genetic pathways as pigmentation production [80-82]. In one study in *Drosophila falleni*, selection

for a reduced number of abdominal spots resulted in higher infection rates by a parasitic nematode [83].

In other insects there are similar links between the levels or patterns of cuticular melanization and

resistance to parasites [84-89]. As such, it has been postulated that *Drosophila* abdominal coloration may

vary in some systems primarily due to selection by parasitism [90, 91], but direct evidence for this is

lacking.

Since adult flies of the *D. antillea* species are among the more highly pigmented flies of the *dunni* group,

we were interested in the possibility that they might be particularly resistant to infection by endemic

parasitoids. However, we instead found the opposite. *L. malgretoutensis* wasps are highly virulent on *D.*

antillea hosts. This observation could indicate that *L. malgretoutensis* is a relatively recent colonizer of

Saint Lucia and that *D. antillea* has not had sufficient time to evolve resistance. In contrast, *L.*

malgretoutensis was less effective at using *D. willistoni* as a host. Our findings open the door to future

studies to explore the potential mechanism of *L. malgretoutensis* virulence on flies of the *dunni* group as

well as the potential mechanisms of resistance for *D. willistoni.* It will also be of interest to test the

virulence of *L. malgretoutensis* on other species from the *dunni* group as well as how virulence tracks with

island of origin of the wasp. For example, are *L. malgretoutensis* wasps as effective when infecting *dunni*

group flies that are native to other islands of the Antilles chain? Are *L. malgretoutensis* wasps that

originate from Guadeloupe as virulent on *D. antillea*? Indeed, the island of Guadeloupe where Carton

collected *L. malgretoutensis* is home to the dunni group fly *Drosophila arawakana.* It would be of interest

to know how the virulence of Guadeloupian wasps relates to Saint Lucian wasps on fly species from the

various islands.

The island biogeography of *L. malgretoutensis* may also play a role in the evolution of their reproductive

mode. Our antibiotic curing experiments indicated that our *L. malgretoutensis* colony derived from Saint

Lucia is unable to revert to sexual reproduction. The inability to restore sex is a strong indicator of a so-

called "virginity mutation", which is well described in several other wasp species with symbiont-mediated

parthenogenesis [58, 92-97]. There are multiple hypotheses for the evolutionary pressures that lead to

virginity mutations [49, 92, 98, 99], but in brief, they are mutations that result in the decay or loss of

sexual function, to the point that the animal becomes reliant upon the PI symbiont for the production of

female offspring. Related to this, it is notable that the colony of *L. malgretoutensis* derived from

Guadeloupe collections in the 1980s were all female. However, we do not know if this PB10-5 colony was

(a) infected with *Wolbachia*, (b) asexual because of *Wolbachia*, or (c) obligately asexual (i.e., unable to

revert to a sexual mode). Indeed, some other populations of parasitic wasps with PI *Wolbachia* have

significant within-species polymorphism and vary in which *Wolbachia* strain they have, if they are

asexually reproducing, and if they have a so-called virginity mutation [49, 100]. Thus, it may be that

populations of *L. malgretoutensis* on different islands vary in whether they are sexually reproducing, and if

asexual reproduction is fixed or reversible. Notably, asexual reproduction across many taxa appears to be

more common on islands [101, 102], and thus *L. malgretoutensis* in the Lesser Antilles may provide a

tractable system to understand the evolution of reproductive mode in the context of biogeography.

While there are many other examples of species with virginity mutations, our findings are quite unusual in

that the *Wolbachia* genome here also contains what appears to be functional genes for inducing

cytoplasmic incompatibility (CI). Paradoxically, the CI phenotype only manifests with sexual reproduction:

males with CI *Wolbachia*-modified sperm cause embryonic lethality in embryos derived from females

without a compatible *Wolbachia* strain [66]. We have several hypotheses for why these *Wolbachia*

encode for CI loci despite inducing parthenogenesis. First, the *cif* loci may have undergone some form of

neofunctionalization: perhaps these proteins are no longer involved in sperm-egg incompatibility as in CI

but are impacting other host processes. Alternatively, it might be that the transition to parthenogenesis is

more recent, and CI is no longer under selection. Perhaps the *cifs* are en route to be eliminated from the

genome in the future. "Recent" is of course difficult to determine quantitatively, especially in island

systems where there are often strong bottlenecks that can drive rapid evolution [103]. Finally, it might be

that if the PI phenotype has lower penetrance and occasional males are produced. These males might

mate with incompatible conspecifics (that either have no *Wolbachia* or an incompatible strain) and induce

CI, perhaps reducing competition for the asexual matriline. However, we see two issues with this

scenario. First, lower penetrance of PI phenotypes is typically due to the loss of *Wolbachia* (e.g., due to

higher temperatures, or in the lab, antibiotics), which leads to the production of males. In this case, the

occasional males produced by PI mothers might not have sufficient *Wolbachia* to induce CI. Second,

because the PI females transmitting *Wolbachia* to the next generation would never need to rescue CI,

there would be no selection pressure to maintain CI function.

Despite the peculiarity of a PI *Wolbachia* strain encoding for CI genes, all four loci associated with host

reproductive manipulation (*cifA, cifB,* and two copies of *pifB*) are found in the EAM of WO: a classic

signature of *Wolbachia* genomes and reproductive effector proteins. The multiple identical *pifB* copies,

genes for CI in combination with a clear *Wolbachia*-mediated PI phenotype, and evidence for multiple

prophage infections all point towards recent dynamism in the *w*Lmal genome. Many *Wolbachia* genes

mediating host reproductive switches are horizontally acquired from other *Wolbachia* strains, likely due to

the WO phage [64, 104]. For this reason, it was not surprising that the two PI *Wolbachia* strains from

Leptopilina both encoded for PifB proteins, despite the strains belonging to different clades. In contrast,

we did not find any evidence in *w*Lmal for the PifA protein, a *transformer* mimic that interacts with the

sexual differentiation pathway [60, 65]. Importantly however, we do not know the exact mechanism of

parthenogenesis induction employed by *w*Lmal (e.g., one versus two step, apomixis versus gamete

duplication, etc [105]). And, given the diversity of PI mechanisms found across *Wolbachia* [49] it is quite

possible that *w*Lcla and *w*Lmal are not using precisely the same toolset. Closer inspection of meiosis and

early embryonic mitoses in *L. malgretoutensis* will be key to understanding how *w*Lmal causes PI, and

what the role of PifB, or other proteins, might be in this process. Altogether, the description of *L.*

malgretoutensis and its reproductive biology will enable future explorations of host-parasitoid and host-

symbiont interactions, especially in the context of speciation.

In summary, this study provides a clear example of how descriptive work in under-studied species can

lead to novel discoveries across a broad set of biological disciplines. Insects generally, but especially

parasitic Hymenoptera, are under described [5]. Furthermore, of the described parasitic hymenopterans,

a significant majority lack biological information such as host associations and ecology. Here,

identification of a new species of parasitoid has provided a unique view into the intricate co-evolution of

host insect, parasitoid, and *Wolbachia* symbiont, with dynamic interactions between genomic evolution,

virulence, and reproductive ecology. We hope our work inspires more integration of descriptions of new

species coupled with investigations into novel aspects of their biology.

DATA AVAILABILITY

BioProject: PRJNA1150626

L. malgretoutensis BioSample: SAMN43291974

*w*Lmal BioSample: SAMN43291989

Oxford Nanopore Technologies Long Reads (all, including <5kbp): SRR30339708

WGS Illumina Reads: SRR30339709

RNA-seq Illumina Reads: SRR30339707

L. malgretoutensis Mitochondrial Genome: #Accession Forthcoming

*w*Lmal *Wolbachia* Genome: CP185405.1

L. malgretoutensis Nuclear Genome Assembly: This Whole Genome Shotgun project has been deposited

at DDBJ/ENA/GenBank under the accession JBMIRX000000000. The version described in this paper is

version JBMIRX010000000.

Supplemental File S1 contains:

Supplemental Text

Figure S1. Western Blot for *Wolbachia* detection.

Figure S2. Draft assembly snail plot and blob plot.

Supplemental Tables are provided in an excel file where each sheet corresponds to a different table

number:

Table S1. Passed reads statistics.

Table S2. All draft assembly contigs statistics, including BlobTools annotations.

Table S3. Purged contigs.

Table S4. Repeat annotations.

Table S5. *Wolbachia* protein queries.

Table S6. wLmal phage annotation.

ACKNOWLEDGEMENTS

The research expedition of the sailing vessel Sea Salt was conceived in conversations with W. Daniel

Tracey Sr. (now deceased), and we dedicate this study to his memory. We are very thankful to Amalia

Rowan the first mate on the Sea Salt without whose assistance our collection expedition could not have

taken place. We are thankful for Dr. Hannah Romain (Saint Lucia Ministry of Agriculture) for granting

permission to collect insects on Saint Lucia. We are also thankful to Michael Gerth for making the WHOP

pipeline available and to Keith Hopper for sharing the *Ganaspis* assembly prior to its release. Research

reported in this publication was supported by the National Institute of General Medical Sciences of the

National Institutes of Health under award number R35GM150991 to ARIL, the National Institute of Food

and Agriculture Predoctoral Fellowship to LCF (NIFA 2023-67034-40496), and the Linda and Jack Gill

Institute for Neuroscience Research Stipend to WDT. Mention of trade names or commercial products in

this publication is solely for the purpose of providing specific information and does not imply

recommendation or endorsement by the USDA. USDA is an equal opportunity provider and employer.

CONFLICTS OF INTEREST

MKD works for Oxford Nanopore Technologies.

CITATIONS

1.   Karvonen A, Seehausen O. The role of parasitism in adaptive radiations—when might parasites
promote and when might they constrain ecological speciation? International Journal of Ecology.
2012;2012(1):280169.

2.   Nyman T, Bokma F, Kopelke J-P. Reciprocal diversification in a complex plant-herbivore-
parasitoid food web. BMC Biol. 2007;5:1-9.

3.   Lue CH, Buffington ML, Scheffer S, Lewis M, Elliott TA, Lindsey AR, et al. DROP: Molecular
voucher database for identification of *Drosophila* parasitoids. Molecular Ecology Resources.
2021;21(7):2437-54.

4.   Saunders TE, Ward DF. Variation in the diversity and richness of parasitoid wasps based on
sampling effort. PeerJ. 2018;6:e4642.

5.   Forbes AA, Bagley RK, Beer MA, Hippee AC, Widmayer HA. Quantifying the unquantifiable: why
Hymenoptera, not Coleoptera, is the most speciose animal order. BMC Ecol. 2018;18:1-11.

6.   Sosa-Calvo J, Forshage M, Buffington ML. Circumscription of the *Ganaspis brasiliensis* (Ihering,
1905) species complex (Hymenoptera, Figitidae), and the description of two new species parasitizing the
spotted wing drosophila, *Drosophila suzukii* Matsumura, 1931 (Diptera, Drosophilidae). J Hymenoptera
Res. 2024;97:441-70.

7.   Coyne JA, Orr HA. Speciation: a catalogue and critique of species concepts. Philosophy of
Biology: An Anthology. 2004:272-92.

8.   Carson HL, Templeton AR. Genetic revolutions in relation to speciation phenomena: the founding
of new populations. Annu Rev Ecol Syst. 1984:97-131.

9.   Kaneshiro K, Gillespie R, Carson H. Chromosomes and male genitalia of Hawaiian *Drosophila*:
tools for interpreting phylogeny and geography. Hawaiian Biogeography: Evolution on a hot spot
archipelago. 1995:57-71.

10.   O'Grady P, DeSalle R. Hawaiian *Drosophila* as an evolutionary model clade: days of future past.
Bioessays. 2018;40(5):1700246.

11.   Dworkin I, Jones CD. Genetic changes accompanying the evolution of host specialization in
Drosophila sechellia. Genetics. 2009;181(2):721-36.

12.   Hollocher H. Island hopping in *Drosophila*: patterns and processes. Philosophical Transactions of
the Royal Society of London Series B: Biological Sciences. 1996;351(1341):735-43.

13.   Heed WB, Krishnamurthy NB. Genetic studies on the cardini group of *Drosophila* in the West
Indies. Univ Texas Publ. 1959;5914:155-79.

14.   Hollocher H, Hatcher JL, Dyreson EG. Genetic and developmental analysis of abdominal
pigmentation differences across species in the *Drosophila dunni* subgroup. Evolution. 2000;54(6):2057-
71.

15.   Lue C-H, Driskell AC, Leips J, Buffington ML. Review of the genus *Leptopilina* (Hymenoptera,
Cynipoidea, Figitidae, Eucoilinae) from the Eastern United States, including three newly described
species. J Hymenoptera Res. 2016;(53).

16.   Stacconi MVR, Buffington M, Daane KM, Dalton DT, Grassi A, Kaçar G, et al. Host stage
preference, efficacy and fecundity of parasitoids attacking *Drosophila suzukii* in newly invaded areas. Biol
Control. 2015;84:28-35.

17.   Abram PK, Franklin MT, Hueppelsheuser T, Carrillo J, Grove E, Eraso P, et al. Adventive larval
parasitoids reconstruct their close association with spotted-wing drosophila in the invaded North
American range. Environ Entomol. 2022;51(4):670-8.

18.   Newton IL, Savytskyy O, Sheehan KB. *Wolbachia* utilize host actin for efficient maternal
transmission in *Drosophila melanogaster*. PLoS Path. 2015;11(4):e1004798. doi:
10.1371/journal.ppat.1004798. PubMed PMID: 25906062; PubMed Central PMCID: PMCPMC4408098.

19.   Lindsey AR, Tennessen JM, Gelaw MA, Jones MW, Parish AJ, Newton IL, et al. The intracellular
symbiont *Wolbachia* alters *Drosophila* development and metabolism to buffer against nutritional stress.
bioRxiv. 2023.

20.   Fricke LC, Lindsey AR. Examining *Wolbachia*-Induced Parthenogenesis in Hymenoptera.
Wolbachia: Methods and Protocols: Springer; 2024. p. 55-68.

21.   Shen W, Le S, Li Y, Hu F. SeqKit: a cross-platform and ultrafast toolkit for FASTA/Q file
manipulation. PLoS One. 2016;11(10):e0163962.

22.   Kolmogorov M, Yuan J, Lin Y, Pevzner PA. Assembly of long, error-prone reads using repeat
graphs. Nat Biotechnol. 2019;37(5):540-6.

23.   Roach MJ, Schmidt SA, Borneman AR. Purge Haplotigs: allelic contig reassignment for third-gen
diploid genome assemblies. BMC Bioinformatics. 2018;19(1):1-10.

24.   Li H. Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM. arXiv
preprint arXiv:13033997. 2013.

25.   Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, et al. The Sequence Alignment/Map
format and SAMtools. Bioinformatics. 2009;25(16):2078-9. doi: 10.1093/bioinformatics/btp352.

26.   Challis R, Richards E, Rajan J, Cochrane G, Blaxter M. BlobToolKit–interactive quality
assessment of genome assemblies. G3: Genes, Genomes, Genetics. 2020;10(4):1361-74. PubMed
Central PMCID: PMCPMC7144090.

27.   Simão FA, Waterhouse RM, Ioannidis P, Kriventseva EV, Zdobnov EM. BUSCO: Assessing
genome assembly and annotation completeness with single-copy orthologs. Bioinformatics. 2015. doi:
10.1093/bioinformatics/btv351.

28.   Astashyn A, Tvedte ES, Sweeney D, Sapojnikov V, Bouk N, Joukov V, et al. Rapid and sensitive
detection of genome contamination at scale with FCS-GX. Genome Biol. 2024;25(1):60.

29.   Bernt M, Donath A, Jühling F, Externbrink F, Florentz C, Fritzsch G, et al. MITOS: improved de
novo metazoan mitochondrial genome annotation. Mol Phylogen Evol. 2013;69(2):313-9.

30.   Wanner N, Larsen PA, McLain A, Faulk C. The mitochondrial genome and Epigenome of the
Golden lion Tamarin from fecal DNA using Nanopore adaptive sequencing. BMC Genomics. 2021;22:1-
11.

31.   Katoh K, Standley DM. MAFFT multiple sequence alignment software version 7: improvements in
performance and usability. Mol Biol Evol. 2013;30(4):772-80. doi: 10.1093/molbev/mst010.

32.   Nguyen L-T, Schmidt HA, Von Haeseler A, Minh BQ. IQ-TREE: a fast and effective stochastic
algorithm for estimating maximum-likelihood phylogenies. Mol Biol Evol. 2015;32(1):268-74.

33.   Flynn JM, Hubley R, Goubert C, Rosen J, Clark AG, Feschotte C, et al. RepeatModeler2 for
automated genomic discovery of transposable element families. Proc Natl Acad Sci. 2020;117(17):9451-
7.

34.   Storer J, Hubley R, Rosen J, Wheeler TJ, Smit AF. The Dfam community resource of
transposable element families, sequence models, and genome annotations. Mobile DNA. 2021;12:1-14.

35.   Keilwagen J, Hartung F, Grau J. GeMoMa: homology-based gene prediction utilizing intron
position conservation and RNA-seq data. Gene Prediction: Methods and Protocols. 2019:161-77.

36.   Dobin A, Gingeras TR. Mapping RNA-seq reads with STAR. Current Protocols in Bioinformatics.
2015;51(1):11.4. 1-.4. 9.

37.   Emms DM, Kelly S. OrthoFinder: phylogenetic orthology inference for comparative genomics.
Genome Biol. 2019;20:1-14.

38.   Gerth M. WHOP (v0.1). Zenodo. 2023. doi: https://doi.org/10.5281/zenodo.8343898.

39.   Seemann T. Prokka: rapid prokaryotic genome annotation. Bioinformatics. 2014;30(14):2068-9.

40.   Bruen T, Bruen T. PhiPack: PHI test and other tests of recombination. McGill University,
Montreal, Quebec. 2005:1-8.

41.   Grant JR, Enns E, Marinier E, Mandal A, Herman EK, Chen C-y, et al. Proksee: in-depth
characterization and visualization of bacterial genomes. Nucleic Acids Res. 2023;51(W1):W484-W92.

42.   Guo J, Bolduc B, Zayed AA, Varsani A, Dominguez-Huerta G, Delmont TO, et al. VirSorter2: a
multi-classifier, expert-guided approach to detect diverse DNA and RNA viruses. Microbiome. 2021;9:1-
13.

43.   Starikova EV, Tikhonova PO, Prianichnikov NA, Rands CM, Zdobnov EM, Ilina EN, et al. Phigaro:
high-throughput prophage sequence annotation. Bioinformatics. 2020;36(12):3882-4.

44.   Brown CL, Mullet J, Hindi F, Stoll JE, Gupta S, Choi M, et al. mobileOG-db: a manually curated
database of protein families mediating the life cycle of bacterial mobile genetic elements. Appl Environ
Microbiol. 2022;88(18):e00991-22.

45.   R Core Team. R: A language and environment for statistical computing. R Foundation for
Statistical Computing, Vienna, Austria: URL http://www.R-project.org/; 2014.

46.   Frerebeau N, Lebrun B, Arel-Bundock V. khroma: Colour Schemes for Scientific Data
Visualization. 2019.

47.   Wachi N, Nomano FY, Mitsui H, Kasuya N, Kimura MT. Taxonomy and evolution of putative
thelytokous species of *Leptopilina* (Hymenoptera: Figitidae) from Japan, with description of two new
species. Entomol Sci. 2015;18(1):41-54.

48.   Chen F, Schenkel M, Geuverink E, van de Zande L, Beukeboom LW. Absence of complementary
sex determination in two *Leptopilina* species (Figitidae, Hymenoptera) and a reconsideration of its
incompatibility with endosymbiont-induced thelytoky. Insect Sci. 2022;29(3):900-14.

49.   Ma WJ, Schwander T. Patterns and mechanisms in instances of endosymbiont-induced
parthenogenesis. J Evol Biol. 2017;30(5):868-88.

50.   Jervis MA, Heimpel GE, Ferns PN, Harvey JA, Kidd NA. Life-history strategies in parasitoid
wasps: a comparative analysis of 'ovigeny'. J Anim Ecol. 2001;70(3):442-58.

51.   Doremus MR, Stouthamer CM, Kelly SE, Schmitz-Esser S, Hunter MS. Quality over quantity:
unraveling the contributions to cytoplasmic incompatibility caused by two coinfecting *Cardinium*
symbionts. Heredity. 2022;128(3):187-95.

52.   Duron O, Bouchon D, Boutin S, Bellamy L, Zhou L, Engelstädter J, et al. The diversity of
reproductive parasites among arthropods: *Wolbachia* do not walk alone. BMC Biol. 2008;6(1):1-12.

53.   Kageyama D, Narita S, Watanabe M. Insect sex determination manipulated by their
endosymbionts: incidences, mechanisms and implications. Insects. 2012;3(1):161-99.

54.   Perlmutter JI, Bordenstein SR. Microorganisms in the reproductive tissues of arthropods. Nat Rev
Micro. 2020;18(2):97-111.

55.   Hopper KR, Wang X, Kenis M, Seehausen ML, Abram PK, Daane KM, et al. Genome divergence
and reproductive incompatibility among populations of *Ganaspis* near *brasiliensis*. G3: Genes, Genomes,
Genetics. 2024:jkae090.

56.   Cameron SL. Insect mitochondrial genomics: implications for evolution and phylogeny. Annu Rev
Entomol. 2014;59(1):95-117.

57.   Khan S, Sowpati DT, Srinivasan A, Soujanya M, Mishra RK. Long-read genome sequencing and
assembly of *Leptopilina boulardi*: a specialist *Drosophila* parasitoid. G3: Genes, Genomes, Genetics.
2020;10(5):1485-94.

58.   Kraaijeveld K, Anvar SY, Frank J, Schmitz A, Bast J, Wilbrandt J, et al. Decay of sexual trait
genes in an asexual parasitoid wasp. Genome Biol Evol. 2016;8(12):3685-95.

59.   Wey B, Heavner ME, Wittmeyer KT, Briese T, Hopper KR, Govind S. Immune suppressive
extracellular vesicle proteins of *Leptopilina heterotoma* are encoded in the wasp genome. G3: Genes,
Genomes, Genetics. 2020;10(1):1-12.

60.   Fricke LC, Lindsey AR. Identification of parthenogenesis-inducing effector proteins in *Wolbachia*.
Genome Biol Evol. 2024:evae036.

61.   Lindsey ARI, Werren JH, Richards S, Stouthamer R. Comparative genomics of a
parthenogenesis-inducing *Wolbachia* symbiont. G3: Genes|Genomes|Genetics. 2016;6(7):2113-23. Epub
2016/05/20. doi: 10.1534/g3.116.028449. PubMed PMID: 27194801; PubMed Central PMCID:
PMCPMC4938664.

62.   Kaur R, Shropshire JD, Cross KL, Leigh B, Mansueto AJ, Stewart V, et al. Living in the
endosymbiotic world of *Wolbachia*: A centennial review. Cell Host & Microbe. 2021;29(6):879-93.
PubMed Central PMCID: PMC8192442.

63.   Lindsey ARI. Sensing, Signaling, and Secretion: A review and analysis of systems for regulating
host interaction in *Wolbachia*. Genes. 2020;11(7):813. PubMed Central PMCID: PMC7397232.

64.   Bordenstein SR, Bordenstein SR. Eukaryotic association module in phage WO genomes from
Wolbachia. Nat Commun. 2016;7.

65.   Li C, Li C-Q, Chen Z-B, Liu B-Q, Sun X, Wei K-H, et al. *Wolbachia* symbionts control sex in a
parasitoid wasp using a horizontally acquired gene. Curr Biol. 2024.

66.   Beckmann JF, Bonneau M, Chen H, Hochstrasser M, Poinsot D, Merçot H, et al. The toxin–
antidote model of cytoplasmic incompatibility: genetics and evolutionary implications. Trends Genet.
2019;35(3):175-85. PubMed Central PMCID: PMC6519454

67.   Schilthuizen M, Nordlander G, Stouthamer R, van Alphen J. Morphological and molecular
phylogenetics in the genus *Leptopilina* (Hymenoptera: Cynipoidea: Eucoilidae). Syst Entomol.
1998;23(3):253-64.

68.   Vet LE, van Alphen JJ. A comparative functional approach to the host detection behaviour of
parasitic wasps. 1. A qualitative study on Eucoilidae and Alysiinae. Oikos. 1985:478-86.

69.   Kanzaki N, Tsai IJ, Tanaka R, Hunt VL, Liu D, Tsuyama K, et al. Biology and genome of a newly
discovered sibling species of *Caenorhabditis elegans*. Nat Commun. 2018;9(1):3216.

70.   Koehler G, Khaing KPP, Than NL, Baranski D, Schell T, Greve C, et al. A new genus and species
of mud snake from Myanmar (Reptilia, Squamata, Homalopsidae). Zootaxa. 2021;4915(3):zootaxa.
4915.3. 1-zootaxa. .3. 1.

71.   Köhler G, Vargas J, Than NL, Schell T, Janke A, Pauls SU, et al. A taxonomic revision of the
genus Phrynoglossus in Indochina with the description of a new species and comments on the
classification within Occidozyginae (Amphibia, Anura, Dicroglossidae). Vertebrate Zoology. 2021;71:1-26.

72.   Köhler G, Zwitzers B, Than NL, Gupta DK, Janke A, Pauls SU, et al. Bioacoustics reveal hidden
diversity in frogs: Two new species of the genus Limnonectes from Myanmar (Amphibia, Anura,
Dicroglossidae). Diversity. 2021;13(9):399.

73.   Sullivan JP, Hopkins CD, Pirro S, Peterson R, Chakona A, Mutizwa TI, et al. Mitogenome
recovered from a 19th Century holotype by shotgun sequencing supplies a generic name for an orphaned
clade of African weakly electric fishes (Osteoglossomorpha, Mormyridae). ZooKeys. 2022;1129:163.

74.   Brandão-Dias PF, Zhang YM, Pirro S, Vinson CC, Weinersmith KL, Ward AK, et al. Describing
biodiversity in the genomics era: A new species of Nearctic Cynipidae gall wasp and its genome. Syst
Entomol. 2022;47(1):94-112.

75.   Heckenhauer J, Razuri-Gonzales E, Mwangi FN, Schneider J, Pauls SU. Holotype sequencing of
Silvatares holzenthali Rázuri-Gonzales, Ngera & Pauls, 2022 (Trichoptera, Pisuliidae). ZooKeys.
2023;1159:1.

76.   Pohl H, Niehuis O, Gloyna K, Misof B, Beutel RG. A new species of *Mengenilla* (Insecta,
Strepsiptera) from Tunisia. ZooKeys. 2012;(198):79.

77.   Rázuri-Gonzales E, Graf W, Heckenhauer J, Schneider JV, Pauls SU.   A new species of
Rhyacophila Pictet, 1834 (Trichoptera, Rhyacophilidae) from Corsica with the genomic characterization of
the holotype. ZooKeys. 2024;1218:295.

78.   Wilder J, Dyreson E, O'Neill R, Spangler M, Gupta R, Wilder A, et al. Contrasting modes of
natural selection acting on pigmentation genes in the *Drosophila dunni* subgroup. Journal of Experimental
Zoology Part B: Molecular and Developmental Evolution. 2004;302(5):469-82.

79.   Sugumaran M, Barek H. Critical analysis of the melanogenic pathway in insects and higher
animals. International Journal of Molecular Sciences. 2016;17(10):1753.

80.   Dolezal T. How to eliminate pathogen without killing oneself? Immunometabolism of
encapsulation and melanization in *Drosophila*. Frontiers in Immunology. 2023;14:1330312.

81.   Carton Y, Poirié M, Nappi AJ. Insect immune resistance to parasitoids. Insect Sci. 2008;15(1):67-
87.

82.   Russo J, Dupas S, Frey F, Carton Y, Brehelin M. Insect immunity: early events in the
encapsulation process of parasitoid (*Leptopilina boulardi*) eggs in resistant and susceptible strains of
Drosophila. Parasitology. 1996;112(1):135-42.

83.   Dombeck I, Jaenike J. Ecological genetics of abdominal pigmentation in *Drosophila falleni*: a
pleiotropic link to nematode parasitism. Evolution. 2004;58(3):587-96.

84.   Siva–Jothy MT. A mechanistic link between parasite resistance and expression of a sexually
selected trait in a damselfly. Proceedings of the Royal Society of London Series B: Biological Sciences.
2000;267(1461):2523-7.

85.   Wilson K, Cotter SC, Reeson AF, Pell JK. Melanism and disease resistance in insects. Ecol Lett.
2001;4(6):637-49.

86.   Tan S, Wang Y, Liu P, Ge Y, Li A, Xing Y, et al. Increase of albinistic hosts caused by gut
parasites promotes self-transmission. Frontiers in Microbiology. 2018;9:1525.

87.   Lindsey E, Altizer S. Sex differences in immune defenses and response to parasitism in monarch
butterflies. Evol Ecol. 2009;23:607-20.

88.   Krams I, Burghardt GM, Krams R, Trakimas G, Kaasik A, Luoto S, et al. A dark cuticle allows
higher investment in immunity, longevity and fecundity in a beetle upon a simulated parasite attack.
Oecologia. 2016;182:99-109.

89.   Dubovskiy I, Whitten M, Kryukov V, Yaroslavtseva O, Grizanova E, Greig C, et al. More than a
colour change: insect melanism, disease resistance and fecundity. Proc R Soc Lond B.
2013;280(1763):20130584.

90.   Wittkopp PJ, Beldade P, editors. Development and evolution of insect pigmentation: genetic
mechanisms and the potential consequences of pleiotropy. Semin Cell Dev Biol; 2009: Elsevier.

91.   Kohler LJ, Carton Y, Mastore M, Nappi AJ. Parasite suppression of the oxidations of eumelanin
precursors in *Drosophila melanogaster*. Archives of Insect Biochemistry and Physiology: Published in
Collaboration with the Entomological Society of America. 2007;66(2):64-75.

92.   Stouthamer R, Russell JE, Vavre F, Nunney L. Intragenomic conflict in populations infected by
Parthenogenesis Inducing *Wolbachia* ends with irreversible loss of sexual reproduction. BMC Evol Biol.
2010;10:12. doi: 10.1186/1471-2148-10-229. PubMed PMID: WOS:000282746700002.

93.   Ma W-J, Pannebakker BA, Li X, Geuverink E, Anvar SY, Veltsos P, et al. A single QTL with large
effect is associated with female functional virginity in an asexual parasitoid wasp. Mol Ecol.
2021;30(9):1979-92. doi: https://doi.org/10.1111/mec.15863.

94.   Ma WJ, Pannebakker BA, Beukeboom LW, Schwander T, van de Zande L. Genetics of decayed
sexual traits in a parasitoid wasp with endosymbiont-induced asexuality. Heredity. 2014;113(5):424-31.
doi: 10.1038/hdy.2014.43. PubMed PMID: 24781809; PubMed Central PMCID: PMC4220718.

95.   Russell JE, Stouthamer R. The genetics and evolution of obligate reproductive parasitism in
Trichogramma pretiosum infected with parthenogenesis-inducing *Wolbachia*. Heredity. 2011;106(1):58-
67. doi: 10.1038/hdy.2010.48. PubMed PMID: WOS:000285336100007.

96.   Gottlieb Y, Zchori-Fein E. Irreversible thelytokous reproduction in *Muscidifurax uniraptor*. Entomol
Exp Appl. 2001;100(3):271-8.

97.   Stouthamer R, Mak F. Influence of antibiotics on the offspring production of the *Wolbachia*-
infected parthenogenetic parasitoid *Encarsia formosa*. J Invertebr Pathol. 2002;80(1):41-5.

98.   van der Kooi CJ, Schwander T. On the fate of sexual traits under asexuality. Biol Rev Camb
Philos Soc. 2014;89(4):805-19. doi: 10.1111/brv.12078. PubMed PMID: 24443922.

99.   Neiman M, Sharbel TF, Schwander T. Genetic causes of transitions from sexual reproduction to
asexuality in plants and animals. J Evol Biol. 2014;27(7):1346-59. doi: 10.1111/jeb.12357. PubMed PMID:
24666600.

100.   van der Kooi CJ, Matthey-Doret C, Schwander T. Evolution and comparative ecology of
parthenogenesis in haplodiploid arthropods. Evolution Letters. 2017;1(6):304-16.

101.   Vrijenhoek RC, Parker ED. Geographical parthenogenesis: general purpose genotypes and
frozen niche variation. Lost sex: the evolutionary biology of parthenogenesis. 2009:99-131.

102.   Hörandl E. Geographical parthenogenesis: opportunities for asexuality. Lost sex: The
evolutionary biology of parthenogenesis. 2009:161-86.

103.   Gillespie RG. Island time and the interplay between ecology and evolution in species
diversification. Evolutionary Applications. 2016;9(1):53-73.

104.   Lindsey ARI, Rice DW, Bordenstein SR, Brooks AW, Bordenstein SR, Newton ILG. Evolutionary
genetics of cytoplasmic incompatibility genes *cifA* and *cifB* in prophage WO of *Wolbachia*. Genome Biol
Evol. 2018;10(2):434-51. doi: 10.1093/gbe/evy012. PubMed Central PMCID: PMCPMC5793819.

105.   Verhulst EC, Pannebakker BA, Geuverink E. Variation in sex determination mechanisms may
constrain parthenogenesis-induction by endosymbionts in haplodiploid systems. Current Opinion in Insect
Science. 2023:101023.
1177