Large adipocytes in a biomimetic adipose tissue model promote breast cancer malignancy

Garrett F Beeghly[1], Marlee I Pincus[1], Rohan R Varshney[2], Dilip D Giri[3], Domenick J Falcone[4], Michael C Rudolph[2], Marc A Antonyak[5], Neil M Iyengar[6,7], and Claudia Fischbach[1,8]*

1. Nancy E. and Peter C. Meinig School of Biomedical Engineering, Cornell University, Ithaca, NY, USA
2. Harold Hamm Diabetes Center and Department of Physiology, University of Oklahoma Health Sciences Center, Oklahoma City, OK, USA
3. Department of Pathology, Memorial Sloan Kettering Cancer Center, New York, NY, USA
4. Department of Pathology and Laboratory Medicine, Weill Cornell Medical College, New York, NY, USA
5. Department of Molecular Medicine, Cornell University, Ithaca, NY, USA
6. Department of Medicine, Memorial Sloan Kettering Cancer Center, New York, NY, USA
7. Department of Medicine, Weill Cornell Medical College, New York, NY, USA
8. Kavli Institute at Cornell for Nanoscale Science, Cornell University, Ithaca, NY, USA

*Correspondence: cf99@cornell.edu

Abstract

Obesity worsens breast cancer survival but how increased adipocyte size, which typifies obese adipose tissue and correlates with poor prognosis, impacts cancer progression remains unclear. Understanding these connections is challenging as adipocyte size is highly heterogeneous and not tunable with traditional experimental approaches. Here, we develop a biomimetic, engineered adipose tissue model to culture size-sorted adipocytes isolated from the same donor and assess their impact on tumor cell behavior. We find that large adipocytes are transcriptionally distinct from small adipocytes and upregulate genes related to adipose tissue dysfunction, including altered lipid processing. In coculture, large adipocytes promote lipid accumulation in cancer cells, increasing their migration and proliferation via enhanced fatty acid oxidation. These changes align with greater extracellular vesicle release by large adipocytes, which transfer lipid to recipient tumor cells. Our findings posit adipocyte size as an independent prognostic biomarker for breast cancer patients and demonstrate the value of biomimetic tissue models for mechanistic studies.

Introduction

Over 40% of adult women in the U.S. are obese (*1*) and obesity is an established prognostic factor associated with poor clinical outcomes across all breast cancer subtypes (*2*). Obesity is a complex disease with multivariate pathophysiology, including systemic metabolic dysfunction and local remodeling of white adipose tissue (WAT), which forms a primary component of the breast stroma (*3*). Recent studies indicate that adipocytes, which are the functional cell type of WAT, are more plastic (*4*, *5*) and heterogenous (*6*, *7*) than once considered and comprise distinct subpopulations which change in obesity. In breast cancer, adipocytes release pro-inflammatory cytokines which promote tumor cell migration (*8*), extracellular vesicles which enhance tumor cell proliferation (*9*), and fatty acids (FAs) which reprogram tumor cell metabolism (*10*). Many of these phenomena are exacerbated in obesity, where adipocyte function is impaired (*11*). Nevertheless, our mechanistic understanding of how obesity impacts breast cancer prognosis remains limited.

Large or hypertrophic adipocytes are one subpopulation of interest, as increased adipocyte size correlates with the transition from pre-malignant to invasive breast cancer regardless of body mass index (BMI) (*12*). However, WAT expands via a combination of hyperplasia (increase in adipocyte number) and hypertrophy (increase in adipocyte size) (*13*), leading to the presence of both small and large cells within obese tissue. Likewise, large adipocytes exist outside of obesity and are especially prevalent in lean individuals with metabolic disorders (*14*, *15*). How small and large adipocytes differ in function, and which effect these

49 changes have on breast cancer cells, remains largely unclear due to insufficient strategies to study adipocyte
50 subpopulations of interest in a tractable manner.
51
Most studies of adipocyte function extrapolate differences by comparing lean and obese animals, which
contain cells of varied size. This approach also introduces additional variables (e.g., diet, genetic differences)
making it difficult to attribute observations to adipocyte size alone. *In vitro* studies could address these
limitations but rely on progenitor cells differentiated on two-dimensional substrates, which are smaller in size
and functionally distinct compared to primary adipocytes (*16*). However, studies with primary adipocytes are
particularly challenging due to the distinct properties of these cells. Primary adipocytes are large (often over
100 µm in diameter), buoyant in aqueous media, post-mitotic, and fragile, which renders them incompatible
with many culture platforms (*17*) and single-cell technologies (*6*). Thus, improving our understanding of how
adipocyte size contributes to cancer progression will require new experimental approaches that enable
comparisons of differently sized primary adipocytes isolated from the same donor tissue.
62
Here, we develop a biomimetic, engineered adipose tissue platform to decouple adipocyte size from other
variables and overcome the limitations of traditional animal models and two-dimensional culture systems.
We first establish our ability to sort primary adipocytes from the same donor tissue into large and small cell
populations across both diet-induced and genetic mouse models of obesity. Using RNA sequencing and
targeted lipidomics, we characterize inherent phenotypic differences between large and small adipocytes
while controlling for other obesity-associated effects. Subsequently, we leverage our engineered tissue model
(*18*) to maintain these adipocytes in coculture with breast cancer cells *ex vivo*. We monitor tumor cell lipid
accumulation, migration, proliferation, and oxidative metabolism as a function of coculture with different
adipocyte subpopulations and assess potential means of intercellular communication which mediated our
observations. We then validate the relevance of our findings with a cohort of human mastectomy patients
(*19*) for which banked histology, paired RNA sequencing, and pre-operative blood panels are available.
Collectively, our results suggest that adipocyte size independently impacts adipocyte phenotype and, in turn,
influences breast cancer progression by altering tumor cell behavior. These findings highlight the potential
value of mammary adipocyte size as a prognostic biomarker and point to the utility of biomimetic tissue
models to provide mechanistic insights not possible with conventional approaches.
78
Results
Hypertrophic adipocytes can be isolated by differential buoyancy and size exclusion.
Although adipocyte hypertrophy is a hallmark of obesity, obese WAT contains both small and large adipocytes
as it expands through a combination of hyperplasia and hypertrophy (**Fig. 1A**). *In vitro* studies fail to resolve
size-dependent differences in adipocyte function due to a reliance on differentiated progenitor cells (**Fig. 1B**),
which are multilocular, smaller in size, and functionally distinct compared to primary adipocytes (*16*). To
address these challenges, we developed experimental strategies to isolate the effect of cell size on adipocyte
function while accounting for other covariates associated with weight gain. First, we assessed the size
distributions of adipocytes in both diet-induced and genetic mouse models of obesity and their respective
lean controls. For diet-induced studies, female C57BL/6 mice were fed a high-fat diet (HFD) or low-fat diet
(LFD) for 12 weeks starting at 8 weeks of age (**Fig. 1C**). For genetic studies, we used female B6.Cg-Lep[Ob]
mice (*ob/ob*) which become morbidly obese due to a loss-of-function mutation in the satiety hormone leptin
compared to wild-type (WT) mice (*20*) (**Fig. 1C**). Consistent with the development of obesity, HFD and *ob/ob*
mice weighed significantly more compared to LFD and WT mice at the time of adipocyte isolation (**Fig. S1**).
As expected, adipocytes in the mammary fat pads of both HFD and *ob/ob* mice were larger relative to
adipocytes in their respective lean controls (**Fig. 1D-E**). Of note, we found that adipocyte hypertrophy was
more pronounced in *ob/ob* mice than HFD mice, likely due to the impaired capacity of *ob/ob* progenitor cells
to undergo adipogenesis (*21*). However, both HFD and *ob/ob* mice contained a broad distribution of cell
sizes, including adipocytes similar in size to those found in LFD and WT mice in addition to hypertrophic cells

characteristic of obesity. We recognized this inherent heterogeneity as an opportunity to examine the impact
of cell size on adipocyte function while controlling for other systemic effects of obesity.

Figure 1. Hypertrophic adipocytes can be isolated by differential buoyancy and size exclusion. (**A**) Schematic of hyperplastic versus hypertrophic adipose tissue expansion. (**B**) Representative images of *in vitro* differentiated and primary adipocytes stained for nuclei (cyan) and neutral lipid (gray). Scale bars = 100 μm. (**C**) Schematic of diet-induced and genetic mouse models of obesity used to isolate primary adipocytes. (**D**) Hematoxylin- and eosin-stained sections of the anterior and inguinal fat pads surrounding the mammary glands of lean and obese mice. Scale bars = 100 μm. (**E**) Quantification of adipocyte diameters estimated from histology. Violin plots show the distribution of all quantified diameters and circles show the average adipocyte diameter per mouse (n = 5). (**F**) Schematic of protocol to separate obese adipocytes by size. (**G**) Representative images of size-sorted adipocytes stained for neutral lipid (gray). Scale bars = 100 μm. (**H**) Quantification of sorted adipocyte diameters. Violin plots show the distribution of all quantified diameters and circles show the average adipocyte diameter per mouse (n = 3–4). Dashed line indicates the average lipid droplet diameter of *in vitro* differentiated adipocytes after two weeks. (**I**) Cell viability of lean and sorted obese adipocytes determined via propidium iodide exclusion (n = 3–4). All data are presented as mean ± standard deviation unless otherwise noted. Statistics were performed with a nested one-way analysis of variance (ANOVA) with Tukey's multiple comparisons test. Compact letter display indicates statistical significance (p = 0.05).

To assess the impact of hypertrophy on adipocyte function independently of other obesity-associated
changes, we separated adipocytes isolated from HFD and *ob/ob* mice into large and small cells by leveraging
size-dependent differences in buoyancy (*22*) (**Fig. 1F**). Notably, this approach enabled us to obtain small
adipocytes from HFD and *ob/ob* mice that were comparable in diameter to adipocytes isolated from LFD and
WT mice, respectively (**Fig. 1G-H**). Consistent with histology, large cells from *ob/ob* mice were greater in size
than large cells from HFD mice. We also confirmed that size sorting adipocytes from HFD and *ob/ob* mice

did not negatively impact cell viability compared to unsorted adipocytes from LFD and WT mice (**Fig. 1I**).
Indeed, we maintained viability around 80% for all primary adipocyte subpopulations, consistent with
previous results from our lab and others (*16, 18*). Collectively, these findings confirm that we can reliably
isolate small and large adipocytes from the mammary fat pads across two mouse models of obesity, enabling
us to analyze the effect of cell size on adipocyte function independently of other factors.
112
Large and small adipocytes are transcriptionally distinct.
To broadly assess how size affects adipocyte phenotype, we first performed RNA sequencing on size-sorted
adipocytes from HFD and *ob/ob* mice and unsorted adipocytes from LFD and WT mice. Principal component
analysis (**Fig. 2A**) and hierarchical clustering (**Fig. 2B**) grouped samples based on condition. For both models
of obesity, small adipocytes clustered in between large adipocytes and adipocytes from their respective lean
controls, which indicates adipocyte dysfunction in obesity may exist along a continuum. Interestingly, small
HFD adipocytes clustered more closely to LFD and WT adipocytes than large HFD adipocytes, suggesting
that changes to cell size rather than global differences due to diet drive the transcriptional response of
adipocytes to obesity. Of note, adipocytes isolated from *ob/ob* mice clustered further away from adipocytes
isolated in all other conditions likely due to transcriptional differences driven by global leptin deficiency.
Indeed, this result is consistent with sequencing of unsorted adipocytes, which demonstrate a greater number
of differentially expressed genes between *ob/ob* and WT adipocytes than HFD and LFD adipocytes (**Fig.**
S2). When comparing large and small adipocytes from HFD and *ob/ob* mice, we found a hypertrophy-
associated gene signature of 821 differentially expressed genes that were shared across both models of
obesity (**Fig. 2C**). These results suggest that altered adipocyte size causes distinct transcriptional changes
that are independent of how obesity develops (e.g., differences in diet composition for LFD versus HFD mice
or genetic differences and quantity of diet consumed for WT versus *ob/ob* mice).
130
Next, we assessed the functional states associated with adipocyte hypertrophy for both obesity models using
gene set enrichment analysis (GSEA). Large adipocytes from both HFD and *ob/ob* mice were enriched for
gene sets related to adipose tissue dysfunction in obesity (*13*) including altered collagen deposition, lipid
processing, pro-inflammatory signaling, response to hypoxia, and angiogenic signaling (**Fig. 2D**).
Interestingly, we also found that large adipocytes were enriched for gene sets related to cytoskeleton
organization, calcium- and integrin-mediated mechanosignaling, and RhoA signal transduction, consistent
with recent studies indicating that adipocytes mechanically respond to WAT expansion (*23–26*). In contrast,
small adipocytes were enriched for gene sets related to lipid catabolism, mitochondrial organization, and
oxidative phosphorylation, consistent with results linking metabolic syndrome to defective aerobic
metabolism in enlarged adipocytes (*27, 28*). To assess the relevance of our findings in humans, we used a
published cohort of 170 mastectomy patients for which banked histology and paired RNA sequencing of
uninvolved mammary WAT were available (*19*) (**Fig. S3A**). We first quantified the mean adipocyte diameter
using histology for each patient to compare the transcriptomes of mammary WAT with larger and smaller
adipocytes on average (e.g., patients with adipocyte sizes in the fourth and first quartiles, respectively) (**Fig.**
S3B-D). Interestingly, patients in the fourth quartile of mean adipocyte diameter were enriched for similar
biological processes from GSEA as large cells across both mouse models of obesity (**Fig. S3E**). Moreover,
mammary WAT collected from patients in the fourth quartile of adipocyte diameter was significantly enriched
for the hypertrophy-associated gene signature we defined from murine adipocytes (**Fig. S3F**). Collectively,
these results indicate that large and small adipocytes are transcriptionally distinct across two mouse models
of obesity, that large adipocytes are enriched for gene sets related to WAT dysfunction, and that key
transcriptional changes associated with altered adipocyte size are shared between mice and humans.

Figure 2. Large and small adipocytes from obese mice are transcriptionally distinct. (**A**) Principal component analysis of gene expression data for lean adipocytes and size-sorted obese adipocytes. Ellipses show the 95% confidence interval for each condition and individual points represent replicates (n = 4–5). (**B**) Heatmap with hierarchical clustering of normalized gene expression data for the top 500 most variable genes between samples. (**C**) Venn diagram of differentially expressed genes (FDR < 0.05) between large and small adipocytes from HFD and *ob/ob* mice, including a hypertrophy-associated gene signature of 821 genes shared across both models. (**D**) Gene sets from Gene Ontology Biological Processes (GOBP) enriched between large and small adipocytes across both models. All comparisons shown meet the criteria set for statistical significance (FDR < 0.05).

Large adipocytes promote breast cancer cell lipid accumulation, migration, and proliferation.

To assess how large versus small adipocytes differentially impact breast cancer cell behavior, we
encapsulated size-sorted adipocytes into our biomimetic, engineered adipose tissue model as described (*18*)
(**Fig. 3A**). This approach mimics the collagen-rich extracellular matrix of WAT (*29*, *30*), achieves a cell volume
fraction similar to native tissue, and maintains the viability of primary adipocytes in three-dimensional culture
for extended periods of time. Moreover, this setup facilitates coculture studies with breast cancer cells by
allowing for the exchange of soluble factors between adipocytes and tumor cells while enabling separation
of both cell types for downstream analysis. Initial studies with four different breast cancer cell lines and
unsorted adipocytes suggested that, regardless of cell line, tumor cells accumulate more lipid when cultured
with adipocytes from *ob/ob* relative to WT mice (**Fig. S4**). Of the cell lines tested, MDA-MB-231s acquired
the greatest amount of lipid, possibly due to associations between altered lipid metabolism and aggressive
mesenchymal cell states (*31*). Given these findings, MDA-MB-231s were selected for future studies. To
account for differences in adipocyte size reflective of hyperplastic versus hypertrophic expansion, we kept

Figure 3. Large adipocytes promote lipid accumulation, migration, and proliferation of breast cancer cells. (**A**) Schematic of coculture setup between primary adipocytes and MDA-MB-231 breast cancer cells with a representative image of the biomimetic adipose tissue model stained for nuclei (cyan) and F-actin (magenta). Confocal reflectance image (right) shows collagen fiber structure between cells. Scale bar = 100 μm. (**B**) Timeline of coculture experiments. Representative images of cancer cells after monoculture (mono) or coculture with lean or size-sorted obese adipocytes for diet-induced (**C**) and genetic (**K**) mouse models of obesity. Scale bars = 100 μm. Insets show nuclei (cyan) and neutral lipid (gray). Inset scale bars = 35 μm. Quantification of neutral lipid staining with BODIPY 493/503 per cancer cell for diet-induced (**D**) and genetic (**L**) models (n = 3–4). Size distribution of quantified lipid droplets for diet-induced (**E**) and genetic (**M**) models. Box and whisker plots show the distribution of all droplets and circles show the average droplet size per replicate (n = 3–4). Schematic of migration and proliferation assays of precultured cancer cells for diet-induced (**F**) and genetic (**N**) models. Number of migrated cancer cells 12 hours after reseeding onto a tissue-culture insert for diet-induced (**G**) and genetic (**O**) models (n = 3). Representative images of cancer cells 14 hours after reseeding and 2 hours after EdU treatment for diet-induced (**H**) and genetic (**P**) models. Quantification of EdU-positive cancer cells as a percent of total cells for diet-induced (**I**) and genetic (**Q**) models (n = 3). Average lipid content for EdU– and EdU+ cancer cells across diet-induced (**J**) and genetic (**R**) models (n = 3). All data are presented as mean ± standard deviation unless otherwise noted. Statistics for (**J**) and (**R**) performed with a paired t-test per condition. All other statistics performed with a one-way analysis of variance (ANOVA) with Tukey's multiple comparisons test. Compact letter display indicates statistical significance (p = 0.05).

the total volume rather than number of adipocytes constant between conditions. After three days of coculture,
MDA-MB-231 breast cancer cells were either fixed and stained for analysis via fluorescence microscopy or
reseeded for functional assays (**Fig. 3B**). By staining for neutral lipid with BODIPY 493/503 (**Fig. 3C, K**), we
found that breast cancer cells cocultured with large adipocytes from HFD or *ob/ob* mice accumulated
significantly more lipid than cancer cells cocultured with small adipocytes from HFD or *ob/ob* mice, adipocytes
from LFD or WT mice, or monocultured controls (**Fig. 3D, L**). Across both models of obesity, this increase in
neutral lipid was due to a greater number of lipid droplets per cell as well as a greater average lipid droplet
size (**Fig. 3E, M**), suggesting that large adipocytes promote lipid uptake or biosynthesis by neighboring
cancer cells.
176
For functional studies, breast cancer cells precultured with size-sorted adipocytes were reseeded into
collagen-coated tissue culture inserts for migration assays or onto glass coverslips for proliferation assays
(**Fig. 3F, N**). We found that cancer cells precultured with large adipocytes from HFD or *ob/ob* mice migrated
significantly more than cancer cells precultured with small adipocytes from HFD or *ob/ob* mice, adipocytes
from LFD or WT mice, or monocultured controls. (**Fig. 3G, O**). Likewise, breast cancer cells precultured with
large adipocytes from HFD or *ob/ob* mice were more proliferative relative to all other conditions as measured
by incorporation of the thymidine analog 5-ethynyl-2'-deoxyuridine (EdU) (**Fig. 3H-I, P-Q**). Interestingly, EdU-
positive cancer cells had more lipid than EdU-negative cancer cells on average across all conditions, but this
difference was most pronounced and only statistically significant for breast cancer cells precultured with large
adipocytes from HFD or *ob/ob* mice (**Fig. 3J, R**). These results suggest that increased cancer cell proliferation
due to preculture with large adipocytes may be mediated by the greater presence of lipid in these cells, and
we aimed to test this hypothesis moving forward. Collectively, our findings indicate that soluble factors
released by large versus small adipocytes promote breast cancer cell lipid accumulation, as well as migration
and proliferation. Since we observed similar trends between experiments performed with HFD and *ob/ob*
mice, we focused our remaining studies on adipocytes isolated from *ob/ob* mice given that these mice do not
require expensive and lengthy feeding periods prior to tissue isolation.
193

Large adipocytes alter breast cancer cell lipid metabolism.
Given that adipocytes are implicated in altering lipid metabolism in melanoma cells (*32, 33*) and that we
observed adipocyte size positively correlated with lipid accumulation in cocultured breast cancer cells (**Fig.**
3D, L), we speculated that adipocyte size promoted tumor cell migration and proliferation in our studies by
affecting lipid metabolism. Interestingly, fluorescence staining revealed that lipid droplets were closely
associated with the mitochondrial network of MDA-MB-231 breast cancer cells cocultured with adipocytes
(**Fig. 4A**), suggestive of altered lipid metabolism given that FAs must be transported from cytoplasmic lipid
droplets into the mitochondria to generate ATP via a process known as FA oxidation (FAO). To test this
possibility, we first measured the oxygen consumption rate of breast cancer cells precultured with different
adipocyte populations as a relative metric of mitochondrial activity in the presence or absence of etomoxir,
which prevents FAO by irreversibly inhibiting carnitine palmitoyltransferase-1 (CPT1A) and precluding the
transport of FAs into the mitochondria (**Fig. 4B**). At baseline, breast cancer cells precultured with large *ob/ob*
adipocytes consumed more oxygen per ng of DNA than those precultured with small *ob/ob* adipocytes, WT
adipocytes, or monocultured controls as determined by metabolic flux analysis (**Fig. 4C**). Treatment with
etomoxir reduced oxygen consumption rates for all precultured conditions to levels comparable to
monocultured controls for both basal respiration (**Fig. 4D**) and maximal respiration (**Fig. 4E**), which was
induced by the addition of the electron transport chain uncoupler FCCP. However, etomoxir had the greatest
relative effect on MDA-MB-231 cells that were precultured with large *ob/ob* adipocytes. These results indicate
that cell-cell communication between large adipocytes and breast cancer cells increases tumor cell oxygen
consumption in a manner dependent on the oxidation of FAs as metabolic substrates. To determine if this
increased mitochondrial activity was responsible for our observed changes in breast cancer cell behavior in
our previous preculture studies, we repeated these experiments but treated cancer cells with or without

etomoxir when replating for functional assays (**Fig. 4F**). Strikingly, the addition of etomoxir prevented
increases to MDA-MB-231 migration (**Fig. 4G**) and proliferation (**Fig. 4H**) following preculture with large *ob/ob*
adipocytes compared to vehicle controls. Etomoxir treatment also negated the correlation between lipid
content and EdU incorporation (**Fig. 4I**), which suggests that an inability to use FAs as metabolic substrates
prevents the growth advantage otherwise conferred to breast cancer cells with greater lipid accumulation.
Collectively, our results indicate that large adipocytes secrete factors which increase the mitochondrial
activity of breast cancer cells by promoting FAO and, in turn, facilitating their migration and proliferation.

Figure 4. Large adipocytes alter breast cancer cell lipid metabolism to promote migration and proliferation. (**A**) Representative image of a precultured MDA-MB-231 breast cancer cell showing colocalization of neutral lipid and mitochondria. Scale bar = 15 μm. (**B**) Schematic of FAO inhibition by etomoxir. (**C**) Normalized oxygen consumption rate of monocultured or precultured cancer cells over time. Dashed lines indicate sequential addition of etomoxir or vehicle, oligomycin, FCCP, and rotenone/antimycin A (n = 5–6). Only vehicle conditions are plotted to improve visibility. (**D**) Quantification of basal respiration rate per condition from (C) (n = 5–6). (**E**) Quantification of maximal respiration rate per condition from (C) (n = 5–6). (**F**) Schematic of migration and proliferation assays for precultured cancer cells treated with etomoxir or vehicle control. (**G**) Number of migrated cancer cells 12 hours after reseeding onto a tissue-culture insert with or without etomoxir (n = 3). (**H**) Quantification of EdU-positive cancer cells as a percent of total cells with or without etomoxir (n = 3). (**I**) Average lipid content for EdU– and EdU+ cancer cells across conditions (n = 3). All data are presented as mean ± standard deviation unless otherwise noted. Statistics for (I) performed with a paired t-test per condition. All other statistics performed with a one-way analysis of variance (ANOVA) with Tukey's multiple comparisons test. Compact letter display indicates statistical significance (p = 0.05).

Large adipocytes increase lipid transfer via vesicle-mediated release.

225
We next aimed to assess if the increased lipid content of breast cancer cells cocultured with large adipocytes
could be due to size-dependent differences in lipid release. To first confirm that cancer cells uptake lipid
released by adipocytes, we prelabeled adipocytes with BODIPY 493/503 prior to coculture. After 48 hours,
we observed the presence of prelabeled, adipocyte-derived lipid in cocultured breast cancer cells (**Fig. 5A**).
These results suggested that adipocytes increase the lipid content of cocultured cancer cells via direct
transfer of lipid rather than by changing tumor cell lipogenesis. We then aimed to determine how large
adipocytes increase lipid transfer given that adipocytes can release lipids via two separate mechanisms.
Canonical lipolysis by adipocytes is mediated by lipase-dependent breakdown of triglycerides into glycerol
and non-esterified FAs, which then bind to carrier proteins for export into the extracellular space (**Fig. 5Bi**).
In addition, adipocytes can release lipid via a lipase-independent mechanism where small lipid droplets
containing intact triglycerides bud from the central lipid droplet and are packaged into extracellular vesicles
for secretion (*34*) (**Fig. 5Bii**). Conversely, cancer cells store triglycerides in their own lipid droplets, which
requires esterification of FAs following uptake or lipogenesis (**Fig. 5C**). To determine which of these two
mechanisms facilitates increased accumulation of lipid in breast cancer cells due to coculture with large
adipocytes, we first measured the glycerol content of media collected from our coculture experiments as a
relative metric of lipolysis. Media collected from cocultures of cancer cells with large *ob/ob* adipocytes
contained less glycerol than cocultures with small *ob/ob* or WT adipocytes (**Fig. 5D**). These results suggest
that the rate of lipolysis, as estimated by glycerol release, cannot explain the observed increase in lipid
content of cancer cells cocultured with large adipocytes. Therefore, we next tested whether adipocytes
transfer lipid to cancer cells via extracellular vesicles and if this transfer differs as a function of adipocyte
size. To this end, we purified extracellular vesicles from the media of a constant volume of small or large
adipocytes in monoculture via ultracentrifugation for downstream analysis (**Fig. 5E**). We found that large
ob/ob adipocytes secreted significantly more vesicles than small *ob/ob* or WT adipocytes (**Fig. 5F**). This
increase was primarily driven by vesicles less than 200 nm in size, suggestive of greater exosome release
(**Fig. 5G**). Moreover, we confirmed the purity of our isolated vesicles by blotting for the cell-specific marker
FAK and the general extracellular vesicle marker FLOT2 (**Fig. 5H**). We then validated these results with
adipocytes isolated from HFD and LFD mice. Similarly, we found that media from cocultures of cancer cells
with large HFD adipocytes contained less glycerol than other conditions and that large HFD adipocytes
secreted significantly more vesicles per cell volume than small HFD or LFD adipocytes (**Fig. S5**).

255
To determine if adipocytes of varying size differed in their lipid composition, which could, in turn, influence
lipid accumulation in cocultured breast cancer cells, we performed targeted lipidomics of total and non-
esterified FAs in WT, small *ob/ob*, and large *ob/ob* adipocytes (**Fig. 5I**). While large *ob/ob* adipocytes
contained more saturated, unsaturated, and total FAs per mg protein relative to small *ob/ob* and WT
adipocytes as expected, we did not observe consistent differences in the relative composition of specific lipid
species in small versus large *ob/ob* adipocytes (**Fig. S6-S7**). However, both small and large *ob/ob* adipocytes
contained a higher ratio of arachidonic acid to eicosapentaenoic and docosahexaenoic acids than WT
adipocytes, indicative of a pro-inflammatory lipid profile and suggestive of altered FA desaturase activity in
ob/ob adipocytes given that *ob/ob* and WT mice received the same diet (**Fig. S6-S7**). These results
underscore the importance of examining adipocytes from the same donor when investigating the impact of
size on adipocyte function, as some differences in adipocyte phenotype are donor-dependent regardless of
size. Interestingly, we found that the percentage of non-esterified to total FAs was significantly decreased in
large versus small *ob/ob* adipocytes (**Fig. 5J**), consistent with our glycerol release data that large cells are
less lipolytic relative to small cells. Finally, to confirm that the increased lipid content of cancer cells cocultured
with large versus small *ob/ob* adipocytes was due to vesicle-mediated transfer, we applied extracellular
vesicles collected and purified from a constant volume of size-sorted adipocytes in monoculture to cancer
cells in monoculture. Moreover, we inhibited DGAT-1 and DGAT-2 (iDGAT) concurrently with vesicle
treatment in a subset of samples to prevent esterification of free FAs and further rule out the contributions of

Figure 5. Large adipocytes promote breast cancer cell lipid accumulation through extracellular vesicle-mediated transfer of triglycerides. (**A**) Schematic of lipid transfer experiment. Adipocytes were pre-labeled with BODIPY 493/503 and washed to remove excess dye prior to the initiation of coculture. Scale bar = 35 μm. (**B**) Schematic of (i) lipase-mediated and (ii) vesicle-mediated lipid release by adipocytes. (**C**) Schematic of lipid storage by cancer cells. (**D**) Concentration of glycerol in conditioned media collected from cancer cells and adipocytes in coculture after 72 hours (n = 3–4). (**E**) Schematic of extracellular vesicle isolation from adipocyte-conditioned media. (**F**) Concentration of extracellular vesicles collected from adipocytes in monoculture after 24 hours (n = 5). (**G**) Size distribution of extracellular vesicles from (F) (n = 5). (**H**) Western blot of cell and extracellular vesicle lysates collected from lean and size-sorted obese adipocytes. (**I**) Heatmap of targeted lipidomics data showing the total FA composition of lean and size-sorted obese adipocytes normalized to protein content (n = 5). (**J**) Percent of non-esterified relative to total FAs from (I) (n = 5). (**K**) Representative images of MDA-MB-231 breast cancer cells 24 hours after treatment with oleic acid as a positive control or extracellular vesicles collected from WT, small *ob/ob*, or large *ob/ob* adipocytes. Cancer cells were also treated concurrently with vehicle or dual inhibitors for DGAT-1 and DGAT-2 (iDGAT). Scale bars = 35 μm. (**L**) Quantification of neutral lipid staining with BODIPY 493/503 from (K) per cancer cell (n = 4). All data are presented as mean ± standard deviation unless otherwise noted. Statistics were performed with a one-way analysis of variance (ANOVA) with Tukey's multiple comparisons test. Compact letter display indicates statistical significance (p = 0.05).

lipogenesis by cancer cells (**Fig. 5C**). After 24 hours, we found that breast cancer cells treated with vesicles
from large *ob/ob* adipocytes contained significantly more lipid than those treated with vesicles from small
ob/ob or WT adipocytes (**Fig. 5K-L**). However, lipid uptake was significantly lower across all conditions
compared to coculture studies, likely due to the shorter period of vesicle collection (24 hours) than coculture
(72 hours) and loss of vesicles during sample processing. Nevertheless, iDGAT prevented lipid storage in
tumor cells treated with exogenous oleic acid as expected, while iDGAT treatment did not affect lipid transfer
via adipocyte-derived vesicles (**Fig. 5K-L**), indicating that these vesicles contain intact triglycerides rather
than non-esterified FAs and that altered lipogenesis does not significantly contribute to cancer cell lipid
content in our coculture system. Collectively, our results indicate that large adipocytes promote breast cancer
cell lipid uptake through vesicle-mediated transfer of intact triglycerides.
284

Adipocyte size is more predictive of lipid dysfunction than body mass index.
Given that we observed greater lipid release by large adipocytes in our coculture studies, we next aimed to
determine if adipocyte size correlated with broader, systemic metabolic changes in humans. To this end, we
compared banked adipose tissue and pre-operative blood panels for each of the 170 mastectomy patients
in our cohort (**Fig. 6A**). Again, we stratified patients into quartiles based on their average adipocyte diameter
determined via paired histology (**Fig. 6B**). Using this approach, we found that average adipocyte diameter
(**Fig. 6C**) more accurately stratified patients diagnosed with dyslipidemia than BMI category (**Fig. 6D**).
Likewise, average adipocyte diameter correlated more significantly with fasting triglycerides (**Fig. 6E**) than
BMI (**Fig. 6F**), even when their interaction effects were considered using multiple linear regression (**Fig. 6G**).
Collectively, these findings suggest that in addition to releasing more lipid to cells locally, as observed in our
coculture studies, large adipocytes may also increase lipid release into circulation with consequences for
whole-body metabolism.
297

Figure 6. Adipocyte size is more predictive of systemic lipid dysfunction than body mass index. (**A**) Schematic of paired mammary adipose tissue histology and pre-operative blood panels from a cohort of 170 mastectomy patients. (**B**) Histogram of patients stratified into quartiles based on their average adipocyte diameter determined from histology. Percent of patients diagnosed with dyslipidemia when classified by (**C**) average adipocyte diameter or (**D**) BMI category (BMI < 25 for normal, BMI > 30 for obese). Simple linear regression of fasting triglycerides versus (**E**) average adipocyte diameter or (**F**) BMI. (**G**) Multiple linear regression of fasting triglycerides as a function of both average adipocyte diameter and BMI following z-score normalization.

Discussion

Obesity is an established risk factor for breast cancer initiation and prognosis but the mechanisms behind these associations remain unclear. Our results suggest that the increased presence of hypertrophic adipocytes in obese WAT contributes to these observations and that adipocyte size could serve as a more specific prognostic biomarker than BMI for breast cancer patients. By using our biomimetic, engineered adipose tissue model, we find that large adipocytes are distinct from small adipocytes isolated from donor-matched tissue and increase lipid accumulation in neighboring breast cancer cells through vesicle-mediated release of triglycerides. In turn, coculture with large adipocytes enhances breast cancer cell migration, proliferation, and oxygen consumption in a manner dependent on FAO. Collectively, our findings suggest that examining mammary WAT morphology during breast cancer biopsy could help identify women at greatest risk for aggressive disease and regions of tissue most susceptible to invasion.

Our understanding of adipocyte heterogeneity and biology remains limited given that single-cell analysis of primary cells is challenging due to their buoyancy and fragility (*6, 17*). Here, we sequenced adipocytes phenotypically sorted by size and identified that large, hypertrophic adipocytes constitute a transcriptionally distinct subpopulation of cells. Our findings are consistent with recent results demonstrating that WAT is more plastic (*4, 5*) and heterogenous (*6, 7*) than once considered. For example, multiple adipocyte subpopulations have been identified in mice with distinct sensitivities to insulin (*7*) and whose relative proportions change with administration of a high-fat diet (*6*)**.** In our data, large adipocytes from both diet-induced and genetic mouse models of obesity were enriched for gene sets related to WAT dysfunction (*13*), as were tissue samples collected from mastectomy patients with increased adipocyte size regardless of BMI. Moreover, large adipocytes also exhibited enrichment for gene sets related to collagen deposition, actin cytoskeleton remodeling, calcium ion transport, and Rho protein signal transduction, consistent with an emerging understanding of the mechanoresponsive adaptations required of adipocytes during drastic cell expansion (*24, 26*). To what extent these transcriptional changes are adipocyte intrinsic or a function of environmental cues in hypertrophic regions of WAT remains to be determined. Re-examining the transcriptome of large adipocytes after extended *ex vivo* culture in our biomimetic tissue model could help make this distinction. Likewise, assessing hypertrophic adipocytes in the absence of obesity, such as those present in lean individuals with metabolic disorders (*14*), could help identify the relative contribution of cell intrinsic factors. Regardless of the cause, our findings underscore the need for a more nuanced view of WAT beyond BMI that accounts for other parameters such as adipocyte size. Indeed, in blood panels collected from our cohort of mastectomy patients, average adipocyte diameter correlated more significantly with dyslipidemia and fasting triglycerides than BMI, even when their interaction effects were considered—findings that are broadly consistent with the increased release of lipid by large adipocytes in our coculture experiments.

Previous studies have demonstrated that adipocytes can influence the behavior of neighboring tumor cells, though this connection is most established for melanoma compared to other carcinomas that develop in WAT-rich stroma such as breast cancer. For example, researchers found that melanoma cells overexpress fatty acid transport protein 1 (FATP1), which facilitates the uptake of adipocyte-derived non-esterified FAs and promotes tumor invasion (*33*). In a separate study, adipocyte-derived exosomes, which were enriched for proteins implicated in FAO, mediated the metabolic reprogramming of melanoma cells (*32*). In the context of breast cancer, several studies have linked adipocyte-derived lipids to altered tumor cell metabolism (*10, 35*). However, these studies used *in vitro* differentiated progenitor cells and thus did not assess obesity- or hypertrophy-dependent effects. Indeed, while we saw trends for cocultures with small adipocytes, we only observed significant changes in breast cancer cell phenotypes (i.e. lipid content, migration, proliferation, oxygen consumption) for cocultures with large adipocytes. This may be due to our experimental setup, as we performed shorter cocultures than some of the cited studies (*10, 33*) or may reflect underlying differences in adipocyte-tumor cell communication when using primary adipocytes. Moreover, our results corroborate previous findings that obese WAT secretes more extracellular vesicles per mg tissue than lean WAT (*34*) and

that this increase is driven by adipocytes rather than cells of the stromal vascular fraction (*32*). Our findings
indicate that the greater rate of extracellular vesicle release by obese WAT likely stems from the increased
presence of hypertrophic adipocytes in obesity, a conclusion not previously possible with gross analysis of
tissue explants or unsorted cells. Moving forward, future studies will be needed to assess the impact of these
adipocyte-derived vesicles on other breast cancer subtypes and stromal cells, such as tissue-resident
macrophages and fibroblasts, within the tumor microenvironment.
354

Across experiments, our findings suggest that large adipocytes shift toward favoring a vesicle-mediated
mode of lipid release, though the lack of endocrine signals which stimulate lipolysis in our system may be a
confounding factor in these observations. Indeed, across both mouse and human sequencing data,
hypertrophic adipocytes and hypertrophic WAT demonstrated positive gene set enrichment for vesicle-
mediated transport and negative gene set enrichment for lipid catabolism. While initially counterintuitive given
the greater degree of lipid transfer observed between large adipocytes and breast cancer cells in coculture,
these findings are consistent with a lipase-independent, vesicle-mediated mode of lipid release by
hypertrophic cells. Direct transfer of adipocyte-derived triglycerides via extracellular vesicles as opposed to
non-esterified FAs via lipolysis could offer several potential advantages to recipient tumor cells. Beyond
serving as metabolic substrates for FAO, a steady supply of vesicle-delivered triglycerides may also provide
a readily available source of FAs for membrane biosynthesis, lipid signaling, and intracellular trafficking. In
particular, tumor cells undergoing rapid division exhibit increased demand for phospholipids to sustain
plasma membrane expansion, and specific lipid classes such as monounsaturated and polyunsaturated FAs
are necessary for membrane fluidity and lipid raft formation (*36*). Our lipidomics data indicate that large
adipocytes contain an increased abundance of these classes of FAs relative to small adipocytes. This finding
suggests that vesicle-mediated lipid transfer could facilitate tumor progression not only by fueling oxidative
metabolism but also by supplying key structural lipids for tumor growth. Future untargeted lipidomics of the
specific lipid species enriched in adipocyte-derived vesicles, as well as the lipids transferred to recipient
tumor cells, will be critical to dissect this potential dual role of FAs in breast cancer malignancy.
374

In addition to triglycerides, previous studies have demonstrated that extracellular vesicles secreted by
adipocytes are also enriched for lipid-droplet-associated proteins required for FAO (*32*, *34*). Thus, by
increasing vesicle release, hypertrophic adipocytes may provide tumor cells not only with FAs for
mitochondrial oxidation but also the molecular machinery needed to metabolize them. Future proteomic
studies of vesicles derived from large versus small adipocytes would reveal if altered protein cargo
contributed to our observed increase in FAO for breast cancer cells precultured with large adipocytes.
Moreover, excess quantities of non-esterified FAs, especially trans- or saturated FAs, can induce lipotoxicity
by activating cell stress responses (*37*). Indeed, inhibiting the ability of tumor cells to esterify free FAs, and
thus induce lipotoxicity, has been explored as a therapeutic strategy for hepatocellular carcinoma (*38*). In
contrast, the direct transfer of intact triglycerides would bypass the need for FA esterification and limit the
potential lipotoxic effects on cancer cells. Conversely, controlled oxidation of FAs by tumor cells has been
shown to mediate chemoresistance to platinating agents (*39*), which are increasingly prescribed for patients
with advanced or triple-negative breast cancer (*40*). Thus, moving forward, the effect of specific adipocyte
populations of interest on breast cancer chemoresistance should be explored. More broadly, this shift in the
mode of lipid release by hypertrophic adipocytes likely produces systemic metabolic alterations outside the
context of cancer by altering the bioavailability and biodistribution of lipids in circulation, as supported by our
patient data.
392

In conclusion, our results suggest that the increased presence of hypertrophic adipocytes in the mammary
WAT of obese breast cancer patients contributes to the poor clinical outcomes seen in these women.
However, risk stratification by BMI alone is a flawed approach given that not all WAT expands equally. Indeed,
our findings posit that adipocyte size could serve as an independent prognostic biomarker for breast cancer

patients regardless of BMI. Determining adipocyte size is clinically tractable given that peritumoral WAT is
often included in routine biopsies used for diagnosis and automated approaches exist to characterize
adipocyte morphology (*41*, *42*). Moreover, our results indicate that biomimetic tissue models can capture
phenomena not assessable with traditional approaches, demonstrating the value of bioengineering strategies
in basic research. Collectively, our findings provide insight into the role of hypertrophic adipocytes as a distinct
cell population with implications for the management of breast cancer and metabolic diseases more broadly.

Resource availability

All source data and code will be provided in the final publication of this manuscript.

Acknowledgments

We thank all members of the Fischbach Lab for valuable discussion and input during the preparation of this manuscript; the Weill Institute of Cell and Molecular Biology and David McDermitt for access to an ultracentrifuge; Rebecca Williams for assistance with confocal microscopy; Tina Abratte for assistance with metabolic flux experiments; Malin Lönn for assistance with developing the adipocyte size-sorting protocol; the Cornell Center for Animal Resources and Education (CARE), Matthew Whitman, and Jessie Gorges for assistance with animal studies; the Cornell Animal Health Diagnostic Core for assistance with paraffin embedding and sectioning; and the Cornell Statistical Consulting Unit for assistance with statistical analysis. We acknowledge financial support from the NSF DGE1650441 (G.F.B.), NCI F31CA278410 (G.F.B.), NCI R01CA259195 (C.F., M.A.A.) and R01CA276392 (C.F.), the Center on the Physics of Cancer Metabolism NCI 1U54CA210184 (C.F.), and the Cornell Engineering Learning Initiatives program (M.I.P.). We also acknowledge support from the Cornell Biotechnology Resource Center Genomics Facility (RRID:SCR_021727) and the Cornell Biotechnology Resource Center Imaging Facility funded by NYSTEM C029155, NIH S10OD018516, and NIH S10RR025502.

Author contributions

G.F.B. and C.F. conceived and designed experiments. G.F.B. performed all experiments with assistance from M.I.P. unless otherwise noted. R.R.V. and M.C.R. performed targeted lipidomics and associated analysis. M.A.A. performed Western blotting of cell and extracellular vesicle lysates. D.D.G., D.J.F., and N.M.I. obtained and analyzed the cohort of human mammary adipose tissue collected from mastectomy patients. G.F.B. prepared all figures and schematics. G.F.B. and C.F. wrote the manuscript with input from all authors.

Declaration of interests

The authors declare no competing interests.

Supplemental data

Figures S1–S7.

Methods

Culture of breast cancer cell lines

MDA-MB-231, MCF7, PY8119, and EO771 breast cancer cell lines were maintained in high-glucose Dulbecco's Modified Eagle Medium (Gibco #12800) supplemented with 10% (v/v) fetal bovine serum (FBS, R&D Systems #S11150) and 1% (v/v) penicillin-streptomycin (ThermoFisher #15070063) unless otherwise noted. Primary adipocytes and cocultures were maintained in a 1:1 mixture of low-glucose Dulbecco's Modified Eagle Medium (Gibco #316000) and Ham's F12 Nutrient Mix (Gibco #21700) supplemented with 10% (v/v) FBS and 1% (v/v) penicillin-streptomycin unless otherwise noted. For functional studies of precultured tumor cells, a 1:1 mixture of low-glucose Dulbecco's Modified Eagle Medium and Ham's F12 Nutrient Mix supplemented with 1% (v/v) FBS and 1% (v/v) penicillin-streptomycin was used. For experiments involving extracellular vesicle collection or treatment, exosome-depleted FBS (ThermoFisher #A2720803) was used to limit the presence of exogenous vesicles.

Maintenance and differentiation of 3T3-L1 preadipocytes

Prior to the induction of adipogenesis, 3T3-L1s were seeded at a density of 5,000 cells per cm^2 onto 12 mm glass coverslips coated with 30 µg/mL collagen type I (VWR #47747-218) and cultured until 80% confluent. Adipogenesis was initiated with induction media consisting of high-glucose DMEM supplemented with 5% FBS, 1% penicillin-streptomycin, 1 mM dexamethasone (Sigma-Aldrich #D4902), 500 mM IBMX (Krackeler Scientific #45-I7018), 20 mM indomethacin (Sigma-Aldrich #405268), and 200 µM insulin (Krackeler Scientific #45-91077C). After 48 hours, cells were changed to maintenance media consisting of high-glucose DMEM supplemented with 5% FBS, 1% penicillin-streptomycin, and 200 µM insulin. Maintenance media was replaced every 48 hours until experimental endpoints were reached.

Animal models

For all animal studies, WAT was isolated from the anterior and inguinal fat pads around the mammary glands of nulliparous female C57BL6 mice (The Jackson Laboratories). For the genetic model of obesity, wildtype C57BL6 mice (strain 000664) and leptin-deficient B6.Cg-Lep[ob] mice (strain 000632) were purchased at 10 weeks of age and fed a standard rodent diet (LabDiet #5053) *ad libitum* until sacrifice at 12 weeks of age. For the dietary model of obesity, wildtype C57BL6 mice were purchased at 6 weeks of age and fed a standard rodent diet for a two-week acclimation period. At 8 weeks of age, mice were randomized and placed on either a low-fat diet (Research Diets #D12450J, 10 kcal% fat) or high-fat diet (Research Diets #D12492, 60 kcal% fat) *ad libitum* until sacrifice at 20 weeks of age. Diets were administered weekly and remaining chow from the previous week was discarded. All mice were housed between 21°C and 23°C. All animal protocols were approved by the Institutional Animal Care and Use Committees at Weill Cornell Medicine and Cornell University.

Histology of resected adipose tissue

Pieces of resected WAT were collected from euthanized C57BL6 mice and fixed in 4% (w/v) paraformaldehyde in 1x PBS for 24 hours at 4°C. Resected tissue was then stored in 70% (v/v) ethanol in water at 4°C until processing. Samples were sent to the Cornell College of Veterinary Medicine Animal Health Diagnostic Center for paraffin embedding, sectioning, and staining with hematoxylin and eosin. Stained sections were imaged and digitized on an Aperio ScanScope CS2 (Leica) with a 40x objective.

Primary adipocyte isolation

Resected WAT was immediately placed in cold DMEM/F12 until processing. Tissue was then minced for several minutes until separated into approximately 1 mm^3 pieces. Floating tissue pieces were decanted and reserved while any remaining pieces were discarded. The reserved tissue was then digested in 1.5 mg/mL collagenase type I (Worthington Biochemical #LS004197) in Hank's Buffered Salt Solution (Gibco #14065056) supplemented with 1% (w/v) bovine serum albumin (ThermoFisher #BP1600-100) in a vertical

tube rotator for 50 minutes at 37°C. The resulting cell suspension was passed through 200 µm cell strainers
(pluriSelect #435020003) to remove undigested pieces of tissue. Adipocytes were allowed to float out of
solution for 10 minutes before the infranatant was aspirated and Krebs-Ringer HEPES buffer (116 mM NaCl,
25 mM HEPES, 4 mM KCl, 2 mM D-glucose, 1.8 mM $CaCl_2$, 1 mM $MgCl_2$) supplemented with 1% (w/v)
bovine serum albumin (KRHB) was added. This process was repeated three times to remove contaminating
stromal vascular cells. To ensure consistent cell seeding, isolated adipocytes were packed via centrifugation
for 3 minutes at 100 g and the resulting layers of aqueous buffer and free lipid were aspirated before use as
described(18). All subsequent handling of isolated adipocytes was performed with wide bore pipette tips.

Size sorting of primary adipocytes

To separate adipocytes by size, isolated cells were placed in a 125 mL separatory funnel (ThermoFisher
#4301-0125) along with 50 mL of KRHB. The funnel was gently inverted several times to resuspend the
adipocytes. After 35 seconds, the bottom 35 mL of cell suspension was eluted and passed through a 50 µm
cell strainer (pluriSelect #435005003). 35 mL of fresh KRHB was added to the funnel and this process was
repeated. Adipocytes that passed through the cell strainers were pooled and used as the "small" fraction for
all experiments. 35 mL of fresh KRHB was then added to the funnel before gently inverting several times.
After 15 seconds, the bottom 35 mL of cell suspension was eluted and discarded. This process was repeated
two additional times to remove cells of intermediate size. The remaining adipocytes were treated as the
"large" fraction for all experiments.

Assessment of primary adipocyte viability

To assess cell viability after isolation and sorting, unfixed adipocytes were stained with 2 µg/mL Hoechst
33342 (ThermoFisher #H3570) and 2 µg/mL propidium iodide (ThermoFisher #P3566) in DMEM/F12 for 30
minutes at 37°C protected from light. To image floating adipocytes, an imaging chamber was created by
stacking two 120 µm-thick imaging spacers (GraceBio #654004) on top of a microscope slide. Primary
adipocytes were diluted in DMEM/F12, added as a 30 µL drop to the center of the spacers, and covered with
a glass coverslip for imaging. The percentage of live adipocytes was quantified as the number of cells with
propidium iodide-negative nuclei over the number of cells with Hoechst-positive nuclei. The imaging chamber
was cleaned with 70% (v/v) ethanol, rinsed with 1x PBS, and dried between samples.

RNA sequencing and analysis

For sample collection, 100 µL of packed adipocytes were lysed in 1 mL Qiazol (Qiagen #79306), incubated
for 5 minutes at room temperature, and then stored at –80°C until RNA extraction. RNA extraction was
performed using an RNeasy Lipid Tissue Mini Kit (Qiagen #74804) according to manufacturer protocols. RNA
concentration, 260/280 purity, and 260/230 purity were assessed via a NanoDrop Spectrophotometer
(ThermoFisher) and RNA concentration and quality were assessed via a 4200 TapeStation System (Agilent).
Libraries were then prepared using a QuantSeq FWD Kit (Lexogen) and sequenced as 86 bp single-end
reads on a NextSeq500 System (Illumina). Reads were trimmed using fastp (v.0.23.4) and aligned to the
mouse reference genome GRCm39 (GENCODE) using STAR (v.2.7.10b). Reads were counted using the
GeneCounts feature of STAR and differential expression analysis was performed with DESeq2 (v.1.40.2).
Differentially expressed genes were defined as those with an absolute $\log_2$ fold-change greater than 1 and
adjusted p-value less than 0.05. Heat maps were generated using pheatmap (v.1.0.12), volcano plots were
generated using EnhancedVolcano (v.1.18.0), and principal component analysis plots were generated using
PCAtools (v.2.12.0). Gene set enrichment analysis (GSEA) was performed using fgsea (v.1.26.0) by
providing a list of ranked genes from the DESeq2 statistic column.

Collection of mastectomy patient data

529   The Institutional Review Boards of Memorial Sloan Kettering Cancer Center (IRB 10-040) and Weill Cornell
530   Medicine (IRB 1004010984-01) approved the collection of non-tumor breast tissue and blood under a

biospecimen acquisition protocol. Informed consent was obtained from all participants. In total, 196 patients underwent mastectomy for breast cancer risk reduction or treatment. Height and weight were recorded prior to surgery and used to calculate body mass index (BMI). Standard definitions were used to categorize BMI as normal weight (BMI < 25), overweight (BMI 25.0–29.9), or obese (BMI ≥ 30). All data were reviewed for accuracy independently by research staff and a physician. On the day of mastectomy, paraffin blocks and snap frozen samples were prepared from breast WAT. If a tumor was present, samples were taken from an uninvolved quadrant of the breast. Two hematoxylin- and eosin-stained sections were generated from formalin-fixed, paraffin-embedded breast tissue in order to measure adipocyte diameters as previously described(43). The sections were photographed with a 20x objective using an Olympus BX43 or BX50 microscope equipped with an Olympus DP27 or MicroFire (Optronics) digital camera, respectively. Mean diameters were calculated using measurements for each patient using the linear dimensional tool in the Canvas 11 Software (ACD Systems International, Inc.), which was calibrated using a microscope stage micrometer.

On the day of surgery, a 30 mL fasting blood sample was obtained preoperatively. Centrifugation was used to separate blood into serum and plasma within 3 hours of collection and stored at −80°C. Serum levels of triglycerides and high-density lipoprotein (HDL) cholesterol were determined in the Memorial Sloan Kettering Cancer Center Clinical Chemistry Lab.

Human RNA sequencing data have been previously published(19) and were deposited to the European Genome-Phenome Archive (EGA) under accession number EGAS00001004665.

Preparation of elastomer microwells
Elastomer microwells for *ex vivo* culture of primary adipocytes were prepared as previously described(18). Briefly, SYLGARD 184 silicone elastomer (Ellsworth Adhesives #4019862) was cast in 1.5 mm-thick slabs according to manufacturer protocols. 10 mm outer diameter, 6 mm inner diameter elastomer rings were cut and bonded to glass coverslips using a plasma cleaner (Harrick Plasma). The rings were treated with 1% (v/v) poly(ethylenimine) (Sigma-Aldrich #181978), 0.1% (v/v) glutaraldehyde (ThermoFisher #ICN19859580), and washed three times with sterile water to facilitate covalent adhesion of adipocyte-collagen gels.

Preparation of adipocyte-collagen gels
For *ex vivo* culture, packed adipocytes were mixed into a collagen gel at a 3:1 ratio (v/v) using wide-bore pipette tips as previously described(18). Briefly, to prepare the collagen gel, collagen type I (VWR #47747-218) was diluted to a final concentration of 2.5 mg/mL in DMEM/F12. This solution was kept on ice and neutralized with 1N NaOH to begin polymerization immediately before the addition of packed adipocytes. During polymerization, the adipocyte-collagen gels were inverted every 2 minutes for 20 minutes at room temperature to ensure an even distribution of cells. The resulting gels were submerged in DMEM/F12 and cultured for up to 5 days at 37°C.

Coculture studies
For coculture studies, 30,000 breast cancer cells (MDA-MB-231, MCF7, PY8119, or EO771) were plated per well in a 24-well plate containing 12 mm glass coverslips (Chemglass #CLS-1760-012) coated with 30 µg/mL collagen type I (VWR #47747-218). The following day, primary adipocytes were isolated and adipocyte-collagen gels in elastomer microwells were prepared. One cancer cell coverslip and one adipocyte-collagen gel were placed into each well of a 12-well plate along with 1.5 mL of DMEM/F12. Cancer cells and primary adipocytes were cocultured together for 72 hours. Cancer cells were then reserved for functional studies or fixed in 4% (w/v) paraformaldehyde for imaging.

EdU proliferation assays
For proliferation studies, precultured cancer cells were trypsinized and reseeded at 30,000 cells per well in
a 24-well plate containing 12 mm glass coverslips coated with 30 µg/mL collagen type I (VWR #47747-218).
Once adherent, replated tumor cells were incubated in low-serum DMEM/F12 for 12 hours. The following
morning, proliferating tumor cells were labeled with the Alexa Fluor 647 EdU Kit (ThermoFisher #C10340)
according to manufacturer protocols. Briefly, a 20 µM EdU solution was prepared in low-serum DMEM/F12
and added 1:1 to existing media to achieve a final concentration of 10 µM EdU. Cancer cells were incubated
for 2 hours at 37°C before proceeding with fixation, permeabilization, and detection of incorporated EdU.
Cancer cells were co-stained with Hoechst 33342 (ThermoFisher #H3570) and BODIPY 493/503
(ThermoFisher #D3922). For inhibitor studies, 40 µM etomoxir (Millipore #236020) or DMSO vehicle were
added with the low-serum DMEM/F12 after precultured cancer cells adhered to collagen-coated glass
coverslips.
592
Tissue-culture insert migration assays
For migration assays, precultured cancer cell coverslips were transferred to a 24-well plate and incubated in
low-serum DMEM/F12 for 12 hours. The following morning, 8 µm pore tissue-culture inserts (VWR #29442-
120) were coated with 30 µg/mL collagen type I (VWR #47747-218) and placed in 24-well plates containing
600 µL of DMEM/F12. Precultured cancer cells were then trypsinized and reseeded at 30,000 cells per tissue-
culture insert in 100 µL of low-serum DMEM/F12 to establish a serum gradient. Cancer cells were allowed to
migrate for 12 hours at 37°C. Cells were then fixed with 4% paraformaldehyde and a cotton swab was used
to gently remove cells from the upper side of the insert membrane. The underside of each membrane was
stained with DAPI (ThermoFisher #D1306) and imaged to quantify the number of transmigrated cancer cells.
For inhibitor studies, 40 µM etomoxir (Millipore #236020) or DMSO vehicle were added to the low-serum
DMEM/F12 when reseeding precultured cancer cells into tissue-culture inserts.
604
Mitochondrial imaging in live breast cancer cells
To assess co-localization of breast cancer cell lipid droplets and mitochondria after coculture with adipocytes,
precultured MDA-MB-231s were reseeded at 100,000 cells per glass-bottom petri dish (World Precision
Instruments #FD35-100) coated with 30 µg/mL collagen type I (VWR #47747-218). Precultured cancer cells
were then stained with 200 nM MitoView 640 (Biotium #70082) for 12 hours. Live samples were co-stained
with 2 µg/mL Hoechst 33342 and 2.5 µg/mL BODIPY 493/503 for 30 minutes, washed twice with DMEM/F12,
and imaged on an inverted Zeiss LSM 880 confocal microscope with an incubation chamber.
612
Pre-labeling lipid to monitor transfer from adipocytes to breast cancer cells
For pre-labeling experiments, primary adipocytes were stained in suspension with 2.5 µg/mL BODIPY
493/503 for 30 minutes and washed three times with KRHB prior to embedding into collagen gels. Pre-labeled
adipocytes were then cocultured with MDA-MB-231 breast cancer cells as described. After 48 hours, breast
cancer cells were fixed with 4% (w/v) paraformaldehyde in 1x PBS and counterstained with 2.5 µg/mL DAPI
and 165 µM Alexa Fluor 568 phalloidin. Breast cancer cells were then imaged on an inverted Zeiss LSM 880
confocal microscope to detect the presence of BODIPY-labeled neutral lipid.
620
Sample fixation, permeabilization, and fluorescence staining
Samples were fixed in cold 4% (w/v) paraformaldehyde (Electron Microscopy Sciences #19208) in 1x PBS
for 15 minutes at room temperature. Samples were washed twice with 1x PBS and permeabilized with 0.1%
Triton X-100 (v/v) (ThermoFisher #A16046-AE) in 1x PBS supplemented with 1% (w/v) bovine serum albumin
(BSA) (ThermoFisher #BP1600-100) for 15 minutes at room temperature. Samples were washed twice with
1x PBS and then stained with one or more of the indicated dyes as follows: 0.165 µM Alexa Fluor 488
phalloidin (ThermoFisher #A12379) or 0.165 µM Alexa Fluor 568 phalloidin (ThermoFisher #A12380) in 1x
PBS with 1% BSA for 1 hour at room temperature; 2.5 µg/mL 4,4-difluoro-1,3,5,7,8-pentamethyl-4-bora-

3a,4a-diaza-*s*-indacene (BODIPY 493/503) in 1x PBS with 1% BSA for 30 minutes at room temperature; 2.5
µg/mL 4',6-diamidino-2-phenylindole, dihydrochloride (DAPI) (ThermoFisher #D1306) in 1x PBS with 1%
BSA for 30 minutes at room temperature. Samples were then washed twice with 1x PBS and stored in 1x
PBS at 4°C until imaged.

Confocal microscopy and image analysis

Images were acquired on an inverted or upright Zeiss LSM 880 confocal microscope using either a 10x/0.45
W C-Apochromat objective, 20x/0.5 EC Plan-Neofluar objective, 20x/1.0 W Plan-Apochromat objective,
40x/1.2 W C-Apochromat objective, or 63x/1.4 O Plan-Apochromat objective. All images analysis was
performed in QuPath(*44*) and ImageJ (National Institutes of Health). For semi-automated quantification of
adipocytes, cancer cell lipid droplets, and 3T3-L1 lipid droplets, the AdipoQ ImageJ plug-in(*45*) was used.
Manuscript figures were prepared using Adobe Photoshop, Adobe Illustrator, and Biorender.com.

Substrate oxidation stress test

To assess ATP production and FAO, cancer cell metabolism was probed using a Seahorse XFe 96
Extracellular Flux Analyzer (Agilent) and Seahorse XF Long Chain Fatty Acid Oxidation Stress Test Kit
(Agilent #103672-100). Control or precultured cancer cells were seeded at 20,000 cells per well in an XFe96
Cell Culture Microplate (Agilent #103794-100) in DMEM/F12 and allowed to adhere overnight. The following
morning, media was changed to Seahorse XF DMEM, pH 7.4 (Agilent #103680-100), and manufacturer
protocols were followed to complete the Long Chain Fatty Acid Oxidation Stress Test Kit. Inhibitor
concentrations of 4 µM for etomoxir, 1.5 µM for oligomycin, 1 µM for FCCP, and 0.5 µM for rotenone/antimycin
A were used. After the stress test was complete, DNA was extracted using lysis buffer (25 mM Tris-HCl, 0.4
M NaCl, 0.5% (w/v) sodium dodecyl sulfate) and total DNA content per well was measured using the
QuantiFluor dsDNA Assay (VWR #PAE2671) for data normalization.

Glycerol assay

655   Conditioned media from cancer cell and adipocyte-collagen gel cocultures was collected after 72 hours and
656   stored at –20°C until processed. Once thawed, the relative degree of lipolysis during culture was estimated
657   by measuring the concentration of glycerol in the conditioned media using the Adipolysis Assay Kit (Sigma-
658   Aldrich #MAK313) according to manufacturer protocols.

Isolation of extracellular vesicles from adipocyte-conditioned media

Adipocyte-collagen gels were cultured in 24-well plates containing 750 µL of exosome-depleted DMEM/F12
to generate conditioned media. Extracellular vesicles were then isolated from conditioned media using
modified versions of published protocols(*46*). Briefly, conditioned media was centrifugated at 300 x g at 4°C
for 10 minutes to remove dead cells and other debris. The resulting supernatant was transferred to a new
tube and centrifuged at 3000 x g at 4°C for 30 minutes to remove apoptotic bodies. The resulting supernatant
was transferred to a new tube and combined at a 1:1 ratio (v/v) with 16% (w/v) PEG 6000 (Sigma-Aldrich
#82160) in 1 M NaCl for a final concentration of 8% PEG 6000 in 0.5 M NaCl. The resulting solution was
inverted several times and incubated at 4°C for at least 12 hours to precipitate soluble factors and
extracellular vesicles. The precipitate was then pelleted via centrifugation at 3220 x g at 4°C for 1 hour. After
aspirating the supernatant containing PEG 6000, the pellet was resuspended in 2 mL of filtered 1x PBS and
ultracentrifuged in a TLS-55 swinging-bucket rotor (Beckman Coulter #346936) at 100,000 x g at 4°C for 4
hours. The supernatant containing soluble factors was discarded and the extracellular vesicle pellet was
resuspended in 1 mL of filtered 1x PBS for nanoparticle tracking analysis.

Nanoparticle tracking analysis of extracellular vesicles

676   Isolated extracellular vesicles were quantified via nanoparticle tracking analysis (NTA) using a Malvern
677   NanoSight NS300. Samples were passed through the NanoSight flow cell using the included syringe pump

678   at a flow rate of 50. At least 10 videos, each 30 seconds in length, were acquired with a 488 nm laser per
679   sample. The concentration and size distribution of vesicles in each sample were then calculated using the
680   Malvern Nanoparticle Tracking Analysis Software.
681
Western blotting of vesicle markers
Sorted adipocytes for each condition, as well as preparations of the extracellular vesicles these cells
generated, were lysed using lysis buffer (25 mM Tris, 100 mM NaCl, 1% Triton X-100, 1 mM EDTA, 1.0 μg/
mL each of aprotinin and leupeptin, 1.0 mM β-glycerol phosphate, and 1.0 mM DTT). Protein concentrations
of the cell (whole cell lysates; WCLs) and extracellular vesicle lysates were determined using the Bio-Rad
Protein Assay Dye (Bio-Rad) and a spectrophotometer at a wavelength of 595. The lysates were normalized
based on protein concentration and resolved on 4-20% SDS-PAGE gels (Invitrogen). The gels were
transferred to PVDF membranes (Thermo-Fisher Scientific), and the membranes were blocked in 5.0%
bovine serum albumin (BSA) in TBST buffer (20 mM Tris, 135 mM NaCl, and 0.02% Tween) for 1 hour at
room temperature. The membranes were then incubated with one of the following primary antibodies diluted
in TBST overnight at 4°C: FAK antibody (1:1000 dilution; Cell Signaling Technologies, #3285) and FLOT2
antibody (1:1000 dilution; Cell Signaling Technologies, #3436). The next day, the membranes were washed
3 times for 5 minutes with TBST before being incubated with HPR-conjugated secondary antibodies diluted
in TBST for 1 hour at room temperature. The membranes were again washed with TBST 3 times for 5
minutes and exposed to ECL reagents (Bio-Rad). Images of the membranes were obtained using X-ray film
(The Lab Depot).
698
Breast cancer cell treatment with adipocyte-derived vesicles
For studies requiring treatment with adipocyte-derived extracellular vesicles, conditioned media was
processed as described for analysis of vesicles through PEG 6000 precipitation and centrifugation at 3220 x
g. The precipitate was then resuspended in 2 mL of exosome-depleted DMEM/F12 instead of 1x PBS. This
solution was ultracentrifuged in a TLS-55 swinging-bucket rotor (Beckman Coulter) at 100,000 x g at 4°C for
4 hours. The resulting extracellular vesicle pellet was resuspended in 2 mL of exosome-depleted DMEM/F12
for downstream use.
706
Extracellular vesicle treatment and DGAT1/2 inhibitor studies
For studies of lipid uptake from adipocyte-derived extracellular vesicles, 30,000 MDA-MB-231 breast cancer
cells were plated per well in a 24-well plate containing 12 mm glass coverslips coated with 30 μg/mL collagen
type I (VWR #47747-218). The following day, vesicles were applied to the cancer cells with 40 μM iDGAT1
(PF 04620110) (Tocris #1109276-89-2) and 40 μM iDGAT2 (PF 06424439) (Tocris #1469284-79-4) or an
equivalent volume of DMSO vehicle. Treatment was applied such that each glass coverslip seeded with
breast cancer cells received vesicles collected from one adipocyte-collagen gel over 24 hours. Breast cancer
cells were incubated with vesicles for 24 hours at 37°C and then fixed with 4% (w/v) paraformaldehyde in 1x
PBS for analysis of lipid uptake via fluorescence microscopy.
716
Lipid extraction and derivatization for lipidomics
100 μL of flash-frozen adipocytes were lysed in 1 mL 66% methanol (v/v) in 0.1 M potassium phosphate
buffer (pH 6.8) by vortexing twice on high for 20 seconds each. A 10 μL aliquot from each lysate was reserved
for total protein quantification using BCA assay (Thermo Fisher, #23225). The remaining lysate was
transferred to 13 x 100 mm borosilicate glass tubes (Fisher Scientific, #14-961-27), acidified with 10 μL of
1N HCl, and briefly vortexed. Total lipids were extracted twice with 1 mL of isooctane:ethyl acetate (3:1 v/v)
and once with 1 mL hexanes. Each extraction was followed by 10-15 seconds of vortexing and then
centrifugation at 2000 x g for 1 minute to ensure phase separation. The upper organic phases from all
extractions were combined in fresh 13 x 100 mm borosilicate glass tubes, evaporated under gaseous
nitrogen, and resuspended in 300 μL isooctane. For the non-esterified fatty acid (NEFA) fraction, 100 μL of

the total lipid extract was transferred to a new glass tube, 25 ng internal stable isotope standard was added,
evaporated under nitrogen, and resuspended in 25 µL of 1% pentafluorobenzyl bromide in acetonitrile (v/v).
An equal volume of 1% diisopropylethylamine in acetonitrile (v/v) was added, followed by a 30-minute
incubation at 25°C. The resulting pentafluorobenzyl-fatty acid derivatives were dried under nitrogen,
resuspended in 100 µL of hexanes, and prepared for injection into the GC-MS. For the total fatty acid (TFA)
fraction, 10 µL of the total lipid extract was transferred to a Teflon-lined glass tube, mixed with 75 ng of the
stable isotope internal standard, and dried under nitrogen. Samples were resuspended in 500 µL of ethanol,
and 500 µL of 1 M NaOH was added to saponify over 30 minutes at 90°C. After cooling, samples were
acidified with 550 µL of 1 M HCl and extracted twice with 1.5 mL of hexanes. The extracts were dried under
nitrogen and derivatized as described above. TFA samples were resuspended in 300 µL of hexanes for GC-
MS injection.
738
GC-MS conditions and data acquisition
For both the NEFA and TFA fractions, 1 µL of the pentafluorobenzyl-fatty acid derivatives was injected into
an Agilent 8890 GC system coupled to a 5977B MS detector, using a DB-1MS UI column (Agilent, #122-
0112UI). The GC temperature program consisted of an initial hold at 80°C for 3 minutes, followed by a ramp
of 30°C per min to 125°C, and a second ramp of 25°C per min to 320°C with a final hold of 2 minutes.
Methane was the carrier gas, flowing at 1.5 mL per min. Data were acquired in full-scan negative chemical
ionization mode, using m/z for acyl chain lengths ranging from 8 to 22 carbons and unsaturation. Peak areas
of analytes and internal standards were measured, and the ratio of analyte-derived ion areas to internal
standard areas was calculated for quantification relative to each standard curve(*47*, *48*).
748
Statistical analysis
All experiments were performed with at least three biological replicates unless otherwise noted. Data with
two conditions were evaluated with a nonparametric Mann-Whitney U test unless otherwise noted. Data with
three or more conditions were evaluated with either a standard or nested one-way ANOVA with Tukey multiple
comparisons test unless otherwise noted. P-values less than 0.05 were considered statistically significant.
Unless otherwise noted, all data are plotted as means with standard deviations. All statistical analysis was
performed using GraphPad Prism (v.10.2.3) or R (v.4.3.1).

References

1. Products - Data Briefs - Number 508 - September 2024 (2024). https://doi.org/10.15620/cdc/159281.

2. Obesity and Cancer Fact Sheet - NCI (2022). https://www.cancer.gov/about-cancer/causes-prevention/risk/obesity/obesity-fact-sheet.

3. P. González-Muniesa, M.-A. Mártinez-González, F. B. Hu, J.-P. Després, Y. Matsuzawa, R. J. F. Loos, L. A. Moreno, G. A. Bray, J. A. Martinez, Obesity. *Nature Reviews Disease Primers* **3**, 1–18 (2017).

4. B. A. Shook, R. R. Wasko, O. Mano, M. Rutenberg-Schoenberg, M. C. Rudolph, B. Zirak, G. C. Rivera-Gonzalez, F. López-Giráldez, S. Zarini, A. Rezza, D. A. Clark, M. Rendl, M. D. Rosenblum, M. B. Gerstein, V. Horsley, Dermal Adipocyte Lipolysis and Myofibroblast Conversion Are Required for Efficient Skin Repair. *Cell Stem Cell* **26**, 880-895.e6 (2020).

5. Q. Zhu, Y. Zhu, C. Hepler, Q. Zhang, J. Park, C. Gliniak, G. H. Henry, C. Crewe, D. Bu, Z. Zhang, S. Zhao, T. Morley, N. Li, D.-S. Kim, D. Strand, Y. Deng, J. J. Robino, O. Varlamov, R. Gordillo, M. G. Kolonin, C. M. Kusminski, R. K. Gupta, P. E. Scherer, Adipocyte mesenchymal transition contributes to mammary tumor progression. *Cell Reports* **40**, 111362 (2022).

6. M. P. Emont, C. Jacobs, A. L. Essene, D. Pant, D. Tenen, G. Colleluori, A. Di Vincenzo, A. M. Jørgensen, H. Dashti, A. Stefek, E. McGonagle, S. Strobel, S. Laber, S. Agrawal, G. P. Westcott, A. Kar, M. L. Veregge, A. Gulko, H. Srinivasan, Z. Kramer, E. De Filippis, E. Merkel, J. Ducie, C. G. Boyd, W. Gourash, A. Courcoulas, S. J. Lin, B. T. Lee, D. Morris, A. Tobias, A. V. Khera, M. Claussnitzer, T. H. Pers, A. Giordano, O. Ashenberg, A. Regev, L. T. Tsai, E. D. Rosen, A single-cell atlas of human and mouse white adipose tissue. *Nature* **603**, 926–933 (2022).

7. J. Bäckdahl, L. Franzén, L. Massier, Q. Li, J. Jalkanen, H. Gao, A. Andersson, N. Bhalla, A. Thorell, M. Rydén, P. L. Ståhl, N. Mejhert, Spatial mapping reveals human adipocyte subpopulations with distinct sensitivities to insulin. *Cell Metabolism*, doi: 10.1016/j.cmet.2021.07.018 (2021).

8. B. Dirat, L. Bochet, M. Dabek, D. Daviaud, S. Dauvillier, B. Majed, Y. Y. Wang, A. Meulle, B. Salles, S. Le Gonidec, I. Garrido, G. Escourrou, P. Valet, C. Muller, Cancer-associated adipocytes exhibit an activated phenotype and contribute to breast cancer invasion. *Cancer Res.* **71**, 2455–2465 (2011).

9. S. Liu, A. Benito-Martin, F. A. Pelissier Vatter, S. Z. Hanif, C. Liu, P. Bhardwaj, P. Sethupathy, A. R. Farghli, P. Piloco, P. Paik, M. Mushannen, X. Dong, D. M. Otterburn, L. Cohen, R. Bareja, J. Krumsiek, L. Cohen-Gould, S. Calto, J. A. Spector, O. Elemento, D. C. Lyden, K. A. Brown, Breast adipose tissue-derived extracellular vesicles from obese women alter tumor cell metabolism. *EMBO reports* **24**, e57339 (2023).

10. S. Balaban, R. F. Shearer, L. S. Lee, M. van Geldermalsen, M. Schreuder, H. C. Shtein, R. Cairns, K. C. Thomas, D. J. Fazakerley, T. Grewal, J. Holst, D. N. Saunders, A. J. Hoy, Adipocyte lipolysis links obesity to breast cancer growth: adipocyte-derived fatty acids drive breast cancer cell proliferation and migration. *Cancer Metab* **5**, 1 (2017).

11. J. Park, T. S. Morley, M. Kim, D. J. Clegg, P. E. Scherer, Obesity and cancer—mechanisms underlying tumour progression and recurrence. *Nature Reviews Endocrinology* **10**, 455–465 (2014).

12. M. M. M. Almekinders, M. Schaapveld, B. Thijssen, L. L. Visser, T. Bismeijer, J. Sanders, E. Isnaldi, I. Hofland, M. Mertz, L. F. A. Wessels, A. Broeks, E. Hooijberg, W. Zwart, E. H. Lips, C. Desmedt, J. Wesseling, Breast adipocyte size associates with ipsilateral invasive breast cancer risk after ductal carcinoma in situ. *npj Breast Cancer* **7**, 1–10 (2021).

13. A. L. Ghaben, P. E. Scherer, Adipogenesis and metabolic health. *Nat. Rev. Mol. Cell Biol.* **20**, 242–258 (2019).

14.   J. R. Acosta, I. Douagi, D. P. Andersson, J. Bäckdahl, M. Rydén, P. Arner, J. Laurencikiene, Increased fat cell
size: a major phenotype of subcutaneous white adipose tissue in non-obese individuals with type 2
diabetes. *Diabetologia* **59**, 560–570 (2016).

15.   N. M. Iyengar, K. A. Brown, X. K. Zhou, A. Gucalp, K. Subbaramaiah, D. D. Giri, H. Zahid, P. Bhardwaj, N. K.
Wendel, D. J. Falcone, H. Wang, S. Williams, M. Pollak, M. Morrow, C. A. Hudis, A. J. Dannenberg,
Metabolic Obesity, Adipose Inflammation and Elevated Breast Aromatase in Women with Normal Body
Mass Index. *Cancer Prevention Research* **10**, 235–243 (2017).

16.   M. J. Harms, Q. Li, S. Lee, C. Zhang, B. Kull, S. Hallen, A. Thorell, I. Alexandersson, C. E. Hagberg, X.-R.
Peng, A. Mardinoglu, K. L. Spalding, J. Boucher, Mature Human White Adipocytes Cultured under
Membranes Maintain Identity, Function, and Can Transdifferentiate into Brown-like Adipocytes. *Cell Rep*
27, 213-225.e5 (2019).

17.   J. Dufau, J. X. Shen, M. Couchet, T. De Castro Barbosa, N. Mejhert, L. Massier, E. Griseti, E. Mouisel, E.-Z.
Amri, V. M. Lauschke, M. Rydén, D. Langin, In vitro and ex vivo models of adipocytes. *American Journal of*
Physiology-Cell Physiology **320**, C822–C841 (2021).

18.   G. F. Beeghly, J. Deng, C. Fischbach, Protocol to fabricate elastomer microwells for three-dimensional
culture of primary adipocytes. *STAR Protoc* **5**, 103264 (2024).

19.   B. A. Cho, N. M. Iyengar, X. K. Zhou, M. Morrow, D. D. Giri, A. Verma, O. Elemento, M. Pollak, A. J.
Dannenberg, Blood biomarkers reflect the effects of obesity and inflammation on the human breast
transcriptome. *Carcinogenesis* **42**, 1281–1292 (2021).

20.   Y. Zhang, R. Proenca, M. Maffei, M. Barone, L. Leopold, J. M. Friedman, Positional cloning of the mouse
obese gene and its human homologue. *Nature* **372**, 425–432 (1994).

21.   K. Velickovic, H. A. L. Leija, B. Kosic, H. Sacks, M. E. Symonds, V. Sottile, Leptin deficiency impairs
adipogenesis and browning response in mouse mesenchymal progenitors. *European Journal of Cell*
Biology **102**, 151342 (2023).

22.   M. Jernås, J. Palming, K. Sjöholm, E. Jennische, P.-A. Svensson, B. G. Gabrielsson, M. Levin, A. Sjögren, M.
Rudemo, T. C. Lystig, B. Carlsson, L. M. S. Carlsson, M. Lönn, Separation of human adipocytes by size:
hypertrophic fat cells display distinct gene expression. *FASEB J* **20**, 1540–1542 (2006).

23.   Y. Hara, S. Wakino, Y. Tanabe, M. Saito, H. Tokuyama, N. Washida, S. Tatematsu, K. Yoshioka, K. Homma,
K. Hasegawa, H. Minakuchi, K. Fujimura, K. Hosoya, K. Hayashi, K. Nakayama, H. Itoh, Rho and Rho-
kinase activity in adipocytes contributes to a vicious cycle in obesity that may involve mechanical stretch.
Sci Signal **4**, ra3 (2011).

24.   B. Hansson, B. Morén, C. Fryklund, L. Vliex, S. Wasserstrom, S. Albinsson, K. Berger, K. G. Stenkula,
Adipose cell size changes are associated with a drastic actin remodeling. *Sci Rep* **9**, 12941 (2019).

25.   S. Wang, S. Cao, M. Arhatte, D. Li, Y. Shi, S. Kurz, J. Hu, L. Wang, J. Shao, A. Atzberger, Z. Wang, C. Wang,
W. Zang, I. Fleming, N. Wettschureck, E. Honoré, S. Offermanns, Adipocyte Piezo1 mediates obesogenic
adipogenesis through the FGF1/FGFR1 signaling pathway in mice. *Nat Commun* **11**, 2303 (2020).

26.   M. C. M. Aboy-Pardal, M. C. Guadamillas, C. R. Guerrero, M. Català-Montoro, M. Toledano-Donado, S.
Terrés-Domínguez, D. M. Pavón, V. Jiménez-Jiménez, D. Jimenez-Carretero, M. Zamai, C. Folgueira, A.
Cerezo, F.-N. Lolo, R. Nogueiras, G. Sabio, M. Sánchez-Álvarez, A. Echarri, R. Garcia, M. A. Del Pozo,
Plasma membrane remodeling determines adipocyte expansion and mechanical adaptability. *Nat*
Commun **15**, 10102 (2024).

27.   W. Xia, P. Veeragandham, Y. Cao, Y. Xu, T. E. Rhyne, J. Qian, C.-W. Hung, P. Zhao, Y. Jones, H. Gao, C.
Liddle, R. T. Yu, M. Downes, R. M. Evans, M. Rydén, M. Wabitsch, Z. Wang, H. Hakozaki, J. Schöneberg, S.
M. Reilly, J. Huang, A. R. Saltiel, Obesity causes mitochondrial fragmentation and dysfunction in white
adipocytes due to RalA activation. *Nat Metab* **6**, 273–289 (2024).

28.   F. Baldini, R. Fabbri, C. Eberhagen, A. Voci, P. Portincasa, H. Zischka, L. Vergani, Adipocyte hypertrophy
parallels alterations of mitochondrial status in a cell model for adipose tissue dysfunction in obesity. *Life*
Sciences **265**, 118812 (2021).

29.   L. Ling, J. A. Mulligan, Y. Ouyang, A. A. Shimpi, R. M. Williams, G. F. Beeghly, B. D. Hopkins, J. A. Spector, S.
G. Adie, C. Fischbach, Obesity-Associated Adipose Stromal Cells Promote Breast Cancer Invasion
through Direct Cell Contact and ECM Remodeling. *Advanced Functional Materials* **n/a**, 1910650 (2020).

30.   B. R. Seo, X. Chen, L. Ling, Y. H. Song, A. A. Shimpi, S. Choi, J. Gonzalez, J. Sapudom, K. Wang, R. C. A.
Eguiluz, D. Gourdon, V. B. Shenoy, C. Fischbach, Collagen microarchitecture mechanically controls
myofibroblast differentiation. *PNAS*, doi: 10.1073/pnas.1919394117 (2020).

31.   S. Y. Loo, L. P. Toh, W. H. Xie, E. Pathak, W. Tan, S. Ma, M. Y. Lee, S. Shatishwaran, J. Z. Z. Yeo, J. Yuan, Y. Y.
Ho, E. K. L. Peh, M. Muniandy, F. Torta, J. Chan, T. J. Tan, Y. Sim, V. Tan, B. Tan, P. Madhukumar, W. S. Yong,
K. W. Ong, C. Y. Wong, P. H. Tan, Y. S. Yap, L.-W. Deng, R. Dent, R. Foo, M. R. Wenk, S. C. Lee, Y. S. Ho, E. H.
Lim, W. L. Tam, Fatty acid oxidation is a druggable gateway regulating cellular plasticity for driving
metastasis in breast cancer. *Science Advances* **7**, eabh2443 (2021).

32.   I. Lazar, E. Clement, S. Dauvillier, D. Milhas, M. Ducoux-Petit, S. LeGonidec, C. Moro, V. Soldan, S. Dalle,
S. Balor, M. Golzio, O. Burlet-Schiltz, P. Valet, C. Muller, L. Nieto, Adipocyte Exosomes Promote Melanoma
Aggressiveness through Fatty Acid Oxidation: A Novel Mechanism Linking Obesity and Cancer. *Cancer*
Res **76**, 4051–4057 (2016).

33.   M. Zhang, J. S. D. Martino, R. L. Bowman, N. R. Campbell, S. C. Baksh, T. Simon-Vermot, I. S. Kim, P.
Haldeman, C. Mondal, V. Yong-Gonzales, M. Abu-Akeel, T. Merghoub, D. R. Jones, X. G. Zhu, A. Arora, C. E.
Ariyan, K. Birsoy, J. D. Wolchok, K. S. Panageas, T. Hollmann, J. J. Bravo-Cordero, R. M. White, Adipocyte-
Derived Lipids Mediate Melanoma Progression via FATP Proteins. *Cancer Discov* **8**, 1006–1025 (2018).

34.   S. E. Flaherty, A. Grijalva, X. Xu, E. Ables, A. Nomani, A. W. Ferrante, A lipase-independent pathway of lipid
release and immune modulation by adipocytes. *Science* **363**, 989–993 (2019).

35.   Y. Y. Wang, C. Attané, D. Milhas, B. Dirat, S. Dauvillier, A. Guerard, J. Gilhodes, I. Lazar, N. Alet, V. Laurent,
S. Le Gonidec, D. Biard, C. Hervé, F. Bost, G. S. Ren, F. Bono, G. Escourrou, M. Prentki, L. Nieto, P. Valet, C.
Muller, Mammary adipocytes stimulate breast cancer invasion through metabolic remodeling of tumor
cells. *JCI Insight* **2**, e87489 (2017).

36.   E. Sezgin, I. Levental, S. Mayor, C. Eggeling, The mystery of membrane organization: composition,
regulation and roles of lipid rafts. *Nat Rev Mol Cell Biol* **18**, 361–374 (2017).

37.   H. Yoon, J. L. Shaw, M. C. Haigis, A. Greka, Lipid metabolism in sickness and in health: Emerging
regulators of lipotoxicity. *Mol Cell* **81**, 3708–3730 (2021).

38.   R. Rudalska, J. Harbig, M. T. Snaebjornsson, S. Klotz, S. Zwirner, L. Taranets, F. Heinzmann, T.
Kronenberger, M. Forster, W. Cui, L. D'Artista, E. Einig, M. Hinterleitner, W. Schmitz, A. Dylawerska, T.-W.
Kang, A. Poso, M. T. Rosenfeldt, N. P. Malek, M. Bitzer, S. Laufer, B. J. Pichler, N. Popov, A. Schulze, L.
Zender, D. Dauch, LXRα activation and Raf inhibition trigger lethal lipotoxicity in liver cancer. *Nat Cancer*
2, 201–217 (2021).

39. Y. Tan, J. Li, G. Zhao, K.-C. Huang, H. Cardenas, Y. Wang, D. Matei, J.-X. Cheng, Metabolic reprogramming
from glycolysis to fatty acid uptake and beta-oxidation in platinum-resistant cancer cells. *Nat Commun*
13, 4554 (2022).

40. F. Poggio, M. Bruzzone, M. Ceppi, N. F. Pondé, G. La Valle, L. Del Mastro, E. de Azambuja, M. Lambertini,
Platinum-based neoadjuvant chemotherapy in triple-negative breast cancer: a systematic review and
meta-analysis. *Annals of Oncology* **29**, 1497–1508 (2018).

41. M. Galarraga, J. Campión, A. Muñoz-Barrutia, N. Boqué, H. Moreno, J. A. Martínez, F. Milagro, C. Ortiz-de-
Solórzano, Adiposoft: automated software for the analysis of white adipose tissue cellularity in
histological sections. *J Lipid Res* **53**, 2791–2796 (2012).

42. K. Sieckmann, N. Winnerling, M. Huebecker, P. Leyendecker, D. Juliana Silva Ribeiro, T. Gnad, A. Pfeifer, D.
Wachten, J. N. Hansen, AdipoQ—a simple, open-source software to quantify adipocyte morphology and
function in tissues and in vitro. *Mol Biol Cell* **33**, br22 (2022).

43. N. M. Iyengar, P. G. Morris, X. K. Zhou, A. Gucalp, D. Giri, M. D. Harbus, D. J. Falcone, M. D. Krasne, L. T.
Vahdat, K. Subbaramaiah, M. Morrow, C. A. Hudis, A. J. Dannenberg, Menopause is a determinant of
breast adipose inflammation. *Cancer Prev Res (Phila)* **8**, 349–358 (2015).

44. P. Bankhead, M. B. Loughrey, J. A. Fernández, Y. Dombrowski, D. G. McArt, P. D. Dunne, S. McQuaid, R. T.
Gray, L. J. Murray, H. G. Coleman, J. A. James, M. Salto-Tellez, P. W. Hamilton, QuPath: Open source
software for digital pathology image analysis. *Sci Rep* **7**, 16878 (2017).

45. K. Sieckmann, N. Winnerling, M. Huebecker, P. Leyendecker, D. Juliana Silva Ribeiro, T. Gnad, A. Pfeifer, D.
Wachten, J. N. Hansen, AdipoQ—a simple, open-source software to quantify adipocyte morphology and
function in tissues and in vitro. *MBoC* **33**, br22 (2022).

46. M. A. Rider, S. N. Hurwitz, D. G. Meckes, ExtraPEG: A Polyethylene Glycol-Based Method for Enrichment of
Extracellular Vesicles. *Sci Rep* **6**, 23978 (2016).

47. M. C. Rudolph, N. K. Maluf, E. A. Wellberg, C. A. Johnson, R. C. Murphy, S. M. Anderson, Mammalian Fatty
Acid Synthase Activity From Crude Tissue Lysates Tracing [13C]-Substrates Using GC Mass Spectrometry.
Anal Biochem **428**, 158–166 (2012).

48. R. Varshney, S. Das, G. D. Trahan, J. W. Farriester, G. P. Mullen, G. Kyere-Davies, D. M. Presby, J. A. Houck,
P. G. Webb, M. Dzieciatkowska, K. L. Jones, M. S. Rodeheffer, J. E. Friedman, P. S. MacLean, M. C.
Rudolph, Neonatal intake of Omega-3 fatty acids enhances lipid oxidation in adipocyte precursors.
iScience **26**, 105750 (2023).