Rare mutations implicate CGE interneurons as a vulnerable axis of cognitive deficits across psychiatric disorders

Stephanie A. Herrlinger[1,2]*, Jiayao Wang[3]*, Bovey Y Rao[1,2,4], Jonathan Chang[3], Joseph A. Gogos [1,2,5,6,7]†, Attila Losonczy[1,2]† and Dennis Vitkup[3]†

[1] Department of Neuroscience, Columbia University, New York, NY, United States
[2] Mortimer B. Zuckerman Mind Brain Behavior Institute, Columbia University, New York, NY, United States
[3] Department of Systems Biology and Department Biomedical Informatics, Columbia University, New York, NY, United States
[4] Doctoral Program in Neurobiology and Behavior, Columbia University, New York, NY, United States
[5] Department of Physiology, Columbia University, New York, NY, United States
[6] Department of Psychiatry, Columbia University, New York, NY, United States
[7] Stavros Niarchos Foundation Center for Precision Psychiatry and Mental Health, Columbia University, New York, NY, United States
*These authors contributed equally to this work
† Correspondence should be addressed to JG (jag90@columbia.edu), AL (al2856@columbia.edu) and DV (dv2121@cumc.columbia.edu)

Abstract

Neuropsychiatric disorders such as autism spectrum disorder (ASD) and schizophrenia (SCZ) share genetic risk factors, including rare high penetrance single nucleotide variants and copy number variants (CNVs), and exhibit both overlapping and distinct clinical phenotypes. Cognitive deficits and intellectual disability—critical predictors of long-term outcomes—are common to both conditions. To investigate shared and disorder-specific neurobiological impact of highly penetrant rare mutations in ASD and SCZ, we analyzed human single-nucleus whole-brain sequencing data to identify strongly affected brain cell types. Our analysis revealed Caudal Ganglionic Eminence (CGE)-derived GABAergic interneurons as a key nexus for cognitive deficits across these disorders. Notably, genes within 22q11.2 deletions, known to confer a high risk of SCZ, ASD, and cognitive impairment, showed a strong expression bias toward vasoactive intestinal peptide-expressing cells (VIP+) among CGE subtypes. To explore VIP+ GABAergic interneuron perturbations in the 22q11.2 deletion syndrome *in vivo,* we examined their activity in the *Df(16)A^{+/-}* mouse model during a spatial navigation task and observed reduced activity along with altered responses to random rewards. At the population level, VIP+ interneurons exhibited impaired spatial encoding and diminished subtype-specific activity suggesting deficient disinhibition in CA1 microcircuits in the hippocampus, a region essential for learning and memory. Overall, these results demonstrate the crucial role of CGE-derived interneurons in mediating cognitive processes that are disrupted across a range of psychiatric and neurodevelopmental disorders.

Introduction

Psychiatric disorders exhibit distinct clinical symptoms yet frequently share genetic risk factors
and underlying cellular and molecular mechanisms[1–6]. For example, schizophrenia (SCZ) is
characterized by psychosis, including hallucinations and delusions, as well as cognitive control
impairments leading to disorganization, impulsivity, and difficulties in working memory. In contrast,
autism spectrum disorder (ASD) is characterized by restricted interests and heightened anxiety,
along with cognitive rigidity and repetitive behaviors[7,8]. Several copy number variations (CNVs),
such as the 22q11.2 deletion, have been implicated in both SCZ and ASD, as well as in other
psychiatric disorders[9,10]. Among their diverse symptoms, cognitive impairments and intellectual
disability (ID), which are frequently comorbid with neuropsychiatric phenotypes[11] and are
especially pronounced in cases with high genetic risks[11–14]. Cognitive deficits in psychiatric
disorders are particularly debilitating, as their severity strongly predicts long-term health deficits
and well-being outcomes[15–17]. These deficits encompass a broad range of functions including
memory, attention, reasoning, decision-making and social cognition[18].

Despite advances in genetic research on disorders such as ASD and SCZ[1,2,19], therapeutic
progress in treating cognitive symptoms remains limited, largely due to gaps in understanding the
neurobiological mechanisms through which genetic risk factors impact cognition. To better
understand the relationships between genetics and psychiatric phenotypes, researchers are
increasingly integrating genetic analyses with single-cell and single-nucleus sequencing data to
identify cell types affected across psychiatric disorders[20–22]. Animal models, particularly mouse
models that replicate rare high-risk genetic insults, play a critical role in this integrative approach.
The evolutionary conservation of brain regions, cell types, and core biological processes between
mice and humans[23–26] facilitates detailed examination of genetic risk factors at both cellular and
circuit levels[27–32].

In this study we investigated the collective neurobiological impact of rare, highly penetrant
mutations in ASD, SCZ, and 22q11.2 deletion syndrome across diverse brain cell types, using
recently released human single-nucleus whole-brain RNA sequencing data[33] (Fig. 1). Our
analyses revealed distinct disorder-specific mutation biases affecting different cell types, with
caudal ganglionic eminence (CGE)-derived interneurons consistently implicated across genetic
insults associated with cognitive dysfunction, including ASD with intellectual disability, SCZ, and
Developmental Disorders and Intellectual Disability (DD/ID). Furthermore, our analysis of genes
within the 22q11.2 deletion revealed that VIP+ CGE-derived interneurons were strongly affected
in this condition. Using *in vivo* two-photon calcium imaging of VIP+ GABAergic interneuron activity
in a mouse model of 22q11.2 deletion, we observed specific alterations in interneuron-selective
interneuron (ISI) cell types, which primarily inhibit other interneurons. These findings implicate ISI
dysfunction in circuit-level abnormalities contributing to cognitive and behavioral phenotypes
characteristic of this syndrome and likely many other psychiatric conditions.

88
Figure 1. Overview of study methodology. Left: Integration of genetic and single-cell transcriptomic data to identify
cell type-specific disorder biases. Top significant genes implicated in ASD and SCZ[1,2] and human brain snRNA-seq
data were used to create cell type mutation bias analysis using 461 human brain cell types[33]. **Middle:** Disorder cell
type bias analysis reveals shared cell type risk patterns between distinct disorders with overlapping cognitive
phenotypes. **Right:** Experimental validation in a relevant mouse model allows us to describe identified cell type specific
deficits *in vivo* using 2-photon calcium imaging to capture neuronal activity during behavior.
95

96 **Results**

97 **Identifying disorder cell type mutation bias leveraging rare variants and snRNA-seq data**
98

In our study we focused on rare mutations, which individually typically have substantially higher
functional impacts on psychiatric phenotypes compared to common variants[1,34]. The availability
of comparably-sized, well-characterized rare variant datasets from both ASD and SCZ cohorts
enabled us to perform direct cross-disorder comparisons. For ASD, we utilized a recent meta-
analysis performed by the SPARK consortium[2], which comprised *de novo* mutations from 42,607
ASD probands. This analysis identified 61 genes with exome-wide significance. For
schizophrenia, we used a case-control study involving 24,248 schizophrenia patients and 97,322
control patients from the SCHEMA consortium[1]. In our main analysis, we selected the top 61 SCZ
genes based on genetic association strength, to match the number of considered ASD genes.
108

A significant fraction of individuals in the ASD cohort exhibit low IQs (IQ ≤ 70, which we refer to
as "ASD with ID")[35,36] (Suppl. Fig. 1). To facilitate a more appropriate comparison with SCZ, where
the average premorbid IQ is approximately 94[37,38], we selected higher IQ (IQ >70) ASD cases
from the SPARK cohort (hereafter referred to as "ASD", with the average IQ ~95). Notably,
individuals with SCZ who carry rare mutations have IQ scores that are only marginally lower than
those in the broader schizophrenia population[39]. Furthermore, genes implicated in higher-IQ ASD
also share more similar developmental expression profiles with SCZ genes compared to genes
associated with lower-IQ ASD (Suppl. Fig. 2). Even though SCZ and ASD cases have comparable
overall IQs, the nature of cognitive dysfunction in these conditions is often distinct. While SCZ is
usually characterized by broad deficits across multiple cognitive domains, such as working
memory, processing speed, and executive function[40,41], high-functioning ASD cases tend to show
more selective, domain-specific cognitive impairments, particularly in areas like social cognition
and communication[42,43].
122

To investigate relative brain cell type impact in ASD and SCZ, we used a recent single-nucleus
sequencing dataset collected from the entire human brain[33]. This dataset encompassed
approximately 105 anatomical locations dissected from three different donors across the forebrain,
midbrain, and hindbrain, with 3,369,219 individual nuclei that were classified into 31 superclusters
and 461 distinct cell types (Fig. 1). To identify which brain cell types are most impacted by
disease-causing mutations, we calculated cell-type-specific mutation biases. Our goal was to
identify cell types where affected mutations showed strong expression specificity relative to other
cell types, indicating their likely crucial role in the disorder's pathophysiology. To quantify the
mutation biases, we first calculated the specificity of each gene's expression to a particular cell
type using a Z-score (see Methods). To account for overall expression differences across different
cell types, we then compared each gene's Z-score to the distribution of Z-scores for random genes
with similar brain expression levels. Finally, we weighted each ASD gene by *de novo* mutation
counts, and SCZ genes by their effective mutation counts in cases, due to differences in the
original genetic study designs (Suppl. Fig. 3, see Methods). The mutation bias towards each cell
types in ASD and SCZ was then calculated as the mutation-weighted average of expression
specificities across cell types for each disorder (Suppl. Table 2, Suppl. Fig. 4). This cell-type

mutation bias thus considers both the degree to which each disease gene is specifically
expressed in each cell type and the strength of its genetic contribution with the disorder.
141

Shared cell type vulnerability signatures across ASD and SCZ
Our analysis revealed a strong correlation between ASD and SCZ mutation biases across cell
type clusters in human brain cell atlas (Fig. 2A), with cell types showing strong mutation biases in
ASD also showing strongly mutation biases in SCZ (Spearman correlation = 0.66, $P<10^{-10}$). This
correlation remains strong when only considering neuronal cell types, indicating that it is not
simply due to a general neuronal bias in both disorders. The strong correlation is also not due to
gene overlapping between ASD and SCZ (Spearman correlation = 0.60, $P<10^{-10}$, after removing
3 overlapping genes). To understand what drives this shared pattern, we investigated the role of
gene constraint, i.e., the intolerance to mutational changes for genes frequently affected across
neuropsychiatric disorders[1,2,35,44,45]. When we progressively removed the most constrained genes
(measured by loss of function Z-score (LoF-Z)[46], which quantifies the deviation of observed likely
gene disrupting (LGD) mutations from expected, in healthy populations) from the gene sets
associated with both disorders, the correlation between ASD and SCZ rapidly decreased,
reaching zero after removing ~25 of the genes, while removing the least constrained genes had
no effect on the correlation (Fig. 2B). We then asked whether gene expression levels might also
explain the shared correlation of mutation biases between ASD and SCZ, as both disorders
involve genes highly expressed in the brain. Interestingly, in contrast to gene constraint, removing
highly expressed genes had little impact on the cell type bias correlation (Suppl. Fig. 5). These
findings suggest that highly constrained genes – those most resistant to (LGD) mutations in
healthy populations – tend to be more specifically expressed in cell types that are mutationally
biased in both disorders.
163

Several specific superclusters (hierarchically organized groups of brain cell type clusters sharing
transcriptional signatures in the human brain cell atlas) are strongly biased by mutations in both
ASD and SCZ, notably including excitatory neurons in the amygdala and hippocampus as well as
Deep-layer intratelencephalic (IT) neurons (Fig. 2C, Suppl. Fig. 6A-D). These findings are
consistent with several previous studies reporting frequent perturbations of these neurons in brain
disorders. For example, amygdalar neurons and their cortical and hippocampal projections are
implicated in a broad range of behavioral disturbances in both SCZ and ASD, including deficits in
emotion, motivation, and social interactions[47,48]. Additionally, the hippocampus is known to play a
key role in social and cognitive deficits in ASD[49], as well as in learning deficits and positive
symptoms associated with SCZ[50,51]. Deep-layer IT neurons, which are central to sensory
processing, decision-making, and learning, are also affected in both disorders[52]. Collectively,
these shared neuronal disruptions may underlie overlapping dysfunction of brain circuits
mediating disorders' phenotypes[53].
177

Distinct cell type biases between ASD and SCZ

To identify disorder-specific mutational biases at the supercluster level, we assessed the
differential bias between ASD and SCZ, quantified by the average difference in mutation bias
across cell type clusters within a supercluster. This measure describes how strongly each
particular supercluster is preferentially affected in one disorder compared to the other, which may

underlie the unique physiological and behavioral phenotypes associated with each disorder. We
observed that mutations in ASD and SCZ are preferentially biased toward several specific neuron
types (Fig. 2C). Among them, medium spiny neurons (MSNs) showed stronger differential biases
in ASD compared to SCZ (Fig. 2D). Nearly all MSN cell types exhibited stronger ASD mutation
biases, with a few also exhibiting SCZ bias. This agrees with previous findings implicating MSNs
in ASD[54] and multiple experimental studies[55,56]. In contrast, GABAergic interneurons, including
MGE-, CGE-derived, and Lamp5-LHX6/Chandelier interneurons showed the strongest differential
biases towards SCZ compared to ASD (Fig. 2 CEF). As mentioned above, while overall IQ may
be similar between SCZ and ASD cohorts, cognitive impairment is thought to be widespread in
schizophrenia while cognitive symptoms in high-functioning ASD are often more localized and
specific. Considering this, we explored what neuron types are more affected in ASD probands
with ID compared to ASD probands without ID. This analysis showed that, CGE interneurons
were also showed stronger differential biases in ASD with ID (Fig. 2G, Suppl. Fig 6G), revealing
the potential role of CGE cell types in cognitive dysfunctions observed in both ASD with ID and
SCZ.
198

Figure 2. ASD and SCZ Mutation bias towards different brain cell types. A. Scatter plot showing correlation of mutation biases between ASD and SCZ across all cell types. Neuronal (blue) and non-neuronal (brown) cell types are indicated. Strong correlation is observed for both all cell types (Spearman R= 0.66) and neuron-specific analysis (Spearman R = 0.62). **B.** Spearman correlation of cell type bias between ASD and SCZ genes, assessed by sequential gene removal. The red line represents the bias correlation when removing genes with the lowest constraint (lofZ) scores, while the blue line represents the correlation when removing genes with the highest constraint scores. The gray line shows the average correlation for random gene removals, with error bars indicating ±1 standard deviation from the mean of random removals. **C.** Bar plot comparing mutation biases across superclusters between ASD (blue) and SCZ (orange). Superclusters showing positive mutation bias for at least one disorder are displayed in the figure and sorted by average differential bias between ASD and SCZ. Error bars represent standard error cross cluster biases from corresponding supercluster. Asterisks indicate significance level between groups (*: P < 0.05. **: P < 0.01. ***: P < 0.001, ****: P < 0.0001, ns=not significant). Detailed data can be found in Supplementary table 4. **D-G.** Detailed comparison of mutation biases in specific neuron superclusters, each point representing a cell type cluster from the corresponding supercluster, P-values were calculated based on Mann–Whitney U test with false discovery rate (FDR) correction. **(D)** Medium spiny neurons show stronger bias in ASD (adjusted P = 1.1e-02), **(E)** MGE interneurons exhibit stronger bias in SCZ (adjusted P = 9.1e-06), **(F)** CGE interneurons display bias toward SCZ (adjusted P = 1.8e-04),

216 and **(G)** CGE interneurons show stronger bias when comparing ASD with ID to ASD (adjusted P = 7.9e-05). Detailed
217 data can be found in Supplementary table 2 and 3.
218
219

VIP+ interneurons are strongly affected in multiple psychiatric phenotypes with cognitive impairments.

To further explore the involvement of CGE interneurons in cognitive function, we analyzed
additional cohorts with cognitive phenotypes (Fig. 3A). In the Developmental Disorders and
Intellectual Disability (DD/ID) cohort[45], which is characterized by severe cognitive impairments,
we also observed a strong bias towards CGE interneurons (Fig. 3A, grey, Suppl. Fig. 6F).
Furthermore, analysis of the UK Biobank rare protein-coding variants associated with cognitive
functions, including Verbal-Numerical Reasoning (VNR), Reaction Time (RT), and Educational
Attainment (EDU))[57], revealed that genes negatively associated with VNR (mutations in these
genes tend to lead to worse VNR performance) exhibited a pronounced bias toward CGE
interneurons compared to genes positively associated with VNR (Suppl. Fig. 6E). The consistent
strong CGE interneuron mutation bias across different cohorts suggests that these neurons play
a key role in cognitive processes. In contrast, MGE and Lamp5-LHX6/Chandelier interneurons
did not demonstrate comparably strong mutation biases across the cognitive phenotype cohorts
we examined (Suppl. Fig. 7).
235

To quantify the relationship between expression biases toward different human brain cell types
and cognitive function, we calculated a Phenotype Bias Sensitivity (PBS) score for each cell type
using data for ASD probands with *de novo* mutations and corresponding full scale IQ (FSIQ)
measured in the SPARK and ASC cohorts[2,58] (Suppl. Fig. 1). For a given cell type, PBS was
calculated as the linear regression slope between the genes' expression bias toward that cell type
and the average FSIQ of probands carrying *de novo* mutations in the gene (Suppl. Fig. 1B, Fig.
3C). In this way, more negative PBS values indicate that stronger gene expression biases towards
a cell type are associated with lower probands' IQ. The statistical significance of PBS for cell
clusters was determined by a permutation approach, where we randomly reassigned FSIQ across
probands and computed PBS for each permuted set, to create a null distribution (see Methods).
The PBS calculations showed that the three major groups of interneurons, including CGE-derived
interneurons, LAMP5-LHX6/Chandelier cells, and MGE-derived interneurons, exhibited the
strongest PBS values (Fig. 3B). Further analysis of CGE interneuron subtypes revealed that
vasoactive intestinal peptide-expressing (VIP+) interneurons consistently demonstrated
significantly stronger PBS values (Fig. 3D). This suggests that mutations affecting VIP+
interneurons may be particularly important in determining cognitive outcomes in ASD probands.
CGE-derived VIP+ interneurons modulate neural circuits by reducing inhibition and enhancing
excitatory activity[59,60]—an essential mechanism for cognitive functions such as perception,
attention, learning, and memory[61]. They achieve this by suppressing Sst+, Cck+, and possibly
Pvalb+ interneurons[62,63], thereby indirectly amplifying pyramidal neuron activity. Moreover, VIP+
interneurons interact with cholinergic, serotonergic, and dopaminergic systems[63–68] to maintain
the excitation-inhibition balance critical for network stability and effective cognitive processing[61].
258

259
Figure 3. CGE interneurons are impacted in multiple cohorts with cognitive abnormalities. A. CGE interneuron
mutation biases across different cohorts shows increasing bias in disorders with cognitive deficits. Each dot represents
a CGE interneuron subtype. Comparison includes ASD (without ID) (adjusted P = 1), verbal-numerical reasoning (VNR-
: adjusted P = 3.3e-09, VNR+: adjusted P = 1), schizophrenia (SCZ) (adjusted P = 0.19), ASD with ID (adjusted P =
7.4e-03), and developmental disorders/intellectual disability (DD/ID) (adjusted P = 7.5e-08). P values were calculated
with Mann-Whitney U test with FDR correction. **B.** Phenotype Bias Sensitivity (PBS) analysis across neuronal cell types
reveals strongest correlation with Full Scale IQ (FSIQ). Negative PBS values indicate genes with stronger cell type bias
are associated with lower FSIQ scores. CGE interneuron average PBS=-6.78, adjusted P=0.014, LAMP5-LHX6 and
Chandelier, average PBS=-6.65, adjusted P=0.03. P values were calculated with mutation permutation test with FDR
correction. Detailed data can be found in supplementary table 5. **C.** Association between cell type bias and FSIQ for
CGE interneuron cluster 295 (VIP+), which is the cell type has strongest PBS. Each dot represents a High confident
ASD gene, with FSIQ averaged across multiple probands with mutation from the gene shown on the y-axis. The x-axis
displays each gene's bias towards cell type 295. **D.** Comparison of PBS values between VIP+ and VIP- CGE
interneuron subtypes shows stronger association with cognitive phenotypes in VIP+ interneurons (PBS difference = -
1.96, P = 9.7e-03, Mann-Whitney U test).
275

VIP+ interneurons are strongly affected by 22q11.2 deletions

Copy number variants (CNVs) consistently stand out as highly penetrant genetic risks for SCZ
and ASD[11,32,69,70]. The 22q11.2 deletion syndrome is a neurodevelopmental disorder marked by a
significantly elevated risk of schizophrenia in adulthood, along with comorbidities such as ASD
and intellectual disability[10,71]. Cognitive impairment is a key feature of 22q11.2 deletion syndrome,
with nearly half of affected individuals displaying an IQ below 70[72,73]. Consistent with most
disease-predisposing CNVs, the full range of phenotypic outcomes and disease symptoms results
from the synergistic effects of multiple genes within the 22q11.2 deletion[10]. Given the significant
SCZ risk and cognitive disability phenotypes associated with the 22q11.2 deletion we investigated

next the bias toward CGE and other interneurons for genes in the deletions. The 22q11.2
deletions occur in two primary forms: a 3 Mb deletion (46 genes) and a smaller 1.5 Mb deletion
(28 genes). Analysis of cell type biases for genes in the 3 Mb deletion (Suppl. Fig. 8) revealed a
substantial impact toward CGE-derived interneurons compared to other major interneuron types
(Fig. 4A), suggesting a crucial link to cognitive function and IQ. Notably, among CGE interneuron
cell type clusters, VIP+ interneurons exhibited significantly stronger biases compared to biased
toward VIP- CGE cell types (Fig. 4B). This pattern remains consistent when examining genes
within the 1.5 Mb deletion. This selective impact on VIP+ interneurons over other CGE interneuron
types suggests that 22q11.2 deletion may exert a particularly pronounced impact on the VIP+
interneurons, potentially contributing to the severe cognitive and behavioral phenotypes
associated with 22q11.2 deletion syndrome.

Figure 4. VIP+ CGE interneurons are preferentially affected by 22q.11.2 deletion genes. (A) Boxplot showing
expression biases for three major interneuron types (CGE interneurons, Lamp5/LHX6 and Chandelier cells, and MGE
interneurons) in the 22q11.2 deletion region. CGE interneurons exhibit the strongest bias compared to other interneuron
populations (P-values shown for pairwise comparisons, Mann-Whitney U-test). Complete data can be found in
Supplementary table 2**. (B)** The CGE interneuron cell type bias crosses different gene sets. including complete 22q.11.2
genes, genes included in 1.5MB deletion which equivalent to mouse 16qA13 deletion. VIP+ CGE types shown in red
and VIP- CGE types shown in blue. Difference between VIP+ and VIP- bias is significant for all 22q.11.2 genes and
suggestive for 1.5MB deletion genes (P=2.3e-03 and 6.7e-02, respectively, Mann-Whitney U-test.)

Altered VIP+ interneuron activity and population specificity in a mouse model for the
22q11.2 deletion syndrome
Having identified a robust genetic association between mutation biases toward CGE and VIP+
cell types and cognitive phenotypes across multiple genetic disorders, including the 22q11.2
deletion syndrome, we next experimentally investigated whether and how this bias is reflected in
altered function and integration of these interneurons within hippocampal microcircuits. To this
end we used the *Df(16)A*[+/-] mouse model, which faithfully recapitulates the human 1.5 Mb 22q11.2
deletion[9,29,31,74]. This model, known for deficits in working and episodic memory, synaptic

plasticity, and cortical and hippocampal development, has proven invaluable for uncovering novel biological pathways relevant to cognitive disfunction and psychiatric disorders[10,31,32,71,74–77]. By complementing our computational analyses with in vivo assessments, we examined the physiological impact of the 22q11.2 deletion on specific CGE/VIP+ subtypes—beyond the resolution of our computational methods—thus advancing our understanding of the cellular mechanisms by which CGE-derived interneurons mediate cognitive processes are disrupted in the 22q11.2 deletion syndrome.

To examine VIP+ interneuron activity *in vivo* we crossed *Df(16)A$^{+/-}$* mice with VIP-Cre mice and locally injected an adenovirus carrying a floxed GCaMP8m or GCaMP7f calcium indicator into CA1 to selectively express GCaMP in VIP+ interneurons (Suppl. Fig. 9A). Using two-photon three-dimensional acousto-optic deflector (3D-AOD)[78–80] microscopy, we then recorded VIP+ interneuron activity in a 3D volume within the dorsal hippocampal CA1 region (Suppl. Fig. 9B/C/D). *Post hoc* immunohistochemistry[28,78,81] (see Methods) allowed us to identify four major VIP+ interneuron subtypes: interneuron-selective interneurons type 2 (ISI2), interneuron-selective interneurons type 3 (ISI3), VIP+ cholecystokinin basket cells (CCKBCs), and a group of VIP-expressing cells, referred to as "VIP+ only" (VIPo) cells, that were immunonegative for all other tested markers[59,82,83]. The distribution of these subtypes across hippocampal layers did not significantly differ between wild type and *Df(16)A$^{+/-}$* mice (Suppl. Fig. 9G/H), and comparable numbers of each interneuron subtype were recorded from both wild type and mutant genotype (Suppl. Fig. 9I). Thus our analysis did not find any evidence suggesting the 22q11.2 deletion affects the number or proportion of VIP+ interneuron subtypes in the hippocampal CA1 region.

As was established previously, global interneuron activity is strongly driven by ambulatory state[84–87] and in treadmill-based experiments we reliably observed that locomotion is the greatest predictor of global interneuron activity[28,78,88]. To investigate locomotion-related activity of hippocampal VIP+ interneurons, mice were next trained to run on a head-fixed, cue-rich 2-meter treadmill for randomly distributed water rewards while undergoing calcium imaging (Fig. 5A)[28,78,81]. Behavioral metrics, including laps run, licking, and region occupancy in the belt during the 10-minute trials, were similar across genotypes (Suppl. Fig. 9J-M). Expected locomotion-related activity patterns were preserved across genotypes, with ISI3, ISI2, and VIPo subtypes positively correlated with movement, and VIP+ CCKBCs anticorrelated (Fig. 5B)[78,89]. However, ISI3 and ISI2 cell activity exhibited a significantly reduced correlation with locomotion (Fig. 5C), and ISI3, ISI2, and VIPo interneurons showed visibly reduced calcium transient amplitudes in *Df(16)A$^{+/-}$* mice (Fig. 5B). Due to the spike-like activity of ISI3, ISI2, and VIPo cells we were able to perform transient waveform analysis (see Methods[90]). This analysis confirmed reduced amplitudes in ISI3, ISI2, and VIPo cells (Fig. 5D-J), while VIPo cell transients uniquely occurred at a higher frequency (Fig. 5J). Reward responses, accessed by Peri-event Time Averages (PETA, see Methods), exhibited genotype-dependent differences, including reduced VIPo response amplitudes and enhanced ISI2 and ISI3 response amplitudes (Fig. 5K,L), suggesting that ISIs in mutant mice exhibit heightened sensitivity to random salient features[88]. Overall, these results show that VIP+ interneurons in the hippocampal CA1 region exhibit reduced *in vivo* activity dynamics in the 22q11.2 deletion model and that reward responses are significantly altered among specific VIP+ subtypes, demonstrating physiological consequences of the genetic perturbation inferred by our computational analysis.

Figure 5: Baseline changes in VIP+ cell type calcium dynamics in mutant mice. A. Imaging and behavior schematic: mice are trained to run on a cue-rich belt to expect random water rewards to be deposited across the belt (bottom) while head-fixed and imaged with 2-photon 3D-AOD microscopy (top). **B.** Example traces from all four VIP+ cell types detected in a single session from a Wildtype littermate (left) and a *Df(16)A+/-* (right) mouse. **C.** Pearson correlation coefficients of mouse velocity against average subtype calcium activity. **D.** Waveform comparison of average

detected calcium transient from wildtype littermates (black) and *Df(16)A$^{+/-}$* (brown) ISI3 cells. **E.** Average amplitude of
detected ISI3 waveforms compared between genotypes. **F.** Waveform comparison of average detected calcium
transients from wildtype littermates (black) and Df(16)A$^{+/-}$ (brown) ISI2 cells. **G.** Average amplitude of detected ISI2
waveforms compared between genotypes. **H.** Waveform comparison of average detected calcium transients from
wildtype littermates (black) and *Df(16)A$^{+/-}$* (brown) VIPo cells. **I.** Average amplitude of detected VIPo waveforms
compared between genotypes. **J.** Number of transients from VIPo cells per cell between genotypes. **K.** Averaged z-
scored reward-onset peri-event time averages (PETAs) between -1.0 and 1.0 z-scores by cell type across all cells
within cell type during the Random Foraging task for 3 seconds pre- and 3 seconds post-event. **L.** Post - Pre PETA
responses per cell type before and after reward onset comparing the 3 seconds pre- to 3 seconds post-event. *CCKBC:*
VIP+ and cholecystokinin + basket cell; *ISI3:* Interneuron-selective interneuron type 3; *ISI2:* Interneuron-selective
interneuron type 2; *VIPo*: VIP+/Cal-/M2R-/CCK- cells in layer oriens or layer pyramidale. *: $P < 0.05$. **: $P < 0.01$. ****:
$P < 0.0001$. See Suppl. Table 1 for detailed statistics.

Hippocampal place cells, a subset of pyramidal neurons, are active in specific environmental
locations and provide cellular substrates for episodic memory and goal-directed spatial
learning[91,92]. In *Df(16)A$^{+/-}$* mice, our previous study revealed significant impairments in learning
performance when both contextual cues and reward locations were altered during a spatial
navigation task[76]. These learning deficits were associated with reduced long-term stability of place
fields, impaired context-related remapping and plasticity, and a lack of reward-related
reorganization. Further investigation showed that parvalbumin- and somatostatin-expressing
GABAergic interneurons, which are directly inhibited by VIP+ cell activity, displayed diminished
place fields and heightened reward-related activity, likely contributing to the observed place cell
deficits[28]. Changes in activity dynamics of VIP+ cells in dorsal CA1 may also reflect altered spatial
information encoding by these cells. To explore this, we next evaluated how well mouse position
could be decoded based on VIP+ subtype activity during random foraging (Fig. 5A) by training a
support vector machine using all recorded IN activity (Fig. 6A). Decoding error did not differ
between genotypes when the total number of imaged cells per session was not normalized.
However, when controlling for the number of cells tested (Fig. 6B) or the number of laps run per
session (Fig. 6C), decoding error was significantly higher in *Df(16)A$^{+/-}$* mice, indicating that
individual mutant cells carry less spatial information. Furthermore, analysis of place preference in
VIP+ cell activity revealed that fewer VIP+ cells in *Df(16)A$^{+/-}$* mice exhibited significant place fields,
with a reduced proportion displaying at least one place field (Fig. 6D). These findings demonstrate
that VIP+ interneurons in the 22q11.2 deletion model exhibit markedly impaired spatial encoding
within the CA1 microcircuit.

Local brain circuits balance coordinated and unique activity depending on input properties,
connectivity, and brain state. Taking advantage of our simultaneous recordings across multiple
VIP+ subtypes in our experiments, we computed pairwise Pearson's correlation coefficients
between calcium activity traces within and between different VIP+ interneuron subtypes (see
Methods) and found that VIP+ cells in *Df(16)A$^{+/-}$* mice displayed unusually high correlations during
random foraging between subtypes. This suggests a loss of subtype-specific heterogeneity of *in*
vivo activity dynamics, particularly among disinhibitory VIP+ subtypes, indicating a potential
microcircuit-level disruption and impaired fine-scale coordination between VIP+ interneuron
subtypes (Fig. 6E/F). These findings reveal that VIP+ interneurons in the CA1 region of *Df(16)A$^{+/-}$*
mice exhibit impaired spatial encoding — reflected in increased decoding errors and reduced
place preference — and that this disruption is also accompanied by a loss of subtype-specific

heterogeneity, as indicated by unusually high inter-subtype correlations during foraging. These
physiological consequences align with our computational analyses and illuminate how
perturbations in specific subtypes of VIP+ interneurons can disrupt network dynamics and
information processing important for cognition.

412
Figure 6. Reduced specificity in *Df(16)A*^{+/-} VIP+ interneuron population activity. A. All recorded VIP+ interneuron
activity was used to train a support vector machine (*SVM*) to decode discrete position bins (*D.E.*: decoding error) across
the belt. Each bin is equivalent to 2 cm. **B.** SVM results bootstrapped 100 times by number of cells used per genotype
per bin. **C.** SVM results bootstrapped 100 times by the number of laps used per genotype per bin. **D.** Proportion of
positive place fields detected using the Peak Place method. **E.** Pairwise Pearson's correlation coefficient value matrix
between cell types of ($\Delta F/F$) traces across the trial during the Random Foraging paradigm from WT (left) and *Df(16)A*^{+/-}
(right). **F.** Comparative differences between cell type correlated activity between genotypes. *CCKBC:* VIP+ and
cholecystokinin + basket cell; *ISI3:* Interneuron-selective interneuron type 3; *ISI2:* Interneuron-selective interneuron
type 2; *VIPo*: VIP+/Cal-/M2R-/CCK- cells in layer oriens or layer pyramidale. *: P < 0.05. **: P < 0.01. ***: P < 0.001.
See Suppl. Table 1 for detailed statistics.
423

Discussion

Cognitive dysfunction is central to psychopathology, as many psychiatric conditions share overlapping yet challenging-to-treat cognitive impairments[15–18,36,72]. For instance, schizophrenia often presents with deficits in executive functioning and working memory[7,8]. Similarly, ASD cases driven by rare damaging mutations[54,93] are often associated with pronounced cognitive deficits. Cognitive disabilities are also common in such genetic conditions as 22q11.2 deletion syndrome, where affected individuals frequently display difficulties with attention, working memory, and executive processes[10,71,72]. In this study we focused on understanding neuronal mutation biases of rare highly penetrant genetic variants. We observed a strong correlation between cell type biases in ASD and SCZ that was primarily driven by mutations in highly constrained genes. We also identified several cell type biases that were more pronounced in each disorder, possibly contributing to disorder-specific symptoms. For example, medium spiny neurons of the striatum showed a significantly stronger mutation bias in ASD compared to SCZ, consistent with ASD's core symptoms of repetitive behaviors and social impairments, which are frequently associated with striatal dysfunction[55,56]. In contrast, multiple types of interneurons exhibited stronger mutation bias in SCZ, consistent with recent GWAS-based studies[21,22], particularly parvalbumin-expressing interneurons crucial for gamma oscillations and cognitive processing[94,95], as well as somatostatin-expressing interneurons important for information processing[96]. Supporting these findings, transcriptomic studies have identified alterations in these interneuron populations in SCZ patients[97].

Our genetic analysis revealed dysfunction in CGE-derived GABAergic interneurons as a central vulnerability contributing to cognitive deficits across multiple psychiatric conditions (Fig. 3). Specifically, we found that CGE interneurons are more affected in SCZ compared to high-functioning ASD, consistent with the cognitive impairments being a core and defining symptom of SCZ[16,17]. We also observed strong genetic impact on CGE interneurons in low-functioning versus high-functioning ASD, in intellectual disability and in 22q11.2 deletion syndrome. Verbal and numerical reasoning phenotypes available in the UK Biobank[57], which generally reflect milder cognitive impairments compared to psychiatric disorders, showed significant association with mutational impact on CGE interneurons and on spiny neurons of the striatum, with the latter previously implicated in studies of human intelligence[98]. Of note, impaired disinhibition and CGE-derived VIP+ interneuron dysfunction have been previously described in other neurodevelopmental disorders[99–101]. In Rett syndrome, for example, selective deletion of Mecp2 in VIP+ interneurons, but not in Pvalb+ or Sst+ interneurons, leads to deficits in cortical state transitions and social memory[100]. In Dravet syndrome, SCN1A mutations lead to persistent VIP+ interneuron dysfunction in juvenile and adult mutant mice, contributing to social and cognitive deficits[99,101].

Because VIP+ interneurons inhibit other major interneuron subtypes, their activation increases pyramidal neuron excitability, thereby improving the detection of low-contrast signals[83,102]. In this way, these interneurons are essential for learning and attention, as they respond to salient stimuli by adjusting overall neural excitability. For instance, during fear conditioning, blocking VIP+ interneurons disrupts projection neuron activity and impairs fear learning[60]. In the hippocampal

468 CA1 region, proper VIP+ interneuron function is vital for spatial learning because excessive
469 inhibition of pyramidal neurons hinders place cell remapping and performance, while their
470 activation enhances learning outcomes[88]. Similarly, in the prefrontal cortex, activation of VIP+
471 interneurons improves working memory and attention, whereas activation of Pvalb+, Sst+, or
472 pyramidal neurons produced detrimental effects[68,88,102,103].
473
Following our computational genetics analysis implicating a strong CGE/VIP+ interneuron
mutation bias in cognitive disorders, including 22q11.2 deletion syndrome, we evaluated VIP+
interneuron activity *in vivo* using the *Df(16)A$^{+/-}$* mouse model. To that end, using advanced two-
photon (2P) calcium imaging in the well-characterized hippocampal CA1 microcircuit we
measured VIP+ interneuron activity during a spatial navigation task. We observed significant
defects at both the cellular and circuit levels, likely contributing to broader hippocampal
dysfunction previously reported in this mouse model[28,76]. Specifically, VIP+ interneurons exhibited
dampened calcium activity, suggesting a reduction in the top-down disinhibition[61,104] normally
exerted by these cells in cortical circuits. In addition, we found a reduction in the spatial
information carried by their activity. Because synaptic plasticity is critical for fine-tuning synaptic
connections and forming dynamic spatial representations and episodic memories, this reduction
likely reflects impaired plasticity mechanisms. Such impairment may hinder MGE-derived
interneurons and pyramidal neurons from effectively encoding and updating spatial
information[28,76], ultimately disrupting the integration of spatial signals within the hippocampal
network. Furthermore, VIP+ interneuron subtypes exhibited increased correlation in their *in vivo*
activity, particularly among interneuron-selective interneuron subtypes. This suggests a loss of
their normally distinct activity patterns, leading to reduced informational diversity in the circuit.
Impaired disinhibition within the CA1 microcircuit in patients with 22q11.2 deletion syndrome
disrupt the processing and integration of salient inputs, resulting in cognitive deficits associated
with hippocampal dysfunction. Given that VIP+ disinhibitory motifs are highly conserved across
cortical circuits[59,60,105], the deficits we uncovered in the CA1 circuit are likely pervasive across
neocortical brain regions.
496
While further research is needed to link specific disease-related behaviors to VIP+ interneuron
circuit dysfunction, our combined computational and experimental results suggest that disruption
in VIP+ interneuron activity may be a key driver of cognitive deficits in 22q11.2 deletion syndrome
and in other neurodevelopmental and psychiatric disorders[106]. These findings position CGE/VIP+
interneurons as a critical and targetable point of dysfunction. Given the persistent cognitive
comorbidities across multiple psychiatric disorders, pharmacologically targeting VIP+ interneuron
dysfunction, especially during the chronic phase, may offer a viable treatment strategy. This
emphasizes the need for cell-type-specific tools to address VIP+ interneuron dysfunction across
a range of brain disorders.
506

508
509

METHODS

Genetic analysis:
Rare mutation genetic data for neuropsychiatric disorders and phenotypes
For the analysis on cell type mutation biases of ASD and SCZ, we primarily used the aggregated mutation data obtained from the SFARI Simons Foundation Powering Autism Research for Knowledge (SPARK) cohort, and The Schizophrenia Exome Sequencing Meta-Analysis (SCHEMA). In the SPARK study[2], de novo mutations were identified through familial sequencing of 42,607 ASD probands and 8,267 unaffected siblings. We focused our analysis on de novo likely gene disrupting (LGD) mutations and damaging missense (Dmis) mutations. Dmis mutations were identified based on the REVEL score[107], which estimates the probability of a missense mutation to be pathogenic; mutations with REVEL scores higher than 0.5 were defined as deleterious. For our main analysis, we used 61 genes that reached exome-wide significance (P-value < 1.3×10^{-6}) identified in the SPARK study, using the DeNovoWEST statistical model[45].

In the SCHEMA study[1], which analyzed 24,248 schizophrenia cases and 97,322 controls, we selected the top 61 genes with strongest association for our primary analysis (p-value < 1.0×10^{-3} or FDR<0.3). Dmis mutations were identified based on the MPC score[108]. two tiers established: MPC score > 3 (mis3) and 3 > MPC > 2 (mis2). We included LGD and mis3 mutations in our main analysis since they are most damaging mutation type.

To analyze cell type biases across different cohorts with cognitive abnormalities, we examined several additional gene sets beyond ASD and SCZ. We included genes within the 22q11.2 deletion region as defined by Karayiorgou et al[10]. We also analyzed genes associated with adult cognitive function from the UK Biobank study by Chen et al[57], which identified rare coding mutations linked to educational attainment (EDU), reaction time (RT), and verbal-numerical reasoning (VNR). For each cognitive phenotype, we selected the top 61 genes based on statistical significance (smallest p-values). We separately analyzed genes with positive and negative impacts (indicated by the beta coefficients of gene-phenotype associations), focusing on European population data for optimal statistical power.

Additionally, we included genes associated with developmental disorders and intellectual disability (DD/ID) from Kaplanis et al[45], selecting the top 61 genes according to DeNovoWEST p-values. To ensure comparable bias magnitudes across analyses, we consistently used 61 genes for each cohort to match the number of genome-wide significant genes identified in the SPARK ASD study, except for 22q11.2 genes, since the deletion region consists of only 46 genes.

Human whole brain single cell data
For our analysis, we used a recent human brain single-nucleus sequencing dataset from the Allen Institute[33], comprising 3,369,219 nuclei sampled from 105 regions across the entire brain of three donors. To the best of our knowledge, this dataset represents the first comprehensive human brain single-cell atlas with extensive regional coverage. This comprehensive sampling is crucial for accurately identifying affected brain cell types without regional sampling biases. The dataset

553   contains 461 clusters and 3,313 subclusters organized into 31 superclusters, reflecting their
554   transcriptional similarities and developmental origins.
555
Cell type mutation bias
We calculated cell type mutation biases at the cluster level (which we refer as cell type throughout
the text) for each disorder or phenotype. First, for each cell type cluster, we computed gene
expression levels by averaging $\log_2 \text{UMI} + 1$ values across all individual cells assigned to that
cluster. $\log_2$ transformation was applied to make expression values more robust to outliers and
extreme values. For each disorder-associated gene, we then characterized its expression
specificity toward each cell type cluster. We calculated expression specificity Z-scores for each
gene in a given cell type by comparing its expression level in that cell type to its expression across
all 461 cell types in the atlas, as shown in equation 1.
565

$$Z_{g,ct} = \frac{E_{g,ct} - <E_g>}{\sigma_g} \qquad (1)$$

567
568   where $E_{g,ct}$ is the expression of a gene g in a cell type ct, $< E_g >$ is the average expression level
569   of the gene g across all cell types, and σ_g is the standard deviation of the gene g expression
570   across all cell types.
571
The high granularity at the cell cluster level combined with the inherent sparsity of single-cell
sequencing data meant that some cell types had too few cells for reliable estimation of expression
levels. To mitigate noise in our cell type bias calculations, we capped the cell type specificity
scores at a maximum of 5 and a minimum of -5, preventing individual genes from having
unreasonably high specificity values that could dominate the mutational bias profile of a disease
cohort. For genes with zero expression in certain cell types, we consistently assigned the
minimum specificity score, ensuring uniform handling of non-expressed genes across all cell
types.
580
To account for variability in expression specificity that may arise simply due to different gene
expression levels across cell types (reflected in total UMI counts), we used a Z'-score, to
characterize how a gene's specificity Z-score (Equation 1) in a given cell type deviates from the
distribution of Z-scores of random genes with similar overall brain expression levels. To identify
genes with similar expression levels, we first ranked all genes based on their overall brain
expression (defined as the sum of expression levels across all individual cells in the human brain
cell atlas). We then uniformly sampled 1,000 genes within ±5% of the target gene's expression
level. Our validation tests confirmed that results were not sensitive to the sampling interval range
or the sampling kernel function. This comparison to genes with similar expression levels ensures
that neither the expression level of target genes nor cell types skews or artificially influences the
specificity measurement. Thus, for each disorder gene and each brain cell type we computed the
Z'-score:
593

$$Z'_{g,ct} = \frac{Z_{g,ct} - <Z_{ct}>_M}{\sigma_{M,ct}} \qquad (2)$$

596 where the label M corresponds to a set of sampled genes with expression levels similar to the
597 gene g, $<Z_{ct}>_M$ represents the average Z-score of genes in the set M for the cell type ct, and
598 $\sigma_{M,ct}$ represents the standard deviation of Z-scores of genes in the set M for the cell type ct.
599
600 Finally, to calculate cell type-specific mutation biases across each disorder cohort, we integrated
601 gene expression specificities (Z'-score, Equation 2) with their corresponding genetic burden. This
602 weighting approach allows us to account for the relative genetic contribution of individual genes
603 to the disorder. For the ASD cohort, we quantified each gene's contribution using a weighting
604 formula where the mutation weight for gene g was determined as follows:
605
606 $$w_g = n_{g,LGD} + n_{g,Dmis} \quad (3)$$
607
608 where $n_{g,LGD}$ and $n_{g,Dmis}$ are the numbers of observed *de novo* LGD and Dmis mutations in gene
609 g.
610
For the SCZ cohort, we calculated mutation biases by integrating both de novo and inherited
mutations identified in case-control studies. Since the genetic evidence comes from these two
distinct sources, we computed "effective mutation counts" to standardize their contributions. For
case-control mutations, the effective count represents the excess mutations in cases after
subtracting the expected counts (calculated by scaling control mutations by the case-control ratio).
We applied this approach separately for each mutation class (LGD and Mis3) and then combined
these values with observed de novo mutations, as shown in Equation 4:
618
619 $$n_{g,mutType}^{eff} = n_{g,mutType}^{case} - \frac{n_{g,LGD}^{ctrl}}{N_{ctrl}} \times N_{case} + n_{g,mutType}^{de\,novo}$$
620 $$w_g = n_{g,LGD}^{eff} + n_{g,Mis3}^{eff} \quad (4)$$
621
For genes associated with UK Biobank cognition phenotypes, we utilized each gene's effect size
(beta coefficient from gene-phenotype association tests) as the gene weight. This approach
assumes that genes with larger phenotypic effects contribute more significantly to cell type biases
relevant to cognitive function. Unlike our approach with case-control studies, mutation counts
were not employed for these analyses since VNR, EDU, and RT are quantitative traits analyzed
through linear regression rather than categorical case-control comparisons.
628
629 $$w_g^{pheno} = beta_g^{pheno} \quad (5)$$
630
631 For genes in 22q.11 deletion, we applied a uniform weight for all genes ($w_g = 1$), as there is no
632 systematic method available to determine the contribution of biases from each individual gene.
633
634 Based on the calculated mutation weight for each gene (Equations 3-5), the mutation bias for all
635 disorder gene sets towards a cell type ct was then calculated as:
636

$$637 \quad Mutation\_Bias_{ct} = \frac{\sum_{g \in G} w_g \cdot Z'_{g,ct}}{\sum_{g \in G} w_g} \quad (6)$$

where the summation is over genes g in the set G containing all implicated genes of a
disorder/phenotype, and w_g is the mutation weight for each gene g. In summary, our cell type
mutation bias approach identifies neuronal populations most strongly implicated in psychiatric
disorders by integrating three key factors: genetic evidence (mutation frequency or association
coefficients), cell type-specific expression patterns, and their statistical relationships with clinical
phenotypes. This comprehensive measurement reveals which brain cell types are preferentially
affected by high-confidence risk genes, providing insight into the neurobiological substrates of
different psychiatric conditions.

Supercluster-level bias significance assessment
To determine the statistical significance of mutation biases for each disorder-supercluster
combination, we employed the Mann-Whitney U test. This non-parametric approach compared
the distribution of mutation biases across all cell type clusters within a given supercluster against
the distribution of mutation biases in all clusters not belonging to that supercluster. This allowed
us to identify superclusters with significantly elevated mutation biases relative to the background.
Multiple comparison correction was performed using the Benjamini-Hochberg procedure to
control the false discovery rate (FDR) across all tested superclusters, ensuring robust statistical
inference.

Supercluster differential bias for a pair of disorders.
To identify disorder-specific mutational biases at the supercluster level, we assessed the
differential bias between disorders (e.g., ASD vs. SCZ), quantified by the difference in mutation
bias across cell type clusters within a supercluster. This differential bias measure describes how
strongly each particular supercluster is preferentially affected in one disorder compared to another,
potentially underlying the unique physiological and behavioral phenotypes associated with each
condition.

$$665 \quad \Delta Bias_s^{Pheno_{AB}} = \frac{\sum_{ct\_1}^{ct\_n}(Bias_{ct}^{Pheno_A} - Bias_{ct}^{Pheno_B})}{n} \quad (7)$$

Where $\Delta Bias_s^{Pheno_{AB}}{}_s$ is differential bias of Supercluster s between phenotype A and B.
$Bias_{ct}^{Pheno_A}$ is mutation bias for cell type (cluster) for phenotype A. ct are n clusters from
corresponding supercluster s.
To test whether these differential biases were statistically significant, we applied the Mann-
Whitney U test to compare bias distributions between disorders for each superclass (e.g., CGE
interneurons, MGE interneurons, medium spiny neurons). Multiple testing correction was
implemented using the Benjamini-Hochberg procedure to control the false discovery rate across
31 superclusters. This approach allowed us to identify neuronal populations consistently showing
disorder-specific vulnerability patterns across their constituent subtypes.

Phenotype bias sensitivity

Proband phenotypic data (Full Scale IQ) were obtained from ASC/SSC and from SPARK
Phenotype Collection V6[58,109]. To investigate quantitative relationships between cell type biases
and cognitive phenotypes, we performed linear regression analyses comparing Full Scale IQ
(FSIQ) scores with cell type-specific expression biases using SPARK probands with FSIQ
measurement (Supp Fig 1). Genes with exome wide significance were included in PBS analysis.
For each gene harboring de novo mutations, we calculated the mean FSIQ across all affected
probands. We then performed linear regression modeling these gene-level mean FSIQ scores as
a function of cell type-specific gene expression biases.

For a given cell type ct the FSIQ of a proband can be represent as a linear function of cell type
bias $Bias_{ct}$. The coefficient PBS_{ct} represent how sensitive the bias towards the cell type can
determine the IQ.

$$FSIQ_g = a_{ct} + PBS_{ct} \times Bias_{ct,g} + e_{ct} \qquad (8)$$

Where $FSIQ_g$ is the average FSIQ for gene g (averaged across probands with de novo mutations
in gene g). a_{ct} is intercept which approximately the average IQ of probands with De novo mutation
in highly penetrate genes. $Bias_{ct,g}$ is expression bias of gene g towards cell type ct, PBS_{ct}, is the
Phenotype Bias Sensitivity coefficient (regression slope), quantifying the relationship between cell
type bias and IQ. e_{ct} represents residual variance in IQ not explained by cell type biases. The
residual variance could be contributed by phenotypical heterogeneity within genes[110], common
genetic variants[98], environmental factors and limitations in our cell type expression bias metrics.
To assess statistical significance of PBS values, we compared observed values to a null
distribution generated through permutation testing. In each permutation, we randomly reassigned
FSIQ scores across probands included in the analysis, recalculated gene-level average IQs using
these permuted values, and computed PBS for each cell type using Equation 7. For supercluster-
level analysis, we used the mean PBS across all cell type clusters within a supercluster as the
test statistic and compared it to the corresponding null distribution from the same permutation
procedure.

Mouse Experiments:
Viruses
Cre-dependent recombinant adeno-associated virus (rAAV) for GCaMP7f (rAAV1-syn-FLEX-
jGCaMP7f-WPRE, Addgene #1944820-AAV1, titer: ≥1:10^13 vg/mL) and GCaMP8m ((rAAV1-
syn-FLEX-jGCaMP8m-WPRE, Addgene # 162378-AAV1, titer: ≥1:10^13 vg/mL) were used to
express GCaMP7f or GCaMP8m in the right, dorsal hippocampus of VIP-Cre mice crossed with
Df(16)A$^{+/-}$ mice. The Cre-dependent recombinase enabled expression of calcium sensor
GCaMP7f or GCaMP8m in VIP+ interneurons (INs).

Viral injections and hippocampal window/headpost implant
During isoflurane anesthesia, Df(16)A$^{+/-}$ x VIP-cre mice were placed onto a stereotaxic surgery
instrument. Meloxicam and bupivacaine were administered subcutaneously to reduce discomfort
and provide analgesia. The skin was cut along the midline to expose the skull, and a craniotomy
was performed over the right hemisphere with a dental drill. A Nanoject injector was loaded with

a sterile glass capillary filled with mineral oil and rAAV1-syn-FLEX-jGCaMP7f-WPRE or rAAV1-
syn-FLEX-jGCaMP8m-WPRE and lowered into dorsal CA1 of the right hemisphere. Injections
were made at AP -2.2, ML -1.75, DV - 1.8, -1.6, -1.4, -1.2, -1.1, -1.0, -0.8 relative to bregma with
64 nl of virus injected at each DV level. The pipette was held within the brain for 5 minutes after
the last injection and slowly raised from the brain. The skin was sutured, and a topical antibiotic
was applied to the suture site. 1.0 mL of saline was subcutaneously given to each mouse, while
the mice recovered in their home cage with a heat pad. Four days after viral injection, a circular
cannula with a glass window was inserted into the cavity produced by craniotomy as previously
described[111]. Similarly as described above, mice were anesthetized with isoflurane and placed on
a stereotaxic surgery instrument. Meloxicam and bupivacaine were subcutaneously given to the
mice. The skull was exposed, and a 3 mm craniotomy was performed over the right hippocampus
at AP -2.2, ML -1.75 relative to bregma, centered around the injection site. After removal of the
dura, the cortex was slowly aspirated with negative pressure and the consistent application of ice-
cold cortex buffer. When the horizontal fibers over CA1 were exposed, aspiration was stopped
and a sterile collagen membrane was applied to stop bleeding. A 3 mm circular cannula with a
glass window was placed into the craniotomy and glued into the site with Vetbond. A headpost
was then attached to the posterior of the skull with layers of dental cement. 1.0 mL of saline was
subcutaneously given to each mouse, while the mice recovered in their home cage with a heat
pad. All mice were monitored for 3 days of post-operative care until behavior training began.
741
Behavioral training and paradigms
Following the post-op recovery period, mice were acclimated to handling and head-fixation for a
few days, while being water restricted to 85-90% of their original weight. Mice were trained to run
on a fabric, cue-free burlap belt with random rewards that were automatically dispensed at a
decreasing number of random locations. Mice were trained for ~1 week on the cue-free belt and
then were transitioned to a cue rich belt during the imaging experiments.
748
749   **Random Foraging Task:**
750   Water restricted mice were imaged on a cue rich familiar belt for 2 days with 3 random rewards
751   distributed across the length of the belt, randomized every lap.
752
AOD-based two-photon calcium imaging
After the mice were acclimated and trained in the respective tasks, a high-resolution structural Z-
stack was captured 24 hours prior to imaging, while the mouse was anesthetized with isoflurane.
The reference Z-stack was used to identify the 3D position of the GCaMP-positive neurons for
access by the custom-modified AOD (acousto-optic deflector) microscope (Femto3D-ATLAS,
Femtomics Ltd). The AOD microscope was used as previously described in Geiller et al. 2020,
and Vancura and Geiller et al. 2023.
760
To provide stable transmission parameters during chronic imaging in the entire 3D scanning
volume, the AOD microscope was extended with a high-speed and precision beam stabilization
unit, which was directly attached to the AOD scan head, sensitive to input beam misalignment.
The beam stabilization unit consisted of two quadrant detectors (PDQ80A and TPA101, Thorlabs)

and two broadband dielectric mirrors (Thorlabs) mounted on motorized mirror mounts
(Femtonics). Beam alignment was performed by the LaserControl software (Femtonics).
767
A water immersion objective (x16 Nikon CF175) was used to focus at the dorsal CA1 pyramidal
layer and then fixed in position. A tunable two-photon laser (Coherent Ultra II) was adjusted to
wavelength = 920 nm. The high-resolution structural Z-stack was captured in CA1 from ~100 um
below to glass to ~500 um to span the entirety of CA1 from the *stratum oriens* to the *stratum
lacunosum-moleculare* with a 800x800 pixel capture with a resolution of 1.25 um per pixel with a
4 um step size. Laser power and photomultiplier tube (PMT) detectors (GaAsP, H10770PA-40,
Hamamatsu) were compensated in power and gain throughout the Z-stack capture. After the Z-
stack was completed, the mouse was returned to its home cage and given 24 hours to recover
before functional imaging.
777
Using the integrated software from Femtonics (MES), the Z-stack was examined, and 100-150
interneurons were manually selected to generate a spreadsheet of the x-y-z positions at the
center of each respective cell. These x-y-z positions served as the regions of interest (ROIs) that
were then used on subsequent days by the AOD microscope to find the position of the
corresponding interneurons. For each day of functional imaging in each mouse, the identical field
of view was identified within the CA1 pyramidal layer and then the ROIs were loaded in using the
software for AOD 3D imaging. Each imaging session was taken for 10 minutes, with
approximately 5-10 Hz for each recording experiment (frame rate depended on the number of
ROIs, ROI size, imaging resolution, and scanning speed).
787
Immunohistochemistry
The immunohistochemical (IHC) staining strategy is consistent with our previously published
methods[78,81]. In brief: mice were collected the last day after imaging and perfused with 20 mL
Phosphate Buffered Saline (PBS) followed by 20 mL of 4% paraformaldehyde (PFA, Electron
Microscopy Sciences) and the brain extracted for overnight fixation in 4% PFA. The next day, the
brains were washed 3 x 5 minutes with 1x PBS and subsequently vibratome sliced for horizontal
sections from the dorsal side of the brain into 75 µM sections. The relevant sections were then
taken through our IHC protocol. Sections were permeabilized for 2 x 20 minutes (0.03% Triton-X
100 (Sigma-Aldrich), 1x PBS), blocked for 45 minutes (0.03% Triton-X 100, 10% Normal Donkey
Serum (NDS), 1x PBS), and incubated with primary antibodies (Table 1) for 1 hour at room
temperature and overnight at 4 degrees with agitation. After 48 hours, primary antibodies were
washed out (3 x 5 minutes 1x PBS) and secondary antibodies were applied for 2 hours (Table 1).
Secondary antibodies were removed and the sections were washed with 1x PBS 5 x 15 minutes,
and the sections were mounted for confocal imaging.
802

Cell Marker	Round	Primary Antibody (dilution)	Secondary Antibody (dilution)	Catalog #
Cholecystokinin (CCK)	1	rabbit anti-proCCK (1:500)	donkey anti-rabbit DyLight 405 (1:300)	Frontier Institute CCK-pro-Rb-Af350

Calretinin (CalR)	1	Guinea pig anti-Calretinin (1:1000)	donkey anti-guinea pig Alexa 568 (1:300)	Swant CRgp7
Muscarinic Cholinergic M2 Receptor (M2R)	1	rat anti-M2R (1:2000)	donkey anti-rat Alexa 647 (1:300)	Millipore MAB367

Table 1. Antibodies used in this study

Confocal imaging

The NikonA1 confocal microscope used to take multi-channel fluorescence images (405 nm, 488 nm, 561 nm, and 640 nm excitation for the four blue, green, red and far-red colored channels) of labeled tissue sections. Image capturing yielded 2048 x 2048 pixel z-stacks of the entire depth of the section (with 3 microns between each image of a given stack). Z-stacks were viewable via Fiji (Image J).

Registration of confocal images to *in vivo* z-stacks and identification of immunopositivity/negativity

Unique groupings of cells from *in vivo* z-stacks were used as visual fiducial markers to manually register *ex vivo* 75 μm z-stack images in ImageJ, section by section. Care was taken to note the orientation of the *ex vivo* sections regarding the *in vivo* z-stack images. The coordinates taken from the AOD software when selecting ROIs for imaging against the *in vivo* z-stack were used to identify ROI ID numbers in the registered immunohistochemically stained images.

Assignment of subtype identity based on immunostaining

Interneuron subtypes were assigned based on our previously published classification strategy[78].

Cholecystokinin-expressing basket cells (CCKBCs):
GCaMP-expressing, immunopositive for proCCK, immunonegative for CalR and M2R[112,113]. This population represents perisomatic-targeting basket cells.

Interneuron-selective interneuron type 2 (ISI2s):
GCaMP-expressing, immunonegative for CalR, CCK and M2R. Present in *stratum radiatum* or *stratum lacunosum moleculare* CA1 layers[112,113].

Interneuron-selective interneuron type 3 (ISI3s):
GCaMP-expressing, immunopositive for CalR and immunonegative for CCK and M2R. Present in all CA1 layers[102,112–115].

VIP only (VIPos):
GCaMP-expressing, immunonegative for CalR, CCK and M2R. Present in *stratum oriens* or *stratum pyamidale* CA1 layers. This category excludes CCKBCs, ISI2s, and ISI3s, and could possibly include long-range projecting interneurons[116], but it may not be restricted to them; therefore we called them VIP only for our purposes in this study.

Pre-processing of Ca2+ imaging data

Raw movies of each individual cell were independently motion corrected using the whole-frame cross registration implemented in the SIMA package[117]. ROIs were then hand-drawn over each imaged cell. Fluorescence was extracted from each drawn ROI using SIMA. DeltaF/F was then

845 computed based on a 1st percentile baseline value over a 30-second rolling window. The DeltaF/F
846 traces were then smoothed through an exponential filter. For analyses involving velocity
847 regressed traces, each animal's velocity was smoothed with a Hanning filter and then covariate
848 linear regression was applied to each fluorescence trace where the smoothed velocity is
849 regressed out.
850
851 **Velocity Correlation**
852 For each cell, velocity correlation was computed by taking Pearson's correlation coefficient
853 between the DeltaF/F trace and animal's smoothed velocity with a -5 second to +5 second rolling
854 lag. The highest absolute value of the correlation coefficients in this time window was taken as
855 the correlation coefficient for the respective cell.
856
857 **Peri-Event Time Averages (PETA)**
858 For salient stimuli such as reward distribution events, a PETA was constructed around each event
859 respectively. These events were z-score and then averaged for each respective cell, and the z-
860 scored averages were visualized using a heatmap or an averaged line-plot. A Post-Pre metric
861 was calculated based on the z-scored averages as a measure for event modulation.
862
863 **Spatial Tuning Curve**
864 The 2 meter belt was divided into 100 bins (~2cm per bin). Spatial tuning curves for each cell
865 during an experiment were generated by averaging the velocity regressed DeltaF/F in each bin,
866 when the animal was running over 5 cm/s.
867
Place field calculation
The spatial tuning curves were shuffled 1000 times by circularly rotating the velocity-regressed
ΔF/F tuning curves around the 100 spatial bins to generate a shuffle distribution. Cells were
classified as being spatially selective (i.e., having a place field) if the cell's tuning curve exceeded
the 95th percentile (or was under the 5th percentile for negatively selective cells) with at least 10
sequential bins over/under the cutoffs from the shuffled distribution. The number of bins over or
under the 95th and 5th percentile curves were visualized and a cutoff of 30 consecutive bins was
used for the figures as it represented the approximate 75th quartile for place field width.
876
Position Decoding
A linear classifier support vector machine (SVM) was trained on the tuning curves of each
respective cell and the decoding error was visualized. The SVM was cross-validated by training
the classifier on (n-1) laps and evaluating on the held-out lap and this process was repeated such
that every lap was utilized as a test lap at least once. Furthermore, the SVM was bootstrapped
with different numbers of input cells and input laps to compare the position decoding efficacy
between genotypes.
884
885 **Cross-correlation Analysis**
886 Pairwise Pearson's correlation coefficients were computed for every pair of cells within a given
887 experiment. The average coefficient value for each subtype was then calculated per trial pair and
888 averaged across mice.

889
890 **Transient Analysis**
891 Online Active Set method to Infer Spikes (OASIS) was used to generate deconvolved spike
892 events based on transient events within the dF/F traces[90]. We understand that these are not true
893 spiking events, and we are using OASIS to algorithmically identify transient events rather than
894 interpret the deconvolved spiking events.
895
896 Frequency: The deconvolved events were filtered to remove events that were within 10 frames.
897 The total number of filtered deconvolved events by OASIS was normalized by the total duration
898 of the recording.
899
For each cell, the average transient was segmented within a 15-second time window and
computed by averaging along aligned deconvolved spike times. If multiple detected events were
within 10 frames, the events were treated as a single transient. A cell was only used if there were
at least 3 detected transients within the total trace to exclude cells without obvious GCaMP-Ca^{2+}
dynamics. For each average transient, we computed the median value based on 5 seconds pre
peak transient and the range 5-10 seconds after the peak transient (given that the average
transient took significantly less than 5 seconds to resolve) to act as the baseline.
907
908 Amplitude: The difference between the max value of the average transient and the baseline value
909 was computed
910
911
912 **QUANTIFICATION AND STATISTICAL ANALYSIS**
913
Statistical analyses
Subtype comparisons were conducted using one-way ANOVA tests with Tukey's range tests with
correction for multiple testing if appropriate. For comparisons across genotypes, unpaired t tests
were utilized if the data followed a normal distribution. Otherwise, for non-normally distributed
comparisons, the Mann-Whitney U test was used. *, $P < 0.05$, **, $P < 0.01$, ***, $P < 0.001$. All data
analysis and visualization were done with custom software built on Python version 2.7.15.
920
Code and Data availability:
All data used in this report are publicly available. ASD genes and mutations counts can be
found in Supplementary table from Zhou et al[2]. SCZ genes and mutation counts can be found in
Supplementary table from Singh et al[1]. DD/ID genes and mutations counts can be found in
Supplementary table from Kaplanis et al[45]. Genes implicated in UKBB adult cognitive functions
can be found in Supplementary table from Chen et al[57]. Genes in 22q11.2 3MB deletion and 1.5
MB deletion can be found in Karayiorgou et al[10]. Human brain cell atlas can be found
(https://cellxgene.cziscience.com/collections/283d65eb-dd53-496d-adb7-7570c7caa443).
Individual genetics and phenotypic data used in our study from the SPARK project are available
through the Simons Foundation (https://base.sfari.org) following the access procedures
described in Zhou et al.
932

933 The code used for our implementation of the Cell type bias with analysis pipeline, and code to
934 generating all figures and tables, is available on GitHub:
935 (https://github.com/explorerwjy/CellTypeBias_VIP.git)
936

Acknowledgements

S.A.H. is supported by the Burroughs Welcome grant. This work has been supported by the grants
NIMHR01MH124923 to D.V. and J.A.G, NIMHR01MH124047 to J.A.G, Stavros Niarchos
Foundation to J.A.G., and NIMHR01MH124047 NIMHR01MH124867 NINDSR01NS133381 to
A.L. We thank members of the Gogos, Losonczy, and Vitkup labs for fruitful scientific discussions.

Author Contributions

S.H., J.W., J.A.G., A.L., and D.V. conceived the study. J.W. and J.C. performed genetic bias and
computational analyses. S.H. and B.R. collected and analyzed experimental data. S.H., J.W.,
J.A.G., A.L., and D.V. wrote the manuscript.
947

948 **Supplemental Figures**

Figure	Statistics
5C	*CCKBC:* CCK basket cells student's t-test: T statistic: -1.313, P = 0.019. *ISI3*: Interneuron-Selective Interneuron Type 3: Student's T test Statistic: -3.284, P = 0.001. *ISI2:* Interneuron-Selective Interneuron Type 2: Student's T test Statistic: 2.191, P = 0.030. *VIP+ Alone:* Student's T-test Statistic: 1.105, P = 0.269
5E	Mann-Whitney test, t = 104931.0, P = 4.1530e-05.
5G	Mann-Whitney test, t =24290.0, P = 0.0388.
5I	Mann-Whitney test, t = 42011.0, P = 2.5817e-13.
5J	Mann-Whitney test, t = 54521.5, P = 0.0013
5L	*CCKBC*: cholecystokinin-expressing basket cells Mann-Whitney U test: P = 0.292, *ISI3*: interneuron-selective interneuron type 3 cells Mann-Whitney U test: P = 0.0029, *ISI2*: interneuron-selective interneuron type 2 cells Mann-Whitney U test: P = 0.0026, *VIP-Alone*: Mann-Whitney U test: P = 0.036. For Mann Whitney U tests: median ± inter-quartile range.
6A	Averaged per animal: Mann-Whitney U test: statistic = 3.5, P = 0.144
6B	10 cells: Mann-Whitney U test: P = 6.67e-10, 30 cells: Mann-Whitney U test: P = 8.32e-11, 50 cells: Mann-Whitney U test: P = 2.75e-16, 70 cells: Mann-Whitney U test: P = 2.67e-18, 90 cells: Mann-Whitney U test: P = 3.00e-21.
6C	5 laps: Mann-Whitney U test: P = 2.66e-6, 10 laps: Mann-Whitney U test: P =2.27e-21, 15 laps: Mann-Whitney U test: P =2.49e-15.
6D	0 fields: Mann-Whitney U test: p = 0.056, 1 field: Mann-Whitney U test: P = 0.030, 2 fields: Mann-Whitney U test: P = 0.097, 3 fields: Mann-Whitney U test: P = 0.062.
6F	Welch's T-test comparisons: ISI2 vs VIPo: P = 0.0067. ISI3 vs ISI3: P = 0.043. ISI3 vs VIPo: P = 0.039. VIPo vs ISI2: P = 0.0068. VIPo vs ISI3: P = 0.038. Welch's T-test.

949 **Supplementary Table 1: Statistical detail summary.**
950 Boxplots show the 25th, 50th (median), and 75th quartile ranges, with the whiskers extending to 1.5 interquartile
951 ranges below or above the 25th or 75th quartiles, respectively. Outliers are defined as values extending beyond. the
952 whisker ranges. For Welch's t tests: mean ± sem. For Mann Whitney U tests: median ± inter-quartile range. n.s.: not
953 significant. *: P < 0.05. **: P < 0.01. ***: P < 0.001.
954

955 **Supplementary Table 2: Human brain Cell type biases (cluster level) for psychiatric disorders.**
956

957 **Supplementary Table 3: Human brain Cell type biases (supercluster level) for psychiatric disorders.**
958

959 **Supplementary Table 4: Human brain differential biases for psychiatric disorders.**
960

961 **Supplementary Table 5: PBS values for ASD FSIQ.**
962

965
Supplemental Figure 1: Full-scale IQ distribution among ASD probands in the SPARK and ASC cohorts. A).
The distribution of FSIQ scores among ASD probands from the SPARK and ASC cohorts. The histogram (blue bars)
shows the FSIQ distribution with an overlaid density plot (red line). A two-component Gaussian mixture model was
fitted to the data (blue and orange dashed curves), revealing the distinct subpopulations within the FSIQ distribution.
The vertical dashed line at FSIQ = 70 indicates a frequently used threshold for intellectual disability (IQ < 70). FSIQ
phenotype data was obtained from Zhou *et al.*[2] and Satterstrom *et al.*[58] **B).** Gene-specific FSIQ patterns across the top
61 high-confidence ASD genes, represented by vertical lines. Individual probands are represented as dots, colored by
the mutation type, with likely gene-disrupting (LGD) mutations in red and damaging missense (Dmis) mutations in blue.
The green diamonds indicate the average FSIQ in the cohorts for each gene. Genes are arranged along the x-axis in
based on *de novo* west P-values from Zhou *et al.*[2].
976

977
978
979
Supplemental Figure 2. Temporal expression patterns of disorder-associated genes across human brain
development. Average expression levels (log10 RPKM) of genes associated with different disorders across human
brain developmental stages, from embryonic to adult periods. Gene sets include ASD genes (green), ASD with ID
genes (orange), SCZ genes (blue), and background genes (dashed yellow). Error bars represent standard error of the
mean. SCZ genes show similar prenatal expression level as ASD genes while ASD with ID genes show higher prenatal
expression level. The vertical dashed line marks the transition from prenatal to postnatal periods. Expression data
derived from the BrainSpan Atlas of Human Brain Development[118].

987
Supplemental Figure 3: Illustration of disorder cell type mutation bias. Illustration of the gene expression analysis
used to determine disorder mutation biases for each cell type. To calculate the biases, gene expression specificities
were first calculated for each gene and for each cell type in the brain; the expression specificity of a gene in a brain cell
type was calculated as the Z-score between the gene expression level in the cell type compared to the distribution of
the gene expression levels in the whole brain (sum of UMI cross all cell). The specificities were calculated for each
disorder gene (red) and for randomly matched genes (green) with similar expression levels in the brain. For each cell
type, expression biases were then calculated for each disorder gene as the Z-score between the expression specificity
of that gene and the distribution of expression specificities of random genes with similar expression level in the brain.
Based on gene expression biases, disorder mutation biases were determined for each cell type by computing a
weighted average of mutation biases across all disorder genes; Gene weights were based on disorder mutation
contribution from each gene (see Methods).

999
Supplemental Figure 4. Comprehensive analysis of mutation biases across brain cell types in psychiatric
disorders. Box plots showing mutation biases for all major brain cell types (superclusters) in **A.)** ASD (top left), **B.)**
SCZ (top right), **C)**. VNR Negative genes (bottom left), and **D.)** DD/ID (bottom right). Each bar represents the bias
distribution of the human adult brain neuron type belonging to a supercluster, X-axis shows the magnitude of biases.
Red diamond shows average bias of each supercluster, red vertical line shows medium bias of each supercluster.
Detailed data can be found in Supplementary table 2.

1006
Supplemental Figure 5. Impact of gene expression levels on ASD-SCZ cell type bias correlation. Effect of
sequentially removing genes based on their expression levels on the cell type mutation bias correlation between ASD
and SCZ. Red line shows correlation changes when removing genes with lowest expression levels first, while blue
dashed line shows removal of highest expressing genes first. Grey line and error bars show mean ± standard deviation
for random gene removal. The correlation remains relatively stable when removing lowly expressed genes or highly
expressed genes. This suggests gene expression levels play a secondary role in determining shared cell type patterns
between disorders compared to gene constraint.
1014

1015
Supplemental Figure 6. ASD and SCZ Mutation bias towards different superclusters. A-D. Detailed comparison
of mutation biases in specific neuron superclusters, P-value was calculated based on Wilcoxon signed-rank test with
FDR correction: **(A)** Hippocampal CA1-3 neurons show similar bias in ASD and SCZ. **(B)** Amygdala excitatory neurons
show similar bias in ASD and SCZ. **(C)** Deep-layer IT neurons show similar bias in ASD and SCZ. **(D)** Deep-layer
corticothalamic and 6b neurons show similar bias in ASD and SCZ. **(E)** CGE interneurons show stronger bias in VNR-
genes compared to VNR+ genes (adjusted P =3.7e-06). **(F)** CGE interneurons show stronger bias in DD/ID genes
compared to ASD genes (adjusted P =1.4e-07). **(G)** Bar plot comparing mutation biases across major neuron types
between ASD (blue) and ASD with ID (orange). Asterisks indicate significance level between groups (*: P < 0.05. **: P
< 0.01. ***: P < 0.001, ****: P < 0.0001, ns=not significant). Neuron types showing positive mutation bias from at least
one disorder are displayed and sorted by bias difference between ASD and ASD with ID. Error bars represent standard
error across clusters from corresponding superclass. Detailed data can be found in Supplementary table 4.
1027

1028

1029

Supplemental Figure 7. Mutation bias comparison across psychiatric disorders for LAMP5-LHX6/Chandelier
and MGE interneurons. A: Mutation bias distribution for LAMP5-LHX6 and Chandelier cells across different cohorts:
ASD (Mean Bias = -0.10, adjusted P = 1), ASD with ID (Mean Bias = 0.2, adjusted P = 4.0e-04), SCZ (Mean Bias =
0.07, adjusted P = 0.7), VNR- (Mean Bias = -0.03, adjusted P = 1), and DD/ID (Mean Bias = 0.22, adjusted P = 1.0e-
03). **B:** Mutation bias distribution for MGE interneurons across different cohorts: ASD (Mean Bias = -0.12, adjusted P
= 1), ASD with ID (Mean Bias = 0.02, adjusted P = 1), SCZ (Mean Bias = 0.11, adjusted P = 0.047), VNR- (Mean Bias
= -0.1, adjusted P = 1), and DD/ID (Mean Bias = 0.06, adjusted P = 0.1). Detailed data can be found in Supplementary
table 2 and 3.

1038
1039 **Supplemental Figure 8. Comprehensive analysis of mutation biases across brain cell types in 22q11.2 deletion.**
1040 Box plots showing mutation biases for all major brain cell types (superclusters). Each bar represents the bias distribution
1041 of the human adult brain neuron type belonging to a superclass, X-axis shows the magnitude of biases. Red diamond
1042 shows average bias of each supercluster, red vertical line shows medium bias of each supercluster. Detailed data can
1043 be found in Supplementary table 2 and 3.

 Supplemental Figure 9. VIP+ Cell imaging and targeting strategy.
 A. Schematic for mouse crossing and viral injection and calcium imaging setup. **B.** 3D reconstruction of Z-stack from
 VIP-cre mice **C.** Schematic of 3D regions of interest (ROIs) imaged with AOD 2-photon microscopy. **D.** Example
 chessboard scan view of imaged ROIs. **E.** Manual registration of *in vivo* Z-stack section with *ex vivo* 75 □m transverse
 section for *post hoc* subtype classification with immunohistochemistry. **F.** Histological strategy for defining VIP-
 expressing cell subtypes. **G.** 3D distribution of all cells imaged from both genotypes. **H.** Distribution of VIP+ subtypes

across CA1 hippocampal layers. **I.** Number of cells recorded per mouse (n=4, ISI3: ISI2: VIP+ Alone: CCKBC:). **J.**
Average velocity in cm per second plotted by location on the belt of WT (black) and *Df(16)A$^{+/-}$* (brown) mice. **K.** Average
percent occupancy on the belt by position per genotype plotted by location on the belt of WT (black) and *Df(16)A$^{+/-}$*
(brown) mice. **L.** Total laps across mice within genotype (WT: 14.4 ± 1.1 laps, *Df(16)A$^{+/-}$*: 16.2 ± 2.0 laps, Mann Whitney
U = 51.5, P = 0.21). **M**. Licking proportion of WT (black) and *Df(16)A$^{+/-}$* (brown). There is no significant difference in
total licks across the belt during Random Foraging in WT (*n* = 4) or *Df(16)A$^{+/-}$* (*n* = 4) depicted by box-and-whisker plot,
WT: 1916 ± 304 licks, *Df(16)A$^{+/-}$*: 2200 ± 489 licks, Mann-Whitney U Test, P = 0.408.

References

1058
1059
1.   Singh, T., Poterba, T., Curtis, D., Akil, H., Al Eissa, M., Barchas, J.D., Bass, N., Bigdeli, T.B.,
Breen, G., Bromet, E.J., et al. (2022). Rare coding variants in ten genes confer substantial
risk for schizophrenia. Nature *604*, 509–516.

2.   Zhou, X., Feliciano, P., Shu, C., Wang, T., Astrovskaya, I., Hall, J.B., Obiajulu, J.U., Wright,
J.R., Murali, S.C., Xu, S.X., et al. (2022). Integrating de novo and inherited variants in 42,607
autism cases identifies mutations in new moderate-risk genes. Nat. Genet. *54*, 1305–1319.

3.   Gilman, S.R., Chang, J., Xu, B., Bawa, T.S., Gogos, J.A., Karayiorgou, M., and Vitkup, D.
(2012). Diverse types of genetic variation converge on functional gene networks involved in
schizophrenia. Nat. Neurosci. *15*, 1723–1728.

4.   Gilman, S.R., Iossifov, I., Levy, D., Ronemus, M., Wigler, M., and Vitkup, D. (2011). Rare de
novo variants associated with autism implicate a large functional network of genes involved
in formation and function of synapses. Neuron *70*, 898–907.

5.   Gandal, M.J., Haney, J.R., Parikshak, N.N., Leppa, V., Ramaswami, G., Hartl, C., Schork,
A.J., Appadurai, V., Buil, A., Werge, T.M., et al. (2018). Shared molecular neuropathology
across major psychiatric disorders parallels polygenic overlap. Science *359*, 693–697.

6.   Rees, E., Creeth, H.D.J., Hwu, H.-G., Chen, W.J., Tsuang, M., Glatt, S.J., Rey, R., Kirov, G.,
Walters, J.T.R., Holmans, P., et al. (2021). Schizophrenia, autism spectrum disorders and
developmental disorders share specific disruptive coding mutations. Nat. Commun. *12*,
5353.

7.   Trevisan, D.A., Foss-Feig, J.H., Naples, A.J., Srihari, V., Anticevic, A., and McPartland, J.C.
(2020). Autism spectrum disorder and schizophrenia are better differentiated by positive
symptoms than negative symptoms. Front. Psychiatry *11*, 548.

8.   Spek, A.A., and Wouters, S.G.M. (2010). Autism and schizophrenia in high functioning
adults: Behavioral differences and overlap. Res. Autism Spectr. Disord. *4*, 709–717.

9.   Schneider, M., Debbané, M., Bassett, A.S., Chow, E.W.C., Fung, W.L.A., van den Bree, M.,
Owen, M., Murphy, K.C., Niarchou, M., Kates, W.R., et al. (2014). Psychiatric disorders from
childhood to adulthood in 22q11.2 deletion syndrome: results from the International
Consortium on Brain and Behavior in 22q11.2 Deletion Syndrome. Am. J. Psychiatry *171*,
627–639.

10. Karayiorgou, M., Simon, T.J., and Gogos, J.A. (2010). 22q11.2 microdeletions: linking DNA
structural variation to brain dysfunction and schizophrenia. Nature Reviews Neuroscience
2010 11:6 *11*, 402–416.

11. Rees, E., Kendall, K., Pardiñas, A.F., Legge, S.E., Pocklington, A., Escott-Price, V.,
MacCabe, J.H., Collier, D.A., Holmans, P., O'Donovan, M.C., et al. (2016). Analysis of
intellectual disability copy number variants for association with schizophrenia. JAMA
Psychiatry *73*, 963–969.

12. Miller, D.T., Adam, M.P., Aradhya, S., Biesecker, L.G., Brothman, A.R., Carter, N.P., Church,
D.M., Crolla, J.A., Eichler, E.E., Epstein, C.J., et al. (2010). Consensus statement:

chromosomal microarray is a first-tier clinical diagnostic test for individuals with
developmental disabilities or congenital anomalies. Am. J. Hum. Genet. *86*, 749–764.

13. Ropers, H.H. (2008). Genetics of intellectual disability. Curr. Opin. Genet. Dev. *18*, 241–250.

14. Vissers, L.E.L.M., Gilissen, C., and Veltman, J.A. (2016). Genetic studies in intellectual
disability and related disorders. Nat. Rev. Genet. *17*, 9–18.

15. Harvey, P.D., and Strassnig, M. (2012). Predicting the severity of everyday functional
disability in people with schizophrenia: cognitive deficits, functional capacity, symptoms, and
health status. World Psychiatry *11*, 73–79.

16. Bowie, C.R., and Harvey, P.D. (2006). Cognitive deficits and functional outcome in
schizophrenia. Neuropsychiatr. Dis. Treat. https://doi.org/10.2147/nedt.s24531.

17. Bowie, C.R., Reichenberg, A., Patterson, T.L., Heaton, R.K., and Harvey, P.D. (2006).
Determinants of real-world functional performance in schizophrenia subjects: Correlations
with cognition, functional capacity, and symptoms. Am. J. Psychiatry *163*, 418–425.

18. Millan, M.J., Agid, Y., Brüne, M., Bullmore, E.T., Carter, C.S., Clayton, N.S., Connor, R.,
Davis, S., Deakin, B., DeRubeis, R.J., et al. (2012). Cognitive dysfunction in psychiatric
disorders: characteristics, causes and the quest for improved therapy. Nat. Rev. Drug
Discov. *11*, 141–168.

19. Trubetskoy, V., Pardiñas, A.F., Qi, T., Panagiotaropoulou, G., Awasthi, S., Bigdeli, T.B.,
Bryois, J., Chen, C.Y., Dennison, C.A., Hall, L.S., et al. (2022). Mapping genomic loci
implicates genes and synaptic biology in schizophrenia. Nature *604*, 502–508.

20. Bryois, J., Skene, N.G., Hansen, T.F., Kogelman, L.J.A., Watson, H.J., Liu, Z., Eating
Disorders Working Group of the Psychiatric Genomics Consortium, International Headache
Genetics Consortium, 23andMe Research Team, Brueggeman, L., et al. (2020). Genetic
identification of cell types underlying brain complex traits yields insights into the etiology of
Parkinson's disease. Nat. Genet. *52*, 482–493.

21. Duncan, L.E., Li, T., Salem, M., Li, W., Mortazavi, L., Senturk, H., Shahverdizadeh, N.,
Vesuna, S., Shen, H., Yoon, J., et al. (2025). Mapping the cellular etiology of schizophrenia
and complex brain phenotypes. Nat. Neurosci., 1–11.

22. Yao, S., Harder, A., Darki, F., Chang, Y.-W., Li, A., Nikouei, K., Volpe, G., Lundström, J.N.,
Zeng, J., Wray, N., et al. (2024). Connecting genomic results for psychiatric disorders to
human brain cell types and regions reveals convergence with functional connectivity.
medRxiv. https://doi.org/10.1101/2024.01.18.24301478.

23. Broglio, C., Martín-Monzón, I., Ocaña, F.M., Gómez, A., Durán, E., Salas, C., and Rodríguez,
F. (2015). Hippocampal pallium and map-like memories through vertebrate evolution. J.
Behav. Brain Sci. *05*, 109–120.

24. Rodríguez, F., López, J.C., Vargas, J.P., Broglio, C., Gómez, Y., and Salas, C. (2002).
Spatial memory and hippocampal pallium through vertebrate evolution: insights from reptiles
and teleost fish. Brain Res. Bull. *57*, 499–503.

25. Buzsáki, G. (2002). Theta oscillations in the hippocampus. Neuron *33*, 325–340.

26. Pelkey, K.A., Vargish, G.A., Pellegrini, L.V., Calvigioni, D., Chapeton, J., Yuan, X., Hunt, S.,
Cummins, A.C., Eldridge, M.A.G., Pickel, J., et al. (2023). Evolutionary conservation of
hippocampal mossy fiber synapse properties. Neuron *111*, 3802-3818.e5.

27. Hamm, J.P., Peterka, D.S., Gogos, J.A., and Yuste, R. (2017). Altered Cortical Ensembles
in Mouse Models of Schizophrenia. Neuron *94*, 153-167.e8.

28. Herrlinger, S.A., Rao, B.Y., Conde Paredes, M.E., Tuttman, A.L., Arain, H., Varol, E., Gogos,
J.A., and Losonczy, A. (2024). Disorganized Inhibitory Dynamics and Functional
Connectivity in Hippocampal area CA1 of 22q11.2 Deletion Mutant Mice. bioRxiv,
2024.04.28.591464. https://doi.org/10.1101/2024.04.28.591464.

29. Gogos, J.A., Crabtree, G., and Diamantopoulou, A. (2020). The abiding relevance of mouse
models of rare mutations to psychiatric neuroscience and therapeutics. Schizophr. Res. *217*,
37–51.

30. Diamantopoulou, A., and Gogos, J.A. (2019). Neurocognitive and Perceptual Processing in
Genetic Mouse Models of Schizophrenia: Emerging Lessons. Neuroscientist *25*, 597–619.

31. Arguello, P.A., and Gogos, J.A. (2006). Modeling madness in mice: one piece at a time.
Neuron *52*, 179–196.

32. Gordon, A., Forsingdal, A., Klewe, I.V., Nielsen, J., Didriksen, M., Werge, T., and Geschwind,
D.H. (2019). Transcriptomic networks implicate neuronal energetic abnormalities in three
mouse models harboring autism and schizophrenia-associated mutations. Molecular
Psychiatry 2019 26:5 *26*, 1520–1534.

33. Siletti, K., Hodge, R., Mossi Albiach, A., Lee, K.W., Ding, S.-L., Hu, L., Lönnerberg, P.,
Bakken, T., Casper, T., Clark, M., et al. (2023). Transcriptomic diversity of cell types across
the adult human brain. Science *382*, eadd7046.

34. Geschwind, D.H., and Flint, J. (2015). Genetics and genomics of psychiatric disease.
Science *349*, 1489–1494.

35. Iossifov, I., O'Roak, B.J., Sanders, S.J., Ronemus, M., Krumm, N., Levy, D., Stessman, H.A.,
Witherspoon, K.T., Vives, L., Patterson, K.E., et al. (2014). The contribution of de novo
coding mutations to autism spectrum disorder. Nature *515*, 216–221.

36. Matson, J.L., and Shoemaker, M. (2009). Intellectual disability and its relationship to autism
spectrum disorders. Res. Dev. Disabil. *30*, 1107–1114.

37. Woodberry, K.A., Giuliano, A.J., and Seidman, L.J. (2008). Premorbid IQ in schizophrenia:
a meta-analytic review. Am. J. Psychiatry *165*, 579–587.

38. Khandaker, G.M., Barnett, J.H., White, I.R., and Jones, P.B. (2011). A quantitative meta-
analysis of population-based studies of premorbid intelligence and schizophrenia. Schizophr.
Res. *132*, 220–227.

39. Creeth, H.D.J., Rees, E., Legge, S.E., Dennison, C.A., Holmans, P., Walters, J.T.R., O'Donovan, M.C., and Owen, M.J. (2022). Ultrarare coding variants and cognitive function in schizophrenia. JAMA Psychiatry *79*, 963–970.

40. Nuechterlein, K.H., Green, M.F., Kern, R.S., Baade, L.E., Barch, D.M., Cohen, J.D., Essock, S., Fenton, W.S., Frese, F.J., 3rd, Gold, J.M., et al. (2008). The MATRICS Consensus Cognitive Battery, part 1: test selection, reliability, and validity. Am. J. Psychiatry *165*, 203–213.

41. Mesholam-Gately, R.I., Giuliano, A.J., Goff, K.P., Faraone, S.V., and Seidman, L.J. (2009). Neurocognition in first-episode schizophrenia: a meta-analytic review. Neuropsychology *23*, 315–336.

42. Minshew, N.J., Goldstein, G., and Siegel, D.J. (1997). Neuropsychologic functioning in autism: profile of a complex information processing disorder. J. Int. Neuropsychol. Soc. *3*, 303–316.

43. Hill, E.L. (2004). Executive dysfunction in autism. Trends Cogn. Sci. *8*, 26–32.

44. Palmer, D.S., Howrigan, D.P., Chapman, S.B., Adolfsson, R., Bass, N., Blackwood, D., Boks, M.P.M., Chen, C.-Y., Churchhouse, C., Corvin, A.P., et al. (2022). Exome sequencing in bipolar disorder identifies AKAP11 as a risk gene shared with schizophrenia. Nat. Genet. *54*, 541–547.

45. Kaplanis, J., Samocha, K.E., Wiel, L., Zhang, Z., Arvai, K.J., Eberhardt, R.Y., Gallone, G., Lelieveld, S.H., Martin, H.C., McRae, J.F., et al. (2020). Evidence for 28 genetic disorders discovered by combining healthcare and research data. Nature *586*, 757–762.

46. Karczewski, K.J., Francioli, L.C., Tiao, G., Cummings, B.B., Alföldi, J., Wang, Q., Collins, R.L., Laricchia, K.M., Ganna, A., Birnbaum, D.P., et al. (2020). The mutational constraint spectrum quantified from variation in 141,456 humans. Nature *581*, 434–443.

47. Aleman, A., and Kahn, R. (2005). Strange feelings: Do amygdala abnormalities dysregulate the emotional brain in schizophrenia? Prog. Neurobiol. *77*, 283–298.

48. Baron-Cohen, S., Ring, H.A., Bullmore, E.T., Wheelwright, S., Ashwin, C., and Williams, S.C. (2000). The amygdala theory of autism. Neurosci. Biobehav. Rev. *24*, 355–364.

49. Banker, S.M., Gu, X., Schiller, D., and Foss-Feig, J.H. (2021). Hippocampal contributions to social and cognitive deficits in autism spectrum disorder. Trends Neurosci. *44*, 793–807.

50. Lieberman, J.A., Girgis, R.R., Brucato, G., Moore, H., Provenzano, F., Kegeles, L., Javitt, D., Kantrowitz, J., Wall, M.M., Corcoran, C.M., et al. (2018). Hippocampal dysfunction in the pathophysiology of schizophrenia: a selective review and hypothesis for early detection and intervention. Mol. Psychiatry *23*, 1764–1772.

51. Zierhut, K., Bogerts, B., Schott, B., Fenker, D., Walter, M., Albrecht, D., Steiner, J., Schütze, H., Northoff, G., Düzel, E., et al. (2010). The role of hippocampus dysfunction in deficient memory encoding and positive symptoms in schizophrenia. Psychiatry Res. *183*, 187–194.

52. Moberg, S., and Takahashi, N. (2022). Neocortical layer 5 subclasses: From cellular
properties to roles in behavior. Front. Synaptic Neurosci. *14*, 1006773.

53. Jutla, A., Foss-Feig, J., and Veenstra-VanderWeele, J. (2022). Autism spectrum disorder
and schizophrenia: An updated conceptual review. Autism Res. *15*, 384–412.

54. Chang, J., Gilman, S.R., Chiang, A.H., Sanders, S.J., and Vitkup, D. (2015). Genotype to
phenotype relationships in autism spectrum disorders. Nat. Neurosci. *18*, 191–198.

55. Peça, J., Feliciano, C., Ting, J.T., Wang, W., Wells, M.F., Venkatraman, T.N., Lascola, C.D.,
Fu, Z., and Feng, G. (2011). Shank3 mutant mice display autistic-like behaviours and striatal
dysfunction. Nature *472*, 437–442.

56. Rothwell, P.E., Fuccillo, M.V., Maxeiner, S., Hayton, S.J., Gokce, O., Lim, B.K., Fowler, S.C.,
Malenka, R.C., and Südhof, T.C. (2014). Autism-associated neuroligin-3 mutations
commonly impair striatal circuits to boost repetitive behaviors. Cell *158*, 198–212.

57. Chen, C.-Y., Tian, R., Ge, T., Lam, M., Sanchez-Andrade, G., Singh, T., Urpa, L., Liu, J.Z.,
Sanderson, M., Rowley, C., et al. (2023). The impact of rare protein coding genetic variation
on adult cognitive function. Nat. Genet. *55*, 927–938.

58. Satterstrom, F.K., Kosmicki, J.A., Wang, J., Breen, M.S., De Rubeis, S., An, J.-Y., Peng, M.,
Collins, R., Grove, J., Klei, L., et al. (2020). Large-Scale Exome Sequencing Study Implicates
Both Developmental and Functional Changes in the Neurobiology of Autism. Cell *180*, 568-
584.e23.

59. Kepecs, A., and Fishell, G. (2014). Interneuron cell types are fit to function. Nature *505*, 318–
326.

60. Chamberland, S., and Topolnik, L. (2012). Inhibitory control of hippocampal inhibitory
neurons. Front. Neurosci. *0*, 165.

61. Letzkus, J.J., Wolff, S.B.E., and Lüthi, A. (2015). Disinhibition, a circuit mechanism for
associative learning and memory. Neuron *88*, 264–276.

62. Pfeffer, C.K., Xue, M., He, M., Huang, Z.J., and Scanziani, M. (2013). Inhibition of inhibition
in visual cortex: the logic of connections between molecularly distinct interneurons. Nat.
Neurosci. *16*, 1068–1076.

63. Fu, Y., Tucciarone, J.M., Espinosa, J.S., Sheng, N., Darcy, D.P., Nicoll, R.A., and Stryker M
(2014). A cortical circuit for gain control by behavioral state. Cell *156*, 1139–1152.

64. Bennett, C., Arroyo, S., Berns, D., and Hestrin, S. (2012). Mechanisms generating dual-
component nicotinic EPSCs in cortical interneurons. J. Neurosci. *32*, 17287–17296.

65. Letzkus, J.J., Wolff, S.B.E., Meyer, E.M.M., Tovote, P., Courtin, J., Herry, C., and Lüthi, A.
(2011). A disinhibitory microcircuit for associative fear learning in the auditory cortex. Nature
480, 331–335.

66. Poorthuis, R.B., Enke, L., and Letzkus, J.J. (2014). Cholinergic circuit modulation through
differential recruitment of neocortical interneuron types during behaviour: Cholinergic circuit
modulation via interneurons. J. Physiol. *592*, 4155–4164.

67. Froemke, R.C., Merzenich, M.M., and Schreiner, C.E. (2007). A synaptic memory trace for
cortical receptive field plasticity. Nature *450*, 425–429.

68. Lee, A.T., Cunniff, M.M., See, J.Z., Wilke, S.A., Luongo, F.J., Ellwood, I.T., Ponnavolu, S.,
and Sohal, V.S. (2019). VIP interneurons contribute to avoidance behavior by regulating
information flow across hippocampal-prefrontal networks. Neuron *102*, 1223-1234.e4.

69. Marshall, C.R., Howrigan, D.P., Merico, D., Thiruvahindrapuram, B., Wu, W., Greer, D.S.,
Antaki, D., Shetty, A., Holmans, P.A., Pinto, D., et al. (2017). Contribution of copy number
variants to schizophrenia from a genome-wide study of 41,321 subjects. Nat. Genet. *49*, 27–
35.

70. Matsunami, N., Hadley, D., Hensel, C.H., Christensen, G.B., Kim, C., Frackelton, E.,
Thomas, K., da Silva, R.P., Stevens, J., Baird, L., et al. (2013). Identification of rare recurrent
copy number variants in high-risk autism families and their prevalence in a large ASD
population. PLoS One *8*, e52239.

71. Robin, N.H., and Shprintzen, R.J. (2005). Defining the clinical spectrum of deletion 22q11.2.
J. Pediatr. *147*, 90–96.

72. Tang, K.L., Antshel, K.M., Fremont, W.P., and Kates, W.R. (2015). Behavioral and
psychiatric phenotypes in 22q11.2 deletion syndrome. J. Dev. Behav. Pediatr. *36*, 639–650.

73. Swillen, A., Devriendt, K., Legius, E., Eyskens, B., Dumoulin, M., Gewillig, M., and Fryns,
J.P. (1997). Intelligence and psychosocial adjustment in velocardiofacial syndrome: a study
of 37 children and adolescents with VCFS. J. Med. Genet. *34*, 453–458.

74. Stark, K.L., Xu, B., Bagchi, A., Lai, W.S., Liu, H., Hsu, R., Wan, X., Pavlidis, P., Mills, A.A.,
Karayiorgou, M., et al. (2008). Altered brain microRNA biogenesis contributes to phenotypic
deficits in a 22q11-deletion mouse model. Nat. Genet. *40*, 751–760.

75. Drew, L.J., Stark, K.L., Fénelon, K., Karayiorgou, M., Macdermott, A.B., and Gogos, J.A.
(2011). Evidence for altered hippocampal function in a mouse model of the human 22q11.2
microdeletion. Mol. Cell. Neurosci. *47*, 293–305.

76. Zaremba, J.D., Diamantopoulou, A., Danielson, N.B., Grosmark, A.D., Kaifosh, P.W.,
Bowler, J.C., Liao, Z., Sparks, F.T., Gogos, J.A., and Losonczy, A. (2017). Impaired
hippocampal place cell dynamics in a mouse model of the 22q11.2 deletion. Nat. Neurosci.
20, 1612–1623.

77. Sun, Z., Williams, D.J., Xu, B., and Gogos, J.A. (2018). Altered function and maturation of
primary cortical neurons from a 22q11.2 deletion mouse model of schizophrenia. Transl.
Psychiatry *8*. https://doi.org/10.1038/s41398-018-0132-8.

78. Geiller, T., Vancura, B., Terada, S., Troullinou, E., Chavlis, S., Tsagkatakis, G., Tsakalides,
P., Ócsai, K., Poirazi, P., Rózsa, B.J., et al. (2020). Large-Scale 3D Two-Photon Imaging of

Molecularly Identified CA1 Interneuron Dynamics in Behaving Mice. Neuron *108*, 968-983.e9.

79. Szalay, G., Judák, L., Katona, G., Ócsai, K., Juhász, G., Veress, M., Szadai, Z., Fehér, A., Tompa, T., Chiovini, B., et al. (2016). Fast 3D imaging of spine, dendritic, and neuronal assemblies in behaving animals. Neuron *92*, 723–738.

80. Katona, G., Szalay, G., Maák, P., Kaszás, A., Veress, M., Hillier, D., Chiovini, B., Vizi, E.S., Roska, B., and Rózsa, B. (2012). Fast two-photon in vivo imaging with three-dimensional random-access scanning in large tissue volumes. Nat. Methods *9*, 201–208.

81. Vancura, B., Geiller, T., Grosmark, A., Zhao, V., and Losonczy, A. (2023). Inhibitory control of sharp-wave ripple duration during learning in hippocampal recurrent networks. Nat. Neurosci. *26*, 788–797.

82. Somogyi, P., and Klausberger, T. (2005). Defined types of cortical interneurone structure space and spike timing in the hippocampus. J. Physiol. *562*, 9–26.

83. Guet-McCreight, A., Skinner, F.K., and Topolnik, L. (2020). Common principles in functional organization of VIP/calretinin cell-driven disinhibitory circuits across cortical areas. Front. Neural Circuits *14*, 32.

84. Varga, C., Golshani, P., and Soltesz, I. (2012). Frequency-invariant temporal ordering of interneuronal discharges during hippocampal oscillations in awake mice. Proc. Natl. Acad. Sci. U. S. A. *109*, E2726-34.

85. Lapray, D., Lasztoczi, B., Lagler, M., Viney, T.J., Katona, L., Valenti, O., Hartwich, K., Borhegyi, Z., Somogyi, P., and Klausberger, T. (2012). Behavior-dependent specialization of identified hippocampal interneurons. Nat. Neurosci. *15*, 1265–1271.

86. Katona, L., Lapray, D., Viney, T.J., Oulhaj, A., Borhegyi, Z., Micklem, B.R., Klausberger, T., and Somogyi, P. (2014). Sleep and movement differentiates actions of two types of somatostatin-expressing GABAergic interneuron in rat hippocampus. Neuron *82*, 872–886.

87. Somogyi, P., Katona, L., Klausberger, T., Lasztóczi, B., and Viney, T.J. (2014). Temporal redistribution of inhibition over neuronal subcellular domains underlies state-dependent rhythmic change of excitability in the hippocampus. Philos. Trans. R. Soc. Lond. B Biol. Sci. *369*, 20120518.

88. Turi, G.F., Li, W.-K., Chavlis, S., Pandi, I., Hare, J.O., Priestley, J.B., Grosmark, A.D., Liao, Z., Ladow, M., Zhang, J.F., et al. (2019). Vasoactive Intestinal Polypeptide-Expressing Interneurons in the Hippocampus Support Goal- Article Vasoactive Intestinal Polypeptide-Expressing Interneurons in the Hippocampus Support Goal-Oriented Spatial Learning. Neuron, 1–16.

89. Dudok, B., Klein, P.M., Hwaun, E., Lee, B.R., Yao, Z., Fong, O., Bowler, J.C., Terada, S., Sparks, F.T., Szabo, G.G., et al. (2021). Alternating sources of perisomatic inhibition during behavior. Neuron *109*, 997-1012.e9.

90. Friedrich, J., Zhou, P., and Paninski, L. (2017). Fast online deconvolution of calcium imaging data. PLoS Comput. Biol. *13*, e1005423.

91. Buzsáki, G., and Moser, E.I. (2013). Memory, navigation and theta rhythm in the hippocampal-entorhinal system. Nature Neuroscience 2013 16:2 *16*, 130–138.

92. O'Keefe, J., and Dostrovsky, J. (1971). The hippocampus as a spatial map. Preliminary evidence from unit activity in the freely-moving rat. Brain Res. *34*, 171–175.

93. Robinson, E.B., Samocha, K.E., Kosmicki, J.A., McGrath, L., Neale, B.M., Perlis, R.H., and Daly, M.J. (2014). Autism spectrum disorder severity reflects the average contribution of de novo and familial influences. Proc. Natl. Acad. Sci. U. S. A. *111*, 15161–15165.

94. Gonzalez-Burgos, G., and Lewis, D.A. (2012). NMDA receptor hypofunction, parvalbumin-positive neurons, and cortical gamma oscillations in schizophrenia. Schizophr. Bull. *38*, 950–957.

95. Buzsáki, G., and Draguhn, A. (2004). Neuronal oscillations in cortical networks. Science *304*, 1926–1929.

96. Van Derveer, A.B., Bastos, G., Ferrell, A.D., Gallimore, C.G., Greene, M.L., Holmes, J.T., Kubricka, V., Ross, J.M., and Hamm, J.P. (2021). A role for somatostatin-positive interneurons in neuro-oscillatory and information processing deficits in schizophrenia. Schizophr. Bull. *47*, 1385–1398.

97. Batiuk, M.Y., Tyler, T., Dragicevic, K., Mei, S., Rydbirk, R., Petukhov, V., Deviatiiarov, R., Sedmak, D., Frank, E., Feher, V., et al. (2022). Upper cortical layer-driven network impairment in schizophrenia. Sci. Adv. *8*, eabn8367.

98. Savage, J.E., Jansen, P.R., Stringer, S., Watanabe, K., Bryois, J., de Leeuw, C.A., Nagel, M., Awasthi, S., Barr, P.B., Coleman, J.R.I., et al. (2018). Genome-wide association meta-analysis in 269,867 individuals identifies new genetic and functional links to intelligence. Nat. Genet. *50*, 912–919.

99. Goff, K.M., and Goldberg, E.M. (2021). A role for vasoactive intestinal peptide interneurons in neurodevelopmental disorders. Dev. Neurosci. *43*, 168–180.

100. Mossner, J.M., Batista-Brito, R., Pant, R., and Cardin, J.A. (2020). Developmental loss of MeCP2 from VIP interneurons impairs cortical function and behavior. Elife *9*. https://doi.org/10.7554/eLife.55639.

101. Goff, K.M., Liebergall, S.R., Jiang, E., Somarowthu, A., and Goldberg, E.M. (2023). VIP interneuron impairment promotes in vivo circuit dysfunction and autism-related behaviors in Dravet syndrome. Cell Rep. *42*, 112628.

102. Luo, X., Guet-McCreight, A., Villette, V., Francavilla, R., Marino, B., Chamberland, S., Skinner, F.K., and Topolnik, L. (2020). Synaptic mechanisms underlying the network state-dependent recruitment of VIP-expressing interneurons in the CA1 hippocampus. Cereb. Cortex *30*, 3667–3685.

103. Kamigaki, T., and Dan, Y. (2017). Delay activity of specific prefrontal interneuron subtypes modulates memory-guided behavior. Nat. Neurosci. *20*, 854–863.

104. Möhler, H., and Rudolph, U. (2017). Disinhibition, an emerging pharmacology of learning
and memory. F1000Res. *6*, 101.

105. Lovett-Barron, M., and Losonczy, A. (2014). Behavioral consequences of GABAergic
neuronal diversity. Curr. Opin. Neurobiol. *26*, 27–33.

106. Murray, J.D., Anticevic, A., Gancsos, M., Ichinose, M., Corlett, P.R., Krystal, J.H., and Wang,
X.-J. (2014). Linking microcircuit dysfunction to cognitive impairment: effects of disinhibition
associated with schizophrenia in a cortical working memory model. Cereb. Cortex *24*, 859–
872.

107. Ioannidis, N.M., Rothstein, J.H., Pejaver, V., Middha, S., McDonnell, S.K., Baheti, S., Musolf,
A., Li, Q., Holzinger, E., Karyadi, D., et al. (2016). REVEL: An Ensemble Method for
Predicting the Pathogenicity of Rare Missense Variants. Am. J. Hum. Genet. *99*, 877–885.

108. Samocha, K.E., Kosmicki, J.A., Karczewski, K.J., O'Donnell-Luria, A.H., Pierce-Hoffman, E.,
MacArthur, D.G., Neale, B.M., and Daly, M.J. (2017). Regional missense constraint improves
variant deleteriousness prediction. bioRxiv, 148353. https://doi.org/10.1101/148353.

109. SPARK https://sparkforautism.org/.

110. Chiang, A.H., Chang, J., Wang, J., and Vitkup, D. (2021). Exons as units of phenotypic
impact for truncating mutations in autism. Mol. Psychiatry *26*, 1685–1695.

111. Lovett-Barron, M., Kaifosh, P., Kheirbek, M.A., Danielson, N., Zaremba, J.D., Reardon, T.R.,
Turi, G.F., Hen, R., Zemelman, B.V., and Losonczy, A. (2014). Dendritic inhibition in the
hippocampus supports fear learning. Science *343*, 857–863.

112. Acsády, L., Görcs, T.J., and Freund, T.F. (1996). Different populations of vasoactive
intestinal polypeptide-immunoreactive interneurons are specialized to control pyramidal cells
or interneurons in the hippocampus. Neuroscience *73*, 317–334.

113. Acsády, L., Arabadzisz, D., and Freund, T.F. (1996). Correlated morphological and
neurochemical features identify different subsets of vasoactive intestinal polypeptide-
immunoreactive interneurons in rat hippocampus. Neuroscience *73*, 299–315.

114. Tyan, L., Chamberland, S., Magnin, E., Camiré, O., Francavilla, R., David, L.S., Deisseroth,
K., and Topolnik, L. (2014). Dendritic inhibition provided by interneuron-specific cells controls
the firing rate and timing of the hippocampal feedback inhibitory circuitry. J. Neurosci. *34*,
4534–4547.

115. Pelkey, K.A., Chittajallu, R., Craig, M.T., Tricoire, L., Wester, J.C., and Mcbain, X.C.J. (2017).
Hippocampal GABAergic inhibitory interneurons. Physiol. Rev. *97*, 1619–1747.

116. Francavilla, R., Villette, V., Luo, X., Chamberland, S., Muñoz-Pino, E., Camiré, O., Wagner,
K., Kis, V., Somogyi, P., and Topolnik, L. (2018). Connectivity and network state-dependent
recruitment of long-range VIP-GABAergic neurons in the mouse hippocampus. Nat.
Commun. *9*, 5043.

117. Kaifosh, P., Zaremba, J.D., Danielson, N.B., and Losonczy, A. (2014). SIMA: Python
software for analysis of dynamic fluorescence imaging data. Front. Neuroinform. *8*, 80.

118. Kang, H.J., Kawasawa, Y.I., Cheng, F., Zhu, Y., Xu, X., Li, M., Sousa, A.M.M., Pletikos, M.,
Meyer, K.A., Sedmak, G., et al. (2011). Spatio-temporal transcriptome of the human brain.
Nature *478*, 483–489.