# Long-lasting coexistence of multiple asexual lineages alongside their
# sexual counterparts in a fungal plant pathogen

**Ammar Abdalrahem[1], Axelle Andrieux[1], Ronan Becheler[1,2], Sébastien Duplessis[1], Pascal**
**Frey[1], Benoit Marçais[1], Kadiatou Schiffer-Forsyth[1], Solenn Stoeckel[3], Fabien Halkett[1]**

[1] Université de Lorraine, INRAE, IAM, F-54000 Nancy, France
[2] INRAE, Agrocampus Ouest, Université de Rennes, IGEPP, Le Rheu, France
[3] DECOD (Ecosystem Dynamics and Sustainability), INRAE, Institut Agro, IFREMER, Rennes, France

# Abstract

Cases of sexual to asexual transitions within species are key to understanding the evolution of
reproductive modes, yet the coexistence of both modes in the same environment is poorly
documented. Here, we reported this transition in a plant pathogen species coexisting in the same
geographical environment. Our biological model is the poplar rust fungus *Melampsora larici-*
*populina,* which displays a complex life cycle typical of rust fungi (Pucciniales). It alternates
between two unrelated hosts to complete an obligate sexual life cycle once a year. We conducted
a comprehensive population genetic analysis, utilizing 21 microsatellite markers and data from
2,122 individuals gathered over 30 years from various locations in France. Our results
demonstrate the existence of many distinct lineages that reproduce asexually through the years,
skipping the sexual phase. A clustering analysis identified a group of multilocus lineages that
displayed all hallmarks of the genetic consequences of asexual reproduction, including highly
negative and large variance among loci of the inbreeding coefficient ($F_{IS}$). This indirect evidence
for asexual reproduction was confirmed by the direct observation of these asexual lineages being
repeatedly sampled across multiple years. This result demonstrates the coexistence of these
lineages with their sexual counterparts in the same ecological niche, challenging conventional
assumptions about geographical sorting of reproductive modes. These considerations are of
paramount importance for understanding the contemporary evolution of major pathogen species.
This switch from sexual to asexual reproduction has contributed to devastating epidemics
worldwide.

**Keywords**: sexual-asexual transition | evolution of reproductive modes | population genetics |
rust fungi | time samples.

# 1 Introduction

The evolution of sex remains a fundamental question and an intriguing topic in evolutionary biology (Culotta & Hines, 1998; Otto, 2009; Hartfield & Keightley, 2012). Sexual reproduction, which has benefits and costs, is widespread in eukaryotic organisms (Schurko et al., 2009). The key benefit is increased genetic diversity, purging deleterious mutations, and enhancing adaptation to changing environments (Pierre et al., 2022). However, it incurs costs such as time and energy to find mates, combined with the biological cost of meiosis (Roughgarden, 1991; Aanen & Hoekstra, 2007; Hörandl, 2013). Asexual reproduction, on the other hand, produces offspring without a mate, allowing for efficient reproduction, maintenance of lineages adapted to local stable conditions, and quick multiplication (Shick et al., 1979; Lodé, 2013). The offspring is genetically identical to the parent, barring somatic mutations, which renders asexual lineages more vulnerable to the accumulation of deleterious mutations due to a lack of recombination (Lynch et al., 1993; Peck, 1994; Keightley & Eyre-Walker, 2000), and more prone to loose allele diversity by genetic drift (Rouger et al., 2016). The study of species displaying both sexual and asexual reproduction enables pinpointing the pros and cons of each mode of reproduction (Butlin, 2002; Simon et al., 2010; Billiard et al., 2012). In particular, the study of cases where there is a dynamic transition from sexual to asexual reproductive modes is likely to highlight scenarios in which each reproductive mode could be favored.

The success of asexual reproduction frequently relies on specific environmental conditions. Remarkably, invasive species often shift towards asexual reproduction in their exotic range, while reproducing sexually in their native environment (Song et al., 2011; Platt & Jeschke, 2014; Tilquin & Kokko, 2016; Gladieux et al., 2015). Ant species can exhibit such diverse reproductive strategies, with asexual reproduction frequently being dominant in invasive populations, as observed in *Wasmannia auropunctata* and *Cerapachys biroi* (Foucaud et al., 2006; Kronauer et al., 2012). Plant pathogenic fungi also adopt diverse reproductive strategies, with asexual reproduction being responsible for the most damaging epidemics (Ashu & Xu, 2015; Gladieux et al., 2015; Drenth et al., 2019). Potato late blight is a devastating disease caused by the oomycete pathogen *Phytophthora infestans.* Both sexual and asexual reproduction take place in the native range, but it is asexual reproduction only that led to the devastating Irish Potato Famine in the 1840s (Gavino et al., 2000; Ristaino, 2002; Fry, 2020). The wheat yellow rust fungus *Puccinia*

*striiformis* f. sp. *tritici* displays a similar pattern, with different reproductive modes depending on

the region. The fungus reproduces sexually in China and Pakistan, which is highlighted by its

high genotypic diversity, whereas it reproduces asexually in the rest of the world (Bahri et al.,

2009; Mboup et al., 2009; Ali et al., 2014). The rice blast fungus *Magnaporthe oryzae* (Syn:

*Pyricularia oryzae*) reproduces sexually in a limited area of Southeast Asia, its native range,

whereas epidemics globally are driven by asexual lineages (Saleh et al., 2012).

The coexistence of both reproductive modes in the same geographical environment is still poorly

documented, despite it is offering a unique opportunity to decipher the evolutionary and

ecological dynamics of sexually or asexually derived lineages. While such studies are rare on

fungal biological models, they have been documented in other taxa, such as in aphids (Halkett et

al., 2006; Simon et al., 2010), daphnia (Lehto & Haag, 2010), and a cyprinid fish *Carassius*

*gibelio* (Barbuti et al., 2012). The phytopathogenic fungus *Melampsora larici-populina*, causing

poplar rust, represents an ideal candidate for elucidating the transition from one reproductive

mode to another, as both modes can be found in the same environment. *M. larici-populina*

belongs to the taxonomic order of Pucciniales (Steenackers et al., 1996). This species follows a

complex life cycle with an obligate sexual reproduction that alternates with rounds of clonal

multiplication (Figure S1). Two unrelated host plants are required to complete the life cycle

(Hacquard et al., 2011). Clonal multiplication occurs on poplar trees (*Populus* spp.), while sexual

reproduction takes place on larch trees (*Larix* spp.) (Lorrain et al., 2019). In the following,

lineages that complete the entire life cycle, including the obligate sexual reproduction phase, are

referred to as sexual lineages. The sexual reproduction on larch ensures genetic recombination

and obliterates all traces of the previous rounds of clonal multiplication (Xhaard et al., 2011;

Rouger et al., 2016). The impact of this fungal disease on larch is insignificant, whereas the

many rounds of clonal multiplication that occur on poplar leaves from spring to autumn can be

responsible for strong epidemics with large nutrient uptake by the fungus and perturbation of

photosynthesis (Jiang et al., 2016). Disease symptoms and damage are greatest on cultivated

poplars and resemble those of cereal rusts caused by several *Puccinia* species. A previous

population genetic study (Xhaard et al., 2011) has provided evidence for some variability in

reproductive mode, with an obligate sexual genetic group coexisting with the genetically distant

group that displays all hallmarks of long-lasting asexual reproduction. This suggests that some

lineages of the poplar rust fungus could survive alone through clonal multiplication on poplar.
Yet, this preliminary study did not provide detailed investigations on the ecological and
evolutionary dynamics of this asexual group, largely because it was based on a single year of
sampling.

In this study, we provide an in-depth characterization of the reproductive strategies of *M. larici-*
*populina*, with an emphasis on the (spatio-temporal) persistence of asexual lineages and their
genetic characteristics. Our study also examines the ecological and spatial factors influencing the
distribution of sexual and asexual populations, thereby offering insights into the evolution of
reproductive modes in rust fungi. This study encompasses the analysis of 2122 individuals
sampled across France over 30 years of population surveys that have been genotyped using 21
microsatellite markers.

We used a two-fold approach to assess the reproductive mode variation among poplar rust
individuals. First, we employed an indirect population genetic approach based on the clustering
among lineages and subsequent evaluation of the variation in population genetic indices known
to be signatures of asexual reproductive mode, namely: repeated genotypes, linkage
disequilibrium, and overall negative $F_{IS}$ value and large variance over loci (Balloux et al., 2003;
Halkett et al., 2005; Stoeckel & Masson, 2014). Second, we benefited from time-samples to
investigate the resampling through years of the same multilocus lineage (MLL), which is direct
evidence for their asexual survival. Furthermore, the genetic relatedness of all individuals was
analyzed to investigate evolutionary relationships among asexual lineages and with their sexual
counterparts, which points to multiple events of the emergence of asexual lineages. Finally,
examining the coexistence of sexual and asexual lineages, their long-term persistence, and their
spatial distribution suggested that environmental and ecological factors potentially play a role in
shaping the survival and maintenance of asexual lineages in this species.

# 2 Material and Methods

**2.1 Isolate collections**

In this study, we examined 2122 individuals of *M. larici-populina* sampled from 61 "départements" across France over 32 years (1992-2024). The dataset (Table S1) includes previously published genotypes from the study by Xhaard et al. (2011), sampled from a countrywide survey conducted in France in 2009 (23 locations | 428 samples), and more recent samples with genotype data original to this study. These recent additions comprise i) samples from a second countrywide survey conducted in 2011 (60 locations | 1374 samples), ii) individuals from the laboratory collection (1992, 1993, 1994, 1996, 1998, 2008, and 2014 | 34 samples), and genotypes from a project monitoring poplar rust epidemics in the Durance River valley (Xhaard et al., 2012; Becheler et al., 2016) that were suspected to belong to the asexual group and thus not included in the previously cited publications (2004, 2008 and 2020 | 246 samples). All these samples were collected in late summer, at the end of the clonal multiplication phase, therefore before leaf fall and lineage engagement into sexual reproduction (Figure S1). In addition, we benefit from the observation of a distinctive poplar stand in Cavalaire-sur-Mer (south of France) that maintains green leaves until mid-winter, where samples were collected in February 2024 (40 samples).

**2.2 Genotyping**

The DNA extraction process was guided by a protocol that employs the BioSprint 96 DNA Plant Kit and the BioSprint 96 automated workstation (Qiagen). Samples were genotyped using 21 microsatellite loci [Mlp_91, Mlp_83, Mlp_50, Mlp_71, Mlp_96, Mlp_95, Mlp_55, Mlp_66, Mlp_12, Mlp_68b Mlp_54, Mlp_57, Mlp_58, Mlp_49, Mlp_93, Mlp_82b Mlp_56, Mlp_87, Mlp_97, Mlp_77, Mlp_94 (Xhaard et al., 2009)]. These loci have been extensively utilized in several population genetic studies of this species (Xhaard et al., 2011; Becheler et al., 2017; Persoons et al., 2017; Louet et al., 2023; Saubin et al., 2024). The amplification through PCR with the Multiplex PCR Kit (Qiagen) was performed as described by Xhaard et al. (2011). It was conducted across three multiplex reactions containing eight, five, and eight loci, ranked by amplification size, respectively. The PCR products were then pooled and analyzed using an ABI 3730 Genetic Analyzers (Applied Biosystems) by the company GenoScreen. Fragment sizing

utilized a LIZ-1200 size standard, and allele scoring was performed with GENEMAPPER V4.0.

Samples that failed to amplify at more than six loci were excluded from further analysis.

**2.3 Population genetic and statistical analyses**

First, to handle genetic data of microsatellite markers in the R programming language v.4.4.1, we

converted the data format from a data frame to a genuine object using the `df2genind` function

from the **adegenet** package v.2.1.10. This function optimizes multilocus genotype data and

accommodates missing data if it is found in some individuals or loci. The automated workflow

code              provides              full              details              on              subsequent              analyses

(https://github.com/ammarabdalrahem/poplar_rust.git).    Moreover,    specific    analyses    were

executed using the GenAPoPop 1.0 software due to specific methodological requirements.

*Identification of repeated genotypes:* The prerequisite to an accurate assessment of reproductive

mode is the proper delineation of repeated genotypes derived from clonal multiplication (Halkett

et al., 2005; Arnaud-Haond et al., 2007). Individuals with the same combination of alleles (the

so-called multilocus genotypes, MLG) were consistently identified through both an ad hoc

procedure in Excel during the manual curation of chromatographs and using `mlg.filter` from

the **poppr** package v.2.9.3 (Kamvar et al., 2014). Genotypes differing by a few alleles were

double checked by returning to the original chromatographs. All remaining differences between

genotypes are therefore confirmed and reliable. We computed (*Pidu*) and (*N_Pidu)* to test for the

discrimination power of our marker set, and to rule out the probability of finding repeated MLGs

by chance (that is deriving from sexual reproduction) using GenAPoPop 1.0 (Evett & Weir,

1998; Waits et al., 2001; Stoeckel et al., 2024).

*Delineation of multilocus lineages:* As the accumulation of somatic mutations over time can lead

to slight differences among multilocus genotypes within a given asexual lineage (Arnaud-Haond

et al., 2007), we subsequently grouped MLGs into multilocus lineages (MLL). So as not to apply

an arbitrary threshold for MLL delineation, we first computed the distribution of pairwise shared

allele distance as advised by Arnaud-Haond et al. (2007), and Bailleul et al. (2016). As intra

lineage differentiation can vary across MLLs, we chose to use the UPGMA clustering method,

which aggregates MLGs one by one by examining the nearest neighbors until the MLL

boundaries are found (Kamvar et al., 2014, 2015). The threshold on genetic distance between

neighbor genotypes was set using the function `cutoff_predictor` from **poppr**.

*Clustering procedure to indirectly identify asexual lineages:* As the asexual reproductive mode

freezes genotypes, it retains ancestral allelic states. Asexual lineages can thus rapidly diverge

from their sexual counterparts, the latter being still influenced by genetic drift (Rouger et al.,

2016). In line with this statement, we previously found large genetic differences between sexual

and asexual groups of *M. larici-populina* (Xhaard et al., 2011). Assigning poplar rust individuals

would thus be an effective approach to identify asexual lineages without any *a priori*. The main

pitfall of this method being that the clustering algorithm should not rely on gathering individuals

according to Hardy-Weinberg expectations, which is the common assumption of any Structure-

like approach (Pritchard et al., 2000). We thus carry out our clustering analysis using a PCA and

discriminant analysis approach based on allelic distribution (Jombart et al., 2010). We used the

`find.clusters` function from the **adegenet** package to reduce the dimensionality of the

dataset. Discriminant analysis of principal components (DAPC) was then used to refine the

identification of clusters by maximizing the variance between two clusters, while minimizing

within cluster variance (Jombart et al., 2010). Nonetheless, 97.86% of genetic variation among

individuals was captured by retaining 90 principal components (n.pca = 90). Individuals were

grouped into two clusters using the `dpca` function with *K*=2. To ensure the reliability of the

assignment, only individuals with a high membership coefficient (≥0.8) were retained for further

analyses of within-cluster genetic characteristics.

*Resampling multilocus lineages (MLLs) as direct evidence for asexual reproduction.* Observing

the same genotype over several years is an indication that this lineage survived asexually - unless

sexual reproduction could have resulted in the same combination of alleles by chance, which can

be ruled out by a *Pidu* analysis (Evett & Weir, 1998; Stoeckel et al., 2024). To track the

persistence of MLLs over time, we recorded the years each MLL was detected and calculated its

frequency across different years. A lineage that persisted for at least two years was considered

asexual. We tested the overlap between the direct and indirect approaches for assessing asexual

reproductive mode by performing Fisher's exact test (Clarkson et al., 1993) on the contingency

table of MLL sorting according to the direct or indirect approach by using the `fisher.test`

function from the **stats** package v.4.4.1. In the following, a given MLL is considered asexual if it

meets at least one of the two criteria: the assignment to an asexual cluster or the inter-annual
resampling over a minimum of two years.

*Computation of classical population genetic indices:* As asexual reproductive mode distorts
allele repartition within and among individuals, we computed several classical population genetic
indices used to characterize within cluster and within MLL genetic signatures. Observed
heterozygosity ($H_O$), unbiased expected heterozygosity ($H_E$) (Nei, 1978), genotype richness ($R$),
and multilocus linkage disequilibrium corrected for the number of loci $\bar{r_d}$ were calculated using
the **poppr** package (basic function). Allelic richness ($A_R$), and *F*-statistics, including the
inbreeding coefficient ($F_{IS}$) (Weir & Cockerham, 1984) were computed using the **hierfstat**
package v.0.5-11. The clonal evenness was measured by calculating the Pareto β index, which
summarizes the distribution in size of MLLs using GenAPoPop 1.0 (Arnaud-Haond et al., 2007;
Stoeckel et al., 2024).

*Spatial analysis of the repartition of sexual and asexual lineages:* To illustrate the geographical
distribution of sexual and asexual individuals across France, the **maps** package v3.4 was utilized.
This enabled us to obtain information not only on spatial distribution, but also on the potential
coexistence of the two reproduction modes within the same geographical area. Using the `glmer`
function from the **lme4** v.1.1-35.5 package, a generalized linear mixed model (GLMM) with a
binomial distribution and a logit as a link function was employed to assess the regression
coefficients (Bolker et al., 2009). To further test the relationship between the reproductive mode
variation and the geographic locations, we performed a regression analysis on the count of sexual
and asexual lineages per site. The independent variables (fixed effect) were the geographic
location represented by latitude and longitude. The effect of the year of sampling was added as a
random effect. This analysis was performed on the 2009 and 2011 data originating from the two
countrywide surveys, so as not to overrepresent certain locations. To further assess the
distribution and relative abundance of asexual MLLs across time and space, we performed
Fisher's exact test for fluctuation in MLL abundance across years and locations. Given the large
size of the data sets, we employed a Monte Carlo simulation approach with 100,000 replicates to
estimate p-values.

*Genetic relationships between and among sexual and asexual lineages:* To further elucidate the
evolutionary trajectories of sexual and asexual individuals, and to address the question of the
dynamics of transition from sexual to asexual reproduction and vice versa, a discrete matrix
based on the relative dissimilarity between individuals was calculated by `diss.dist` with the
**poppr** package to create a neighbor-joining tree (Saitou & Nei, 1987) using `nj` from the **ape**
package v5.8.1.

*Inference of the rate of clonality:* Finally, to estimate the rate of clonality in the *M. larici-*
*populina* populations, we employed the ClonEstiMate Bayesian method. This method analyzes
the change of genotype frequencies from the same population over generations (Becheler et al.,
2017). We calculated based on two scenarios: i) merging all individuals within a mixed
population (that is any individuals can survive asexually by chance), or ii) considering two
distinct sexual and asexual populations, based on the lineage sorting detailed above. Since the
ClonEstiMate method should be applied on temporal samplings of the same kind of populations,
we used the data collected during the countrywide surveys carried out in the years 2009 and
2011.

# 3 Results

## 3.1 Delineation of multilocus lineages

We identified 1475 multilocus genotypes (MLGs) over the 2121 genotyped individuals. The 21 markers we used provided enough statistical power to ensure unambiguous discrimination between genotypes. The overall probability of obtaining the same combination of alleles by chance under panmictic sexual reproduction (*Pidu*) is approximately $10^{-16}$. This implies that 118 million individuals would have to be genotyped before observing two sexually produced descendants with identical genotypes (*N_Pidu*). We thus confidently considered that all the repeated genotypes we found were clones. We then grouped closely related MLGs into MLLs to account for the few somatic mutations that could occur during clonal multiplication. The delineation of MLLs was undertaken using a genetic distance threshold of 4.6, as determined by the UPGMA clustering algorithm. This grouping led to the delineation of 1384 MLLs, so slightly fewer than the number of MLGs, showing that the vast majority of MLGs have no closely related genotype.

## 3.2 Reproductive mode assessment

*The clustering approach* indicated that almost all individuals exhibited a high probability of membership in one of two genotypic clusters: 445 individuals were unambiguously assigned to cluster one, while 1676 individuals were assigned to cluster two (Figure S2). Only one isolate had membership coefficients below the threshold and was excluded from further analyses.

A comprehensive population genetic analysis of each cluster (Table 1) was conducted to assess the value of key population genetic indices, thereby revealing a contrast between the two clusters. Cluster 1 exhibits a highly negative $F_{IS}$ value, low genotypic diversity, and high linkage disequilibrium (LD), all of which are indicative of a strong rate of clonal reproduction (Balloux et al., 2003; Halkett et al., 2005; Arnaud-Haond et al., 2007). The high variance of $F_{IS}$ across loci further points to rare events of recombination within a population reproducing essentially asexually over the long term. In contrast, cluster 2 exhibited slightly positive $F_{IS}$ values and higher genetic diversity, suggesting a higher degree of allelic recombination, a feature of sexual reproduction. The reduced linkage disequilibrium in this cluster further supported this

classification. Consequently, cluster 1 was labeled as asexual, and cluster 2 as sexual. It is

noteworthy that similar levels of observed heterozygosity were found in both clusters, but a

much reduced allelic richness and gene diversity in the asexual cluster 1. Further, the *Pareto β*

index was low (~ 0.2), indicating a small number of dominant MLGs, which was consistent with

high clonality in cluster 1. In the sexual cluster 2, the *Pareto β* was greater than 1, reflecting a

more even distribution of genotypes due to genetic recombination.

**Table 1.** Characteristics of the two genetic clusters identified in *M. larici-populina* individuals.

| | MLG | MLL | R | *Pareto β* | $A_R$ | $H_O$ | $H_E$ | $F_{IS}$ |
|---|---|---|---|---|---|---|---|---|
| ı | | | | | | | | |
| | 64 | 11 | 0.142 | 0.226 | 4.372 | 0.525 | 0.360 | -0.461 |
| 5 | 1411 | 1373 | 0.842 | 1.078 | 9.529 | 0.506 | 0.543 | 0.070 |
| ch | | | | | | | | |
| | 90 | 19 | 0.183 | 0.248 | 5.364 | 0.516 | 0.390 | -0.322 |
| 3 | 1385 | 1365 | 0.848 | 1.057 | 9.700 | 0.508 | 0.545 | 0.068 |
| ɔroaches | | | | | | | | |
| | 96 | 25 | 0.191 | 0.252 | 5.414 | 0.516 | 0.393 | -0.315 |
| 2 | 1379 | 1359 | 0.850 | 1.051 | 9.749 | 0.508 | 0.544 | 0.068 |

MLG*:* number of multilocus genotypes, MLL*:* number of multilocus lineages, *N*: number of individuals, *R*: genotypic diversity, *Pareto β*: evenness among repeated genotypes; $A_R$: allelic richness, $H_O$*:* observed heterozygosity, $H_E$*:* expected heterozygosity, $F_{IS}$: inbreeding coefficient, Var ($F_{IS}$): variance of inbreeding coefficient, and $\bar{r}_d$: multilocus linkage disequilibrium.

*The resampling approach* revealed two groups: 488 individuals with lineages spanning more
than one year due to asexual reproduction and 1,633 sexual individuals without resampled
lineages exceeding two years. The contingency table (Table 2), which compares the clustering
and resembling approaches, shows a discrepancy. Among the 54 individuals identified as sexual
through cluster analysis, subsequent resampling evidence validated their clonality. In contrast, 11
individuals were clustered as asexual; however not resampled.

**Table 2.** Contingency table showing the relationship between clustering and resampling
approaches for reproductive mode definition in the dataset. Values represent the number of
individuals assigned to each category.

|  | Asex (Resampling approach) | Sex (Resampling approach) |
|---|---|---|
| **Asex (Clustering approach)** | 434 | 11 |
| **Sex (Clustering approach)** | 54 | 1622 |

A Fisher's exact test was performed to evaluate the correlation between the assignment of
reproductive modes from clustering and resampling analyses. The results were highly significant
(p-value $< 2.2 \times 10^{-1\Box}$), thereby rejecting the null hypothesis of no association, with a 95%
confidence interval. This indicated a robust relationship between the two classification methods.
Therefore, a combination of clustering and resampling approaches was used to determine the
reproductive mode of each individual. The subsequent analysis involved the calculation of the
population genetic indices for both groups. The inference of reproductive mode using two
approaches as well as a combination of both, yielded consistent results over all population
genetic indices, confirming the robustness of the classification (Table 1). However, there are
slight differences in the values of population genetic indices among classification methods, with
the asexual group determined by the clustering approach revealing a stronger signature of
asexual reproduction.

## 3.3 Spatial distribution of reproductive modes

The spatial distribution of both sexual and asexual lineages of *M. larici-populina* individuals in France revealed distinct patterns (Figure 1). Sexual lineages were widespread across most regions, predominating in the northern and central parts of the country. In contrast, asexual lineages were more concentrated in the southern regions.

**Figure 1**

Geographic distribution of sexual and asexual lineages of *Melampsora larici-populina* individuals collected in 2009 and 2011 across France. The pie charts show the proportions of asexual (red) and sexual (blue) lineages at each sampling location. The size of each pie chart corresponds to the total number of individuals collected from that site.

To further investigate the factors that influence the prevalence of asexual lineages, a GLMM

binomial regression analysis was conducted. The analysis (Table 3) revealed a significant effect

of both latitude and longitude (Figure S3) on the proportion of sexual and asexual lineages, with

positive coefficients indicating that the relative abundance of asexual individuals increases with

lower latitude (the closer we get to the south of the country). Additionally, the random effect of

years (variance = 18.78, sd = 4.333) indicated that the temporal distribution of individuals was

not associated with a specific reproductive mode.

**Table 3.** Results of the binomial regression model predicting the effect of spatial distribution on

the reproductive mode of *M. larici-populina*

| *Predictor* | *Estimate* | *Std. Error* | *z-value* | *p-value* | *Significance* |
|---|---|---|---|---|---|
| (Intercept) | -22.99815 | 2.66959 | -8.615 | < 2e-16 | *** |
| Lat | 0.46035 | 0.04730 | 9.732 | < 2e-16 | *** |
| Long | 0.09671 | 0.02912 | 3.321 | 0.000896 | *** |

Estimate (regression coefficient), Std. Error (standard error of the estimation), z-value (z-statistic), p-value

(statistical significance), and Significance power indicated by asterisks (0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1

' ' 1).

The ClonEstiMate Bayesian analysis yielded insights into the rate of asexual reproduction in *M.*

*larici-populina*. The 99% credible interval from the posterior probability obtained for each

group's clonal rates. In the context of a mixed group, where sexual and asexual lineages are

merged into one population, the estimated clonal rate was between 0.22 to 0.31, indicating that

about 27% of the population reproduces asexually. When the analysis was performed under the

assumption of two distinct populations, the clonal rate was between 0.93 to 0.99 for the asexual

group and between 0 to 0.21 for the sexual group (Figure 2).

**Figure 2**

Ternary plot representing the relative proportions of selfing, outcrossing, and clonality rates in three different groups of mixed reproductive, asexual reproductive, and sexual reproductive in France (2009–2011). The color gradient reflects the posterior probability estimates.

**3.4 Genetic relationships between sexual and asexual lineages**

The dendrogram tree shows different clades for sexual and asexual lineages (Figure 3), highlighting the seven main asexual lineages. However, some asexual lineages also appeared as closely related sister branches from the sexual clade. Barring MLL9, one of the most abundant asexual lineages, all of these asexual MLLs, closely related to the sexual clade are of small size. These small asexual lineages were thus sporadic, unlike the seven most abundant asexual MLLs that show persistence through time (Figure 3, Table 4). These main asexual MLLs also displayed some reticulation, indicating they consist of different MLGs. In contrast, MLL5 (a most prevalent MLL but classified as sexual) consists of individuals sampled only in 2024 that have exactly the same genotype. These observations underscored the evolutionary relationships and temporal dynamics between reproductive modes within and among *M. larici-populina* lineages.

 **Figure 3**

Circular Neighbor-joining (NJ) tree based on microsatellite data (21 markers) illustrating the genetic
relationships among *M. larici-populina* individuals. The color of the branches indicates the reproductive mode,
with blue representing sexual and red representing asexual modes. The outer rings represent the year of sample
collection, with each height corresponding to a year. The nine most abundant multilocus lineages (MLLs) are
color-coded.

**3.5 Genetic structure and persistence of asexual lineages**

A more detailed analysis of the asexual group identified several distinct genetic lineages. Among
these, seven lineages were found to be the most abundant. The dominant lineages exhibited
varying levels of persistence over the years, as indicated by their range of collection years (Table
4). In particular, MLL1 was consistently detected in almost all collection years and across all
regions, highlighting its long-term persistence and widespread occurrence.

**Table 4.** Characteristics of the seven most abundant asexual multilocus lineages (MLLs) of *M.*
*larici-populina*.

| MLL | MLG | N | Year range | R | $A_R$ | $H_O$ | $H_E$ |
|-----|-----|-----|------------|-------|-------|-------|-------|
| 1 | 36 | 280 | 1992:2024 | 0.125 | 1.274 | 0.519 | 0.274 |
| 2 | 3 | 49 | 2014:2020 | 0.042 | 1.290 | 0.571 | 0.287 |
| 3 | 4 | 44 | 1996:2009 | 0.070 | 1.316 | 0.619 | 0.312 |
| 4 | 9 | 38 | 1996:2020 | 0.216 | 1.266 | 0.499 | 0.263 |
| 6 | 6 | 23 | 1996:2020 | 0.227 | 1.196 | 0.354 | 0.192 |
| 8 | 5 | 14 | 1993:2014 | 0.308 | 1.306 | 0.575 | 0.296 |
| 9 | 3 | 14 | 2009:2011 | 0.154 | 1.189 | 0.350 | 0.183 |

*MLL:* number of multilocus lineages, *MLG:* number of multilocus genotypes, N: number of individuals, *R*:
Genotypic diversity, $A_R$: Allelic richness, $H_O$: Observed heterozygosity, $H_E$: Expected heterozygosity.

The distribution of these lineages in France differed with some lineages being widespread and
others more restricted to some areas (Figure 4).

**Figure 4**

Distribution of asexual lineages over time and across locations. (A) The abundance of asexual MLLs in
different French administrative regions. (B) The abundance of asexual MLLs in different years.

Fisher's exact test revealed a highly significant association between MLL distribution and both
temporal and spatial factors (P = 1e-05). The analysis revealed that certain MLLs were
repeatedly observed in the same geographic regions and over multiple sampling years,
suggesting a pattern of persistence. Conversely, other MLLs appeared sporadic.

## 4 Discussion

The poplar rust fungus *M. larici-populina* is commonly regarded as an obligate sexual species that needs the presence of both poplars and larches to perform its annual life cycle (Pinon & Frey, 2005; Hacquard et al., 2011). However, a previous population genetic study suggested the existence of a distinct group of individuals that would survive asexually only on poplar hosts (Xhaard et al., 2011). Here i) we characterized the existence of an asexual group maintaining on the poplar hosts over the years, ii) we found that this asexual group is made up of several genetic lineages. We documented that some of these asexual lineages have persisted for nearly three decades despite their coexistence with sexual lineages in some places. These findings challenge conventional assumptions about the evolutionary dominance of sexual recombination in this species and offer a unique opportunity to investigate the transitions from sexual to asexual reproduction in a plant pathogenic fungus.

In the present study, two complementary approaches were employed to reveal and characterize asexual individuals. A clustering approach based on differences in allelic frequency and probability of membership was combined with a resampling approach aiming to identify the persistence of lineages over the years. We observed a strong overlap between these indirect and direct approaches, which enabled a sound delineation of asexual lineages among *M. larici-populina* individuals. This result further supports the reliability of the clustering approach performed in the initial study of (Xhaard et al., 2011) to identify asexual lineages indirectly. Applying such an individual based clustering approach is all the more relevant that sexual and asexual lineages can be admixed in the same location (Halkett et al., 2006). Consequently, the clustering approach can be regarded as a robust and independent method for detecting cryptic population genetic structures. In addition to the genetic differentiation between sexual and asexual clusters caused by the lack of gene flow, several genetic signatures clearly distinguished asexual lineages thereby reflecting the fundamental differences in evolutionary trajectories between these reproductive modes (Balloux et al., 2003; Halkett et al., 2005; Rouger et al., 2016). Sexual reproduction ensures the convergence toward Hardy-Weinberg equilibrium (HWE) with recombination among loci that reduces linkage disequilibrium (LD). Asexual reproduction, on the other hand, distorts both allelic and genotypic frequencies mostly because it freezes genotypes (Stoeckel & Masson, 2014; Rouger et al., 2016). As a result, the best way to

identify asexual lineages is to apply a multi-criteria method (Kronauer et al., 2012) in which the
(diploid) population displays the following hallmarks (Halkett et al., 2005; Arnaud-Haond et al.,
2007): a reduced genotypic diversity, combined with a significant excess of heterozygotes
(negative $F_{IS}$ values). It should be noted that we also observed a very high variance of $F_{IS}$ across
loci, as expected when some recombination events seldom occur (Balloux et al., 2003; Stoeckel
& Masson, 2014; Stoeckel et al., 2021). Here, the values found for these indices are consistent
with a very high asexuality rate as inferred by ClonEstiMate analysis. The asexual group also
displayed significant linkage disequilibrium among loci, which is often reported in asexual and
partially asexual species, despite not being consistently expected in neutrally evolving
populations (Stoeckel et al., 2021). Linkage disequilibrium was also observed in the sexual group
along with a positive $F_{IS}$ value as the result of a Wahlund effect. Indeed, the sexual group
encompasses different genetic subgroups that have been identified by previous population
genetics studies (Xhaard et al., 2011; Persoons et al., 2017) and which are not considered here.
We observed clone mates in the sexual group too, as these can form lineages as abundant as
those in the asexual group, with the notable difference that they occur always at a unique
sampling site and that, above all, they are never resampled over the years in spite of their local
abundance. This local abundance of clone-mate lineages is in accordance with the sampling
mostly done in the autumn, at the end of the clonal epidemic phase of *M. larici-populina* life
cycle. With regards to this, it should be recalled that the mandatory sexual reproduction that
takes place on larch resets the population genetic indices in the sexual group each year (Xhaard
et al., 2011; Rouger et al., 2016).The contrast in population genetic indices provide a sound
framework for distinguishing reproductive modes, supporting the reliable assignment of
individuals into either sexual or asexual groups and providing the first insights into their
respective evolutionary history (Xhaard et al., 2011).

In accordance with the previous study of Xhaard et al. (2011), we found that the asexual group
was unevenly distributed across the country, and mostly occurred in southern France. Asexual
lineages have often been described as distributed at the margins of a species' natural range
(Simon et al., 1999; Halkett et al., 2005; Drenth et al., 2019; Pereyra et al., 2023), a phenomenon
called the geographic parthenogenesis (Peck et al., 1998). Poplar rust is intimately dependent
upon the presence of its two hosts. Although the native range of European Larch (*Larix decidua*)

is restricted to the Alps and some other Central-European Mountain ranges, it has been widely
planted in many parts of France in mountains and plains since the 18th century. Nevertheless, it
is less frequent in the Mediterranean area, which can therefore be considered as a margin for the
poplar rust fungus. If we compare our results with those of other rust fungi, particularly those
infecting wheat, we can see a larger spatial segregation between the sexual lineages found only
in the disease's center of origin and the asexual lineages that have invaded the crops worldwide.
Beyond rusts, this observation is common to many pathogenic species as for the ascomycete
fungus responsible for rice blast (Saleh et al., 2014), or for the oomycete responsible for potato
late blight (Ristaino, 2002). Interestingly, asexual lineages in these cases are specialized on
different cultivars, and therefore on different host genotypes (Goyeau et al., 2007), and the
relative abundance of asexual lineages therefore fluctuates over time with the use or disuse of
these cultivars (Fontyn et al., 2022). This shows to which extent the asexual evolution of these
plant pathogen species is associated with human practices. By contrast, in *M. larici-populina*,
asexual lineages are sampled mostly on wild poplar trees and are thus found on different host
genotypes. We thus can argue that this rust species enables the incipient stage of sexual to
asexual transition to be documented.

A puzzling observation is that asexual lineages of *M. larici-populina* coexist with their sexual
counterparts in most sampling sites. The absence of larch trees in some areas can be
compensated for by the very high dispersal abilities of this species (Barrès et al., 2008), which
can be wind dispersed over hundreds of kilometers in a year  (Saubin et al., 2024).This can
readily explain why sexual and asexual lineages are admixed even if no larch tree exists in the
close vicinity of the sampling site. Another non-exclusive hypothesis to account for the long-
lasting coexistence of sexual and asexual lineages is that of a temporal fluctuation in the
environmental conditions that favor the ones over the others. This temporal fluctuation could be
responsible for the fluctuations in abundance we observed, with almost half of the asexual
lineages sampled in the year 2020 alone. These fluctuations in relative sexual vs. asexual
abundances could be linked to peculiar meteorological events that would desynchronize the
alternation between poplar and larch during the sexual life cycle and disadvantage sexual
lineages that have to complete this complex life cycle. An outstanding example of the effect of a
hazardous event on the fate of sexual and asexual lineages was documented in the red alga

*Agarophyton chilense* along the Chilean coasts (Becheler et al., 2020). This study monitored wild
(mostly sexual) and cultivated (asexually propagated) populations of this red alga before and
after a devastating earthquake. Although sexual populations paid a large tribute immediately
after the catastrophe, the artificial maintenance of an asexual population caused the algae culture
to collapse in the long term, suggesting that each mode of reproduction plays a role in
maintaining a dynamic equilibrium.

The distribution of asexual lineages of *M. larici-populina* prevailing in southern France points to
the first mechanism that can explain their maintenance and diversification. In milder climates the
asexual survival could be facilitated by a longer growing period of poplars, creating a so-called
green bridge that would enable the rust lineages to jump over the winter from poplar leaves to
poplar leaves. According to this hypothesis, the asexual lineages would passively bypass the
sexual phase on larch, without the need for any genetic determinism. In support of this
hypothesis, it should be noted that *M. larici-populina* urediniospores can survive for 10 months
and still be able to infect poplar leaves (Taris, 1968). But in these experiments, rust-infected
poplar leaves were stored either in a greenhouse, or in a wire cage 2 meters above ground level,
which does not mimic real natural overwintering conditions. According to this neutral
hypothesis, both the onset and the survival of asexual lineages would be a matter of chance only.
In support of this hypothesis, we observed that some asexual lineages branched among the sexual
cluster and that these lineages have only been revealed through the resampling over years,
indicating they are in very close genetic proximity with the lineages that undergo obligate sexual
reproduction each year.

The persistence of several asexual lineages over decades, attesting to their evolutionary
maintenance and relative success, could also be indicative of some genetic determinism favoring
their survival over winter. In support of this hypothesis, the trade-off between overwintering
ability and multiplication during the epidemic season has been documented in other fungal
species (Hamelin et al., 2016; Suffert et al., 2018), and proven to account for the coexistence and
evolutionary maintenance of sexual and asexual lineages (Castel et al., 2014). It is also likely that
those long-lasting asexual lineages could have lost, at least to some degree, their ability to invest
in sexual reproduction, maybe to the benefit of greater asexual survival. Our population genetic
analysis, and especially the high variance of $F_{IS}$, would nonetheless indicate that recombination

does not occur among asexual lineages and thus that they do not evolve under strict asexuality
but rather as a succession of clones maintaining on poplar leaves over winter, renewed over the
years. Furthermore, the rare engagement of asexual lineages into the sexual reproduction can
bring additional explanations for the diversity of asexual lineages we observed. In line with the
concept of contagious asexuality observed in other organisms with mixed reproductive modes,
such as aphids and daphnia (Simon et al., 2002, 2003; Halkett et al., 2006, 2008), asexual
lineages that invest in sexual reproduction could be able to produce offspring that inherit their
propensity to survive asexually (Halkett et al., 2008). Deciphering further the history and
evolutionary success of these asexual lineages calls for a dedicated experiment to test their
performance and investment throughout the life cycle. While the complex life cycle of *M. larici-*
*populina* can be implemented in the laboratory (Pernaci et al., 2014), it remains a challenging
experiment that is still ongoing.

The main original result of our study was to document the diversity among asexual lineages in
*M. larici-populina*, with some lineages persisting over decades and others appearing more
sporadically. The fact that two sampling campaigns covered the entire French territory, and that
other sites were sampled regularly over more than 30 years is evidence for this pattern not being
due to sampling bias. Notably, differences in abundance correlate with differences in the
genotypic structure between lineages. We documented that asexual MLLs can consist of
different genotypes (MLGs), unlike lineages within the sexual group that primarily consist of a
singular repeated MLG. While some persistent asexual lineages, such as MLL1, showed
relatively high genotypic diversity over time. Here it is important to state that only slight
differences distinguish MLGs within asexual MLLs, in the vast majority of cases up to three loci
differ only. Moreover, the changes consist in either heterozygous to homozygous conversion or
to increase or decrease in allele length by a one step mutation. This regular pattern of allele
changes, found across all MLGs within MLLs, supports the hypothesis that the asexual lineages
evolved through single mutation steps.  Although mutation rates for the microsatellite markers
used in *M. larici-populina* have not yet been estimated, this regular pattern could present an
opportunity to apply a tip calibration approach of molecular clock analysis to estimate the timing
of the emergence and persistence of these asexual lineages (Rieux & Balloux, 2016). Yet, the
differential abundance and genotypic diversity among asexual lineages reflects heterogeneity

within asexual population dynamics and emphasizes the necessity to consider each lineage
independently, rather than presuming that the whole asexual population follows the same
evolutionary trajectory. The most parsimonious hypothesis is that this diversity has been
generated by the repeated emergence of asexual lineages through the evolutionary mechanisms
previously discussed (chance of survival or contagious asexuality). Among all asexual lineages,
MLL1 stood out as the most abundant lineage, being repetitively sampled from 1992 and
showing an outbreak in 2020. It was detected in all regions where asexual lineages were
sampled, and forms the only contribution to the unexpected occurrence of asexual lineages
northward in the center of France, nearby Paris. The genome of a reference asexual isolate,
representative of this MLL, has recently been sequenced, and analysis of its genome structure,
polymorphisms, and gene content will further document the tale of this dominant asexual
lineage.

## Data

To ensure reproducibility, all analyses were executed using R version 4.2.2. The relevant scripts and data can be accessed at https://github.com/ammarabdalrahem/poplar_rust.git.

## Author Contributions

Sébastien Duplessis, Pascal Frey, and Fabien Halkett conceived and designed the study. Axelle Andrieux designed and performed most of the molecular experiments. Ronan Becheler, Ammar Abdalrahem, Axelle Andrieux, and Fabien Halkett performed the manual curation of genotyping results. Axelle Andrieux, Ronan Becheler, and Fabien Halkett did the preliminary analyses of the data. Ammar Abdalrahem performed the final data analyses with help and advice from Benoit Marçais, Ronan Becheler, Solenn Stoeckel, and Fabien Halkett. Ammar Abdalrahem, Kadiatou Schiffer-Forsyth, and Fabien Halkett prepared the manuscript. All authors revised and approved the manuscript.

## Acknowledgments

We thank Jean Pinon for having created the historical poplar rust collection at INRAE Nancy in the 1980s and for having sampled the Cavalaire-sur-Mer population in 2014 and 2024. We warmly thank Asmaa A. Eltahlawy and Scott Heslop for proofreading an earlier version of the manuscript. This long-term research program was supported by several grants from the French National Research Agency (ANR-11-BSV7-0007, Clonix project; ANR-12-ADAP-0009, GANDALF project; ANR-13-BSV7-0011, FUNFIT project; ANR-18-CE32-0001, Clonix2D project; ANR-23-CE20-0032, ENDURANCE project). Ronan Becheler was supported by a postdoctoral fellowship from both the Region Lorraine and the French National Research Agency (CLONIX project, ANR-11-BSV7-007). Ammar Abdalrahem was supported by a PhD fellowship from the French Ministry of Education and Research (MESR).

561 We have used DeepL, Grammarly, and ChatGPT to proofread, suggest synonyms, check
562 grammar and spelling, and debug R script. All suggestions made by these tools have been
563 scrutinized and cross-checked by the authors.

## Conflicts of Interest

565 The authors declare no conflicts of interest

## Benefit-Sharing Statement

567 The advantage of this study is that its analysis code is an automated workflow and data are
568 publicly accessible.

# References

Aanen, D. K., & Hoekstra, R. F. (2007). Why Sex Is Good: On Fungi and Beyond. In *Sex in Fungi* (pp. 527–534). John Wiley & Sons, Ltd. https://doi.org/10.1128/9781555815837.ch32

Ali, S., Gladieux, P., Rahman, H., Saqib, M. S., Fiaz, M., Ahmad, H., Leconte, M., Gautier, A., Justesen, A. F., Hovmøller, M. S., Enjalbert, J., & de Vallavieille-Pope, C. (2014). Inferring the contribution of sexual reproduction, migration and off-season survival to the temporal maintenance of microbial populations: A case study on the wheat fungal pathogen uccinia striiformis f.sp. tritici. *Molecular Ecology*, *23*(3), 603–617. https://doi.org/10.1111/mec.12629

Arnaud-Haond, S., Duarte, C. M., Alberto, F., & Serrão, E. A. (2007). Standardizing methods to address clonality in population studies. *Molecular Ecology*, *16*(24), 5115–5139. https://doi.org/10.1111/j.1365-294X.2007.03535.x

Ashu, E. E., & Xu, J. (2015). The roles of sexual and asexual reproduction in the origin and dissemination of strains causing fungal infectious disease outbreaks. *Infection, Genetics and Evolution*, *36*, 199–209. https://doi.org/10.1016/j.meegid.2015.09.019

Bahri, B., Leconte, M., Ouffroukh, A., De Vallavieille-Pope, C., & Enjalbert, J. (2009). Geographic limits of a clonal population of wheat yellow rust in the Mediterranean region. *Molecular Ecology*, *18*(20), 4165–4179. https://doi.org/10.1111/j.1365-294X.2009.04267.x

Bailleul, D., Stoeckel, S., & Arnaud-Haond, S. (2016). RClone: A package to identify
MultiLocus Clonal Lineages and handle clonal data sets in r. *Methods in Ecology and*
*Evolution*, *7*(8), 966–970. https://doi.org/10.1111/2041-210X.12550

Balloux, F., Lehmann, L., & de Meeûs, T. (2003). The Population Genetics of Clonal and
Partially Clonal Diploids. *Genetics*, *164*(4), 1635–1644.
https://doi.org/10.1093/genetics/164.4.1635

Barbuti, R., Mautner, S., Carnevale, G., Milazzo, P., Rama, A., & Sturmbauer, C. (2012).
Population dynamics with a mixed type of sexual and asexual reproduction in a
fluctuating environment. *BMC Evolutionary Biology*, *12*(1), 49.
https://doi.org/10.1186/1471-2148-12-49

Barrès, B., Halkett, F., Dutech, C., Andrieux, A., Pinon, J., & Frey, P. (2008). Genetic structure
of the poplar rust fungus Melampsora larici-populina: Evidence for isolation by distance
in Europe and recent founder effects overseas. *Infection, Genetics and Evolution*, *8*(5),
577–587. https://doi.org/10.1016/j.meegid.2008.04.005

Becheler, R., Guillemin, M.-L., Stoeckel, S., Mauger, S., Saunier, A., Brante, A., Destombe, C.,
& Valero, M. (2020). After a catastrophe, a little bit of sex is better than nothing: Genetic
consequences of a major earthquake on asexual and sexual populations. *Evolutionary*
*Applications*, *13*(8), 2086–2100. https://doi.org/10.1111/eva.12967

Becheler, R., Masson, J.-P., Arnaud-Haond, S., Halkett, F., Mariette, S., Guillemin, M.-L.,
Valero, M., Destombe, C., & Stoeckel, S. (2017). ClonEstiMate, a Bayesian method for

609 quantifying rates of clonality of populations genotyped at two-time steps. *Molecular*

610 *Ecology Resources*, *17*(6), e251–e267. https://doi.org/10.1111/1755-0998.12698

611 Becheler, R., Xhaard, C., Klein, E. K., Hayden, K. J., Frey, P., De Mita, S., & Halkett, F. (2016).

612 Genetic signatures of a range expansion in natura: When clones play leapfrog. *Ecology*

613 *and Evolution*, *6*(18), 6625–6632. https://doi.org/10.1002/ece3.2392

614 Billiard, S., López-Villavicencio, M., Hood, M. E., & Giraud, T. (2012). Sex, outcrossing and

615 mating types: Unsolved questions in fungi and beyond. *Journal of Evolutionary Biology*,

616 *25*(6), 1020–1038. https://doi.org/10.1111/j.1420-9101.2012.02495.x

617 Bolker, B. M., Brooks, M. E., Clark, C. J., Geange, S. W., Poulsen, J. R., Stevens, M. H. H., &

618 White, J.-S. S. (2009). Generalized linear mixed models: A practical guide for ecology

619 and evolution. *Trends in Ecology & Evolution*, *24*(3), 127–135.

620 https://doi.org/10.1016/j.tree.2008.10.008

621 Butlin, R. (2002). The costs and benefits of sex: New insights from old asexual lineages. *Nature*

622 *Reviews Genetics*, *3*(4), 311–317. https://doi.org/10.1038/nrg749

623 Castel, M., Mailleret, L., Andrivon, D., Ravigné, V., & Hamelin, F. M. (2014). Allee Effects and

624 the Evolution of Polymorphism in Cyclic Parthenogens. *The American Naturalist*,

625 *183*(3), E75–E88. https://doi.org/10.1086/674828

626 Clarkson, D. B., Fan, Y., & Joe, H. (1993). A remark on algorithm 643: FEXACT: an algorithm

627 for performing Fisher's exact test in r x c contingency tables. *ACM Trans. Math. Softw.*,

628 *19*(4), 484–488. https://doi.org/10.1145/168173.168412

Culotta, E., & Hines, P. (1998). The Evolution of Sex. *Science*, *281*(5385), 1979–1979.

https://doi.org/10.1126/science.281.5385.1979

Drenth, A., McTaggart, A. R., & Wingfield, B. D. (2019). Fungal clones win the battle, but

recombination wins the war. *IMA Fungus*, *10*(1), 18. https://doi.org/10.1186/s43008-019-

0020-8

Evett, I., & Weir, B. S. (Bruce S. ). (1998). *Interpreting DNA evidence: Statistical genetics for*

*forensic scientists*. Sinauer Associates. https://cir.nii.ac.jp/crid/1130282269131400192

Fontyn, C., Zippert, A.-C., Delestre, G., Marcel, T. C., Suffert, F., & Goyeau, H. (2022). Is

virulence phenotype evolution driven exclusively by Lr gene deployment in French

Puccinia    triticina    populations?    *Plant    Pathology*,    *71*(7),    1511–1524.

https://doi.org/10.1111/ppa.13599

Foucaud, J., Jourdan, H., Breton, J. L., Loiseau, A., Konghouleux, D., & Estoup, A. (2006).

RARE SEXUAL REPRODUCTION EVENTS IN THE CLONAL REPRODUCTION

SYSTEM OF INTRODUCED POPULATIONS OF THE LITTLE FIRE ANT.

*Evolution*, *60*(8), 1646–1657. https://doi.org/10.1111/j.0014-3820.2006.tb00509.x

Fry, W. E. (2020). Phytophthora infestans: The itinerant invader; "late blight": the persistent

disease. *Phytoparasitica*, *48*(1), 87–94. https://doi.org/10.1007/s12600-019-00778-3

Gavino, P. D., Smart, C. D., Sandrock, R. W., Miller, J. S., Hamm, P. B., Lee, T. Y., Davis, R.

M., & Fry, W. E. (2000). Implications of Sexual Reproduction for Phytophthora infestans

in the United States: Generation of an Aggressive Lineage. *Plant Disease*, *84*(7), 731–

735. https://doi.org/10.1094/PDIS.2000.84.7.731

Gladieux, P., Feurtey, A., Hood, M. E., Snirc, A., Clavel, J., Dutech, C., Roy, M., & Giraud, T.

(2015). The population biology of fungal invasions. *Molecular Ecology*, *24*(9), 1969–

1986. https://doi.org/10.1111/mec.13028

Goyeau, H., Halkett, F., Zapater, M.-F., Carlier, J., & Lannou, C. (2007). Clonality and host

selection in the wheat pathogenic fungus *Puccinia triticina*. *Fungal Genetics and*

*Biology*, *44*(6), 474–483. https://doi.org/10.1016/j.fgb.2007.02.006

Hacquard, S., Petre, B., Frey, P., Hecker, A., Rouhier, N., & Duplessis, S. (2011). The Poplar-

Poplar Rust Interaction: Insights from Genomics and Transcriptomics. *Journal of*

*Pathogens*, *2011*, 716041. https://doi.org/10.4061/2011/716041

Halkett, F., Kindlmann, P., Plantegenest, M., Sunnucks, P., & Simon, J. C. (2006). Temporal

differentiation and spatial coexistence of sexual and facultative asexual lineages of an

aphid species at mating sites. *Journal of Evolutionary Biology*, *19*(3), 809–815.

https://doi.org/10.1111/j.1420-9101.2005.01055.x

Halkett, F., Plantegenest, M., Bonhomme, J., & Simon, J.-C. (2008). Gene flow between sexual

and facultatively asexual lineages of an aphid species and the maintenance of

reproductive    mode    variation.    *Molecular    Ecology*,    *17*(12),    2998–3007.

https://doi.org/10.1111/j.1365-294X.2008.03798.x

Halkett, F., Simon, J.-C., & Balloux, F. (2005). Tackling the population genetics of clonal and

partially clonal organisms. *Trends in Ecology & Evolution*, *20*(4), 194–201.

https://doi.org/10.1016/j.tree.2005.01.001

Hamelin, F. M., Bisson, A., Desprez□Loustau, M., Fabre, F., & Mailleret, L. (2016). Temporal

niche differentiation of parasites sharing the same plant host: Oak powdery mildew as a

case study. *Ecosphere*, *7*(11), e01517. https://doi.org/10.1002/ecs2.1517

Hartfield, M., & Keightley, P. D. (2012). Current hypotheses for the evolution of sex and

recombination. *Integrative Zoology*, *7*(2), 192–209. https://doi.org/10.1111/j.1749-

4877.2012.00284.x

Hörandl, E. (2013). Meiosis and the Paradox of Sex in Nature. In *Meiosis*. IntechOpen.

https://doi.org/10.5772/56542

Jiang, Y., Ye, J., Veromann, L.-L., & Niinemets, Ü. (2016). Scaling of photosynthesis and

constitutive and induced volatile emissions with severity of leaf infection by rust fungus

(Melampsora larici-populina) in Populus balsamifera var. Suaveolens. *Tree Physiology*,

*36*(7), 856–872. https://doi.org/10.1093/treephys/tpw035

Jombart, T., Devillard, S., & Balloux, F. (2010). Discriminant analysis of principal components:

A new method for the analysis of genetically structured populations. *BMC Genetics*,

*11*(1), 94. https://doi.org/10.1186/1471-2156-11-94

Kamvar, Z. N., Brooks, J. C., & Grünwald, N. J. (2015). Novel R tools for analysis of genome-
wide population genetic data with emphasis on clonality. *Frontiers in Genetics*, *6*.
https://www.frontiersin.org/articles/10.3389/fgene.2015.00208

Kamvar, Z. N., Tabima, J. F., & Grünwald, N. J. (2014). Poppr: An R package for genetic
analysis of populations with clonal, partially clonal, and/or sexual reproduction. *PeerJ*, *2*,
e281. https://doi.org/10.7717/peerj.281

Keightley, P. D., & Eyre-Walker, A. (2000). Deleterious Mutations and the Evolution of Sex.
*Science*, *290*(5490), 331–333. https://doi.org/10.1126/science.290.5490.331

Kronauer, D. J. C., Pierce, N. E., & Keller, L. (2012). Asexual reproduction in introduced and
native populations of the ant Cerapachys biroi. *Molecular Ecology*, *21*(21), 5221–5235.
https://doi.org/10.1111/mec.12041

Lehto, M. P., & Haag, C. R. (2010). Ecological differentiation between coexisting sexual and
asexual strains of Daphnia pulex. *Journal of Animal Ecology*, *79*(6), 1241–1250.
https://doi.org/10.1111/j.1365-2656.2010.01726.x

Lodé, T. (2013). Adaptive Significance and Long-Term Survival of Asexual Lineages.
*Evolutionary Biology*, *40*(3), 450–460. https://doi.org/10.1007/s11692-012-9219-y

Lorrain, C., Gonçalves dos Santos, K. C., Germain, H., Hecker, A., & Duplessis, S. (2019).
Advances in understanding obligate biotrophy in rust fungi. *New Phytologist*, *222*(3),
1190–1206. https://doi.org/10.1111/nph.15641

Louet, C., Saubin, M., Andrieux, A., Persoons, A., Gorse, M., Pétrowski, J., Fabre, B., De Mita,

S., Duplessis, S., Frey, P., & Halkett, F. (2023). A point mutation and large deletion at

the candidate avirulence locus AvrMlp7 in the poplar rust fungus correlate with poplar

RMlp7 resistance breakdown. *Molecular Ecology*, *23*(10), 2472–2483.

https://doi.org/10.1111/mec.16294

Lynch, M., Bürger, R., Butcher, D., & Gabriel, W. (1993). The Mutational Meltdown in Asexual

Populations. *Journal of Heredity*, *84*(5), 339–344.

https://doi.org/10.1093/oxfordjournals.jhered.a111354

Mboup, M., Leconte, M., Gautier, A., Wan, A. M., Chen, W., De Vallavieille-Pope, C., &

Enjalbert, J. (2009). Evidence of genetic recombination in wheat yellow rust populations

of a Chinese oversummering area. *Fungal Genetics and Biology*, *46*(4), 299–307.

https://doi.org/10.1016/j.fgb.2008.12.007

Nei, M. (1978). Estimation of Average Heterozygosity and Genetic Distance from a Small

Number of Individuals. *Genetics*, *89*(3), 583–590.

Otto, S. P. (2009). The Evolutionary Enigma of Sex. *The American Naturalist*, *174*(S1), S1–S14.

https://doi.org/10.1086/599084

Peck, J. R. (1994). A Ruby in the Rubbish: Beneficial Mutations, Deleterious Mutations and the

Evolution of Sex. *Genetics*, *137*(2), 597–606.

Peck, J. R., Yearsley, J. M., & Waxman, D. (1998). Explaining the geographic distributions of

sexual and asexual populations. *Nature*, *391*(6670), 889–892.

https://doi.org/10.1038/36099

Pereyra, R. T., Rafajlović, M., De Wit, P., Pinder, M., Kinnby, A., Töpel, M., & Johannesson, K.

(2023). Clones on the run: The genomics of a recently expanded partially clonal species.

*Molecular Ecology*, *32*(15), 4209–4223. https://doi.org/10.1111/mec.16996

Pernaci, M., De Mita, S., Andrieux, A., Pétrowski, J., Halkett, F., Duplessis, S., & Frey, P.

(2014). Genome-wide patterns of segregation and linkage disequilibrium: The

construction of a linkage genetic map of the poplar rust fungus Melampsora larici-

populina. *Frontiers in Plant Science*, *5*.

https://www.frontiersin.org/articles/10.3389/fpls.2014.00454

Persoons, A., Hayden, K. J., Fabre, B., Frey, P., De Mita, S., Tellier, A., & Halkett, F. (2017).

The escalatory Red Queen: Population extinction and replacement following arms race

dynamics in poplar rust. *Molecular Ecology*, *26*(7), 1902–1918.

https://doi.org/10.1111/mec.13980

Pierre, J.-S., Stoeckel, S., & Wajnberg, E. (2022). The advantage of sex: Reinserting fluctuating

selection in the pluralist approach. *PLOS ONE*, *17*(8), e0272134.

https://doi.org/10.1371/journal.pone.0272134

Pinon, J., & Frey, P. (2005). Interactions between poplar clone and Melampsora populations and

their implications for breeding for durable resistance. In *Rust Diseases of Willow and*

*Poplar* (pp. 139–154). https://doi.org/10.1079/9780851999999.0139

Platt, V., & Jeschke, J. M. (2014). Are exotic species red queens? *Ethology Ecology &*

*Evolution*, *26*(2–3), 101–111. https://doi.org/10.1080/03949370.2014.898699

Pritchard, J. K., Stephens, M., & Donnelly, P. (2000). Inference of Population Structure Using

Multilocus Genotype Data. *Genetics*, *155*(2), 945–959.

https://doi.org/10.1093/genetics/155.2.945

Rieux, A., & Balloux, F. (2016). Inferences from tip□calibrated phylogenies: A review and a

practical guide. *Molecular Ecology*, *25*(9), 1911–1924.

https://doi.org/10.1111/mec.13586

Ristaino, J. B. (2002). Tracking historic migrations of the Irish potato famine pathogen,

Phytophthora infestans. *Microbes and Infection*, *4*(13), 1369–1377.

https://doi.org/10.1016/S1286-4579(02)00010-2

Rouger, R., Reichel, K., Malrieu, F., Masson, J. P., & Stoeckel, S. (2016). Effects of complex

life cycles on genetic diversity: Cyclical parthenogenesis. *Heredity*, *117*(5), Article 5.

https://doi.org/10.1038/hdy.2016.52

Roughgarden, J. (1991). The Evolution of Sex. *The American Naturalist*, *138*(4), 934–953.

https://doi.org/10.1086/285261

Saitou, N., & Nei, M. (1987). The neighbor-joining method: A new method for reconstructing

phylogenetic trees. *Molecular Biology and Evolution*, *4*(4), 406–425.

https://doi.org/10.1093/oxfordjournals.molbev.a040454

Saleh, D., Milazzo, J., Adreit, H., Fournier, E., & Tharreau, D. (2014). South-East Asia is the
center of origin, diversity and dispersion of the rice blast fungus, Magnaporthe oryzae.
*New Phytologist*, *201*(4), 1440–1456. https://doi.org/10.1111/nph.12627

Saleh, D., Xu, P., Shen, Y., Li, C., Adreit, H., Milazzo, J., Ravigné, V., Bazin, E., Nottéghem, J.-
L., Fournier, E., & Tharreau, D. (2012). Sex at the origin: An Asian population of the rice
blast fungus Magnaporthe oryzae reproduces sexually. *Molecular Ecology*, *21*(6), 1330–
1344. https://doi.org/10.1111/j.1365-294X.2012.05469.x

Saubin, M., Coville, J., Xhaard, C., Frey, P., Soubeyrand, S., Halkett, F., & Fabre, F. (2024). A
mechanistic-statistical approach to infer dispersal and demography from invasion
dynamics, applied to a plant pathogen. *Peer Community Journal*, *4*.
https://doi.org/10.24072/pcjournal.356

Schurko, A., Neiman, M., & Logsdon, J. (2009). Signs of sex: What we know and how we know
it. *Trends in Ecology & Evolution*, *24*, 208–217.
https://doi.org/10.1016/j.tree.2008.11.010

Shick, J. M., HOFFMANN, R. J., & LAMB, A. N. (1979). Asexual Reproduction, Population
Structure, and Genotype-Environment Interactions in Sea Anemones. *American
Zoologist*, *19*(3), 699–713. https://doi.org/10.1093/icb/19.3.699

Simon, J.-C., Baumann, S., Sunnucks, P., Hebert, P. D. N., Pierre, J.-S., Le Gallic, J.-F., &
Dedryver, C.-A. (1999). Reproductive mode and population genetic structure of the
cereal aphid Sitobion avenae studied using phenotypic and microsatellite markers.
*Molecular Ecology*, *8*(4), 531–545. https://doi.org/10.1046/j.1365-294x.1999.00583.x

Simon, J.-C., Delmotte, F., Rispe, C., & Crease, T. (2003). Phylogenetic relationships between

parthenogens and their sexual relatives: The possible routes to parthenogenesis in

animals: ROUTES TO PARTHENOGENESIS IN ANIMALS. *Biological Journal of the*

*Linnean Society*, *79*(1), 151–163. https://doi.org/10.1046/j.1095-8312.2003.00175.x

Simon, J.-C., Rispe, C., & Sunnucks, P. (2002). Ecology and evolution of sex in aphids. *Trends*

*in Ecology & Evolution*, *17*(1), 34–39. https://doi.org/10.1016/S0169-5347(01)02331-X

Simon, J.-C., Stoeckel, S., & Tagu, D. (2010). Evolutionary and functional insights into

reproductive strategies of aphids. *Comptes Rendus. Biologies*, *333*(6–7), 488–496.

https://doi.org/10.1016/j.crvi.2010.03.003

Song, Y., Scheu, S., & Drossel, B. (2011). Geographic parthenogenesis in a consumer-resource

model for sexual reproduction. *Journal of Theoretical Biology*, *273*(1), 55–62.

https://doi.org/10.1016/j.jtbi.2010.12.020

Steenackers, J., Steenackers, M., Steenackers, V., & Stevens, M. (1996). Poplar diseases,

consequences on growth and wood quality. *Biomass and Bioenergy*, *10*(5), 267–274.

https://doi.org/10.1016/0961-9534(95)00121-2

Stoeckel, S., Becheler, R., Bocharova, E., & Barloy, D. (2024). GenAPoPop 1.0: A user-friendly

software to analyse genetic diversity and structure from partially clonal and selfed

autopolyploid organisms. *Molecular Ecology Resources*, *24*(1), e13886.

https://doi.org/10.1111/1755-0998.13886

Stoeckel, S., & Masson, J.-P. (2014). The Exact Distributions of FIS under Partial Asexuality in
Small    Finite    Populations    with    Mutation.    *PLOS    ONE*,    *9*(1),    e85228.
https://doi.org/10.1371/journal.pone.0085228

Stoeckel, S., Porro, B., & Arnaud-Haond, S. (2021). The discernible and hidden effects of
clonality on the genotypic and genetic states of populations: Improving our estimation of
clonal    rates.    *Molecular    Ecology    Resources*,    *21*(4),    1068–1084.
https://doi.org/10.1111/1755-0998.13316

Suffert, F., Goyeau, H., Sache, I., Carpentier, F., Gélisse, S., Morais, D., & Delestre, G. (2018).
Epidemiological trade☐off between intra☐ and interannual scales in the evolution of
aggressiveness in a local plant pathogen population. *Evolutionary Applications*, *11*(5),
768–780. https://doi.org/10.1111/eva.12588

Taris, B. (1968). Contribution à l'étude des rouilles des Populus observées en France. *Annales
Des Épiphyties*, *19*, 5–54.

Tilquin, A., & Kokko, H. (2016). What does the geography of parthenogenesis teach us about
sex? *Philosophical Transactions of the Royal Society B: Biological Sciences*, *371*(1706),
20150538. https://doi.org/10.1098/rstb.2015.0538

Waits, L. P., Luikart, G., & Taberlet, P. (2001). Estimating the probability of identity among
genotypes in natural populations: Cautions and guidelines. *Molecular Ecology*, *10*(1),
249–256. https://doi.org/10.1046/j.1365-294X.2001.01185.x

Weir, B. S., & Cockerham, C. C. (1984). Estimating F-Statistics for the Analysis of Population

Structure. *Evolution*, *38*(6), 1358–1370. https://doi.org/10.2307/2408641

Xhaard, C., Andrieux, A., Halkett, F., & Frey, P. (2009). Characterization of 41 microsatellite

loci developed from the genome sequence of the poplar rust fungus, Melampsora larici-

populina. *Conservation Genetics Resources*, *1*, 21–25. https://doi.org/10.1007/s12686-

009-9005-z

Xhaard, C., Barrès, B., Andrieux, A., Bousset, L., Halkett, F., & Frey, P. (2012). Disentangling

the genetic origins of a plant pathogen during disease spread using an original molecular

epidemiology approach. *Molecular Ecology*, *21*(10), 2383–2398.

https://doi.org/10.1111/j.1365-294X.2012.05556.x

Xhaard, C., Fabre, B., Andrieux, A., Gladieux, P., Barrès, B., Frey, P., & Halkett, F. (2011). The

genetic structure of the plant pathogenic fungus Melampsora larici-populina on its wild

host is extensively impacted by host domestication. *Molecular Ecology*, *20*(13), 2739–

2755. https://doi.org/10.1111/j.1365-294X.2011.05138.x

**Supplementary figures**

**Figure S1**

The *Melampsora larici-populina* life cycle. The blue arrows indicate the heteroecious life cycle, the clonal
multiplication phase on poplar tree leaves during summer, and the sexual phase on the larch tree. The dashed red
arrow indicates pure asexual multiplication on poplar, skipping the sexual phase.

**Figure S2**

844 Scatter plot illustrating the membership probability for individuals within each genotypic cluster, with the names of
845 individuals of lower probability.

**Figure S3**

The regression plot illustrates the relationship between latitude and the proportion of sexual reproduction. This
relationship was analyzed using a binomial generalized linear mixed model (GLMM). It was based on data from
2009 and 2011. The size of the points represents sample sizes.

# Supplementary Tables

**Table S1.** The data set for 2,122 *Melampsora larici-populina* individuals used in population
genetic analyses includes collection year, geographic location, and genotypes across 21
microsatellite markers.

Aecial host:
*Larix* sp.
(larch)

Telial host:
*Populus* sp.
(poplar)

Aeciospores (n+n)

Uredinia (n+n)

Urediniospores (n+n)

Aecia (n+n)

Plasmogamy

Spermatia (n)

Spring

Summer

Telia (n+n)
Teliospores (n+n)

Spermogonia (n)

Autumn & Winter

+

−

Basidiospores (n)

Karyogamy
(2n)

Meiosis

Basidia