# Elimination of senescent cells with senolytic host-directed therapy reduces tuberculosis progression in mice

List of authors.

Somnath Shee[1], Yazmin B. Martinez-Martinez[1], Benjamin Koleske[1], Shivraj Yabaji[2], Lester Kobzik[3], Igor Kramnik[2], William Bishai[1]

[1]Center for Tuberculosis Research, Johns Hopkins University, School of Medicine, Baltimore, MD, USA.

[2] The National Emerging Infectious Diseases Laboratories (NEIDL), Boston University, Boston, MA, USA.

[3] Cellecta, Inc, Mountain View, CA, USA

Short title:  Senolytic therapy for TB

Keywords.

Senescence, aging, Tuberculosis, Mycobacterium, Senolytics, Dasatinib, Quercetin, Fisetin, granuloma, necrosis, SASP, DNA damage

**Abstract**

By eliciting lung necrosis, which enhances aerosol transmission, *Mycobacterium tuberculosis* (*Mtb*) sustains its long-term survival as a human pathogen. In studying the human-like necrotic granuloma lesions characteristic of *Mtb*-infected *B6.Sst1S* mice, we found that lung myeloid cells display elevated senescence markers- cell cycle arrest proteins p21 and p16, the DNA damage marker γH2A.X, senescence-associated β-Galactosidase activity, and senescence-associated secretory phenotype (SASP). These markers were also elevated in *Mtb*-infected aged wild type (WT) mice but not in young WT mice. Global transcriptomics data revealed activation of pro-survival (PI3K, MAPK) and anti-apoptotic pathways in *Mtb*-infected *B6.Sst1S* macrophages. As senescent cells are long-lived, non-dividing cells that release tissue-damaging SASP, we treated *Mtb*-infected mice with a cocktail of three senolytic drugs (dasatinib, quercetin, and fisetin) designed to kill senescent cells. Senolytic drug treatment prolonged survival and reduced *Mtb* lung counts in *B6.Sst1S* and aged WT mice to a greater degree than young WT mice and concomitantly reduced lung senescence markers. These findings indicate that (1) *Mtb* infection may induce lung myeloid cells to enter a senescent state and that these cells play a causal role in disease progression, and (2) Senolytics merit consideration for human clinical trials against tuberculosis (TB).

## Introduction

Despite the introduction of 3 new anti-tuberculosis (TB) drugs since 2012, the global TB pandemic remains poorly controlled.[1] Most humans are resistant to TB and are capable of limiting the bacterial growth and inflammatory damage caused by Mycobacterium tuberculosis (*Mtb*) infection.[2–4] However, susceptible individuals develop active disease characterized by necrotic granulomatous lung lesions that may further transform into lung cavities, leading to lung damage and morbidity even with appropriate anti-TB treatment, and ultimately TB transmission.[5,6] Understanding the molecular basis for failed granuloma control and necrotic lesion formation is a major unmet need in TB research.

While it is well known that aging is a significant risk factor for progression from contained *Mtb* infection to active TB disease in both high and low-level income countries,[7–11] there have been relatively few studies investigating the mechanistic basis for age-associated active TB.[12–20] Senescence is a rapidly expanding research discipline as many investigators have sought to understand the mechanism of gradual tissue failure in age-associated chronic diseases such as dementia, heart failure, arthritis, and renal failure.[21] The accumulation of senescent cells is a principal mechanism of functional tissue impairment in aging.[22,23] A key lesson from the senescence literature is that aged tissues are populated by long-lived, non-dividing, functionally impaired senescent cells. These cells release deleterious signaling molecules- collectively termed senescence-associated secretory phenotype (SASP), that locally contribute to inflammation and tissue failure.[24] Indeed, transplantation of isolated senescent cells led to premature aging in mice.[25,26] In this context, small molecule senolytic drugs—designed to induce apoptosis

in the non-dividing senescent cells— prolong mice lifespans, and improve organ function

in humans with chronic disease.[27,28]

To gain insight into the factors that lead to the loss of immune containment of *Mtb*, we

study the *B6.Sst1S* mice, which develop human-like necrotic lung granulomas following

*Mtb* infection in contrast to most wild type (WT) mouse strains.[29–35] These mice harbor

the *Sst1S* allele, which has loss of function mutations in the Sp140 chromatin remodeling

protein that plays a key role in immune cell gene regulation.[36,37] Recently Kotov *et al.*

described a population of interstitial macrophages (IM) and plasmacytoid dendritic cells

(pDC) that aberrantly secrete type I IFN (associated with TB exacerbation)[38–40], are poorly

responsive to IFNγ and may play a key role in granuloma breakdown in the lungs of the

closely related Sp140$^{-/-}$ mouse strain.[41] In this paper, we report that lung myeloid cells in

*Mtb*-infected *B6.Sst1S* mice displayed high levels of senescence markers prior to necrotic

granuloma formation and subsequently appeared closely associated with non-controlling,

early necrotic granulomas. Similar senescence marker increases were found in *Mtb*-

infected aged WT C57BL/6 (B6) mice, but not young WT B6 mice. Treatment with a three-

drug cocktail of FDA-approved senolytic drugs to eliminate these senescent cells restored

containment of *Mtb* proliferation, prolonged mouse survival, and reduced senescence

markers. Thus, myeloid cell entry into a senescent state may drive the failure of immune

containment and progression to necrosis during pulmonary TB.

## Results

**Elevated senescence markers *in vitro* were detected in BMDMs from *B6.Sst1S* and aged WT B6 but not young WT B6 mice.**

We previously observed that TNFα exposure induced a stress response characterized by overexpression of IFNβ (Supplementary Fig. 1a) and the integrated stress response (ISR) in *B6.Sst1S* bone marrow-derived macrophages (BMDMs).[34,35] Because chronic ISR activation is a potential driver of cellular senescence,[42,43] we tested TNFα-stimulated BMDMs from young *B6.Sst1S* mice (8-12 weeks old) alongside BMDMs from young and aged (>72 weeks old) WT B6 mice (Fig. 1a). Not only did *B6.Sst1S* and aged WT B6 BMDMs showed poorer *Mtb* H37Rv containment following TNFα stress induction than young WT B6 (Fig. 1b), they also showed greater activity of senescence-associated β-galactosidase (SA-βGal) than young WT B6 BMDMs, although not to the level seen with etoposide, an inducer of DNA damage-mediated senescence (Fig. 1c, 1d). In addition, we observed the induction of SASP genes (*Il1β, Il1α, Il6, Mmp12, Mmp13, Ccl2, Ccl8, Pai1*) transcription in TNFα-treated *B6.Sst1S* BMDMs compared to similarly treated WT B6 cells (Fig. 1e). Moreover, TNFα- treated *B6.Sst1S* BMDMs and aged WT B6 BMDMs showed the presence of established biomarkers of senescence- higher levels of the cyclin-dependent kinase inhibitor P16 (cell cycle arrest), phospho(S139)-H2A.X (DNA damage marker) and lower protein levels of Lamin B1 (structural constituent of the nuclear lamina) (Supplementary Fig. 1b), and displayed reduced cell proliferation as measured by BrdU-incorporation assay (Fig. 1f) without a significant reduction in viability (Supplementary Fig. 1c). In sum, TNFα-treated *B6.Sst1S* BMDMs exhibited numerous

markers of senescence similar to BMDMs derived from aged WT B6 mice or etoposide-

treated WT B6 BMDMs.

**Transcriptomic signatures show upregulation of senescence markers in BMDMs**

**and cells from non-controlling lung lesions in *B6.Sst1S* mice.**

Reanalysis of transcriptomic signatures in *B6.Sst1S* versus WT B6 BMDMs with and

without TNFα treatment (GSE99456)[35] suggested upregulation of cellular senescence

pathways, enhanced DNA damage responses, activation of mitotic checkpoints, pro-

survival ERK1/ERK2 protein kinase cascade, and cellular response to IFNβ (by

Metascape Analysis;[44] Supplementary Fig. 2, Supplementary File S1) in *B6.Sst1S*

BMDMs. To examine whether these signatures appear in the context of active lung

infection, we performed spatial transcriptomics to profile myeloid cells associated with

controlling (C) versus non-controlling (NC) lesions in *B6.Sst1S* mouse lungs (Fig. 2a). To

do this, we first administered $1 \times 10^6$ CFU of *Mtb* H37Rv subcutaneously in the hock, and

lung samples were harvested at 14 weeks post-infection. We categorized the lesions into

non-controlling (NC; advanced multibacillary lesions with necrotic core) and controlling

(C; non-necrotic paucibacillary lesions) granulomas. We identified 734 differentially

expressed genes (DEGs; $log_2$fold change>1 or <-1, adjusted *p*< 0.05) in NC vs C lesions

(Fig. 2b, Supplementary File S2), and functional pathway enrichment analysis[44] identified

inflammatory response, extracellular matrix organization, lung fibrosis and negative

regulation of cell proliferation pathways to be significantly higher in NC lesions

(Supplementary Fig. 3a). NC lesion DEGs also showed a high degree of overlap with the

SenMayo[45] senescence-associated gene set (Supplementary Fig. 3b, 3c). Taken

together, our spatial transcriptomics data further supports the association of senescence

markers with progression to necrotic granuloma formation in *Mtb*-infected *B6.Sst1S* mouse lungs.

***In vivo* senescence markers are elevated in *Mtb*-infected young *B6.Sst1S* and aged WT B6 compared to young WT B6 mice, predominantly on myeloid cells at earlier time points.**

We infected young *B6.Sst1S* and WT B6 mice, aged WT B6 mice with a low-dose *Mtb* aerosol infection route (Fig. 2c). *Mtb* proliferation was significantly higher uniquely in young *B6.Sst1S* mice, with necrosis appearing as early as 4 weeks post-infection (wpi) (Fig. 2d, 2g). Both young *B6.Sst1S* and aged WT B6 showed increased dissemination to the spleen at 6 wpi, suggesting that aged WT B6 mice have a defect in containment compared to young WT B6 (Fig. 2e). At 2 wpi, when lung CFU counts were equivalent across all three groups, we observed a higher degree of immunopathology as measured by area of lung consolidation in the H&E-stained lungs of aged WT B6 (*p*= 0.03), and *B6.Sst1S* mice showed no difference in H&E stained area than in young WT B6 mice (Fig. 2f, 2h). Surprisingly, at 4 wpi we observed decreased pathology in aged WT B6 mice lungs (Fig. 2g, 2h).

To follow up on our *in vitro* BMDM findings with senescence markers, we sought to determine whether markers of senescence were elevated during *in vivo* mouse infections. We evaluated lung levels of γH2A.X and p21 at the 2 wpi-timepoint when no differences in CFU were noted (Fig. 2d) and observed that both young *B6.Sst1S* and aged WT B6 mice showed qualitatively higher immunostaining for γH2A.X and p21 by IHC (Fig. 2f), and these levels continued to increase at 4 wpi in *B6.Sst1S* mice but not aged WT B6 (Fig. 2g). Quantification of γH2A.X staining confirmed elevated levels for both *B6.Sst1S*

and aged WT B6 mice relative to young WT B6 mice at 2 wpi (Fig. 2i). To refine this

analysis, we turned to flow cytometric measurements of widely accepted senescence

markers (p21, p16 and SA-βGal)[46,47] with lung and spleen cells (Fig. 3a). To examine the

temporal relationship between senescence and elevated *Mtb* burden, we measured the

senescent cell populations at an early time point, 10 days post-infection (dpi) when lung

CFU burdens are the same across all groups (Supplementary Fig. 4a). We found that the

percent double positive p16$^+$p21$^+$ cells and the percent SA-βGal$^+$ cells out of all live cells

were significantly higher in the lungs of *Mtb*-infected young *B6.Sst1S* and aged WT B6

mice than in young WT B6 (Fig. 3b, 3c). The elevation of senescence markers in young

*B6.Sst1S* and aged WT B6 mice were not seen in either uninfected lung or spleen cells

(Fig. 3b, 3c), likely because at 10 dpi there was little or no *Mtb* dissemination to spleen

(Supplementary Fig. 4b). Importantly, myeloid cells— predominantly alveolar (AM) and

interstitial macrophages (IM)- accounted for the majority of the enrichment in senescence

markers (double positive p16$^+$p21$^+$ cells and SA-βGal$^+$ cells) as seen in young *B6.Sst1S*

and aged WT B6 mice lung cells (Fig. 3d, 3e). No differences in senescence marker

abundance between the three types of mice were observed for B cells, and with T cells

we found a modest elevation of SA-βGal$^+$ senescence marker in aged WT B6 mice which

was not seen with the double positive p16$^+$p21$^+$ marker (Fig. 3d, 3e). Similar data were

observed at 20 dpi (Supplementary Fig. 5). Lastly, we evaluated lung levels of SASP

cytokines and found that five of the eight tested (IL-1β, IL-6, TNFα, IL-17a, and GM-CSF)

were significantly elevated in young *B6.Sst1S* mouse lungs compared to young WT B6,

although no differences were seen in aged WT B6 (Fig. 3f). Thus during *in vivo* mouse

infections, certain senescence markers— most notably macrophages doubly positive

p16$^+$p21$^+$ or for SA-βGal$^+$-are elevated in young *B6.Sst1S* and aged WT B6 mice

compared to young WT B6 mice, at a time point where no differences in CFU or pathology

are detected.

**'Omics studies identify the upregulation of pro-survival phosphatidylinositol 3-**

**kinase signaling (PI3K)- ERK1/2 cascade and anti-apoptotic pathways in *Mtb*-**

**infected BMDMs from *B6.Sst1S* mice.**

Senescent cells characteristically express cell cycle arrest and pro-inflammatory

transcriptional profiles. To compare the gene expression in stressed *B6.Sst1S* compared

to young WT B6 BMDMs, we performed bulk RNA- sequencing on BMDMs subjected to

10 ng/ml TNFα pretreatment, followed by *Mtb* infection in the continued presence of TNFα

(Fig. 4a). Despite increased *Mtb* proliferation in TNFα-treated *B6.Sst1S* BMDMs, cell

viability remained intact across all groups (Supplementary Fig. 6a, 6b). The dataset

revealed 1177 DEGs (*p. adj.*< 0.05, log$_2$fold change >1 or <-1):  647 upregulated and 530

downregulated genes in *B6.Sst1S* relative to similarly treated WT B6 BMDMs

(Supplementary File S3). Gene ontology (GO) analysis for biological processes identified

receptor tyrosine kinase (RTK) signaling pathway (GO:0007169; *p.adj.*= 0.004), positive

regulation of PI3K signaling (GO:0014068; *p.adj.*= 0.005), negative regulation of cell

proliferation (GO:0008285; *p.adj.*= 0.02), and cellular response to interferon-β

(GO:0035458; *p.adj.*= 0.01) to be significantly enriched in the *B6.Sst1S* BMDMs (Fig. 4b,

Supplementary File S3). As expected, the genes absent in *B6.Sst1S* cells (*Sp140, Sp110*)

were among the top downregulated genes in *B6.Sst1S* BMDMs (Fig. 4c). Interestingly,

we observed cyclin-dependent kinase inhibitor *Cdkn1c,* senoantigen[48] *Gpnmb* and

several genes included in SenMayo[45] gene set (*Arg2, Ptger2, Iqgap2, Selplg, Mmp10,*

*Mmp12, Gem, Ets2*), to be upregulated in *B6.Sst1S* BMDMs (Fig. 4c). In addition to the

pro-survival (PI3K-ERK1/2/ MAPK cascade) and SASP expression regulatory pathways

(RTKs) known to be up-regulated in senescent cells, the stressed *B6.Sst1S* BMDMs

exhibited upregulation of anti-apoptotic genes (*Bcl2a1b, Bcl2a1d, Itga6, Tnfaip8*)

suggesting that these senescent cells are resistant to apoptosis, and thereby may provide

a growth-permissive condition for *Mtb*.

**A senolytic drug cocktail (DQF) improves the survival of the host in *Mtb*-infected**

***B6.Sst1S* mice.**

A potential strategy to combat aging has been to employ senolytic drugs designed to kill

senescent cells and reverse their SASP-mediated local deleterious effects. Indeed,

several senolytic drugs have been shown to prolong lifespan in rodent models,[49] and

reduce age-associated disease morbidity in human clinical trials.[21,50–53] We selected a

cocktail of three most prevalent senolytic agents: dasatinib (D), quercetin (Q), and fisetin

(F).[54] Dasatinib is a tyrosine kinase inhibitor active against Src family and other tyrosine

kinases.[55] Inhibiting multiple kinases promotes apoptotic cell death in senescent cells by

disrupting pro-survival signaling networks such as the PI3K-ERK1/2- MAPK pathway.[49]

Quercetin and fisetin are flavonoid antioxidants found in common fruits and vegetables;

they inhibit both the anti-apoptotic BCL-2 pathway and the pro-survival PI3K-ERK1/2

pathway.[56–59] Since these pathways are upregulated in *Mtb*-infected *B6.Sst1S*

macrophages (Fig. 4b, 4c), we reasoned that targeting these pathways by the DQF

cocktail may alleviate *Mtb*-induced pathology in mice.

First, we confirmed that D, Q, and F do not exert direct anti-microbial activity against *Mtb*

at pharmacologically relevant concentrations (Supplementary Fig. 7a, 7b). We then

evaluated the host-directed effect of DQF alone or DQF in combination with partial

antimicrobial therapy comprised of ethambutol and rifampicin (DQF + Eth/Rif + Eth)

following an aerosol *Mtb* infection (Supplementary Fig. 8a). Three weeks post-infection,

when the lung CFU burden was 7.18 $\log_{10}$CFU units (Supplementary Fig. 8b), young

*B6.Sst1S* mice were randomly apportioned to four oral gavage treatment groups for a

period of 28 days: (i) vehicle, (ii) DQF (d21-49), (iii) Eth/Rif (d21-28) + Eth (d29-49) and

(iv) DQF (d21-49) +Eth/Rif (d21-28) + Eth (d29-49) (Supplementary Fig. 8a). Compared

with vehicle, mice receiving DQF alone showed a trend towards prolonged survival in a

subset of mice (*p*= 0.09) with some degree of weight stabilization (Fig. 5a, 5b). Mice

receiving DQF and partial antimicrobial therapy (DQF +Eth/Rif + Eth) showed a median

time to death (MTD) of 98 days and displayed significantly more weight gain in

comparison to those receiving partial antimicrobial therapy alone (Eth/Rif + Eth, MTD of

68 days, *p*= 0.05) (Fig. 5a, 5b). As has been noted by our group and others, [31,35,37,60]

*B6.Sst1S* mice display a bimodal mortality curve irrespective of therapeutic intervention

with these challenge doses within all groups, with 40-50% of all animals succumbing over

the first 35 days, and thus the impact of our drug regimens was only apparent in the

remaining 50-60% of surviving mice.

**The DQF senolytic drug cocktail reduces *Mtb* proliferation to a greater degree in**

**young *B6.Sst1S* and aged WT B6 mice than in young WT B6 mice.**

Next, we evaluated the anti-TB effect of the DQF cocktail with or without ethambutol

antimicrobial therapy in a lung CFU burden study of young *B6.Sst1S*, aged WT B6, and

young WT B6 mice following *Mtb* aerosol challenge. Because we had previously observed

poorer containment of *Mtb* proliferation in BMDMs from young *B6.Sst1S* and from aged

WT B6 mice compared to young WT B6 mice, we started drug treatment at 2 wpi for

young *B6.Sst1S* and aged WT B6 mice, and at 4 wpi in young WT B6 mice (Fig. 5c).

Compared to the vehicle-treated group, DQF therapy significantly reduced lung CFU

counts in *B6.Sst1S* mouse lungs by 1.3 ($p$= 0.03) and 2.2 ($p$= 0.05) $log_{10}$CFU units at 5

and 8 wpi, respectively, and in aged WT B6 mouse lungs by 0.5 ($p$= 0.11) $log_{10}$ units at 5

wpi (Fig. 5d). We did not find a detectable difference in lung CFU counts of young WT B6

mice at 8 wpi (Fig. 5d). The addition of DQF to ethambutol significantly lowered lung CFU

counts in all mouse strains compared to ethambutol alone, however, the magnitude of the

reduction was most profound in *B6.Sst1S* mice at 0.31 and 1 $log_{10}$CFU units at 5 and 8

wpi, respectively, and in aged WT B6 mice at 0.6 $log_{10}$CFU units at 5 wpi compared to

young WT B6 mice (Fig. 5d). Young WT B6 mice showed minimal changes with a

reduction of only 0.3 $log_{10}$CFU units at 8 wpi upon adding DQF to ethambutol (Fig. 5d).

With spleen CFU counts, the addition of DQF to ethambutol lowered bacterial burden in

*B6.Sst1S* mice by 1.0 ($p$= 0.16) and 0.6 ($p$=0.05) $log_{10}$CFU units at 5 and 8 wpi,

respectively, and in aged WT B6 mice by 1.1 ($p$= 0.06) $log_{10}$CFU units at 5 wpi compared

to ethambutol alone (Fig. 5e). There was no significant reduction observed between these

groups for young WT B6 mice (Fig. 5e).

**The DQF cocktail reduces lung pathology and senescence markers in *Mtb*-infected**

**lungs of young *B6.Sst1S* and aged WT B6 mice.**

To determine if the improved survival and *Mtb* containment observed in DQF-treated mice

correlated with the removal of senescent cells, we evaluated markers of lung damage

and senescence in these same mouse strains (Fig. 5c) treated with DQF in the presence

or absence of ethambutol. Considering overall lung immunopathology, as measured by

the area of consolidated lesions on H&E histology slides, all three mouse strains showed

greater reductions with DQF+ Eth as compared with Eth alone (*B6.Sst1S*: *p*= 0.02, aged

WT B6: *p*= 0.13, and young WT B6: *p*= 0.26), and the overall degree of lung consolidation

remaining (% lesion area) was lower in young *B6.Sst1S* (5 wpi- 8.4%, 8 wpi- 9.5%) and

aged WT B6 mice (5 wpi- 15.4%) receiving this maximal treatment (DQF+ Eth) than in

young WT B6 (8 wpi) mice (26.6%) (Fig. 6a, Supplementary Fig. 9a-c, 10a).

Immunohistochemistry for the DNA damage marker γH2A.X showed a significant

reduction in DQF+ Eth- treated *B6.Sst1S* and aged WT B6 mice with mice receiving both

drugs showed a near absence of immunoreactivity (Fig. 6b, Supplementary Fig. 10b).

More notably when we evaluated live lung cells by flow cytometry for dual staining with

$p21^+\beta Gal^+$ and for $p16^+\beta Gal^+$, there were significant reductions in both pairs of

senescence markers with adding DQF alone or with DQF+ Eth in young *B6.Sst1S* and

aged WT B6 mice (Fig. 6c, Supplementary Fig. 10c). The lung cells showing the highest

reductions in dual positivity for p21+βGal+ or for p16+βGal+ were from the myeloid

lineage (Fig. 6d, 6e). Interestingly, Eth monotherapy also decreased the senescent cell

population in *B6.Sst1S* mice (*p*< 0.0001) and aged WT B6 mice (*p*= 0.08-0.09), indicating

that inhibiting *Mtb* replication can also impede the generation of senescent cells (Fig. 6c).

In contrast, we saw no changes for dual positivity in young WT B6 mice with any of the

DQF and/or Eth regimens compared to vehicle (Fig. 6c).

In sum, we observed that the senolytic drug cocktail DQF prolonged survival and reduced

mouse lung CFU counts with a concomitant reduction in senescence markers and

pathology, with TB-susceptible young *B6.Sst1S* and aged WT B6 mice generally showed

the largest changes compared with young WT B6.

**Discussion**

This study provides proof of concept data that lung myeloid cells acquire a senescence

phenotype during TB and that these senescent cells are causal drivers of disease

progression. In this work, we used BMDMs from aged WT B6 mice as well as young mice

to establish that TNFα treatment in aged and *B6.Sst1S* BMDMs are not only associated

with poorer *Mtb* containment but also the induction of classic senescence markers

including SA-βGal and SASP gene expression, cell cycle arrest (reduced BrdU

incorporation, Lamin B1 and increased p16), and DNA damage (γH2A.X). Our previous

work with the *B6.Sst1S* macrophages revealed up-regulation of many SASP genes (*Il6,*

*Ccl2, Cxcl11, Tgfβ3, Mmp13, Tnfsf10*) (Supplementary File S1) and induction of the anti-

apoptotic ISR pathway after TNFα- stimulation at an early time point (18 hours).[35] Using

*Mtb*-infected aged WT B6 mice as a benchmark model of senescence during TB, we

sought to determine whether young *B6.Sst1S* mice resembled aged WT B6 or young WT

B6. Our studies revealed that both young *B6.Sst1S* mice and their BMDMs displayed the

same panel of senescence markers during stress or *Mtb* infection as observed in aged

WT B6 samples. The senescence phenotypes seen in the *B6.Sst1S* model resembled

those of aged WT B6 samples to a much greater degree than those of young WT B6.

These data suggest that the *B6.Sst1S* mouse may serve as an accelerated model of

myeloid cell senescence exhibiting canonical senescence markers γH2A.X, p16, p21, and

SA-βGal activity alongside SASP enrichment in TB.

Advanced age is a well-known TB risk factor,[7,62,63] early mouse studies by Orme and

Turner showed that *Mtb*-infected old mice have accelerated lethality post-*Mtb*

challenge.[14,61,62] The macrophages in aged mice were *Mtb*-infected in higher numbers

and expressed higher levels of the IFN-I responsive genes IRGM-1 and IRF-1.[63]

Interestingly, DiNardo *et al.* recently found that active TB leads to accelerated aging in

humans and in a guinea pig acute TB model using the DNA methylation clock.[64,65] Using

combined hypermethylation and gene expression clock algorithms, they found that active

TB adds 12.7 years (by the Horvath DNA hypermethylation clock) or 14.4 years (by the

Transcriptomic 'RNAAgeCalc' clock) of cellular aging compared to latent TB-infected

humans (LTBI).[64] Consistent with this, TB patients have been reported to show

significantly shorter telomere length and increased mitochondrial DNA copy number

(markers of senescence) relative to healthy controls in a diverse population.[66–68]

Additional studies have indeed demonstrated that *Mtb* infection induces DNA damage,

manipulates host cell cycle checkpoints, and promotes senescence-like features in

macrophages,[69–72] or T cells.[73]

The senescence literature has demonstrated that pathways required for homeostatic

development and wound healing in young age become dysregulated during aging.[27]

These include fibrosis with stem cell exhaustion in the setting of chronic mTOR activation,

persistent inflammation with excessive SASP cytokine release due to chronic NF-κB

activation, and DNA damage and mitochondrial dysfunction in the setting of declining

sirtuin activity.[74,75] A hallmark of senescent cells is entry into a non-dividing, long-lived

state with chronic expression of locally deleterious signals that promote age-associated

disease processes. In TB, the majority of infected humans adequately contain *Mtb*

infection within immune-balanced quiescent lung granulomas. Our study lends support to

the hypothesis that failed *Mtb* containment and progression to active TB may be promoted

by the presence of non-dividing senescent lung myeloid cells which may suppress the

necessary homeostatic immune containment required for TB control.

We found that senescence markers in aged WT B6 and young *B6.Sst1S* appear as

early as 10 days after *Mtb* infection, well before the appearance of granuloma formation

and lung immunopathology. This early chronology suggests that myeloid cell senescence

may drive the more severe TB pathology in aged WT B6 and young *B6.Sst1S* animals.

To establish causality, we used a cocktail of the senolytic drugs DQF- which eliminate

senescent cells by inhibiting the kinase (RTK/ PI3K/ ERK)-dependent survival pathways

and anti-apoptotic mechanisms that sustain their long-lived, non-dividing state. These

drugs have been shown to prolong lifespan in rodents and to prevent the progression of

age-associated diseases in humans.[21,49–51] We found that the DQF cocktail with or without

partial antimicrobial therapy was sufficient to prolong mouse survival and reduce *Mtb*

organ burden. Moreover, the impact of the senolytic DQF cocktail was more prominent in

mice that showed elevated senescence markers during *Mtb* infection, specifically young

*B6.Sst1S* and aged WT B6 mice. The fact that senolytic drugs improve TB disease

outcome parameters is a potent argument that senescent lung myeloid cells are causal

drivers of TB progression in these animal models.  As major risk factors for TB (age,

malnutrition, diabetes, pollution, and smoking) have been shown to induce senescence,

[62,76–81] our evidence that senescent cells drive TB progression may also offer a

mechanistic link for how these risk factors promote TB progression.

Considerable recent attention has been directed to the problem of post-TB lung disease.

Despite adequate and durable cures, many TB patients are left with serious pulmonary

disease, including bronchiectasis and either obstructive or restrictive lung disease.[82–85]

Senescent cells have been strongly associated with chronic tissue inflammation and

inappropriate fibrotic scarring.[86] Our studies with senolytic drugs indicated that lung

immunopathology was reduced to a greater degree by senolytic drugs in young *B6.Sst1S*

and aged WT B6 mice, which have high senescent cell burdens. Thus, in addition to

serving as promising adjunctive host-directed therapies for active TB, senolytic drugs may

also reduce post-TB lung disease.

In sum, by demonstrating the efficacy of Senolytics in reducing senescent cell burden

and improving TB outcomes in our preclinical model, we provide the rationale for testing

senescence-targeting strategies in human TB clinical trials.

**Material and Methods**

**Animals used in this study.**

Young C57BL/6J female mice (6 weeks old; Strain: 000664) and Aged C57BL/6J females (>68 weeks old) were bought from The Jackson Laboratory (Bar Harbor, ME, USA). The B6.C3H-sst1(B6J.C3-sst1C3HeB/FejKrmn) congenic mice were generated previously by transferring the *sst1S* allele from C3HeB/FeJ mouse strain to the B6 (C57BL/6J) genetic background using 12 backcrosses (*B6.Sst1S*)[29,35] and confirmed by Genotyping. The oligonucleotides used for the PCR are shown in Supplementary Table 1. This mice strain was bred and maintained at the Rodent Vivarium of Johns Hopkins University, School of Medicine and was used in this study. All experiments were conducted under protocols approved by the Johns Hopkins Animal Care and Use Committee (Protocol Number: MO22M466) and followed established ethical animal handling and care guidelines. Animals were housed in individually ventilated cages under controlled conditions, including a 12-hour light/dark cycle, ambient temperatures ranging from 20 to 24°C, and relative humidity between 45% and 65%. Each cage contained no more than five mice of the same strain/sex to ensure proper animal welfare. Mice were randomly assigned to experimental groups to minimize bias. Humane and painless euthanasia procedures were employed in compliance with AVMA (American Veterinary Medical Association) guidelines.

All experiments involving *Mycobacterium tuberculosis* H37Rv were conducted in Biosafety Level 3 (BSL3) and Animal Biosafety Level 3 (ABSL3) facilities at the Johns Hopkins University, School of Medicine, following protocols approved by the Institutional Biosafety Committee. Aerosol infections were performed using a Glas-Col aerosolization

system, allowing the mice to move freely in the chamber and minimizing stress or

discomfort. Anesthesia was not required for this procedure. Mice were briefly restrained

during oral gavage to ensure accurate delivery. Infected animals were monitored for signs

of illness, such as weight loss or lethargy, and those that became moribund before the

experiment's conclusion were promptly euthanized, except in survival studies.

**BMDMs isolation, culture, and treatment with murine-TNFα.**

BMDMs were isolated from mice at the indicated age, as described previously[87]. Briefly,

BMDMs were isolated from the tibias and femurs of mice and were differentiated into

macrophages in RPMI-Glutamax (Thermo Fisher Scientific; 61870127) containing 10%

heat-inactivated FBS (Millipore Sigma, Catalog no.- F4135), 1X antibiotic-antimycotic

Solution (Sigma Aldrich; A5955) and 10% (v/v) L929 fibroblast-conditioned medium (as a

source of M-CSF(macrophage colony-stimulating factor)) for 5 days. L929 fibroblast cell

lines were cultured in DMEM-Glutamax (Thermo Fisher Scientific; 10566016) containing

10% heat-inactivated FBS, 10 mM HEPES, and 1X antibiotic- antimycotic Solution. L929

cells were mycoplasma-negative (Venorgem Mycoplasma detection kit; Millipore Sigma-

MP0025). BMDMs were either left untreated or treated with 10 ng/ml of murine-TNFα

(Peprotech; 315-01A) in RPMI-Glutamax containing 10% heat-inactivated FBS and 10%

(v/v) L929 fibroblast-conditioned medium, for indicated time durations at 37°C in 5% $CO_2$.

BMDMs were washed, scraped out, counted, and replated overnight prior to the

experiments. All experiments were done with confluency of BMDMs 60-75%.

**Senescence associated β-Galactosidase assay (SA-βGal assay).**

SA-βGal assays at indicated time points were done with both paraformaldehyde- fixed

(Senescence Cells Histochemical Staining Kit; Sigma; CS0030) and live BMDMs

(Senescence β-Galactosidase Activity Assay Kit (Fluorescence, Flow Cytometry); Cell

Signaling Technology; 35302), using a modified manufacturer's protocol. At indicated time

points, BMDMs were washed twice with PBS and fixed with 1X fixation buffer for 10

minutes. BMDMs were washed thrice, the staining mixture was added to each well and

incubated for 48 h at 37°C without $CO_2$. We used 0.04X the X-gal Solution (10 μl in 1ml

of Staining mixture), to avoid saturation of the signal. The plate was sealed with Parafilm

to prevent evaporation of the staining mixture. After the incubation, cells were observed

under a bright field microscope and images were captured from the center of the well. For

experiments with live cells, adherent BMDMs were washed twice with warm PBS, and

incubated in Bafilomycin A1 (freshly prepared; 100 nM) containing complete RPMI

medium for 2 hours. Next, 0.05X of the recommended concentration of the SA-βGal

Substrate Solution (0.66 μM) was added to the cells and incubated for another 2- 2.5

hours at 37°C and 5% $CO_2$. The cells were washed thrice with PBS, scraped out in 1ml

RPMI (without phenol red) (Gibco; 11835), and LIVE/DEAD Fixable Near-IR Dead Cell

Stain (Thermo Fisher Scientific; L10119) was added for the detection of dead cells. FACS

was immediately done using CytoFLEX cytometer (Beckman Coulter), and 10000

events/sample were recorded. BMDMs from aged WT B6 mice (>72 weeks) were used

as a positive control. For etoposide (Millipore Sigma; E1383)- induced senescence

control, WT B6 BMDMs were treated with 10 μM etoposide for 2 days in complete RPMI

media, followed by washing to remove DNA-damaging molecules, and further incubation

for 2 days, in complete RPMI media only (no loss of viability under these conditions, as

detected by Alamar HS assay).

**Cell viability assay.**

The viability of cells was measured using an Alamar Blue HS Cell Viability Reagent

(Thermo Fisher Scientific; A50100), according to the manufacturer's protocol. Briefly, at

the indicated point, the (warmed) viability reagent was added to each well at 1/10[th] the

volume of cells, incubated for 6-8 hours and the fluorescence intensity was measured

using a FLUOstar Omega Multidetection Microplate Reader (BMG Labtech) in the bottom-

reading mode with excitation at 544 nm and emission at 590 nm. Untreated cells and cells

treated with 0.06% SDS, were used as positive and negative control, respectively.

Percent viability was calculated from the relative fluorescence units compared with

positive and negative control cells, to which the experimental values were normalized.

**Isolation of RNA and quantitative RT-PCR analysis to detect SASP expression.**

At the indicated time points, BMDMs were lysed in TRIzol (Thermo Fisher Scientific;

15596026), and RNA was extracted using Qiagen RNeasy Mini Kit (74104) and RNase-

free DNase Set (Qiagen; 79254), following the manufacturer's protocol. RNA purity and

concentration were quantified using Nanodrop One (Thermo Fisher Scientific). 1 µg of

RNA was used for cDNA synthesis using the iScript cDNA Synthesis Kit (BioRad;

1708891) and quantitative RT-PCR was done with QuantStudio 3 (Applied Biosystems;

Thermo Fisher Scientific) using gene-specific primers (Supplementary Table 1) and iTaq

Universal SYBR Green Supermix (BioRad; 1725121). Expression of genes was

normalized with Ct value for *ActB* (internal housekeeping control).

**BrdU Cell proliferation assay.**

The capacity of BMDMs to proliferate was measured by using the BrdU Cell Proliferation Assay Kit (Cell Signaling Technology; 6813), following the manufacturer's protocol. Briefly, 1-day post seeding of 50000 BMDMs in a 48-well plate, growth media (RPMI-Glutamax+ FBS) was removed, and RPMI-only was added to the cells to synchronize their cell cycle (upon serum starvation) and incubated for 18 hours at 37°C and 5% $CO_2$. After this synchronization-step, complete RPMI media was added to all wells and incubated for another 24 hours. The cells were either left untreated or treated with 10ng/ml TNFα, and BrdU incorporation into the cells was measured following the manufacturer's recommendations. Treatment of WT B6 BMDMs with 10 µM etoposide acted as a positive control. Absorbance at 450 nm was measured using the Bio-Rad iMark microplate reader.

**Protein isolation and Western Blotting.**

BMDMs were washed twice with ice-cold PBS, and the total protein was extracted by lysing cells in radioimmunoprecipitation (RIPA) Buffer (Millipore Sigma; R0278) containing Protease/Phosphatase Inhibitor Cocktail (1X; Cell Signaling Technology; 5872). After incubating on ice for 15 min with gentle vortexing, the lysates were centrifuged at 14000 RPM, 4°C for 15 min. The clarified supernatant was taken, and protein concentrations were quantified using the BCA Assay kit (Thermo Fisher Scientific; 23225). Approximately 50 µg of protein were mixed with Laemmli sample buffer (Bio-Rad; 1610747) and heated at 95 °C for 5 min before loading onto 4-15% MP TGX polyacrylamide Gel (Bio-Rad; 45610848) and transferring to the PVDF membranes (Bio-Rad; 1620177). The PVDF membranes were blocked using 5% Blotting-grade blocker (Bio-Rad; 1706404) in TBST buffer- Tris-buffered saline (Quality Biological; 351086101)+ 0.1% Tween 20 (Sigma;

P2287) for 1-2 hour and incubated overnight at 4ºC with the primary antibodies from

Mouse Reactive Senescence Marker Antibody Sampler Kit (Cell Signaling Technology-

78551)- Phospho-Histone H2A.X (Ser139) (20E3) Rabbit mAb (1: 1000 dilution), Lamin

B1 (E6M5T) Rabbit mAb (1: 1000 dilution), HMGB1 (D3E5) Rabbit mAb (1: 1000 dilution),

p16 INK4A (E5F3Y) Rabbit mAb (1: 1000 dilution) and β-Actin (D6A8) Rabbit mAb (Cell

Signaling Technology; 8457) (1:5000 dilution). Following three 15 min washing steps with

TBST, the membranes were incubated with secondary antibody- Anti-rabbit IgG, HRP-

linked Antibody (1:1000 dilution) for 2 hours at room temperature (RT). Following three

15 min washing steps, the membranes were developed with Pierce™ ECL Western

Blotting Substrate (Thermo Fisher Scientific; 32109) and visualized using a KwikQuant

Digital Western Blot Detection System (Kindle Bioscience; D1001). Images were

processed in Adobe Photoshop Software. Stripping Buffer (Restore™ Western Blot

Stripping Buffer; Thermo Fisher Scientific; 21059) was used to redevelop PVDF

membranes, following the manufacturer's protocol.

**Bulk RNA-Sequencing and analysis.**

BMDMs were lysed in TRIzol (Thermo Fisher Scientific; 15596026), and RNA was

extracted using Qiagen RNeasy Mini Kit (74104) and RNase-free DNase Set (Qiagen;

79254), following the manufacturer's protocol. RNA purity and concentration was

quantified using Nanodrop One (Thermo Fisher Scientific) and stored at -80 ºC. RNA

samples were shipped on dry ice to Genewiz, Azenta Life Sciences (South Plainfield, NJ,

USA) for standard bulk RNA- sequencing and downstream analysis. Sample QC, library

preparations, sequencing reactions, and initial bioinformatic analysis were conducted as

follows: Total RNA samples were quantified using Qubit 2.0 Fluorometer (Life

Technologies, Carlsbad, CA, USA) and RNA integrity was checked with 4200 TapeStation

(Agilent Technologies, Palo Alto, CA, USA). Samples were initially treated with TURBO

DNase (Thermo Fisher Scientific, Waltham, MA, USA) to remove DNA contaminants. The

next steps included performing rRNA depletion using QIAseq® FastSelect™−rRNA HMR

kit (Qiagen, Germantown, MD, USA), which was conducted following the manufacturer's

protocol. RNA sequencing libraries were generated with the NEBNext Ultra II RNA Library

Preparation Kit for Illumina by following the manufacturer's recommendations. Briefly,

enriched RNAs were fragmented for 15 minutes at 94°C. First strand and second strand

cDNA were subsequently synthesized. cDNA fragments were end-repaired and

adenylated at 3'ends, and universal adapters were ligated to cDNA fragments, followed

by index addition and library enrichment with limited cycle PCR. Sequencing libraries

were validated using the Agilent Tapestation 4200 (Agilent Technologies, Palo Alto, CA,

USA), and quantified using Qubit 2.0 Fluorometer (ThermoFisher Scientific, Waltham,

MA, USA) as well as by quantitative PCR (KAPA Biosystems, Wilmington, MA, USA). The

sequencing libraries were multiplexed and clustered on one lane of a flow cell. After

clustering, the flow cell was loaded on the Illumina HiSeq 4000 instrument according to

the manufacturer's instructions. The samples were sequenced using a 2x150 Pair-End

(PE) configuration. Image analysis and base calling were conducted by HiSeq Control

Software (HCS). Raw sequence data (.bcl files) generated from Illumina HiSeq was

converted into FASTQ files and de-multiplexed using Illumina's bcl2fastq 2.17 software.

One mismatch was allowed for index sequence identification.

Raw sequencing reads were trimmed to remove adapter sequences and low-quality

bases using Trimmomatic v0.36, and the filtered reads were aligned to the *Mus musculus*

GRCm38 reference genome (ENSEMBL) using the splice-aware STAR aligner v2.5.2b,

generating BAM files. Gene hit counts were quantified using featureCounts from the

Subread package v1.5.2, counting only uniquely mapped reads overlapping annotated

exon regions, with strand specificity applied where appropriate. Differential expression

analysis was performed using the DESeq2 package[88]. Genes with an adjusted *p*-value

<0.05 and a $log_2$ fold change > 1 or <-1 were deemed differentially expressed between

groups. Statistically significant differentially expressed genes (DEGs) were analyzed for

biological process enrichment through gene ontology analysis using the software

GeneSCF v1.1-p2 with the MGI GO database, generating clusters of functionally relevant

GO terms. Raw data were deposited in NCBI's Gene Expression Omnibus (GEO,

accession number-GSE292458). Reviewers can access this data using the following

token number- ihyhoqmkbhwxxmp.

**Minimum inhibitory concentration (MIC) and Minimum bactericidal concentration**

**(MBC) determination against *Mycobacterium tuberculosis* H37Rv.**

*Mycobacterium tuberculosis* strain H37Rv (*Mtb*) was obtained from the Johns Hopkins

University, Center for Tuberculosis Research (CTBR) and sequence-verified regularly.

*Mtb* was grown in 7H9 broth supplemented with 0.2% glycerol, 0.05% Tyloxapol (Millipore

Sigma; T0307), and OADC (BD; 212351) with shaking at 180 rpm in a rotary shaker

incubator or on 7H11 agar supplemented with OADC at 37°C. MIC was determined by a

resazurin microtiter assay (REMA) using 96-well flat-bottom plates[89]. *Mtb* H37Rv was

cultured in 7H9+OADC medium and grown to an exponential phase (optical density at

600 nm = 0.6-0.8). Approximately $1 \times 10^5$ bacteria per well were added in a total volume

of 200 µL of 7H9+OADC medium. Wells lacking *Mtb* served as controls. Additional

controls consisted of wells containing *Mtb* cells without drug treatment (growth control).

After 5 days of incubation at 37°C in the presence of drugs, 30 µL of 0.02% resazurin

(Sigma-Aldrich; R7017) was added, and plates were incubated for an additional 24-48

hours. The fluorescence intensity was measured using a FLUOstar Omega Multidetection

Microplate Reader (BMG Labtech) in the bottom-reading mode with excitation at 544 nm

and emission at 590 nm. Percent inhibition was calculated from the relative fluorescence

units compared with an untreated control culture; the MIC was taken as the lowest drug

concentration that resulted in at least 90% reduction in fluorescence compared to the

untreated growth control. To determine the MBC, 20 µl of cells from the respective MIC

dilutions (from the 96-well plate) were added onto the 7H11+OADC agar plate. Following

8-10 days of incubation at 37 °C, the colonies were checked for *Mtb* growth. Three wells

each from media containing no inoculum and, media containing *Mtb* but no drug, were

used as blank and growth controls, respectively. MBC was defined as the lowest drug

concentration at which no *Mtb* colonies appeared on the 7H11 plates.

**Spatial Transcriptomics of Lung from *Mtb* infected *B6.Sst1S* mice.**

To characterize gene expression in controlled vs uncontrolled lesions, we performed

spatial transcriptomics analysis using the Nanostring GeoMX Digital Spatial Profiler

(DSP) system (Nanostring, Seattle, WA).[90,91] The lung sections we processed for spatial

transcriptomics were obtained from *B6.Sst1S* mice. We used $10^6$ CFU of *Mtb* H37Rv for

infection, administered subcutaneously in the hock. The samples were collected at 14

weeks post-infection. Based on routine hematoxylin and eosin histopathology, we

selected lungs from 2 mice with paucibacillary controlled  lesions and 2 mice with

advanced uncontrolled multibacillary lesions with necrotic areas. Additional FFPE

sections were stained with fluorescent CD45-, pan-keratin-specific antibodies and DAPI.

Diseased regions of interest (ROI) were selected to focus on myeloid-rich areas avoiding

necrosis and tertiary lymphoid tissue. Eight ROI each of controlled and uncontrolled

lesions (respectively) were studied. The profiling used the Mouse Whole Transcriptome

Atlas (WTA) panel which targets ~21,000+ transcripts for mouse protein coding genes

plus ERCC ( External RNA Controls Consortium) negative controls to profile the whole

transcriptome, excluding uninformative high expressing targets such as ribosomal

subunits.

Samples from each ROI were packaged into library for sequencing (NextSeq550,

Illumina) following the procedure recommended by Nanostring. After sequencing, the

data analysis workflow began with QC evaluation of each ROI based on thresholds for

number of raw and aligned reads, sequencing saturation, negative control probe means,

and number of nuclei and surface area. Background correction is performed using

subtraction of the mean of negative probe counts. Q3 normalization (recommended by

Nanostring) results in scaling of all ROIs to the same value for their 3rd quartile value.

Statistical analysis was then performed to identify differentially expressed genes, using

an unpaired t-test with Benjamini-Hochberg adjustment for *p* values. Raw data were

deposited in NCBI's Gene Expression Omnibus (GEO, accession number- GSE292392).

Reviewers can access this data using the following token number-  clozaycmjjivduj.

**Animal infection studies.**

Mice of the specified age and sex were infected via aerosol exposure using the Glas-Col

Inhalation Exposure System (Terre Haute, IN, USA). Fresh cryo-aliquots of *Mtb* H37Rv

were used for each infection, diluted in sterile 7H9 broth to empirically determined

concentrations to achieve the desired bacterial load on day 1 post-infection. To minimize

variability between groups, mice from all experimental groups were infected concurrently

and randomly allocated to experimental arms. The bacterial implantation in the lungs was

assessed one day after infection by sacrificing 3–5 mice per group and determining

colony-forming units (CFUs). Mice were euthanized at specified time points (as outlined

in the Results section), and the lungs and spleen were aseptically harvested, weighed

and either homogenized by bead-beating for bacterial quantification (Right lung) or stored

in 10% neutral-buffered formalin for histopathological analysis (Left lung). CFU

enumeration was performed by plating serial dilutions on 7H11 agar supplemented with

OADC, followed by incubation at 37°C for 3 to 4 weeks. Bacterial counts were normalized

to the weight of the total organ, as previously described[92].

**Histopathology to estimate lung inflammation.**

For histology, left lungs were fixed by immersion in 10% neutral-buffered formalin (Thermo

Fisher Scientific; 5705) for at least 2 days, before it was submitted to the SKCCC

Oncology Tissue and Imaging Service Core Laboratory (Johns Hopkins University, School

of Medicine) for downstream processing. The lungs were paraffin-embedded, 4 µm-

sectioned, and stained with hematoxylin and eosin (H&E). Slides were digitally scanned

at 40× using a Hamamatsu Nanozoomer S210 digital slide scanner (Hamamatsu

Photonics, Shizuoka, Japan) at 40X magnification (0.23 microns/pixel). Image files were

submitted in https://digital.pathology.johnshopkins.edu/ using Concentriq LS platform

(Proscia, PA, USA). Lung inflammation was quantified as a percentage of total lung area

using an in-house protocol developed with ImageJ software version 1.54m (NIH, USA) to

minimize human bias, through standardized software-based settings. Images were first

opened in ImageJ, and the scale was set by drawing a line over the scale bar using the

line tool and defining the known distance in millimeters under Analyze> Set Scale. Images

were then cropped to exclude the ruler (Image > Crop) and saved. For area quantification,

the Image> Adjust> Color Threshold tool was used, with consistent settings for hue (0–

255), saturation (0–255), and brightness (set based on analysis type)- full lung area=(0-

250) and inflamed area= (0-160). After selecting the ROI, the threshold window was

closed, and images were converted to binary format using Process> Binary> Make

Binary. The Signal to Noise ratio was improved by removing the Outliers- using Process>

Noise> Remove Outliers, adjusting the radius (=4) and the Threshold (=50).

Measurements were extracted (Analyze> Measure), with the initial setup requiring the

selection of "area" and "limit to threshold" under Set Measurements. Processed images

were saved, and the steps were repeated for all images with identical parameters to

ensure consistency. ROIs were then divided by the total lung area per slide to calculate

the ratio of inflamed to lung tissue- area.

**Multicolor Flow cytometry to quantify senescent cell population in *Mtb*-infected**

**mice lungs and spleen.**

To assess markers of senescence (p16+, p21+, and increased SA-βGalactosidase

activity) in lung and spleen tissues from *Mtb*-infected mice, organs were harvested at

indicated time points and dissociated using the Lung Dissociation Kit (Miltenyi Biotec;130–

095-927), and the gentleMACS Octo Dissociator (program m_lung_02), according to the

manufacturer's protocol. Single-cell suspensions were obtained by filtering dissociated

tissue through 70 μm mesh strainers (Corning; 431751), washing with 5 mL complete

RPMI media, and centrifuging at 300g for 5 min at RT. Red blood cells were lysed using

ACK lysis buffer (Quality Biological; 118156101), neutralized with complete RPMI, and

centrifuged. Cells were stained for viability using a live-dead near-IR dye (1:1000 dilution

in PBS; Thermo Fisher Scientific; L10119) and blocked with Rat Anti-Mouse CD16/CD32

Fc Block (Clone 2.4G2, BD Pharmingen; 553142). The panel of antibodies to detect the

senescent cell population (SenPanel) was designed using FluoroFinder

(https://fluorofinder.com/). For extracellular staining, cells were incubated with 50 µL of an

antibody cocktail in Brilliant Stain Buffer (BD Biosciences; 563794) containing anti- mouse

CD45 (Brilliant Violet 480; BD Biosciences;566095), CD11b (Brilliant Violet 650;

BioLegend; 101239), CD45R/B220 (Brilliant Violet 421; BioLegend; 103239), CD3

(StarBright Violet 610; BioRad-MCA500SBV610), EpCAM (PerCP-Cy5.5; BioLegend;

118220), F4/80 (PE-Dazzle 594; BioLegend; 123145), CD11c (PE-Fire 810; BioLegend;

161105), and Siglec-F (Alexa Fluor 700; BioLegend; 155533) for 30 minutes at 4°C in the

dark. After extracellular staining, cells were fixed with 4% formaldehyde, washed with

PBS, and incubated with the CellEvent™ Senescence Green Flow Cytometry Assay Kit

(Thermo Fisher Scientific; C10841; 1:250 dilution) in the senescence reaction buffer for 4

hours at 37°C, protected from light and without $CO_2$. Following incubation, cells were

washed and permeabilized using Perm/Wash buffer (BD Biosciences; 554723).

Intracellular staining was performed by incubating cells with a cocktail containing

p14ARF/CDKN2A Antibody (PE; Novus Biologicals; NB200-111PE), p21/CIP1/CDKN1A

Antibody (HJ21) (APC; Novus Biologicals; NBP2-47896APC), and CD68 (Brilliant Violet

785; BioLegend; 137035) for 60 minutes at 4°C in the dark. After staining, cells were

washed, filtered through strainer-capped BD FACS tubes (BD; 352235), and analyzed via

flow cytometry. $1×10^6$ events per sample were acquired using a BD LSR Fortessa™ Cell

Analyzer (BD Biosciences), and the data were analyzed with FlowJo software version

10.8.1 (BD Biosciences). Single stain controls and compensation matrices were prepared

using UltraComp eBeads Plus Compensation Beads (Thermo Fisher Scientific; 01-3333-

41) and cells, as required. All flow antibodies were titrated to determine optimal staining

concentrations, and only manufacturer-validated antibodies with demonstrated specificity

were used. For the analysis, cells were first gated on singlets (FSC-H vs. FSC-A) and live

cells (live-dead near-IR negative) before further selection of cell types. T cells are

CD45+CD3+ and B cells are CD45+B220+. Myeloid cells: Tissue-resident Macrophages

are CD45+CD68+F4/80+CD11B-CD11C+, Interstitial macrophages are CD45+CD68+

F4/80+CD11B+CD11C+SiglecF-, Monocytes are CD45+CD68+F4/80+CD11B+CD11C-,

and Dendritic cells are CD45+CD68+F4/80-CD11B+CD11C+. Gating for $p16^+$, $p21^+$, and

SA-βGal$^+$ events for each cell type were established using reference cells from *Mtb*-

infected WT B6 mice (set at 1 %).

**Immunohistochemistry against p21 and γH2A.X (phosphor S139) in mice lungs.**

Immunostaining was conducted at the SKCCC Oncology Tissue and Imaging Service

Core Laboratory, Johns Hopkins University, School of Medicine. Immunolabeling for

γH2A.X (phospho S139) and p21 was performed on 4 µm-thick formalin-fixed, paraffin-

embedded (FFPE) tissue sections using the Ventana Discovery Ultra autostainer (Roche

Diagnostics). Sections underwent onboard dewaxing and rehydration, followed by

epitope retrieval in Ventana Ultra CC1 buffer (Roche Diagnostics; 6414575001) at 96°C

for 64 minutes. Primary antibodies- anti-gamma H2A.X (phospho S139) [EP854(2)Y]

(Abcam; ab81299; 1:1000 dilution) and Recombinant Anti-p21 antibody [EPR18021]

(Abcam; ab188224; 1:1000 dilution) were applied as indicated, at 36°C for 60 minutes.

Detection of the primary antibodies was achieved using the anti-rabbit HQ detection

system (Roche Diagnostics; Catalog 7017936001 and 7017812001), followed by

visualization with the Chromomap DAB IHC detection kit (Roche Diagnostics;

5266645001). Sections were counterstained with Mayer's hematoxylin, dehydrated, and

mounted. Whole-slide imaging was performed at the Oncology Tissue Services Core of

Johns Hopkins University using a Hamamatsu Nanozoomer S210 digital slide scanner

(Hamamatsu Photonics, Shizuoka, Japan) at 40X magnification (0.23 microns/pixel).

Images were visualized using the Concentriq digital pathology platform (Proscia,

Philadelphia, PA) and analyzed in ImageJ following a previously established in- house

protocol. Area quantification was performed using the Image> Adjust> Color Threshold

tool in ImageJ, applying consistent hue (0–255), saturation (0–255), and brightness

thresholds based on analysis type: full lung area (0–250) and DAB+ area (0–120).

Identical thresholds and detection settings were applied for all slides to ensure

consistency across analyses.

**Senescence-associated secretory phenotype (SASP)- cytokine quantification in**

*Mtb*- **infected mice lungs.**

To estimate SASP- cytokine levels, lung homogenates were centrifuged at 14,000 RPM

for 10 minutes at 4°C, and the resulting supernatants were filtered by a 0.2 µm filter, prior

to removal from the BSL3 facility. Supernatants were flash-frozen in liquid nitrogen and

stored at –80°C until cytokine quantification analysis. Cytokine levels in the lung

homogenate were measured using the LEGENDplex Mouse Inflammation Panel 13-plex

kit (BioLegend; 740446) following the manufacturer's protocol. Data acquisition was

performed on the CytoFLEX Flow Cytometer (Beckman Coulter), and cytokine

concentrations were analyzed using the LEGENDplex data analysis software Suite

(BioLegend). The amount of SASP cytokines were normalized to the protein

concentration of the homogenate (as measured by the BCA Assay kit).

**Drug preparation and study arms to test Senolytics as host- directed therapy *in***

***vivo*.**

Dasatinib (Cayman Chemical; 11498), Fisetin (Cayman Chemical;15246), and Quercetin

(Millipore Sigma; Q4951) at required concentrations were first dissolved in 10ml  DMSO

(Millipore Sigma; D2650) to make a stock solution. This mixture was then added to 190

ml of the Vehicle (60% Phosal 50 PG (Lipoid LLC; 368315), 30% PEG 400 (Sigma Aldrich;

202398), and 10% Ethanol) to prepare the final Senolytics reagent. Ethambutol (Millipore

Sigma; E4630) and Rifampicin (Millipore Sigma; R3501) were used as anti-*Mtb* drug

control. Solutions of Ethambutol and Rifampicin in distilled water were prepared and

stored at 4 °C.

After the day of aerosol infection, female mice were randomly divided into 4 groups (n = 4-

8 per group, as indicated in the Results). Feed and water were given *ad libitum*. Treatment

with drugs started at 2 weeks (*B6.Sst1S* mice and WT B6 Old), or 4 weeks (for WT B6

mice) post-infection. The drug treatment continued for another 6 weeks (for *B6.Sst1S*

mice), 3 weeks (for WT B6 Old) or 4 weeks (for WT B6). Treated mice were administered

Ethambutol (100 mg/kg of body weight) by oral gavage, once daily, 5 times a week.

Vehicle or Senolytics (Dasatinib: 5 mg/kg, Fisetin: 100 mg/kg, and Quercetin: 50 mg/kg

of body weight of mice) were administered by oral gavage, once daily, 3 times a week.

The fourth group received both the Ethambutol and Senolytics combination. All drugs

were administered in a total volume of 0.2 ml per treatment. Three infected mice were

sacrificed at the start of treatment as pretreatment controls. At indicated time points, mice

were sacrificed, and the lungs and spleen were harvested for the measurement of

bacterial burden, IHC, FACS and histopathology.

**Time to death assay.**

To examine if the Senolytics administration can improve the survival and body weight, the

survival of both male and female *B6.Sst1S* mice (8-10 weeks old) following *Mtb* infection

were assessed. Mice were infected via the aerosol route using a Glas-Col aerosolization

instrument to deliver a high dose of *Mtb* to the *B6.Sst1S* lungs (approximately 100 CFU).

After 3 weeks post-infection, infected mice were randomly divided into 4 groups and the

drug was administered as mentioned above, with slight modifications. The drug treatment

continued for another 4 weeks, where mice were administered Ethambutol (50 mg/kg of

body weight) by oral gavage, once daily, 5 times a week. In the second week, Ethambutol

(50 mg/kg of body weight) was combined with Rifampicin (10 mg/kg of body weight) for 1

week. Vehicle or Senolytics (Dasatinib: 5 mg/kg, Fisetin: 100 mg/kg, and Quercetin: 50

mg/kg of body weight of mice) were administered by oral gavage, once daily, 3 times a

week. The fourth group received both the Ethambutol and Senolytics combination. All

drugs were administered in a total volume of 0.2 ml per treatment. All mice were monitored

daily for signs of disease progression, including weight loss, lethargy, hunched posture,

and reduced activity. Survival data were analyzed using Kaplan-Meier survival curves to

determine the median time to death and assess differences between treatment groups.

**Statistical analysis.**

Statistical analyses were performed using GraphPad Prism version 10.4.0 software.

Normality distribution was first assessed using the Shapiro-Wilk test. For comparisons of

lung CFUs, data was $log_{10}$-transformed and checked for normality before the analysis.

Statistical significance was determined using unpaired two-tailed Student's t-tests or

Mann Whitney test for two-group comparisons, and one-way or two- way ANOVA with

post hoc tests for multiple- group comparisons. Kaplan-Meier survival analysis was

conducted for time-to-death (TTD) studies, with group differences assessed by the Log-

rank (Mantel-Cox) test. In animal experiments, each data point represented a single

mouse, and experiments were randomized, though blinding was not implemented.

Results are presented as mean ± SEM, with error bars indicating SEM. *p*-values less than

0.05 were considered statistically significant.

**Contact for reagent and resource sharing.**

All data associated with this study are present in the paper- either in the Results section

or in the Supplementary files. Requests for further details on resources, protocol, and

reagents can be directed to the corresponding author- Prof. William R Bishai. Raw data

generated during this study is provided as a separate Excel file labeled "Source Data file".

The RNA sequencing and Spatial transcriptomics datasets are available on NCBI's Gene

Expression Omnibus (GEO). The *B6.Sst1S* mice used in this study are available for

academic and non-commercial research upon request to Prof. Igor Kramnik, under a

Material Transfer Agreement.

**Acknowledgements**

We gratefully acknowledge the support of the NIH grants AI152688, AI155602, CA006973, and R01HL126066 (to IK). We extend our gratitude to the SKCCC Core centers for their assistance with Histology and Flow cytometry. The funders have no role in study-design, data collection and interpretation, and the decision to submit the work for publication or preparation of the manuscript. We sincerely thank all the members of Prof. Bishai lab and Prof. Kramnik lab, for their valuable suggestions and discussions throughout this project. The schematic figures in this study were created using BioRender (with Academic Lab License).

**Author contributions.**

Funding Acquisition was contributed by- WB, IK. Conceptualization- WB, IK, SS. Methodology- WB, IK, SS, YMM, BK, SY, LK. Investigation and Formal Analysis- WB, IK, SS. Data curation and Validation- SS, SY, LK. Writing original draft- WB, IK, SS, SY. Review and editing manuscript- WB, IK, SS, YMM, BK, SY, LK.

**Conflict of interest statement.**

The authors declare that they have no competing or conflicting interests.

## References

1.  *2024 Global Tuberculosis Report*. https://www.who.int/teams/global-tuberculosis-programme/tb-reports/global-tuberculosis-report-2024.

2.  Horton, K. C., Richards, A. S., Emery, J. C., Esmail, H. & Houben, R. M. G. J. Reevaluating progression and pathways following Mycobacterium tuberculosis infection within the spectrum of tuberculosis. *Proc Natl Acad Sci U S A* **120**, e2221186120 (2023).

3.  Sossen, B. *et al.* The natural history of untreated pulmonary tuberculosis in adults: a systematic review and meta-analysis. *Lancet Respir Med* **11**, 367–379 (2023).

4.  Richards, A. S. *et al.* Quantifying progression and regression across the spectrum of pulmonary tuberculosis: a data synthesis study. *Lancet Glob Health* **11**, e684–e692 (2023).

5.  Nguyen, H. V., Tiemersma, E., Nguyen, N. V., Nguyen, H. B. & Cobelens, F. Disease Transmission by Patients With Subclinical Tuberculosis. *Clin Infect Dis* **76**, 2000–2006 (2023).

6.  Chen, R. Y. *et al.* Radiological and functional evidence of the bronchial spread of tuberculosis: an observational analysis. *Lancet Microbe* **2**, e518–e526 (2021).

7.  GBD Tuberculosis Collaborators. The global burden of tuberculosis: results from the Global Burden of Disease Study 2015. *Lancet Infect Dis* **18**, 261–284 (2018).

8.  Olmo-Fontánez, A. M. & Turner, J. Tuberculosis in an Aging World. *Pathogens* **11**, (2022).

9.  Negin, J., Abimbola, S. & Marais, B. J. Tuberculosis among older adults--time to take notice. *Int J Infect Dis* **32**, 135–7 (2015).

10. Caraux-Paz, P., Diamantis, S., de Wazières, B. & Gallien, S. Tuberculosis in the Elderly. *J Clin Med* **10**, 5888 (2021).

11. Teo, A. K. J. *et al.* Tuberculosis in older adults: challenges and best practices in the Western Pacific Region. *Lancet Reg Health West Pac* **36**, 100770 (2023).

12. Orme, I. M. Aging and immunity to tuberculosis: prolonged survival of old mice infected with Mycobacterium tuberculosis by adoptive immunization with memory-immune T lymphocytes. *Cell Immunol* **118**, 229–33 (1989).

13. Orme, I. M. A mouse model of the recrudescence of latent tuberculosis in the elderly. *Am Rev Respir Dis* **137**, 716–8 (1988).

14. Turner, J., Frank, A. A. & Orme, I. M. Old mice express a transient early resistance to pulmonary tuberculosis that is mediated by CD8 T cells. *Infect Immun* **70**, 4628–37 (2002).

15. Torrelles, J. B. *et al.* The Impact of Aging on the Lung Alveolar Environment,
Predetermining Susceptibility to Respiratory Infections. *Frontiers in Aging* **3**, 1–9
(2022).

16. Canan, C. H. *et al.* Characterization of lung inflammation and its impact on
macrophage function in aging. *J Leukoc Biol* **96**, 473–480 (2014).

17. Lafuse, W. P. *et al.* Identification of an Increased Alveolar Macrophage
Subpopulation in Old Mice That Displays Unique Inflammatory Characteristics and
Is Permissive to Mycobacterium tuberculosis Infection. *J Immunol* **203**, 2252–
2264 (2019).

18. Olmo-Fontánez, A. M. *et al.* Impact of the elderly lung mucosa on Mycobacterium
tuberculosis transcriptional adaptation during infection of alveolar epithelial cells.
*Microbiol Spectr* **12**, e0179024 (2024).

19. Olmo-Fontánez, A. M. *et al.* Human alveolar lining fluid from the elderly promotes
Mycobacterium tuberculosis intracellular growth and translocation into the cytosol
of alveolar epithelial cells. *Mucosal Immunol* **17**, 155–168 (2024).

20. Moliva, J. I. *et al.* The Lung Mucosa Environment in the Elderly Increases Host
Susceptibility to Mycobacterium tuberculosis Infection. *Journal of Infectious
Diseases* **220**, 514–523 (2019).

21. Di Micco, R., Krizhanovsky, V., Baker, D. & d'Adda di Fagagna, F. Cellular
senescence in ageing: from mechanisms to therapeutic opportunities. *Nat Rev
Mol Cell Biol* **22**, 75–95 (2021).

22. Wissler Gerdes, E. O., Zhu, Y., Tchkonia, T. & Kirkland, J. L. Discovery,
development, and future application of senolytics: theories and predictions. *FEBS
J* **287**, 2418–2427 (2020).

23. Kennedy, B. K. *et al.* Geroscience: linking aging to chronic disease. *Cell* **159**,
709–13 (2014).

24. Wang, B., Han, J., Elisseeff, J. H. & Demaria, M. The senescence-associated
secretory phenotype and its physiological and pathological implications. *Nat Rev
Mol Cell Biol* **25**, 958–978 (2024).

25. Xu, M. *et al.* Transplanted Senescent Cells Induce an Osteoarthritis-Like
Condition in Mice. *J Gerontol A Biol Sci Med Sci* **72**, 780–785 (2017).

26. Franco, A. C. *et al.* Senescent cell transplantation into the skin induces age-
related peripheral dysfunction and cognitive decline. *Aging Cell* **24**, (2025).

27. Childs, B. G., Durik, M., Baker, D. J. & van Deursen, J. M. Cellular senescence in
aging and age-related disease: from mechanisms to therapy. *Nat Med* **21**, 1424–
35 (2015).

28.   Justice, J. N. *et al.* Senolytics in idiopathic pulmonary fibrosis: Results from a first-
in-human, open-label, pilot study. *EBioMedicine* **40**, 554–563 (2019).

29.   Pichugin, A. V., Yan, B. S., Sloutsky, A., Kobzik, L. & Kramnik, I. Dominant role of
the sst1 locus in pathogenesis of necrotizing lung granulomas during chronic
tuberculosis infection and reactivation in genetically resistant hosts. *American*
*Journal of Pathology* **174**, 2190–2201 (2009).

30.   Yan, B. S., Kirby, A., Shebzukhov, Y. V., Daly, M. J. & Kramnik, I. Genetic
architecture of tuberculosis resistance in a mouse model of infection. *Genes*
*Immun* **7**, 201–210 (2006).

31.   Ji, D. X. *et al.* Type I interferon-driven susceptibility to Mycobacterium tuberculosis
is mediated by IL-1Ra. *Nat Microbiol* **4**, 2128–2135 (2019).

32.   Yabaji, S. M. *et al.* Cell state transition analysis identifies interventions that
improve control of Mycobacterium tuberculosis infection by susceptible
macrophages. *Sci Adv* **9**, eadh4119 (2023).

33.   Pan, H. *et al.* Ipr1 gene mediates innate immunity to tuberculosis. *Nature* **434**,
767–772 (2005).

34.   Krug, S. *et al.* Adjunctive Integrated Stress Response Inhibition Accelerates
Tuberculosis Clearance in Mice. *mBio* **14**, e0349622 (2023).

35.   Bhattacharya, B. *et al.* The integrated stress response mediates necrosis in
murine Mycobacterium tuberculosis granulomas. *Journal of Clinical Investigation*
**131**, (2021).

36.   Fraschilla, I., Amatullah, H., Rahman, R. U. & Jeffrey, K. L. Immune chromatin
reader SP140 regulates microbiota and risk for inflammatory bowel disease. *Cell*
*Host Microbe* **30**, 1370-1381.e5 (2022).

37.   Ji, D. X. *et al.* Role of the transcriptional regulator SP140 in resistance to bacterial
infections via repression of type I interferons. *Elife* **10**, (2021).

38.   Yabaji, S. M. *et al.* Lipid Peroxidation and Type I Interferon Coupling Fuels
Pathogenic Macrophage Activation Causing Tuberculosis Susceptibility. *bioRxiv*
(2024) doi:10.1101/2024.03.05.583602.

39.   Moreira-Teixeira, L. *et al.* Type I IFN exacerbates disease in tuberculosis-
susceptible mice by inducing neutrophil-mediated lung inflammation and NETosis.
*Nat Commun* **11**, 5566 (2020).

40.   Berry, M. P. R. *et al.* An interferon-inducible neutrophil-driven blood transcriptional
signature in human tuberculosis. *Nature* **466**, 973–7 (2010).

41.   Kotov, D. I. *et al.* Early cellular mechanisms of type I interferon-driven
susceptibility to tuberculosis. *Cell* **186**, 5536-5553.e22 (2023).

42.  Kalinin, A., Zubkova, E. & Menshikov, M. Integrated Stress Response (ISR)
Pathway: Unraveling Its Role in Cellular Senescence. *Int J Mol Sci* **24**, (2023).

43.  Payea, M. J. *et al.* Senescence suppresses the integrated stress response and
activates a stress-remodeled secretory phenotype. *Mol Cell* **84**, 4454-4469.e7
(2024).

44.  Zhou, Y. *et al.* Metascape provides a biologist-oriented resource for the analysis of
systems-level datasets. *Nat Commun* **10**, 1523 (2019).

45.  Saul, D. *et al.* A new gene set identifies senescent cells and predicts senescence-
associated pathways across tissues. *Nat Commun* **13**, (2022).

46.  Sharpless, N. E. & Sherr, C. J. Forging a signature of in vivo senescence. *Nat
Rev Cancer* **15**, 397–408 (2015).

47.  González-Gualda, E., Baker, A. G., Fruk, L. & Muñoz-Espín, D. A guide to
assessing cellular senescence in vitro and in vivo. *FEBS Journal* **288**, 56–80
(2021).

48.  Suda, M. *et al.* Senolytic vaccination improves normal and pathological age-
related phenotypes and increases lifespan in progeroid mice. *Nat Aging* **1**, 1117–
1126 (2021).

49.  Zhu, Y. *et al.* The Achilles' heel of senescent cells: from transcriptome to senolytic
drugs. *Aging Cell* **14**, 644–58 (2015).

50.  Zhang, L. *et al.* Cellular senescence: a key therapeutic target in aging and
diseases. *J Clin Invest* **132**, (2022).

51.  Raffaele, M. & Vinciguerra, M. The costs and benefits of senotherapeutics for
human health. *Lancet Healthy Longev* **3**, e67–e77 (2022).

52.  Robbins, P. D. *et al.* Senolytic Drugs: Reducing Senescent Cell Viability to Extend
Health Span. *Annu Rev Pharmacol Toxicol* **61**, 779–803 (2021).

53.  Chaib, S., Tchkonia, T. & Kirkland, J. L. Cellular senescence and senolytics: the
path to the clinic. *Nat Med* **28**, 1556–1568 (2022).

54.  Lee, E. *et al.* Exploring the effects of Dasatinib, Quercetin, and Fisetin on DNA
methylation clocks: a longitudinal study on senolytic interventions. *Aging* **16**,
3088–3106 (2024).

55.  Shah, N. P. *et al.* Overriding imatinib resistance with a novel ABL kinase inhibitor.
*Science* **305**, 399–401 (2004).

56.  Yousefzadeh, M. J. *et al.* Fisetin is a senotherapeutic that extends health and
lifespan. *EBioMedicine* **36**, 18–28 (2018).

57. Zhu, Y. *et al.* New agents that target senescent cells: the flavone, fisetin, and the BCL-XL inhibitors, A1331852 and A1155463. *Aging* **9**, 955–963 (2017).

58. Lim, J. Y., Lee, J. Y., Byun, B. J. & Kim, S. H. Fisetin targets phosphatidylinositol-3-kinase and induces apoptosis of human B lymphoma Raji cells. *Toxicol Rep* **2**, 984–989 (2015).

59. Kim, J. A. *et al.* Fisetin, a dietary flavonoid, induces apoptosis of cancer cells by inhibiting HSF1 activity through blocking its binding to the hsp70 promoter. *Carcinogenesis* **36**, 696–706 (2015).

60. Amaral, E. P. *et al.* BACH1 promotes tissue necrosis and Mycobacterium tuberculosis susceptibility. *Nat Microbiol* **9**, 120–135 (2024).

61. Orme, I. Mechanisms underlying the increased susceptibility of aged mice to tuberculosis. *Nutr Rev* **53**, S35-8; discussion S38-40 (1995).

62. Vesosky, B. & Turner, J. The influence of age on immunity to infection with Mycobacterium tuberculosis. *Immunol Rev* **205**, 229–43 (2005).

63. Rottinghaus, E. K., Vesosky, B. & Turner, J. TLR-2 independent recognition of Mycobacterium tuberculosis by CD11c+ pulmonary cells from old mice. *Mech Ageing Dev* **131**, 405–14 (2010).

64. Bobak, C. A. *et al.* Increased DNA methylation, cellular senescence and premature epigenetic aging in guinea pigs and humans with tuberculosis. *Aging* **14**, 2174–2193 (2022).

65. DiNardo, A. R. *et al.* DNA hypermethylation during tuberculosis dampens host immune responsiveness. *J Clin Invest* **130**, 3113–3123 (2020).

66. Udomsinprasert, W. *et al.* Leukocyte telomere length as a diagnostic biomarker for anti-tuberculosis drug-induced liver injury. *Sci Rep* **10**, (2020).

67. Freimane, L. *et al.* Telomere length and mitochondrial DNA copy number in multidrug-resistant tuberculosis. *Tuberculosis* **131**, (2021).

68. Katoto, P. D. M. C. *et al.* Telomere length and outcome of treatment for pulmonary tuberculosis in a gold mining community. *Sci Rep* **11**, (2021).

69. Lochab, S., Singh, Y., Sengupta, S. & Nandicoori, V. K. Mycobacterium tuberculosis exploits host ATM kinase for survival advantage through secA2 secretome. *Elife* **9**, (2020).

70. Cumming, B. M., Addicott, K. W., Adamson, J. H. & Steyn, A. J. C. Mycobacterium tuberculosis induces decelerated bioenergetic metabolism in human macrophages. *Elife* **7**, 1–28 (2018).

71. Cumming, B. M. *et al.* Mycobacterium tuberculosis arrests host cycle at the G1/S transition to establish long term infection. *PLoS Pathog* **13**, (2017).

72.   Cyktor, J. C. *et al.* Killer cell lectin-like receptor G1 deficiency significantly
enhances survival after Mycobacterium tuberculosis infection. *Infect Immun* **81**,
1090–9 (2013).

73.   Chang, E., Cavallo, K. & Behar, S. M. CD4 T cell dysfunction is associated with
bacterial recrudescence during chronic tuberculosis. *Nat Commun* **16**, 2636
(2025).

74.   Chien, Y. *et al.* Control of the senescence-associated secretory phenotype by NF-
κB promotes senescence and enhances chemosensitivity. *Genes Dev* **25**, 2125–
36 (2011).

75.   Miwa, S., Kashyap, S., Chini, E. & von Zglinicki, T. Mitochondrial dysfunction in
cell senescence and aging. *J Clin Invest* **132**, (2022).

76.   VanValkenburg, A. *et al.* Malnutrition leads to increased inflammation and
expression of tuberculosis risk signatures in recently exposed household contacts
of pulmonary tuberculosis. *Front Immunol* **13**, 1011166 (2022).

77.   Cottage, C. T. *et al.* Targeting p16-induced senescence prevents cigarette smoke-
induced emphysema by promoting IGF1/Akt1 signaling in mice. *Commun Biol* **2**,
307 (2019).

78.   Marino, F. *et al.* Diabetes-Induced Cellular Senescence and Senescence-
Associated Secretory Phenotype Impair Cardiac Regeneration and Function
Independently of Age. *Diabetes* **71**, 1081–1098 (2022).

79.   Topiwala, A. *et al.* Alcohol consumption and telomere length: Mendelian
randomization clarifies alcohol's effects. *Mol Psychiatry* **27**, 4001–4008 (2022).

80.   Restrepo, B. I. *et al.* Differential Role of Type 2 Diabetes as a Risk Factor for
Tuberculosis in the Elderly versus Younger Adults. *Pathogens* **11**, 1551 (2022).

81.   Auld, E. *et al.* HIV Infection Is Associated with Shortened Telomere Length in
Ugandans with Suspected Tuberculosis. *PLoS One* **11**, e0163153 (2016).

82.   Auld, S. C. *et al.* Pathogenesis of Post-Tuberculosis Lung Disease: Defining
Knowledge Gaps and Research Priorities at the Second International Post-
Tuberculosis Symposium. *Am J Respir Crit Care Med* **210**, 979–993 (2024).

83.   Ravimohan, S., Kornfeld, H., Weissman, D. & Bisson, G. P. Tuberculosis and lung
damage: from epidemiology to pathophysiology. *Eur Respir Rev* **27**, (2018).

84.   Auld, S. C. *et al.* Pulmonary restriction predicts long-term pulmonary impairment
in people with HIV and tuberculosis. *BMC Pulm Med* **21**, 19 (2021).

85.   Allwood, B. W. *et al.* Perspectives from the 2nd International Post-Tuberculosis
Symposium: mobilising advocacy and research for improved outcomes. *IJTLD*
*open* **1**, 111–123 (2024).

86.    Schafer, M. J. *et al.* Cellular senescence mediates fibrotic pulmonary disease. *Nat*
1023           *Commun* **8**, 14532 (2017).

1024    87.    Parveen, S. *et al.* Glutamine metabolism inhibition has dual immunomodulatory
1025           and antibacterial activities against Mycobacterium tuberculosis. *Nat Commun* **14**,
1026           7427 (2023).

1027    88.    Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and
1028           dispersion for RNA-seq data with DESeq2. *Genome Biol* **15**, 1–21 (2014).

1029    89.    Shee, S. *et al.* Moxifloxacin-Mediated Killing of Mycobacterium tuberculosis
1030           Involves Respiratory Downshift, Reductive Stress, and Accumulation of Reactive
1031           Oxygen Species. *Antimicrob Agents Chemother* **66**, (2022).

1032    90.    Merritt, C. R. *et al.* Multiplex digital spatial profiling of proteins and RNA in fixed
1033           tissue. *Nat Biotechnol* **38**, 586–599 (2020).

1034    91.    Danaher, P. *et al.* Advances in mixed cell deconvolution enable quantification of
1035           cell types in spatial transcriptomic data. *Nat Commun* **13**, 385 (2022).

1036    92.    Krug, S. *et al.* Inhibition of host PARP1 contributes to the anti-inflammatory and
1037           antitubercular activity of pyrazinamide. *Nat Commun* **14**, 8161 (2023).

1038

1039

 **Figure legends.**

 **Fig. 1. Evidence of senescence-like phenotype in BMDMs derived from young *B6.***

 ***Sst1S* and aged WT B6 mice. (a)** Schematic to determine senescence associated

 markers in BMDMs. **(b)** BMDMs were infected for 4 hours with *Mtb* H37Rv at MOI=10,

 followed by removal of extracellular bacteria by extensive washing with warm media.

 Intramacrophage survival was determined by enumerating CFU at 2 days post-infection.

 The data are means ± SEM of two independent experiments (n=4). **(c)** Median

 Fluorescence intensity of fluorescent SA-βGal substrate by live BMDMs in presence or

 absence of TNFα for 48 h. 10 µM Etoposide treated WT B6 BMDMs or BMDMs from aged

 WT B6 mice are used as positive control. **(d)** Representative images of SA-βGal- positive

 BMDMs (PFA-fixed) located at the center of the TC-treated well at 20X magnification by

 bright field microscopy. **(e)** mRNA levels of SASP-associated genes normalized to *ActB*

 are shown as determined by qPCR. **(f)** BrdU incorporation by BMDMs in absence or

 presence of TNFα, respectively with time. The data are means ± SEM. Data is

 representative of two independent experiments. (*p*>0.05: non- significant (ns), *p*<0.05: *,

 *p*<0.01: **, *p*<0.001: ***, *p*<0.0001: ***, one-way ANOVA with Tukey's (b, c, f) or Sidak's

 (e) multiple comparisons test).

 **Fig. 2. Detection of Senescence- associated markers during early stages of disease**

 **progression in *Mtb* infected mice. (a)** Schematics of Spatial transcriptomics using a

 Nanostring GeoMX Digital Spatial Profiler to determine differential gene expression in

 cells at non-controlling multibacillary granulomatous lesions (NC) relative to controlling

 lesions (C) in *B6.Sst1S* mice lungs. (n= 8 region of interests (ROI) from 2 mice). **(b)**

 Volcano plot to show differentially expressed genes between controlling and non-

controlling granulomatous lesions. Significantly upregulated (fold change> 2) and downregulated (fold change< 2) genes (*p.adj*< 0.05) in NC relative to C granulomas are shown in red and cyan, respectively. Top senescence associated genes in upregulated-DEGs are labelled. **(c)** Schematics of the experimental strategy to measure senescence markers *in vivo* at 2- and 4-weeks post infection (p.i.). Day1 CFU=39. *Mtb* H37Rv burden at indicated time points in WT B6, *B6.Sst1S* and WT B6 old mice **(d)** lungs and **(e)** spleen. The data are means ± SEM. Each data point represents a mouse. *p* was determined by two-way ANOVA with Tukey's multiple comparisons test. Representative H&E-stained and Immunohistochemistry against γH2A.X and p21 of mice lungs at **(f)** 2 weeks p.i. and **(g)** 4 weeks p.i. **(h)** ImageJ quantification of lung pathology (% inflamed area) in uninfected mice and *Mtb* H37Rv infected mice lungs at 2 weeks and 4 weeks p.i. (n=4-6). Squares indicate necrotic granulomas. **(i)** ImageJ quantification of γH2A.X-stained area in uninfected mice and *Mtb* H37Rv infected mice lungs at 2 weeks and 4 weeks p.i. (n=4-6). The data are means ± SEM. Each data point represents a mouse. *p* values are indicated as determined by statistical analysis between two groups as measured by unpaired two- tailed Student's t test (normal distribution) or Mann Whitney test.

**Fig. 3. Detection of senescent cell population and SASP cytokines by flow cytometry during early stages of disease progression in *Mtb* infected mice. (a)** Schematics of the experimental strategy to measure senescence markers (p21, p16 and SA- βGal activity) and SASP cytokines levels in mice lungs at 10 and 20 days p.i. Day 1 CFU= 55. **(b)** %p21+p16+ and **(c)** %SA-βGal+ (CellEvent Senescence green+) cells out of all live lung (uninfected and *Mtb*-infected mice) and Spleen (*Mtb*-infected mice) cells at 10 days p.i. as determined by multicolor flow cytometry. **(d)** %p21+p16+ and **(e)** %SA-

βGal+ cells out of different lung- cell types at 10 days p.i., as determined by multicolor flow cytometry. Gating for p16$^+$, p21$^+$, and SA-βGal$^+$ events for each cell type were established using reference cells from *Mtb*-infected WT B6 mice (set at 1 %). (AMs: Alveolar macrophages, IMs: Interstitial macrophages) **(f)** Normalized concentration of SASP- cytokines in lung homogenates at 20 dpi and 4 wpi as measured by a LEGENDplex™ Mouse Inflammation Panel (13-plex). The data are means ± SEM. Each data point represents a mouse (n=11-12). (two-way ANOVA with Tukey's (b- e) or Dunnett's (f) multiple comparisons test).

**Fig. 4. Global transcriptomics data by bulk-RNA Sequencing revealed upregulation of pro-survival pathways- PI3K-ERK1/2 and MAPK cascade in BMDMs derived from young *B6.Sst1S* relative to young WTB6 BMDMs, after treatment with TNFα and *Mtb* infection. (a)** Schematic of the experiment. **(b)** Selective representation of Gene-Ontology (GO) analysis for the differentially expressed genes (= 1177) indicating all enriched (*p.adj*< 0.05) signaling pathways. The size of the bubble indicates the percentage of DEGs out of total genes in that GO class and color indicates number of DEGs in the respective GO class. **(c)** Volcano plot of DEGs with upregulated genes in stressed (TNFα- treated and *Mtb*- infected) *B6.Sst1S* BMDMs (log$_2$ fold change >1; *p. adj.*< 0.05) shown in light gray and downregulated genes (log$_2$ fold change <1; *p. adj.*< 0.05) shown in light pink, relative to stressed WT B6 BMDMs. Genes representative of top significantly enriched GO class- Receptor tyrosine kinase signaling/ PI3K pathway (shown in orange), MAPK cascade (blue), anti-apoptotic genes (red) and SenMayo gene set (pink) are shown. Top expected downregulated genes- *Sp140* and *Sp110* are shown in yellow.

**Fig. 5. Senolytics DQF administration with or without anti-TB drug improved survival, body weight and reduced _Mtb_ burden in mice. (a)** Both male and female _B6.Sst1S_ mice were aerosol infected with 91 CFU of _Mtb_ H37Rv and at 3 weeks p.i., mice were oral gavaged with vehicle (n=16 mice) or indicated drug regimens (n=11 mice) for the next 4 weeks. Mice were monitored for changes in body weight and survival after the start of drug treatment. **(a)** Kaplan-Meir survival curve of _B6.Sst1S_ mice given varying drug regimens. Line indicates duration of oral gavage therapy. Indicated _p_ values are measured by log-rank (Mantel-Cox) test. **(b)** Changes in body weight are plotted as a % change in body weight of surviving mice relative to Day 1 p.i. over time. The data are means ± SEM. ($p < 0.05$: *, two-way ANOVA with Tukey's multiple comparisons test). **(c)** The experimental strategy. Day 1 CFU in _B6.Sst1S_ mice= 11; WT B6 young and old mice= 126. _Mtb_ H37Rv burden in **(d)** lung and **(e)** spleen of _B6.Sst1S_ (left), old WT B6 mice (center), and WT B6 (right), at indicated time points (n=4-7). Square indicates CFU below the limit of detection (=10). The green arrow indicates the start of drug treatment. The data are means ± SEM. Statistical analysis between two groups was done by unpaired two- tailed Student's t test.

**Fig. 6. Senolytics administration alone, and in combination with ethambutol reduced lung pathology and senescence-associated markers in _Mtb_ H37Rv infected _B6.Sst1S_ and WT B6 old mice. (a)** Representative H&E-stained images of _Mtb_-infected mice lungs treated with Vehicle or drugs, and respective ImageJ-based quantification. Yellow arrow indicates necrotic granuloma in Vehicle treated _B6.Sst1S_ mice at 5 wpi. The data are means ± SEM. Each data point represents a mouse. Statistical analysis between two groups was done by unpaired two- tailed Student's t test. Square

represents necrotic granuloma formation in the lung. **(b)** Representative γH2A.X-

Immunohistochemistry images of mice after lungs vehicle/ drugs treatment, and

respective Violin plots to show ImageJ quantification of γH2A.X-stained area. Each data

point represents a mouse. Statistical analysis between two groups was done by unpaired

two- tailed Student's t test. **(c)** %p21+βGal+ and %p16+βGal+ subpopulation observed

in all live lung cells of *Mtb*-infected mice (n= 5-8). The data are means ± SEM. Each data

point represents a mouse. Statistical analysis was calculated by two-way ANOVA with

Dunnett's multiple comparisons test.

Bubble plot to show %p21+βGal+ and %p16+βGal+ subpopulation in different lung cell

types in *Mtb*-infected **(d)** *B6.Sst1S* mice (n= 8) and **(e)** WT B6 old mice (n= 5) at indicated

time point and treatment groups. The size of the bubble indicates %subpopulation out of

all of particular cell types and color is adjusted *p* values relative to Veh group. Statistics

are calculated by one-way ANOVA with Tukey's multiple comparisons test (*p*> 0.15: ns).

1145

1146

1147

1148
1149
1150

**Fig. 1. Evidence of senescence-like phenotype in BMDMs derived from young *B6. Sst1S* and aged WT B6 mice. (a)** Schematic to determine senescence associated markers in BMDMs. **(b)** BMDMs were infected for 4 hours with *Mtb* H37Rv at MOI=10, followed by removal of extracellular bacteria by extensive washing with warm media. Intramacrophage survival was determined by enumerating CFU at 2 days post-infection. The data are means ± SEM of two independent experiments (n=4). **(c)** Median Fluorescence intensity of fluorescent SA-βGal substrate by live BMDMs in presence or absence of TNFα for 48 h. 10 μM Etoposide treated WT B6 BMDMs or BMDMs from aged WT B6 mice are used as positive control. **(d)** Representative images of SA-βGal- positive BMDMs (PFA-fixed) located at the center of the TC-treated well at 20X magnification by bright field microscopy. **(e)** mRNA levels of SASP-associated genes normalized to *ActB* are shown as determined by qPCR. **(f)** BrdU incorporation by BMDMs in absence or presence of TNFα, respectively with time. The data are means ± SEM. Data is representative of two independent experiments. ($p$>0.05: non- significant (ns), $p$<0.05: *, $p$<0.01: **, $p$<0.001: ***, $p$<0.0001: ***, one-way ANOVA with Tukey's (b, c, f) or Sidak's (e) multiple comparisons test).

**Fig. 2. Detection of Senescence- associated markers during early stages of disease progression in**
***Mtb* infected mice. (a)** Schematics of Spatial transcriptomics using a Nanostring GeoMX Digital Spatial
Profiler to determine differential gene expression in cells at non-controlling multibacillary granulomatous

lesions (NC) relative to controlling lesions (C) in *B6.Sst1S* mice lungs. (n= 8 region of interests (ROI) from 2 mice). **(b)** Volcano plot to show differentially expressed genes between controlling and non-controlling granulomatous lesions. Significantly upregulated (fold change> 2) and downregulated (fold change< 2) genes (*p.adj*< 0.05) in NC relative to C granulomas are shown in red and cyan, respectively. Top senescence associated genes in upregulated-DEGs are labelled. **(c)** Schematics of the experimental strategy to measure senescence markers *in vivo* at 2- and 4-weeks post infection (p.i.). Day1 CFU=39. *Mtb* H37Rv burden at indicated time points in WT B6, *B6.Sst1S* and WT B6 old mice **(d)** lungs and **(e)** spleen. The data are means ± SEM. Each data point represents a mouse. *p* was determined by two-way ANOVA with Tukey's multiple comparisons test. Representative H&E-stained and Immunohistochemistry against γH2A.X and p21 of mice lungs at **(f)** 2 weeks p.i. and **(g)** 4 weeks p.i. **(h)** ImageJ quantification of lung pathology (% inflamed area) in uninfected mice and *Mtb* H37Rv infected mice lungs at 2 weeks and 4 weeks p.i. (n=4-6). Squares indicate necrotic granulomas. **(i)** ImageJ quantification of γH2A.X-stained area in uninfected mice and *Mtb* H37Rv infected mice lungs at 2 weeks and 4 weeks p.i. (n=4-6). The data are means ± SEM. Each data point represents a mouse. *p* values are indicated as determined by statistical analysis between two groups as measured by unpaired two- tailed Student's t test (normal distribution) or Mann Whitney test.

**Fig. 3. Detection of senescent cell population and SASP cytokines by flow cytometry during early stages of disease progression in *Mtb* infected mice. (a)** Schematics of the experimental strategy to measure senescence markers (p21, p16 and SA- βGal activity) and SASP cytokines levels in mice lungs at 10 and 20 days p.i. Day 1 CFU= 55. **(b)** %p21+p16+ and **(c)** %SA-βGal+ (CellEvent Senescence green+) cells out of all live lung (uninfected and *Mtb*-infected mice) and Spleen (*Mtb*-infected mice) cells at 10 days p.i. as determined by multicolor flow cytometry. **(d)** %p21+p16+ and **(e)** %SA-βGal+ cells out of different lung- cell types at 10 days p.i., as determined by multicolor flow cytometry. Gating for p16[+], p21[+], and SA-βGal[+] events for each cell type were established using reference cells from *Mtb*-infected WT B6 mice (set at 1 %). (AMs: Alveolar macrophages, IMs: Interstitial macrophages) **(f)** Normalized concentration of SASP-cytokines in lung homogenates at 20 dpi and 4 wpi as measured by a LEGENDplex™ Mouse Inflammation Panel (13-plex). The data are means ± SEM. Each data point represents a mouse (n=11-12). (two-way ANOVA with Tukey's (b- e) or Dunnett's (f) multiple comparisons test).

Fig. 4. Global transcriptomics data by bulk-RNA Sequencing revealed upregulation of pro-survival pathways- PI3K-ERK1/2 and MAPK cascade in BMDMs derived from young *B6.Sst1S* relative to young WTB6 BMDMs, after treatment with TNFα and *Mtb* infection. (a) Schematic of the experiment. (b) Selective representation of Gene-Ontology (GO) analysis for the differentially expressed genes (= 1177) indicating all enriched (*p.adj*< 0.05) signaling pathways. The size of the bubble indicates the percentage of DEGs out of total genes in that GO class and color indicates number of DEGs in the respective GO class. (c) Volcano plot of DEGs with upregulated genes in stressed (TNFα- treated and *Mtb*- infected) *B6.Sst1S* BMDMs (log$_2$ fold change >1; *p. adj.*< 0.05) shown in light gray and downregulated genes (log$_2$ fold change <1; *p. adj.*< 0.05) shown in light pink, relative to stressed WT B6 BMDMs. Genes representative of top significantly enriched GO class- Receptor tyrosine kinase signaling/ PI3K pathway (shown in orange), MAPK cascade (blue), anti-apoptotic genes (red) and SenMayo gene set (pink) are shown. Top expected downregulated genes- *Sp140* and *Sp110* are shown in yellow.

**Fig. 5. Senolytics DQF administration with or without anti-TB drug improved survival, body weight and reduced *Mtb* burden in mice. (a)** Both male and female *B6.Sst1S* mice were aerosol infected with 91 CFU of *Mtb* H37Rv and at 3 weeks p.i., mice were oral gavaged with vehicle (n=16 mice) or indicated drug regimens (n=11 mice) for the next 4 weeks. Mice were monitored for changes in body weight and survival after the start of drug treatment. **(a)** Kaplan-Meir survival curve of *B6.Sst1S* mice given varying drug regimens. Line indicates duration of oral gavage therapy. Indicated *p* values are measured by log-rank (Mantel-Cox) test. **(b)** Changes in body weight are plotted as a % change in body weight of surviving mice relative to Day 1 p.i. over time. The data are means ± SEM. ($p<0.05$: *, two-way ANOVA with Tukey's multiple comparisons test). **(c)** The experimental strategy. Day 1 CFU in *B6.Sst1S* mice= 11; WT B6 young and old mice= 126. *Mtb* H37Rv burden in **(d)** lung and **(e)** spleen of *B6.Sst1S* (left), old WT B6 mice (center), and WT B6 (right), at indicated time points (n=4-7). Square indicates CFU below the limit of detection (=10). The green arrow indicates the start of drug treatment. The data are means ± SEM. Statistical analysis between two groups was done by unpaired two-tailed Student's t test.

**Fig. 6. Senolytics administration alone, and in combination with ethambutol reduced lung pathology and senescence-associated markers in *Mtb* H37Rv infected *B6.Sst1S* and WT B6 old mice. (a)** Representative H&E-stained images of *Mtb*-infected mice lungs treated with Vehicle or drugs, and respective ImageJ-based quantification. Yellow arrow indicates necrotic granuloma in Vehicle treated *B6.Sst1S* mice at 5 wpi. The data are means ± SEM. Each data point represents a mouse. Statistical analysis between two groups was done by unpaired two- tailed Student's t test. Square represents necrotic

granuloma formation in the lung. **(b)** Representative γH2A.X- Immunohistochemistry images of mice after
lungs vehicle/ drugs treatment, and respective Violin plots to show ImageJ quantification of γH2A.X-stained
area. Each data point represents a mouse. Statistical analysis between two groups was done by unpaired
two- tailed Student's t test. **(c)** %p21+βGal+ and %p16+βGal+ subpopulation observed in all live lung cells
of *Mtb*-infected mice (n= 5-8). The data are means ± SEM. Each data point represents a mouse. Statistical
analysis was calculated by two-way ANOVA with Dunnett's multiple comparisons test.

Bubble plot to show %p21+βGal+ and %p16+βGal+ subpopulation in different lung cell types in *Mtb*-
infected **(d)** *B6.Sst1S* mice (n= 8) and **(e)** WT B6 old mice (n= 5) at indicated time point and treatment
groups. The size of the bubble indicates %subpopulation out of all of particular cell types and color is
adjusted *p* values relative to Veh group. Statistics are calculated by one-way ANOVA with Tukey's multiple
comparisons test ($p > 0.15$: ns).