Layers of immunity: Deconstructing the *Drosophila* effector response

Ryckebusch F[1], Yao Tian[1], Rapin M[1], Schüpfer F[1], Hanson MA[1,2]*, Lemaitre B[1]*

1. Global Health Institute, Ecole Polytechnique Fédérale de Lausanne (EPFL), Lausanne, Switzerland
2. Centre for Ecology and Conservation, University of Exeter, Penryn, United Kingdom

These authors contributed equally to this work

Corresponding authors:
Bruno Lemaitre: bruno.lemaitre@epfl.ch
Hanson, Mark: m.hanson@exeter.ac.uk

Abstract

The host innate immune response relies on the cooperation of multiple defence modules. In insects and other arthropods, which have only innate immune mechanisms, four main immune-specific modules have been described in the defence against microbial invaders: the Toll pathway, the Imd pathway, the melanization response, and phagocytosis by plasmatocytes. Our present understanding of their relative importance remains fragmented as their contribution to host defence has never been simultaneously assessed across a large panel of pathogens. Here, we have taken advantage of newly-described immune mutants in a controlled genetic background to systematically delete these four immune modules individually, in pairs, or even all four simultaneously. Surprisingly, immune deficient flies simultaneously deficient in all four immune modules are viable, homozygous fertile, and display no overt morphological defects, suggesting these immune mechanisms are not strictly required for organismal development. With this new set of lines, we assessed the contribution of each module individually and collectively to host defense against five viruses, three fungi, eight Gram-positive bacteria, and eight Gram-negative bacteria. Our findings show that these four modules largely function independently and additively in host defense, although synergistic effects can occur for select pairs of modules. Our study confirmed the importance of the Imd pathway against Gram-negative bacteria and the Toll pathway against Gram-positive bacteria and fungi, largely via the induction of effectors such as antimicrobial peptides (AMPs) and Bomanins (Bom), but also reveals the important role of melanization against viruses, and the contribution of phagocytosis against various germs. Additionally, by examining microbial load kinetics in different mutants, we provide insight into how these modules contribute to tolerance or resistance against specific microbes. Our study provides insights into the architecture of the *Drosophila* immune system, revealing differential requirements of immune modules according to each pathogen. The set of immune deficient lines provided here offer tools to better assess the role of these immune modules in host defense.

Keywords: Drosophila, innate immunity, effectors, phagocytosis, melanization, Imd, Toll

Introduction

The immune system is made up of a network of tissues and circulating cells whose primary function is to prevent and limit pathogenic infections (Danilova, 2006; Pradeu et al., 2024; Schmid-Hempel, 2021). Although immunity has been studied in many animal species, it has been extensively characterized in only a subset of them such as the nematode *Caenorhabditis elegans,* the fruit fly *Drosophila melanogaster*, zebrafish, rodents, and humans. The traditional understanding of the immune system pictures it as a specialized group of cells and tissues, but this has gradually changed over recent years to a more complex picture of highly redundant multilevel defense modules. We have learned a great deal in the last few decades on the molecular mechanisms regulating each of these immune modules from recognition and signaling to effector action (ex. phagocytosis, antimicrobial peptides) (Pradeu et al., 2024). However, an immune response encompasses the concomitant activation of several modules and there has been little investigation of how each module collectively contributes to host defense. Indeed, the immune research field has traditionally focused on one module at a time. This article aims to fill this gap by collectively analyzing the specific contribution of the main immune modules involved in the *Drosophila* systemic response.

The *Drosophila* systemic immune response consists of a set of humoral and cellular reactions in the hemolymph (insect blood) when flies are systemically infected by bacteria, viruses, or fungi (Liegeois and Ferrandon, 2022; Westlake et al., 2024). A first facet of the systemic immune response is the production of host-defense effectors by the fat body and hemocytes that are secreted into the hemolymph. Among the best-characterized effectors are the antimicrobial peptides (AMPs), whose expression is induced to incredibly high levels upon infection in a race to control invading microbes (Hanson et al., 2019; Imler and Bulet, 2005). AMPs are small cationic peptides that exhibit antimicrobial activity that directly participate in microbial killing. Beyond AMPs, many other proteins and host defense peptides are secreted into the hemolymph, creating a hostile environment for invading pathogens. The systemic antimicrobial response is predominantly regulated at the transcriptional level by the Toll and Imd pathways (De Gregorio et al., 2002; Lemaitre et al., 1996). In *Drosophila*, the Toll pathway is activated by cell wall components (fungal glucans and Lysine-type peptidoglycan), as well as microbial proteases (Gottar et al., 2006; Issa et al., 2018; Leulier et al., 2003; Vaz et al., 2019). Toll signaling culminates in the activation of two NF-κB transcription factors, Dif and Dorsal, which regulate a large set of immune genes (Ip, 1993; Lemaitre et al., 1996). These encode small peptides such as the antifungal peptide Drosomycin, the Toll-regulated Bomanin peptides, and many other proteins (*e.g.* serine proteases, serpins, lipases) (Clemmons et al., 2015; De Gregorio et al., 2002; Fehlbaum et al., 1994; Ligoxygakis et al., 2003). Flies lacking the Toll pathway are viable but highly susceptible to infection by Gram-positive bacteria and fungi (Buchon et al., 2009; Lemaitre et al., 1996; Rutschmann et al., 2002). The Imd pathway is activated by DAP-type peptidoglycan produced by Gram-negative bacteria and a subset of Gram-positive bacteria (*e.g. Bacillus*) (Kaneko et al., 2004; Lemaitre et al., 1995; Leulier et al., 2003). The binding of peptidoglycan to receptors of the Peptidoglycan Recognition Protein family (PGRP-SD, PGRP-LC, PGRP-LE) initiates an intracellular signaling cascade, which ultimately activates the NF-κB factor Relish (Choe et al., 2005; Iatsenko et al., 2016; Kaneko et al., 2006; Kaneko et al., 2006; Westlake et al., 2024). The Imd pathway regulates the expression of many genes encoding antibacterial peptides, and serine proteases (De Gregorio et al., 2002; Lemaitre et al., 1997). Imd deficient flies are viable but susceptible to Gram-negative bacterial infection.

In addition to humoral pathways, melanization, and phagocytosis are two complementary mechanisms that contribute to host survival upon systemic infection. Melanization is an arthropod-specific immune response resulting in the rapid deposition of the black pigment melanin at wound or infection sites and the concomitant production of microbicidal reactive oxygen species (Cerenius et al., 2008; Nappi et al., 2009; Westlake et al., 2024). The melanization and Toll pathways are co-activated at the level of the serine proteases Hayan and Persephone, which in turn activate Spatzle-processing

enzyme (SPE) of the Toll pathway and serine proteases critical to cleavage of prophenoloxidases (PPOs) (Dudzic et al., 2019; Nakano et al., 2023; Shan et al., 2023), collectively referred to as the Toll-PO SP cascade (Westlake et al., 2024). The melanization reaction itself relies on the oxidation of phenols, resulting in the polymerization of melanin. Two PPOs (PPO1 and PPO2) are the primary catalysts for the synthesis of melanin during the melanization response upon systemic infection. Flies deficient for both *PPO1* and *PPO2* are viable, but lack hemolymph phenoloxidase activity and exhibit susceptibility to certain Gram-positive bacteria and fungi (Binggeli et al., 2014; Dudzic et al., 2015). In addition, a third PPO gene (PPO3) is specifically expressed by lamellocytes, specialized hemocytes that differentiate in larvae responding to and enveloping invading parasites (Dudzic et al., 2015).

Phagocytosis by both sessile and circulating plasmatocytes (*Drosophila* equivalent of macrophages) is thought to provide a complementary and important host defense (Melcarne et al., 2019; Ulvila et al., 2011). The use of hemocyte-deficient flies, via the targeted expression of a pro-apoptotic gene in plasmatocytes only, has shown that hemocytes contribute to survival upon systemic infection to certain bacterial species (Charroux and Royet, 2009; Defaye et al., 2009; Stephenson et al., 2022). Phagocytosis of bacteria greatly relies on two transmembrane receptors, NimC1 and Eater (Bretscher et al., 2015; Kocks et al., 2005; Kurucz et al., 2007; Melcarne et al., 2019). *NimC1, Eater* double mutant larvae are viable, have elevated hemocyte numbers at the larval stage, and their hemocytes can encapsulate and melanize macroparasites; however, their hemocytes are non-sessile and nearly totally phagocytosis deficient. *NimC1, Eater* adult flies have decreased hemocyte number and fail to perform phagocytosis, providing a good tool to assess the role of cellular response (Melcarne et al., 2019; Melcarne, 2020).

We have circumstantial evidence that these four immune modules are critical for defense against certain pathogens. However, how, when, and to what extent each of these four immune modules contribute to host defense against specific pathogens is fragmented as they have never been simultaneously assessed. Moreover, the extent to which these findings can be generalized to broad classes of pathogens remains unclear.

In this article, we have used a specific set of mutations principally affecting phagocytosis *(NimC1, Eater),* melanization *(PPO1, PPO2),* the Toll pathway *(spz),* and the Imd pathway *(Rel)* in a controlled genetic background to address their role in host defense. We then recombined these and other mutations to generate double and even quadruple mutants of the four immune modules. This set of tools allowed us to compare the survival of flies deficient in these responses against a broad panel of viruses, fungi, Gram-positive bacteria, and Gram-negative bacteria, to reveal the relative contribution of each immune mechanism to resistance against these pathogens.

Material and methods

Insect stocks

Precise details of *Drosophila* stocks used in this study are provided in **Table 1**. To minimize the influence of genetic background for the mutations used in this study, mutations were partially isogenized into the *DrosDel iso w^{1118}* genetic background (BDSC #5905) as described by Ferreira et al. (Ferreira et al., 2014; Ryder et al., 2004). Isogenized lines were generated and used for every mutant except the double mutant Toll and Imd *(Rel^{E20}, spz^{rm7})* for which homozygous isogenized flies were not viable in our hands. Double mutants for *NimC1^1; Eater^1* were poorly viable when isogenized in the *DrosDel iso w^{1118}* background, and so we used both isogenic and non-isogenic lines over the course of our study to facilitate investigation. Furthermore, the *NimC1^1; spz^{rm7}, Eater^1* triple mutants were homozygous lethal isogenized or not, and so could not be included. In addition, the line Δ*AMP14* bears a knockout of 14 different AMP genes, which includes the AMP families Drosomycin, Metchnikowin,

Cecropin, Defensin, Drosocin, Attacin, and Diptericin, described previously (Carboni et al., 2022;
Hanson et al., 2019). The $Bom^{\Delta 55C}$ mutation was characterized in Clemmons et al. (Clemmons et al.,
2015) and isogenic flies generated in Hanson et al. (Hanson et al., 2019). For larval experiments, use of
GFP-labelled CyO or TM6,Tb balancers were used to enable picking of homozygous mutant larvae.
Of note, the strains used in this study differ in their presence/absence of the $white^+$ gene, present
in the $PPO1^{\Delta}$, $NimC1^1$ and $eater^1$ mutations. In addition to its well established function in eye
pigmentation, the *white* gene can also impact host neurology and intestinal stem cell proliferation
(Ferreiro et al., 2017; Sasaki et al., 2021). We did not observe any obvious correlations between $white^+$
gene status and susceptibilities in this study. Moreover, in a previous study looking at the cumulative
effects of AMP mutations on lifespan, *white* gene status and fluorescent markers did not readily explain
differences in longevity (Hanson and Lemaitre, 2023). We therefore believe that the extreme immune
susceptibility we have created through deficiencies for pathways regulating hundreds of genes, or major
immune modules, overwhelms the potential effects of $white^+$ and other transgenic markers. For
additional information on which stocks bear which markers, see discussion in Supplementary file 1.
152
Microorganisms culture
Microbe strain information, microorganism classifications, and culture conditions are listed in **Table**
S1. Bacterial cultures and *C. albicans* were grown overnight shaking at 200rpm. *Aspergillus fumigatus*
fungus was grown on Malt Agar at room temperature until sporulation. The entomopathogenic fungus
Beauveria bassiana strain R444 was provided by Andermatt AG as spore preparations (BB-PROTEC)
which were directly used in natural infections or dissolved in PBS for septic injuries. Viruses DCV and
FHV were kindly provided by Carla Saleh (Pasteur Institute, Paris), produced in S2 cells. The
supernatant of S2 cells was titrated before being used for fly infections. Viruses SINV, DXV and IIV-6
were kindly provided by Ronald van Rij (Radboud University Medical Center, Nijmegen).
162
163 Heat-killed microbes were prepared by two repeats of boiling at 95°C for 30 minutes then freezing at -
164 20°C for 30 minutes, before storage long term at -20°C. Microbe preparations were streaked onto agar
165 plates to check for full efficiency of heat-killing before use in experiments.
166
Gene expression
Flies were inoculated by pricking in the junction of thoracic pleura with a needle dipped in a mixed
pellet containing a 1:1 mixture of OD600 = 200 heat-killed *E. coli* and *M. luteus* (final OD600 = 100
for each), and frozen at -20°C 6h, 12h and 24h post-infection. Total RNA was then extracted from pooled
samples of five flies using TRIzol reagent per manufacturer's protocol and resuspended in MilliQ dH2O.
Reverse transcription was performed using PrimeScript RT (Takara) with random hexamer and oligo
dT primers. Quantitative PCR was performed on a LightCycler 480 (Roche) in 96-well plates using
Applied Biosystems PowerUP SYBR green Master Mix (Applied Biosystems). Data points represent
pooled samples from three replicate experiments. Error bars represent one standard deviation from the
mean. Statistical analyses were performed using one-way ANOVA with Holms-Sidak multiple test
correction. Primers used in this study were:
178
Drs-F 5'- CGTGAGAACCTTTTCCAATATGAT-3'
Drs-R 5'- TCCCAGGACCACCAGCAT-3'
DptA-F: 5'-GCTGCGCAATCGCTTCTACT-3'
DptA-R: 5'-TGGTGGAGTGGGCTTCATG-3'
RPL32-F: 5'-GACGCTTCAAGGGACAGTATCTG-3'
RPL32-R: 5'-AAACGCGGTTCTGCATGAG-3'
185

Ex vivo larval hemocyte phagocytosis assays

Ex vivo phagocytosis assays were performed using a mix of equal volumes of *E. coli* and *S. aureus* AlexaFluorTM488 BioParticlesTM (Invitrogen), following manufacturer's instructions; and see (Melcarne et al., 2019). Five L3 wandering larvae from our newly generated mutant lines, or carrying the *Hml-Gal4, UAS-GFP* hemocytes marker as a control, were bled into 150µL of Schneider's insect medium (Sigma-Aldrich) containing 1µM phenylthiourea (PTU; Sigma-Aldrich). The hemocyte suspension was then transferred to 1.5 mL low binding tubes (Eppendorf, SigmaAldrich), and AlexaFluorTM488 bacteria BioParticlesTM were added. The samples were vortexed, incubated at room temperature for 2 hours to allow phagocytosis and then placed on ice to stop the reaction. The fluorescence of extracellular particles was quenched by adding 0.4% trypan blue (Sigma-Aldrich) diluted to 1/3 concentration. Phagocytosis was quantified using a flow cytometer (BD Accuri C6) to measure the fraction of cells phagocytosing, and their fluorescence intensity. Isogenic wild-type iso w^{1118} larvae and *Hml-Gal4, UAS-GFP* larvae with or without bacterial particles were used to define the gates for hemocyte counting and the thresholds for phagocytosed particle emission. The phagocytic index was calculated as follows:

$$\text{Fraction of haemocytes phagocytosing}(f) = \frac{[\text{number of haemocytes in fluorescence positive gate}]}{[\text{total number of haemocytes}]}$$

$$\text{Phagocytic index}(PI) = [\text{Mean fluorescence intensity of haemocytes in fluorescence positive gate}] \times f$$

Finally, due to experimenter and experiment batch differences in total hemocytes collected and knock-on effects to calculating phagocytic index, phagocytic index was normalised with the wild-type as a reference set to 100% within experiment batch.

Wounding experiment

Clean injury was performed with a needle sterilized with an EtOH and PBS wash. For imaging of the melanization reaction upon pricking, the thorax of 3-8 days old flies was pricked using a sterile needle (diameter ~0.1mm). Pictures were taken 24h post-pricking and categorized into normal, weak, or no melanization blackening seen at the injury site per Dudzic et al. (Dudzic et al., 2019).

Systemic infections and survival

Systemic infections were performed by pricking 3- to 5-day-old adult males in the thorax with a 0.1mm-thick needle dipped into a concentrated pellet of bacteria, yeast, or fungal spores. For natural infections with *A. fumigatus,* flies were anesthetized and then shaken on a sporulating plate of fungi for 30 seconds. For *B. bassiana*, flies were shaken for 30s in a tube containing an excess of commercial spores sufficient to coat all flies uniformly, of which the flies remove the vast majority later through grooming. At least two replicate survival experiments were performed for each infection, though in specific cases some genotypes may have been included in only one experiment. 20-35 flies were included per vial when possible, kept on cornmeal fly media containing (per 1L): 6.2g agar, 58.8g cornmeal, 58.8g yeast, 60mL grape fruit juice, 4.83mL propionic acid, and 26.5mL 100g/L moldex in pure ethanol. Survivals were scored daily, and flies were flipped to new food vials 3 times per week. Due to challenges working with genotypes of different viability, we could not include all genotypes in all experimental replicates. Moreover, comparisons across genotypes required multiple levels of consideration. Given the complexity of genotype-vs-reference-by-pathogen comparisons in our study, needing to compare infections both to clean injury within genotype, and to wild-type or *ΔITPM* across treatments, we chose to focus on major differences apparent in summary statistics, highlighting survival differences only

when they were: **i)** consistent across experimental replicates; **ii)** of a consistent logic across comparable
genotypes; for instance, compound mutants containing *ΔMel* (e.g. *ΔIMD, ΔMel*) should be as or more
susceptible to infections as *ΔMel* alone if melanization is truly critical for defence; and **iii)** of a mean
lifespan difference ≥1.0 days after accounting for comparisons with unchallenged or clean-injury data.
Total experiments (N exp) and total flies per genotype are reported to provide summary statistics per
pathogen-by-genotype interaction. The mean lifespan reported in survival summary data has a maximum
of 7 or 10 days depending on experimental conditions. Kaplan-Meier survival curves for all experiments
are provided in the main text or supplementary information.

Quantification of microbial load for growth kinetics
3-8 days old flies were infected with the indicated microbe and concentration per OD600 as described
in **Table S1**, and allowed to recover. At the indicated time post-infection, flies were anesthetized using
CO_2 and surface sterilized by washing them in 70% ethanol. Ethanol was removed, and then flies were
homogenized using a Precellys bead beater at 6500 rpm for 30 s in LB broth (*Pr. rettgeri*), BHI (*S.*
aureus), or YPG (*C. albicans*) with 500 µL as pools of 5 flies. These homogenates were serially diluted
and 100µL was plated on LB, BHI or YPG agar. Plates were incubated overnight, and colony-forming
units (CFUs) were counted using a Interscience Scan 500 plate scanning colony counter, and validated
independently by manual counts for a subset of plates to ensure accuracy. Statistical differences were
tested using Brown-Forsythe and Welch ANOVA tests with Dunnett's multiple test correction, which
consider unequal variance.

Results
A set of immune-deficient lines in the iso Drosdel background
Mutations affecting only one of the four main modules of the *Drosophila* systemic immune
response have been described. These include *Relish^{E20}* (*Rel^{E20}*, deficient for the Imd pathway, referred
to as *ΔIMD*), *spatzle^{rm7}* (*spz^{rm7}*, deficient for the Toll pathway, referred to as *ΔTOLL*), *PPO1^Δ, PPO2^Δ*
(no melanization, referred to as *ΔMel*), and *NimC1^1; eater^1* (reduced phagocytosis, referred to as
ΔPhag). In addition, to blocking phagocytosis, *ΔPhag* flies have a defect in hemocyte sessility, an
increased total number of larval hemocytes, and a decreased number of hemocytes at the adult stage
(Bretscher et al., 2015; Melcarne et al., 2019; Melcarne, 2020). We have preferred to use *NimC1^1; eater^1*
to study the contribution of hemocytes rather than other approaches with important indirect effects such
as the pre-injection of latex beads to saturate phagocytes (Elrod-Erickson et al., 2000), or expression of
a pro-apoptotic gene in plasmatocytes (Charroux and Royet, 2009; Defaye et al., 2009). All the
mutations in this study were partly isogenized for seven or more generations into the DrosDel *iso w^{1118}*
genetic background (Ferreira et al., 2014; Ryder et al., 2004). We then generated by recombination six
out of the seven combinations of flies simultaneously lacking two immune modules with the exception
of *ΔTOLL, ΔPhag* as *NimC1^1; eater^1, spz^{rm7}* flies were not viable (see fly strains in **Table 1**). Of note,
in our hands recombinant *Rel^{E20}, spz^{rm7}* flies were not viable in the DrosDel isogenic background forcing
us to use non-iso flies to assess the host defense of *ΔIMD, ΔTOLL* flies. Over the course of our study,
isogenic *NimC1^1;eater^1* flies also displayed viability issues, and specific experiments used non-isogenic
flies to improve confidence in phenotypes; these experiments are noted in figure captions. We used two
approaches to simultaneously delete the Toll and Melanization modules: **i)** *PPO1^Δ, PPO2^Δ; spz^{rm7}* flies
(referred to as *ΔTOLL, ΔMel^{PPS}*) and **ii)** flies carrying a small viable deletion *Hayan-psh^{Def}* (referred to
as *ΔTOLL, ΔMel^{HP}*) that removes two clustered serine protease genes, *Hayan,* and *persephone,* which
blocks the TOLL-PO SP cascade that regulates both humoral-Toll and melanization (Dudzic et al., 2019;
Westlake et al., 2024). This allowed us to delete the Toll and melanization pathways using only one
deletion event on the *X* chromosome. Excitingly, we succeeded in generating a fly line deficient for all

four immune modules with the genotype: *Hayan-psh^{Def}; NimC1^{1}; eater^{1}, Rel^{E20}* hereafter named *ΔITPM*
(ΔIMD, ΔTOLL, ΔPhag, ΔMel). These flies were homozygous viable with very poor viability, and
homozygous fertile, despite major deficiencies in these four important systemic immune mechanisms.
These flies allow us to monitor survival kinetics and pathogen growth in the near-total absence of an
immune system. Finally, we added two iso fly lines in our analysis that delete key host defense peptides
downstream of the Toll and Imd pathways: **i)** *Bom^{Δ55C}*, a short deletion removing ten Bomanins at
position 55C, which causes a major susceptibility to infection by Gram-positive bacteria and fungi
(Clemmons et al., 2015), and **ii)** *ΔAMP14*, a compound fly line of eight mutations removing 14 AMP
genes (4 Cecropins, 4 Attacins, 2 Diptericins, Drosocin, Defensin, Drosomycin, and Metchnikowin),
which causes a major susceptibility to infection by Gram-negative bacteria (Carboni et al., 2022; Hanson
et al., 2019). This set of immune-deficient fly lines in a defined genetic background provides a unique
tool kit to address the function of immune modules individually or collectively.
286
Imd, Toll, Phagocytosis, and Melanization modules function largely independently
The Imd, Toll, Phagocytosis, and Melanization modules are interconnected (Westlake et al.,
2024). For instance, the Toll and Imd pathways both regulate a subset of genes involved in melanization
(De Gregorio et al., 2002; Ligoxygakis et al., 2002). Moreover, one module can indirectly affect another.
For instance, a lack of melanization could increase the bacterial load after infection, resulting in higher
activation of the Toll and Imd pathways (Binggeli et al., 2014). In the first step, we analyzed the
activation of each of the four modules in single- and multiple-module deficient flies using classical
readouts of these modules. This also allowed monitoring of the possible effects of deleting one module
on the activity of another. We analyzed the immune response in single module deficient flies by
monitoring **i)** the expression of the antibacterial peptide gene *Diptericin* (*DptA* - a target of the Imd
pathway), and **ii)** the antifungal peptide gene *Drosomycin* (*Drs* - a target of the Toll pathway) upon
systemic infection with a mixture of heat-killed *Escherichia coli* and *Micrococcus luteus*, **iii)** ex vivo
phagocytosis of *E. coli* and S*taphylococcus aureus* coated bio-particles incubated with larval hemocytes,
and **iv)** melanization at the injury site of adult flies. This analysis first confirmed that Δ*IMD*, Δ*TOLL*,
Δ*Phag,* and Δ*Mel* flies are deficient for Imd, Toll, phagocytosis, and melanization respectively (**Figure
1A, B, C, D**). We note that the expression level of *DptA* was roughly similar to the wild-type in *ΔTOLL*
and Δ*Mel* flies, however, *ΔPhag* flies show reduced *DptA* induction at 6h compared to WT (~50%)
(**Figure 1A**). This still represents an immense induction of *DptA* after infection, but may suggest a lag
or lesser overall Imd response in *ΔPhag* flies. *DptA* expression in *ΔPhag* was, however, comparable to
wild-type at 12h and 24h. On the other hand, the *Drs* induction at 6h of *ΔMel* flies was significantly
higher than WT (~1.6x), as found previously using live *M. luteus* infection (Binggeli et al., 2014)
(**Figure 1B**). The phagocytic assay confirmed that the Δ*Phag* mutant was significantly impaired in its
ability to phagocytose bacterial bio-particles. Furthermore, Δ*IMD,* and Δ*Mel* mutant larvae had similar
levels of phagocytic index to the wild-type (**Figure 1C**), while Δ*TOLL* had a higher phagocytic index,
if anything. As expected, there was no apparent melanization at the site of the injury in Δ*Mel* flies while
melanization blackening was readily observed in the wild-type and in other single-module mutants
(**Figure 1D**). Collectively, this analysis indicates that each of the four immune modules can be
selectively blocked in the isogenic DrosDel background without interfering with the competence of the
others. We sometimes noticed a modest difference of module activity in the absence of another (higher
Toll activity in Δ*Mel,* lower IMD activity in Δ*Phag* and higher phagocytic activity in Δ*TOLL),* findings
that require further exploration to be fully established.
We then repeated our analysis using our set of module compound mutants. Not surprisingly, all
compound flies display the expected immune deficiencies validating the lines we constructed. This
analysis confirms that *ΔITPM* flies, although viable, are indeed fully immune deficient for the four host
defense modules. We did not detect any striking synergistic or antagonistic effects in which deleting

two modules has a vastly different impact compared to the additive effect of deleting the two modules individually. Notably, a small level of *Drs* expression is retained in Δ*TOLL* mutant flies, but was fully abolished in Δ*IMD, ΔTOLL* flies (**Figure 1A vs 1E**), confirming the *Drs* gene receives a small input from the Imd pathway (Leulier et al., 2000). The Δ*TOLL, ΔMel^PPS* and, Δ*TOLL, ΔMel^HP* flies display *Drs* induction at 6h that is resolved to unchallenged WT levels by 12h, consistent with this induction coming from an early input from the IMD pathway in these backgrounds. Taken together, all the compound mutant flies behave largely as expected given the sum effects of their individual mutations, confirming that these modules largely function independently.

Both the Melanization and TOLL modules contribute to wounding and lifespan

Before assessing the resistance to infection, we monitored fly survival at 25°C and 29°C to determine if these mutations impact viability in the absence of challenge. At 25°C, no single mutants died significantly more than the wild-type in the absence of challenge within the ten-day window (**Figure 2A**). However, looking at compound mutants, even in the absence of challenge, each compound mutant other than Δ*IMD, ΔTOLL* suffered 25% cumulative mortality by 10 days post-eclosion, while Δ*TOLL, ΔMel^PPS* and Δ*ITPM* flies suffered even greater cumulative mortality. Similar trends were observed at 29°C with slightly elevated mortality (**Figure 2supp1**). The observation that Δ*TOLL, ΔMel^PPS* (*PPO1[1],PPO2[1], spz^{rm7}*) flies have shorter lifespan compared to Δ*TOLL, ΔMel^HP* (*Hayan-psh^{Def}*) could be explained by residual Toll and/or PO activity in *Hayan-psh^{Def}* or additional indirect effects in Δ*TOLL, ΔMel^PPS* flies. As such, the combined effect of Toll and Melanization deficiencies in unchallenged and clean injury survival is best assessed by considering signals from both genotypes. Δ*ITPM* flies (*Hayan-psh^{Def}; NimC1[1]; Eater[1], Rel^{E20}*) have the lowest unchallenged survival rate of genotypes included in this study. Taken together, the Imd, Toll, Melanization, and Phagocytosis modules contribute little to lifespan maintenance individually, but are crucial to maintain lifespan collectively (**Figure 2B**).

Systemic infections were performed by septic injury (pricked with a contaminated needle). We therefore monitored the survival of single and double mutant flies after a clean injury to disentangle the effects of injury from those of infection. We found a cumulative mortality of ~50% for Δ*Mel* flies by day 10 (**Figure 2C**), while other single module mutants displayed relatively little mortality over the 10-day window; Δ*IMD* had notable late-onset mortality in a minority of experimental replicates, which may be due to stochastic dysbiosis effects during aging (Hanson and Lemaitre, 2023; Marra et al., 2021). Double mutant Δ*IMD, ΔTOLL* and Δ*IMD, ΔPhag* flies retained a high survival rate. Consistent with an importance of melanization in the injury response (Binggeli et al., 2014), all the compound mutants affected in melanization were susceptible to clean injury, with similar mortality seen in Δ*Phag, ΔMel*, Δ*IMD, ΔMel*, Δ*TOLL, ΔMel^PPS*, Δ*TOLL, ΔMel^HP*, and Δ*ITPM* flies (**Figure 2D**). Thus, while these genotypes differ in their base lifespan, their viability upon injury is remarkably similar. These results emphasize an interconnectedness of the melanization response with each other immune module for maintaining health after injury.

Similar results were obtained for both unchallenged and clean injury lifespans at 29°C. Of note, Δ*TOLL, ΔMel^PPS* suffered significantly greater mortality in unchallenged conditions than other genotypes, even including Δ*ITPM* flies (**Figure 2supp1**). Other compound mutant genotypes displayed a continuum of mortality in unchallenged conditions between Δ*ITPM* (~50% cumulative mortality) and Δ*TOLL, ΔMel^HP* (~15% cumulative mortality). Upon clean injury, results were broadly consistent with 25°C trends, with a more uniform mortality across all genotypes.

Collectively, our study highlights that all these immune modules contribute somewhat to the survival of flies upon clean injury, with a major role of the melanization response in maintaining lifespan and survival to injury. In the next steps of our study, we compared survival upon infection to both

369 unchallenged and clean injured flies to distinguish the impact of microbial infection from the injury
370 itself.
371

Multiple mechanisms contribute to resistance to microbial infection

To compare the contribution of each module to host defense, we performed infection
experiments of wild-type and single- and multiple-module mutant flies following septic
infection with five viruses, eight Gram-positive bacteria, eight Gram-negative bacteria, septic
injury for two fungal species, and natural infection (i.e. spore deposition on the cuticle) for two
fungal species. Temperature, dose, and days monitored are shown in **Figure 3**, and see **Table
S1** for other key parameters. Survival curves are shown in **Supplementary file 2** for each
pathogen and results of these survivals are summarized in **Figure 3** (and **Supplementary file
3**) as mean lifespans given 7 or 10 days as the maximum lifespan per treatment, with number
of tested flies indicated on the right and darker blue intensity as indicative of lower lifespan.

We performed infection experiments with the following viruses and microbial species: DCV,
FHV, DXV, IIV-6, SINV, *Aspergillus fumigatus*, *Beauveria bassiana*, *Candida albicans*,
Mycobacterium marinum, *Corynebacterium diphtheriae*, *Micrococcus luteus*, *Streptococcus
pneumoniae*, *Enterococcus faecalis*, *Staphylococcus aureus*, *Listeria monocytogenes*, *Bacillus
subtilis*, *Vibrio parahemolyticus*, *Providencia rettgeri*, *Providencia burhodogranariea*,
Pectobacterium carotavorum Ecc15, *Klebsiella pneumoniae*, *Enterobacter cloacae*,
Escherichia coli, and *Salmonella enterica ser. typhimurium*. Collectively, this represents
infections with 24 pathogens across 14 genotypes and two infection routes (336 genotype-by-
pathogen interactions). When combined with comparisons to clean injury and unchallenged
controls to validate if a lifespan difference is meaningful, and to wild-type or $\Delta ITPM$ treatments,
this inflates to 1000+ interactions. We therefore focused our survival analysis on major effects.
We report the core summary statistics and overt trends, and comment only where difference in
mean lifespan between reference genotypes or control treatments was a minimum of 1 day.

Our study confirms that the Imd pathway plays a major role against all tested Gram-negative
bacteria, as ΔIMD flies succumb more rapidly than wild-type flies to all of them. This pathway
also contributes, albeit to a lesser extent, to survival against the Gram-positive bacteria *B.
subtilis* and *L. monocytogenes,* which bear DAP-type peptidoglycan at their membrane that
triggers Imd activation (Kaneko et al., 2004; Leulier et al., 2003). We were interested to see
the impact of our ΔIMD flies in host defense to viruses as the specific mutation we used (Rel^{E20})
deletes a transcription factor common to both the canonical Imd pathway and the recently-
appreciated cGLR-Sting-Relish pathway (Goto et al., 2018). We found that ΔIMD mutant flies
were somewhat susceptible to the four viruses DCV, FHV, DXV, and IIV6, but not SINV,
agreeing with previous studies showing a reduction in lifespan of 1-2 days for *Rel* mutant flies
after viral infection (Costa et al., 2009; Goto et al., 2018; Sansone et al., 2015).

This study also confirms a prominent role of the Toll pathway in survival upon infection by all
virulent Gram-positive bacteria and fungi upon septic injury. Less virulent microbes and
infectious routes (such as *A. fumigatus* by natural infection) did not result in increased mortality
in individual $\Delta TOLL$ mutant flies. A contribution of the Toll pathway to survival was also

observed for some Gram-negative bacteria, including a minor contribution to survival upon *Pr.*
rettgeri infection, and a major contribution after *V. parahemolyticus* infection. In addition, we
found a susceptibility of ΔTOLL flies to FHV, DXV, and IIV6, suggesting Toll-regulated
effectors mediate defence against viral infection.

Use of *ΔPhag* mutants reveals a role of the cellular response in survival against the fungus *B.*
bassiana. *ΔPhag* flies further show a minor susceptibility to *S. aureus,* and a subset of Gram-
negative bacteria (*Pr. rettgeri*, *Ecc15,* and *E. coli*). Susceptibilities to Gram-negative bacteria
could result from the observed lag in Imd pathway activation (Figure 1A); indeed, previous
studies have shown a full *DptA* response is pivotal for defence against *Pr. rettgeri* (Hanson et
al., 2023, 2019; Unckless et al., 2016), and lifespans are greatly impacted by subtle differences
when using pathogens with intermediate levels of virulence (Duneau et al., 2017a). A role for
Imd in explaining *ΔPhag* susceptibilities does not detract from the importance of the cellular
response, but instead offers a mechanistic route to explore for these phenotypes.

Surprisingly, our study reveals a relatively consistent role of melanization against all the tested
viruses, notably so for DXV, IIV-6, and SINV. Δ*Mel* flies also display a minor increased
susceptibility to *B. bassiana* natural infection and septic injury. As previously described
(Dudzic et al., 2019), *ΔMel* flies are highly susceptible to the Gram-positive bacterium *S.*
aureus, rivaling the susceptibility of *ΔITPM* flies. *ΔMel* flies further show a strong
susceptibility to *Pr. rettgeri* rivalling *ΔIMD* flies, and also a minor susceptibility to infection
by other Gram-negative bacteria (*Pr. burhodogranariea, E. coli*).

Collectively, the use of single module mutants confirms the major role of the Imd pathway
against Gram-negative bacteria and Toll against virulent Gram-positive bacteria and fungi.
Interestingly, melanization was consistently important to survive virus infection, and plays a
prominent and important role in defense against specific bacterial species. While it was almost
never the most critical module, phagocytosis and the cellular response contributes to survival
against various germs. Use of single module deficient flies alongside *ΔITPM* flies reveals that
survival to some of the microbes tested can rely almost entirely on one pathway (Imd – *Ecc15*,
K. pneumoniae, E. cloacae, Pr. burhodogranariea; Toll – *E. faecalis*), or can receive notable
contributions from two pathways (Toll and IMD – *L. monocytogenes, B. subtilis, V.*
parahemolyticus; Toll and Phagocytosis – *A. fumigatus* and *B. bassiana* natural infection*;* Imd
and Melanization - DCV). Strikingly, many pathways contribute to host defense to *S. aureus*
and *Pr. rettgeri*.

Contribution of AMPs and Bomanins to Toll and Imd activities

Recent studies have described a prime importance of antimicrobial peptides (AMPs) and host
defense peptides (HDPs) in responding to infection (reviewed in Hanson (2024)). These
peptides are the principal effectors of the Toll and Imd pathways, yet mutations for AMPs and
HDPs typically have not been tested alongside mutations for both the Imd and the Toll pathways
across pathogens, as studies have typically included just the one relevant pathway mutation as
a positive control. Thus, the importance of AMPs and HDPs relative to the pathways that

regulate them has been only partially investigated. We used *ΔAMP14* and *ΔBom* flies to assess
the extent by which AMPs and Bomanins contribute to Toll and Imd activities (**Figure 3**). Here
we found that AMPs are critical for survival to infection against all Gram-negative bacteria
tested, often rivaling the susceptibility of *ΔIMD* flies. Notably, *ΔAMP14* flies were also
somewhat susceptible to the yeast *C. albicans,* and the Gram-positive bacteria *B. subtilis* and
M. luteus, reinforcing minor susceptibilities seen previously (Carboni et al., 2022; Hanson et
al., 2019). Interestingly, we found that *ΔAMP14* survivals to DCV, FHV, DXV, and IIV6,
largely paralleled those of *ΔIMD* flies (Figure 3, Supplemental File 1), suggesting Imd-
mediated survival phenotypes rely greatly on the presence of the seven classical AMP families.

On the other hand, *ΔBom* flies were susceptible to septic injury by *B. bassiana* and also
somewhat to *C. albicans* consistent with previous studies (**Supplementary file 2**), as well as
all virulent Gram-positive bacteria, consistent with their important role downstream of the Toll
pathway (Clemmons et al., 2015; Lindsay et al., 2018; Xu et al., 2023). In addition, *ΔBom* flies
paralleled the *ΔTOLL* susceptibility to DCV, FHV, and DXV, but were not susceptible to IIV6
like *ΔTOLL*.

Collectively, our study confirms the key role of antimicrobial peptides downstream of the Imd
pathway in the defense against Gram-negative bacteria, and a primary role of Bomanins
downstream of Toll in defense against fungi and Gram-positive bacteria, with little input to
survival against Gram-negative bacteria. We additionally find these secreted peptides can
contribute to survival after certain viral infections.

Immune modules mostly contribute additively to host defense

The results described above revealed the key contribution of single modules and immune
effectors to host defense. However, they did not assess the possible additive, synergistic, or
antagonistic contribution of these four modules. Moreover, they did not monitor to which extent
these modules were important in the absence of other modules. To address this, we compared
the survival of single-module deficient flies to double module-deficient flies lacking two
modules, and *ΔITPM* flies lacking all four modules (**Figure 3**).

We first observed that *ΔITPM* flies, which lack the four modules, were always as or more
susceptible than single and double module deficient flies. In multiple cases, single or double
mutants already rivalled the susceptibility of *ΔITPM* flies in microbe-specific ways. In
particular, the susceptibility of *ΔIMD*-flies to most Gram-negative bacteria rivalled *ΔITPM,* and
for some germs *ΔTOLL* (*L. monocytogenes*, *E. faecalis*, *B. bassiana* injury) or *ΔMel* (*S. aureus*)
alone were comparable to *ΔITPM*. In other cases, however, combined loss of two pathways was
necessary to rival *ΔITPM* susceptibilities. For instance, *ΔIMD, ΔTOLL* deletion causes
complete *ΔITPM*-like susceptibility to *B. subtilis*, *M. luteus*, and *E. coli*, and near-complete
susceptibility to *Sa. typhimurium*. *ΔIMD, ΔTOLL* further causes increased susceptibility to most
viruses compared to their individual mutant strains. In these cases, there is clear synergy or
additivity of the contributions of the two pathways, as either pathway alone results in only a
minor or no susceptibility. We also observed intriguing differences between our two versions

of *ΔTOLL, ΔMel* flies, as *ΔTOLL, ΔMel^{PPS}* tended to succumb to infection to a greater extent
than *ΔTOLL, ΔMel^{HP}*, consistent with the differences observed upon clean injury (**Figure 2**).
ΔTOLL, ΔMel^{PPS} displayed a minor increase in susceptibility compared to *ΔTOLL, ΔMel^{HP}* to
A. fumigatus natural infection, and more prominent susceptibilities to the yeast *C. albicans*, the
Gram-positive bacteria *M. marinum*, *M. luteus* and also to Gram-negative bacteria *K.*
pneumoniae, *E. coli*, *Pr. rettgeri* and *Pr. burhodogranariea*.

Collectively, use of double module deficient flies uncovers the role of immune modules to host
defense that were masked by other modules. While *ΔIMD* and *ΔTOLL* deficiencies sometimes
displayed a synergistic effect (i.e. little mortality of single mutants, but *ΔITPM*-like mortality
of double mutants), for combinations of other pathways the results were more often minor with
a cumulative effect on susceptibility, suggesting most modules contribute to defence
independently. The use of two variations of *ΔTOLL, ΔMel*, with *ΔTOLL, ΔMel^{PPS}* having a
stronger impact than *ΔTOLL, ΔMel^{HP},* also highlights how the level of pathway disruption can
affect infection outcomes, particularly for certain fungal or bacterial pathogens like *Providencia*
species that may cleave host proteases (suggested by Duneau et al., 2017b; Issa et al., 2018).

Timing of Toll, Imd, Phagocytosis, and Melanization contribution to host defense

Host defense programs can rely on resistance mechanisms that directly target or limit growth
of pathogens (Howick and Lazzaro, 2017; Lafont et al., 2021; Medzhitov et al., 2012). When
effectors reach a critical concentration threshold that promotes resistance, pathogen growth is
inhibited; for instance, the time to pathogen control of *Pr. rettgeri* is ~7 hours, relying heavily
on the expression and production of DptA (Duneau et al., 2017a; Hanson et al., 2023).
Phagocytosis and the melanization response are often described as providing near-immediate
immune protection (Haine et al., 2008), with reactions ex vivo progressing within minutes.
Meanwhile activation of the Imd and Toll pathways takes hours to reach peak concentrations
of host defense peptides (Uttenweiler-Joseph et al., 1998). Thus, we were curious if different
mutations would display differences in their time of action to control microbial growth. We
therefore measured the microbial growth rate for *Pr. rettgeri, S. aureus,* and *C. albicans* at
different time points post-infection in wild-type and single module mutant flies as these
pathogens were combatted by several immune modules in our survival data.

Agreeing with contributions to survival, flies singly deficient for IMD, Phagocytosis, or
Melanization modules display a higher bacterial load than wild-type when infected with *Pr.*
rettgeri (**Figure 4A**), which was apparent after 12 hours post-infection. Our observation that
ΔPhag showed impaired IMD signalling at 6h (**Figure 1A**) might explain the contribution of
phagocytosis to *Pr. rettgeri* microbial growth control with the same timing as *ΔIMD* flies. We
observed that *ΔIMD* flies, *ΔMel* flies, and *ΔITPM* flies that each succumb completely to *Pr.*
rettgeri infection, also show high and consistent microbe loads at 24h indicative of sepsis-
induced death. On the other hand, wild-type, *ΔTOLL*, and *ΔPhag* flies show a greater
stochasticity in bacterial load at 24h, consistent with a fraction of individuals surviving the
infection by controlling the pathogen. This indicates deletion of the Toll pathway and
phagocytosis does not fully ablate essential resistance mechanisms for combatting *Pr. rettgeri*.

In the case of *S. aureus* **(Figure 4B)**, we observed some individuals with higher bacterial loads in ΔPhag, and ΔIMD flies at 2h and 6h, which was not seen in wild-type, ΔTOLL, or ΔMel flies. This suggests the Imd pathway can suppress early *S. aureus* growth, and possibly also phagocytosis, although this ΔPhag could ultimately stem from lesser Imd induction. Interestingly, flies lacking melanization displayed the highest susceptibility, but did not depart from wild-type or other modules in pathogen load until 12h post-infection. While the melanization reaction is a rapid response, these results suggest that the killing activity of the melanization response acts with slower kinetics. In ΔITPM flies, we further observed earlier *S. aureus* growth at 2h and 6h like ΔIMD and ΔPhag flies. However, *S. aureus* loads at 12-24h were more comparable in ΔITPM flies to ΔMel flies. These trends agree both with survival data kinetics, and an independent action of each of these pathways in their resistance effects against *S. aureus*. This suggests that ΔITPM flies succumbs even more quickly than individual modules due to loss of multiple resistance mechanisms with different kinetics of activation that independently contribute to defense.

For *C. albicans* **(Figure 4C)**, mortality begins 2-3 days post-infection for ΔTOLL mutants, but takes place at later time points for ΔIMD, ΔPhag, and ΔMel flies. In microbial growth kinetics, ΔTOLL already shows high *C. albicans* loads at 24h consistent with a previous study (Hanson et al., 2019) and observations done with *Candida glabrata* (Quintin et al., 2013). Interestingly, a few ΔIMD flies also showed elevated growth of *C. albicans* at early time points. This suggests Imd-regulated effectors could help to suppress initial *C. albicans* growth, while Toll-regulated genes (e.g. Bomanins, Drs) contribute to this pathogen suppression to a greater extent at later time points. On the other hand, ΔPhag, and ΔMel flies did not show any increased *C. albicans* loads within the 24h time window despite onset of mortality at later time points. Finally, ΔITPM flies displayed more consistent and rapid growth of *C. albicans* than ΔTOLL or ΔIMD alone, particularly visible at 12-24h post-infection, consistent with independent contributions of both Toll and Imd in resistance to this yeast.

All survival experiments to this point were done with males. We therefore assessed key survival trends for these infections in females to learn whether the dynamics we observed were consistent across sexes **(Figure 4supp1)**. For all three pathogens (*Pr. rettgeri*, *S. aureus*, *C. albicans)* the rank order of susceptibility was broadly similar between males and females, with higher rates of mortality in females overall. Thus, we found no marked sex-by-genotype interaction. Interestingly, the greater susceptibility of females in our hands is true even for ΔITPM flies, although there were only a few surviving flies on which we can base these conclusions. However, these data may suggest the sexual dimorphism in defense against infection that we see against these pathogens is due to factors independent of the immune modules we disrupted.

Collectively, we observed that microbial growth parallels the susceptibility of modules indicating that they work in resistance. Surprisingly we did not observe an early role in pathogen killing for melanization despite melanization regulating a rapid blackening response in ex vivo assays (Nakhleh et al., 2017). We further recover stochasticity in microbe loads across genotypes as expected when only a subset of individuals suppress the pathogen and survive the

569 infection (Duneau et al., 2017a). The more rapid and more consistent microbial growth kinetics
570 of *ΔITPM* flies, particularly visible by 18h, further demonstrate that these pathways contribute
571 to resistance collectively.

Contributions of individual modules to disease tolerance

An increasingly used metric to delineate roles of resistance and tolerance is the Pathogen Load
Upon Death (PLUD) (Duneau et al., 2017a; Duneau et al., 2025). The PLUD is determined by
the virulence of the pathogen to set an upper limit, but also the tolerance of the host in surviving
to a given pathogen burden before succumbing to infection. We monitored PLUDs for the same
pathogens: *Pr. rettgeri, S. aureus,* and *C. albicans*. The PLUD of both *Pr. rettgeri* and *S. aureus*
upon death was overall similar in wild-type and single-module mutant flies **(Figure 5A-B)**.
Notably, there was a greater stochasticity in *Pr. rettgeri* infections trending towards lower
PLUD in *ΔITPM* flies, suggesting a stochastic reduction in disease tolerance of *ΔITPM* flies to
this bacterium, although this was not significant compared to wild-type (P>.05). We saw no
statistical differences in PLUD across genotypes for *S. aureus*.

For *C. albicans*, the distribution of PLUD values for *ΔPhag* flies was markedly different from
wild-type, also seen to some extent for *ΔMel*. Interestingly, flies with these genotypes suppress
C. albicans growth well **(Figure 4C)**. Yet here we recovered many individuals from both of
these module-deficient lines that died with a far lower PLUD **(Figure 5C)**. Taken together, this
suggests that the humoral immune pathways, particularly Toll, contribute to resistance against
C. albicans. However, the cellular response and melanization reaction instead regulate
tolerance to *C. albicans* infection. The use of *ΔITPM* flies further emphasizes that when both
resistance (Toll, Imd) and tolerance (Phagocytosis, Melanization) mechanisms are deficient,
the loss of resistance plays a primary importance, and can mask the loss of tolerance from being
observed.

Collectively, monitoring of PLUD suggests that some modules, specifically phagocytosis and
melanization, contribute to tolerance of certain infections. Importantly, this tolerance effect was
only seen in genotypes that can resist infection. These findings demonstrate the collective
contribution to resistance and/or tolerance of the four immune modules studied here.

Discussion

In this article, we have generated a set of single and compound immune module deficient fly
lines in a controlled genetic background. Using various assays, we confirmed the validity of
our lines, revealing that each module can be activated independently. We did, however, observe
a higher Toll activation in *ΔMel* flies upon systemic infection with dead bacteria. Future studies
may reveal if this higher Toll activity in *ΔMel* flies involves a regulatory pathway, or perhaps
reflects higher persistence of microbial elicitors (e.g. peptidoglycan) that activate the Toll
pathway. We also observed a lesser Imd activation in *ΔPhag* flies at 6 hours post-infection,
which could reflect a role for hemocytes to stimulate the Imd systemic response. In this study,
we used *NimC1[1]; eater[1]* double mutants to assess the cellular response. These flies have defects
in phagocytosis, but also adhesion and sessility (Melcarne et al., 2019). *NimC1[1]; eater[1]* larvae

also have increased hemocyte number at the larval stage, but adults rapidly lose Hml positive
cells (Melcarne, 2020). Despite these limitations, we believe that the mutations we have used
here are among the best available to assess the role of these four modules to host defense.
Surprisingly, we could produce a fly line lacking the four main defense mechanisms of the
systemic immune response, indicating that none of these modules is essential for survival.
Theoretically, $\Delta ITPM$ flies are almost completely immune deficient but they can still clot
wounds, activate the JNK and JAK-STAT pathways that mediate the wound healing response,
and retain constitutive immune defense molecules that could also provide a certain degree of
protection. Consistent with several studies (Capilla et al., 2017; Carvalho et al., 2014; Rämet et
al., 2002), we show that Toll and melanization synergistically contribute to wound healing in
adults. However, the observation that $\Delta Phag$, ΔMel or ΔIMD, ΔMel double mutant flies are also
more susceptible to clean injury than ΔMel flies indicates that the Imd pathway and
phagocytosis also contribute to wound healing. Thus, although immune deficient fly lines for
the four modules are viable, the use of compound mutation and $\Delta ITPM$ flies reveals a clear role
of the immune system in response to wounding and lifespan maintenance.

Use of single and double module-deficient flies provides key insights on the mechanisms used
by *Drosophila* to combat infection. We could confirm previous studies revealing the role of
Imd against Gram-negative bacteria, Toll against Gram positive bacteria and virulent fungi, and
an importance of melanization and phagocytosis against specific pathogens (Charroux and
Royet, 2009; Defaye et al., 2009; Garg and Wu, 2014; Nehme et al., 2011). Surprisingly,
melanization was a consistently important module in survival to virus infection. It seems
unlikely that prophenoloxidase activity taking place in the hemolymph could combat viral
agents that are intracellular. We instead speculate this contribution of melanization to surviving
virus infection could be due to its role in wound healing, or perhaps autotoxic contributions of
melanization reaction intermediates that fail to convert in phenoloxidase-deficient flies. We
additionally recovered a role of Imd signalling (i.e. *Relish*) in antiviral defense, which was
expected given recent characterizations of cGLR-Sting-Relish antiviral immunity (Ai et al.,
2024; Cai et al., 2022; Goto et al., 2018). However, susceptibilities of Rel^{E20} flies were
paralleled by $\Delta AMP14$ in all cases, suggesting AMPs contribute to this susceptibility. While
Sting regulates a number of genes likely important for antiviral defense (Goto et al., 2018), the
susceptibility we observe here could be a direct action of AMPs on enveloped viruses as
described in some studies (Feng et al., 2020; Huang et al., 2013; Yasin et al., 2004), or an
indirect effect, such as a failure to regulate gut microbes after the damage induced by viral
replication (Marra et al., 2021). We additionally used dual modes of infection for the fungus *B.*
bassiana, finding an importance of melanization and Toll signalling in both infection modes.
However, our study reveals that Bomanin effectors explain most of the Toll contribution upon
septic injury but not natural infection, for *B. bassiana*. This observation is in line with different
importance of effectors or modules according to the route of infection (Martins et al., 2013).

Our double module mutant analysis reveals that most modules contribute additively to host
defense. This is consistent with these modules functioning independently. However, we noticed
instances of synergy between two pathways, notably including Toll and Imd. Synergy between
Toll and Imd can be explained by the fact that many immune-inducible regulated genes receive

input from both the Imd and Toll pathways, including genes like *Metchnikowin*, *Drosomycin*
and *Transferrin1* (De Gregorio et al., 2002). Finally, we also observed rarer cases of synergistic
susceptibility in flies deficient for Toll and Melanization or Imd and Melanization.

Our study confirms that the Toll and Imd humoral modules provide a broad role against certain
classes of pathogens, Imd for Gram-negative bacteria and DAP-type containing Gram positive
bacteria, and Toll for Fungi and Gram-positive bacteria. Use of AMP and Bomanin mutants
revealed that this can be largely explained by the effectors they control. In contrast, the
contribution of phagocytosis and melanization appears to be critical to a more specific and
diverse set of pathogens. We speculate that phagocytes or melanization are critical to handle
bacteria that resist host defense peptides (Hanson et al., 2019), or can hide from them (Touré et
al., 2023). The melanization reaction is a source of ROS that is potent against pathogens such
as *S. aureus* that have been shown to be sensitive to ROS, and resistant to Toll and Imd defenses
(Dudzic et al., 2019; Ford and King, 2021; Needham et al., 2004; Ramond et al., 2021). These
two modules play important roles in immune related processes such as encapsulation
(melanization), the uptake of bacteria escaping from the gut, or tissue homeostasis
(phagocytosis) (Braun et al., 1998; C. Melcarne et al., 2019; Nehme et al., 2007) that were not
assessed here. Collectively, our study validates, with minor discrepancies, many studies that
have assessed the contribution of these modules individually (ex. (Apidianakis et al., 2005;
Binggeli et al., 2014; Lamiable et al., 2016; Lemaitre et al., 1996, 1995; Nehme et al., 2011).
Our mutant lines can now be used to analyze the contribution of these immune modules in
resistance to other pathogens, notably wasps, nematodes, microsporidia and protozoans or in
other contexts such as mating and local infection.

While some immune modules play a predominant role against some pathogens, other pathogens
are handled by multiple modules. We hypothesize that pathogens that have intermediate levels
of virulence, killing only a fraction of wild-type flies, may better reveal the role of multiple
modules. Indeed, the stochasticity in survival analysis partly stems from the arms-race
occurring between the pathogen and host immunity, as shown for *Pr. rettgeri* (Duneau et al.,
2017a). In this condition, any small change in the immune system may tip the outcome of the
arm-race toward lethality or survival. In this line, it is notable that multiple modules were
important to survive *Pr. rettgeri* infection. Previous studies have revealed a major role of the
Imd pathway-regulated AMP Diptericin A against this bacterium (Hanson et al., 2023, 2019;
Unckless et al., 2016). However, Duneau et al. showed that survival patterns to *Pr. rettgeri*
bifurcate into two outcomes based on time taken to fully activate systemic defenses (Duneau
et al., 2017a), and showed a role for a Toll-PO SP cascade regulating serine protease in defense
against this microbe (Duneau et al., 2017b). We may speculate that melanization, although not
as uniquely critical as Diptericin, might tip the balance of this arms race towards host lethality,
resulting in comparable survival phenotypes. The observation that *ΔPhag* flies expressed less
DptA (**Figure 1**) can also explain their susceptibility. Future studies will clarify how these
multiple modules can contribute to host survival according to additive or achilles dynamics –
the concept that microbes have generic or specific weaknesses that host effectors target
(Hanson, 2024). It will also be important to consider the distinct roles of resistance, tolerance,

and resilience in host defense (Howick and Lazzaro, 2017; Wukitch et al., 2023; Duneau et al.,
2025).

We observed a good correlation between survival analysis and pathogen growth in single
module flies for *Pr. rettgeri*, *S. aureus,* and *C. albicans*. This indicates that these immune
modules mostly contribute to resistance mechanisms that target pathogen growth. Our study
did not reveal key early contributions of phagocytosis or melanization to control pathogen
growth, despite their quasi-immediate activation; higher *S. aureus* growth at 2 hours in *ΔPhag*
flies was paralleled by *ΔImd* flies, and we observed lesser Imd activation in *ΔPhag* flies.
Melanization, while being critical to survive *S. aureus* infection, impacts bacterial growth
beginning only at the 12 hour time point. This indicates that the Melanization microbicidal
activity in vivo takes place slower than the blackening reaction seen from bled hemolymph.
Interestingly, ΔPhag and also ΔMel flies could suppress *C. albicans* yeast growth, but
ultimately some individuals succumb to infection with lower PLUD levels. We confirmed
ΔPhag PLUD results using both an isogenic and a second wild-type genetic background,
suggesting this lower PLUD is genuine. Susceptibility to fungal infection independent of fungal
proliferation has also been reported using an *A. fumigatus* septic infection model, and relies on
the protection offered by Bomanins from pathogen-secreted toxins (Xu et al., 2023). It is
tempting to speculate, based on the modules involved, that the loss of tolerance to *C. albicans*
we observe has to do with wound repair or accumulating damage, such as what has been
reported in beetles that suffer renal failure (Khan et al., 2017; Li et al., 2020), or flies with
autotoxic trachea degradation upon stress pathway disruption (Rommelaere et al., 2024). Thus,
this tolerance effect could rely on pathogen-mediated or autotoxic damage, which may be
elucidated in a future study.

Here we have provided a single and double mutant analysis of *Drosophila* immune module
functions. Our study provides several insights on what modules are most important to survive
infection by defined pathogens. However, we also highlight the collective contribution of
modules to defence even when one module is of an outsized importance. We extend our
comprehension of innate immune responses by revealing higher complexity, implicating
multiple host defense modules in survival to various germs, including some with more cryptic
contributions. As illustrated by our previous characterizations of AMP function (Carboni et al.,
2022; Hanson et al., 2019), the melanization response (Dudzic et al., 2015), and stress-induced
Turandot proteins (Rommelaere et al., 2024), a combinatorial mutation approach to deciphering
immune functions can be extended even to the broad level of immune modules. Of note, we
were unable to systematically sample all genotype-by-pathogen interactions equally. We have
therefore been highly conservative in our reporting of major effects. There are likely many
important interactions not discussed in our study. Future investigations may highlight important
biology that is apparent in our data, but which we may not have mentioned here. To this end,
we have deposited our isogenic immunity fly stocks in the Vienna Drosophila Resource Centre
to facilitate their use. Beyond immunity, our tools can also be of use to study various questions
at the cutting edge of aging, memory, neurodegeneration, cancer, and more, where immune
genes have been implicated repeatedly. We hope that this set of lines will be useful to the
community to better characterize the *Drosophila* host defense.

Acknowledgement:

We thank Mélanie Blokesch (EPFL), Jan-Willem Veening (Unil), Vivek Thacker (Heidelberg University), Carla Saleh (Pasteur Institut) and Ronald P van Rij (Radboud University Medical Center) for key reagents. We thank Luis Teixeira for iso Drosdel wild-type, rel and spz flies. We thank Elodie Koenig, Prince Kumar Sah, and Hannah Westlake for experimental help and Hannah Westlake for editing. The AMP, Bom and single and quadruple module mutants were deposited at the Vienna *Drosophila* Research Center. This project was supported by the SNSF grant 310030_215073 awarded to BL, and Wellcome Trust grant 227559/Z/23/Z awarded to MAH.

Author contributions

Conceptualization: FR, MAH, BL
Data curation: FR, MAH
Funding acquisition: BL, MAH
Investigation: FR, YT, MR, FS
Methodology: FR
Project administration: BL
Validation: YT, MAH
Resources: FR, MAH
Supervision: MAH, BL
Writing – original draft: FR, BL
Writing – review & editing: MAH, BL

References

Ai X, Deng H, Li X, Wei Z, Chen Y, Yin T, Zhang J, Huang J, Li H, Lin X, Tan L, Chen D, Zhang Xiaohan, Zhang Xiuqing, Meignin C, Imler J-L, Cai H. 2024. cGAS-like receptors drive a systemic STING-dependent host response in Drosophila. *Cell Reports* **43**:115081. doi:10.1016/j.celrep.2024.115081

Apidianakis Y, Mindrinos MN, Xiao W, Lau GW, Baldini RL, Davis RW, Rahme LG. 2005. Profiling early infection responses: Pseudomonas aeruginosa eludes host defenses by suppressing antimicrobial peptide gene expression. *Proc Natl Acad Sci U S A* **102**:2573–8.

Binggeli O, Neyen C, Poidevin M, Lemaitre B. 2014. Prophenoloxidase Activation Is Required for Survival to Microbial Infections in Drosophila. *PLoS Pathogens* **10**. doi:10.1371/journal.ppat.1004067

Braun A, Hoffmann JA, Meister M. 1998. Analysis of the Drosophila host defense in domino mutant larvae, which are devoid of hemocytes. *Proc Natl Acad Sci U S A* **95**:14337–42.

Bretscher AJ, Honti V, Binggeli O, Burri O, Poidevin M, Kurucz E, Zsamboki J, Ando I, Lemaitre B. 2015. The Nimrod transmembrane receptor Eater is required for hemocyte attachment to the sessile compartment in Drosophila melanogaster. *Biology open* **4**:355–63. doi:10.1242/bio.201410595

Buchon N, Poidevin M, Kwon H-M, Guillou A, Sottas V, Lee B-L, Lemaitre B. 2009. A
single modular serine protease integrates signals from pattern-recognition receptors
upstream of the Drosophila Toll pathway. *Proc Natl Acad Sci U S A* **106**:12442–12447.
doi:10.1073/pnas.0901924106

Cai H, Meignin C, Imler J-L. 2022. cGAS-like receptor-mediated immunity: the insect
perspective. *Curr Opin Immunol* **74**:183–189. doi:10.1016/j.coi.2022.01.005

Capilla A, Karachentsev D, Patterson RA, Hermann A, Juarez MT, McGinnis W. 2017. Toll
pathway is required for wound-induced expression of barrier repair genes in the
Drosophila epidermis. *Proceedings of the National Academy of Sciences* **114**:E2682–
E2688. doi:10.1073/pnas.1613917114

Carboni AL, Hanson MA, Lindsay SA, Wasserman SA, Lemaitre B. 2022. Cecropins
contribute to *Drosophila* host defense against a subset of fungal and Gram-negative
bacterial infection. *Genetics* **220**. doi:10.1093/genetics/iyab188

Carvalho L, Jacinto A, Matova N. 2014. The Toll/NF-κB signaling pathway is required for
epidermal wound repair in Drosophila. *PNAS* **111**:E5373–E5382.
doi:10.1073/pnas.1408224111

Cerenius L, Lee BL, Söderhäll K. 2008. The proPO-system: pros and cons for its role in
invertebrate immunity. *Trends in Immunology* **29**:263–271. doi:10.1016/j.it.2008.02.009

Charroux B, Royet J. 2009. Elimination of plasmatocytes by targeted apoptosis reveals their
role in multiple aspects of the Drosophila immune response. *Proc Natl Acad Sci U S A*
106:9797–802. doi:10.1073/pnas.0903971106

Choe K-M, Lee H, Anderson KV. 2005. Drosophila peptidoglycan recognition protein LC
(PGRP-LC) acts as a signal-transducing innate immune receptor. *Proceedings of the
National Academy of Sciences* **102**:1122–1126. doi:10.1073/pnas.0404952102

Clemmons AW, Lindsay SA, Wasserman SA. 2015. An Effector Peptide Family Required for
Drosophila Toll-Mediated Immunity. *PLoS Pathogens* **11**.
doi:10.1371/journal.ppat.1004876

Costa A, Jan E, Sarnow P, Schneider D. 2009. The Imd pathway is involved in antiviral
immune responses in Drosophila. *PLoS ONE* **4**:e7436.
doi:10.1371/journal.pone.0007436

Danilova N. 2006. The evolution of immune mechanisms. *J Exp Zool* **306B**:496–520.
doi:10.1002/jez.b.21102

De Gregorio E, Spellman PT, Tzou P, Rubin GM, Lemaitre B. 2002. The Toll and Imd
pathways are the major regulators of the immune response in Drosophila. *Embo J*
21:2568–79.

Defaye A, Evans I, Crozatier M, Wood W, Lemaitre B, Leulier F. 2009. Genetic Ablation of
Drosophila Phagocytes Reveals Their Contribution to Both Development and Resistance
to Bacterial Infection. *Journal of Innate Immunity* **1**:322–334. doi:10.1159/000210264

Dudzic JP, Hanson MA, Iatsenko I, Kondo S, Lemaitre B. 2019. More Than Black or White:
Melanization and Toll Share Regulatory Serine Proteases in Drosophila. *Cell Reports*
27:1050-1061.e3. doi:10.1016/j.celrep.2019.03.101

Dudzic JP, Kondo S, Ueda R, Bergman CM, Lemaitre B. 2015. Drosophila innate immunity:
regional and functional specialization of prophenoloxidases. *BMC biology* **13**:81–97.
doi:10.1186/s12915-015-0193-6

Duneau D, Ferdy J-B, Revah J, Kondolf H, Ortiz G, Lazzaro B, Buchon N. 2017a. Stochastic
variation in the initial phase of bacterial infection predicts the probability of survival in
D. melanogaster. *eLife* **6**. doi:10.7554/eLife.28298

Duneau DF, Kondolf H, Im J, Ortiz G, Chow C, Fox M, Eugénio A, Revah J, Buchon N,
Lazzaro B. 2017b. The Toll pathway underlies host sexual dimorphism in resistance to
both Gram-negative and Gram-positive bacteria in mated Drosophila. *BMC Biology* **15**.
doi:10.1186/s12915-017-0466-3

Duneau D, Lafont PD, Lauzeral C, Parthuisot N, Faucher C, Jin X, Buchon N, Ferdy J-B.
2025. A within-host infection model to explore tolerance and resistance. *eLife*
14:e104052. doi:10.7554/eLife.104052

Elrod-Erickson M, Mishra S, Schneider D. 2000. Interactions between the cellular and
humoral immune responses in drosophila. *Curr Biol* **10**:781–4.

Fehlbaum P, Bulet P, Michaut L, Lagueux M, Broekaert WF, Hetru C, Hoffmann JA. 1994.
Insect immunity: Septic injury of drosophila induces the synthesis of a potent antifungal
peptide with sequence homology to plant antifungal peptides. *Journal of Biological*
Chemistry **269**:33159–33163.

Feng M, Fei S, Xia J, Labropoulou V, Swevers L, Sun J. 2020. Antimicrobial Peptides as
Potential Antiviral Factors in Insect Antiviral Immune Response. *Frontiers in*
Immunology **11**.

Ferreira AG, Naylor H, Esteves SS, Pais IS, Martins NE, Teixeira L. 2014. The Toll-dorsal
pathway is required for resistance to viral oral infection in Drosophila. *PLoS pathogens*
10:e1004507. doi:10.1371/journal.ppat.1004507

Ferreiro MJ, Pérez C, Marchesano M, Ruiz S, Caputi A, Aguilera P, Barrio R, Cantera R.
2017. Drosophila melanogaster White Mutant w1118 Undergo Retinal Degeneration.
Front Neurosci **11**:732. doi:10.3389/fnins.2017.00732

Ford SA, King KC. 2021. In Vivo Microbial Coevolution Favors Host Protection and Plastic
Downregulation of Immunity. *Molecular Biology and Evolution* **38**:1330–1338.
doi:10.1093/molbev/msaa292

Garg A, Wu LP. 2014. *D rosophila* Rab14 mediates phagocytosis in the immune response to
S taphylococcus aureus: Rab14 is critical for cellular immune responses to *S. aureus*.
Cellular Microbiology **16**:296–310. doi:10.1111/cmi.12220

Goto A, Okado K, Martins N, Cai H, Barbier V, Lamiable O, Troxler L, Santiago E, Kuhn L,
Paik D, Silverman N, Holleufer A, Hartmann R, Liu J, Peng T, Hoffmann JA, Meignin
C, Daeffler L, Imler JL. 2018. The Kinase IKKβ Regulates a STING- and NF-κB-

Dependent Antiviral Response Pathway in Drosophila. *Immunity* **49**:225-234.e4. doi:10.1016/j.immuni.2018.07.013

Gottar M, Gobert V, Matskevich AA, Reichhart J-M, Wang C, Butt TM, Belvin M, Hoffmann JA, Ferrandon D. 2006. Dual Detection of Fungal Infections in Drosophila via Recognition of Glucans and Sensing of Virulence Factors. *Cell* **127**:1425–1437. doi:10.1016/j.cell.2006.10.046

Haine ER, Moret Y, Siva-Jothy MT, Rolff J. 2008. Antimicrobial Defense and Persistent Infection in Insects. *Science* **322**:1257–1259. doi:10.1126/science.1165265

Hanson MA. 2024. When the microbiome shapes the host: immune evolution implications for infectious disease. *Phil Trans R Soc B* **379**:20230061. doi:10.1098/rstb.2023.0061

Hanson MA, Dostálová A, Ceroni C, Poidevin M, Kondo S, Lemaitre B. 2019. Synergy and remarkable specificity of antimicrobial peptides in vivo using a systematic knockout approach. *eLife* **8**. doi:10.7554/elife.44341

Hanson MA, Grollmus L, Lemaitre B. 2023. Ecology-relevant bacteria drive the evolution of host antimicrobial peptides in *Drosophila*. *Science* **381**:eadg5725. doi:10.1126/science.adg5725

Hanson MA, Lemaitre B. 2023. Antimicrobial peptides do not directly contribute to aging in *Drosophila* , but improve lifespan by preventing dysbiosis. *Disease Models & Mechanisms* **16**:dmm049965. doi:10.1242/dmm.049965

Howick VM, Lazzaro BP. 2017. The genetic architecture of defence as resistance to and tolerance of bacterial infection in Drosophila melanogaster. *Molecular Ecology* **26**:1533–1546. doi:10.1111/mec.14017

Huang Z, Kingsolver MB, Avadhanula V, Hardy RW. 2013. An Antiviral Role for Antimicrobial Peptides during the Arthropod Response to Alphavirus Replication. *Journal of Virology* **87**:4272–4280. doi:10.1128/jvi.03360-12

Iatsenko I, Kondo S, Mengin-Lecreulx D, Lemaitre B. 2016. PGRP-SD, an Extracellular Pattern-Recognition Receptor, Enhances Peptidoglycan-Mediated Activation of the Drosophila Imd Pathway. *Immunity* **45**:1013–1023. doi:10.1016/j.immuni.2016.10.029

Imler J-L, Bulet P. 2005. Antimicrobial peptides in Drosophila: structures, activities and gene regulation. *Chemical immunology and allergy* **86**:1–21. doi:10.1159/000086648

Ip Y. 1993. Dif, a dorsal-related gene that mediates an immune response in Drosophila. *Cell* **75**:753–763. doi:10.1016/0092-8674(93)90495-C

Issa N, Guillaumot N, Lauret E, Matt N, Schaeffer-Reiss C, Van Dorsselaer A, Reichhart JM, Veillard F. 2018. The Circulating Protease Persephone Is an Immune Sensor for Microbial Proteolytic Activities Upstream of the Drosophila Toll Pathway. *Molecular Cell* **69**:648-663.e7. doi:10.1016/j.molcel.2018.01.029

Kaneko T, Goldman WE, Mellroth P, Steiner H, Fukase K, Kusumoto S, Harley W, Fox A, Golenbock D, Silverman N. 2004. Monomeric and polymeric gram-negative

peptidoglycan but not purified LPS stimulate the Drosophila IMD pathway. *Immunity*
20:637–49.

Kaneko Takashi, Yano T, Aggarwal K, Lim J-H, Ueda K, Oshima Y, Peach C, Erturk-
Hasdemir D, Goldman WE, Oh B-H, Kurata S, Silverman N. 2006. PGRP-LC and
PGRP-LE have essential yet distinct functions in the drosophila immune response to
monomeric DAP-type peptidoglycan. *Nat Immunol* **7**:715–723. doi:10.1038/ni1356

Kaneko T., Yano T, Aggarwal K, Lim JH, Ueda K, Oshima Y, Peach C, Erturk-Hasdemir D,
Goldman WE, Oh BH, Kurata S, Silverman N. 2006. PGRP-LC and PGRP-LE have
essential yet distinct functions in the drosophila immune response to monomeric DAP-
type peptidoglycan. *Nat Immunol* **7**:715–23.

Khan I, Agashe D, Rolff J. 2017. Early-life inflammation, immune response and ageing.
Proceedings of the Royal Society B: Biological Sciences **284**:20170125.
doi:10.1098/rspb.2017.0125

Kocks C, Cho JH, Nehme N, Ulvila J, Pearson AM, Meister M, Strom C, Conto SL, Hetru C,
Stuart LM, Stehle T, Hoffmann JA, Reichhart J-M, Ferrandon D, Rämet M, Ezekowitz
RAB. 2005. Eater, a Transmembrane Protein Mediating Phagocytosis of Bacterial
Pathogens in Drosophila. *Cell* **123**:335–346. doi:10.1016/j.cell.2005.08.034

Kurucz É, Márkus R, Zsámboki J, Folkl-Medzihradszky K, Darula Z, Vilmos P, Udvardy A,
Krausz I, Lukacsovich T, Gateff E, Zettervall C-J, Hultmark D, Andó I. 2007. Nimrod, a
Putative Phagocytosis Receptor with EGF Repeats in Drosophila Plasmatocytes. *Current*
Biology **17**:649–654. doi:10.1016/j.cub.2007.02.041

Lafont PDM, Lauzeral C, Parthuisot N, Faucher C, Duneau D, Ferdy J-B. 2021. A within-host
infection model to explore tolerance and resistance (preprint). Evolutionary Biology.
doi:10.1101/2021.10.19.464998

Lamiable O, Arnold J, de Faria IJ da S, Olmo RP, Bergami F, Meignin C, Hoffmann JA,
Marques JT, Imler J-L. 2016. Analysis of the Contribution of Hemocytes and Autophagy
to Drosophila Antiviral Immunity. *Journal of Virology* **90**:5415–5426.
doi:10.1128/JVI.00238-16

Lemaitre B, Kromer-Metzger E, Michaut L, Nicolas E, Meister M, Georgel P, Reichhart JM,
Hoffmann JA. 1995. A recessive mutation, immune deficiency (imd), defines two
distinct control pathways in the Drosophila host defense. *Proc Natl Acad Sci U S A*
92:9465–9469. doi:10.1073/pnas.92.21.9465

Lemaitre B, Nicolas E, Michaut L, Reichhart J-M, Hoffmann JA. 1996. The Dorsoventral
Regulatory Gene Cassette spätzle/Toll/cactus Controls the Potent Antifungal Response
in Drosophila Adults. *Cell* **86**:973–983. doi:10.1016/S0092-8674(00)80172-5

Lemaitre B, Reichhart JM, Hoffmann JA. 1997. Drosophila host defense: differential
induction of antimicrobial peptide genes after infection by various classes of
microorganisms. *Proceedings of the National Academy of Sciences of the United States*
of America **94**:14614–9. doi:10.1073/pnas.94.26.14614

Leulier F, Parquet C, Pili-Floury S, Ryu J-H, Caroff M, Lee W-J, Mengin-Lecreulx D,
Lemaitre B. 2003. The Drosophila immune system detects bacteria through specific
peptidoglycan recognition. *Nat Immunol* **4**:478–484. doi:10.1038/ni922

Leulier F, Rodriguez A, Khush RS, Abrams JM, Lemaitre B. 2000. The Drosophila caspase
Dredd is required to resist gram-negative bacterial infection. *EMBO Rep* **1**:353–8.
doi:10.1093/embo-reports/kvd073

Li X, Rommelaere S, Kondo S, Lemaitre B. 2020. Renal Purge of Hemolymphatic Lipids
Prevents the Accumulation of ROS-Induced Inflammatory Oxidized Lipids and Protects
Drosophila from Tissue Damage. *Immunity* **52**:374-387.e6.
doi:10.1016/j.immuni.2020.01.008

Liegeois S, Ferrandon D. 2022. Sensing microbial infections in the Drosophila melanogaster
genetic model organism. *Immunogenetics* **74**:35–62. doi:10.1007/s00251-021-01239-0

Ligoxygakis P, Pelte N, Ji C, Leclerc V, Duvic B, Belvin M, Jiang H, Hoffmann JA,
Reichhart JM. 2002. A serpin mutant links Toll activation to melanization in the host
defence of Drosophila. *Embo J* **21**:6330–7.

Ligoxygakis P, Roth S, Reichhart J-M. 2003. A Serpin Regulates Dorsal-Ventral Axis
Formation in the Drosophila Embryo. *Current Biology* **13**:2097–2102.
doi:10.1016/j.cub.2003.10.062

Lindsay SA, Lin SJH, Wasserman SA. 2018. Short-Form Bomanins Mediate Humoral
Immunity in Drosophila. *J Innate Immun* **10**:306–314. doi:10.1159/000489831

Marra A, Hanson MA, Kondo S, Erkosar B, Lemaitre B. 2021. Drosophila Antimicrobial
Peptides and Lysozymes Regulate Gut Microbiota Composition and Abundance. *mBio*
e0082421. doi:10.1128/mBio.00824-21

Martins NE, Faria VG, Teixeira L, Magalhães S, Sucena É. 2013. Host Adaptation Is
Contingent upon the Infection Route Taken by Pathogens. *PLoS Pathogens* **9**:e1003601.
doi:10.1371/journal.ppat.1003601

Medzhitov R, Schneider DS, Soares MP. 2012. Disease Tolerance as a Defense Strategy.
Science **335**:936–941. doi:10.1126/science.1214935

Melcarne C. 2020. Role of Nimrod receptors in the Drosophila melanogaster cellular immune
response. EPFL. doi:10.5075/epfl-thesis-9933

Melcarne C., Lemaitre B, Kurant E. 2019. Phagocytosis in Drosophila: From molecules and
cellular machinery to physiology. *Insect Biochemistry and Molecular Biology* **109**:1–12.
doi:10.1016/j.ibmb.2019.04.002

Melcarne C, Ramond E, Dudzic J, Bretscher A, Kurucz E, Andó I, Lemaitre B. 2019. Two
Nimrod receptors, NimC1 and Eater, synergistically contribute to bacterial phagocytosis
in *Drosophila melanogaster*. *The FEBS Journal*. doi:10.1111/febs.14857

Nakano S, Kashio S, Nishimura K, Takeishi A, Kosakamoto H, Obata F, Kuranaga E,
Chihara T, Yamauchi Y, Isobe T, Miura M. 2023. Damage sensing mediated by serine

proteases Hayan and Persephone for Toll pathway activation in apoptosis-deficient flies. *PLoS Genet* **19**:e1010761. doi:10.1371/journal.pgen.1010761

Nakhleh J, El Moussawi L, Osta MA. 2017. The Melanization Response in Insect ImmunityAdvances in Insect Physiology. Elsevier. pp. 83–109. doi:10.1016/bs.aiip.2016.11.002

Nappi A, Poirié M, Carton Y. 2009. The role of melanization and cytotoxic by-products in the cellular immune responses of Drosophila against parasitic wasps. *Adv Parasitol* **70**:99–121. doi:10.1016/S0065-308X(09)70004-1

Needham AJ, Kibart M, Crossley H, Ingham PW, Foster SJ. 2004. Drosophila melanogaster as a model host for Staphylococcus aureus infection. *Microbiology* **150**:2347–2355. doi:10.1099/mic.0.27116-0

Nehme NT, Liégeois S, Kele B, Giammarinaro P, Pradel E, Hoffmann JA, Ewbank JJ, Ferrandon D. 2007. A Model of Bacterial Intestinal Infections in Drosophila melanogaster. *PLoS Pathogens* **3**:e173. doi:10.1371/journal.ppat.0030173

Nehme NT, Quintin J, Cho JH, Lee J, Lafarge M-C, Kocks C, Ferrandon D. 2011. Relative Roles of the Cellular and Humoral Responses in the Drosophila Host Defense against Three Gram-Positive Bacterial Infections. *PLOS ONE* **6**:e14743. doi:10.1371/journal.pone.0014743

Pradeu T, Thomma BPHJ, Girardin SE, Lemaitre B. 2024. The conceptual foundations of innate immunity: Taking stock 30 years later. *Immunity* 613–31.

Quintin J, Asmar J, Matskevich AA, Lafarge M-C, Ferrandon D. 2013. The Drosophila Toll Pathway Controls but Does Not Clear Candida glabrata Infections. *The Journal of Immunology* **190**:2818–2827. doi:10.4049/jimmunol.1201861

Rämet M, Lanot R, Zachary D, Manfruelli P. 2002. JNK Signaling Pathway Is Required for Efficient Wound Healing in Drosophila. *Developmental Biology* **241**:145–156. doi:10.1006/dbio.2001.0502

Ramond E, Jamet A, Ding X, Euphrasie D, Bouvier C, Lallemant L, He X, Arbibe L, Coureuil M, Charbit A. 2021. Reactive Oxygen Species-Dependent Innate Immune Mechanisms Control Methicillin-Resistant Staphylococcus aureus Virulence in the *Drosophila* Larval Model. *mBio* **12**:e00276-21. doi:10.1128/mBio.00276-21

Rommelaere S, Carboni A, Bada Juarez JF, Boquete J-P, Abriata LA, Meireles FTP, Rukes V, Vincent C, Kondo S, Dionne MS, Dal Peraro M, Cao C, Lemaitre B. 2024. A humoral stress response protects *Drosophila* tissues from antimicrobial peptides. *Current Biology* 1–12. doi:10.1101/2023.07.24.550293

Rutschmann S, Kilinc A, Ferrandon D. 2002. Cutting edge: the toll pathway is required for resistance to gram- positive bacterial infections in Drosophila. *J Immunol* **168**:1542–6.

Ryder E, Blows F, Ashburner M, Bautista-Llacer R, Coulson D, Drummond J, Webster J, Gubb D, Gunton N, Johnson G, O'Kane CJ, Huen D, Sharma P, Asztalos Z, Baisch H, Schulze J, Kube M, Kittlaus K, Reuter G, Maroy P, Szidonya J, Rasmuson-Lestander \AAsa, Ekström K, Dickson B, Hugentobler C, Stocker H, Hafen E, Lepesant JA,

Pflugfelder G, Heisenberg M, Mechler B, Serras F, Corominas M, Schneuwly S, Preat T, Roote J, Russell S. 2004. The DrosDel collection: A set of P-element insertions for generating custom chromosomal aberrations in Drosophila melanogaster. *Genetics* **167**:797–813. doi:10.1534/genetics.104.026658

Sansone CL, Cohen J, Yasunaga A, Xu J, Osborn G, Subramanian H, Gold B, Buchon N, Cherry S. 2015. Microbiota-dependent priming of antiviral intestinal immunity in Drosophila. *Cell host & microbe* **18**:571–581.

Sasaki A, Nishimura T, Takano T, Naito S, Yoo SK. 2021. white regulates proliferative homeostasis of intestinal stem cells during ageing in Drosophila. *Nat Metab* **3**:546–557. doi:10.1038/s42255-021-00375-x

Schmid-Hempel P. 2021. Evolutionary parasitology: the integrated study of infections, immunology, ecology, and genetics. Oxford University Press.

Shan T, Wang Y, Bhattarai K, Jiang H. 2023. An evolutionarily conserved serine protease network mediates melanization and Toll activation in *Drosophila*. *Sci Adv* **9**:eadk2756. doi:10.1126/sciadv.adk2756

Stephenson HN, Streeck R, Grüblinger F, Goosmann C, Herzig A. 2022. Hemocytes are essential for *Drosophila melanogaster* post-embryonic development, independent of control of the microbiota. *Development* **149**:dev200286. doi:10.1242/dev.200286

Touré H, Durand N, Guénal I, Herrmann J-L, Girard-Misguich F, Szuplewski S. 2023. Mycobacterium abscessus Opsonization Allows an Escape from the Defensin Bactericidal Action in *Drosophila*. *Microbiol Spectr* e00777-23. doi:10.1128/spectrum.00777-23

Ulvila J, Vanha-Aho L-M, Rämet M. 2011. Drosophila phagocytosis - still many unknowns under the surface. *APMIS* **119**:651–662. doi:10.1111/j.1600-0463.2011.02792.x

Unckless RL, Howick VM, Lazzaro BP. 2016. Convergent Balancing Selection on an Antimicrobial Peptide in Drosophila. *Current Biology* **26**:257–262. doi:10.1016/j.cub.2015.11.063

Uttenweiler-Joseph S, Moniatte M, Lagueux M, Van Dorsselaer A, Hoffmann JA, Bulet P. 1998. Differential display of peptides induced during the immune response of Drosophila: a matrix-assisted laser desorption ionization time-of- flight mass spectrometry study. *Proc Natl Acad Sci U S A* **95**:11342–7.

Vaz F, Kounatidis I, Covas G, Parton RM, Harkiolaki M, Davis I, Filipe SR, Ligoxygakis P. 2019. Accessibility to Peptidoglycan Is Important for the Recognition of Gram-Positive Bacteria in Drosophila. *Cell Reports* **27**:2480-2492.e6. doi:10.1016/j.celrep.2019.04.103

Westlake H, Hanson MA, Lemaitre B. 2024. The Drosophila immunity handbook, First. ed. EPFL Press. doi:10.55430/6304TDIHVA01

Wukitch AM, Lawrence MM, Satriale FP, Patel A, Ginder GM, Van Beek EJ, Gilani O, Chambers MC. 2023. Impact of Chronic Infection on Resistance and Tolerance to Secondary Infection in Drosophila melanogaster. *Infect Immun* e00360-22. doi:10.1128/iai.00360-22

Xu R, Lou Y, Tidu A, Bulet P, Heinekamp T, Martin F, Brakhage A, Li Z, Liégeois S,
Ferrandon D. 2023. The Toll pathway mediates Drosophila resilience to Aspergillus
mycotoxins through specific Bomanins. *EMBO Rep* **24**:e56036.
doi:10.15252/embr.202256036

Yasin B, Wang W, Pang M, Cheshenko N, Hong T, Waring AJ, Herold BC, Wagar EA,
Lehrer RI. 2004. θ Defensins Protect Cells from Infection by Herpes Simplex Virus by
Inhibiting Viral Adhesion and Entry. *Journal of Virology* **78**:5147–5156.
doi:10.1128/jvi.78.10.5147-5156.2004

1056

1057

1058　**Figures**

1059　**Table 1: List of Mutants used in this study.** "-" indicates the deleted immune module.

Drosophila lines	Reference	Name	Deficient pathway			
			IMD	Toll	Phagocytosis	Melanization
iso ; iso ; Rel^{E20}	Hedengren et al., 1999	*ΔIMD*	-	+	+	+
iso ; iso ; spz^{rm7}	Lemaitre et al., 1996	*ΔTOLL*	+	-	+	+
iso ; NimC1^1 ; Eater^1	Melcarne et al., 2019	*ΔPhag*	+	+	-	+
iso ; PPO1Δ, PPO2Δ; iso	Bingelli et al., 2014	*ΔMel*	+	+	+	-
+ ; + ; Rel^{E20}, spz^{rm7}	this paper	*ΔIMD, ΔTOLL*	-	-	+	+
iso ; PPO1Δ, PPO2Δ ; NimC1^1,Eater^1	this paper	*ΔPhag, ΔMel*	+	+	-	-
iso ; NimC1^1 ; Eater^1, Rel^{E20}	this paper	*ΔIMD, ΔPhag*	-	+	-	+
iso ; PPO1Δ PPO2Δ ; Rel^{E20}	this paper	*ΔIMD, ΔMel*	-	+	+	-
iso ; PPO1Δ PPO2Δ ; spz^{rm7}	this paper	*ΔTOLL, ΔMel^{PPS}*	+	-	+	-
iso Hay-psh^{Def} ; iso ; iso	Dudzic et al., 2019	*ΔTOLL, ΔMel^{HP}*	+	-	+	-
iso Hay-psh^{Def} ; NimC1^1 ; Eater^1 Rel^{E20}	this paper	*ΔITPM*	-	-	-	-

1060

1061

1062
Figure 1: Each immune module functions largely independently from other immune
modules. **A)** Activation of the Imd pathway, as revealed by *DptA* expression following
infection with a mixture of heat-killed *E. coli* and *M. luteus,* is broadly wild-type in single
module mutants other than Imd. In the case of *ΔPhag* flies, we observed a lesser induction at
6h compared to wild-type. **B)** Activation of the Toll pathway, as revealed by the expression of
Drs is broadly wild-type in single module mutants other than *ΔTOLL* flies. In the case of *ΔMel*,

we observed a slightly greater Toll pathway activity. **C)** The ability of plasmatocytes to phagocytose bacterial particles is not negatively affected in single module mutants except *ΔPhag* (also see Figure 1supp1) **D)** Cuticle blackening after clean injury is not impaired except in *ΔMel* flies. **E)** Activation of the Imd pathway remains strongly inducible in compound mutants except when deficient for the Imd pathway. **F)** Activation of the Toll pathway remains strongly inducible in compound mutants except when flies were deficient of the Toll pathway. Note that *Drs* receives a minor input from Imd signaling (Leulier et al., 2000), explaining minor induction of *Drs* in *ΔToll* flies at early time points. **G)** The ability of plasmatocytes to phagocytose bacterial particles is not significantly affected in compound mutants except those including *ΔPhag*. **H)** Cuticle blackening after clean injury is not impaired in compound mutants except in *ΔMel* flies.

Figure 2: Lifespans of mutants used in this study in unchallenged flies (A-B) and upon clean injury (C-D) conditions at 25°C. See Figure 2supp1 for 29°C comparisons.

Figure 3: Heatmap of lifespans of immune module-deficient flies upon infection by various pathogens. Darker blue indicates lower survival, while white indicates maximum survival. Experiments used either 7 or 10 days as a maximum lifespan/time course, and the heatmap is adjusted accordingly per row to have white fill for the maximum possible lifespan of that row. Heatmap colours: blue (low survival), white (high survival). Survival curves are presented in **Supplementary file 2**, and a summary of susceptibilities is provided in Figure 3supp1. Small numbers beside mean lifespans indicate total sample size per genotype per treatment.

1093

Figure 4: Growth kinetics of *Pr. rettgeri* (A), *S. aureus* (B), and *C. albicans* (C) in wild-type, single module mutant and *ΔITPM* flies. Survival curves underlying data in Figure 3 are shown for comparison. Each data point reflects a pooled sample of 5 flies. Error bars reflect 1 standard deviation from the mean.

1099

Figure 5: Measurement of PLUD upon infection with *Pr. rettgeri*, *S. aureus*, and *C. albicans*. A) The PLUD of *Pr. rettgeri* in individual module mutant flies is not significantly different from wild-type. The PLUD of *ΔITPM* flies was not significantly different, although censoring of a single low-PLUD outlier in the wild-type would result in a significant difference between wild-type and *ΔITPM* flies ($p < .01$). **B)** The PLUD of *S. aureus*-infected flies is not different across any genotype. **C)** The PLUD of *C. albicans*-infected flies was significantly lower in *ΔPhag* flies with a notably lower mean PLUD. This difference was robust to use of different *ΔPhag* genetic backgrounds (merged data shown here). *ΔMel* flies and *ΔITPM* flies also had significantly lower PLUD. Error bars reflect 1 standard deviation from the mean.

1109

1110

Figure 1supp1: We used *ΔPhag* mutants in the genetic background of Melcarne et al. (2019) *(+; ΔPhag)* in some experiments, and confirm here that these mutants are equally deficient in phagocytic capacity to *ΔPhag* flies from the iso DrosDel genetic background *(ΔPhag)*.

1114

Figure 2supp1: Lifespans of mutants used in this study in unchallenged flies and upon clean injury conditions at 29°C.

Figure 3supp1: Summary table of susceptibilities per the three conditions outlined in section "Systemic infections and survival." A ✓ indicates the pathway contributes to defence against a given pathogen. In the event that a single module mutant is not more susceptible to the infection, but contributes to increased mortality in a double mutant line, a ✓ is given and the co-deleted relevant pathways highlighted by their first letter. Black boxes indicate no difference to wild-type or to clean injury.

Figure 4supp1: Survival to infection of females against *P. rettgeri* **(A)**, *S. aureus* **(B)**, and *C.*
albicans **(C)**. Survival trends largely parallel the rank order of males (**Figure 4**).

Table S1: Microbe characteristics and growth conditions.

Supplementary file 1: Supplemental discussion of fly genetics.

Supplementary file 2: Survival curves for all data summarized in Figure 3.

Supplementary file 3: Supplemental discussion of fly genetics and considerations for the use
of genotypes described in this study.

Type	Species	Temp	Dose	Days	N exp	WT	ΔAMP14	ΔBom	ΔIMD	ΔTOLL	ΔPhag	ΔMel	ΔPhag, ΔMel	ΔTOLL, ΔMel^nrs	ΔTOLL, ΔMel^spz	ΔIMD, ΔMel	ΔIMD, ΔPhag	ΔIMD, ΔTOLL	ΔITPM
Cuticle	Unchallenged	25°C	n/a	7	13	6.9		6.9		6.8		6.6	7.0		5.3	6.0			5.5
	Clean Injury	25°C	n/a	7	36	6.7		6.6		6.7	6.6	6.8		5.3	4.6	5.4	6.5	6.5	6.2	5.0
	Clean Injury	29°C	n/a	7	11	6.6		6.1		6.5		6.5	6.6		4.9	5.0				3.9
	Clean Injury	25°C	n/a	10	36	9.6		9.2		9.2	9.3	9.6		8.5	6.2	5.2	9.1	8.7	6.3	5.0
	Unchallenged	29°C	n/a	10	11	9.9		9.3		9.5		8.9	9.1		8.7				7.7
Viruses	Drosophila C virus	25°C	2000 TCID90	10	5	8.3		6.7		7.9	7.7	6.8	6.1	6.8	6.8	5.1	5.7	0.4	3.8	4.1
	Flock House virus	25°C	2500000 TCID90	10	5	9.5		7.1		7.7	7.2	7.3	7.3	8.1	8.0	6.2	6.1	5.4	6.5	5.1
	Drosophila X virus	25°C	431000 TCID50	10	4	6.9		8.2		7.1	7.9	7.4	7.8	5.4	7.4	4.9	5.1	6.0	6.2	5.6
	Invertebrate iridescent 6	28°C	31700 TCID90	10	3	7.5		9.4		8.1	5.8	8.4	8.8	5.6	5.8	5.9	4.8	8.3	5.9	4.4
	Sindbis virus	25°C	50000 TCID50	10	3	9.4		8.4		8.6	8.9	8.6	8.9	5.4	4.8	4.5	4.5	4.5	4.3
Fungi	Aspergillus fumigatus (natural)	25°C	not on plate	5	10	10.0		9.2		9.9	9.3	6.7	7.6	4.1	9.1	8.2	7.0	5.6	2.7
	Beauveria bassiana (natural)	29°C	30mg	10	6	5.9		6.9		9.4	7.3	7.8	4.6	7.7	5.0	6.9	6.4	5.0	4.4
	Beauveria bassiana	25°C	OD=5	7	6	6.4		6.0		3.6	5.1	5.7	3.6	5.2	4.4	2.6	4.7	3.1	3.3
	Candida albicans	29°C	OD=5	7	8	6.7		5.6		5.0	5.1	5.3	5.9	5.0	5.6	1.7	5.7	5.9	3.2	2.4
Gram positive bacteria	Mycobacterium marinum	25°C	OD=0.1	7	8	6.6		6.5		5.8	6.5	6.1	5.9	5.7	5.8	4.1	5.1	5.1	5.4	3.9
	Corynebacterium diphtheriae	25°C	OD=200	7	3	6.0		6.2		6.5	5.6	5.8	6.6	4.8	3.6	3.3	4.6	3.4	5.3	3.8
	Micrococcus luteus	25°C	OD=200	7	3	5.7		6.8		5.7	6.4	5.9	6.3	5.2	5.9	4.8	6.3	2.1	1.9
	Streptococcus pneumoniae	25°C	OD=5	7	8	6.1		6.4		5.9	6.0	6.5	6.5	5.1	5.1	6.0	5.5	4.5	1.8
	Enterococcus faecalis	25°C	OD=10	7	10	5.0		4.5		4.2	3.2	1.8	5.2	5.1	3.9	1.8	2.2	1.3
	Staphylococcus aureus	25°C	OD=0.5	7	8	5.1		4.8		4.2	3.4	3.6	2.4	1.9	2.0	4.1	3.4	1.9
	Listeria monocytogenes	25°C	OD=200	7	8	5.5		5.2		4.2	3.3	3.9	3.1	2.1	2.3	3.3	3.3	2.0	1.4
	Bacillus subtilis	25°C	OD=200	7	7	5.7		5.3		4.1	4.0	4.5	4.5	3.8	2.8	2.7	4.8	2.0
Gram negative bacteria	Vibrio parahaemolyticus	25°C	OD=10	7	2	3.1		1.1		2.6	1.3	2.8	2.3	3.2	1.8	1.0	1.1	1.0	1.0
	Providencia rettgeri	25°C	OD=0.1	7	3	2.0		4.4		5.1	3.9	3.9	4.3	2.9	4.9	2.1	1.4	1.3	1.4
	Providencia burhodogranariea	25°C	OD=10	7	7	5.1		1.7		4.6	4.1	4.9	4.0	5.4	3.9	1.3	1.0	1.4	1.2
	Pectobacterium carotovorum	25°C	OD=10	7	4	1.5		5.6		4.9	1.8	4.6	4.6	1.2	1.8	1.4	1.4	1.6
	Klebsiella pneumoniae	25°C	OD=200	7	4	6.7		1.0		6.0	1.1	6.0	5.7	6.9	6.0	1.4	4.9	5.8	1.8
	Enterobacter cloacae	25°C	OD=200	7	4	6.6		1.0		6.6	1.1	6.7	6.4	4.1	4.0	5.7	1.5
	Escherichia coli	25°C	OD=200	7	4	5.0		6.9		6.8	3.8	6.2	7.0	3.3	3.8	4.6	1.6	1.9
	Salmonella enterica ser. typhimurium	25°C	OD=200	7	4	6.6		6.0		4.6	5.2	5.3	5.3	6.0	4.3	4.8	4.3	1.8

Drosophila lines	Reference	Name	Deficient pathway			
			IMD	Toll	Phagocytosis	Melanization
iso ; iso ; Rel^{E20}	Hedengren et al., 1999	ΔIMD	-	+	+	+
iso ; iso ; spz^{rm7}	Lemaitre et al., 1996	ΔTOLL	+	-	+	+
iso ; NimC1^{1} ; Eater^{1}	Melcarne et al., 2019	ΔPhag	+	+	-	+
iso ; PPO1Δ, PPO2Δ; iso	Bingelli et al., 2014	ΔMel	+	+	+	-
+ ; + ; Rel^{E20}, spz^{rm7}	this paper	ΔIMD, ΔTOLL	-	-	+	+
iso ; PPO1Δ, PPO2Δ ; NimC1^{1},Eater^{1}	this paper	ΔPhag, ΔMel	+	+	-	-
iso ; NimC1^{1} ; Eater^{1}, Rel^{E20}	this paper	ΔIMD, ΔPhag	-	+	-	+
iso ; PPO1Δ PPO2Δ ; Rel^{E20}	this paper	ΔIMD, ΔMel	-	+	+	-
iso ; PPO1Δ PPO2Δ ; spz^{rm7}	this paper	ΔTOLL, ΔMel^{PPS}	+	-	+	-
iso Hay-psh^{Def} ; iso ; iso	Dudzic et al., 2019	ΔTOLL, ΔMel^{HP}	+	-	+	-
iso Hay-psh^{Def} ; NimC1^{1} ; Eater^{1} Rel^{E20}	this paper	ΔITPM	-	-	-	-

P. rettgeri

S. aureus

C. albicans