Layers of immunity: Deconstructing the *Drosophila* effector response

Ryckebusch F[1], Yao Tian[1], Rapin M[1], Schüpfer F[1], Hanson MA[1,2]*, Lemaitre B[1]*

1. Global Health Institute, Ecole Polytechnique Fédérale de Lausanne (EPFL), Lausanne, Switzerland
2. Centre for Ecology and Conservation, University of Exeter, Penryn, United Kingdom
   * *These authors contributed equally to this work*

Corresponding authors:
Bruno Lemaitre (lead): bruno.lemaitre@epfl.ch
Hanson, Mark: M.Hanson@exeter.ac.uk

Abstract

The host innate immune response relies on the cooperation of multiple defense modules. In insects and other arthropods, which have only innate immune mechanisms, four main immune-specific modules contribute to defense against microbial invaders: the Toll pathway, the Imd pathway, the melanization response, and phagocytosis by plasmatocytes. Our present understanding of their relative importance remains fragmented, as their contribution to host defense has never been simultaneously assessed across a large panel of pathogens. Here, we have taken advantage of newly-described immune mutants in a controlled genetic background to systematically delete these four immune modules individually, in pairs, or all four simultaneously. Surprisingly, flies simultaneously deficient in all four immune modules are viable, homozygous fertile, and display no overt morphological defects, suggesting these immune mechanisms are not strictly required for organismal development. With this new set of lines, we assessed the individual and collective contribution of each module to host defense against five viruses, three fungi, eight Gram-positive bacteria, and eight Gram-negative bacteria. Our findings show that these four modules largely function independently or additively in host defense, although synergistic effects can occur for select pairs of modules. Our study confirmed the importance of the Imd pathway against Gram-negative bacteria and the Toll pathway against Gram-positive bacteria and fungi, largely via the induction of effectors such as antimicrobial peptides (AMPs) and Bomanins (Bom), but also reveals an important role of melanization against viruses, and a contribution of phagocytosis against various germs. Additionally, by examining microbial load kinetics in different mutants, we provide insight into how these modules contribute to tolerance or resistance against specific microbes. Our study provides insights into the architecture of the *Drosophila* immune system, revealing differential requirements of immune modules according to each pathogen. The set of immune deficient lines provided here offers tools to better assess the role of these immune modules in host defense.

Keywords: Drosophila, innate immunity, effectors, phagocytosis, melanization, Imd, Toll

Introduction

The immune system is made up of a network of tissues and circulating cells whose primary function is to prevent and limit pathogenic infections (Danilova, 2006; Pradeu et al., 2024; Schmid-Hempel, 2021). Although immunity has been studied in many animal species, it has been extensively characterized in only a few, such as the nematode *Caenorhabditis elegans,* the fruit fly *Drosophila melanogaster*, zebrafish, rodents, and humans. In recent years, this traditional understanding of the immune system as a specialized group of cells and tissues has gradually changed to a more complex picture of highly redundant multilevel defense modules. The last few decades have revealed a great deal about the molecular mechanisms regulating each of these immune modules, from recognition and signaling to effector production (*e.g.* phagocytosis, antimicrobial peptides) (Pradeu et al., 2024). However, an immune response encompasses the concomitant activation of several modules and there has been little investigation of how modules contribute collectively to host defense. Immune research has traditionally focused on modules in isolation due to technical limitations. This article aims to fill this gap by analyzing the specific contributions of the main immune modules to the collective *Drosophila* systemic response.

The *Drosophila* systemic immune response consists of a set of humoral and cellular reactions in the hemolymph (insect blood) that occur when flies are systemically infected by bacteria, viruses, or fungi (Liegeois and Ferrandon, 2022; Westlake et al., 2024). A first facet of the systemic immune response is the production of host-defense effectors by the fat body and hemocytes that are secreted into the hemolymph. Among the best-characterized effectors are the antimicrobial peptides (AMPs), whose expression is rapidly induced to incredibly high levels upon infection in a race to control invading microbes (Hanson et al., 2019; Imler and Bulet, 2005). AMPs are small cationic peptides that exhibit antimicrobial activity and directly participate in microbial killing. Beyond AMPs, many other proteins and host defense peptides are secreted into the hemolymph, creating a hostile environment for invading pathogens through a variety of mechanisms. The systemic antimicrobial response is predominantly regulated at the transcriptional level by the Toll and Imd pathways (De Gregorio et al., 2002; Lemaitre et al., 1996). In *Drosophila*, the Toll pathway is activated by cell wall components (fungal glucans and Lysine-type peptidoglycan), as well as microbial proteases (Gottar et al., 2006; Issa et al., 2018; Leulier et al., 2003; Vaz et al., 2019). Toll signaling culminates in the activation of two NF-κB transcription factors, Dif and Dorsal, which regulate a large set of immune genes (Ip, 1993; Lemaitre et al., 1996). These encode small peptides such as the antifungal peptide Drosomycin, the Toll-regulated Bomanin peptides, and many other proteins (*e.g.* serine proteases, serpins, lipases) (Clemmons et al., 2015; De Gregorio et al., 2002; Fehlbaum et al., 1994; Ligoxygakis et al., 2003). Flies lacking the Toll pathway are viable but highly susceptible to infection by Gram-positive bacteria and fungi (Buchon et al., 2009; Lemaitre et al., 1996; Rutschmann et al., 2002). The Imd pathway is activated by DAP-type peptidoglycan produced by Gram-negative bacteria and a subset of Gram-positive bacteria (*e.g. Bacillus*) (Kaneko et al., 2004; Lemaitre et al., 1995; Leulier et al., 2003). The binding of peptidoglycan to receptors

of the Peptidoglycan Recognition Protein family (PGRP-SD, PGRP-LC, PGRP-LE) initiates an intracellular signaling cascade, which ultimately activates the NF-κB factor Relish (Choe et al., 2005; Iatsenko et al., 2016; Kaneko et al., 2006; Kaneko et al., 2006; Westlake et al., 2024). The Imd pathway regulates the expression of many genes, including those encoding antibacterial peptides and serine proteases (De Gregorio et al., 2002; Lemaitre et al., 1997). Imd deficient flies are viable but susceptible to Gram-negative bacterial infection.

In addition to humoral pathways, melanization and phagocytosis are two complementary mechanisms that contribute to host survival upon systemic infection. Melanization is an arthropod-specific immune response resulting in rapid deposition of the black pigment melanin at wound or infection sites and the concomitant production of microbicidal reactive oxygen species (Cerenius et al., 2008; Nappi et al., 2009; Westlake et al., 2024). The melanization and Toll pathways are co-activated at the level of the serine proteases Hayan and Persephone, which in turn activate Spatzle-processing enzyme (SPE) and serine proteases critical to cleavage of prophenoloxidases (PPOs) (Dudzic et al., 2019; Nakano et al., 2023; Shan et al., 2023). This is collectively referred to as the Toll-PO SP cascade (Westlake et al., 2024). The melanization reaction itself relies on the oxidation of phenols, resulting in the polymerization of melanin. Two PPOs (PPO1 and PPO2) are the primary catalysts for melanin synthesis during the melanization response to systemic infection. Flies deficient for both *PPO1* and *PPO2* are viable, but lack hemolymph phenoloxidase (PO) activity and exhibit susceptibility to certain Gram-positive bacteria and fungi (Binggeli et al., 2014; Dudzic et al., 2015).

Phagocytosis by both sessile and circulating plasmatocytes (*Drosophila* macrophage-equivalent cells) is thought to provide a complementary and important host defense (Melcarne et al., 2019; Ulvila et al., 2011). The use of hemocyte-deficient flies (via targeted expression of a pro-apoptotic gene in plasmatocytes) has shown that hemocytes contribute to survival upon systemic infection to certain bacterial species (Charroux and Royet, 2009; Defaye et al., 2009; Stephenson et al., 2022). Phagocytosis of bacteria greatly relies on two transmembrane receptors, NimC1 and Eater (Bretscher et al., 2015; Kocks et al., 2005; Kurucz et al., 2007; Melcarne et al., 2019). *NimC1, Eater* double mutant larvae are viable and their hemocytes can effectively encapsulate and melanize macroparasites; however, their hemocytes are non-sessile and nearly totally phagocytosis deficient. While *NimC1, Eater* larvae have elevated hemocyte numbers, adult flies have few hemocytes and fail to perform phagocytosis, providing a good tool to assess the role of cellular response (Melcarne et al., 2019; Melcarne, 2020).

We have scattered circumstantial evidence that each of these four immune modules are critical for defense against certain pathogens. However, we lack a global view of how, when, and to what extent each of these four immune modules contribute to host defense against specific pathogens, as they have never been simultaneously assessed. Moreover, the extent to which individual findings can be generalized to broad classes of pathogens remains unclear.

In this article, we have used a specific set of mutations principally affecting phagocytosis *(NimC1, Eater),* melanization *(PPO1, PPO2),* the Toll pathway *(spz),* and the Imd pathway *(Rel)* in a controlled genetic background to address their individual roles in host defense. We then recombined these and other mutations to generate double and even quadruple mutants of the four immune modules. This set of tools allowed us to systematically compare the survival of flies deficient in these responses against a broad panel of viruses, fungi, Gram-

130 positive bacteria, and Gram-negative bacteria, to reveal the relative contributions of each
131 immune mechanism to resistance against these pathogens.
132

Materials and methods
133

Insect stocks
134

Precise details of *Drosophila* stocks used in this study are provided in **Table 1**. To minimize
the influence of genetic background as a variable in this study, mutations were isogenized into
the *DrosDel iso w^1118^* genetic background (BDSC #5905) as described by Ferreira et al.
(Ferreira et al., 2014; Ryder et al., 2004). Isogenized lines were generated and used for every
mutant except the double mutant Toll and Imd *(Rel^E20^, spz^rm7^)* line, for which homozygous
isogenized flies were not viable in our hands. Double mutants for *NimC1^1^; Eater^1^* were poorly
viable when isogenized in the *DrosDel iso w^1118^* background, and so we used both isogenic and
non-isogenic lines over the course of our study to facilitate investigation. Furthermore, *NimC1^1^;*
*spz^rm7^, Eater^1^* triple mutants were homozygous lethal regardless of isogenization, and so could
not be included. In addition to the immune module mutants, the Δ*AMP14* line bears a knockout
of 14 different AMP genes, which includes the AMP families Drosomycin, Metchnikowin,
Cecropin, Defensin, Drosocin, Attacin, and Diptericin, as described previously (Carboni et al.,
2022; Hanson et al., 2019). The *Bom^Δ55C^* mutation was characterized in Clemmons et al.
(Clemmons et al., 2015) and isogenic flies were generated in Hanson et al. (Hanson et al., 2019).
For larval experiments, GFP-labelled CyO or TM3 balancers were used to enable selection of
homozygous mutant larvae.
151

Microorganism cultures
152

Microbe strain information, microorganism classifications, and culture conditions are listed in
Table S1. Bacterial cultures and *Candida albicans* were grown overnight shaking in
appropriate media at 200rpm. *Aspergillus fumigatus* fungus was grown on Malt Agar at room
temperature until sporulation. The entomopathogenic fungus *Beauveria bassiana* strain R444
was provided by Andermatt as spore preparations (BB-PROTEC) which were directly used in
natural infections or dissolved in PBS for septic injury. Viruses DCV and FHV were kindly
provided by Carla Saleh (Pasteur Institute, Paris), produced in S2 cells. The supernatant of S2
cells was titrated before being used for fly infections. Viruses SINV, DXV and IIV-6 were
kindly provided by Ronald van Rij (Radboud University Medical Center, Nijmegen).
162

163 Heat-killed microbes were prepared by two repeats of boiling at 95°C for 30 minutes then
164 freezing at -20°C for 30 minutes, before storage long term at -20°C. Microbe preparations were
165 streaked onto agar plates to verify full efficiency of heat-killing before use in experiments.
166

Gene expression
167

Flies were inoculated by pricking at the junction of thoracic pleura with a needle dipped in a
mixed pellet containing a 1:1 mixture of OD_{600} = 200 heat-killed *Escherichia coli* and
Micrococcus luteus (final OD_{600} = 100 for each), and frozen at -20°C 6h, 12h and 24h post-
infection. Total RNA was then extracted from pooled samples of five flies using TRIzol reagent
per manufacturer's protocol and resuspended in MilliQ dH2O. Reverse transcription was

performed using PrimeScript RT (Takara) with random hexamer and oligo dT primers.
Quantitative PCR was performed on a LightCycler 480 (Roche) in 96-well plates using Applied
Biosystems PowerUP SYBR green Master Mix (Applied Biosystems). Data points represent
pooled samples from three replicate experiments. Error bars represent one standard deviation
from the mean. Statistical analyses were performed using one-way ANOVA with Holms-Sidak
multiple test correction. Primers used in this study were:
179
Drs-F 5'- CGTGAGAACCTTTTCCAATATGAT-3'
Drs-R 5'- TCCCAGGACCACCAGCAT-3'
DptA-F: 5'-GCTGCGCAATCGCTTCTACT-3'
DptA-R: 5'-TGGTGGAGTGGGCTTCATG-3'
RPL32-F: 5'-GACGCTTCAAGGGACAGTATCTG-3'
RPL32-R: 5'-AAACGCGGTTCTGCATGAG-3'
186

Ex vivo larval hemocyte phagocytosis assays
Ex vivo phagocytosis assays were performed using a mix of equal volumes of *E. coli* and
Staphylococcus aureus AlexaFluorTM488 BioParticlesTM (Invitrogen), following
manufacturer's instructions; and see (Melcarne et al., 2019). Five L3 wandering larvae from
our newly generated mutant lines, or carrying the Hml-Gal4, UAS-GFP hemocyte marker as a
control, were bled into 150 uL of Schneider's insect medium (Sigma-Aldrich) containing 1 uM
phenylthiourea (PTU; Sigma-Aldrich). The hemocyte suspension was then transferred to 1.5
mL low binding tubes (Eppendorf, SigmaAldrich), and AlexaFluorTM488 bacteria
BioParticlesTM were added. The samples were vortexed, incubated at room temperature for 2
hours to allow phagocytosis and then placed on ice to stop the reaction. The fluorescence of
extracellular particles was quenched by adding 0.4% trypan blue (Sigma-Aldrich) diluted to 1/3
concentration. Phagocytosis was quantified using a flow cytometer (BD Accuri C6) to measure
the fraction of cells phagocytosing, and their fluorescence intensity. Isogenic wild-type iso w^{1118}
larvae and Hml-Gal4, UAS-GFP larvae with or without bacterial particles were used to define
the gates for hemocyte counting and the thresholds for phagocytosed particle emission. The
phagocytic index was calculated as follows:

$$\text{Fraction of haemocytes phagocytosing}(f) = \frac{[\text{number of haemocytes in fluorescence positive gate}]}{[\text{total number of haemocytes}]}$$

$$\text{Phagocytic index}(PI) = [\text{Mean fluorescence intensity of haemocytes in fluorescence positive gate}] \times f$$

203
204   Finally, due to experimenter differences in total hemocytes collected and knock-on effects to
205   calculating phagocytic index, phagocytic index was normalized with the wild-type as a
206   reference set to 100% within experiment batch.
207

Wounding experiment
Clean injury was performed with a needle sterilized with an EtOH and PBS wash. For
imaging of the melanization reaction upon pricking, the thorax of 3-8 days old flies was
pricked using a sterile needle (diameter ~0.1mm). Pictures were taken 24h post-pricking and

categorized into normal, weak, or no melanization blackening seen at the injury site per
Dudzic et al. (Dudzic et al., 2019).
214

Systemic infections and survival
Systemic infections were performed by pricking 3- to 5-day-old adult males in the thorax with
a 0.1mm-thick needle dipped into a concentrated pellet of bacteria, yeast, or fungal spores. For
natural infections with *A. fumigatus,* flies were anesthetized and then shaken on a sporulating
plate of fungi for 30 seconds. For *B. bassiana*, flies were shaken for 30s in a tube containing an
excess of commercial spores sufficient to coat all flies uniformly, of which the flies later remove
the vast majority through grooming. At least two replicate survival experiments were performed
for each infection, though in specific cases some genotypes may have been included in only
one or no experiments. 20-35 flies were included per vial when possible, and maintained on
cornmeal fly media containing (per 1L): 6.2g agar, 58.8g cornmeal, 58.8g yeast, 60mL grape
fruit juice, 4.83mL propionic acid, and 26.5mL 100g/L moldex in pure ethanol. Survivals were
scored daily, and flies were flipped to new food vials 3 times per week. Due to challenges
working with genotypes of different viability, we could not include all genotypes in all
experimental replicates. Moreover, comparisons across genotypes required multiple levels of
consideration. Given the complexity of genotype-vs-reference-by-pathogen comparisons in our
study, needing to compare infections both to clean injury within genotype, and to wild-type
(WT, *DrosDel iso w^{1118}*) or *ΔITPM* (flies lacking all four immune modules) across treatments,
we chose to highlight survival differences only when they were: **i)** consistent across
experimental replicates; **ii)** of a consistent logic across comparable genotypes; for instance,
compound mutants containing *ΔMel* (*e.g. ΔIMD, ΔMel*) should be as or more susceptible to
infections as *ΔMel* alone if melanization is truly critical for defense; and **iii)** of a mean lifespan
difference ≥1.0 days after accounting for comparisons with unchallenged or clean-injury data.
Total experiments (N exp) and total flies per genotype are reported to provide summary
statistics per pathogen-by-genotype interaction. The mean lifespan reported in survival
summary data has a maximum of 7 or 10 days depending on experimental conditions. Kaplan-
Meier survival curves are provided in the supplementary information.
241

Quantification of microbial load for growth kinetics
3-8 days old flies were infected with the indicated microbe and concentration per OD_{600} as
described in **Table S1**, and allowed to recover. At the indicated time post-infection, flies were
anesthetized using CO_2 and surface sterilized by washing them in 70% ethanol. Ethanol was
removed, and then flies were homogenized using a Precellys bead beater at 6500 rpm for 30 s
in LB broth (*Providencia rettgeri*), BHI (*S. aureus*), or YPG (*C. albicans*) with 500 μL as pools
of 5 flies. These homogenates were serially diluted and 100uL was plated on LB, BHI or YPG
agar. Plates were incubated overnight, and colony-forming units (CFUs) were counted using a
Interscience Scan 500 plate scanning colony counter, and validated independently by manual
counts for a subset of plates to ensure accuracy. Statistical differences were tested using Brown-
Forsythe and Welch ANOVA tests with Dunnett's multiple test correction, which consider
unequal variance.
254

Results

A set of immune-deficient lines in the iso Drosdel background

Mutations affecting only one of the four main modules of the *Drosophila* systemic immune response have been described. These include *Relish*^{E20} (*Rel*^{E20}, deficient for the Imd pathway, referred to here as *ΔIMD*), *spatzle*^{rm7} (*spz*^{rm7}, deficient for the Toll pathway, referred to here as *ΔTOLL*), *PPO1*^{Δ}, *PPO2*^{Δ} (no melanization, referred to here as *ΔMel*), and *NimC1*^{1}; *eater*^{1} (reduced phagocytosis, referred to here as *ΔPhag*). In addition to blocking phagocytosis, *ΔPhag* flies have a defect in hemocyte sessility and a decreased number of hemocytes at the adult stage (Bretscher et al., 2015; Melcarne et al., 2019; Melcarne, 2020). We have preferred the use of *NimC1*^{1}; *eater*^{1} to study the contribution of hemocytes to defense rather than other approaches with important indirect effects such as the pre-injection of latex beads to saturate phagocytes (Elrod-Erickson et al., 2000), or expression of pro-apoptotic genes in plasmatocytes (Charroux and Royet, 2009; Defaye et al., 2009).

All the mutations in this study were isogenized for at least seven generations into the DrosDel *iso w*^{1118} genetic background (Ferreira et al., 2014; Ryder et al., 2004). We then generated by recombination six of the seven possible combinations of flies simultaneously lacking two immune modules, with the exception of *ΔTOLL, ΔPhag* as *NimC1*^{1}; *eater*^{1}, *spz*^{rm7} flies were not viable (see fly strains in **Table 1**). Of note, in our hands recombinant *Rel*^{E20}, *spz*^{rm7} flies were not viable in the DrosDel isogenic background, necessitating use of non-iso flies to assess host defense in *ΔIMD, ΔTOLL* flies. Over the course of our study, isogenic *NimC1*^{1};*eater*^{1} flies also displayed viability issues, and specific experiments used non-isogenic flies to improve confidence in phenotypes; these experiments are noted in figure captions. We used two approaches to simultaneously delete the Toll and melanization modules: **i)** *PPO1*^{Δ}, *PPO2*^{Δ}; *spz*^{rm7} flies (referred to as *ΔTOLL, ΔMel*^{PPS}) and **ii)** flies carrying a small viable deletion *Hayan-psh*^{Def} (referred to as *ΔTOLL, ΔMel*^{HP}) which removes two clustered serine protease genes, *Hayan,* and *persephone,* blocking the TOLL-PO SP cascade that regulates both humoral-Toll and melanization (Dudzic et al., 2019; Westlake et al., 2024). This effectively deletes the Toll and melanization pathways using only one deletion event on the *X* chromosome. Excitingly, we succeeded in generating a fly line deficient for all four immune modules with the genotype: *Hayan-psh*^{Def}; *NimC1*^{1}; *eater*^{1}, *Rel*^{E20} hereafter named *ΔITPM (ΔIMD, ΔTOLL, ΔPhag, ΔMel)*. These flies were homozygous viable (very poor viability), and fertile, despite major deficiencies in these four important systemic immune mechanisms. These flies allow us to monitor survival kinetics and pathogen growth in the near-total absence of an immune system. Finally, we added two iso fly lines in our analysis that delete key host defense peptides downstream of the Toll and Imd pathways: **i)** *Bom*^{Δ55C}, a short deletion removing ten Bomanins at position 55C, which causes a major susceptibility to infection by Gram-positive bacteria and fungi (Clemmons et al., 2015), and **ii)** *ΔAMP14*, a compound fly line of eight mutations removing 14 AMP genes (4 Cecropins, 4 Attacins, 2 Diptericins, Drosocin, Defensin, Drosomycin, and Metchnikowin), which causes a major susceptibility to infection by Gram-negative bacteria (Carboni et al., 2022; Hanson et al., 2019). This set of immune-deficient fly lines in a defined genetic background provides a unique tool kit to systematically address the individual and collective contributions of immune modules to defense against a variety of pathogens.

Imd, Toll, Phagocytosis, and Melanization modules function largely independently

The Imd, Toll, Phagocytosis, and Melanization modules are interconnected (Westlake et al., 2024). For instance, the Toll and Imd pathways both regulate a subset of genes involved in melanization (De Gregorio et al., 2002; Ligoxygakis et al., 2002). Moreover, one module can indirectly affect another. For instance, a lack of melanization could increase the bacterial load after infection, resulting in higher activation of the Toll and Imd pathways (Binggeli et al., 2014). In the first step, we analyzed the activation of each of the four modules in single- and multiple-module deficient flies using classical readouts of these modules. This also allowed monitoring of the possible effects of deleting one module on the activity of another. We analyzed the immune response in single module deficient flies by monitoring **i)** the expression of the antibacterial peptide gene *Diptericin* (*DptA* - a target of the Imd pathway), and **ii)** the antifungal peptide gene *Drosomycin* (*Drs* - a target of the Toll pathway) upon systemic infection with a mixture of heat-killed *Escherichia coli* and *Micrococcus luteus*, **iii)** ex vivo phagocytosis of *E. coli* and *Staphylococcus aureus* coated bioparticles incubated with larval hemocytes, and **iv)** melanization at the injury site of adult flies. This analysis first confirmed that Δ*IMD*, Δ*TOLL*, Δ*Phag,* and Δ*Mel* flies are deficient for Imd, Toll, phagocytosis, and melanization respectively (**Figure 1A-D**). We note that the expression level of *DptA* was roughly similar to the wild-type in Δ*TOLL* and Δ*Mel* flies, however, *ΔPhag* flies show reduced *DptA* induction at 6h compared to WT (~50%) (**Figure 1A**). This represents an immense induction of *DptA* after infection, but may suggest either a lag or lesser overall Imd response in *ΔPhag* flies. *DptA* expression in *ΔPhag* was, however, comparable to WT at 12h and 24h. On the other hand, *Drs* induction at 6h in *ΔMel* flies was significantly higher than WT (~1.6x), as found previously using live *M. luteus* infection (Binggeli et al., 2014) (**Figure 1B**). The phagocytic assay confirmed that the Δ*Phag* mutant was significantly impaired in its ability to phagocytose bacterial bioparticles. Furthermore, Δ*IMD,* and Δ*Mel* mutant larvae had a phagocytic index similar to WT (**Figure 1C**), while Δ*TOLL* had a higher phagocytic index, but with high variance. As expected, there was no apparent melanization at the site of the injury in Δ*Mel* flies while blackening was readily observed in the wild-type and in other single-module mutants (**Figure 1D**). Collectively, this analysis indicates that each of the four immune modules can be selectively blocked in the isogenic DrosDel background without interfering with the competence of the others. We did note in some cases a modest change in module activity in the absence of another (higher Toll activity in Δ*Mel,* lower IMD activity in Δ*Phag,* and higher phagocytic activity in Δ*TOLL),* findings that require further exploration to establish consistency.

      We then repeated our analysis using our set of module compound mutants. Unsurprisingly, all compound flies displayed the expected immune deficiencies, validating the lines we constructed. This analysis confirms that *ΔITPM* flies, although viable, are indeed fully immune deficient for the four host defense modules. We did not detect any striking synergistic or antagonistic effects where deleting two modules had a vastly different impact compared to the additive effect of deleting the modules individually. Notably, a small level of *Drs* expression is retained in Δ*TOLL* mutant flies, but was fully abolished in Δ*IMD,* Δ*TOLL* flies (**Figure 1A vs 1E**), confirming observations that the *Drs* gene receives a small input from the Imd pathway (Leulier et al., 2000). The Δ*TOLL,* Δ*Mel^{PPS}* and, Δ*TOLL,* Δ*Mel^{HP}* flies display *Drs* induction at 6h that is resolved to unchallenged WT levels by 12h, an expression pattern consistent with

mediation of this induction by early input from the IMD pathway in these backgrounds. Taken together, all the compound mutant flies behave largely as expected given the sum effects of their individual mutations, confirming that these modules largely function independently.

Both the Melanization and TOLL modules contribute to lifespan and survival to wounding

Before assessing resistance to infection, we monitored fly survival at 25°C and 29°C to determine if module mutations impact viability in the absence of challenge. At 25°C, no single module mutants died significantly more than wild-type in the absence of challenge within the ten-day window (**Figure 2A**). However, even in the absence of challenge all compound mutants other than Δ*IMD, ΔTOLL* suffered at least 25% cumulative mortality by 10 days post-eclosion, while *ΔTOLL, ΔMel^{PPS}* and Δ*ITPM* flies suffered even greater cumulative mortality. Similar trends were observed at 29°C with slightly elevated mortality (**Figure 2supp1**). The observation that *ΔTOLL, ΔMel^{PPS}* (*PPO1^1,PPO2^1, spz^{rm7}*) flies have shorter lifespan compared to *ΔTOLL, ΔMel^{HP}* (*Hayan-psh^{Def}*) could be explained by residual Toll and/or PO activity in *Hayan-psh^{Def}* or additional indirect effects in *ΔTOLL, ΔMel^{PPS}* flies. As such, the combined effect of Toll and Melanization deficiencies in unchallenged and clean injury survival is best assessed by considering signals from both genotypes. *ΔITPM* flies (*Hayan-psh^{Def}; NimC1^1; Eater^1, Rel^{E20}*) have the lowest unchallenged survival rate of genotypes included in this study. Taken together, the Imd, Toll, Melanization, and Phagocytosis modules contribute little to lifespan maintenance individually, but are crucial to maintain lifespan collectively (**Figure 2B**).

Systemic infections were performed by septic injury (pricking with a contaminated needle). We therefore monitored the survival of single and double mutant flies after a clean injury to disentangle the effects of injury from those of infection. We found a cumulative mortality of ~50% for *ΔMel* flies by day 10 (**Figure 2C**), while other single module mutants displayed relatively little mortality over the 10-day window; *ΔIMD* had notable late-onset mortality in a minority of experimental replicates, which may be due to stochastic dysbiosis effects during aging (Hanson and Lemaitre, 2023; Marra et al., 2021). Double mutant *ΔIMD, ΔTOLL* and Δ*IMD, ΔPhag* flies retained a high survival rate. Consistent with an importance of melanization in the injury response (Binggeli et al., 2014), all the compound mutants affected in melanization were susceptible to clean injury, with similar mortality seen in *ΔPhag, ΔMel*, *ΔIMD, ΔMel, ΔTOLL, ΔMel^{PPS}, ΔTOLL, ΔMel^{HP}*, and *ΔITPM* flies. Thus, while these genotypes differ in their base lifespan, their viability upon injury is remarkably similar. These results emphasize the interconnectedness of the melanization response with every other immune module for maintaining health after injury.

Similar results were obtained for both unchallenged and clean injury lifespans at 29°C. Of note, *ΔTOLL, ΔMel^{PPS}* suffered significantly greater mortality in unchallenged conditions than other genotypes, even including *ΔITPM* flies (**Figure 2supp1**). Other compound mutant genotypes displayed a continuum of mortality in unchallenged conditions between *ΔITPM* (~50% cumulative mortality) and *ΔTOLL, ΔMel^{HP}* (~15% cumulative mortality). Upon clean injury, results were broadly consistent with 25°C trends, with a more uniform mortality across all genotypes.

385   Collectively, our study highlights that all these immune modules contribute somewhat
386 to the survival of flies upon clean injury, with a major role of the melanization response in
387 maintaining lifespan and survival to injury. In the next steps of our study, we compared survival
388 upon infection to both unchallenged and clean injured flies to distinguish the impact of
389 microbial infection from the injury itself.
390
Multiple mechanisms contribute to resistance to microbial infection
To compare the contribution of each module to host defense, we performed infection
experiments of wild-type and single- and multiple-module flies following septic infection with
five viruses, eight Gram-positive bacteria, eight Gram-negative bacteria, septic injury for two
fungal species, and natural infection (*i.e.* spore deposition on the cuticle) for two fungal species.
Temperature, dose, and days monitored are shown in **Figure 3**, and see **Table S1** for other key
parameters. Survival curves are shown in **Supplementary file 1** for each pathogen and results
of these survivals are summarized in **Figure 3**.
We performed infection experiments with 24 viruses and microbial species: DCV, FHV,
DXV, IIV-6, SINV, *Aspergillus fumigatus*, *Beauveria bassiana*, *Candida albicans*,
Mycobacterium marinum, *Corynebacterium diphtheriae*, *Micrococcus luteus*, *Streptococcus*
pneumoniae, *Enterococcus faecalis*, *Staphylococcus aureus*, *Listeria monocytogenes*, *Bacillus*
subtilis, *Vibrio parahemolyticus*, *Providencia rettgeri*, *Providencia burhodogranariea*,
Pectobacterium carotovorum Ecc15, *Klebsiella pneumoniae*, *Enterobacter cloacae*,
Escherichia coli, and Salmonella enterica ser. typhimurium. Collectively, this represents
infections with 24 pathogens across 14 genotypes and two infection routes (~350 genotype-by-
pathogen interactions). When combined with comparisons to clean injury and unchallenged
controls to validate if a lifespan difference is meaningful, and to wild-type or $\Delta ITPM$ treatments,
this inflates to 1000+ interactions. We therefore focused our survival analysis on major effects.
We report the core summary statistics and overt trends, and comment only where difference in
mean lifespan between treatments and reference genotypes or controls was a minimum of 1
day.
Our study confirms that the Imd pathway plays a major role against all tested Gram-
negative bacteria, as ΔIMD flies succumb more rapidly than wild-type flies to all of them. This
pathway also contributes, albeit to a lesser extent, to survival against the Gram-positive bacteria
B. subtilis and *L. monocytogenes,* which bear DAP-type peptidoglycan at their membrane that
triggers Imd activation (Kaneko et al., 2004; Leulier et al., 2003). We were interested to see
the impact of our ΔIMD flies in host defense to viruses as the specific mutation we used (Rel^{E20})
deletes a transcription factor common to both the canonical Imd pathway and the recently-
appreciated cGLR-Sting-Relish pathway with roles in viral defense (Goto et al., 2018). We
found that ΔIMD mutant flies were somewhat susceptible to the four viruses DCV, FHV, DXV,
and IIV6, but not SINV, agreeing with previous studies showing a reduction in lifespan of 1-2
days for *Rel* mutant flies after viral infection (Costa et al., 2009; Goto et al., 2018; Sansone et
al., 2015).
This study also confirms a prominent role of the Toll pathway in survival upon infection
by all virulent Gram-positive bacteria and fungi upon septic injury. Less virulent microbes and
infectious routes (such as *A. fumigatus* by natural infection) did not result in increased mortality
in individual $\Delta TOLL$ mutant flies. A contribution of the Toll pathway to survival was also

observed for some Gram-negative bacteria, including a minor contribution to survival upon *P. rettgeri* infection, and a major contribution after *V. parahemolyticus* infection. In addition, we found a susceptibility of *ΔTOLL* flies to FHV, DXV, and IIV6, suggesting Toll-regulated effectors mediate defense against some viral infections.

Use of *ΔPhag* mutants reveals a role of the cellular response in survival against the fungus *B. bassiana*. *ΔPhag* flies further show a minor susceptibility to *S. aureus*, and a subset of Gram-negative bacteria (*P. rettgeri*, *Ecc15*, and *E. coli*). Susceptibilities to Gram-negative bacteria could result from the observed lag in Imd pathway activation (Figure 1A); indeed, previous studies have shown a full *DptA* response is pivotal for defense against *P. rettgeri* (Hanson et al., 2023, 2019; Unckless et al., 2016), and lifespans are greatly impacted by subtle difference for pathogens with intermediate levels of virulence (Duneau et al., 2017). A role for Imd in explaining *ΔPhag* susceptibilities does not detract from the importance of the cellular response, but instead offers a route to explore mechanism for these phenotypes.

Surprisingly, our study reveals a relatively consistent role of melanization against all the tested viruses, notably so for DXV, IIV-6, and SINV. *ΔMel* flies also display a minor increased susceptibility to *B. bassiana* natural infection and septic injury. As previously described (Dudzic et al., 2019), *ΔMel* flies are highly susceptible to the Gram-positive bacterium *S. aureus*, rivaling the susceptibility of *ΔITPM* flies. *ΔMel* flies further show a strong susceptibility to *P. rettgeri* rivalling *ΔIMD* flies, and also a minor susceptibility to infection by other Gram-negative bacteria (*P. burhodogranariea, E. coli*).

Collectively, the use of single module mutants confirms the major role of the Imd pathway against Gram-negative bacteria and Toll against virulent Gram-positive bacteria and fungi. Interestingly, melanization was consistently important to survive viral infection, and plays a prominent and important role in defense against specific bacterial species. While it was almost never the most critical module, phagocytosis and the cellular response contributes to survival against various germs. Use of single module deficient flies alongside *ΔITPM* flies reveals that survival to some of the microbes tested relies almost entirely on one pathway (Imd – *Ecc15, K. pneumoniae, E. cloacae, P. burhodogranariea*; Toll – *E. faecalis*), or can receive notable contributions from two pathways (Toll and IMD – *L. monocytogenes, B. subtilis, V. parahemolyticus*; Toll and Phagocytosis – *A. fumigatus* and *B. bassiana* natural infection; Imd and Melanization - DCV). Strikingly, many pathways contribute to host defense to *S. aureus* and *P. rettgeri*.

Contribution of AMP and Bomanins to Toll and Imd activities

Recent studies have described a prime importance of antimicrobial peptides (AMPs) and host defense peptides (HDPs) in responding to infection (Clemmons et al., 2015; Hanson, 2024; Hanson et al., 2019). These peptides are the principal effectors of the Toll and Imd pathways, yet mutations for AMPs and HDPs typically have not been tested alongside mutations for both the Imd and the Toll pathways across pathogens, as studies have typically included just the single relevant pathway mutation as a positive control. Thus, the importance of AMPs and HDPs relative to the pathways that regulate them has been only partially investigated. We used *ΔAMP14* and *ΔBom* flies to assess the extent to which AMPs and Bomanins mediate Toll and Imd contributions to defense (**Figure 3**). Here we found that AMPs are critical for survival to

infection against all Gram-negative bacteria tested, often rivaling the susceptibility of *ΔIMD* flies. Notably, *ΔAMP14* flies were also somewhat susceptible to the yeast *C. albicans,* and the Gram-positive bacteria *B. subtilis* and *M. luteus*, reinforcing minor susceptibilities seen previously (Carboni et al., 2022; Hanson et al., 2019). Interestingly, we found that *ΔAMP14* survivals to DCV, FHV, DXV, and IIV6, largely paralleled those of *ΔIMD* flies (Figure 3, Supplemental File 1), suggesting Imd-mediated survival phenotypes rely greatly on the presence of the seven classical AMP families.

On the other hand, *ΔBom* flies were susceptible to septic injury by *B. bassiana* and also somewhat to *C. albicans,* consistent with previous studies (**Supplementary file 1**), as well as all virulent Gram-positive bacteria, consistent with their important role downstream of the Toll pathway (Clemmons et al., 2015; Lindsay et al., 2018; Xu et al., 2023). In addition, *ΔBom* flies paralleled the *ΔTOLL* susceptibility to DCV, FHV, and DXV, but were not susceptible to IIV6 like *ΔTOLL*.

Collectively, our study confirms the key role of antimicrobial peptides downstream of the Imd pathway in the defense against Gram-negative bacteria and a primary role of Bomanins downstream of Toll in defense against fungi and Gram-positive bacteria, with little input to survival against Gram-negative bacteria. We additionally find these secreted peptides can contribute to survival after certain viral infections.

Immune modules mostly contribute additively to host defense

The results described above reveal the key contribution of single modules and immune effectors to host defense. However, they did not assess possible additive, synergistic, or antagonistic contribution of these four modules. Moreover, they did not monitor to what extent these modules are important in the absence of other modules. To address this, we compared the survival of single-module deficient flies to flies lacking two modules, and *ΔITPM* flies lacking all four modules (**Figure 3**).

We first observed that *ΔITPM* flies, which lack the four modules, were always as or more susceptible than single- and double-module deficient flies. In multiple microbe-specific cases, single or double mutants rivalled the susceptibility of *ΔITPM* flies. In particular, the susceptibility of *ΔIMD*-flies to most Gram-negative bacteria rivalled *ΔITPM,* and for some germs *ΔTOLL* (*L. monocytogenes*, *E. faecalis*, *B. bassiana* injury) or *ΔMel* (*S. aureus*) alone was comparable to *ΔITPM*. In other cases, however, combined loss of two pathways was necessary to rival *ΔITPM* susceptibilities. For instance, *ΔIMD, ΔTOLL* deletion causes complete *ΔITPM*-like susceptibility to *B. subtilis*, *M. luteus*, and *E. coli*, and near-complete susceptibility to *S. typhimurium*. *ΔIMD, ΔTOLL* further causes increased susceptibility to most viruses compared to its single module counterparts. In these cases, there is clear synergy or additivity of the contributions of the two modules, as either pathway alone results in only minor or no susceptibility. We also observed intriguing differences between our two versions of *ΔTOLL, ΔMel* flies, as *ΔTOLL, ΔMel^{PPS}* tended to succumb to infection to a greater extent than *ΔTOLL, ΔMel^{HP}*, consistent with the differences observed upon clean injury (**Figure 2**). The *ΔTOLL, ΔMel^{PPS}* displayed a minor increase in susceptibility compared to *ΔTOLL, ΔMel^{HP}* to *A. fumigatus* natural infection, and more prominent susceptibilities to the yeast *C.*

514   *albicans*, the Gram-positive bacteria *M. marinum*, *M. luteus* and also to Gram-negative

515   bacteria *K. pneumoniae*, *E. coli*, *P. rettgeri* and *P. burhodogranariea*.

516   Collectively, use of double-module deficient flies uncovers contributions of immune

modules to host defense that were masked by other modules. While *ΔIMD* and *ΔTOLL*

deficiencies sometimes displayed a synergistic effect (*i.e.* little mortality of single mutants,

but *ΔITPM*-like mortality of double mutants), for combinations of other pathways the results

were more often minor with a cumulative effect on susceptibility, suggesting most modules

contribute to defense independently. The use of two variations of *ΔTOLL*, *ΔMel*, with *ΔTOLL*,

ΔMel^{PPS} having a stronger impact than *ΔTOLL*, *ΔMel^{HP},* also highlights how the level of

pathway disruption can affect infection outcomes, particularly for certain fungal or bacterial

pathogens like *Providencia* species that may cleave host immune proteases (suggested by

Duneau et al., 2017; Issa et al., 2018).

Timing of Toll, Imd, Phagocytosis, and Melanization contribution to host defense

Host defense programs can rely on resistance mechanisms that directly target or limit growth

of pathogens (Howick and Lazzaro, 2017; Lafont et al., 2021; Medzhitov et al., 2012). When

effectors reach a critical concentration threshold that promotes resistance, pathogen growth is

inhibited; for instance, the time to pathogen control of *P. rettgeri* is ~7 hours in *Drosophila*,

relying heavily on the expression and production of DptA (Duneau et al., 2017; Hanson et al.,

2023). Phagocytosis and the melanization response are often described as providing near-

immediate immune protection (Haine et al., 2008), with reactions ex vivo progressing within

minutes. Meanwhile activation of the Imd and Toll pathways takes hours to reach peak

concentrations of host defense peptides (Uttenweiler-Joseph et al., 1998). Thus, we were

curious if different mutations would alter time of action to control microbial growth. We

therefore measured the microbial growth rate for *P. rettgeri, S. aureus,* and *C. albicans* at

different time points post-infection in wild-type and single module mutant flies as these

pathogens were combatted by several immune modules in our survival data.

541   Agreeing with contributions to survival, flies singly deficient for IMD, Phagocytosis,

or Melanization modules display a higher bacterial load than wild-type when infected with *P.*

rettgeri **(Figure 4A)**, which was apparent at 12 hours post-infection. Our observation that

ΔPhag showed impaired IMD signaling at 6h (**Figure 1A**) may explain the contribution of

phagocytosis to *P. rettgeri* microbial growth control with the same timing as *ΔIMD* flies. We

observed that *ΔIMD* flies, *ΔMel* flies, and *ΔITPM* flies, each of which succumb completely to

P. rettgeri infection, also show high and consistent microbe loads at 24h that are indicative of

sepsis-induced death. On the other hand, wild-type, *ΔTOLL*, and *ΔPhag* flies show a greater

stochasticity in bacterial load at 24h that is consistent with a fraction of individuals surviving

the infection by controlling the pathogen. This indicates deletion of the Toll pathway and

phagocytosis does not fully ablate resistance mechanisms essential for combatting *P. rettgeri*.

552   In the case of *S. aureus* (**Figure 4B**), we observed some individuals with higher bacterial

553   loads in ΔPhag, and ΔIMD flies at 2h and 6h, which was not seen in wild-type, ΔTOLL, or ΔMel

554   flies. This suggests that the Imd pathway can suppress early *S. aureus* growth, and possibly also

555   phagocytosis, although this difference in ΔPhag could ultimately stem from reduced IMD

556   induction. Interestingly, flies lacking melanization displayed the highest susceptibility, but did

not depart from wild-type or other modules in pathogen load until 12h post-infection. While the melanization reaction is a rapid response, these results suggest that the killing activity of the melanization response acts with slower kinetics. In *ΔITPM* flies, we observed earlier *S. aureus* growth at 2h and 6h as in *ΔIMD* and *ΔPhag* flies. However, *S. aureus* loads at 12-24h were more comparable in *ΔITPM* flies to *ΔMel* flies. These trends agree both with survival data kinetics, and an independent action of each of these pathways in their resistance effects against *S. aureus*. This suggests that *ΔITPM* flies succumb even more quickly than individual module mutants due to loss of multiple resistance mechanisms with different kinetics of activation that independently contribute to defense.

For *C. albicans* (**Figure 4C**), mortality begins 2-3 days post-infection in ΔTOLL mutants, but takes place at later time points for Δ*IMD,* Δ*Phag,* and *ΔMel* flies. Microbial growth kinetics show high *C. albicans* loads at 24h in Δ*TOLL,* consistent with a previous study (Hanson et al., 2019) and observations done with *Candida glabrata* (Quintin et al., 2013). Interestingly, a few ΔIMD flies also showed elevated growth of *C. albicans* at early time points. This suggests Imd-regulated effectors may help to suppress initial *C. albicans* growth, while Toll-regulated genes (*e.g.* Bomanins, Drs) contribute to this pathogen suppression to a greater extent at later time points. On the other hand, ΔPhag and ΔMel flies did not show any increased *C. albicans* load within the 24h time window, despite onset of mortality at later time points. Finally, *ΔITPM* flies displayed more consistent and rapid growth of *C. albicans* than *ΔTOLL* or *ΔIMD* alone, particularly visible at 12-24h post-infection, consistent with independent contributions of both Toll and Imd in resistance to this yeast.

Collectively, we observed that microbial growth in each module mutant parallels susceptibility, indicating that they contribute to resistance. Surprisingly we did not observe an early role in pathogen killing for melanization despite the rapid blackening response mediated by melanization in ex vivo assays (Nakhleh et al., 2017). We further recover stochasticity in microbe loads across genotypes, as expected when only a subset of individuals suppress the pathogen and survive the infection, resulting in bimodal outcomes (Duneau et al., 2017). The more rapid and consistent microbial growth kinetics of *ΔITPM* flies, particularly visible at 18h, further demonstrate that the modules contribute to resistance collectively.

Contributions of individual modules to disease tolerance

A metric increasingly used to delineate roles of resistance and tolerance is the Pathogen Load Upon Death (PLUD) (Duneau et al., 2017; Duneau et al., 2025). PLUD is affected by both the virulence of the pathogen and tolerance of the host to survive to a given pathogen burden before succumbing to infection. We monitored PLUD for the same pathogens used to determine the timing of defense modules: *P. rettgeri, S. aureus,* and *C. albicans.* The PLUD of both *P. rettgeri* and *S. aureus* was overall similar in wild-type and single-module mutant flies **(Figure 5A-B)**. Notably, there was a greater stochasticity in *P. rettgeri* infections trending towards lower PLUD in *ΔITPM* flies, suggesting a stochastic reduction in disease tolerance of *ΔITPM* flies to this bacterium, although this was not significant compared to wild-type (*P > .05*).

For *C. albicans*, the distribution of PLUD values for *ΔPhag* flies was markedly different from wild-type, also seen to some extent for *ΔMel*. Interestingly, flies with these genotypes suppress *C. albicans* growth well (**Figure 4C**). Yet here we recovered many individuals from

both of these module-deficient lines that died with a far lower PLUD (**Figure 5C**). Taken
together, this suggests that the humoral immune pathways, particularly Toll, contribute to
resistance against *C. albicans*. However, the cellular response and melanization reaction instead
regulate tolerance to *C. albicans* infection. The use of Δ*ITPM* flies further emphasizes that
when both resistance (Toll, Imd) and tolerance (Phagocytosis, Melanization) mechanisms are
deficient, the loss of resistance plays a primary importance, and can mask observations of
reduced tolerance.
Collectively, monitoring of PLUD suggests that some modules, specifically
phagocytosis and melanization, contribute to tolerance of certain infections. Importantly, this
tolerance effect was only seen in genotypes that are able to resist infection. These findings
demonstrate the collective contribution to resistance and/or tolerance of the four immune
modules studied here.
612

Discussion

In this article, we have generated a set of single and compound immune module deficient fly
lines in a controlled genetic background. Using various assays, we confirmed the validity of
our lines, revealing that each module can be activated independently. We did, however, observe
higher Toll activation in Δ*Mel* flies upon systemic infection with dead bacteria. Future studies
may reveal that this higher Toll activity in Δ*Mel* flies is due to interaction with a regulatory
pathway, or reflects higher persistence of microbial elicitors (*e.g.* peptidoglycan) that activate
the Toll pathway in melanization-deficient flies. We also observed reduced Imd activation in
Δ*Phag* flies at 6 hours post-infection, which could reflect a role for hemocytes in stimulation
of the Imd systemic response. In this study, we used *NimC1[1]; eater[1]* double mutants to assess
the cellular response. These flies have defects in phagocytosis, but also hemocyte adhesion and
sessility (Melcarne et al., 2019). *NimC1[1]; eater[1]* larvae also have increased hemocyte number
at the larval stage, but Hml positive cells are rapidly lost in adults (Melcarne, 2020). Despite
these limitations, we believe that the mutations we have used here are among the best available
to assess the roles of these four modules to host defense. Surprisingly, we were able to produce
a fly line lacking all four main defense mechanisms of the systemic immune response,
indicating that none of these modules is essential for survival. Theoretically, Δ*ITPM* flies are
almost completely immune deficient, but they are still able to clot wounds, activate the JNK
and JAK-STAT pathways that mediate the wound healing response, and retain constitutive
immune defense molecules that could provide a certain degree of protection. Consistent with
several studies (Capilla et al., 2017; Carvalho et al., 2014; Rämet et al., 2002), we show that
Toll and melanization contribute synergistically to wound healing in adults. However, the
observation that Δ*Phag, ΔMel* or Δ*IMD, ΔMel* double mutant flies are also more susceptible to
clean injury than Δ*Mel* flies indicates that the Imd pathway and phagocytosis also contribute to
wound healing. Thus, although immune deficient fly lines for the four modules are viable, the
use of compound module mutants and Δ*ITPM* flies reveals a clear role of the immune system
in response to wounding and lifespan maintenance.
Use of single- and double module-deficient flies provides key insights on the
mechanisms used by *Drosophila* to combat infection. We confirm previous studies revealing
the role of Imd against Gram-negative bacteria, Toll against Gram positive bacteria and virulent
fungi, and an importance of Melanization and Phagocytosis against specific pathogens

(Charroux and Royet, 2009; Defaye et al., 2009; Garg and Wu, 2014; Nehme et al., 2011).
Surprisingly, the Melanization module was consistently important in survival to viral infection.
It seems unlikely that prophenoloxidase activity in the hemolymph can combat intracellular
viral agents. We instead speculate this contribution of melanization to viral defense could be
due to its role in wound healing, or perhaps autotoxic contributions of melanization reaction
intermediates that fail to convert in phenoloxidase-deficient flies. We additionally recovered a
role of Imd signaling (i.e. *Relish*) in antiviral defense, which was expected given recent
characterizations of cGLR-Sting-Relish antiviral immunity (Ai et al., 2024; Cai et al., 2022;
Goto et al., 2018). However, susceptibilities of *Rel^{E20}* flies were paralleled by Δ*AMP14* in all
cases, suggesting AMPs largely mediate this susceptibility. While Sting regulates a number of
genes likely important for antiviral defense (Goto et al., 2018), the susceptibility we observe
here could be a direct action of AMPs on enveloped viruses as described in some studies (Feng
et al., 2020; Huang et al., 2013; Yasin et al., 2004), or an indirect effect, such as a need to
regulate gut microbes after the damage induced by viral replication (Marra et al., 2021). We
additionally used dual modes of infection for the fungus *B. bassiana,* finding that both
Melanization and Toll modules have a role in both infection modes. However, our study reveals
that Bomanin effectors explain most of the Toll contribution upon septic injury but not natural
infection by *B. bassiana*. This observation is in line with previous observations showing
differential importance of effectors or modules according to infection route (Martins et al.,
2013).

Our double module mutant analysis reveals that most modules contribute additively to
host defense. This is consistent with largely independent function of these modules. However,
we noted some instances of synergy between two pathways, notably Toll and Imd. Synergy
between Toll and Imd can be explained by the fact that many immune-inducible regulated genes
receive input of both the Imd or Toll pathways; some like *Metchnikowin*, *Drosomycin* and
Transferrin1 that can be induced through either Toll or Imd (De Gregorio et al., 2002). Finally,
we observed rarer cases of synergy between Toll and Melanization, or Imd and Melanization.

Our study confirms that the Toll and Imd humoral modules have a defined roles against
broad classes of pathogens, Imd for Gram-negative bacteria and DAP-type containing Gram-
positive bacteria, and Toll for fungi and Gram-positive bacteria. Use of AMP and Bomanin
mutants revealed that this can be largely explained by the effectors they control. In contrast, the
contributions of Phagocytosis and Melanization appear to be critical to more specific and
diverse sets of pathogens. We speculate that Phagocytosis and/or Melanization have critical
roles in defense against bacteria that resist host defense peptides (Hanson et al., 2019), or can
hide from them (Touré et al., 2023). The melanization reaction is a source of ROS that is a
potent defense against pathogens such as *S. aureus* resistant to Toll and Imd defense (Dudzic
et al., 2019; Ford and King, 2021; Needham et al., 2004; Ramond et al., 2021). These two
modules play important roles in immune-related processes such as encapsulation
(Melanization), the uptake of bacteria escaping from the gut, or tissue homeostasis
(Phagocytosis) (Braun et al., 1998; C. Melcarne et al., 2019; Nehme et al., 2007) that were not
assessed here. Collectively, our study validates, with minor discrepancies, many studies that
have assessed the individual contributions of these modules (*e.g.* Apidianakis et al., 2005;
Binggeli et al., 2014; Lamiable et al., 2016; Lemaitre et al., 1996, 1995; Nehme et al., 2011).
Our mutant lines can now be used to analyze the contribution of these immune modules in

resistance to other pathogens, notably wasps, nematodes, microsporidia and protozoans, or in
other contexts such as mating and local infection.

While defense against some pathogens is largely mediated by single immune modules,
other pathogens are handled by multiple modules. We hypothesize that pathogens that have
intermediate virulence levels (kill only a fraction of wild-type flies), may better reveal the roles
of multiple modules. Indeed, stochasticity in survival analyses partly stems from the arms-race
occurring between the pathogen and host immunity, as shown for *P. rettgeri* (Duneau et al.,
2017). In this scenario, any small change in immune competence may tip the outcome of the
arms-race toward lethality or survival. Here, it is notable that multiple modules contribute to
survival in *P. rettgeri* infection. Previous studies have revealed a major role of the Imd pathway
AMP Diptericin against this bacterium (Hanson et al., 2023, 2019; Unckless et al., 2016).
However, Duneau et al. (Duneau et al., 2017) showed that survival patterns to *P. rettgeri*
bifurcate into two outcomes based on time taken to fully activate systemic defenses, and showed
a role for a Toll-PO SP cascade regulating serine proteases in defense against this microbe
(Duneau et al., 2017). We may speculate that Melanization, although not as great a contributor
as Diptericin, may be a factor that tips the balance of this precarious arms race towards host
lethality. The observation that *ΔPhag* flies expressed less *DptA* (**Figure 1**) can explain their
susceptibility to *P. rettgeri*. Future studies should investigate how multiple modules can
contribute to host survival according to additive or Achilles dynamics – the concept that
microbes have specific weaknesses available for host effectors to target (Hanson, 2024). It will
also be important to consider the distinct roles of resistance, tolerance, and resilience in host
defense (Howick and Lazzaro, 2017; Wukitch et al., 2023).

We observed a good correlation between survival analysis and pathogen growth in
single module flies for *P. rettgeri*, *S. aureus,* and *C. albicans*. This indicates that these immune
modules mostly contribute to resistance mechanisms that suppress pathogen growth. Our study
did not reveal key early contributions of phagocytosis or melanization in pathogen growth
control, despite quasi-immediate activation of these modules; higher *S. aureus* growth at 2
hours in *ΔPhag* flies was paralleled by *ΔIMD* flies, and we observed reduced Imd activation in
ΔPhag flies. Melanization, while critical to surviving *S. aureus* infection, impacts bacterial
growth beginning only at the 12-hour time point. This indicates that the microbicidal activity
associated with Melanization in vivo acts more slowly than the blackening reaction observed
in bled hemolymph. Interestingly, ΔPhag and also ΔMel flies suppress *C. albicans* yeast
growth, but some individuals ultimately succumb to infection with lower PLUD levels. We
confirmed ΔPhag PLUD results using both an isogenic and a second wild-type genetic
background, suggesting that this lower PLUD is genuine. Susceptibility to fungal infection
independent of fungal proliferation has also been reported using an *A. fumigatus* septic infection
model, and relies on the protection offered by Bomanins from pathogen-secreted toxins (Xu et
al., 2023). It is tempting to speculate, based on the modules involved, that the loss of tolerance
to *C. albicans* we observe is related to wound repair or accumulating damage, similar to reports
of renal failure in beetles (Khan et al., 2017; Li et al., 2020), or flies with autotoxic trachea
degradation upon stress pathway disruption (Rommelaere et al., 2024). Thus, this tolerance
effect could rely on pathogen-mediated or autotoxic damage, which may be elucidated in a
future study.

Here we have provided a single and double mutant analysis of *Drosophila* immune module functions. Our study provides several insights showing which modules are most important to survive infection by defined pathogens. However, we also highlight collective contribution of modules to defense, even when one module is of outsized importance. We extend our comprehension of innate immune responses by revealing higher complexity in defense, implicating multiple host immune modules in survival to various germs, including some with more cryptic contributions. As illustrated by our previous characterizations of AMP function (Carboni et al., 2022; Hanson et al., 2019), the melanization response (Dudzic et al., 2015), and stress-induced Turandot proteins (Rommelaere et al., 2024), a combinatorial mutation approach to deciphering immune function can be extended even to the broad level of whole immune modules. Together with these previous studies and others, our study provides an immunity toolkit, including flies simultaneously deficient for four immune modules, which can be used to better characterize the contributions of immune processes to host defense. Our tools can also be of use to study various questions at the cutting edge of aging, memory, neurodegeneration, cancer, and more, where immune genes are repeatedly implicated. We hope that this set of lines will be useful to the community to better characterize the *Drosophila* host defense.

Acknowledgement:
We thank Mélanie Blokesch (EPFL), Jan-Willem Veening (Unil), Vivek Thacker (Heidelberg University), Carla Saleh (Pasteur Institut) and Ronald P van Rij (Radboud University Medical Center) for key reagents. We thank Luis Teixeira for iso Drosdel wild-type, rel and spz flies. We thank Koenig E, Prince Kumar Sah for experimental help and Hannah Westlake for editing. The AMP, Bom and single and quadruple module mutants were deposited at the Vienna *Drosophila* Research Center. This project was supported by the SNSF grant 310030_215073 awarded to B.L. and Wellcome Trust grant 227559/Z/23/Z awarded to M.H.

Author contributions
Conceptualization: FR, MAH, BL
Data curation: FR, MAH
Funding acquisition: BL, MAH
Investigation: FR, YT, MR, FS
Methodology: FR
Project administration: BL
Validation: YT, MAH
Resources: FR, MAH
Supervision: MAH, BL
Writing – original draft: FR, BL
Writing – review & editing: MAH, BL

References

Ai X, Deng H, Li X, Wei Z, Chen Y, Yin T, Zhang J, Huang J, Li H, Lin X, Tan L, Chen D, Zhang Xiaohan, Zhang Xiuqing, Meignin C, Imler J-L, Cai H. 2024. cGAS-like receptors drive a systemic STING-dependent host response in Drosophila. *Cell*

Reports **43**:115081. doi:10.1016/j.celrep.2024.115081

Apidianakis Y, Mindrinos MN, Xiao W, Lau GW, Baldini RL, Davis RW, Rahme LG. 2005.
Profiling early infection responses: Pseudomonas aeruginosa eludes host defenses by
suppressing antimicrobial peptide gene expression. *Proc Natl Acad Sci U S A*
102:2573–8.

Binggeli O, Neyen C, Poidevin M, Lemaitre B. 2014. Prophenoloxidase Activation Is
Required for Survival to Microbial Infections in Drosophila. *PLoS Pathogens* **10**.
doi:10.1371/journal.ppat.1004067

Braun A, Hoffmann JA, Meister M. 1998. Analysis of the Drosophila host defense in domino
mutant larvae, which are devoid of hemocytes. *Proc Natl Acad Sci U S A* **95**:14337–
42.

Bretscher AJ, Honti V, Binggeli O, Burri O, Poidevin M, Kurucz E, Zsamboki J, Ando I,
Lemaitre B. 2015. The Nimrod transmembrane receptor Eater is required for
hemocyte attachment to the sessile compartment in Drosophila melanogaster. *Biology*
open **4**:355–63. doi:10.1242/bio.201410595

Buchon N, Poidevin M, Kwon H-M, Guillou A, Sottas V, Lee B-L, Lemaitre B. 2009. A
single modular serine protease integrates signals from pattern-recognition receptors
upstream of the Drosophila Toll pathway. *Proc Natl Acad Sci U S A* **106**:12442–
12447. doi:10.1073/pnas.0901924106

Cai H, Meignin C, Imler J-L. 2022. cGAS-like receptor-mediated immunity: the insect
perspective. *Curr Opin Immunol* **74**:183–189. doi:10.1016/j.coi.2022.01.005

Capilla A, Karachentsev D, Patterson RA, Hermann A, Juarez MT, McGinnis W. 2017. Toll
pathway is required for wound-induced expression of barrier repair genes in the
Drosophila epidermis. *Proceedings of the National Academy of Sciences* **114**:E2682–
E2688. doi:10.1073/pnas.1613917114

Carboni AL, Hanson MA, Lindsay SA, Wasserman SA, Lemaitre B. 2022. Cecropins
contribute to *Drosophila* host defense against a subset of fungal and Gram-negative
bacterial infection. *Genetics* **220**. doi:10.1093/genetics/iyab188

Carvalho L, Jacinto A, Matova N. 2014. The Toll/NF-κB signaling pathway is required for
epidermal wound repair in Drosophila. *PNAS* **111**:E5373–E5382.
doi:10.1073/pnas.1408224111

Cerenius L, Lee BL, Söderhäll K. 2008. The proPO-system: pros and cons for its role in
invertebrate immunity. *Trends in Immunology* **29**:263–271.
doi:10.1016/j.it.2008.02.009

Charroux B, Royet J. 2009. Elimination of plasmatocytes by targeted apoptosis reveals their
role in multiple aspects of the Drosophila immune response. *Proc Natl Acad Sci U S A*
106:9797–802. doi:10.1073/pnas.0903971106

Choe K-M, Lee H, Anderson KV. 2005. Drosophila peptidoglycan recognition protein LC
(PGRP-LC) acts as a signal-transducing innate immune receptor. *Proceedings of the*
National Academy of Sciences **102**:1122–1126. doi:10.1073/pnas.0404952102

Clemmons AW, Lindsay SA, Wasserman SA. 2015. An Effector Peptide Family Required for
Drosophila Toll-Mediated Immunity. *PLoS Pathogens* **11**.
doi:10.1371/journal.ppat.1004876

Costa A, Jan E, Sarnow P, Schneider D. 2009. The Imd pathway is involved in antiviral

immune responses in Drosophila. *PLoS ONE* **4**:e7436.
doi:10.1371/journal.pone.0007436

Danilova N. 2006. The evolution of immune mechanisms. *J Exp Zool* **306B**:496–520.
doi:10.1002/jez.b.21102

De Gregorio E, Spellman PT, Tzou P, Rubin GM, Lemaitre B. 2002. The Toll and Imd
pathways are the major regulators of the immune response in Drosophila. *Embo J*
21:2568–79.

Defaye A, Evans I, Crozatier M, Wood W, Lemaitre B, Leulier F. 2009. Genetic Ablation of
Drosophila Phagocytes Reveals Their Contribution to Both Development and
Resistance to Bacterial Infection. *Journal of Innate Immunity* **1**:322–334.
doi:10.1159/000210264

Dudzic JP, Hanson MA, Iatsenko I, Kondo S, Lemaitre B. 2019. More Than Black or White:
Melanization and Toll Share Regulatory Serine Proteases in Drosophila. *Cell Reports*
27:1050-1061.e3. doi:10.1016/j.celrep.2019.03.101

Dudzic JP, Kondo S, Ueda R, Bergman CM, Lemaitre B. 2015. Drosophila innate immunity:
regional and functional specialization of prophenoloxidases. *BMC biology* **13**:81–97.
doi:10.1186/s12915-015-0193-6

Duneau D, Ferdy J-B, Revah J, Kondolf H, Ortiz G, Lazzaro B, Buchon N. 2017. Stochastic
variation in the initial phase of bacterial infection predicts the probability of survival
in D. melanogaster. *eLife* **6**. doi:10.7554/eLife.28298

Duneau DF, Kondolf H, Im J, Ortiz G, Chow C, Fox M, Eugénio A, Revah J, Buchon N,
Lazzaro B. 2017. The Toll pathway underlies host sexual dimorphism in resistance to
both Gram-negative and Gram-positive bacteria in mated Drosophila. *BMC Biology*
15. doi:10.1186/s12915-017-0466-3

Duneau D, Lafont PD, Lauzeral C, Parthuisot N, Faucher C, Jin X, Buchon N, Ferdy J-B.
2025. A within-host infection model to explore tolerance and resistance. *eLife*
14:e104052. doi:10.7554/eLife.104052

Elrod-Erickson M, Mishra S, Schneider D. 2000. Interactions between the cellular and
humoral immune responses in drosophila. *Curr Biol* **10**:781–4.

Fehlbaum P, Bulet P, Michaut L, Lagueux M, Broekaert WF, Hetru C, Hoffmann JA. 1994.
Insect immunity: Septic injury of drosophila induces the synthesis of a potent
antifungal peptide with sequence homology to plant antifungal peptides. *Journal of*
Biological Chemistry **269**:33159–33163.

Feng M, Fei S, Xia J, Labropoulou V, Swevers L, Sun J. 2020. Antimicrobial Peptides as
Potential Antiviral Factors in Insect Antiviral Immune Response. *Frontiers in*
Immunology **11**.

Ferreira AG, Naylor H, Esteves SS, Pais IS, Martins NE, Teixeira L. 2014. The Toll-dorsal
pathway is required for resistance to viral oral infection in Drosophila. *PLoS*
pathogens **10**:e1004507. doi:10.1371/journal.ppat.1004507

Ford SA, King KC. 2021. In Vivo Microbial Coevolution Favors Host Protection and Plastic
Downregulation of Immunity. *Molecular Biology and Evolution* **38**:1330–1338.
doi:10.1093/molbev/msaa292

Garg A, Wu LP. 2014. *D rosophila* Rab14 mediates phagocytosis in the immune response to
S taphylococcus aureus: Rab14 is critical for cellular immune responses to *S. aureus*.

Cellular Microbiology **16**:296–310. doi:10.1111/cmi.12220

Goto A, Okado K, Martins N, Cai H, Barbier V, Lamiable O, Troxler L, Santiago E, Kuhn L, Paik D, Silverman N, Holleufer A, Hartmann R, Liu J, Peng T, Hoffmann JA, Meignin C, Daeffler L, Imler JL. 2018. The Kinase IKKβ Regulates a STING- and NF-κB-Dependent Antiviral Response Pathway in Drosophila. *Immunity* **49**:225-234.e4. doi:10.1016/j.immuni.2018.07.013

Gottar M, Gobert V, Matskevich AA, Reichhart J-M, Wang C, Butt TM, Belvin M, Hoffmann JA, Ferrandon D. 2006. Dual Detection of Fungal Infections in Drosophila via Recognition of Glucans and Sensing of Virulence Factors. *Cell* **127**:1425–1437. doi:10.1016/j.cell.2006.10.046

Haine ER, Moret Y, Siva-Jothy MT, Rolff J. 2008. Antimicrobial Defense and Persistent Infection in Insects. *Science* **322**:1257–1259. doi:10.1126/science.1165265

Hanson MA. 2024. When the microbiome shapes the host: immune evolution implications for infectious disease. *Phil Trans R Soc B* **379**:20230061. doi:10.1098/rstb.2023.0061

Hanson MA, Dostálová A, Ceroni C, Poidevin M, Kondo S, Lemaitre B. 2019. Synergy and remarkable specificity of antimicrobial peptides in vivo using a systematic knockout approach. *eLife* **8**. doi:10.7554/elife.44341

Hanson MA, Grollmus L, Lemaitre B. 2023. Ecology-relevant bacteria drive the evolution of host antimicrobial peptides in Drosophila. *Science* **381**:eadg5725. doi:10.1126/science.adg5725

Hanson MA, Lemaitre B. 2023. Antimicrobial peptides do not directly contribute to aging in *Drosophila* , but improve lifespan by preventing dysbiosis. *Disease Models & Mechanisms* **16**:dmm049965. doi:10.1242/dmm.049965

Howick VM, Lazzaro BP. 2017. The genetic architecture of defense as resistance to and tolerance of bacterial infection in Drosophila melanogaster. *Molecular Ecology* **26**:1533–1546. doi:10.1111/mec.14017

Huang Z, Kingsolver MB, Avadhanula V, Hardy RW. 2013. An Antiviral Role for Antimicrobial Peptides during the Arthropod Response to Alphavirus Replication. *Journal of Virology* **87**:4272–4280. doi:10.1128/jvi.03360-12

Iatsenko I, Kondo S, Mengin-Lecreulx D, Lemaitre B. 2016. PGRP-SD, an Extracellular Pattern-Recognition Receptor, Enhances Peptidoglycan-Mediated Activation of the Drosophila Imd Pathway. *Immunity* **45**:1013–1023. doi:10.1016/j.immuni.2016.10.029

Imler J-L, Bulet P. 2005. Antimicrobial peptides in Drosophila: structures, activities and gene regulation. *Chemical immunology and allergy* **86**:1–21. doi:10.1159/000086648

Ip Y. 1993. Dif, a dorsal-related gene that mediates an immune response in Drosophila. *Cell* **75**:753–763. doi:10.1016/0092-8674(93)90495-C

Issa N, Guillaumot N, Lauret E, Matt N, Schaeffer-Reiss C, Van Dorsselaer A, Reichhart JM, Veillard F. 2018. The Circulating Protease Persephone Is an Immune Sensor for Microbial Proteolytic Activities Upstream of the Drosophila Toll Pathway. *Molecular Cell* **69**:648-663.e7. doi:10.1016/j.molcel.2018.01.029

Kaneko T, Goldman WE, Mellroth P, Steiner H, Fukase K, Kusumoto S, Harley W, Fox A, Golenbock D, Silverman N. 2004. Monomeric and polymeric gram-negative peptidoglycan but not purified LPS stimulate the Drosophila IMD pathway. *Immunity*

20:637–49.

Kaneko Takashi, Yano T, Aggarwal K, Lim J-H, Ueda K, Oshima Y, Peach C, Erturk-
Hasdemir D, Goldman WE, Oh B-H, Kurata S, Silverman N. 2006. PGRP-LC and
PGRP-LE have essential yet distinct functions in the drosophila immune response to
monomeric DAP-type peptidoglycan. *Nat Immunol* **7**:715–723. doi:10.1038/ni1356

Kaneko T., Yano T, Aggarwal K, Lim JH, Ueda K, Oshima Y, Peach C, Erturk-Hasdemir D,
Goldman WE, Oh BH, Kurata S, Silverman N. 2006. PGRP-LC and PGRP-LE have
essential yet distinct functions in the drosophila immune response to monomeric DAP-
type peptidoglycan. *Nat Immunol* **7**:715–23.

Khan I, Agashe D, Rolff J. 2017. Early-life inflammation, immune response and ageing.
Proceedings of the Royal Society B: Biological Sciences **284**:20170125.
doi:10.1098/rspb.2017.0125

Kocks C, Cho JH, Nehme N, Ulvila J, Pearson AM, Meister M, Strom C, Conto SL, Hetru C,
Stuart LM, Stehle T, Hoffmann JA, Reichhart J-M, Ferrandon D, Rämet M, Ezekowitz
RAB. 2005. Eater, a Transmembrane Protein Mediating Phagocytosis of Bacterial
Pathogens in Drosophila. *Cell* **123**:335–346. doi:10.1016/j.cell.2005.08.034

Kurucz É, Márkus R, Zsámboki J, Folkl-Medzihradszky K, Darula Z, Vilmos P, Udvardy A,
Krausz I, Lukacsovich T, Gateff E, Zettervall C-J, Hultmark D, Andó I. 2007.
Nimrod, a Putative Phagocytosis Receptor with EGF Repeats in Drosophila
Plasmatocytes. *Current Biology* **17**:649–654. doi:10.1016/j.cub.2007.02.041

Lafont PDM, Lauzeral C, Parthuisot N, Faucher C, Duneau D, Ferdy J-B. 2021. A within-host
infection model to explore tolerance and resistance (preprint). Evolutionary Biology.
doi:10.1101/2021.10.19.464998

Lamiable O, Arnold J, de Faria IJ da S, Olmo RP, Bergami F, Meignin C, Hoffmann JA,
Marques JT, Imler J-L. 2016. Analysis of the Contribution of Hemocytes and
Autophagy to Drosophila Antiviral Immunity. *Journal of Virology* **90**:5415–5426.
doi:10.1128/JVI.00238-16

Lemaitre B, Kromer-Metzger E, Michaut L, Nicolas E, Meister M, Georgel P, Reichhart JM,
Hoffmann JA. 1995. A recessive mutation, immune deficiency (imd), defines two
distinct control pathways in the Drosophila host defense. *Proc Natl Acad Sci U S A*
92:9465–9469. doi:10.1073/pnas.92.21.9465

Lemaitre B, Nicolas E, Michaut L, Reichhart J-M, Hoffmann JA. 1996. The Dorsoventral
Regulatory Gene Cassette spätzle/Toll/cactus Controls the Potent Antifungal Response
in Drosophila Adults. *Cell* **86**:973–983. doi:10.1016/S0092-8674(00)80172-5

Lemaitre B, Reichhart JM, Hoffmann JA. 1997. Drosophila host defense: differential
induction of antimicrobial peptide genes after infection by various classes of
microorganisms. *Proceedings of the National Academy of Sciences of the United*
States of America **94**:14614–9. doi:10.1073/pnas.94.26.14614

Leulier F, Parquet C, Pili-Floury S, Ryu J-H, Caroff M, Lee W-J, Mengin-Lecreulx D,
Lemaitre B. 2003. The Drosophila immune system detects bacteria through specific
peptidoglycan recognition. *Nat Immunol* **4**:478–484. doi:10.1038/ni922

Leulier F, Rodriguez A, Khush RS, Abrams JM, Lemaitre B. 2000. The Drosophila caspase
Dredd is required to resist gram-negative bacterial infection. *EMBO Rep* **1**:353–8.
doi:10.1093/embo-reports/kvd073

Li X, Rommelaere S, Kondo S, Lemaitre B. 2020. Renal Purge of Hemolymphatic Lipids
952   Prevents the Accumulation of ROS-Induced Inflammatory Oxidized Lipids and
953   Protects Drosophila from Tissue Damage. *Immunity* **52**:374-387.e6.
954   doi:10.1016/j.immuni.2020.01.008
955   Liegeois S, Ferrandon D. 2022. Sensing microbial infections in the Drosophila melanogaster
956   genetic model organism. *Immunogenetics* **74**:35–62. doi:10.1007/s00251-021-01239-0
957   Ligoxygakis P, Pelte N, Ji C, Leclerc V, Duvic B, Belvin M, Jiang H, Hoffmann JA,
958   Reichhart JM. 2002. A serpin mutant links Toll activation to melanization in the host
959   defense of Drosophila. *Embo J* **21**:6330–7.
960   Ligoxygakis P, Roth S, Reichhart J-M. 2003. A Serpin Regulates Dorsal-Ventral Axis
961   Formation in the Drosophila Embryo. *Current Biology* **13**:2097–2102.
962   doi:10.1016/j.cub.2003.10.062
963   Lindsay SA, Lin SJH, Wasserman SA. 2018. Short-Form Bomanins Mediate Humoral
964   Immunity in Drosophila. *J Innate Immun* **10**:306–314. doi:10.1159/000489831
965   Marra A, Hanson MA, Kondo S, Erkosar B, Lemaitre B. 2021. Drosophila Antimicrobial
966   Peptides and Lysozymes Regulate Gut Microbiota Composition and Abundance. *mBio*
967   e0082421. doi:10.1128/mBio.00824-21
968   Martins NE, Faria VG, Teixeira L, Magalhães S, Sucena É. 2013. Host Adaptation Is
969   Contingent upon the Infection Route Taken by Pathogens. *PLoS Pathogens*
970   **9**:e1003601. doi:10.1371/journal.ppat.1003601
971   Medzhitov R, Schneider DS, Soares MP. 2012. Disease Tolerance as a Defense Strategy.
972   *Science* **335**:936–941. doi:10.1126/science.1214935
973   Melcarne C. 2020. Role of Nimrod receptors in the Drosophila melanogaster cellular immune
974   response. EPFL. doi:10.5075/epfl-thesis-9933
975   Melcarne C., Lemaitre B, Kurant E. 2019. Phagocytosis in Drosophila: From molecules and
976   cellular machinery to physiology. *Insect Biochemistry and Molecular Biology* **109**:1–
977   12. doi:10.1016/j.ibmb.2019.04.002
978   Melcarne C, Ramond E, Dudzic J, Bretscher A, Kurucz E, Andó I, Lemaitre B. 2019. Two
979   Nimrod receptors, NimC1 and Eater, synergistically contribute to bacterial
980   phagocytosis in *Drosophila melanogaster*. *The FEBS Journal*. doi:10.1111/febs.14857
981   Nakano S, Kashio S, Nishimura K, Takeishi A, Kosakamoto H, Obata F, Kuranaga E,
982   Chihara T, Yamauchi Y, Isobe T, Miura M. 2023. Damage sensing mediated by serine
983   proteases Hayan and Persephone for Toll pathway activation in apoptosis-deficient
984   flies. *PLoS Genet* **19**:e1010761. doi:10.1371/journal.pgen.1010761
985   Nakhleh J, El Moussawi L, Osta MA. 2017. The Melanization Response in Insect
986   ImmunityAdvances in Insect Physiology. Elsevier. pp. 83–109.
987   doi:10.1016/bs.aiip.2016.11.002
988   Nappi A, Poirié M, Carton Y. 2009. The role of melanization and cytotoxic by-products in the
989   cellular immune responses of Drosophila against parasitic wasps. *Adv Parasitol*
990   **70**:99–121. doi:10.1016/S0065-308X(09)70004-1
991   Needham AJ, Kibart M, Crossley H, Ingham PW, Foster SJ. 2004. Drosophila melanogaster
992   as a model host for Staphylococcus aureus infection. *Microbiology* **150**:2347–2355.
993   doi:10.1099/mic.0.27116-0
Nehme NT, Liégeois S, Kele B, Giammarinaro P, Pradel E, Hoffmann JA, Ewbank JJ,

Ferrandon D. 2007. A Model of Bacterial Intestinal Infections in Drosophila
melanogaster. *PLoS Pathogens* **3**:e173. doi:10.1371/journal.ppat.0030173

Nehme NT, Quintin J, Cho JH, Lee J, Lafarge M-C, Kocks C, Ferrandon D. 2011. Relative
Roles of the Cellular and Humoral Responses in the Drosophila Host Defense against
Three Gram-Positive Bacterial Infections. *PLOS ONE* **6**:e14743.
doi:10.1371/journal.pone.0014743

Pradeu T, Thomma BPHJ, Girardin SE, Lemaitre B. 2024. The conceptual foundations of
innate immunity: Taking stock 30 years later. *Immunity* 613–31.

Quintin J, Asmar J, Matskevich AA, Lafarge M-C, Ferrandon D. 2013. The Drosophila Toll
Pathway Controls but Does Not Clear Candida glabrata Infections. *The Journal of*
Immunology **190**:2818–2827. doi:10.4049/jimmunol.1201861

Rämet M, Lanot R, Zachary D, Manfruelli P. 2002. JNK Signaling Pathway Is Required for
Efficient Wound Healing in Drosophila. *Developmental Biology* **241**:145–156.
doi:10.1006/dbio.2001.0502

Ramond E, Jamet A, Ding X, Euphrasie D, Bouvier C, Lallemant L, He X, Arbibe L,
Coureuil M, Charbit A. 2021. Reactive Oxygen Species-Dependent Innate Immune
Mechanisms Control Methicillin-Resistant Staphylococcus aureus Virulence in the
Drosophila Larval Model. *mBio* **12**:e00276-21. doi:10.1128/mBio.00276-21

Rommelaere S, Carboni A, Bada Juarez JF, Boquete J-P, Abriata LA, Meireles FTP, Rukes
V, Vincent C, Kondo S, Dionne MS, Dal Peraro M, Cao C, Lemaitre B. 2024. A
humoral stress response protects *Drosophila* tissues from antimicrobial peptides.
Current Biology 1–12. doi:10.1101/2023.07.24.550293

Rutschmann S, Kilinc A, Ferrandon D. 2002. Cutting edge: the toll pathway is required for
resistance to gram- positive bacterial infections in Drosophila. *J Immunol* **168**:1542–6.

Ryder E, Blows F, Ashburner M, Bautista-Llacer R, Coulson D, Drummond J, Webster J,
Gubb D, Gunton N, Johnson G, O'Kane CJ, Huen D, Sharma P, Asztalos Z, Baisch H,
Schulze J, Kube M, Kittlaus K, Reuter G, Maroy P, Szidonya J, Rasmuson-Lestander
\AAsa, Ekström K, Dickson B, Hugentobler C, Stocker H, Hafen E, Lepesant JA,
Pflugfelder G, Heisenberg M, Mechler B, Serras F, Corominas M, Schneuwly S, Preat
T, Roote J, Russell S. 2004. The DrosDel collection: A set of P-element insertions for
generating custom chromosomal aberrations in Drosophila melanogaster. *Genetics*
167:797–813. doi:10.1534/genetics.104.026658

Sansone CL, Cohen J, Yasunaga A, Xu J, Osborn G, Subramanian H, Gold B, Buchon N,
Cherry S. 2015. Microbiota-dependent priming of antiviral intestinal immunity in
Drosophila. *Cell host & microbe* **18**:571–581.

Schmid-Hempel P. 2021. Evolutionary parasitology: the integrated study of infections,
immunology, ecology, and genetics. Oxford University Press.

Shan T, Wang Y, Bhattarai K, Jiang H. 2023. An evolutionarily conserved serine protease
network mediates melanization and Toll activation in *Drosophila*. *Sci Adv*
9:eadk2756. doi:10.1126/sciadv.adk2756

Stephenson HN, Streeck R, Grüblinger F, Goosmann C, Herzig A. 2022. Hemocytes are
essential for *Drosophila melanogaster* post-embryonic development, independent of
control of the microbiota. *Development* **149**:dev200286. doi:10.1242/dev.200286

Touré H, Durand N, Guénal I, Herrmann J-L, Girard-Misguich F, Szuplewski S. 2023.

Mycobacterium abscessus Opsonization Allows an Escape from the Defensin

Bactericidal Action in *Drosophila*. *Microbiol Spectr* e00777-23.

doi:10.1128/spectrum.00777-23

Ulvila J, Vanha-Aho L-M, Rämet M. 2011. Drosophila phagocytosis - still many unknowns

under the surface. *APMIS* **119**:651–662. doi:10.1111/j.1600-0463.2011.02792.x

Unckless RL, Howick VM, Lazzaro BP. 2016. Convergent Balancing Selection on an

Antimicrobial Peptide in Drosophila. *Current Biology* **26**:257–262.

doi:10.1016/j.cub.2015.11.063

Uttenweiler-Joseph S, Moniatte M, Lagueux M, Van Dorsselaer A, Hoffmann JA, Bulet P.

1998. Differential display of peptides induced during the immune response of

Drosophila: a matrix-assisted laser desorption ionization time-of- flight mass

spectrometry study. *Proc Natl Acad Sci U S A* **95**:11342–7.

Vaz F, Kounatidis I, Covas G, Parton RM, Harkiolaki M, Davis I, Filipe SR, Ligoxygakis P.

2019. Accessibility to Peptidoglycan Is Important for the Recognition of Gram-

Positive Bacteria in Drosophila. *Cell Reports* **27**:2480-2492.e6.

doi:10.1016/j.celrep.2019.04.103

Westlake H, Hanson MA, Lemaitre B. 2024. The Drosophila immunity handbook, First. ed.

EPFL Press. doi:10.55430/6304TDIHVA01

Wukitch AM, Lawrence MM, Satriale FP, Patel A, Ginder GM, Van Beek EJ, Gilani O,

Chambers MC. 2023. Impact of Chronic Infection on Resistance and Tolerance to

Secondary Infection in Drosophila melanogaster. *Infect Immun* e00360-22.

doi:10.1128/iai.00360-22

Xu R, Lou Y, Tidu A, Bulet P, Heinekamp T, Martin F, Brakhage A, Li Z, Liégeois S,

Ferrandon D. 2023. The Toll pathway mediates Drosophila resilience to Aspergillus

mycotoxins through specific Bomanins. *EMBO Rep* **24**:e56036.

doi:10.15252/embr.202256036

Yasin B, Wang W, Pang M, Cheshenko N, Hong T, Waring AJ, Herold BC, Wagar EA,

Lehrer RI. 2004. θ Defensins Protect Cells from Infection by Herpes Simplex Virus by

Inhibiting Viral Adhesion and Entry. *Journal of Virology* **78**:5147–5156.

doi:10.1128/jvi.78.10.5147-5156.2004

1069

1070 **Figure Legends**

1071 **Table 1: List of mutants used in this study.** "-" indicates the deleted immune module.

1072

Figure 1: Each immune module functions largely independently from other immune

modules. **A)** Activation of the Imd pathway, as revealed by *DptA* expression following

infection with a mixture of heat-killed *E. coli* and *M. luteus,* is broadly wild-type (WT) in single

module mutants other than Imd. In the case of *ΔPhag* flies, we observed a lesser induction at

6h compared to wild-type. **B)** Activation of the Toll pathway as revealed by the expression of

Drs is broadly wild-type in single module mutants other than *ΔTOLL* flies. Note that *Drs*

receives a minor input from Imd signaling (Leulier et al., 2000), explaining minor induction of

Drs in *ΔTOLL* flies at early time points. In the case of *ΔMel*, we observed a slightly greater Toll

pathway activity. **C)** The ability of plasmatocytes to phagocytose bacterial particles is not

negatively affected in single module mutants except *ΔPhag* (also see Figure 1supp1) **D)** Cuticle blackening after clean injury is not impaired except in *ΔMel* flies. **E)** Activation of the Imd pathway remains strongly inducible in compound mutants except when deficient for the Imd pathway. **F)** Activation of the Toll pathway remains strongly inducible in compound mutants except when flies were deficient of the Toll pathway. **G)** The ability of plasmatocytes to phagocytose bacterial particles is not significantly affected in compound mutants except those including *ΔPhag*. **H)** Cuticle blackening after clean injury is not impaired in compound mutants except in those including *ΔMel*.

Figure 2: Lifespans of mutants used in this study in unchallenged flies and upon clean injury conditions at 25°C. A-B) Single module mutants; **C-D)** compound module mutants. See Figure 2supp1 for 29°C comparisons.

Figure 3: Heatmap of lifespans of immune module-deficient flies upon infection by various pathogens. Darker blue indicates lower survival, while white indicates maximum survival. Experiments used either 7 or 10 days as a maximum lifespan/time course, and the heatmap is adjusted accordingly per row to have white fill for the maximum possible lifespan of that row. Small text to right of mean lifespan indicates total flies. Survival curves are presented in **Supplementary file 1**, and a summary of susceptibilities is provided in Figure 3supp1.

Figure 4: Growth kinetics of *P. rettgeri* (A), *S. aureus* (B), and *C. albicans* (C) in wild-type, single module mutant and *ΔITPM* flies. Survival curves from **Supplementary file 1** underlying data in **Figure 3** are shown for context. Each data point reflects a pooled sample of 5 flies.

Figure 5: Measurement of PLUD upon infection with *P. rettgeri*, *S. aureus*, and *C. albicans*. A) The PLUD of *P. rettgeri* in individual module mutant flies is not significantly different from wild-type. The PLUD of *ΔITPM* flies was not significantly different, although censoring of a single low-PLUD outlier in the wild-type would result in a significant difference between wild-type and *ΔITPM* flies ($p < .01$). **B)** The PLUD of *S. aureus*-infected flies is not different across any genotype. **C)** The PLUD of *C. albicans*-infected flies was significantly lower in *ΔPhag* flies with a notably lower mean PLUD. This difference was robust to use of different *ΔPhag* genetic backgrounds (merged data shown here). *ΔMel* flies and *ΔITPM* flies also had significantly lower PLUD. *, $P < 0.05$; **, $P < 0.01$

Table S1: Microbe characteristics and growth conditions.

Figure 1supp1: We used *ΔPhag* mutants in the genetic background of Melcarne et al. (2019) *(+; ΔPhag)* in some experiments, and confirm here that these mutants are equally deficient in phagocytic capacity to *ΔPhag* flies in the iso DrosDel genetic background (*ΔPhag)*.

Figure 2supp1: Lifespans of mutants used in this study in unchallenged flies and upon clean injury conditions at 29°C. **A-B)** Single module mutants; **C-D)** compound module mutants.

1125

1126 **Supplementary file 1:** Survival curves for all data summarized in **Figure 3**.

1127

Figure 3supp1: Summary table of susceptibilities per the three conditions outlined in section
"Systemic infections and survival." ✓ indicates that the pathway contributes to defense against
a given pathogen. In the event that a single module mutant is not more susceptible to the
infection, but contributes to increased mortality in a double mutant line, a ✓ is given and the
co-deleted relevant pathways highlighted by their first letter. Black boxes indicate no difference
to wild-type or to clean injury. NI, natural infection; SI, septic infection.

Drosophila lines	Reference	Name	Deficient pathway			
			IMD	Toll	Phagocytosis	Melanization
iso ; iso ; Rel^{E20}	Hedengren et al., 1999	ΔIMD	-	+	+	+
iso ; iso ; spz^{rm7}	Lemaitre et al., 1996	$\Delta TOLL$	+	-	+	+
iso ; NimC1^{1} ; Eater^{1}	Melcarne et al., 2019	$\Delta Phag$	+	+	-	+
iso ; PPO1Δ, PPO2Δ; iso	Bingelli et al., 2014	ΔMel	+	+	+	-
+ ; + ; Rel^{E20}, spz^{rm7}	this paper	$\Delta IMD, \Delta TOLL$	-	-	+	+
iso ; PPO1Δ, PPO2Δ ; NimC1^{1},Eater^{1}	this paper	$\Delta Phag, \Delta Mel$	+	+	-	-
iso ; NimC1^{1} ; Eater^{1}, Rel^{E20}	this paper	$\Delta IMD, \Delta Phag$	-	+	-	+
iso ; PPO1Δ PPO2Δ ; Rel^{E20}	this paper	$\Delta IMD, \Delta Mel$	-	+	+	-
iso ; PPO1Δ PPO2Δ ; spz^{rm7}	this paper	$\Delta TOLL, \Delta Mel^{PPS}$	+	-	+	-
iso Hay-psh^{Def} ; iso ; iso	Dudzic et al., 2019	$\Delta TOLL, \Delta Mel^{HP}$	+	-	+	-
iso Hay-psh^{Def} ; NimC1^{1} ; Eater^{1} Rel^{E20}	this paper	$\Delta ITPM$	-	-	-	-

Table 1

Figure 1

Figure 2

Type	Species	Temp	Dose	Days	N exp	WT	ΔAMP14	ΔBom	ΔIMD	ΔTOLL	ΔPhag	ΔMel	ΔPhag, ΔMel	ΔTOLL, ΔMel^{PPS}	ΔTOLL, ΔMel^{HP}	ΔIMD, ΔMel	ΔIMD, ΔPhag	ΔIMD, ΔTOLL	ΔITPM
Controls	Unchallenged	25°C	n/a	7	13	6.9 (177)	6.9 (60)	6.9 (60)	6.8 (60)	6.8 (80)	7.0 (60)	6.6 (60)	6.6 (60)	7.0 (116)	6.8 (239)	6.0 (60)	6.6 (60)	6.8 (100)	5.8 (80)
	Clean Injury	25°C	n/a	7	36	6.7 (709)	6.6 (60)	6.5 (60)	6.7 (543)	6.6 (260)	6.8 (248)	6.4 (526)	5.3 (57)	4.6 (71)	5.4 (70)	4.7 (67)	6.5 (60)	6.2 (60)	5.5 (445)
	Clean Injury	29°C	n/a	7	11	6.6 (211)	6.5 (60)	5.9 (60)	6.3 (100)	6.1 (75)	6.6 (60)	6.4 (119)	4.9 (96)	3.0 (67)	4.8 (56)	3.7 (57)	6.0 (64)	5.6 (60)	3.9 (94)
	Clean Injury	25°C	n/a	10	36	9.6 (709)	9.1 (60)	9.2 (60)	9.2 (543)	9.3 (260)	9.6 (248)	8.5 (526)	6.2 (57)	5.2 (71)	6.3 (70)	5.6 (67)	9.1 (60)	8.7 (60)	6.3 (445)
	Unchallenged	29°C	n/a	10	11	9.9 (139)	9.3 (60)	9.5 (60)	9.4 (49)	7.1 (34)	8.7 (60)	8.5 (60)	9.1 (60)	9.2 (60)	9.3 (40)	5.9 (64)	9.3 (40)	8.9 (60)	7.7 (61)
Viruses	Drosophila C virus	25°C	2000 TCID50	10	5	8.3 (100)	6.7 (89)	7.7 (85)	6.8 (73)	7.8 (83)	8.1 (78)	6.3 (86)	6.3 (88)	5.5 (56)	5.0 (92)	5.1 (60)	6.7 (64)	3.8 (53)	4.1 (78)
	Flock House virus	25°C	250000 TCID50	10	5	8.5 (122)	7.1 (73)	7.2 (84)	7.3 (79)	7.3 (77)	7.3 (81)	6.9 (82)	6.2 (82)	5.1 (56)	6.0 (85)	4.9 (61)	6.8 (73)	5.2 (59)	5.1 (84)
	Drosophila X virus	25°C	431000 TCID50	10	2	9.0 (40)	6.9 (41)	8.2 (20)	7.1 (45)	7.9 (34)	7.8 (41)	5.4 (38)	7.4 (46)	4.9 (57)	5.1 (38)	6.0 (15)	6.2 (40)	5.6 (19)	4.6 (43)
	Invertebrate iridescent-6 virus	25°C	31700 TCID50	10	3	9.4 (59)	7.5 (50)	9.4 (64)	8.1 (58)	8.4 (54)	8.8 (60)	5.6 (58)	6.0 (61)	5.8 (59)	5.8 (58)	6.1 (48)	8.3 (54)	5.9 (47)	4.4 (69)
	Sindbis virus	25°C	50000 TCID50	10	3	9.4 (58)	8.4 (61)	8.6 (61)	8.9 (60)	8.8 (60)	8.3 (60)	5.4 (65)	5.9 (66)	4.9 (51)	4.8 (59)	4.5 (60)	7.0 (50)	4.5 (45)	4.3 (61)
Fungi	Aspergillus fumigatus (natural)	29°C	rolled on plate	10	5	7.0 (80)	6.8 (81)	7.0 (20)	6.8 (56)	6.5 (81)	6.8 (41)	6.7 (60)	6.5 (60)	5.7 (62)	6.5 (60)	6.0 (60)	6.5 (66)	5.9 (60)	4.5 (92)
	Beauvaria bassiana (natural)	29°C	30mg	10	5	9.6 (120)	8.2 (61)	9.7 (60)	9.4 (60)	7.3 (70)	7.8 (43)	8.7 (76)	7.7 (60)	6.3 (64)	6.0 (36)	7.3 (66)	7.1 (40)	6.5 (51)	4.4 (62)
	Beauvaria bassiana	29°C	OD=5	7	6	6.4 (81)	6.0 (74)	3.6 (59)	5.7 (58)	3.6 (66)	5.2 (39)	5.3 (62)	4.4 (57)	2.6 (63)	3.4 (63)	4.4 (21)	4.7 (51)	3.1 (56)	3.3 (95)
	Candida albicans	29°C	OD=200	7	8	6.7 (160)	5.6 (45)	6.0 (102)	5.9 (94)	5.0 (89)	6.4 (65)	5.9 (60)	6.4 (102)	1.7 (20)	3.1 (156)	5.7 (36)	5.9 (89)	3.2 (27)	2.4 (82)
Gram positive bacteria	Mycobacterium marinum	25°C	OD=1	7	6	6.1 (132)	6.0 (50)	6.5 (51)	6.5 (111)	5.9 (108)	5.2 (77)	5.8 (99)	5.7 (108)	4.1 (62)	5.8 (84)	5.1 (51)	5.1 (64)	6.3 (56)	3.9 (117)
	Corynebacterium diphteriae	25°C	OD=100	7	3	6.0 (73)	6.2 (84)	6.5 (97)	5.6 (79)	5.1 (80)	6.6 (87)	4.6 (82)	5.8 (69)	3.2 (47)	3.3 (103)	4.9 (72)	5.9 (88)	4.5 (61)	3.0 (75)
	Micrococcus luteus	29°C	OD=200	7	8	6.7 (121)	5.7 (80)	6.8 (60)	6.2 (120)	6.4 (148)	6.5 (71)	6.5 (81)	5.6 (81)	4.6 (68)	6.5 (80)	4.8 (77)	6.3 (84)	2.1 (102)	2.1 (91)
	Streptococcus pneumoniae	25°C	OD=0.3	7	3	6.8 (41)	6.1 (60)	6.4 (60)	6.7 (61)	6.6 (70)	6.3 (60)	6.2 (60)	6.4 (60)	5.7 (60)	5.0 (61)	5.1 (60)	6.0 (60)	5.5 (60)	4.5 (50)
	Enterococcus faecalis	25°C	OD=5	7	10	5.0 (130)	4.5 (100)	2.1 (62)	4.2 (122)	1.8 (126)	5.2 (79)	5.1 (122)	3.9 (76)	1.6 (51)	1.8 (102)	4.2 (62)	4.0 (79)	1.4 (61)	1.3 (101)
	Staphylococcus aureus	25°C	OD=0.5	7	8	5.1 (119)	4.8 (58)	5.7 (21)	4.2 (118)	3.4 (73)	3.6 (79)	2.4 (73)	1.9 (78)	2.0 (80)	2.0 (112)	2.1 (78)	4.1 (59)	3.4 (40)	1.9 (62)
	Listeria monocytogenes	25°C	OD=0.5	7	5	4.8 (91)	4.7 (78)	2.2 (60)	2.3 (62)	1.6 (112)	4.6 (71)	5.3 (76)	3.9 (78)	2.1 (70)	2.2 (73)	3.5 (88)	3.2 (60)	2.0 (62)	1.6 (62)
	Bacillus subtilis	25°C	OD=5	7	7	5.7 (120)	3.3 (40)	4.1 (20)	4.0 (123)	3.8 (119)	6.0 (55)	5.4 (110)	4.5 (98)	2.8 (69)	2.7 (84)	4.5 (76)	5.4 (20)	2.1 (41)	2.0 (81)
Gram negative bacteria	Vibrio parahemolyticus	25°C	OD=10	7	2	3.1 (58)	1.1 (61)	2.9 (62)	1.1 (62)	1.3 (70)	2.8 (64)	2.3 (70)	3.2 (57)	1.8 (68)	1.9 (41)	1.0 (59)	1.1 (63)	1.0 (58)	1.0 (64)
	Providencia rettgeri	25°C	OD=0.1	7	7	5.5 (146)	1.1 (61)	5.1 (58)	2.0 (78)	4.4 (70)	3.6 (100)	1.8 (95)	1.5 (61)	1.5 (75)	5.3 (40)	1.3 (75)	2.1 (62)	1.0 (21)	1.4 (123)
	Providencia burhodogranaria	25°C	OD=10	7	7	5.1 (161)	1.7 (76)	4.6 (40)	1.6 (58)	4.6 (88)	4.9 (63)	4.0 (124)	4.2 (84)	2.9 (61)	4.9 (60)	1.3 (78)	1.0 (61)	1.4 (40)	1.3 (68)
	Pectobacterium carotovorum Ecc15	29°C	OD=10	7	7	6.4 (131)	1.5 (57)	4.8 (20)	1.9 (91)	5.8 (108)	4.4 (120)	5.8 (100)	5.2 (102)	4.7 (60)	5.2 (60)	1.8 (72)	- (-)	1.4 (62)	1.4 (106)
	Klebsiella pneumoniae	25°C	OD=1	7	3	6.8 (60)	1.4 (68)	6.5 (60)	1.5 (77)	6.8 (60)	6.9 (60)	5.8 (60)	6.3 (60)	3.9 (80)	6.9 (60)	1.3 (59)	2.0 (56)	- (-)	1.3 (64)
	Enterobacter cloacae	25°C	OD=10	7	8	6.6 (149)	1.3 (83)	6.7 (60)	1.0 (100)	6.6 (90)	6.7 (60)	6.4 (60)	6.1 (80)	4.9 (61)	5.8 (99)	1.3 (73)	1.0 (37)	1.6 (61)	1.2 (83)
	Escherichia coli	25°C	OD=200	7	8	6.6 (141)	5.0 (79)	6.9 (40)	3.8 (60)	6.2 (70)	5.4 (60)	5.5 (43)	5.5 (85)	3.9 (63)	5.3 (63)	3.8 (60)	4.6	1.7 (70)	1.6 (60)
	Salmonella enterica ser. typhimurium	25°C	OD=200	7	4	6.7 (80)	4.8 (40)	6.6 (80)	5.1 (60)	6.3 (40)	6.7 (20)	6.3 (40)	6.6 (63)	5.5 (40)	4.8 (64)	4.7 (61)	3.9 (18)	2.7 (56)	1.8 (68)

Figure 3

Figure 4

Figure 5

Type	Kingdom	Phylum	Class	Order	Family	Nucleic acid / Membrane	Species	Strain, Source, DOI of previous study	Inf. mode	Infection temperature	Used concentration	Growth temperature	Medium of culture
Viruses	Bamfordvirae	Nucleocytoviricota	Megaviricetes	Pimascovirales	Iridoviridae	dsDNA+	*Invertebrate iridescent-6 virus*	gift from Ronald van Rij	intra-thoracique	25°C	31 700 TCID50	provided	provided
	Orthornavirae	nd	nd	nd	Bimaviridae	ssRNA+	*Drosophila X virus*	gift from Ronald van Rij	intra-thoracique	25°C	431 000 TCID50	provided	provided
	Orthornavirae	Pisuviricota	Pisoniviricetes	Picornavirales	Dicistroviridae	ssRNA+	*Drosophila C virus*	gift from Carla Saleh	intra-thoracique	25°C	2 000 TCID50	26°C	S2 cells in Schneider
	Orthornavirae	Kitrinoviricota	Magsaviricetes	Nodamuvirales	Nodaviridae	dsRNA+	*Flock House virus*	gift from Carla Saleh	intra-thoracique	25°C	250 000 TCID50	26°C	S2 cells in Schneider
	Orthornavirae	Kitrinoviricota	Alsuviricetes	Martelliirales	Togaviridae	ssRNA+	*Sindbis virus*	gift from Ronald van Rij	intra-thoracique	25°C	50 000 TCID50	provided	provided
Fungi	Fungi	Ascomycota	Sacharomycete	Sacharomycetales	Sacharomycetacae	*Beta-glucans*	**Candida albicans**	ATCC 2001	septic	29°C	OD_{600}= 200	37°C	YPG
		Ascomycota	Eurotiomycete	Eurotiales	Trychocomacae	*Beta-glucans*	**Aspergillus fumigatus**	*doi: 10.7554/eLife.44341*	natural	29°C	rolled on plate	37°C	Malt-agar
		Ascomycota	Sordaryomycete	Coricypitacae	Ophiocordycipitacae	*Beta-glucans*	**Beauvaria bassiana**		septic	29°C	OD_{600}= 5	commercial	commercial
						Beta-glucans		R444	natural	29°C	30mg	commercial	commercial
Gram positive bacteria	Bacteria	Actinomycetota	Actinomycetia	Mycobacteriales	Mycobacteriaceae	mAGP-type	**Mycobacterium marinum**	1218R WT	intra-thoracique	25°C	OD_{600}= 1	37°C	LB
		Actinomycetota	Actinomycetia	Mycobacteriales	Corynebacteriaceae	Lys-type	**Corynebacterium diphtheriae**	*DSM 44123*	septic	25°C	OD_{600}= 100	37°C	LB
		Actinomycetota	Actinomycetia	Actinomycetales	Micrococcacae	Lys-type	**Micrococcus luteus**	*doi: 10.7554/eLife.44341*	septic	25°C	OD_{600}= 200	29°C	LB
		Bacillota	Bacilli	Lactobacilliales	Streptococcacae	Lys-type	**Streptococcus pneumoniae**	D39V WT	septic	25°C	OD_{600}= 0.3	37°C	LB
		Bacillota	Bacilli	Lactobacilliales	Enterococcacae	Lys-type	**Entecoccus faecalis**	254	septic	25°C	OD_{600}= 5	37°C	LB
		Bacillota	Bacilli	Caryophanale	Staphylococcacae	Lys-type	**Staphylococcus aureus**	*doi: 10.7554/eLife.44341*	septic	25°C	OD_{600}= 0.5	37°C	BHI
		Bacillota	Bacilli	Bacilliales	Listeriacae	DAP-type	**Listeria monocytogenes**	BUG 1600	septic	25°C	OD_{600}= 0.5	37°C	LB
		Bacillota	Bacilli	Bacilliales	Bacillacae	DAP-type	**Bacillus subtilis**	168	septic	25°C	OD_{600}= 5	37°C	BHI
Gram negative bacteria	Bacteria	Pseudomonadota	Gammaproteobacteriacae	Pseudomonadales	Pseudomonadacae	DAP-type	**Pseudomonas aeruginosa**	PAO1	septic	25°C	OD_{600}= 0.01	37°C	LB
		Pseudomonadota	Gammaproteobacteriacae	Vibrionales	Vibrionacae	DAP-type	**Vibrio parahemolyticus**		septic	25°C	OD_{600}= 10	37°C	LB
		Pseudomonadota	Gammaproteobacteriacae	Enterobacteriales	Morgenallacae	DAP-type	**Providencia rettgeri**	Dmel	septic	25°C	OD_{600}= 0.1	37°C	LB
		Pseudomonadota	Gammaproteobacteriacae	Enterobacteriales	Morgenallacae	DAP-type	**Providencia burhodogranaria**	B	septic	25°C	OD_{600}= 10	37°C	LB
		Pseudomonadota	Gammaproteobacteriacae	Enterobacteriales	Enterobacteriacae	DAP-type	**Pectobacterium carotovorum**	Ecc15	septic	25°C	OD_{600}= 10	37°C	LB
		Pseudomonadota	Gammaproteobacteriacae	Enterobacteriales	Enterobacteriacae	DAP-type	**Klebsiella pneumoniae**		septic	25°C	OD_{600}= 1.0	37°C	LB
		Pseudomonadota	Gammaproteobacteriacae	Enterobacteriales	Enterobacteriacae	DAP-type	**Enterobacter cloacae**	B12	septic	25°C	OD_{600}= 10	37°C	LB
		Pseudomonadota	Gammaproteobacteriacae	Enterobacteriales	Enterobacteriacae	DAP-type	**Escherichia coli**	1106	septic	25°C	OD_{600}= 200	37°C	LB
		Pseudomonadota	Gammaproteobacteriacae	Enterobacteriales	Enterobacteriacae	DAP-type	**Salmonella enterica ser. typhimurium**	*doi: 10.7554/eLife.44341*	septic	25°C	OD_{600}= 200	37°C	LB

Table S1

Figure 1supp1

Figure 2supp1

WT

ΔAMP14

Bom^{Δ55C}

ΔIMD

ΔTOLL

ΔPhag

ΔMel

ΔPhag, ΔMel

ΔTOLL, ΔMel^{PPS}

ΔTOLL, ΔMel^{HP}

ΔIMD, ΔMel

ΔIMD, ΔPhag

ΔIMD, ΔTOLL

ΔITPM

Figure 3supp1

	Microbe	AMP14	Bom	IMD	TOLL	Phag	Mel
Viruses	DCV	✓		✓	(✓ I, M)		✓
	FHV	✓	✓	✓	✓	✓	✓
	DXV	✓		✓	✓	✓	✓
	IIV-6	✓		✓	(✓ I)		✓
	SINV			(✓ T, P)	(✓ I)	✓	✓
Fungi	*A. fumigatus (NI)*			(✓ T)	(✓ I, M)		(✓ T)
	B. bassiana (NI)	✓		(✓ P)	(✓ I, M)	(✓ I, M)	(✓ T, P)
	B. bassiana (SI)		✓		✓	✓	
	C. albicans (SI)	✓		(✓ T)	✓		(✓ T)
Gram positive bacteria	*M. marinum*						
	C. diphteriae			(✓ T)	(✓ I, M)		✓
	M. luteus			(✓ T)	(✓ I)		
	S. pneumoniae						
	E. faecalis		✓		✓	(✓ M)	(✓ P)
	S. aureus				✓	✓	✓
	L. monocytogenes		✓	✓	✓		
	B. subtilis	✓	✓	✓	✓		(✓ T)
Gram negative bacteria	*V. parahemolyticus*	✓		✓	✓		
	P. rettgeri	✓		✓	✓	✓	✓
	P. burhodogranaria	✓		✓	(✓ M)		✓
	Ecc15	✓	✓	✓		✓	
	K. pneumoniae	✓		✓			
	E. cloacae	✓		✓			
	E. coli	✓		✓	(✓ I)	✓	
	S. enterica typhimurium	✓		✓	(✓ I)	(✓ I)	

Supplementary file 1
Fungi

Beauveria bassiana
OD600 = 5, 29ºC

○	*WT*
△	*ΔAMP14*
+	*ΔBom*
✳	*ΔIMD*
◇	*ΔTOLL*
▽	*ΔPhag*
⊠	*ΔMel*
✱	*ΔPhag, ΔMel*
◈	*ΔTOLL, ΔMel^{PPS}*
⊕	*ΔTOLL, ΔMel^{HP}*
⊗	*ΔIMD, ΔMel*
⊞	*ΔIMD, ΔPhag*
⊠	*ΔIMD, ΔTOLL*
⊟	*ΔITPM*

Candida albicans
OD600 = 200, 29ºC

○	*WT*
△	*ΔAMP14*
+	*ΔBom*
✳	*ΔIMD*
◇	*ΔTOLL*
▽	*ΔPhag*
⊠	*ΔMel*
✱	*ΔPhag, ΔMel*
◈	*ΔTOLL, ΔMel^{PPS}*
⊕	*ΔTOLL, ΔMel^{HP}*
⊗	*ΔIMD, ΔMel*
⊞	*ΔIMD, ΔPhag*
⊠	*ΔIMD, ΔTOLL*
⊟	*ΔITPM*

Gram-negative bacteria

Enterobacter cloacae
OD600 = 10, 25°C

Escherichia coli
OD600 = 200, 25°C

Klebsiella pneumoniae
OD600 = 1.0, 25°C

Pectobacterium carotovorum
OD600 = 10, 25°C

Providencia burhodogranariea
OD600 = 10, 25°C

Legend:
- WT
- ΔAMP14
- ΔBom
- ΔIMD
- ΔTOLL
- ΔPhag
- ΔMel
- ΔPhag, ΔMel
- ΔTOLL, ΔMel^{PPS}
- ΔTOLL, ΔMel^{HP}
- ΔIMD, ΔMel
- ΔIMD, ΔPhag
- ΔIMD, ΔTOLL
- ΔITPM

Providencia rettgeri
OD600 = 0.1, 25°C

Legend:
- WT
- ΔAMP14
- ΔBom
- ΔIMD
- ΔTOLL
- ΔPhag
- ΔMel
- ΔPhag, ΔMel
- ΔTOLL, ΔMel^{PPS}
- ΔTOLL, ΔMel^{HP}
- ΔIMD, ΔMel
- ΔIMD, ΔPhag
- ΔIMD, ΔTOLL
- ΔITPM

Salmonella typhimurium
OD600 = 200, 25ºC

Legend:
- WT
- ΔAMP14
- ΔBom
- ΔIMD
- ΔTOLL
- ΔPhag
- ΔMel
- ΔPhag, ΔMel
- ΔTOLL, ΔMel^{PPS}
- ΔTOLL, ΔMel^{HP}
- ΔIMD, ΔMel
- ΔIMD, ΔPhag
- ΔIMD, ΔTOLL
- ΔITPM

Vibrio parahemolyticus
OD600 = 10, 25ºC

Gram-positive bacteria

Enterococcus faecalis
OD600 = 5, 25°C

Listeria monocytogenes
OD600 = 0.5, 25°C

Micrococcus luteus
OD600 = 200, 25°C

WT	
ΔAMP14	
ΔBom	
ΔIMD	
ΔTOLL	
ΔPhag	
ΔMel	
ΔPhag, ΔMel	
ΔTOLL, ΔMel^{PPS}	
ΔTOLL, ΔMel^{HP}	
ΔIMD, ΔMel	
ΔIMD, ΔPhag	
ΔIMD, ΔTOLL	
ΔITPM	

Mycobacterium marinum
OD600 = 1, 25°C

WT	
ΔAMP14	
ΔBom	
ΔIMD	
ΔTOLL	
ΔPhag	
ΔMel	
ΔPhag, ΔMel	
ΔTOLL, ΔMel^{PPS}	
ΔTOLL, ΔMel^{HP}	
ΔIMD, ΔMel	
ΔIMD, ΔPhag	
ΔIMD, ΔTOLL	
ΔITPM	

Staphylococcus aureus
OD600 = 0.5, 25°C

Legend:
- WT
- ΔAMP14
- ΔBom
- ΔIMD
- ΔTOLL
- ΔPhag
- ΔMel
- ΔPhag, ΔMel
- ΔTOLL, ΔMel^{PPS}
- ΔTOLL, ΔMel^{HP}
- ΔIMD, ΔMel
- ΔIMD, ΔPhag
- ΔIMD, ΔTOLL
- ΔITPM

Streptococcus pneumoniae
OD600 = 0.3, 25°C

Legend:
- WT
- ΔAMP14
- ΔBom
- ΔIMD
- ΔTOLL
- ΔPhag
- ΔMel
- ΔPhag, ΔMel
- ΔTOLL, ΔMel^{PPS}
- ΔTOLL, ΔMel^{HP}
- ΔIMD, ΔMel
- ΔIMD, ΔPhag
- ΔIMD, ΔTOLL
- ΔITPM

Viruses

Drosophila C virus, 25ºC

Drosophila X virus, 25ºC

Flock house virus, 25°C

─○─	*WT*
─△─	*ΔAMP14*
─+─	*ΔBom*
─✕─	*ΔIMD*
─◇─	*ΔTOLL*
─▽─	*ΔPhag*
─⊠─	*ΔMel*
─✳─	*ΔPhag, ΔMel*
─◈─	*ΔTOLL, ΔMel^{PPS}*
─⊕─	*ΔTOLL, ΔMel^{HP}*
─✶─	*ΔIMD, ΔMel*
─⊞─	*ΔIMD, ΔPhag*
─⊠─	*ΔIMD, ΔTOLL*
─⊟─	*ΔITPM*

Invertebrate iridescent-6 virus, 25°C

─○─	*WT*
─△─	*ΔAMP14*
─+─	*ΔBom*
─✕─	*ΔIMD*
─◇─	*ΔTOLL*
─▽─	*ΔPhag*
─⊠─	*ΔMel*
─✳─	*ΔPhag, ΔMel*
─◈─	*ΔTOLL, ΔMel^{PPS}*
─⊕─	*ΔTOLL, ΔMel^{HP}*
─✶─	*ΔIMD, ΔMel*
─⊞─	*ΔIMD, ΔPhag*
─⊠─	*ΔIMD, ΔTOLL*
─⊟─	*ΔITPM*

Sindbis virus, 25°C