ECHOES OF 1816: MICROBIAL FOOTPRINTS IN HERITAGE ARTIFACTS FROM ARGENTINA'S MUSEUM OF INDEPENDENCE

Daniel Gonzalo Alonso-Reyes[1,2]; Fátima Silvina Galván[1,2]; Natalia Noelia Alvarado[1,2], María Cecilia D´Arpino[1,3], Luciano José Martinez[1], Hernán José Esquivel[1], Cecilia Aymara Gallardo[1], María Julia Silva Manco[1,2], Virginia Helena Albarracín[1, 2, 3,4]*

[1]Laboratorio de Microbiología Ultraestructural y Molecular, Centro Integral de Microscopía Electrónica (CIME), Universidad Nacional de Tucumán. CONICET NOA SUR, CONICET, Tucumán, Argentina.

[2]Facultad de Agronomía, Zootecnia y Veterinaria, Universidad Nacional de Tucumán, Tucumán, Argentina.

[3]Instituto de Salud y Calidad de Vida, Universidad de San Pablo-Tucumán.

[4]Facultad de Ciencias Naturales e Instituto Miguel Lillo, Universidad Nacional de Tucumán, Tucumán, Argentina.

Running headline: MICROBIAL BIODIVERSITY OF CASA HISTORICA MUSEUM

Keywords: BIODIVERSITY, CULTURAL HERITAGE, MUSEUM, EXTREMOPHILES, ELECTRON MICROSCOPY, TUCUMAN

*Corresponding author:

Virginia Helena Albarracín, Centro Integral de Microscopía Electrónica (CIME, CONICET, UNT) Camino de Sirga s/n. FAZyV, Finca El Manantial, Yerba Buena (4107). Tucumán, Argentina. E-mail: cime@conicet.gov.ar

ABSTRACT

Historical artifacts preserved in museums are invaluable cultural treasures, yet they are often vulnerable to biodeterioration caused by microbial colonization. This study presents the first comprehensive investigation of microbial communities inhabiting heritage artifacts from Argentina's Museum of Independence. By integrating advanced microscopy with phenotypic and genomic characterization, we analyzed samples collected from wooden objects, textiles, architectural elements, and exterior walls. Scanning electron microscopy revealed diverse and well-structured biofilms, with intricate three-dimensional arrangements embedded in extracellular polymeric substances. A total of 49 bacterial strains were isolated and identified via VITEK MALDI-TOF mass spectrometry, with a predominance of Gram-positive genera such as *Bacillus*, *Micrococcus*, and *Kocuria*. Remarkably, the 19th-century albumen print photograph emerged as the most biodiverse artifact, yielding 21 distinct strains, including *Streptomyces, Oceanobacillus, and thermophilic Caldibacillus thermoamylovorans.* The protein-rich and halophilic environment of the albumen layer likely facilitated microbial colonization and persistence. Pseudomonas species were exclusively associated with this photographic substrate, further underscoring its niche specificity. Human-associated taxa like *Staphylococcus epidermidis* and *Staphylococcus equorum* were prevalent in high-contact areas, while exterior surfaces displayed distinct microbial signatures, including potential pathogens linked to environmental exposure. These findings reveal a complex and artifact-specific microbial landscape, emphasizing the need for tailored, bio-informed conservation strategies. This work advances the field of heritage microbiology and supports efforts to safeguard culturally significant objects through a deeper understanding of their microbial ecosystems.

INTRODUCTION

The National Museum "Casa Histórica de la Independencia" (CHM), located in San Miguel de Tucumán, Argentina, holds a vast and culturally significant historical collection from the 19th and 20th centuries. The CHM is one of the most visited historical museums in the country (Chatruc, 2021) and is renowned as the site where the independence of the United Provinces of the Río de la Plata was declared on July 9, 1816 — a date later recognized as Argentine Independence Day.

The colonial mansion housing the Museum was built in 1760 and has undergone modifications over time (Marinsalda, 2015). While the original building was demolished in 1874, its 20th-century reconstruction has solidified its status as a national symbol of unity. The "Salón de la Jura" or Hall of the Oath, where the Declaration of Independence was redacted, remains in its original state. Apart from historical documents and furniture, the Museum conserves an extensive book library and a photographic archive. Among the highlights is Provincia de Tucumán by Arsenio Granillo, a 213-page book (Granillo, 1872) commissioned by Governor Federico Helguera, features descriptive articles and news about Tucumán, accompanied by original albumen print photographs depicting the city and its sugar mills. Notably, the book includes a unique photograph of the CHM's original façade, considered an invaluable contribution to Argentine documentary photography and essential for the building's restoration in 1943.

Museums serve a fundamental role in the conservation of cultural heritage (CH), safeguarding irreplaceable artifacts that embody humanity's identity, memory, and history. Preventive conservation is vital because cultural assets are inherently vulnerable to deterioration processes that threaten their integrity (Gutarowska et al., 2014; Grabek-Lejko

et al., 2017; Negi and Sarethy, 2019). Microbiological contamination is a common issue, as many artifacts are of organic origin and prone to colonization by bacteria and fungi. These microorganisms thrive in museum environments, adapting their versatile metabolisms to withstand extreme conditions, including fluctuations in temperature, humidity, and pH. Indeed, microbes may have been present in the original piece since its first production, as invisible part of the materials used to craft the artifact. Microbes then settled into specific ecological niches, often remaining undetected until visible biofilm formation, discoloration, or structural weakening became apparent. Their activity can cause significant damage to textiles, wood, paper, and other organic materials, manifesting as discoloration, structural weakening, odor, and cracking (Skóra et al., 2015; Grabek-Lejko et al., 2017; Negi and Sarethy, 2019). The consequences of microbial growth extend beyond artifact damage, potentially diminishing monetary value, and bringing about costly decontamination procedures (Sterflinger and Piñar, 2013). Despite these challenges, the cultural and historical value of many CH objects is immeasurable and cannot be quantified solely in monetary terms.

Biodeterioration is defined as "any undesirable change in a material brought about by the vital activities of organisms"(Perito and Cavalieri, 2018). Although abiotic processes were historically viewed as the primary cause of CH decay, the role of microorganisms has gained recognition only recently. While biodeterioration cannot be entirely halted, advances in microbial ecology and technological interventions offer strategies to mitigate its effects. A deeper understanding of microbial communities on CH objects is critical for identifying their deteriorative mechanisms and implementing effective control measures (Perito and Cavalieri, 2018). These aspects have become an important research field for both,

conservators and microbiologists (Adamiak et al., 2018). This issue also raises occupational

health concerns, as museum staff may be exposed to harmful biological agents during the

handling of contaminated materials, such as paper, wood, and textiles (Borrego et al., 2010;

Gutarowska et al., 2014). Indeed, many studies indicated that conservators, librarians, and

archivists face risks related to microbial exposure (Zielińska-Jankiewicz et al., 2008;

Wiszniewska et al., 2009; Karbowska-Berent et al., 2011).

Traditional methods of investigating microorganisms on CH objects have primarily

been culture-dependent, offering insights into their deteriorative roles. Recently, amplicon

sequencing has emerged as a powerful yet resource-intensive tool for microbial

identification (Kraková et al., 2012a, 2018). An alternative approach combines the

automated VITEK system for rapid biochemical identification (Książczyk et al., 2016),

with whole-genome sequencing (WGS) for functional profiling of key strains. Scanning

electron microscopy (SEM) has also proven effective in biodeterioration studies, providing

visual insights into microbial activity on CH surfaces (Díaz-Herráiz, 2015; Puškárová et al.,

2016; Elhagrassy, 2018; Liu et al., 2018).

To better understand the role of microbial communities in the biodeterioration of

cultural heritage artifacts, we employed a multidisciplinary approach integrating advanced

microscopy, culture-dependent techniques, and genomic analyses. Scanning electron

microscopy (SEM) was chosen to visualize microbial colonization and biofilm structures

on historical surfaces at high resolution, while culture-based methods allowed us to isolate

and characterize viable bacterial strains. The application of MALDI-TOF mass

spectrometry enabled rapid and accurate taxonomic identification, and whole-genome

sequencing provided insights into the functional potential of selected isolates, particularly

their capacity for biodegradation and resistance to environmental stressors. This

multidisciplinary approach allows us to analyze the microbial communities in detail,

revealing their influence on the artifacts' preservation and contributing to informed

conservation practices.

2. MATERIALS AND METHODS

2.1. Selection of heritage artifacts of the Museum.

The CHM is located in San Miguel de Tucumán, the capital of Tucumán Province,

Argentina at a pedestrian zone with significant foot traffic. (https://sketchfab.com/3d-

models/museo-nacional-casa-historica-5b37f88491c0456bbd15fe87f4b8b171). The Casa

Histórica is a preserved colonial-style building and functions as a museum, offering guided

tours and exhibitions related to Argentina's independence and the revolutionary era. The

museum features four courtyards, a cistern, and native trees from the region. In 1941, the

house was declared a National Historic Monument.

CHM currently holds over 700 items in its collection. This includes the original Act

of the Declaration of Independence and various other pieces such as period furniture,

weapons, personal items, dishware, religious artifacts, paintings, portraits, coins, and

commemorative plaques and medals from the 18th and 19th centuries. The museum's library

contains classic texts, recent historiographical debates, museum studies, conservation,

restoration, literature, journals and early provincial newspapers. With over 500 volumes,

the collection covers diverse topics such as ecclesiastical history, the Wars of

Independence, civil wars, art, the sugar industry, economics, travel narratives, Jesuit

chronicles, and journalism, with a focus on the province of Tucumán. Before sampling, an

extensive survey and photographic documentation were conducted to capture the condition

of cultural heritage items affected by different types of deterioration. This preliminary

assessment informed the selection of specific sampling locations within the building and on

key artifacts of cultural significance.

The initial phase of sampling took place in August 2019. Samples were taken from

the museum's exterior, including facade elements like the front door and painted walls (Fig.

1A), which are subjected to annual repainting, weathering, and urban pollution; they were

thus considered a control zone related to outdoor urban microbiome (OUM). The front was

built as part of the last reconstruction in 1943 including the wooden door initially coated

with flax oil. Within the museum, samples were taken from the window and furniture in

the "Salón de la Jura". Additional historical artifacts (points 2.2 and 2.3) that required

preservation evaluations were also sampled (Table 1, Fig. 1). All these samples were

considered as specimens of the indoors museum microbiome (IMM).

2.2. Salón de la Jura

The Salón de la Jura or Hall of the Oath (SJ) is a central room within the CHM (Fig.

1B). This site holds historical significance as it is where the Congress of Tucumán

proclaimed Argentina's independence from Spanish rule on July 9, 1816. The hall is

renowned for its role as the setting of intense debates and the eventual drafting of the "Acta

de la Independencia," marking the foundation of the Argentine nation.SJ is the only room

of the house that has been conserved intact since then. The room features period furnishings

and artifacts that transport visitors back to the momentous occasion when representatives

from the United Provinces of the Río de la Plata gathered to declare sovereignty. Samples

for our study were collected from one chair where congressmen sat and from the table

where the Declaration of Independence was signed; this furniture remains in its original

state, dating back to 1810.The wooden frame from the armrest and backrest and the velvet-

covered seat of one of the chairs were sampled to assess the microbial communities present

on these distinct materials (Fig. 1C-D). The table's wooden surface was swabbed in

multiple locations, especially in the areas less exposed to regular cleaning, to try to sample

the original microbiome in the furniture (Fig. 1E). Additionally, we took samples from

biodeteriored zones of one of the windows of the room, subjected to infestation by

woodworm eggs (1F-G).

2.3. Child Suit from Juan Bautista Alberdi, the father of Argentinean Constitution.

Juan Bautista Alberdi (1810-1884) was a distinguished statesman, diplomat, and

writer who played a pivotal role in shaping Argentina's political landscape. Born in San

Miguel de Tucumán, he was known for his influential work in drafting the Argentine

Constitution of 1853. Thus, Alberdi's legacy as a thinker and leader has left a lasting

impact on the country's history and development.

The Museum holds a captivating collection of artifacts attributed to Alberdi,

including an item that offers a rare and personal look at 19[th]-century life: a childhood suit

made from natural red silk (Catalog Number: CH321; 53 cm x 29 cm; sewing technique

manufactured in Argentina [Fig. 1H-I]), This garment reflects the fashion of adult male

clothing of the period, characterized by its fabric, design, and color. The suit comprises two

main pieces: a jacket featuring a draped front center with a faux belt detail and a pleated

back skirt; the trousers are adorned with decorative lace and bows at the side hems. These

garments serve as a tangible connection to Alberdi's early life and provide a unique

perspective on the sartorial and social customs of the era. Its historical significance is enriched by its association with one of Argentina's key historical figures, illustrating how clothing can encapsulate cultural identity and societal norms. The Museum has employed preventive conservation strategies, including controlled humidity, light, and temperature, to maintain the suit's condition and ensure its longevity. https://sketchfab.com/3d-models/traje-de-juan-baustista-alberdi-037b9c3badd54d21b330a3d25e2b91d3

2.4. An iconic photograph on the Historical Book *Provincia de Tucumán*

The second phase of sampling in December 2021 targeted a key artifact from the CHM: the 1872 book *Provincia de Tucumán. Serie de artículos descriptivos y noticiosos*, authored by Dr. Arsenio Granillo (Fig. 1J-L). Commissioned by Governor Federico Helguera, this work is historically notable for its pioneering inclusion of albumen prints captured by Italian-born photographer Ángel Paganelli. These early photographs, which include a depiction of the Casa Histórica façade, provide rare insights into Tucumán's 19th-century landscapes and urban spaces.

Stored in a metallic safe case to preserve its condition, this book has survived with minimal light exposure, thermal fluctuations, and limited handling. Of the estimated 300 original copies printed, only about 100 featured the albumen prints, and few remain today—primarily in the Casa Histórica and private collections.

The albumen prints stand out both for their historical value and the meticulous craftsmanship involved in their creation. Paganelli used large glass negatives (approximately 15 x 12 cm) to produce contact prints on albumen-coated paper. This egg-white and sodium chloride mixture served as a binder for silver salts, enabling high-quality

image transfer. The labor-intensive process involved exposure to sunlight under negatives,

toning, fixing, and washing the images (Reilly, 1980) before affixing them to book pages

by hand. Each of the 100 photograph-included books contained 21 images, totaling over

2,100 individually prepared prints—a remarkable testament to 19th-century photographic

artistry.

Due to their production method, we considered albumen prints as halophilic

microenvironments, which may influence microbial colonization. The unique

environmental conditions within the book highlight the delicate balance required for

preserving such historical artifacts while exploring their microbiological significance.

2.5. Sampling procedures for direct material observation and bacterial isolation.

To conduct the sampling, a non-invasive approach was employed, using precision

tools to extract material samples without compromising the structural integrity of the

pieces. Micro-sampling techniques, supported by visual documentation and environmental

controls, were used to assess the physical condition of the albumen prints and the rest of the

elements identify potential areas of degradation, such as yellowing, flaking, or damage

from previous handling.

For the direct observation of microorganisms on heritage objects, freshly-open

3M™ adhesive tape was used (Michaelsen et al., 2009). Briefly, tape was cut into strips (2-

3 cm), subjected to 15-min sterilization under UV-C. Subsequently, the strips were gently

adhered to the different surfaces and then transferred to 2 ml tubes containing Karnovsky's

fixative solution (2.66% w/v paraformaldehyde and 1.66% w/v glutaraldehyde in 0.1 M

phosphate buffer, pH 7.2), for transportation to the lab. The tape sampling method was not

used for the book of Granillo, to avoid any undesired consequences over its conservation.

For isolation of individual bacterial strains, sample collection was performed with

dry and sterilized cotton swabs. The swab was passed over the entire area of visibly

damaged material; each swab covered approximately a 10 cm² area to standardize sample

collection. Immediately after sampling, the swabs were used to inoculate plates with Luria-

Bertani (LB) agar media. Parallel agar plates were supplemented with nalidixic acid (10 µg

mL-1) and cycloheximide (10 µg mL-1) to inhibit the growth of Gram-negative bacteria

and fungi, respectively (Rasuk et al., 2017). This non-invasive method is considered

essential for the study of collections in museum environments (Michaelsen et al., 2009;

Piñar et al., 2015).Once the colonies appeared, they were collected with a sterile loop for

purification and subsequent streaking allows generating axenic cultures. We stored the cells

in glycerol 20% at -20°C. Subcultures were performed in LB broth for morphological and

molecular identification analysis. From the pure cultures of the bacterial isolates,

macroscopic practices were carried out to determine the structure of the colonies and their

pigmentation, and microscopic practices (optical and electronic) of individual cells to

establish shapes, sizes, and spatial arrangement. Additionally, the Gram reactions of all

isolates were recorded.

All isolated strains in this work are kept and curated in the CEBAC collection

(Cepario de Bacterias Ambientales del CIME), Catalogue Urban Microbiome Strains, under

the corresponding codes indicated in Table 1.

2.6. Scanning Electron Microscopy (SEM)

For SEM studies, the adhesive tapes or pure axenic cultures of isolated strains were

fixed in a mixture of aldehydes (Karnovsky fixative) overnight at 4ºC, and were prepared

following the processing method already previously described in Alonso-Reyes et al

(2021). During post-fixation, samples were washed with 0.1 M phosphate buffer twice and

dehydrated in a graded concentration series of ethanol (50, 70, 90, and 100%), following

acetone 100%. This was followed by critical point drying (Denton Vacuum model DCP-1).

The samples were then mounted on aluminum supports and subjected to gold sputtering

(JEOL model JFC-1100) and examined using a Zeiss Supra 55VP scanning electron

microscope (Carl Zeiss NTS GmbH, Germany), in the Electron Microscopy Core Facility

(CIME-CONICET-UNT).

2.7. Identification of microbial strains by MALDI-TOF MS (VITEK)

MALDI-TOF analysis was performed with a bioMérieux VITEK MALDI-TOF mass

spectrometer (bioMérieux, France), and the spectra were compared to the VITEK MS

SARAMIS database for research use only (RUO), version 4.09, using the SuperSpectra

algorithm (referred to here as MALDI-TOF MS). For MALDI-TOF MS analysis, a colony

from a pure plate with 24h growth was applied to a slide and covered with 1 µl of matrix

solution (α-cyano-4-hydroxycinnamic acid) and dried completely before analysis. The

identification was considered definitive when the probability provided was greater than

70%. This service was provided by Laboratorios CIBIC in Rosario, Argentina.

2.8. DNA extraction

DNA from eight isolated strains was obtained using a commercial extraction kit (DNA

Puriprep B-kit for bacteria, Inbio Highway, Argentina); following the manufacturer's

instructions. For Gram-positive bacteria, the pellet was suspended in 162 µl of BRB buffer

and 18 µl of lysozyme, and then incubated for 30 min at 37°C. For gram-negative bacteria,

the use of lysozyme was omitted. Next, 200 µl of BT buffer and 20 ul of proteinase K were

added to each sample. It was incubated for 30 min at 56°C. After, RNaseA (10 mg/ml), was

added and incubated for 30 min at 37°C. Then the lysis buffer (200 µl) was added, vortex

pulsed for 10 sec and the homogenate was incubated for 10 min at 56°C. Then, 200 µl of

100% ethanol was added to each sample, shaken by inversion, and transferred to a

microcolumn. It was centrifuged for 1 min at 12,000 g, and the filtrate was separated. Then,

two column washes were performed each with 500 µl of Blav1 and Blav2 buffer

respectively. Finally, the column was dried by centrifuging for 3 min at 12,000 g. DNA

elution was carried out with 60 µl of pH 9 elution buffer, equilibrated at 70°C. DNA

concentration and quality were measured using a µDrop plate (Thermo Scientific).

2.9. Genome sequencing, assembly, and analysis.

Whole genome sequencing service was performed by Novogene UK services

(https://www.novogene.com/amea-en/) using the IlluminaNovaseq platform with pair-end

150 strategy. Quality control of the reads was also performed by Novogene. The Illumina

reads were assembled using SPAdes assembler v. 3.15.4 (Bankevich et al., 2012), with --

careful and --cov-cutoff auto options. The resulting eight assemblies were uploaded to the

NCBI database under the Bioproject accession number PRJNA1233209, and are also

available at Zenodo (https://doi.org/10.5281/zenodo.15169295). The functional annotation

was performed using Arche v. 1.0.1 (Alonso-Reyes and Albarracín, 2022) pipeline with --

kegg option, and the resultant KEGG identifiers (KO) were mapped through the KEGG

mapper reconstruct tool (https://www.genome.jp/kegg/mapper/reconstruct.html).   The

genomic potential for the synthesis of secondary metabolites was elucidated through

antiSMASH v. 7.1.0 (Blin et al., 2023). To estimate the taxonomic identity of each strain at

genus level, 16s-rDNA and DNA gyrase B genes were retrieved from the genomes using

barrnap and blast tools, and then blasted against the NCBI NR database. For complete taxa

identification, average nucleotide identity (ANI) analysis was performed using pyani

(Pritchard et al., 2016), against genus-specific genomes deposited in NCBI database.

3. RESULTS

3.1. SEM analysis of microbial communities on heritage objects and façade surfaces.

Biofilms on heritage surfaces including furniture, wooden objects (chair, table, window

and washing trough) and textile garments were analyzed using scanning electron

microscopy (Fig. 2). The freshly painted building façade, entrance door and exterior walls,

were considered as control samples of the OUM for comparison with the IMM. In all cases,

the adhesive tape sampling method proved to be a useful, simple, and non-destructive

technique for collecting microbial communities for immediate imaging analysis.

Interestingly, the surfaces of objects within and outside the museum were highly colonized

with microbes forming well-structured biofilms.

Most observed biofilms exhibited three-dimensional cellular organization and were

surrounded by substantial amounts of extracellular material. For instance, the wooden chair

showed spindle or hazelnut-shaped bacterial cells embedded in an abundant

exopolysaccharide (EPS) matrix (Fig. 2A). The wooden table displayed a homogeneous

bacterial biofilm with coccoid units connected by thin, short filaments (Fig. 2B). In the

wooden washing trough, compact coconut-shaped biofilms were observed attached to

wooden rests (Fig. 2C), while forming an interconnected by extracellular filaments or

vesicles. Other areas revealed a filamentous bacterial mantle (Fig. 2D) likely belonging to

the genus *Streptomyces* or related Actinobacteria. On garments, microbial filaments were

much scarce (Fig. 2G-2H); they were adhered tightly to deteriorate fibers of the clothing,

with fibrous materials and deposits visible at higher magnifications. In this sample, diverse

fungi spores were also found. In the window sample, we observed a biofilm formed by

heterogeneous microbial community dominated by large cocci and compacted bacilli, with

minimal extracellular material facilitating cell interactions (Fig. 2I).

The façade also showed high colonization of the surfaces by biofilms; the entrance door

exhibited a thick biofilm structure composed of cocci, dividing bacilli, larger rough-

surfaced bacilli, and filamentous bacteria, all forming strongly interconnected aggregates

(Fig. 2J). The exterior walls demonstrated consolidated coccoid biofilms held together by

EPS, likely offering protection against environmental factors, while other regions showed

sparse adherence of bacilli in simple distributions.

3.2. VITEK-MS-based Identification of Isolated Bacteria

A total of 49 bacterial strains were isolated from various samples (Table 1, Fig. 3-

4), with 12 strains attributed to the OUM and 37 strains linked to the IMM. Each sampled

niche exhibited a distinct profile of bacterial taxa, reflecting the contrasting conditions of

outdoor and indoor environments. Moreover, each artifact sampled from within the Casa

Histórica de la Independencia can be regarded as an isolated microbial island, shaped by its

unique environmental conditions, material composition, age, usage and historical context.

The microbial communities inhabiting these artifacts form a complex and dynamic

microecosystem, where each niche harbors a distinctive collection of bacteria specifically

adapted to its substrate (Fig. 3). Gram-positive bacteria dominated the isolates, as expected

given the use of selective isolation media supplemented with antibiotics to exclude fungi

and gram-negative bacteria. Nevertheless, five strains of *Pseudomonas* were successfully

isolated under these same conditions from the Granillo Book—one from the paper and four

from the photograph itself. Indeed, the most prolific source of isolates was the albumen

photograph, from which 21 distinct strains were recovered, predominantly from the genera

Bacillus, Oceanobacillus, and Streptomyces. In contrast, the Alberdi´s garment gave a

unique isolate: *Actinomyces odontolyticus*. Morphological characteristics of each strain

(texture, color, spore production) were recorded to differentiate isolates (Table 1).

Four strains were isolated from the textile upholstery of a chair (*Micrococcus*

luteus, Staphylococcus equorum, Bacillus licheniformis and Bacillus simplex) while three

strains were found on the wooden frame of the same chair (*Micrococcus* sp., *Kocuria* sp.

and one unidentified strain). Additional 15 isolates were recovered from other wooden

elements, including the SJ table (*Kocuria rosea, Micrococcus luteus, Bacillus*

altitudinis/pumilus), the SJ window frame (*Micrococcus luteus, Microbacterium aurum* and

one unidentified strain), the museum's entrance door (*Microbacterium aurum*, *Bacillus*

altitudinis/pumilus, *Clostridium subterminale*, *Micrococcus luteus*, *Corynebacterium*

pseudodiphtheriticum, Kocuria rosea and one unidentified strain) and a washing trough

(*Staphylococcus equorum* and *Bacillus* sp.) where meat of slaughter animals was prepared.

Finally, five strains were isolated from the front wall of the building made of white painted

bricks (*Corynebacterium pseudodiphtheriticum, Kocuria rosea, Turicella otitidis, Bacillus*

megaterium, Kocuria rosea).

3.3. Genome-based taxonomical identification.

The genomes of eight representative strains were sequenced and analyzed for

functional characterization. Three of them correspond to bacterial isolates from the front

and will be considered as representatives of the OUM while the rest are considered

representatives of the IMM (Table 2). The sequencing process and quality control yielded

approximately 100 million reads and 15 Gb of data, with an average of 12 million reads and

1.88 Gb per genome. Assembly resulted in fewer than 115 contigs for all strains, except for

the CH-041 strain, which had 200 contigs. The G + C content varied between strains,

ranging from ~35% in CH-031 to ~72% in CH-021. The ARCHE pipeline, employing

GeneMarkS-2 for gene prediction, achieved high annotation rates for protein-coding

sequences (CDS). The lowest annotation rate was observed in CH-026, with ~85% of CDS

annotated, while CH-036 reached approximately 97% annotation.

Species identification, initially conducted using the automated VITEK system

(Książczyk et al., 2016) was supplemented with 16S-rDNA and DNA gyrase B marker

genes (MGs) against NCBI databases. The gyrB gene was used when the 16S-rDNA gene

was incomplete or truncated. ANI analyses were performed using genus-specific genomes

from the NCBI database for more precise identification (Table 2).

VITEK identification results for most strains differed from those obtained through

MGs. Only strains CH-041 and CH-015 maintained the same species classification by both

methods. CH-036 was identified as belonging to the genus *Streptomyces* by VITEK, MGs,

and ANI, but the species differed between methods. VITEK accurately classified CH-031 at

the family level (Bacillaceae), but MGs identified it at the genus level as *Bacillus*, whereas

ANI suggested *Oceanobacillus*. CH-021 and CH-026 only matched at the class level

(Actinomycetes) by both methods, while CH-019 matched at the phylum level (Bacillota).

CH-007 could not be classified by VITEK but was identified as *Microbacterium* by MGs

and ANI. Significant differences were evident in species-level classifications; MGs and

ANI agreed on the species for CH-041 (*Caldibacillus thermoamylovorans*) and CH-015

(*Microbacterium aurum*). In CH-031, where MGs did not yield a species classification,

ANI clustered it with *Oceanobacillus kimchi* due to >95% similarity to a reference genome

in the NCBI database (Richter and Rosselló-Móra, 2009). Notably, CH-007, and CH-026

did not align with any reference genome above the 95% ANI threshold, suggesting they

might belong to novel species.

Pairwise comparisons of the CHM genomes with public database genomes revealed

the closest matches. Most strains shared >97% similarity, except for CH-026 (87.11%) and

CH-007 (87.82%). For example, CH-031 had a 99.47% sequence identity with

Oceanobacillus kimchii p3-SID1558 (Accession No. NZ_JALXUQ000000000), isolated

from human back skin (Hannah et al., 2023). CH-019 showed a 99.32% identity with

Bhargavaea massiliensis Marseille-Q1000 (Accession No. NZ_JAHBCJ000000000), from

a urine sample of patients with urinary tract infections (Olowo-okere et al., 2021). CH-021

matched with *Kocuria* sp. CD08_4 (Accession No. NZ_LQBK00000000) at 98.71%, an

isolate from the duodenal mucosa of a celiac disease patient, which carries genes linked to

virulence and oxidative stress resistance (Munish et al., 2017). As expected, CH-036

matched a *Streptomyces* strain with 98.48% of similarity, *Streptomyces* sp. f150, which was

isolated from soil at Florida, USA (Accession No. NTHG00000000.1). CH-015 showed a

97.98% similarity with the metagenome-assembled genome (MAG) of *Microbacterium* sp.

DE14.009 (Accession No. JACXUU000000000.1) from geothermal springs in Greece. CH-

041's genome matched *Caldibacillus thermoamylovorans* MSL 185.1 (Accession No.

NZ_JAMAXM000000000), a thermophile isolated from a spacecraft environment, capable

of enduring heat-shock at 80°C for 15 minutes (Tran et al., 2023). In contrast, CH-007 and

CH-026 were related at a distance to *Microbacterium arborescens* RCB1 (87.82%

similarity; Accession No. QNUW00000000) and *Pseudoclavibacter helvolus* PS02339

(87.11% similarity; Accession No.NZ_JASCYB000000000), respectively, and do not meet

the 95% threshold for known species. Genetic comparisons indicated approximately 12%

divergence from their closest known relatives in the NCBI database, indicating that both

may be novel species.

3.4. Functional Annotation of Genomes: Insights into Biodeterioration and

Environmental Resilience

The functional annotation carried out using ARCHE led to useful information

regarding the genomic potential of the strains (Fig. 5.A-E, Supplementary Tables S1-15).

Genes involved in resistances to antibiotics, osmotic stress, and heavy metals were reported

from the annotation. Furthermore, relevant genes related to the biodeterioration of the

cultural heritage were found, such as those required for the degradation of wood and paper,

and the production of pigments. *Streptomyces* sp. CH-036 revealed the highest numbers of

genes in all the systems mentioned before; except for heavy metal resistance (the numbers

are still relatively high). Moreover, CH-036 reported more than twice as many genes

related to biodeterioration (Fig. 5.D-E) as the rest of the strains studied. Particularly, those

genes related to the degradation of cellulose and production of pigments, which are

potentially harmful to paper documents, were highly abundant in CH-036. *Kocuria* sp.

CH-021 strain displayed a high potential to resist both the osmotic and heavy metal

stresses. A relatively elevated number of cation transporters were annotated in CH-021's

genome, and genes related to copper and chromate homeostasis were also noticeable. Genes

for the degradation of cellulose and hemicellulose are also present, together with the

capability to produce some pigments. On the other hand, both *Ocenobacillus* sp. CH-031

and *Caldibacillus* sp. CH-041 had a reduced biodeteriorative potential; a tiny group of

genes devoted to pigment production and degradation of wood and paper were found (none

of the latter are related to cellulose degradation). Finally, CH-041 showed a clear heavy

metal resistance profile considering the coding-sequences found in the genome.

Further functional capabilities of relevance were studied, such as the potential for

the synthesis of secondary metabolites, and degradation of xenobiotics (Fig. 5.F). The

results of the screening with antiSMASH revealed *Streptomyces* sp. CH-036 as the most

diverse of the strains regarding to the hosting of biosynthetic gene clusters (BGCs) devoted

to the production of secondary metabolites. Among the putative products of these BCGs are

lanthipeptides, and ε-poly-L-lysines, both compounds with reported antimicrobial

properties (Repka et al., 2017; Wang et al., 2021). Other compounds reported from CH-036

genome were desferrioxamin (Holden and Nair, 2019), and isorenieratene, which is a light-

induced yellow pigment (Takano et al., 2005). On the other hand, KEGG identifiers

retrieved from the annotation of *Kocuria* sp. CH-021 genome revealed several complete

pathways for xenobiotic degradation, including those for the degradation of benzoate,

phenylacetate, xylene, and toluene.

4.   DISCUSSION

4.1. Microbial Niches in the Museum and Their Impact on Heritage Conservation

This study revealed significant microbial colonization on historical artifacts and

architectural surfaces, highlighting both the conservation challenges and the potential

scientific value of these microbial communities. SEM analysis of historical artifacts showed

extensive biofilm development in most samples. In some cases, like the Alberdi´s garment,

it was observed tightly adhered microbial filaments to altered original material of the

pieces, providing direct evidence of biodeterioration processes that threaten cultural

heritage.

Microbial colonization in the museum environment is shaped by distinct ecological

niches, with significant differences observed between the outdoor urban microbiome

(OUM) and the indoor museum microbiome (IMM). The OUM, primarily represented by

microbial communities on the museum's façade, entrance door, and exterior walls, is

continuously exposed to fluctuating environmental conditions such as temperature

variations, UV radiation, precipitation, and air pollutants. This exposure selects for stress-

resistant taxa thriving under changing conditions, including endospore-forming *Bacillus*

spp., UV-resistant *Micrococcus luteus*, and environmental actinobacteria, which form

resilient biofilms. For instance, *Microbacterium aurum* is known for its environmental

robustness and ability to survive desiccation and nutrient limitations, while *Bacillus*

altitudinis/pumilus forms endospores that protect against UV radiation and other stressors.

The presence of species such as *Kocuria rosea*, known for pigment production and

environmental stress resistance, further reflects the selective pressures of urban niches.

Additionally, airborne contaminants and human activity in urban settings introduce

anthropogenic microbes, some of which may be opportunistic pathogens

(*Pseudoclavibacter* sp., *Kocuria* sp., *Bhargavaea* sp.) or biofilm producers that facilitate

475 biodeterioration (Stackebrandt et al., 1995; Bartlett et al., 2022; Zhou et al., 2022; Ahamad

476 et al., 2023). Furthermore, when ANI comparisons are used the genome of the strain CH-

477 019 shares a 99,32% of similarity with *Bhargavaea massiliensis* Marseille-Q1000 isolated

478 from a urine sample in patients with urinary tract infections (Table 2, Section 3.3).

479 Likewise, ANI showed a 98,71% of similarity between the genomes of CH-021 and

480 *Kocuria sp.* CD08_4 an isolate from the duodenal mucosa of a celiac disease patient.

481 Finally, CH-017 and CH-027 were both classified by VITEK as *B. pumilus* and *B.*

482 *megaterium* which have been reported as plant pathogens in other work (Bull et al., 2010;

483 Yuan and Gao, 2015).

484 In contrast, the IMM comprises microbial communities inhabiting historical

485 furniture, textiles, paper artifacts, and architectural surfaces within the controlled indoor

486 environment of the museum. These niches are characterized by relatively stable

487 temperature and humidity conditions, lower UV exposure, and reduced air circulation,

488 which favors specialized microbial assemblages. Here, bacterial communities may be

489 defined by the original organic and inorganic substrates of the artifact and the use given to

490 the museum piece. Wooden furniture, including the chair and table, was dominated by

491 *Bacillus pumilus*, *Micrococcus* sp., and *Kocuria rosea*, which are adapted to nutrient-

492 limited and stable conditions. Alberdi's suit yielded only one strain, *Actinomyces*

493 *odontolyticus*, a filamentous bacterium well-suited to organic substrates like fabric fibers

494 and related with dental infections. The albumen photograph exhibited exceptional diversity,

495 including *Caldibacillus thermoamylovorans*, *Oceanobacillus kimchii*, and *Streptomyces*

496 *rubiginosohelvolus*, showcasing the unique properties of albumen as a protein-rich

497 substrate conducive to microbial colonization. The dominance of *Actinomyces*,

Staphylococcus, and *Bacillus* species in the IMM suggests a strong association with human

contact, dust deposition, and material degradation. Notably, *Pseudomonas* spp., exclusively

isolated from the albumen photograph, may play a role in gelatin and paper

biodeterioration, highlighting the niche specificity of microbial colonization. These

findings emphasize the IMM's potential as a window into historical microbial communities

and their interactions with cultural heritage substrates, offering valuable insights into

microbial persistence and biodeterioration. Simultaneously, they highlight the need for

targeted conservation strategies to address microbial degradation while preserving the

historical significance of these microbial communities.

Indeed, it was pointed out the importance of regular disinfection routines, after

several *Bacillus* spp., *Staphylococcus* spp., and *Streptomyces* spp. with potentially

pathogenic characteristics were reported in museum collections (Gutarowska et al., 2014).

Furthermore, it has been suggested that some *Streptomyces* spp. present in museums could

be noxious to the human respiratory system (Borrego et al., 2010). In our work, we found

the strains CH-036 and CH-042 classified as *Streptomyces* spp. by different methods.

Surprisingly, the genome of CH-036 showed a strong antibiotic resistance profile of genes

(Fig. 5.A). It is thus important keep away threats to health from the CHM's staff and

visitors, especially those who are immunocompromised. Protection elements should be

used in the manipulation of museum artifacts. Also, we encourage the adoption of

appropriate measures to avoid contamination of the CHM's elements with new infectious

agents coming from the outside of the CHM's. Proof of the unprotected handling of the

book of albumen print is the presence of the commensal species *Staphylococcus epidermis*,

related to the human skin microbiome (Villalba et al., 2004). Also, a 99.47% of similarity

was found between the genomes of CH-031 and *Oceanobacillus kimchii* p3-SID1558

isolated from the human back skin (Table 2). In this case, as photographs were hand-crafted

prepared and then pasted in the book, an initial inoculation of the strain from the original

photographer ´skin microbiome cannot be rule out.

The interaction between OUM and IMM microbiomes is a critical yet

underexplored aspect of microbial dynamics in heritage conservation. Airborne microbial

transfer, visitor-mediated contamination, and ventilation systems could facilitate cross-

colonization between outdoor and indoor environments. Additionally, spores and biofilm

fragments from external surfaces may infiltrate indoor spaces, potentially influencing

microbial succession on artifacts. Understanding these interactions is essential for

designing targeted conservation strategies that mitigate microbial spread while preserving

the unique ecological stability of indoor heritage collections. Further studies incorporating

metagenomic and metatranscriptomic analyses could provide deeper insights into microbial

connectivity and functional roles in museum environments.

4.2. Photographic heritage as novel source for polyextremophilic bacteria

The albumen prints in the book Provincia de Tucumán represent a unique

microenvironment that may influence microbial colonization and biodeterioration

processes. Albumen, composed primarily of denatured egg white proteins, creates a

proteinaceous, nitrogen-rich substrate with high organic content, potentially favoring the

growth of microorganisms capable of metabolizing proteins. This composition makes

albumen prints particularly susceptible to colonization by proteolytic bacteria and fungi,

which may exploit albumen as a carbon and nitrogen source. Moreover, the high

hygroscopicity of albumen paper, resulting from both the organic matrix and the

hydrophilic nature of its gelatinous binder, creates microclimatic conditions conducive to

microbial proliferation. Fluctuations in relative humidity (RH) can lead to the absorption

and retention of moisture, facilitating the activation of dormant microbial spores and the

formation of biofilms. The presence of *Pseudomonas* spp., a known gelatin and protein

degrader, exclusively in albumen-coated prints, supports this hypothesis, suggesting that

this niche selects for microbial taxa with proteolytic capabilities.

Another key factor is the interaction between albumen, silver salts, and

environmental pollutants. Albumen prints were historically prepared with silver nitrate,

which reacts with the albumen matrix to form light-sensitive silver salts. While silver ions

possess antimicrobial properties, over time, their oxidized and degraded forms may lose

effectiveness, allowing microbial colonization in aged prints. Additionally, sulfur-

containing air pollutants (e.g., hydrogen sulfide) could interact with residual silver, forming

silver sulfide (Ag□S), which is less antimicrobial and contributes to the well-known

yellowing and fading of albumen photographs. We hypothesize that the combination of

albumen's organic composition, moisture retention, and silver-based chemical interactions

creates a selective microbial niche that differs from other paper-based heritage objects. The

albumen photograph provided a particularly compelling example of a microbial

microenvironment, serving as a protein-rich niche that supported a diverse microbial

consortium. Notably, isolates such as *Bacillus altitudinis/pumilus, Oceanobacillus kimchii,*

and *Streptomyces rubiginosohelvolus* were identified, all of which thrive in high-salt, low-

nutrient conditions, suggesting that albumen prints present unique biochemical properties

favorable for microbial colonization. The interaction between microorganisms and the

albumen layer, which contains organic compounds such as egg whites and silver salts, may

play a significant role in the biodeterioration of photographic materials, potentially leading

to discoloration, fading, or other forms of structural damage over time. Other strains

isolated from the CHM's photo book were classified by VITEK, MGs, and ANI as

Microccocus, *Staphylococcus* and *Kocuria* (Table 1, Fig. 3). These taxa were consistently

reported as microbial contamination in historical archives and audiovisual materials

(Villalba et al., 2004; Puškárová et al., 2016; Mazzoli et al., 2018; Branysova et al., 2023).

They have proven amylolytic, cellulolytic, proteolytic, and/or lipolytic capacities related

tobiodeterioration of paper, photographs, and/or other visual material which were mainly

determined by culture-dependent methods (Villalba et al., 2004; Branysova et al., 2023).

The amylolytic capacities can be used for the consumption of starch from paper (used as a

glue in its manufacturing) while the proteolytic activity presents a threat especially to visual

materials with a gelatinous binder. Some species with cellulolytic properties pose a risk to

collodion visual materials (as the main component of collodion is cellulose nitrate) while

others have the ability to produce alkaline serine proteases which are harmful to albumin

materials (Villalba et al., 2004; Branysova et al., 2023). Functional genomic analysis of

Streptomyces sp. CH-036 revealed strong capacities for biodeterioration, including

cellulose degradation and pigment biosynthesis, such as isorenieratene, a light-induced

pigment. These findings highlight the critical need for targeted conservation measures,

particularly avoiding exposure to light and maintaining stable environmental conditions to

prevent microbial proliferation. Further experimental validation through controlled

humidity exposure tests and protease activity assays would be necessary to confirm this

hypothesis and refine conservation strategies for photographic materials.

Notably, *Pseudomonas* strains were exclusively isolated from the photograph,

aligning with its role as a frequent contaminant in photographic manufacturing due to its

ability to metabolize gelatin and resist silver nitrate (Gadd et al., 1989; Borrego et al., 2010;

Kraková et al., 2012b; Sterflinger and Piñar, 2013; Madigan et al., 2014; Grabek-Lejko et

al., 2017; Mazzoli et al., 2018; Warner Marien, 2022). Additional research will be required

to evaluate the relationship between the *Pseudomonas* isolates of the book and these

inorganic compounds.

Pseudomonas spp. and *Streptomyces* spp. poses a risk not only because of its

potential catalytic properties, but also because of the ability of some members to produce

deteriorative pigments. For example, some *Streptomyces* spp. ruined graphic art inside

some Etruscan and Roman tombs due to the production of violet pigments, while members

of *Pseudomonas* produced both green (in an alkaline environment) and red (in acidic

conditions) pigments during wool degradation of historical textiles (Díaz-Herráiz, 2015;

Gutarowska et al., 2016; Mazzoli et al., 2018). In accordance with culture-dependent

methods, our present genomic analysis over the strain *Streptomyces* sp. CH-036 (Table 2),

revealed strong capabilities to degrade components of paper (Fig. 5.D), plus a high number

of genes for the biosynthesis of several pigments (Fig. 5.E) including isorenieratene (Fig.

5.F), a light-induced pigment (Takano et al., 2005). These findings should encourage the

managers of the museum to take the necessary precautions to avoid exposition of the book

to light or favorable conditions for the spreading of the microbe (moist, warm temperatures,

etc.).

Some microbes having their optimal growth during the different manufacturing

processes of the photo book, they could have been preserved via spores once the

manufacturing conditions changed. Indeed, 7 of the 17 strains from the book identified

taxonomically by the VITEK method are producers of endospores (*Bacillus* spp.) or

resistance spores (*Streptomyces* spp.). Both endospores and resistance spores are

metabolically inactive structures, extremely resistant to adverse environmental conditions,

such as high concentrations of salts, absence of nutrients or low levels of humidity. The

adaptive advantage conferred on bacteria is to allow them to remain as spores for as long as

necessary until the conditions are suitable (Madigan et al., 2014). Apparently, this could

explain the isolation of the thermophilic strain *Caldibacillus thermoamylovorans* CH-041

which grows optimally at 50°C in the book which is stored at room temperature (20-30°C).

The most likely hypothesis is that the thermophile proliferated during the initial

manufacturing. Different methods were used to harden the albumen layers, including

steaming and storing the albumen-coated papers inside a warm hayloft for half a year. The

temperature of a hayloft can sometimes reach 50 °C or more, providing an optimal

environment for the growth of CH-041 and other thermophiles like *Geobacillus* reported

from a previous albumen photography study (Puškárová et al., 2016). An evolutionary

significance of CH-041 is that its genome may have remained unchanged over the past 160

years. A deeper look in to the genomics of CH-041 revealed a scarce amount of genes for

antibiotic resistance (Fig. 5.A), and biodeterioration (Fig. 5.D-E), making it the most

environment friendly strain reported for the CHM.

4.3. Textiles as profilic microenvironments for bacteria proliferation.

A particularly intriguing case was observed in Alberdi's suit, a 19th-century silk

and lace garment, where we identified a unique bacterial strain, classified by VITEK as

Actinomyces odontolyticus. This filamentous bacterium is typically found in human

microbiomes, particularly in the oral cavity, and profusely isolated from dental caries (Batty, 1958). Manifestations of infection by *A. odontolyticus* include thoracic, abdominal, pelvic and central nervous system disease (Razok et al., 2022). Recently, it was proposed as driver of Colorectal Cancer (Breau, 2024). Its presence on the textile suggests potential contamination from human handling or storage conditions, a closed cartoon box that avoids contamination and humidity changes. Notably, the possibility that the original owner of the suit, Juan Bautista Alberdi, introduced this bacterium as a child cannot be ruled out, given the high prevalence of dental infections in toddlers and young children. Once established on the fabric, the bacterium's filamentous morphology may have facilitated its survival by intertwining with textile fibers, contributing to both microbial persistence and the progressive degradation of the material. This hypothesis aligns with scanning electron microscopy (SEM) observations, which revealed extensive colonization of textile fibers by filamentous bacteria (Fig. 2A). The genus *Actinomyces sp.* was previously cited in a list of the most common bacteria responsible for biodeterioration of CH (Grabek-Lejko et al., 2017). A more detailed study of the strain, including genome sequencing should reveal with certainty its taxonomic identity and toxicity, considering the hypothesis that it could have originated from an infected person.

The presence of *Micrococcus luteus* on textiles, a microorganism commonly associated with the accumulation of biological material and potential biodeterioration. Additionally, *Staphylococcus equorum*, found on both textiles and wooden surfaces, is typically linked to environments with human interaction, possibly contributing to the degradation of protein-based materials. Likewise, the *Bacillus* genus, including *Bacillus*

simplex and Bacillus licheniformis, was primarily found on wooden structures, suggesting

their potential role in the breakdown of cellulose and lignocellulosic compounds.

4.4. Furniture and wooden objects associated bacteria

The museum's wooden furniture subjected to sampling is currently exposed to the

air during the year, and suffers the dynamics in temperature and humidity that can enhance

bacterial growth. However, they also are affected by frequent procedures of cleaning and

disinfection, as well as repairs and preventive actions against wood decay. Frequently, the

wooden material in the museum is sensible to other pests, such as insects. For this reason, it

is not expected to these wooden items to suffer biodeterioration driven primarily by

bacteria. In spite of this, the isolates from the wooden parts of the chair, table, window,

washing trough, and main access door could have cellulolytic or lignocellulolytic

capabilities required for the degradation of wood. The most commonly reported bacteria

associated with wood-decay environments are genera such as *Clostridium*, and

cosmopolitan taxa such as *Bacillus* (Liu et al., 2018; Pyzik et al., 2021), both isolated from

the aforementioned elements (Table 1). A notable high number of genes related to the

breakup of cellulose and hemicellulose were found in the genomes of *Microbacterium* sp.

CH-007, *Kocuria* sp. CH-021, and *Microbacterium* sp. CH-015 (Fig. 5.D), which were

isolated from the chair wood, the entrance door and the window respectively (Table 1, Fig.

3). This could be supporting some kind metabolic activity over the wooden compounds of

the furniture. Moreover, the strongest profiles of heavy metal resistances belong to CH-021

and CH-015, and are probably the reflection of the chemical treatments used over the years

to preserve the wood from decay. Most of these wood preservatives recruit heavy metals in

their formulas, like copper chromate arsenate (CCA) which was widely used in the past but

now restricted due its toxicity (Woźniak, 2022). Finally, SEM micrographs of the washing

trough showed the presence of filamentous bacteria probably from the genus *Streptomyces*

or related (Fig. 2.D). Previous studies showed that filamentous actinomycetes are

associated with wood decay (Antai and Crawford, 1981; Jayasinghe and Parkinson, 2008),

while other claim that their presence in wood could prevent worst scenarios of

biodeterioration like those caused by fungi (Mortabit et al., 2015; Cai and Kuo, 2022).

One of the strains we isolated from the chair textile was classified as *Bacillus*

licheniformis, a species which is list of pathogens infecting humans (Bartlett et al., 2022).

B. licheniformis was reported in several cases of infection, even in a patient with no history

of any immune deficiency (Haydushka et al., 2012). Another intriguing species is

Staphylococcus equorum, isolated from both the textile chair and the washing trough. *S.*

equorum is a commensal and obligate aerobe usually found on the skin of various farm

animals, such as horses, dairy cattle and goats (Vermeersch et al., 2023; Reydams et al.,

2024; Senoussi et al., 2024). This microorganism probably has colonized the seat and meat

washing trough during the nineteenth century, when people of the city interacted routinely

with farm animals and used horses as their primary vehicle or cattle as meat source.

4.5. Metabolic traits and biotechnological potential of heritage bacteria

The identification of biosynthetic gene clusters (BGCs) related to pigment

production and antimicrobial compound synthesis in several isolates carries significant

implications for the preservation and study of heritage materials. Pigment biosynthesis

pathways—such as those detected in *Streptomyces* sp. CH-036, which includes genes for

the light-induced pigment isorenieratene—may contribute to visible discoloration and

irreversible staining of culturally significant surfaces. These pigments, often photoreactive

and chemically stable, can cause localized chromatic alterations, especially in substrates

such as textiles, photographs, and historical paper that are sensitive to light exposure.

In parallel, the detection of BGCs encoding antimicrobial compounds—such as

lanthipeptides and ε-poly-L-lysine—suggests that certain strains may exert ecological

pressure on their microbial surroundings. These bioactive compounds can inhibit

susceptible taxa and promote the survival of resistant microorganisms, potentially

influencing the microbial succession and stability on heritage objects. This microbial

antagonism may contribute to the reduced diversity observed in specific niches, such as the

surfaces of textiles or wooden elements with limited nutrient input.

Together, these metabolic features reveal a dual impact on conservation: while some

microbial metabolites contribute directly to biodeterioration through staining or enzymatic

degradation, others modulate microbial community dynamics, potentially altering the long-

term colonization patterns on heritage materials. These insights underscore the importance

of incorporating functional microbial traits—not just taxonomic identification—into the

design of informed, preventive conservation strategies.

Moreover, the genomic profiles of the isolated bacteria reveal a promising

biotechnological potential beyond their role in biodeterioration. Several *Bacillus* and

Pseudomonas strains identified in this study have previously been proposed as candidates

for bioprecipitation of calcium carbonate crystals, a process with practical applications in

biorestoration and consolidation of stone and plaster monuments. Notably, *Bacillus*

megaterium and *Bacillus simplex*, both detected among the isolates, are among the most

effective carbonate bioprecipitators and are used in protective treatments for historical

architecture (Paramo-Aguilera, 2012; Pyzik et al., 2021). In addition, members of the

Bacillus genus have been suggested for use in biotreatments of textiles and paper artifacts,

particularly for their enzymatic capabilities in cleaning and stabilizing cellulosic and

proteinaceous materials. Similarly, *Micrococcus* spp. are known for their capacity to

produce bioactive molecules and enzymes of interest to the pharmaceutical and

bioremediation industries, including antimicrobial compounds and pollutant-degrading

enzymes (Díaz-Herráiz, 2015).

Importantly, our genomic analyses revealed that *Streptomyces* sp. CH-036 and

Kocuria sp. CH-021 harbor genes involved in the synthesis of valuable secondary

metabolites and the degradation of xenobiotic compounds (see Fig. 5.F and section 3.4).

These findings position these strains as promising sources of novel biotechnological

resources and reinforce the concept of cultural heritage environments as overlooked

reservoirs of useful microorganisms. Further exploration of these isolates could yield new

applications in conservation, biotechnology, and environmental remediation.

CONCLUDING REMARKS AND PROSPECT

This study provides the first comprehensive microbiological survey of the Casa

Histórica de la Independencia Museum, revealing a rich and complex microbial landscape

shaped by historical, material, and environmental factors. Through the integration of

scanning electron microscopy, microbial isolation, MALDI-TOF MS, and genome

sequencing, we uncovered distinct microbial communities inhabiting both outdoor and
indoor surfaces of cultural heritage significance.

The comparative analysis between the Outdoor Urban Microbiome (OUM) and the
Indoor Museum Microbiome (IMM) revealed striking contrasts. Exterior surfaces, such as
the entrance door and walls, were dominated by stress-tolerant and resilient taxa—primarily
from the phyla Actinobacteriota and Firmicutes—reflecting their adaptation to fluctuating
environmental conditions, UV radiation, and atmospheric pollutants. In contrast, the IMM
showed greater taxonomic richness and niche specialization, shaped by more stable
temperature, humidity, and limited light exposure. Indoor isolates exhibited strong
substrate-specific associations, particularly in organic-rich materials like textiles and
albumen-based photographic prints, where human contact and historical use played an
important role in microbial transmission.

Artifacts such as the 19th-century albumen print photograph and Alberdi's silk suit
emerged as microecological hotspots, each serving as distinct reservoirs of specialized
bacteria. These niches supported extremophilic and potentially novel strains, including
Caldibacillus thermoamylovorans, *Streptomyces rubiginosohelvolus*, and *Oceanobacillus*
kimchii. Their presence reinforces the idea that heritage materials not only face
biodeterioration risks but also harbor microbial biodiversity of potential biotechnological
value.

Functional genome annotation highlighted a dual role of these microbes: some
strains possess genes linked to biodeterioration (e.g., cellulose degradation, pigment
biosynthesis), while others exhibit traits with positive applications, such as enzymatic

degradation of environmental pollutants. These findings illustrate how cultural heritage

microbiomes represent both a conservation challenge and an untapped biological resource.

Understanding microbial colonization patterns at the interface of cultural heritage

and environmental microbiology is critical for designing more informed, bio-aware

conservation strategies. This includes tailored preservation protocols that address not only

visible deterioration but also the metabolic capacities and ecological interactions of

microbial communities. Future work should expand to metagenomic and

metatranscriptomic approaches, allowing for a deeper exploration of microbial dynamics,

functional activity, and long-term succession on heritage substrates.

Ultimately, this study highlights the importance of heritage sites not only as

repositories of national memory but also as living ecosystems—microbial archive that

reflect centuries of human history, environmental exposure, and material transformation.

Protecting this biological dimension of cultural heritage will be key to its sustainable

preservation.

AKNOWLEDGEMENTS

The authors sincerely thank the Museo Casa Histórica de la Independencia staff,

especially former Director Cecilia Guerra Orozco and Cecilia Barrionuevo and Ana Oliva

from the restoration, conservation and museology department, for kindly granting access to

the museum's collections. All authors are members of the National Research Council

(CONICET) in Argentina. VHA was a recipient of a Georg Foster Scholarship for

Experienced Researchers, Alexander von Humboldt Foundation (2021-2023) and of a grant

of the Williams Foundation (2023-2024). Electron micrographs used in this study were

taken at the Center for Electron Microscopy (CIME) belonging to UNT and CONICET, in

Tucumán, Argentina.

FUNDING

This work was funded by National University of Tucuman with Projects PIUNT

G603 (2017-2022) and G703 (2023-2027). Funding was also available through research

Project in Thematic Areas of Museums and National Institutes 2021-2022 (CONICET and

Ministry of Culture of Argentina) and Universidad San Pablo-Tucumán (Project ISYCAV-

842 2019-2021).

AUTHOR CONTRIBUTIONS

VHA had the original project idea, designed and conceptualized the research. DAR

and VHA wrote the manuscript. FSG, NA, MCG and VHA performed the sampling. FSG,

NA, MCD and MJSM worked in the isolation and characterization of strains and curated

the CEBAC collection. MCG performed the photo documentation during the sampling in

the museum. FSG, LM, HE performed the electron microscopy analyses. DGAR performed

the genomic and bioinformatic analysis. VHA also provided funding, infrastructure,

consumables and equipment for the experiments. All authors read and approved the final

manuscript, which is available at biorxiv preprint server

https://doi.org/10.1101/2025.03.28.645971

REFERENCES

Adamiak, J., Otlewska, A., Tafer, H., Lopandic, K., Gutarowska, B., Sterflinger, K., et al.
(2018). First evaluation of the microbiome of built cultural heritage by using the Ion
Torrent next generation sequencing platform. *Int. Biodeterior. Biodegradation* 131,
11–18. doi: https://doi.org/10.1016/j.ibiod.2017.01.040

Ahamad, A., Yuan, C., Chung, C., Blair, B., Tran, A., and Tehreem, B. (2023). Metabolism
and gene sequence variation in Turicella otitidis implies its adaptability and
pathogenicity in extra-otic infection: a systematic review. *BMC Infect. Dis.* 23, 735.
doi: 10.1186/s12879-023-08721-y

Alonso-Reyes, D. G., and Albarracín, V. H. (2022). Arche: a Hierarchical, Easy to
Download, Functional-optimized Annotator for Microbial Meta(genomes). *bioRxiv*,
2022.11.28.518280. doi: 10.1101/2022.11.28.518280

Alonso-Reyes, D. G., Galván, F. S., Portero, L. R., Alvarado, N. N., Farías, M. E.,
Vazquez, M. P., et al. (2021). Genomic insights into an andean multiresistant soil
actinobacterium of biotechnological interest. *World J. Microbiol. Biotechnol.* 37, 166.
doi: 10.1007/s11274-021-03129-9

Antai, S. P., and Crawford, D. L. (1981). Degradation of Softwood, Hardwood, and Grass
Lignocelluloses by Two Streptomyces Strains . *Appl. Environ. Microbiol.* 42, 378–
380. doi: 10.1128/aem.42.2.378-380.1981

Bankevich, A., Nurk, S., Antipov, D., Gurevich, A. A., Dvorkin, M., Kulikov, A. S., et al.

(2012). SPAdes: a new genome assembly algorithm and its applications to single-cell

sequencing. *J. Comput. Biol.* 19, 455–477. doi: 10.1089/cmb.2012.0021

Bartlett, A., Padfield, D., Lear, L., Bendall, R., and Vos, M. (2022). A comprehensive list

of bacterial pathogens infecting humans. *Microbiology* 168. doi:

https://doi.org/10.1099/mic.0.001269

Batty, I. (1958). Actinomyces odontolyticus, a new species of actinomycete regularly

isolated from deep carious dentine. *J. Pathol. Bacteriol.* 75, 455–459. doi:

https://doi.org/10.1002/path.1700750225

Blin, K., Shaw, S., Augustijn, H. E., Reitz, Z. L., Biermann, F., Alanjary, M., et al. (2023).

antiSMASH 7.0: new and improved predictions for detection, regulation, chemical

structures and visualisation. *Nucleic Acids Res.* 51, W46–W50. doi:

10.1093/nar/gkad344

Borrego, S., Guiamet, P., Gómez de Saravia, S., Batistini, P., Garcia, M., Lavin, P., et al.

(2010). The quality of air at archives and the biodeterioration of photographs. *Int.*

Biodeterior. Biodegrad. 64, 139–145. doi: 10.1016/j.ibiod.2009.12.005

Branysova, T., Limpouch, O., Durovic, M., Demnerova, K., and Stiborova, H. (2023).

Bacterial Diversity on Historical Audio-Visual Materials and in the Atmosphere of

Czech Depositories. *Microbiol. Spectr.* 11. doi: 10.1128/spectrum.01176-23

Breau, K. A. (2024). Actinomyces odontolyticus: From Carries to Colorectal

Cancer. *Cell. Mol. Gastroenterol. Hepatol.* doi: 10.1016/j.jcmgh.2024.02.009

Bull, C. T., de Boer, S. H., Denny, T. P., Firrao, G., Fischer-Le Saux, M., Saddler, G. S., et

al. (2010). Comprehensive list of names of plant pathogenic bacteria, 1980-2007 . *J.*

Plant Pathol. 92, 551–592. Available at:

http://www.scopus.com/inward/record.url?scp=79955133206&partnerID=8YFLogxK

Cai, L., and Kuo, C.-J. (2022). Epsilon poly-L-lysine as a novel antifungal agent for

sustainable wood protection. *Front. Microbiol.* 13. Available at:

https://www.frontiersin.org/journals/microbiology/articles/10.3389/fmicb.2022.90854

1

Chatruc, C. (2021). Los diez museos más visitados de la Argentina. *La Nación*. Available

at: https://www.lanacion.com.ar/cultura/los-diez-museos-mas-visitados-de-la-

argentina-nid20092021/

Díaz-Herráiz, M. (2015). Caracterización de Comunidades Microbianas en Tumbas

Etruscas y Romanas. Universidad de Sevilla.

Elhagrassy, A. F. (2018). Isolation and characterization of actinomycetes from Mural

paintings of Snu- Sert-Ankh tomb, their antimicrobial activity, and their

biodeterioration. *Microbiol. Res.* 216, 47–55. doi: 10.1016/j.micres.2018.08.005

Gadd, G. M., Laurence, O. S., Briscoe, P. A., and Trevors, J. T. (1989). Silver

accumulation inPseudomonas stutzeri AG259. *Biol. Met.* 2, 168–173. doi:

10.1007/BF01142556

Grabek-Lejko, D., Tekiela, A., and Kasprzyk, I. (2017). Risk of biodeterioration of cultural

heritage objects, stored in the historical and modern repositories in the Regional

Museum in Rzeszow (Poland). A case study. *Int. Biodeterior. Biodegrad.* 123, 46–55.

doi: 10.1016/j.ibiod.2017.05.028

Granillo, A. (1872). *Provincia de Tucumán*. San Miguel de Tucumán.

Gutarowska, B., Pietrzak, K., Machnowski, W., and Milczarek, J. M. (2016). Historical

textiles – a review of microbial deterioration analysis and disinfection methods. *Text.*

Res. J. 87, 2388–2406. doi: 10.1177/0040517516669076

Gutarowska, B., Pietrzak, K., and Skóra, J. (2014). Disinfection as a factor reducing

microbial threat at workposts in museum and library - a comparison of the

effectiveness of photocatalytic ionization , UV irradiation and chemical misting. *Int. J.*

Curr. Microbiol. Appl. Sci. 3, 945–959.

Haydushka, I. A., Markova, N., Kirina, V., and Atanassova, M. (2012). Recurrent Sepsis

Due To Bacillus Licheniformis. *J. Glob. Infect. Dis.* 4. Available at:

https://journals.lww.com/jgid/fulltext/2012/04010/recurrent_sepsis_due_to_bacillus_li

cheniformis.17.aspx

Holden, P., and Nair, L. S. (2019). Deferoxamine: An Angiogenic and Antioxidant

Molecule for Tissue Regeneration. *Tissue Eng. Part B Rev.* 25, 461–470. doi:

10.1089/ten.teb.2019.0111

Jayasinghe, B. A. T. D., and Parkinson, D. (2008). Actinomycetes as antagonists of litter

decomposer fungi. *Appl. Soil Ecol.* 38, 109–118. doi:

https://doi.org/10.1016/j.apsoil.2007.09.005

Karbowska-Berent, J., Górny, R. L., Strzelczyk, A. B., and Wlazło, A. (2011). Airborne

and dust borne microorganisms in selected Polish libraries and archives. *Build.*

Environ. 46, 1872–1879. doi: https://doi.org/10.1016/j.buildenv.2011.03.007

Kraková, L., Chovanová, K., Selim, S. A., Šimonovičová, A., Puškarová, A., Maková, A.,

et al. (2012a). A multiphasic approach for investigation of the microbial diversity and

its biodegradative abilities in historical paper and parchment documents. *Int.*

Biodeterior. Biodegrad. 70, 117–125. doi: 10.1016/j.ibiod.2012.01.011

Kraková, L., Chovanová, K., Selim, S. A., Šimonovičová, A., Puškarová, A., Maková, A.,

et al. (2012b). A multiphasic approach for investigation of the microbial diversity and

its biodegradative abilities in historical paper and parchment documents. *Int.*

Biodeterior. Biodegradation 70, 117–125. doi:

https://doi.org/10.1016/j.ibiod.2012.01.011

Kraková, L., Šoltys, K., Otlewska, A., Pietrzak, K., Purkrtová, S., Savická, D., et al. (2018).

Comparison of methods for identification of microbial communities in book

collections: Culture-dependent (sequencing and MALDI-TOF MS) and culture-

independent (Illumina MiSeq). *Int. Biodeterior. Biodegrad.* 131, 51–59. doi:

10.1016/j.ibiod.2017.02.015

Książczyk, M., Kuczkowski, M., Dudek, B., Korzekwa, K., Tobiasz, A., Korzeniowska-

Kowal, A., et al. (2016). Application of Routine Diagnostic Procedure, VITEK 2

Compact, MALDI-TOF MS, and PCR Assays in Identification Procedure of Bacterial

Strain with Ambiguous Phenotype. *Curr. Microbiol.* 72, 570–582. doi:

10.1007/s00284-016-0993-0

Liu, Z., Fu, T., Hu, C., Shen, D., Macchioni, N., Sozzi, L., et al. (2018). Microbial

community analysis and biodeterioration of waterlogged archaeological wood from

the Nanhai No. 1 shipwreck during storage. *Sci. Rep.* 8, 1–11. doi: 10.1038/s41598-

018-25484-8

Madigan, M. T., Martinko, J. M., Bender, K. S., Buckley, D. H., and Stahl, D. A. (2014).

Brock Biology of Microorganisms., 14th Edn. Benjamin Cummings.

Marinsalda, J. C. (2015). La Casa Histórica de la Independencia Argentina. Universidad de

Sevilla. Available at: https://idus.us.es/items/8bf776a0-ce72-442c-9ac4-

91c634d76d33

Mazzoli, R., Giuffrida, M. G., and Pessione, E. (2018). Back to the past: "find the guilty

bug—microorganisms involved in the biodeterioration of archeological and historical

artifacts." *Appl. Microbiol. Biotechnol.* 102, 6393–6407. doi: 10.1007/s00253-018-

9113-3

Michaelsen, A., Piñar, G., Montanari, M., and Pinzari, F. (2009). Biodeterioration and

restoration of a 16th-century book using a combination of conventional and molecular

techniques: A case study. *Int. Biodeterior. Biodegrad.* 63, 161–168. doi:

10.1016/j.ibiod.2008.08.007

Mortabit, D., Zyani, M., and Koraichi, S. I. (2015). Biological control of wood decay fungi

by Streptomyces strains's extract. *Int. J. Innov. Sci. Res.* 14, 100–103.

Munish, C. A., Munesh, K., Rakesh, K., Kumar, D. D., Kumar, B. S., and Shanmugam, M.

(2017). Genome Sequence of Kocuria polaris Strain CD08_4, an Isolate from the

Duodenal Mucosa of a Celiac Disease Patient. *Genome Announc.* 5,

10.1128/genomea.01158-17. doi: 10.1128/genomea.01158-17

Negi, A., and Sarethy, I. P. (2019). Microbial Biodeterioration of Cultural Heritage: Events,

Colonization, and Analyses. *Microb. Ecol.* 78, 1014–1029. doi: 10.1007/s00248-019-

01366-y

Olowo-okere, A., Ibrahim, Y. K. E., Lo, C. I., Olayinka, B. O., Yimagou, E. K., Yacouba,

A., et al. (2021). Bhargavaea massiliensis sp. nov. and Dietzia massiliensis sp. nov.,

Novel Bacteria Species Isolated from Human Urine Samples in Nigeria. *Curr.*

Microbiol. 79, 18. doi: 10.1007/s00284-021-02721-4

Paramo-Aguilera, L. A. (2012). Caracterización de Comunidades Microbianas con

Potencial Biotecnológico Para la Prevención del Deterioro Estructural. Instituto

Politecnico Nacional, Mexico.

Perito, B., and Cavalieri, D. (2018). Innovative metagenomic approaches for detection of

microbial communities involved in biodeteriorattion of cultural heritage. *IOP Conf.*

Ser. Mater. Sci. Eng. 364. doi: 10.1088/1757-899X/364/1/012074

Piñar, G., Tafer, H., Sterflinger, K., and Pinzari, F. (2015). Amid the possible causes of a

very famous foxing: molecular and microscopic insight into Leonardo da Vinci's self-

portrait. *Environ. Microbiol. Rep.* 7, 849–859. doi: https://doi.org/10.1111/1758-

2229.12313

Pritchard, L., Glover, R. H., Humphris, S., Elphinstone, J. G., and Toth, I. K. (2016).

Genomics and taxonomy in diagnostics for food security: soft-rotting enterobacterial

plant pathogens. *Anal. Methods* 8, 12–24. doi: 10.1039/C5AY02550H

Puškárová, A., Bučková, M., Habalová, B., Kraková, L., Maková, A., and Pangallo, D.

(2016). Microbial communities affecting albumen photography heritage: A

methodological survey. *Sci. Rep.* 6, 1–14. doi: 10.1038/srep20810

Pyzik, A., Ciuchcinski, K., Dziurzynski, M., and Dziewit, L. (2021). The bad and the good-

microorganisms in cultural heritage environments-an update on biodeterioration and

biotreatment approaches. *Materials (Basel).* 14, 1–15. doi: 10.3390/ma14010177

Rasuk, M. C., Ferrer, G. M., Kurth, D., Portero, L. R., Farías, M. E., and Albarracín, V. H.

(2017). UV-Resistant Actinobacteria from High-Altitude Andean Lakes: Isolation,

Characterization and Antagonistic Activities. *Photochem. Photobiol.* 93, 865–880. doi:

10.1111/php.12759

Razok, A., Ali, M., Aker, L., and Ziglam, H. (2022). Actinomyces odontolyticus

bacteraemia associated with cervical and mediastinal abscesses in an

immunocompetent patient: First reported case in Qatar. *New Microbes New Infect.* 45,

100956. doi: https://doi.org/10.1016/j.nmni.2022.100956

Reilly, J. M. (1980). *The Albumen & Salted Paper Book: The History and Practice of*

Photographic Printing, 1840-1895. Rochester: Light Impressions Corp.

Repka, L. M., Chekan, J. R., Nair, S. K., and van der Donk, W. A. (2017). Mechanistic

Understanding of Lanthipeptide Biosynthetic Enzymes. *Chem. Rev.* 117, 5457–5520.

doi: 10.1021/acs.chemrev.6b00591

Reydams, H., Toledo-Silva, B., Mertens, K., Piepers, S., Vereecke, N., Souza, F. N., et al.

(2024). Phenotypic and genotypic assessment of iron acquisition in diverse bovine-

associated non-aureus staphylococcal strains. *Vet. Res.* 55, 6. doi: 10.1186/s13567-

023-01260-z

Richter, M., and Rosselló-Móra, R. (2009). Shifting the genomic gold standard for the

prokaryotic species definition. *Proc. Natl. Acad. Sci. U. S. A.* 106, 19126–31. doi:

10.1073/pnas.0906412106

Senoussi, A., Aissaoui-Zitoun, O., Chenchouni, H., Senoussi, S., Saoudi, Z., Pediliggieri,

C., et al. (2024). Microbial screening of animal skin bags used in traditional

cheesemaking. *Int. J. Food Microbiol.* 411, 110549. doi:

https://doi.org/10.1016/j.ijfoodmicro.2023.110549

Skóra, J., Gutarowska, B., Pielech-Przybylska, K., Stępień, Ł., Pietrzak, K., Piotrowska,

M., et al. (2015). Assessment of microbiological contamination in the work

environments of museums, archives and libraries. *Aerobiologia (Bologna).* 31, 389–

401. doi: 10.1007/s10453-015-9372-8

Stackebrandt, E., Koch, C., Gvozdiak, O., and Schumann, P. (1995). Taxonomic dissection

of the genus Micrococcus: Kocuria gen. nov., Nesterenkonia gen. nov., Kytococcus

gen. nov., Dermacoccus gen. nov., and Micrococcus cohn 1872 gen. emend. *Int. J.*

Syst. Bacteriol. 45, 682–692. doi: 10.1099/00207713-45-4-682

Sterflinger, K., and Piñar, G. (2013). Microbial deterioration of cultural heritage and works

of art - Tilting at windmills? *Appl. Microbiol. Biotechnol.* 97, 9637–9646. doi:

10.1007/s00253-013-5283-1

Takano, H., Obitsu, S., Beppu, T., and Ueda, K. (2005). Light-induced carotenogenesis in

Streptomyces coelicolor A3(2): Identification of an extracytoplasmic function sigma

factor that directs photodependent transcription of the carotenoid biosynthesis gene

cluster. *J. Bacteriol.* 187, 1825–1832. doi: 10.1128/JB.187.5.1825-1832.2005

Tran, M. T., Seuylemezian, A., Wright, A., Coil, D., Eisen, J., and Guan, L. (2023). Draft

Genome Sequences of Spacecraft-Associated Microbes Isolated from Six NASA

Missions. *Microbiol. Resour. Announc.* 12. doi: 10.1128/mra.01011-22

Vermeersch, A. S., Ali, M., Gansemans, Y., Van Nieuwerburgh, F., Ducatelle, R., Geldhof,

P., et al. (2023). An in-depth investigation of the microbiota and its virulence factors

associated 11 with severe udder cleft dermatitis lesions. *J. Dairy Sci.* doi: 10.3168/jds.2023-24180

Villalba, L. S., Mikán, J. F., and Sánchez, J. (2004). Actividades hidrolíticas y caracterización isoenzimática de poblaciones microbianas aisladas del patrimonio documental del Archivo General de Colombia. *Nova* 2, 50–58. doi: 10.22490/24629448.7

Wang, L., Zhang, C., Zhang, J., Rao, Z., Xu, X., Mao, Z., et al. (2021). Epsilon-poly-L-lysine: Recent Advances in Biomanufacturing and Applications. *Front. Bioeng. Biotechnol.* 9. Available at: https://www.frontiersin.org/journals/bioengineering-and-biotechnology/articles/10.3389/fbioe.2021.748976

Warner Marien, M. (2022). *Photography: A Cultural History*., 5th Edn. London: Laurence King Publishing.

Wiszniewska, M., Walusiak-Skorupa, J., Pannenko, I., Draniak, M., and Palczynski, C. (2009). Occupational exposure and sensitization to fungi among museum workers. *Occup. Med. (Chic. Ill).* 59, 237–242. doi: 10.1093/occmed/kqp043

Woźniak, M. (2022). Antifungal Agents in Wood Protection—A Review. *Molecules* 27. doi: 10.3390/molecules27196392

Yuan, Y., and Gao, M. (2015). Genomic analysis of a ginger pathogen Bacillus pumilus providing the understanding to the pathogenesis and the novel control strategy. *Sci. Rep.* 5, 10259. doi: 10.1038/srep10259

Zhou, Y., Wang, S., and Weng, X. B. (2022). Clostridium subterminale infection in a patient with diffuse large B-cell lymphoma and haemophagocytic syndrome: A case

report and literature review. *J. Int. Med. Res.* 50. doi: 10.1177/03000605221129558

Zielińska-Jankiewicz, K., Kozajda, A., Piotrowska, M., and Szadkowska-Stańczyk, I.

(2008). Microbiological Contamination With Moulds in Work Environment in

Libraries and Archive Storage Facilities. *Ann. Agric. Environ. Med.* 15, 71–78.

Available at: https://www.aaem.pl/MICROBIOLOGICAL-CONTAMINATION-

WITH-MOULDS-IN-WORK-ENVIRONMENT-nIN-LIBRARIES-AND-

ARCHIVE,90139,0,2.html

Figure and table legends.

Figure 1. Sampled locations of the CHM. **A.** Facade. **B.** "Salon de la Jura". **C-E**. Chair

and table from the "Salón de la Jura". **F-G.** Window Salon de la Jura. **H-I.** Alberdi's

garments. **J-L.** Albumen print book. **M.** Washing trough. **N.** Main door.

Figure 2. Biofilms on heritage surfaces including furniture, wooden objects (chair, table,

window and washing trough) and textile garments. **A.** Chair. **B.** Table. **C-D.** Washing

trough. **G-H.** Garments. **I.** Window. **J.** Entrance door.

Figure 3. Illustration of the CHM and sampling sites. Several taxonomical groups assigned

to the isolates through VITEK are represented by symbols.

Figure 4. Subcultures performed in LB broth for some CHM strains, complemented by

gram staining and SEM microscopy. **A.** CH-005. **B.** CH-003. **C.** CH-010 **D.** CH-011. **E.**

CH-013 **F.** CH-015. **G.** CH-021 **H.** CH-026.

Figure 5. The genomic potential of eight sequenced strains unveiled by functional

annotation. **A-E.** In barplots are depicted the abundances for those genes involved in

resistances to antibiotics, osmotic stress, heavy metals and those required for the

degradation of wood and paper, and the production of pigments. **F.** Potential for the

synthesis of secondary metabolites, and degradation of xenobiotics.

1052

1053 **Table 1.** Taxonomical, macroscopical and microscopical details for most of the 49 strains
1054 isolated from the CHM.

ID	TAXONOMICAL IDENTIFICATION BY VITEK	TAXONOMICAL CLASSIFICATION	SOURCE	MACROSCOPICAL CHARACTERIZATION		MICROSCOPICAL CHARACTERIZATION	
				COLOUR	SURFACE AND TEXTURE	MORPHOLOGY	GRAM
CH - 001	*Micrococcus luteus*	*Actinobacteriota-Actinomycetia-Actinomycetales-Micrococcaceae*	SJ Chair (textile)	Intense-yellow	Smooth and creamy	Coccoid	(+)
CH - 002	*Staphylococcus equorum*	*Firmicutes-Bacilli-Staphylococcales-Staphylococcaceae*	SJ Chair (textile)	Pale-yellow	Smooth and creamy	Coccoid	(+)
CH - 003	*Bacillus simplex*	*Firmicutes-Bacilli-Bacillales-Bacillaceae*	SJ Chair (textile)	Cream-colored	Smooth and creamy	Bacillus	(+)
CH - 004	*Bacillus licheniformis*	*Firmicutes-Bacilli-Bacillales-Bacillaceae*	SJ Chair (textile)	Cream-colored	Smooth and creamy	Bacillus	(+)
CH - 005	*Micrococcus sp.*	*Actinobacteriota-Actinomycetia-Actinomycetales-Micrococcaceae*	SJ Chair (wood)	Orange	Smooth and creamy	Coccoid	(+)
CH - 006	*Kocuria rosea*	*Actinobacteriota-Actinomycetia-Actinomycetales-Micrococcaceae*	SJ Chair (wood)	Orange	Smooth and creamy	Coccoid	(+)
CH - 007	Unidentified	-	SJ Chair (wood)	Pale-yellow	Smooth and creamy		
CH - 008	*Kocuria rosea*	*Actinobacteriota-Actinomycetia-Actinomycetales-Micrococcaceae*	SJ Table (wood)	Orange	Smooth and creamy	Coccoid	(+)
CH - 009	*Micrococcus luteus*	*Actinobacteriota-Actinomycetia-Actinomycetales-Micrococcaceae*	SJ Table (wood)	Intense-yellow	Smooth and creamy	Coccoid	(+)
CH - 010	*Bacillus altitudinis/pumilus*	*Firmicutes-Bacilli-Bacillales-Bacillaceae*	SJ Table (wood)	Cream-colored	Rough and mucoid	Bacillus	(+)
CH - 011	*Staphylococcus equorum*	*Firmicutes-Bacilli-Staphylococcales-Staphylococcaceae*	Washing trough (wood)	Pale-yellow	Smooth and creamy	Coccoid	(+)
CH - 012	*Bacillus sp.*	*Firmicutes-Bacilli-Bacillales-Bacillaceae*	Washing trough (wood)	Cream-colored	Smooth and creamy	Bacillus	(+)
CH - 013	*Actinomyces odontolyticus*	*Actinobacteriota-Actinomycetia-Actinomycetales-Actinomycetaceae*	Alberdi's Suit (silk and lace)	White	Dry and hard	Mycelium	(+)
CH - 014	*Micrococcus luteus*	*Actinobacteriota-Actinomycetia-Actinomycetales-Micrococcaceae*	SJ Window (blue painted wood)	Intense-yellow	Smooth and creamy	Coccoid	(+)
CH - 015	*Microbacterium aurum*	*Actinobacteriota-Actinomycetia-Actinomycetales-Microbacteriaceae*	SJ Window (blue painted wood)	Yellow	Smooth and creamy	Bacillus	(+)
CH - 016	Unidentified	-	SJ Window (blue painted wood)	Yellow	Smooth and creamy		
CH - 017	*Microbacterium aurum*	*Actinobacteriota-Actinomycetia-Actinomycetales-Microbacteriaceae*	Entrance door (blue painted wood)	Yellow	Smooth and creamy	Bacillus	(+)
CH - 018	*Bacillus altitudinis/pumilus*	*Firmicutes-Bacilli-Bacillales-Bacillaceae*	Entrance door (blue painted wood)	Pale-yellow	Smooth and creamy	Bacillus	(+)
CH - 019	*Clostridium subterminale*	*Firmicutes-Clostridia-Clostridiales-Clostridiaceae*	Entrance door (blue painted wood)	Yellow	Smooth and creamy	Bacillus	(+)
CH - 020	*Micrococcus luteus*	*Actinobacteriota-Actinomycetia-Actinomycetales-Micrococcaceae*	Entrance door (blue painted wood)	Intense-yellow	Smooth and creamy	Coccoid	(+)
CH - 021	*Corynebacterium pseudodiphtheriticum*	*Actinobacteriota-Actinomycetia-Mycobacteriales-Mycobacteriaceae*	Entrance door (blue painted wood)	Orange	Smooth and creamy	Bacillus	(+)
CH - 022	Unidentified	-	Entrance door (blue painted wood)	Yellow	Smooth and creamy		
CH - 023	*Kocuria rosea*	*Actinobacteriota-Actinomycetia-Actinomycetales-Micrococcaceae*	Entrance door (blue painted wood)	Orange	Smooth and creamy	Coccoid	(+)
CH - 024	*Corynebacterium pseudodiphtheriticum*	*Actinobacteriota-Actinomycetia-Mycobacteriales-Mycobacteriaceae*	Exterior walls (white painted bricks)	Intense-orange	Smooth and creamy	Bacillus	(+)
CH - 025	*Kocuria rosea*	*Actinobacteriota-Actinomycetia-Actinomycetales-Micrococcaceae*	Exterior walls (white painted bricks)	Orange	Smooth and creamy	Coccoid	(+)
CH - 026	*Turicella otitidis*	*Actinobacteriota-Actinomycetia-Actinomycetales-Corynebacteriaceae*	Exterior walls (white painted bricks)	Yellow	Smooth and creamy	Bacillus	(+)
CH - 027	*Bacillus megaterium*	*Firmicutes-Bacilli-Bacillales-Bacillaceae*	Exterior walls (white painted bricks)	Cream-colored	Smooth and hard	Bacillus	(+)
CH - 028	*Kocuria rosea*	*Actinobacteriota-Actinomycetia-Actinomycetales-Micrococcaceae*	Exterior walls (white painted bricks)	Orange	Smooth and creamy	Coccoid	(+)
CH - 029	*Bacillus altitudinis/pumilus*	*Firmicutes-Bacilli-Bacillales-Bacillaceae*	GB Albumen print	Cream-colored	Smooth and creamy	Bacillus	(+)
CH - 030	Unidentified	-	GB Albumen print	Intense-orange	Smooth and mucoid	Coccoid	(+)
CH - 031	*Bacillus galactosidilyticus*	*Firmicutes-Bacilli-Bacillales-Bacillaceae*	GB Albumen print	Cream-colored	Smooth and creamy	Bacillus	(+)
CH - 032	*Pseudomonas sp*	*Proteobacteria-Gammaproteobacteria-Pseudomonadales-Pseudomonadaceae*	GB Albumen print	Cream-colored	Smooth and creamy	Rod-shaped	(-)
CH - 033	*Pseudomonas sp*	*Proteobacteria-Gammaproteobacteria-Pseudomonadales-Pseudomonadaceae*	GB Albumen print	Cream-colored	Smooth and creamy	Rod-shaped	(-)
CH - 034	*Pseudomonas sp*	*Proteobacteria-Gammaproteobacteria-Pseudomonadales-Pseudomonadaceae*	GB Albumen print	Cream-colored	Smooth and creamy	Rod-shaped	(-)
CH - 035	*Kocuria rosea*	*Actinobacteriota-Actinomycetia-Actinomycetales-Micrococcaceae*	GB Albumen print	Pale-yellow	Smooth and creamy	Coccoid	(+)
CH - 036	*Streptomyces griseus*	*Actinobacteriota-Actinomycetia-Streptomycetales-Streptomycetaceae*	GB Albumen print	White	Dry and hard	Mycelium	(+)
CH - 037	Unidentified	-	GB Albumen print	Cream-colored	Smooth and creamy	Bacillus	(+)
CH - 038	*Kocuria rosea*	*Actinobacteriota-Actinomycetia-Actinomycetales-Micrococcaceae*	GB Albumen print	Intense-orange	Smooth and creamy	Coccoid	(+)
CH - 039	*Bacillus galactosidilyticus*	*Firmicutes-Bacilli-Bacillales-Bacillaceae*	GB Albumen print	Cream-colored	Smooth and mucoid	Bacillus	(+)
CH - 040	*Pseudomonas sp*	*Proteobacteria-Gammaproteobacteria-Pseudomonadales-Pseudomonadaceae*	GB Albumen print	Pale-yellow	Smooth and mucoid	Bacillus	(+)
CH - 041	*Bacillus thermoamylovorans*	*Firmicutes-Bacilli-Bacillales-Bacillaceae*	GB Albumen print	Cream-colored	Smooth and creamy	Bacillus	(+)
CH - 042	*Streptomyces sp*	*Actinobacteriota-Actinomycetia-Streptomycetales-Streptomycetaceae*	GB Albumen print	Pale-cream	Smooth and hard	Mycelium	(+)
CH - 043	*Micrococcus luteus*	*Actinobacteriota-Actinomycetia-Actinomycetales-Micrococcaceae*	GB Albumen print	Intense-yellow	Smooth and creamy	Coccoid	(+)
CH - 044	Unidentified	-	GB Albumen print	Pale-yellow	Smooth and mucoid		(+)
CH - 045	*Bacillus cereus group*	*Firmicutes-Bacilli-Bacillales-Bacillaceae*	GB Albumen print	Cream-colored	Smooth and creamy	Bacillus	(+)
CH - 046	*Staphylococcus epidermis*	*Firmicutes-Bacilli-Staphylococcales-Staphylococcaceae*	GB Albumen print	Cream-colored	Smooth and dry	Coccoid	(+)
CH - 047	*Micrococcus luteus*	*Actinobacteriota-Actinomycetia-Actinomycetales-Micrococcaceae*	GB Albumen print	Intense-yellow	Smooth and creamy	Coccoid	(+)
CH - 048	Unidentified	-	GB Albumen print	Orange	Smooth and mucoid	Coccoid	(+)
CH - 049	*Pseudomonas sp*	Gammaproteobacteria-Pseudomonadales-Pse...	GB Paper Sheet	Cream-colored	Smooth and creamy	Rod-shaped	(-)

1055

1056 **Table 2.** Data of the sequenced isolates, including the isolation place, sequencing results,
1057 de novo assembly, annotation, and taxonomical identification by several methods. In grey,
1058 are the Indoors Museum Microbiome (IMM) isolated strains, in white the Outdoors Urban
1059 Microbiome (OMM) ones.

1060

Strain code	CH-041	CH-031	CH-036	CH-007	CH-015	CH-021	CH-019	CH-026
Isolation place	Albumen picture	Albumen picture	Albumen picture	SJ Chair (wood)	SJ Window	Main door	Main door	Outside wall
Raw data (Gb)	1,4	1,5	1,6	2,3	2,1	2,2	1,8	2,2
Length after assembly (bp)	4288172	3929860	7613203	3319824	3252758	3911307	3193378	3689050
Contigs	200	64	114	24	57	102	73	85
CDS	4067	3925	6773	3143	3063	3584	3223	3416
Annotated genes	3864	3737	6577	2887	2766	3208	2988	2909
Taxonomy with Vitek	*Bacillus thermoamylovorans*	*Bacillus galactosidilyticus*	*Streptomyces griseus*	*Unidentified*	*Microbacterium aurum*	*Corynebacterium pseudodiphtheriticum*	*Clostridium subterminale*	*Turicella otitidis*
Taxonomy with 16S/gyrB	*Caldibacillus thermoamylovorans (gyrB)*	*Oceanobacillus sp. (gyrB)*	*Streptomyces globisporus (16S)*	*Microbacterium imperiale (16S)*	*Microbacterium aurum (16S)*	*Kocuria rosea (16S)*	*Bhargavaea ginsengi (16S)*	*Pseudoclavibacter sp. (16S)*
Taxonomy with ANI	*Caldibacillus thermoamylovorans*	*Oceanobacillus kimchii*	*Streptomyces rubiginosohelvolus*	*Microbacterium sp.*	*Microbacterium aurum*	*Kocuria sp.*	*Bhargavaea massiliensis*	*Pseudoclavibacter sp.*
Closest match in NCBI genomes	*Caldibacillus thermoamylovorans* MSL 185.1 Spacecraft Cleanroom ASM2371276v1 (97,93%)	*Oceanobacillus kimchii* p3-SID1558 Human Back Skin ASM2514966v1 (99,47%)	*Streptomyces* sp. f150 Soil ASM255143v1 (98,48%)	***Microbacterium arborescens Ricinus comunis* Leaves ASM333966v1 (87,82%)**	*Microbacterium* sp. DE14.009 Geothermal springs ASMf476380v1 (97,98%)	*Kocuria* sp. Duodenal mucosa of Celiac disease patient ASM148375v1 (98,71%)	*Bhargavaea massiliensis* Marseille-Q1000 Human Urine GCA_902825275.1 (99,32%)	***Pseudoclavibacter helvolus* PS02339 Danish Slaughterhouse Wall ASM3001191v1 (87,11%)**

1061