Mapping cellular targets of covalent cancer drugs in the entire mammalian body

Authors: Zhengyuan Pang[1, 2,10], Verina H. Leung[1,10], Cailynn C. Wang[1], Ahmadreza Attarpour[3,4], Anthony Rinaldi[3], Hanbing Shen[1], Maria Dolores Moya-Garzon[5, 6], Logan H. Sigua[1], Claire Rammel[7], Alexandra Selke[1], Christopher Glynn[1], Melaina Yender[1], Senhan Xu[1], Javid J. Moslehi[7], Peng Wu[8], Jonathan Z. Long[5], Maged Goubran[3,4,9], Benjamin F. Cravatt[2], Li Ye [1, 11]*

[1]Howard Hughes Medical Institute, Department of Neuroscience, Dorris Neuroscience Center, The Scripps Research Institute, La Jolla, California, 92037, United States.

[2]Department of Chemistry, The Scripps Research Institute, La Jolla, California, 92037, United States.

[3]Physical Sciences Platform, Sunnybrook Research Institute, Toronto, Ontario, Canada.

[4]Department of Medical Biophysics, University of Toronto, Toronto, Ontario, Canada.

[5]Department of Pathology, Stanford School of Medicine, Stanford, CA, USA

[6]Sarafan ChEM-H, Stanford University, Stanford, CA, USA

[7]Section of Cardio-Oncology & Immunology, Cardiovascular Research Institute (CVRI), University of California San Francisco, School of Medicine, San Francisco, California, 94158, United States.

[8]Department of Molecular and Cellular Biology, The Scripps Research Institute, La Jolla, California, 92037, United States.

[9]Hurvitz Brain Sciences, Sunnybrook Health Sciences Centre, Toronto, Ontario, Canada.

[10]These authors contributed equally.

[11]Lead contact

*Correspondence: liye@scripps.edu

SUMMARY:

As our understanding of biological systems reaches single-cell and high spatial resolutions, it
becomes imperative that pharmacological approaches match this precision to understand drug
actions. This need is particularly urgent for the targeted covalent inhibitors that are currently re-
entering the stage for cancer treatments. By leveraging the unique kinetics of click reactions, we
developed volumetric clearing-assisted tissue click chemistry (vCATCH) to enable deep and
homogeneous click labeling across the 3D mammalian body. With simple and passive incubation
steps, vCATCH offers cellular resolution drug imaging in the entire adult mouse. We combined
vCATCH with HYBRiD imaging and virtual reality to visualize and quantify *in vivo* targets of
two clinical cancer drugs, afatinib and ibrutinib, which recapitulated their known
pharmacological distribution and revealed previously unreported tissue and cell type engagement
potentially linked to off-target effects. vCATCH provides a body-wide, unbiased platform to map
covalent drug engagements at unprecedented scale and precision.

KEYWORDS:

drug targets, tissue clearing, whole-body imaging, targeted covalent inhibitors

INTRODUCTION

The primary binding sites for drug molecules are critical determinants of drug efficacy and toxicity. Although molecular target identification strategies, such as chemoproteomics[1-5], chemogenomic[6-9], and thermal shift profiling[10-13], have revolutionized our understanding of drug activities, mapping *in vivo* drug targets with spatial and cellular resolution remains challenging. This shortfall is even more pressing in light of the innovations in genomics and imaging techniques over the past decade that have enabled the characterization of tissue architecture at the single-cell level with spatial resolution[14-20]. This has led to a widening gap between our understanding of native biological systems and our ability to track *in vivo* drug actions within these systems, as the latter still primarily rely on low-resolution methods such as radioactive imaging. It is imperative that we develop methods to visualize *in situ* drug engagement with cellular resolution across tissue compartments and organ systems to better understand *in vivo* drug actions.

One promising avenue for imaging drug targets is fluorescence imaging, which has been widely used for visualizing endogenous biomolecules like DNA, mRNA, and proteins. However, directly imaging drug molecules conjugated with a fluorescent tag is often not suitable as a larger tag could distort drug activity and distribution *in vivo*. As an alternative approach, we recently developed a method to visualize drug molecules in tissue sections by combining tissue clearing with *in situ* click chemistry (CATCH)[21]. However, efforts to extend CATCH from thin sections to large, heterogeneous 3D tissue environments face significant challenges similar to those typically encountered by whole-mount immunostaining or *in situ* hybridization methods, such as the accumulation of probes at the tissue surface, uneven staining across the tissue depth, and insufficient labeling at the center of large samples[22-26].

Here, we develop new CATCH capacities to enable rapid and homogenous *in situ* click labeling in ultra-large 3D tissues at the scale of a whole mouse body. This new technology leverages the unique characteristics of click chemistry by introducing pre-reaction copper saturation (PRCS) and repeated iterations of reaction (RIR). The resulting volumetric CATCH (vCATCH) method, when combined with HYBRiD (hydrogel-based reinforcement of three-dimensional imaging solvent-cleared organs (DISCO)) clearing[27], is further highlighted by its marked simplicity as it only relies on easy and passive incubation steps to achieve whole-body labeling and imaging.

RESULTS

Establishing the principles for deep tissue click reactions

To achieve 3D click labeling, we turned to recent advances in whole-brain and whole-body immunostaining techniques[24,28-35]. As summarized by previous works[36-39], two main strategies have been used to achieve deep tissue antibody penetration: (1) enhancing diffusion of the probe (e.g., antibody) into the tissue; and (2) minimizing probe-target interactions (e.g., antibody-antigen binding) during the diffusion stage while maximizing reaction efficiency at the staining stage, such that the large difference in reaction rates between the two stages will prevent probe depletion at the tissue surface and facilitate homogeneous tissue labeling[23,33,34]. Biophysics studies have demonstrated that changing the pH, salt concentration, and other physicochemical parameters can modulate antibody-antigen binding kinetics by 4-400 fold[40,41]. We hypothesized that we could adopt a similar strategy to enable 3D click reactions. CATCH relies on copper-

catalyzed alkyne azide cycloaddition (CuAAC) to label target-bound drug molecules with
fluorescent signals. As its name indicates, the kinetics of CuAAC strictly depend on the catalytic
Cu(I)-ligand complex, but for ease of handling, a Cu(II) compound, typically $CuSO_4$, is used
during the diffusion stage and then a reducing agent (e.g., sodium ascorbate) is added *in situ* to
generate Cu(I) at the binding stage. Compared to uncatalyzed conditions, the generation of
catalytic Cu(I) can lower the energy barrier by 14.9-18.4 kcal/mol, or an equivalent of 7-8 orders
of magnitude increase favoring the cycloaddition that attaches the fluorescent probe to the drug
in situ[42]. Thus, by temporarily preventing the generation of Cu(I) from Cu(II) through
withholding the reducing agent sodium ascorbate, in principle, we could modulate the probe-
target reaction by 10^{7-8} fold between the diffusion and binding stages, which could improve the
penetration depth of CATCH in large tissues.

For method development, we used the covalent monoamine oxidase (MAO) inhibitor pargyline,
whose alkyne probe pargyline-yne has previously been characterized by activity-based protein
profiling (ABPP) and 2D-CATCH studies (Figure S1A)[21,43]. Although CATCH has demonstrated
homogeneous click labeling of N_3-dye in 500 μm tissues[21], when tested in mouse hemispheres,
even with the aforementioned catalyst withholding strategy, only the surface of the mouse brain
was labeled, a depth of 0.4 mm (Figure 1A). First, we used mouse brains treated with the fatty
acid amide hydrolase (FAAH) inhibitor PF7845-yne (Figure S1A), another CATCH probe we
previously established, to determine whether other CATCH components might have trouble
diffusing through the tissue[21,44]. We removed individual reagents from the diffusion step and
evaluated the resulting labeling depth. For instance, eliminating the N_3-dye, the largest molecular
weight component, in the diffusion step did not affect labeling depth, suggesting N_3-dye
diffusion is not a limiting factor (Figure S1B and S1C). By contrast, removing $CuSO_4$ or ligand
strongly reduced labeling depth in these conditions, suggesting that the catalyst is still likely the
diffusion bottleneck (Figure S1B and S1C).

Cu is well-known to bind multiple amino acid side chains (i.e., cysteine and histidine)[45-48]. We
therefore hypothesized that the ubiquitous copper binding sites could consume free Cu(II) as it
diffuses into the sample, resulting in insufficient catalyst conversion in deeper tissues (Figure
1B). Unlike typical staining in which increasing dye or antibody concentrations tends to result in
nonspecific labeling and high background, high Cu(II) alone, as a catalytic precursor, should not
lead to nonspecific labeling. Based on this theoretical prediction, we reasoned that excessive
Cu(II) could be used to pre-saturate potential copper binding sites (therefore "blocking" them) to
boost catalytic activity in deeper tissues. Importantly, this strategy should not increase
nonspecific labeling or lead to high background. Indeed, we observed that incorporating a 10x
Cu excess pre-reaction copper saturation (PRCS) step increased the reaction depth to 1 mm while
maintaining the signal-to-noise ratio after a single one-hour reaction (Figure 1A-1C).

We next sought to modify CATCH to reach targets in tissues deeper than 1 mm. Although
increasing reaction time could potentially increase labeling depth, prolonged reactions would
deplete the reducing agent in the buffer, which could eventually lead to reactive oxygen species
(ROS) build-up and side reactions[49-51]. Instead, we refreshed the click reaction cocktail every
hour, a procedure we termed repeated iterations of reactions (RIR). In practice, PRCS plus two
rounds of RIR resulted in full-depth, homogenous CATCH labeling across the mouse hemisphere
(Figure 1A-1C, Supplementary Video 1). We term this approach volumetric CATCH (vCATCH).

132 **vCATCH quantitatively profiles brain-wide targets of Pargyline**

133 After achieving brain-wide labeling, we employed vCATCH to unbiasedly map drug targets of
134 pargyline. After intraperitoneal (i.p.) injection of pargyline-yne (10 mg/kg) or vehicle (non-drug
135 treated) in wild-type mice, mouse hemispheres were harvested and cleared by HYBRiD.
136 Hemispheres then underwent one day of PRCS and two rounds of RIR before imaging with
137 lightsheet fluorescence microscopy (Figure 1D). Importantly, vCATCH allowed the use of high-
138 throughput, paralleled passive processing to generate cohort-scale pargyline-labeled brains with
139 marked reproducibility, critical for performing biologically meaningful analysis of its cellular
140 targets across the whole brain (Figure 1E-1H, S1D, and S1E).

141 First, we manually verified pargyline probe binding in the brain based on previously reported
142 regional targets[21]. These regions included distinct cellular structures in the pons and
143 hypothalamus (Figure 1H and S1E). In addition, we found pargyline-yne-enriched cells in the
144 choroid plexus (in the ventricle) and the medulla (Figure 1H and S1E). We also observed
145 additional neuron-like structures in the midbrain (Figure S1E). These findings demonstrated that
146 vCATCH maintained high fidelity in large tissue volumes and across the whole brain.

147 The throughput and reproducibility of vCATCH across cohort-scale samples also allowed
148 automatic and unbiased quantification of drug targets beyond visual inspection. We applied a
149 recently developed AI-based Cartography of Ensembles (ACE) pipeline to analyze these datasets
150 (Figure 2A-2C, Supplementary Table 1)[52,53]. After registering the 3D volumes onto the Allen
151 Brain Atlas (ABA)[54] (Figure S2A), we first quantified the fluorescence intensity (indicating drug
152 binding abundance) across all brain regions (Figure 2B). Intensity quantifications revealed high
153 drug abundance in the pons, thalamus, and midbrain (summarized in Figure 2D and S2B), for
154 example, in the locus coeruleus (LC) and nucleus accumbens (ACB) (Figure 2D).

155 Leveraging the cellular resolution imaging provided by vCATCH, we next sought to unbiasedly
156 quantify the cell targets of pargyline. Empowered by a large training dataset and AI-based
157 segmentation algorithms, the ACE pipeline was used to identify brain-wide pargyline-yne-
158 positive cells (Figure 2C, S2C, and S2D). After segmentation, the cell density patterns generally
159 agreed with fluorescence intensity profiles. Since pargyline targets both MAO-A and MAO-B
160 with similar affinities[43], we performed MAO-A and MAO-B immunofluorescence and confirmed
161 that pargyline-yne-targeted cells overlap with either or both MAO-A/B protein expression,
162 consistent with the known affinity of pargyline on these two targets (Figure S2E)[43].

163 Interestingly, unbiased ACE analysis also revealed unexpected differences between fluorescence
164 intensity and cell counts. For example, the ACB, bed nuclei of the stria terminals (BST), and the
165 medial group of the dorsal thalamus (MED), showed extensive drug intensity, but almost no
166 neuronal structures were detected in these areas, indicating drug binding may be associated with
167 non-soma structures (Figure S1E, 2D, and 2E). Conversely, MAO-B positive drug-bound cells
168 were identified in the dorsal raphe (DR) and the anterior tegmental nucleus (AT) of the pons
169 (Figure S1E, 2D, 2E, and S2E) despite the lack of overall fluorescence intensity increase,
170 suggesting sparse neuronal populations were targeted by pargyline in these regions.

171 Taken together, by using a pargyline probe as a proof-of-concept, we demonstrated that vCATCH
172 is compatible with clearing methods and high-throughput computational pipelines to unbiasedly
173 quantify regional enrichment and cellular targets of covalent drugs in the mouse brain, which
174 was not possible with traditional methods.

vCATCH profiles whole-body targets of covalent cancer drugs

We next sought to expand vCATCH target mapping to whole animals across all organ systems. We selected two widely used cancer drugs for this study: afatinib (Gilotrif®) and ibrutinib (Imbruvica®), both covalent targeted kinase inhibitors (TKIs)[55,56]. Since their landmark approval in 2013, the epidermal growth factor receptor (EGFR) inhibitor afatinib and Bruton's tyrosine kinase (BTK) inhibitor ibrutinib, respectively, have transformed the clinical landscape of non-small cell lung cancer (NSCLC)[57,58] and B cell malignancies[59-61]. Despite their popularity, however, there are several well-documented side effects of TKIs that are poorly understood. For example, ibrutinib is associated with cardiovascular and bleeding risks[62,63]. Although many studies have been conducted to understand the protein targets (or off-targets) of ibrutinib[64-68], the *in vivo* tissue(s) or cell types responsible for such toxicities remain elusive. The ability to unbiasedly profile TKI targets *in vivo* at cellular resolution could provide crucial insight to bridge this gap.

We injected previously established clickable probes of ibrutinib-yne (10 mg/kg, i.p.) and afatinib-yne (10 mg/kg, i.p.) in wild-type mice (Figure S3A)[69]. These probes largely retain the pharmacokinetics and tissue distribution of the parent compounds (Figure S3B and S3C). We established a vCATCH protocol with 4-day PRCS and 5 rounds of RIR to homogenously label a whole HYBRiD-cleared, one-week-old mouse with a total volume of more than 38x18x12mm (Figure 3A-3D). Labeling specificity was again confirmed by controlling for each vCATCH component in the lung or the spleen (canonical EGFR+ or BTK+ tissue, respectively, Figure S3D-S3G).

We then performed whole-body lightsheet imaging of these 1-week-old mouse bodies (Figure 3B-3H, S3H, S3J-S3M) and visualized them through virtual reality (VR) (Supplementary Video 2 and 3)[70]. The performance of vCATCH in young mice motivated us to expand its capacity to adult bodies. Enhanced by 7-day PRCS and 8 rounds of RIR (Figure 3A), we labeled ibrutinib-yne in an adult mouse with a total volume of 70x30x20 mm. Although this whole volume exceeded the field of view (FOV) of most commercial lightsheet systems, we were able to focus on the center of the torso (37x30x8 mm, Figure 3I) and examined drug distribution of major organs (Figure 3J-3M, and S3I).

Globally, the binding of both TKIs was highly abundant in the liver and the gastrointestinal tract (Figure 3F, 3K, S3K and S4). In the upper chest, ibrutinib showed elevated heart binding compared to the lung, whereas the opposite enrichment was observed with afatinib (Figure 4A, 4F and 4K). These findings are consistent with the general distributions of these two drugs based on established autoradiography data from the Food and Drug Administration[71,72].

The resolution of vCATCH further allowed us to examine intra-organ drug distribution in detail. In several organs, afatinib and ibrutinib displayed similar intra-organ distributions; for example, both drugs were selectively enriched in the villus in the small intestine (Figure S4). Higher-magnification confocal imaging further revealed detailed drug-bound structures across organs (Figure 5A-5F). For example, both drugs strongly bound to circular structures in the bone (Figure 5A and 5D), which were also observed in the spleen (Figure S4). Subsequent lectin staining identified these structures as vasculatures (Figure S5A and S5B). Considering that BTK expression is generally absent in blood vessels[73], we suspect the vasculature associated binding may be implicated in the off-target toxicity associated with ibrutinib[62,63].

In other organs, we observed distinct target distributions between these two drugs. For example,
robust tissue enrichments in the inner stomach lining and along the nasal tract were associated
with ibrutinib but not afatinib (Figure S4A and S4B). Furthermore, we discovered widely spread
afatinib targets in the lung, whereas pulmonary ibrutinib labeling was restricted along the
bronchi (Figure 4B, 4C, 4G, 4H, 4L, and 4M).

Interestingly, although both drugs were bound to the kidney, they displayed different intra-organ
patterns (Figure S4). Afatinib was associated with glomeruli in the cortex and near the cortex-
medulla interface, whereas ibrutinib was more localized to cortical tubular-like structures (Figure
5B and 5E). Similarly, our global organ-level imaging (Figure 4D, 4I, and 4N) showed that both
TKIs were enriched in the liver, in agreement with the literature[71,72]. However, higher resolution
vCATCH revealed differences in their intra-organ distribution: afatinib showed crown-like
signals throughout the hepatic tissue, whereas ibrutinib was associated with sparsely scattered
individual cells, as demonstrated by both lightsheet (Figure 4E, 4J, and 4O) and confocal
imaging (Figure 5C and 5F).

Multiplexed molecular characterizations of drug-enriched cellular structures

Guided by the drug distribution patterns, we next sought to molecularly characterize drug-
targeted cells. We first focused on afatinib binding in the lung and ibrutinib binding in the spleen
based on their canonical "on-target" engagements[73]. As expected[69], afatinib-yne and ibrutinib-
yne signals were primarily on EGFR+ and BTK+ cells, respectively, as determined by
immunostaining (Figure 5G and 5H). Importantly, pretreating the animals with the parent
compound of each probe robustly reduced probe labeling (Figure 5G-5J), further confirming the
specificity of these probes.

In the liver, we observed enriched yet distinct binding patterns of afatinib-yne and ibrutinib-yne
(Figure 5C and 5F). We next performed molecular analysis to determine the targeted cell types of
each probe. First, consistent with high EGFR expression in the hepatocytes[73], we found that the
majority (59.17±10.91%) of afatinib-bound cells were hepatocytes, followed by small fractions
of macrophages (F4/80+) and T cells (CD3e+) (Figure 5K-5M, and S5C). By contrast, only a
small fraction (13.95±1.32%) of ibrutinib-bound cells were hepatocytes, while a larger
proportion (39.53±4.42%) of these cells were F4/80+ macrophages (Figure 5N-5P), consistent
with known BTK expression in liver macrophages[73]. Interestingly, the largest fraction
(52.78±3.25%) of ibrutinib-bound cells were CD3e+ T cells despite their low BTK expression
(Figure 5O and S5C-E)[73], suggesting non-BTK engagement could be involved (see Discussion).

Beyond the liver, we were also able to identify and quantify afatinib-enriched immune cell
populations in other tissues such as the lung (Figure S5F-S5H). In summary, our vCATCH data
revealed that the two TKIs are enriched in distinct cell types within individual tissues and organs,
highlighting vCATCH's capacity to profile drug targets across global, organ, tissue, and cellular
scales.

DISCUSSION

Covalent TKIs have reentered the stage for cancer treatment and broadened the horizon for the
druggable space as demonstrated by the recent developments of mutant selective EGFR inhibitor
osimertinib, and KRAS G12C inhibitors sotorasib and adagarasib[55]. However, despite the

reignited enthusiasm, the concern for toxicity (for example, cardiotoxicity and bleeding for
ibrutinib) calls for continuous innovations in characterizing their *in vivo* targets to guide future
covalent drug development.

Here, by treating samples with excessive copper (PRCS) and performing multiple rounds of click
reactions (RIR), we established a new method, vCATCH, which enabled robust covalent TKI
mapping across the whole mouse body. This method only requires simple passive incubations
and therefore offers reliability, reproducibility, and high throughput for cohort-scale biological
and pharmacological analyses. Using existing pharmacokinetic data from the Food and Drug
Administration (FDA) as a reference, vCATCH confirmed organ-level drug distribution.
Critically, vCATCH enabled us to identify drug engagement with intra-organ cellular resolution
that is typically inaccessible through conventional methods. The capacity to resolve drug
engagement across molecularly defined cell populations, tissue architectures, and organ systems,
once combined with single-cell and spatial omics tools, presents transformative opportunities for
spatially defined, cell-type-specific drug action characterization.

Our work so far has focused on wild-type mice for method development. By examining drug
binding in healthy mice, we uncovered patterns potentially associated with off-target engagement
and toxicities. For example, ibrutinib is well known to be associated with cardiac toxicity such as
atrial fibrillation (AF), but the mechanism behind this association remains incompletely
understood[64-67]. Although animal studies suggested AF induction would likely require chronic
administration[66], our study revealed clear ibrutinib enrichment in the heart after a single
injection, suggesting an intrinsic enrichment of ibrutinib for the cardiovascular system. It would
be of great interest to further examine ibrutinib engagement in AF models to better understand its
cardiotoxicity mechanism. In addition to AF, bleeding is another major ibrutinib risk factor[62,63].
Despite low BTK expression in endothelial cells[73], we observed ibrutinib engagement in the
vasculature, a finding that could have implications for ibrutinib-associated hypertension and help
guide subsequent molecular identification work to further characterize unique ibrutinib targets in
endothelial cells[75].

Moreover, we observed ibrutinib binding in T cells that normally do not express BTK. This
binding suggested non-BTK targets of ibrutinib at the dose we used in our studies. Indeed,
previous ABPP studies in Ramos cells have identified human epidermal growth factor receptor 2
(HER2)[69] as an ibrutinib target. *In vitro* kinase activity profiling also identified interleukin-2
inducible T-cell kinase (ITK) and Janus kinase 3 (JAK3) with nM affinity to ibrutinib[68]. All of
these non-BTK affinities could contribute to our observed T cell binding.

On the technology side, vCATCH represents a new strategy to use biorthogonal click chemistry
to label large tissues, in contrast to most mainstream labeling methods (e.g., immunostaining,
DNA/RNA hybridization, etc) that primarily rely on biological affinities, such as antibody-
antigen, biotin-streptavidin, and complementary DNA/RNA interactions. Click chemistry allows
us to easily control labeling kinetics by simple buffer exchanges, achieving uniform probe
penetration in ultra-large tissue without delicate procedures or apparatus. The covalent click
chemistry linkage also means that the same sample can withstand harsh and prolonged
processing and imaging operations, opening numerous possibilities to multiplex vCATCH with
other spatial omics methods.

Finally, it is a major challenge in whole-mount tissue staining to balance the high probe
concentration (such as antibody) required for tissue penetration and its associated background

resulting from the nonspecific interactions between the probe and endogenous biomolecules. In this regard, the biorthogonal nature of click chemistry could better tolerate a wide range of labeling conditions without increasing noise. Currently, click chemistry-based imaging studies are primarily associated with examining transcription/translation[77,78]. By demonstrating vCATCH's capacity to map covalent drug binding across the entire mouse body, we propose that similar principles can be applied in ultra-large 3D tissues to visualize substrates including but not limited to protein[79], nucleic acid[80], glycan[81], lipid[82], and drug molecules[21]. For example, a recent study (Click3D) has demonstrated that the original CATCH or similar strategies can label endogenous RNA and dyes in the mouse brain and tumor[83], suggesting such approaches could be adopted in future biomedical studies, although it was limited to small, individual organs.

LIMITATIONS OF THE STUDY

Focusing on method development, our current study used healthy wild-type mice, in which the engagement of TKIs could differ from tumor-bearing animals. Given the robustness and simplicity of the vCATCH pipeline, however, we believe other researchers with oncology expertise could readily apply vCATCH in various cancer models. We also note that due to the size limit of the current lightsheet microscopes, our imaging scope is limited to the center of adult mouse torsos. We foresee that the challenge can be addressed with emerging hardware and reconstruction pipelines[31,84,85]. Moreover, all clickable TKI probes used in this study have been rigorously validated by published medicinal chemistry and chemoproteomics studies[69]. However, we acknowledge that generating alkyne derivatives with similar absorption, distribution, metabolism, and excretion (ADME) properties of the original drugs requires dedicated efforts. We note that our ibrutinib-yne probe has a higher liver distribution than the parent drug, despite the latter being known to be enriched in the liver based on the FDA[72]. This difference could confound our interpretations of ibrutinib-bound immune cells (Figure 5). Lastly, as CATCH readout can potentially reflect both on- and off-target binding and metabolite intermediates (i.e., glutathione conjugation for cysteine-targeting drugs), it is important to note that the design of future CATCH probes for other drugs must be verified by similar PK and target identification approaches; although it is becoming common to generate and characterize clickable alkyne analogs during the development of new covalent inhibitors.

RESOURCE AVAILABILITY

Lead Contact

Further information and requests for resources and reagents should be directed to and will be fulfilled by the lead contact, Li Ye (liye@scripps.edu)

Materials availability

All unique/stable reagents generated in this study are available from the lead contact with a completed materials transfer agreement.

Data and code availability

All data are available in the main text or the supplementary materials. Any additional information required to reanalyze the data reported in this paper is available from the lead contact upon request. Relevant code for the study is available from https://github.com/yelabscripps/vCATCH.

ACKNOWLEDGMENTS

We thank all members of the Ye lab and the Dorris Neuroscience Center for their support and feedback. We thank Barry Sharpless for suggestions. This work was supported by NCI IMAT program (CA281918 to L.Y. and B.F.C), NIH Director's New Innovator Award (DK128800 to L.Y.), and NIDA (DA059393 to L.Y.). L.Y. was also supported by the HHMI, NIDDK, NIMH/BRAIN, Chan Zuckerberg Initiative, and the Dana, Whitehall, Baxter, and Abide-Vividion Foundations. J.J.M received funding from the NIH (R01HL141466, R01HL155990, R01HL156021, P01HL141084). Z.P. and H.S. were supported by the Dorris Scholar Award. L.H.S is supported by an NIH Medical Scientist Training Grant T32GM007198-49. M.D.M.-G. was supported by Stanford Wu Tsai Human Performance Alliance. P.W. is supported by the NIH (R35GM139643).

AUTHOR CONTRIBUTIONS

L.Y., Z.P. conceived the study. L.Y., Z.P., V.H.L., P.W., J.J.M., J.Z.L., M.G., and B.F.C. designed the experiments. Z.P., V.H.L., C.C.W., C.G., L.H.S., M.Y., A.S., and S.X. performed the experiments and analyzed the data. A.A., A.R., H.S., and M.G. performed the ACE analysis. Z.P., V.H.L., and M.D.M.-G. performed the PK study. Z.P., V.H.L., and C.R. performed the parental drug blocking study. L.Y. and B.F.C. acquired funding. L.Y. supervised the study. L.Y., Z.P., V.H.L., and C.C.W. wrote the manuscript with input from all authors. All authors reviewed and provided feedback on the manuscript.

DECLARATION OF INTERESTS

The design, steps, and applications of vCATCH are covered in pending patent application material from The Scripps Research Institute.

FIGURE LEGENDS:

Figure 1. Establishing the principles for deep tissue click reactions
(A) Brain-wide characterization of click labeling in pargyline-yne-treated hemispheres (10 mg/kg, 1 h, i.p.). Pre-reaction copper saturation (PRCS) and repeated iterations of reactions (RIR, 1 hour per iteration) enhanced labeling depth and generated homogeneous click labeling across the whole hemisphere.
(B) Schematics of copper competition in tissue. Copper binding sites could chelate copper and limit the available free copper for CuAAC catalyst conversion, thereby impairing click labeling efficiency deep in tissues.
(C) Quantification of click labeling depth along the marked lines in Figure 1A. Four hemispheres in each condition.
(D) Representative 3D rendering of vehicle and pargyline-yne treated hemispheres.
(E) 3D rendering of pargyline-yne treated hemispheres showing the total imaging volume.
(F) Schematics of imaging depth shown in Figure 1G, highlighting changes in different brain regions covered.
(G) Sagittal 2D views of pargyline-yne treated hemispheres at imaging depth Z=1.5 mm and 3.0 mm as indicated in Figure 1F.
(H) Zoomed-in views showing pargyline-yne enriched neurons and brain regions across the hemisphere.
Data are plotted as mean±SEM. Scale bars: 1000 μm (A); 2000 μm (D and G); 100 μm (H)

Figure 2. ACE analysis of pargyline-yne targets across the hemisphere.
(A) Schematics of the ACE analysis pipeline for brain wide pargyline-yne (10 mg/kg, 1 h, i.p.) target identification.
(B) Representative regional drug intensity quantification in atlas space. White boundary indicates brain regions by Allen Brain Atlas (ABA). The LC showed the highest overall labeling intensity in pargyline-yne hemispheres. Raw intensity values indicated by scale bar.
(C) Representative segmented neurons in atlas space. Known pargyline-yne-bound neurons in the pons and hypothalamus were used as positive controls. Additional neuronal structures (i.e. in the medulla) were identified. The scale bar indicates raw segmented neuron density in each 10x10x10 μm voxel.
(D-E) Heatmap of normalized pargyline-yne labeling intensity (D) and normalized pargyline-yne positive neuron density (E). A total of 5 pargyline-yne-treated mice (rows) and 184 brain regions (columns) were included for analysis. For individual mice, intensity and density values were normalized by the average intensity and density of all the regions from each mouse. Purple labels indicate regions with only elevated intensity. Blue labels indicate regions with only elevated neuronal density. Red labels indicate regions with shared elevated intensity and neuronal density. Barrington's nucleus (B*) was later found to be a false positive (See Figure S2F). The full dataset and acronyms can be found in Supplementary Table 1. EPI, epithalamus; PMD, dorsal premammillary nucleus; PMV, ventral premammillary nucleus; AT, anterior tegmental nucleus; SLC, subceruleus nucleus; SLD, sublaterodorsal nucleus; AMB, nucleus ambiguus; VS, ventricular system.

Figure 3. vCATCH reveals whole-body TKI distribution with high spatial resolution.
(A) Workflow of vCATCH whole-body labeling with estimated time of each step.
(B) Schematic of imaging depth.

(C) Coronal (x-y) views of vCATCH whole-body labeling in ibrutinib-yne (10 mg/kg, 1 h,
i.p.) treated juvenile mouse across different z-depths.
(D) Schematic of lightsheet acquisition FOV, represented by red dotted box, for juvenile
whole body.
(E) Representative ibrutinib-yne distribution overviews in a juvenile mouse with digital plane
slicing in 3D volume.
(F-H) Coronal (F), sagittal (G), and transverse (H) views of a juvenile mouse as shown in
Figure 3E.
(I) Schematic of lightsheet acquisition FOV, represented by a red dotted box, for adult mouse
torso.
(J) Representative ibrutinib-yne distribution overview in an adult mouse torso with digital
plane slicing in 3D volume image.
(K-M) Coronal (K), sagittal (L), and transverse (M) views of adult mouse as shown in Figure
3J.
Dashed lines indicate body boundary. Scale bars: 4000 µm (C, E, F, J, K); 2000 µm (G, H, L,
M).

Figure 4. vCATCH reveals intra-organ TKI distribution

(A-E) Afatinib-yne binding in the upper chest (A), lung (B and C), and liver (D and E) of
juvenile mice.
(F-J) Ibrutinib-yne binding in the upper chest (F), lung (G and H), and liver (I and J) of
juvenile mice.
(K-O) Ibrutinib-yne binding in adult mice. Figures showing the upper chest (K), lung (L and
M), and liver (N and O).
Arrows indicate drug-enriched structures. Dashed lines indicate tissue boundary. Scale bars:
1000 µm (A, B, F, G); 500 µm (C, D, H, I, M, O); 100 µm (E, J); 2000 µm (K, L, N).

Figure 5. Cell type characterization of TKI engagement in drug-enriched tissues.

(A-F) Afatinib-yne (A-C) and ibrutinib-yne (D-F) characterizations in the bone (A, D), kidney
(B, E), and liver (C, F).
(G) Afatinib pretreatment (25 mg/kg, 1 h, i.p.) in the lung co-stained with EGFR.
(H) Ibrutinib pretreatment (25 mg/kg, 1 h, i.p.) in the spleen co-stained with BTK.
(I-J) Quantification of afatinib-yne labeling intensity of EGFR-positive pixels in the lung (I)
and ibrutinib labeling intensity of BTK-positive pixels in the spleen (J). n = 4 mice for each
condition. One-way ANOVA with Tukey's multiple comparisons test. p-values plotted in the
graph.
(K) Representative images of afatinib-yne+ hepatocytes (top row) and afatinib-yne+
macrophages (F4/80+, bottom row).
(L-M) Afatinib-yne positive cell type (L) and cell type specific intensity (M) quantifications.
n=4 biological replicates, 4-6 FOV were acquired for each mouse.
(N) Representative images of ibrutinib-yne+ hepatocytes (top row) and ibrutinib-yne+
macrophages (F4/80+, bottom row).
(O-P) Ibrutinib-yne positive cell type (O) and cell type specific intensity (P) quantifications.
n=4 biological replicates, 4-6 FOV were acquired for each mouse.
All values are shown as mean ± SEM. Dashed lines represent the cell boundary. Detailed cell
type quantification is in Supplementary Table 1. Scale bars: 200 µm (A, C, global view; B, E,
zoomed-in view; G, H); 20 µm (A, C, zoomed-in view); 1000 µm (B, E, global view); 500
µm (C, F, global view); 50 µm (C, F, zoomed-in view); 10 µm (K, N).

Figure S1. vCATCH development, related to Figure 1.

(A) Chemical structure of the monoamine oxidase A/B (MAO A/B) inhibitor pargyline, fatty acid amide hydrolase (FAAH) inhibitor PF7845, and their alkyne analogs used in the study. Both analogs have been verified in previous CATCH and ABPP studies.

(B) CATCH in 100-micron vehicle or PF7845-yne treated (1 mg/kg, 1 h, i.p.) brain sections. Individual components (Cu, ligand, N_3) were removed during the click reaction incubation. Representative images were taken at the primary somatosensory cortex (S1, imaging depth Z=10 and 50 microns).

(C) Quantification of CATCH intensity in Figure S1B. Six tissue sections (one field of view per section) were used for each condition. Intensity at Z=10 or 50 microns was measured and then normalized to the average vehicle labeling intensity at Z=10 or 50 microns, respectively. Paired t-test in each condition. P-values are plotted in the graph. n = 6 tissue sections for each condition.

(D) Transverse 2D views of hemispheres in Figure 1E.

(E) Coronal 2D views of hemispheres in Figure 1E. Zoomed-in views showing pargyline-yne enriched structures across the whole brain.

Statistics determined by two-tailed paired t-test in (C). Dashed lines indicate tissue boundary.

Scale bars: 20 µm (B); 2000 µm (D); 1000 µm (E, coronal view); 100 µm (E, zoom-in view)

ACB, nucleus accumbens; MED, medial group of the dorsal thalamus; PMV, ventral premammillary nucleus; DR, dorsal nucleus raphe; AT, anterior tegmental nucleus; LC, locus coeruleus.

Figure S2. ACE analysis and validation, related to Figure 2.

(A) Evaluation of ACE registration and warping algorithms. Lightsheet data were registered to the ABA using ACE. Left and right panels show data overlaid on labels for two subjects in each group. Segmentation maps from ACE were voxelized and warped to ABA 25-µm resolution. Panels show a randomly selected subject from the pargyline group.

(B) Atlas plot of the average intensity difference of vCATCH labeling intensity of vehicle vs. pargyline-yne treated samples. The greatest labeling intensity elevation is denoted in yellow.

(C) Atlas plot of the average segmented neuronal density (number of cells per mm^3) of vehicle vs. pargyline-yne treated samples. The greatest density elevation is denoted in yellow.

(D) Atlas plot of the average [-log(p-value)] of segmented neuronal density in vehicle vs. pargyline-yne treated samples. The greatest statistical significance is denoted in yellow.

(E) MAO-A and -B immunostaining in pargyline-yne positive regions.

(F) Histology validation of pargyline-yne binding in the LC. The boundary of LC and B is identified by NeuN and tyrosine hydroxylase (TH) staining using a published protocol[86]. Pargyline-yne binding is restricted in LC, suggesting that "B" was likely a false positive from ACE due to its small size.

Scale bars: 200 µm (E and F).

Figure S3: Whole body vCATCH visualization of TKI binding, related to Figure 3.

(A) Chemical structures of afatinib, ibrutinib, and their alkyne analogs.

(B) PK studies of plasma drug concentrations. Plasma samples taken 15, 30 and 60 min after drug injection (10 mg/kg). Two parental afatinib-60 min samples were excluded as outliers. n=4 mice for all time points except parental afatinib at 60 min (n = 2 mice).

(C) PK studies of tissue drug concentrations at 60 min post injection (10 mg/kg). n=4 mice for afatinib-yne, ibrutinib, and ibrutinib-yne; n=2 mice for afatinib.

(D-E) Click reaction specificity validation in afatinib-yne treated lung (D) and ibrutinib-yne
treated spleen (E).
(F-G) Quantifications of click labeling intensity in afatinib-yne-treated lung (F) and ibrutinib-
yne treated spleen (G). Intensity normalized to vehicle full click condition. Three independent
tissues were used for quantification. Two-way ANOVA with Šidák multiple comparisons test.
(H) Representative 3D lightsheet image volume of whole-body drug distribution (10 mg/kg,
1h, i.p.) in juvenile vehicle, afatinib-yne, and ibrutinib-yne mice. D, dorsal; V, ventral; S,
superior; I, inferior; R, right; L, left.
(I) Adult whole torso 3D overview of mice treated with vehicle and ibrutinib-yne (10 mg/kg,
1h, i.p.).
(J) Afatinib-yne distribution overviews with digital plane slicing in 3D volume image.
(K-M) Coronal (K), sagittal (L), and transverse (M) views of mice injected with afatinib-yne
as shown in Figure S3J.
Data are plotted as mean ± SEM. Dashed lines indicate tissue boundary. Scale bars: 200 μm
(D, E); 4000 μm (H, I, J, K); 2000 μm (L, M).

Figure S4: Organ level TKI binding, related to Figure 4.
(A) Representative organ views from afatinib-yne treated mice. Afatinib-yne enrichment was
observed in small intestine villi, spleen central arterioles (orange arrows), and kidney cortex.
No significant drug enrichment observed in the stomach lining or nasal tract. Scale bars: small
intestine, kidney, stomach, and nasal tract, 500 μm; spleen, 250 μm.
(B) Representative organ views from ibrutinib-yne treated mice. Ibrutinib-yne enrichment was
observed in small intestine villi, spleen central arterioles (green arrows), kidney, and inner
stomach lining. Scale bars: kidney and stomach, 500 μm; small intestine and nasal tract, 300
μm; spleen, 200 μm.
(C) Representative organ views from ibrutinib-yne treated adult mice. Ibrutinib-yne
enrichment was observed in small intestine villi, spleen central arterioles (green arrows), and
kidney. Scale bars: small intestine, 1000 μm; spleen, 500 μm; kidney, 1500 μm kidney.

Figure S5: TKI positive cell type characterization, related to Figure 5.
(A-B) Lectin staining in bone (A) and spleen (B) sections after CATCH labeling.
Representative images showing lectin overlaps TKI enriched labeling structures.
(C) High resolution afatinib-yne and ibrutinib T cell characterization in the liver.
Representative images show CD3e+ cells that are afatinib- (left) and ibrutinib+ (right).
Arrows indicate CD3e+ cells.
(D) Cell type characterizations of ibrutinib-yne positive liver T cells. Samples are co-stained
for helper T cells (CD4) and cytotoxic T cells (CD8).
(E) Quantifications ibrutinib-yne positive T cell subtypes in the liver. n=4 biological
replicates, 11-17 FOV were acquired for each mouse.
(F) Global characterization of afatinib-yne binding in the lung. Zoom-in views showing
discrete afatinib-yne enriched structures scattered throughout the lung.
(G) High resolution afatinib-yne cell type characterization in the lung. Representative images
showing afatinib-yne enriched F4/80 positive macrophages (top) and CD19 positive B cells
(bottom).
(H) Cell type characterization of afatinib-yne enriched cells in the lung. n=4 biological
replicates, 5-6 FOV were acquired for each mouse.

All values are shown as mean ± SEM. Detailed cell type quantification is in Supplementary
Table 1. Scale bars: 20 µm (A and B); 10 µm (C, D, and G); 1000 µm (F, global view); 50 µm
(F, zoomed-in view).

STAR METHODS

Key Resources Table

EXPERIMENTAL MODEL AND SUBJECT DETAILS

Mouse model

Mice were group-housed on a 12-hr light-dark cycle and fed a standard rodent chow diet. Both
male and female mice were used for the study. Six to nine-week-old WT C57BL6J mice were
used for vCATCH method development in 100-micron sections and hemispheres. One-week-old
and eight-week-old WT C57BL6J mice were used for vCATCH in the whole mouse body. All
experimental protocols were approved by the Scripps Research Institute Institutional Animal
Care and Use Committee and were in accordance with the guidelines from the NIH.

Sample collection

Alkyne drugs were administered to mice in a vehicle of 10% DMSO and 2% Tween-80 in saline
for i.p. injections at the indicated dose. 1 hour (for most direct labeling experiments) or 4 hour
(for pretreatment test) after injections, mice were heavily anesthetized with isoflurane and then
transcardially perfused with ice-cold PBS, followed by ice-cold 4% PFA solution. Mouse organs
were carefully dissected and then post-fixed in 4% PFA solution overnight at 4℃. Tissues were
washed with PBS. Mouse brains were embedded in 2% agarose and sectioned with a vibratome
(Leica VT1000S) at 100-micron thickness. For hemispheres, brains were cut along the midline
with a razor knife. Peripheral organs from one-week-old mice were dehydrated in 30% sucrose at
4℃ prior to sectioning. Samples were then carefully dried with a kimwipe and embedded in
O.C.T. compound. After reaching equilibrium at -20℃, samples were cut at 100-micron
thickness with a cryostat (Leica CM 1860). All obtained samples were stored in PBS with 0.02%
sodium azide at 4℃ for further processing.

For mice used in whole-body drug distribution labeling, the skin was carefully removed
immediately after perfusion, and the peritoneum was cut down the midline to facilitate clearing
and click reactions. Mice were post-fixed in 4% PFA solution for two days at 4℃ followed by 7-
day decalcification treatment in 10% EDTA/15% Imidazole at 4℃. Then, samples underwent
decoloring treatment with 25% N, N, N', N'-Tetrakis(2 Hydroxypropyl)ethylenediamine
(Quadrol) in 1x PBS at 37°C with gentle agitation for 7 days. After PBS washes to remove
Quadrol, mouse whole bodies were stored in PBS with 0.02% sodium azide at 4℃ for further
processing.

For mice used in PK studies, after anesthesia, blood was collected via cardiac puncture and
immediately stored in a heparin tube and placed on ice. Plasma was separated by centrifugation
at 5,000 rpm for 5 min at 4°C. Next, 150 µL of a 2:1 mixture of acetonitrile/methanol was added
to 50 µL of plasma. The mixture was then centrifuged at 15,000 rpm for 10 min at 4°C, and the
supernatant was transferred to a liquid chromatography-mass spectrometry (LC-MS) vial. Tissue

samples were dissected out and weighed. Next, tissues were homogenized in water at 250 mg/ml
using a Homogenizer at 4°C. The mixture was then centrifuged at 15,000 rpm for 10 min at 4°C.
50 μL of the supernatant was then collected, and 150 μL of 2:1 mixture of acetonitrile/methanol
was added to the supernatant. The mixture was then centrifuged at 4°C at 15,000 rpm for 10 min,
and the supernatant was transferred to an LC-MS vial. To generate a standard curve, compounds
were dissolved in a 2:1:1 mixture of acetonitrile/methanol/water at 10, 1, 0.1, 0.01, 0.001 μM for
MS analysis.

Tissue clearing

CLARITY

100-micron sections were cleared with CLARITY as previously described[22,87]. Briefly, samples
were incubated in A1P4 hydrogel (1% acrylamide, 0.125% Bis, 4% PFA, 0.025% VA-044
initiator (w/v), in 1X PBS at 4°C for CLARITY embedding. After overnight incubation with
gentle agitation, samples were then degassed and polymerized at 37°C for 4 hours with gentle
agitation. Samples were removed from hydrogel and washed with 6% PBS-SDS (pH=7.0) at
37°C for two days. After clearing, samples were washed with PBST (pH=7.0 with 0.2% Triton-
X100, same for the following) three times (10 min each time) at RT to remove residue SDS.
Samples were briefly washed with PBS and then stored in PBS with 0.02% sodium azide at 4°C.

HYBRiD

Mouse hemispheres and bodies were cleared with HYBRiD as previously described[27]. In brief,
samples were dehydrated in an increasing THF/25% Quadrol gradient (50%, 70%, 80%, 100%,
100%; 1 hour per step for hemispheres, 3-12 hours per step for whole bodies), delipidated three
times in DCM (1 hour each for hemispheres, 3-12 hours each for whole bodies), and rehydrated
in a decreasing THF/25% Quadrol gradient (100%, 100%, 80%, 70%, 50%; 0.5 hour per step for
hemispheres, 1.5-6 hours per step for whole bodies). After clearing, samples were extensively
washed with PBS to remove residual organic solvent before hydrogel embedding. Samples were
equilibrated in A1P4 hydrogel (3 days for hemispheres, 7-10 days with one refreshment for
whole bodies) at 4°C on a shaker, then degassed and polymerized at 37°C for 4 hours with gentle
agitation. Thereafter, samples were passively cleared with LiOH Boric Buffer with 6% SDS
(pH=9) until transparent (1 week for hemispheres, 6-10 weeks for whole bodies). Samples were
extensively washed in PBST to remove residual SDS and then stored in PBS with 0.02% sodium
azide at 4°C.

Click labeling in tissues

CATCH in 100-micron sections

100-micron sections were labeled with an updated CATCH protocol. Unless otherwise noted,
sections were incubated in 10 mM CuSO$_4$ in H$_2$O at RT overnight. Sections were then transferred
to CATCH reaction buffer containing 5 μM AZDye 647-picolyl azide, 150 μM CuSO4, 300 μM
BTTP, 2.5 mM NaAsb, and 10% DMSO in 1X PBS. After 1 hour of reaction in the dark with
minor agitation, samples were washed with buffer containing 4 mM EDTA in PBST (PBST-
EDTA, same for the following) at RT for 3x10 min. Tissue samples were then stained with DAPI
(1:3000 dilution in PBST from 10 μM stock) for 15 mins at RT. Samples were then either used
for refractive index (RI) matching by RapiClear and confocal imaging or proceeded to further
immunolabeling for cell-type registration.

 vCATCH in hemispheres

 For hemisphere labeling, each hemisphere was first pre-incubated with 25 mL of 10 mM CuSO4
 in H_2O for one day at RT for PRCS. Hemispheres were transferred to vCATCH incubation buffer
 (5 μM AZDye 647-picolyl azide, 1 mM CuSO4, 2 mM BTTP, 10% DMSO in 1X PBS, 3 mL for
 each hemisphere) for 1 day at RT. Hemispheres were then transferred to the newly prepared full
 vCATCH reaction buffer (5 μM AZDye 647-picolyl azide, 1 mM CuSO4, 2 mM BTTP, 25 mM
 NaAsb, 10% DMSO in 1X PBS, 3 mL for each hemisphere) for click labeling. The vCATCH
 reaction buffer was refreshed every hour for two rounds of RIR. Hemispheres were washed with
 PBST-EDTA at RT. The washing buffer was refreshed at the end of the day and then daily until it
 became colorless. The hemispheres were then washed with PBST for 1 hour at RT. After
 washing, hemispheres were cut into 100-micron coronal sections for labeling depth
 characterizations. Alternatively, hemispheres were carefully dried with a Kimwipe and then
 incubated in 3 mL EasyIndex for RI matching and lightsheet imaging.

 vCATCH in the whole mouse

 Cleared mouse bodies were placed in 500 mL per mouse of 10 mM $CuSO_4$ in H_2O with agitation
 for 4-7 days at RT with daily buffer refreshes. After PRCS, whole-body samples were
 individually incubated in 10-40 mL of vCATCH incubation buffer for 3-5 days on a shaker at RT
 with daily buffer changes. Samples were then transferred to vCATCH reaction buffer (10-40 mL
 each) to initiate the reaction. Five to eight successive rounds of 1-hour click reactions were
 performed to thoroughly label the entire tissue depth. After click labeling, samples were washed
 with PBST-EDTA with daily refreshments until the buffer was colorless. After an additional day
 of PBST wash, mouse bodies were carefully dried with a Kimwipe and then thoroughly
 immersed in 25-40 mL EasyIndex at 37°C for thorough RI matching in preparation for lightsheet
 imaging.

 Immunostaining

 CATCH-labeled 100-micron sections were incubated with primary antibodies 1:200 diluted in
 PBST overnight at 4°C. Samples were then washed with PBST 3 x 30 min at RT prior to
 incubation with secondary antibodies diluted 1:600 in PBST. Samples were then incubated with
 secondary antibodies 1:600 diluted in PBST. Samples were incubated with secondary antibodies
 overnight and then washed with PBST 3 x 30 min at RT.

 Confocal microscopy

 100-micron samples were mounted with RapiClear for RI matching and imaged with an
 Olympus FV3000 confocal microscope. To characterize CATCH labeling depth in 100-micron
 sections, samples were imaged at the primary somatosensory cortex layer V with a 10X, 0.6 NA,
 water immersion objective (XLUMPlanFI, Olympus) at 0.414 micron/pixel resolution and a step
 size of 10 microns. To examine hemisphere labeling depth and global drug distribution within
 individual organs, samples were imaged with a 10X, 0.6 NA, water immersion objective
 (XLUMPlanFI, Olympus) at 2.49 microns/pixel resolution and a step size of 10 microns. For cell
 type characterizations, samples were imaged at the top surface with a 40X, 1.25 NA, silicone oil
 objective (UPlanSApo, Olympus) at 0.094 micron/pixel resolution and a step size of 3 microns.
 All confocal data was saved in TIFF format for further analysis.

 Lightsheet microscopy

Whole hemisphere/body samples were imaged using a lightsheet microscope from LifeCanvas
Technologies. RI-matched whole hemisphere and body samples were mounted in 1%
agarose/Easy Index for imaging. Lightsheet imaging was performed in a SmartSPIM chamber
with Easy Index matched immersion oil (RI=1.52). A 3.6x objective was used with 0.28 NA,
1.8μm/1.8μm/2μm *xyz* voxel size. Imaging was performed by illuminating samples along the
bilateral plane of symmetry. Lightsheet raw data was then destriped, stitched, and saved as TIFF
format for further conversion.

683 Image processing and analysis

Hemisphere labeling depth characterization

Hemispheres were sectioned coronally as 100 micron sections and imaged to characterize
vCATCH labeling depth. Images were first processed as maximum intensity projection. A
rectangular area (805 x 50 pixels, 2001.17 x 136.73 microns) was drawn from the cortical outer
layer inwards (from tissue surface to tissue center) as indicated in Figure 1A. The signal profile
was plotted using the 'Plot Profile' function in ImageJ as reported[27,88,89]. The fluorescence signal
profile was then normalized by the intensity value at the start of the line.

Thin section labeling depth characterization

CATCH labeling intensity across the 100-micron section was quantified as previously
reported[21,27,88]. Briefly, a 150 x 150 pixel (62.1 x 62.1 micron) ROI surrounding a labeled neuron
was generated at z = 10 μm and 50 μm at cortex layer V, and an auto threshold was applied to
measure the mean drug-positive pixel intensity as I_{signal}. The mean intensity of the remaining
pixels was used $I_{background}$. The mean labeling intensity was defined as $I_{labeling}=I_{signal}-I_{background}$.

Cell type-specific characterization

CD3e, F4/80, CD4, CD8, and CD19 positive cells were characterized by immune staining.
Hepatocytes were identified based on their large size (20-30 microns), round/oval-shaped
nucleus, and hexagonal morphology by well-trained experts. To minimize bias, individual fields
of view (FOV) were first acquired based on the DAPI and the immunostaining channel without
knowledge of the CATCH/drug channel (with the exception of CD3e+ibrutinib+ cells, where the
CD4 and CD8 channel were kept blind during imaging acquisition) to assign cell types. The
CATCH positivity was post-hoc assigned to individual cells after their cell types have been
determined, followed by overlapping quantifications. For intensity characterizations, individual
cells were selected along the boundary and cropped out for quantification. A threshold was
applied in the drug channel to analyze CATCH labeling intensity in individual cells of interest.
The full dataset of each cell population can be found in Supplementary Table 1.

Global drug intensity characterization

To measure global drug intensity across intact thin tissue sections, confocal images were first
processed as maximum intensity projections. A threshold was then applied in the counterstain
channel (DAPI or target immunostaining) to cover the whole tissue or target positive pixels. The
threshold area was selected as tissue containing pixels. The average drug channel intensity in the
selected pixels was then quantified as global drug intensity.

Volumetric drug binding visualization

Lightsheet imaged hemispheres and whole body images were converted into Imaris files and
visualized via Imaris 10.2.0. The hemisphere drug labeling video was generated by Imaris. For
VR-based whole-body drug visualization, data was converted by syGlass, and videos were made
with a Meta Quest 3 glass. For hemisphere coronal and transverse views, TIFF files were
imported as image sequences and then resliced by Fiji ImageJ. For drug distribution visualization
across organs, TIFF files were imported as image sequences and sampled at 100-micron step
size. Individual organ of interest was then cropped out for visualization.

Automated Brain-wide Cell Mapping & Signal Intensity Calculation.

We employed an AI-based Cartography of Ensembles (ACE) pipeline to generate whole-brain
segmentation maps of pargyline-yne+ cells[53]. ACE is an open-source 3D deep learning pipeline
(available through the MIRACL platform[90]; https://miracl.readthedocs.io/) to map local cell
activations using tera-voxel size light sheet microscopy datasets. Given the significant
differences in morphological features between pargyline-yne+ cells and cFos+ cells (which the
ACE models were originally trained on), we first fine-tuned the ACE vision transformer (ViT)
model. For fine-tuning, we randomly selected 225 image patches of 963 voxels each
(0.17x0.17x0.19 mm^3) from two subjects in the dataset and generated ground truth using the
Labkit annotator[91] in ImageJ/Fiji software[92]. We fine-tuned the ViT model on half of these
patches and used the other half for evaluation. The model was fine-tuned for 200 epochs using
the Adam optimizer[93] with an initial learning rate of 0.0001 and an equally weighted Dice-Cross
Entropy loss function. To account for artifacts in the dataset, the cross-entropy calculation for
"background" voxels was weighted 2× higher than for voxels annotated as pargyline-yne+ cells.
The best model, which achieved an average Dice score of 0.7 on the validation dataset, was used
to generate segmentation maps for the entire dataset. Subsequently, a shape filter was applied to
the segmentation maps to eliminate false positives such as vasculature traces in cortical regions
using the shape filter plugin in ImageJ/Fiji[94]. To distinguish individual cells, a connected
component analysis was applied to the 3D segmentation maps of each subject. Following
segmentation, we automatically registered the light sheet hemisphere data to the ABA using the
MIRACL platform[54,90]. The accuracy of the registration and warping were evaluated using
quality control checkpoints. Deformation fields obtained from the registration algorithm were
used to warp the ABA labels to the native space of each subject. Finally, we computed the
number of cells within each region at depth 6 (labels grouped to a maximum atlas ontology depth
of 6, by combining finer labels under their parent labels), by identifying the coordinates of the
center of each cell.

To map significant localized group-wise differences in pargyline-yne+ cells in an atlas-agnostic
manner, we employed ACE's cluster-wise threshold-free cluster enhancement permutation test
using a group-wise ANOVA. The segmentation maps obtained by the deep learning models were
downsampled to ABA 10-μm resolution while minimizing information loss by applying a
convolution-based voxelization procedure to the whole-brain segmentation maps (based on our
prior work[90]). Following voxelization, segmentation maps were aligned with the ABA at a 10-
μm resolution using deformation fields obtained via the registration module. The voxelated and
warped segmentation maps were then passed to the ACE statistical module (using default
parameters), which produced a cluster-wise p-value map representing significant (alpha < 0.05
after correction) localized cell activation between the pargyline-yne treated and control groups.

We computed signal intensity information for each ABA label using the "*miracl lbls stats*"
function in our MIRACL platform, which processes NIfTI images and their registered warped

ABA labels for each subject. This function extracts key intensity metrics, such as the average,
minimum, and maximum voxel intensity, across the input volumes for each label. We further
summarized it by grouping labels at depth 6 using the atlas ontology.

Brain-wide drug binding heatmap visualization

Brain regions with positive cell counts in all 5 pargyline-yne-treated hemispheres were included
for heatmap generation. Cell density (number of cells per mm^3) was obtained by dividing cell
count values by the volume of the brain region. Intensity and cell density were then normalized
by the average values in individual hemispheres. Heatmaps were generated using the Python
function 'sns.heatmap'[95], with each region displaying the normalized intensity and density across
different subjects. The height of each subplot, corresponding to different regions, is proportional
to the number of areas within that region. Code for generating heatmaps is available via
https://github.com/yelabscripps/vCATCH.

Drug PK analysis

Targeted measurements were performed using an Agilent 6470 triple quadrupole LC-MS
instrument. MS analysis was performed using AJS in positive mode. The AJS source parameters
were set as follows: the dry gas temperature was set at 250 °C with a gas flow of 12 l/min and the
nebulizer pressure at 25 psi; the sheath gas temperature was set to 300 °C with the sheath gas
flow set at 12 l/min; and the capillary voltage was set to 3,500 V. Separation of compound was
conducted using a ZORBAX RR Eclipse Plus C18 95Å LC column (Agilent 959961-902) with
reversed-phase chromatography and the column temperature was maintained at 30 °C. Mobile
phases were as follows: buffer A, 100% water with 0.1% formic acid; buffer B, 100%
acetonitrile with 0.1% formic acid. The flow rate for each run started at 95% A for 9 minutes at
0.7 ml/min, followed by a gradient starting at 95% A, changing linearly to 5% A / 95% B over
the course of 12 minutes at 0.7 ml/min. The flow rate was maintained at 5% A / 95% for 6
minutes at 0.7 ml/min. The last 3 minutes consisted of a re-equilibration back to 95% A / 5% B at
0.7 ml/min. Multiple reaction monitoring was performed for the indicated chemicals with the
listed dwell times, fragmentor voltage, collision energies, cell accelerator voltages and polarity.
We selected previously reported transitions for the quantification of ibrutinib[96] and afatinib[97].
The MS ionization parameters for the targeted metabolomics are presented in Supplemental
Table 1. Quantification of the compound concentrations was performed by generating an external
standard curve with known concentrations of each compound. Compound standards were
analyzed alongside the biological samples using the same targeted triple quadrupole LC/MS
method in the same run. A calibration standard curve generated from the compound standard
concentrations and total peak areas were used to calculate the concentrations of each compound.
It is worth noting that, contrary to the high liver distribution from the autoradiography study[72],
our PK data showed overall lesser ibrutinib liver enrichment (Figure S3C). The caveat may be
due to the different detection analytes (isotope tag vs. intact free drug). And the observed
ibrutinib liver enrichment may reflect strong local metabolism and potential off-target binding.

SUPPLEMENTAL INFORMATION

Supplementary Table 1: Raw data summary and key MS parameters. Related to Figure
1C, 2D, 2E, 5I, 5J, 5L, 5M, 5O, 5P, S1C, S3B, S3C, S3F, S3G, S5E, S5H.

Supplementary Video 1: 3D rendering of pargyline-yne binding across the whole mouse
hemisphere. Related to Figure 1D.

Supplementary Video 2: 3D rendering of afatinib-yne binding across the whole mouse
body. Related to Figure S3H, S3J.

Supplementary Video 3: 3D rendering of ibrutinib-yne binding across the whole mouse
body. Related to Figure S3H, 3E, S3H.

REFERENCES

1. Huang, Z., Ogasawara, D., Seneviratne, U.I., Cognetta, A.B., 3rd, Am Ende, C.W., Nason, D.M., Lapham, K., Litchfield, J., Johnson, D.S., and Cravatt, B.F. (2019). Global Portrait of Protein Targets of Metabolites of the Neurotoxic Compound BIA 10-2474. ACS Chem Biol *14*, 192-197. 10.1021/acschembio.8b01097.

2. Chang, J.W., Cognetta, A.B., 3rd, Niphakis, M.J., and Cravatt, B.F. (2013). Proteome-wide reactivity profiling identifies diverse carbamate chemotypes tuned for serine hydrolase inhibition. ACS Chem Biol *8*, 1590-1599. 10.1021/cb400261h.

3. Van Esbroeck, A.C., Janssen, A.P., Cognetta, A.B., Ogasawara, D., Shpak, G., Van Der Kroeg, M., Kantae, V., Baggelaar, M.P., De Vrij, F.M., and Deng, H. (2017). Activity-based protein profiling reveals off-target proteins of the FAAH inhibitor BIA 10-2474. Science *356*, 1084-1087. 10.1126/science.aaf7497.

4. Reinecke, M., Brear, P., Vornholz, L., Berger, B.T., Seefried, F., Wilhelm, S., Samaras, P., Gyenis, L., Litchfield, D.W., Medard, G., et al. (2024). Chemical proteomics reveals the target landscape of 1,000 kinase inhibitors. Nat Chem Biol *20*, 577-585. 10.1038/s41589-023-01459-3.

5. Offensperger, F., Tin, G., Duran-Frigola, M., Hahn, E., Dobner, S., Ende, C.W.A., Strohbach, J.W., Rukavina, A., Brennsteiner, V., Ogilvie, K., et al. (2024). Large-scale chemoproteomics expedites ligand discovery and predicts ligand behavior in cells. Science *384*, eadk5864. 10.1126/science.adk5864.

6. Shalem, O., Sanjana, N.E., Hartenian, E., Shi, X., Scott, D.A., Mikkelson, T., Heckl, D., Ebert, B.L., Root, D.E., Doench, J.G., and Zhang, F. (2014). Genome-scale CRISPR-Cas9 knockout screening in human cells. Science *343*, 84-87. 10.1126/science.1247005.

7. Zhao, J., Tang, Z., Selvaraju, M., Johnson, K.A., Douglas, J.T., Gao, P.F., Petrassi, H.M., Wang, M.Z., and Wang, J. (2022). Cellular Target Deconvolution of Small Molecules Using a Selection-Based Genetic Screening Platform. ACS Cent Sci *8*, 1424-1434. 10.1021/acscentsci.2c00609.

8. Li, H., Ma, T., Remsberg, J.R., Won, S.J., DeMeester, K.E., Njomen, E., Ogasawara, D., Zhao, K.T., Huang, T.P., Lu, B., et al. (2023). Assigning functionality to cysteines by base editing of cancer dependency genes. Nat Chem Biol *19*, 1320-1330. 10.1038/s41589-023-01428-w.

9. Shi, J., Wang, E., Milazzo, J.P., Wang, Z., Kinney, J.B., and Vakoc, C.R. (2015). Discovery of cancer drug targets by CRISPR-Cas9 screening of protein domains. Nat Biotechnol *33*, 661-667. 10.1038/nbt.3235.

10. Perrin, J., Werner, T., Kurzawa, N., Rutkowska, A., Childs, D.D., Kalxdorf, M., Poeckel, D., Stonehouse, E., Strohmer, K., Heller, B., et al. (2020). Identifying drug targets in tissues and whole blood with thermal-shift profiling. Nat Biotechnol. 10.1038/s41587-019-0388-4.

11. Liu, Z., Wang, P., Wold, E.A., Song, Q., Zhao, C., Wang, C., and Zhou, J. (2021). Small-Molecule Inhibitors Targeting the Canonical WNT Signaling Pathway for the Treatment of Cancer. J Med Chem *64*, 4257-4288. 10.1021/acs.jmedchem.0c01799.

850   12.   Ji, H., Lu, X., Zhao, S., Wang, Q., Liao, B., Bauer, L.G., Huber, K.V.M., Luo, R., Tian,
851   R., and Tan, C.S.H. (2023). Target deconvolution with matrix-augmented pooling strategy
852   reveals cell-specific drug-protein interactions. Cell Chem Biol *30*, 1478-1487 e1477.
853   10.1016/j.chembiol.2023.08.002.

854   13.   Savitski, M.M., Reinhard, F.B., Franken, H., Werner, T., Savitski, M.F., Eberhard, D.,
855   Martinez Molina, D., Jafari, R., Dovega, R.B., Klaeger, S., et al. (2014). Tracking cancer
856   drugs in living cells by thermal profiling of the proteome. Science *346*, 1255784.
857   10.1126/science.1255784.

858   14.   Stickels, R.R., Murray, E., Kumar, P., Li, J., Marshall, J.L., Di Bella, D.J., Arlotta, P.,
859   Macosko, E.Z., and Chen, F. (2020). Highly sensitive spatial transcriptomics at near-
860   cellular resolution with Slide-seqV2. Nature Biotechnology. 10.1038/s41587-020-0739-1.

861   15.   Rodriques, S.G., Stickels, R.R., Goeva, A., Martin, C.A., Murray, E., Vanderburg, C.R.,
862   Welch, J., Chen, L.M., Chen, F., and Macosko, E.Z. (2019). Slide-seq: A scalable
863   technology for measuring genome-wide expression at high spatial resolution. Science
864   *363*, 1463-1467. 10.1126/science.aaw1219.

865   16.   Bhatia, H.S., Brunner, A.-D., Öztürk, F., Kapoor, S., Rong, Z., Mai, H., Thielert, M., Ali,
866   M., Al-Maskari, R., Paetzold, J.C., et al. (2022). Spatial proteomics in three-dimensional
867   intact specimens. Cell *185*, 5040-5058.e5019. 10.1016/j.cell.2022.11.021.

868   17.   Unterauer, E.M., Shetab Boushehri, S., Jevdokimenko, K., Masullo, L.A., Ganji, M.,
869   Sograte-Idrissi, S., Kowalewski, R., Strauss, S., Reinhardt, S.C.M., Perovic, A., et al.
870   (2024). Spatial proteomics in neurons at single-protein resolution. Cell *187*, 1785-
871   1800.e1716. 10.1016/j.cell.2024.02.045.

872   18.   Hickey, J.W., Neumann, E.K., Radtke, A.J., Camarillo, J.M., Beuschel, R.T., Albanese,
873   A., McDonough, E., Hatler, J., Wiblin, A.E., Fisher, J., et al. (2021). Spatial mapping of
874   protein composition and tissue organization: a primer for multiplexed antibody-based
875   imaging. Nat Methods. 10.1038/s41592-021-01316-y.

876   19.   Shi, H., He, Y., Zhou, Y., Huang, J., Maher, K., Wang, B., Tang, Z., Luo, S., Tan, P., Wu,
877   M., et al. (2023). Spatial atlas of the mouse central nervous system at molecular
878   resolution. Nature. 10.1038/s41586-023-06569-5.

879   20.   Zeng, H., Huang, J., Ren, J., Wang, C.K., Tang, Z., Zhou, H., Zhou, Y., Shi, H., Aditham,
880   A., Sui, X., et al. (2023). Spatially resolved single-cell translatomics at molecular
881   resolution. Science *380*, eadd3067. 10.1126/science.add3067.

882   21.   Pang, Z., Schafroth, M.A., Ogasawara, D., Wang, Y., Nudell, V., Lal, N.K., Yang, D.,
883   Wang, K., Herbst, D.M., Ha, J., et al. (2022). In situ identification of cellular drug targets
884   in mammalian tissue. Cell *185*, 1793-1805 e1717. 10.1016/j.cell.2022.03.040.

885   22.   Sylwestrak, E.L., Rajasethupathy, P., Wright, M.A., Jaffe, A., and Deisseroth, K. (2016).
886   Multiplexed Intact-Tissue Transcriptional Analysis at Cellular Resolution. Cell *164*, 792-
887   804. 10.1016/j.cell.2016.01.038.

888   23.   Murray, E., Cho, J.H., Goodwin, D., Ku, T., Swaney, J., Kim, S.Y., Choi, H., Park, Y.G.,
889   Park, J.Y., Hubbert, A., et al. (2015). Simple, Scalable Proteomic Imaging for High-
890   Dimensional Profiling of Intact Systems. Cell *163*, 1500-1514.
891   10.1016/j.cell.2015.11.025.

892   24.   Mai, H., Luo, J., Hoeher, L., Al-Maskari, R., Horvath, I., Chen, Y., Kofler, F., Piraud, M.,
893   Paetzold, J.C., Modamio, J., et al. (2023). Whole-body cellular mapping in mouse using
894   standard IgG antibodies. Nat Biotechnol. 10.1038/s41587-023-01846-0.

895   25.   Wang, X., Allen, W.E., Wright, M.A., Sylwestrak, E.L., Samusik, N., Vesuna, S., Evans,
896   K., Liu, C., Ramakrishnan, C., Liu, J., et al. (2018). Three-dimensional intact-tissue
897   sequencing of single-cell transcriptional states. Science *361*. 10.1126/science.aat5691.

898   26.   Tyler, D.S., Vappiani, J., Cañeque, T., Lam, E.Y., Ward, A., Gilan, O., Chan, Y.-C.,
899   Hienzsch, A., Rutkowska, A., and Werner, T. (2017). Click chemistry enables preclinical
900   evaluation of targeted epigenetic therapies. Science *356*, 1397-1401.

901   27.   Nudell, V., Wang, Y., Pang, Z., Lal, N.K., Huang, M., Shaabani, N., Kanim, W., Teijaro,
902   J., Maximov, A., and Ye, L. (2022). HYBRiD: hydrogel-reinforced DISCO for clearing
903   mammalian bodies. Nat Methods. 10.1038/s41592-022-01427-0.

904   28.   Murakami, T.C., Mano, T., Saikawa, S., Horiguchi, S.A., Shigeta, D., Baba, K., Sekiya,
905   H., Shimizu, Y., Tanaka, K.F., Kiyonari, H., et al. (2018). A three-dimensional single-cell-
906   resolution whole-brain atlas using CUBIC-X expansion microscopy and tissue clearing.
907   Nat Neurosci *21*, 625-637. 10.1038/s41593-018-0109-1.

908   29.   Kim, S.Y., Cho, J.H., Murray, E., Bakh, N., Choi, H., Ohn, K., Ruelas, L., Hubbert, A.,
909   McCue, M., Vassallo, S.L., et al. (2015). Stochastic electrotransport selectively enhances
910   the transport of highly electromobile molecules. Proc Natl Acad Sci U S A *112*, E6274-
911   6283. 10.1073/pnas.1510133112.

912   30.   Tainaka, K., Murakami, T.C., Susaki, E.A., Shimizu, C., Saito, R., Takahashi, K.,
913   Hayashi-Takagi, A., Sekiya, H., Arima, Y., Nojima, S., et al. (2018). Chemical Landscape
914   for Tissue Clearing Based on Hydrophilic Reagents. Cell Rep *24*, 2196-2210 e2199.
915   10.1016/j.celrep.2018.07.056.

916   31.   Park, J., Wang, J., Guan, W., Gjesteby, L.A., Pollack, D., Kamentsky, L., Evans, N.B.,
917   Stirman, J., Gu, X., Zhao, C., et al. (2024). Integrated platform for multiscale molecular
918   imaging and phenotyping of the human brain. Science *384*, eadh9979.
919   10.1126/science.adh9979.

920   32.   Ku, T., Guan, W., Evans, N.B., Sohn, C.H., Albanese, A., Kim, J.-G., Frosch, M.P., and
921   Chung, K. (2020). Elasticizing tissues for reversible shape transformation and accelerated
922   molecular labeling. Nat Methods *17*, 609-613.

923   33.   Lai, H.M., Tang, Y., Lau, Z.Y.H., Campbell, R.A.A., Yau, J.C.N., Chan, C.C.Y., Chan,
924   D.C.W., Wong, T.Y., Wong, H.K.T., Yan, L.Y.C., et al. (2022). Antibody stabilization for
925   thermally accelerated deep immunostaining. Nat Methods *19*, 1137-1146.
926   10.1038/s41592-022-01569-1.

927   34.   Susaki, E.A., Shimizu, C., Kuno, A., Tainaka, K., Li, X., Nishi, K., Morishima, K., Ono,
928   H., Ode, K.L., Saeki, Y., et al. (2020). Versatile whole-organ/body staining and imaging
929   based on electrolyte-gel properties of biological tissues. Nat Commun *11*, 1982.
930   10.1038/s41467-020-15906-5.

931   35.   Zhao, S., Todorov, M.I., Cai, R., Maskari, R.A.I., Steinke, H., Kemter, E., Mai, H., Rong,
932   Z., Warmer, M., Stanic, K., et al. (2020). Cellular and Molecular Probing of Intact Human
933   Organs. Cell. 10.1016/j.cell.2020.01.030.

36. Choi, S.W., Guan, W., and Chung, K. (2021). Basic principles of hydrogel-based tissue
transformation technologies and their applications. Cell *184*, 4115-4136.
10.1016/j.cell.2021.07.009.

37. Yau, C.N., Lai, H.M., Lee, K., Kwok, A.J., Huang, J., and Ko, H. (2023). Principles of
deep immunohistochemistry for 3D histology. Cell Rep Methods *3*, 100458.
10.1016/j.crmeth.2023.100458.

38. Richardson, D.S., Guan, W., Matsumoto, K., Pan, C., Chung, K., Erturk, A., Ueda, H.R.,
and Lichtman, J.W. (2021). Tissue Clearing. Nat Rev Methods Primers *1*.
10.1038/s43586-021-00080-9.

39. Ueda, H.R., Erturk, A., Chung, K., Gradinaru, V., Chedotal, A., Tomancak, P., and Keller,
P.J. (2020). Tissue clearing and its applications in neuroscience. Nat Rev Neurosci *21*, 61-
79. 10.1038/s41583-019-0250-1.

40. Dejaegere, A., Choulier, L., Lafont, V., De Genst, E., and Altschuh, D. (2005). Variations
in Antigen− Antibody Association Kinetics as a Function of pH and Salt Concentration: A
QSAR and Molecular Modeling Study. Biochemistry *44*, 14409-14418.

41. Cann, J.R., and Clark, E.W. (1956). Kinetics of the Antigen-Antibody Reaction. Effect of
Salt Concentration and pH on the Rate of Neutralization of Bacteriophage by Purified
Fractions of Specific Antiserum1a, b. J Am Chem Soc *78*, 3627-3631.

42. Himo, F., Lovell, T., Hilgraf, R., Rostovtsev, V.V., Noodleman, L., Sharpless, K.B., and
Fokin, V.V. (2005). Copper (I)-catalyzed synthesis of azoles. DFT study predicts
unprecedented reactivity and intermediates. J Am Chem Soc *127*, 210-216.

43. Krysiak, J.M., Kreuzer, J., Macheroux, P., Hermetter, A., Sieber, S.A., and Breinbauer, R.
(2012). Activity-based probes for studying the activity of flavin-dependent oxidases and
for the protein target profiling of monoamine oxidase inhibitors. Angew Chem Int Ed
Engl *51*, 7035-7040. 10.1002/anie.201201955.

44. Niphakis, M.J., Johnson, D.S., Ballard, T.E., Stiff, C., and Cravatt, B.F. (2012). O-
hydroxyacetamide carbamates as a highly potent and selective class of endocannabinoid
hydrolase inhibitors. ACS Chem Neurosci *3*, 418-426. 10.1021/cn200089j.

45. Festa, R.A., and Thiele, D.J. (2011). Copper: an essential metal in biology. Curr Biol *21*,
R877-883. 10.1016/j.cub.2011.09.040.

46. Inesi, G. (2017). Molecular features of copper binding proteins involved in copper
homeostasis. IUBMB Life *69*, 211-217. 10.1002/iub.1590.

47. Andreini, C., Banci, L., Bertini, I., and Rosato, A. (2008). Occurrence of copper proteins
through the three domains of life: a bioinformatic approach. The Journal of Proteome
Research *7*, 209-216.

48. Selden, C.R., Schilling, K., Godfrey, L., and Yee, N. (2024). Metal-binding amino acid
ligands commonly found in metalloproteins differentially fractionate copper isotopes. Sci
Rep *14*, 1902. 10.1038/s41598-024-52091-7.

49. Li, S., Cai, H., He, J., Chen, H., Lam, S., Cai, T., Zhu, Z., Bark, S.J., and Cai, C. (2016).
Extent of the Oxidative Side Reactions to Peptides and Proteins During the CuAAC
Reaction. Bioconjug Chem *27*, 2315-2322. 10.1021/acs.bioconjchem.6b00267.

50.   Abel, G.R., Jr., Calabrese, Z.A., Ayco, J., Hein, J.E., and Ye, T. (2016). Measuring and
976   Suppressing the Oxidative Damage to DNA During Cu(I)-Catalyzed Azide-Alkyne
977   Cycloaddition. Bioconjug Chem *27*, 698-704. 10.1021/acs.bioconjchem.5b00665.

978   51.   Wiest, A., and Kielkowski, P. (2024). Cu-Catalyzed Azide-Alkyne-Thiol Reaction Forms
979   Ubiquitous Background in Chemical Proteomic Studies. J Am Chem Soc.
980   10.1021/jacs.3c11780.

981   52.   Attarpour, A., Osmann, J., Rinaldi, A., Qi, T., Lal, N., Patel, S., Rozak, M., Yu, F., Cho,
982   N., Squair, J., et al. (2025). A deep learning pipeline for three-dimensional brain-wide
983   mapping of local neuronal ensembles in teravoxel light-sheet microscopy. Nat Methods
984   *22*, 600-611. 10.1038/s41592-024-02583-1.

985   53.   Attarpour, A., Osmann, J., Rinaldi, A., Qi, T., Lal, N., Patel, S., Rozak, M., Yu, F., Cho,
986   N., Squair, J., et al. A deep learning pipeline for 3D brain-wide mapping of local neuronal
987   ensembles in tera-voxel light sheet microscopy. Nat Methods *in press*.

988   54.   Wang, Q., Ding, S.-L., Li, Y., Royall, J., Feng, D., Lesnar, P., Graddis, N., Naeemi, M.,
989   Facer, B., and Ho, A. (2020). The Allen mouse brain common coordinate framework: a
990   3D reference atlas. Cell *181*, 936-953. e920.

991   55.   Boike, L., Henning, N.J., and Nomura, D.K. (2022). Advances in covalent drug
992   discovery. Nat Rev Drug Discov. 10.1038/s41573-022-00542-z.

993   56.   Ghosh, A.K., Samanta, I., Mondal, A., and Liu, W.R. (2019). Covalent Inhibition in Drug
994   Discovery. ChemMedChem *14*, 889-906. 10.1002/cmdc.201900107.

995   57.   Miller, V.A., Hirsh, V., Cadranel, J., Chen, Y.-M., Park, K., Kim, S.-W., Zhou, C., Su, W.-
996   C., Wang, M., Sun, Y., et al. (2012). Afatinib versus placebo for patients with advanced,
997   metastatic non-small-cell lung cancer after failure of erlotinib, gefitinib, or both, and one
998   or two lines of chemotherapy (LUX-Lung 1): a phase 2b/3 randomised trial. The Lancet
999   Oncology *13*, 528-538. 10.1016/s1470-2045(12)70087-6.

58.   Sequist, L.V., Yang, J.C., Yamamoto, N., O'Byrne, K., Hirsh, V., Mok, T., Geater, S.L.,
Orlov, S., Tsai, C.M., Boyer, M., et al. (2013). Phase III study of afatinib or cisplatin plus
pemetrexed in patients with metastatic lung adenocarcinoma with EGFR mutations. J
Clin Oncol *31*, 3327-3334. 10.1200/jco.2012.44.2806.

59.   Byrd, J.C., Furman, R.R., Coutre, S.E., Flinn, I.W., Burger, J.A., Blum, K.A., Grant, B.,
Sharman, J.P., Coleman, M., Wierda, W.G., et al. (2013). Targeting BTK with ibrutinib in
relapsed chronic lymphocytic leukemia. N Engl J Med *369*, 32-42.
10.1056/NEJMoa1215637.

60.   Treon, S.P., Tripsas, C.K., Meid, K., Warren, D., Varma, G., Green, R., Argyropoulos,
K.V., Yang, G., Cao, Y., Xu, L., et al. (2015). Ibrutinib in previously treated
Waldenstrom's macroglobulinemia. N Engl J Med *372*, 1430-1440.
10.1056/NEJMoa1501548.

61.   Wang, M.L., Rule, S., Martin, P., Goy, A., Auer, R., Kahl, B.S., Jurczak, W., Advani,
R.H., Romaguera, J.E., Williams, M.E., et al. (2013). Targeting BTK with ibrutinib in
relapsed or refractory mantle-cell lymphoma. N Engl J Med *369*, 507-516.
10.1056/NEJMoa1306220.

62.   Yun, S., Vincelette, N.D., Acharya, U., and Abraham, I. (2017). Risk of Atrial Fibrillation
and Bleeding Diathesis Associated With Ibrutinib Treatment: A Systematic Review and
Pooled Analysis of Four Randomized Controlled Trials. Clin Lymphoma Myeloma Leuk
17, 31-37 e13. 10.1016/j.clml.2016.09.010.

63.   Lipsky, A.H., Farooqui, M.Z., Tian, X., Martyr, S., Cullinane, A.M., Nghiem, K., Sun, C.,
Valdez, J., Niemann, C.U., Herman, S.E., et al. (2015). Incidence and risk factors of
bleeding-related adverse events in patients with chronic lymphocytic leukemia treated
with ibrutinib. Haematologica *100*, 1571-1578. 10.3324/haematol.2015.126672.

64.   Berglof, A., Hamasy, A., Meinke, S., Palma, M., Krstic, A., Mansson, R., Kimby, E.,
Osterborg, A., and Smith, C.I. (2015). Targets for Ibrutinib Beyond B Cell Malignancies.
Scand J Immunol *82*, 208-217. 10.1111/sji.12333.

65.   Tuomi, J.M., Xenocostas, A., and Jones, D.L. (2018). Increased Susceptibility for Atrial
and Ventricular Cardiac Arrhythmias in Mice Treated With a Single High Dose of
Ibrutinib. Can J Cardiol *34*, 337-341. 10.1016/j.cjca.2017.12.001.

66.   Xiao, L., Salem, J.E., Clauss, S., Hanley, A., Bapat, A., Hulsmans, M., Iwamoto, Y.,
Wojtkiewicz, G., Cetinbas, M., Schloss, M.J., et al. (2020). Ibrutinib-Mediated Atrial
Fibrillation Attributable to Inhibition of C-Terminal Src Kinase. Circulation *142*, 2443-
2455. 10.1161/CIRCULATIONAHA.120.049210.

67.   McMullen, J.R., Boey, E.J., Ooi, J.Y., Seymour, J.F., Keating, M.J., and Tam, C.S.
(2014). Ibrutinib increases the risk of atrial fibrillation, potentially through inhibition of
cardiac PI3K-Akt signaling. Blood, The Journal of the American Society of Hematology
124, 3829-3830.

68.   Honigberg, L.A., Smith, A.M., Sirisawad, M., Verner, E., Loury, D., Chang, B., Li, S.,
Pan, Z., Thamm, D.H., Miller, R.A., and Buggy, J.J. (2010). The Bruton tyrosine kinase
inhibitor PCI-32765 blocks B-cell activation and is efficacious in models of autoimmune
disease and B-cell malignancy. Proc Natl Acad Sci U S A *107*, 13075-13080.
10.1073/pnas.1004594107.

69.   Lanning, B.R., Whitby, L.R., Dix, M.M., Douhan, J., Gilbert, A.M., Hett, E.C., Johnson,
T.O., Joslyn, C., Kath, J.C., Niessen, S., et al. (2014). A road map to evaluate the
proteome-wide selectivity of covalent kinase inhibitors. Nat Chem Biol *10*, 760-767.
10.1038/nchembio.1582.

70.   Kaltenecker, D., Al-Maskari, R., Negwer, M., Hoeher, L., Kofler, F., Zhao, S., Todorov,
M., Rong, Z., Paetzold, J.C., Wiestler, B., et al. (2024). Virtual reality-empowered deep-
learning analysis of brain cells. Nat Methods *21*, 1306-1315. 10.1038/s41592-024-02245-
2.

71.   CENTER FOR DRUG EVALUATION AND RESEARCH APPLICATION NUMBER:
201292Orig1s000. (2013).
https://www.accessdata.fda.gov/drugsatfda_docs/nda/2013/201292Orig1s000PharmR.pdf
.

72.   CENTER FOR DRUG EVALUATION AND RESEARCH APPLICATION NUMBER:
205552Orig2s000. (2014).

https://www.accessdata.fda.gov/drugsatfda_docs/nda/2013/205552Orig1s000PharmR.pdf
.

73.   Karlsson, M., Zhang, C., Méar, L., Zhong, W., Digre, A., Katona, B., Sjöstedt, E., Butler,
L., Odeberg, J., Dusart, P., et al. (2021). A single–cell type transcriptomics map of human
tissues. Science Advances *7*, eabh2169. doi:10.1126/sciadv.abh2169.

74.   Xie, J., Zhang, J., Wu, H., Tang, X., Liu, J., Cheng, G., and Li, P. (2017). The influences
of age on T lymphocyte subsets in C57BL/6 mice. Saudi J Biol Sci *24*, 108-113.
10.1016/j.sjbs.2016.09.002.

75.   Fleming, M.R., Xiao, L., Jackson, K.D., Beckman, J.A., Barac, A., and Moslehi, J.J.
(2021). Vascular Impact of Cancer Therapies: The Case of BTK (Bruton Tyrosine Kinase)
Inhibitors. Circ Res *128*, 1973-1987. 10.1161/CIRCRESAHA.121.318259.

76.   Ryan, C.E., Sahaf, B., Logan, A.C., O'Brien, S., Byrd, J.C., Hillmen, P., Brown, J.R.,
Dyer, M.J., Mato, A.R., Keating, M.J., et al. (2016). Ibrutinib efficacy and tolerability in
patients with relapsed chronic lymphocytic leukemia following allogeneic HCT. Blood
128, 2899-2908. 10.1182/blood-2016-06-715284.

77.   Tom Dieck, S., Muller, A., Nehring, A., Hinz, F.I., Bartnik, I., Schuman, E.M., and
Dieterich, D.C. (2012). Metabolic labeling with noncanonical amino acids and
visualization by chemoselective fluorescent tagging. Curr Protoc Cell Biol *Chapter 7*, 7
11 11-17 11 29. 10.1002/0471143030.cb0711s56.

78.   Jao, C.Y., and Salic, A. (2008). Exploring RNA transcription and turnover in vivo by
using click chemistry. Proc Natl Acad Sci U S A *105*, 15779-15784.

79.   Nikic, I., Kang, J.H., Girona, G.E., Aramburu, I.V., and Lemke, E.A. (2015). Labeling
proteins on live mammalian cells using click chemistry. Nat Protoc *10*, 780-791.
10.1038/nprot.2015.045.

80.   El-Sagheer, A.H., and Brown, T. (2010). Click chemistry with DNA. Chemical Society
Reviews *39*, 1388-1405.

81.   Soriano del Amo, D., Wang, W., Jiang, H., Besanceney, C., Yan, A.C., Levy, M., Liu, Y.,
Marlow, F.L., and Wu, P. (2010). Biocompatible copper (I) catalysts for in vivo imaging
of glycans. J Am Chem Soc *132*, 16893-16899.

82.   Hulce, J.J., Cognetta, A.B., Niphakis, M.J., Tully, S.E., and Cravatt, B.F. (2013).
Proteome-wide mapping of cholesterol-interacting proteins in mammalian cells. Nat
Methods *10*, 259-264. 10.1038/nmeth.2368.

83.   Tamura, I., Sakamoto, D.M., Yi, B., Saito, Y., Yamada, N., Morimoto, J., Takakusagi, Y.,
Kuroda, M., Kubota, S.I., and Yatabe, H. (2024). Click3D: Click reaction across deep
tissues for whole-organ 3D fluorescence imaging. Science Advances *10*, eado8471.

84.   Shi, M.Y., Yao, Y., Wang, M., Yang, Q., Ding, L., Li, R., Li, Y., Huang, H., Yang, C.Y.,
Zhou, Z., et al. (2025). High-speed mapping of whole-mouse peripheral nerves at
subcellular resolution. Cell *188*, 3897-3915 e3820. 10.1016/j.cell.2025.06.011.

85.   Xu, F., Shen, Y., Ding, L., Yang, C.Y., Tan, H., Wang, H., Zhu, Q., Xu, R., Wu, F., Xiao,
Y., et al. (2021). High-throughput mapping of a whole rhesus monkey brain at
micrometer resolution. Nat Biotechnol *39*, 1521-1528. 10.1038/s41587-021-00986-5.

86. Verstegen, A.M.J., Vanderhorst, V., Gray, P.A., Zeidel, M.L., and Geerling, J.C. (2017).
Barrington's nucleus: Neuroanatomic landscape of the mouse "pontine micturition
center". J Comp Neurol *525*, 2287-2309. 10.1002/cne.24215.

87. Chung, K., Wallace, J., Kim, S.Y., Kalyanasundaram, S., Andalman, A.S., Davidson, T.J.,
Mirzabekov, J.J., Zalocusky, K.A., Mattis, J., Denisin, A.K., et al. (2013). Structural and
molecular interrogation of intact biological systems. Nature *497*, 332-337.
10.1038/nature12107.

88. Qi, Y., Yu, T., Xu, J., Wan, P., Ma, Y., Zhu, J., Li, Y., Gong, H., Luo, Q., and Zhu, D.
(2019). FDISCO: Advanced solvent-based clearing method for imaging whole organs.
Science Advances *5*, eaau8355. 10.1126/sciadv.aau8355.

89. Park, Y.G., Sohn, C.H., Chen, R., McCue, M., Yun, D.H., Drummond, G.T., Ku, T.,
Evans, N.B., Oak, H.C., Trieu, W., et al. (2018). Protection of tissue physicochemical
properties using polyfunctional crosslinkers. Nat Biotechnol *37*, 73-83. 10.1038/nbt.4281.

90. Goubran, M., Leuze, C., Hsueh, B., Aswendt, M., Ye, L., Tian, Q., Cheng, M.Y., Crow,
A., Steinberg, G.K., and McNab, J.A. (2019). Multimodal image registration and
connectivity analysis for integration of connectomic data from microscopy to MRI.
Nature communications *10*, 5504.

91. Arzt, M., Deschamps, J., Schmied, C., Pietzsch, T., Schmidt, D., Tomancak, P., Haase, R.,
and Jug, F. (2022). LABKIT: labeling and segmentation toolkit for big image data.
Frontiers in computer science *4*, 777728.

92. Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T.,
Preibisch, S., Rueden, C., Saalfeld, S., Schmid, B., et al. (2012). Fiji: an open-source
platform for biological-image analysis. Nat Methods *9*, 676-682. 10.1038/nmeth.2019.

93. Kinga, D., and Adam, J.B. (2015). A method for stochastic optimization. (San Diego,
California;), pp. 6.

94. Wagner, T., and Lipinski, H.-G. (2013). IJBlob: an ImageJ library for connected
component analysis and shape analysis. Journal of Open Research Software *1*.

95. Waskom, M.L. (2021). Seaborn: statistical data visualization. Journal of Open Source
Software *6*, 3021.

96. Tang, P.-f., Bao, S.-s., Xiao, Z.-x., Xie, W.-f., Wu, X.-m., Ge, H.-l., and Shao, C.-f.
(2023). A novel UHPLC–MS/MS method for quantitative analysis of zanubrutinib in rat
plasma: application to an in vivo interaction study between zanubrutinib and triazole
antifungal. BMC chemistry *17*, 107.

97. Lu, X., Liu, S., Yang, X., Han, M., and Sun, K. (2019). Determination of tyrosine kinase
inhibitor afatinib in rat plasma using LC–MS/MS and its application to in vivo
pharmacokinetic studies of afatinib liposomes. Journal of Pharmaceutical and Biomedical
Analysis *164*, 181-186.

1135

Figure 1

Figure 2

A

Passive clearing — Passive labeling

Drug-Injected Brains → HYBRiD Clearing → Whole Hemisphere *vCATCH* (PRCS / 2X RIR) → Lightsheet Imaging

Deep learning based segmentation ⟷ Allen Brain Atlas registration → *ACE* (AI-based Cartography of Ensembles)

Regional and cellular statistics
Intensity / Regions
Density / Regions

B — *Regional drug intensity* in atlas space

Vehicle
Pargyline-yne
Pargyline-yne
Pargyline-yne
280 / 160 / 40
LC

C — Segmented *drug-bound neurons* in atlas space

Vehicle
Pargyline-yne
i Pons
ii Hypothalamus
iii Medulla

D — Normalized drug intensity (A.U.)

indiv. mouse

ACB BST   EPI MED   PMD PMV   AT DR   B* LC   SLC SLD   AMB

Cerebral cortex | Cerebral nuclei | Thalamus | Hypothalamus | Midbrain | Pons | Cerebellum | Medulla | VS | fiber tracts

2.0 / 1.5 / 1.0 / 0.5

E — Normalized drug-bound cell density (A.U.)

indiv. mouse

ACB BST   EPI MED   PMD PMV   AT DR   B* LC   SLC SLD   AMB

Cerebral cortex | Cerebral nuclei | Thalamus | Hypothalamus | Midbrain | Pons | Cerebellum | Medulla | VS | fiber tracts

4 / 3 / 2 / 1 / 0

Figure 3

A

Passive clearing — *Passive labeling*

6-10 weeks / 4-7 days / 3-5 days / 1h 1h ... 1h 1h

In vivo Drug Injection — HYBRiD Clearing — PRCS — Whole-body **vCATCH** — 5X RIR — Lightsheet Imaging — Whole-body Drug Mapping

Organ-level — Cellular-level — Molecular-level

B x-y

Z: Imaging depth — Dorsal to Ventral

C Z=4.4 mm Z=5.9 mm Z=7.3 mm Z=8.8 mm Z=9.9 mm

x-y in-plane views

Ibrutinib-yne (juvenile mouse)

D Juvenile (1 week old)   18 mm   38 mm

E Transverse (ii) Transverse (i) Sagittal Coronal — Ibrutinib-yne

F Lung Heart Liver Gut — Coronal

G Lung Gut Liver Heart — Sagittal

H Lung Heart — Transverse (i)   Transverse (ii)

Ibrutinib-yne (adult mouse)

I Adult (8 week old)   30 mm   37 mm

J Sagittal Transverse Coronal — Ibrutinib-yne

K Lung Heart Liver Gut — Coronal

L Lung Liver Gut Heart — Sagittal

M Liver Stomach Esophagus Small intestine — Transverse

Figure 4

Figure 5

Figure S1

A

pargyline   pargyline-yne   PF7845   PF7845-yne

B

Vehicle   PF7845-yne

Full buffer | Full buffer | No N₃ | No ligand | No CuSO₄

z=10 micron

z=50 micron

CATCH in 100 micron brain sections

C

Normalized fluorescence intensity

0.2861   0.2112   0.9571   0.0008   0.0023

Z=10 micron   Z=50 micron

Full   Full   no N3   no BTTP   no Cu

Vehicle   PF7845-yne

D

Vehicle   Transverse

Pargyline-yne   Transverse

E

Vehicle

Coronal Bregma 1.5 | Coronal Bregma -0.2 | Coronal Bregma -1.5 | Coronal Bregma -3.5 | Coronal Bregma -4.7 | Coronal Bregma -5.4

Pargyline-yne

Coronal Bregma 1.5 | Coronal Bregma -0.2 | Coronal Bregma -1.5 | Coronal Bregma -3.5 | Coronal Bregma -4.7 | Coronal Bregma -5.4

ACB | Choroid plexus | MED | PMV | DR / AT | LC

Figure S2

A

Registration   Voxelization and Warping

Pargyline-yne   Vehicle   Pargyline-yne

B

Average *regional intensity* difference in atlas space

C

Average *regional neuronal density* differences in atlas space

D

Average *regional neuronal density -log(p-value)* in atlas space

E

| | *MAO-A* enriched | | | | *MAO-B* enriched | | | | *Dual MAO-A/B* enriched | |
| | SLC | SLD | LC/B | AMB | ACB | BST | DR | AT | PMD | PMV |

Pargyline-yne

MAO-A

Nissl

Pargyline-yne

MAO-B

Nissl

F

Pargyline-yne binding characterizations in LC/B

| Pargyline-yne | TH | NeuN | Nissl |

Figure S3

A

Afatinib   Afatinib-yne   Ibrutinib   Ibrutinib-yne

B

Plasma concentration (uM) vs Time post injection
- Afatinib
- Afatinib-yne

Plasma concentration (uM) vs Time post injection
- Ibrutinib
- Ibrutinib-yne

C

Tissue concentration (ng/g wet tissue): Heart, Lung, Liver, Kidney, Spleen
- Afatinib
- Afatinib-yne

Tissue concentration (ng/g wet tissue): Heart, Lung, Liver, Kidney, Spleen
- Ibrutinib
- ibrutinib-yne

D Afatinib-yne **lung**

Azide only   No Cu   No Azide   Full click

Vehicle

Afatinib-yne

E Ibrutinib-yne **spleen**

Azide only   No Cu   No Azide   Full click

Vehicle

Ibrutinib-yne

F

Normalized fluorescence intensity
- Vehicle
- Afatinib-yne

Azide only 0.9998   No Cu >0.9999   No Azide >0.9999   Full click <0.0001

G

Normalized fluorescence intensity
- Vehicle
- Ibrutinib-yne

Azide only 0.7174   No Cu 0.8537   No Azide 0.9954   Full click <0.0001

H Juvenile mouse 3D views

Vehicle   Afatinib-yne   Ibrutinib-yne

I Adult mouse 3D views

Vehicle   Ibrutinib-yne

J Afatinib-yne (juvenile mouse)

Afatinib-yne
Sagittal
Transverse (ii)
Transverse (i)
Coronal

K Afatinib-yne

Heart
Lung
Liver
Stomach
Gut
Coronal

L

Gut
Lung
Liver
Heart
Sagittal

M

Transverse (i)   Transverse (ii)

Figure S4

A

Afatinib-yne (juvenile mouse)

B

Ibrutinib-yne (juvenile mouse)

C

Ibrutinib-yne (adult mouse)

Figure S5