Mapping cellular targets of covalent cancer drugs in the entire mammalian body

Authors: Zhengyuan Pang[1, 2, 6], Verina H. Leung[1, 6], Cailynn C. Wang[1], Ahmadreza Attarpour[3,4], Anthony Rinaldi[3], Hanbing Shen[1], Christopher Glynn[1], Logan H. Sigua[1], Melaina Yender[1], Alexandra Selke[1], Senhan Xu[1], Peng Wu[5], Maged Goubran[3,4], Benjamin F. Cravatt[2,] Li Ye [1, 7*]

[1]Howard Hughes Medical Institute, Department of Neuroscience, Dorris Neuroscience Center, Scripps Research, La Jolla, CA, United States.

[2]Department of Chemistry, The Scripps Research Institute; La Jolla, California, 92037, United States.

[3]Sunnybrook Research Institute, University of Toronto, Toronto, ON, Canada.

[4]Department of Medical Biophysics, University of Toronto, Toronto, ON, Canada.

[5]Department of Molecular and Cellular Biology, The Scripps Research Institute; La Jolla, California, 92037, United States.

[6]These authors contributed equally.

[7]Lead contact

*Correspondence: liye@scripps.edu

SUMMARY:

As our understanding of biological systems reaches single-cell and high spatial resolutions, it becomes imperative that pharmacological approaches match this precision to understand drug actions. This need is particularly urgent for the targeted covalent inhibitors that are currently re-entering the stage for cancer treatments. Therefore, we developed 3D volumetric clearing-assisted tissue click chemistry (vCATCH) to image *in vivo* target-bound drugs by leveraging the unique kinetics of click reactions. In mice, vCATCH overcomes limitations of existing methods to achieve high-throughput volumetric chemical labeling through simple and passive incubation steps, thus enabling unbiased cellular resolution drug imaging in entire animals. We combined vCATCH with HYBRiD imaging and virtual reality to visualize and quantify *in vivo* targets of two clinical cancer drugs, afatinib and ibrutinib, which recapitulated their known pharmacological distribution and revealed previously unreported tissue and cell type engagement potentially linked to off-target effects. vCATCH provides a body-wide, unbiased platform to map covalent drug engagements at unprecedented scale and precision.

KEYWORDS:

drug targets, tissue clearing, whole-body imaging, targeted covalent inhibitors

INTRODUCTION

The primary binding sites for drug molecules are critical determinants of drug efficacy and toxicity. Although molecular target identification strategies, such as chemoproteomics[1-5], chemogenomic[6-9], and thermal shift profiling[10-13], have revolutionized our understanding of drug activities, mapping *in vivo* drug targets with spatial and cellular resolution remains challenging. This shortfall is even more pressing in light of the innovations in genomics and imaging techniques over the past decade that have enabled the characterization of tissue architecture at the single-cell level with spatial resolution[14-20]. This has led to a widening gap between our understanding of native biological systems and our ability to track *in vivo* drug actions within these systems, as the latter still primarily rely on low-resolution methods such as radioactive imaging. It is imperative that we develop methods to visualize *in situ* drug engagement with cellular resolution across tissue compartments and organ systems to better understand *in vivo* drug actions.

One promising avenue for imaging drug targets is fluorescence imaging, which has been widely used for visualizing endogenous biomolecules like DNA, mRNA, and proteins. However, directly imaging drug molecules conjugated with a fluorescent tag is often not suitable as a larger tag could distort drug activity and distribution *in vivo*. As an alternative approach, we recently developed a method to visualize drug molecules in tissue sections by combining tissue clearing with *in situ* click chemistry (CATCH)[21]. However, efforts to extend CATCH from thin sections to large, heterogeneous 3D tissue environments face significant challenges similar to those typically encountered by whole-mount immunostaining or *in situ* hybridization methods, such as the accumulation of probes at the tissue surface, uneven staining across the tissue depth, and insufficient labeling at the center of large samples[22-26].

Here, we develop new CATCH capacities to enable rapid and homogenous *in situ* click labeling in ultra-large 3D tissues at the scale of a whole mouse body. This new technology leverages the unique characteristics of click chemistry by introducing pre-reaction copper saturation (PRCS) and repeated iterations of reaction (RIR) strategies. The resulting volumetric CATCH (vCATCH) method, when combined with HYBRiD (hydrogel-based reinforcement of three-dimensional imaging solvent-cleared organs (DISCO)) clearing[27], is further highlighted by its marked simplicity as it only relies on easy and passive incubation steps to achieve whole-body labeling and imaging.

RESULTS

Establishing the principles for deep tissue click reactions

To achieve 3D click labeling, we turned to recent advances in whole-brain and whole-body immunostaining techniques[24,28-35]. As summarized by previous works[36-39], two main strategies have been used to achieve deep tissue antibody penetration: (1) enhancing diffusion of the probe (e.g., antibody) into the tissue; and (2) minimizing probe-target interactions (e.g., antibody-antigen binding) during the diffusion stage while maximizing reaction efficiency at the staining stage, such that the large difference in reaction rates between the two stages will prevent probe depletion at the tissue surface and facilitate homogeneous tissue labeling[23,33,34]. Biophysics studies have demonstrated that changing the pH, salt concentration, and other physicochemical parameters can modulate antibody-antigen binding kinetics by 4-400 fold[40,41]. We hypothesized

that we could adopt a similar strategy to enable 3D click reactions. CATCH relies on copper-
catalyzed alkyne azide cycloaddition (CuAAC) to label target-bound drug molecules with
fluorescent signals. As its name indicates, the kinetics of CuAAC strictly depend on the catalytic
Cu(I)-ligand complex, but for ease of handling, a Cu(II) compound, typically $CuSO_4$, is used
during the diffusion stage and then a reducing agent (e.g., sodium ascorbate) is added *in situ* to
generate Cu(I) at the binding stage. Compared to uncatalyzed conditions, the generation of
catalytic Cu(I) can lower the energy barrier by 14.9-18.4 kcal/mol, or an equivalent of 7-8 orders
of magnitude increase favoring the cycloaddition that attaches the fluorescent probe to the drug
in situ[42]. Thus, by temporarily preventing the generation of Cu(I) from Cu(II) through
withholding the reducing agent sodium ascorbate, in principle, we could modulate the probe-
target reaction by 10^{7-8} fold between the diffusion and binding stages, which could improve the
penetration depth of CATCH in large tissues.

For method development, we used the covalent monoamine oxidase (MAO) inhibitor pargyline,
whose alkyne probe pargyline-yne has previously been characterized by activity-based protein
profiling and 2D-CATCH studies (Figure S1A)[21,43]. Although CATCH has demonstrated
homogeneous click labeling of N_3-dye in 500 μm tissues[21], when tested in mouse hemispheres,
even with the aforementioned catalyst withholding strategy, only the surface of the mouse brain
was labeled (up to 0.4 mm, Figure 1A). First, to exclude the possibility that other CATCH
components might have trouble diffusing through the tissue, we removed individual reagents
from the diffusion step and evaluated the resulting labeling depth. For instance, eliminating the
N_3-dye, the largest molecular weight component, in the diffusion step did not affect labeling
depth, suggesting N_3-dye diffusion is not a limiting factor (Figure S1B and S1C). By contrast,
removing $CuSO_4$ or ligand strongly reduced labeling depth in these conditions, suggesting that
the catalyst is still likely the diffusion bottleneck (Figure S1B and S1C).

Cu is well-known to bind multiple amino acid side chains (i.e., cysteine and histidine)[44-47]. We
therefore hypothesized that the ubiquitous copper binding sites could consume free Cu(II) as it
diffuses into the sample, resulting in insufficient catalyst conversion in deeper tissues (Figure
1B). Unlike typical staining in which increasing dye or antibody concentrations tends to result in
nonspecific labeling and high background, high Cu(II) alone, as a catalytic precursor, should not
lead to nonspecific labeling. Based on this theoretical prediction, we reasoned that excessive
Cu(II) could be used to pre-saturate potential copper binding sites (therefore "blocking" them) to
boost catalytic activity in deeper tissues. Importantly, this strategy should not increase
nonspecific labeling or lead to high background. Indeed, we observed that incorporating a 10x
Cu excess pre-reaction copper saturation (PRCS) step increased the reaction depth to 1 mm while
maintaining the signal-to-noise ratio after a single one-hour reaction (Figure 1A-1C).

We next sought to modify CATCH to reach targets in tissues even deeper than 1 mm. Although
increasing reaction time could potentially increase labeling depth, prolonged reactions would
deplete the reducing agent in the buffer, which could eventually lead to reactive oxygen species
(ROS) build-up and side reactions[48-50]. Instead, we refreshed the click reaction cocktail every
hour, a procedure we termed repeated iterations of reactions (RIR). In practice, PRCS plus two
rounds of RIR resulted in full-depth, homogenous CATCH labeling across the mouse hemisphere
(Figure 1A-1C, Supplementary Video 1). We term this approach volumetric CATCH (vCATCH).

vCATCH quantitatively profiles brain-wide targets of Pargyline

After achieving brain-wide labeling, we employed vCATCH to unbiasedly map drug targets of
pargyline. After intraperitoneal (i.p.) injection of pargyline-yne (10 mg/kg) in wild-type mice,
mouse hemispheres were harvested and cleared by HYBRiD. Hemispheres then underwent one
day of PRCS and two rounds of RIR before imaging with lightsheet fluorescence microscopy
(Figure 1D). Importantly, vCATCH allowed the use of high-throughput, paralleled passive
processing to generate cohort-scale pargyline-labeled brains with marked reproducibility, critical
for performing biologically meaningful analysis of its cellular targets across the whole brain
(Figure 1E-1H, S1D, and S1E).

First, we manually verified pargyline probe binding in the brain based on previously reported
regional targets[21]. These regions included distinct cellular structures in the pons and
hypothalamus (Figure 1H). In addition, we found pargyline-yne-enriched cells in the choroid
plexus (in the ventricle) and the medulla (Figure 1H). We also observed additional neuron-like
structures in the midbrain (Figure S1E). These findings demonstrated that vCATCH maintained
high fidelity in large tissue volumes and across the whole brain.

The throughput and reproducibility of vCATCH across cohort-scale samples also allowed
automatic and unbiased quantification of drug targets beyond visual inspection. We applied a
recently developed AI-based Cartography of Ensembles (ACE) pipeline to analyze these datasets
(Figure 2A-2C, Supplementary Table 1)[51]. After registering the 3D volumes onto the Allen Brain
Atlas (ABA)[52], we first quantified the fluorescence intensity (indicating drug binding abundance)
across all brain regions (Figure 2B). Intensity quantifications revealed high drug abundance in
the pons, thalamus, and midbrain (Figure 2D and S2A). Additionally, we found the highest drug
abundance in the locus coeruleus (LC), followed by the subceruleus nucleus (SLC) and
Barrington's nucleus (B). Interestingly, we observed high drug abundance in the nucleus
accumbens (ACB), an important limbic region known to receive extensive monoaminergic
inputs[53], potentially reflecting high local MAO expression to degrade monoamine
neurotransmitters (Figure 2D and S2A). Moreover, in the medulla, we found drug enrichment in
the nucleus ambiguous (AMB), which sends motor outputs to muscles (Figure 2D)[54].

Leveraging the cellular resolution imaging provided by vCATCH, we next sought to unbiasedly
quantify the cell targets of pargyline. Empowered by a large training dataset and AI-based
segmentation algorithms, the ACE pipeline was used to identify brain-wide pargyline-yne-
positive cells (Figure 2C). The segmentation models were further fine-tuned to optimize their
performance and generalizability on our dataset. After segmentation, the cell density patterns
mostly agreed with fluorescence intensity profiles. For example, in the pons, higher cell density
was observed in the LC, B, SLC, and sublaterodorsal nucleus (SLD) (Figure 2E, S2B and S2C).
Similarly, in the hypothalamus, the dorsal and ventral premammillary nucleus (PMD and PMV)
showed elevated cell density (Figure 2E). In addition, the epithalamus (EPI) was found to have a
high cell density (Figure 2E).

However, unbiased ACE analysis also revealed unexpected differences between fluorescence
intensity and cell counts. For example, the ACB and bed nuclei of the stria terminals (BST), as
well as the medial group of the dorsal thalamus (MED), showed extensive drug binding by
intensity, but almost no neuronal structures were detected in this area, indicating drug binding
may be associated with non-soma structures (Figure S1E, 2D, and 2E). Conversely, drug-bound
cells were identified in the dorsal nucleus raphe (DR) and the anterior tegmental nucleus (AT) of
the pons (Figure S1E, 2D, and 2E) despite the lack of overall fluorescence intensity increase,
suggesting sparse neuronal populations were targeted by pargyline in these regions.

Taken together, by using a pargyline probe as a proof-of-concept, we demonstrated that vCATCH
is compatible with clearing methods and high-throughput computational pipelines to unbiasedly
quantify regional enrichment and cellular targets of covalent drugs in the mouse brain, which
was not possible with traditional methods.

vCATCH profiles whole-body targets of covalent cancer drugs

We next sought to expand vCATCH target mapping to whole animals across all organ systems.
We selected two widely used cancer drugs for this study: afatinib (Gilotrif) and ibrutinib
(Imbruvica), both covalent targeted kinase inhibitors (TKIs)[55,56]. Since their landmark approval
in 2013, the epidermal growth factor receptor (EGFR) inhibitor afatinib and Bruton's tyrosine
kinase (BTK) inhibitor ibrutinib, respectively, have transformed the clinical landscape of non-
small cell lung cancer (NSCLC)[57,58] and B cell malignancies[59-61]. Despite their popularity,
however, there are several well-documented side effects of TKIs that are poorly understood. For
example, ibrutinib is associated with cardiovascular and bleeding risks[62,63]. Although many
studies have been conducted to understand the protein targets (or off-targets) of ibrutinib[64-67], the
in vivo tissue(s) or cell types responsible for such toxicities remain elusive. The ability to
unbiasedly profile TKI targets *in vivo* at cellular resolution could provide crucial insight to
bridge this gap.

We injected previously established clickable probes of ibrutinib-yne (10 mg/kg, i.p.) and
afatinib-yne (10 mg/kg, i.p.) in wild-type mice (Figure S3A)[68]. We established a vCATCH
protocol with 4-day PRCS and 5 rounds of RIR to homogenously label a whole HYBRiD-cleared
juvenile mouse with a total volume of more than 18x38x12mm (Figure 3A). We then performed
whole-body lightsheet imaging of the entire mouse bodies (Figure 3B-3O, S3B-S3G) and
visualized them through virtual reality (VR) (Supplementary Video 2 and 3)[69]. Globally, the
binding of both TKIs was highly abundant in the liver and the gastrointestinal tract (Figure 3D
and 3J). In the upper chest, ibrutinib showed elevated heart binding compared to the lung,
whereas the opposite enrichment was observed with afatinib (Figure 3E and 3K). These findings
are consistent with general distributions of these two drugs based on previous autoradiography
data from the Food and Drug Administration[70,71].

The high resolution of vCATCH further allowed us to examine the detailed drug distribution
within the organs using lightsheet fluorescence microscopy (Figure 3F-3I, 3L-3O, S3H, and S3I).
In several organs, afatinib and ibrutinib displayed similar intra-organ distributions; for example,
both drugs were selectively enriched in the villus in the small intestine (Figure S3H and S3I).
Higher magnitude confocal imaging further revealed that both drugs strongly bound to circular
structures in the bone (Figure 4A and 4D) and spleen (Figure S3H and S3I). Subsequent lectin
staining identified these structures as vasculatures (Figure S4A and S4B). Considering that BTK
expression is generally absent in blood vessels[72], we suspect the vasculature enrichment may be
implicated in the off-target toxicity associated with ibrutinib[62,63].

In other organs, we observed distinct target distributions between these two drugs. For example,
robust tissue enrichments in the inner stomach lining and along the nasal tract were associated
with ibrutinib but not afatinib (Figure S3H and S3I). Furthermore, we discovered widely spread
afatinib targets in the lung whereas pulmonary ibrutinib labeling was highly restricted along the
bronchi (Figure 3F, 3G, 3L, and 3M).

Interestingly, although both drugs were bound to the kidney, they displayed different intra-organ
patterns (Figure S3H and S3I). Afatinib was associated with glomeruli in the cortex and near the

cortex-medulla interface, whereas ibrutinib was more localized to cortical tubular-like structures
(Figure 4B and 4E).

Molecular quantification of afatinib and ibrutinib targets in the liver

Our global organ-level imaging (Figure 3D and 3J) showed that both TKIs were enriched in the
liver, in agreement with the literature[70,71]. However, vCATCH imaging revealed differences in
their intra-organ distribution: afatinib showed crown-like signals throughout the hepatic tissue
whereas ibrutinib was associated with sparsely scattered individual cells (Figure 3H, 3I, 3N, and
3O).

To test whether afatinib and ibrutinib shared the same cellular targets in the liver, we leveraged
the compatibility of CATCH with molecular phenotyping and performed immunostaining to
examine the cellular targets of both TKIs (Figure 4C, 4F-4L). Consistent with high hepatocyte
EGFR expression[72], we found that the majority (59.17±10.91%) of afatinib-bound cells were
hepatocytes, followed by 21.10±5.11% being F4/80-positive macrophages (Figure 4G-4I). By
contrast, only a small fraction (2.61±1.65%) were identified as CD3e-positive T cells (Figure
4H). In contrast to afatinib, only a small fraction (13.95±1.32%) of ibrutinib-bound cells were
hepatocytes, while a large proportion (39.53±4.42%) of these cells were F4/80 positive
macrophages (Figure 4F and 4J-4L). This is consistent with a modest BTK expression in liver
macrophages[72]. However, unexpectedly, the largest fraction (52.78±3.25%) of ibrutinib-bound
cells were CD3e-positive T cells despite their low BTK expression (Figure 4J and 4K)[72]. This
suggests potential ibrutinib off-target engagement in hepatic T cells.

Beyond the liver, we were also able to identify and quantify afatinib-enriched immune cell
populations in the lung (Figure S4C-S4E). In summary, our vCATCH data revealed that the two
TKIs are enriched in distinct cell types within individual tissues and organs, highlighting
vCATCH's capacity to profile drug targets across global, organ, tissue, and cellular scales.

DISCUSSION

Covalent TKIs have reentered the stage for cancer treatment and broadened the horizon for the
druggable space as demonstrated by the recent developments of mutant selective EGFR inhibitor
osimertinib, and KRAS G12C inhibitors sotorasib and adagarasib[55]. However, despite the
reignited enthusiasm, the concern for toxicity (for example, cardiotoxicity and bleeding for
ibrutinib) calls for continuous innovations in characterizing their *in vivo* targets to guide future
covalent drug development.

Here, by treating samples with excessive copper (PRCS) and performing multiple rounds of click
reactions (RIR), we established a new method, vCATCH, which enabled robust covalent TKI
mapping across the whole mouse body. This method only requires simple passive incubations
and therefore offers reliability, reproducibility, and high throughput for cohort-scale biological
and pharmacological analyses. Using existing pharmacokinetic data from the FDA as a reference,
vCATCH confirmed organ-level drug distribution. Critically, vCATCH enabled us to identify
drug engagement with intra-organ cellular resolution that is typically inaccessible through
conventional methods. The capacity to resolve drug engagement across molecularly defined cell
populations, tissue architectures, and organ systems, once combined with single-cell and spatial
omics tools, presents transformative opportunities for spatially defined, cell-type-specific drug
action characterization.

Our work so far has focused on wild-type mice for method development. By examining drug
binding in healthy mice, we uncovered patterns potentially associated with off-target engagement
and toxicities. For example, ibrutinib is well known to be associated with cardiac toxicity such as
atrial fibrillation (AF), but the mechanism behind this association remains incompletely
understood[64-67]. Although animal studies suggested AF induction would likely require chronic
administration[66], our study revealed clear ibrutinib enrichment in the heart after a single
injection, suggesting an intrinsic enrichment of ibrutinib for the cardiovascular system. It would
be of great interest to further examine ibrutinib engagement in AF models to better understand its
cardiotoxicity mechanism. In addition to AF, bleeding is another major ibrutinib risk factor[62,63].
Despite low BTK expression in endothelial cells[72], we observed ibrutinib engagement in the
vasculature, a finding that could guide subsequent molecular identification work to further
characterize unique ibrutinib targets in endothelial cells.

On the technology side, vCATCH represents a new strategy to use biorthogonal click chemistry
to label large tissues, in contrast to most mainstream labeling methods (e.g., immunostaining,
DNA/RNA hybridization, etc) that primarily rely on biological affinities, such as antibody-
antigen, biotin-streptavidin, and complementary DNA/RNA interactions. Click chemistry allows
us to easily control labeling kinetics by simple buffer exchanges, achieving uniform probe
penetration in ultra-large tissue without delicate procedures or apparatus. The covalent click
chemistry linkage also means that the same sample can withstand harsh and prolonged
processing and imaging operations, opening numerous possibilities to multiplex vCATCH with
other spatial omics methods.

Finally, it is a major challenge in whole-mount tissue staining to balance the high probe
concentration (such as antibody) required for tissue penetration and its associated background
resulting from the nonspecific interactions between the probe and endogenous biomolecules. In
this regard, the biorthogonal nature of click chemistry could better tolerate a wide range of
labeling conditions without increasing noise. Currently, click chemistry-based imaging studies
are primarily associated with examining transcription/translation[73,74]. By demonstrating
vCATCH's capacity to map covalent drug binding across the entire mouse body, we propose that
similar principles can be applied in ultra-large 3D tissues to visualize substrates including but not
limited to protein[75], nucleic acid[76], glycan[77], lipid[78], and drug molecules[21]. For example, a recent
study has demonstrated the original CATCH method or similar click labeling strategies can be
used to image endogenous RNA and hypoxia dyes in 3D organs (such as mouse brain and
tumor)[79], suggesting such approaches could be broadly adopted in future biomedical studies.

Limitations of the study

Given our focus here on method development, our current study used healthy wild-type mice, in
which the engagement of TKIs could differ from tumor-bearing animals. Given the robustness
and simplicity of the vCATCH pipeline, however, we believe other researchers with oncology
expertise could readily apply vCATCH in various *in vivo* cancer models. We also note that due to
the size limit of the current lightsheet microscopes, we chose to demonstrate whole-body
imaging capacity in young mice (1 week old). Our previous studies have demonstrated HYBRiD
can reliably clear and image parts of adult mice (i.e., mouse torso)[27,80]. With the deep tissue click
reaction mechanism described here, we foresee vCATCH being readily applied to image drug
binding in adult mouse bodies with more advanced microscope systems in the future. Finally, all
clickable TKI probes used in this study have been rigorously validated by published medicinal
chemistry and chemoproteomics studies[68]; it is important to note that the design of future

CATCH probes for other drugs must be verified by similar approaches; although it is becoming common to generate and characterize clickable alkyne analogs during the development of new covalent inhibitors.

ACKNOWLEDGMENTS

We thank all members of the Ye lab and the Dorris Neuroscience Center for their support and feedback. We thank Barry Sharpless for suggestions. This work was supported by NCI IMAT program (CA281918 to L.Y. and B.F.C), NIH Director's New Innovator Award (DK128800 to L.Y.), and NIDA (DA059393 to L.Y.). L.Y. was also supported by the HHMI, NIDDK, NIMH/BRAIN, Chan Zuckerberg Initiative, and the Dana, Whitehall, Baxter, and Abide-Vividion Foundations. Z.P. and H.S. were supported by the Dorris Scholar Award. L.H.S is supported by an NIH Medical Scientist Training Grant T32GM007198-49. P.W. is supported by the NIH (R35GM139643).

AUTHOR CONTRIBUTIONS

L.Y., Z.P., conceived the study. L.Y., Z.P., P.W., M.G., and B.F.C. designed the experiments. Z.P., V.H.L., C.C.W., C.G., L.H.S., M.Y., A.S., and S.X. performed the experiments and analyzed the data. A.A., A.R., H.S., and M.G. performed ACE analysis. L.Y. and B.F.C. acquired funding. L.Y. supervised the study. L.Y., Z.P., V.H.L., and C.C.W. wrote the manuscript with input from all authors. All authors reviewed and provided feedback on the manuscript.

DECLARATION OF INTERESTS

The design, steps, and applications of vCATCH are covered in pending patent application material from The Scripps Research Institute.

Figure legends:

Figure 1. Establishing the principles for deep tissue click reactions

(A) Brain-wide characterization of click labeling in pargyline-yne-treated hemispheres (10 mg/kg, 1 h, i.p.). After HYBRiD clearing and click labeling, hemispheres were post-sliced into coronal sections for penetration characterizations. Regular CATCH failed to label the whole hemisphere depth. Pre-reaction copper saturation (PRCS) and repeated iterations of reactions (RIR, 1 hour per iteration) enhanced labeling depth and generated homogenous click labeling across the whole hemisphere. Scale bar, 1000 μm.

(B) Schematics of copper competition in tissue. Copper binding sites could chelate copper and limit the available free copper for CuAAC catalyst conversion, thereby impairing click labeling efficiency deep in tissues.

(C) Quantification of click labeling depth along the marked lines in Figure 1A. Labeling intensity is normalized to the surface intensity of individual samples. Four hemispheres in each condition were used for quantification. Data are shown as mean±SEM.

(D) Representative 3D rendering of vehicle and pargyline-yne treated hemispheres. Scale bar, 2000 μm.

(E) 3D rendering of pargyline-yne treated hemispheres showing the total imaging volume.

(F) Schematics of imaging depth for 1G highlighting changes in different brain regions covered.

(G) Sagittal 2D views of pargyline-yne treated hemispheres at imaging depth Z=1.5 mm and 3.0 mm as indicated in Figure 1F. Scale bar, 2000 μm

(H) Zoomed-in views showing pargyline-yne enriched neurons and brain regions across the hemisphere. Scale bar, 100 μm.

Figure 2. ACE analysis of pargyline-yne targets across the hemisphere.

(A) Schematics of the ACE analysis pipeline. vCATCH labeled hemispheres were imaged under the lightsheet microscope. Whole hemisphere images were first registered onto the ABA whereby the average intensity of each region was quantified. Known pargyline-yne positive neurons in the hypothalamus and pons (as indicated in Figure 1H) were used to train the ACE model for subsequent neuronal segmentation across the hemisphere. Segmented images were down-sampled to 10x10x10 μm voxels. The density (number of cells/mm^3) of identified pargyline-yne positive neurons was then quantified across different brain regions.

(B) Representative regional drug intensity quantification of the vehicle and pargyline-yne-treated (10 mg/kg, 1 h, i.p.) hemispheres in atlas space. Hemispheres were registered based on the ABA as indicated by the white boundary, and the average fluorescence intensity in each region was quantified. The LC and Barrington's nucleus (B) showed the highest overall labeling intensity in pargyline-yne hemispheres. The scale bar indicates raw intensity values.

(C) Representative segmented pargyline-yne enriched neuron in atlas space. Known pargyline-yne-bound neurons in the pons and hypothalamus were used as positive controls. Additional neuronal structures (i.e. in the medulla) were successfully identified. The scale bar indicates raw segmented neuron density in each 10x10x10 μm voxel.

(D-E) Heatmap of normalized pargyline-yne labeling intensity (D) and normalized pargyline-yne positive neuron density (E). A total of 5 pargyline-yne-treated mice (rows) and 184 brain regions (columns) were included for analysis. For individual mice, intensity and density values were normalized by the average intensity and density of all the regions from each mouse. Purple labels indicate regions with only elevated intensity. Blue labels indicate

regions with only elevated neuronal density. Red labels indicate regions with shared elevated intensity and neuronal density. Gray labels indicate regions without significant intensity or density elevation. The full dataset and acronyms can be found in Supplementary Table 1. VS, ventricular system.

Figure 3. vCATCH reveals whole-body TKI distribution with high spatial resolution.
(A) Workflow of vCATCH whole-body labeling in one-week old mice. HYBRiD cleared whole mouse bodies were placed in $CuSO_4$ solution for PRCS. After equilibrating samples in click incubation buffer, mice underwent five RIRs for click labeling and were imaged by lightsheet microscopy.
(B) Schematic of imaging depth across whole-body vCATCH labeling.
(C) Coronal (x-y) views of vCATCH whole-body labeling of ibrutinib-yne (10 mg/kg, 1 h, i.p.) treated mouse at different z-depths. Scale bar, 4000 µm.
(D) Characterization of afatinib-yne whole-body drug distribution. Representative 3D lightsheet image volume of afatinib-yne (10 mg/kg, 1 h, i.p.) treated mouse. D, dorsal; V, ventral; S, superior; I, inferior; R, right; L, left. Scale bar, 4000 µm.
(E) Afatinib-yne binding in the upper chest, heart and lung, revealing greater drug enrichment in lung compared to heart tissue. RA, right atrium; RV, right ventricle; LA, left atrium; LV, left ventricle. Scale bar, 1000 µm.
(F-G) Overview of afatinib-yne engagement in lung (F) and zoom in views (G). Zoom-in view shows distinct afatinib-yne positive cellular structures in the lung. Scale bar, lung overview, 1000 µm; lung zoom-in, 300 µm.
(H-I) Afatinib-yne enrichment in liver (H) with zoom-in view (I) revealing cluster-like drug positive structures. Scale bar, liver overview, 500 µm; liver zoom-in, 100 µm.
(J) Characterization of ibrutinib-yne whole-body drug distribution. Representative 3D lightsheet image volume of ibrutinib-yne treated mouse. Scale bar, 4000 µm.
(K) Upper chest ibrutinib-yne binding characterizations reveal greater drug enrichment in heart tissue compared to lung. Scale bar, 1000 µm.
(L-M) Overview of ibrutinib-yne engagement in lung (L) and zoom in view (M). Ibrutinib-yne enrichment is primarily localized along the lining of bronchiole, with no discernable drug enriched structures observed in alveoli. Scale bar, lung overview, 1000 µm; lung zoom-in, 250 µm.
(N-O) Overview of ibrutinib-yne engagement in the liver (N). Liver zoom-in view (O) reveals distinct drug positive cellular structures. Scaler bar, liver overview, 500 µm; liver zoom-in, 100 µm.

Figure 4. Cell type characterization of TKI engagement in drug-enriched tissues.
(A) Afatinib-yne characterization in leg bone section (top). Bone zoom-in views (bottom) show drug enrichment in central arteriole-like structures. Scale bar, bone global views, 200 µm; bone zoomed-in views, 20 µm.
(B) Representative view of afatinib-yne binding in kidney section. Zoom-in views (i and ii) reveal afatinib-yne drug enrichment in glomeruli. Scale bar, kidney overview view, 1000 µm; kidney zoomed-in views, 200 µm.
(C) Overview of afatinib-yne engagement in liver section. Cluster-like afatinib-yne enriched structures are observed in liver zoom-in views (i and ii). Scale bar, liver global views, 500 µm; liver zoomed-in views, 50 µm.

(D) Ibrutinib-yne characterization in leg bone (top), with zoom-in view (bottom) showing
drug enrichment in central arteriole-like structures. Scale bar, bone global views, 200 µm;
bone zoomed-in views, 20 µm.
(E) Distinct ibrutinib-yne binding pattern observed in kidney section. Zoom-in views (i and
ii) reveal drug localization in cortical tubules. Scale bar, kidney overview view, 1000 µm;
kidney zoomed-in views, 200 µm.
(F) Overview of ibrutinib-yne engagement in liver section. Discrete ibrutinib-yne-enriched
structures scattered throughout the liver are observed in liver zoom-in views. Scale bar: liver
global views, 500 µm; liver zoomed-in views, 50 µm.
(G) High resolution afatinib-yne cell type characterization in the liver. Representative images
reveal afatinib-yne+ hepatocytes are F4/80- (top row) and CD3e- (bottom row). The dashed
line represents the cell boundary. Scale bar, 10 µm.
(H) Cell type characterization of afatinib-yne enriched cells in the liver. 59.17±10.91% of
cells were identified as hepatocytes, followed by 21.10±5.11% as F4/80+ macrophages. A
small fraction (2.61±1.65%) was identified as CD3e positive T cells. Data are shown as
mean±SEM.
(I) Quantification of cell type specific afatinib-yne labeling intensity. Tissue samples from
four mice with 4-6 field of views acquired per mouse for analysis. A total of 23 afatinib-yne
negative hepatocytes, 23 afatinib-yne positive hepatocytes, 23 afatinib-yne negative
macrophages, 20 afatinib-yne positive macrophages, 26 afatinib-yne negative T cells, and 9
afatinib-yne positive T cells were included in the analysis. Intensity was normalized to the
average intensity of non-enriched hepatocytes. Data are shown as mean±SEM with individual
value plotted.
(J) High resolution ibrutinib-yne cell type characterization in the liver. Representative
images showing ibrutinib-yne enriched cells are F4/80+ macrophages (top row) or CD3e+ T
cells (bottom row). The dashed line represents the cell boundary. Scale bar, 10 µm.
(K) Cell type characterization of ibrutinib-yne enriched cells in the liver. 52.78±3.25% of
cells were identified as CD3e positive T cells, followed by 39.3±4.42% being F4/80+
macrophages and 13.95±1.32% being hepatocytes. Data are shown as mean±SEM.
(L) Quantification of cell type specific ibrutinib-yne labeling intensity. Tissue samples from
four mice with 4-6 fields of view acquired per mouse for analysis. A total of 20 ibrutinib-yne
negative hepatocytes, 6 ibrutinib -yne positive hepatocytes, 20 ibrutinib-yne negative
macrophages, 17 ibrutinib-yne positive macrophages, 33 ibrutinib-yne negative T cells, and
19 ibrutinib-yne positive T cells were included in the analysis. Intensity was normalized to the
average intensity of non-enriched hepatocytes. Data are shown as mean±SEM with individual
value plotted.

Figure S1. vCATCH development, related to Figure 1.
(A) Chemical structure of the monoamine oxidase A/B (MAO A/B) inhibitor pargyline and
their alkyne analogs used in the study.
(B) CATCH in 100-micron vehicle or PF7845-yne treated (1 mg/kg, 1 h, i.p.) brain sections.
Individual components (Cu, ligand, N3) were removed during the click reaction incubation.
Disruption of the Cu-ligand complex significantly impaired click labeling performance.
Representative images were taken at the primary somatosensory cortex (S1, imaging depth
Z=10 and 50 microns). Scale bar, 20 µm.

(C) Quantification of CATCH intensity in Figure S1B. Six tissue sections (one field of view per section) were used for each condition. Intensity at Z=10 or 50 microns was measured and then normalized to the average vehicle labeling intensity at Z=10 or 50 microns, respectively. Paired t test in each condition. p-values are listed in the graph.

(D) Transverse 2D views of hemispheres in Figure 1E. Scale bar, 2000 μm.

(E) Coronal 2D views of hemispheres in Figure 1E. Zoomed-in views showing pargyline-yne enriched structures across the whole brain. Scale bar: coronal views, 1000 μm; zoomed-in views, 100 μm. ACB, nucleus accumbens; MED, medial group of the dorsal thalamus; PMV, ventral premammillary nucleus; DR, dorsal nucleus raphe; AT, anterior tegmental nucleus; LC, locus coeruleus; B, Barrington's nucleus.

Figure S2. ACE analysis and validation, related to Figure 2.

(A) Atlas plot of the average intensity difference of vCATCH labeling intensity of vehicle vs. pargyline-yne treated samples. Five vehicle and five pargyline-yne-treated mice were included in the analysis. The greatest labeling intensity elevation is denoted in yellow.

(B) Atlas plot of the average segmented neuronal density (number of cells per mm^3) of vehicle vs. pargyline-yne treated samples. Five vehicle and five pargyline-yne-treated mice were included in the analysis. The greatest density elevation is denoted in yellow.

(C) Atlas plot of the average [-log(p-value)] of segmented neuronal density in vehicle vs. pargyline-yne treated samples. Five vehicle and 5 pargyline-yne-treated mice were included in the analysis. The greatest statistical significance is denoted in yellow.

Figure S3: whole body vCATCH visualization of TKI binding, related to Figure 3.

(A) Chemical structures of afatinib, ibrutinib, and their alkyne analogs.

(B) Representative 3D view of vehicle treated mice. Scale bar, 4000 μm.

(C) Afatinib-yne and ibrutinib-yne distribution overviews with digital plane slicing in 3D volume image. Scale bars, 3000 μm.

(D-E) Coronal (D), sagittal (E), and transverse (E) views of mice injected with afatinib-yne as shown in Figure S3B. Scale bars: coronal, 3000 μm; sagittal, 2000 μm; transverse (i and ii), 1000 μm.

(F-G) Coronal (D), sagittal (E), and transverse (E) views of mice injected with ibrutinib-yne as shown in Figure S3B. Scale bars: coronal, 3000 μm; sagittal, 2000 μm; transverse (i), 1500 μm; ibrutinib transverse (ii), 1000 μm.

(H) Representative organ views from afatinib-yne treated mice. Afatinib-yne enrichment was observed in small intestine villi, spleen central arterioles (white arrows), and kidney cortex. No significant drug enrichment observed in the stomach lining or nasal tract. Scale bars: small intestine, kidney, stomach, and nasal tract, 500 μm; spleen, 250 μm.

(I) Representative organ views from ibrutinib-yne treated mice. Ibrutinib-yne enrichment was observed in small intestine villi, spleen central arterioles (white arrows), kidney, and inner stomach lining. Scale bars: kidney and stomach, 500 μm; small intestine and nasal tract, 300 um; spleen, 200 μm.

Figure S4: TKI positive cell type characterization, related to Figure 4.

(A-B) Lectin staining in bone (A) and spleen (B) sections after CATCH labeling. Representative images showing lectin overlaps TKI enriched labeling structures. Scale bar, 20 μm.

(C) Global characterization of afatinib-yne binding in the lung. Zoomed-in views showing discrete afatinib-yne enriched structures scattered throughout the lung. Scale bar: global views, 1000 μm; zoomed-in views, 50 μm.

(D) High resolution afatinib-yne cell type characterization in the lung. Representative images showing afatinib-yne enriched macrophages (top) and CD19 positive B cells (bottom). Scale bar, 10 μm.

(E) Cell type characterization of afatinib-yne enriched cells in the lung. 24.15±4.19% of cells were identified as B cells, and 23.58±8.53% as macrophages. Data are shown as mean±SEM.

Supplementary Table 1: Brain-wide pargyline-yne binding intensity and cell count analysis.

Supplementary Video 1: 3D rendering of pargyline-yne binding across the whole mouse hemisphere.

Supplementary Video 2: 3D rendering of afatinib-yne binding across the whole mouse body.

Supplementary Video 3: 3D rendering of ibrutinib-yne binding across the whole mouse body.

STAR METHODS

Key Resources Table

Resource availability

Lead Contact

Further information and requests for resources and reagents should be directed to and will be fulfilled by the lead contact, Li Ye (liye@scripps.edu)

Materials availability

All unique/stable reagents generated in this study are available from the lead contact with a completed materials transfer agreement.

Data and code availability

All data are available in the main text or the supplementary materials. Any additional information required to reanalyze the data reported in this paper is available from the lead contact upon request. Relevant code for the study is available from https://github.com/yelabscripps/vCATCH.

EXPERIMENTAL MODEL AND SUBJECT DETAILS

Mouse model

Mice were group-housed on a 12-hr light-dark cycle and fed a standard rodent chow diet. Both male and female mice were used for the study. Six to nine-week-old WT C57BL6J mice were used for vCATCH method development in 100-micron sections and hemispheres. One-week-old WT C57BL6J mice were used for vCATCH in the whole mouse body. All experimental protocols were approved by the Scripps Research Institute Institutional Animal Care and Use Committee and were in accordance with the guidelines from the NIH.

Sample collection

Alkyne drugs were administered to mice in a vehicle of 10% DMSO and 2% Tween-80 in saline for i.p. injections. After drug injection at the designated dose and time, mice were heavily anesthetized with isoflurane and then transcardially perfused with ice-cold PBS, followed by ice-cold 4% PFA solution.

Mouse organs were carefully dissected and then post-fixed in 4% PFA solution overnight at 4℃. Tissues were washed with PBS. Mouse brains were embedded in 2% agarose and sectioned with a vibratome (Leica VT1000S) at 100-micron thickness. For hemispheres, brains were cut along the midline with a razor knife. Peripheral organs from one-week-old mice were dehydrated in 30% sucrose at 4℃ prior to sectioning. Samples were then carefully dried with a kimwipe and embedded in O.C.T. compound. After reaching equilibrium at -20℃, samples were cut at 100-micron thickness with a cryostat (Leica CM 1860). All obtained samples were stored in PBS with 0.02% sodium azide at 4℃ for further processing.

For mice used in whole-body drug distribution labeling, the skin was carefully removed immediately after perfusion, and the peritoneum was cut down the midline to facilitate clearing and click reactions. Mice were post-fixed in 4% PFA solution for two days at 4℃ followed by 7-day decalcification treatment in 10% EDTA/15% Imidazole at 4℃. Then, samples underwent decoloring treatment with 25% N, N, N', N'-Tetrakis(2 Hydroxypropyl)ethylenediamine (Quadrol) in 1x PBS at 37°C with gentle agitation for 7 days. After PBS washes to remove Quadrol, mouse whole bodies were stored in PBS with 0.02% sodium azide at 4℃ for further processing.

Tissue clearing

CLARITY

100-micron sections were cleared with CLARITY as previously described[22,81]. Briefly, samples were incubated in A1P4 hydrogel (1% acrylamide, 0.125% Bis, 4% PFA, 0.025% VA-044 initiator (w/v), in 1X PBS at 4℃ for CLARITY embedding. After overnight incubation with gentle agitation, samples were then degassed and polymerized at 37℃ for 4 hours with gentle agitation. Samples were removed from hydrogel and washed with 6% PBS-SDS (pH=7.0) at 37℃ for two days. After clearing, samples were washed with PBST (pH=7.0 with 0.2% Triton-X100, same for the following) three times (10 min each time) at RT to remove residue SDS. Samples were briefly washed with PBS and then stored in PBS with 0.02% sodium azide at 4℃.

HYBRiD

Mouse hemispheres and bodies were cleared with HYBRiD as previously described[27]. In brief, samples were dehydrated in an increasing THF/25% Quadrol gradient (1 hour per step for

hemispheres, 3 hours per step for whole bodies), delipidated in DCM (1 hour for hemispheres,
3x3 hours for whole bodies), and rehydrated in a decreasing THF/25% Quadrol gradient (0.5
hour per step for hemispheres, 1.5 hour per step for whole bodies). After clearing, samples were
extensively washed with PBS to remove residual organic solvent before hydrogel embedding.
Samples were equilibrated in A1P4 hydrogel (3 days for hemispheres, 7 days for whole bodies)
at 4°C on a shaker, then degassed and polymerized at 37°C for 4 hours with gentle agitation.
Thereafter, samples were passively cleared with LiOH Boric Buffer with 6% SDS (pH=9) until
transparent (1 week for hemispheres, 6 weeks for whole bodies). Samples were extensively
washed in PBST to remove residual SDS and then stored in PBS with 0.02% sodium azide at
4°C.

Click labeling in tissues

CATCH in 100-micron sections

100-micron sections were labeled with an updated CATCH protocol. Unless otherwise noted,
sections were incubated in 10 mM $CuSO_4$ in H_2O at RT overnight. Sections were then transferred
to CATCH reaction buffer containing 5 uM AZDye 647-picolyl azide, 150 μM CuSO4, 300 μM
BTTP, 2.5 mM NaAsb, and 10% DMSO in 1X PBS. After 1 hour of reaction in the dark with
minor agitation, samples were washed with buffer containing 4 mM EDTA in PBST (PBST-
EDTA, same for the following) at RT for 3x10 min. Tissue samples were then stained with DAPI
(1:3000 dilution in PBST from 10 μM stock) for 15 mins at RT. Samples were then either used
for refractive index (RI) matching by RapiClear and confocal imaging or proceeded to further
immunolabeling for cell-type registration.

vCATCH in hemispheres

For hemisphere labeling, each hemisphere was first pre-incubated with 25 mL of 10 mM CuSO4
in H_2O for one day at RT for PRCS. Hemispheres were transferred to vCATCH incubation buffer
(5 μM AZDye 647-picolyl azide, 1 mM CuSO4, 2 mM BTTP, 10% DMSO in 1X PBS, 3 mL for
each hemisphere) for 1 day at RT. Hemispheres were then transferred to the newly prepared full
vCATCH reaction buffer (5 μM AZDye 647-picolyl azide, 1 mM CuSO4, 2 mM BTTP, 25 mM
NaAsb, 10% DMSO in 1X PBS, 3 mL for each hemisphere) for click labeling. The vCATCH
reaction buffer was refreshed every hour for two rounds of RIR. Hemispheres were washed with
PBST-EDTA at RT. The washing buffer was refreshed at the end of the day and then daily until it
became colorless. The hemispheres were then washed with PBST for 1 hour at RT. After
washing, hemispheres were cut into 100-micron coronal sections for labeling depth
characterizations. Alternatively, hemispheres were carefully dried with a Kimwipe and then
incubated in 3 mL EasyIndex for RI matching and lightsheet imaging.

vCATCH in the whole mouse

Cleared mouse bodies were placed in 500 mL per mouse of 10 mM CuSO4 in H_2O with agitation
for 4 days at RT with daily buffer refreshes. After PRCS, whole-body samples were individually
incubated in 10 mL of vCATCH incubation buffer for 3 days on a shaker at RT with daily buffer
changes. Samples were then transferred to vCATCH reaction buffer (10 mL each) to initiate the
reaction. Five successive rounds of 1-hour click reactions were performed to thoroughly label the
entire tissue depth. After click labeling, samples were washed with PBST-EDTA with daily
refreshments until the buffer was colorless. After an additional day of PBST wash, mouse bodies

were carefully dried with a Kimwipe and then placed in 25 mL EasyIndex at 37°C for thorough
RI matching in preparation for lightsheet imaging.

Immunostaining

632   CATCH-labeled 100-micron sections were incubated with primary antibodies 1:200 diluted in
633   PBST overnight at 4°C. Samples were then washed with PBST 3 x 30 min at RT prior to
634   incubation with secondary antibodies diluted 1:600 in PBST. Samples were then incubated with
635   secondary antibodies 1:600 diluted in PBST. Samples were incubated with secondary antibodies
636   overnight and then washed with PBST 3 x 30 min at RT.

Confocal microscopy

100-micron samples were mounted with RapiClear for RI matching and imaged with an
Olympus FV3000 confocal microscope. To characterize CATCH labeling depth in 100-micron
sections, samples were imaged at the primary somatosensory cortex layer V with a 10X, 0.6 NA,
water immersion objective (XLUMPlanFI, Olympus) at 0.414 micron/pixel resolution and a step
size of 10 microns. To examine hemisphere labeling depth and global drug distribution within
individual organs, samples were imaged with a 10X, 0.6 NA, water immersion objective
(XLUMPlanFI, Olympus) at 2.49 microns/pixel resolution and a step size of 10 microns. For cell
type characterizations, samples were imaged at the top surface with a 40X, 1.25 NA, silicone oil
objective (UPlanSApo, Olympus) at 0.094 micron/pixel resolution and a step size of 3 microns.
All confocal data was saved in TIFF format for further analysis.

Lightsheet microscopy

Whole hemisphere/body samples were imaged using a lightsheet microscope from LifeCanvas
Technologies. RI-matched whole hemisphere and body samples were mounted in 1%
agarose/Easy Index for imaging. Light-sheet imaging was performed in a SmartSPIM chamber
with Easy Index matched immersion oil (RI=1.52). A 3.6x objective was used with 0.28 NA,
1.8μm/1.8μm/2μm xyz voxel size. Imaging was performed by illuminating samples along the
bilateral plane of symmetry. Lightsheet raw data was then destriped, stitched, and saved as TIFF
format for further conversion.

Image processing and analysis

Hemisphere labeling depth characterization

Images were processed as z projections with maximum intensity. A rectangle (805 x 50 pixels,
2001.17 x 136.73 microns) was drawn from the tissue surface as indicated. The signal profile
was plotted along the rectangle. The plotted data was then normalized by dividing each value by
the initial intensity value (distance=0 micron).

Thin section labeling depth characterization

CATCH labeling intensity across the 100-micron section was quantified as previously reported
[82,83]. Briefly, a 150 x 150 pixel (62.1 x 62.1 micron) ROI was generated at z=10μm and 50μm at
cortex layer V, and an auto threshold was applied to measure the mean drug-positive pixel
intensity as I_{signal}. A random line was then drawn in drug-negative areas to measure $I_{background}$. The
mean labeling intensity was defined as $I_{labeling}=I_{signal}-I_{background}$.

Cell type-specific intensity characterization

CD3e, F4/80 positive cells were characterized by respective immune staining. Hepatocytes were
identified based on their morphology and size. Individual cells were then cropped out, and a
threshold was applied to analyze CATCH labeling intensity in different cells of interest.

Volumetric drug binding visualization

Lightsheet imaged hemispheres and whole body images were converted into Imaris files and
visualized via Imaris 10.2.0. The hemisphere drug labeling video was generated by Imaris. For
VR-based whole-body drug visualization, data was converted by syGlass, and videos were made
with a Meta Quest 3 glass. For hemisphere coronal and transverse views, TIFF files were
imported as image sequences and then resliced by Fiji ImageJ. For drug distribution visualization
across organs, TIFF files were imported as image sequences and sampled at 100-micron step
size. Individual organ of interest was then cropped out for visualization.

Automated Brain-wide Cell Mapping & Signal Intensity Calculation.

We employed an AI-based Cartography of Ensembles (ACE) pipeline to generate whole-brain
segmentation maps of pargyline-yne+ cells[51]. ACE is an open-source 3D deep learning pipeline
(available through the MIRACL platform[84]; https://miracl.readthedocs.io/) to map local cell
activations using tera-voxel size light sheet microscopy datasets. Given the significant
differences in morphological features between pargyline-yne+ cells and cFos+ cells (which the
ACE models were originally trained on), we first fine-tuned the ACE vision transformer (ViT)
model. For fine-tuning, we randomly selected 225 image patches of 963 voxels each
$(0.17 \times 0.17 \times 0.19 \text{ mm}^3)$ from two subjects in the dataset and generated ground truth using the
Labkit annotator[85] in ImageJ/Fiji software[86]. We fine-tuned the ViT model on half of these
patches and used the other half for evaluation. The model was fine-tuned for 200 epochs using
the Adam optimizer[87] with an initial learning rate of 0.0001 and an equally weighted Dice-Cross
Entropy loss function. To account for artifacts in the dataset, the cross-entropy calculation for
"background" voxels was weighted 2× higher than for voxels annotated as pargyline-yne+ cells.
The best model, which achieved an average Dice score of 0.7 on the validation dataset, was used
to generate segmentation maps for the entire dataset. Subsequently, a shape filter was applied to
the segmentation maps to eliminate false positives such as vasculature traces in cortical regions
using the shape filter plugin in ImageJ/Fiji[88]. To distinguish individual cells, a connected
component analysis was applied to the 3D segmentation maps of each subject. Following
segmentation, we automatically registered the light sheet hemisphere data to the ABA using the
MIRACL platform[52,84]. The accuracy of the registration and warping were evaluated using
quality control checkpoints. Deformation fields obtained from the registration algorithm were
used to warp the ABA labels to the native space of each subject. Finally, we computed the
number of cells within each region at depth 6 (labels grouped to a maximum atlas ontology depth
of 6, by combining finer labels under their parent labels), by identifying the coordinates of the
center of each cell.

To map significant localized group-wise differences in pargyline-yne+ cells in an atlas-agnostic
manner, we employed ACE's cluster-wise threshold-free cluster enhancement permutation test
using a group-wise ANOVA. The segmentation maps obtained by the deep learning models were
downsampled to ABA 10-μm resolution while minimizing information loss by applying a
convolution-based voxelization procedure to the whole-brain segmentation maps (based on our
prior work[84]). Following voxelization, segmentation maps were aligned with the ABA at a 10-
μm resolution using deformation fields obtained via the registration module. The voxelated and

warped segmentation maps were then passed to the ACE statistical module (using default
parameters), which produced a cluster-wise p-value map representing significant (alpha < 0.05
after correction) localized cell activation between the pargyline-yne treated and control groups.

We computed signal intensity information for each ABA label using the "*miracl lbls stats*"
function in our MIRACL platform, which processes NIfTI images and their registered warped
ABA labels for each subject. This function extracts key intensity metrics, such as the average,
minimum, and maximum voxel intensity, across the input volumes for each label. We further
summarized it by grouping labels at depth 6 using the atlas ontology.

Brain-wide drug binding heatmap visualization

Brain regions with positive cell counts in all 5 pargyline-yne-treated hemispheres were included
for heatmap generation. Cell density (number of cells per mm^3) was obtained by dividing cell
count values by the volume of the brain region. Intensity and cell density were then normalized
by the average values in individual hemispheres. Heatmaps were generated using the Python
function 'sns.heatmap', with each region displaying the normalized intensity and density across
different subjects. The height of each subplot, corresponding to different regions, is proportional
to the number of areas within that region. Code for generating heatmaps is available via
https://github.com/yelabscripps/vCATCH.

REFERENCES

1. Huang, Z., Ogasawara, D., Seneviratne, U.I., Cognetta, A.B., 3rd, Am Ende, C.W., Nason, D.M., Lapham, K., Litchfield, J., Johnson, D.S., and Cravatt, B.F. (2019). Global Portrait of Protein Targets of Metabolites of the Neurotoxic Compound BIA 10-2474. ACS Chem Biol *14*, 192-197. 10.1021/acschembio.8b01097.

2. Chang, J.W., Cognetta, A.B., 3rd, Niphakis, M.J., and Cravatt, B.F. (2013). Proteome-wide reactivity profiling identifies diverse carbamate chemotypes tuned for serine hydrolase inhibition. ACS Chem Biol *8*, 1590-1599. 10.1021/cb400261h.

3. Van Esbroeck, A.C., Janssen, A.P., Cognetta, A.B., Ogasawara, D., Shpak, G., Van Der Kroeg, M., Kantae, V., Baggelaar, M.P., De Vrij, F.M., and Deng, H. (2017). Activity-based protein profiling reveals off-target proteins of the FAAH inhibitor BIA 10-2474. Science *356*, 1084-1087. 10.1126/science.aaf7497.

4. Reinecke, M., Brear, P., Vornholz, L., Berger, B.T., Seefried, F., Wilhelm, S., Samaras, P., Gyenis, L., Litchfield, D.W., Medard, G., et al. (2024). Chemical proteomics reveals the target landscape of 1,000 kinase inhibitors. Nat Chem Biol *20*, 577-585. 10.1038/s41589-023-01459-3.

5. Offensperger, F., Tin, G., Duran-Frigola, M., Hahn, E., Dobner, S., Ende, C.W.A., Strohbach, J.W., Rukavina, A., Brennsteiner, V., Ogilvie, K., et al. (2024). Large-scale chemoproteomics expedites ligand discovery and predicts ligand behavior in cells. Science *384*, eadk5864. 10.1126/science.adk5864.

6. Shalem, O., Sanjana, N.E., Hartenian, E., Shi, X., Scott, D.A., Mikkelson, T., Heckl, D., Ebert, B.L., Root, D.E., Doench, J.G., and Zhang, F. (2014). Genome-scale CRISPR-Cas9 knockout screening in human cells. Science *343*, 84-87. 10.1126/science.1247005.

7. Zhao, J., Tang, Z., Selvaraju, M., Johnson, K.A., Douglas, J.T., Gao, P.F., Petrassi, H.M., Wang, M.Z., and Wang, J. (2022). Cellular Target Deconvolution of Small Molecules Using a Selection-Based Genetic Screening Platform. ACS Cent Sci *8*, 1424-1434. 10.1021/acscentsci.2c00609.

8. Li, H., Ma, T., Remsberg, J.R., Won, S.J., DeMeester, K.E., Njomen, E., Ogasawara, D., Zhao, K.T., Huang, T.P., Lu, B., et al. (2023). Assigning functionality to cysteines by base editing of cancer dependency genes. Nat Chem Biol *19*, 1320-1330. 10.1038/s41589-023-01428-w.

9. Shi, J., Wang, E., Milazzo, J.P., Wang, Z., Kinney, J.B., and Vakoc, C.R. (2015). Discovery of cancer drug targets by CRISPR-Cas9 screening of protein domains. Nat Biotechnol *33*, 661-667. 10.1038/nbt.3235.

10. Perrin, J., Werner, T., Kurzawa, N., Rutkowska, A., Childs, D.D., Kalxdorf, M., Poeckel, D., Stonehouse, E., Strohmer, K., Heller, B., et al. (2020). Identifying drug targets in tissues and whole blood with thermal-shift profiling. Nat Biotechnol. 10.1038/s41587-019-0388-4.

11. Liu, Z., Wang, P., Wold, E.A., Song, Q., Zhao, C., Wang, C., and Zhou, J. (2021). Small-Molecule Inhibitors Targeting the Canonical WNT Signaling Pathway for the Treatment of Cancer. J Med Chem *64*, 4257-4288. 10.1021/acs.jmedchem.0c01799.

771   12.   Ji, H., Lu, X., Zhao, S., Wang, Q., Liao, B., Bauer, L.G., Huber, K.V.M., Luo, R., Tian,
772   R., and Tan, C.S.H. (2023). Target deconvolution with matrix-augmented pooling strategy
773   reveals cell-specific drug-protein interactions. Cell Chem Biol *30*, 1478-1487 e1477.
774   10.1016/j.chembiol.2023.08.002.

775   13.   Savitski, M.M., Reinhard, F.B., Franken, H., Werner, T., Savitski, M.F., Eberhard, D.,
776   Martinez Molina, D., Jafari, R., Dovega, R.B., Klaeger, S., et al. (2014). Tracking cancer
777   drugs in living cells by thermal profiling of the proteome. Science *346*, 1255784.
778   10.1126/science.1255784.

779   14.   Stickels, R.R., Murray, E., Kumar, P., Li, J., Marshall, J.L., Di Bella, D.J., Arlotta, P.,
780   Macosko, E.Z., and Chen, F. (2020). Highly sensitive spatial transcriptomics at near-
781   cellular resolution with Slide-seqV2. Nature Biotechnology. 10.1038/s41587-020-0739-1.

782   15.   Rodriques, S.G., Stickels, R.R., Goeva, A., Martin, C.A., Murray, E., Vanderburg, C.R.,
783   Welch, J., Chen, L.M., Chen, F., and Macosko, E.Z. (2019). Slide-seq: A scalable
784   technology for measuring genome-wide expression at high spatial resolution. Science
785   *363*, 1463-1467. 10.1126/science.aaw1219.

786   16.   Bhatia, H.S., Brunner, A.-D., Öztürk, F., Kapoor, S., Rong, Z., Mai, H., Thielert, M., Ali,
787   M., Al-Maskari, R., Paetzold, J.C., et al. (2022). Spatial proteomics in three-dimensional
788   intact specimens. Cell *185*, 5040-5058.e5019. 10.1016/j.cell.2022.11.021.

789   17.   Unterauer, E.M., Shetab Boushehri, S., Jevdokimenko, K., Masullo, L.A., Ganji, M.,
790   Sograte-Idrissi, S., Kowalewski, R., Strauss, S., Reinhardt, S.C.M., Perovic, A., et al.
791   (2024). Spatial proteomics in neurons at single-protein resolution. Cell *187*, 1785-
792   1800.e1716. 10.1016/j.cell.2024.02.045.

793   18.   Hickey, J.W., Neumann, E.K., Radtke, A.J., Camarillo, J.M., Beuschel, R.T., Albanese,
794   A., McDonough, E., Hatler, J., Wiblin, A.E., Fisher, J., et al. (2021). Spatial mapping of
795   protein composition and tissue organization: a primer for multiplexed antibody-based
796   imaging. Nat Methods. 10.1038/s41592-021-01316-y.

797   19.   Shi, H., He, Y., Zhou, Y., Huang, J., Maher, K., Wang, B., Tang, Z., Luo, S., Tan, P., Wu,
798   M., et al. (2023). Spatial atlas of the mouse central nervous system at molecular
799   resolution. Nature. 10.1038/s41586-023-06569-5.

800   20.   Zeng, H., Huang, J., Ren, J., Wang, C.K., Tang, Z., Zhou, H., Zhou, Y., Shi, H., Aditham,
801   A., Sui, X., et al. (2023). Spatially resolved single-cell translatomics at molecular
802   resolution. Science *380*, eadd3067. 10.1126/science.add3067.

803   21.   Pang, Z., Schafroth, M.A., Ogasawara, D., Wang, Y., Nudell, V., Lal, N.K., Yang, D.,
804   Wang, K., Herbst, D.M., Ha, J., et al. (2022). In situ identification of cellular drug targets
805   in mammalian tissue. Cell *185*, 1793-1805 e1717. 10.1016/j.cell.2022.03.040.

806   22.   Sylwestrak, E.L., Rajasethupathy, P., Wright, M.A., Jaffe, A., and Deisseroth, K. (2016).
807   Multiplexed Intact-Tissue Transcriptional Analysis at Cellular Resolution. Cell *164*, 792-
808   804. 10.1016/j.cell.2016.01.038.

809   23.   Murray, E., Cho, J.H., Goodwin, D., Ku, T., Swaney, J., Kim, S.Y., Choi, H., Park, Y.G.,
810   Park, J.Y., Hubbert, A., et al. (2015). Simple, Scalable Proteomic Imaging for High-
811   Dimensional Profiling of Intact Systems. Cell *163*, 1500-1514.
812   10.1016/j.cell.2015.11.025.

24.   Mai, H., Luo, J., Hoeher, L., Al-Maskari, R., Horvath, I., Chen, Y., Kofler, F., Piraud, M.,
Paetzold, J.C., Modamio, J., et al. (2023). Whole-body cellular mapping in mouse using
standard IgG antibodies. Nat Biotechnol. 10.1038/s41587-023-01846-0.

25.   Wang, X., Allen, W.E., Wright, M.A., Sylwestrak, E.L., Samusik, N., Vesuna, S., Evans,
K., Liu, C., Ramakrishnan, C., Liu, J., et al. (2018). Three-dimensional intact-tissue
sequencing of single-cell transcriptional states. Science *361*. 10.1126/science.aat5691.

26.   Tyler, D.S., Vappiani, J., Cañeque, T., Lam, E.Y., Ward, A., Gilan, O., Chan, Y.-C.,
Hienzsch, A., Rutkowska, A., and Werner, T. (2017). Click chemistry enables preclinical
evaluation of targeted epigenetic therapies. Science *356*, 1397-1401.

27.   Nudell, V., Wang, Y., Pang, Z., Lal, N.K., Huang, M., Shaabani, N., Kanim, W., Teijaro,
J., Maximov, A., and Ye, L. (2022). HYBRiD: hydrogel-reinforced DISCO for clearing
mammalian bodies. Nat Methods. 10.1038/s41592-022-01427-0.

28.   Murakami, T.C., Mano, T., Saikawa, S., Horiguchi, S.A., Shigeta, D., Baba, K., Sekiya,
H., Shimizu, Y., Tanaka, K.F., Kiyonari, H., et al. (2018). A three-dimensional single-cell-
resolution whole-brain atlas using CUBIC-X expansion microscopy and tissue clearing.
Nat Neurosci *21*, 625-637. 10.1038/s41593-018-0109-1.

29.   Kim, S.Y., Cho, J.H., Murray, E., Bakh, N., Choi, H., Ohn, K., Ruelas, L., Hubbert, A.,
McCue, M., Vassallo, S.L., et al. (2015). Stochastic electrotransport selectively enhances
the transport of highly electromobile molecules. Proc Natl Acad Sci U S A *112*, E6274-
6283. 10.1073/pnas.1510133112.

30.   Tainaka, K., Murakami, T.C., Susaki, E.A., Shimizu, C., Saito, R., Takahashi, K.,
Hayashi-Takagi, A., Sekiya, H., Arima, Y., Nojima, S., et al. (2018). Chemical Landscape
for Tissue Clearing Based on Hydrophilic Reagents. Cell Rep *24*, 2196-2210 e2199.
10.1016/j.celrep.2018.07.056.

31.   Park, J., Wang, J., Guan, W., Gjesteby, L.A., Pollack, D., Kamentsky, L., Evans, N.B.,
Stirman, J., Gu, X., Zhao, C., et al. (2024). Integrated platform for multiscale molecular
imaging and phenotyping of the human brain. Science *384*, eadh9979.
10.1126/science.adh9979.

32.   Ku, T., Guan, W., Evans, N.B., Sohn, C.H., Albanese, A., Kim, J.-G., Frosch, M.P., and
Chung, K. (2020). Elasticizing tissues for reversible shape transformation and accelerated
molecular labeling. Nat Methods *17*, 609-613.

33.   Lai, H.M., Tang, Y., Lau, Z.Y.H., Campbell, R.A.A., Yau, J.C.N., Chan, C.C.Y., Chan,
D.C.W., Wong, T.Y., Wong, H.K.T., Yan, L.Y.C., et al. (2022). Antibody stabilization for
thermally accelerated deep immunostaining. Nat Methods *19*, 1137-1146.
10.1038/s41592-022-01569-1.

34.   Susaki, E.A., Shimizu, C., Kuno, A., Tainaka, K., Li, X., Nishi, K., Morishima, K., Ono,
H., Ode, K.L., Saeki, Y., et al. (2020). Versatile whole-organ/body staining and imaging
based on electrolyte-gel properties of biological tissues. Nat Commun *11*, 1982.
10.1038/s41467-020-15906-5.

35.   Zhao, S., Todorov, M.I., Cai, R., Maskari, R.A.I., Steinke, H., Kemter, E., Mai, H., Rong,
Z., Warmer, M., Stanic, K., et al. (2020). Cellular and Molecular Probing of Intact Human
Organs. Cell. 10.1016/j.cell.2020.01.030.

36.   Choi, S.W., Guan, W., and Chung, K. (2021). Basic principles of hydrogel-based tissue
transformation technologies and their applications. Cell *184*, 4115-4136.
10.1016/j.cell.2021.07.009.

37.   Yau, C.N., Lai, H.M., Lee, K., Kwok, A.J., Huang, J., and Ko, H. (2023). Principles of
deep immunohistochemistry for 3D histology. Cell Rep Methods *3*, 100458.
10.1016/j.crmeth.2023.100458.

38.   Richardson, D.S., Guan, W., Matsumoto, K., Pan, C., Chung, K., Erturk, A., Ueda, H.R.,
and Lichtman, J.W. (2021). Tissue Clearing. Nat Rev Methods Primers *1*.
10.1038/s43586-021-00080-9.

39.   Ueda, H.R., Erturk, A., Chung, K., Gradinaru, V., Chedotal, A., Tomancak, P., and Keller,
P.J. (2020). Tissue clearing and its applications in neuroscience. Nat Rev Neurosci *21*, 61-
79. 10.1038/s41583-019-0250-1.

40.   Dejaegere, A., Choulier, L., Lafont, V., De Genst, E., and Altschuh, D. (2005). Variations
in Antigen− Antibody Association Kinetics as a Function of pH and Salt Concentration: A
QSAR and Molecular Modeling Study. Biochemistry *44*, 14409-14418.

41.   Cann, J.R., and Clark, E.W. (1956). Kinetics of the Antigen-Antibody Reaction. Effect of
Salt Concentration and pH on the Rate of Neutralization of Bacteriophage by Purified
Fractions of Specific Antiserum1a, b. J Am Chem Soc *78*, 3627-3631.

42.   Himo, F., Lovell, T., Hilgraf, R., Rostovtsev, V.V., Noodleman, L., Sharpless, K.B., and
Fokin, V.V. (2005). Copper (I)-catalyzed synthesis of azoles. DFT study predicts
unprecedented reactivity and intermediates. J Am Chem Soc *127*, 210-216.

43.   Krysiak, J.M., Kreuzer, J., Macheroux, P., Hermetter, A., Sieber, S.A., and Breinbauer, R.
(2012). Activity-based probes for studying the activity of flavin-dependent oxidases and
for the protein target profiling of monoamine oxidase inhibitors. Angew Chem Int Ed
Engl *51*, 7035-7040. 10.1002/anie.201201955.

44.   Festa, R.A., and Thiele, D.J. (2011). Copper: an essential metal in biology. Curr Biol *21*,
R877-883. 10.1016/j.cub.2011.09.040.

45.   Inesi, G. (2017). Molecular features of copper binding proteins involved in copper
homeostasis. IUBMB Life *69*, 211-217. 10.1002/iub.1590.

46.   Andreini, C., Banci, L., Bertini, I., and Rosato, A. (2008). Occurrence of copper proteins
through the three domains of life: a bioinformatic approach. The Journal of Proteome
Research *7*, 209-216.

47.   Selden, C.R., Schilling, K., Godfrey, L., and Yee, N. (2024). Metal-binding amino acid
ligands commonly found in metalloproteins differentially fractionate copper isotopes. Sci
Rep *14*, 1902. 10.1038/s41598-024-52091-7.

48.   Li, S., Cai, H., He, J., Chen, H., Lam, S., Cai, T., Zhu, Z., Bark, S.J., and Cai, C. (2016).
Extent of the Oxidative Side Reactions to Peptides and Proteins During the CuAAC
Reaction. Bioconjug Chem *27*, 2315-2322. 10.1021/acs.bioconjchem.6b00267.

49.   Abel, G.R., Jr., Calabrese, Z.A., Ayco, J., Hein, J.E., and Ye, T. (2016). Measuring and
Suppressing the Oxidative Damage to DNA During Cu(I)-Catalyzed Azide-Alkyne
Cycloaddition. Bioconjug Chem *27*, 698-704. 10.1021/acs.bioconjchem.5b00665.

50.   Wiest, A., and Kielkowski, P. (2024). Cu-Catalyzed Azide-Alkyne-Thiol Reaction Forms
Ubiquitous Background in Chemical Proteomic Studies. J Am Chem Soc.
10.1021/jacs.3c11780.

51.   Attarpour, A., Osmann, J., Rinaldi, A., Qi, T., Lal, N., Patel, S., Rozak, M., Yu, F., Cho,
N., Squair, J., et al. A deep learning pipeline for 3D brain-wide mapping of local neuronal
ensembles in tera-voxel light sheet microscopy. Nat Methods *in press*.

52.   Wang, Q., Ding, S.-L., Li, Y., Royall, J., Feng, D., Lesnar, P., Graddis, N., Naeemi, M.,
Facer, B., and Ho, A. (2020). The Allen mouse brain common coordinate framework: a
3D reference atlas. Cell *181*, 936-953. e920.

53.   Floresco, S.B. (2015). The nucleus accumbens: an interface between cognition, emotion,
and action. Annual review of psychology *66*, 25-52.

54.   Petko, B., and Tadi, P. (2019). Neuroanatomy, nucleus ambiguus.

55.   Boike, L., Henning, N.J., and Nomura, D.K. (2022). Advances in covalent drug
discovery. Nat Rev Drug Discov. 10.1038/s41573-022-00542-z.

56.   Ghosh, A.K., Samanta, I., Mondal, A., and Liu, W.R. (2019). Covalent Inhibition in Drug
Discovery. ChemMedChem *14*, 889-906. 10.1002/cmdc.201900107.

57.   Miller, V.A., Hirsh, V., Cadranel, J., Chen, Y.-M., Park, K., Kim, S.-W., Zhou, C., Su, W.-
C., Wang, M., Sun, Y., et al. (2012). Afatinib versus placebo for patients with advanced,
metastatic non-small-cell lung cancer after failure of erlotinib, gefitinib, or both, and one
or two lines of chemotherapy (LUX-Lung 1): a phase 2b/3 randomised trial. The Lancet
Oncology *13*, 528-538. 10.1016/s1470-2045(12)70087-6.

58.   Sequist, L.V., Yang, J.C., Yamamoto, N., O'Byrne, K., Hirsh, V., Mok, T., Geater, S.L.,
Orlov, S., Tsai, C.M., Boyer, M., et al. (2013). Phase III study of afatinib or cisplatin plus
pemetrexed in patients with metastatic lung adenocarcinoma with EGFR mutations. J
Clin Oncol *31*, 3327-3334. 10.1200/jco.2012.44.2806.

59.   Byrd, J.C., Furman, R.R., Coutre, S.E., Flinn, I.W., Burger, J.A., Blum, K.A., Grant, B.,
Sharman, J.P., Coleman, M., Wierda, W.G., et al. (2013). Targeting BTK with ibrutinib in
relapsed chronic lymphocytic leukemia. N Engl J Med *369*, 32-42.
10.1056/NEJMoa1215637.

60.   Treon, S.P., Tripsas, C.K., Meid, K., Warren, D., Varma, G., Green, R., Argyropoulos,
K.V., Yang, G., Cao, Y., Xu, L., et al. (2015). Ibrutinib in previously treated
Waldenstrom's macroglobulinemia. N Engl J Med *372*, 1430-1440.
10.1056/NEJMoa1501548.

61.   Wang, M.L., Rule, S., Martin, P., Goy, A., Auer, R., Kahl, B.S., Jurczak, W., Advani,
R.H., Romaguera, J.E., Williams, M.E., et al. (2013). Targeting BTK with ibrutinib in
relapsed or refractory mantle-cell lymphoma. N Engl J Med *369*, 507-516.
10.1056/NEJMoa1306220.

62.   Yun, S., Vincelette, N.D., Acharya, U., and Abraham, I. (2017). Risk of Atrial Fibrillation
and Bleeding Diathesis Associated With Ibrutinib Treatment: A Systematic Review and
Pooled Analysis of Four Randomized Controlled Trials. Clin Lymphoma Myeloma Leuk
17, 31-37 e13. 10.1016/j.clml.2016.09.010.

63.   Lipsky, A.H., Farooqui, M.Z., Tian, X., Martyr, S., Cullinane, A.M., Nghiem, K., Sun, C.,
Valdez, J., Niemann, C.U., Herman, S.E., et al. (2015). Incidence and risk factors of
bleeding-related adverse events in patients with chronic lymphocytic leukemia treated
with ibrutinib. Haematologica *100*, 1571-1578. 10.3324/haematol.2015.126672.

64.   Berglof, A., Hamasy, A., Meinke, S., Palma, M., Krstic, A., Mansson, R., Kimby, E.,
Osterborg, A., and Smith, C.I. (2015). Targets for Ibrutinib Beyond B Cell Malignancies.
Scand J Immunol *82*, 208-217. 10.1111/sji.12333.

65.   Tuomi, J.M., Xenocostas, A., and Jones, D.L. (2018). Increased Susceptibility for Atrial
and Ventricular Cardiac Arrhythmias in Mice Treated With a Single High Dose of
Ibrutinib. Can J Cardiol *34*, 337-341. 10.1016/j.cjca.2017.12.001.

66.   Xiao, L., Salem, J.E., Clauss, S., Hanley, A., Bapat, A., Hulsmans, M., Iwamoto, Y.,
Wojtkiewicz, G., Cetinbas, M., Schloss, M.J., et al. (2020). Ibrutinib-Mediated Atrial
Fibrillation Attributable to Inhibition of C-Terminal Src Kinase. Circulation *142*, 2443-
2455. 10.1161/CIRCULATIONAHA.120.049210.

67.   McMullen, J.R., Boey, E.J., Ooi, J.Y., Seymour, J.F., Keating, M.J., and Tam, C.S.
(2014). Ibrutinib increases the risk of atrial fibrillation, potentially through inhibition of
cardiac PI3K-Akt signaling. Blood, The Journal of the American Society of Hematology
124, 3829-3830.

68.   Lanning, B.R., Whitby, L.R., Dix, M.M., Douhan, J., Gilbert, A.M., Hett, E.C., Johnson,
T.O., Joslyn, C., Kath, J.C., Niessen, S., et al. (2014). A road map to evaluate the
proteome-wide selectivity of covalent kinase inhibitors. Nat Chem Biol *10*, 760-767.
10.1038/nchembio.1582.

69.   Kaltenecker, D., Al-Maskari, R., Negwer, M., Hoeher, L., Kofler, F., Zhao, S., Todorov,
M., Rong, Z., Paetzold, J.C., Wiestler, B., et al. (2024). Virtual reality-empowered deep-
learning analysis of brain cells. Nat Methods *21*, 1306-1315. 10.1038/s41592-024-02245-
2.

70.   CENTER FOR DRUG EVALUATION AND RESEARCH APPLICATION NUMBER:
201292Orig1s000. (2013).
https://www.accessdata.fda.gov/drugsatfda_docs/nda/2013/201292Orig1s000Approv.pdf.

71.   CENTER FOR DRUG EVALUATION AND RESEARCH APPLICATION NUMBER:
205552Orig2s000. (2014).
https://www.accessdata.fda.gov/drugsatfda_docs/nda/2014/205552Orig2s000MedR.pdf.

72.   Karlsson, M., Zhang, C., Méar, L., Zhong, W., Digre, A., Katona, B., Sjöstedt, E., Butler,
L., Odeberg, J., Dusart, P., et al. (2021). A single–cell type transcriptomics map of human
tissues. Science Advances *7*, eabh2169. doi:10.1126/sciadv.abh2169.

73.   Tom Dieck, S., Muller, A., Nehring, A., Hinz, F.I., Bartnik, I., Schuman, E.M., and
Dieterich, D.C. (2012). Metabolic labeling with noncanonical amino acids and
visualization by chemoselective fluorescent tagging. Curr Protoc Cell Biol *Chapter 7*, 7
11 11-17 11 29. 10.1002/0471143030.cb0711s56.

74.   Jao, C.Y., and Salic, A. (2008). Exploring RNA transcription and turnover in vivo by
using click chemistry. Proc Natl Acad Sci U S A *105*, 15779-15784.

75. Nikic, I., Kang, J.H., Girona, G.E., Aramburu, I.V., and Lemke, E.A. (2015). Labeling
proteins on live mammalian cells using click chemistry. Nat Protoc *10*, 780-791.
10.1038/nprot.2015.045.

76. El-Sagheer, A.H., and Brown, T. (2010). Click chemistry with DNA. Chemical Society
Reviews *39*, 1388-1405.

77. Soriano del Amo, D., Wang, W., Jiang, H., Besanceney, C., Yan, A.C., Levy, M., Liu, Y.,
Marlow, F.L., and Wu, P. (2010). Biocompatible copper (I) catalysts for in vivo imaging
of glycans. J Am Chem Soc *132*, 16893-16899.

78. Hulce, J.J., Cognetta, A.B., Niphakis, M.J., Tully, S.E., and Cravatt, B.F. (2013).
Proteome-wide mapping of cholesterol-interacting proteins in mammalian cells. Nat
Methods *10*, 259-264. 10.1038/nmeth.2368.

79. Tamura, I., Sakamoto, D.M., Yi, B., Saito, Y., Yamada, N., Morimoto, J., Takakusagi, Y.,
Kuroda, M., Kubota, S.I., and Yatabe, H. (2024). Click3D: Click reaction across deep
tissues for whole-organ 3D fluorescence imaging. Science Advances *10*, eado8471.

80. Wang, Y., Leung, V.H., Zhang, Y., Nudell, V.S., Loud, M., Servin-Vences, M.R., Yang,
D., Wang, K., Moya-Garzon, M.D., Li, V.L., et al. (2022). The role of somatosensory
innervation of adipose tissues. Nature *609*, 569-574. 10.1038/s41586-022-05137-7.

81. Chung, K., Wallace, J., Kim, S.Y., Kalyanasundaram, S., Andalman, A.S., Davidson, T.J.,
Mirzabekov, J.J., Zalocusky, K.A., Mattis, J., Denisin, A.K., et al. (2013). Structural and
molecular interrogation of intact biological systems. Nature *497*, 332-337.
10.1038/nature12107.

82. Qi, Y., Yu, T., Xu, J., Wan, P., Ma, Y., Zhu, J., Li, Y., Gong, H., Luo, Q., and Zhu, D.
(2019). FDISCO: Advanced solvent-based clearing method for imaging whole organs.
Science Advances *5*, eaau8355. 10.1126/sciadv.aau8355.

83. Pan, C., Cai, R., Quacquarelli, F.P., Ghasemigharagoz, A., Lourbopoulos, A., Matryba, P.,
Plesnila, N., Dichgans, M., Hellal, F., and Erturk, A. (2016). Shrinkage-mediated imaging
of entire organs and organisms using uDISCO. Nat Methods *13*, 859-867.
10.1038/nmeth.3964.

84. Goubran, M., Leuze, C., Hsueh, B., Aswendt, M., Ye, L., Tian, Q., Cheng, M.Y., Crow,
A., Steinberg, G.K., and McNab, J.A. (2019). Multimodal image registration and
connectivity analysis for integration of connectomic data from microscopy to MRI.
Nature communications *10*, 5504.

85. Arzt, M., Deschamps, J., Schmied, C., Pietzsch, T., Schmidt, D., Tomancak, P., Haase, R.,
and Jug, F. (2022). LABKIT: labeling and segmentation toolkit for big image data.
Frontiers in computer science *4*, 777728.

86. Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T.,
Preibisch, S., Rueden, C., Saalfeld, S., Schmid, B., et al. (2012). Fiji: an open-source
platform for biological-image analysis. Nat Methods *9*, 676-682. 10.1038/nmeth.2019.

87. Kinga, D., and Adam, J.B. (2015). A method for stochastic optimization. (San Diego,
California;), pp. 6.

88.   Wagner, T., and Lipinski, H.-G. (2013). IJBlob: an ImageJ library for connected
1019   component analysis and shape analysis. Journal of Open Research Software *1*.

1020

Figure 1

Figure 2

A Passive clearing — Passive labeling

Drug-Injected Brains → HYBRiD Clearing → Whole Hemisphere *vCATCH* (PRCS, 2X R/R) → Lightsheet Imaging

ACE (AI-based Cartography of Ensembles)

Deep learning based segmentation ↔ Allen brain atlas registration

Regional and cellular statistics — Intensity / Regions — Density / Regions

B *Regional drug intensity* in atlas space

Vehicle / Pargyline-yne / Pargyline-yne / Pargyline-yne

LC / B

C Segmented *drug-bound neurons* in atlas space

Vehicle / Pargyline-yne

i Pons / ii Hypothalamus / iii Medulla

D Normalized drug intensity (A.U.)

ACB BST EPI MED PMD PMV AT DR B LC SLC SLD AMB

indiv. mouse

Cerebral cortex | Cerebral nuclei | Thalamus | Hypo-thalamus | Midbrain | Pons | Cerebellum | Medulla | VS | fiber tracts

2.0 / 1.5 / 1.0 / 0.5

E Normalized drug-bound cell density (A.U.)

ACB BST EPI MED PMD PMV AT DR B LC SLC SLD AMB

indiv. mouse

Cerebral cortex | Cerebral nuclei | Thalamus | Hypo-thalamus | Midbrain | Pons | Cerebellum | Medulla | VS | fiber tracts

4 / 3 / 2 / 1 / 0

Figure 3

A Passive click labeling

in vivo Drug Injection — HYBRiD Clearing — Whole-body *vCATCH* (4 x 1 days, 3 x 1 days, 1h 1h 1h 1h 1h, PRCS, incubation, Cu II, Cu I, 5X RIR) — Lightsheet Imaging — Whole-body Drug Mapping (Organ-level, Cellular-level, Molecular-level)

B x-y, Z: Imaging depth, Dorsal to Ventral

C Z=4.4 mm Z=5.9 mm Z=7.3 mm Z=8.8 mm Z=9.9 mm
x-y in-plane views

Afatinib-yne

D Whole-body View

E Upper chest — LA, RA, Heart, LV, RV, Lung

F Lung

G

H Liver

I

Ibrutinib-yne

J Whole-body View

K Upper Chest — LA, RA, Heart, LV, RV, Lung

L Lung

M

N Liver

O

Figure 4

Figure S1

A

pargyline   pargyline-yne   PF7845   PF7845-yne

B

Vehicle   PF7845-yne

z=10 micron   Full buffer | Full buffer | No N₃ | No ligand | No CuSO₄

z=50 micron

CATCH in 100 micron brain sections

C

Normalized fluorescence intensity

0.9729   0.2352   0.6230   0.0001   0.0014

Z=10 micron   Z=50 micron

Full | Full | no N3 | no BTTP | no Cu

Vehicle   PF7845-yne

D

Vehicle   Transverse   Pargyline-yne   Transverse

E

Vehicle

Coronal Bregma 1.5 | Coronal Bregma -0.2 | Coronal Bregma -1.5 | Coronal Bregma -3.5 | Coronal Bregma -4.7 | Coronal Bregma -5.4

Pargyline-yne

Coronal Bregma 1.5 | Coronal Bregma -0.2 | Coronal Bregma -1.5 | Coronal Bregma -3.5 | Coronal Bregma -4.7 | Coronal Bregma -5.4

ACB | Choroid plexus | MED | PMV | DR / AT | B / LC

Figure S2

A

Average *regional intensity* difference in atlas space

B

Average *regional neuronal density* differences in atlas space

C

Average *regional neuronal density -log(p-value)* in atlas space

Figure S3

Figure S4

A Afatinib-yne **Bone** | Ibrutinib-yne **Bone**

Afatinib-yne | Lectin | DAPI | Ibrutinib-yne | Lectin | DAPI

B Afatinib-yne **Spleen** | Ibrutinib-yne **Spleen**

Afatinib-yne | Lectin | DAPI | Ibrutinib-yne | Lectin | DAPI

C Afatinib-yne lung | i | ii

D Afatinib-yne | F4/80 | DAPI

Afatinib-yne | CD19 | DAPI

E Afatinib-yne+ cells (100%)

Afatinib-yne+ B cells (24.15±4.19%)

Afatinib-yne+ macrophages (23.58±8.53%)