Memory's avalanche: Recent feelings of familiarity and resultant DA nuclei activity enable subsequent memory reactivation

Matthew R. Dougherty[1], Anuya Patil[1], Katherine D. Duncan

Department of Psychology, University of Toronto
100 St. George Street, 4th floor,
Toronto, ON, Canada, M5S 3G3

Corresponding Author: Matthew Dougherty
Department of Psychology, University of Toronto
100 St. George Street, Room 627
Toronto, ON, Canada, M5S 3GS
416-946-0207
Matthew.dougherty@mail.utoronto.ca

1: denotes equal contribution.

Abstract

Memory retrieval is notoriously variable. Various neurocognitive states have been theorized to

affect retrieval success from moment to moment, but the presence and catalysts of these states in

the human brain remain largely understudied. Building on previous work, we studied how recent

memory experiences (i.e. novelty vs. familiarity) and their corresponding neural activity prepare

the brain to reactivate memories. Our data uncover a pronounced neural bias, whereby regions

representing episodic information fail to reactivate memories following recent experiences of

novelty. Drawing on computational models, we hypothesized that novelty-evoked activity in

cholinergic nuclei underlie this bias. The data revealed an unexpected alternative mediator of

memory reactivation in response to recent novelty, namely recent dopaminergic nuclei activity.

By identifying dopamine as a potential mediator of retrieval variability, our results challenge

existing models and open new avenues for investigating how neuromodulatory states

dynamically shape memory accessibility.

Keywords: episodic memory, human, fMRI, retrieval, novelty, familiarity, dopamine,

acetylcholine, neuromodulators, reinstatement

Introduction

Making effective decisions hinges on recalling relevant experiences, but memory retrieval is

notoriously inconsistent. At times, past events vividly spring to mind with little effort; at other

times, we fail to even recall a colleague's name. What drives this frustrating variability? Decades

of memory research have highlighted the quality of memory formation as a major determinant

(1–5). When a well-learned memory is cued with a congruent reminder, successful retrieval often

follows (6, 7). However, even well-formed memories can fail to surface when cued, suggesting

that factors beyond initial learning shape variability in moment-to-moment retrieval success (7,

8). Here, we investigate the cognitive and neural mechanisms that contribute to this variability,

revealing factors that prepare our brains to recall past experiences.

The idea that retrieval depends on fluctuating preparatory states of mind was first proposed

by Endel Tulving in 1983. He theorized that successful episodic retrieval requires proactively

recruiting the retrieval mode, a neurocognitive state including the goal to retrieve memories and

the orientation of attention towards reactivated information (9). Recent theoretical and empirical

work has revived and extended Tulving's ideas, characterizing neural states contributing to

retrieval goals and internal attention (10–12). For the retrieval mode to be fruitful, though, the

brain must reactivate memories, a process termed ecphory by Tulving (1983), and more recently

formalized as pattern completion (13–15). Specifically, the hippocampus can reactivate full

memories in response to partial retrieval cues, which in turn reinstates memory components in

corresponding neocortical regions (16, 17) that can be observed through analyses of functional

magnetic resonance imaging (fMRI) patterns (17–20). While hippocampal pattern completion is

conventionally construed as automatic, it might be possible to prime this process as well (12, 21,

22). Hinting at this, Patil and Duncan (2018) behaviourally showed that people were less

accurate at retrieving associations after detecting a novel image than they were after recognizing

a familiar but unrelated image.

Here we leverage novelty's impact on memory to explore neural mechanisms underlying

retrieval variability. One possible mechanism is that novelty engages neuromodulatory systems

that suppress upcoming retrieval. Dopamine (DA), acetylcholine (ACh), and norepinephrine

(NE)—each of which is released in response to novelty (23–28)—have lingering effects on

attention (29–34), goal maintenance (34–38), and memory (22, 30, 33, 38–41). DA and NE are

well-positioned to engage the retrieval mode by enhancing goal maintenance and attention (29,

32–34, 36–38), making them unlikely to underpin novelty's suppression of retrieval. A more

plausible candidate is ACh. Cholinergic neurons in the septum (CH1 and CH2) and portions of

the diagonal band of Broca (CH3) project to the hippocampus(42), where increased ACh levels

are hypothesized to suppress retrieval-based processing (i.e. pattern completion) in favour of

encoding-based processing (i.e. pattern separation(22)). This cholinergically mediated

hippocampal switch would reduce brain-wide memory reactivation and retrieval success. Thus,

comparatively, recent familiarity would beget more fluid pattern completion and memory

reactivation than recent novelty, consistent with the behavioral results of Patil and Duncan

(2018).

In this study, we used functional magnetic resonance imaging (fMRI) to test this possibility.

Specifically, we measured activity across regions known to display memory reactivation along

with neuromodulatory nuclei while participants learned and retrieved associations. We predict

that, if Patil and Duncan's (2018) findings reflect a novelty-evoked bias away from hippocampal

pattern completion, then neural reinstatement of memories will also be lower following novelty

compared to familiarity. We also hypothesized that preceding activity in cholinergic nuclei would

mediate recent novelty's effect on reinstatement. This pattern of results would be consistent with

the idea that ongoing fluctuations in ACh release could explain retrieval variability. While our

results confirmed that neural reinstatement is robustly modulated by recent memory experiences,

they also revealed an unexpected set of neuromodulatory mechanisms mediating this effect.

Results

Creating an fMRI paradigm suited to investigate the preparatory states affecting memory

reactivation

We designed a memory task to evaluate how novelty, and the neuromodulatory nuclei

activity it evokes, influences the upcoming neural reinstatement of memories. As this

neuromodulatory influence is expected to last for only a few seconds (43, 44), we carefully

designed our experiment to overcome the slow nature of the blood-oxygen-level-

dependent (BOLD) signal (45). Specifically, we limited inter-trial-intervals (ITIs) to 1 second to

measure these short-lived neuromodulatory biases, and we used representationally distinct

stimulus categories to disambiguate signals arising from consecutive trials. In retrieval blocks,

trials were presented in a series of pairs, such that participants retrieved different stimulus

categories from trial to trial (Fig. 1a). The first trial of each pair was designed to induce

novelty/familiarity, in which participants viewed a trial-unique new or previously studied object

(*induction trials*). When they recognized an object, they attempted to recall the word associated

with it during encoding ("Plain" or "Safe"). The second trial of each pair was designed to assess

associative retrieval performance and neural reinstatement following novelty/familiarity, in

which participants viewed trial-unique new or previously studied words (*probe trials*). When

they recognized a word, they attempted to recall the stimulus category of the image associated

with it during encoding ("Face" or "Scene"; see *Methods: Procedure* for full design

specifications). For analyses, we categorized induction trials according to participants' subjective

memory judgements. If they responded with "new," trials were considered inductions of novelty;

if they recognized the objects, regardless of their associative retrieval success, trials were

considered inductions of familiarity (see *Methods: Retrieval Session* for response options). We

extended ITIs between trial pairs to 5 seconds to increase the sensitivity of our reinstatement

analyses.

First, we assessed how our induction trial manipulation influences memory retrieval in

this unique paradigm. After replicating Patil and Duncan (2018), we assessed how this

manipulation influenced the neural reinstatement of memories using encoding-retrieval similarity

analyses. We then measured how neuromodulatory nuclei responded to novelty on induction

trials, before determining if these responses mediated novelty's impact on neural reinstatement.

Fig. 1 | Task Design & Behavioral Analyses. **a**, Schematic of encoding & retrieval tasks.

During Object Encoding, participants associated trial-unique objects with repeating words.

During Word Encoding, participants associated trial-unique words with repeating face and scene

images (see *Methods: Encoding Blocks*). At retrieval, participants were shown the trial-unique

half of an encoded pair (object image or non-repeating word) and asked to retrieve its associate.

Trials were structured in pairs, with object trials acting as inductions of novelty/familiarity and

word trials acting as probes to measure neural reinstatement & associative memory accuracy (see

Methods: Retrieval Session). **b**, Associative memory accuracy on probe trials as a function of

preceding memory experiences. Points represent the estimated marginal means and error bars

represent their standard error from a generalized linear mixed effects model (see *Methods:*

Behavioral data analyses). **c**, Item-only recognition memory (d') on probe trials as a function

preceding memory experiences. Points represent mean d' across participants for trials preceded

by familiarity and novelty, respectively. Error bars represent the standard error of the mean,

applying the Morey, 2008 correction for within-participant comparisons. **d**, Response time on

probe trials as a function of preceding memory experiences. Points represent the estimated

marginal means and error bars represent their standard error from a linear mixed effects model

(see *Methods: Behavioral data analyses*). ~ indicates $0.10 > p > 0.05$, *** indicates $p < 0.001$.

Experiences of novelty compared to familiarity reduce successive associative memory, but

not recognition memory

We first sought to replicate and extend Patil & Duncan's (2018) demonstration that

associative memory accuracy was lower following experiences of novelty compared to

familiarity. In this experiment, associative memory accuracy was, indeed, significantly lower

following experiences of novelty ($b = 0.184$, $SE = 0.0674$, $p < 0.001$, 95% CI = [0.059, 0.310];

Fig. 1b). This effect held when controlling for the associate's stimulus class (i.e. face vs. scene)

and the preceding trial's response time (SI Appendix, Table S1). Like Patil & Duncan (2018), we

also found that item-only recognition memory accuracy (i.e., recognition in the absence of

associative memory) was not significantly influenced by preceding experiences of novelty (t(26)

= -1.55, p = 0.134, d = -0.3, 95% CI = [-0.68, 0.09]; Fig. 1c). Numerically, this recognition

accuracy was even higher following novelty. Our novelty manipulation's specific influence on

associative memory is consistent with our hypothesis that the hippocampus is biased away from

pattern completion following novelty compared to familiarity (21, 47).

However, two trends raise the possibility that another mechanism could also contribute to

these sequential memory effects. First, participants were marginally slower to correctly retrieve

associations following experiences of familiarity compared to novelty (b = 0.013, SE = 0.015, p

= 0.063, 95% CI = [-0.001, 0.026], Fig. 1d). This potential speed-accuracy trade-off suggests that

associative accuracy benefits could reflect participants being more willing to engage in effortful

processing following familiarity. We further probed this possibility in an exploratory analysis; we

asked if the relative benefits of recent familiarity extend beyond retrieval to also foster memory

formation. At the end of the experiment, participants completed a short recognition test (see

Methods: Localizer Tasks), which assessed their memory for 16 words first shown during

retrieval blocks. Critically, these novel foils were presented on probe trials following either the

induction of novelty or familiarity. Participants were marginally better at recognizing words

encoded following experiences of familiarity compared to novelty (b = 0.223, SE = 0.121, p =

0.065, 95% CI = [-0.014, 0.461]; SI Appendix, Table S2). This encoding effect could be

explained by participants adopting more effortful processing following familiarity compared to

novelty, but it would be hard to explain with a pattern completion bias, as this process is unlikely

to facilitate memory formation. We caution over-interpretation of these marginal findings, but the

new ideas they raise merit additional testing.

Experiences of novelty compared to familiarity reduce neural reinstatement across high-

level ventral stream regions

To more directly test our hypothesis that hippocampal pattern completion would be

reduced following experiences of novelty compared to familiarity, we used encoding-retrieval

similarity (ERS(48)) to measure memory reinstatement. While more invasive methods

convincingly demonstrate pattern completion in the hippocampus (14), this is inconsistently

observed with fMRI (49–52). Fortunately, the cortical reinstatement of memories, which depends

on the hippocampus (16, 17), can be robustly measured with fMRI. Accordingly, in addition to

the hippocampus, we analyzed probe trial ERS in high-level ventral stream and frontoparietal

ROIs that display reliable signatures of memory reinstatement (*a priori:* hippocampus, medial

temporal lobe (MTL) cortex, fusiform cortex (fusiform), medial parietal cortex; *post-hoc:* dorsal

& lateral prefrontal cortex (d&lPFC), ventromedial prefrontal cortex (vmPFC), and lateral

parietal cortex(52–57)), referred to as *reinstatement ROIs* (SI Appendix, Table S3). We

calculated ERS in three ways, beginning with the most inclusive definition (*broad ERS*). This

calculation captures reinstatement of information about the category of the to-be-retrieved item

(Scene or Face, necessary for the judgement) and information about the trial-specific association

(e.g. Ryan Gosling + tissue). Broad ERS values above zero suggest that a region reinstated traces

present during encoding at retrieval (see Fig. 2a for schematic and *Methods: fMRI data analysis*

for calculation specifics).

Medial parietal, MTL, and fusiform cortices all demonstrated above-chance levels of

broad ERS (MTLcortex: b = 0.005, SE = 0.002, p = 0.006, CI = [0.002, 0.009]; fusiform: b =

0.008, SE = 0.003, p = 0.006, 95% CI = [0.002, 0.014]; mPar: b = 0.011, SE = 0.002, p < 0.001,

95% CI = [0.006, 0.015]). Unsurprisingly, the hippocampus did not show evidence of

reinstatement (b < 0.001, SE = 0.003, p = 0.956, 95% CI = [-0.005, 0.005]; Fig. 2c; SI Appendix,

Table S4). Importantly, preceding novelty influenced ERS much like it did memory accuracy.

ROIs across the ventral processing stream all demonstrated significantly lower levels of broad

ERS following experiences of novelty compared to familiarity (Hipp: b = 0.005, SE = 0.002, p =

0.027, 95% CI = [0.001, 0.010]; MTLcortex: b = 0.006, SE = 0.002, p = 0.005, 95% CI = [0.002,

0.010]; fusiform: b = 0.009, SE = 0.003, p = 0.003, 95% CI = [0.003, 0.015]; mPar: b = 0.002,

SE = 0.003, p = 0.538, 95% CI = [-0.004, 0.007]; Fig. 2c; SI Appendix, Table S5). The influence

of the preceding experience was so pronounced that broad ERS was only observed following

familiarity in the MTL and fusiform cortices, not following novelty (MTLcortex following

familiar: estimated marginal mean (EMM) = 0.014, p < 0.001; MTL cortex following novel:

EMM = 0.002, p = 0.358; fusiform following familiar: EMM = 0.021, p < 0.001; fusiform

following novel: EMM = 0.002, p = 0.554; p-values FDR corrected for multiple comparisons;

Fig. 2d; SI Appendix, Table S6). These effects all held in control models accounting for response

time and accuracy (SI Appendix, Table S5 & S6), suggesting that the reinstatement effects are

not secondary to the observed behavioural biases. Analyses of frontoparietal *post-hoc* ROIs

demonstrated similar trends to those seen in the medial parietal cortex, with the d&lPFC and

lateral parietal cortex demonstrating significant broad ERS insensitive to the preceding trial

experience (Fig. 2c & 2d; SI Appendix, Table S4, S7 & S8). Taken together, these results suggest

that memory reinstatement is reduced across high-level ventral processing regions following

experiences of novelty compared to familiarity, but that reinstatement in frontoparietal regions is

insensitive to these preceding experiences.

Fig. 2 | Broad ERS. **a**, *A priori* and *post-hoc* reinstatement ROIs selected for ERS analyses. **b**,

Schematic of broad ERS calculation. Activity patterns from probe trials were correlated with

patterns from corresponding encoding trials (r_{same}). r_{same} was corrected by subtracting retrieval

trials' average correlation with encoding trials that included the unprobed image category (r_{diff};

See *Methods: fMRI data analysis*). **c**, Broad ERS on probe trials against chance. Points represent

intercepts and error bars represent their standard error from a linear mixed effects model (see

Methods: fMRI data analyses). **d**, Broad ERS on probe trials as a function of preceding memory

experiences. Points represent estimated marginal means and error bars represent their standard

error from a linear mixed effects model (see *Methods: fMRI data analyses*). Black stars

represent significance of the main effect of preceding memory experiences on broad ERS, while

dark and light blue stars represent significance of broad ERS against chance following

familiarity and novelty, respectively. * indicates $p < 0.05$, ** indicates $p < 0.01$, *** indicates $p <$

0.001. Post-hoc ROI p-values are FDR corrected for multiple comparisons.

Category & Trial-Specific Reinstatement

We designed our task to also evaluate whether ROIs reinstate information about the

category of the to-be-retrieved image (Scene or Face; *category ERS*) or information about the

trial-specific association (e.g. Ryan Gosling with tissue*; trial-specific ERS*). Of note, participants

only needed to reinstate categories to accurately perform the retrieval task (judgment of Scene

vs. Face). In fact, category ERS was higher for source-correct compared to incorrect trials in

both the medial parietal and MTL cortices (MTLcortex: $b = 4.715$, SE $= 2.130$, $p = 0.027$, 95%

CI $= [0.541, 8.890]$; mPar: $b = 4.144$, SE $= 1.758$, $p = 0.018$, 95% CI $= [0.698, 7.590]$). Among

both our *a priori* and *post-hoc* ROIs, only the fusiform cortex significantly reinstated category

information, and the medial parietal cortex demonstrated a marginal effect (fusiform: $b = 0.002$,

SE $= 0.001$, $p = 0.035$, 95% CI $= [0.000, 0.003]$; mPar: $b = 0.002$, SE $= 0.001$, $p = 0.050$, 95% CI

$= [<0.001, 0.004]$; Fig. 3c; SI Appendix, Table S9). As faces were one of the two reinstated

categories, it is unsurprising that the fusiform cortex demonstrated robust category ERS (58).

The fusiform cortex was also sensitive to preceding memory experiences, so much so that it only

showed significant category ERS following familiarity (fusiform: $b = 0.004$, SE $= 0.000$, $p <$

0.001, 95% CI $= [0.002, 0.005]$; fusiform following familiar: EMM $= 0.007$, $p < 0.001$; fusiform

following novel: EMM $= -0.001$, $p = 0.437$; Fig. 3d see SI Appendix, Table S10 & S11). On the

whole, the reinstatement of task-relevant category information was only modestly influenced by

preceding memory experiences.

We next used trial-specific ERS to identify regions where reinstatement extends beyond

task-relevant category information. Medial parietal and MTL cortices both showed significant

trial-specific ERS, while this metric was marginal in fusiform cortex (Hipp: $b < 0.001$, SE =

0.002, $p = 0.984$, 95% CI = [-0.005, 0.005]; MTLcortex: $b = 0.004$, SE = 0.002, $p = 0.028$, 95%

CI = [<0.001, 0.008]; fusiform: $b = 0.006$, SE = 0.003, $p = 0.054$, 95% CI = [0.000, 0.012];

mPar: $b = 0.008$, SE = 0.002, $p = 0.002$, 95% CI = [0.003, 0.012]; Fig. 3c, SI Appendix, Table

S14). Notably, preceding memory experiences influenced trial-specific ERS across all our

ventral stream ROIs (Hipp: $b = 0.005$, SE = 0.002, $p = 0.029$, 95% CI = [0.001, 0.010];

MTLcortex: $b = 0.006$, SE = 0.002, $p = 0.005$, 95% CI = [0.002, 0.010]; fusiform: $b = 0.008$, SE

= 0.003, $p = 0.011$, 95% CI = [0.002, 0.014]; mPar: $b = 0.003$, SE = 0.003, $p = 0.209$, 95% CI =

[-0.002, 0.008]; Fig. 3d, SI Appendix, Table S15), with above-chance reinstatement in MTL and

fusiform cortices only following familiarity (MTLcortex following familiar: EMM = 0.013, $p =$

0.001; MTLcortex following novel: EMM = 0.001, $p = 0.655$; fusiform following familiar: EMM

= 0.016, $p = 0.004$; fusiform following novel: EMM < 0.001, $p = 0.910$; p-values FDR corrected

for multiple comparisons; Fig. 3d; see SI Appendix, Table S15 & S16). As with broad ERS,

frontoparietal *post-hoc* ROIs once again displayed a similar pattern to that seen in the medial

parietal cortex, with the dorsal & lateral PFC and the lateral parietal cortex both demonstrating

above-chance levels of trial-specific ERS insensitive to preceding experiences (Fig. 3d; see SI

Appendix, Table S14, S17 & S18). Overall, preceding memory experiences had a more

pronounced impact on the reinstatement of trial-specific episodic details as compared to task-

relevant category information across high-level dorsal stream regions. The influence also closely

tracked that observed in broad, ERS suggesting that reinstatement of trial-specific largely

underpinned those effects.

These ERS analyses illuminate how preceding memory experiences prepare the brain to

retrieve. We primarily observed modulation of memory reactivation in the ventral stream and not

in frontoparietal regions. Additionally, recent memory experiences tended to modulate granular

trial-specific reinstatement, rather than more general and task-relevant category reinstatement.

These results are consistent with the hypothesis that hippocampal pattern completion may be

biased by recent novelty. As the ventral stream ROIs are robustly connected with the

hippocampus, their reinstatement patterns may more closely reflect hippocampal pattern

completion than frontoparietal regions. Perhaps this is why their trial-specific ERS, similar to the

hippocampus, was analogously modulated by the prior memory experience. These results are less

consistent with the retrieval mode account. Category information was most task-relevant in this

experiment, and if reinstatement was modulated by goal-maintenance, one would anticipate

modulation of category rather than trial information. Further, the retrieval mode is thought to be

instantiated by frontoparietal networks, which were insensitive to recent memory experiences.

While these results provide some support for our hypotheses, what remains unclear is whether

this memory reinstalment bias is mediated through basal forebrain responses.

Fig. 3 | Category and Trial-Specific ERS. **a**, Schematic of Category ERS calculation. Activity

patterns from probe trials were correlated with encoding trials that included the unprobed

associate from the same image category (r_{same}). r_{same} was corrected by subtracting retrieval trials'

average correlation with encoding trials that included the unprobed image category (r_{diff}; See

Methods: fMRI data analysis). **b**, Schematic of Trial-Specific ERS calculation. Activity patterns

from probe trials were correlated with patterns from corresponding encoding trials (r_{same}). r_{same}

was corrected by subtracting retrieval trials' average correlation with all other encoding that

included the same associate (r_{diff}; See *Methods: fMRI data analysis*). **c**, Category and Trial-

Specific ERS on probe trials against chance. Points represent intercepts and error bars represent

their standard error from a linear mixed effects model (see *Methods: fMRI data analyses*). **d**,

Category and Trial-Specific ERS on probe trials as a function of preceding memory experiences.

Points represent estimated marginal means and error bars represent their standard error from a

linear mixed effects model (see *Methods: fMRI data analyses*). Black stars represent

significance of the main effect of preceding memory experiences on broad ERS, while dark and

light blue stars represent significance of broad ERS against chance following familiarity and

novelty, respectively. ~ indicates $0.10 > p > 0.05$, * indicates $p < 0.05$, ** indicates $p < 0.01$, ***

indicates $p < 0.001$. Post-hoc ROI p-values are FDR corrected for multiple comparisons.

Neuromodulatory regions signal familiarity, rather than novelty, that indirectly & directly

predicts subsequent neural reinstatement of memories across the brain

We next turned to the neural mechanisms that incite these reinstatement and retrieval

biases by analyzing the preceding induction trials. Based on computational models (22, 59), we

hypothesized that novelty would evoke stronger basal forebrain activity than familiarity,

resulting in reduced reinstatement on successive probe trials. However, to remain open to other

mechanisms, we selected four nuclei that contain dense concentrations of neurons that project

neuromodulators throughout the brain (dopaminergic neurons in SN/VTA, noradrenergic neurons

in LC, and cholinergic neurons in hippocampally projecting CH123 and neocortically projecting

CH4(42, 60–62)). As induction trials consisted of object stimuli, we also analyzed activity in the

anterior hippocampus and perirhinal cortex, regions associated with contextual and object

novelty signalling (63, 64).

We predicted that each of these ROIs (*induction ROIs*) would respond more when

detecting novelty compared to familiarity. This, however, was far from the case. Counter to our

hypothesis, CH123 was insensitive to the manipulation (CH123: $b = -0.008$, SE $= 0.020$, p $=$

0.698, 95% CI $= [-0.047, 0.031]$). The anterior hippocampus displayed the expected novelty

preference ($b = -0.085$, SE $= 0.024$, p $= 0.001$, 95% CI $= [-0.132, -0.038]$), though it was most

active during the implicit baseline. This resulted in negative beta estimates, which are common

when passive fixation separates memory trials (65). The perirhinal cortex and LC were

insensitive to the manipulation (PRC: $b = 0.028$, SE $= 0.022$, p $= 0.214$, 95% CI $= [-0.015,$

0.071]; LC: $b = 0.008$, SE $= 0.021$, p $= 0.711$, 95% CI $= [-0.033, 0.048]$). Most surprisingly, the

opposite was seen in SN/VTA and the CH4, which responded more when identifying familiarity

compared to novelty (SN/VTA: $b = 0.086$, SE $= 0.023$, p $= 0.001$, 95% CI $= [0.043, 0.130]$; CH4:

$b = 0.043$, SE $= 0.020$, p $= 0.040$, 95% CI $= [0.004, 0.083]$; Fig 4b; SI Appendix, Table S19).

While unexpected from a novelty signalling perspective, a related literature might explain

the observed SN/VTA and CH4 familiarity preferences. SN/VTA and CH4 responses have

respectively been linked to motivational salience (66, 67) and shifting attentional focus towards

motivationally-relevant information (68–70). This begs the question; what goal orientation do

participants adopt during retrieval tasks? It has been argued that people see identifying "old"

trials as their goal and are less focused on correctly rejecting "new" trials (71, 72) in recognition

tasks. Notably, in these tasks, SN/VTA and the highly interconnected striatum preferentially

respond to the most motivationally salient trials, regardless of novelty (73, 74). In addition to

being the implicit targets in our task, old trials also required more effort than new trials because

participants were tasked with recalling associates. Accordingly, elevated SN/VTA and CH4

responses to familiarity compared to novelty could reflect participants' feeling of goal attainment

and attentional recruitment for effortful memory search. Regardless of the cause, these nuclei's

preferential responses to familiar trials position them to mediate the impact of recent familiarity

on successive memory reinstatement.

To identify potential mediators, we focus on ROIs sensitive to the induction trial

manipulation: CH4, anterior hippocampus, and SN/VTA. We asked whether any of these regions

predicted subsequent ERS in the ventral stream ROIs that were biased by preceding familiarity.

Here, we focus on trial-specific ERS, which appeared to underpin broad ERS effects. We

replicate these analyses with broad ERS in the supplement (see SI Appendix, Table S20 & S21).

CH4 responses on induction trials did not predict successive ERS, but responses in the two other

ROIs did. MTL cortex ERS tended to be weaker following induction trials with high anterior

hippocampus activity (b = -0.004, SE = 0.002, $p < 0.05$, 95% CI [-0.008, 0.000]) and stronger

following induction trials with high SN/VTA activity (b = 0.006, SE = 0.002, p = 0.001, 95% CI

= [0.002, 0.010]). Moreover, SN/VTA univariate activity was also marginally predictive of trial-

specific ERS in the hippocampus (b = 0.005, SE = 0.002, p = 0.058, 95% CI = [<0.001, 0.009];

SI Appendix, Table S20). We also explored reinstatement ROIs that did not demonstrate a

preceding familiarity bias, and found that SN/VTA activity predicted trial-specific ERS in the

d&lPFC (b = 0.007, SE = 0.002, p = 0.023, 95% CI = [0.002, 0.012]; SI Appendix, Table S21).

These results reveal that the SN/VTA may play an important role in preparing ventral and

frontoparietal regions to reactivate the details of individual experiences.

338   We next asked whether anterior hippocampus and SN/VTA univariate activity also

mediated the effect of preceding familiarity on memory reinstatement. Here, we again focus on

trial-specific ERS (SI Appendix, Table S22 for broad ERS mediation findings). The mediation

analyses did not yield significant findings for two of the three pairs: anterior hippocampus

activity did not significantly mediate preceding familiarity's effect on MTL cortex ERS (indirect

effect (ab) b = 0.003, p = 0.354, 95% CI = [-0.002, 0.010]; total effect (c) b = 0.066, p = 0.020,

95% CI = [0.009, 0.120]; direct effect (c') b = 0.063 p = 0.026, 95% CI = [0.008, 0.110]), nor did

SN/VTA activity significantly mediate preceding familiarity's effect on hippocampal ERS

(indirect effect (ab) b = 0.004, p = 0.154, 95% CI = [-0.001, 0.010]; total effect (c) b = 0.051, p =

0.078, 95% CI = [-0.007, 0.011]; direct effect (c') b = 0.047, p = 0.100, 95% CI = [-0.010,

0.110]). However, SN/VTA univariate activity partially mediated preceding familiarity's effect

on MTL cortex ERS (indirect effect (ab) b = 0.004, p = 0.038, 95% CI = [0.000, 0.010]; total

effect (c) b = 0.067, p = 0.020, 95% CI = [0.013, 0.120]; direct effect (c') b = 0.063, p = 0.034,

95% CI = [0.009, 0.120]; Fig 4c; SI Appendix, Table S23). This finding suggests that preceding

SN/VTA activity not only predicts upcoming memory reinstatement, it also partially accounts for

the power that recent familiarity exerts over memory reactivation.

Fig. 4 | Induction ROI Univariate Activity & ERS Relationships. a, Induction ROIs. **b,**

Univariate activity on induction trials separated by memory judgments. Points represent mean

beta estimates across participants for trials given familiar and novel judgements. Error bars

represent the standard error of the mean, applying the Morey (2008) correction for within-

participant comparisons. **c,** Analyses testing whether univariate activity in induction trial ROIs

mediate the impact of recent memory experiences on reinstatement ROI ERS. Models with

circled beta values indicate significant mediation (See *Methods: fMRI data analyses*). ~ indicates

$0.10 > p > 0.05$, * indicates $p < 0.05$, ** indicates $p < 0.01$, *** indicates $p < 0.001$.

One lingering question is whether the SN/VTA is particularly involved in preparing

ventral and frontoparietal regions to reinstate memories, or whether concurrent interactions may

also be at play. Accordingly, we conducted a final set of exploratory analyses investigating

whether univariate activity during probe trials related to simultaneous reinstatement. None of our

induction ROIs demonstrated significant preferences for experiences of novelty or familiarity (SI

Appendix, Table S24), perhaps because of design differences between induction and probe trials

(see Discussion). Critically, activity in induction ROIs did not significantly relate to concurrent

broad or trial-specific ERS after correcting for multiple comparisons (see SI Appendix, Table

S25 & S26). Of note, however, the negative relation between SN/VTA responses and

simultaneous hippocampal trial-specific ERS was significant before this correction (SI

Appendix, Table S26), in stark contrast to the positive predictive relationship reported above. On

the whole, the null concurrent relations provide important context, highlighting how SN/VTA

specifically prepares the brain for memory reactivation, rather than facilitating it in the moment.

Discussion

Here, we explored the cognitive and neural mechanisms that contribute to the variability of memory retrieval. First, we replicated prior reports that recent experiences of novelty compared to familiarity reduce people's upcoming ability to recall associations(47). Then, we asked how novelty influenced the fidelity of upcoming memory reinstatement in the brain. When associative memory was tested following novelty compared to familiarity, patterns of activity in high-level ventral stream regions less closely resembled patterns evoked during corresponding encoding trials. In fact, these regions did not display evidence of memory reinstatement following novelty. Notably, recent novelty tended to modulate granular trial-specific reinstatement, rather than more general and task-relevant category reinstatement. Lastly, we explored whether neuromodulatory nuclei activity on the preceding trial set the stage for upcoming memory reinstatement. Counter to our predictions, no neuromodulatory nuclei of interest signaled novelty. Rather, the SN/VTA and CH4 both preferentially responded to familiar trials. Moreover, SN/VTA responses significantly predicted upcoming reinstatement across frontal and ventral stream ROIs. Most notably, SN/VTA activity partially mediated the effect of recent memory experiences on MTL cortex reinstatement, uncovering a mechanism through which dopamine may prepare the brain to remember.

Our findings revealed how preceding memory experiences powerfully modulate neural memory reinstatement. We used ERS analyses to characterize reinstatement in regions previously shown to reinstate category and/or trial-specific information during retrieval (52–57). Reinstatement in all high-level ventral stream ROIs was biased by preceding memory experiences. The MTL and fusiform cortices both reliably reinstated memories only following familiarity, and while hippocampal reinstatement was stronger following familiarity compared to

novelty, it did not reliably exceed chance levels. FMRI signatures of hippocampal reinstatement

are inconsistently observed (49–52), despite this region being the source of associative retrieval

(75). Some studies suggest that this inconsistency is rooted in the hippocampus only reinstating

trial-specific details (50, 52, 76, 77), leaving designs targeting the reinstatement of

decontextualized content insensitive to the episodic representations contained in the

hippocampus. Our findings somewhat align with this proposal, in that only trial-specific

hippocampal ERS was influenced by our novelty manipulation. Our findings also provide new

insights into the inconsistency of hippocampal reinstatement. Two of the first studies to find

reliable hippocampal reinstatement (78, 79) both only included previously studied items in their

retrieval tests. Our findings suggest that omitting novel foils may have amplified hippocampal

reinstatement, as each trial in these experiments was experienced following a series of familiar

stimuli. More broadly, our findings highlight how neural reinstatement of memories can be

surprisingly state dependent, with important implications for the interpretation of previous data,

the design of future experimental paradigms, and the understanding of preparatory memory

states.

Motivated by computational models (44, 80, 81), we hypothesized that preceding

novelty-evoked cholinergic nuclei activity would mediate novelty's impact on memory

reinstatement. Anchored in rodent physiology, the models propose that increases in cholinergic

innervation of the hippocampus following novelty induce a lingering bias away from pattern

completion and towards pattern separation (30, 44, 80–82). These predictions have been

indirectly supported by prior behavioural research: people are better at forming memories (83)

and making fine-grained memory discriminations (84) in the seconds following novelty,

consistent with a pattern separation bias. Conversely, people are better at retrieving (47) and

using (83) associative memories in the seconds following familiarity, consistent with a pattern

completion bias. Our initial results more directly support prediction: pattern completion-

dependent signatures of memory reinstatement were reduced following novelty compared to

familiarity in the hippocampus and interconnected ventral stream regions.

However, we failed to observe comprehensive evidence for this mechanism, as

cholinergic nuclei did not mediate reinstatement biases. The hippocampally projecting CH123

ROI did not preferentially respond to novel induction trials, and its activity did not predict

hippocampal ERS on the following trial ($b = -0.002$, $SE = 0.002$, $p = 0.380$, 95% CI = [-0.006,

0.002]). Several methodological limitations, though, caution against an overinterpretation of

these null effects. First, CH3—a region that does not strongly project to the hippocampus—was

combined with hippocampally projecting CH1 and CH2 regions in the available probabilistic

ROI (62). Second, the compositions of these nuclei are heterogeneous, with cholinergic neurons

outnumbered by functionally distinct GABA-ergic and glutaminergic neurons (85). Accordingly,

while BOLD responses in CH123 should correlate with hippocampal cholinergic innervation, the

relationship is not direct and will not be very strong. By contrast, CH4—the largest basal

forebrain nucleus—is composed of approximately 90% cholinergic neurons (86). BOLD activity

in this region was sensitive to our task manipulations, showing similar preferences to the

SN/VTA, adding to a growing literature that CH4 can be a fruitful target for MRI imaging (87–

89). Taken together, while our imaging results did not support the hypothesis that dips in basal

forebrain activity prepare the hippocampus to retrieve memories, they also do not undermine

decades of relevant research using invasive methods in animals (90–94).

Our imaging results did, however, uncover that DA may play a role in preparing the brain

to retrieve. Animal research has convincingly demonstrated that DA is critical for the formation

of memories in the hippocampus; pharmacological and optogenetic activation of hippocampal

dopamine receptors around the time of memory formation enhances later expression of these

memories (95–99). By contrast, similar manipulations applied around the time of retrieval do not

influence memory performance (98), leading to the general consensus that DA is important for

the formation, but not retrieval of hippocampal memories (100). However, studies that measure

or manipulate DA more systemically—rather than targeting the hippocampus—have yielded

mixed results. In both humans and animals, retrieval performance has been positively linked to

pharmacological manipulations and SN/VTA activity in some studies (39, 101–105), while others

found no relation (95, 100, 106–109). These findings suggest that DA may facilitate retrieval

under some circumstances via extra-hippocampal targets. Our results further inform this mixed

literature. We observed that SN/VTA responses positively predict memory reinstatement on

successive, but not concurrent, trials. While pharmacological manipulations can more directly

assess DA, they lack the temporal specificity to delineate these complex relations across time. By

providing temporally precise evidence linking SN/VTA activity to retrieval dynamics, our results

refine existing models of DA's role in memory and suggest future avenues for investigation using

invasive and non-invasive techniques.

Elements of how the SN/VTA prepares the brain to remember may align with retrieval

mode mechanisms. Firstly, this region was more responsive on trials that strongly recruit the

retrieval mode, preferentially responding to familiar induction trials. These trials likely elicited

strong retrieval goal-maintenance, as participants commonly see detecting familiarity as their

primary job in recognition tasks (71, 72). Further, familiar trials also required strong attentional

and control orientations toward retrieved content, as participants needed to recall associations.

This preferential responsivity, however, was not seen on probe trials. Multiple experimental

design factors might explain why probe trials were less sensitive than induction trials, including

stimulus class differences, shorter anticipatory delays blunting DA neuron ramping (110, 111),

and the predominance of old trials. The latter point could have both reduced statistical power and

made the rare new trials more salient. Secondly, unlike the novelty/familiarity manipulation,

SN/VTA activity predicted upcoming reinstatement in both high-level ventral and frontal

regions. In particular, SN/VTA activity predicted reinstatement in the dorsal and lateral portions

of the PFC. These regions are heavily involved in executive functioning, particularly attending to

goal-relevant information and shifting between internally and externally oriented attention (112,

113), and have been implicated in the retrieval mode (114). SN/VTA's preferential signaling to

motivationally relevant familiar experiences, as well as its relationship with upcoming

reinstatement in both ventral and frontal regions, suggest that it may play a role as a catalyst of

the retrieval mode.

Pulling these potential mechanisms together, our results raise the possibility that

variability in memory retrieval may be best understood by considering interactions between

pattern completion and retrieval mode biases. At any given moment, ongoing hippocampal states

may make it easier or harder for the region to reinstate specific episodic memory traces. For

example, the details of specific memories were more fluently reactivated across connected

ventral stream regions when in a state triggered by recent familiar trials. But reinstatement also

depends on recent SN/VTA engagement. These dopaminergic nuclei may directly facilitate

hippocampal pattern completion, though there is little physiological evidence for this mechanism

(100). Alternatively, or additionally, SN/VTA engagement could support goal maintenance and

appropriate attentional orientations. In this way, SN/VTA activity, and the retrieval mode it

potentially fosters, could boost retrieval-relevant input to the hippocampus as well as its pattern

completed output. Future research designed to measure the retrieval mode could explore this

possibility in more detail.

In sum, our findings reveal that the fidelity of memory reinstatement is dynamically

shaped by recent experiences and neuromodulatory states, providing new insights into the long

standing puzzle why our memory retrieval abilities are so variable across time. We show that

preceding familiarity, compared to novelty, enhances neural reinstatement across high-level

ventral stream regions. While we found limited evidence for cholinergic modulation of

reinstatement, our results highlight a potential role for dopamine in fostering a retrieval mode

that facilitates memory reactivation. These findings advance our understanding of the state-

dependent nature of memory retrieval and point to an emerging framework in which

hippocampal pattern completion and retrieval goal maintenance are jointly modulated by recent

experience and neuromodulatory tone. We hope that our work will inspire new research into this

dynamic account of episodic memory retrieval, probing how this foundational ability is deeply

shaped by ongoing mental and neural states.

Resource Availability

All post-processed data and code used to run analyses can be found at https://

doi.org/10.17605/OSF.IO/UY9ZV.

Acknowledgments and Funding Sources

We gratefully acknowledge Justine Vorvis and Hannah Cho for their assistance with data

collection, as well as all participants for their time, attention, and willingness to be a part of the

research process. This work was supported by the Natural Sciences and Engineering Research

Council of Canada (NSERC) Discovery Grant (RGPIN-2023-04231 to Katherine D. Duncan);

Canada Research Chair Program (CRC-2023-00363 to Katherine D. Duncan); Ontario Research

and Innovation (ER18-14-139 to Katherine D. Duncan); Canada Foundation for Innovation

JELF & Ontario Research Fund (34479 to Katherine D. Duncan); and the Ontario Graduate

Scholarship (to Anuya Patil).

Author Contributions

MD led formal analysis, visualization, data curation, writing- draft. and writing- review &

editing. AP led investigation, methodology, and project administration. AP & MD contributed

equally to software. KDD led supervision, funding acquisition, and resources. AP & KDD led

conceptualization. MD, AP, and KDD contributed equally to validation.

Declaration of Interests

The authors declare no competing interests.

Materials & Methods

Participants

Forty-four adults were recruited via convenience sampling from the University of

Toronto community through online advertisements and flyers. Two participants withdrew from

the study before completing all experiment tasks. Data from 13 participants were excluded from

analyses due to previously participating in the study ($n = 1$), showing signs of drowsiness ($n = 3$),

failing to follow instructions to stay still inside the scanner ($n = 1$), moving excessively during

the study (framewise displacement > 0.3 mm for more than 10% of the volumes on the retrieval

and word encoding tasks and 20% of the volumes on the localizer task; $n = 1$), vision issues ($n =$

1), or lower than chance performance on the associative memory task ($n = 6$). Participants were

categorized as having lower than chance performance if a binomial test demonstrated that

significantly over half of associative judgments were source errors. The final analyses included

29 participants (21 self-identified females, 7 self-identified males, 1 declined to answer, mean

age = 22.9 years [19, 30], mean years of education = 16.43 [13, 22]). Participants identified their

race and ethnicity as follows: Japanese/Korean/Chinese (10), Indian/Pakistani/Sri Lankan (8),

White (5), Bangladeshi (1), Vietnamese (1), Mixed (Black & White 1, White & Chinese 1),

Euroasian (1), and Declined to Answer (1). Participants were right-handed, fluent in English, had

normal or corrected-to-normal vision, and had no history of psychiatric, neurological, or learning

disorders. Demographic information regarding gender, socioeconomic status, and ancestry were

not collected. Convenience sampling resulted in uneven groupings of participants by

race/ethnicity and self-identified sex, and as such, we were not sufficiently statistically powered

to analyze whether these variables influenced our study's main results. The racial and ethnic

heterogeneity of the University of Toronto community resulted in a diverse sample, increasing

generalizability across race and ethnicity. However, the skew of self-identified sex towards

female may limit sex-based generalizability. Participants gave written informed consent and were

compensated $20-25 CAD/hour for their participation. All experimental procedures were

approved by the University of Toronto Research Ethics Board (REB).

Stimuli

Stimuli consisted of colored images of 144 common objects (115), two faces of famous

individuals (Justin Trudeau, Ryan Gosling, derived from Google Search), two famous Canadian

landmarks (CN Tower, Niagara Falls, derived from Google Search), 192 object nouns, and two

adjectives ("Plain" and "Safe"). The set of object nouns were selected to not overlap with the

images of common objects. For example, if an image of a violin was a part of the 144 object

images, the word "violin" was not included in the set of 192 object nouns. Stimuli were

presented using PsychoPy (116) on an MR-compatible BOLDscreen monitor (32", 1920 x 1080)

placed behind the scanner. Participants viewed the stimuli through a system of mirrors installed

on the head coil. Another image set of objects, faces, places, and scrambled objects were

presented in a functional localizer with Matlab & PsychToolbox, but this data is not reported

here (see Localizer Tasks).

Procedure

Patil & Duncan (2018) demonstrated that the effect of recent novelty on associative

memory declines over seconds. Studying this effect necessitates the use of short intertrial

intervals (ITIs) during retrieval. However, consistently short ITIs pose challenges for statistically

disambiguating responses to consecutive trials due to the slow BOLD response (117). To resolve

this issue, we (a) arranged trials into pairs of novelty/familiarity induction trials and successive

probe trials, on which we measured neural reinstatement, (b) fixed the ITIs within trial pairs to 1

second, and (c) orthogonalized content across these two kinds of trials. Specifically, we used

images of objects to induce novelty/familiarity. During probe trials, we tested how well

participants retrieved a face or a scene when probed with a previously associated word. This

alternating pattern, where each pair of trials consisted of independent content, allowed us to

estimate how preceding memory experiences influence successive neural reinstatement (see Fig.

1a). To increase the sensitivity of our reinstatement analyses, we included 5 second fixation

periods between retrieval pairs.

The experiment consisted of multiple blocks of encoding, retrieval, and localizer tasks.

All tasks were completed inside the MRI scanner. Participants were given instructions and

practiced all tasks outside of the scanner until they were comfortable with each task. Participants

were aware that their memory for stimuli would be tested before they started the experiment.

Encoding Blocks

Participants completed two encoding tasks, corresponding to the induction and probe

retrieval trials: object encoding and word encoding, respectively. Their order was

counterbalanced across participants.

Object Encoding

Participants were shown a series of object images (72 randomly selected from a pool of

144) paired with one of two repeating words, "Plain" or "Safe," and asked to imagine a scenario

in which the word could describe the image. This task was chosen to encourage deep encoding of

the object-word pairs. The presentation order of objects was intermixed such that the repeating

word associate would not repeat across more than three consecutive trials, and the pairing of

each object image with "Plain" or "Safe" was randomized for each participant. Each pair was

shown for 2.5 seconds. Participants were then given 1.5 seconds to rate the vividness of the

association on a 4-point scale of "very vivid" to "not vivid at all," followed by a 1-second

fixation cross. Participants completed one 6.2-minute block of the object encoding task. The

memories formed during object encoding were always used to induce novelty/familiarity during

retrieval. fMRI data from the object encoding blocks were not analyzed.

Word Encoding

Participants were shown a series of words (96 object-nouns randomly selected from a

pool of 192) paired with one of four repeating images of famous landmarks (CN Tower, Niagara

Falls) or celebrities (Ryan Gosling, Justin Trudeau). Each image was presented twenty-four times

across the two encoding blocks, and the presentation order of images was intermixed such that

the image category would not repeat across more than three consecutive trials. The pairing of

each word with the repeating images was randomized for each participant. The word was

presented on the left side of the screen for 1.5 seconds, followed by one of the four repeating

images on the right. The pair together was shown for 2.5 seconds, resulting in 4 second trials.

For words paired with a scene, participants were asked to imagine themselves in that scene using

that object from a first-person perspective. For words paired with a face, participants were asked

to imagine the celebrity using that object against a white background. Participants were then

given 1.5 seconds to provide their vividness rating, followed by a 4.5 second fixation cross.

Participants completed two 8.2-minute blocks of the word encoding task. Encoding trials in each

block were evenly split between scenes and faces. The memories formed during word encoding

were used to measure reinstatement during retrieval, and as such, we used a longer ITI than in

the object encoding task to facilitate estimates of single-trial fMRI responses.

Retrieval Session

Participants completed an associative retrieval session with alternating induction (72 old

and 72 new objects) and probe (48 old face-associated, 48 old scene-associated, and 48 new

words) trials. Note that we include twice as many old probe trials compared to new because

reinstatement can only be measured on old trials. New stimuli were also randomly selected from

the same stimulus pools for each participant, excluding the encoded, old stimuli. Trial order was

pseudorandomized such that an equal number of "old" and "new" induction trials preceded "old"

probe trials, with no more than three consecutive "old" or "new" trials. During each retrieval

trial, an object or word was shown on the screen for 2 seconds. Participants were asked to

indicate if the object/word was new (by pressing the thumb key) or old, in which case, they were

asked to recall the paired associate. For the object trials, the associate responses were Plain

(middle finger key) or Safe (ring finger key). For the word trials, the associate responses were

Scene (middle finger key) or Face (ring finger key). Participants were asked to press the index

finger key if they recognized the object/word but were unable to recall its associate (labelled

"old?"). Response options were presented at the bottom of the screen in an intuitive order to help

participants remember the mappings. A 1-second and 5-second fixation cross followed each

induction and probe trial, respectively. Participants completed four 6.2-minute retrieval blocks.

Localizer tasks

After the retrieval blocks, participants completed two functional localizer tasks. Both are

beyond the scope of the current project, but we analyzed behavioural data from one condition of

the second localizer. In this retrieval mode localizer, participants made one of three judgments in

response to a presented word: whether the real-world cost of the object was expensive or cheap

(semantic), whether the object word was previously shown in the preceding retrieval session or

novel (recognition), and whether the font color changed or remained the same (sensory). Across

tasks, half of the words were old (i.e., first seen as novel foils during retrieval probe trials) and

half were new (i.e., the remaining 48 words from the pool of 192). Words were presented for a

maximum of 2 seconds. At the end of the 2 seconds, participants saw a fixation cross for 3

seconds before beginning the next trial. We analyzed the 32 recognition judgments (16 old and

16 new words) made across two blocks.

Behavioural data analysis

All analyses were conducted in R (version 4.0.2). Linear mixed effects models were used

to reflect the hierarchical nature of our data in which multiple trials are clustered within

participants. These models were run with unstructured covariance matrices (lmer function in

lme4 package, version 1.1-23(118) with uncorrelated random intercepts and slopes for each

within-participant fixed effect varying across participants. We estimated the marginal means of

predictors to evaluate their simple effects (*emmeans* package, version 1.11.0(119)).

We categorized induction trials according to participants' subjective memory judgements.

If they responded with "new," trials were considered inductions of novelty; if they recognized

the objects, regardless of their associative retrieval success, trials were considered inductions of

familiarity. We also divided probe trials to focus on our pattern completion hypothesis.

Specifically, we obtained independent measures of associative memory (pattern completion

dependent) and item-only recognition memory (comparatively pattern completion independent).

To do this, we calculated associative accuracy only on probe trials where participants correctly

recognized words as "old," and item-only recognition memory on trials where participants failed

to recall the correct association.

We ran a binomial linear mixed effect model to examine the influence of preceding

experiences (novelty vs. familiarity on induction trials) on successive probe trial associative

memory accuracy. Associative accuracy was dummy coded in these models, where correct

associative judgments were coded as 1 and incorrect associative and "old?" judgments were

coded as 0. For these and all following models, we effect coded the preceding judgment predictor

(*preceding judgement*: novelty as -1; familiarity as 1). We added a covariate to correct for

potential response priming, indicating whether the correct associative response on the probe trial

was the same as the participants' response on the preceding trial (*response bias*: dummy coded,

unaffected probe trials as 0; affected probe trials as 1). In addition, we ran a separate model to

control for the effects of current trial stimulus class ("Scene" or "Face", *stimulus class*) and the

preceding trial response time (*preceding response time*). Stimulus class was effect coded

("Scene" as -1; "Face" as 1). Preceding response time was z-scored across all induction trials

within each participant. We used a paired sample t-test to assess the impact of preceding

judgement on item-only recognition accuracy. We calculated d' (z(hit rate) – z(false alarm rate))

for each participant separately for probe trials preceded by novelty and familiarity.

We ran a linear mixed effect model to examine the influence of preceding judgment on

accurate probe trial response time. We used an inverse gaussian log linking function to account

for the shape of the response time distribution. In addition to preceding judgement, we included

four covariates: Response bias, stimulus class, retrieval block number (*centered block number*)

and trial number (*scaled trial number;* z-scored within block*)*. The latter two were included to

control for task fatigue/learning.

As previously stated, recognition for a subset of probe trial novel foils was tested in a

retrieval mode localizer. We ran a binomial linear mixed effect model to determine if judgments

made on preceding induction trials affected participants' learning of these words. "Old" trial

recognition accuracy during the retrieval mode localizer was dummy coded (misses as 0; hits as

1). Preceding judgment and response bias were included as predictors.

fMRI data acquisition

Whole-brain imaging data were acquired on a Siemens Prisma 3T MRI scanner at the

Toronto Neuroimaging Facility (ToNI) of the University of Toronto using a 32-channel head coil.

Functional images were collected using a T2*-weighted multi-band accelerated EPI sequence

(TR = 1.5 s, TE = 26 ms, flip angle = 70º, GRAPA parallel imaging factor = 2, multiband

acceleration factor = 2, distance factor = 20%, field of view = 220 mm, matrix size = 88 x 88, 54

slices oriented parallel to the anterior commissure-posterior commissure line, voxel size = 2.5 x

2.5 x 2.5 mm). One field map was collected (TR = 888 ms, TE 1 = 4.92 ms, TE 2 = 7.38 ms, flip

angle = 60º, voxel size = 2.5 x 2.5 x 2.5 mm) to correct EPI images for susceptibility-induced

distortions. In addition, one anatomical T1-weighted 3D magnetization-prepared rapid gradient

echo (MPRAGE) image (TR = 2 s, TE = 2.67 ms, flip angle = 9º, field of view = 256 mm, matrix

= 256 x 256, voxel size = 1 x 1 x 1 mm) was also collected.

Neuroimaging data processing

Preprocessing was performed using fMRIPrep 1.2.5 (120, 121), which is based

on Nipype 1.1.6 (122, 123).

Anatomical data preprocessing

The T1-weighted (T1w) image was corrected for intensity non-uniformity (INU)

using N4BiasFieldCorrection (ANTs 2.2.0(124)) and used as T1w-reference throughout the

workflow. The T1w-reference was then skull-stripped using antsBrainExtraction.sh (ANTs

2.2.0), using OASIS as the target template. Spatial normalization to the ICBM 152 Nonlinear

Asymmetrical template version 2009c (125, 126) was performed through nonlinear registration

with antsRegistration (ANTs 2.2.0(127)), using brain-extracted versions of both T1w volume and

template. Brain tissue segmentation of cerebrospinal fluid (CSF), white-matter (WM) and gray-

matter (GM) was performed on the brain-extracted T1w using FAST (FSL 5.0.9(128, 129)).

Functional data preprocessing

The first 4 volumes (6 s) of each of the task BOLD scans were removed to allow for

scanner magnet stabilization. For each of the BOLD runs per participant (across all tasks), the

following preprocessing was performed. First, a reference volume and its skull-stripped version

were generated using a custom methodology of fMRIPrep. A deformation field to correct for

susceptibility distortions was estimated based on a field map that was co-registered to the BOLD

reference, using a custom workflow of fMRIPrep derived from D.

Greve's epidewarp.fsl script (https://www.nmr.mgh.harvard.edu/~greve/fbirn/b0/epidewarp.fsl)

and further improvements of HCP Pipelines (130). Based on the estimated susceptibility-induced

distortion, an unwarped BOLD reference was calculated for a more accurate co-registration with

the anatomical reference. The BOLD reference was then co-registered to the T1w reference

using FLIRT (FSL 5.0.9(131)) with the boundary-based registration (132) cost-function. Co-

registration was configured with nine degrees of freedom to account for distortions remaining in

the BOLD reference. Head-motion parameters with respect to the BOLD reference

(transformation matrices, and six corresponding rotation and translation parameters) were

estimated before any spatial transformations using MCFLIRT (FSL 5.0.9(133)). The BOLD

time-series were resampled onto their original, native space by applying a single, composite

transform to correct for head-motion and susceptibility distortions.

The BOLD time-series were resampled to MNI152NLin2009cAsym standard space,

generating a preprocessed BOLD run in MNI152NLin2009cAsym space. All resamplings were

performed with a single interpolation step by composing all the pertinent transformations

(i.e. head-motion transform matrices, susceptibility distortion correction, and co-registrations to

anatomical and template spaces). Gridded (volumetric) resamplings were performed using

antsApplyTransforms (ANTs (134)), configured with Lanczos interpolation to minimize the

smoothing effects of other kernels (135). Lastly, functional images were further spatially

smoothed using a Gaussian kernel of 2mm full-width at half-maximum in FSL.

Several confounding time-series were calculated based on the preprocessed BOLD.

Framewise displacement (FD) and DVARS were calculated for each functional run, both using

their implementations in Nipype(136). Additionally, a set of physiological regressors were

extracted to allow for component-based noise correction (CompCor(137)). Principal components

were estimated after high-pass filtering the preprocessed BOLD time-series (using a discrete

cosine filter with 128s cut-off) for anatomical CompCor (aCompCor). Firstly, a subcortical mask

was obtained by heavily eroding the brain mask, which ensured it did not include cortical grey

matter (GM) regions. Six aCompCor components were then calculated within the intersection of

the aforementioned mask and the union of cerebrospinal fluid (CSF) and white matter (WM)

masks calculated in T1w space, after their projection to the native space of each functional run

(using the inverse BOLD-to-T1w transformation). The head-motion estimates calculated in the

correction step were also placed within the corresponding confounds file.

Many internal operations of fMRIPrep use Nilearn 0.5.0 (138), mostly within the

functional processing workflow. For more details of the pipeline, see the section corresponding

to workflows in fMRIPrep's documentation (https://fmriprep.org/en/1.2.5/workflows.html).

ROI selection

To measure reinstatement, we defined four *a priori* ROIs (Fig. 2a) that have previously

been shown to reinstate trial-specific and category information (52, 53, 55, 56): whole

hippocampus, medial temporal lobe cortex (MTLcortex), fusiform cortex (fusiform), and medial

parietal cortex (mPar). After conversations with colleagues, we added three additional

frontoparietal *post-hoc* ROIs (Fig. 2a) that have previously been shown to reinstate trial-specific

or category information (54, 55, 57): lateral parietal cortex (LPC), dorsal & lateral prefrontal

cortex (d&lPFC), and ventromedial prefrontal cortex (vmPFC). We refer to all seven of these

regions as *reinstatement ROIs*. To measure novelty/familiarity responses during the induction

trials, we evaluated six *a priori induction ROIs* (Fig. 4a): basal forebrain medial septum and the

diagonal band of Broca (CH123), basal forebrain nucleus basalis (CH4), substantia nigra and

ventral tegmental area (SN/VTA), locus coeruleus (LC), anterior hippocampus (aHipp), and

perirhinal cortex (PRC). The probabilistic atlases and thresholds used for each brain region are

listed in SI Appendix, Table S3. All masks were defined in T1 MNI space and resampled to the

functional MNI space.

Generating Single Trial Estimates

GLMs were constructed for each of the 96 word encoding and 288 retrieval trials to

estimate single-trial beta values. This was done using the LS-S method (139) implemented in

FSL with *Nipype* (version 1.8.5) and in custom Python (version 2.7) scripts in which separate

GLMs were conducted for each trial. These GLMs included a regressor for the trial of interest,

and responses to all other trials were modelled together with a second regressor. Each trial was

modelled at stimulus onset (for word encoding, the time when the word and the image were first

displayed together) using a 2.0-second (retrieval task) or a 2.5-second (word encoding task)

boxcar function convolved with a double-gamma hemodynamic response function. In addition to

these primary regressors of interest, each GLM also included the following nuisance regressors:

the temporal derivatives of six motion estimates and six aCompCor motion components.

Volumes for which the framewise displacement was greater than 0.3 mm were added as spike

regressors in the GLMs.

fMRI data analysis

We obtained three estimates of neural reinstatement that differed in their levels of

specificity. The first (broad ERS) estimated both category information of the trial (required to

make a retrieval judgment; "Scene" vs. "Face"), as well as specific information about the trial

itself (the specific scene/face and its association with the probed word). The second (category

ERS) estimated only the reinstatement of the trial category, while the third (trial-specific ERS)

estimated only the reinstatement of each trial's specific association. For each ERS measure, we

calculated two Pearson correlations and operationally defined ERS as the difference between

them ($r_{same} - r_{diff}$). For broad ERS, activity patterns from probe trials were correlated with

patterns from corresponding encoding trials (r_{same}) and corrected by subtracting retrieval trials'

average correlation with encoding trials that included the unprobed image category (e.g., scenes

if the retrieval trial probed a face; r_{diff}; Fig. 2b). For category ERS, activity patterns from probe

trials were correlated with encoding trials that included the unprobed associate from the same

image category (e.g. if the retrieval probe was studied with Ryan Gosling, all Justin Trudeau

encoding trials; r_{same}) and corrected by subtracting retrieval trials' average correlation with

encoding trials that included the unprobed image category (e.g., scenes if the retrieval trial

probed a face; r_{diff}; Fig. 3a). For trial-specific ERS, activity patterns from probe trials were

correlated with patterns from corresponding encoding trials (r_{same}) and corrected by subtracting

retrieval trials' average correlation with all other encoding that included the same associate (e.g.,

if tissue were the probe, all Ryan Gosling encoding trials except for Ryan Gosling + tissue; r_{diff};

Fig. 3b). For all ERS analyses, we Fisher z-transformed correlation coefficients and only

included probe trials on which participants correctly recognized words as "old."

We ran a linear mixed effect model for each ERS measure to examine whether regions

significantly reinstated learned information. We included response bias as a covariate in all ERS

analyses, as it significantly related to ERS in several reinstatement ROIs (SI Appendix, Table

S5).

We tested whether preceding memory experiences (novelty vs. familiarity) influenced

ERS on probe trials in each of the reinstatement ROIs using linear mixed effects models.

Preceding judgment and response bias were included as covariates. We ran an additional model

for each ROI to control for the effects of each trial's *accuracy* and *response time*. Accuracy was

effect coded (accurate trials as 1; inaccurate trials as -1), while response time was z-scored across

all probe trials for each participant.

We were interested in whether category ERS significantly related to participants'

associative accuracy, as category information was needed for correct judgments ("Scene" vs.

"Face"). We ran a generalized linear mixed effects model for each of our *a priori* reinstatement

ROIs predicting whether associative accuracy significantly related to category ERS. Each model

included Fisher z transformed category ERS, response bias, and preceding judgment as

covariates.

To evaluate whether induction ROIs preferentially responded to experiences of

familiarity or novelty, we ran linear mixed effects models predicting average univariate signal, z-

scored within participant. Each model included the current trial judgment (*judgment*) as a

predictor, effect coded with novelty as -1 and familiarity as 1. We ran additional follow up

analyses exploring this on probe trials using models with identical structure.

To determine whether induction trial univariate signal in our induction ROIs predicted

successive reinstatement, we ran linear mixed effect models predicting trial-by-trial ERS

estimates in each of our reinstatement ROIs. Each model included univariate signal on the

preceding induction trial in each of the induction ROIs, z-scored within participant, and response

bias as predictors. We ran additional follow up analyses using models with identical structure

evaluating the relationship between concurrent probe trial univariate activity in induction ROIs

and ERS, rather than successive relationships.

We then performed multilevel mediation analyses using *lme4* (version 1.1-23) and

mediation (version 4.5.0) packages in R (version 4.0.2) to assess if univariate signal in induction

ROIs (mediator) mediated the effects of preceding judgments (independent variable; IV) on

successive ERS (outcome). To establish mediation, there must be evidence for all pair-wise

relations. Only three pairs of ROIs met this criteria: SN/VTA induction trial activity as a

mediator for preceding judgement on hippocampal and MTL cortex ERS, as well as anterior

hippocampus induction trial activity as a mediator for preceding judgement on MTL cortex ERS

(ultimately, these models were run for both broad and trial-specific ERS estimates). The

following multilevel models were run as part of the mediation:

Model (1): Influence of preceding judgment on univariate activity in induction ROI

(mediator ~ IV)

Model (2): Influence of preceding judgment and univariate activity in all of the induction

ROIs on successive ERS (outcome ~ IV + mediator).

These models were run on data that included correctly recognized probe trials only. Response

bias was included as a covariate in Model (2). Both univariate activity and ERS estimates were

z-scored within participant for these models.

Works Cited

1. J. W. Alba, L. Hasher, Is memory schematic? *Psychol. Bull.* **93**, 203–231 (1983).

2. J. R. Anderson, L. M. Reder, "An Elaborative Processing Explanation of Depth of
Processing" in *Levels of Processing in Human Memory (PLE: Memory)*, (Psychology
Press, 1979).

3. S. Bertsch, B. J. Pesta, R. Wiscott, M. A. McDaniel, The generation effect: A meta-
analytic review. *Mem. Cognit.* **35**, 201–210 (2007).

4. F. I. M. Craik, R. S. Lockhart, Levels of processing: A framework for memory research.
J. Verbal Learn. Verbal Behav. **11**, 671–684 (1972).

5. N. J. Slamecka, P. Graf, The generation effect: Delineation of a phenomenon. *J. Exp.*
Psychol. [Hum. Learn.] **4**, 592–604 (1978).

6. C. D. Morris, J. D. Bransford, J. J. Franks, Levels of processing versus transfer
appropriate processing. *J. Verbal Learn. Verbal Behav.* **16**, 519–533 (1977).

7. E. Tulving, D. M. Thomson, Encoding specificity and retrieval processes in episodic
memory. *Psychol. Rev.* **80**, 352–373 (1973).

8. E. Butowska-Buczyńska, P. Kliś, K. Zawadzka, M. Hanczakowski, The role of variable
retrieval in effective learning. *Proc. Natl. Acad. Sci.* **121**, e2413511121 (2024).

9. E. Tulving, *Elements of Episodic Memory* (Oxford University Press, 1983).

10. N. M. Long, The intersection of the retrieval state and internal attention. *Nat.*
Commun. **14**, 3861 (2023).

11. K. P. Madore, A. D. Wagner, Readiness to remember: predicting variability in episodic
memory. *Trends Cogn. Sci.* **26**, 707–723 (2022).

12. H. Tarder-Stoll, M. Jayakumar, H. R. Dimsdale-Zucker, E. Günseli, M. Aly, Dynamic
internal states shape memory retrieval. *Neuropsychologia* **138**, 107328 (2020).

13. K. Y. Liu, R. L. Gould, M. C. Coulson, E. V. Ward, R. J. Howard, Tests of pattern
separation and pattern completion in humans—A systematic review. (2016).
Available at: https://onlinelibrary.wiley.com/doi/epdf/10.1002/hipo.22561 [Accessed
24 March 2025].

14. E. Rolls, The mechanisms for pattern completion and pattern separation in the
hippocampus. *Front. Syst. Neurosci.* **7** (2013).

15. A. Treves, E. T. Rolls, Computational constraints suggest the need for two distinct
input systems to the hippocampal CA3 network. *Hippocampus* **2**, 189–199 (1992).

16.   K. Z. Tanaka, *et al.*, Cortical Representations Are Reinstated by the Hippocampus
903   during Memory Retrieval. *Neuron* **84**, 347–354 (2014).

904   17.   S. Theves, X. Grande, E. Duzel, C. F. Doeller, "Pattern Completion and the Medial
905   Temporal Lobe Memory System" (Oxford University Press, 2024).

906   18.   L. Davachi, J. Danker, 18 Cognitive Neuroscience of Episodic Memory. *Oxf. Handb.*
907   *Cogn. Neurosci. Vol. 1 Core Top.* **1**, 375 (2013).

908   19.   B. P. Staresina, R. N. A. Henson, N. Kriegeskorte, A. Alink, Episodic Reinstatement in
909   the Medial Temporal Lobe. *J. Neurosci.* **32**, 18150–18156 (2012).

910   20.   A. N. Trelle, *et al.*, Hippocampal and cortical mechanisms at retrieval explain
911   variability in episodic remembering in older adults. *eLife* **9**, e55335 (2020).

912   21.   K. D. Duncan, M. L. Schlichting, Hippocampal representations as a function of time,
913   subregion, and brain state. *Neurobiol. Learn. Mem.* **153**, 40–56 (2018).

914   22.   M. E. Hasselmo, J. M. Bower, Acetylcholine and memory. *Trends Neurosci.* **16**, 218–
915   222 (1993).

916   23.   N. Bunzeck, *et al.*, Mesolimbic Novelty Processing in Older Adults. *Cereb. Cortex* **17**,
917   2940–2948 (2007).

918   24.   M. G. Giovannini, *et al.*, Effects of novelty and habituation on acetylcholine, GABA,
919   and glutamate release from the frontal cortex and hippocampus of freely moving
920   rats. *Neuroscience* **106**, 43–53 (2001).

921   25.   R. McQuade, D. Creton, S. C. Stanford, Effect of novel environmental stimuli on rat
922   behaviour and central noradrenaline function measured by in vivo microdialysis.
923   *Psychopharmacology (Berl.)* **145**, 393–400 (1999).

924   26.   D. Moncada, H. Viola, Induction of long-term memory by exposure to novelty requires
925   protein synthesis: Evidence for a behavioral tagging. *J. Neurosci.* **27**, 7476–7481
926   (2007).

927   27.   P. C. Prokopiou, *et al.*, Lower novelty-related locus coeruleus function is associated
928   with Aβ-related cognitive decline in clinically healthy individuals. *Nat. Commun.* **13**,
929   1571 (2022).

930   28.   K. Zhang, C. D. Chen, I. E. Monosov, Novelty, Salience, and Surprise Timing Are
931   Signaled by Neurons in the Basal Forebrain. *Curr. Biol.* **29**, 134-142.e3 (2019).

932   29.   L. C. Dang, J. P. O'Neil, W. J. Jagust, Dopamine Supports Coupling of Attention-Related
933   Networks. *J. Neurosci.* **32**, 9582–9587 (2012).

30. A. L. Decker, K. D. Duncan, Acetylcholine and the complex interdependence of
memory and attention. *Curr. Opin. Behav. Sci.* **32**, 21–28 (2020).

31. I. Klinkenberg, A. Sambeth, A. Blokland, Acetylcholine and attention. *Behav. Brain Res.*
221, 430–442 (2011).

32. A. Nieoullon, Dopamine and the regulation of cognition and attention. *Prog.*
Neurobiol. **67**, 53–83 (2002).

33. N. Unsworth, M. K. Robison, A locus coeruleus-norepinephrine account of individual
differences in working memory capacity and attention control. *Psychon. Bull. Rev.*
24, 1282–1311 (2017).

34. Y. Zhang, Y. Chen, Y. Xin, B. Peng, S. Liu, Norepinephrine system at the interface of
attention and reward. *Prog. Neuropsychopharmacol. Biol. Psychiatry* **125**, 110751
(2023).

35. A. L. Collins, T. J. Aitken, V. Y. Greenfield, S. B. Ostlund, K. M. Wassum, Nucleus
Accumbens Acetylcholine Receptors Modulate Dopamine and Motivation.
Neuropsychopharmacology **41**, 2830–2838 (2016).

36. L. J. Herberg, D. N. Stephens, K. B. J. Franklin, Catecholamines and self-stimulation:
Evidence suggesting a reinforcing role for noradrenaline and a motivating role for
dopamine. *Pharmacol. Biochem. Behav.* **4**, 575–582 (1976).

37. R. Ventura, E. C. Latagliata, C. Morrone, I. L. Mela, S. Puglisi-Allegra, Prefrontal
Norepinephrine Determines Attribution of "High" Motivational Salience. *PLOS ONE* **3**,
e3044 (2008).

38. R. A. Wise, Dopamine, learning and motivation. *Nat. Rev. Neurosci.* **5**, 483–494 (2004).

39. M. Clos, N. Bunzeck, T. Sommer, Dopamine Enhances Item Novelty Detection via
Hippocampal and Associative Recall via Left Lateral Prefrontal Cortex Mechanisms.
J. Neurosci. **39**, 7920–7933 (2019).

40. C. W. Harley, Norepinephrine and Dopamine as Learning Signals. *Neural Plast.* **11**,
360813 (2004).

41. M. E. Hasselmo, The role of acetylcholine in learning and memory. *Curr. Opin.*
Neurobiol. **16**, 710–715 (2006).

42. M.-M. Mesulam, E. J. Mufson, A. I. Levey, B. H. Wainer, Cholinergic innervation of
cortex by the basal forebrain: Cytochemistry and cortical connections of the septal
area, diagonal band nuclei, nucleus basalis (Substantia innominata), and
hypothalamus in the rhesus monkey. *J. Comp. Neurol.* **214**, 170–197 (1983).

43. M. E. Hasselmo, B. P. Fehlau, Differences in time course of ACh and GABA modulation
of excitatory synaptic potentials in slices of rat hippocampus. *J. Neurophysiol.* **86**,
1792–1802 (2001).

44. M. Meeter, J. m. j. Murre, L. m. Talamini, Mode shifting between storage and recall
based on novelty detection in oscillating hippocampal circuits. *Hippocampus* **14**,
722–741 (2004).

45. M. A. Burock, R. L. Buckner, M. G. Woldorff, B. R. Rosen, A. M. Dale, Randomized
event-related experimental designs allow for extremely rapid presentation rates using
functional MRI. *NeuroReport* **9**, 3735 (1998).

46. R. D. Morey, Confidence Intervals from Normalized Data: A correction to Cousineau
(2005) in *Tutorials in Quantitative Methods for Psychology*, (2008), pp. 61–64.

47. A. Patil, K. D. Duncan, Lingering Cognitive States Shape Fundamental Mnemonic
Abilities. *Psychol. Sci.* **29**, 45–55 (2018).

48. M. Ritchey, E. A. Wing, K. S. LaBar, R. Cabeza, Neural similarity between encoding and
retrieval is related to memory via hippocampal interactions. *Cereb. Cortex N. Y. N*
1991 **23**, 2818–2828 (2013).

49. D. J. Huffman, C. E. L. Stark, Multivariate pattern analysis of the human medial
temporal lobe revealed representationally categorical cortex and representationally
agnostic hippocampus. *Hippocampus* **24**, 1394–1403 (2014).

50. S.-H. Lee, D. J. Kravitz, C. I. Baker, Differential Representations of Perceived and
Retrieved Visual Information in Hippocampus and Cortex. *Cereb. Cortex N. Y. N 1991*
29, 4452–4461 (2019).

51. J. C. Liang, A. R. Preston, Medial temporal lobe reinstatement of content-specific
details predicts source memory. *Cortex* **91**, 67–78 (2017).

52. X. Xiao, *et al.*, Individual-specific and shared representations during episodic memory
encoding and retrieval. *NeuroImage* **217**, 116909 (2020).

53. J. F. Danker, A. Tompary, L. Davachi, Trial-by-Trial Hippocampal Encoding Activation
Predicts the Fidelity of Cortical Reinstatement During Subsequent Retrieval. *Cereb.*
Cortex N. Y. N 1991 **27**, 3515–3524 (2017).

54. B. A. Kuhl, M. K. Johnson, M. M. Chun, Dissociable Neural Mechanisms for Goal-
Directed Versus Incidental Memory Reactivation. *J. Neurosci.* **33**, 16099–16109
(2013).

55. B. A. Kuhl, M. M. Chun, Successful Remembering Elicits Event-Specific Activity
Patterns in Lateral Parietal Cortex. *J. Neurosci.* **34**, 8051–8060 (2014).

56. E. A. Wing, M. Ritchey, R. Cabeza, Reinstatement of Individual Past Events Revealed
by the Similarity of Distributed Activation Patterns during Encoding and Retrieval. *J.*
Cogn. Neurosci. **27**, 679–691 (2015).

57. X. Xiao, *et al.*, Transformed Neural Pattern Reinstatement during Episodic Memory
Retrieval. *J. Neurosci.* **37**, 2986–2998 (2017).

58. N. Kanwisher, G. Yovel, The fusiform face area: a cortical region specialized for the
perception of faces. *Philos. Trans. R. Soc. B Biol. Sci.* **361**, 2109–2128 (2006).

59. M. E. Hasselmo, Neuromodulation: acetylcholine and memory consolidation. *Trends*
Cogn. Sci. **3**, 351–359 (1999).

60. C. W. Berridge, B. D. Waterhouse, The locus coeruleus-noradrenergic system:
modulation of behavioral state and state-dependent cognitive processes. *Brain Res.*
Brain Res. Rev. **42**, 33–84 (2003).

61. O. Lindvall, A. Björklund, I. Divac, Organization of mesencephalic dopamine neurons
projecting to neocortex and septum. *Adv. Biochem. Psychopharmacol.* **16**, 39–46
(1977).

62. L. Zaborszky, *et al.*, Stereotaxic probabilistic maps of the magnocellular cell groups in
human basal forebrain. *NeuroImage* **42**, 1127–1141 (2008).

63. R. A. Diana, A. P. Yonelinas, C. Ranganath, Adaptation to cognitive context and item
information in the medial temporal lobes. *Neuropsychologia* **50**, 3062–3069 (2012).

64. B. A. Strange, A. Duggins, W. Penny, R. J. Dolan, K. J. Friston, Information theory,
novelty and hippocampal responses: unpredicted or unpredictable? *Neural Netw.*
18, 225–230 (2005).

65. C. E. L. Stark, L. R. Squire, When zero is not zero: The problem of ambiguous baseline
conditions in fMRI. *Proc. Natl. Acad. Sci.* **98**, 12760–12766 (2001).

66. E. S. Bromberg-Martin, M. Matsumoto, O. Hikosaka, Dopamine in Motivational
Control: Rewarding, Aversive, and Alerting. *Neuron* **68**, 815–834 (2010).

67. V. P. Murty, R. A. Adcock, Enriched encoding: reward motivation organizes cortical
networks for hippocampal detection of unexpected events. *Cereb. Cortex* **24**, 2160–
2168 (2014).

68. M. Voytko, *et al.*, Basal forebrain lesions in monkeys disrupt attention but not learning
and memory. *J. Neurosci.* **14**, 167–186 (1994).

69. G. L. Wenk, The nucleus basalis magnocellularis cholinergic system: One hundred
years of progress. *Neurobiol. Learn. Mem.* **67**, 85–95 (1997).

70. F. Tashakori-Sabzevar, R. D. Ward, Basal Forebrain Mediates Motivational Recruitment
of Attention by Reward-Associated Cues. *Front. Neurosci.* **12** (2018).

71. H. J. Neville, M. Kutas, G. Chesney, A. L. Schmidt, Event-related brain potentials during
initial encoding and recognition memory of congruous and incongruous words. *J.*
Mem. Lang. **25**, 75–92 (1986).

72. K. L. Vilberg, M. D. Rugg, Functional significance of retrieval-related activity in lateral
parietal cortex: Evidence from fMRI and ERPs. *Hum. Brain Mapp.* **30**, 1490–1501
(2009).

73. S. Han, S. A. Huettel, A. Raposo, R. A. Adcock, I. G. Dobbins, Functional Significance
of Striatal Responses during Episodic Decisions: Recovery or Goal Attainment? |
Journal of Neuroscience. (2010). Available at:
https://www.jneurosci.org/content/30/13/4767.short [Accessed 27 March 2025].

74. N. A. Herweg, T. Sommer, N. Bunzeck, Retrieval Demands Adaptively Change Striatal
Old/New Signals and Boost Subsequent Long-Term Memory. *J. Neurosci.* **38**, 745–754
(2018).

75. N. J. Cohen, H. Eichenbaum, *Memory, Amnesia, and the Hippocampal System* (MIT
Press, 1993).

76. Z. M. Reagh, C. Ranganath, Flexible reuse of cortico-hippocampal representations
during encoding and recall of naturalistic events. *Nat. Commun.* **14**, 1279 (2023).

77. A. Tompary, L. Davachi, Consolidation Promotes the Emergence of Representational
Overlap in the Hippocampus and Medial Prefrontal Cortex. *Neuron* **105**, 199–200
(2020).

78. Michael. L. Mack, Alison. R. Preston, Decisions about the past are guided by
reinstatement of specific memories in the hippocampus and perirhinal cortex.
NeuroImage **127**, 144–157 (2016).

79. A. Tompary, K. D. Duncan, L. Davachi, High-resolution investigation of memory-
specific reinstatement in the hippocampus and perirhinal cortex. *Hippocampus* **26**,
995–1007 (2016).

80. M. E. Hasselmo, E. Schnell, E. Barkai, Dynamics of learning and recall at excitatory
recurrent synapses and cholinergic modulation in rat hippocampal region CA3. *J.*
Neurosci. Off. J. Soc. Neurosci. **15**, 5249–5262 (1995).

81. M. E. Hasselmo, E. Schnell, Laminar selectivity of the cholinergic suppression of
synaptic transmission in rat hippocampal region CA1: computational modeling and
brain slice physiology. *J. Neurosci. Off. J. Soc. Neurosci.* **14**, 3898–3914 (1994).

82. A. Easton, V. Douchamps, M. Eacott, C. Lever, A specific role for septohippocampal
acetylcholine in memory? *Neuropsychologia* **50**, 3156–3168 (2012).

83. K. D. Duncan, D. Shohamy, Memory states influence value-based decisions. *J. Exp.*
Psychol. Gen. **145**, 1420–1426 (2016).

84. K. D. Duncan, A. Sadanand, L. Davachi, Memory's Penumbra: Episodic Memory
Decisions Induce Lingering Mnemonic Biases. *Science* **337**, 485–487 (2012).

85. S.-C. Lin, R. E. Brown, M. G. Hussain Shuler, C. C. H. Petersen, A. Kepecs, Optogenetic
Dissection of the Basal Forebrain Neuromodulatory Control of Cortical Activation,
Plasticity, and Cognition. *J. Neurosci. Off. J. Soc. Neurosci.* **35**, 13896–13903 (2015).

86. E. J. Mufson, S. D. Ginsberg, M. D. Ikonomovic, S. T. DeKosky, Human cholinergic basal
forebrain: chemoanatomy and neurologic dysfunction. *J. Chem. Neuroanat.* **26**, 233–
242 (2003).

87. F. Gargouri, *et al.*, Multimodal magnetic resonance imaging investigation of basal
forebrain damage and cognitive deficits in Parkinson's disease. *Mov. Disord.* **34**, 516–
525 (2019).

88. X. Liu, *et al.*, Subcortical evidence for a contribution of arousal to fMRI studies of brain
activity. *Nat. Commun.* **9**, 395 (2018).

89. J. Turchi, *et al.*, The Basal Forebrain Regulates Global Resting-State fMRI Fluctuations.
Neuron **97**, 940-952.e4 (2018).

90. V. Douchamps, A. Jeewajee, P. Blundell, N. Burgess, C. Lever, Evidence for Encoding
versus Retrieval Scheduling in the Hippocampus by Theta Phase and Acetylcholine.
J. Neurosci. **33**, 8689–8704 (2013).

91. J. Haam, J. L. Yakel, Cholinergic modulation of the hippocampal region and memory
function. *J. Neurochem.* **142 Suppl 2**, 111–121 (2017).

92. M. Vandecasteele, *et al.*, Optogenetic activation of septal cholinergic neurons
suppresses sharp wave ripples and enhances theta oscillations in the hippocampus.
Proc. Natl. Acad. Sci. **111**, 13535–13540 (2014).

93. L. Záborszky, *et al.*, Specific Basal Forebrain–Cortical Cholinergic Circuits Coordinate
Cognitive Operations. *J. Neurosci.* **38**, 9446–9458 (2018).

94. Y. Zhang, *et al.*, Cholinergic suppression of hippocampal sharp-wave ripples impairs
working memory. *Proc. Natl. Acad. Sci. U. S. A.* **118**, e2016432118 (2021).

95. I. Bethus, D. Tse, R. G. M. Morris, Dopamine and Memory: Modulation of the
Persistence of Memory for Novel Hippocampal NMDA Receptor-Dependent Paired
Associates. *J. Neurosci.* **30**, 1610–1618 (2010).

96. K. A. Kempadoo, E. V. Mosharov, S. J. Choi, D. Sulzer, E. R. Kandel, Dopamine release
from the locus coeruleus to the dorsal hippocampus promotes spatial learning and
memory. *Proc. Natl. Acad. Sci. U. S. A.* **113**, 14835–14840 (2016).

97. C. G. McNamara, Á. Tejero-Cantero, S. Trouche, N. Campo-Urriza, D. Dupret,
Dopaminergic neurons promote hippocampal reactivation and spatial memory
persistence. *Nat. Neurosci.* **17**, 1658–1660 (2014).

98. C. M. O'Carroll, S. J. Martin, J. Sandin, B. Frenguelli, R. G. M. Morris, Dopaminergic
modulation of the persistence of one-trial hippocampus-dependent memory. *Learn.*
Mem. **13**, 760–769 (2006).

99. T. Takeuchi, *et al.*, Locus coeruleus and dopaminergic consolidation of everyday
memory. *Nature* **537**, 357–362 (2016).

100. S. A. Thomas, Hippocampus | Neuroscience Journal | Wiley Online Library. (2015).
Available at:
https://onlinelibrary.wiley.com/doi/full/10.1002/hipo.22394?casa_token=lNzIWlrgT6k
AAAAA%3Av2cSBQDNQFn9FVSl3HqkUHqS0LnUiN0mY3xIjkGXC4zUa_rxk5XSN_Dx5
Fz1XrUEysZ2SJJpas_fCQ [Accessed 27 March 2025].

101. S. B. Floresco, A. G. Phillips, Delay-dependent modulation of memory retrieval by
infusion of a dopamine D1 agonist into the rat medial prefrontal cortex. *Behav.*
Neurosci. **115**, 934–939 (2001).

102. M. Hotte, *et al.*, D1 receptor modulation of memory retrieval performance is
associated with changes in pCREB and pDARPP-32 in rat prefrontal cortex. *Behav.*
Brain Res. **171**, 127–133 (2006).

103. D. Quartermain, H. J. Altman, Facilitation of retrieval by d-amphetamine following
anisomycin-induced amnesia. *Physiol. Psychol.* **10**, 283–292 (1982).

104. S. J. Sara, B. Deweer, Memory retrieval enhanced by amphetamine after a long
retention interval. *Behav. Neural Biol.* **36**, 146–160 (1982).

105. W. S. Stone, R. J. Rudd, P. E. Gold, Amphetamine, epinephrine, and glucose
enhancement of memory retrieval. *Psychobiology* **18**, 227–230 (1990).

106. M. E. Ballard, D. A. Gallo, H. de Wit, Amphetamine Increases Errors During Episodic
Memory Retrieval. *J. Clin. Psychopharmacol.* **34**, 85 (2014).

107. R. Pakdel, A. Rashidy-Pour, Glucocorticoid-induced impairment of long-term
memory retrieval in rats: An interaction with dopamine D2 receptors. *Neurobiol.*
Learn. Mem. **85**, 300–306 (2006).

108. J. Weafer, D. A. Gallo, H. de Wit, Amphetamine Fails to Alter Cued Recollection of
Emotional Images: Study of Encoding, Retrieval, and State-Dependency. *PLOS ONE*
9, e90423 (2014).

109. I. Zeeuws, E. Soetens, Verbal memory performance improved via an acute
administration of D-amphetamine. *Hum. Psychopharmacol.* **22**, 279–287 (2007).

110. J. E. Elum, *et al.*, Distinct dynamics and intrinsic properties in ventral tegmental area
populations mediate reward association and motivation. *Cell Rep.* **43** (2024).

111. R. M. Krebs, D. Heipertz, H. Schuetze, E. Duzel, Novelty increases the mesolimbic
functional connectivity of the substantia nigra/ventral tegmental area (SN/VTA)
during reward anticipation: Evidence from high-resolution fMRI. *NeuroImage* **58**,
647–655 (2011).

112. C. L. Asplund, J. J. Todd, A. P. Snyder, R. Marois, A central role for the lateral prefrontal
cortex in goal-directed and stimulus-driven attention. *Nat. Neurosci.* **13**, 507–512
(2010).

113. J. W. Y. Kam, A.-K. Solbakk, T. Endestad, T. R. Meling, R. T. Knight, Lateral prefrontal
cortex lesion impairs regulation of internally and externally directed attention.
NeuroImage **175**, 91–99 (2018).

114. M. Lepage, O. Ghaffar, L. Nyberg, E. Tulving, Prefrontal cortex and episodic memory
retrieval mode. *Proc. Natl. Acad. Sci.* **97**, 506–511 (2000).

115. M. B. Brodeur, K. Guérard, M. Bouras, Bank of Standardized Stimuli (BOSS) Phase II:
930 New Normative Photos. *PLOS ONE* **9**, e106953 (2014).

116. J. W. Peirce, PsychoPy--Psychophysics software in Python. *J. Neurosci. Methods* **162**,
8–13 (2007).

117. A. M. Dale, Optimal experimental design for event-related fMRI. *Hum. Brain Mapp.* **8**,
109–114 (1999).

118. D. Bates, M. Mächler, B. Bolker, S. Walker, Fitting Linear Mixed-Effects Models Using
lme4. *J. Stat. Softw.* **67**, 1–48 (2015).

119. R. V. Lenth, emmeans: Estimated Marginal Means, aka Least-Squares Means.
https://doi.org/10.32614/CRAN.package.emmeans. Deposited 2025.

120. O. Esteban, *et al.*, poldracklab/fmriprep: 1.2.5. (2018).
https://doi.org/10.5281/zenodo.1942006. Deposited 4 December 2018.

121. O. Esteban, *et al.*, FMRIPrep: a robust preprocessing pipeline for functional MRI.
bioRxiv 306951 (2018). https://doi.org/10.1101/306951.

122. K. Gorgolewski, *et al.*, Nipype: A Flexible, Lightweight and Extensible Neuroimaging
Data Processing Framework in Python. *Front. Neuroinformatics* **5** (2011).

123. K. J. Gorgolewski, *et al.*, nipy/nipype: 1.1.6. (2018).
https://doi.org/10.5281/zenodo.1560596. Deposited 26 November 2018.

124. N. J. Tustison, *et al.*, N4ITK: Improved N3 Bias Correction. *IEEE Trans. Med. Imaging*
29, 1310–1320 (2010).

125. V. Fonov, A. Evans, R. McKinstry, C. Almli, D. Collins, Unbiased nonlinear average
age-appropriate brain templates from birth to adulthood. *NeuroImage* **47**, S102
(2009).

126. V. Fonov, *et al.*, Unbiased average age-appropriate atlases for pediatric studies.
NeuroImage **54**, 313–327 (2011).

127. B. B. Avants, C. L. Epstein, M. Grossman, J. C. Gee, Symmetric diffeomorphic image
registration with cross-correlation: Evaluating automated labeling of elderly and
neurodegenerative brain. *Med. Image Anal.* **12**, 26–41 (2008).

128. Y. Zhang, M. Brady, S. Smith, Segmentation of brain MR images through a hidden
Markov random field model and the expectation-maximization algorithm. *IEEE Trans.*
Med. Imaging **20**, 45–57 (2001).

129. M. Jenkinson, C. F. Beckmann, T. E. J. Behrens, M. W. Woolrich, S. M. Smith, FSL.
NeuroImage **62**, 782–790 (2012).

130. M. F. Glasser, *et al.*, The minimal preprocessing pipelines for the Human
Connectome Project. *NeuroImage* **80**, 105–124 (2013).

131. M. Jenkinson, S. Smith, A global optimisation method for robust affine registration of
brain images. *Med. Image Anal.* **5**, 143–156 (2001).

132. D. N. Greve, B. Fischl, Accurate and robust brain image alignment using boundary-
based registration. *NeuroImage* **48**, 63–72 (2009).

133. M. Jenkinson, P. Bannister, M. Brady, S. Smith, Improved Optimization for the Robust
and Accurate Linear Registration and Motion Correction of Brain Images.
NeuroImage **17**, 825–841 (2002).

134.   A. Klein, *et al.*, Evaluation of 14 nonlinear deformation algorithms applied to human
brain MRI registration. *NeuroImage* **46**, 786–802 (2009).

135.   C. Lanczos, Evaluation of Noisy Data. *J. Soc. Ind. Appl. Math. Ser. B Numer. Anal.* **1**,
76–85 (1964).

136.   J. D. Power, *et al.*, Methods to detect, characterize, and remove motion artifact in
resting state fMRI. *NeuroImage* **84**, 320–341 (2014).

137.   Y. Behzadi, K. Restom, J. Liau, T. T. Liu, A component based noise correction method
(CompCor) for BOLD and perfusion based fMRI. *NeuroImage* **37**, 90–101 (2007).

138.   A. Abraham, *et al.*, Machine learning for neuroimaging with scikit-learn. *Front.*
Neuroinformatics **8** (2014).

139.   J. A. Mumford, B. O. Turner, F. G. Ashby, R. A. Poldrack, Deconvolving BOLD activation
in event-related designs for multivoxel pattern classification analyses. *NeuroImage*
59, 2636–2643 (2012).

1209