A minimal CRISPR polymerase produces decoy cyclic nucleotides to detect phage anti-defense proteins

Ashley E. Sullivan[1], Ali Nabhani[2], Kate Schinkel[1], David M. Dinh[2], Melissa L. Duncan[1], Eirene Marie Q. Ednacot[2,3], Charlotte R.K. Hoffman[1], Daniel S. Izrailevsky[2], Emily M. Kibby[1], Toni A. Nagy[1], Christy M. Nguyen[4,5], Uday Tak[1], A. Maxwell Burroughs[6], L. Aravind[6], Aaron T. Whiteley[1]*, Benjamin R. Morehouse[2,3,7,8]*

[1] Department of Biochemistry, University of Colorado Boulder, Boulder, CO, USA
[2] Department of Molecular Biology and Biochemistry, University of California Irvine, Irvine, CA, USA
[3] Department of Pharmaceutical Sciences, University of California Irvine, Irvine, CA, USA
[4] Department of Biological Chemistry, University of California Irvine, Irvine, CA, USA
[5] Center for Epigenetics and Metabolism, University of California Irvine, Irvine, CA, USA
[6] National Center for Biotechnology Information, National Library of Medicine, National Institutes of Health, Bethesda, MD, USA
[7] Institute for Immunology, University of California Irvine, Irvine, CA, USA
[8] Center for Virus Research, University of California Irvine, Irvine, CA, USA

* To whom correspondence should be addressed: aaron.whiteley@colorado.edu (A.T.W) and b.morehouse@uci.edu (B.R.M.)

26 **Abstract**
27
Bacteria use antiphage systems to combat phages, their ubiquitous competitors, and evolve new
defenses through repeated reshuffling of basic functional units into novel reformulations. A
common theme is generating a nucleotide-derived second messenger in response to phage that
activates an effector protein to halt virion production. Phages respond with counter-defenses that
deplete these second messengers, leading to an escalating arms race with the host. Here we
discover a novel antiphage system we call Panoptes that detects phage infection by surveying
the cytosol for phage proteins that antagonize the nucleotide-derived second messenger pool.
Panoptes is a two-gene operon, *optSE*. OptS is predicted to synthesize a second messenger
using a minimal CRISPR polymerase (mCpol) domain, a version of the polymerase domain found
in Type III CRISPR systems (Cas10) that is distantly related to GGDEF and Thg1 tRNA repair
polymerase domains. OptE is predicted to be a transmembrane effector protein that binds cyclic
nucleotides. *optSE* potently restricted phage replication but mutant phages that had loss-of-
function mutations in anti-CBASS protein 2 (Acb2) escaped defense. These findings were
unexpected because Acb2 is a nucleotide "sponge" that antagonizes second messenger
signaling. Using genetic and biochemical assays, we found that Acb2 bound the OptS-
synthesized nucleotide, 2′,3′-cyclic adenosine monophosphate (2′,3′-c-di-AMP); however, 2′,3′-c-
di-AMP was synthesized constitutively by OptS and inhibited OptE. Nucleotide depletion by Acb2
released OptE toxicity thereby initiating abortive infection to halt phage replication. These data
demonstrate a sophisticated immune strategy that hosts use to guard their second messenger
pool and turn immune evasion against the virus.
48
49
50 **Keywords**: Acb2, CBASS, cyclic oligonucleotide based antiphage signaling system, S-2TMβ
51

52 **Manuscript Text**
53
54 **Introduction**
55
Bacteria are under constant assault from their viruses, bacteriophages (phages), and have
evolved sophisticated mechanisms to protect themselves[1,2]. The immune system of any
bacterium is a combination of individual pathways called antiphage systems that limit phage
replication. Bacteria encode an average of 5.8 antiphage systems per strain[3]. However, each
bacterial strain within a species dynamically exchanges antiphage systems through horizontal
gene transfer giving each a different suite of defenses[1].
62
Antiphage systems must rapidly sense phage infection and transduce an activation signal to an
effector protein to stop virion production. Recent progress cataloging antiphage systems has
identified that many systems transduce their activation signal through nucleotide-derived second
messengers. The cyclic oligonucleotide-based antiphage signaling system (CBASS) synthesizes
cyclic di– and trinucleotides[4–6], the pyrimidine cyclase system for antiphage resistance (Pycsar)
synthesizes cyclic mononucleotides[7], and the Thoeris system synthesizes cyclic ADP-ribose
derivatives in response to phage[8]. The advantage of nucleotide-derived second messengers is
they can be synthesized from abundant precursors and amplify a rare phage-sensing event into
stoichiometrically more activated effector proteins. The success of these strategies is further
underscored by their remarkable spread through lateral transfer: homologs of CBASS
components can be found in the cGAS-STING pathway of metazoans and homologs of Thoeris
components can be found in the immune system of plants and metazoans[9,10].
75
Nucleotide-derived second messengers are also a vulnerability because viruses use
phosphodiesterases and "sponge proteins" to interrupt signaling[11–18]. Just as the nucleotide
second messenger strategy is pervasive from bacteria to eukaryotes, so too are some of the
immune evasion strategies from viruses that infect bacteria and metazoans[19,20]. Bacterial
antiphage systems synthesize a wide variety of chemical variations of their class of second
messenger, often using different nucleotide bases, forming different phosphodiester linkages, or
even incorporating amino acids[6,7,21]. This variability is likely the result of the arms race between
host and pathogen. Nevertheless, most viral proteins that antagonize second messengers can
interact with a wide range of second messenger variants suggesting that changing the second
messenger is not sufficient to out-pace the virus[11–14,17,18,22–24].
86
Here we investigated a two-gene operon (*optSE*) that appeared similar to CBASS, Pycsar, and
Thoeris systems with a predicted nucleotide second messenger synthase (OptS) and a
nucleotide-binding effector protein (OptE). However, unlike systems that synthesize signaling
nucleotides in response to phage, we found that OptS constitutively produces a signaling
nucleotide to repress growth inhibition by OptE. The OptS nucleotide, 2′,3′-c-di-AMP, is similar to
nucleotides found in CBASS systems, which is synthesized by a SMODS domain containing
protein called a CD-NTase. Phages that encoded anti-defense proteins that deplete cyclic
nucleotides to circumvent CBASS signaling inadvertently depleted OptS-synthesized 2′,3′-c-di-
AMP and unmasked OptE toxicity, which interrupted the phage replication cycle through
destruction of the host cell, a process called abortive infection. In this way, OptSE guards the
cyclic dinucleotide pool of the cell and detects phage infection by sensing immune evasion. We
named OptSE the Panoptes antiphage system for Argus Panoptes, the all-seeing, many-eyed
giant in Greek mythology who was a faithful watchman to Hera.

Results

The Panoptes system is antiphage

We investigated a candidate two-gene Panoptes operon from *Vibrio navarrensis*. The first gene (*optS,* S = Synthase) encoded a minimal CRISPR polymerase (mCpol) domain that is predicted to synthesize a nucleotide second messenger[25]. The second gene (*optE*, E = Effector) encoded two predicted transmembrane domains and a SMODS-associating 2TM, β-strand rich (S-2TMβ) domain (**Fig. 1a**)[25]. OptE is homologous to Cap15 proteins within CBASS that disrupt membrane integrity in response to nucleotide[25,26]. OptSE resembled CBASS[5], Pycsar[7], and Theoris[8] systems with a nucleotide-derived second messenger-generating protein and a second messenger-responsive effector protein. We expressed the operon from its endogenous promoter on a medium-copy plasmid in *E. coli* MG1655 and challenged these bacteria with a panel of diverse phages. The *optSE* operon specifically defended against phages from the Straboviridae (**Fig. 1b)**. We focused on the *Tequatroviruses* genus that includes the T-even phages and found *optSE* provided over 100-fold protection against phage BASEL 38, 41, 42, and 44, and over 500-fold protection for BASEL36, 37, 39, 40, and 43 (**Fig. 1b and c**).

We selected the best characterized T-even phage, phage T4, for additional characterization. We confirmed that *optSE* provided defense against phage T4 in both soft-agar and liquid cultures (**Fig. 1c**; **ED Fig. 1**). A somewhat surprising outlier was phage T2, which was not restricted by *optSE*. These findings were unexpected as phage T2 is generally much more sensitive to phage defense systems than phage T4[27,28]. Taken together, the results support that Panoptes is an antiphage system.

OptS is a minimal CRISPR polymerase

To better understand the possible enzymatic function of the OptS protein, we expressed and purified an ortholog from *K. pneumoniae* KP67 (*Kp*OptS) for *in vitro* biochemical and structural characterization experiments. While *Kp*OptS is a small protein (~13 kDa) based on primary amino acid sequence alone, size exclusion chromatography supported the existence of a stable tetrameric species in solution (**Fig. 2a**). We next determined the crystal structure of *Kp*OptS, which could be refined at a nominal resolution of 1.75 Å (**ED Table 1, See methods**). The crystallographic asymmetric unit contained four copies of *Kp*OptS, forming an apparent tetrameric complex, which further supports that *Kp*OptS likely exists as a tetramer (**Fig. 2b; ED Fig. 2a-c**). Minimal conformational and B-factor variation was observed across the protomers and superposition revealed RMSD values of ~0.3-0.4 Å on average between pairs.

The protomer of *Kp*OptS contains a RNA-recognition motif (RRM, sometimes referred to as ferredoxin) polymerase palm-domain fold composed of four primary α-helices surrounding a five-stranded β-sheet (**Fig. 2c**)[25]. Although quite different in quaternary structure, the protomeric unit of *Kp*OptS aligns well with the palm domains of the Cas10 (Csm1) from *Thermococcus onnurineus* and the GGDEF diguanylate cyclase PleD from *Caulobacter crescentus* with RMSD values of 3.86 Å and 2.39 Å, respectively (**Fig. 2d and e**)[29,30]. These polymerase domains were also earlier unified with the histidinyl tRNA repair polymerase Thg1[31]. Indeed, the OptS fold observed in the crystal structure previously predicted by Burroughs et al. and described as a minimal CRISPR polymerase (mCpol) owing to predicted homology to Cas10 enzymes from type III CRISPR immune systems[25]. GGDEF diguanylate cyclases synthesize cyclic diguanosine monophosphate (c-di-GMP) from GTP precursors and Cas10 enzymes synthesize cyclic oligoadenylate molecules (between 2–6 AMPs) from ATP precursors[32–34]. The high degree of structural similarity to GGDEF, Cas10, and other nucleotide polymerases (palm domains are

151 found in a variety of nucleotidyltransferases such as the SMODS of CBASS and cUMP/cCMP
152 synthases of Pycsar defense systems) strongly implicates Panoptes OptS proteins in the
153 enzymatic processing of nucleotides.
154
155 *OptS synthesizes 2′,3′-c-di-AMP*
156

Purified *Kp*OptS was incubated with ribonucleotide triphosphates (ATP, GTP, CTP, UTP) and the
reactions were separated and monitored by high-performance liquid chromatography (HPLC) to
detect any synthesis activity (**ED Fig. 3a-d**). The largest abundance product observed was
derived from ATP with minor products requiring the coincubation of both ATP and GTP (**Fig. 3a,**
ED Fig. 3b,c). We sought to determine the identity of the ATP-derived *Kp*OptS product and found
that it had a similar retention time to an isomer of cyclic diadenosine monophosphate with mixed
phosphodiester linkages, 2′,3′-c-di-AMP (c[A(2′–5′)pA(3′–5′)p]) (**Fig. 3b**). Additional biochemical
evidence for both the cyclic and mixed phosphodiester linkages of the product came from
nuclease treatment experiments (**ED Fig. 3e**). The product was resistant to calf-intestinal
phosphatase (CIP, cleaves terminal phosphates) but was partially susceptible to P1 nuclease
(cleaves 3′–5′ phosphodiester bonds) shifting to a new retention time. The combination CIP and
P1 treatment led to loss of signal at retention times observed in the untreated and P1-alone
treatment conditions and the appearance of a new, late eluting, peak that was consistent with
A(2′–5′)pA generated by CIP treatment of commercially available linear pppA(2′–5′)pA.

Based on conservation with diverse nucleotidyltransferases, we expected that OptS Asp6 and
Asp57 would be involved in catalysis as they likely coordinate Mg^{2+} ions for triphosphate
stabilization in the binding pocket (**Fig. 3c and ED Fig. 3f**). Accordingly, mutation of these amino
acids (either to Ala or Asn) alone or in tandem led to complete loss of cyclic dinucleotide
production (**Fig. 3d**). Finally, liquid chromatography-mass spectrometry (LC-MS) analysis of the
*Kp*OptS reaction with ATP confirmed that the main product is indeed 2′,3′-c-di-AMP (**Fig. 3e and**
ED Fig. 3g).

178
179
180 *Acb2 is necessary for activation of the Panoptes system*
181
Phage T4 formed irregular plaques in soft-agar overlays of *optSE*-expressing bacteria (**ED Fig.**
4a). At higher concentrations of phage, large clear plaques appeared and we hypothesized that
these large plaques might be caused by mutant phages capable of escaping *optSE*-mediated
defense. We isolated escaper mutants from three unrelated, clonal T4 lysates. 15 candidate
escapers were isolated, plaque purified, and genome sequenced along with their parent wild-type
phages (**ED Fig. 4b**). Surprisingly, 14 of 15 escaper phages encoded mutations exclusively in the
anti-CBASS 2 gene (*acb2/vs.*4; **Fig. 4a and ED Table 2**). Mutations included amino acid
substitutions (E26K, C78W, F82C, or P84Q), insertions, and deletions that caused frame shifts,
premature stop codons, and large-scale deletions.
191
Acb2 was recently discovered as an anti-defense protein that antagonizes antiphage immunity by
acting as a nucleotide "sponge," sequestering CBASS-derived cyclic oligonucleotide signaling
molecules[12,13,22]. Acb2 binding of CBASS-derived nucleotides interrupts downstream effector
protein activation, allowing the phage to evade defense. We were surprised to find that escaper
phages harbored obvious loss-of-function mutations in Acb2 because this ruled out that our
escapers had simply mutated Acb2 to increase affinity for the OptS-derived nucleotide. Instead,
these data suggested that Acb2 was necessary for OptSE-mediated defense.
199

We tested whether Acb2 was necessary for OptSE-mediated defense by challenging empty vector (EV) or *optSE*-expressing bacteria with either wild-type or Δ*acb2* T4 phages. Wild-type T4 efficiently replicated on EV-expressing bacteria and was restricted by *optSE*. T4 Δ*acb2*, however, was no longer restricted by OptSE and similarly replicated in both the EV- and *optSE*-expressing bacteria (**Fig. 4b**). Phage T4 also encodes anti-CBASS 1 (*acb1*), a phosphodiesterase that degrades cyclic oligonucleotides to similarly antagonize CBASS signaling[11]. We constructed a T4Δ*acb1* phage and found no impact on *optSE*-mediated defense (**Fig. 4b; ED Fig 4c**). Nevertheless, the T4Δ*acb1*Δ*acb2* double mutant appeared similar to T4 Δ*acb2* alone. Taken together, these data suggest that the phage gene *acb2*, but not *acb1*, is necessary for *optSE*-mediated defense.

Acb2 is sufficient for activation of the Panoptes system

We next explored whether *acb2* is sufficient to activate Panoptes in the absence of phage. OptE is a member of the S-2TMβ family of proteins that are comprised of a two-pass transmembrane element obligately fused to a soluble β-barrel nucleotide-binding domain. Another member of this family, Cap15, is a CBASS effector that disrupts the bacterial inner membrane to initiate abortive infection[26]. We hypothesized that OptE might similarly disrupt bacterial host membranes and inhibit colony formation. Therefore, we co-expressed the *optSE* operon with *acb2* and assayed for colony formation. T4 *acb2* selectively inhibited colony formation only when co-expressed with the *optSE* operon (**Fig. 4c**). We observed the same phenotype when we expressed *acb2* from phage BASEL 38, which is restricted by the *optSE* operon by over 400-fold. These data show that Acb2 is both necessary and sufficient to activate OptSE.

Using the same co-expression assay, we found that the mutant *acb2* allele encoding Acb2[F82C] identified in our escaper phage did not inhibit colony formation. The structure of Acb2 from phage T4 shows the F82 residue base-stacking with bound cyclic dinucleotides[13]. These data, along with other identified escaper mutations, suggested that Acb2 nucleotide binding was essential to activation of OptSE. We tested this hypothesis using Acb2[Y8A], a mutation that was previously shown to disrupt cyclic dinucleotide binding (**Fig. 4c**)[12,13]. Acb2[Y8A] co-expression with *optSE* also failed to inhibit colony formation. These findings suggest that Acb2 binds nucleotides to activate OptSE signaling.

Given that Acb2 activates the OptSE system and is a phage "sponge" that binds a wide array of nucleotide-derived products, we confirmed that Acb2 can bind the *Kp*OptS product, 2′,3′-c-di-AMP. HPLC experiments showed that incubation of purified T4 Acb2 with *Kp*OptS-derived 2′,3′-c-di-AMP depleted detectable signal (**ED Fig. 5a,b**). When treated with proteinase K, Acb2 was degraded, and 2′,3′-c-di-AMP was once again detected by HPLC, indicating that Acb2 functions as a sponge of 2′,3′-c-di-AMP. As expected, the escaper Acb2[F82C] showed diminished binding (**ED Fig. 5c**). Consistent with these results, isothermal titration calorimetry (ITC) demonstrated that Acb2 binds 2′,3′-c-di-AMP with an apparent K_d of 63.7 nM (**Fig. 4d; ED Fig. 5d-f**).

Our findings were initially paradoxical. Phage defense systems such as CBASS, Pycsar, and Thoeris each generate nucleotide second messengers to activate defense. OptS and OptE are predicted to generate and respond to cyclic oligonucleotides, respectively. Yet Acb2 nucleotide binding activated, rather than inhibited, OptSE signaling. To explain these observations, we hypothesized that rather than OptS-derived nucleotides *activating* OptE, instead, OptS-derived second messengers *inhibit* OptE. If this were the case, OptE should inhibit growth unless an OptS-derived nucleotide is present.

We tested this hypothesis using plasmids that expressed *optS* and *optE* from IPTG- and arabinose-inducible promoters. Expression of *optS* alone did not impact growth, however,

252 expression of *optE* alone potently inhibited colony formation (**Fig. 4e**). Growth of *optE*-expressing
253 strains could be restored by expressing wild-type *optS* but required the predicted OptS active site
254 residues.
255
To further confirm our hypothesis, we measured OptS synthesis of 2′,3′-c-di-AMP *in vivo* using a
biosensor to interrogate bacterial cell lysates (**ED Fig. 6**). 2′,3′-c-di-AMP could not be detected in
E. coli expressing empty vector, however, the nucleotide was robustly detected in *optSE*-
expressing cells. We estimate the intracellular concentration of 2′,3′-c-di-AMP in *optSE*-
expressing cells to be between 165 and 330 nM. These data support a model where, under
steady-state conditions, OptS constitutively produces 2′,3′-c-di-AMP to restrain the toxicity of
OptE and, thus, depletion of nucleotide by Acb2 unleashes OptE-mediated growth inhibition.
263
264 *Diverse phage proteins activate Panoptes*
265
Phages encode a wide variety of genes that allow evasion of bacterial immune pathways that use
nucleotide-derived signaling molecules. Thoeris anti-defense proteins 1 (Tad1) and 2 (Tad2)
sequester both glycocyclic ADP-Ribose (gcADPR) molecules synthesized by TIR domains during
Thoeris defense and cyclic oligonucleotides synthesized by CBASS systems[14,17,24]. Another
mechanism that phages use to bypass bacterial immune pathways is through degradation of the
nucleotide signaling molecule. Anti-Pycsar 1 (Apyc1), anti-CBASS 1 (Acb1), and type III anti-
CRISPR (AcrIII-1) are phage-encoded phosphoesterases that degrade cyclic nucleotide
molecules[11,18]. We hypothesized that these phage anti-defense proteins might also activate the
OptSE system through either sequestration or degradation of 2′,3′-c-di-AMP molecules
synthesized by OptS.
276
We co-expressed four Tad1 alleles from diverse phages with *optSE* in *E. coli* and observed
inhibition of colony formation when either *Cb*Tad1 *or Coli*Tad1 were expressed (**Fig. 5a**). We also
tested four different Tad2 proteins and only observed a slight reduction in colony formation when
SptTad2 was expressed in the presence of OptSE. These data suggest that other phage "sponge"
proteins bind and sequester 2′,3′-c-di-AMP, thereby activating the Panoptes system. On the other
hand, we found the phage phosphoesterases Apyc1 or AcrIII-1 did not activate the Panoptes
system (**Fig. 5a**), and that *acb1* did not alter Panoptes restriction of T4 phage (**Fig. 4b**). These
data indicate that a subset of phage immune evasion proteins can be detected by the Panoptes
system, which may be dictated by their nucleotide preference.
286
287 *Phyletic patterns and gene-neighborhood linkages of the mCpol-dependent prokaryotic immune*
288 *systems*
289
mCpol-containing systems are relatively rare but widely distributed across the two prokaryotic
superkingdoms (**Extended Data Fig. 7a**). Like the GGDEF and CRISPR polymerase domains,
they are strictly excluded from eukaryotes[31]. 53% of all mCpol systems are encoded in genomes
also coding for systems centered on SMODS enzymes, a highly significant linkage (p=3.7 x 10^{-12})
indicative of functional coupling between them (**Extended Data Fig. 7b and c**). Further, the
mCpol genes show a similarly significant coupling with genes coding for members of the S-2TMβ
family, like OptE (p=6.9 x 10^{-25}). Notably, beyond co-occurring in the same genome as SMODS
domain proteins, in roughly half of these cases, *optSE* is also frequently coupled to CBASS
systems in the same gene neighborhood as immediately adjacent operons (**Extended Data Fig.**
7). In these instances, the coupled CBASS systems typically encode an effector with either the
S-2TMβ or patatin-like phospholipase domain and a Ub-conjugation-like system implicated in the
generation of CD-NTase-target protein adducts[13,28,35]. Taken together, these observations
strongly suggest that Panoptes systems are predominantly guardians of CBASS systems whose
effectors primarily act by compromising the integrity of the cell membrane. Frequently, these

coupled systems display two distinct S-2TMβ effectors – one related to Cap15 and one related to
OptE (**Extended Data Fig. 7d**) – suggesting an interplay between the versions that are positively
and negatively controlled by the nucleotides generated by their respective regulatory synthases.
To emphasize the selective benefit or disadvantage of phage T4 expressing *acb2*, we showed
that while expressing *acb2* is harmful for phage replication in the presence of the Panoptes
defense system, it is required for circumventing a *Vibrio cholerae* CBASS system that synthesizes
3′,3′-cGAMP (**Fig. 5b**).
311
Beyond coupling with the S-2TMβ, 40% of the mCpol genes are coupled in operons with a gene
coding for a protein with a 2TM segment coupled with a CARF domain and a smaller set with a
comparable gene encoding a similar 2TM protein with a SAVED domain in lieu of the CARF
domain (**Extended Data Fig. 7d**). In these contexts, the CARF and SAVED domains are related
Rossmannoid second-messenger sensor domains that take the place of the nucleotide-binding
β-barrel domains of the S-2TMβ effectors[36]. Hence, these are predicted to be comparable
effectors to the S-2TMβ proteins that are regulated by second-messenger nucleotides. Further,
more infrequently, the mCpol genes are coupled in operons or fused to HEPN RNases and an
array of predicted membrane perforating 2TM domains. This suggests that the mCpol-generated
nucleotides might regulate a wider spectrum of effectors; however, it remains to be seen if they
are negative regulators as in the Panoptes system or positive regulators as in the CRISPR and
SMODS-based CBASS systems.
324
325 **Discussion**
326
Here we discovered that the Panoptes antiphage system flips the liability of a nucleotide-derived
second messenger in immune signaling against the phage by detecting the activity of counter-
defense proteins. OptS constitutively synthesized 2′,3′-c-di-AMP to hold OptE in an inactive state.
During phage infection, Acb2 or similar immune evasion proteins are produced that sequester the
OptS-derived signaling molecule, leading to a population of OptE that is no longer bound to cyclic
dinucleotide and is free to become activated, in turn aborting infection. While both OptE and
Cap15 are closely related members of the S-2TMβ family, they appear to use opposite signaling
modalities. OptE inhibited growth in the absence of cyclic nucleotide, whereas Cap15 inhibited
growth in the presence of cyclic nucleotide. We anticipate that the growth-inhibiting state of both
proteins is the oligomer, which disrupts membrane integrity, but that subtle differences alter
whether nucleotide binding stabilizes that monomer or oligomer.
338
The Panoptes system forces a dilemma for phage: should the phage lose Acb2 and be susceptible
to CBASS or evade CBASS and risk agonizing Panoptes. Phage T4 has maintained a wild-type
Acb2, however, Phage T2 encodes a mutation that likely selectively disrupts the Acb2 cyclic
dinucleotide binding site and the related *Pseudomonas aeruginosa* phage PaMx41 encodes a
premature stop codon in *acb2*[12]. Phage can solve this dilemma by tuning the specificity of Acb2
away from the OptS-derived nucleotide, using an alternative CBASS-evasion protein such as
Acb1 with different nucleotide specificity, or employing CBASS antagonists that cripple the
nucleotide synthase, called a CD-NTase, directly[37]. However, these responses by phage are
limited by a key feature of the bacterial immune system: antiphage systems are distributed
throughout that bacterial pangenome. In this way, Panoptes and CBASS can be encoded in the
same genome so long as they use slightly different nucleotides or as different genes and use the
identical nucleotide. The latter circumstance would result in antiphage system incompatibility–an
interesting additional layer to the ecology of phage defense gene distribution. Undoubtedly,
Panoptes is partially responsible for the evolutionary pressures that have selected for an
astonishing diversity of CD-NTase products found in CBASS systems.
354

Panoptes joins a growing list of antiphage systems that detect immune evasion, such as PARIS and PrrC[38,39]. However, a key difference is that Panoptes surveys the cell using a decoy molecule to sense the activity of anti-defense, as opposed to directly binding the proteins of anti-defense. We speculate that bacteria must encode yet-to-be-discovered OptSE-like systems that guard Pycsar- and Thoeris-derived second messenger pools too. The discovery of these systems has the exciting potential to identify further protein folds capable of synthesizing diverse nucleotide-derived second messengers. In terms of domain architecture and structure, the minimal polymerase catalytic palm domain seen in mCpols is in stark contrast to the incorporation of the cognate domain in complex domain architectures both in Type III CRISPR Cas10 polymerases and GGDEF domains involved in physiological signaling. This is in keeping with the hypothesis that mCpol-centered systems like Panoptes are surviving remnants of ancient antiviral systems that were subsequently elaborated and incorporated as the second messenger signaling arm of a subset of the CRISPR systems[31,36].

Like bacteria, eukaryotes also guard immune pathways by surveying for immune evasion often in a process called "effector-triggered immunity"[40–42]. Mammals use nucleotide-derived second messengers for immune signaling and mammalian viruses degrade second messengers in the cGAS-STING and OAS-RNaseL pathways using poxin and 2′,5′-phosphodiesterases, respectively[15,16]. It is highly likely that further mechanisms can be used by these viruses to disrupt second messenger pools, including sponge proteins analogous to Acb2. Undiscovered layers of human immune pathways may also exist that detect these viral evasion mechanisms in a manner similar to OptSE defense systems. Intriguingly, there are many SMODS and TIR domain enzymes of unknown function in several metazoans that could mediate comparable activities[43,44].

The Panoptes system represents a novel reformulation of basic functional units that are used in other antiphage systems. However, subtle molecular changes have enabled a new mechanism for sensing phage infection through detecting immune evasion. Although the phage would appear to be powerless to escape the combination of CBASS and Panoptes, future research will surely uncover the inevitable phage-encoded anti-anti-anti-defense.

References

385   **References**
386
1. Bernheim, A. & Sorek, R. The pan-immune system of bacteria: antiviral defence as a

community resource. *Nat. Rev. Microbiol.* **18**, 113–119 (2020).

2. Georjon, H. & Bernheim, A. The highly diverse antiphage defence systems of bacteria. *Nat.*

Rev. Microbiol. **21**, 686–700 (2023).

3. Millman, A. *et al.* An expanded arsenal of immune systems that protect bacteria from phages.

Cell Host Microbe S1931312822004735 (2022) doi:10.1016/j.chom.2022.09.017.

4. Ye, Q. *et al.* HORMA Domain Proteins and a Trip13-like ATPase Regulate Bacterial cGAS-

like Enzymes to Mediate Bacteriophage Immunity. *Mol. Cell* **77**, 709-722.e7 (2020).

5. Cohen, D. *et al.* Cyclic GMP–AMP signalling protects bacteria against viral infection. *Nature*

574, 691–695 (2019).

6. Whiteley, A. T. *et al.* Bacterial cGAS-like enzymes synthesize diverse nucleotide signals.

Nature **567**, 194–199 (2019).

7. Tal, N. *et al.* Cyclic CMP and cyclic UMP mediate bacterial immunity against phages. *Cell*

184, 5728-5739.e16 (2021).

8. Ofir, G. *et al.* Antiviral activity of bacterial TIR domains via immune signalling molecules.

Nature **600**, 116–120 (2021).

9. Bonhomme, D. *et al.* A human homolog of SIR2 antiphage proteins mediates immunity via

the TLR pathway. 2024.09.18.613514 Preprint at https://doi.org/10.1101/2024.09.18.613514

(2024).

10.   Ledvina, H. E. & Whiteley, A. T. Conservation and similarity of bacterial and eukaryotic

innate immunity. *Nat. Rev. Microbiol.* **22**, 420–434 (2024).

11.   Hobbs, S. J. *et al.* Phage anti-CBASS and anti-Pycsar nucleases subvert bacterial

immunity. *Nature* **605**, 522–526 (2022).

12.   Huiting, E. *et al.* Bacteriophages inhibit and evade cGAS-like immune function in

bacteria. *Cell* **186**, 864-876.e21 (2023).

13.   Jenson, J. M., Li, T., Du, F., Ea, C.-K. & Chen, Z. J. Ubiquitin-like conjugation by

bacterial cGAS enhances anti-phage defence. *Nature* **616**, 326–331 (2023).

14.   Leavitt, A. *et al.* Viruses inhibit TIR gcADPR signalling to overcome bacterial defence.

Nature **611**, 326–331 (2022).

15.   Eaglesham, J. B., Pan, Y., Kupper, T. S. & Kranzusch, P. J. Viral and metazoan poxins

are cGAMP-specific nucleases that restrict cGAS–STING signalling. *Nature* **566**, 259–263

(2019).

16.   Zhang, R. *et al.* Homologous 2',5'-phosphodiesterases from disparate RNA viruses

antagonize antiviral innate immunity. *Proc. Natl. Acad. Sci. U. S. A.* **110**, 13114–13119

(2013).

17.   Yirmiya, E. *et al.* Phages overcome bacterial immunity via diverse anti-defence proteins.

Nature **625**, 352–359 (2024).

18.   Athukoralage, J. S. *et al.* An anti-CRISPR viral ring nuclease subverts type III CRISPR

immunity. *Nature* **577**, 572–575 (2020).

19.   Hobbs, S. J., Nomburg, J., Doudna, J. A. & Kranzusch, P. J. Animal and bacterial

viruses share conserved mechanisms of immune evasion. *Cell* **187**, 5530-5539.e8 (2024).

20.   Nomburg, J., Price, N. & Doudna, J. A. *Birth of New Protein Folds and Functions in the*

Virome. http://biorxiv.org/lookup/doi/10.1101/2024.01.22.576744 (2024)

doi:10.1101/2024.01.22.576744.

21.   Sabonis, D. *et al.* TIR domains produce histidine-ADPR conjugates as immune signaling

molecules in bacteria. 2024.01.03.573942 Preprint at

https://doi.org/10.1101/2024.01.03.573942 (2024).

22.   Cao, X. *et al.* Phage anti-CBASS protein simultaneously sequesters cyclic trinucleotides

and dinucleotides. *Mol. Cell* **84**, 375-385.e7 (2024).

23.   Chang, R. B. *et al.* A widespread family of viral sponge proteins reveals specific

inhibition of nucleotide signals in anti-phage defense. 2024.12.30.630793 Preprint at

https://doi.org/10.1101/2024.12.30.630793 (2024).

24.   Li, D. *et al.* Single phage proteins sequester signals from TIR and cGAS-like enzymes.

Nature **635**, 719–727 (2024).

25.   Burroughs, A. M., Zhang, D., Schäffer, D. E., Iyer, L. M. & Aravind, L. Comparative

genomic analyses reveal a vast, novel network of nucleotide-centric systems in biological

conflicts, immunity and signaling. *Nucleic Acids Res.* **43**, 10633–10654 (2015).

26.   Duncan-Lowey, B., McNamara-Bordewick, N. K., Tal, N., Sorek, R. & Kranzusch, P. J.

Effector-mediated membrane disruption controls cell death in CBASS antiphage defense.

Mol. Cell **81**, 5039-5051.e5 (2021).

27.   Gao, L. *et al.* Diverse enzymatic activities mediate antiviral immunity in prokaryotes.

Science **369**, 1077–1084 (2020).

28.   Ledvina, H. E. *et al.* An E1–E2 fusion protein primes antiviral immune signalling in

bacteria. *Nature* (2023) doi:10.1038/s41586-022-05647-4.

29.   Jia, N., Jones, R., Sukenick, G. & Patel, D. J. Second Messenger cA4 Formation within

the Composite Csm1 Palm Pocket of Type III-A CRISPR-Cas Csm Complex and Its Release

Path. *Mol. Cell* **75**, 933-943.e6 (2019).

30.   Chan, C. *et al.* Structural basis of activity and allosteric control of diguanylate cyclase.

Proc. Natl. Acad. Sci. **101**, 17084–17089 (2004).

31.   Anantharaman, V., Iyer, L. M. & Aravind, L. Presence of a classical RRM-fold palm

domain in Thg1-type 3'- 5'nucleic acid polymerases and the origin of the GGDEF and

CRISPR polymerase domains. *Biol. Direct* **5**, 43 (2010).

32.   Kazlauskiene, M., Kostiuk, G., Venclovas, Č., Tamulaitis, G. & Siksnys, V. A cyclic

oligonucleotide signaling pathway in type III CRISPR-Cas systems. *Science* **357**, 605–609

(2017).

33.   Niewoehner, O. *et al.* Type III CRISPR–Cas systems produce cyclic oligoadenylate

second messengers. *Nature* **548**, 543–548 (2017).

34.   Hengge, R., Gründling, A., Jenal, U., Ryan, R. & Yildiz, F. Bacterial Signal Transduction

by Cyclic Di-GMP and Other Nucleotide Second Messengers. *J. Bacteriol.* **198**, 15–26

(2015).

35.   Krüger, L. *et al.* Reversible conjugation of a CBASS nucleotide cyclase regulates

bacterial immune response to phage infection. *Nat. Microbiol.* **9**, 1579–1592 (2024).

36.   Burroughs, A. M. & Aravind, L. Identification of Uncharacterized Components of

Prokaryotic Immune Systems and Their Diverse Eukaryotic Reformulations. *J. Bacteriol.* **202**,

10.1128/jb.00365-20 (2020).

37.   Yirmiya, E. *et al.* Structure-guided discovery of viral proteins that inhibit host immunity.

Cell S0092867424014788 (2025) doi:10.1016/j.cell.2024.12.035.

38.   Rousset, F. *et al.* Phages and their satellites encode hotspots of antiviral systems. *Cell*

Host Microbe **30**, 740-753.e5 (2022).

39.   Penner, M., Morad, I., Snyder, L. & Kaufmann, G. Phage T4-coded Stp: Double-Edged

Effector of Coupled DNA and tRNA-Restriction Systems. *J. Mol. Biol.* **249**, 857–868 (1995).

40.   Jones, J. D. G., Staskawicz, B. J. & Dangl, J. L. The plant immune system: From

discovery to deployment. *Cell* **187**, 2095–2116 (2024).

41.   Remick, B. C., Gaidt, M. M. & Vance, R. E. Effector-Triggered Immunity. *Annu. Rev.*

Immunol. **41**, 453–481 (2023).

42.   Gaidt, M. M. *et al.* Self-guarding of MORC3 enables virulence factor-triggered immunity.

Nature **600**, 138–142 (2021).

43.   Kranzusch, P. J. cGAS and CD-NTase enzymes: structure, mechanism, and evolution.

Curr. Opin. Struct. Biol. **59**, 178–187 (2019).

44.   Aravind, L., Nicastro, G. G., Iyer, L. M. & Burroughs, A. M. The Prokaryotic Roots of

Eukaryotic Immune Systems. *Annu. Rev. Genet.* **58**, 365–389 (2024).

45.   Kibby, E. M. *et al.* Bacterial NLR-related proteins protect against phage. *Cell* **186**, 2410-
2424.e18 (2023).

46.   Gibson, D. G. *et al.* Enzymatic assembly of DNA molecules up to several hundred
kilobases. *Nat. Methods* **6**, 343–345 (2009).

47.   Maffei, E. *et al.* Systematic exploration of Escherichia coli phage–host interactions with
the BASEL phage collection. *PLOS Biol.* **19**, e3001424 (2021).

48.   Kropinski, A. M., Mazzocco, A., Waddell, T. E., Lingohr, E. & Johnson, R. P.
Enumeration of Bacteriophages by Double Agar Overlay Plaque Assay. in *Bacteriophages:*
Methods and Protocols, Volume 1: Isolation, Characterization, and Interactions (eds. Clokie,
M. R. J. & Kropinski, A. M.) 69–76 (Humana Press, Totowa, NJ, 2009). doi:10.1007/978-1-
60327-164-6_7.

49.   Millman, A. *et al.* Bacterial Retrons Function In Anti-Phage Defense. *Cell* **183**, 1551-
1561.e12 (2020).

50.   Baym, M. *et al.* Inexpensive Multiplexed Library Preparation for Megabase-Sized
Genomes. *PLOS ONE* **10**, e0128036 (2015).

51.   Adler, B. A. *et al.* Broad-spectrum CRISPR-Cas13a enables efficient phage genome
editing. *Nat. Microbiol.* **7**, 1967–1979 (2022).

52.   Volkmer, B. & Heinemann, M. Condition-Dependent Cell Volume and Concentration of
Escherichia coli to Facilitate Data Conversion for Systems Biology Modeling. *PLoS ONE* **6**,
e23126 (2011).

53.   Kabsch, W. XDS. *Acta Crystallogr. D Biol. Crystallogr.* **66**, 125–132 (2010).

54.   Liebschner, D. *et al.* Macromolecular structure determination using X-rays, neutrons and
electrons: recent developments in Phenix. *Acta Crystallogr. Sect. Struct. Biol.* **75**, 861–877
(2019).

55.   Mirdita, M. *et al.* ColabFold: making protein folding accessible to all. *Nat. Methods* **19**,
679–682 (2022).

56.   Emsley, P. & Cowtan, K. Coot: model-building tools for molecular graphics. *Acta*

Crystallogr. D Biol. Crystallogr. **60**, 2126–2132 (2004).

Acknowledgements

The authors would like to thank Annette Erbse and the Shared Instruments Pool (RRID: SCR_018986) of the Department of Biochemistry at the University of Colorado Boulder for providing access to the Avanti JXN 26 Super Speed centrifuges and rotors, which are funded by NIH Grant R24OD033699-01, as well as the ITC 200, funded by the NIH Shared Instrumentation Grant S10RR026516; and members of the Whiteley and Morehouse labs for their advice and helpful discussion. We wish to thank the UCI Mass Spectrometry Facility for assistance with small molecule analyses and accurate mass measurements. Use of the Stanford Synchrotron Radiation Lightsource, SLAC National Accelerator Laboratory, is supported by the U.S. Department of Energy, Office of Science, Office of Basic Energy Sciences under Contract No. DE-AC02-76SF00515. The SSRL Structural Molecular Biology Program is supported by the DOE Office of Biological and Environmental Research, and by the National Institutes of Health, National Institute of General Medical Sciences (P30GM133894). This work was funded by the National Institute of General Medical Sciences of the National Institutes of Health under award number R35GM157311 (B.R.M); the National Institutes of Health through the NIH Director's New Innovator award DP2AT012346 (A.T.W.); the PEW Charitable Trust Biomedical Scholars Award (A.T.W.); the Boettcher Foundation Webb-Waring Biomedical Research Award (A.T.W.); the Burroughs Wellcome Fund PATH Award (A.T.W.); a University Colorado ABNexus Grant (joint with A.T.W and Kelly Doran); and a Mallinckrodt Foundation Grant (A.T.W.). A.E.S was supported in part by the NIH Interdisciplinary Predoctoral Training in Molecular Biophysics grant (T32GM145437); K.S. was supported in part by the Undergraduate Research Opportunities Program Individual Grant funded by CU Boulder; and C.R.K.H was supported in part by the Boettcher Foundation Collaboration Grant. U.T. was supported as a fellow of the Cancer Research Institute Irvington Postdoctoral Fellowship (CRI4043). A.N. was supported by Graduate Assistance in Areas of National Need (GAANN) Fellowship #P200A240034 through the University of California Irvine Department of Molecular Biology and Biochemistry. D.S.I. was supported in part by funding through a University of California Irvine Undergraduate Research Opportunities Program Fellowship. A.M.B. and L.A. are supported by the intramural funds of the National Library of Medicine, NIH, DHHS (ZIA LM594244).

Author contributions

Conceptualization, A.E.S., A.T.W., and B.R.M.; Methodology, A.E.S., A.T.W., and B.R.M.; Investigation, A.E.S., A.N., K.S., D.M.D., M.L.D., E.M.Q.E., C.R.K.H., D.S.I., T.A.N., C.M.N., A.M.B., L.A., A.T.W., and B.R.M.; Resources, U.T. and E.M.K.; Writing – Original Draft, A.E.S., A.T.W, and B.R.M.; Writing – Review & Editing, A.E.S., A.T.W, and B.R.M.; Visualization, A.E.S., D.S.I., and B.R.M.; Supervision, A.T.W. and B.R.M; Funding Acquisition, A.T.W. and B.R.M. Note: The order of authors listed 4th–11th is alphabetical by last name.

Declaration of interests

The University of Colorado Boulder and the University of California Irvine have patents pending for OptSE and related technologies on which A.E.S., A.T.W., and B.R.M. are listed as inventors.

Data availability

566 The crystal structure data for OptS have been deposited in the PDB (9MNR). All other data are
567 available in the manuscript or the supplementary materials. Source data are provided.
568
569 **Materials and Methods**
570
571 *Bacterial strains and growth conditions*
572
E. coli strains that were used in this study are listed in **Supplementary Table 1**. All bacterial
cultures were grown in 3.5 mL of media in 14 mL culture tubes shaking at 220 rpm at 37 °C,
unless otherwise indicated. "Overnight" cultures are defined as those that were grown for 16–20
hours following inoculation from a single colony or glycerol stock. Where applicable, culture media
was supplemented with carbenicillin (100 µg/mL) and/or chloramphenicol (20 µg/mL) for plasmid
maintenance or strain selection. *E. coli* OmniPir[45] was used for strain construction and storage of
plasmids and *E. coli* MG1655 (CGSC6300) was used for all phage, colony formation, and
nucleotide extraction experiments.
581
All *E. coli* cultures used for cloning, strain construction, protein expression, and indicated colony
formation assays were grown in LB medium (1 % tryptone, 0.5 % yeast extract, and 0.5 % NaCl).
All strains were frozen for long-term storage in LB plus 30 % glycerol (v/v) at −70 °C. Strains used
to perform phage amplification, phage infection assays, 2′,3′-c-diAMP extraction and
measurement, and indicated colony formation assays were cultivated in "MMCG" minimal medium
(47.8 mM Na_2HPO_4, 22 mM KH_2PO_4, 18.7 mM NH_4Cl, 8.6 mM NaCl, 22.2 mM glucose, 2 mM
$MgSO_4$, 100 mM $CaCl_2$, 3 mM thiamine, Trace Metals at 0.1× (Trace Metals mixture T1001,
Teknova, final concentration: 8.3 µM $FeCl_3$, 2.7 µM $CaCl_2$, 1.4 µM $MnCl_2$, 1.8 µM $ZnSO_4$, 370 nM
$CoCl_2$, 250 nM $CuCl_2$, 350 nM $NiCl_2$, 240 nM Na_2MoO_4, 200 nM Na_2SeO_4, 200 nM H_3BO_3)). When
a strain with two plasmids was cultivated in MMCG medium, bacteria were grown in carbenicillin
(20 µg/mL) and chloramphenicol (4 µg/mL). When growing strains that required induction, 500
µM IPTG or 0.2 % arabinose was used to induce, as appropriate. MMCG and LB agar plates
contain 1.6 % agar and media components described above.
595
596 *Plasmid construction*
597
The plasmids used in this study are listed in **Supplementary Table 1**. Cloning and plasmid
construction were performed as previously described[6]. Briefly, genes of interest were amplified
from phage genomic DNA or previously constructed plasmids using Q5 Hot Start High Fidelity
Master Mix (NEB, M0494L), or were synthesized as FragmentGENEs (Genewiz). Gene inserts
were flanked by at least 18 base pairs of homology to the vector backbone outside of the
restriction digest sites. Ligation of genes into the digested, linearized backbone vector was done
using modified Gibson Assembly[46] with HiFi DNA Assembly Master Mix (NEB, E2621L). Gibson
assemblies were transformed by electroporation into competent OmniPir and plated onto LB (1.6
% agar) plates with appropriate antibiotics to select for successful transformations. Phage genes
and *OptS* with point mutations were generated by amplifying the gene of interest in two parts from
a plasmid template, with the desired mutation occurring in the overlapping region between the
two amplicons. Unless otherwise indicated, all enzymes were purchased from New England
Biolabs.
611
612 For the *OptSE* operon in a pLOCO3 backbone, complete vectors with the indicated operons were
613 generated as ValueGENEs (Genewiz). The pLOCO3 vector (including sfGFP) was initially
614 constructed using Gibson assembly to join and circularize two FragmentGENEs, one with the
615 pLOCO3 backbone, and one with the sfGFP gene (Genewiz; gene fragment sequences listed in
616 **Supplementary Table 1**).
617

618   For all vectors using the pTACxc backbone, pAW1608 was amplified and purified from OmniPir.
619   Purified plasmid was then linearized using BamHI-HF and NotI-HF, or EcoRV-HF and PstI-HF.
620   Gibson ligation was used to circularize the plasmid with the new insert.
621
622   For all vectors using the pBAD30 backbone, pAW1640 was amplified and purified from OmniPir.
623   Purified plasmid was then linearized using EcoRI-HF and NotI-HF. Gibson ligation was used to
624   circularize the plasmid with the new insert.
625
626   For all vectors using the pET16SUMO2 backbone, pAW1123 was amplified and purified from
627   Sure1. Purified plasmid was then linearized using BamHI-HF and NotI-HF. Gibson ligation was
628   used to circularize the plasmid with the new insert.
629
630   Plasmid sequences were verified with Sanger sequencing (Quintara Biosciences, Azenta, and/or
631   Plasmidsaurus). Reads were mapped to the predicted plasmid sequence using the Map to
632   Reference feature of Geneious Prime (default settings).
633
634   *Phage amplification and storage*
635
The phages used in this study are listed in **Supplementary Table 1**. Information on the BASEL
collection used in this study is reported in Maffei et al. 2021[47]. Phage lysates were generated via
plate amplification using a modified double agar overlay[48]. For plate amplification, 400 mL of mid-
log MG1655 were mixed with 3.5 mL MMCG soft agar mix (MMCG with 0.35 % agar and 10 mM
$MgCl_2$, 10 mM $CaCl_2$, and 100 mM $MnCl_2$) and 100-1,000 PFU. Plates were then incubated
overnight at 37 °C. Phages were collected by adding 5 mL of SM buffer (100 mM NaCl, 8 mM
$MgSO_4$, 50 mM Tris-HCl pH 7.5, 0.01 % gelatin) to the plate and incubating for 1 hour at room
temperature. To increase phage titers, the top agar overlay was scraped and harvested along
with the SM buffer. The SM buffer and top agar mixture was centrifuged at 4,000 × g for 10
minutes and the supernatant was transferred to a new tube. The resulting liquid was passed
through a 0.2 mm filter or treated with 2–3 drops of chloroform, followed by vortexing, to remove
viable bacteria. All amplified phages were stored at 4 °C in SM buffer.
648
649   *Phage infection assays*
650
Phage infection assays and phage titer quantifications were performed using a modified double
agar overlay technique[48]. Strains containing the indicated plasmids were cultivated overnight in
MMCG medium (including appropriate antibiotics) and were diluted 1:10 in fresh medium the
following day. The bacteria were grown until they reached mid-logarithmic phase (OD_{600} 0.1-0.8).
400 μL of mid-log bacteria were mixed with 3.5 ml 0.35 % MMCG agar (plus 5 mM $MgCl_2$ and 0.1
mM $MnCl_2$) and poured on top of a MMCG (1.6 % agar) plate. The plate was allowed to cool for
15 minutes. Once cooled, 2 μL of a phage 10-fold serial dilution series was spotted onto the soft
agar overlay and allowed to dry, after which the plates were incubated at 37 °C overnight. Plates
were imaged ~18-24 hr after infection.
660
The resulting phage titer was quantified in PFU/mL for each phage lysate tested. PFU were
enumerated based on the lowest phage dilution spot with individual, quantifiable PFU. The dilution
at that spot was used to calculate the PFU/mL appropriately. When there was a hazy zone of
clearance rather than identifiable plaques, the lowest phage concentration at which this was seen
was counted as ten plaques. When there was no clearance observed, the least dilute spot was
counted as 0.9 plaques, and this was used as the limit of detection for the assay. Phage infection
data was reported as PFU/mL ± standard error of the mean (S.E.M) of n = 3 biological replicates.
668
669   *Phage infection time course in liquid culture*

670
Bacterial strains containing the indicated plasmid were grown overnight in 25 mL MMCG media
plus appropriate antibiotics. Cultures were diluted to an OD_{600} of 0.1 in fresh media without
antibiotics (30 mL total volume) and were grown for two hours at 37 °C with shaking at 220 rpm.
After two hours, the cultures were infected with phage at the indicated MOI. The OD_{600} of the
cultures was measured at the indicated time points after infection. To enumerate PFU at each
time point, 250 µL of culture was harvested and centrifuged at 20,000 × g for 5 min at 4 °C. The
supernatant was transferred to a new tube and 3–5 drops of chloroform were added, followed by
vortexing, to kill any bacteria that remained. The resulting lysates were titered using the phage
infection assay protocol explained above.
680
681 *Escaper phage generation and amplification*
682
T4 escaper phages were generated from three unrelated, clonal T4 lysates ("parents") that were
separately plate amplified on wild-type *E. coli* MG1655. To make the T4 escaper phages
("daughters"), 400 µL of mid-log bacteria expressing the *OptSE* operon (in MMCG plus 100 µg/mL
carbenicillin) was mixed with 100 µL of parent T4 lysate (~4 × 10^5 PFU) and 3.5 mL MMCG top
agar and poured onto a MMCG agar plate. The plate was allowed to dry and was incubated
overnight at 37 °C. The next day, five single escaper plaques were individually isolated from each
parent T4 plate using a Pasteur pipette and soaked in 500 µL SM buffer in an Eppendorf tube. A
dilution series of each T4 escaper phage was spot plated onto *E. coli* MG1655 expressing the
OptSE system to confirm replication in the presence of OptSE. From this plate, single plaques
from each escaper were individually purified and plate amplified (as described in the phage
amplification and storage protocol above) before storage.
694
695 *Phage genome sequencing and escaper analysis*
696
The genomes of the parent and escaper T4 phages were purified as previously described[49]. To
do this, 450 mL of phage lysate (>10^8 PFU/mL) was mixed with 50 µL 10× DNAse I buffer (100
mM Tris-HCl pH 7.6, 25 mM $MgCl_2$, and 5 mM $CaCl_2$) and treated with DNAse I (final concentration
2 × 10^{-3} U/µL) and RNAse A (final concentration 2 × 10^{-2} mg/mL). This mixture was incubated for
1.5 hours at 37 °C to remove extracellular nucleic acids. After, EDTA was added to a final
concentration 20 mM to stop the reaction. Each parent and escaper phage genome was then
isolated and purified using the Qiagen DNeasy cleanup kit, starting at the proteinase K digestion
step[49].
705
The purified phage genomes were prepared for Illumina sequencing using a modification of the
Nextera kit protocol as previously described[50]. Illumina sequencing was performed using a MiSeq
V2 Micro 300-cycle kit (SeqCenter, LLC). The resulting reads were mapped to Genome accession
AF158101.6 using Geneious software's Map to Reference feature. Each read was trimmed to
remove the Nextera adapter sequences before mapping (sequence trimmed:
AGATGTGTATAAGAGACAG) using the "Trim sequences" option; otherwise, Geneious default
settings were used. The trimmed sequences were mapped to the phage genome using default
settings with the "Map to Reference(s)" feature. The Geneious feature "Find Variations/SNPs"
was used to identify variants in daughter phage genomes. Called variants were identified as
escaper mutations if they were present in ≥75% of reads and were not present in parent phage
genomes.
717
Construction of phage gene deletions
T4 knockout phages were generated as previously described[51]. Briefly, 5 mL of *E. coli* MG1655
expressing a pET vector encoding the template for homologous repair were grown in LB to mid-

log phase. Bacteria were then infected with ~4 x 10^8 PFU of T4 and grown for 2–3 hours before
harvesting lysate. Phage lysates amplified in this way were then mixed with 400 µL mid-log
bacteria expressing eLbuCas13a constructs with spacers targeting the gene of interest and
poured onto an MMCG agar plate as in the method described in solid plate amplification detailed
above. Individual plaques were isolated and spot-plated on *E. coli* MG1655 expressing the same
spacer to confirm mutation and to plaque-purify each clone. Target gene deletion was validated
using PCR.

Homologous repair templates encoded 250 bp of homology on either side of the target gene. In-
frame deletions retained the first and last 6 amino acids of the target gene, but deleted the
intervening sequence. Two 31 nt spacers were selected to target the beginning of each gene and
induced as needed using anhydrotetracycline at 5 nM in the top agar of the soft agar overlay.

The *acb1* knockout was constructed with repair template pEK0220 and spacers pEK0223 and
pEK0224. The knockout was PCR verified using the primers oEK0490 and oEK0491.

The *acb2* knockout was constructed with repair template pEK0221 and spacers pEK0225 and
pEK0226. The knockout was PCR verified using the primers oEK0492 and oEK0493.

The double knockout was created by doing the ∆*acb2* knockout steps on a confirmed ∆*acb1*
knockout phage.
738
739 *Colony formation assays for measuring bacterial growth inhibition*
740
Bacterial growth inhibition was tested using colony formation assays. Bacterial strains with
indicated plasmids were grown overnight in MMCG media plus appropriate antibiotics. The
cultures were then 10-fold serially diluted in fresh MMCG media (without antibiotics) and 5 µL of
each dilution was spotted onto an MMCG agar plate containing the appropriate antibiotics, as well
as IPTG (500 µM; induced condition) as indicated. Data in **Fig. 4e** was collected using LB media
with appropriate antibiotics, with or without glucose (0.2% w/v; uninduced condition), IPTG (500
µM; induced condition), or arabinose (0.2% w/v; induced condition), as indicated. After the spotted
bacteria was allowed to dry, plates were incubated at 37 °C for ~16–18 hr for LB plates or ~24 hr
for MMCG plates. Growth inhibition was quantified the next day by counting the number of colony
forming units (CFU) of the lowest dilution that had individual colonies. When no individual colonies
could be counted, the lowest bacterial concentration at which growth was observed was counted
as ten CFU. In instances where no growth was visible, the least dilute spot was counted as 0.9
CFU and used as the limit of detection. Colony formation data was reported as CFU/mL ± S.E.M
of n = 3 biological replicates.
755
756 *Acb2 protein expression and Isothermal titration calorimetry*
757
The vector expressing WT 6xHis-hSUMO2-Acb2 was transformed into Rosetta2 expressing the
pRARE2 plasmid and plated onto 1.6% MMCG agar plates + 100 µg/mL carbenicillin and 20
µg/mL chloramphenicol. An individual colony was picked the following day and inoculated into
100 mL of M9ZB media (47.8 mM Na_2HPO_4, 22 mM KH_2PO_4, 18.7 mM NH_4Cl, 85.6 mM NaCl,
1% Casamino acids (VWR), 0.5% v/v Glycerol, 2 mM $MgSO_4$, Trace Metals at 0.5 × (Trace Metals
Mixture) plus 100 µg/mL carbenicillin and 20 µg/mL chloramphenicol. The culture was then grown
overnight shaking at 37 °C and 220 rpm. The following day, the culture was used to inoculate 2 L
of the same, fresh media to an OD_{600} of 0.05, then grown to an OD_{600} of ~1.5. Cultures were
crash-cooled on ice for 30 minutes before IPTG was added to 500 µM to induce protein
expression. The culture was then moved to a 16 °C shaking incubator and allowed to grow
overnight.
769

Cultures were harvested by centrifugation for 30 minutes at 5,000 rpm and 4 °C in an Avanti JXN-
26 Floor Centrifuge using the JXN 12.500 rotor (Beckman). The resulting pellets were
resuspended in 40 mL Lysis buffer (20 mM HEPES pH 7.5, 400 mM NaCl, 10% v/v Glycerol, 20
mM Imidazole, 0.1 mM Dithiothreitol (DTT)). After resuspension, cells were lysed by sonication at
80% amplitude, with 15 second on, 45 second off pulses for a total processing time of 10 minutes
using a Q500 sonicator (Qsonica). Cellular debris was removed from sonicated lysates by
centrifugation for 45 minutes at 4 °C and 16,000 × g in an Avanti JXN-26 Floor Centrifuge using
the JA 25.50 rotor (Beckman). The soluble lysate was then decanted and protein was purified
using immobilized metal affinity chromatography. Briefly, the soluble lysate was run over 2 mL of
HisPure cobalt slurry (Fisher Sci) equilibrated in Lysis Buffer. The resin was then washed with 2
× 25 mL of Wash Buffer (20 mM HEPES pH 7.5, 1 M NaCl, 10% v/v glycerol, 20 mM Imidazole,
0.1 mM DTT) and protein was eluted in 10 mL of Elution Buffer (20 mM HEPES pH 7.5, 400 mM
NaCl, 10% v/v glycerol, 300 mM Imidazole, 0.1 mM DTT). Proteins were then dialyzed against 2
× 1 L of Dialysis Buffer (20 mM HEPES pH 7.5, 250 mM KCl, 0.1 mM DTT), overnight at 4 °C
using 3.5 kDa MWCO Snakeskin Dialysis Tubing (VWR). The 6×His-SUMO-tag was cleaved
using 6×His-hSENP2 (produced in-house; final concentration of 1:100 hSENP2:protein w/w)
during the overnight dialysis step. After dialysis, proteins were run over 2 mL HisPure cobalt slurry
equilibrated in dialysis buffer to remove any 6×His-SUMO tagged proteins.

After dialysis, the protein was concentrated as needed using 3 kDa MWCO Nanosep spin
concentration columns (Pall Labs) and stored 200–500 µL aliquots in Dialysis Buffer at –70 °C.
Protein concentrations were measured using A_{280} on a Nanodrop OneC (Thermo) and protein
purity was visualized using SDS-PAGE followed by Coomassie staining.
792
ITC assays were adapted from the protocol described in Huiting et al., 2023. Briefly, the K_d, ΔH,
and ΔS for the binding of WT Acb2 with 2′,3′-c-di-AMP were determined using a MicroCal ITC200
calorimeter. Purified Acb2 and 2′,3′-c-di-AMP were dialyzed into the ITC buffer (20 mM HEPES
pH 7.5 and 200 mM NaCl) at 4 °C overnight. The titration was carried out with 27 successive, 1.5
uL injections of 150 µM 2′,3′-c-di-AMP into the sample cell containing 50 µM WT Acb2. Each
injection was spaced by 180 seconds and the cell was kept at 25 °C and stirred at 750 rpm. Origin
software was used for integration and global curve fitting with a single site binding model.
800
801 *AbCap5 protein expression and purification*
802
The vector for expressing 6×His-hSUMO2-AbCap5 was transformed into Rosetta2 expressing
the pRARE2 plasmid and plated onto 1.6 % LB agar plates (plus 100 µg/mL carbenicillin and 20
µg/mL chloramphenicol). The plate was allowed to dry and was incubated at 37 °C overnight. The
next day, an individual colony was picked to inoculate 25 mL of liquid LB media (plus 100 µg/mL
carbenicillin and 20 µg/mL chloramphenicol), which was then grown overnight shaking at 220 rpm
and 37 °C. The following day, the overnight culture was diluted 1:100 into fresh media plus
antibiotics, then grown to an OD_{600} of ~0.6 prior to IPTG addition to a final concentration of 500
µM to induce protein expression. Each culture flask was then moved to a 16 °C shaking incubator
(220 rpm) and allowed to grow overnight.
812
The next day, cultures were harvested by centrifugation for 30 minutes at 5,000 rpm and 4 °C in
an Avanti JXN-26 Centrifuge using the JXN 12.500 rotor (Beckman Coulter). Each bacterial pellet
was resuspended to a total volume of 40 mL lysis buffer (20 mM HEPES pH 7.5, 400 mM NaCl,
10% v/v glycerol, 30 mM imidazole, and 1 mM dithiothreitol (DTT)) plus 1 µL of Pierce Universal
Nuclease (Thermo Fisher Sci.). The resuspended bacterial pellets were kept chilled and were
lysed by sonication at 80% amplitude, with 30 seconds on, followed by 30 seconds off, for a total
processing time of 30 minutes using a Sonicator 4000 (Misonix). The lysed bacteria were

centrifuged at 4 °C for 1 hr at 14,000 xg in a 5910 R tabletop centrifuge (Eppendorf) to pellet
cellular debris left over from sonication.

The resulting soluble lysate was transferred to a new conical tube and kept on ice. To purify
protein, the entire soluble lysate was run over 1 mL of lysis buffer-equilibrated Ni-NTA resin
(Thermo Fisher Sci.). The flow through was collected and reapplied to the resin. The resin was
then washed with 5 × 25 mL of wash Buffer (20 mM HEPES pH 7.5, 1 M NaCl, 10% v/v glycerol,
30 mM imidazole, and 1 mM DTT). The protein was eluted in 10 mL of elution Buffer (20 mM
HEPES pH 7.5, 400 mM NaCl, 10% v/v glycerol, 300 mM imidazole, and 1 mM DTT). The eluted
protein was added to 10 kDa MWCO tubing (VWR) and then dialyzed in 1 L of Dialysis Buffer (20
mM HEPES pH 7.5, 250 mM KCl, and 1 mM DTT) for 1 hr at 4 °C. Afterwards, the dialysis tubing
and protein were placed into 1 L of fresh Dialysis Buffer and allowed to dialyze overnight at 4 °C.
The 6×His-SUMO-tag was cleaved from the N-terminus of AbCap5 using 6×His-hSENP2 (purified
in-house; final concentration of 1:100 hSENP2:protein w/w), which was added to the purified
protein immediately before it was placed in the dialysis tubing. After dialysis and 6x-His-SUMO
cleavage, the purified proteins were applied to 1 mL of dialysis buffer-equilibrated Ni-NTA and the
flow through was collected and reapplied to the resin to remove any uncleaved 6×His-SUMO
tagged proteins.

The purified AbCap5 was concentrated as needed using 30 kDa MWCO Macrosep spin
concentration columns (Pall Labs) and stored in 1 mL aliquots in 50% glycerol v/v at –20 °C. A
Nanodrop OneC (Thermo Fisher Sci.) was used to measure A_{280} to determine the protein
concentration. Protein purity was determined by using SDS-PAGE followed by Coomassie
staining. The AbCap5 protein purified in this way was used in nuclease assays for the
measurement of intracellular 2',3'-c-di-AMP.

Bacterial lysate preparation and nucleotide extraction

Bacterial strains expressing either the *OptSE* operon or an empty vector were grown overnight in
25 mL MMCG plus carbenicillin (100 µg/mL). The following day, the overnight culture was diluted
1:10 into 100 mL (total volume) of fresh media (without antibiotics) and was allowed to grow to an
OD_{600} of ~0.6–0.7 (noting the specific OD_{600} at harvest). Once the appropriate OD_{600} was reached,
the cultures were centrifuged at 4,000 × g for 10 min at 4 °C in a 5910 R tabletop centrifuge
(Eppendorf) to pellet the bacteria. The resulting supernatant was discarded, and the bacterial
pellet was resuspended in 500 µL of lysis Buffer (10 mM Tris-HCl pH 7.5 and 25 mM KCl) and
transferred to a new Eppendorf tube. Hen-lysozyme (0.2 mg/mL final concentration; VWR) and
~200 µL of zirconium beads were added prior to boiling at 95 °C for 2.5 min, and then vortexed
at max speed for 2 min. The samples were boiled and vortexed again and then treated with
proteinase K (0.03 mg/mL final concentration; Qiagen) and 1 µL Pierce Universal Nuclease
(Thermo Fisher Sci.) for 30 min at 37 °C. After treatment, the samples were boiled and vortexed
once again and Triton-X was added to a final concentration of 0.5 % v/v. The samples underwent
a final boil and vortex, and were centrifuged at 17,500 × g for 10 min at 4 °C. The resulting
lysate/supernatant was transferred to a new Eppendorf tube and kept on ice until ready for
nucleotide extraction.

Nucleotides were extracted from bacterial lysates using a modified version of a previously
described phenol-chloroform/chloroform extraction protocol[12]. Briefly, 450 µL of phenol-
chloroform (Thermo Fisher Sci.) was added to 450 µL of bacterial lysate (prepared above),
vortexed for 30 sec, and then centrifuged at 17,500 × g for 45 min at 4 °C. 400 µL of the top
aqueous layer was carefully removed and added to 400 µL of fresh phenol-chloroform in a new
tube. This mixture was vortexed for 30 seconds and centrifuged at 17,500 × g for 10 min at 4 °C.
350 µL of the resulting top aqueous layer was carefully removed and added to 350 µL of

chloroform (VWR) in a new tube. The mixture was vortexed again for 30 sec and centrifuged one
final time at 17,500 × g for 10 min at 4 °C. The resulting top aqueous layer was removed and
placed in a new tube and for storage at –20 °C. Nucleotide extraction from indicated strains was
carried out for a total of n = 3 biological replicates.
876
877
878 *Intracellular 2′,3′-c-di-AMP measurements*
879
Commercially available 2′,3′-c-di-AMP (Enzo) was used to make a standard curve by 2-fold
serially diluting the nucleotide in nuclease-free water. A dilution series of nucleotide extracts from
the tested bacterial strains were made by 2-fold serially diluting the extracts in nuclease-free
water. For each 25 µL reaction, AbCap5 (final concentration 50 nM) was incubated with 500 ng
of linear PCR-amplified DNA and 5 µL of one dilution of nucleotide extract, 2′,3′-c-di-AMP, or
water in Reaction Buffer (final concentrations: 10 mM Tris-HCl pH 7.5, 25 mM KCl, and 10 mM
$MgCl_2$). The mixture was incubated for 2 hrs at 37 °C, followed by a 5 min boil at 95 °C. Reactions
were mixed with 5 µL of 6× Gel Loading Dye plus SDS (NEB) and 20 µL were loaded and
visualized on a 2% (w/v) agarose gel stained with SYBR Safe (Thermo Fisher Sci.).
889
890 The concentration of intracellular 2′,3′-c-di-AMP ([n.t.]) was calculated using the following
891 equation:
892

$$[n.t.] = \frac{(SC) * (lysate\ volume)}{(CV) * (OD_{600}) * (culture\ volume)}$$

894
SC is the upper or lower 2′,3′-c-di-AMP standard curve concentration (in [M]) that corresponds to
a similar DNA degradation amount as the last visible nucleotide extract dilution reaction (from
OptSE-expressing cells) that contains non- or partially-degraded DNA PCR product, respectively.
Lysate volume is the volume of Lysis Buffer (in L) that was used to resuspend the bacterial pellet
before the lysis steps. CV is the OD_{600}-specific cell volume (in L * OD^{-1} * mL^{-1}) of the bacteria
based on the *E. coli* strain and growth conditions[52]. OD_{600} is the optical density of the bacterial
culture at the time of harvest (before pelleting). Culture volume is the volume of bacterial culture
(in mL) at the time of harvest. Intracellular [2′,3′-c-di-AMP] reported as an average for the upper
and lower range of concentrations (in nM) for n = 3 biological replicates of the nucleotide
extraction.
905
906 *Recombinant protein expression and purification*
907
The genes for full-length *OptS* from *Klebsiella pneumoniae* strain KP67, and bacteriophage T4
Acb2 were codon optimized and synthesized as dsDNA fragments (gBlocks; Integrated DNA
Technologies). The genes were cloned using Gibson assembly into a linearized (restriction
enzyme digested with BamHI and NotI) in-house pET16 expression vector modified to allow in-
frame cloning of an N-terminal hexa-histidine tag with or without a hSUMO2 tag with a Gly-Ser
linker. After Sanger sequencing confirmation, the resulting plasmids were subsequently
transformed into BL21-RIL *E. coli* cells (BL21 derivative, DE3; Invitrogen, Life Technologies) for
protein expression.
915
Transformant colonies were grown overnight at 37 °C on MDG media (0.5 % glucose, 25 mM
Na_2HPO_4, 25 mM KH_2PO_4, 50 mM NH_4Cl, 5 mM Na_2SO_4, 2 mM $MgSO_4$, 0.25 % aspartic acid and
trace metals) plates supplemented with 100 µg/mL ampicillin and 34 µg/mL chloramphenicol for
selection. Single colonies were used to inoculate liquid MDG media cultures (same components
without agar) which were grown overnight ~16–20 hours at 37 °C and 230 rpm shaking. Overnight
cultures were used to inoculate one-liter of liquid M9ZB expression media (0.5 % glycerol, 1 %
Cas-amino acids, 47.8 mM Na_2HPO_4, 22 mM KH_2PO_4, 18.7 mM NH_4Cl, 85.6 mM NaCl, 2 mM

MgSO$_4$ and trace metals) contained in 2.5-liter flasks. After cultures reached an OD$_{600nm}$ >2.5,
flasks were placed on ice for 15 minutes followed by addition of 0.5 mM IPTG (final) and then
incubated at 16 °C with 230 rpm shaking overnight (~16–20 hours). Cells were harvested by
centrifugation, washed with PBS buffer, and flash frozen with liquid N$_2$ and stored at −80 °C until
needed.

Conventional nickel-affinity chromatography was carried out using gravity flow at 4 °C. Briefly, *E.*
coli cell pellets were resuspended with lysis buffer (20 mM HEPES-KOH pH 7.5, 400 mM NaCl,
10% glycerol, 30 mM Imidazole pH 7.5, 1 mM DTT) and subjected to sonication to release cellular
contents (10 sec on, 20 sec off, 70% amplitude, 5 min total on time). The lysate was clarified with
centrifugation and the supernatant loaded onto 4–6 mL of packed Ni-NTA resin pre-equilibrated
with lysis buffer. The column was then sequentially washed with 20 mL of lysis buffer, 70 mL of
wash buffer (20 mM HEPES-KOH pH 7.5, 1 M NaCl, 10% glycerol, 30 mM Imidazole pH 7.5, 1
mM DTT), and a final 35 mL of lysis buffer to remove high salt. Protein was eluted with 20 mL of
elution buffer (20 mM HEPES-KOH pH 7.5, 400 mM NaCl, 10% glycerol, 300 mM Imidazole pH
7.5, 1 mM DTT). The eluent was dialyzed overnight in 20 mM HEPES-KOH pH 7.5, 250 mM KCl,
and 1 mM DTT at 4°C with gentle stirring. In the case of Acb2, hSENP2 protease (D364–L589,
M497A) was added to the dialysis tubing to cleave the N6xHis-SUMO2 tag. The dialyzed eluent
was concentrated and further purified via size exclusion chromatography (SEC) using a HiLoad™
16/600 Superdex™ 200 pg column (Cytiva) equilibrated with 20 mM HEPES (pH 7.5), 250 mM
KCl, and 1 mM TCEP-KOH. SEC fractions were analyzed with SDS-PAGE and fractions that
contained recombinant protein of interest were pooled and concentrated to >10 mg/mL and were
flash frozen in liquid N$_2$ followed by storage at −80 °C until needed.

Protein crystallization and structure determination

Apoprotein crystals for OptS from *Klebsiella pneumoniae* strain KP67 were obtained at room
temperature by mixing 1 µL of reservoir solution consisting of 8 % (w/v) PEG 4000 with an equal
volume of 7 mg/mL of the protein solution using the hanging-drop vapor diffusion method. The
crystals were soaked in mother liquor with an additional 15 % ethylene glycol as cryoprotectant
and subsequently plunged into liquid N2 and shipped for remote data collection.

X-ray diffraction data for *Kp*OptS were collected at experimental station 12-2 at the Stanford
Synchrotron Radiation Lightsource (SSRL) using a Dectris Pilatus® 16M PAD detector. Beamline
12-2 was set at 0.97946 Å and the crystal data was collected at 100 K. The data set was
processed using the X-ray Detector Software (XDS) package[53] that is incorporated into the
automated processing pipeline used at SSRL. The structure was determined using molecular
replacement conducted by the Phaser-MR program within the PHENIX suite[54] using a predicted
structural model of *Kp*OptS generated by ColabFold v1.5.5 which uses a homology search by
MMseqs2 with AlphaFold2[55]. The structure of *Kp*OptS was iteratively refined using the
phenix.refine program and the residue positions were manually adjusted in Coot[56]. The final
refined structure had the following Ramachandran statistics- 98.09 % favored, 1.91% allowed,
0.00% outlier. The full data collection and refinement statistics are summarized in **ED Table 1**.
The models were visualized and analyzed using PyMOL (The PyMOL Molecular Graphics
System, version 3.1.3, Schrödinger, LLC). The atomic coordinates have been deposited to the
Protein Data Bank (PDB ID: 9MNR).
964

A*nalysis of enzyme activity with high-performance liquid chromatography*

Enzymatic reactions were performed in 100–600 µL total volume with typically 250 µM total
equimolar ribonucleotide triphosphates (ATP, GTP, CTP, UTP), 1 mM MnCl$_2$, 10 mM MgSO$_4$, 20
mM HEPES-KOH pH 7.4, 100 mM NaCl, and 100 µM OptS protein added last. Reactions were
transferred to a 10 kDa-cutoff filter and subjected to centrifugation for 10 minutes at 13,500 x g.

When appropriate, purified Acb2was added at 250 µM and incubated for an additional 1 hr at 37
°C. 1 µL of ~20 mg/mL Proteinase K (NEB, cat no. P8107S) was added to degrade Acb2 and
release cyclic dinucleotide before filtering. Likewise, samples were at times treated with P1
nuclease and/or calf-intestinal phosphatase as indicated. All samples were injected at 10 µL onto
an Agilent 1200 Series HPLC equipped with a 4.6 × 150 mm and 5 µm particle-size Zorbax Bonus-
RP C18 column using an isocratic elution method including 97% 50 mM NaH_2PO_4 pH 6.8 and 3%
acetonitrile buffer system held at 40 °C. Nucleotide separation was monitored at 254 nm using a
multiwavelength detector. For mass spectrometry analysis, enzymatic reactions were prepared
using 20 mM ammonium acetate pH 8.0 instead of HEPES-KOH and NaCl but otherwise treated
the same with or without an LC fractionation step to isolate specific peaks. For fractionation on
HPLC, temperature was kept at 23 °C with a gradient elution method as follows: Solvent A- 20
mM ammonium acetate pH 8.0; Solvent B- Methanol. 0–2 min 100% A, 2–8 min 0–100% B, 8–10
min 100% B, 10–11 min 0–100% A, 11–17 min 100% A. ~1 mL fractions were collected and
concentrated via speed-vac (1–3 hr, 30 °C) prior to injection on a Waters Acquity UPLC-QDA
instrument equipped with a C18 50 mm UPLC column and single quadrupole MS detector (QDA;
m/z 50-1250) and photodiode array detector (PDA; 200-500 nm). Standard gradient elution was
used from 0–100% 0.1 % formic acid:acetonitrile over 5 minutes. In ESI+ mode, analyte m/z ions
(M+H)+ or (M+NH3)+ were validated to less than ± 5 ppm relative to the nearest sodiated
polyethylene glycol (CAS: 25322-68-3, av. Mwt 400)) or sodiated methoxypolyethyleneglycol
(CAS: 990-74-4, av. Mwt 350) calibrant peak lockmass.

Synthetic nucleotide ligands

Synthetic cyclic dinucleotide ligands used for HPLC and mass spectrometry analysis were
purchased from Biolog Life Science Institute: 3′,3′-c-di-AMP (cat no. C 088), 2′,3′-c-di-AMP (cat
no. C 187), 2′,2′-c-di-AMP (cat no. C 188), 3′,3′-c-di-GMP (cat no. C 057), 2′,3′-c-di-GMP (cat no.
C 182), 2′,2′-c-di-GMP (cat no. C 162), 3′,3′-cGAMP (cat no. C 117), 2′,3′-cGAMP (cat no. C 161),
3′,2′-cGAMP (cat no. C 238), 2′,2′-cGAMP (cat no. C 210), pppA(2′,5′)pA (cat no. T 073).

Accession numbers

The structure of *Kp*OptS has been deposited in the Protein Data Bank (PDB ID: 9MNR). All
other relevant accession numbers can be found in **Supplementary Table 1C**.

Statistics and reproducibility
1000
Each experiment presented was performed in independent biological duplicate or triplicate using
bacterial cultures grown on two or three separate days, respectively. Data was plotted using
Graphpad Prism 9 at an n = 2 or n = 3 with error bars indicating the standard error of the mean.
Illumina sequencing results were analyzed using Geneious Prime Software.

Figure 1. The Panoptes system is antiphage
(A) Domain architectures of Panoptes operon genes, *optS* and *optE.*
(B) Heatmap of fold defense provided by the Panoptes system for a panel of diverse phages
from the BASEL collection. *E. coli* expressing the Panoptes system were challenged with
phages and fold defense was calculated for each phage by dividing the efficiency of plating (in
PFU/mL) on empty vector by the efficiency of plating on Panoptes system-expressing bacteria.
An "X" represents uncountable plaques. Family names are above each indicated set of phages.
(C) Efficiency of plating of indicated phages infecting *E. coli* expressing a plasmid with either the
Panoptes system or an empty vector. Data represent the mean ± standard error of the mean
(SEM) of n = 3 biological replicates (except for BASEL45, n=1), shown as individual points.

a

b

c

d

PleD

GGDEF domain

PDB 1W25

e

Csm1

Cas10 (Type III CRISPR)

PDB 6O75

Figure 2. OptS is a minimal CRISPR polymerase
(A) Size exclusion chromatography reveals a monodisperse signal for *Kp*OptS with an elution
volume consistent with an oligomer of at least a trimer. Grey dots indicate molecular weight
standards: Blue dextran, 46.2 mL, 2000 kDa; Ferritin, 59.8 mL, 440 kDa; Aldolase, 71.0 mL, 158
kDa; Conalbumin, 78.9 mL, 75 kDa; Ovalbumin, 85.3 mL, 43 kDa; Carbonic anhydrase, 91.5
mL, 29 kDa; Ribonuclease A, 98.6 mL, 13.7 kDa; Aprotinin, 104.4 mL, 6.5 kDa.
(B) Crystal structure of the *Kp*OptS apoprotein reveals a tetrameric architecture. Two rotated
views are depicted with each protomer shown in a different color.
(C) Structure of the isolated protomer (monomer) highlighting overall domain fold.
(D) Superposition of GGDEF-containing enzyme PleD (PDB 1W25, light blue) with *Kp*OptS.
(E) Superposition of Type III Cas10 CRISPR polymerase palm domain (PDB 6O75, light
orange) with *Kp*OptS.
1027

Figure 3. OptS is capable of synthesizing 2′,3′-c-di-AMP
(A) HPLC analysis of an ATP chemical standard compared with the product of *Kp*OptS when
incubated with ATP alone. Traces representative of at least three replicates.
(B) HPLC analysis of 2',2'-c-di-AMP, 2',3'-c-di-AMP, and 3',3'-c-di-AMP chemical standards
compared with the product of *Kp*OptS. Traces representative of at least three replicates.
(C) Magnified view of the *Kp*OptS putative active site highlighting catalytic Asp6 and Asp57
(orange).

(D) HPLC analysis of the reaction products when wild type or mutant *Kp*OptS is incubated with
ATP, compared to an ATP chemical standard. Traces representative of at least three replicates.
(E) Chemical structure of 2',3'-cyclic di-adenosine monophosphate (2',3'-c-di-AMP).
1038

Figure 4. Acb2 is necessary and sufficient for activating the Panoptes system
(A) Schematic of phage T4 escape mutations from this paper (top) and a previously validated
binding site mutation (bottom).
(B) Efficiency of plating of T4 phages with the indicated genotype on *E. coli* expressing an
empty vector or the Panoptes operon from a plasmid. Images are representative of n=3
biological replicates.
(C) Colony formation of *E. coli* expressing an empty vector or Panoptes on a low copy plasmid
and indicated empty vector or Acb2 allele on an IPTG-inducible plasmid. Data represent the
mean ± SEM of n = 3 biological replicates, shown as individual points.
(D) Isothermal titration calorimetry (ITC) to test binding of 2′,3′-c-di-AMP to Acb2. The individual
data points of three technical replicates are shown. The K_d, ΔH, and ΔS were determined by
generating a global fit for the three indicated replicates. Raw data for these curves are shown in
Extended Data Figure 5.
(E) Colony formation of *E. coli* co-expressing indicated genes from IPTG-inducible (Plasmid 1)
and arabinose-inducible (Plasmid 2) plasmids. Data represent the mean ± SEM of n = 3
biological replicates, shown as individual points.
1055

Figure 5. Phage anti-defense proteins that antagonize phage defense activate Panoptes
(A) Colony formation of *E. coli* expressing an empty vector or the Panoptes operon on a low
copy plasmid and indicated empty vector or phage protein on an IPTG-inducible plasmid. Data
represent the mean ± SEM of n = 3 biological replicates, shown as individual points.
(B) Efficiency of plating of T4 WT or T4 Δ*acb2* phage infecting *E. coli* expressing a plasmid with
either an empty vector, Panoptes operon, or CBASS operon. Data represent the mean ±
standard error of the mean (SEM) of n = 3 biological replicates, shown as individual points.

1064 **Extended Data Figure 1.** Panoptes protects against phage in liquid culture
1065 **(A)** Above: growth curve of *E. coli* expressing the indicated plasmid. Arrows indicate the time
1066 each culture was infected with phage T4 at the indicated multiplicity of infection (MOI). Below:
1067 efficiency of plating of the phage present in each sample at the indicated time points. Data are
1068 representative of n = 2 biological replicates.
1069
1070

a

OptS1

OptS2

Dimer 1 Interface

b

OptS1   OptS3

Dimer 2 Interface

c

OptS1   OptS4

Dimer 3 Interface

1071 **Extended Data Figure 2.** The *Kp*OptS tetramer contains three dimers (pair of protomers)
1072 interfaces
1073 **(A)** Dimer 1 interface
1074 **(B)** Dimer 2 interface
1075 **(C)** Dimer 3 interface
1076

Extended Data Figure 3. Characterization of *Kp*OptS enzymatic activity *in vitro*
(A) HPLC analysis of NTP chemical standards compared with the product of *Kp*OptS when
incubated with a mixture of NTPs. Traces representative of at least three replicates.
(B) HPLC analysis of a GTP chemical standard compared with the product of *Kp*OptS when
incubated with GTP alone. Traces representative of at least three replicates.
(C) HPLC analysis of ATP and GTP chemical standards compared with the product of *Kp*OptS
when incubated with ATP and GTP. Traces representative of at least three
replicates.
(D) HPLC analysis of 2',2'-cGAMP, 2',3'-cGAMP, 3',2'-cGAMP, and 3',3'-cGAMP chemical
standards compared with the product of *Kp*OptS. Traces representative of at least three
replicates.
(E) P1 nuclease and CIP treatment of *Kp*OptS ATP-only product. Traces representative of at
least three replicates.
(F) A less magnified view of the putative catalytic site highlighting catalytic Asp6 and Asp57
(orange)
(G) MS spectra of a 2',3'-c-di-AMP standard (bottom) and the *Kp*OptS product (top).
1093

Extended Data Figure 4. Phage T4 escapers evade Panoptes-mediated defense
(A) Efficiency of plating of T4 parent and escaper phages on *E. coli* expressing an empty vector
or the Panoptes operon from a plasmid. Images are representative of n=3 biological replicates.
(B) Efficiency of plating of T4 parent phages and corresponding escaper mutant phages
(Escaper, T4.X.Y, where X indicates the parent phage number and Y indicates the escaper
phage number) on *E. coli* expressing an empty vector or the Panoptes operon from a plasmid.
Data represent the mean ± standard error of the mean (SEM) of n = 3 biological replicates,
shown as individual points.
(C) Efficiency of plating of T4 phages with the indicated genotype on *E. coli* expressing an
empty vector or the Panoptes operon from a plasmid. Data represent the mean ± standard error
of the mean (SEM) of n = 3 biological replicates, shown as individual points.
1105

(A) HPLC analysis of the ability of Acb2 to bind the main products of *Kp*OptS (ATP and GTP
reaction). Traces representative of at least three replicates.
(B) HPLC analysis of the ability of Acb2 to bind and release the product of *Kp*OptS when treated
with proteinase K (ATP only reaction). Traces representative of >3 replicates.
(C) HPLC analysis of the ability of mutant Acb2 to bind the ATP-derived product of *Kp*OptS.
Traces representative of at least three replicates.
(D-F) Replicates of ITC assays to test binding of 2′,3′-c-di-AMP to Acb2.
1114

Extended Data Figure 6. *Ab*Cap5 is a sensor of 2',3'-c-di-AMP
(A) Visualization of DNA degradation by *Ab*Cap5 when incubated with either 2',3'-c-di-AMP,
3',3'-c-di-AMP, 2',2'-cGAMP, 2',3'-cGAMP, 3',2'-cGAMP, or 3',3'-cGAMP. Data are
representative images of n=2 technical replicates.
(B) Visualization of DNA degradation by *Ab*Cap5 when incubated with a dilution series of 2',3'-c-
di-AMP or extracted nucleotides from *E. coli* expressing an empty vector or the Panoptes
operon. Data are representative images of n=3 biological replicates.

B

Pairwise overlap counts for systems and components, n=13710

	R-M	CRISPR (Cas1)	CARF	CATRA	GSDM	SMODS	S-2TMB	mCpol
R-M	79.30%	31.12%	10.07%	0.34%	0.53%	12.76%	2.23%	0.39%
CRISPR (Cas1)		36.18%	7.96%	0.08%	0.18%	4.75%	0.77%	0.22%
CARF			12.33%	0.04%	0.07%	1.80%	0.34%	0.17%
CATRA				0.36%	0	0.10%	0	0.01%
GSDM					0.62%	0.13%	0.04%	0
SMODS						14.48%	2.22%	0.23%
S-2TMB							2.58%	0.18%
mCpol								0.42%

p-value results for pairwise Fisher's exact test

	R-M	CRISPR (Cas1)	CARF	CATRA	GSDM	SMODS	S-2TMB	mCpol
R-M	0	1.236e-50	0.010	7.762e-03	0.282	2.656e-28	5.301e-04	8.335e-03
CRISPR (Cas1)		0	8.918e-142	0.039	0.141	7.109e-04	9.880e-03	0.019
CARF			0	0.829	0.742	0.854	0.567	1.249e-07
CATRA				0	1.000	0.013	0.641	0.191
GSDM					0	0.088	0.069	1
SMODS						0	3.574e-208	3.715e-12
S-2TMB							0	6.891e-25
mCpol								0

Extended Data Figure 7. mCpol distribution and gene co-occurrences

(A) Heat map depicting genome presence/absence of selected conflict systems in prokaryotic lineages.

(B) Tables depicting percentages of genome occupancy and pairwise co-occupancy for selected systems across prokaryotes (top) and p-values representing the significance of pairwise co-occurrences from exact Fisher's tests (bottom). Co-occurrences with p-values of less than 1e-05 are shaded in red.

(C) Venn diagram depictions of genome overlap between selected pairs, significant p-values are provided below the diagram where applicable.

1132   **(D)** Conserved mCpol gene neighborhood associations, with individual genes depicted as boxed
1133   arrows. "Core" mCpol-containing systems are separated from discrete associating systems by
1134   blue circles.
1135

List of documents and display items:

Cover letter
Manuscript

Figures:

Figure 1. The Panoptes system is antiphage
Figure 2. OptS is a minimal CRISPR polymerase
Figure 3. OptS is capable of synthesizing 2′,3′-c-di-AMP
Figure 4. Acb2 is necessary and sufficient for activating the Panoptes system
Figure 5. Phage anti-defense proteins that antagonize phage defense activate Panoptes

Extended Data Figures:

Extended Data Figure 1. Panoptes protects against phage in liquid culture
Extended Data Figure 2. The *Kp*OptS tetramer contains three dimer interfaces
Extended Data Figure 3. Characterization of *Kp*OptS enzymatic activity *in vitro*
Extended Data Figure 4. Phage T4 escapers evade Panoptes-mediated defense
Extended Data Figure 5. Acb2 binds the product of *Kp*OptS
Extended Data Figure 6. *Ab*Cap5 is a sensor of 2',3'-c-di-AMP
Extended Data Figure 7. mCpol distribution and gene co-occurrences

Extended Data Tables:

Extended Data Table 1. X-ray crystallography data collection and refinement statistics
Extended Data Table 2. T4 phage escaper mutations

Supplementary Information:

Supplementary Table 1. Bacterial strains, plasmids, phages, and proteins used in this study