Learning interpretable representation for context-specific transcription regulatory networks using a foundation model

Zhaowei Yu[1,2], Dongxu Yang[1,2], Qianqian Chen[1,2], Yuxuan Zhang[1,2], Zhanhao Li[1], Yucheng Wang[1], Chenfei Wang[1], Yong Zhang[1*]

[1]State Key Laboratory of Cardiovascular Diseases and Medical Innovation Center, Institute for Regenerative Medicine, Department of Neurosurgery, Shanghai East Hospital, Shanghai Key Laboratory of Signaling and Disease Research, Frontier Science Center for Stem Cell Research, School of Life Sciences and Technology, Tongji University, Shanghai, 200092, China

[2]These authors contributed equally to this work

*Correspondence: Yong Zhang Ph.D., 1239 Siping Road, Shanghai 200092, China; Tel: 86-21-65981196; Fax: 86-21-65981041; E-mail: yzhang@tongji.edu.cn

Abstract

Gene expression is shaped by transcription regulatory networks (TRNs), where transcription regulators interact within regulatory elements in a context-specific manner. Despite significant efforts, understanding the intricate interactions of transcription regulators across different genomic regions and cell types remains a major challenge, largely due to data sparsity. Here, we introduce ChromBERT, a foundation model pre-trained on large-scale human ChIP-seq datasets. ChromBERT effectively captures the interaction syntax of approximately one thousand transcription regulators across diverse genomic contexts, generating interpretable representations of context-specific TRNs and their constituent regulators. Fine-tuned for various transcription regulation tasks, ChromBERT demonstrates superior performance in imputing previously unseen cistromes and modeling regulatory effects and dynamics in specific cell types. Notably, ChromBERT adapts its TRN representations for cell-type-specific tasks, providing deep insights into the roles of transcription regulators underlying observed regulatory effects or dynamics without requiring cell-type-specific genomic data for each regulator. By

31   overcoming the limitations of sparse transcription regulator data, ChromBERT

32   significantly enhances our ability to model, interpret, and predict transcriptional regulation

33   across diverse biological contexts.

Introduction

Gene expression patterns are shaped by the intricate interactions among transcription

regulators, regulatory elements, and target genes, collectively forming transcription

regulatory networks (TRNs)[1-4]. A central goal in biology is to understand how transcription

regulators interact within specific genomic contexts, how distinct regulatory architectures

are formed in different genomic regions, and how these interactions vary across diverse

cell types. These interactions underlie cell-type-specific transcriptional programs, which

are critical for maintaining cellular identity and function. Landmark projects like the

Encyclopedia of DNA Elements (ENCODE) have provided invaluable insights into

transcription regulatory landscapes by generating extensive data on transcription factor

binding and chromatin states[2,4]. However, these efforts have primarily focused on a

limited number of well-studied cell types. Consequently, the context-dependent

interactions of transcription regulators and their hierarchical roles in modulating gene

expression remain poorly understood in most biological settings, particularly in dynamic

processes such as cellular transitions and disease progression.

Pre-trained foundation models have achieved remarkable success in genomics studies

by leveraging large-scale heterogeneous datasets to learn fundamental knowledge that

can be fine-tuned for a wide range of downstream tasks[5-13]. These models, often

implemented as deep language models, adept at data-scarce scenarios where traditional

approaches struggle, offering both the improvement on predictive performance and the

ability to capture biologically meaningful representations. Specifically, the representations

learned during pre-training can be interpreted using post hoc analysis or by-design

strategies to uncover valuable biological insights[14]. Fine-tuning expands this capability by

enabling these models to adjust their learned representations for specific tasks,

generating biological insights that are directly relevant to the given fine-tuning

objectives[6,10,15]. By combining broad generalization from pre-training with task-specific

adaptability through fine-tuning, pre-trained foundation models are particularly suitable

for representing and interpreting the complex dynamics of TRNs, especially in the settings

where transcription regulator landscapes are highly incomplete and poorly characterized.

Foundation models pre-trained on DNA sequences have demonstrated remarkable

capabilities in deciphering the regulatory architecture of genomes. They effectively

capture the organizational syntax of genomic sequences and decode the *cis*-regulatory

grammar that governs gene transcription[7,8,11,12,16]. However, these models often fall short

in addressing the roles of *trans*-acting transcription regulators, which are essential for a

comprehensive understanding of gene transcription. A recent study seeks to bridge this

gap by integrating DNA-binding motifs of transcription regulators and chromatin

accessibility to improve gene expression predictions[10], and another study also integrates

motifs with multi-omics data to infer gene regulatory architecture[17]. However, these

approaches are limited as accessible motifs account for only a fraction of actual binding

events, and many transcription regulators lack well-characterized motifs[18,19]. Binding

events are often modulated by co-factors, chromatin context, and epigenetic

modifications, which extend beyond simple motif recognition, influencing how and where

transcription factors bind to their target regions. Consequently, motif-dependent

approaches are insufficient to capture the intricate and context-specific interactions

among transcription regulators, highlighting the need for foundation models that directly

represent and learn these complex interactions.

Here, we introduce ChromBERT, a foundation model specifically designed to directly

model genome-wide combinatorial binding patterns of transcription regulators. Pre-

trained on the Cistrome-Human-6K dataset, comprising large-scale, heterogeneous

ChIP-seq data, ChromBERT effectively captures the interaction syntax of transcription

regulators across diverse genomic contexts (Fig. 1a). By leveraging advanced

transformer architectures, ChromBERT generates context-specific, biologically

interpretable representations of TRNs and their constituent transcription regulators,

enabling precise biological interpretation of regulatory roles and the functional

collaborations of these regulators. ChromBERT demonstrates superior performance in

cistrome imputation for unseen cell types through prompt-enhanced fine-tuning,
outperforming previous methods, and showcasing robust generalization across a diverse
range of specific cell types. More importantly, when fine-tuned for cell-type-specific
downstream tasks, ChromBERT not only achieves superior predictive performance but
also adapts its representations to reflect the unique regulatory architecture of each cell
type. This adaptability provides clear, interpretable insights into the roles of transcription
regulators in driving observed regulatory effects and dynamic changes, without requiring
additional cell-type-specific genomic data for each regulator (Fig. 1b). This capability is
particularly significant in addressing the challenges posed by sparse transcription
regulator data, a common limitation in most biological settings, where comprehensive
datasets for transcription regulators are often unavailable.

Results

Overview of ChromBERT

ChromBERT distinguishes itself from other genomics foundation models that primarily
focus on DNA sequence organization[7,8] by modeling the combinatorial binding patterns
of thousands of transcription regulators and learning their intricate interaction syntax (Fig.
1a). To enable this, ChromBERT was pre-trained on a large-scale corpus of transcription
regulator landscapes, leveraging self-attention mechanisms within a transformer
architecture[20] to learn how transcription regulators interact with others at each genomic
region. Using a masked learning strategy akin to BERT[21], the model captures context-
dependent interactions of transcription regulators across diverse genomic contexts in a
self-supervised manner. These pre-training techniques equip ChromBERT with a
foundational understanding of TRNs, enabling strong generalization across a variety of
downstream transcription regulation tasks.

The pre-training dataset for ChromBERT was constructed by compiling and filtering nearly
all publicly available human ChIP-seq, DNase-seq, and ATAC-seq data from the Cistrome
Data Browser[22] (see Methods for details). This process resulted in the Cistrome-Human-
6K dataset, comprising 6,391 qualified cistromes (genome-wide maps of the *cis-*

regulatory binding sites of *trans*-acting regulators[23]) for 991 transcription regulators, 76

histone modifications, and chromatin accessibility from various cell types (Fig. 1a,

Supplementary Table. S1). Binding signals from all cistromes were discretized into

binding statuses, which were then aggregated for each 1-kilobase (kb) genomic interval

to construct input features for ChromBERT (Supplementary Fig. S1a, b). This approach

enabled the model to learn context-dependent interactions among transcription regulators.

Given the limited number of cistromes in individual cell types (Supplementary Fig. S1c),

we pooled all qualified human cistromes from different cell types to enrich the diversity of

input features. While this cell-type-agnostic strategy introduces variability in

representation frequencies for transcription regulators and cell types, it maximizes the

use of all available data and supports the model's ability to generalize. Pre-training on

over two million such 1-kb genomic intervals enhanced ChromBERT's capacity to capture

intricate interaction patterns and achieve strong awareness to different genomic contexts.

For each genomic region presented to ChromBERT, the pre-trained model embeds it into

a TRN embedding that represents the combinatorial interactions among over a thousand

transcription regulators within the given region (Fig. 1a). Each regulator's embedding,

encoded as a 768-dimensional vector within the TRN embedding, represents its

contextual interactions with other regulators in the same genomic region.

Pre-trained embeddings reveal human regulatory architecture

After pre-training, we interpreted the pre-trained embeddings from ChromBERT to check

whether they could effectively reveal the transcription regulatory architectures specific to

their genomic regions. Given the diversity in transcription regulatory architectures across

different genomic regions, we hypothesized that pre-trained TRN embeddings would

cluster in similar patterns for different groups of regulatory elements. Applying

ChromBERT to different groups of regulatory elements on chromosome one, as defined

by ChromHMM states[24], we observed that TRN embeddings generally clustered

separately (Supplementary Fig. S2a), suggesting that they effectively delineate distinct

regulatory architectures. We also hypothesized that TRN embeddings of nearby genomic

regions would display higher similarity due to shared local chromatin structure and

transcription regulators. As expected, we noted that the similarity in TRN embeddings
decreased with increasing distance between regions (Supplementary Fig. S2b).

Considering the impact of 3D chromatin organization, which segregates the genome into
potential functional units[25], we examined the association between TRN embeddings and
3D genome architecture. Our results showed that regions within the same topologically
associating domains (TAD) exhibited higher similarity in their embeddings compared to
those separated by TAD boundaries (Supplementary Fig. S2c). Furthermore, long-range
genomic region pairs with chromatin contacts showed significantly higher embedding
similarity than those without contacts (Supplementary Fig. S2d). To further evaluate how
TRN embeddings relate to 3D genome organization, we trained a deep learning model
that integrates embedding similarity with Hi-C contact data from long-range genomic
region pairs (see Methods for details). This approach successfully imputed high-
resolution chromatin contacts (Micro-C) in human embryonic stem cells (hESCs) and
demonstrated robust generalization to unseen cell type (Supplementary Fig. S2e). These
findings demonstrate that ChromBERT's pre-trained TRN embeddings effectively
represent the complex regulatory architecture of the human genome, providing a powerful
framework for characterizing transcriptional regulatory architecture and chromatin
organization.

Next, we explored the effectiveness of ChromBERT's regulator embeddings in
representing the functional collaborations among transcription regulators. Each regulator
embedding at a given genomic region encapsulates its contextual relationship with other
regulators. Intuitively, regulator pairs with strong functional collaborations should exhibit
greater embedding similarity. To validate this, we averaged the embeddings of each
regulator across genomic regions on chromosome one and calculated the pairwise
similarities. This analysis revealed a highly heterogeneous pattern of co-association
among transcription regulators, forming distinct clusters (Supplementary Fig. S2f). We
identified a specific cluster involving well-characterized transcription factors and co-
activators, such as SMARCA4, EP300, BRD4, MED1, FOXM1, and MYC (Supplementary

Fig. S2f), indicating functional associations among transcription regulators with high
similarity in embeddings.

Moreover, pairs of transcription regulators with high embedding similarity correspond to
increased protein-protein interaction frequencies (as defined by affinity-purification mass
spectrometry data from BioPlex 3.0[26]) and functional association (as defined by hallmark
gene sets and ontology gene sets from the Molecular Signatures Database[27])
(Supplementary Fig. S2g). This provides strong evidence that pre-trained regulator
embeddings can reflect the functional collaborations of transcription regulators.
Considering the context-specific nature of TRNs, we also examined the embedding
similarities of regulator pairs across distinct genomic regions. RNF2, a classical
component of the Polycomb Repressive Complex 1 (PRC1)[28], exhibited high embedding
similarity with factors typically associated with transcription repression (such as PCGF2)
at some genomic regions, as well as with regulators involved in transcription activation
(such as EP300 and MYC) at others (Supplementary Fig. S2h). This pattern largely
corresponds to RNF2's dual functional roles in chromatin[29]. These findings demonstrate
the robust ability of ChromBERT to represent context-dependent functional collaborations
among transcription regulators.

ChromBERT boosts cistrome imputation in unseen cell types
Despite extensive efforts in experimental profiling, the genomic landscapes of
transcription regulators remain far from comprehensive, particularly given their vast
number. To address this limitation, we fine-tuned ChromBERT to impute cistromes of
transcription regulators, *i.e.*, to predict the presence or absence of binding events, in
previously unseen cell types. A significant challenge in this task lies in effectively
integrating cell-type-specific information to facilitate transfer across diverse cell types.
Inspired by the advances in prompt-enhanced fine-tuning, a highly efficient and flexible
approach that has proven transformative in pre-trained deep language models[30-33], we
adapted this technique for cell-type-specific cistrome imputation. Specifically, we
designed cell-type-specific prompts derived from chromatin accessibility profiles or
transcriptome profiles to guide the model (see Methods for details). These prompts

served as cell-type-specific cues, seamlessly integrating with pre-trained regulator
embeddings from ChromBERT to support flexible and accurate predictions for a wide
range of transcription regulators across diverse cell types (Fig. 2a). To align these different
prompts and embeddings within a unified feature space, we fine-tuned the model using
approximately one hundred cistromes from multiple cell types, supplemented by cell-type-
specific DNase-seq or transcriptome data (see Method for details). This integration
ensured that the model could effectively reconcile variability across cell types and
transcription regulators. After this alignment, ChromBERT has the capability to
systematically impute cistromes for any given cell type-regulator combination without
requiring additional cell-type-specific or regulator-specific fine-tuning.

ChromBERT demonstrated superior performance in cistrome imputation across a wide
range of bulk cell types by leveraging DNase-seq data as prompts. It achieved a mean
area under the precision-recall curve (AUPRC) of 0.552 on imputed cistromes,
significantly outperforming the baseline approach, which achieved a mean AUPRC of
0.270. The baseline method, which integrates regulator-specific putative peaks and
DNase-seq profiles, is a common strategy in the absence of ChIP-seq data. This
performance advantage was consistent across both cell lines and primary cells (Fig. 2b).
ChromBERT also exhibited exceptional robustness, with its performance largely
unaffected by the representation frequency of imputed cell types or transcription
regulators in the pre-training dataset (Supplementary Fig. S3a, b). This robustness
underscores ChromBERT's reliability across diverse biological contexts. Notably, one of
the most challenging aspects of cistrome imputation is identifying cell-type-specific
binding sites, which are biologically significant yet computationally challenging, especially
when chromatin accessibility is similar across cell types. ChromBERT demonstrated
superior performance in classifying cell-type-specific binding sites that could not be
explained by differential chromatin accessibility alone (Fig. 2c-e). To further validate its
effectiveness, ChromBERT was compared to an XGBoost model trained directly on the
same input cistromes used during pre-training. ChromBERT consistently outperformed
XGBoost (Fig. 2f), underscoring the predictive power of its pre-training approach.
Furthermore, when benchmarked against Avocado, a state-of-the-art model for cistrome

imputation[34], ChromBERT achieved superior accuracies across applicable test datasets

(Fig. 2g).

Imputing single-cell cistromes is significantly more challenging than bulk-cell cistromes

due to the limited availability of paired single-cell omics data[35,36]. Despite the rapid

expansion of single-cell transcriptomics over the past decade, the lack of corresponding

transcription regulator data impedes our understanding of transcription regulation at the

single-cell level, where cell-type and state-specific dynamics play crucial roles. Building

on the successful alignment of bulk-cell RNA-seq-based prompts from scGPT[6] with pre-

trained ChromBERT embeddings for bulk-cell cistromes imputation (Supplementary Fig.

S4a), we evaluated its potential to impute cistromes at single-cell resolution by replacing

bulk-cell RNA-seq prompts with single-cell prompts (see Method for details). We

evaluated ChromBERT's single-cell imputation performance using a human peripheral

blood mononuclear cells (PBMCs) single-cell multi-omics dataset. Two key aspects were

assessed: predictive accuracy in B cells, using ChIP-seq data from GM12878 as ground

truth, and the single-cell-specificity of prediction, using motif accessibility inferred from

single-cell ATAC-seq data as ground truth (see Methods for details). ChromBERT

achieved significantly higher predictive performance for eight regulators in B cells,

outperforming a combination of DNA-binding motifs and imputed chromatin accessibility

(Supplementary Fig. S4b). Additionally, ChromBERT exhibited robust single-cell-specific

performance (Supplementary Fig. S4c, d). These results underscore the utility of

ChromBERT in cistrome imputation and highlight its potential to extend transcription

regulatory studies by leveraging diverse prompts tailored to a range of biological settings.

ChromBERT reveals key regulators in specific cell types

Transcription regulation is highly complex, as regulatory effects vary across both cell

types and genomic contexts. Within each cell type, genomic regions exhibit specific

regulatory effects (e.g., active enhancers or promoters), but the roles of transcription

regulators mediating these behaviors are often unclear due to limited cell-type-specific

data. To address this, we fine-tuned ChromBERT for cell-type-specific tasks, enabling it

to learn regulatory effects (such as enhancer activity or gene expression level) for diverse

genomic regions within each cell type (Fig. 3a). By interpreting these fine-tuned TRN

representations, ChromBERT offers insights into the roles of transcription regulators in

mediating cell-type-specific regulatory effects. Specifically, a regulator's contextualized

embedding can be interpreted to reflect its context-dependent roles across genomic

regions[15,37]. When the differences between a regulator's embeddings for distinct groups

of genomic regions correspond to observed regulatory effects, the regulator can be

identified as a key contributor in the biological context of interest (see Method for details).

This approach is particularly beneficial for rare or hard-to-study samples with limited

experimental profiling. We fine-tuned ChromBERT for three downstream tasks to

demonstrate its ability.

The first task involved tissue-specific fine-mapping of expression quantitative trait loci

(eQTL), a specific type of genetic variants that influence the expression levels of genes.

Fine-mapping within eQTLs helps to pinpoint the exact causal variants affecting gene

expression, which is crucial for understanding the genetic basis of complex traits and

diseases[38]. To address this, we fine-tuned ChromBERT to classify causal and non-causal

variants using the latest data from the eQTL Catalogue. We incorporated DNA sequence

variation by integrating a DNA sequence prompt from DNABERT-2 into ChromBERT (see

Methods for details). As shown in Fig. 3b, fine-tuned ChromBERT achieved a mean area

under receiver operating characteristic (AUROC) of 0.828, outperforming Enformer

(mean AUROC = 0.770), DNABERT-2 model alone (mean AUROC = 0.788), ChromBERT

without pre-training (mean AUROC = 0.804), and an XGBoost model across 49 human

tissues. Fine-tuned ChromBERT generated contextualized embedding for transcription

regulators across eQTLs, allowing the identification of differential regulators that exhibited

the significant embedding distance between casual and non-causal eQTLs (see Methods

for details).

Chromatin accessibility, indicated by DNase-seq signals, varied the most on average

across 49 tissues (Supplementary Fig. S5a), aligning with its known significant role in

transcription regulation[39,40]. Further analysis of public DNase-seq profiles across 10

tissues confirmed that causal eQTLs exhibited elevated chromatin accessibility

(Supplementary Fig. S5b). Additionally, we also investigated transcription factors that
exhibited significant distance in contextualized embeddings between the two groups of
eQTLs (Supplementary Fig. S5b). These differential factors not only have a higher motif
presence at causal QTLs but also demonstrated significant alternations in motif affinity
following single-nucleotide genetic variations at these loci (Fig. 3c). For instance, in a
case study of causal eQTL rs17079281 influencing the transcription of *DCBLD1* in lung,
we found that a C to T variation led to increased motif affinity of YY1 (Fig. 3d), a differential
regulator identified by ChromBERT. It is consistent with a previous study demonstrating
that rs17079281 decreases lung cancer risk by creating an YY1-binding site to suppress
DCBLD1 expression[41]. These results indicate that ChromBERT is effective in fine-
mapping eQTLs, providing valuable insights into the genetic variants that drive gene
expression changes.
319
The second task was to fine-tune ChromBERT for modeling genome-wide enhancer
activity quantified from STARR-seq[42]. In this task, ChromBERT showed superior
prediction performance than Enformer, DNABERT-2, ChromBERT without pre-training,
and XGBoost model in predicting STARR-seq signals in HCT116 cells (Supplementary
Fig. S5c). However, although numerous transcription regulators and cofactors are known
to co-bound at active enhancers, perturbation experiments for regulators suggests that
differential cofactor dependencies define distinct types of human enhancer[43,44]. To
explore the cofactor dependencies in enhancer activity, we further fine-tuned ChromBERT
to model perturbation effects of regulators using a STARR-seq dataset for
EP300/CREBBP, BRD2 and CDK7 perturbations[44]. The model performed well in
predicting both wild-type and perturbed STARR-seq signals, capturing at least a fraction
of perturbation effects (Fig. 3e, Supplementary Fig. S5d-f). The perturbation model
enables us to capture key regulators dependent on perturbated regulators. We interpreted
regulators with significant embedding shifts upon EP300 perturbation at enhancers with
decreased activity as key cofactors of EP300 (see Methods for details). This approach
identified cofactors such as FOSL1 and JUND having strong functional associations with
EP300 (Fig. 3f), which are consistent with the findings of the previous study[44].
Interestingly, while other well-characterized cofactors like BRD4 were also enriched at

these regions, they did not show a close association with EP300 in our interpretation
analysis (Fig. 3f, g). To validate this finding, we re-analyzed STARR-seq data following
BRD4 perturbation and confirmed the weak functional association between BRD4 and
EP300 in enhancer activities (Fig. 3h). These results demonstrate that fine-tuning
ChromBERT can help to investigate the complex regulatory mechanisms governing
enhancer activity.

The third task was to distinguish distinct functional loci of a multi-function transcription
regulator. EZH2, a core component in Polycomb Repressive Complex 2 (PRC2)[45], has
been reported to have two context-dependent functions: a classical function involving
H3K27me3 and a non-classical function independent of H3K27me3[29,46]. We fine-tuned
ChromBERT to classify classical and non-classical sites of EZH2, where ChromBERT
showed a better performance in this task than other methods (Supplementary Fig. S5g).
Further interpretation of the fine-tuned representations revealed their functional
collaborations with EZH2 in determining EZH2's context-dependent functions (see
Methods for details). Two distinct groups of transcription regulators showed high co-
association with EZH2 (Fig. 3i). The first group is strongly associated with regulators
involved in repressive function, aligning with the classical function of EZH2. In contrast,
the second group is related to active function, including E2F1 and STAT3, two known
cofactors in EZH2's non-classical function[29,47,48]. These results demonstrate the ability of
ChromBERT to decipher the context-dependent functions of transcription regulators.

ChromBERT uncovers key drivers in cell state transition
Understanding the transcription regulatory dynamics underlying cellular heterogeneity
and cellular reprogramming is crucial for gaining deep insights into development, disease
processes, and drug discovery. To investigate these dynamics and identify key regulators
involved in cell state transitions, we fine-tuned ChromBERT to model genome-wide
dynamics of regulatory effects, *i.e.*, to capture transcriptomic and chromatin accessibility
changes, between two cell types. By interpreting the fine-tuned representation adapted
to these tasks, we identified driver regulators responsible for the transition between two
cell states (Fig. 4a; see Methods for details).

We first examined the transcription regulatory dynamics within the human PBMC dataset, which includes five cell types: B cells, T cells, natural killer (NK) cells, dendritic cells, and monocytes, based on marker genes. ChromBERT was fine-tuned to predict the genome-wide changes in the transcriptome or chromatin accessibility between these cell types in a pairwise manner (see Methods for details), enabling the model to capture the dynamics of TRNs. In this regression task, ChromBERT outperformed other methods (Supplementary Fig. S6). To evaluate the impact of transcription regulators, we analyzed the embedding distance between two groups of genomic regions characterized by increased or unchanged gene transcription or chromatin accessibility between two cell types. This interpretation analysis revealed the contribution of specific regulators to cellular heterogeneity and identified key regulators unique to each cell type (see Methods for details). Notably, 37 out of 84 identified key regulators have previously been reported to play important functions in the corresponding cell types (Fig. 4b). For instance, BACH2, NCOR2, PAX5, BCL6, BCOR and IRF4 are well-characterized factors involved in the transcription regulation of memory B cell differentiation[49]. Similarly, FOXP3 and TBX21 are known regulators in T cell lineage commitment[50,51], CEBPA, CEBPB, and SPI1 are well-characterized regulators in monocytes[52-54]. Strikingly, many of these known functional regulators recapitulated by ChromBERT could not be detected through differential expression analysis or motif enrichment analysis in differentially accessible chromatin (Fig. 4c), two commonly used methods for key regulator identification[55]. Additionally, SCENIC+, a computational framework for single-cell multi-omics inference of gene regulatory networks[56], also failed to predict most known functional regulators in this dataset (Supplementary Fig. S7a). Remarkably, these known functional regulators displayed greater distances and higher rankings in embeddings in the fine-tuned model than the pre-trained model (Supplementary Fig. S7b, c), confirming the effectiveness of fine-tuning and representation adaption in uncovering important regulators in cellular heterogeneity. These findings suggest that ChromBERT can accurately capture the dynamics of TRNs between different cell types and highlight important transcription regulators.

Next, we focused on transdifferentiation, which involves directly reprogramming from
starting cells to target cells. In this scenario, ChromBERT was fine-tuned to predict
genome-wide transcriptome and chromatin accessibility changes across five
transdifferentiation processes. ChromBERT consistently outperformed other methods
(Supplementary Fig. S8a, b). Driver regulators were identified by ranking the embedding
distances between genomic regions with increased or unchanged transcription or
chromatin accessibility during transitions from starting to target cell states (see Methods
for details). Most known driver regulators involved in these transitions were successfully
recapitulated by this interpretation analysis (Fig. 4d, Supplementary Fig. S8c). Moreover,
the remaining identified key regulators showed higher embedding similarity with known
driver regulators (Supplementary Fig. S8d), indicating that newly identified transcription
regulators may function collaboratively with known drivers. For instance, identified key
regulators in three cell state transition processes, from fibroblast to iPSCs, from B cells
to macrophage, and from fibroblast to myoblast, were highly enriched in gene ontology
(GO) terms related to the target cell fate determination (Fig. 4e). Additionally, we observed
that these known driver regulators have higher rankings and more significant shifts in
regulator embeddings in the fine-tuned model than the pre-trained model (Supplementary
Fig. S8e, f), also confirming the effectiveness of fine-tuning and representation adaption.
These results demonstrate ChromBERT's ability to identify key transcription regulators
driving transdifferentiation processes, providing valuable insights for targeted cell fate
manipulation.
421

Discussion

An essential aspect of studying transcription regulation is elucidating context-specific
TRNs across diverse genomic contexts and cell types. These TRNs are crucial for
explaining differences in gene expression, cellular functions, and phenotypes. This study
introduces ChromBERT, a deep language model pre-trained on a large-scale human
ChIP-seq dataset to directly model genome-wide combinatorial binding patterns of
transcription regulators. Through fine-tuning, ChromBERT demonstrates superior
performance in a variety of downstream tasks related to transcription regulation, including
cistrome imputation in previously unseen cell types or single cells, as well as modeling

cell-type-specific regulatory effects and dynamics. In addition, by interpreting transformers-encoded TRN embeddings, ChromBERT sheds light on the roles and functional collaborations of transcription regulators in regulating gene expression and chromatin states. Notably, ChromBERT adapts these embeddings to the specific biological contexts associated with downstream objectives, offering cell-type-specific insights into the regulatory architectures without requiring cell-type-specific genomic data on transcription regulators.

ChromBERT's architecture and training paradigm offer significant opportunities for expanding its applications to additional downstream tasks in transcription regulation. For example, ChromBERT could play a transformative role in single-cell omics studies by identifying context-specific transcriptional regulatory architectures and their dynamics across heterogeneous cell populations. It could also enhance efforts to understand tissue-specific regulatory mechanisms, aiding studies that investigate the interplay between transcription factors, chromatin states, and lineage-specific differentiation processes. Although ChromBERT primarily focuses on the interactions of transcription regulators within each genomic context, integrating its TRN embeddings from multiple genomic contexts with long-range genomic distances could address emerging challenges, such as enhancer-promoter interaction prediction and the dissection of long-range chromatin interactions. ChromBERT's potential can be further amplified through integration with other foundation models via prompt-enhanced fine-tuning. By leveraging prompts that encode biological information, ChromBERT can synergistically interact with DNA-sequence-based foundation models, which focus on DNA sequence grammar, or single-cell foundation models, which specialize in single-cell transcriptomics. For instance, prompts can guide the integration of DNA sequence features from DNA-sequence-based foundation models with ChromBERT's TRN embeddings, enabling joint analyses of sequence and regulatory patterns. Similarly, combining ChromBERT with single-cell foundation models through prompts could yield insights into the gene-regulatory networks underlying single-cell transcriptomic data, enhancing the understanding of dynamic cell states and transitions.

Despite its promising potential, ChromBERT has limitations that hinder its full utility in
modeling transcriptional regulation. These challenges arise primarily from the diversity
and representation frequency of transcription regulators during pre-training and the
complexities of aligning different foundation models for downstream tasks. ChromBERT
relies on publicly available cistromes for approximately one thousand transcription
regulators, but many regulators lack such data, limiting its representation of the full
transcriptional regulatory spectrum. Additionally, the scarcity of cell-type-specific
cistromes necessitates pooling data across cell types, which, while broadening regulator
inclusion, compromises cell type specificity. Expanding ChIP-seq datasets and
incorporating cell type information as metadata could enable ChromBERT to analyze
transcription regulation in a cell-type-specific manner and leverage cell-type-specific
prompts for downstream tasks. Integration with other foundation models poses further
challenges due to differences in latent spaces across modalities, such as DNA sequences
and single-cell transcriptomics. For example, accurately predicting variant effects requires
better alignment between ChromBERT embeddings and sequence-based embeddings,
as the current model struggles to distinguish distinct single-nucleotide variants within the
same genomic region. A more comprehensive solution involves pre-training a multi-modal
foundation model from scratch, combining diverse data types like DNA sequences, long-
range chromatin interactions, and gene expression profiles. This approach could provide
a unified framework for transcription regulation, offering systematic and comprehensive
insights while overcoming the limitations of post hoc model alignment.

Methods

Assembly and binding status encoding of cistromes

Assembly of cistromes in Cistrome-Human-6K

487 To create a comprehensive pre-training corpus, we gathered data from a wide range of
488 cell types using publicly available sources. Initially, we collected all human ChIP-seq,
489 DNase-seq, and ATAC-seq data (cistromes) from the Cistrome Data Browser[22]. We then
490 applied several filtration steps to select qualified datasets:

- Quality control metrics: We included datasets that met at least four out of five quality control metrics provided by the Cistrome Data Browser: sequence median quality score ≥ 25, uniquely mapped ratio ≥ 60%, PCR bottleneck coefficient ≥ 80%, fraction of reads in peaks ≥ 1% and number of 10-fold confident peaks ≥ 500.

- Peak counts: We recorded datasets with more than 100 called peaks for transcription factors and over 1000 called peaks for other transcription regulators.

- Genome coverage: We ensured that more than 60% of genome-wide regions without called peaks were covered by reads.

- Replicate removal: To ensure accuracy in our annotations, we followed a structured approach to associate datasets with cell populations. Specifically, when a dataset explicitly included a cell line annotation, we used the cell line as the primary label. For datasets without cell line annotations, we assigned labels based on a combination of cell type and tissue to avoid conflating datasets with similar cell types derived from different tissues. This approach ensures consistent and biologically meaningful categorization. To avoid redundancy, we retained the highest-ranking dataset for each transcription regulator in the same cell line / cell type, based on the combined score of all quality control metrics. Given the inherent heterogeneity of tissues, all datasets derived from tissues that met our quality control criteria were retained for analysis.

Following these filtration steps, we assembled a dataset Cistrome-Human-6K consisting of 6,391 ChIP-seq, DNase-seq and ATAC-seq cistromes. Additionally, we manually curated its metadata to ensure its accuracy (Supplementary Table. S1). The dataset was subsequently processed into a signal matrix, $S \in \mathbb{R}^{N \times P}$, where each element $S_{i,j}$ represents average signal (reads per million mapped reads, RPM) of cistromes $c_j \in \{c_1, c_2, \cdots, c_P\}$ ($P = 6,391$) at the i-th bin $b_i \in \{b_1, b_2, \cdots, b_N\}$ in the genome. The set $\{b_1, b_2, \cdots, b_N\}$ denotes consecutive 1-kb bins split from the human build genome hg38. To avoid potential sequencing bias, blacklist regions from ENCODE were excluded from the analysis (https://www.encodeproject.org/files/ENCFF356LFX/). To focus on co-binding events at genomics regions, only bins with peak presence of more than two cistromes

among all cistromes were kept in the analysis. Ultimately, a total of 2,137,894 1-kb bins

were retained in the matrix.

Binding status encoding of cistromes

Transcription regulators often co-bind to chromatin and function in a combinatorial

manner[1]. In ChromBERT, we model the combinatorial binding patterns of cistromes to

generate representations for context-specific TRNs. Therefore, the input into ChromBERT

is the binding statuses of all assembled cistromes at each individual 1-kb bin b_i, which

were processed from signal vector $S_i = \left[S_{i,1}, S_{i,2}, \cdots, S_{i,P} \right]$ at genomic bin b_i.

To ensure the comparability and suitability of signals from different cistromes for input into

the deep language model, we first discretized these signals into distinct categories

representing different binding statuses. Specifically, for each cistrome c_j, we categorized

its signals across the entire genome into two groups, "binding" or "non-binding". The

threshold was defined as the 10-th percentile of signals from 1-kb bins with peak presence.

Peak files were downloaded from Cistrome Data Browser[22]. This categorization is

designed to highlight the biologically meaningful but sparse binding events of transcription

regulators in the genome. The "binding" category was further divided into three equal

portions, "slightly positive", "moderately positive" and "strongly positive", based on the

signals within this group. Similarly, the "non-binding" group was divided into two equal

portions, "slightly negative" and "strongly negative" (Supplementary Fig. S1a). This finer

classification aims to incorporate the impact of binding strength in modeling the co-

association pattern of transcription regulators. After binding status categorization for all

cistromes, the signal vector can be converted to a binding status vector $X_i =$

$\left[X_{i,1}, X_{i,2}, \cdots, X_{i,P} \right]$, where $X_{i,j}$ corresponding to the defined five categories of binding

statuses from "strongly negative", "slightly negative", "slightly positive", "moderately

positive" to "strongly positive".

Architecture and pre-training of ChromBERT

The architecture of ChromBERT

ChromBERT architecture includes three parts: (1) an input embedding layer to generate

input embeddings from input features; (2) eight transformer blocks (encoder); and (3)

output heads adapted to different tasks (decoder).

The input embedding layer converts the binding status vector of each 1-kb bin into a high-

dimension representation. The binding status vector in each 1-kb bin, X_i, is analogous to

a sentence in natural language processing, where binding status of each cistrome, $X_{i,j}$,

is considered as a word in the sentence. Therefore, we utilize the similar input

representation approach in BERT model[21], to generate input embeddings from X_i. The

input embeddings include position embeddings that represent different cistromes and

token embeddings that represent different binding statuses. Each word in X_i is assigned

a unique integer position identifier based on its originating cistrome, resulting in a position

vector for all words, denoted as $v = [id(1), id(2) \cdots, id(P)]$. And token vector is X_i, where

each item $X_{i,j}$ belongs to one of five binding status categories. Then ChromBERT employ

the commonly used embedding layers in PyTorch (v2.0.1)[57], $Emb_{position}$ and Emb_{token},

for two vectors respectively, to map each position and token to a fixed-length embedding

vector of dimension $d_{input} = 768$. Consequently, the resultant summarized input

embeddings for each bin b_i, $E_i^{(input)} \in \mathbb{R}^{P \times d_{input}}$, is defined as:

$$E_i^{(input)} = Emb_{position}(v) + Emb_{token}(X_i) \tag{1}$$

$E_i^{(input)}$ represents the binding statuses of P cistromes at the same genomic bin,

facilitating the following modeling of their co-association patterns with transformers.

ChromBERT employs a transformer encoder to convert input embeddings $E_i^{(input)}$ into

TRN embeddings $E_i^{(TRN)} \in \mathbb{R}^{P \times d_{model}}$, which represents context-specific TRN, where

$d_{model} = 768$. The encoder comprises eight self-attention transformer blocks[20,21], each

consisting of a multi-head self-attention layer and a feed forward neural network layer

(Supplementary Fig. S1b). These self-attention mechanisms operate on the input

embeddings, allowing the model to capture intricate interactions among all cistromes. The

self-attention mechanism of l-th block operates as follows:

$$Q = E_i^{(l-1)}W_q^{(l)}, K = E_i^{(l-1)}W_k^{(l)}, V = E_i^{(l-1)}W_v^{(l)} \tag{2}$$

$$Attention(Q,K,V) = softmax\left(\frac{QK^T}{\sqrt{d_{model}}}\right)V \tag{3}$$

Here, $E_i^{(l-1)}$ is the output of the $(l-1)$-th block, and $E_i^{(0)}$ is $E_i^{(input)}$, Q, K and V are the

query, key and value vectors respectively, each with dimension $P \times d_{model}$. The weight

matrices $W_q^{(l)}, W_k^{(l)}, W_v^{(l)}$ are of dimensions $d_{model} \times d_{model}$, and $W_q^{(0)}, W_k^{(0)}, W_v^{(0)}$ are of

dimensions $d_{input} \times d_{model}$. Additionally, ChromBERT splits the computation of self-

attention into eight attention heads to allow the model to jointly attend to information from

different representation subspaces at different positions. To enhance processing

efficiency for the large sequence length $(P = 6,391)$, ChromBERT incorporates

FlashAttention-2 (v2.3.2)[58], which optimizes attention computation through improved

parallelism and work partitioning.

Due to the lack of explicit ground truth for genome-wide combinatorial interactions among

transcription regulators, ChromBERT's pre-training is performed in a self-supervised

manner. This involves a decoder architecture that follows the encoder to reconstruct input

binding status vector X_i from the TRN embedding $E_i^{(TRN)}$, producing an output $\hat{Y}_i \in \mathbb{R}^{P \times 5}$.

The model setup and training processes are facilitated using PyTorch, ensuring robust

configuration, data handling, and computational efficiency.

Pre-training with masked learning

ChromBERT employs a masked learning approach, widely used in pre-trained models, to

enhance the generalizability of fundamental knowledge acquired during pre-training,

which allows for efficient and accurate transfer learning in a variety of downstream fine-

tuning tasks[5,6,8,21,59]. In the pre-training of ChromBERT, 15% of the binding statuses in the

input sequence X_i were randomly selected. The model's pre-training objective is to

predict these masked statuses based on the context provided by the remaining cistromes.

For these 15% binding statuses, we applied a similar masked learning strategy used in

BERT[21]. In details, we replaced the binding statuses $X_{i,j}$ with (1) the $[MASK]$ in 80% of

the time or (2) a random binding status in 10% of the time or (3) the unchanged $X_{i,j}$ in 10%

of the time. Through this process, ChromBERT gained deep insights into the grammar
that governs the combinatorial interactions among transcription regulators, effectively
learning to infer masked elements.

Due to the severe class imbalance among the five binding status categories, focal loss[60]
was employed to refine the training process. This loss function modifies the standard
cross-entropy loss to focus more on difficult-to-classify instances by decreasing the
weight of easily classified examples. In the pre-training stage, only masked objectives
were taken account in the loss computation. Let $y_j = Y_{i,j}$ denote as the true binding
status category of the cistrome c_j at the i-th bin b_i, then the focal loss function is defined
as:

$$FL_{multi}(\hat{y}, y) = -\sum_{c_j \in c_{masked}} \sum_{t=1}^{5} (1 - \hat{y}_{j,t})^{\gamma} \cdot y_{j,t} \cdot log(\hat{y}_{j,t}) \qquad (4)$$

In this formula, γ is a focusing parameter, set to 2 by default, which scales the contribution
of each example based on the ease of its classification. $y_{j,t}$ is an indicator variable equals
to 1 if $X_{i,j} = t$, and equals to 0 for otherwise, where t is one of the five binding status
categories. $\hat{y}_{j,t}$ is the predicted probability for the binding status cistrome c_j being t. This
adjustment ensures that the model pays greater attention to learning from challenging
cases, thereby addressing the imbalance in the training data effectively.

Specific hyperparameters during the pre-training of ChromBERT were empirically
selected as follows. The learning rate scheduler was configured to a linear schedule,
incorporating a warm-up period that comprised 10% of all learning steps, increasing from
0 to the maximum learning rate 1×10^{-4}, and a decayed period that comprised remain
90%, decreasing from the maximum learning rate 1×10^{-4} to 0. The AdamW optimizer
was employed with its default settings to facilitate efficient optimization. The training batch
size was set to 32, with gradient accumulation occurring every eight batches to manage
memory effectively while achieving more stable gradient estimates. To speed up the
training process, PyTorch Lightning (v2.0.4) (https://lightning.ai/pytorch-lightning) was
implemented, enabling distributed computation across multiple GPUs. This framework
significantly enhanced the training efficiency by parallelizing computations. ChromBERT

underwent a rigorous training stage, spanning 100 epochs, which was completed in
approximately 23 days with four NVIDIA A800 80GB GPUs. This intensive training period
was necessary to ensure that the model adequately learned the complex patterns in the
training data, setting a foundation for the subsequent fine-tuning stage.

Representations of ChromBERT

TRN embeddings

ChromBERT is designed to represent context-specific TRNs by modeling the
combinatorial binding patterns of transcription regulators across diverse genomic
contexts. For each genomic region b_i, ChromBERT encodes it into TRN embeddings
$E_i^{(TRN)} \in \mathbb{R}^{P \times d_{model}}$. These embeddings are generated through transformer blocks, which
provide a flexible architecture for fine-tuning the model's parameters to adapt to diverse
downstream tasks. By adjusting attention mechanisms between transcription regulators
during fine-tuning, ChromBERT captures the architectures of TRNs specific to the given
biological contexts. These embeddings offer crucial insights and task-specific biological
interpretations, underpinning the model's utility in various transcription regulation-related
applications.

Specifically, the $E_i^{(TRN)}$ focused on the context-specific interactions of P (6,391) cistromes,
which are pooled ChIP-seq profiles of M (1,073) regulators. To improve the model's utility
and interpretability, we transformed the cistrome-based TRN embeddings $E_i^{(TRN)}$ to
regulator-based TRN embeddings $E_i'^{(TRN)} \in \mathbb{R}^{M \times d_{model}}$. This transformation involves
characterizing the role of individual transcription regulators within the TRN. For regulators
associated with multiple cistromes, we computed the average of all cistromes
embeddings linked to a single regulator r_m across different cell types at genomic bin b_i,
resulting in $E_{i,m}'^{(TRN)} \in \mathbb{R}^{d_{model}}$, which is defined as follows:

$$E_{i,m}'^{(TRN)} = \frac{1}{|c_{r_m}|} \sum\nolimits_{c_j \in c_{r_m}} E_{i,j}^{(TRN)} \tag{5}$$

where c_{r_m} represent all cistromes associated with r_m. These regulator-based embeddings

were consistently applied in subsequent fine-tuning tasks and interpretability analysis by

default.

Regulator embeddings

After transformation from cistrome-based TRN embedding to regulator-based TRN

embedding, ChromBERT can output regulator embedding $E_{i,m}'^{(TRN)}$ for each regulator r_m

at genomic bin b_i

Inference for functional collaborations of transcription regulators

In assessing the functional collaborations of transcription regulators, we inferred their

interactions based on the similarity of their regulator embeddings within TRN

presentations, derived from either a pre-trained model or a fine-tuned model adapting to

specific downstream tasks. Specifically, for a given genomic region b_i, we computed the

embedding similarity between pairs of regulators, denoted as r_j and r_k. This was

quantified using the cosine similarity formula:

$$CS_{b_i}(r_j, r_k) = \frac{E_{i,j}'^{(TRN)} \cdot E_{i,k}'^{(TRN)}}{\left\| E_{i,j}'^{(TRN)} \right\| \left\| E_{i,k}'^{(TRN)} \right\|} \tag{6}$$

where $E_{i,j}'^{(TRN)}$ and $E_{i,k}'^{(TRN)}$ represents regulator embeddings for r_j and r_k at bin b_i from

the TRN embeddings $E_{i,j}'^{(TRN)}$. Furthermore, for functional collaborations across a specific

set of bins, we averaged the regulator embeddings across all bin in the set. The

embedding similarity was then calculated in a similar manner, enabling a broader

understanding of regulator interactions across multiple genomic regions.

In silico perturbation and omission

In silico perturbation

ChromBERT takes the genome-wide binding events of around one thousand transcription

regulators as input, which can be in silico perturbed for analysis exploring the impact of

binding events of individual regulator on the model output. This involved modifying the

binding events of cistromes associated with each transcription regulator r_m across the

genome. Specifically, we transformed the original input $X_{i,j}$ into an in silico knockout input

$X'_{i,j}$, where converted all categories with positive or "slightly negative" binding status to

"strongly negative" category. For each perturbed regulator r_m, this in silico perturbation

process was applied to all associated cistromes $c_j \in c_{r_m}$.

Omission of cistromes

Generally, ChromBERT processes binding status vector $X_i = [X_{i,1}, X_{i,2}, \cdots, X_{i,P}](P =$

$6{,}391)$. However, in certain analyses, highly dominant cistromes may overshadow the

contributions of cofactors related to the learning objectives. For example, the presence of

H3K27me3 is strongly correlated with EZH2's classical role in gene expression repression.

This dominance often leads to the exclusion of other regulators during fine-tuning tasks

of classifying classical versus non-classical EZH2 binding sites. As a result, omitting

H3K27me3 cistromes can enhance the identification of other key regulators. Formally, we

yield a new binding status vector $X'_i = [X_{i,1}, X_{i,2}, \cdots, X_{i,P}]$ where $c_j \notin c_{r_m}$ by omitting all

cistrome related with r_m.

Fine-tuning of ChromBERT

Basic principle of fine-tuning

The fine-tuning of ChromBERT was conducted using a selective training strategy to

effectively leverage its pre-trained knowledge. Initially, ChromBERT was loaded with pre-

trained weights to retain the general representations learned during pre-training. To

preserve this foundational knowledge, the early layers of the encoder were frozen, while

the final layers of the encoder and all layers of the task-specific decoder were made

trainable. This approach ensures that ChromBERT retains its pre-trained knowledge while

adapting effectively to the specific requirements of diverse downstream tasks. By

selectively adjusting the model, ChromBERT can adapt its TRN embeddings to align with

task-specific and cell-type-specific objectives. This capability not only enables the model

to perform well across a range of downstream tasks but also facilitates the interpretation

of the roles and functional collaborations of transcription regulators in cell-type-specific

transcription regulation.

Fine-tuning for cell-type-specific regulatory effects

ChromBERT was fine-tuned to model regulatory effects specific to cell types by learning genomic-context-specific labels across diverse genomic regions within each cell type. These labels correspond to the regulatory effects of specific genomic regions (e.g., active or inactive enhancers) in the given cell type. Generally, ChromBERT has adopted a universal design of decoder header architecture for these, eliminating the need for task-specific configurations. This design includes a convolutional layer followed by a multi-layer perceptron (MLP), which efficiently transforms TRN embeddings into the requisite outputs. The output can be a single probability for binary classification tasks or a single value for regression tasks, depending on the needs of the downstream tasks. Notably, this design also supports tasks with inputs subjected to in silico perturbation or omission. Further details on this kind of downstream tasks were described in the following downstream tasks section, and the frozen number of transformer blocks, loss functions and training hyperparameters were summarized in Supplementary Table S3.

Identification of key regulators in cell-type-specific regulatory effects

After fine-tuning ChromBERT to model regulatory effects in specific cell types, we analyzed the task-adapted TRN embeddings to identify key transcription regulators driving these cell-type-specific regulatory effects. Transformer-based deep language models, such as BERT, are particularly well-suited for this purpose due to their ability to generate contextualized embeddings that capture the context-dependent roles of individual elements[15,37]. ChromBERT leverages this capability to generate regulator embeddings that vary across genomic contexts (e.g., 1-kb genomic bins), allowing for a detailed analysis of transcription regulators' roles within diverse genomic regions.

To identify key regulators, genomic regions are classified into two groups characterized by distinct regulatory effects. For each transcription regulator, the average embedding is computed across the regions in each group using the formula:

$$\bar{E}_{G,m}^{\prime(TRN)} = \frac{1}{|G|} \sum_{i \in G} E_{i,m}^{\prime(TRN)} \tag{7}$$

749   where G represent the set of 1-kb genomic bins and $\bar{E}'^{(TRN)}_{G,m}$ denotes the average

750   embedding for the regulator r_m across the set of loci G derived from the fine-tuned model.

751   Then, we evaluate the distance between embeddings of the two groups of loci for each

752   transcription regulator as follows:

$$D_m(G_1, G_2) = 1 - \frac{\bar{E}'^{(TRN)}_{G_1,m} \cdot \bar{E}'^{(TRN)}_{G_2,m}}{\left\| \bar{E}'^{(TRN)}_{G_1,m} \right\| \left\| \bar{E}'^{(TRN)}_{G_2,m} \right\|} \tag{8}$$

A greater distance D_m indicates higher contextual variability for the regulator, signifying

its functional importance in distinguishing the regulatory effects between two groups.

Transcription regulators with the highest embedding distances are identified as key

regulators of context-specific regulatory effects within each task. This approach highlights

the strength of ChromBERT, and other BERT-based architectures, in capturing the

nuanced, context-dependent roles of transcription regulators. By interpreting regulator

embeddings across functionally distinct loci, ChromBERT enables the identification of key

regulators underlying the given genomic diversity. This provides valuable insights into

transcriptional regulation within cell-type-specific settings, particularly in the absence of

extensive experimental data for individual regulators.

Fine-tuning for regulatory dynamics between cell types

In our study, we utilized ChromBERT to understand transcription regulatory dynamics in

single cellular heterogeneity and during cell state transitions. Specifically, the model was

fine-tuned to predict genome-wide changes in transcriptome and chromatin accessibility

between pairwise cell types. The fine-tuning process for chromatin accessibility followed

the same objectives as those used for determining cell-type-specific regulatory effects.

However, the objectives for transcriptome changes differed due to the potential

uncertainty in the influence of regions adjacent to transcription start sites (TSSs) on gene

expression. For each gene, we included both upstream and downstream regions relative

to the TSS to capture a comprehensive genomic context. We selected four genomic bins

on either side of the TSS, as well as the bin containing the TSS itself, denoted as

$[b_{i-4}, b_{i-3}, b_{i-2}, b_{i-1}, b_i, b_{i+1}, b_{i+2}, b_{i+3}, b_{i+4}]$. To derive the most informative TRN

embeddings from these regions, we then employed a max pooling strategy. This method

ensures that the most significant signals across these genomic bins are captured,
enhancing the accuracy of our predictions regarding gene expression regulation. A
universal decoder architecture was followed to transform the TRN embeddings to the
requisite output. Further details on this kind of downstream tasks were also described in
the following downstream tasks section, and Supplementary Table S3.

Identification of key regulators in regulatory dynamics between cell types

Following fine-tuning for regulatory dynamics, ChromBERT identifies key transcription
regulators critical to changes in regulatory effects between cell types. This process
parallels the identification of key regulators in cell-type-specific regulatory effects, with the
difference being that the embedding distances are calculated between groups of regions
categorized as having changed or unchanged regulatory effects between cell types. This
enables ChromBERT to pinpoint regulators driving transcriptional dynamics across cell
types, providing insights into the mechanisms underlying cellular transitions.

Fine-tuning with prompts for additional information incorporation

Due to the mixed cell type pooling operation and reliance on cistromes involved in pre-
training, the encoder of ChromBERT exhibits a limited capacity to generate variable TRN
embeddings in response to specific inputs. In this study, we employed a prompt-enhanced
fine-tuning approach, integrating TRN representations with additional relevant information,
including cell-type-specific data or genetic variations. This approach successfully
addresses these limitations and broadens the model's applicability across diverse
objectives, including cistromes imputation and the identification of causal eQTLs. By
annotating cistromes, cell-type-specific chromatin accessibility representation derived
from pre-trained ChromBERT embedding can be utilized as cell-type-specific information
for cistromes imputation. Additionally, transcriptome embeddings from scGPT were also
employed as cell-type-specific information in this study. Genetic variation information we
used was sourced from DNABERT-2 for identification of causal eQTLs. Vectors
containing additional information were denoted as prompts in this study. To utilize the TRN
representations and prompts, we concatenated each TRN embeddings $E_i'^{(TRN)}$ with
respective prompts $E_i^{(prompts)}$. In this concatenation, we performed a mean pooling on

sequence length dimension to convert the TRN embeddings into a 768-dimensional

vector for convenient concatenation with prompts. This concatenation produced a prompt-

enhanced embeddings $E_i''^{(TRN)}$, which was subsequently processed by an MLP decoder

to generate the necessary outputs. Further details on this kind of downstream tasks were

described in the following downstream tasks section.

Downstream tasks

Hi-C imputation

We employ pre-trained TRN embeddings to enhance Hi-C contact map imputation,

effectively capturing spatial chromatin regulation. We employed a CNN model utilizing

three input channels: a O/E normalized Hi-C contact map, a pairwise genomic distance

map expressed as $(1 + D_g)^{-0.75}$, where D_g represents genomic distance, and a pairwise

TRN embedding distance map defined by cosine similarity. The input window is set at

1000 kb with a 1-kb resolution and moves diagonally in 200-kb steps. Hi-C contact maps

were initially processed at 5-kb resolution and then linearly interpolated to a 1-kb

resolution. We used an O/E normalized Micro-C contact map as the ground truth, dividing

the chromosomes into three sets for training, validation, and testing. The training set

includes chromosomes 1, 4, 7, 10, 13, 17, and 18; the validation set consists of

chromosomes 3, 6, 9, 12, 16, 19, 20, and X; and the test set contains chromosomes 2, 5,

8, 11, 14, 15, 21, and 22. Specially, 1-kb bins excluded during ChromBERT's pre-training

were assumed to lack TF binding, generating embeddings with an all-zero binding status

vector. To minimize potential data bias, we adjusted the cosine similarity values for

contacts involving these excluded regions by substituting them with the mean cosine

similarity value of contacts at the same distance.

Cistrome imputation using cell-type-specific prompts (fine-tuning with prompts)

The current matrix representing cell types and transcription regulators is highly sparse

due to the lack of available cistromes for many transcription regulators across diverse cell

types. To address this issue, our method integrates cell-type-specific prompts based on

chromatin accessibility into the model to impute the matrix by predicting new cistromes,

leveraging chromatin accessibility's established role in predicting transcription regulator

binding statuses. Alternatively, cell-type-specific prompts based on transcriptomic profiles

can be incorporated, as these profiles are also linked to transcription regulator binding

statuses and are abundantly available. Furthermore, our approach incorporates a

regulator prompt to differentiate among various transcription regulators, facilitating a

universal generative model architecture capable of generating cistromes for distinct

transcription regulators and performing binary classification to determine their presence

or absence. These prompts improve the model's ability to adapt its predictions to specific

cell types.

• Chromatin accessibility prompt: We leveraged DNase-seq data extensively

available for various cell types in the Cistrome-Human-6K. For each cell type,

ChromBERT's pre-trained embeddings specifically for DNase-seq data were used,

denoted as $E_{i,j}^{(DNase)} \in \mathbb{R}^{d_{model}}$, where i represents the i-th bin and j indicates the

position index of DNase-seq data within the Cistrome-Human-6K dataset for the

given cell type.

• Transcriptome prompt: Transcriptome prompts were derived from cell embeddings

of scGPT[6] following zero-shot learning on transcriptomic profiles, either from

single-cell or bulk cell data, specific to the cell type, denoted as $E^{(scGPT)} \in \mathbb{R}^{d_{scGPT}}$,

where $d_{scGPT} = 512$. For single-cell RNA-seq data, cells covering fewer than 200

or more than 7,000 genes were filtered out, and the sum of read counts in each

cell were normalized to 1×10^4 followed by a log1p transformation using Scanpy

(v1.9.5)[61]. scGPT then processed these profiles, only considering genes with non-

zero expression levels for the computation of attention. For bulk cell RNA-seq data,

we utilized the same normalization and transformation procedure and used top

7,000 highly-expressed genes to obtain cell embeddings.

• The regulator prompt was pre-trained regulator embedding $E_{i,m}^{\prime(TRN)}$ for given

regulator r_m at the i-th 1-kb bin.

Then for this fine-tuning task, the prompts can be:

$$E_i^{(prompt)} = concat(E_{i,m}^{\prime(TRN)}, E_{i,j}^{(DNase)}) \tag{9}$$

867 or

$$E_i^{(prompt)} = concat(E_{i,m}'^{(TRN)}, E'^{(scGPT)}) \tag{10}$$

based on the choice of cell-type-specific information representation, where $E'^{(scGPT)}$
represents transformed $E^{(scGPT)}$ by a linear layer to match the hidden dimensions for
concatenation. Remarkably, $E'^{(scGPT)}$ is cell representations and is consistent across all
bins in a certain cell type.

To fine-tune the model for cistrome prediction across various cell types and transcription
regulators using chromatin accessibility as cell-type-specific prompts, the model was
trained with DNase-seq data from 22 cell types and utilized 104 cistromes corresponding
to 29 transcription regulators within these cell types (Supplementary Table S4). For the
model fine-tuned with transcriptomic prompts as cell-type-specific indicators, training was
conducted using scGPT cell prompts derived from the RNA-seq dataset, encompassing
20 cell types and 102 cistromes for 29 transcription regulators within these cell types
(Supplementary Table S4). After the training, the fine-tuned model based on chromatin
accessibility prompts or transcriptomic prompts were both applied to perform prediction
in bulk cell types for transcription regulators.

And the fine-tuned model based on transcriptome prompts was further applied to human
PBMC 10x single-cell multi-omics dataset. Paired single-cell RNA-seq and ATAC-seq
data were collected from the 10x Genomics multi-omics repository
(https://www.10xgenomics.com/datasets/pbmc-from-a-healthy-donor-granulocytes-
removed-through-cell-sorting-10-k-1-standard-2-0-0). Initial data processing and quality
control for single-cell RNA-seq data were performed according to protocols in muon-
tutorials (https://muon-tutorials.readthedocs.io/en/latest/single-cell-rna-atac/pbmc10k/1-
Gene-Expression-Processing.html). We then processed to cluster cells and annotated
different cell clusters based on marker gene expression profiles as describe in the same
tutorial. Similar data processing and quality control steps were performed for the single-
cell ATAC-seq data according to the muon-tutorials (https://muon-
tutorials.readthedocs.io/en/latest/single-cell-rna-atac/pbmc10k/2-Chromatin-
Accessibility-Processing.html). After these initial quality checks, we further refined the

dataset by excluding single-cell ATAC-seq reads detected in fewer than 10 cells and cells

with fewer than 1,000 detected genes in single-cell RNA-seq data. With this workflow, we

obtained a final 9,493 cells from an initial pool of 11,898 cells. After data processing and

cell clustering, we applied fine-tuned ChromBERT with single-cell RNA-seq embeddings

as transcriptome prompts to perform prediction in single cells for transcription regulators.

Specifically, in this prediction section, we removed all cell clusters related to T cells as we

observed that scGPT cell embeddings could not distinguish T cells from other cell clusters

well.

Fine-mapping of eQTLs with DNA sequence prompt (fine-tuning with prompts)

To fine-tune ChromBERT for fine-mapping of eQTLs with DNA sequence prompt, we

concatenated mean pooled TRN embeddings with DNA sequence prompt to be fed into

an MLP decoder to classify the causal and non-causal eQTLs. For each eQTL, we

obtained DNA sequence embeddings with DNABERT-2 for a 1-kb window (from upstream

500 bp to downstream 499 bp of the coordinate of the eQTL). The variant DNA sequence

embeddings $E_i^{(variant)}$ were used as the DNA sequence prompt.

For this classification task, we collected eQTLs from the EBI eQTL catalogue

(https://www.ebi.ac.uk/eqtl) which utilized the SuSiE protocol as described in the

Enformer study[7,62]. We systematically analyzed eQTLs across 49 different tissues. For

each tissue, variants demonstrating a high likelihood of influence on gene expression,

indicated by a posterior inclusion probability (PIP) > 0.9, were categorized as causal

variants. Conversely, for each identified causal variant, we sought a corresponding non-

causal variant. This non-causal variant was chosen from the set with a PIP < 0.01 but an

absolute Z-score > 4 tested for the same gene, wherever available. In cases where a

matched gene variant was not available, we instead selected a random variant from the

genome-wide locus with a PIP < 0.01 and an absolute Z-score > 6, to serve as the non-

causal counterpart. Following the approach used in Enformer study[7], we fine-tuned the

model in each tissue by conducting 10 iterations of cross validation, maintaining a

training/testing ratio of 6:4.

Prediction for genome-wide STARR-seq signals (fine-tuning for cell-type-specific regulatory effects)

For this task, we utilized STARR-seq data from wild type HCT116 cells available in GSE156740[44]. This involved downloading both the STARR-seq signals and a reference set of enhancers, which are based on the human genome build hg19. To align this with ChromBERT, which uses hg38, we converted hg38 1-kb bins to hg19 using the liftOver tool (https://hgdownload.soe.ucsc.edu/admin/exe/linux.x86_64/liftOver). We then associated the human hg38 1-kb bins with the reference STARR-seq enhancers and processed the STARR-seq signals using a $\log_2$-transformed fold changes with the pseudo count one. The model was trained specifically on regions that included these reference enhancers and additional open chromatin regions identified by DNase-seq data in HCT116 cells (ENCFF240LRP). This training approach was designed to enhance robustness of our model for variance of STARR-seq signals. Importantly, we excluded all cistromes related to histone modifications or chromatin accessibility to focus specifically on the influence of transcription factors on enhancer activity.

Prediction for perturbation effect on STARR-seq signals (fine-tuning for cell-type-specific regulator effects)

For predicting the perturbation effects on STARR-seq signals, we adapted ChromBERT to predict changes in control and 3-indoleacetic acid (IAA)-treated HCT116 cells. IAA treatment is known to rapidly deplete tagged transcription factors[44]. We used the original reference cistromes for control cell predictions and applied in silico perturbations for binding status of depleted factors in treated cells (see the in silico perturbation section above). Training for this in silico perturbation model included STARR-seq signals in both control (-IAA) and treated (+IAA) conditions for three transcription factors: BRD2, CDK7, and EP300/CREBBP. Unlike the model for wild type signals, this in silico perturbation model was trained using reference STARR-seq enhancers that show significant variance in response to treatment.

Classification of EZH2's classical and non-classical sites (fine-tuning for cell-type-specific regulatory effects)

For this classification task, we used EZH2 (GSE29611[2]) and H3K27me3 (GSE61176)
ChIP-seq data from human embryonic stem cells. We labeled bins overlapping with both
EZH2 and H3K27me3 as positive and those only overlapping with EZH2 as negative to
create a clear binary classification dataset. As H3K27me3 was included in the pre-trained
dataset and was proposed to have dominant effect on the model output, we omitted
cistromes associated with H3K27me3 from the input reference cistromes.

Prediction for genome-wide changes in transcriptome or chromatin accessibility
(fine-tuning for regulatory dynamics between cell types)
In this task, we fine-tuned ChromBERT to predict genome-wide changes in transcriptome
or chromatin accessibility in cellular heterogeneity and during transdifferentiation.
Specifically, we quantified genome-wide changes in transcriptome with fold changes of
gene expression levels, and quantified genome-wide changes in chromatin accessibility
through fold changes of DNase-seq or ATAC-seq signals.

After data processing and cell clustering for human PBMC 10x single-cell multi-omics
dataset mentioned in the above sections in Methods. We specifically assigned cell
clusters into five major cell types for this prediction section, including T cells, B cells,
monocytes, dendritic cells (DCs) and natural killer (NK) cells. This assignment helps us
to collected well-known key regulators for each cell type from the public studies. For each
cell type, single-cell RNA-seq data were aggregated into pseudo-bulk counts and
normalized using a factor of 1×10^6. Single-cell ATAC-seq data were also compiled into
pseudo-bulk reads and analyzed for peak calling using pycisTopic (v1.0.3)[56], and these
reads were normalized to a count of 1×10^6. Then we fine-tuned ChromBERT to predict
changes in transcriptome or chromatin accessibility between cell types in a pairwise
manner. For transcriptome profiles, we calculated changes using differences of log1p-
transformed pseudo-bulk expression level across all genes. For chromatin accessibility,
changes were calculated by $\log_2$-transformed fold changes in pseudo-bulk reads density
at merged peaks and background regions (spanning 10 kb upstream and downstream
around all TSSs), with a pseudo count one in fold change calculation.

For transcriptome changes in transdifferentiation, we collected transcriptome profiles
derived from CAGE-seq in seven cell types available in the FANTOM5 database[63],
including fibroblasts, induced pluripotent stem cells (iPSCs), myoblasts, hepatocytes,
macrophages, cardiac myocytes, and B cells. We focused on five transdifferentiation
processes: fibroblast to iPSCs, fibroblast to myoblast, fibroblast to hepatocyte, fibroblast
to cardiac myocyte, and B cells to macrophage. Bulk-cell transcriptome changes across
different cell states were quantified with differences of log1p-transformed gene
expression level (TPM) across all genes.

For chromatin accessibility changes in transdifferentiation, we utilized DNase-seq in
fibroblast (ENCFF184KAM), iPSCs (ENCFF540VPT), myoblast (ENCFF647RNC),
cardiac myocyte (ENCFF389SOW), B cells (ENCFF650BNV) and macrophage
(ENCFF580ICE), and ATAC-seq in fibroblast and hepatocyte (GSE179011[64],
GSE185358[65]). Peak calling and reads density normalization were performed in the
approaches similar to those used in our single-cell ATAC-seq pseudo-bulk processing.
And bulk-cell chromatin accessibility changes were also quantified by $\log_2$-transformed
fold changes in reads density at merged peaks and background regions (spanning 10 kb
upstream and downstream around all TSSs), with a pseudo count one in fold change
calculation.

1009

We performed interpretation analysis to identify key regulators in cellular heterogeneity
and during cell state transitions. In the realm of cellular heterogeneity, we categorized
genes and loci based on elevated or stable expression and chromatin accessibility.
Specifically, genes exhibiting a log1p-transformed expression difference greater than 1
were classified as having increased expression, while those with a difference ranging
from -0.5 to 0.5 were considered stable. In chromatin accessibility analysis using ATAC-
seq, regions with a $\log_2$-transformed increase greater than 2 were marked as having
increased chromatin accessibility. After excluding regions that had no coverage, we
selected the top 40,000 regions showing the minimal absolute fold changes as having
unchanged chromatin accessibility. The method for defining genes and regions with

increased or unchanged expression or accessibility during cell state transitions was
analogous to that used for analyzing cellular heterogeneity.

1022

Each analysis between two single-cell pseudo-bulk cell types in cellular heterogeneity or
two cell states in a transdifferentiation process involved ranking transcription factors
based on their embedding distance in two set of loci for transcriptome or chromatin
accessibility dynamics, respectively. Epigenetic modifications were not considered in this
analysis. We averaged ranking scores from these two modalities to compute a composite
ranking score for 991 transcription factors, identifying those most influential in driving
cellular state transition.

1030

In our analysis of cell state transition in each transdifferentiation process, the top 25
transcription factors were recognized as key regulators of this transdifferentiation process.
For cellular heterogeneity, we computed an average ranking score for each regulator by
averaging ranking scores for the given cell type relative to each other (defined as cell-
type-specific ranking scores). This detailed evaluation helps us understand the
importance of these key regulators in differentiating the given cell type from all other cell
types and identify the top 25 key regulators for each cell type, 84 key regulators in total.
And cell-type-specific ranking scores for all 84 key regulators help us to further assign the
cell-type-specificity of identified key regulators, which were generally classified into four
categories, B cells-specific, T cells/NK cells-specific, monocytes-specific and DCs-
specific, as illustrated in Fig. 4b.

1042

Benchmarks

Benchmark for cistrome imputation

We assessed the performance of ChromBERT on the cistrome imputation task by
comparing it with both Avocado and a baseline approach based on putative peaks and
chromatin accessibility. For the Avocado comparison, we used imputed tracks from
ENCODE (ENCSR481OSA) and restricted our analysis to cistromes withheld from
training. For benchmarking, imputed signals derived from BigWig files were aggregated
into 1-kb bins.

1051

The baseline method was established by integrating putative peaks with chromatin
accessibility. For each test cistrome, putative peaks were identified as follows. If at least
five cistromes (excluding the test cistrome) were present in our Cistrome-Human-6K
dataset, related peaks were collected and merged into 1-kb bins to define putative peaks.
If fewer than five were available, putative peaks were instead derived from 1-kb bins
based on the corresponding regulator's motif. The prediction probability in the benchmark
analysis was then computed by multiplying the average DNase-seq signal (aggregated
over 1-kb bins) by a binary indicator (0 or 1) denoting the presence of putative peaks.

1060

We evaluated ChromBERT's performance to predict transcription regulator binding sites
that are specific to a particular cell type, even when chromatin accessibility is similar
across cell types. To define these sites, we performed pairwise comparisons between
paired cistromes from different cell types but associated with the same transcription
regulator. A binding site was classified as cell-type-specific if it was present in one cell
type but absent in the other, represented as 1-kb bins. For a site to be considered peak-
present, it had to meet two criteria: (1) the signal intensity had to exceed 3; (2) the signal
had to be at least threefold higher than in the peak-absent cell type. To control for potential
differences in chromatin accessibility, we applied additional constraints: (1) the absolute
$\log_2$-transformed fold change in DNase-seq signal between the two cell types had to be
less than 0.5; (2) the DNase-seq signal in both cell types had to exceed 3, ensuring that
both sites were located in accessible chromatin. ChIP-seq and DNase-seq signals were
normalized to the genomic background based on read densities. To ensure statistical
robustness, we included only cistrome pairs with at least 50 cell-type-specific binding
events, while also requiring that the ratio of peak-present to peak-absent bins remained
balanced, with a maximum difference of less than fivefold. ChromBERT's performance to
distinguish cell-type-specific binding events was assessed by comparing its predicted
probability differences between paired cell types against the ground truth—defined by the
actual presence or absence of transcription regulator binding based on ChIP-seq data.

1080

Benchmark with XGBoost

The XGBoost model (implemented by Python package XGBoost, version 2.1.3) was

utilized as a baseline to demonstrate the effectiveness of TRN embeddings generated by

ChromBERT across various downstream tasks, using the same input as ChromBERT.

While other tasks can directly utilize the XGBoost model, fine-tuning tasks with prompts

require specific configurations, as direct training is not practical.

1087

For cistrome imputation tasks based on DNase-seq prompts, the input vector is

constructed by concatenating the following three components: a discrete signal token for

the DNase-seq in the target cell type, serving as a chromatin accessibility prompt; a

discrete signal token for cistromes associated with the target transcription regulator

indicated by the most frequent one, functioning as the regulator prompt; and the cistromes

binding status vector $X_i = [X_{i,1}, X_{i,2}, \cdots, X_{i,P}]$. The training and validation of the XGBoost

model was conducted on the same datasets used by ChromBERT, following an

incremental learning strategy with the binary cross-entropy (BCE) loss function. The

training dataset was divided into 356 randomly assigned blocks, processed sequentially.

Initially, the first block permitted up to 50 rounds of boosting, implementing early stopping

if no improvement was detected on the validation dataset for 10 consecutive rounds. For

subsequent blocks, the model employed a number of boosting rounds equivalent to the

number of trees in the prior block, maintaining the same early stopping methodology.

1101

1102   For the task of fine-mapping of eQTLs, the input vector fed into XGBoost is a

1103   concatenated vector comprising variant DNA sequence embeddings $E_i^{(variant)}$ and the

1104   cistromes binding status vector X_i. All other parameters use the default settings.

1105

Benchmark with Enformer and DNABERT-2

In our study, we also compared the performance of ChromBERT with two advanced deep

language models, Enformer[7] and DNABERT-2[8], in a subset of our downstream tasks

which can also be accomplished by the two models based on DNA sequences (genome

build hg38). These models, which generate contextual representations for input DNA

sequences, were fine-tuned using the established method in the original studies with

slight modification to suit specific requirements of the tasks. For DNABERT-2, we

implemented a fine-tuning protocol available from the study (https://github.com/MAGICS-

LAB/DNABERT_2/blob/main/finetune/train.py). In the case of Enformer, we used a

random forest approach, either as a classifier or a regressor, to perform predictions on

downsteam tasks based on the embeddings generated from DNA sequences. The

random forest classifier and regressor were implemented using the package scikit-learn

(v1.3.0)[66], with the "max_features" parameter set to "log2". The number of frozen

transformer layers, loss functions and training hyperparameters for DNABERT-2 and

Enformer were summarized in Supplementary Table S3.

1121

Both models were trained using the same dataset as ChromBERT. DNABERT-2

processed the entire DNA sequences in 1-kb bins, similar to ChromBERT. If ChromBERT

was set to consider multiple adjacent bins (e.g., $n = 9$ bins for changes in the cellular

transcriptome), DNABERT-2 followed the same approach and then applied a max pooling

strategy to the embeddings. Enformer, designed to handle long-range DNA sequences,

required inputs of 196,608 bp. For this purpose, we used one-hot-encoded DNA

sequences centered around the 1-kb bins used in ChromBERT. These inputs were then

transformed into 896 sequence vectors by Enformer, corresponding to 114,689 bp in 128-

bp resolution. To match the lengths of input regions from different models, we selected

the central 8 out of 896 sequence vectors from Enformer's output, corresponding to 1,024

bp, as Enformer's embeddings for the given 1-kb bin. For these 8 sequence vectors, we

performed a mean pooling strategy to generate an averaged sequence vector for the

following random forest.

1135

1136 For tasks where ChromBERT used multiple adjacent bins (e.g., $n = 9$ bins for changes in

1137 the cellular transcriptome), we selected the central 70 out of 896 sequence vectors from

1138 Enformer's output, corresponding to 8,960 bp, and applied a similar mean pooling

1139 strategy.

1140

1141 **Gene ontology enrichment analysis**

1142 We conducted Gene Ontology (GO) enrichment analysis using the DAVID online tool

1143 (https://david.ncifcrf.gov/tools.jsp)[67]. This analysis assessed the enrichment of human

1144 gene ontology biological process (BP) terms.

1145

1146 **Motif scan**

1147 Motif scans were performed using FIMO (v5.0.5)[69] against the JASPAR core 2024

1148 vertebrates database[70] with the following parameters "--max-stored-scores 1000000".

1149 Motifs with p-value $\leq 1 \times 10^{-5}$ were used for the following analysis.

1150

Data availability

Cistrome-Human-6K dataset is available on the Hugging Face Dataset Hub

(https://huggingface.co/datasets/TongjiZhanglab/chrombert). The blacklist of genomic

regions was obtained from the ENCODE project (https://www.encodeproject.org) under

accession code ENCFF356LFX. ChIP-seq data used for analysis in Supplementary

Fig.S2h was obtained from GEO accession code GSE62562 (H3K27me3)[68] and

GSE39912 (H3K4me3)[69]. The Hi-C dataset used for the analysis of Supplementary Fig.

S2c was obtained from GEO accession GSE63525[70]. The Micro-C and Hi-C data in hESC

cells (4DNES21D8SP8, 4DNES2M5JIGV) and HFF cells (4DNESWST3UBH,

4DNES2R6PUEK[71]) used for Hi-C imputation in Supplementary Fig. S2d-e were obtained

from 4DN consortium (https://data.4dnucleome.org/). The ChIP-seq of BRD4

(GSE89128[72], GSE99178[73]) and DNase-seq data (GSE26328[74], GSE25344[75]) in A549

and K562 cells used in Fig. 2c-d are obtained from GEO. The ChIP-seq of EP300

(GSE51176[76], GSE59681[77]) and DNase-seq data (GSE50610[78], GSM723024[79]) in

HCT116 and IMR-90 cells used in Fig. S3c are obtained from GEO. The data used in Fig

2b, Fig 2e-g and Supplementary Fig. S4a-b were obtained from ENCODE and GEO,

whose file accession codes were available in Supplementary Table S2. eQTLs are

available from the EBI eQTL catalogue (https://www.ebi.ac.uk/eqtl). ChIP-seq data of YY1

and DNase-seq data used for the analysis in Fig. 3d was obtained from GEO accession

GSE32465[2] and GSE18927[79]. The DNase-seq in HCT116 used to define open regions

were downloaded from (https://www.encodeproject.org) with file accession code

ENCFF240LRP and the list of enhancer regions for analysis in Fig.3e-h is available from

the supplementary material of the previous study[44]. The STARR-seq data used for analysis in Fig.3e-h were downloaded from accession code GSE156740[44]. ChIP-seq data of EP300 and BRD4 used in Fig. 3g were from accession codes GSE51176[76] and GSE57628[80]. ChIP-seq data in hESC used for analysis in Fig. 3i was obtained from GEO accession codes GSE61176[81] (H3K27me3) and GSE29611[2] (EZH2). The RNA-seq data used as transcriptome prompt of ChromBERT in the cistrome imputation task were obtained from ENCODE and the file accession codes were listed in Supplementary Table S4. The single-cell RNA-seq and ATAC-seq data used in cistrome imputation task of Supplementary Fig. S4 was obtained from PBMC 10x single-cell multi-omics dataset (https://www.10xgenomics.com/datasets/pbmc-from-a-healthy-donor-granulocytes-removed-through-cell-sorting-10-k-1-standard-2-0-0). The DNase-seq data used for modeling the chromatin accessibility changes in transdifferentiation was obtained from (https://www.encodeproject.org) with file accession codes ENCFF184KAM (fibroblast), ENCFF540VPT (iPSC), ENCFF647RNC (myoblast), ENCFF389SOW (cardiac myocyte), ENCFF650BNV (B cell) and ENCFF580ICE (macrophage). The ATAC-seq data used for modeling the chromatin accessibility changes in transdifferentiation was downloaded from accession codes GSE179011[64] and GSE185358[82]. The CAGE-seq data used for modeling the transcriptome changes in transdifferentiation was obtained from (https://fantom.gsc.riken.jp/5/datafiles/reprocessed/hg38_latest/extra/CAGE_peaks_exp ression/hg38_fair+new_CAGE_peaks_phase1and2_tpm.osc.txt.gz).

Code availability

The pre-trained model, reference datasets and all source codes of ChromBERT are available on the GitHub repository (https://github.com/TongjiZhanglab/ChromBERT).

Acknowledgments

We would like to thank Prof. Rui Jiang for comments on designing the research. This work was supported by the National Natural Science Foundation of China (32325012 (Y(Yong).Z.), 32030022 (Y(Yong).Z.), 32488101 (Y(Yong).Z.), 32400522 (Z.Y.)), the National Key Research and Development Program of China (2021YFA1302500

1203 (Y(Yong).Z.)), the Science and Technology Commission of Shanghai Municipality
1204 (23JS1401200 (Y(Yong).Z.)), the Postdoctoral Innovation Talents Support Program
1205 (BX20230265 (Z.Y.)), the China Postdoctoral Science Foundation (2022M722423 (Z.Y.))
1206 and the GHfund C (202302033256 (Z.Y.)).

1207

Author Contributions

1209 Y(Yong).Z. conceived the research, Y(Yong).Z. and Z.Y. designed the research. Z.Y. and
1210 D.Y. developed the model, Q.C. and Y(Yuxuan).Z. performed computational analysis with
1211 the help of Y.W., Z.L. and C.W., Z.Y., D.Y., Q.C., Y(Yuxuan).Z. and Y(Yong).Z. wrote the
1212 manuscript.

1213

Competing interests declaration

1215 The authors declare they have no competing interests.

1216

References

1.   Gerstein, M.B. *et al.* Architecture of the human regulatory network derived from ENCODE
data. *Nature* **489**, 91-100 (2012).
2.   Consortium, E.P. An integrated encyclopedia of DNA elements in the human genome.
Nature **489**, 57-74 (2012).
3.   Stampfel, G. *et al.* Transcriptional regulators form diverse groups with context-dependent
regulatory functions. *Nature* **528**, 147-51 (2015).
4.   Consortium, E.P. *et al.* Expanded encyclopaedias of DNA elements in the human and
mouse genomes. *Nature* **583**, 699-710 (2020).
5.   Theodoris, C.V. *et al.* Transfer learning enables predictions in network biology. *Nature* **618**,
616-624 (2023).
6.   Cui, H. *et al.* scGPT: toward building a foundation model for single-cell multi-omics using
generative AI. *Nat Methods* (2024).
7.   Avsec, Z. *et al.* Effective gene expression prediction from sequence by integrating long-
range interactions. *Nat Methods* **18**, 1196-1203 (2021).
8.   Zhou, Z. *et al.* Dnabert-2: Efficient foundation model and benchmark for multi-species
genome. *arXiv preprint arXiv:2306.15006* (2023).
9.   Hao, M. *et al.* Large-scale foundation model on single-cell transcriptomics. *Nat Methods*
21, 1481-1491 (2024).
10.   Fu, X. *et al.* A foundation model of transcription across human cell types. *Nature* (2025).
11.   Nguyen, E. *et al.* Sequence modeling and design from molecular to genome scale with Evo.
Science **386**, eado9336 (2024).

12. Dalla-Torre, H. *et al.* Nucleotide Transformer: building and evaluating robust foundation
models for human genomics. *Nat Methods* (2024).
13. Gururangan, S. *et al.* Don't stop pretraining: Adapt language models to domains and tasks.
arXiv preprint arXiv:2004.10964 (2020).
14. Chen, V. *et al.* Applying interpretable machine learning in computational biology-pitfalls,
recommendations and opportunities for new developments. *Nat Methods* **21**, 1454-1461
(2024).
15. Ethayarajh, K. How contextual are contextualized word representations? Comparing the
geometry of BERT, ELMo, and GPT-2 embeddings. *arXiv preprint arXiv:1909.00512* (2019).
16. Yu, Z. & Zhang, Y. Foundation model for comprehensive transcriptional regulation analysis.
Natl Sci Rev **11**, nwae355 (2024).
17. Kamimoto, K. *et al.* Dissecting cell identity via network inference and in silico gene
perturbation. *Nature* **614**, 742-751 (2023).
18. Slattery, M. *et al.* Absence of a simple code: how transcription factors read the genome.
Trends Biochem Sci **39**, 381-99 (2014).
19. Yu, Z., Wang, Q., Zhu, G., Zhu, J. & Zhang, Y. Decoding the genomic landscape of chromatin-
associated biomolecular condensates. *bioRxiv*, 2023.08. 23.554542 (2023).
20. Vaswani, A. *et al.* Attention is all you need. *Advances in neural information processing*
systems **30**(2017).
21. Devlin, J., Chang, M.-W., Lee, K. & Toutanova, K. Bert: Pre-training of deep bidirectional
transformers for language understanding. *arXiv preprint arXiv:1810.04805* (2018).
22. Taing, L. *et al.* Cistrome Data Browser: integrated search, analysis and visualization of
chromatin data. *Nucleic Acids Res* **52**, D61-D66 (2024).
23. Liu, T. *et al.* Cistrome: an integrative platform for transcriptional regulation studies.
Genome Biol **12**, R83 (2011).
24. Vu, H. & Ernst, J. Universal annotation of the human genome through integration of over
a thousand epigenomic datasets. *Genome Biol* **23**, 9 (2022).
25. Bonev, B. & Cavalli, G. Organization and function of the 3D genome. *Nat Rev Genet* **17**,
772 (2016).
26. Huttlin, E.L. *et al.* Dual proteome-scale networks reveal cell-specific remodeling of the
human interactome. *Cell* **184**, 3022-3040 e28 (2021).
27. Liberzon, A. *et al.* The Molecular Signatures Database (MSigDB) hallmark gene set
collection. *Cell Syst* **1**, 417-425 (2015).
28. Wang, H. *et al.* Role of histone H2A ubiquitination in Polycomb silencing. *Nature* **431**, 873-
8 (2004).
29. Hu, S. *et al.* ncHMR detector: a computational framework to systematically reveal non-
classical functions of histone modification regulators. *Genome Biol* **21**, 48 (2020).
30. Liu, P. *et al.* Pre-train, prompt, and predict: A systematic survey of prompting methods in
natural language processing. *ACM Computing Surveys* **55**, 1-35 (2023).
31. Kirillov, A. *et al.* Segment anything. *arXiv preprint arXiv:2304.02643* (2023).
32. Chowdhery, A. *et al.* Palm: Scaling language modeling with pathways. *Journal of Machine*
Learning Research **24**, 1-113 (2023).
33. Wei, J. *et al.* Finetuned language models are zero-shot learners. *arXiv preprint*
arXiv:2109.01652 (2021).

34. Schreiber, J., Durham, T., Bilmes, J. & Noble, W.S. Avocado: a multi-scale deep tensor
factorization method learns a latent representation of the human epigenome. *Genome*
Biol **21**, 81 (2020).
35. Grosselin, K. *et al.* High-throughput single-cell ChIP-seq identifies heterogeneity of
chromatin states in breast cancer. *Nat Genet* **51**, 1060-1066 (2019).
36. Bartosovic, M., Kabbe, M. & Castelo-Branco, G. Single-cell CUT&Tag profiles histone
modifications and transcription factors in complex tissues. *Nat Biotechnol* **39**, 825-835
(2021).
37. Wiedemann, G., Remus, S., Chawla, A. & Biemann, C. Does BERT make any sense?
Interpretable word sense disambiguation with contextualized embeddings. *arXiv preprint*
arXiv:1909.10430 (2019).
38. Uffelmann, E. *et al.* Genome-wide association studies. *Nature Reviews Methods Primers*
1, 59 (2021).
39. Boyle, A.P. *et al.* High-resolution mapping and characterization of open chromatin across
the genome. *Cell* **132**, 311-22 (2008).
40. Song, L. & Crawford, G.E. DNase-seq: a high-resolution technique for mapping active gene
regulatory elements across the genome from mammalian cells. *Cold Spring Harb Protoc*
2010, pdb prot5384 (2010).
41. Wang, Y. *et al.* SNP rs17079281 decreases lung cancer risk through creating an YY1-binding
site to suppress DCBLD1 expression. *Oncogene* **39**, 4092-4102 (2020).
42. Arnold, C.D. *et al.* Genome-wide quantitative enhancer activity maps identified by STARR-
seq. *Science* **339**, 1074-7 (2013).
43. Krebs, A.R., Karmodiya, K., Lindahl-Allen, M., Struhl, K. & Tora, L. SAGA and ATAC histone
acetyl transferase complexes regulate distinct sets of genes and ATAC defines a class of
p300-independent enhancers. *Mol Cell* **44**, 410-423 (2011).
44. Neumayr, C. *et al.* Differential cofactor dependencies define distinct types of human
enhancers. *Nature* **606**, 406-413 (2022).
45. Margueron, R. & Reinberg, D. The Polycomb complex PRC2 and its mark in life. *Nature* **469**,
343-9 (2011).
46. Xu, K. *et al.* EZH2 oncogenic activity in castration-resistant prostate cancer cells is
Polycomb-independent. *Science* **338**, 1465-9 (2012).
47. Xu, H. *et al.* Integrative analysis reveals the transcriptional collaboration between EZH2
and E2F1 in the regulation of cancer-related gene expression. *Molecular Cancer Research*
14, 163-172 (2016).
48. Kim, E. *et al.* Phosphorylation of EZH2 activates STAT3 signaling via STAT3 methylation and
promotes tumorigenicity of glioblastoma stem-like cells. *Cancer Cell* **23**, 839-52 (2013).
49. Laidlaw, B.J. & Cyster, J.G. Transcriptional regulation of memory B cell differentiation. *Nat*
Rev Immunol **21**, 209-220 (2021).
50. Fontenot, J.D. *et al.* Regulatory T cell lineage specification by the forkhead transcription
factor foxp3. *Immunity* **22**, 329-41 (2005).
51. Fang, D. *et al.* Differential regulation of transcription factor T-bet induction during NK cell
development and T helper-1 cell differentiation. *Immunity* **55**, 639-655 e7 (2022).

52. Chopin, M. *et al.* Transcription Factor PU.1 Promotes Conventional Dendritic Cell Identity
and Function via Induction of Transcriptional Regulator DC-SCRIPT. *Immunity* **50**, 77-90 e5
(2019).
53. Welner, R.S. *et al.* C/EBPalpha is required for development of dendritic cell progenitors.
Blood **121**, 4073-81 (2013).
54. Scholz, F. *et al.* The transcription factor C/EBPbeta orchestrates dendritic cell maturation
and functionality under homeostatic and malignant conditions. *Proc Natl Acad Sci U S A*
117, 26328-26339 (2020).
55. Hammelman, J., Patel, T., Closser, M., Wichterle, H. & Gifford, D. Ranking reprogramming
factors for cell differentiation. *Nat Methods* **19**, 812-822 (2022).
56. Bravo Gonzalez-Blas, C. *et al.* SCENIC+: single-cell multiomic inference of enhancers and
gene regulatory networks. *Nat Methods* **20**, 1355-1367 (2023).
57. Paszke, A. *et al.* Pytorch: An imperative style, high-performance deep learning library.
Advances in neural information processing systems **32**(2019).
58. Dao, T. Flashattention-2: Faster attention with better parallelism and work partitioning.
arXiv preprint arXiv:2307.08691 (2023).
59. Yang, F. *et al.* scBERT as a large-scale pretrained deep language model for cell type
annotation of single-cell RNA-seq data. *Nature Machine Intelligence* **4**, 852-+ (2022).
60. Lin, T.-Y., Goyal, P., Girshick, R., He, K. & Dollár, P. Focal loss for dense object detection. in
Proceedings of the IEEE international conference on computer vision 2980-2988 (2017).
61. Wolf, F.A., Angerer, P. & Theis, F.J. SCANPY: large-scale single-cell gene expression data
analysis. *Genome biology* **19**, 1-5 (2018).
62. Wang, G., Sarkar, A., Carbonetto, P. & Stephens, M. A simple new approach to variable
selection in regression, with application to genetic fine mapping. *J R Stat Soc Series B Stat*
Methodol **82**, 1273-1300 (2020).
63. Noguchi, S. *et al.* FANTOM5 CAGE profiles of human and mouse samples. *Sci Data* **4**,
170112 (2017).
64. Zhang, X. *et al.* MYOCD is Required for Cardiomyocyte-like Cells Induction from Human
Urine Cells and Fibroblasts Through Remodeling Chromatin. *Stem Cell Rev Rep* **18**, 2414-
2430 (2022).
65. Ma, H. *et al.* The nuclear receptor THRB facilitates differentiation of human PSCs into more
mature hepatocytes. *Cell Stem Cell* **29**, 1611 (2022).
66. Pedregosa, F. *et al.* Scikit-learn: Machine learning in Python. *the Journal of machine*
Learning research **12**, 2825-2830 (2011).
67. Sherman, B.T. *et al.* DAVID: a web server for functional enrichment analysis and functional
annotation of gene lists (2021 update). *Nucleic Acids Res* **50**, W216-W221 (2022).
68. Vallot, C. *et al.* Erosion of X Chromosome Inactivation in Human Pluripotent Cells Initiates
with XACT Coating and Depends on a Specific Heterochromatin Landscape. *Cell Stem Cell*
16, 533-46 (2015).
69. Akdemir, K.C. *et al.* Genome-wide profiling reveals stimulus-specific functions of p53
during differentiation and DNA damage of human embryonic stem cells. *Nucleic Acids Res*
42, 205-23 (2014).
70. Rao, S.S. *et al.* A 3D map of the human genome at kilobase resolution reveals principles of
chromatin looping. *Cell* **159**, 1665-80 (2014).

71. Krietenstein, N. *et al.* Ultrastructural Details of Mammalian Chromosome Architecture.
Mol Cell **78**, 554-565.e7 (2020).
72. Rusan, M. *et al.* Suppression of Adaptive Responses to Targeted Cancer Therapy by
Transcriptional Repression. *Cancer Discov* **8**, 59-73 (2018).
73. Liu, X. *et al.* In Situ Capture of Chromatin Interactions by Biotinylated dCas9. *Cell* **170**,
1028-1043.e19 (2017).
74. Maurano, M.T. *et al.* Large-scale identification of sequence variants influencing human
transcription factor occupancy in vivo. *Nat Genet* **47**, 1393-401 (2015).
75. Song, L. *et al.* Open chromatin defined by DNaseI and FAIRE identifies regulatory elements
that shape cell-type identity. *Genome Res* **21**, 1757-67 (2011).
76. Hu, D. *et al.* The MLL3/MLL4 branches of the COMPASS family function as major histone
H3K4 monomethylases at enhancers. *Mol Cell Biol* **33**, 4745-54 (2013).
77. Ferrari, R. *et al.* Adenovirus small E1A employs the lysine acetylases p300/CBP and tumor
suppressor Rb to repress select host genes and promote productive virus infection. *Cell*
Host Microbe **16**, 663-76 (2014).
78. Maurano, M.T. *et al.* Role of DNA Methylation in Modulating Transcription Factor
Occupancy. *Cell Rep* **12**, 1184-95 (2015).
79. Bernstein, B.E. *et al.* The NIH Roadmap Epigenomics Mapping Consortium. *Nat Biotechnol*
28, 1045-8 (2010).
80. Baranello, L. *et al.* RNA Polymerase II Regulates Topoisomerase 1 Activity to Favor Efficient
Transcription. *Cell* **165**, 357-71 (2016).
81. Singh, Amar M. *et al.* Cell-Cycle Control of Bivalent Epigenetic Domains Regulates the Exit
from Pluripotency. *Stem Cell Reports* **5**, 323-336 (2015).
82. Ma, H. *et al.* The nuclear receptor THRB facilitates differentiation of human PSCs into more
mature hepatocytes. *Cell Stem Cell* **29**, 795-809.e11 (2022).
83. Grant, C.E., Bailey, T.L. & Noble, W.S. FIMO: scanning for occurrences of a given motif.
Bioinformatics **27**, 1017-8 (2011).
84. Gertz, J. *et al.* Distinct properties of cell-type-specific and shared transcription factor
binding sites. *Mol Cell* **52**, 25-36 (2013).
85. Huang da, W., Sherman, B.T. & Lempicki, R.A. Systematic and integrative analysis of large
gene lists using DAVID bioinformatics resources. *Nat Protoc* **4**, 44-57 (2009).
1400

1401 **Fig. 1**

1402
Fig. 1. The ChromBERT architecture.

a. Pre-training schematic of ChromBERT. ChromBERT was pre-trained on the Cistrome-
Human-6K dataset, which encompasses cistromes derived from over 100 tissues,
sourced from three primary categories: cell lines, primary cancer cells, and naturally
occurring primary cells. The input consisted of a binding status matrix, representing the
binding statuses of 6,391 cistromes (*Y*-axis in the heatmap, input features) across
genome-wide 1-kb intervals (*X*-axis in the heatmap, training samples). The pre-training
process employed a masked learning strategy. ChromBERT employed eight self-

attention transformer blocks in an encoder to convert masked binding status matrix (where grey in heatmap indicates masked values and other colors denote discretized binding statuses) into TRN representations, complemented by a decoder to reconstruct the binding statuses. During pre-training stage, binding statuses of 15% cistromes were randomly masked, and ChromBERT was trained to predict the masked objectives using the remaining unmasked binding statuses (see Methods for details), which reinforces the model's ability to capture intricate interaction syntax of these cistromes. ChromBERT generates TRN embeddings (where colors indicate different feature values) representing context-specific TRN, which can be can be further decomposed into regulator embeddings that capture the roles of individual regulators within the TRN. **b.** Fine-tuning schematic of ChromBERT. *Task-adapted fine-tuning*: ChromBERT can be fine-tuned for various transcription regulation-related downstream tasks, such as predicting regulatory effects in specific cell types or regulatory dynamics between cell types. The learning objectives in these tasks are genomic-context-specific labels, which represent the regulatory effects or dynamics of different genomic regions in the given cell type. *Task-adapted interpretation*: Fine-tuning adapts ChromBERT's TRN embeddings to these task objectives, allowing for the interpretation of transcription regulators' roles. The key principle is that regulators with significant differences in their embeddings across distinct groups of genomic regions (classified by their regulatory effects) can be identified as key regulators for the task. This approach helps uncover key transcription regulators driving cell-type-specific regulatory effects and mediating cell state transitions.

 Fig. 2

 Fig. 2. ChromBERT boosts cistrome imputation.

 a. Schematic of cistrome imputation in new cell types or single cells by prompt-enhanced

 fine-tuning of ChromBERT. This process utilizes TRN embedding along with cell-type-

 specific prompts and regulator embeddings (see Methods for details). **b.** Performance

 comparison of cistrome imputation tasks in cell lines and primary cells. The comparison

 is between ChromBERT with DNase-seq prompts, and a baseline method that utilize

 DNase-seq signals and regulator-specific putative peaks (see Methods for details). The

 top heatmap shows performances using the AUPRC metric for both approaches, while

 the bottom bar plot illustrates performance differences. Red bars indicate ChromBERT's

superiority, while blue bars indicate the baseline method's advantage. All test cistromes

were excluded from the training dataset for prompt-enhanced fine-tuning. **c.** Heatmaps

showing predictive probability of ChromBERT with DNase-seq as prompts (red),

alongside ChIP-seq signals (green) and DNase-seq signals (blue) on cell-type-specific

peaks of BRD4 that are accessible in both K562 and A549 (see Methods for details). The

BRD4 ChIP-seq and DNase-seq signals were normalized to the genome background.

The black dashed lines divide the test regions into K562 highly-specific, weakly-specific,

and A549 highly-specific peaks, arranged from bottom to top. **d.** The UCSC genome

browser view illustrates the representative BRD4 binding sites (green) in A549 and K562

cell lines, at consistent peaks (chr17:28,351,650-28,365,238), A549-specific peaks

(chr1:220,683,084-220,697,918), and K562-specific peaks (chr3:14,334,778-14,370,222)

arranged from left to right. Predictive probabilities of ChromBERT with DNase-seq

prompts for BRD4 (red) are shown separately for each cell type, alongside the

corresponding DNase-seq signals (blue). **e.** The histogram illustrates ChromBERT's

performance with DNase-seq as prompts on cistrome imputation. Only regions exhibiting

cell-type-specific binding events and having similar accessibility in both cell types were

evaluated (see Methods for details). The metrics were calculated by using the difference

in predictive probabilities between two cell types to predict the ground truth of cell-type-

specificity obtained from ChIP-seq peaks. This evaluation included 186 cell-type pairs for

24 regulators. **f.** Scatter plots show the performance comparison between ChromBERT

with DNase-seq prompts and XGBoost (see Methods for details) for cistromes in cell lines

(left) or primary cells (right). **g.** Scatter plots show the performance comparison between

ChromBERT utilizing DNase-seq as prompts and Avocado (see Methods for details). For

f and **g**, the diagonal gray line indicates equal performance, with percentages of cistromes

where ChromBERT outperforms annotated alongside the statistical significance

calculated by a two-sided Student's *t*-test.

 Fig. 3

 Fig. 3. ChromBERT reveals key regulators in cell-type-specific regulatory effects

a. Schematic of ChromBERT's fine-tuning in modeling cell-type-specific regulatory effects

and interpreting the underlying roles of transcription regulators. From left to upper right,

ChromBERT is fine-tuned to generate fine-tuned TRN embeddings (colors in the heatmap

indicate feature values) to predict observed regulatory effects in specific cell types or

tissues, where the task objective is context-specific labels across diverse genomic loci.

From left to lower right, fine-tuned TRN embeddings are interpreted to reveal the roles of

transcription regulators corresponding to observed regulatory effects (see Methods for

details). **b.** Box plots comparing the performance in classifying causal versus non-causal

eQTLs across 49 human tissues for ChromBERT, ChromBERT (without pre-training),

DNABERT-2, Enformer, and XGBoost, as indicated by AUROC (see Methods for details).

c. Box plots comparing the motif presence fraction and motif affinity alternation between

differential regulators (top 20% showing the highest embedding shifts between two

groups of eQTLs in each tissue) and other regulators. The motif presence and affinity

(log-odds scores) around +/- 10bp causal eQTLs were determined using FIMO (v5.5.5)[83].

For each regulator, motif presence was identified by p-value < 0.01 in either wild type or

mutant form, and only sites with motif affinity > 0 in either wild type or mutant form were

analyzed for motif affinity alternation. For **b** and **c**, the statistical significance was

performed by a two-sided Mann-Whitney U-test and **** represents p-value $< 1 \times 10^{-4}$.

The center lines mark the median, the box limits indicate the 25th and 75th percentiles,

and the whiskers extend to $1.5\times$ the interquartile range from the 25th and 75th

percentiles. **d.** The UCSC genome browser view shows YY1 signals (blue) near eQTL

rs1707928, along with DNase-seq signals in lung (green). The YY1 ChIP-seq signals

were from GSE32465[84], and DNase-seq signals were from GSE18927[79]. The DNA

sequence in reference genome and genetic variants around the eQTL were shown in the

upper right, and the YY1 binding motif from JASPAR was also shown. **e.** Scatter plots

showing the ground truth of $\log_2$-transformed fold change of STARR-seq signals upon

EP300 perturbation and predictions by ChromBERT at test enhancers. The Pearson's

correlation coefficient is annotated in the plot, and the color represents the density of

points. STARR-seq signals were from previous study (GSE156741[44]). **f.** Heatmap

showing the distance in regulator embeddings of potential cofactors of EP300 at the test

enhancers before and after in silico perturbation for EP300/CREBBP. Two groups of

enhancers were depicted: those with decreased activity (log$_2$-transformed fold change <

-1, $n = 164$) and those unchanged (-0.5 < log$_2$-transformed fold change < 0.5, $n = 343$)

after IAA-treated EP300/CREBBP depletion. Potential cofactors of EP300 were defined

as top 50 regulators that showing the highest embedding similarity with EP300 at all test

enhancers. The color represents the shifts (1 - cosine similarity) in embeddings.

Representative regulators showing high embedding shift at decreased enhancers are

highlighted in blue and representative regulators showing low embedding shift at

decreased enhancers are highlighted in red. **g.** Line charts showing average ChIP-seq

profile of BRD4 around test enhancers. BRD4 ChIP-seq data was from previous study

(GSE57628[80]). **h.** Heatmap showing log$_2$-transformed fold changes of STARR-seq signals

after IAA-treated EP300/CREBBP or BRD4 depletion, test enhancers were divided into

two groups with decreased or unchanged activity after IAA-treated EP300/CREBBP

depletion. **i.** Circos plot showing embedding similarities with EZH2 at classical (w/

H3K27me3) and non-classical (w/o H3K27me3) genomic loci, highlighting two groups of

regulators having locus-preferential embedding similarities with EZH2. The classical

group shows higher embedding similarity with EZH2 at classical loci compared to non-

classical loci (embedding similarity with EZH2 at classical loci ranking the top 5% among

all regulators and the embedding similarity difference between classical loci and non-

classical loci > 0.1), and the non-classical group exhibits the converse pattern. The

classical group was marked in green and the non-classical group was marked in yellow.

The known cofactors in classical function of EZH2 were highlighted in blue, and the known

cofactors in non-classical function of EZH2, E2F1 and STAT3, were highlighted in red.

Each node represents a regulator, and the transparency of edges linking two nodes

represent the embedding similarity of two regulators, only edges with high pairwise

embedding similarity (> 0.8) were plotted.

 Fig. 4

 Fig. 4. ChromBERT uncovers key regulators in cell state transition

a. Schematic of the fine-tuning of ChromBERT in modeling regulatory dynamics between cell types and interpreting underlying roles of transcription regulators. From left to upper right, ChromBERT is fine-tuned to generated fine-tuned TRN embeddings (colors in heatmap indicate feature values) to predict observed regulatory dynamics between cell types, where the task objective is genome-wide changes in transcriptome or chromatin accessibility. From left to lower right, fine-tuned TRN embeddings are interpreted to reveal roles of transcription regulators corresponding to observed regulatory dynamics during cell state transition (see Methods for details). **b**. Heatmap showing cell-type-specific ranking scores of all key regulators identified in five major cell types: B cells, T cells, NK cells, monocytes and DCs. The cell-type-specific ranking scores evaluate the importance of regulators in differentiating the given cell type from all other cell types (see Methods for details). All key regulators identified were generally classified into four categories, B cells-specific, T cells/NK cells-specific, monocytes-specific and DCs-specific according to the cell-type-specific ranking scores for all 84 identified key regulators. Reported key regulators in each cell type were highlighted in red and specifically annotated below the heatmap. **c**. Dot plots showing the low coverage of reported important regulators in each cell type by cell-type-specific differential expression analysis or motif analysis. Reported key regulators covered by differential expression analysis were highlighted in red, while those covered by differential motif analysis were highlighted in blue. Regulators showing differential expression are those with notably higher expression in a specific cell type compared to others (log1p-transformed expression level in the given cell type − average expression level in other cell types > 1). Regulators showing differential motif enrichment are those with notably higher motif enrichment at regions with increased chromatin accessibility in the specific cell type compared to others (observed/expected motif occurrence frequency in the given cell type − average observed/expected motif occurrence frequency in other cell types > 1). The sizes of points indicate the expression level or motif enrichment, and the colors indicate the difference between the given cell type and mean value in other cell types. **d**. The extent to which ChromBERT recapitulates known driver regulators across five transdifferentiation processes. Known driver regulators recapitulated by ChromBERT were highlighted in red. MEF2C, not recapitulated by ChromBERT, was marked in blue.

1562 And FOXA3, lacking related cistromes in our model, was marked in brown. **e**. Gene
1563 ontology (GO) enrichment analysis for 25 identified key regulators using DAVID online
1564 tool (https://david.ncifcrf.gov/tools.jsp)[85]. The top ten human gene ontology biological
1565 process (BP) terms with a false discovery rate (FDR) < 0.05 and high fold enrichment
1566 were displayed. The GO terms involved in the terminal cell states were highlighted in red.