Modeling viral and bacterial infections in human lung organotypic systems reveals strain specific host responses

Bárbara Faria Fonseca[1], Jérôme Wong-Ng[1], Michael Connor[2], Héloïse Mary[1], Min Hee Kim[1], Rémy Yim[1], Vincent Bondet[3], Vincent Michel[4], Hélène Strick Marchand[5], James Di Santo[5], Darragh Duffy[3], Mélanie Hamon[2], Nathalie Sauvonnet[1], Lisa A. Chakrabarti[6], Samy Gobaa[1*]

[1]Institut Pasteur, Université Paris Cité, Biomaterials and Microfluidics core facility, France

[2]Institut Pasteur, Université Paris Cité, Chromatin and Infection laboratory, France

[3]Institut Pasteur, Université Paris Cité, Translational Immunology Unit, France

[4]Institut Pasteur, Université Paris Cité, Pathogenesis of Vascular Infections Unit, France

[5]Institut Pasteur, Université Paris Cité, Innate Immunity Unit, France

[6]Institut Pasteur, Université Paris Cité, Control of Chronic Viral Infections Group, Virus and Immunity Unit, France.

* Correspondence: samy.gobaa@pasteur.fr

Keywords: lung; organoid; organ-on-chip; pneumococcus; SARS-CoV-2

Highlights

- Human fetal epithelial lung stem cells can be expanded as multipotent organoids and differentiated into both airway or alveolar organoids
- Multipotent lung organoids efficiently produce functional epithelia of small airway or alveoli when grown on-chip.
- *Streptococcus pneumoniae* inoculation in alveoli-on-chip mimics the early stages of bacterial colonization in lung epithelia
- Alveoli on-chip system recapitulates variant-specific interactions. SARS-CoV-2 Delta replicates but not Omicron BA.5.
- Robust interferon response upon SARS-CoV-2 infection shows Alveoli on-chip can model innate immune responses.

Abstract

In this study, we developed novel lung organoid-on-chip models that elucidate differential human tissue response to various strains of respiratory pathogens: *Streptococcus pneumoniae* and SARS-CoV-2. We show that human fetal-derived distal lung epithelial cells are readily expandable in 3D as organoids, thereby providing a highly sustainable source of lung progenitor cells. These 3D organoid progenitors can then be induced to produce airway and alveolar organoids on microfluidic devices. Upon challenge with *Streptococcus pneumoniae*, a bacterium known to cause pneumonia, a rapid and strain-dependent colonization was observed at the epithelial surface of alveolar chips. We also assessed SARS-CoV-2 infection in the alveoli-on-chip system and observed that the Delta variant exhibited greater infectivity as compared to the

Omicron BA.5. Both SARS-CoV-2 variants induced potent interferon responses and triggered the expression of different interferon-stimulated genes. Our results demonstrate that strain-specific host defense mechanisms can be recapitulated in human-organoid-based microfluidic systems, paving the way for the use of such platforms for more targeted assessments of human response to novel emergent pathogen strains.

Introduction

The epithelial lining of the human respiratory tract constitutes the major entry point for airborne pathogens. At the tissue level, lung epithelia are supported by a highly specialized stroma, a richly vascularized network and numerous populations of specialized immune cells. Collectively, this complex system ensures tissue homeostasis, barrier function and efficient gas exchange while facing constant viral and bacterial challenges. Host-microbe interactions start at the epithelial apical surface. Resulting infections trigger early defense mechanisms including mucociliary clearance and mucosal innate immunity [1,2]. The complexity of the mounted responses is in part determined by the heterogeneous histological composition of the tissues along the respiratory tract. Ciliated and mucus-producing cells in the airways work together to remove airborne pollutants and microbes. At the distal end of the tract, the alveolar epithelium can also mount a complex immune response in response to pathogen invasion. Disentangling the lung tissue response to airborne pathogens hinges on the establishment of advanced tissue models that extend beyond what can be recapitulated in standard 2D, cell line-based assays.

The major challenge in devising advanced human lung disease models resides in establishing a sustainable source of human cells. Ideally, the starting cell material should be amenable to *in vitro* expansion while still capable of producing the complete range of cell types that constitute the tissue of interest. Immortalized cell lines, such as A549 and Calu-3, can be useful for drug screening and infection studies, but as they originate from lung carcinoma, they poorly represent the biology of bona fide lung cells [3]. On the other hand, primary adult cells can be used to reconstitute nasopharyngeal, bronchial and alveolar tissues *in vitro* [4–6]. However, these largely postmitotic cells can be difficult to obtain and rapidly dedifferentiate [7]. To overcome these limitations, culturing primary lung cells as organoids has proven highly effective. Both human bronchiolar and alveolar organoids can today be derived from human adult biopsies or from iPSC cell culture [8–15]. Lung organoids can be maintained for several passages and be used for lung disease modeling [16,17].

Reproducing the apicobasal polarity of respiratory epithelia is crucial for correctly modeling the entry of an airborne pathogen in a human lung cell. Most current lung organoid protocols lead to the production of apical surfaces facing inward (lumen). Loading any inoculum on the apical side can therefore be challenging. Various strategies were employed to partially overcome this limitation including microinjection [18,19], inverted-polarity techniques [10] and organoid fragmentation and re-formation [20]. However, these approaches often compromise throughput and reproducibility. Alternatively, organ-on-chip technology was shown to be a powerful technique to reconstruct accessible human epithelial tissues in an environment where tissue perfusion of media, drugs and pathogens can be precisely controlled [4,21,22]. This microfluidic-based approach also enables the creation of interconnected channels and can allow the stretching of cultured epithelia [4,23]. Additionally, organ-on-chip systems allow human lung

epithelial cells to be cultured at an air-liquid interface (ALI), further mimicking the function of the
human respiratory tract. Leveraging these technologies for investing host-pathogen interactions
is gaining traction. However, new developments remain pathogen-dependent.

In this study, we combined two organotypic models, organoids and organ-on-chips, to
build novel small airway and alveolar lung infection models. We expanded human distal lung bud
organoids [24–26] and steered them to differentiate, on-chip, into sample-matched airway or
alveolar tissues. We demonstrate that such advanced organ-on-chip devices can successfully
model viral and bacterial infections. We found that the engineered lung models can be used to
identify cell type-related susceptibilities to bacterial infection. We also show that the presented
approaches can be used to characterize variant-dependent human innate immune responses
upon SARS-CoV-2 infection.

Results

Derivation of human multipotent lung organoids

During human lung development, the epithelial lung bud tip forms a pool of multipotent stem
cells capable of generating both small airways and alveoli tissues. A complex interplay of
signaling pathways, primarily driven by WNT and fibroblast growth factor (FGF) activation,
ensure the plasticity of this fetal structure along the pseudoglandular and early-canalicular
stages of lung development [2]. In vitro, reconstructing this signaling cocktail allows long-term
expansion of lung multipotent progenitors [24]. In this study, we derived a stable culture of human
fetal-derived lung bud tip organoids that can be amplified and differentiated into either alveolar or
small airway epithelial cells (Figure 1A). Lung organoids cultured in self-renewing conditions
grow as simple epithelia that are SOX2+ and SOX9+ (Figure 1B & 1D) [2]. These organoids could
be split every 10 to 14 days, multiple times, to expand the number of available cells (Figure 1C).
We also found that stable multipotent organoid cultures can be successfully reestablished after
multiple freeze-thaw cycles, enabling the banking of this initial material, which greatly simplifies
its use and dissemination.

To steer lung multipotent organoids towards alveolar differentiation, we modified the
culture conditions by withdrawing FGF, maintaining WNT activation and adding NOTCH and
TGF-β inhibitors, as described in Lim et al [26]. After two weeks in these differentiation conditions,
immunostaining revealed the presence of HT2-280 positive cells, a human lung-specific type II
pneumocytes (AT2) marker [27]. Pro-surfactant protein C (pro-SFTPC), a typical surfactant protein
produced by bipotent alveolar progenitors and fully differentiated AT2 cells [2], was also identified
in several cells. These observations confirm the differentiation into AT2 cells (Figure 1E). Here,
we did not identify cells with positive labeling for the HT1-56, AQP5 and RAGE, markers of type I
pneumocytes (ATI). While extensive differentiation dynamics were not investigated, we
speculate that terminal AT1 differentiation within organoids requires additional extrinsic factors
including morphogens or physical cues to support their development. Previous studies with
alveolar organoids derived from primary adult lung cells also highlighted an absence of AT1 cells
[8,28]. Interestingly, recent reports with iPSC-derived and fetal-derived alveolar organoids show

that withdrawal of WNT agonists, combined with YAP activators in the differentiation medium, is
necessary for AT1 differentiation [14,29].

Next, we sought to evaluate the capacity of multipotent organoids to differentiate into
small airway tissues, lung organoids were treated with a pro- airway differentiation medium
where WNT agonists were removed and FGF activation maintained [9]. A two-week treatment was
sufficient to induce the differentiation into typical airway cell types such as beating multiciliated
cells that are positive for acetylated tubulin (ac-TUB+) and basal cells expressing cytokeratin 5
(KRT5+). We also identified secretory cells, including goblet cells producing mucin (MUC5AC+)
and club cells expressing secretoglobin family 1A member 1 (SCGB1A1+) (Figure 1F;
Supplementary Movie 1). Our observations show that the morphology of differentiated organoids
showed varying lumen sizes and different extents of folding but systematically appeared as
pseudostratified when compared to multipotent organoids.

Altogether, we demonstrated that human fetal distal lung buds can be efficiently
harvested and used as a reliable source material for the derivation of multipotent human lung
organoids. We also showed that these organoids have the capacity to differentiate into either
small airways or alveolar epithelia.

135 On-chip differentiation of lung reconstituted epithelia

Devising human lung organotypic models to investigate infectious diseases relies first on the
reconstitution of faithful mimics of human epithelia and second on straightforward access to the
luminal compartment that is the classical entry point of airborne pathogens. While organoids can
efficiently ensure an appropriate cell composition, their inward polarization renders infectious
studies labor intensive [18,19]. Here we engineered an organ-on-chip approach that relies on the
use of human organoids as a source of cells and where both the perfusion of fluids and the
inoculum loading are fully controlled [4,21,22].

Briefly, we seeded fragments of multipotent organoids on a commercial S1™ organ-on-
chip device from emulate™ (Figure 2A). Two days of culture under self-renewing conditions
were sufficient to produce a confluent epithelium on the chips. We also observed that adding
sample-matched mesenchymal cells to the lower channel of the chip drastically improved lung
epithelial cell survival and differentiation as previously reported [6,8,26,30,31]. Next, we induced
differentiation by perfusing either airway or alveolar media optimized for organoid differentiation.
Finally, air-liquid interface (ALI) was implemented to promote differentiation (Figure 2A). Both
types of organoid-derived chips formed a simple epithelium expressing E-cadherin and ZO-1
proteins (Supplementary Figure 1). The produced epithelia maintained ALI for multiple days,
suggesting that the junctions are tight.

Next, we analyzed the cell composition of chips induced to differentiate toward the
alveolar lineage (Figure 2B-F). Immunostaining analysis revealed that approximately one third of
the cells expressed pro-SFTPC (31.6% ±19.6), (Figure 2B, 2C). HT2-280, stained an average of
14.4% ±6.4 of cells, indicating that only 12 days in differentiation medium was sufficient to drive
the differentiation of a substantial population of AT2 cells on-chip. Interestingly, AT1 cells could
be identified with RAGE and HT1-56 antibodies, the latter labelling 9.0 ± 4.8% of the cells
(Figure 2B, 2C). RT-qPCR analyses confirmed a down-regulation in the expression of bud tip

markers (SOX2 and SOX9, Figure 2D) and an up-regulation on the expression of mature alveolar markers including LAMP3 (a member of the lysosomal-associated membrane protein family and marker of mature AT2 cells [32]) and chloride intracellular channel 5 (CLIC5), expressed by bona fide AT1 cells [33] (Figure 2E, 2F). Finally, we checked for incidental airway differentiation on alveolar chips and found no evidence of the presence of ciliated, goblet or club cells presence. Basal cells, positive for cytokeratin 5 (KRT5) and transcription factor SOX2 but negative for SOX9, were very seldomly observed. We termed these devices AlveoChips.

When analyzing the cellular composition of small airway chips, immunostaining showed a standard profile of cells including ciliated cells (acTUB+), club cells (SCGB1A1+), goblet cells (MUC5AC+) and basal cells (KRT5+) (Figure 2G, 2H). In addition, a clearly polarized, pseudo-stratified epithelium was observed (Figure 2G). Time-lapse microscopy also showed the presence of beating cilia at the surface of the reconstituted airway epithelium (Supplementary Movie 2) after 20 days of differentiation. Basal cells accounted for 5.3% ± 4.3 and multi-ciliated cells for 3%± 1.6 of the total cells in the epithelium (Figure 2H). These data show that, as for the organoid cultures, chips seeded with human fetal cells derived from multipotent organoids can be guided to produce either airway or alveolar microphysiological systems.

AlveoChip models *Streptococcus pneumoniae* alveolar colonization

S. pneumoniae colonizes the human upper respiratory tract where it is carried asymptomatically. However, when the bacterium invades the alveolar sacs, it can cause severe pneumoniae [34]. Our understanding of the infectious processes, at early stages of pneumococcal colonization, that drive asymptomatic carriage versus pathogenic invasion is limited. Therefore, studying the early events of *S. pneumoniae* in reconstituted human primary lung tissue is highly relevant. Due to the accurate epithelial cell composition of AlveoChips and the capacity to finely control the inoculum administration and subsequent infectious events through microfluidics, we aimed to model *S. pneumoniae* colonization on the alveolar epithelium. Here we used two *S. pneumoniae* strains, TIGR4 and 6B ST90, which are known to cause disease in human and in murine models, with TIGR4 being more invasive than 6B ST90 [35]. For comparison, we used *Corynebacterium accolens* (*C. accolens*), a commensal nasal microbiont that does not colonize the lungs [36].

Briefly, we inoculated AlveoChips TIGR4, 6B ST90 or *C.accolens* bacteria for 6 min under flow (1000ul/h). We then removed all the medium in the top channel and re-established an air-liquid interface (Figure 3A). A dose of 3.0×10^4 colony-forming units (CFU) was used on every AlveoChip (Figure 3B). At 24 hours post infection (hpi), we observed an approximate 1.5 order of magnitude increase of CFU counts in top channel effluents collected from both TIGR4 (median: $5.0.10^5$ total CFU) and 6B ST90 infected chips (median: $7.9.10^5$ total CFU) (Figure 3C). Interestingly, a 2-log decrease in the number of CFU counts was observed in effluents of chips treated with *C. accolens* (median: $3.3.10^2$ total CFU). These results show that the two pneumococcus strains can successfully multiply in the modeled alveolar microenvironment. In contrast, *C. accolens* does not replicate and its inoculated population declines rapidly within the first 24hpi.

We next asked whether these microorganisms could be found in effluents of the bottom channel, suggesting a possible transmigration towards the mesenchyme-filled compartment.

High levels of TIGR4 pneumococcus counts were detected at the bottom channel effluents
(median: $4.1.10^6$ total CFU). Interestingly, while the commensal-like strain 6B ST90 could also
be detected, it was recovered at lower levels than that of TIGR4 (median: $3.9.10^5$ total CFU).
CFU counts from *C. accolens* treated chips were below the limit of detection, confirming the
absence of growth of this pathogen in alveolar chips (Figure 3D). CFU assays revealed that both
pneumococcus strains, which have contrasted effects in mice [35], were able to proliferate in the
AlveoChips.

Finally, immunolabeling after 24hpi revealed the presence of numerous TIGR4 and 6B
ST90 bacteria but no *C. accolens*, at the surface of the epithelial cells corroborating the CFU
results (Figure 3E). We could also detect TIGR4 and 6B ST90 within the pores of the chip
membrane, suggesting a transmigration across the barrier to the bottom chamber
(Supplementary Figure 2A). Interestingly when assessing the frequency of each cell type in
infected chips, we measured a strong decrease in HT2-280+ cell occurrence upon
pneumococcal invasion (Figure 3F). This decrease seemed specific to type II pneumocytes as
no variation in ATI or in total cell numbers were observed after TIGR4 or 6B ST90 inoculation
(Figure 3G, Supplementary Figure 2B). Our data demonstrate that *S. pneumococcus* is readily
capable of colonizing the AlveoChip model and of crossing the epithelial barrier within the first
24hpi.

220 AlveoChip unveils distinct replicative fitness of SARS-CoV-2 variants

To investigate the replication competence of SARS-CoV-2 on-chip, we infected AlveoChips with
the SARS-CoV-2 Delta variant. Chips where then analyzed at 4 days post infection (4dpi)
(Figure 4A). Immunostaining experiments detected 1.9 ± 5.5 % of SARS-CoV-2 spike+ cells
(Figure 4B and 4C). Quantifications showed that 34.1 ± 27 % of the HT2-280+ were positive for
the spike viral protein (Figure 4D). Interestingly, in our experimental conditions, Delta infection
did not trigger major cell death as no significant differences were observed in the total number of
HT2-280+ cells or in the overall cell count on chip post-infection (Supplementary Figure 3A and
3B).

To compare the fitness of different SARS-CoV-2 variants, we infected AlveoChips with
equal titers of SARS-CoV-2 Delta or Omicron BA.5 variants and then quantified the apical
release of viral RNA over 4 days (Figure 4A). RT-qPCR data showed a steady replication of the
Delta variant that starts at 1dpi and reaches a peak around 4dpi (Figure 4E). In contrast,
Omicron BA.5 did not replicate efficiently in our alveolar model: the corresponding viral RNA
load, measured in chip lavages, remained at significantly lower levels (Figure 4E). At the end of
the experiment, the apically released viral RNA for the two variants differed by four orders of
magnitude ($4.4.10^5$ copies per mL for Delta versus 3.10^1 copies per mL for Omicron BA.5).
Interestingly this significant difference ($p < 0.0001$) was consistently observed across all the
used organoid lines, demonstrating the high reproducibility of the system (Figure 4F). At the end
of the experiment, a higher amount of viral RNA could also be detected in total cell RNA
extracted from Delta-infected chips (Figure 4G). Viral particles could be identified by scanning
electron microscopy in chips infected with Delta, but not in chips in mock-infected samples
(Supplementary Figure 4). In addition, immunostainings performed with an anti-viral
nucleocapsid (NCP1) antibody, further confirmed that SARS-CoV-2 infected cells could be

detected in Delta-infected chips, but not in Omicron BA.5 and mock-infected chips
(Supplementary Figure 5). Taken together, these results show that the AlveoChip model is
susceptible to Delta infection, but not to Omicron BA.5, recapitulating the lower replicative
potential of Omicron in an alveolar environment [37,38]. To our knowledge, Alveochip is the first in
vitro alveolar organ-on-chip system that recapitulates differential variant sensitivity.

Finaly, following viral infection, we quantified type I and III interferon (IFN) release on the
luminal surface of lung epithelial cells to verify whether the reconstituted alveolar epithelia can
mount an innate host-defense response. A significant increase in IFNα protein levels (expressed
as IFNa17 equivalents) was observed at 4dpi in Delta-infected chips, compared to uninfected
controls. Interestingly, despite the lower infectivity of Omicron BA.5 in this system, IFNα
secretion was similarly high in epithelial cells challenged with this variant (Figure 4H). For IFNβ,
average protein levels were higher in Omicron BA.5-infected chips, while no significant IFNλ3
production was detected at 4dpi for either variant (Supplementary Figure 6A and 6B). When
looking at the transcription of interferon-stimulated genes (ISGs), both Delta and Omicron BA.5
infections triggered the upregulation of standard ISG such as IFIT1, ISG15 and MX2 (Figure 4I).
Overall, these data show that, in our AlveoChip model, host epithelial cells respond to SARS-
CoV-2 viral inoculation by secreting IFN molecules and expressing ISGs, thereby initiating an
innate immune response. Interestingly, despite the lower replication competence of Omicron
BA.5 compared to the Delta, the former still triggers a strong IFN response.

Discussion

In this work, we assessed the potential of human fetal-derived lung organoids as starting
material for building respiratory infection models. We showed that multipotent fetal lung
organoids can be largely expanded *in vitro* while preserving their capacity to differentiate into
small airway or alveolar epithelia. Using these lung organoids, we generated airway- and alveoli-
on-chip systems which grant easy access to the apical side of epithelial cells and thus can be
readily used for addressing airborne pathogen infections in well-controlled conditions.

Culturing primary adult lung cells as organoids has been extensively explored in the past
decade [8–10,39]. Both biochemical and mechanical cues provided by the extracellular matrix
facilitate the self-organization and differentiation of stem cells and progenitors into faithful tissue
mimics. However, several limitations, including dedifferentiation, limited supply, and donor-to-
donor variations, remain to be solved. While fetal lung tissue is undeniably in very short supply,
fetal lung bud tips can be expanded *in vitro* to form self-renewing organoids that preserve key
markers of multipotent progenitors [24]. On the other hand, induced pluripotent stem cells (iPSCs)
can also generate lung organoids for infection studies [40,41]. However, because it is necessary to
recapitulate all the differentiation steps to reach the lung progenitor state, iPSC protocols tend to
be longer and more labor-intensive than fetal lung cultures [11–15]. Consequently, our approach
provides a valuable addition to existing lung-on-chip models [4,23,42–45], as our system recapitulates
lung epithelia composition *in vitro* without constantly relying on primary cell sourcing or on less
physiological immortalized cell lines.

Airborne pathogens typically infect through the apical region of epithelial cells. To mimic
this aspect *in vitro*, it is necessary to precisely control the inoculum amount, timing and cell
targeting, which can be highly challenging in organoid systems [18,19]. To address this, we
employed an organ-on-chip technology to gain inoculation precision and reproducibility. When
installed onto microfluidic chips, fetal-derived lung cells proliferated, differentiated and polarized
in response to signaling cues present in the differentiation media. Multipotent organoid-derived
cells differentiated on the airway chips and produced the expected goblet, club, basal, and
multiciliated cell populations. The number of multiciliated cells derived on-chip was consistent
with previous reports on primary cells cultured in this medium [9]. Further optimization could
include adding Notch inhibitors and BMP activators, as demonstrated by Van der Vaart et al. [46],
to better control the ratio between the different cell types. In the case of alveolar chips, we
observed a robust production of bona fide AT2 cells (SFTPC+, HT2-280+ and LAMP3+). This is
in line with recent advancements for the generation of mature AT2 cells derived from fetal
material [6,26,47]. Previous efforts struggled with sustaining differentiated AT2 cells for more than
few days in culture [7]. Additionally, we observed a significant number of AT1 cells, detected with
characteristic markers such as HT1-56, RAGE and CLIC5. AT1 cells were less abundant than
AT2 cells in this model, with a 1:1.6 ratio based on HT1-56 and HT2-280 markers; however, this
ratio is consistent with *in vivo* findings (1:2 or 1:3 ratio, as reported by Crapo et al. and Foster et
al. [48,49]). While increasing the AT1 population was not the focus of this study, it may be
achievable on-chip by inducing AT2-to-AT1 transdifferentiation, as demonstrated in other studies
[14,29,43,48]. Overall, our characterization of the Alveochips validates their use for modeling airborne
infections from the colonization steps to the innate immune response.

Previous studies have demonstrated the effects of *S. pneumoniae* infection in human
immortalized cell lines [35], bronchial epithelial cultures [50] and alveolar organoids [41]. Our work
confirms the capacity of TIGR4 and 6B ST90 strains to colonize human alveolar epithelia. In
murine models, the 6B ST90 strain does not cause disease whereas TIGR4 presents a highly
invasive phenotype [35]. Interestingly, in our model TIGR4 crosses the epithelial barrier to a
greater extent than the 6B ST90 strain, partially phenocopying in vivo observations [35,51]. In our
experimental conditions, limited barrier damage was observed upon TIGR4 infection. While
TIGR4 infection is commonly associated with epithelial cell death in most model systems [34], our
AlveoChips experiments reveal a different dynamic. After 24hpi, high levels of TIGR4 counts
were recovered from the chip bottom channel without a visible effect on the epithelial barrier,
which was confirmed by cell counting and ZO-1 staining. These observations suggest that, at an
early time point, TIGR4 can cross the epithelium without causing widespread damage.
Altogether, our model provides a unique model for future investigations into how barrier level
destruction may occur as disease progresses and whether recruitment of innate immune cells
such as macrophages and neutrophils exacerbates this. Indeed, it is known that murine
pneumococcal challenge with invasive disease favouring isolates leads to two waves of
neutrophil infiltration within the lungs. Interestingly, the second wave coincides with rapid
increase in bacterial burden and tissue damage, as opposed to the first being able to partially
control pneumococci [52]. On the other hand, we also found that *C. accolens*, a commensal nasal
species, did not infect alveolar chips. Previous studies have shown that *C. accolens* is highly
abundant in the upper airways, where it can inhibit pneumococcal growth and colonization of the
respiratory tract [36,53]. Future work could explore whether co-culturing *C. accolens* and *S.*

pneumoniae before chip inoculation affects pneumococcal infection of alveolar cells. The
engineered AlveoChip system is capable of supporting investigations on pneumococcal
transmigration events and host responsiveness within an alveolar epithelium, which are still
poorly understood.

The COVID-19 pandemic showcased the continuous emergence of novel variants of
concern, each using different mechanisms for viral spread and immune evasion to increase
infectivity within the respiratory tract [54,55]. SARS-CoV-2 effects have been widely investigated in
airway and alveolar-like *in vitro* systems such as reconstructed airway epithelia in transwells
[5,30,56–58], organoids [10,39,57,59] and organ-on-chip models [45,60,61]. The general consensus is that
novel variants, from the Omicron sublineage, replicate faster and more efficiently in the upper
respiratory tract epithelium compared to earlier variants, which likely contributes to its increased
transmissibility [37,56,62]. In contrast, in *ex vivo* explant cultures of human lungs, Omicron replicates
less efficiently than earlier variants such as Delta [37]. Other studies using an immortalized
alveolar cell line [63], iPSC-derived alveolar cells in micropatterned wells [64] and adult-derived
alveolar organoids [38] also report the lower infectivity of Omicron in the alveolar
microenvironment when compared to Delta. In this work we found that the AlveoChip model can
recapitulate these replication differences between Delta and Omicron BA.5 variants.

Interestingly, we observed a marked increase in IFNα secretion following SARS-CoV-2
infection in AlveoChips. Both Delta and Omicron BA.5 triggered a robust release of IFNα,
detectable at 4dpi. Our data also showed that IFN-β was also secreted mainly in Omicron BA.5-
infected chips. The expression of ISGs following Delta and Omicron infection reflected the
increased IFN secretion. Despite similar induction of the IFN response, the two variants
displayed different replication efficiencies, suggesting that they vary in their ability to evade IFN
signaling. In the airway microenvironment, Omicron is known to be highly replicative and
appears to be resistant to IFN signaling. These characteristics seem to underlie its higher
transmissibility and fitness [62,65]. Recent single-cell RNA sequencing findings suggest that IFN
resistance may be cell-type dependent [57]. For instance, ciliated cells infected with SARS-CoV-2
elicit a clear IFN response, but secretory cells, which can also become infected, produce an
even stronger IFN response, impeding viral replication at later stages [57]. In pediatric nasal
epithelial cells, goblet cells show a robust IFN expression, and consistently produce fewer viral
particles than ciliated cells [58]. Our findings that the strong IFN induction in alveolar cells exposed
to Omicron viruses correlates with Omicron's lower replication in this tissue suggest that SARS-
CoV-2 exhibits cell- or tissue-specific IFN susceptibility.

In summary, our study presents a novel approach to building organ-on-chip models
derived from human multipotent lung organoids. This strategy allowed the production of airway
and alveolar faithful mimics. We showed that our model can recapitulate the early events
following bacterial or viral infection in vitro. Our work also provides mechanistic understandings
on early pathogenesis events and on human immune responses, that are otherwise
inaccessible.

Methods

Procurement of human fetal lung tissue

Human fetal lung tissues were obtained from Advanced Bioscience Resources Inc. (CA, USA) and Cercle Allocation Services (CA, USA). Sourcing, import and organoid derivation were carried out as per national and institutional regulations and guidelines (import authorization number IE-2022-2353 and file number: 2022-075_Alveolar organoids at Institut Pasteur**)**. Samples aged from 15 to 24 post-conception weeks. All the samples used for the current study had no known genetic abnormalities and were tested negative for Syphilis HIV, HBV, HCV, HSV, and CMV.

Primary culture of human fetal lungs and establishment of organoid cultures

The procedure is based on protocols described by Nikolic et al. and Miller et al. [24,25]. Briefly, distal regions of fetal lung lobes were cut into small 2 mm^3 pieces in washing medium (100 units.mL^{-1} Penicillin, 100 µg.mL^{-1} Streptomycin and 0.25 µg.mL^{-1} Amphotericin B in HBSS). After several washes in the same medium, lung fragments were dissociated for 1h at 37 °C with intermittent agitation in a dissociation medium consisting of 1mg/ml Collagenase/Dispase (Roche), 40 U Dnase I (Thermo), 100 units.ml^{-1} Penicillin, 100 µg.ml^{-1} Streptomycin and 0.25 µg.mL^{-1} Amphotericin B in HBSS in RPMI base medium. The enzymatic reaction was inactivated by adding 10% Fetal Bovine Serum (FBS) and the samples were filtered through a 70 µm cell strainer, to filter out non-dissociated lung pieces. After centrifugation for 5min at 250 g, the samples were resuspended in a small volume of Advanced DMEM/F12 base medium with 100 units.ml^{-1} Penicillin, 100 µg.mL^{-1} Streptomycin and 1x Glutamax, then mixed with Geltrex (Thermo) in a 1:3 ratio. 40 µL drops were placed in 24 well plates and incubated at 37 °C for 30 min to solidify the gel. Finally, 400 µL of human self-renewal medium (hSRm) (Supplementary Table 1) were added to each drop and plates incubated under standard cell culture conditions (37 °C; 5% CO2). Medium was changed twice a week.

After 10-14 days of primary culture, lung organoids are ready for passage or cryopreservation. For passage, the medium was removed and 500 µL of cold Organoid Harvesting Solution (Cultrex) added to each well. Geltrex drops were removed, mechanically dissociated and collected in a 15 mL tube. Cold Advanced DMEM/F12 base medium was added up to 10 mL and the samples were centrifugated for 5 min at 100 g. The pellet was resuspended in undiluted Geltrex according to the desired splitting ratio (1:2 to 1:4). After gel solidification, 400 µL of hSRm medium was added in each well.

For freezing, Geltrex drops were detached and dissociated as described for passaging. After centrifugation, the pellets were resuspended in cold freezing medium (50% hSRm + 40% FBS + 10% DMSO), transferred overnight to the -80 °C freezer then to the liquid nitrogen for long term storage.

Human fetal lung mesenchyme culture

During the first passaging of lung organoids, lung mesenchymal cells that had attached to the bottom of the well were recovered using 0.25% Trypsin-EDTA solution, followed by enzyme inactivation with 10% FBS in PBS. These cells were also retrieved after organoid centrifugation by collecting part of the supernatant above the organoid pellet. The cells were then incubated in 25 cm^2 flasks coated with type I collagen (Sigma) and cultured in Advanced DMEM/F12 base

410   medium supplemented with 10% FBS. Medium was changed twice a week until confluence.
411   Lung mesenchymal cells were passaged several times and cryopreserved in the freezing
412   medium detailed above.

Airway and alveolar differentiation of lung organoids

414   Lung organoids were maintained in hSRm medium up to passage 5. Just after passaging this
415   medium was replaced with either airway [9] or alveolar differentiation medium [26] (Supplementary
416   Table 1). Organoids were then cultured for an additional 2 weeks until fixation and
417   immunostaining.

Whole mount immunostaining of organoids

Organoids immunostaining was performed following the protocol described in Dekkers et al [66].
Lung organoids were recovered from Geltrex drops and washed with cold DPBS
($MgCl_2$+/$CaCl_2$+) after centrifugation. Organoids were gently transferred to 1% bovine serum
albumin (BSA) coated 15 mL tubes and fixed with 4% formaldehyde overnight at 4 °C. On the
next day, the 15 mL tube was filled with 1% Tween in DPBS ($MgCl_2$+/$CaCl_2$+) (PBT) up to 10 mL,
gently mixed, centrifuged at 70 g for 5 min at 4 °C. Organoids were either stored in PBT or
used for immunolabeling.

For immunolabeling, organoids were first blocked for 1h in organoid washing buffer
(OWB: 0.1% Triton + 0.2% BSA in DPBS) and then incubated overnight with primary antibodies
using the same solution, with mild shaking at 4 °C. On the next day, the samples were washed
with OWB for approximately 8 hours. The solution was changed every 2 hours. Secondary
antibodies were incubated overnight in OWB with mild shaking at 4 °C. Samples were
counterstained with 4',6-diamidino-2-phenylindole (DAPI), transferred to imaging dishes (Ibidi)
and imaged with a spinning disk confocal microscope Nikon Ti2E (Nikon). Images acquisition
was performed with the NIS-Elements software (Nikon). The list of the primary and secondary
antibodies used are shown in Supplementary Table 2.

Lung on chip culture

Microfluidic S1™ chips were purchased from Emulate Inc (Boston, USA). Chip handling and
activation was performed as per the manufacturer's indications. Briefly, one day before chip
seeding, the devices were activated using ER1/ER2 reagents. Subsequently, top and bottom
channels were incubated with 200 µg.mL^{-1} of human Collagen IV (C5533, Sigma) and 30 µg.mL^{-1}
^{1} of Fibronectin (354008, Corning) at 37 °C overnight. The next day (day 0, see Figure 2A), the
coating solution was removed and washed with DPBS, the chips were then ready for cell
seeding.

Lung organoids up to passage 3 were recovered from Geltrex matrix as performed for
passaging. The equivalent of 2 drops containing organoids were used for each chip. The
organoids pellet was resuspended with an appropriate amount of hSRm medium (40 µL per
chip) and mechanically fragmented using a P200 pipette. The level of fragmentation was verified
by a standard benchtop microscope, to avoid the seeding of isolated cells. Chips were seeded
with few microliters of cell suspension in the top channel and placed in an incubator at 37 °C
overnight. The next day, hSRm medium was added to the cells. Non adherent cells were gently

washed away. On day 2, sample-matched lung mesenchymal cells were detached from the
culture flasks and seeded on the bottom channel at a density of 0.5 x 10^6 cells per mL. In the
meantime, Alveolar or Airway differentiation media (Supplementary Table 1) were added into top
and bottom channel inlet reservoirs of the Pods (Emulate). When good cell attachment was
reached, the chips were inserted into the Pods, placed into a ZOE CM1 device (Emulate) and
top and bottom channels were perfused with medium at flow rate of 30 µL.h^{-1}. On day 7, when
the epithelial layer had reached 100% confluence, an air-liquid interface (ALI) was established
by emptying media from the top channel. Thereafter, lung epithelial cells were fed through the
bottom channel, perfused by the same differentiation medium. Chips were finally used for
experiments on day 14 (alveolar chips) and day 21 (airway chips).

Immunostaining on-chip

Chips were unmounted froms the Pods and fixed in formaldehyde 4% for 30 min, rinsed with
DPBS and carefully cut in half using a scalpel blade. The samples were then placed in 24-well
plates and treated with a permeabilization solution (0.1% Triton in DPBS) for 30 min, followed by
blocking (2% BSA in DPBS) for 1 hour at room temperature. Next, primary antibodies diluted in
1% BSA in DPBS were incubated overnight at 4 °C. On the following day, secondary antibodies,
diluted in 1% BSA in DPBS, were added for 2 hours. Samples were counterstained with DAPI
and mounted in FluoromountG (Thermo Fisher Scientific) before observation with a spinning
disk confocal microscope Nikon Ti2E (Nikon). Images were acquired with the NIS-Elements
software (Nikon). Details of the primary and secondary antibodies and corresponding dilutions
used are listed in Supplementary Table 2.

To quantify the percentage of positively stained cells, maximal intensity projections were
generated using Fiji [67] and analyzed with QuPath version 0.5.1 [68] softwares. QuPath built-in
tools were used for nuclei segmentation followed by cell classification using an interactive
machine learning based approach. Percentage of positively stained cells was calculated as
(number of stained cells/total nuclei number) x 100. Quantification of cell number per µm^2
(Supplementary Figures 2A and 3A) was based on the ratio of counted nuclei per area of the
chip.

RNA extraction and quantitative real-time PCR

For RNA preparation, cells on chips were lysed with 350 µL of RNeasy Lysis buffer + 2% DTT
(w/v). Total RNA was isolated using RNeasy microRNA kit according to the manufacturer's
instructions (Qiagen). RT-qPCR was performed in technical triplicates using the Luna Universal
One-Step RT-qPCR Kit (New England Biolabs), following the manufacturer's instructions. qPCR
was performed on a QuantStudio 6 Flex thermocycler (Applied Biosystems, Thermo). Results
are presented as relative RNA levels quantified by the 2-ΔΔCt method [69] and normalized to the
endogenous control beta-actin. The gene expression profile was obtained from at least three
independent experiments. Primer sequences are shown in Supplementary Table 3.

Bacterial challenge of alveolar chips and bacterial enumeration

Inoculates of the Corynebacterium accolens (Ca) strain C781 were grown overnight in BHI broth
supplemented with 1% Tween 80 [70]. Experimental starter cultures of Streptococcus pneumoniae

(Spn) isolates TIGR4 and 6B ST90 were prepared and grown as previously described in ref [35].
Briefly, stationary phase Ca or midlog Spn were collected by centrifugation, washed three times
in DPBS followed by a 1:100 dilution in DPBS and dilution to 500 organisms per microliter using
isolate specific CFU OD_{600} per mL conversion factors to make a challenge inoculum of 800 µL.
Alveolar chips were challenged with 100 µl (50,000 organisms) of the challenge inoculum by
perfusion through the apical input reservoir of the Emulate Pod at 1000 µL.h^{-1}. Chips where
exposed to the inoculum for 6 min.

Post-challenge inoculate was recovered from the output reservoir of the Emulate pod,
pelleted, and suspended in 200 µL DPBS for CFU serial dilution plating (1:6). Bacterial
adherence within the device was determined by subtracting the recovered post-challenge
inoculate CFU counts from the counts of the challenge inoculum. Apical bacterial burden of the
alveolar chips was determined by flushing the apical side with 500 µL of DPBS at 1000 µL.h^{-1}
flow rate and collection of the perfused DPBS from the output reservoir. This was then pelleted
and suspended in 200 µL DPBS for CFU serial dilution plating (1:6). Transmigrated bacteria
across the alveolar barrier into the bottom chamber perfusion was determined by collecting total
media from the bottom chamber output reservoir of the Emulate pod. This sample was pelleted
by centrifugation and suspended in 200 µl DPBS for CFU serial dilution plating (1:6). All bacterial
dilutions were plated on 5% Columbia blood agar plates and grown overnight at 37 °C with 5%
CO_2.

SARS-CoV-2 infection of alveolar chips

Alveolar chips were cultured for 7 days in ALI conditions prior to SARS-CoV-2 infection. Chips
and pods were then transferred from the ZOE CM1 device to a biosafety cabinet within a BSL3
laboratory. For each chip, 10^6 RNA viral copies per µL of Delta and Omicron BA.5 viral stocks
were used [71,72]. Viral inputs were diluted in DMEM/F12 medium to a final volume of 200 µL and
added into the pod's top channel inlet reservoir. A 1 min pulse at a flow rate of 1000 µL.h^{-1} was
programmed on the ZOE CM1 machine for viral inoculation. Chips were then incubated for 3 h at
37 °C, 5% CO2 under static conditions. Control chips were mock-treated with DMEM/F12 for the
same duration. Post-infection, the remaining viral inoculum in the pods were collected for RT-
qPCR analysis, and the chips were subjected to a series of washes. Briefly, 200 µL of DPBS
was added to the top channel inlet reservoirs, followed by three 5 min pulses at 1000 µL.h^{-1} on a
ZOE CM1 machine. In between each pulse we let the chips 5 min without perfusion, resulting in
a total washing time of 30 min. A last pulse with DMEM/F12 was performed, followed by 20 min
incubation at 37 °C. After this step, the media was collected in the outlet for RT-qPCR analysis.
ALI conditions were then re-established. Apically released viruses were collected daily by
loading 200 µL Advanced DMEM/F12 in the pod top channel inlet reservoir, followed by 1 min
pulse at 1000 µL.h^{-1} in ZOE and by 20 min incubation at 37 °C. ALI was re-established after this
step. On day 4, chips were either fixed in formaldehyde 4% for 30 min or the cellular RNA was
isolated with RNeasy microRNA kit (Qiagen). Samples were stored until further analysis.

Viral RNA quantification

Top and bottom channel effluents collected at days 0, 1 2 and 4 post-infection were inactivated
for 20 min at 80 °C and subsequently subjected to RT-qPCR analysis with the Luna Universal
One-Step RT-qPCR Kit (New England Biolabs), following the manufacturer's instructions. SARS-

CoV-2 RNA was quantified in a final volume of 5 μL per reaction in 384-well plates using SARS-
CoV-2 N-specific primers on a QuantStudio 6 Flex thermocycler (Applied Biosystems, Thermo).
Standard curve was performed in parallel using purified SARS-CoV-2 viral RNA [5].

Scanning electron microscopy

Samples were fixed with 2.5% glutaraldehyde in 0.1 M cacodylate buffer for 1 h at room
temperature and then for 12 h at 4 °C to inactivate SARS-CoV-2. Samples were then washed in
0.1 M cacodylate buffer and several times in water and incubated for 1 h at room temperature in
1% osmium tetraoxide. After dehydration by incubation in increasing concentrations of ethanol,
samples were critical point dried with hexamethyldisilazabe, mounted on a stub, and analyzed
by field emission scanning electron microscopy with a Jeol IT700HR microscope operating at 3
kV.

Quantification of Interferons using ultra-sensitive digital ELISA (Simoa) assays

Prior to protein analysis, samples were treated in a BSL3 laboratory for viral inactivation as
described by Robinot et al [5]. Briefly, 2.5 μL of 10% Triton X-100 solution (Sigma) were added to
25 μL of each lung-on-chip apical lavage supernatant sample (1% final concentration) to be
analyzed for IFNα, IFNβ and IFNλ3. Samples were then incubated 1 h at room temperature.
Prior to Simoa quantification, inactivated samples were diluted 1:5 by adding 110 μL of Diluent B
(Quanterix, Billerica, MA, USA) containing 11 μL of 10% Triton X-100 solution (1% final
concentration). To estimate the limit of quantification, 8 blanks containing only detector/sample
diluent were treated similarly.

IFNα, IFNβ and IFNl3 protein supernatant concentrations were quantified by Simoa
digital ELISA assays developed with Quanterix Homebrew kits as previously described [73–75].
After checking the absence of cross-reactivity, the three assays were multiplexed to decrease
sample volumes needed. The SBG enzyme concentration was 150 pM.

For the IFNα assay, which quantifies all IFNα subtypes with a similar sensitivity, the 8H1
antibody clone was used as a capture antibody after coating on paramagnetic beads (0.3
mg.mL^{-1}), the 12H5 clone was biotinylated (biotin/antibody ratio = 30/1) and used as the detector
at a concentration of 0.3 μg.mL^{-1}, and the 11150 recombinant human IFNα17 protein (PBL Assay
Science, Piscataway, NJ, USA) was used as the standard. The limit of detection of the assay
was 0.022 fg.mL^{-1} and the mean of the blanks was 0.46 fg.mL^{-1}, including the dilution factor used
(5.5).

For the IFNβ assay, the 710906-9 IgG1, kappa, mouse monoclonal antibody (PBL Assay
Science) was used as a capture antibody after coating on paramagnetic beads (0.3 mg.mL^{-1}),
the 710323-9 IgG1, kappa, mouse monoclonal antibody (PBL Assay Science) was biotinylated
(biotin/antibody ratio = 40/1) and used as the detector antibody at a concentration of 1 μg.mL^{-1},
and the 11415 recombinant human IFNβ protein (PBL Assay Science) was used to quantify IFNβ
concentrations. The limit of detection of the assay was 0.029 pg.mL^{-1} and the mean of the blanks
was 0.17 pg.mL^{-1}, including the dilution factor used (5.5).

For the IFNλ3 assay, the 21730-1 IgG1, kappa, mouse monoclonal antibody (PBL Assay
Science) was used as a capture antibody after coating on paramagnetic beads (0.3 mg.mL^{-1}),

the MAB52591R IgG2a, mouse monoclonal antibody (R&D Systems, Minneapolis, MN, USA) was biotinylated (biotin/antibody ratio = 60/1) and used as the detector antibody at a concentration of 0.3 µg.mL^{-1}, and the 5259-IL-025/CF recombinant human IFNλ3 protein (R&D Systems) was used as calibrator. The limit of detection of the assay was 0.006 pg.mL^{-1} and the mean of the blanks was 0.82 pg.mL^{-1}, including the dilution factor used (5.5).

Statistics and reproducibility

Statistical analyses were performed with the Prism software v8.4.3 (GraphPad). Unless stated otherwise, we used the nonparametric Mann–Whitney or Kruskal-Wallis tests. Statistical significance was assigned when p values were <0.05. The nature of statistical tests and the number of experiments (n) are reported in the figure legends.

References

1. Davis, J. D. & Wypych, T. P. Cellular and functional heterogeneity of the airway epithelium. *Mucosal Immunology* **14**, 978–990 (2021).
2. Nikolić, M. Z., Sun, D. & Rawlins, E. L. Human lung development: recent progress and new challenges. *Development* **145**, dev163485 (2018).
3. Mahieu, L., Van Moll, L., De Vooght, L., Delputte, P. & Cos, P. In vitro modelling of bacterial pneumonia: a comparative analysis of widely applied complex cell culture models. *FEMS Microbiology Reviews* **48**, fuae007 (2024).
4. Huh, D. *et al.* Reconstituting Organ-Level Lung Functions on a Chip. *Science* **328**, 1662–1668 (2010).
5. Robinot, R. *et al.* SARS-CoV-2 infection induces the dedifferentiation of multiciliated cells and impairs mucociliary clearance. *Nat Commun* **12**, 4354 (2021).
6. Sucre, J. M. S. *et al.* Successful Establishment of Primary Type II Alveolar Epithelium with 3D Organotypic Coculture. *Am J Respir Cell Mol Biol* **59**, 158–166 (2018).
7. Dobbs, L. G. Isolation and culture of alveolar type II cells. *American Journal of Physiology-Lung Cellular and Molecular Physiology* **258**, L134–L147 (1990).
8. Barkauskas, C. E. *et al.* Type 2 alveolar cells are stem cells in adult lung. *J Clin Invest* **123**, 3025–3036 (2013).
9. Sachs, N. *et al.* Long-term expanding human airway organoids for disease modeling. *The EMBO Journal* **38**, 1–20 (2019).
10. Salahudeen, A. A. *et al.* Progenitor identification and SARS-CoV-2 infection in human distal lung organoids. *Nature* **588**, 670–675 (2020).
11. Dye, B. R. *et al.* In vitro generation of human pluripotent stem cell derived lung organoids. *eLife* **4**, 1–25 (2015).
12. Yamamoto, Y. *et al.* Long-term expansion of alveolar stem cells derived from human iPS cells in organoids. *Nature Methods* **14**, 1097–1106 (2017).
13. Jacob, A. *et al.* Differentiation of Human Pluripotent Stem Cells into Functional Lung Alveolar Epithelial Cells. *Cell Stem Cell* **21**, 472-488.e10 (2017).
14. Burgess, C. L. *et al.* Generation of human alveolar epithelial type I cells from pluripotent stem cells. *Cell Stem Cell* **31**, 657-675.e8 (2024).
15. Chen, Y.-W. *et al.* A three-dimensional model of human lung development and disease from pluripotent stem cells. *Nature Cell Biology* **19**, 542–549 (2017).

16. Dichtl, S., Posch*, W. & Wilflingseder, D. The breathtaking world of human respiratory in vitro
models: Investigating lung diseases and infections in 3D models, organoids, and lung-on-
chip. *European Journal of Immunology* **54**, 2250356 (2024).

17. Vazquez-Armendariz, A. I. & Tata, P. R. Recent advances in lung organoid development and
applications in disease modeling. *J Clin Invest* **133**, (2023).

18. Kim, M., Fevre, C., Lavina, M., Disson, O. & Lecuit, M. Live Imaging Reveals Listeria
Hijacking of E-Cadherin Recycling as It Crosses the Intestinal Barrier. *Current Biology* **31**,
1037-1047.e4 (2021).

19. Iakobachvili, N. *et al.* Mycobacteria–host interactions in human bronchiolar airway
organoids. *Molecular Microbiology* **117**, 682–692 (2022).

20. Youk, J. *et al.* Three-Dimensional Human Alveolar Stem Cell Culture Models Reveal
Infection Response to SARS-CoV-2. *Cell Stem Cell* **27**, 905-919.e10 (2020).

21. Walocha, R., Kim, M., Wong-Ng, J., Gobaa, S. & Sauvonnet, N. Organoids and organ-on-
chip technology for investigating host-microorganism interactions. *Microbes and Infection*
105319 (2024) doi:10.1016/j.micinf.2024.105319.

22. Grassart, A. *et al.* Bioengineered Human Organ-on-Chip Reveals Intestinal
Microenvironment and Mechanical Forces Impacting *Shigella* Infection. *Cell Host & Microbe*
26, 435-444.e4 (2019).

23. Stucki, A. O. *et al.* A lung-on-a-chip array with an integrated bio-inspired respiration
mechanism. *Lab Chip* **15**, 1302–1310 (2015).

24. Nikolić, M. Z. *et al.* Human embryonic lung epithelial tips are multipotent progenitors that can
be expanded in vitro as long-term self-renewing organoids. *eLife* **6**, 1–33 (2017).

25. Miller, A. J. *et al.* In Vitro Induction and In Vivo Engraftment of Lung Bud Tip Progenitor Cells
Derived from Human Pluripotent Stem Cells. *Stem Cell Reports* **10**, 101–119 (2018).

26. Lim, K. *et al.* Organoid modeling of human fetal lung alveolar development reveals
mechanisms of cell fate patterning and neonatal respiratory disease. *Cell Stem Cell* **30**, 20-
37.e9 (2023).

27. Gonzalez, R. F., Allen, L., Gonzales, L., Ballard, P. L. & Dobbs, L. G. HTII-280, a biomarker
specific to the apical plasma membrane of human lung alveolar type II cells. *Journal of*
Histochemistry and Cytochemistry **58**, 891–901 (2010).

28. van Riet, S. *et al.* Organoid-based expansion of patient-derived primary alveolar type 2 cells
for establishment of alveolus epithelial Lung-Chip cultures. *Am J Physiol Lung Cell Mol*
Physiol **322**, L526–L538 (2022).

29. Lim, K. *et al.* A novel human fetal lung-derived alveolar organoid model reveals mechanisms
of surfactant protein C maturation relevant to interstitial lung disease. *The EMBO Journal* 1–
26 (2025).

30. Lamers, M. M. *et al.* An organoid-derived bronchioalveolar model for SARS-CoV-2 infection
of human alveolar type II-like cells. *The EMBO Journal* **40**, 1–19 (2021).

31. Goltsis, O. *et al.* Influence of mesenchymal and biophysical components on distal lung
organoid differentiation. *Stem Cell Research & Therapy* **15**, 273 (2024).

32. Salaun, B. *et al.* CD208/dendritic cell-lysosomal associated membrane protein is a marker of
normal and transformed type II pneumocytes. *Am J Pathol* **164**, 861–871 (2004).

33. Treutlein, B. *et al.* Reconstructing lineage hierarchies of the distal lung epithelium using
single-cell RNA-seq. *Nature* **509**, 371–375 (2014).

34. Weiser, J. N., Ferreira, D. M. & Paton, J. C. Streptococcus pneumoniae: transmission,
colonization and invasion. *Nat Rev Microbiol* **16**, 355–367 (2018).
35. Connor, M. G. *et al.* The histone demethylase KDM6B fine-tunes the host response to
Streptococcus pneumoniae. *Nat Microbiol* **6**, 257–269 (2021).
36. Bomar, L., Brugger, S. D., Yost, B. H., Davies, S. S. & Lemon, K. P. Corynebacterium
accolens Releases Antipneumococcal Free Fatty Acids from Human Nostril and Skin
Surface Triacylglycerols. *mBio* **7**, e01725-01715 (2016).
37. Hui, K. P. Y. *et al.* SARS-CoV-2 Omicron variant replication in human bronchus and lung ex
vivo. *Nature* **603**, 715–720 (2022).
38. Lamers, M. M. *et al.* SARS-CoV-2 Omicron efficiently infects human airway, but not alveolar
epithelium. 2022.01.19.476898 Preprint at https://doi.org/10.1101/2022.01.19.476898
(2022).
39. Chiu, M. C. *et al.* A bipotential organoid model of respiratory epithelium recapitulates high
infectivity of SARS-CoV-2 Omicron variant. *Cell Discov* **8**, 1–15 (2022).
40. Huang, J. *et al.* SARS-CoV-2 Infection of Pluripotent Stem Cell-Derived Human Lung
Alveolar Type 2 Cells Elicits a Rapid Epithelial-Intrinsic Inflammatory Response. *Cell Stem
Cell* **27**, 962-973.e7 (2020).
41. Sempere, J. *et al.* Minilungs from Human Embryonic Stem Cells to Study the Interaction of
Streptococcus pneumoniae with the Respiratory Tract. *Microbiology Spectrum* **10**, e00453-
22 (2022).
42. Stucki, J. D. *et al.* Medium throughput breathing human primary cell alveolus-on-chip model.
Sci Rep **8**, 14359 (2018).
43. Bai, H. *et al.* Mechanical control of innate immune responses against viral infection revealed
in a human lung alveolus chip. *Nat Commun* **13**, 1928 (2022).
44. Barkal, L. J. *et al.* Microbial volatile communication in human organotypic lung models. *Nat
Commun* **8**, 1770 (2017).
45. Huang, D. *et al.* Reversed-engineered human alveolar lung-on-a-chip model. *Proceedings of
the National Academy of Sciences* **118**, e2016146118 (2021).
46. van der Vaart, J. *et al.* Modelling of primary ciliary dyskinesia using patient-derived airway
organoids. *EMBO reports* **22**, e52058 (2021).
47. Gonzales, L. W., Guttentag, S. H., Wade, K. C., Postle, A. D. & Ballard, P. L. Differentiation
of human pulmonary type II cells in vitro by glucocorticoid plus cAMP. *American Journal of
Physiology-Lung Cellular and Molecular Physiology* **283**, L940–L951 (2002).
48. Foster, C. D., Varghese, L. S., Skalina, R. B., Gonzales, L. W. & Guttentag, S. H. In Vitro
Transdifferentiation of Human Fetal Type II Cells Toward a Type I–like Cell. *Pediatr Res* **61**,
404–409 (2007).
49. Crapo, J. D., Barry, B. E., Gehr, P., Bachofen, M. & Weibel, E. R. Cell number and cell
characteristics of the normal human lung. *Am Rev Respir Dis* **126**, 332–337 (1982).
50. Karwelat, D. *et al.* Influenza virus-mediated suppression of bronchial Chitinase-3-like 1
secretion promotes secondary pneumococcal infection. *The FASEB Journal* **34**, 16432–
16448 (2020).
51. Sandgren, A. *et al.* Effect of Clonal and Serotype-Specific Properties on the Invasive
Capacity of Streptococcus pneumoniae. *The Journal of Infectious Diseases* **189**, 785–796
(2004).

52. Ghanem, E. N. B. *et al.* Extracellular Adenosine Protects against Streptococcus pneumoniae
Lung Infection by Regulating Pulmonary Neutrophil Recruitment. *PLOS Pathogens* **11**,
e1005126 (2015).
53. Horn, K. J. *et al.* Corynebacterium Species Inhibit Streptococcus pneumoniae Colonization
and Infection of the Mouse Airway. *Front Microbiol* **12**, 804935 (2021).
54. Lamers, M. M. & Haagmans, B. L. SARS-CoV-2 pathogenesis. *Nat Rev Microbiol* **20**, 270–
284 (2022).
55. Zhang, Y., Chen, S., Tian, Y. & Fu, X. Host factors of SARS-CoV-2 in infection, pathogenesis,
and long-term effects. *Front Cell Infect Microbiol* **14**, 1407261 (2024).
56. Planas, D. *et al.* Distinct evolution of SARS-CoV-2 Omicron XBB and BA.2.86/JN.1 lineages
combining increased fitness and antibody evasion. *Nat Commun* **15**, 2254 (2024).
57. Meganck, R. M. *et al.* SARS-CoV-2 variant of concern fitness and adaptation in primary
human airway epithelia. *Cell Reports* **43**, 114076 (2024).
58. Woodall, M. N. J. *et al.* Age-specific nasal epithelial responses to SARS-CoV-2 infection. *Nat
Microbiol* **9**, 1293–1311 (2024).
59. Katsura, H. *et al.* Human Lung Stem Cell-Based Alveolospheres Provide Insights into SARS-
CoV-2-Mediated Interferon Responses and Pneumocyte Dysfunction. *Cell Stem Cell* **27**,
890-904.e8 (2020).
60. Thacker, V. V. *et al.* Rapid endotheliitis and vascular damage characterize SARS-CoV-2
infection in a human lung-on-chip model. *EMBO Rep* **22**, e52744 (2021).
61. Domizio, J. D. *et al.* The cGAS-STING pathway drives type I IFN immunopathology in
COVID-19. *Nature* 145–151 (2022).
62. Shi, G. *et al.* Omicron Spike confers enhanced infectivity and interferon resistance to SARS-
CoV-2 in human nasal tissue. *Nat Commun* **15**, 889 (2024).
63. Mache, C. *et al.* SARS-CoV-2 Omicron variant is attenuated for replication in a polarized
human lung epithelial cell model. *Commun Biol* **5**, 1–8 (2022).
64. Masui, A. *et al.* Micro-patterned culture of iPSC-derived alveolar and airway cells
distinguishes SARS-CoV-2 variants. *Stem Cell Reports* 1–17 (2024).
65. Otter, C. J. *et al.* Interferon signaling in the nasal epithelium distinguishes among lethal and
common cold coronaviruses and mediates viral clearance. *Proceedings of the National
Academy of Sciences* **121**, e2402540121 (2024).
66. Dekkers, J. F. *et al.* High-resolution 3D imaging of fixed and cleared organoids. *Nat Protoc
14, 1756–1771 (2019).
67. Schindelin, J. *et al.* Fiji: an open-source platform for biological-image analysis. *Nature
Methods 2012 9:7* **9**, 676–682 (2012).
68. Bankhead, P. *et al.* QuPath: Open source software for digital pathology image analysis. *Sci
Rep* **7**, 16878 (2017).
69. Livak, K. J. & Schmittgen, T. D. Analysis of Relative Gene Expression Data Using Real-Time
Quantitative PCR and the 2–ΔΔCT Method. *Methods* **25**, 402–408 (2001).
70. Menberu, M. A. *et al. In vitro* and *in vivo* evaluation of probiotic properties of
Corynebacterium accolens isolated from the human nasal cavity. *Microbiological Research
255, 126927 (2022).
71. Planas, D. *et al.* Reduced sensitivity of SARS-CoV-2 variant Delta to antibody neutralization.
Nature **596**, 276–280 (2021).

72. Bruel, T. *et al.* Longitudinal analysis of serum neutralization of SARS-CoV-2 Omicron BA.2,
BA.4, and BA.5 in patients receiving monoclonal antibodies. *Cell Reports Medicine* **3**,
100850 (2022).
73. Smith, N. *et al.* Distinct systemic and mucosal immune responses during acute SARS-CoV-2
infection. *Nat Immunol* **22**, 1428–1439 (2021).
74. Llibre, A. *et al.* Plasma Type I IFN Protein Concentrations in Human Tuberculosis. *Front Cell
Infect Microbiol* **9**, 296 (2019).
75. Rodero, M. P. *et al.* Detection of interferon alpha protein reveals differential levels and
cellular sources in disease. *J Exp Med* **214**, 1547–1555 (2017).

Acknowledgements

The authors would like to thank Olivier Schwartz and Florence Guivel-Benhassine for providing
SARS-CoV-2 virus samples, Hugo Mouquet for providing anti-spike and anti-N antibodies and
Sarah Vreugde for sharing bacterial strains. We thank the Single Cell Biomarkers UTechS of the
Institut Pasteur for support in conducting this study. We acknowledge the Flow Cytometry
Platform at Institut Pasteur for support in conducting this work. We thank the Biomics Platform
(C2RT, Institut Pasteur) that is supported by France Génomique (ANR-10-INBS-09) and IBISA.
We gratefully acknowledge the UtechS Photonic BioImaging, (C2RT, Institut Pasteur) supported
by the French National Research Agency (France BioImaging; ANR-10–INBS–04; Investments
for the Future). Experimental work benefited from the support of ultrastructure bioimaging core
facilities of Institut Pasteur.

Funding

769   SG and LAC were supported by *the Pasteur COVID-19 RP call* for the COROCHIP project and
770   by the ANRS-23-PEPR-MIE-0006 *call in the frame of the* France 2030 program for the 3D-
771   LUNGO PEPR project. SG funded the purchase of chips with the support of *Institut Carnot
772   Pasteur Microbes et Santé (ANR 20 CARN 0023-01)* program. LAC was also supported by *the
773   Urgence COVID-19 Fundraising Campaign of Institut Pasteur (PFR4 project).*

Author contributions

BFF designed and conducted most of the experiments including organoid, organ-on-chip and
SARS-CoV-2 infections; JWG derived lung bud organoid cultures and performed organoid
differentiation experiments; MC carried out and analyzed bacterial infection experiments; HM
and MHK assisted with organ-on-chip experiments; RY contributed to organoid and organ-on-
chip experiments; VB performed Simoa analyses; VM performed electron microscopy; JDS and
HSM contributed to the organoid derivation protocol; DD supervised the Simoa experiments; MH
supervised the bacterial experiments, helped secure funding and edited the manuscript; NS
assisted with the experimental designs and reviewed the data; LAC contributed to project
conceptualization, SARS-CoV-2 infections and acquired funding for the project; SG

conceptualized the project, secured funding, designed experiments and analyzed the data; BFF
and SG wrote the manuscript. All authors read, edited and approved the final version.

Figures

Figure 1. Derivation of three types of human fetal lung organoids for the production of airway and alveolar cell types. (A) Schematic representation of the cell culture pipeline used to produce different fetal lung organoid types. (B) Brightfield images of multipotent lung organoids in self-renewing conditions at passage 1 (left) and passage 2 (right). (C) Expansion of the organoid cultures over multiple passages. Each line corresponds to one experiment with a distinct organoid line. (D) Multipotent lung organoids showing positive immunostainings for SOX2 and SOX9 markers (E) Alveolar organoids showing typical alveolar type 2 cells identified with HT2-280 and anti-pro-SFTPC immunolabeling. (E') Higher magnification of the region depicted in (E). (F) Immunostaining of airway organoids showing typical cell types including basal (KRT5+), ciliated

(ac-TUB+), club (SCGB1a1+) and goblet (MUC5AC+) cells. Scale bars: 75 µm (B), 50 µm (D and E), 20 µm (F).

Figure 2: Characterization of human airway-on-chip and alveoli-on-chip systems. (A) Schematic representation of the protocol used to derive alveolar and airway organoid-on-chip devices. (B) Immunolabelling of the alveoli-on-chip system. Left: AT1 (HT1-56+), AT2 (HT2-280+); middle: AT1 (RAGE+); right: AT2 (HT2-280+, pro-SFTPC+). (C) Percentage of cells labelled with the following alveolar markers: HT1-56, HT2-280 and pro-SFTPC. Each dot color corresponds to a different organoid line. Mean +/- standard deviation (SD) are plotted. (D-F) Comparative RT-qPCR analyses between multipotent organoids versus lung cells in the alveolar chips showing the relative expression of typical lung multipotent markers (SOX2 and SOX9) (D), the mature AT2 marker LAMP3 (E) and the mature AT1 marker CLIC5 (F) (D-F: Median +/- interquartile range are plotted and statistically significant differences are assessed with the Mann-Whitney test. n=4

independent experiments with 4 different samples). (G) Immunolabelling of airway cell types in the airway-on-chip system. Left: ciliated (ac-TUB+) and goblet (MUC5AC+) cells; middle: ciliated (ac-TUB+) and club (SCGB1A1+) cells; right: ciliated (ac-TUB+) and basal (KRT5+) cells. (H) Percentage of ciliated (ac-TUB+) and basal (KRT5+) cells in airway chips. Mean +/- SDs are plotted. (B, G) Scale bars: 50µm.

A
① Innoculation ② ALI ③ Lavages ④ IF

TIGR4
6B ST90
C. accolens

CFU (adhesion)
CFU (Prolif.)
CFU (Cross.)

6 min   24h

B *Inoculum*

Total CFU (log10)

ns
ns

TIGR4   6B ST90   C. accolens

C *Top channel eff. (24hpi)*

Total CFU (log10)

*
ns

TIGR4   6B ST90   C. accolens

D *Bottom channel eff. (24hpi)*

Total CFU (log10)

***
*

TIGR4   6B ST90   C. accolens

E

HT1-56/DAPI   HT2-280/DAPI   Bacteria/DAPI

Uninfected

TIGR4   *Streptococcus*

6B ST90   *Streptococcus*

C.accolens   *Coryne*

F

% AT2 cells

Uninfected   TIGR4   6B ST90   C.accolens

G

% AT1 cells

Uninfected   TIGR4   6B ST90   C.accolens

Figure 3: *Streptococcus pneumoniae* strains infect the human alveoli on-chip system. (A) Schematic representation of the infection protocol with *S. pneumoniae* strains (TIGR4 and 6B ST90) or *C. accolens* (commensal control). ALI: air-liquid-interface, IF: immunofluorescence (B) Colony forming unit (CFU) counts the bacteria inoculum. (C) CFU counts of respective bacteria collected after 24h in the top channel effluent. (D) CFU counts of respective bacteria collected after 24h in the bottom channel effluent. Median +/- interquartile range are plotted and statistically significant differences between bacteria were compared with Kruskal-Wallis test. (n= 3 independent experiments with 3 different organoid lines; geometric means of 2 technical counts per chip are shown). Dotted green in lines B-D: limit of detection value (20 total CFU). (E) Representative images of bacterium infection in alveolar chips. HT1-56 and HT2-280 immunolabeling identifies AT1 and AT2 cells, respectively. hpi: hours post-infection. Scale bars: 50µm. (F and G) Percentages of AT2 and AT1 cells upon bacterial infections on-chip. Mean +/- standard deviation, n=2 independent experiments. Minimum cell count per experiment above 20000.

Figure 4: SARS-CoV-2 variant Delta, but not Omicron BA.5, replicates in human alveolar chips. (A) Schematic representation of SARS-CoV-2 infection protocol in alveoli-on-chip system. ALI: air-liquid-interface (B) Detection of SARS-CoV-2 (Delta) infected cells by immunolabeling with anti-spike antibodies (red staining). HT2-280 antibody (in cyan) stains AT2 cells. Counterstaining with DAPI and F-actin (phalloidin, in gray). Orange arrows: spike and HT2-280 co-labelled cells. Scale bars: 50 μm. (C) Percentage of spike+ cells in Delta versus mock-infected conditions. Median +/- interquartile range (IQR) are plotted and statistically significant differences are compared with the Mann-Whitney test. (n= 4 independent experiments with 3 organoid lines). (D) Proportion of HT2-280+ cells among spike+ cells. Mean +/- standard deviation (SD) are plotted. (E) Kinetics of SARS-CoV-2 viral load in top channel effluents measured by RT-qPCR. Mean +/- SD are plotted. (F) Viral RNA load in top channel effluent at 4 dpi. Median +/- IQR are plotted and statistically significant differences are compared with the Mann-Whitney test. Each dot color represents a different organoid line. (E-F, n= 6 independent experiments with 6 organoid lines). (G) Viral RNA detection in cellular extracts at 4 dpi. Median +/- IQR are plotted and statistically significant differences are compared with the Mann-Whitney test. Each dot color represents a different organoid line. (n= 3 independent experiments with 3 organoid lines). (H) IFNα protein concentrations, expressed as IFNα17 equivalents, detected in top channel washes recovered at 4 dpi (n= 5 independent experiments with 5 organoid lines. Each dot color represents an organoid line. Median +/- interquartile range are plotted and Kruskal-Wallis tests are used for addressing statistically significant differences). (I) Relative RNA expression of ISG15, MX2 and IFIT1 at 4 dpi (n=3 independent experiments with 3 organoid lines Median +/- IQR are plotted and statistically significant differences are compared with the Kruskal-Wallis test. Each dot color represents a different organoid line).

Supplementary Figure 1: Multipotent lung organoids form a simple and tight epithelium on-chip. Cell adherent and tight junctions revealed with E-cadherin and ZO-1 immunolabeling, respectively. DAPI was used to counterstain nuclei. Scale bars: 100 µm.

Supplementary Figure 2: *Streptococcus pneumoniae* **infection in human alveolar chips.** (A) Quantification of number of cells per chip based on the ratio of nuclei number per um^2 chip area. (B) Streptococcus immunolabeling of bacteria specimens surrounding the bottom channel lung mesenchyme at 24 hours postinfection (hpi). Scale bars: 100 µm.

Supplementary Figure 3: Effects of SARS-CoV-2 infection on alveolar cell populations. (A) Quantification of number of cells per chip based on the ratio of nuclei number per um^2 chip area. (B) Percentage of total HT2-280+ cells per condition. Median +/- interquartile range are plotted and statistically significant differences are compared with the Kruskal-Wallis test.

Supplementary Figure 4: Ultrastructural features of AT2-like cells upon SARS-CoV-2 infection on-chip. (A) Scanning electron microscopy images evidence abundant small projections similar to microvilli in cells of an uninfected alveolar chip (arrows in A and A'). (B) Scanning electron microscopy images of a SARS-CoV-2 infected chip. (C) Viral particles observed at a higher resolution image than B.

Supplementary Figure 5: Comparison of SARS-CoV-2 variant infections in human alveolar chips. (A) Immunolabeling of SARS-CoV-2 infected cells with anti-NCP1-antibody (green). Alveolar cells are identified with an anti-pro-SFTPC antibody (magenta). DAPI is used to counterstain nuclei. Scale bars: 50 µm (B) Percentage of NCP1+ cells in different conditions at 4 dpi. (C) Percentage of pro-SFTPC+ cells in different conditions at 4 dpi. Median +/- interquartile range are plotted and statistically significant differences are compared with the Kruskal-Wallis test.

Supplementary Figure 6. IFN host-response to SARS-CoV-2 infection in alveolar chips. (A) IFNβ and (B) IFNλ3 proteins detected in top channel washes recovered at 4 days post-infection. (n=5 independent experiments with 5 different organoid lines. Each dot color represents an organoid line. Median +/- interquartile range are plotted and Kruskal-Wallis tests are used for addressing statistically significant differences).