Concurrent multimodal estimation of cortical excitation and inhibition shows that EEG-based estimates are linked to the ratio of glutamate to GABA

1   Aaron Cochrane[1], Luke Rosedahl[1], Masako Tamaki[2], Takeo Watanabe[1], Yuka Sasaki[1]

4   [1] Brown University

5   [2] RIKEN

Abstract

Neural excitation/inhibition (E/I) balance is dynamically regulated on multiple timescales. Adaptive changes in E/I balance changes can support healthy development, learning, and cognition, while disordered E/I balance is implicated in neurodevelopmental disorders, neurodegenerative disorders, and states of impaired vigilance. There has been growing interest in inferring E/I balance from efficient and noninvasive measurements such as electroencephalography (EEG), and several algorithms have been proposed to estimate E/I balance from EEG recordings. Despite promising results, there has been a lack of validation studies testing the underlying neurochemical changes leading to increased or decreased EEG-based E/I. Here we assess E/I balance with concurrent EEG and magnetic resonance spectroscopy (MRS) in humans. We estimate a standard measure of E/I balance, glutamate concentration divided by GABA concentration, and assess the correspondence between each candidate EEG-based E/I algorithm to this MRS-based estimate. Due to the methodological interest in both between-subjects research (e.g., comparing disordered to healthy E/I) as well as within-subjects research (e.g., comparing pre-intervention to post-intervention E/I), we quantify the associations between EEG-based E/I and MRS-based E/I separately for between-subjects and within-subjects comparisons. We find that each EEG-based E/I algorithm shows reliable and positive associations with MRS-based E/I. While these associations are evident for between-subjects comparisons, they are quite weak for within-subjects comparisons. Candidate EEG-based E/I algorithms are thus likely to be reflecting, at least in part, relative concentrations of cortical glutamate and GABA, although poor signal quality appears to limit these methods when attempting to increase temporal precision and identify within-subjects variations.

Introduction

Brain functioning is largely regulated through a balance, or dynamically changing imbalance, of excitatory and inhibitory (E/I) activities (Hensch et al. 1998; Shew et al. 2011). Such dynamics are closely linked to a variety of phenomena including neural plasticity (Bavelier et al. 2010; Hensch et al. 1998; Vogels et al. 2011), brain development (Hensch 2004; Takesian and Hensch 2013) disordered functioning (Bruining et al. 2020; Geertjens et al. 2022; Nifterick et al. 2023), and cognition-related topics (Hellyer et al. 2016) such as vigilance (Yang et al. 2023) and decision-making (Amil and Verschure 2021). Unfortunately, E/I is extremely difficult to measure non-invasively (Ahmad et al. 2022; Craven et al. 2022), and continued refinement of E/I measurement is a core component of advancing these disparate fields. A variety of methods are used to assess E/I in humans. These include the use of electrophysiological signatures of E/I (e.g., using electroencephalography, EEG; Ahmad et al. (2022) and using magnetic resonance spectroscopy (MRS) to identify concentrations of excitatory and inhibitory neurotransmitters (e.g., Glutamate and GABA, respectively; Shibata et al. (2017). Each of these approaches holds great promise and yet faces serious limitations.

EEG-based approaches to quantifying E/I come from several theoretical and computational sources, including scale-free dynamics (Gao, Peterson, and Voytek 2017), criticality theory (Poil et al. 2012), and neural mass modeling (Yokoyama and Kitajo 2023), and have been used to investigate physiological underpinnings of phenomena ranging from vigilance (Yang et al. 2023) to Alzheimer's disease (Nifterick et al. 2023). Each of these has proposed mechanistic bases for the links between a scalp-based electrical signal and cortical excitation or inhibition, but empirical data tracking brain regions' neurochemistry to EEG measures has been largely lacking (Ahmad et al. 2022; Geertjens et al. 2022). The difficulty in establishing the biological bases of EEG-based E/I measures is offset, however, in their relative advantages: EEG is widely available, is inexpensive to implement, and provides data from multiple brain regions simultaneously.

In contrast, the use of MRS to study E/I presents a different set of trade-offs. Studies of E/I using MRS have been conducted to study brain plasticity, such as that involved in sleep-dependent consolidation of perceptual learning (Tamaki et al. 2020) or developmental differences in GABA and plasticity (Frank et al. 2022). While MRS-based E/I has the distinct benefit of having a clear biological basis (i.e., neurotransmitter concentrations such as Glutamate and GABA), it is limited to a single cortical location with fairly low temporal resolution, is limited to measurement inside a MRI scanner (thereby precluding, for example, measurement in certain populations or during certain behaviors), and is relatively expensive.

The field of E/I research using both EEG and MRS would benefit from linking the two domains. There have been multiple reports of associations between MRS-measured GABA concentrations with electrophysiological frequency bands (e.g., Balz et al. (2016); Edden et al. (2009)), but each of these studies measured

electrophysiology and MRS in separate sessions and were therefore limited to stable inter-individual correlations

and were insensitive to functional changes that may occur on short timescales. Shorter timescales of change are

necessary to understand in the context of many fields, including neuroplasticity, pharmacological interventions,

and cognitive consequences of varying E/I.

There remains a clear need, then, for multimodal investigations of EEG-E/I in conjunction with MRS-E/I. Only

by simultaneously measuring the two forms of E/I can both within-subject and between-subject variations in each

of the two measures be assessed. It is possible that the two forms of E/I are empirically distinct from each other,

for example, if EEG-E/I reflects rapidly-changing excitation and inhibition while MRS-E/I reflects E/I that only

changes on a slower timescale, if EEG-E/I reflects specific neural subpopulations generating oscillatory signatures

(Scheeringa and Fries 2019; Shafiei et al. 2023) while MRS-EI reflects the aggregate concentration of certain

molecules in a large voxel, or if the relatively superficial neural populations generating scalp-measured

electrophysiological changes display different dynamics than the relatively deep aggregate population contained

within a MRS voxel. In contrast, it is possible that the two forms of E/I are indexing the same underlying

biological processes, that is, the relative concentrations of the primary excitatory vs. inhibitory neurotransmitters

in a large region of cortex. Here we address these questions by linking visual-area MRS and EEG recorded in

participants who had completed a visual perceptual learning task.

Materials and Methods

Experimental Design and Statistical Analysis

We re-analyzed data from Study 1 of Tamaki et al. (2020), with 18 subjects each sleeping for between 60 and 90

minutes after completing a perceptual learning task. EEG (polysomnography; PSG) and MRS were recorded

throughout the sleep. To assess the extent to which candidate EEG-based measures of E/I were reflecting the

changes in neurotransmitter concentrations measured by MRS-based E/I, we first estimated a single E/I value

using each method for each 5-minute period of the recorded data. We then used cross-validation to estimate out-

of-sample variance explained when predicting MRS-based E/I using EEG-based E/I ($dRsq_{oos}$; see below). We

separated these estimates into between-subjects and within-subjects analyses; while much previous work has

focused on the relatively stable differences between individuals' levels of excitation and inhibition (Balz et al.

2016; Edden et al. 2009), within-subjects variation in E/I are central to many research domains such as vigilance

(Yang et al. 2023) and learning-related neuroplasticity (Bavelier et al. 2010; Tamaki et al. 2020) because these

topics are inherently concerned with intra-individual changes. Separating these results helps identify both the

overall associations between MRS-E/I and EEG-E/I as well as their relative isomorphism in the context of

different applications.

For within-subjects and between-subjects analyses we used two statistical approaches in conjunction. In the first,

in order to best estimate the effect size (i.e., correlation coefficient) between E/I measures, we fit a Bayesian

multivariate linear mixed-effects model (BMVLMEM) using **brms** in R (Bürkner 2017). Using this approach we

specified each of the candidate E/I measures as variables to predict (i.e., "response" variables). Each response

variable was fit with an overall intercept, a random-effects by-participant intercept, and a fixed-effect

autoregressive term with an order of one to account for non-independence of temporally proximal measurements.

Within the multivariate model the correlations across the response variables of the by-participant intercepts were

estimated (in brms "Wilkensen" notation the formula for each E/I measure was `response ~ 1 + ar(time,`

`order = 1) + (1 | subjectR | subjectID)`). In this way we estimated simultaneously the between-

subjects effects, in the form of the correlations between different response variables' by-participant intercepts, as

well as within-subjects effects, in the form of the residual correlations remaining after accounting for the

autoregressive and by-subject terms. We report these two correlations, both between-subjects and within-subjects,

as our best estimate of the links between each measure. In addition, this model included an estimation of the

remainder of the correlation matrix (e.g., between all EEG-E/I measures as well as estimates of Glx and GABA+);

we report these correlations to determine the extent to which our EEG measures are isomorphic and are

potentially more reflective of Glx or GABA+ variations (see Figure S1).

As tests of reliability, and in consideration of the possibility of overfitting our data, we augmented our

BMVLMEM with a maximum-likelihood cross-validation approach. This approach involves iteratively and

randomly partitioning the data into "training" and "test" sets, fitting a maximum-likelihood regression model

predicting MRS-E/I to the training set, and assessing the R-squared (i.e., proportional reduction in sum of squared

error) on the test set. Between-subjects and within-subjects analyses were necessarily implemented somewhat

differently. Between-subjects cross-validation involved first calculating the subject-level means for each value,

and then using simple random partitioning of this aggregated dataset into 80%/20% subsets, with data being fit

with a bivariate robust linear model to 80% of the data and out-of-sample delta R-squared ($dRsq_{oos}$) assessed on

the remaining 20%. In contrast, within-subjects cross-validation involved randomly selecting a single datapoint

from each participant to keep as a test set, fitting a linear model including a participant covariate (i.e., to control

for participant-level means in MRS-E/I) to the remaining training set, and then assessing the proportional change

in out-of-sample squared error ($dRsq_{oos}$) when compared to a model with only the participant covariate as a

predictor. To control for between-subjects variations in EEG-based E/I measures, for each of these within-subjects

cross-validation models the EEG-E/I measure was first normalized within participants (i.e., Yeo-Johnson

transformed to minimize univariate skew, and then z-scored). As will be seen, fairly large between-subjects

effects are evident, which limits the amount of remaining variance left to be explained by within-subjects effects.

We report the median $dRsq_{oos}$ of 500 resampled models for each EEG-E/I measure.

MRS-based E/I

MRS-based E/I is typically implemented by identifying a brain region of interest (e.g., early visual areas) and placing a single voxel in that region. Using magnetic resonance, the spectroscopic signal is then measured in that voxel over the course of several minutes and the relative concentrations of excitatory and inhibitory neurotransmitters are estimated (Shibata et al. 2017). These concentrations can then be examined separately or combined into a single measure of E/I. Using such methods it would be considered high spatial resolution to have a single voxel that is 15mm per side and a high temporal resolution to have one measurement every 2 minutes. The relative trade-offs of MRS measures of E/I are clear, then: While MRS-E/I is a relatively direct measure of actual molecules of interest within the brain, this measurement technique is resource-intensive, spatially and temporally imprecise, and the measured spectrum is itself a noisy measure. Indeed, the estimation of neurotransmitter concentrations has no consensus standard and results from a single MR sequence may vary depending on the estimation algorithm (e.g., inter-algorithm correlations between GABA concentration estimates ranging from below .1 to above .8; Craven et al. (2022).

Our primary results use the originally-published MRS data (Tamaki et al. (2020); Study 1), wherein a MEGA-PRESS sequence was used (1.25s TR; $2.2 \times 2.2 \times 2.2$ cm voxel, placed on midline in early visual cortices; 3 Tesla Siemens scanner). We made two modifications to the data reported in the previous publication, namely, partial volume correction (PVC) to account for relative proportions of grey matter (i.e., to mitigate the influence of grey matter proportion on associations with EEG measures) and separation into 5-minute sections of time instead of 10-minute time sections (mean per participant = 13.1, sd = 3.6). As with the originally-published work, we used LCModel software (Provencher 1993) to identify Glx and GABA+ concentrations normalized to NAA. We then estimate MRS-E/I ratio by dividing Glx by GABA+. We chose to continue using LCModel to estimate Glx and GABA+, rather than other candidate softwares, for several reasons: To maintain congruence with the original findings, because reliable results in the original paper indicate that E/I from LCModel was reflecting true underlying cortical changes, and because LCModel has been found to provide the most "typical" estimates of neurotransmitter concentrations (i.e., highest correlation of GABA estimates with the median of estimates from other algorithms; Craven et al. (2022).

EEG-based measures of E/I

We assessed four approaches to inferring E/I ratio from EEG (i.e., EEG-E/I), with three of these approaches being separated into three canonical frequency bands, leading to 10 EEG-E/I measures total to test associations with MRS-E/I. The first, overall amplitude, is not a measure of E/I *per se*, but associations between MRS-E/I within specific frequency bands have been observed previously (e.g., theta and low beta; Tamaki et al. (2020)). Following amplitude, we assessed EEG-E/I derived from scale-free dynamics reflected by the 1/f power slope (Gao, Peterson, and Voytek 2017), an extension of this scale-free perspective that derives from criticality theory

(Bruining et al. 2020), and a set of estimates utilizing a neural mass model of excitatory and inhibitory dynamics

(Yokoyama and Kitajo 2023).

For details of processing steps such as MR artifact rejection see Methods in Tamaki et al. (2020). In all analyses

we report EEG measures estimated from the closest 7 electrodes to the MRS voxel (i.e., most posterior; Oz, O1,

O2, PO3, PO4, & PO7; we replicated the overall pattern of our results when examining only Oz, O1, & O2). We

applied each of the following analysis approaches to each 5-minute measurement period separately.

Frequency band amplitude

As one of the most basic measures identifiable within EEG signals, we included simple amplitude (i.e., square

root of power) measures from several frequency bands. We used finite impulse response (FIR) bandpass filtering

to isolate theta (5-8Hz), alpha (8-12Hz), and low beta (12-20Hz) bands, estimated the amplitude in 5-second

windows (see section *EEG-E/I from criticality theory*), and then calculated the trimmed mean amplitude for each

5-minute measurement period (i.e., after removing the 5% lowest and 5% highest amplitude windows, calculating

the arithmetic mean of the amplitudes within the remaining windows). We would not expect this approach to

produce diverging results from, for example, simply averaging the amplitude across the entire 5-minute period.

While there have been some indications of links between frequency bands and E/I balance (Balz et al. 2016;

Duncan, Wiebking, and Northoff 2014; Porjesz et al. 2002), we include amplitude here primarily as "a baseline"

approach. That is, as an algorithmically simple derivation from resting-state EEG, we consider amplitudes to be

valuable measures to assess overall data reliability (e.g., see section *Across-channel correlations*), test the

divergence of EEG-E/I measures from amplitudes (see section *Associations among EEG measures*), as well as

providing parsimonious reflections of MRS measures if the relevant correlations were to be found.

EEG-E/I from 1/f slope ("1/f")

Another approach derives from the widespread observation that neural activity, in temporal, spatial, and

frequency domains, displays scale-free (power-law, or "1/f" when in the frequency domain) dynamics (Hahn et al.

2017; He 2014; Linkenkaer-Hansen et al. 2001). Computational derivations of the slopes of these 1/f distributions

have been linked to cortical E/I balance (Gao, Peterson, and Voytek 2017; Lendner et al. 2020). As with previous

work (Gao, Peterson, and Voytek 2017; Yokoyama and Kitajo 2023) we estimated the 1/f slope coefficient

between 30Hz and 50Hz (bandpass filtered) for each window of interest; we implemented this approach with

Python scripts available in conjunction with Yokoyama and Kitajo (2023). The following approach, based in

criticality theory, is an elaboration of these principles of scale-free dynamics.

EEG-E/I from criticality theory ("fE/I")

A third approach is grounded in the computational biology framework of criticality, which characterizes the

nature of many biological systems as existing at the cusp of state transitions (Bak 1999; Márton et al. 2014; Poil et

al. 2012). One application of this theoretical approach is to neural criticality, which proposes that neural activity

exists approximately at a balance between excitation and inhibition in order to maximize the flexibility of macro-

scale neural computations (Avramiea et al. 2022; Cocchi et al. 2017; Shew et al. 2011). To estimate E/I EEG, we

use an algorithm designed to index excitation/inhibition ratio (Juarez Martinez et al. 2019; Ahmad et al. 2022),

which is based in criticality theory and has been validated in computational, pharmacological (Bruining et al.

2020), and animal models (Kat et al. 2024). We applied this approach to the same three bandpass-filtered

frequency bands we considered when examining simple amplitude measures. In brief, this algorithm first

determines the degree of *criticality* (here, linked to the exponent of Detrended Fluctuation Analysis (DFA) Long-

Range Temporal Correlations (LRTC)). This uses links between the window size of an analyzed signal and the

variability in its cumulative detrended amplitude profile. Second, this degree of criticality is linked to the signal

amplitude; in inhibitory-dominant regimes these two values should be positively correlated, while in excitation-

dominant regimes they should be anticorrelated (Bruining et al. 2020). Using MATLAB code published in

conjunction with Bruining et al. (2020), fE/I values were estimated for each of the above-defined frequency bands

and then within- and between-subject correlations with MRS-E/I were estimated. As in the publication

establishing the method, estimation of fE/I used a window size of 5 seconds with an inter-window overlap of .8

(Bruining et al. 2020); we also tested the effects of increasing the window sizes to 15 seconds and found minimal

changes, and here report only 5-second window sizes. For each 5-minute measurement period we also estimated

the DFA coefficient and excluded measurement periods with DFA below .6, as recommended by the developers

of this approach (Bruining et al. 2020; Diachenko et al. 2024).

EEG-E/I from neural mass model ("NMM-E/I")

A fourth approach comes from a body of work instantiating neural mass models, that is, computational models of

small-scale neural circuitry such as cortical columns; such models can be scaled up and can simulate complex

dynamics and characterize dynamic and nonlinear features of neural activity (Cooray, Rosch, and Friston 2023;

Jansen and Rit 1995). Yokoyama and Kitajo (2023) extended earlier neural mass models (Jansen and Rit 1995),

which had been developed for purposes such as modeling epileptic dynamics, with a key insight: such models can

include estimates of the relative activity of circuits' excitatory and inhibitory neural populations. Following this,

and so they could compare to the 1/f approach described above, Yokoyama and Kitajo (2023) showed that the E/I

ratio derived from such models better captured the fluctuating activity of humans' electrophysiological activity

during sleep. This approach to estimating E/I ratio is quite new, however, and to our knowledge has not been

widely assessed. We estimated this NMM-E/I using Python code available in conjunction with Yokoyama and

Kitajo (2023) for each 5-minute measurement period, and for each of the canonical frequency bands described

above (ie., theta, alpha, & low Beta). We also tested the NMM-E/I on a single wider set of frequencies, more

similar to the approach implemented in Yokoyama and Kitajo (2023), on bandpass-filtered data between 5Hz and

20Hz (i.e., the lower and upper limits of the three frequency bands considered in the above analyses). The

resulting NMM-E/I values did not improve correpondence to MRS-E/I relative to the values estimated from

distinct canonical frequency bands.

EEG-E/I reliability assessment

We implemented a cross-channel correlation approach to estimate a heuristic reliability of our EEG-based

measures. Due to basic principles of volume conduction (Burle et al. 2015), as well as our intention to measure

relatively large-scale variations in E/I, we expected highly homogeneous estimates of E/I across our measured

channels. Because each of our EEG-E/I methods involved first estimating E/I for each channel we were able to,

within each method, estimate the cross-channel correlations. We tested these correlations within subjects by

finding the pairwise cross-channel partial rank correlations in estimated E/I values, partialling out participant-

level intercepts. We then report the median cross-channel partial rank correlation. While we do not have a strict

reliability criterion for these correlation measures, we believe that the relative cross-channel correlations should

provide a general sense of the computational stability of the measures (see Figure S2).

Between-subjects variability relative to within-subjects variability

In light of differences in between-subjects relative to within-subjects correspondences between EEG-based and

MRS-based measures of E/I, we sought to quantify the relative between-subjects and within-subjects variances for

each measure. To determine between-subjects variances we first found each participant's median value for each

measure, then calculated the variances of these values. To determine within-subjects variances we first calculated

each participant's variance for each measure, then found the median of these variances. The ratio of between-

subjects variance to within-subjects variances is reported in Figure S3.

<div align="center">**Results**</div>

We first assessed data quality and excluded 5-minute windows in which at least one E/I measure demonstrated

high noise relative to signal. MRS data quality was quantified using the ratio of the uncertainty of molecules'

estimated concentration relative to the corresponding estimate; as in the originally published work, a 20% level of

this "Cramer-Rao lower bound" was used for MRS exclusion. We also assessed EEG quality using the detrended

fluctuation analysis (DFA) exponent coefficient criterion which Bruining et al. (2020) describe in conjunction

with the fE/I method; this involves estimating the DFA coefficient and excluding windows in which the

coefficient is less than .6. While the original sample included 18 participants' datasets, three participants had only

three or fewer remaining MRS values and were excluded from further analysis. This left 15 participants data, with

a mean number of 5-minute samples of 13.1.

Between-subjects associations between EEG measures and MRS-based E/I

Inter-individual differences in certain measures of EEG-E/I were correlated with MRS-E/I (see Figure 1). Each of

the candidate algorithms produced E/I estimates with more than 5% of predictive variance explained in at least

one frequency band (i.e., $dRsq_{oos} > .05$); we consider each of the links with positive $dRsq_{oos}$ values to be reliable.

The criticality-based fE/I approach derived from alpha oscillatory activity was most closely linked to MRS-E/I

across participants, followed by 1/f slope, then NMM-E/I derived from low beta activity, and then fE/I derived

from theta band. All other measures, including overall amplitudes, were not reliably predictive of inter-individual

differences in MRS-E/I

Figure 1. Between-subject associations between EEG-E/I measures and MRS-E/I. (A) Out-of-sample [cross-

validated predictive] proportion of variance explained ($dRsq_{oos}$). (B) Correlation of by-participant random

intercepts from the Bayesian multivariate mixed-effects model. Each panel is sorted by descending $dRsq_{oos}$.

Dashed line separates variables explaining reliable out-of-sample variance with those that do not; we consider

all associations above the dashed line to be reliable.

Within-subjects associations between EEG measures and MRS-based E/I

After controlling for participant-level effects, several approaches to EEG-E/I were reliably associated with MRS-

E/I (i.e., $dRsq_{oos} > 0$; see Figure 2). While this wide range of EEG-E/I approaches providing within-subjects out-

of-sample predictiveness for MRS-E/I is promising, a closer investigation of the effects tempers this promise

somewhat. First, although out-of-sample variance explained was positive, it was quite small, with only simple

amplitude measures providing more than 1% of predictive value. Second, of the EEG measures with positive dRsq$_{oos}$ values, only theta amplitude had a positive correlation with MRS-E/I (r = 0.082). These results are in striking contrast to the between-subjects findings, where a relatively large amount of variance was explained with positive relations between EEG-E/I and MRS-E/I measures.

Figure 2. Within-subject associations between EEG-E/I measures and MRS-E/I. (A) Out-of-sample [cross-validated predictive] proportion of variance explained (dRsq$_{oos}$). (B) Correlation of by-participant random intercepts from the Bayesian multivariate mixed-effects model. Each panel is sorted by descending dRsq$_{oos}$. Dashed line indicates whether variables explained out-of-sample variance or did not; we consider all associations above the dashed line to be reliable.

Discussion

We have seen that EEG could be used to estimate E/I in a manner that is compatible with, but not wholly overlapping with, variations in E/I estimated from MRS. This mechanistic basis in the ratio between glutamate and GABA was true for several distinct approaches to EEG-E/I. In particular, EEG-E/I between-subjects was associated with MRS-E/I between-subjects, to the extent that they are comparable to (or even larger than) the correlations between MRS neurotransmitter concentrations found when comparing different estimation algorithms (Craven et al. 2022). In particular, the "functional E/I" algorithm derived from criticality theory was most closely associated with between-subjects differences in MRS-E/I; this reliance on temporal variations in alpha power

corroborates the work of the developers of the fE/I algorithm (Bruining et al. 2020; Diachenko et al. 2024).

Beyond this criticality-based EEG-E/I measure, each of the other EEG-E/I approaches were also positively

associated with between-subjects variations in MRS-E/I. These results showed broad support for each of the

approaches, which were themselves not highly correlated (see Figure 3). Future work may benefit from

combining these approaches to best characterize cortical E/I. The relative contribution of estimates from separate

frequencies is of particular interest; the generative mechanisms underlying variations in theta, alpha, and beta

rhythms provides the eletrophysiologist with a rich and multidimensional signal from which to infer shifts in

cortical states. Indeed, independent frequency bands have been previously associated with glutamate and/or

GABA levels (Porjesz et al. 2002; Tamaki et al. 2020), and a broad spectrum of frequencies showed links to

MRS-E/I when analyzed with different algorithms in the current work (i.e., theta and alpha analyzed with fE/I;

low beta analyzed with a neural mass model; gamma when calculating a 1/f slope). We believe that combining

amplitude and E/I estimates from several frequency bands may provide additional precision, but we are hesitant to

attempt such a combinatorial search within our limited dataset.

In contrast to between-subjects patterns of E/I balance, within-subjects associations between MRS-E/I and EEG-

E/I were much weaker; indeed, only simple theta amplitude was reliably and positively associated with within-

subjects variations in MRS-E/I. The reasons for the weakness of these associations cannot be known with

certainty, but it is likely to arise partly to a decrease in signal strength and an increase in noise when compared to

between-subject comparisons. That is, E/I is likely to vary a small amount within the course of 60-90 minutes of

sleep, when compared to inter-individual differences in E/I, thereby limiting the magnitude of the variations we

were able to observe. At the same time, estimating EEG-E/I as well as MRS-E/I on windows of only 5 minutes is

likely to lead to relatively noisy estimates; while shorter measurement windows measuring GABA with MRS

have been used in the past (Frank et al. 2022), longer windows (e.g., 10 minutes) are more standard.

Unfortunately our available data is limited enough that estimating such within-subjects associations between

EEG-E/I and MRS-E/I appears to be largely overwhelmed by the reduced signal-to-noise ratio; the MR context of

these estimates may also have itself reduced data quality in estimating EEG-based E/I.

There may be alternative sources of divergences between EEG-based and MRS-based E/I ratios. On one hand, the

neural mechanisms giving rise to each of our measured signals may be distinct; while MRS measured bulk

concentrations of molecules in a relatively deep and spatially precise portion of posterior cortex, our EEG

measurements were reflecting temporal variations in the activity of relatively superficial and widespread cortical

neurons. Similarly, the causal determinants of oscillatory activities, and their variations over time, are not yet

known with temporal or molecular precision; it may be the case that cell types, neuromodulators, or long-range

connections are influencing EEG signals which would not be reflected in the measurement of Glx and GABA+ in

a single voxel within posterior cortices (Ahmad et al. 2022).

Our results also reinforce the fact that the phenomena we are considering may be varying over time on

fundamentally different timescales. One of the most notable functions of variations in excitation and inhibition are

their changes over developmental time, that is, years or even decades (Hensch 2004; Takesian and Hensch 2013),

which in contrast, cognition-related modulation of E/I would necessarily operate on sub-minute timescales

(Hellyer et al. 2016). The methods we considered provide clear tendencies toward longer or shorter timescales,

with MRS-based measures needing a minimum of five minutes, NMM-based measures providing estimates which

are nominally relevant on the sub-second level, and the other methods falling somewhere in between. It may be

possible, then, that these measures simply do not reflect underlying cortical changes that operate on similar

timescales, leading to a lack of evident relations between them.

Robust links between EEG-E/I and MRS-E/I at a between-subjects level provides validation for EEG-based

measures to be used in certain areas of research for which they have developed, namely, in investigating special

populations such as individuals with ASD (Bruining et al. 2020), Alzheimer's disease (Nifterick et al. 2023), or

monogenetic neurodevelopmental disorders (Geertjens et al. 2022). Our results indicate that the between-person

differences in E/I revealed by contrasting these groups with healthy controls are likely to have common

mechanistic bases whether they are measured using EEG or MRS. In contrast, for research which relies heavily on

both within-person and between-person contrasts, such as the study of neuroplasticity (Shibata et al. 2017) or

vigilance (Yang et al. 2023), EEG-E/I measures may not allow the assumption of mechanistic isomorphism.

While our results indicate that several recently developed algorithms for measuring E/I using EEG are indeed

reflecting variations in the cortical ratio of glutamate to GABA, several limitations our present in our findings.

The size of our dataset does not allow for precise estimates of effect sizes nor compelling rejection of candidate

EEG-E/I measures, and the study of sleeping humans may lead to idiosyncratic results when contrasted to

wakefulness; each of these limitations motivates future comparisons using concurrent multimodal imaging in

awake humans. Further, with larger sample sizes, it would be possible to identify several additional useful aspects

of links between EEG-E/I and neurotransmitter concentrations, such as functional nonlinearities or optimal

combinations of EEG-E/I measures.

References

Ahmad, Jumana, Claire Ellis, Robert Leech, Bradley Voytek, Pilar Garces, Emily Jones, Jan Buitelaar, et al. 2022. "From Mechanisms to Markers: Novel Noninvasive EEG Proxy Markers of the Neural Excitation and Inhibition System in Humans." *Translational Psychiatry* 12 (1): 467. https://doi.org/10.1038/s41398-022-02218-z.

Amil, Adrián F, and Paul F M J Verschure. 2021. "Supercritical Dynamics at the Edge-of-Chaos Underlies Optimal Decision-Making." *Journal of Physics: Complexity* 2 (4): 045017. https://doi.org/10.1088/2632-072X/ac3ad2.

Avramiea, Arthur-Ervin, Anas Masood, Huibert D. Mansvelder, and Klaus Linkenkaer-Hansen. 2022. "Long-Range Amplitude Coupling Is Optimized for Brain Networks That Function at Criticality." *The Journal of Neuroscience: The Official Journal of the Society for Neuroscience* 42 (11): 2221–33. https://doi.org/10.1523/JNEUROSCI.1095-21.2022.

Bak, Per. 1999. *How Nature Works: The Science of Self-Organized Criticality*. 1., softcover pr. New York [Heidelberg]: Copernicus.

Balz, Johanna, Julian Keil, Yadira Roa Romero, Ralf Mekle, Florian Schubert, Semiha Aydin, Bernd Ittermann, Jürgen Gallinat, and Daniel Senkowski. 2016. "GABA Concentration in Superior Temporal Sulcus Predicts Gamma Power and Perception in the Sound-Induced Flash Illusion." *NeuroImage* 125 (January): 724–30. https://doi.org/10.1016/j.neuroimage.2015.10.087.

Bavelier, Daphne, Dennis M. Levi, Roger W. Li, Yang Dan, and T. K. Hensch. 2010. "Removing Brakes on Adult Brain Plasticity: From Molecular to Behavioral Interventions." *Journal of Neuroscience* 30 (45): 14964–71. https://doi.org/10.1523/JNEUROSCI.4812-10.2010.

Bruining, Hilgo, Richard Hardstone, Erika L. Juarez-Martinez, Jan Sprengers, Arthur-Ervin Avramiea, Sonja Simpraga, Simon J. Houtman, et al. 2020. "Measurement of Excitation-Inhibition Ratio in Autism Spectrum Disorder Using Critical Brain Dynamics." *Scientific Reports* 10 (1): 9195. https://doi.org/10.1038/s41598-020-65500-4.

Burle, Borís, Laure Spieser, Clémence Roger, Laurence Casini, Thierry Hasbroucq, and Franck Vidal. 2015. "Spatial and Temporal Resolutions of EEG: Is It Really Black and White? A Scalp Current Density View." *International Journal of Psychophysiology* 97 (3): 210–20. https://doi.org/10.1016/j.ijpsycho.2015.05.004.

Bürkner, Paul-Christian. 2017. "**Brms** : An *r* Package for Bayesian Multilevel Models Using *Stan*." *Journal of Statistical Software* 80 (1). https://doi.org/10.18637/jss.v080.i01.

Cocchi, Luca, Leonardo L. Gollo, Andrew Zalesky, and Michael Breakspear. 2017. "Criticality in the Brain: A Synthesis of Neurobiology, Models and Cognition." *Progress in Neurobiology* 158 (November): 132–52. https://doi.org/10.1016/j.pneurobio.2017.07.002.

Cooray, Gerald Kaushallye, Richard Ewald Rosch, and Karl John Friston. 2023. "Global Dynamics of Neural Mass Models." *PLoS Computational Biology* 19 (2): e1010915. https://doi.org/10.1371/journal.pcbi.1010915.

Craven, Alexander R., Pallab K. Bhattacharyya, William T. Clarke, Ulrike Dydak, Richard A. E. Edden, Lars Ersland, Pravat K. Mandal, et al. 2022. "Comparison of Seven Modelling Algorithms for γ-Aminobutyric Acid-Edited Proton Magnetic Resonance Spectroscopy." *NMR in Biomedicine* 35 (7): e4702. https://doi.org/10.1002/nbm.4702.

Diachenko, Marina, Additya Sharma, Dirk J. A. Smit, Huibert D. Mansvelder, Hilgo Bruining, Eco De Geus, Arthur-Ervin Avramiea, and Klaus Linkenkaer-Hansen. 2024. "Functional Excitation-Inhibition Ratio Indicates Near-Critical Oscillations Across Frequencies." *Imaging Neuroscience* 2 (October): 1–17. https://doi.org/10.1162/imag_a_00318.

Duncan, Niall W., Christine Wiebking, and Georg Northoff. 2014. "Associations of Regional GABA and Glutamate with Intrinsic and Extrinsic Neural Activity in Humans—a Review of Multimodal Imaging Studies." *Neuroscience and Biobehavioral Reviews* 47 (November): 36–52. https://doi.org/10.1016/j.neubiorev.2014.07.016.

Edden, Richard A. E., Suresh D. Muthukumaraswamy, Tom C. A. Freeman, and Krish D. Singh. 2009. "Orientation Discrimination Performance Is Predicted by GABA Concentration and Gamma Oscillation Frequency in Human Primary Visual Cortex." *The Journal of Neuroscience: The Official Journal of the Society for Neuroscience* 29 (50): 15721–26. https://doi.org/10.1523/JNEUROSCI.4426-09.2009.

Frank, Sebastian M., Markus Becker, Andrea Qi, Patricia Geiger, Ulrike I. Frank, Luke A. Rosedahl, Wilhelm M. Malloni, Yuka Sasaki, Mark W. Greenlee, and Takeo Watanabe. 2022. "Efficient Learning in Children with Rapid GABA Boosting During and After Training." *Current Biology: CB* 32 (23): 5022–5030.e7. https://doi.org/10.1016/j.cub.2022.10.021.

Gao, Richard, Erik J. Peterson, and Bradley Voytek. 2017. "Inferring Synaptic Excitation/Inhibition Balance from Field Potentials." *NeuroImage* 158 (September): 70–78. https://doi.org/10.1016/j.neuroimage.2017.06.078.

Geertjens, Lisa, Torben W. van Voorst, Arianne Bouman, Maaike A. van Boven, Tjitske Kleefstra, Matthijs Verhage, Klaus Linkenkaer-Hansen, Nael Nadif Kasri, L. Niels Cornelisse, and Hilgo Bruining. 2022. "Following Excitation/Inhibition Ratio Homeostasis from Synapse to EEG in Monogenetic Neurodevelopmental Disorders." *Genes* 13 (2): 390. https://doi.org/10.3390/genes13020390.

Hahn, Gerald, Adrian Ponce-Alvarez, Cyril Monier, Giacomo Benvenuti, Arvind Kumar, Frédéric Chavane, Gustavo Deco, and Yves Frégnac. 2017. "Spontaneous Cortical Activity Is Transiently Poised Close to Criticality." *PLoS Computational Biology* 13 (5): e1005543. https://doi.org/10.1371/journal.pcbi.1005543.

He, Biyu J. 2014. "Scale-Free Brain Activity: Past, Present, and Future." *Trends in Cognitive Sciences* 18 (9): 480–87. https://doi.org/10.1016/j.tics.2014.04.003.

Hellyer, Peter J., Barbara Jachs, Claudia Clopath, and Robert Leech. 2016. "Local Inhibitory Plasticity Tunes Macroscopic Brain Dynamics and Allows the Emergence of Functional Brain Networks." *NeuroImage* 124 (Pt A): 85–95. https://doi.org/10.1016/j.neuroimage.2015.08.069.

Hensch, T. K. 2004. "Critical Period Regulation." *Annual Review of Neuroscience* 27 (1): 549–79.

https://doi.org/10.1146/annurev.neuro.27.070203.144327.

Hensch, T. K., M. Fagiolini, N. Mataga, M. P. Stryker, S. Baekkeskov, and S. F. Kash. 1998. "Local GABA

Circuit Control of Experience-Dependent Plasticity in Developing Visual Cortex." *Science (New York, N.Y.)* 282

(5393): 1504–8. https://doi.org/10.1126/science.282.5393.1504.

Jansen, B. H., and V. G. Rit. 1995. "Electroencephalogram and Visual Evoked Potential Generation in a

Mathematical Model of Coupled Cortical Columns." *Biological Cybernetics* 73 (4): 357–66.

https://doi.org/10.1007/BF00199471.

Juarez Martinez, Erika L., Hilgo Bruining, Jan Sprengers, Sonja Simpraga, Simon Houtman, Simon-Shlomo Poil,

Huibert D. Mansvelder, et al. 2019. "O-20 Epileptiform and Non-Epileptiform EEG Abnormalities in Autism

Spectrum Disorder Relate to Opposite Excitation-Inhibition Balance Disturbances." *Clinical Neurophysiology*

130 (7): e27–28. https://doi.org/10.1016/j.clinph.2019.04.336.

Kat, Renate, Klaus Linkenkaer-Hansen, Marthe A. Koopmans, Simon J. Houtman, Hilgo Bruining, and Martien J.

H. Kas. 2024. "Assessment of the Excitation-Inhibition Ratio in the Fmr1 Ko2 Mouse Using Neuronal Oscillation

Dynamics." *Cerebral Cortex (New York, N.Y.: 1991)* 34 (5): bhae201. https://doi.org/10.1093/cercor/bhae201.

Lendner, Janna D, Randolph F Helfrich, Bryce A Mander, Luis Romundstad, Jack J Lin, Matthew P Walker, Pal

G Larsson, and Robert T Knight. 2020. "An Electrophysiological Marker of Arousal Level in Humans." *eLife* 9

(July): e55092. https://doi.org/10.7554/eLife.55092.

Linkenkaer-Hansen, K., V. V. Nikouline, J. M. Palva, and R. J. Ilmoniemi. 2001. "Long-Range Temporal

Correlations and Scaling Behavior in Human Brain Oscillations." *The Journal of Neuroscience: The Official*

Journal of the Society for Neuroscience 21 (4): 1370–77. https://doi.org/10.1523/JNEUROSCI.21-04-01370.2001.

Márton, L. F., S. T. Brassai, L. Bakó, and L. Losonczi. 2014. "Detrended Fluctuation Analysis of EEG Signals."

Procedia Technology 12: 125–32. https://doi.org/10.1016/j.protcy.2013.12.465.

Nifterick, Anne M. van, Danique Mulder, Denise J. Duineveld, Marina Diachenko, Philip Scheltens, Cornelis J.

Stam, Ronald E. van Kesteren, Klaus Linkenkaer-Hansen, Arjan Hillebrand, and Alida A. Gouw. 2023. "Resting-

State Oscillations Reveal Disturbed Excitation-Inhibition Ratio in Alzheimer's Disease Patients." *Scientific*

Reports 13 (1): 7419. https://doi.org/10.1038/s41598-023-33973-8.

Poil, Simon-Shlomo, Richard Hardstone, Huibert D. Mansvelder, and Klaus Linkenkaer-Hansen. 2012. "Critical-

State Dynamics of Avalanches and Oscillations Jointly Emerge from Balanced Excitation/Inhibition in Neuronal

Networks." *The Journal of Neuroscience: The Official Journal of the Society for Neuroscience* 32 (29): 9817–23.

https://doi.org/10.1523/JNEUROSCI.5990-11.2012.

Porjesz, Bernice, Laura Almasy, Howard J. Edenberg, Kongming Wang, David B. Chorlian, Tatiana Foroud,

Alison Goate, et al. 2002. "Linkage Disequilibrium Between the Beta Frequency of the Human EEG and a

GABAA Receptor Gene Locus." *Proceedings of the National Academy of Sciences of the United States of*

America 99 (6): 3729–33. https://doi.org/10.1073/pnas.052716399.

Provencher, Stephen W. 1993. "Estimation of Metabolite Concentrations from Localized *in Vivo* Proton NMR

Spectra." *Magnetic Resonance in Medicine* 30 (6): 672–79. https://doi.org/10.1002/mrm.1910300604.

Scheeringa, René, and Pascal Fries. 2019. "Cortical Layers, Rhythms and BOLD Signals." *NeuroImage* 197

(August): 689–98. https://doi.org/10.1016/j.neuroimage.2017.11.002.

Shafiei, Golia, Ben D. Fulcher, Bradley Voytek, Theodore D. Satterthwaite, Sylvain Baillet, and Bratislav Misic.

2023. "Neurophysiological Signatures of Cortical Micro-Architecture." *Nature Communications* 14 (1): 6000.

https://doi.org/10.1038/s41467-023-41689-6.

Shew, Woodrow L., Hongdian Yang, Shan Yu, Rajarshi Roy, and Dietmar Plenz. 2011. "Information Capacity

and Transmission Are Maximized in Balanced Cortical Networks with Neuronal Avalanches." *The Journal of*

Neuroscience: The Official Journal of the Society for Neuroscience 31 (1): 55–63.

https://doi.org/10.1523/JNEUROSCI.4637-10.2011.

Shibata, Kazuhisa, Yuka Sasaki, Ji Won Bang, Edward G Walsh, Maro G Machizawa, Masako Tamaki, Li-Hung

Chang, and Takeo Watanabe. 2017. "Overlearning Hyperstabilizes a Skill by Rapidly Making Neurochemical

Processing Inhibitory-Dominant." *Nature Neuroscience* 20 (3): 470–75. https://doi.org/10.1038/nn.4490.

Takesian, Anne E., and T. K. Hensch. 2013. "Balancing Plasticity/Stability Across Brain Development." In

Progress in Brain Research, 207:3–34. Elsevier. https://doi.org/10.1016/B978-0-444-63327-9.00001-1.

Tamaki, Masako, Zhiyan Wang, Tyler Barnes-Diana, DeeAnn Guo, Aaron V. Berard, Edward Walsh, Takeo

Watanabe, and Yuka Sasaki. 2020. "Complementary Contributions of Non-REM and REM Sleep to Visual

Learning." *Nature Neuroscience* 23 (9): 1150–56. https://doi.org/10.1038/s41593-020-0666-y.

Vogels, T. P., H. Sprekeler, F. Zenke, C. Clopath, and W. Gerstner. 2011. "Inhibitory Plasticity Balances

Excitation and Inhibition in Sensory Pathways and Memory Networks." *Science (New York, N.Y.)* 334 (6062):

1569–73. https://doi.org/10.1126/science.1211095.

Yang, Binghao, Haoran Zhang, Tianzi Jiang, and Shan Yu. 2023. "Natural Brain State Change with E/I Balance

Shifting Toward Inhibition Is Associated with Vigilance Impairment." *iScience* 26 (10): 107963.

https://doi.org/10.1016/j.isci.2023.107963.

Yokoyama, Hiroshi, and Keiichi Kitajo. 2023. "A Data Assimilation Method to Track Excitation-Inhibition

Balance Change Using Scalp EEG." *Communications Engineering* 2 (1): 92. https://doi.org/10.1038/s44172-023-

00143-7.

Associations among EEG measures

In addition to our primary questions regarding links between EEG-E/I measures and MRS-based E/I, we additionally explored two additional questions. First, we assessed the extent to which different EEG-based measures of E/I were themselves collinear; second, we assessed the associations between each EEG-E/I measure was associated with either Glx or GABA+, as opposed to these molecules' ratio in the form of MRS-E/I. To assess these relations we extracted the by-participant random intercepts and the residual correlations from the Bayesian multivariate linear mixed-effects model (see Figure S1 see also Figures 1b, 2b).

Figure S1. Correlations among measures apart from MRS-E/I (i.e., EEG measures as well as separated Glx and GABA+ concentration estimates). Above the diagonal are between-subjects correlations (see Figure 1b); below the diagonal are within-subjects correlations (see Figure 2b). Cells of EEG measures with reliable out-of-sample predictiveness of MRS-E/I are outlined. All estimates are drawn from a single multivariate mixed-effects model.

Across-channel correlations

One way to assess the reliability of EEG-based measures, which were each estimated separately for each channel,

was to examine the correlations between the same measure across the posterior channels we analyzed. Due to our

interest in relatively macro-scale changes in E/I, as well as the basic properties of volume conduction (Burle et al.

2015), we expected that our EEG-based E/I estimates would have highly positive correlations with one another;

this would indicate that the underlying algorithms are computationally stable. For most measures we do observe

the expected large positive correlations between channels, although the NMM-based estimates are much more

weakly inter-correlated (See Figure S2). Broadly, then, we see that measures of amplitude, fE/I, and 1/f slope all

demonstrate expected cross-channel correlations indicating reliability, while the low correlations of NMM

measures may be due to true underlying ability to measure independent measures from nearby electrodes, or may

be due to computational instability of the NMM approach; further investigation of this topic may be of interest for

future research.

Figure S2. Median rank correlations between analyzed posterior channels' estimates of EEG-E/I. Most estimates

were highly correlated across channels, although NMM-based estimates had relatively low cross-channel

reliability.

Ratio of between-subjects to within-subjects variance

As a second exploration into the consistency of within-subjects estimates, which may influence the reliability of

within-subjects E/I measures, we examined the ratio of between-subjects variance to within-subjects variance.

Figure S3. Ratios of between-subjects variances to within-subjects variances. Most measure had less than a 3:1

ratio of these variances, with MRS-based E/I having between-subjects variances approximately 1.45 times as

large as within-subjects variances.