Evidence of off-target probe binding in the 10x Genomics Xenium v1 Human Breast Gene

Expression Panel compromises accuracy of spatial transcriptomic profiling

Caleb Hallinan[1,2], Hyun Joo Ji[1,3], Steven L Salzberg[1,2,3,4], Jean Fan[1,2,3]

[1]Center for Computational Biology, Whiting School of Engineering, Johns Hopkins University,

Baltimore, MD 21211, USA

[2]Department of Biomedical Engineering, Johns Hopkins University, Baltimore, MD 21218, USA

[3]Department of Computer Science, Johns Hopkins University, Baltimore, MD 21218, USA

[4]Department of Biostatistics, Johns Hopkins University, Baltimore, MD

Correspondence should be addressed to:

Jean Fan (jeanfan@jhu.edu)

14   **Abstract**

15

16   The accuracy of spatial gene expression profiles generated by probe-based *in situ* spatially-

17   resolved transcriptomic technologies depends on the specificity with which probes bind to their

18   intended target gene. Off-target binding, defined as a probe binding to something other than the

19   target gene, can distort a gene's true expression profile, making probe specificity essential for

20   reliable transcriptomics. Here, we investigate off-target binding in the 10x Genomics Xenium v1

21   Human Breast Gene Expression Panel. We developed a software tool, Off-target Probe Tracker

22   (OPT), to identify putative off-target binding via alignment of probe sequences and found at least

23   21 out of the 280 genes in the panel impacted by off-target binding to protein-coding genes. To

24   substantiate our predictions, we leveraged a previously published Xenium breast cancer dataset

25   generated using this gene panel and compared results to orthogonal spatial and single-cell

26   transcriptomic profiles from Visium CytAssist and 3' single-cell RNA-seq derived from the same

27   tumor block. Our findings indicate that for some genes, the expression patterns detected by

28   Xenium demonstrably reflect the aggregate expression of the target and predicted off-target

29   genes based on Visium and single-cell RNA-seq rather than the target gene alone. Overall, this

30   work enhances the biological interpretability of spatial transcriptomics data and improves

31   reproducibility in spatial transcriptomics research.

Introduction

Recent advances in high-throughput spatially-resolved transcriptomic profiling technologies have

enabled the investigation of gene expression with high spatial resolution within tissues. One such

commercially available spatial transcriptomics platform is Xenium from 10x Genomics, a publicly

traded company with a market capitalization exceeding $1 billion as of March 2025 (Yahoo

Finance 2025). Xenium achieves spatial gene expression profiling at single-cell resolution for

targeted genes using a probe-based *in situ* detection approach. 10x Genomics currently offers

targeted gene panels with pre-designed probe sets, including their 280-gene Xenium v1 Human

Breast Gene Expression Panel. As of December 2024, over 16,000 Xenium consumable reactions

have been sold, with each tissue slide profiled costing approximately $5,000, underscoring the

platform's widespread use and high commercial value (10x Genomics).

Briefly, Xenium uses padlock probes that include sequences complementary to the RNA of target

genes. Once a padlock probe binds to its target, it is ligated and subsequently amplified via rolling

circle amplification (RCA). Fluorescently labeled decoder probes then hybridize to the amplified

RCA product, enabling the simultaneous detection and decoding of the optical signature, or

codeword, specific to each gene in the panel through successive rounds of fluorescence imaging.

When combined with cell-segmentation, this approach allows for spatially-resolved single-cell

quantification of gene expression.

The accuracy of these gene expression measurements thus relies on the specificity of the probes

to bind to their intended target gene. We define off-target binding as when a probe binds to

something other than the RNA sequence intended to correspond to the target gene (Figure 1).

We note once ligation and RCA occur, the resulting fluorescent signal cannot be distinguished

between on-target and off-target binding. As such, off-target binding can distort the

quantification of the intended target gene's expression, as the observed expression would

represent a combination of the target as well as off-target expression.

Figure 1. Schematic of potential off-target binding in 10x Xenium. In this illustration, the arms

of the padlock probes were designed to bind an RNA sequence intended to correspond to a target

gene (green). However, these probes exhibit off-target binding and bind to an RNA sequence in

a different off-target gene (red). The probe is circularized and subsequently amplified via rolling

circle amplification (RCA). Hybridization of fluorescent probes to the RCA product enables the

generation a fluorescent signal that is used to quantify RNA expression within cells.

To predict for such potential off-target binding, we developed Off-target Probe Tracker (OPT), a

software tool that aligns probe sequences to an annotated transcriptome with the option to

allow for mismatches that may still permit probe binding. Using OPT, we identify putative off-

target probe binding to protein-coding genes affecting at least 21 out of 280 genes in the 10x

Genomics Xenium v1 Human Breast Gene Expression Panel, compromising the accuracy of their

spatial transcriptomic profiles. We substantiate our predictions using data from orthogonal

spatial and single-cell gene expression profiling technologies. By facilitating a more rigorous

evaluation of probe sequence specificity, tools like OPT can aid in future probe design decisions

to help ensure that probes are optimized to minimize off-target binding based on current

transcriptome annotations.

Results

<u>OPT predicts potential off-target probe binding</u>

To identify potential off-target binding impacting the 10x Genomics Xenium v1 Human Breast

Gene Expression Panel, we first downloaded the FASTA file containing the probe sequences from

the 10x Genomics Website (Methods, Supplementary Data). This file includes 4,809 probe

sequences that are 40bp in length and designed to target 280 genes, averaging 17 probe

sequences per gene (ranging from 2 to 50 probe sequences per gene).

We developed a software tool called OPT (Methods) that uses nucmer (Marçais et al. 2018) to

predict off-target binding by aligning probe sequences to a reference transcriptome, which

comprises a curated collection of transcript isoforms for all genes in a species. OPT features

adjustable parameters for binding strictness (e.g., number of mismatches) and generates a

summary file that details all targeted genes along with their potential off-targets based on the

sequence alignments. 10x Genomics designed its probe sequences using the GENCODE "basic"

annotation (Mudge et al. 2025), so initially we also used the latest GENCODE "basic" annotation

(v47) to predict off-target binding for these probes.

We first sought to predict if a probe has off-target binding based on perfect sequence homology

(i.e., if it aligns with 100% identity) with any annotated transcripts other than those that belong

to the intended target gene. Of the 4,809 probe sequences in this gene panel, using GENCODE

v47, OPT identified 180 probe sequences across 45 genes as having off-target binding based on

perfect sequence homology (Table 1). Among the 45 genes with predicted off-target binding, the

number of affected probe sequences per gene ranged from 1 to 14. Overall, these off-target

probes matched 90 other genes, including 34 protein-coding genes, 34 pseudogenes, 12 long

non-coding RNA, 9 transcripts labeled as nonsense-mediated decay, and one microRNA gene.

Target gene	Number of probes	Predicted binding genes	Number of probes aligned	Gene types - GENCODE (v47)
ADGRE5	19	ADGRE5,ADGRE2	19,1	PC,PC
ADH1B	23	ADH1B,ADH1A,ADH1C	22,3,3	PC,PC,PC
AKR1C1	21	AKR1C1,AKR1C5P,AKR1C2,AKR1C3, AKR1C4	21,1,1,1,1	PC,PC,PC,PC,PG
AKR1C3	17	AKR1C3,AKR1C1,AKR1C2,AKR1C4	17,1,1,2	PC,PC,PC,PC
APOBEC3A	10	APOBEC3A,APOBEC3B	8,4	PC,PC
APOBEC3B	14	APOBEC3B,APOBEC3D,ENSG00000284554, APOBEC3F,APOBEC3A	14,2,2,2,2	PC,PC,PC,PC,PC
AQP1	19	AQP1,ENSG00000250424	19,4	PC,PC
C15orf48	6	C15orf48,MIR147B	6,1	PC,miRNA
C1QA	4	C1QA,ENSG00000289692	4,2	PC,PC
CD68	7	CD68,ENSG00000264772	7,6	PC,lncRNA
CD79B	5	CD79B,ENSG00000285947	5,3	PC,PC
CEACAM6	13	CEACAM6,ENSG00000267881,CEACAM3	13,1,1	PC,PC,PC
CLECL1; CLECL1P	7	ENSG00000293488,CLECL1P	3,3	lncRNA,PG
DPT	18	DPT,LINC00970	18,13	PC,lncRNA
EPCAM	15	EPCAM,ENSG00000225356	15,1	PC,PG
FCGR3A	20	FCGR3A,FCGR3B,ENSG00000289768	20,3,3	PC,PC,PC
HMGA1	7	HMGA1,HMGA1P3,HMGA1P1,HMGA1P2	7,1,1,1	PC,PG,PG,PG
IL2RG	14	IL2RG,ENSG00000285171	14,11	PC,NMD
KLRC1	15	KLRC1,KLRC2,KLRC4-KLRK1,KLRC3, KLRC4,ENSG00000255641	15,3,1,1,1, 1	PC,PC,PC,PC,NMD,PC
KRT14	6	KRT14,KRT16P6,ENSG00000290977	6,1,1	PC,lncRNA,PG
KRT6B	15	KRT6B,KRT6C,KRT75,KRT6A	15,4,1,1	PC,PC,PC,PC
KRT8	18	KRT8,KRT8P11,KRT8P2,KRT8P3,KRT8P36, KRT8P45,KRT8P33,KRT8P37, ENSG00000304440,CDK5R2-AS1,KRT8P30, KRT8P17, KRT8P22,KRT8P32,KRT8P42	18,1,2,3,1, 2, 2,1,1,1,1, 1,1,1	PC,lncRNA,lncRNA,PG,PG,PG, PG,PG,PG,PG,PG,PG,PG,PG,PG
LDHB	18	LDHB,ENSG00000285854,LDHBP2	18,12,1	PC,NMD,PG
LILRA4	16	LILRA4,ENSG00000275210	16,3	PC,lncRNA
MYLK	35	MYLK,MYLKP1	35,1	PC,PG
MYO5B	20	MYO5B,MYO5BP1,MYO5BP2, ENSG00000266997	20,2,2,5	PC,PG,PG,NMD
PCLAF	9	PCLAF,ENSG00000259316	9,1	PC,NMD
PDGFRA	29	PDGFRA,ENSG00000282278	29,9	PC,PC
PTGDS	5	PTGDS,ENSG00000284341	5,3	PC,NMD
SCD	20	SCD,SCDP1	20,3	PC,PG
SERHL2	8	SERHL2,SERHL	8,7	PC,PG
SERPINA3	27	SERPINA3,ENSG00000273259	27,14	PC,NMD
SLAMF1	24	SLAMF1,ENSG00000228863	23,1	PC,lncRNA
SMS	17	SMS,ENSG00000213080,ENSG00000249779, ENSG00000232389,ENSG00-000251342	17,7,1,3,1	PC,PG,PG,PG,PG
SVIL	20	SVIL,SVIL2P	20,2	PC,PG
THAP2	25	THAP2,ENSG00000258064	25,2	PC,NMD
TPD52	21	TPD52,ENSG00000276418	21,5	PC,NMD
TPSAB1	2	TPSAB1,TPSB2,TPSD1	2,2,1	PC,PC,PC
TRAF4	14	TRAF4,ENSG00000225869	14,1	PC,PG,PG,NMD
TUBA4A	12	TUBA4A,TUBA4B	12,1	PC,PC
TUBB2B	11	TUBB2B,TUBB2BP1,TUBB2A,TUBB4A,TUBB3, ENSG00000198211	11,2,3,1,1, 1	PC,PC,PC,PC,PG,PC
VOPP1	24	VOPP1,ENSG00000223612	24,1	PC,PG
VWF	20	VWF,VWFP1,LINC01665	20,2,1	PC,PG,lncRNA
ZEB2	47	ZEB2,ZEB2P1	47,1	PC,PG
ZNF562	22	ZNF562,ENSG00000291231,ZNF812P	22,3,3	PC,lncRNA,PG

112 **Table 1. OPT output of genes with predicted off-target binding based on perfect sequence**

113 **homology in GENCODE v47.** This table shows the 45 genes whose probes in the 10x Genomics

114 Xenium v1 Human Breast Gene Expression Panel exhibit predicted off-target probe binding,

115 where each off-target alignment involves a perfect 40bp match to the probe sequence. Although

116 OPT predicted off-target binding of CCPG1 probe sequences to the DNAAF1-CCPG1 gene, we

117 manually excluded it from our list because DNAAF1-CCPG1 is a read-through gene containing

118 portions of both DNAAF1 and CCPG1. The final column shows the gene types, in order, of each

119 of the off-target genes shown in column 3. Abbreviations: PC = protein-coding; PG = pseudogene;

120 NMD = nonsense-mediated decay; lncRNA = long non-coding RNA.

121

122

123 <u>Variation in off-target binding predictions across different annotations</u>

124

125 OPT relies on alignments to an annotated transcriptome, which ideally reflects all genes and gene

126 variants stably transcribed in a given species. However, genome annotation is still an active area

127 of research (Varabyou et al. 2023), with discrepancies across annotations in gene counts,

128 isoforms, and many other features. We therefore further used OPT to predict for off-target

129 binding and affected genes using two additional human genome annotation sets, RefSeq (v110),

130 (O'Leary et al. 2016) and CHESS (v3.1.3) (Varabyou et al. 2023), and compared to our previous

131 results from GENCODE (v47) (Mudge et al. 2025) (Methods).

132

When considering only perfect sequence homology, while we previously found 45 affected genes

using GENCODE, we found 22 when using RefSeq and 33 when using CHESS (Supplementary Table

1, 2). Given that RefSeq and CHESS have more transcripts than GENCODE, these discrepancies in

off-target binding predictions was not simply an artifact of the difference in transcript set sizes.

While the human annotation databases mostly agree on the number of protein-coding genes in

the genome, they remain widely divergent on pseudogenes and lncRNA genes. Therefore, we

focused on how the results change when we restrict our analysis to only protein-coding genes.

By excluding pseudogenes (which are presumably not expressed), lncRNAs, and other non-

protein-coding RNAs when using OPT, the number of affected genes fell to 17 for GENCODE and

RefSeq and 16 for CHESS (Supplementary Table 3). Again, these discrepancies reflect annotation

differences. For example, the probe sequence (ENSG00000196154|S100A4|ab4e3dc), which was

designed to target *S100A4* based on GENCODE annotations, also aligns to *S100A5* in RefSeq

(Supplementary Figure 1A). We reason that if a probe sequence aligns off-target to a protein-

coding gene based on any of these annotations, it could result in off-target binding. We therefore

focused further analysis on the union of genes with predicted off-target binding to protein-coding

genes across the 3 annotations, resulting in 21 genes: *ADGRE5*, *ADH1B*, *AKR1C1*, *AKR1C3*,

APOBEC3A, *APOBEC3B*, *AQP1*, *C1QA*, *CD79B*, *CEACAM6*, *FCGR3A*, *FLNB*, *KLRC1*, *KRT6B*, *PDGFRA*,

S100A4, TOMM7, TPD52, TPSAB1, TUBA4A, TUBB2B (Supplementary Table 4).

<u>Comparison with Visium CytAssist reveals spatial gene expression patterns consistent with off-</u>

<u>target binding</u>

To investigate the potential effects of our predicted off-target binding for the Xenium v1 Human

Breast Gene Expression Panel in experimental settings, we compared spatial gene expression

patterns detected in two previously published spatial transcriptomics datasets from serial

sections of the same breast cancer tissue: one section assayed with Xenium using the Xenium v1

Human Breast Gene Expression Panel, and another assayed using Visium CytAssist, an orthogonal

spatial transcriptomics platform (Janesick et al. 2023). Briefly, Visium is a sequencing-based

spatial transcriptomics technology in which RNA is hybridized to spatially barcoded capture spots

on a slide, enabling spatial transcriptomic mapping after sequencing. However, while Xenium

offers single-cell resolution gene expression quantification, Visium quantifies gene expression

within 55μm spots. To enable direct comparison, we first structurally aligned the Xenium and

Visium tissue sections using STalign (Clifton et al. 2023), restricting our analysis to overlapping

regions since different parts of the tissue were profiled (Supplementary Figure 2A). To enhance

visual comparison, we aggregate the Xenium gene expression data at these aligned locations to

match the Visium resolution (Supplementary Figure 2B) (Methods).

Among the 21 genes exhibiting off-target binding to protein-coding genes, 12 were present in

the Visium dataset that had at least one corresponding off-target gene also detected in the

dataset. For genes with no predicted off-target binding based on perfect sequence homology

such as *MS4A1*, we observed a visually similar spatial pattern between the two technologies

(Figure 2A), suggesting that spatially aligned groups of cells across the two technologies express

this gene at comparable relative magnitudes. However, for genes with predicted off-target

binding based on perfect sequence homology such as *TUBB2B*, we observed a visually dissimilar

spatial pattern between the two technologies (Figure 2B). Importantly, its predicted off-target

gene, *TUBB2A*, in Visium shows a visually more similar spatial pattern to the Xenium *TUBB2B*. To

better visualize the effect of off-target binding within the Xenium data, we aggregated the

expression of each gene along with its predicted off-targets found in the Visium dataset and

visually compared across spatial locations. Notably, the spatial pattern of the aggregated

expression of *TUBB2B* and *TUBB2A* in Visium is visually more similar to the spatial pattern of

TUBB2B in Xenium, consistent with the prediction that Xenium *TUBB2B* probes exhibit off-target

binding to *TUBB2A*. We further visually confirmed using the Integrative Genomics Viewer (IGV)

that a probe sequence intending to bind to *TUBB2B* also perfectly aligns to a sequence in *TUBB2A*,

consistent across all annotations evaluated (Supplementary Figure 1B).

Overall, when comparing the total gene expression between the two technologies, we observed

a generally strong positive correlation, consistent with the previously published work (Janesick

et al. 2023). We do not observe an obvious trend between gene expression magnitude and the

presence of predicted off-target probes (Figure 2C), suggesting that off-target binding prediction

alone does not explain the observed higher expression magnitude in Xenium compared to

Visium, which may still be attributed to variation in detection efficiency, sequencing depth, and

other factors.

Figure 2. Comparison of spatial gene expression patterns between Xenium and Visium. (A)

Spatial gene expression of *MS4A1* overlaid on the corresponding histological images for Xenium

and Visium. **(B)** Gene expression patterns for *TUBB2B*: Xenium expression, Visium expression, the

aggregated Visium expression combining *TUBB2B* and its predicted off-target gene's expression

TUBB2A, and Visium expression of *TUBB2B*'s predicted off-target *TUBB2A*. **(C)** Scatterplot of log-

transformed total expression counts (with a pseudocount) for 307 genes comparing Visium and

Xenium data. The dotted line represents X = Y, and points (genes) are colored by probe

information.

<u>Comparison with scRNA-seq reveals single-cell gene expression patterns consistent with off-</u>

<u>target binding</u>

To further investigate the potential effects of our predicted off-target binding for Xenium probes,

we compared the detected single-cell gene expression patterns in the same previously published

work: again, the one assayed with Xenium using the Xenium v1 Human Breast Gene Expression

Panel, and another assayed using Chromium Next GEM Single Cell 3' (Janesick et al. 2023). Briefly,

single-cell RNA sequencing (scRNA-seq) with 3' end capture is a technique used to profile gene

expression at the single-cell level by profiling the 3' ends of mRNA transcripts with sequencing

followed by alignment to a genome or transcriptome for quantification. While this approach

provides single-cell resolution gene expression quantification, it lacks spatial information. To

enable a single-cell comparison with Xenium, we use Harmony (Korsunsky et al. 2019) to remove

batch effects and project cells into a shared UMAP embedding (Supplementary Figure 3)

(Methods).

Among the 21 genes exhibiting off-target binding to protein-coding genes, 16 were present in

the sc-RNAseq dataset that had at least one corresponding off-target gene also detected in the

dataset. Again, for genes with no predicted off-target binding based on perfect sequence

homology such as *MS4A1*, we observed a visually similar gene expression pattern in the

harmonized UMAP across both technologies (Figure 3A), suggesting that transcriptionally similar

clusters of cells or cell-types across the two technologies express this gene at comparable relative

magnitudes. Likewise, again, for genes with predicted off-target binding based on perfect

sequence homology such as *TUBB2B*, we observed a visually dissimilar gene expression pattern

on the harmonized UMAP (Figure 3B). Again, its predicted off-target gene, *TUBB2A,* in scRNA-seq

showed a visually more similar expression pattern in the harmonized UMAP embedding to the

Xenium *TUBB2B*. To better illustrate the impact of the off-target probes, we again aggregated

the expression of a gene and its predicted off-target genes present in the scRNA-seq data and

visually compared across the harmonized UMAP embedding. The aggregated expression of

TUBB2B and *TUBB2A* in the scRNA-seq data shows a visually more similar gene expression

pattern in the harmonized UMAP embedding to *TUBB2B* in the Xenium, consistent with the

prediction that Xenium *TUBB2B* probes exhibit off-target binding with these paralogs (Figure 3B,

Supplementary Figure 4).

Overall, when comparing total gene expression between the two technologies, we again

observed a generally strong positive correlation (Figure 3C), similar to the Visium comparison

results and consistent with the previously published work (Janesick et al. 2023).

Figure 3. Comparison of single-cell gene expression patterns between Xenium and scRNA-seq.

(A) Harmonized UMAP visualization of *MS4A1* expression for Xenium and scRNA-seq data. **(B)**

Comparison of *TUBB2B* expression patterns on harmonized UMAP: Xenium expression, scRNA-

seq expression, an aggregated scRNA-seq profile combining *TUBB2B* and its predicted off-target

gene's expression *TUBB2A*, and scRNA-seq expression of *TUBB2B*'s potential off-target *TUBB2A*.

Note that *TUBB2B* also has two additional off-targets present in the scRNA-seq data

(Supplementary Figure 4). **(C)** Scatterplot of log-transformed total expression counts (with a

pseudocount) for 313 genes between Visium and scRNA-seq data. The dotted line represents X =

Y, and points (genes) are colored by probe information.

<u>OPT results when allowing mismatches at the terminal ends of the probe sequences identifies</u>

<u>additional off-target candidates</u>

Thus far, we have focused on predicting off-target binding based on perfect sequence homology.

However, we reason that imperfect sequence matching could still result in off-target binding.

Specifically for the Xenium v1 Human Breast Gene Expression Panel, padlock probes with two 20

base pair (bp) arms bind complementary mRNA regions, forming a 40bp probe sequence. A ligase

then circularizes the padlock probe, favoring specific 2-bp junctions. Importantly, if there is a

sequence mismatch, particularly outside the ligation site towards the terminal ends of the probe

sequence, hybridization may still occur and result in off-target binding, albeit with reduced

hybridization efficiency (Supplementary Figure 5A). We therefore added an option in OPT to

allow imperfect alignments at the ends of the probe sequences, specifying the sequence length

at either ends, where mismatches, insertions, deletions, or clipping can occur (Methods).

Allowing for 6bp mismatches on either end of the 40bp probe sequence (i.e., requiring a 28bp

match covering the middle of the probe sequence including the ligation site) revealed nine

additional genes with potential off-target binding when using GENCODE v47 (Supplementary

Table 5). Additionally, 18 of the 45 genes previously predicted to be affected by off-target binding

based on perfect sequence homology were now predicted to have additional off-target binding,

either through the accumulation of new probes with predicted off-target binding or from the

identification of additional predicted off-target genes (Supplementary Table 6). Among these

additional genes included *CEACAM8*, for which we observed a visually dissimilar spatial pattern

between Xenium and Visium (Supplementary Figure 6A). Furthermore, the spatial pattern of the

aggregate of *CEACAM8* with its predicted off-target genes (*CEACAM5*, *CEACAM6*, *CEACAM7*,

PSG6) in Visium more closely resembled the spatial pattern of *CEACAM8* in Xenium. Likewise, a

similar trend is observed in the scRNA-seq comparison (Supplementary Figure 6B). Ultimately,

these findings suggest that off-target binding, even with imperfect sequence matching, can

contribute to the expression patterns observed in Xenium.

Discussion

Our study presents evidence of off-target probe binding that may distort gene expression profiles

affecting at least 21 out of the 280 genes in the Xenium v1 Human Breast Gene Expression Panel.

This finding is supported by spatial and single-cell comparative analyses using Xenium with serial

section datasets from Visium CytAssist and 3' single-cell RNA-seq respectively. Although we focus

here on the Xenium v1 Human Breast Gene Expression Panel from 10x Genomics, we do not

exclude the possibility that off-target binding may similarly affect other probe-based gene

detection approaches. To assist in the interpretation of existing probe-based gene expression

data as well as future probe design, we provide OPT as a software tool for predicting potential

off-target probe binding.

Although we have predicted off-target binding based on sequence alignment, its effect on gene

expression quantification may still vary. One reason is that the off-target protein-coding or non-

protein-coding gene may not be expressed. For example, although *CEACAM6* probes have

predicted off-target binding to *CEACAM3* based on perfect sequence homology, the sparse

expression of *CEACAM3* in both the Visium and scRNA-seq breast cancer data led to only a minor

difference in the aggregated expression (Supplementary Figure 7). As such, the extent of the

effect of off-target binding depends on the expression of the off-target gene and will therefore

vary across tissues. On the other hand, self-hybridization or interactions between probes

themselves may occur to induce non-specific signals (Supplementary Fig 5B). In general, probe

binding specificity is influenced by numerous factors, with many methods previously developed

to aid in the design of probe sequences while taking these factors into consideration (Wang and

Seed 2003; Rouillard, Zuker, and Gulari 2003; Wernersson and Nielsen 2005; Chou 2010; Li et al.

2011; Hu et al. 2020; Hershberg et al. 2021; Fornace et al. 2022).

Notably, our evaluation was limited to the base 280-gene Xenium v1 Human Breast Gene

Expression Panel even though the expression of 313 total genes were quantified in these datasets

from Janesick et al. This is because 33 of the 313 targeted genes were probed using custom

sequences that were not publicly disclosed. Our comparative analysis suggests that *HDC*, a gene

with custom probe sequences mentioned in the paper, may also be subject to off-target binding.

In Xenium, *HDC* exhibited a distinct spatial pattern and high global expression level, whereas both

Visium and scRNA-seq data showed a minimal spatial pattern and sparse expression level

respectively (Supplementary Figure 8). We therefore speculate that *HDC* probes may exhibit off-

target binding but cannot fully validate without the probe sequences used for *HDC*. As such, we

emphasize the importance of publishing full probe sequences to enable reproducible science.

We also observed that 34 / 4,809 breast panel probe sequences did not align to any reference

transcripts in the GENCODE v47 annotation. 33 of these unmapped probe sequences matched

intronic sequences, which would not normally appear in mature transcripts. We manually aligned

the one remaining probe sequence (ENSG00000198851|CD3E|02369ef) to the GRCh38 genome

and found that it matched to chr11: 118304577 – 118304616, an intergenic region just upstream

of *CD3E*, the target gene for this probe. We suspect that these intron and intergenic panel probe

sequences were designed using an earlier version of the GENCODE annotation in which those

regions were annotated as exonic. This finding illustrates the importance of disclosing the specific

annotation version to promote reproducibility, as well as the ongoing variability of human gene

annotation. Likewise, as evidenced by our analysis across GENCODE, RefSeq, and CHESS, we

emphasize the variation across these reference annotations and therefore recommend

evaluating probe sequences for off-target effects using multiple annotations to be more

comprehensive.

Given these challenges, we advise probes with predicted off-target binding to protein-coding

genes based on perfect sequence homology be discarded in future experiments. Likewise, we

encourage the use of tools like OPT to aid in future probe design decisions and help ensure that

probes are optimized to minimize off-target binding based on current transcriptome annotations.

For datasets that have already been generated using probes with predicted off-target binding,

we generally recommend taking into consideration these predictions to avoid drawing misleading

conclusions. For example, we recommend expression measurements for genes with predicted

off-target binding be omitted from training foundation models to avoid error propagation.

Alternatively, when performing integrative analyses that compare or align gene expression with

measurements across technologies, it may be necessary to incorporate off-target binding

predictions. For example, integration could be performed between the observed Xenium gene

expression and the aggregated expression of the target and predicted off-target genes for the

orthogonal technology. Finally, existing literature that base conclusions on genes with predicted

off-target binding should be interpreted with caution.

We emphasize that these findings were missed in the previous Janesick et al. publication from

10x Genomics (Janesick et al. 2023). Consistent with previously published observations, we

observed a highly correlated total gene expression magnitude between Xenium and Visium as

well as scRNA-seq. However, a notable exception is *APOBEC3B*, which is not expressed according

to both Visium and scRNA-seq but highly expressed according to Xenium (Supplementary Figure

9) – a discrepancy that Janesick et al. omitted. We emphasize that positive significant average

gene expression correlation is a necessary but not sufficient metric for consistency across

technologies and that individual data points should be scrutinized. Likewise, validation with

orthogonal technologies could have helped identify discrepancies suggestive of off-target effects.

We note Janesick et al. only use immunofluorescence to validate two genes, *ERBB2* and *MS4A1*,

which by our analysis were predicted to exhibit no off-target binding. We therefore stress the

importance of more thorough orthogonal validations. Although Xenium incorporates blank and

negative control probes that are intended to help quantify the rate of non-specific and potential

off-target binding, our findings suggest that relying solely on such probes for error detection may

be insufficient. Implementing probe redundancy, where the same gene is targeted using different

codewords, could provide an additional internal control to enable the detection of off-target

binding.

This is not the first instance in which a commercially available platform has encountered

challenges in probe design (McCartney et al. 2016; Harbig 2005; Mecham et al. 2004; Liu, Bebu,

and Li 2014). These findings underscore the critical role of academic researchers towards

ensuring the robustness of industry-led product development by providing oversight, free of

financial conflicts of interest through independent federal funding. This complementarity

between industry and academia fosters a more rigorous, transparent, and reliable scientific

process, ultimately to the benefit of consumers and the public. By shedding light on putative off-

target probe binding as well as by providing a tool to enable such off-target binding predictions,

this work will help enhance the quality of spatial transcriptomics data and improve the overall

reproducibility in spatial transcriptomics research.

Methods

<u>Off-target Probe Tracker Tool (OPT)</u>

OPT (<u>O</u>ff-target <u>P</u>robe <u>T</u>racker) is a Python program that runs nucmer (Marçais and Kingsford

2011) for alignment and then processes the results to predict probe binding based on sequence

homology. OPT is available as an open-source Python toolkit at <u>https://github.com/JEFworks-</u>

<u>Lab/off-target-probe-tracker</u>. When a user provides a query probe sequence file, a target

transcript sequence file, and the annotation used to extract these transcripts, OPT outputs which

gene each probe is likely to bind to. Nucmer is a fast nucleotide sequence aligner that uses

maximal exact matches as anchors which it then joins together to find longer alignments. By

default, OPT saves nucmer results in SAM format and finds perfect sequence matches between

a query probe and a target transcript, requiring that alignments consist of only matches and cover

the entirety of the query. OPT consists of three modules: (1) `flip` for reverse complementing

probe sequences aligned to the opposite strand of their target genes; (2) `track` for aligning

probe sequences and processing alignment results; and (3) `stat` for compiling summary

statistics on the number of off-target binding probes and affected genes.

In the case that a probe's target gene has synonyms, we consider alignments to genes annotated

with one of its synonym to still be on-target. For example, if a probe that targets *NARS* shows

alignments to a gene called *NARS1*, we don't consider it to be off-target binding. We gathered

relevant gene synonym relationships using the GeneCards and HGNC online database.

OPT also provides a "pad" mode in which imperfect alignments are allowed at either end of the

query (i.e., probe sequence). The -pl parameter sets the pad length at either end of the query

and OPT allows for any number of mismatches in these padded regions. For example, if the pad

length is 6 and the probe sequence length is 40, then the middle 28bp are the only part of the

probe sequence required to match. As long as the critical region is intact, OPT reports an off-

target binding site based on this alignment. By default, -pl is set to 0, and the pad mode is

activated by providing a non-zero integer to -pl.

<u>Obtaining probe sequences for the Xenium v1 Human Breast Gene Expression Panel</u>

To identify potential off-target binding impacting the 10x Genomics Xenium v1 Human Breast

Gene Expression Panel, we obtained the FASTA file of probe sequences from the 10x Genomics

Website (https://cdn.10xgenomics.com/raw/upload/v1684948097/software-support/Xenium-

panels/hBreast_panel_files/xenium_human_breast_gene_expression_panel_probe_sequences.

fasta) and available as Supplementary Data for preservation.

The target gene name and ID was extracted from the probe IDs of the following format:

```
    > gene_id|gene_name|accession
    ```

426   We expected the provided FASTA file to contain probe sequences to be the reverse

427   complemented sequence of their intended target genes and hence align to the reverse strand of

428   their target isoforms. However, when we aligned the breast panel probe sequences to the

429   GENCODE basic (v47) reference transcripts using nucmer, we found that only 2,508 / 4,809

430   (52.2%) of probe sequences aligned on the reverse strand of their target transcripts (i.e., isoforms

431   of their target genes). For consistency, we enforced that all probe sequences be oriented in the

432   same direction and align to the forward strand of their target genes and transcripts. As such, we

433   reverse-complemented these 2,508 probe sequences. We also added this functionality as an OPT

434   module called "flip" in which probe sequences aligned to the reverse strand of their targets are

435   reverse complemented. We expect probe sequences to align to the forward strand of transcripts

436   (i.e., both oriented in the same direction) during the downstream probe sequence binding

437   prediction step.

439   Visium Comparison

441   The Visium CytAssist dataset, collected from a breast cancer tissue block utilized in Janesick et

442   al.,   was   downloaded   from   the   10x   Genomics   website

443   (https://www.10xgenomics.com/products/xenium-in-situ/preview-dataset-human-breast). This

444 dataset originally contained 4,992 spots with x-y coordinates and includes 18,085 genes per spot.

445 Six genes (*AKR1C1*, *ANGPT2*, *BTNL9*, *CD8B*, *POLR2J3*, *TPSAB1*) were excluded from the analysis

446 since they were not present in the Visium dataset.

448 To compare spatial gene expression patterns from Visium and Xenium technologies, we first

449 mapped all the data to the same coordinate space. We used STalign (v1.0.1), a computational

450 tool that utilizes affine transformations along with diffeomorphic metric mapping to align target

451 and source datasets (Clifton et al. 2023). The initial alignment involved only affine

452 transformations and eight manually determined landmarks to align the Visium histology image

453 (source) to the Xenium histology image (target). This transformation brought the Visium image

454 into the coordinate space of the higher-resolution Xenium image. We then applied this learned

455 transformation to the Visium spots, ensuring that they were correctly positioned relative to both

456 histology images. Next, we used STalign to map the Xenium transcripts (source) onto their

457 corresponding Xenium histology image (target) using both affine and diffeomorphic metric

458 mapping. The transcripts were rasterized at 30μm resolution, with an initial affine transformation

459 guided by four manually defined landmarks. Diffeomorphic metric mapping was then performed

460 with the following parameters: a = 2500, epV = 1, niter = 2000, sigmaA = 0.11, sigmaB = 0.10,

461 sigmaM = 0.15, sigmaP = 50, muA = [1, 1, 1], muB = [0, 0, 0], with all other settings left at their

462 defaults. We extracted the overlapping regions between the two datasets (Supplementary Figure

463 2A), which reduced the total spots in the Visium dataset to 3,958. Finally, we aggregated the

464 Xenium gene expression data to ~55μm x 55μm patches that correspond to the spatial locations

465   of the Visium spots, resulting in matched-resolution spatial gene expression for both

466   technologies (Supplementary Figure 2B).

469   <u>Single-cell RNA-seq Comparison</u>

471   The Chromium Next GEM 3' scRNA-seq dataset, collected from a breast cancer tissue block

472   utilized   in   Janesick   et   al.,   was   downloaded   from   the   10x   Genomics   website

473   (https://www.10xgenomics.com/products/xenium-in-situ/preview-dataset-human-breast).   This

474   dataset contained 12,388 cells with 36,601 genes per cell. All 313 unique genes present in the

475   Xenium dataset are also in the scRNA-seq data, hence both datasets were subsetted to these

476   genes for this analysis.

478   Both scRNA-seq and Xenium provide single-cell resolution data. To integrate these datasets, we

479   first removed cells lacking detectable gene expression. We the normalized the combined gene

480   expression data using counts per million (CPM) and applied a log transformation with a

481   pseudocount of 1. Principal component analysis (PCA) is then applied to the normalized data, and

482   batch effects are corrected using Harmony (v1.2.3) on the top 30 principal components (PCs)

483   using default parameters except for theta, which was set to 8, to promote further mixing with

484   clusters across technologies. Finally, Uniform Manifold Approximation and Projection (UMAP) is

485   performed   on   the   harmonized   PCs,   generating   a   shared   2D   embedding   across   the   two

486   technologies, and the data is further facetted by technology for visualization (Supplementary

487   Figure 3).

490   Cross-annotation analysis

492   To compare OPT's results with different reference annotations, we used the most recent releases

493   of GENCODE basic (v47), GENCODE comprehensive (v47), RefSeq (v110), and CHESS (v3.1.3)

494   annotation of the GRCh38 genome. Note that GENCODE "basic" is the more reliable version of

495   the annotation and is much closer to RefSeq and CHESS. GENCODE "comprehensive" includes

496   hundreds of thousands of low-quality annotations, which we included in some of our analyses

497   for completeness. Note also that GRCh38 has many non-reference sequences called "alternative

498   scaffolds"; we removed these for our analysis. We then used gffread to extract transcripts as

499   defined in these annotations by running:

501   ```
$ gffread -w transcripts.fa -g grch38.p12/14.fa annotation.gff
```

The GRCh38.p14 assembly was used during transcript sequence extraction for all reference

annotations, except for CHESS which specifies that the annotation maps genes and transcripts

onto the GRCh38.p12 assembly. For RefSeq, we renamed the VD(J) segment features as

transcript features to ensure consistency, and we also removed transcript sequences with the

gene_biotype "pseudogene." RefSeq has a separate biotype called "transcribed_pseudogene,"

but doesn't annotate transcripts for these features. We considered transcripts annotated for a

small subset of just pseudogenes an error in the annotation.

**Conflicting interests**

The authors declare no conflicting interests.

**Acknowledgements**

We thank Reza Kalhor for his input on the project and feedback on the manuscript. Research

reported in this publication was supported by the National Institute of General Medical Sciences

of the National Institutes of Health under Awards R35-GM142889 and R35-GM130151, and the

HuBMAP Integration, Visualization, and Engagement (HIVE) Initiative under Award Number OT2-

OD033760.

## References

10x Genomics. 2025a. "10x Genomics Product Catalog." March 19, 2025.
www.10xgenomics.com/store/product-catalog?query=xenium.

———. 2025b. "TXG_Q4-and-Full-Year-2024-Earnings-Press-Release." 10x Genomics Reports
Fourth Quarter and Full Year 2024 Financial Results and Provides Outlook for 2025.
March 19, 2025.
https://s28.q4cdn.com/592666581/files/doc_financials/2024/q4/TXG_Q4-and-Full-
Year-2024-Earnings-Press-Release.pdf.

Chou, Hui-Hsien. 2010. "Shared Probe Design and Existing Microarray Reanalysis Using PICKY."
*BMC Bioinformatics* 11 (1): 196. https://doi.org/10.1186/1471-2105-11-196.

Clifton, Kalen, Manjari Anant, Gohta Aihara, Lyla Atta, Osagie K. Aimiuwu, Justus M. Kebschull,
Michael I. Miller, Daniel Tward, and Jean Fan. 2023. "STalign: Alignment of Spatial
Transcriptomics Data Using Diffeomorphic Metric Mapping." *Nature Communications* 14
(1): 8123. https://doi.org/10.1038/s41467-023-43915-7.

Fornace, Mark E., Jining Huang, Cody T. Newman, Nicholas J. Porubsky, Marshall B. Pierce, and
Niles A. Pierce. 2022. "NUPACK: Analysis and Design of Nucleic Acid Structures, Devices,
and Systems." Chemistry. https://doi.org/10.26434/chemrxiv-2022-xv98l.

Harbig, J. 2005. "A Sequence-Based Identification of the Genes Detected by Probesets on the
Affymetrix U133 plus 2.0 Array." *Nucleic Acids Research* 33 (3): e31–e31.
https://doi.org/10.1093/nar/gni027.

Hershberg, Elliot A., Conor K. Camplisson, Jennie L. Close, Sahar Attar, Ryan Chern, Yuzhen Liu,
Shreeram Akilesh, Philip R. Nicovich, and Brian J. Beliveau. 2021. "Author Correction:
PaintSHOP Enables the Interactive Design of Transcriptome- and Genome-Scale
Oligonucleotide FISH Experiments." *Nature Methods* 18 (10): 1265–1265.
https://doi.org/10.1038/s41592-021-01273-6.

Hu, Mengwei, Bing Yang, Yubao Cheng, Jonathan S. D. Radda, Yanbo Chen, Miao Liu, and Siyuan
Wang. 2020. "ProbeDealer Is a Convenient Tool for Designing Probes for Highly
Multiplexed Fluorescence in Situ Hybridization." *Scientific Reports* 10 (1): 22031.
https://doi.org/10.1038/s41598-020-76439-x.

Janesick, Amanda, Robert Shelansky, Andrew D. Gottscho, Florian Wagner, Stephen R. Williams,
Morgane Rouault, Ghezal Beliakoff, et al. 2023. "High Resolution Mapping of the Tumor
Microenvironment Using Integrated Single-Cell, Spatial and in Situ Analysis." *Nature
Communications* 14 (1): 8353. https://doi.org/10.1038/s41467-023-43458-x.

Korsunsky, Ilya, Nghia Millard, Jean Fan, Kamil Slowikowski, Fan Zhang, Kevin Wei, Yuriy
Baglaenko, Michael Brenner, Po-ru Loh, and Soumya Raychaudhuri. 2019. "Fast,
Sensitive and Accurate Integration of Single-Cell Data with Harmony." *Nature Methods*
16 (12): 1289–96. https://doi.org/10.1038/s41592-019-0619-0.

Li, Qiyuan, Nicolai J Birkbak, Balazs Gyorffy, Zoltan Szallasi, and Aron C Eklund. 2011. "Jetset:
Selecting the Optimal Microarray Probe Set to Represent a Gene." *BMC Bioinformatics*
12 (1): 474. https://doi.org/10.1186/1471-2105-12-474.

Liu, Hongfang, Ionut Bebu, and Xin Li. 2014. "Microarray Probes and Probe Sets." *Front Biosci* 2
(1): 325–38. https://doi.org/10.2741/e93.
Marçais, Guillaume, Arthur L. Delcher, Adam M. Phillippy, Rachel Coston, Steven L. Salzberg,
and Aleksey Zimin. 2018. "MUMmer4: A Fast and Versatile Genome Alignment System."
Edited by Aaron E. Darling. *PLOS Computational Biology* 14 (1): e1005944.
https://doi.org/10.1371/journal.pcbi.1005944.
Marçais, Guillaume, and Carl Kingsford. 2011. "A Fast, Lock-Free Approach for Efficient Parallel
Counting of Occurrences of *k* -Mers." *Bioinformatics* 27 (6): 764–70.
https://doi.org/10.1093/bioinformatics/btr011.
McCartney, Daniel L., Rosie M. Walker, Stewart W. Morris, Andrew M. McIntosh, David J.
Porteous, and Kathryn L. Evans. 2016. "Identification of Polymorphic and Off-Target
Probe Binding Sites on the Illumina Infinium MethylationEPIC BeadChip." *Genomics Data*
9 (September):22–24. https://doi.org/10.1016/j.gdata.2016.05.012.
Mecham, Brigham H., Daniel Z. Wetmore, Zoltan Szallasi, Yoel Sadovsky, Isaac Kohane, and
Thomas J. Mariani. 2004. "Increased Measurement Accuracy for Sequence-Verified
Microarray Probes." *Physiological Genomics* 18 (3): 308–15.
https://doi.org/10.1152/physiolgenomics.00066.2004.
Mudge, Jonathan M, Sílvia Carbonell-Sala, Mark Diekhans, Jose Gonzalez Martinez, Toby Hunt,
Irwin Jungreis, Jane E Loveland, et al. 2025. "GENCODE 2025: Reference Gene
Annotation for Human and Mouse." *Nucleic Acids Research* 53 (D1): D966–75.
https://doi.org/10.1093/nar/gkae1078.
O'Leary, Nuala A., Mathew W. Wright, J. Rodney Brister, Stacy Ciufo, Diana Haddad, Rich
McVeigh, Bhanu Rajput, et al. 2016. "Reference Sequence (RefSeq) Database at NCBI:
Current Status, Taxonomic Expansion, and Functional Annotation." *Nucleic Acids*
*Research* 44 (D1): D733–45. https://doi.org/10.1093/nar/gkv1189.
Rouillard, J.-M., Micahel Zuker, and Erdogan Gulari. 2003. "OligoArray 2.0: Design of
Oligonucleotide Probes for DNA Microarrays Using a Thermodynamic Approach."
*Nucleic Acids Research* 31 (12): 3057–62. https://doi.org/10.1093/nar/gkg426.
Varabyou, Ales, Markus J. Sommer, Beril Erdogdu, Ida Shinder, Ilia Minkin, Kuan-Hao Chao,
Sukhwan Park, et al. 2023. "CHESS 3: An Improved, Comprehensive Catalog of Human
Genes and Transcripts Based on Large-Scale Expression Data, Phylogenetic Analysis, and
Protein Structure." *Genome Biology* 24 (1): 249. https://doi.org/10.1186/s13059-023-
03088-4.
Wang, Xiaowei, and Brian Seed. 2003. "Selection of Oligonucleotide Probes for Protein Coding
Sequences." *Bioinformatics* 19 (7): 796–802.
https://doi.org/10.1093/bioinformatics/btg086.
Wernersson, R., and H. B. Nielsen. 2005. "OligoWiz 2.0--Integrating Sequence Feature
Annotation into the Design of Microarray Probes." *Nucleic Acids Research* 33 (Web
Server): W611–15. https://doi.org/10.1093/nar/gki399.
Yahoo Finance. 2025. "10x Genomics, Inc. (TXG)." 10x Genomics, Inc. (TXG). March 19, 2025.
https://finance.yahoo.com/quote/TXG/.
608
609

## Supplementary Figures

Red: RefSeq    Blue: CHESS    Green: GENCODE

probe ID: ENSG00000137285|TUBB2B|1dec8c0

**Supplementary Figure 1. (A)** Screenshot from the Integrated Genome Viewer (IGV) showing the 40bp probe sequence (ID: ENSG00000196154|S100A4|ab4e3dc) that matches both *S100A5* and *S100A4*. Shown are 6 isoforms from the CHESS v3.1 annotation, 4 from GENCODE Basic v47, and 3 from RefSeq v110 for *S100A4*, as well as 2 RefSeq isoforms for the neighboring *S100A5*. The probe sequence aligns to the overlapping region between *S100A5* and *S100A4* gene loci. Matching probe shown in a zoomed-in view below. The forward- and reverse-strand sequences of the probe are shown, and the highlighted area indicates approximately where the probe falls within the gene. **(B)** Screenshot from the Integrated Genome Viewer (IGV) showing the 40bp probe sequence (ENSG00000137285|TUBB2B|1dec8c0) that matches both *TUBB2A* and *TUBB2B*. Shown are the 2 isoforms of *TUBB2A* from the RefSeq v110 annotation, 3 isoforms from the CHESS v3.1 annotation, and 9 from GENCODE Basic v47. Matching probe shown in a zoomed-in view below. The forward- and reverse-strand sequences of the probe are shown, and the highlighted area indicates approximately where the probe falls within the gene. **(A)** and **(B)** share a figure legend.

**Supplementary Figure 2. (A)** Overlap regions between Visium (orange outline) and Xenium (blue outline) data, shown on the Xenium histological image and the Visium histological image, respectively. **(B)** Log transformed aggregated total gene counts for spots (~55$\mu$m x 55$\mu$m) in both Xenium and Visium datasets, overlaid on their corresponding histological image.

612
613
614

**Supplementary Figure 3.** UMAP visualization of integrated scRNA-seq and Xenium datasets: **(A)** before harmony batch correction and **(B)** after harmony batch correction.

**Supplementary Figure 4.** Harmonized UMAP visualization of *TUBB3* and *TUBB4A* expression in the scRNA-seq dataset, as well as the aggregated scRNA-seq expression of *TUBB2B* and all predicted off-targets (*TUBB2A*, *TUBB3*, *TUBB4A*).

**Supplementary Figure 5. (A)** Schematic illustrating that hybridization may still occur even when there is a sequence mismatch at the non-ligated ends of the probe sequence. **(B)** Schematic depicting how probes could bind to each other instead of to their intended target.

**Supplementary Figure 6. (A)** Spatial gene expression patterns for *CEACAM8*: Xenium expression, Visium expression, the aggregated Visium expression combining *CEACAM8* with all predicted off-target binding gene expression, and Visium expression of *CEACAM8*'s potential off-targets (*CEACAM5*, *CEACAM6*, *CEACAM7*, and *PSG6*). **(B)** Comparison of *CEACAM8* expression patterns on harmonized UMAP: Xenium expression, scRNA-seq expression, an aggregated scRNA-seq profile combining *CEACAM8* with all predicted off-target binding gene expression, and scRNA-seq expression of *CEACAM8*'s potential off-targets (*CEACAM5*, *CEACAM6*, *CEACAM7*, and *PSG6*).

**Supplementary Figure 7. (A)** Spatial gene expression patterns for *CEACAM6*: Xenium expression, Visium expression, the aggregated Visium expression combining *CEACAM6* with all predicted off-target binding gene expression, and Visium expression of *CEACAM6* potential off-target (*CEACAM3*). **(B)** Comparison of *CEACAM6* expression patterns on harmonized UMAP: Xenium expression, scRNA-seq expression, an aggregated scRNA-seq profile combining *CEACAM6* with all predicted off-target binding gene expression, and scRNA-seq expression of *CEACAM6*'s potential off-target (*CEACAM3*).

**Supplementary Figure 8. (A)** Spatial gene expression of *HDC* overlaid on the corresponding histological images for Xenium and Visium. **(B)** Harmonized UMAP visualization of *HDC* expression for Xenium and scRNA-seq data.

638
639
640

**Supplementary Figure 9. (A)** Spatial gene expression patterns of: *APOBEC3B* Xenium expression, *APOBEC3B* Visium expression, *APOBEC3F* Visium expression, the aggregated Visium expression combining *APOBEC3B* with all predicted off-target binding gene expression (*APOBEC3F, APOBEC3A, APOBEC3D*), *APOBEC3A* Visium expression, and *APOBEC3D* Visium expression. **(B)** Harmonized UMAP visualization of: *APOBEC3B* Xenium expression, *APOBEC3B* scRNA-seq expression, *APOBEC3F* scRNA-seq expression, the aggregated scRNA-seq expression combining *APOBEC3B* with all predicted off-target binding gene expression (*APOBEC3F, APOBEC3A, APOBEC3D*), *APOBEC3A* scRNA-seq expression, and *APOBEC3D* scRNA-seq expression.

641
642

645    **Supplementary Tables**

| Target gene | Number of probes | Predicted binding genes | Number of probes aligned | Gene types |
|---|---|---|---|---|
| ADGRE5 | 19 | ADGRE5,ADGRE2 | 18,1 | PC,PC |
| ADH1B | 23 | ADH1B,ADH1A,ADH1C | 22,3,3 | PC,PC,PC |
| AKR1C1 | 21 | AKR1C1,AKR1C2,AKR1C3,AKR1C4 | 20,1,1,1 | PC,PC,PC,PC |
| AKR1C3 | 17 | AKR1C3,AKR1C1,AKR1C2,AKR1C4 | 16,1,1,2 | PC,PC,PC,PC |
| APOBEC3A | 10 | APOBEC3A,APOBEC3B | 7,4 | PC,PC |
| APOBEC3B | 14 | APOBEC3B,APOBEC3A, APOBEC3D,APOBEC3F | 13,2,2,2 | PC,PC,PC,PC |
| C15orf48 | 6 | C15orf48,MIR147B | 6,1 | PC, precursor_RNA |
| CEACAM6 | 13 | CEACAM6,CEACAM3 | 13,1 | PC,PC |
| FCGR3A | 20 | FCGR3A,FCGR3B | 20,3 | PC,PC |
| FLNB | 33 | FLNB,LOC124909485 | 33,1 | PC,PC |
| KLRC1 | 15 | KLRC1,KLRC2,KLRC3, KLRC4-KLRK1,KLRC4 | 15,3,1,1,1 | PC,PC,PC,PC,PC |
| KRT6B | 15 | KRT6B,KRT6A,KRT6C, KRT75 | 15,1,4,1 | PC,PC,PC,PC |
| S100A4 | 5 | S100A4,S100A5 | 3,1 | PC,PC |
| SERHL2 | 8 | SERHL2,SERHL | 8,7 | PC,misc_RNA |
| SLAMF1 | 24 | SLAMF1,LOC107985220 | 23,1 | PC,ncRNA |
| SVIL | 20 | SVIL,SVIL2P | 20,2 | PC,misc_RNA |
| TOMM7 | 10 | TOMM7,LOC107986346 | 3,2 | PC,PC |
| TPD52 | 21 | TPD52,TPD52-MRPS28 | 21,5 | PC,PC |
| TPSAB1 | 2 | TPSAB1,TPSB2,TPSD1 | 2,2,1 | PC,PC,PC |
| TUBA4A | 12 | TUBA4A,TUBA4B | 12,1 | PC,PC |
| TUBB2B | 11 | TUBB2B,TUBB2A,TUBB3, TUBB4A | 11,3,1,1 | PC,PC,PC,PC |
| ZNF562 | 22 | ZNF562,ZNF812P | 22,3 | PC,misc_RNA |

647
**Supplementary Table 1.** OPT output of genes with predicted off-target binding based on perfect
sequence homology in RefSeq. This table shows the 22 genes whose probes in the Xenium breast cancer
panel exhibit predicted off-target probe binding, where each off-target alignment involves a perfect
40bp match to the probe sequence. The final column shows the gene types, in order, of each of the off-
target genes shown in column 3. Abbreviations: PC = protein-coding; PG = pseudogene; precursor_RNA
= precursor RNA; misc_RNA = miscellaneous RNA; ncRNA = non-coding RNA.

| Target gene | Number of probes | Predicted binding genes | Number of probes aligned | Gene types |
|---|---|---|---|---|
| ADGRE5 | 19 | ADGRE5, ADGRE2 | 19,1 | PC,PC |
| ADH1B | 23 | ADH1B,ADH1A,ADH1C | 22,3,3 | PC,PC,PC |
| AKR1C1 | 21 | AKR1C1,AKR1C2,AKR1C3, AKR1C4 | 20,1,1,1 | PC,PC,PC,PC |
| AKR1C3 | 17 | AKR1C3,AKR1C1,AKR1C2, AKR1C4 | 17,1,1,2 | PC,PC,PC,PC |
| APOBEC3A | 14 | APOBEC3B,APOBEC3A,APOBEC3D, APOBEC3F | 14,2,2,2 | PC,PC,PC,PC |
| APOBEC3B | 10 | APOBEC3A,APOBEC3B | 8,4 | PC,PC |
| C15orf48 | 6 | C15orf48,MIR147B | 6,1 | PC,miRNA |
| CEACAM6 | 13 | CEACAM6,CEACAM3 | 13,1 | PC,PC |
| EPCAM | 15 | EPCAM,LOC132386 | 14,1 | PC,PG |
| FCGR3A | 20 | FCGR3A,FCGR3B | 20,3 | PC,PC |
| FLNB | 33 | FLNB,LOC124909485 | 33,1 | PC,PC |
| HMGA1 | 7 | HMGA1,HMGA1P1,HMGA1P2, HMGA1P3,HMGA1P6 | 7,1,3,2,2 | PC,PG,PG,PG, PG |
| KLRC1 | 15 | KLRC1,KLRC2,KLRC3,KLRC4-KLRK1,KLRC4 | 15,3,1,1,1 | PC,PC,PC,PC, PC |
| KRT14 | 6 | KRT14,KRT16P6 | 6,1 | PC,PG |
| KRT6B | 15 | KRT6B,KRT6A,KRT6C,KRT75 | 15,1,4,1 | PC,PC,PC,PC |
| KRT8 | 18 | KRT8,KRT8P36,KRT8P11,KRT8P, KRT8P2,KRT8P33,KRT8P45, KRT8P37,KRT8P30,KRT8P17, KRT8P22,KRT8P32,KRT8P42 | 9,1,1,3,2,2, 2,1,1,1,1,1,1 | PC,PG,PG,PG, PG,PG,PG,PG, PG,PG,PG,PG, PG |
| LDHB | 18 | LDHB,LDHBP2 | 18,1 | PC,PG |
| MYLK | 35 | MYLK,MYLKP1 | 29,1 | PC,PG |
| MYO5B | 20 | MYO5B,MYO5BP1,MYO5BP2, LOC102724787 | 20,2,3,2 | PC,PG,PG,PG |
| SCD | 20 | SCD,SCDP1 | 20,6 | PC,PG |
| SERHL2 | 8 | SERHL2,SERHL | 8,7 | PC,PG |
| SMS | 17 | SMS,LOC646347,LOC100419037, LOC642590 | 17,8,1,3 | PC,PG,PG,PG |
| SVIL | 20 | SVIL,SVIL2P | 20,2 | PC,PG |
| TOMM7 | 10 | TOMM7,LOC107986346 | 3,2 | PC,PC |
| TPD52 | 21 | TPD52,TPD52-MRPS28 | 21,5 | PC,PC |
| TPSAB1 | 2 | TPSAB1,TPSB2,TPSD1 | 2,2,1 | PC,PC,PC |
| TRAF4 | 14 | TRAF4,LOC100420513 | 14,1 | PC,PG |
| TUBA4A | 12 | TUBA4A,TUBA4B | 12,1 | PC,PC |
| TUBB2B | 11 | TUBB2B,TUBB2A,TUBB3, TUBB4A | 11,3,1,1 | PC,PC,PC,PC |
| VOPP1 | 24 | VOPP1,LOC729086 | 22,16 | PC,PG |

| VWF | 20 | VWF,VWFP1 | 20,2 | PC,PG |
| ZEB2 | 47 | ZEB2,ZEB2P1 | 28,1 | PC,PG |
| ZNF562 | 22 | ZNF562,ZNF812P | 22,3 | PC,PG |

**Supplementary Table 2.** OPT output of genes with predicted off-target binding based on perfect sequence homology in CHESS. This table shows the 33 genes whose probes in the Xenium breast cancer panel exhibit predicted off-target probe binding, where each off-target alignment involves a perfect 40bp match to the probe sequence. The final column shows the gene types, in order, of each of the off-target genes shown in column 3. Abbreviations: PC = protein-coding; PG = pseudogene; miRNA = microRNA.

| Perfect match | | GENCODE Basic | GENCODE Comprehensive | RefSeq | CHESS |
|---|---|---|---|---|---|
| Number of transcripts | | 158,339 | 387,944 | 181,627 | 158,363 |
| All genes | Off-target probes | 180 | 185 | 60 | 112 |
| | Affected genes | 45 | 45 | 22 | 33 |
| Pseudogenes removed | Off-target probes | 151 | 156 | 60 | 46 |
| | Affected genes | 36 | 36 | 22 | 17 |
| Protein coding genes only | Off-target probes | 58 | 58 | 46 | 45 |
| | Affected genes | 17 | 17 | 17 | 16 |

700
701 **Supplementary Table 3.** The number of off-target probes and affected genes (from the set of 280 genes
702 in the Xenium panel)) found when looking for perfect matches between probe sequences and
703 transcripts in four different reference annotations: GENCODE basic, GENCODE comprehensive, RefSeq,
704 and CHESS. Off-target alignments between CCPG1 probes and DNAAF1-CCPG1 were excluded.
705

| Gene | GENCODE | RefSeq | CHESS |
|---|---|---|---|
| ADGRE5 | ✓ | ✓ | ✓ |
| ADH1B | ✓ | ✓ | ✓ |
| AKR1C1 | ✓ | ✓ | ✓ |
| AKR1C3 | ✓ | ✓ | ✓ |
| APOBEC3A | ✓ | ✓ | ✓ |
| APOBEC3B | ✓ | ✓ | ✓ |
| CEACAM6 | ✓ | ✓ | ✓ |
| FCGR3A | ✓ | ✓ | ✓ |
| KLRC1 | ✓ | ✓ | ✓ |
| KRT6B | ✓ | ✓ | ✓ |
| TPSAB1 | ✓ | ✓ | ✓ |
| TUBA4A | ✓ | ✓ | ✓ |
| TUBB2B | ✓ | ✓ | ✓ |
| AQP1 | ✓ | x | x |
| C1QA | ✓ | x | x |
| CD79B | ✓ | x | x |
| FLNB | x | ✓ | ✓ |
| PDGFRA | ✓ | x | x |
| S100A4 | x | ✓ | x |
| TOMM7 | x | ✓ | ✓ |
| TPD52 | x | ✓ | ✓ |

**Supplementary Table 4.** Union set of protein coding genes that OPT predicts to be affected by off-target binding, across three different reference annotations: GENCODE basic and comprehensive, RefSeq, and CHESS. Genes not shared across all four annotations are colored in red.

| Target gene | Number of probes | Binding genes | Number of probes aligned | Gene types – GENCODE (v47) |
|---|---|---|---|---|
| ACTG2 | 18 | ACTBP2,ACTG1P11,ACTG1P2 | 1,2,2 | PG,PG,PG |
| ANKRD30A | 20 | ENSG00000215398, ENSG00000215512, ENSG00000290354 | 1,1,1 | PG,PG,lncRNA |
| APOC1 | 2 | APOC1P1 | 1 | lncRNA |
| CD83 | 20 | CD83P1 | 1 | PG |
| CEACAM8 | 13 | CEACAM5,CEACAM6,CEACAM7, CEACAMP8,PSG6 | 1,1,1,1,1 | PC,PC,PC,PG,PC |
| CXCL5 | 16 | CXCL6 | 1 | PC |
| ENAH | 20 | ENAHP1 | 2 | PG |
| FLNB | 33 | ENSG00000263343 | 1 | PG |
| KRT5 | 13 | KRT6B,KRT75 | 1,1 | PC,PC |

722 **Supplementary Table 5.** Nine additional genes that were identified to exhibit potential off-target probe
723 binding when allowing for a 6bp error margin on either side of the binding site. Abbreviations: PC =
724 protein-coding; PG = pseudogene; lncRNA = long non-coding RNA.
725
726

| Target gene | Number of probes | Additional binding genes | Number of additional probes aligned | Gene types – GENCODE (v47) |
|---|---|---|---|---|
| ADH1B | 22 | ADH1A,ADH1C | 1,2 | PC,PC |
| AKR1C1 | 21 | AKR1C3,AKR1C2 | 1,1 | PC,PC |
| AKR1C3 | 17 | AKR1C2,AKR1C1,AKR1C5P | 1,2,1 | PC,PC,PG |
| APOBEC3A | 8 | APOBEC3D | 1 | PC |
| CEACAM6 | 13 | CEACAM5,CEACAM7,CEACAM8 | 1,1,1 | PC,PC,PC |
| FCGR3A | 20 | FCGR3B,ENSG00000289768 | 8,1 | PC,PC |
| KRT14 | 6 | ENSG00000290977,KRT16P6, ENSG00000290978,KRT16,KRT16P2 | 1,1,1,1,1 | lncRNA,PG, lncRNA,PC,PG |
| KRT6B | 15 | KRT6A,KRT5 | 2,1 | PC,PC |
| KRT8 | 18 | KRT8P32,KRT8P36, ENSG00000235286,GFAP,KRT8P14, KRT8P15,KRT8P31,KRT8P35, KRT8P43,KRT8P46,KRT8P47, KRT8P48,KRT8P49,KRT8P7,KRT8P9 | 1,1,1,1,1, 1,2,1,1,1, 1,1,1,1,1 | PG,PG,PG,PC, PG,PG,PG,PG, PG,PG,PG,PG, PG,PG,PG |
| LDHB | 18 | LDHBP1 | 1 | PG |
| LILRA4 | 16 | ENSG00000299952,LILRA6,LILRB3, LILRP1,LILRP2 | 1,2,2,1,2 | lncRNA,PC,PC, PG,lncRNA |
| MYO5B | 20 | MYO5BP3 | 1 | PG |
| PTGDS | 5 | ENSG00000284341 | 1 | NMD |
| SMS | 17 | ENSG00000213080, ENSG00000232389, ENSG00000251342 | 2,4,1 | PG,PG,PG |
| SVIL | 20 | ENSG00000290952, ENSG00000304099 | 1,1 | lncRNA, lncRNA |
| TUBB2B | 11 | ENSG00000198211,TUBB3,TUBB4A, TUBB4B,TUBB4BP3 | 1,1,1,1,1 | PC,PC,PC,PC,PG |
| VWF | 20 | VWFP1 | 1 | PG |
| ZNF562 | 22 | ENSG00000291231,ZNF812P, ENSG00000226810,ENSG00000267490, ZNF561 | 1,1,1,1,1 | lncRNA,PG, PG,PG,PC |

**Supplementary Table 6**. Eighteen genes that were previously predicted to be affected by off-target binding based on perfect matching now show additional predicted off-target interactions when a 6bp error margin is allowed on either side of the binding site. This effect is observed either through the accumulation of new probes with predicted off-target binding or via the identification of additional predicted off-target genes per probe. Abbreviations: PC = protein-coding; PG = pseudogene; lncRNA = long non-coding RNA.

736    **Supplementary Data**

737    To access a copy of the Xenium v1 Breast Gene Expression Panel fasta file, open the attached

738    Supplementary           Data           zip           file

739    'xenium_human_breast_gene_expression_panel_probe_sequences.fasta.zip'