Evidence of off-target probe binding affecting 10x Genomics Xenium gene panels compromise

accuracy of spatial transcriptomic profiling

Caleb Hallinan[1,2], Hyun Joo Ji[1,3], Edmund Tsou[1,2,3], Steven L Salzberg[1,2,3,4], Jean Fan[1,2,3]

[1]Center for Computational Biology, Whiting School of Engineering, Johns Hopkins University,

Baltimore, MD 21211, USA

[2]Department of Biomedical Engineering, Johns Hopkins University, Baltimore, MD 21218, USA

[3]Department of Computer Science, Johns Hopkins University, Baltimore, MD 21218, USA

[4]Department of Biostatistics, Johns Hopkins University, Baltimore, MD 21205, USA

Correspondence should be addressed to:

Jean Fan (jeanfan@jhu.edu)

Abstract

The accuracy of spatial gene expression profiles generated by probe-based *in situ* spatially-

resolved transcriptomic technologies depends on the specificity with which probes bind to their

intended target gene. Off-target binding, defined as a probe binding to something other than the

target gene, can distort a gene's true expression profile, making probe specificity essential for

reliable transcriptomics. Here, we investigated off-target binding affecting the 10x Genomics

Xenium technology. We developed a software tool, Off-target Probe Tracker (OPT), to identify

putative off-target binding via alignment of probe sequences and assessing whether mapped loci

corresponded to the intended target gene across multiple reference annotations. Applying OPT

to a Xenium human breast gene panel, we identified at least 14 out of the 313 genes in the panel

potentially impacted by off-target binding to protein-coding genes. To substantiate our

predictions, we leveraged a Xenium breast cancer dataset generated using this gene panel and

compared results to orthogonal spatial and single-cell transcriptomic profiles from Visium

CytAssist and 3' single-cell RNA-seq derived from the same tumor block. Our findings indicate

that for some genes, the expression patterns detected by Xenium demonstrably reflect the

aggregate expression of the target and predicted off-target genes based on Visium and single-

cell RNA-seq rather than the target gene alone. We further applied OPT to identify potential off-

target binding in custom gene panels and integrate tissue-specific RNA-seq data to assess effects.

Overall, this work enhances the biological interpretability of spatial transcriptomics data and

improves reproducibility in spatial transcriptomics research.

Introduction

Recent advances in high-throughput spatially-resolved transcriptomic profiling technologies have enabled the investigation of gene expression with high spatial resolution within tissues. One such commercially available spatial transcriptomics platform is Xenium from 10x Genomics, a publicly traded company with a market capitalization exceeding $2 billion as of November 2025 (Yahoo Finance 2025). Xenium achieves spatial gene expression profiling at single-cell resolution for targeted genes using a probe-based *in situ* detection approach. 10x Genomics currently offers targeted gene panels with pre-designed probe sets. As of December 2024, over 16,000 Xenium consumable reactions have been sold, with each tissue slide profiled costing approximately $5,000, underscoring the platform's widespread use and high commercial value (10x Genomics 2025b; 2025a).

Briefly, Xenium uses padlock probes that include sequences complementary to the RNA of target genes. Once a padlock probe binds to its target, it is ligated and subsequently amplified via rolling circle amplification (RCA). Fluorescently labeled decoder probes then hybridize to the amplified RCA product, enabling the simultaneous detection and decoding of the optical signature, or codeword, specific to each gene in the panel through successive rounds of fluorescence imaging. When combined with cell-segmentation, this approach allows for spatially-resolved single-cell quantification of gene expression.

The accuracy of these gene expression measurements thus relies on the specificity of the probes

to bind to their intended target gene. We define off-target binding as when a probe binds to

something other than the RNA sequence intended to correspond to the target gene (Figure 1).

We note once ligation and RCA occur, the resulting fluorescent signal cannot be distinguished

between on-target and off-target binding. As such, off-target binding can distort the

quantification of the intended target gene's expression, as the observed expression would

represent a combination of the target as well as off-target expression.

Figure 1. Schematic of potential off-target binding in 10x Genomics Xenium. In this illustration,

the arms of the padlock probes were designed to bind an RNA sequence intended to correspond

to a target gene (green). However, these probes exhibit off-target binding and bind to an RNA

sequence in a different off-target gene (red). The probe is circularized and subsequently

amplified via rolling circle amplification (RCA). Hybridization of fluorescent probes to the RCA

product enables the generation a fluorescent signal that is used to quantify RNA expression

within cells.

To predict for such potential off-target binding, we developed Off-target Probe Tracker (OPT), a

software tool that aligns probe sequences to an annotated transcriptome with the option to

allow for mismatches that may still permit probe binding. Using OPT, we identify putative off-

target probe binding to protein-coding genes affecting at least 14 out of 313 genes in a 10x

Genomics Xenium human breast gene panel, compromising the accuracy of their spatial

transcriptomic profiles. We substantiate our predictions using data from orthogonal spatial and

single-cell gene expression profiling technologies. We further apply OPT to identify potential off-

target binding in custom gene panels and integrate tissue-specific RNA-seq data from The Human

BioMolecular Atlas Program (HuBMAP) to assess whether such off-target binding could

meaningfully affect assayed expression patterns in specific tissues. By facilitating a more rigorous

evaluation of probe sequence specificity, tools like OPT can aid in future probe design decisions

to help ensure that probes are optimized to minimize off-target binding based on current

transcriptome annotations.

Results

OPT predicts potential off-target probe binding

To identify potential off-target binding impacting the 10x Genomics Xenium technology, we

require the probe sequences for a specific gene panel of interest, generally represented in a

FASTA file. To this end, we first focus on a human breast gene panel used in the Janesick et al.

publication, courtesy of 10x Genomics (Methods; Supplementary Table 1). This file includes 2,582

probe sequences that are 40bp in length and designed to target 313 genes, including 33 genes

targeted by custom probes, with an average of 8 probe sequences per gene (ranging from 2 to

21 probe sequences per gene). We note this panel represents an earlier iteration of and is highly

similar to the commercially available pre-designed Xenium v1 Human Breast Gene Expression

Panel (Supplementary Note).

To enable the prediction of potential off-target binding, we developed a software tool called OPT

(Methods) that uses nucmer (Marçais et al. 2018) to align probe sequences to various reference

transcriptomes, which comprise curated collections of transcript isoforms for all genes in a

species. OPT features adjustable parameters for binding strictness (e.g., number of mismatches)

and generates a summary file that details all targeted genes along with their potential off-targets

based on the sequence alignments. 10x Genomics designed its probe sequences using the

GENCODE "basic" annotation (Mudge et al. 2025), so initially we also used the latest GENCODE

"basic" annotation (v47) to predict off-target binding for these probes.

We first sought to predict if a probe has off-target binding based on perfect sequence homology

(i.e., if it aligns with 100% identity) with any annotated transcripts other than those that belong

to the intended target gene. Of the 2,582 probe sequences in this gene panel, using GENCODE

v47, OPT identified 121 probe sequences across 37 genes as having off-target binding based on

perfect sequence homology (Table 1). Among the 37 genes with predicted off-target binding, the

number of affected probe sequences per gene ranged from 1 to 8. Overall, these off-target

probes matched 71 other genes, including 20 protein-coding genes, 31 pseudogenes, 10 long

non-coding RNA, 9 transcripts labeled as nonsense-mediated decay, and one microRNA gene.

Target gene	Number of probes	Predicted binding genes	Number of probes aligned	Gene types - GENCODE (v47)
ADH1B	8	ADH1B,ADH1A,ADH1C	8,2,1	PC,PC,PC
AKR1C1	9	AKR1C1,AKR1C2,AKR1C3,AKR1C4,AKR1C5P	9,1,1,1,1	PC,PC,PC,PC,PG
APOBEC3A	8	APOBEC3A,APOBEC3B	8,2	PC,PC
APOBEC3B	8	APOBEC3B,APOBEC3D,APOBEC3F,ENSG00000284554	8,2,2,2	PC,PC,PC,PC
AQP1	10	AQP1,ENSG00000250424	10,4	PC,PC
C15orf48	6	C15orf48,MIR147B	6,1	PC,miRNA
C1QA	4	C1QA,ENSG00000289692	4,2	PC,PC
CD68	7	CD68,ENSG00000264772	7,6	PC,lncRNA
CD79B	5	CD79B,ENSG00000285947	5,3	PC,PC
CD8B	16	CD8B,CD8B2	16, 2	PC,PC
CEACAM6	8	CEACAM6,ENSG00000267881	8,1	PC,PC
CLECL1; CLECL1P	3	CLECL1P,ENSG00000293488	3,3	PG,lncRNA
DPT	8	DPT,LINC00970	8,8	PC,lncRNA
EPCAM	8	EPCAM,ENSG00000225356	8,1	PC,PG
HMGA1	7	HMGA1,HMGA1P1,HMGA1P2,HMGA1P3	7,1,1,1	PC,PG,PG,PG
IL2RG	9	IL2RG,ENSG00000285171	9,8	PC,NMD
KRT14	6	KRT14,KRT16P6,ENSG00000290977	6,1,1	PC,PG,lncRNA
KRT8	16	KRT8,KRT8P3,KRT8P2,KRT8P33,KRT8P45,CDK5R2-AS1,ENSG00000304440,KRT8P11,KRT8P17,KRT8P22,KRT8P30,KRT8P32,KRT8P36,KRT8P37,KRT8P42	16,3,2,2,2,1,1,1,1,1,1,1,1,1,1	PC,PG,PG,PG,PG,lncRNA,lncRNA,PG,PG,PG,PG,PG,PG,PG,PG
LDHB	8	LDHB,ENSG000002854	8,5	PC,NMD
LILRA4	8	LILRA4,ENSG00000275210	8,1	PC,lncRNA
MYLK	11	MYLK,MYLKP1	11,1	PC,PG
MYO5B	8	MYO5B,MYO5BP1,MYO5BP2, ENSG00000266997	8,1,1, 4	PC,PG,PG,NMD
PCLAF	8	PCLAF,ENSG00000259316	8,1	PC,NMD
POLR2J3	10	POLR2J3,POLR2J4,POLR2J,ENSG00000270249,POLR2J2,POLR2J2-UPK3BL1,ENSG00000291154	10,4,3,2,2,2,1	PC,lncRNA,PG,PC,PC,PC,NMD,lncRNA
PTGDS	5	PTGDS,ENSG00000284341	5,3	PC,NMD
SCD	8	SCD,SCDP1	8,2	PC,PG
SERHL2	8	SERHL2,SERHL	8,7	PC,PG
SERPINA3	8	SERPINA3,ENSG00000273259	8,8	PC,NMD
SLAMF1	10	SLAMF1, ENSG00000228863	10, 1	PC, lncRNA
SMS	8	SMS,ENSG00000213080,ENSG00000232389,ENSG00000249779	8,3,1,1	PC,PG,PG,PG
THAP2	13	THAP2,ENSG00000258064	13,2	PC,NMD
TPD52	8	TPD52,ENSG00000276418	8,5	PC,NMD
TPSAB1	2	TPSAB1,TPSB2,TPSD1	2,2,1	PC,PC,PC
TRAF4	9	TRAF4,ENSG00000225869	9,1	PC,PG
TUBB2B	8	TUBB2B,TUBB2BP1	8,1	PC,PG
VOPP1	11	VOPP1,ENSG00000223612	11,1	PC,PG
VWF	8	VWF,VWP1	8,1	PC,PG

Table 1. OPT output of genes with predicted off-target binding based on perfect sequence

homology using GENCODE v47. This table shows the 37 genes whose probes in the 10x Genomics

Xenium v1 Human Breast Gene Expression Panel exhibit predicted off-target probe binding,

where each off-target alignment involves a perfect 40bp match to the probe sequence. Although

OPT predicted off-target binding of *CCPG1* probe sequences to the *DNAAF1-CCPG1* gene, we

manually excluded it from our list because *DNAAF1-CCPG1* is a read-through gene containing

portions of both *DNAAF1* and *CCPG1*. The final column shows the gene types, in order, of each

of the off-target genes shown in column 3. Abbreviations: PC = protein-coding; PG = pseudogene;

NMD = nonsense-mediated decay; lncRNA = long non-coding RNA.

<u>Off-target binding predictions vary across different annotations</u>

OPT relies on alignments to an annotated transcriptome, which ideally reflects all genes and gene

variants stably transcribed in a given species. However, genome annotation is still an active area

of research (Varabyou et al. 2023), with discrepancies across annotations in gene counts,

isoforms, and many other features. We therefore further used OPT to predict for off-target

binding and affected genes using two additional human genome annotation sets, RefSeq (v110),

(O'Leary et al. 2016) and CHESS (v3.1.3) (Varabyou et al. 2023), and compared to our previous

results from GENCODE (v47) (Mudge et al. 2025) (Methods).

When considering only perfect sequence homology, while we previously found 37 affected genes

using GENCODE, we found 14 when using RefSeq and 23 when using CHESS (Supplementary Table

2, 3). Given that RefSeq and CHESS have more transcripts than GENCODE, these discrepancies in

off-target binding predictions was not simply an artifact of the difference in transcript set sizes.

While the human annotation databases mostly agree on the number of protein-coding genes in

the genome, they remain widely divergent on pseudogenes and lncRNA genes. Therefore, we

focused on how the results change when we restrict our analysis to only protein-coding genes.

By excluding pseudogenes (which are presumably not expressed), lncRNAs, and other non-

protein-coding RNAs when using OPT, the number of affected genes fell to 11 for GENCODE, 10

for RefSeq and 9 for CHESS (Supplementary Table 4). Again, these discrepancies reflect

annotation   differences.   For   example,   the   probe   sequence

(ENSG00000196154|S100A4|ab4e3dc), which was designed to target *S100A4* based on

GENCODE annotations, also aligns to *S100A5* in RefSeq (Supplementary Figure 1). We reason that

if a probe sequence aligns off-target to a protein-coding gene based on any of these annotations,

it could result in off-target binding. We therefore focused further analysis on the union of genes

with predicted off-target binding to protein-coding genes across the 3 annotations, resulting in

14 genes: *ADH1B, AKR1C1, APOBEC3A, APOBEC3B, AQP1, C1QA, CD79B, CD8B, CEACAM6,*

POLR2J3, S100A4, TOMM7, TPD52, and *TPSAB1* (Supplementary Figure 2; Supplementary Table

5).

<u>Comparison with Visium CytAssist reveals spatial gene expression patterns consistent with off-</u>

<u>target binding</u>

To investigate the potential effects of our predicted off-target binding for this Xenium human

breast gene panel in experimental settings, we compared spatial gene expression patterns

detected in two previously published spatial transcriptomics datasets from serial sections of the

same breast cancer tissue: one section assayed with Xenium using this gene panel, and another

assayed using Visium CytAssist, an orthogonal spatial transcriptomics platform (Janesick et al.

2023). Briefly, Visium CytAssist is a sequencing-based spatial transcriptomics technology in which

RNA is hybridized to spatially barcoded capture spots on a slide, enabling spatial transcriptomic

mapping after sequencing. However, while Xenium offers single-cell resolution gene expression

quantification, Visium quantifies gene expression within 55μm spots. To enable direct

comparison, we first structurally aligned the Xenium and Visium tissue sections using STalign

(Clifton et al. 2023), restricting our analysis to overlapping regions since different parts of the

tissue were profiled (Supplementary Figure 3A). To improve visual comparability, we aggregated

the Xenium gene expression data at the aligned locations to match the Visium spatial resolution

and visualized using resolution-matched tiles (Supplementary Figure 3B; Methods).

Among the 14 genes exhibiting off-target binding to protein-coding genes, 4 were present in the

Visium dataset that had at least one corresponding off-target gene also detected in the dataset.

For genes with no predicted off-target binding based on perfect sequence homology such as

MS4A1, we observed a visually similar spatial pattern between the two technologies (Figure 2A),

suggesting that spatially aligned groups of cells across the two technologies express this gene at

comparable relative magnitudes. This can be quantitively assessed by comparing the pseudo-log

gene expression values from both Visium and Xenium at matched spatial locations and

computing the RMSE and Pearson correlation in a manner similar to STcompare (Clifton et al.

2025). For *MS4A1*, the RMSE is relatively low at 3.746, and the Pearson correlation of 0.382

indicates a moderate degree of concordance between the two technologies. However, for genes

with predicted off-target binding based on perfect sequence homology such as *APOBEC3B*, we

observed a visually dissimilar spatial pattern between the two technologies (Figure 2B).

Consistent with this, the RMSE is relatively high at 5.452, and the Pearson correlation is *nan*

because *APOBEC3B* is not expressed in the Visium dataset. Importantly, its predicted off-target

genes, *APOBEC3D* and *APOBEC3F*, in Visium shows a visually more similar spatial pattern to the

Xenium *APOBEC3B*. To better visualize the effect of off-target binding within the Xenium data,

we aggregated the expression of each gene along with its predicted off-targets found in the

Visium dataset and visually compared across spatial locations. Notably, the spatial pattern of the

aggregated expression of *APOBEC3B, APOBEC3D* and *APOBEC3F* in Visium is visually more similar

to the spatial pattern of *APOBEC3B* in Xenium. Quantitatively, comparing this aggregated

expression results in a decrease in RMSE to 4.465 and a non-*nan* Pearson correlation of 0.160,

consistent with the prediction that Xenium *APOBEC3B* probes exhibit off-target binding to

APOBEC3D and *APOBEC3F*. We further visually confirmed using the Integrative Genomics Viewer

(IGV) where two probe sequences intending to bind to *APOBEC3B* were found to perfectly align

to sequences in both *APOBEC3D* and *APOBEC3F*, consistent across all annotations evaluated

(Supplementary Figure 4).

Overall, when comparing the total gene expression between the two technologies, we observed

a generally strong positive correlation, consistent with the previously published work (Janesick

et al. 2023). We do not observe an obvious trend between gene expression magnitude and the

presence of predicted off-target probes (Figure 2C), suggesting that off-target binding prediction

alone does not explain the observed higher expression magnitude in Xenium compared to

Visium, which may still be attributed to variation in detection efficiency, sequencing depth, and

other factors.

Figure 2. Comparison of spatial gene expression patterns between Xenium and Visium. (A)

Spatial gene expression of *MS4A1* overlaid on the corresponding histological images for Xenium

and Visium, accompanied by a density plot comparing Xenium vs. Visium *MS4A1* expression. The

dotted line indicates the identity line (X = Y), and the solid line represents the line of best fit. **(B)**

Gene expression patterns for *APOBEC3B*: Xenium expression, Visium expression, the aggregated

Visium expression combining *APOBEC3B* and its predicted off-target gene's expression

APOBEC3D and *APOBEC3F*, and Visium expression of *APOBEC3B*'s predicted off-targets

APOBEC3D and *APOBEC3F*. Two density plots are shown: one comparing Xenium vs. Visium for

APOBEC3B alone, and one comparing Xenium vs. the aggregated Visium expression of *APOBEC3B*

with all off-targets. The dotted line indicates the identity line (X = Y), and the solid line represents

the line of best fit. (**C**) Scatterplot of log-transformed total expression counts (with a

pseudocount) for 307 genes comparing Visium and Xenium data. The dotted line indicates the

identity line (X = Y), and points (genes) are colored by probe information.

<u>Comparison with scRNA-seq reveals single-cell gene expression patterns consistent with off-</u>

<u>target binding</u>

To further investigate the potential effects of our predicted off-target binding for this Xenium

human breast gene panel, we compared the detected single-cell gene expression patterns in the

same previously published work using Chromium Next GEM Single Cell 3' (Janesick et al. 2023).

Briefly, single-cell RNA sequencing (scRNA-seq) with 3' end capture is a technique used to profile

gene expression at the single-cell level by profiling the 3' ends of mRNA transcripts with

sequencing followed by alignment to a genome or transcriptome for quantification. While this

approach provides single-cell resolution gene expression quantification, it lacks spatial

information. To enable a single-cell comparison with Xenium, we use Harmony (Korsunsky et al.

2019) to remove batch effects and project cells into a shared UMAP embedding (Supplementary

Figure 5A-B). We also performed Leiden clustering on the harmonized principal components to

quantitively compare cluster expression (Supplementary Figure 5C; Methods).

Among the 14 genes exhibiting off-target binding to protein-coding genes, 10 were present in

the scRNA-seq dataset that had at least one corresponding off-target gene also detected in the

dataset. Again, for genes with no predicted off-target binding based on perfect sequence

homology such as *MS4A1*, we observed a visually similar gene expression pattern in the

harmonized UMAP across both technologies (Figure 3A), suggesting that transcriptionally similar

clusters of cells or cell-types across the two technologies express this gene at comparable relative

magnitudes. We further quantitatively assessed the data by comparing the pseudo-log gene

expression values obtained from the clusters within the clustered harmonized UMAP and

computed the RMSE and Pearson correlation. For *MS4A1*, the RMSE is relatively low at 0.479,

and the Pearson correlation of 0.991 indicates a strong degree of concordance between the two

technologies. Likewise, again, for genes with predicted off-target binding based on perfect

sequence homology such as *APOBEC3B*, we observed a visually dissimilar gene expression

pattern on the harmonized UMAP (Figure 3B). Consistent with this, the RMSE is relatively high at

0.829, and the Pearson correlation is *nan* because *APOBEC3B* is not expressed in the scRNA-seq

dataset. Again, its predicted off-target genes, *APOBEC3D* and *APOBEC3F,* in scRNA-seq showed a

visually more similar expression pattern in the harmonized UMAP embedding to the Xenium

APOBEC3B. To better illustrate the impact of the off-target probes, we again aggregated the

expression of a gene and its predicted off-target genes present in the scRNA-seq data and visually

compared across the harmonized UMAP embedding. The aggregated expression of *APOBEC3B*,

APOBEC3D and *APOBEC3F* in the scRNA-seq data shows a visually more similar gene expression

pattern in the harmonized UMAP embedding to *APOBEC3B* in the Xenium data (Figure 3B).

Quantitatively, comparing this aggregated expression results in a decrease in RMSE to 0.596 and

a non-nan in Pearson correlation to 0.417, consistent with the prediction that Xenium *APOBEC3B*

probes exhibit off-target binding with these paralogs.

Overall, when comparing total gene expression between the two technologies, we again

observed a generally strong positive correlation (Figure 3C), similar to the Visium comparison

results and consistent with the previously published work (Janesick et al. 2023).

Figure 3. Comparison of single-cell gene expression patterns between Xenium and scRNA-seq.

(A) Harmonized UMAP visualization of *MS4A1* expression for Xenium and scRNA-seq data,

accompanied by a scatterplot comparing Xenium vs. scRNA-seq *MS4A1* cluster expression. The

dotted line indicates the identity line (X = Y), and the solid line represents the line of best fit. **(B)**

Comparison of *APOBEC3B* expression patterns on harmonized UMAP: Xenium expression, scRNA-

seq expression, an aggregated scRNA-seq profile combining *APOBEC3B* and its predicted off-

target gene's expression *APOBEC3D* and *APOBEC3F*, and scRNA-seq expression of *APOBEC3B*'s

predicted off-targets *APOBEC3D* and *APOBEC3F*. Two scatterplots are shown: one comparing

Xenium vs. scRNA-seq for *APOBEC3B* cluster expression alone, and one comparing Xenium vs. the

aggregated scRNA-seq cluster expression of *APOBEC3B* and it's predicted off-targets. The dotted

line indicates the identity line (X = Y), and the solid line represents the line of best fit. **(C)**

Scatterplot of log-transformed total expression counts (with a pseudocount) for 313 genes

between Visium and scRNA-seq data. The dotted line indicates the identity line (X = Y), and points

(genes) are colored by probe information.

<u>OPT results when allowing mismatches at the terminal ends of the probe sequences identifies</u>

<u>additional off-target candidates</u>

Thus far, we have focused on predicting off-target binding based on perfect sequence homology.

However, we reason that imperfect sequence matching could still result in off-target binding.

Specifically, for Xenium v1, which underlies the Xenium human breast gene panel, padlock probes

with two 20 base pair (bp) arms bind complementary mRNA regions, forming a 40bp probe

sequence. A ligase then circularizes the padlock probe, favoring specific 2-bp junctions.

Importantly, if there is a sequence mismatch, particularly outside the ligation site towards the

terminal ends of the probe sequence, hybridization may still occur and result in off-target

binding, albeit with reduced hybridization efficiency (Supplementary Figure 6A). We therefore

added an option in OPT to allow imperfect alignments at the ends of the probe sequences,

specifying the sequence length at either ends, where mismatches, insertions, deletions, or

clipping can occur (Methods).

Allowing for 10bp mismatches on either end of the 40bp probe sequence (i.e., requiring a 20bp

match covering the middle of the probe sequence including the ligation site) revealed 18

additional genes, including to protein-coding genes, with potential off-target binding when using

GENCODE v47 (Supplementary Table 6), among which *ACTG2* was included. Additionally, 10 of

the 37 genes previously predicted to be affected by off-target binding based on perfect sequence

homology were now predicted to have additional off-target genes, including to protein-coding

genes (Supplementary Table 7), among which *TUBB2B* was included. Both genes showed visually

dissimilar spatial patterns between Xenium and Visium, accompanied by comparably high RMSE

and low Pearson correlation (Supplementary Figure 7A-B). In contrast, the spatial pattern of the

aggregate of *TUBB2B* and *ACTG2* with their predicted off-target protein-coding genes (*TUBB2A*

and *ACTB*/*ACTA1*/*POTEM* respectively) in Visium more closely resembled the spatial pattern of

TUBB2B and *ACTG2* observed in Xenium, accompanied by a corresponding decrease in RMSE for

both genes and increase in Pearson correlation for *TUBB2B*. Likewise, a similar trend is observed

in the scRNA-seq comparison (Supplementary Figure 8A-B). Ultimately, these findings suggest

that off-target binding, even with imperfect sequence matching, can contribute to the expression

patterns observed in Xenium.

RNA-seq reference atlases suggest off-target binding can variably impact results in Xenium

custom probe panels

Our analyses to this point focused on the Xenium human breast gene panel from Janesick et al.,

a predecessor to and thus similar to the commercially available pre-designed Xenium v1 Human

Breast Gene Expression Panel (Supplementary Note). We next sought to determine whether such

off-target binding could affect custom Xenium gene panels, in which gene selection is specified

by the user rather than provided as pre-designed panels by 10x Genomics. To investigate this, we

leveraged two custom Xenium gene panels used by the Human BioMolecular Atlas Program

(HuBMAP) (Jain et al. 2023): one designed for the placenta and another designed for the kidney,

lung, and heart (i.e. multi-organ). Applying OPT to the associated FASTA files using the GENCODE

v47 annotation and allowing for 10bp mismatches on either end of the 40bp probes, we found

that 49 genes out of the 300 targeted genes in the placenta panel (Supplementary Table 8) and

24 genes out of the 300 targeted genes in the multi-organ (kidney, lung, heart) panel had

predicted off-target binding (Supplementary Table 9). Of these, 30 of the 49 placenta panel genes

and 11 of the 24 multi-organ panel genes had predicted off-targets that were protein-coding

genes.

To assess the potential effects of our predicted off-target binding for these custom Xenium gene

panels in experimental settings, we examined the expression of the corresponding predicted off-

target genes in matched scRNA-seq or bulk RNA-seq data from the HuBMAP consortium for each

of the relevant tissue types. For the placenta custom panel, 34 of the 49 genes with predicted

off-target genes were detected with non-zero expression magnitudes in the placenta bulk RNA-

seq dataset (Supplementary Figure 9). For the multi-organ panel, of the 21 genes with predicted

off-target genes, 13 were detected with non-zero expression magnitudes in the heart, 12 in the

kidney, and 13 in the lung (Supplementary Figure 10). Together, these results suggest that off-

target binding can impact custom Xenium gene panels by distorting observed Xenium gene

expression measurements in a tissue-dependent manner, particularly where the off-target gene

is expressed at a higher magnitude compared to the target gene within the tissue of interest.

Discussion

Our study presents evidence of off-target probe binding that may distort gene expression profiles

affecting the 10x Genomics Xenium spatial transcriptomics technology. We identified at least 14

out of the 313 genes in a Xenium human breast gene panel, which is highly similar to the

commercially-available pre-designed Xenium v1 Human Breast Gene Expression Panel

(Supplementary Note), that may be affected by off-target probe binding based on sequence

similarity, supported by spatial and single-cell comparative analyses using Xenium with serial

section datasets from Visium CytAssist and 3' single-cell RNA-seq respectively. We further

identified potential off-target probe binding affecting custom Xenium gene panels. To assist in

the interpretation of existing probe-based gene expression data as well as future probe design,

we provide OPT as a software tool for predicting potential off-target probe binding. For future

reference, we have run OPT on all publicly available 10x Genomics pre-designed Xenium gene

panels and supply them as a ZIP file (Supplementary Table 10).

Although we have predicted off-target binding based on sequence alignment, its effect on gene

expression quantification may still vary. One reason is that the off-target protein-coding or non-

protein-coding gene may not be expressed (Supplementary Figure 11). For example, in the

Xenium human breast gene panel, although *ADH1B* probes have predicted off-target binding to

ADH1A and *ADH1C* based on perfect sequence homology, the sparse expression of *ADH1A* and

ADH1C in both the Visium and scRNA-seq breast cancer data led to only a minor difference in the

aggregated expression and quantitative results (Supplementary Figure 12). Our analysis of

HuBMAP custom probe panels demonstrated how to evaluate for the potential impact of

predicted off-targets by integrating tissue-specific single-cell or bulk RNA-seq data from

reference atlases (Supplementary Figures 9, 10). Overall, we anticipate evaluating whether

predicted off-target genes are expressed in a tissue-specific manner will be useful for gauging

whether predicted off-target binding is likely to meaningfully affect observed gene expression

and interpretation when applied to a tissue of interest.

Other sources of non-specific signal may also arise, including probe self-hybridization or probe-

probe interactions (Supplementary Fig 6B). In general, probe binding specificity is influenced by

numerous factors, with many methods previously developed to aid in the design of probe

sequences while taking these factors into consideration (Wang and Seed 2003; Rouillard et al.

2003; Wernersson and Nielsen 2005; Chou 2010; Li et al. 2011; Hu et al. 2020; Hershberg et al.

2021; Fornace et al. 2022; Stenberg et al. 2005; Kuemmerle et al. 2024). For example, in the

Xenium human breast gene panel, *HDC*, a custom gene not included in the pre-designed Xenium

v1 Human Breast Gene Expression Panel, did not have any off-targets predicted by OPT. Yet in

Xenium, *HDC* exhibited a distinct spatial pattern and high global expression level, whereas in both

Visium and scRNA-seq, *HDC* showed a minimal spatial pattern and sparse expression level

respectively (Supplementary Figure 13). This illustrates how discrepancies across platforms can

signal potential off-target activity not captured by alignment-based predictions alone and

highlights the general importance of experimental validation with orthogonal technologies since

sequence alignment-based tools such as OPT may not flag all potential discrepancies.

We note that most off-target binding impacts paralogs, homologous genes that have diverged

following gene duplication events. Paralogs often belong to large gene families whose members

can share high sequence similarity, increasing the risk of off-target probe activity. Unlike

orthologs with conserved functions across species, paralogs are additional copies that can

acquire function-altering mutations (Platt et al. 2000; Pevny et al. 1991; Tsai et al. 1994). Pooling

expression signals across paralogs can therefore prevent researchers from capturing their distinct

functional roles.

We also found that 10 of the 2,582 Xenium human breast gene panel probe sequences did not

align to any reference transcripts in the GENCODE v47 annotation. Upon manually aligning these

probes to the GRCh38 genome, we determined that each unmapped sequence corresponded

either to regions immediately upstream or downstream of annotated transcripts, or to intronic

sequences that would not typically be present in mature RNA. Interestingly, when we aligned

several of these probes, such as (ENSG00000125878|TCF15|5d3cbc2) and

(ENSG00000169083|AR|a0c6719), to an earlier annotation (GENCODE v28), they instead

mapped to exonic regions (Supplementary Figure 14). This suggests that these intronic or

intergenic probe sequences were likely designed using older GENCODE versions in which those

regions were annotated as exons. This finding illustrates the importance of disclosing the specific

annotation version to promote reproducibility, as well as the ongoing variability of human gene

annotation. Likewise, as evidenced by our analysis across GENCODE, RefSeq, and CHESS, we

emphasize the variation across these reference annotations and therefore recommend using

multiple annotations when designing probe sequences and evaluating them for off-target effects

to ensure a more comprehensive assessment.

Given these challenges, we advise probes with predicted off-target binding to protein-coding

genes based on high sequence homology be avoided in future experiments. Likewise, we

encourage the use of tools like OPT to aid in future probe design decisions and help ensure that

probes are optimized to minimize off-target binding based on the most current transcriptome

annotations. When probes with predicted off-targets cannot be avoided, we encourage the

integration of tissue-specific RNA-seq data from HuBMAP and other reference atlases to evaluate

for its potential impact. Further, such integration of tissue-specific RNA-seq data from reference

atlases into the probe design process itself may offer a data-driven opportunity to minimize the

impact of potential off-target binding by enforcing stricter probe-design constraints only where

potential off-target genes are highly expressed in the tissue of interest. For datasets that have

already been generated using probes with predicted off-target binding, we generally recommend

taking into consideration these predictions to avoid drawing misleading conclusions. For

example, we recommend expression measurements for genes with predicted off-target binding

be omitted from training foundation models to avoid error propagation. Alternatively, when

performing integrative analyses that compare or align gene expression with measurements

across technologies, it may be necessary to incorporate off-target binding predictions. For

instance, integration could be performed between the observed Xenium gene expression and the

aggregated expression of the target and predicted off-target genes for the orthogonal

technology. Finally, existing literature that base conclusions on genes with predicted off-target

binding should be interpreted with caution.

We emphasize that these findings were missed in the previous Janesick et al. publication from

10x Genomics (Janesick et al. 2023). Consistent with previously published observations, we

observed a highly correlated total gene expression magnitude between Xenium and Visium as

well as scRNA-seq. However, a notable exception is *APOBEC3B*, which is not expressed according

to both Visium and scRNA-seq but highly expressed according to Xenium (Figure 2B; Figure 3B) –

a discrepancy that Janesick et al. omitted. We emphasize that positive significant average gene

expression correlation is a necessary but not sufficient metric for consistency across technologies

and that individual data points should be scrutinized. Likewise, validation with orthogonal

technologies could have helped identify discrepancies suggestive of off-target effects. We note

Janesick et al. used immunofluorescence to validate two genes, *ERBB2* and *MS4A1*, which by our

analysis were predicted to exhibit no off-target binding. Although Xenium incorporates blank and

negative control probes that are intended to help quantify the rate of non-specific and potential

off-target binding, our findings suggest that relying solely on such probes for error detection may

be insufficient. Implementing probe redundancy, where the same gene is targeted using different

codewords, could provide an additional internal control to enable the detection of off-target

binding.

Although we focus here on the 10x Genomics Xenium technology, we do not exclude the

possibility that off-target binding may similarly affect other probe-based gene detection

approaches from other commercial vendors. Any technology that relies on hybridization-based

detection is inherently susceptible to off-target probe binding when sequence similarity exists.

Further, hybridization-based detection often inherently involves a trade-off between sensitivity

and specificity. Given these inherent technological limitations, we therefore emphasize the

importance of transparency through sharing probe sequences. However, many companies do not

release the probe sequences used in their assays, limiting the consumer's ability to fully interpret

their results as well as the community's ability to effectively characterize and benchmark

performance variation across platforms. Therefore, we strongly recommend that companies

publish probe sequences for pre-designed panels and likewise that researchers using these

technologies should obtain and publish probe sequences used in their studies to support

transparent and reproducible science.

This is not the first instance in which a commercially available platform has encountered

challenges in probe design (McCartney et al. 2016; Harbig 2005; Mecham et al. 2004; Liu et al.

2014). These findings underscore the critical role of academic researchers towards ensuring the

robustness of industry-led product development by providing oversight, free of financial conflicts

of interest through independent federal funding. This complementarity between industry and

academia fosters a more rigorous, transparent, and reliable scientific process, ultimately to the

benefit of consumers and the public. By shedding light on putative off-target probe binding as

well as by providing a tool to enable such off-target binding predictions, this work will help

enhance the quality of spatial transcriptomics data and improve the overall reproducibility in

spatial transcriptomics research.

Methods

Off-target Probe Tracker Tool (OPT)

OPT (Off-target Probe Tracker) is a Python program that runs nucmer (Marçais and Kingsford

2011) for alignment and then processes the results to predict probe binding based on sequence

homology. OPT is available as an open-source Python toolkit at https://github.com/JEFworks-

Lab/off-target-probe-tracker. When a user provides a query probe sequence file, a target

transcript sequence file, and the annotation used to extract these transcripts, OPT outputs which

gene each probe is likely to bind to. Nucmer is a fast nucleotide sequence aligner that uses

maximal exact matches as anchors which it then joins together to find longer alignments. By

default, OPT saves nucmer results in SAM format and finds perfect sequence matches between

a query probe and a target transcript, requiring that alignments consist of only matches and cover

the entirety of the query. OPT consists of four modules: (1) `flip` for reverse complementing

probe sequences aligned to the opposite strand of their target genes; (2) `track` for aligning

probe sequences and processing alignment results; (3) `stat` for compiling summary statistics on

the number of off-target binding probes and affected genes; and (4) `all` for running the `flip`,

`track`, and `stat` modules at once.

In the case that a probe's target gene has synonyms, we consider alignments to genes annotated

with one of its synonym to still be on-target. For example, if a probe that targets *NARS* shows

alignments to a gene called *NARS1*, we don't consider it to be off-target binding. We gathered

relevant gene synonym relationships using the GeneCards and HGNC online database.

OPT also provides a "pad" mode in which imperfect alignments are allowed at either end of the

query (i.e., probe sequence). The -pl parameter sets the pad length at either end of the query

and OPT allows for any number of mismatches in these padded regions. For example, if the pad

length is 10 and the probe sequence length is 40bp, then the middle 20bp are the only part of

the probe sequence required to match. As long as the critical region is intact, OPT reports an off-

target binding site based on this alignment. By default, -pl is set to 0, and the pad mode is

activated by providing a non-zero integer to -pl.

<u>Obtaining probe sequences for the Xenium v1 Human Breast Gene Expression Panel</u>

To identify potential off-target binding impacting the 10x Genomics Xenium v1 Human Breast

Gene Expression Panel, we obtained the FASTA file of probe sequences from the Janesick et al.

publication courtesy of 10x Genomics and available as Supplementary Table 1 for preservation.

Notably, this panel slightly deviates from the commercially available Xenium v1 Human Breast

Gene Expression Panel (Supplementary Note).

The target gene names and IDs were extracted from the probe IDs of the following format:

`> gene_id|gene_name|accession`

We expected the provided FASTA file to contain probe sequences to be the reverse

complemented sequence of their intended target genes and hence align to the reverse strand of

their target isoforms. However, when we aligned the breast panel probe sequences to the

GENCODE basic (v47) reference transcripts using nucmer, we found that 2,563 / 2,582 of probe

sequences aligned on the reverse strand of their target transcripts (i.e., isoforms of their target

genes). For consistency, we enforced that all probe sequences be oriented in the same direction

and align to the forward strand of their target genes and transcripts. As such, we reverse-

complemented these 2,563 probe sequences. We also added this functionality as an OPT module

called "flip" in which probe sequences aligned to the reverse strand of their targets are reverse

complemented. We expect probe sequences to align to the forward strand of transcripts (i.e.,

both oriented in the same direction) during the downstream probe sequence binding prediction

step.

Visium Comparison

The Visium CytAssist dataset, collected from a breast cancer tissue block utilized in Janesick et

al., was downloaded from the 10x Genomics website

(https://www.10xgenomics.com/products/xenium-in-situ/preview-dataset-human-breast). This

dataset originally contained 4,992 spots with x-y coordinates and included 18,085 genes per spot.

Of the 313 unique genes in the Xenium dataset, 307 were shared with the Visium dataset; the

other six genes (*AKR1C1*, *ANGPT2*, *BTNL9*, *CD8B*, *POLR2J3*, and *TPSAB1*) were excluded from the

analysis because they were absent from the Visium dataset.

To compare spatial gene expression patterns from Visium and Xenium technologies, we first

mapped all the data to the same coordinate space. We used STalign (v1.0.1), a computational

tool that utilizes affine transformations along with diffeomorphic metric mapping to align target

and source datasets (Clifton et al. 2023). The initial alignment involved only affine

transformations and eight manually determined landmarks to align the Visium histology image

(source) to the Xenium histology image (target). This transformation brought the Visium image

into the coordinate space of the higher-resolution Xenium image. We then applied this learned

transformation to the Visium spots, ensuring that they were correctly positioned relative to both

histology images. Next, we used STalign to map the Xenium transcripts (source) onto their

corresponding Xenium histology image (target) using both affine and diffeomorphic metric

mapping. The transcripts were rasterized at 30μm resolution, with an initial affine transformation

guided by four manually defined landmarks. Diffeomorphic metric mapping was then performed

with the following parameters: a = 2500, epV = 1, niter = 2000, sigmaA = 0.11, sigmaB = 0.10,

sigmaM = 0.15, sigmaP = 50, muA = [1, 1, 1], muB = [0, 0, 0], with all other settings left at their

defaults. We extracted the overlapping regions between the two datasets (Supplementary Figure

3A), which reduced the total spots in the Visium dataset to 3,958. Finally, we aggregated the

Xenium gene expression data to ~55μm x 55μm patches that correspond to the spatial locations

of the Visium spots, resulting in matched-resolution spatial gene expression for both

technologies (Supplementary Figure 3B). The Visium spatial gene expression data is displayed as

patches rather than spots to enhance visual saliency and ensure consistency with the Xenium

spatial gene expression plots.

After obtaining matched-resolution spatial gene expression matrices, we quantified agreement

between the Visium and Xenium data. For each gene shared between the two platforms, we

constructed expression vectors across the aligned spatial spots, where each element

corresponded to the log-normalized gene expression at its matched location in the tissue section.

We then compared the Visium and Xenium vectors for each gene using two metrics: root-mean-

square error (RMSE), computed relative to the line y = x, and Pearson correlation (r). To assess

whether discrepancies in Xenium measurements could be attributed to predicted off-target

genes, we compared the Xenium data of a target gene to the aggregated Visium expression of

the target genes with its predicted off-targets. We summed the raw Visium counts of all off-target

genes associated with each Xenium gene with predicted off-targets, re-normalized the

aggregated counts using CPM followed by $\log(x + 1)$, and again calculated the RMSE and Pearson

correlation. This approach enabled us to evaluate whether Xenium expression patterns aligned

more closely with the intended target gene alone or with the combined expression of its

predicted off-target genes.

<u>Single-cell RNA-seq Comparison</u>

The Chromium Next GEM 3' scRNA-seq dataset, collected from a breast cancer tissue block

utilized in Janesick et al., was downloaded from the 10x Genomics website

(https://www.10xgenomics.com/products/xenium-in-situ/preview-dataset-human-breast). This

dataset contained 12,388 cells with 36,601 genes per cell. All 313 unique genes present in the

Xenium dataset are also in the scRNA-seq dataset, hence both datasets were subsetted to these

genes for the analysis.

Both scRNA-seq and Xenium provide single-cell resolution data. To integrate these datasets, we

first removed cells lacking detectable gene expression. We then normalized the combined gene

expression data using counts per million (CPM) and applied a log transformation with a

pseudocount of 1. Principal component analysis (PCA) is then applied to the normalized data, and

batch effects are corrected using Harmony (v1.2.3) on the top 30 principal components (PCs)

using default parameters except for theta, which was set to 8, to promote further mixing with

clusters across technologies. Finally, Uniform Manifold Approximation and Projection (UMAP) is

performed on the harmonized PCs, generating a shared 2D embedding across the two

technologies, and the data is further facetted by technology for visualization (Supplementary

Figure 5).

After generating a shared embedding, we quantified differences in gene expression patterns

between scRNA-seq and Xenium. We first computed Leiden clusters on the harmonized principal

components (resolution = 1.0) to identify transcriptionally similar groups of cells shared across

both technologies. For each Leiden cluster, we calculated the mean expression of every gene

present in both datasets. Clusters containing fewer than ten cells from either modality were

excluded to ensure robust gene-level estimates. To compare expression patterns between

scRNA-seq and Xenium for a given gene, we constructed vectors of cluster-level mean expression

from both technologies and evaluated their similarity using two metrics: root-mean-square error

(RMSE), computed relative to the line y = x, and Pearson correlation (r). To investigate whether

predicted off-target genes may contribute to the observed target-gene expression in Xenium, we

compared the Xenium target gene to the aggregated scRNA-seq expression of itself and its

predicted off-targets. We summed the raw counts of a gene and all of its predicted off-targets in

the scRNA-seq dataset, re-normalized the aggregated counts using CPM followed by log(x + 1),

and again calculated the RMSE and Pearson correlation. This allowed us to test whether the

Xenium gene expression more closely reflected the intended gene alone or the combined

expression of the intended gene and its predicted off-targets.

<u>Obtaining custom probe panels from HuBMAP</u>

To evaluate potential off-target binding in the HuBMAP placenta and multi-tissue (heart, kidney,

and lung) custom probe panels, we first downloaded the corresponding BED files from the

HuBMAP   portal

(https://portal.hubmapconsortium.org/browse/dataset/28fe8e4ac8a4193f82fdd9f4d4eb0bb2;

https://portal.hubmapconsortium.org/browse/dataset/6f597ca43db80f2499443f5c5bfac97c).

Using pyfaidx and pandas in Python, we extracted each probe's target gene name, gene identifier,

and genomic coordinates from the BED files, and then generated FASTA files by retrieving the

corresponding sequences from the reference genome (GRCh38). These FASTA files were then

used as input to OPT to predict potential off-target binding.

<u>HuBMAP custom probe-panel evaluation</u>

To assess whether predicted off-target genes were likely to confound Xenium results in the

HuBMAP custom probe panels, we evaluated the expression of the predicted off-target genes in

matched HuBMAP RNA-seq datasets corresponding to the tissues for which each panel was

designed. Four RNA-seq datasets were downloaded from the HuBMAP portal: a bulk RNA-seq

dataset   for   the   placenta

(https://portal.hubmapconsortium.org/browse/dataset/cad68e5e910ca90bda79cc0d6aa586d1

)   and   scRNA-seq   datasets   for   the   heart

(https://portal.hubmapconsortium.org/browse/dataset/e62b7f592efe97894de86cfabbfddde8),

kidney

(https://portal.hubmapconsortium.org/browse/dataset/43f83f08bedac2bc4d58767362dfebe2)

,   and   lung

(https://portal.hubmapconsortium.org/browse/dataset/6c2b7f1b8bc7e4fb6a1cadf29b09b83a).

For each tissue, raw count matrices were normalized to counts per million (CPM) and averaged

across all cells to obtain a bulk-like mean expression profile. Log1p-transformed mean CPM

values were used for all downstream comparisons. OPT was run on the two HuBMAP custom

probe panels with all RNA species included and a pad length of 10. Since OPT reports transcript-

level identifiers, we removed transcript-specific suffixes from Ensembl IDs to align them with

gene symbols present in the RNA-seq datasets. For every target gene in the custom panels, we

then compiled its predicted off-target genes based on OPT results and evaluated whether these

off-targets were expressed in the matched tissue's RNA-seq profile. To visualize these results, we

generated heatmaps in which rows correspond to intended target genes and columns represent

their predicted off-targets ordered by decreasing expression. This enabled direct comparison of

the magnitude and tissue specificity of potential off-target expression across the HuBMAP

datasets.

Cross-annotation analysis

To compare OPT's results with different reference annotations, we used the most recent releases

of GENCODE basic (v47), GENCODE comprehensive (v47), RefSeq (v110), and CHESS (v3.1.3)

annotation of the GRCh38 genome. Note that GENCODE "basic" is the more reliable version of

the annotation and is much closer to RefSeq and CHESS. GENCODE "comprehensive" includes

hundreds of thousands of low-quality annotations, which we included in some of our analyses

for completeness. Note also that GRCh38 has many non-reference sequences called "alternative

scaffolds"; we removed these for our analysis. We then used gffread to extract transcripts as

defined in these annotations by running:

`$ gffread -w transcripts.fa -g grch38.p12/14.fa annotation.gff`

The GRCh38.p14 assembly was used during transcript sequence extraction for all reference

annotations, except for CHESS which specifies that the annotation maps genes and transcripts

onto the GRCh38.p12 assembly. For RefSeq, we renamed the VD(J) segment features as

transcript features to ensure consistency, and we also removed transcript sequences with the

gene_biotype "pseudogene." RefSeq has a separate biotype called "transcribed_pseudogene,"

but doesn't annotate transcripts for these features. We considered transcripts annotated for a

small subset of just pseudogenes an error in the annotation.

Conflicting interests

The authors declare no conflicting interests.

Acknowledgements

We thank Reza Kalhor for his input on the project and feedback on the manuscript. We thank Ian

Fiddes for sharing the Xenium gene panel from the Janesick et al. publication and explaining its

relation to the commercially available pre-designed Xenium v1 Human Breast Gene Expression

Panel. We thank Sergii Domanskyi, Scott Lindsay-Hewett, and Chenchen Zhu for sharing the

HuBMAP custom gene panels. Research reported in this publication was supported by the

National Institute of General Medical Sciences of the National Institutes of Health under Awards

R35-GM142889 and R35-GM130151, the HuBMAP Integration, Visualization, and Engagement

(HIVE) Initiative under Award Number OT2-OD033760, and the National Science Foundation

under Grant No. 2047611.

References

10x Genomics. 2025a. "10x Genomics Product Catalog." March 19. www.10xgenomics.com/store/product-catalog?query=xenium.

10x Genomics. 2025b. "TXG_Q4-and-Full-Year-2024-Earnings-Press-Release." 10x Genomics Reports Fourth Quarter and Full Year 2024 Financial Results and Provides Outlook for 2025, March 19. https://s28.q4cdn.com/592666581/files/doc_financials/2024/q4/TXG_Q4-and-Full-Year-2024-Earnings-Press-Release.pdf.

Chou, Hui-Hsien. 2010. "Shared Probe Design and Existing Microarray Reanalysis Using PICKY." *BMC Bioinformatics* 11 (1): 196. https://doi.org/10.1186/1471-2105-11-196.

Clifton, Kalen, Manjari Anant, Gohta Aihara, et al. 2023. "STalign: Alignment of Spatial Transcriptomics Data Using Diffeomorphic Metric Mapping." *Nature Communications* 14 (1): 8123. https://doi.org/10.1038/s41467-023-43915-7.

Clifton, Kalen, Vivien Jiang, Rafael dos Santos Peixoto, et al. 2025. "STcompare: Comparative Spatial Transcriptomics Data Analysis of Structurally Matched Tissues to Characterize Differentially Spatially Patterned Genes." Preprint, bioRxiv, November 25. https://doi.org/10.1101/2025.11.21.689847.

Fornace, Mark E., Jining Huang, Cody T. Newman, Nicholas J. Porubsky, Marshall B. Pierce, and Niles A. Pierce. 2022. "NUPACK: Analysis and Design of Nucleic Acid Structures, Devices, and Systems." Preprint, Chemistry, November 11. https://doi.org/10.26434/chemrxiv-2022-xv98l.

Harbig, J. 2005. "A Sequence-Based Identification of the Genes Detected by Probesets on the Affymetrix U133 plus 2.0 Array." *Nucleic Acids Research* 33 (3): e31–e31. https://doi.org/10.1093/nar/gni027.

Hershberg, Elliot A., Conor K. Camplisson, Jennie L. Close, et al. 2021. "Author Correction: PaintSHOP Enables the Interactive Design of Transcriptome- and Genome-Scale Oligonucleotide FISH Experiments." *Nature Methods* 18 (10): 1265–1265. https://doi.org/10.1038/s41592-021-01273-6.

Hu, Mengwei, Bing Yang, Yubao Cheng, et al. 2020. "ProbeDealer Is a Convenient Tool for Designing Probes for Highly Multiplexed Fluorescence in Situ Hybridization." *Scientific Reports* 10 (1): 22031. https://doi.org/10.1038/s41598-020-76439-x.

Jain, Sanjay, Liming Pei, Jeffrey M. Spraggins, et al. 2023. "Advances and Prospects for the Human BioMolecular Atlas Program (HuBMAP)." *Nature Cell Biology* 25 (8): 1089–100. https://doi.org/10.1038/s41556-023-01194-w.

Janesick, Amanda, Robert Shelansky, Andrew D. Gottscho, et al. 2023. "High Resolution Mapping of the Tumor Microenvironment Using Integrated Single-Cell, Spatial and in Situ Analysis." *Nature Communications* 14 (1): 8353. https://doi.org/10.1038/s41467-023-43458-x.

Korsunsky, Ilya, Nghia Millard, Jean Fan, et al. 2019. "Fast, Sensitive and Accurate Integration of Single-Cell Data with Harmony." *Nature Methods* 16 (12): 1289–96. https://doi.org/10.1038/s41592-019-0619-0.

Kuemmerle, Louis B., Malte D. Luecken, Alexandra B. Firsova, et al. 2024. "Probe Set Selection
for Targeted Spatial Transcriptomics." *Nature Methods* 21 (12): 2260–70.
https://doi.org/10.1038/s41592-024-02496-z.
Li, Qiyuan, Nicolai J Birkbak, Balazs Gyorffy, Zoltan Szallasi, and Aron C Eklund. 2011. "Jetset:
Selecting the Optimal Microarray Probe Set to Represent a Gene." *BMC Bioinformatics*
12 (1): 474. https://doi.org/10.1186/1471-2105-12-474.
Liu, Hongfang, Ionut Bebu, and Xin Li. 2014. "Microarray Probes and Probe Sets." *Front Biosci* 2
(1): 325–38. https://doi.org/10.2741/e93.
Marçais, Guillaume, Arthur L. Delcher, Adam M. Phillippy, Rachel Coston, Steven L. Salzberg,
and Aleksey Zimin. 2018. "MUMmer4: A Fast and Versatile Genome Alignment System."
PLOS Computational Biology 14 (1): e1005944.
https://doi.org/10.1371/journal.pcbi.1005944.
Marçais, Guillaume, and Carl Kingsford. 2011. "A Fast, Lock-Free Approach for Efficient Parallel
Counting of Occurrences of k-Mers." *Bioinformatics* 27 (6): 764–70.
https://doi.org/10.1093/bioinformatics/btr011.
McCartney, Daniel L., Rosie M. Walker, Stewart W. Morris, Andrew M. McIntosh, David J.
Porteous, and Kathryn L. Evans. 2016. "Identification of Polymorphic and Off-Target
Probe Binding Sites on the Illumina Infinium MethylationEPIC BeadChip." *Genomics Data*
9 (September): 22–24. https://doi.org/10.1016/j.gdata.2016.05.012.
Mecham, Brigham H., Daniel Z. Wetmore, Zoltan Szallasi, Yoel Sadovsky, Isaac Kohane, and
Thomas J. Mariani. 2004. "Increased Measurement Accuracy for Sequence-Verified
Microarray Probes." *Physiological Genomics* 18 (3): 308–15.
https://doi.org/10.1152/physiolgenomics.00066.2004.
Mudge, Jonathan M, Sílvia Carbonell-Sala, Mark Diekhans, et al. 2025. "GENCODE 2025:
Reference Gene Annotation for Human and Mouse." *Nucleic Acids Research* 53 (D1):
D966–75. https://doi.org/10.1093/nar/gkae1078.
O'Leary, Nuala A., Mathew W. Wright, J. Rodney Brister, et al. 2016. "Reference Sequence
(RefSeq) Database at NCBI: Current Status, Taxonomic Expansion, and Functional
Annotation." *Nucleic Acids Research* 44 (D1): D733–45.
https://doi.org/10.1093/nar/gkv1189.
Pevny, Larysa, M. Celeste Simon, Elizabeth Robertson, et al. 1991. "Erythroid Differentiation in
Chimaeric Mice Blocked by a Targeted Mutation in the Gene for Transcription Factor
GATA-1." *Nature* 349 (6306): 257–60. https://doi.org/10.1038/349257a0.
Platt, Adam, Helen C. Ross, Steven Hankin, and Richard J. Reece. 2000. "The Insertion of Two
Amino Acids into a Transcriptional Inducer Converts It into a Galactokinase."
Proceedings of the National Academy of Sciences 97 (7): 3154–59.
https://doi.org/10.1073/pnas.97.7.3154.
Rouillard, J.-M., Micahel Zuker, and Erdogan Gulari. 2003. "OligoArray 2.0: Design of
Oligonucleotide Probes for DNA Microarrays Using a Thermodynamic Approach."
Nucleic Acids Research 31 (12): 3057–62. https://doi.org/10.1093/nar/gkg426.
Stenberg, Johan, Mats Nilsson, and Ulf Landegren. 2005. "ProbeMaker: An Extensible
Framework for Design of Sets of Oligonucleotide Probes." *BMC Bioinformatics* 6 (1): 229.
https://doi.org/10.1186/1471-2105-6-229.

Tsai, Fong-Ying, Gordon Keller, Frank C. Kuo, et al. 1994. "An Early Haematopoietic Defect in
Mice Lacking the Transcription Factor GATA-2." *Nature* 371 (6494): 221–26.
https://doi.org/10.1038/371221a0.
Varabyou, Ales, Markus J. Sommer, Beril Erdogdu, et al. 2023. "CHESS 3: An Improved,
Comprehensive Catalog of Human Genes and Transcripts Based on Large-Scale
Expression Data, Phylogenetic Analysis, and Protein Structure." *Genome Biology* 24 (1):
249. https://doi.org/10.1186/s13059-023-03088-4.
Wang, Xiaowei, and Brian Seed. 2003. "Selection of Oligonucleotide Probes for Protein Coding
Sequences." *Bioinformatics* 19 (7): 796–802.
https://doi.org/10.1093/bioinformatics/btg086.
Wernersson, R., and H. B. Nielsen. 2005. "OligoWiz 2.0--Integrating Sequence Feature
Annotation into the Design of Microarray Probes." *Nucleic Acids Research* 33 (Web
Server): W611–15. https://doi.org/10.1093/nar/gki399.
Yahoo Finance. 2025. "10x Genomics, Inc. (TXG)." 10x Genomics, Inc. (TXG), November 20.
https://finance.yahoo.com/quote/TXG/.

809
810

Red: RefSeq   Blue: CHESS   Green: GENCODE

Supplementary Figure 1. Screenshot from the Integrated Genome Viewer (IGV) showing the 40bp probe sequence (ID: ENSG00000196154|S100A4|ab4e3dc) that matches both *S100A5* and *S100A4*. Shown are 6 isoforms from the CHESS v3.1 annotation, 4 from GENCODE basic v47, and 3 from RefSeq v110 for *S100A4*, as well as 2 RefSeq isoforms for the neighboring *S100A5*. The probe sequence aligns to the overlapping region between *S100A5* and *S100A4* gene loci. Matching probe shown in a zoomed-in view below. The forward- and reverse-strand sequences of the probe are shown, and the highlighted area indicates approximately where the probe falls within the gene.

Supplementary Figure 2. UpSet plot illustrating the overlap of protein-coding genes across three genome annotations: GENCODE basic, RefSeq, and CHESS.

Supplementary Figure 3. (A) Overlap regions between Visium (orange outline) and Xenium (blue outline) data, shown on the Xenium histological image and the Visium histological image, respectively. **(B)** Log transformed aggregated total gene counts for spots ($\sim$55μm x 55μm) in both Xenium and Visium datasets, overlaid on their corresponding histological image.

Red: RefSeq   Blue: CHESS   Green: GENCODE

Supplementary Figure 4. Screenshots from the Integrated Genome Viewer (IGV) showing alignments of **(A)** four 40bp probes targeting *APOBEC3B* (ENSG00000179750|APOBEC3B|17d76bb, ENSG00000179750|APOBEC3B|e03f8ab, ENSG00000179750|APOBEC3B|5991db9, and ENSG00000179750|APOBEC3B|59c9349). **(B)** All four probes align to their intended target gene *APOBEC3B* while two of the four probes align to each off-target gene: **(C)** *APOBEC3D* and **(D)** *APOBEC3F*. The forward- and reverse-strand sequences of the probe are shown, and the highlighted areas indicate approximately where the probe falls within the gene. Panels **(B-D)** share a common legend.

Supplementary Figure 5. UMAP visualization of integrated scRNA-seq and Xenium datasets: **(A)** before harmony batch correction and **(B)** after harmony batch correction. **(C)** Leiden clustering results on the harmonized UMAP.

Supplementary Figure 6. (A) Schematic illustrating that hybridization may still occur even when there is a sequence mismatch at the non-ligated ends of the probe sequence. **(B)** Schematic depicting how probes could bind to each other instead of to their intended target.

Supplementary Figure 7. (A) Gene expression patterns for *ACTG2*: Xenium expression, Visium expression, the aggregated Visium expression combining *ACTG2* and its predicted off-target gene's expression ACTA1, ACTB, and *POTEM*, and Visium expression of *ACTG2*'s predicted off-targets ACTA1, ACTB, and *POTEM*. Two density plots are shown: one comparing Xenium vs. Visium for *ACTG2* alone, and one comparing Xenium vs. the aggregated Visium expression. The dotted line indicates the identity line (X = Y), and the solid line represents the line of best fit. **(B)** Gene expression patterns for *TUBB2B*: Xenium expression, Visium expression, the aggregated Visium expression combining *TUBB2B* and its predicted off-target gene's expression *TUBB2B* and *TUBB2A*, and Visium expression of *TUBB2B*'s predicted off-target *TUBB2A*. Two density plots are shown: one comparing Xenium vs. Visium for *TUBB2B* alone, and one comparing Xenium vs. the aggregated Visium expression. The dotted line indicates the identity line (X = Y), and the solid line represents the line of best fit.

Supplementary Figure 8. (A) Comparison of *ACTG2* expression patterns on harmonized UMAP: Xenium expression, scRNA-seq expression, an aggregated scRNA-seq profile combining *ACTG2* and its predicted off-target gene's expression *ACTB*, *POTEM*, *POTEE*, *POTEF*, *POTEI*, *POTEJ*, and *ACTA1*, and scRNA-seq expression of *ACTG2*'s potential off-targets. Two scatterplots are shown: one comparing Xenium vs. scRNA-seq for *ACTG2* cluster expression alone, and one comparing Xenium vs. the aggregated scRNA-seq cluster expression. The dotted line indicates the identity line (X = Y), and the solid line represents the line of best fit. **(B)** Comparison of *TUBB2B* expression patterns on harmonized UMAP: Xenium expression, scRNA-seq expression, an aggregated scRNA-seq profile combining *TUBB2B* and its predicted off-target gene's expression *TUBB2A*, and scRNA-seq expression of *TUBB2B*'s potential off-target *TUBB2A*. Two scatterplots are shown: one comparing Xenium vs. scRNA-seq for *TUBB2B* cluster expression alone, and one comparing Xenium vs. the aggregated scRNA-seq cluster expression. The dotted line indicates the identity line (X = Y), and the solid line represents the line of best fit.

Supplementary Figure 9. Heatmap visualization of target genes and their predicted off-target genes of the HuBMAP placenta custom probe panel using a corresponding placenta bulk RNA-seq dataset. Gene expression values are counts-per-million (CPM) normalized, with a pseudocount added prior to log transformation. The first column shows the expression of each target gene in the placenta bulk RNA-seq dataset, while the remaining columns display the expression of the corresponding predicted off-target genes.

Supplementary Figure 10. Heatmap visualizations of target genes and their predicted off-target genes of the HuBMAP multi custom probe panel using a corresponding **(A)** kidney, **(B)** lung, and **(C)** heart scRNA-seq datasets. Gene expression values are counts-per-million (CPM) normalized, with a pseudocount added prior to log transformation. The first column shows the expression of each target gene in their respective scRNA-seq dataset, while the remaining columns display the expression of the corresponding predicted off-target genes.

Supplementary Figure 11. Heatmap visualizations of target genes and their predicted off-target genes for the Janesick et al. probes using **(A)** pseudo-bulked scRNA-seq and **(B)** Visium data. Gene expression values are counts-per-million (CPM) normalized, with a pseudocount added prior to log transformation. The first column shows the expression of each target gene in their respective scRNA-seq dataset, while the remaining columns display the expression of the corresponding predicted off-target genes.

Supplementary Figure 12. (A) Gene expression patterns for *ADH1B*: Xenium expression, Visium expression, the aggregated Visium expression combining *ADH1B* and its predicted off-target gene's expression *ADH1A* and *ADH1C*, and Visium expression of *ADH1B*'s predicted off-targets *ADH1A* and *ADH1C*. Two density plots are shown: one comparing Xenium vs. Visium for *ADH1B* alone, and one comparing Xenium vs. the aggregated Visium expression. The dotted line indicates the identity line (X = Y), and the solid line represents the line of best fit. **(B)** Comparison of *ADH1B* expression patterns on harmonized UMAP: Xenium expression, scRNA-seq expression, an aggregated scRNA-seq profile combining *ADH1B* and its predicted off-target gene's expression *ADH1A* and *ADH1C*, and scRNA-seq expression of *ADH1B*'s potential off-targets *ADH1A* and *ADH1C*. Two scatterplots are shown: one comparing Xenium vs. scRNA-seq for *ADH1B* cluster expression alone, and one comparing Xenium vs. the aggregated scRNA-seq cluster expression. The dotted line indicates the identity line (X = Y), and the solid line represents the line of best fit.

Supplementary Figure 13. (A) Spatial gene expression of *HDC* overlaid on the corresponding histological images for Xenium and Visium. **(B)** Harmonized UMAP visualization of *HDC* expression for Xenium and scRNA-seq data.

probe ID: ENSG00000125878|TCF15|5d3cbc2

Red: GENCODE v47   Black: GENCODE v28

probe ID: ENSG00000169083|AR|a0c6719

Supplementary Figure 14. Screenshots from the Integrated Genome Viewer (IGV) illustrating annotation-dependent differences in probe alignment. **(A)** A 40bp probe (ID: ENSG00000125878|TCF15|5d3cbc2) aligns to an exonic region in GENCODE v28 but to an upstream region in GENCODE v47. **(B)** A 40bp probe (ID: ENSG00000169083|AR|a0c6719) aligns to an exonic region in GENCODE v28 but to an intronic region in GENCODE v47. Matching probe shown in a zoomed-in view below. The forward- and reverse-strand sequences of the probe are shown, and the highlighted areas indicate approximately where the probe falls within the gene. Panels **(A-B)** share a common legend.

872   **Supplementary Tables**

FASTA FILE

Supplementary Table 1. Fasta file of probe sequences from 10x Genomics corresponding to the 10x Genomics Xenium v1 Human Breast Gene Expression Panel used in Janesick et al.

Target gene	Number of probes	Predicted binding genes	Number of probes aligned	Gene types
ADH1B	8	ADH1B,ADH1A,ADH1C	8,2,1	PC,PC,PC
AKR1C1	9	AKR1C1,AKR1C2,AKR1C3,AKR1C4	8,1,1,1	PC,PC,PC,PC
APOBEC3A	10	APOBEC3A,APOBEC3B	7,2	PC,PC
APOBEC3B	8	APOBEC3B,APOBEC3D,APOBEC3F	8,2,2	PC,PC,PC
C15orf48	6	C15orf48,MIR147B	6,1	PC, precursor_RNA
CD8B	16	CD8B,CD8B2	15,2	PC,PC
IGF1	13	IGF1,LINC02456	11,1	PC,ncRNA
POLR2J3	12	SPDYE2,SPDYE18,SPDYE5,SPDYE21, SPDYE7P,SPDYE2B,SPDYE16,POLR2J, POLR2J3-UPK3BL2,POLR2J2, POLR2J2-UPK3BL1, POLR2J3,POLR2J4	2,1,1,1,1, 1,3,3,2,2,3,2	PC,PC,PC,PC , misc_RNA, PC,PC,PC,ncRNA,PC, ncRNA,PC,ncRNA
S100A4	5	S100A4,S100A5	3,1	PC,PC
SERHL2	8	SERHL2,SERHL	8,7	PC,misc_RNA
SLAMF1	11	SLAMF1,LOC107985220	10,1	PC,ncRNA
TOMM7	10	TOMM7,LOC107986346	3,2	PC,PC
TPD52	8	TPD52,TPD52-MRPS28	8,5	PC,PC
TPSAB1	2	TPSAB1,TPSB2,TPSD1	2,2,1	PC,PC,PC

911
Supplementary Table 2. OPT output of genes with predicted off-target binding based on perfect
sequence homology in RefSeq. This table shows the 14 genes whose probes in the 10x Genomics
Xenium v1 Human Breast Gene Expression Panel that exhibit predicted off-target probe binding, where
each off-target alignment involves a perfect 40bp match to the probe sequence. The final column shows
the gene types, in order, of each of the off-target genes shown in column 3. Off-target alignments
between *CCPG1* probes and *DNAAF1-CCPG1* were excluded. Abbreviations: PC = protein-coding; PG =
pseudogene; precursor_RNA = precursor RNA; misc_RNA = miscellaneous RNA; ncRNA = non-coding
RNA.

Target gene	Number of probes	Predicted binding genes	Number of probes aligned	Gene types
ADH1B	8	ADH1B,ADH1A,ADH1C	8,2,1	PC,PC,PC
AKR1C1	9	AKR1C1,AKR1C2,AKR1C3, AKR1C4	8,1,1,1	PC,PC,PC,PC
APOBEC3A	10	APOBEC3A,APOBEC3B	8,2	PC,PC
APOBEC3B	8	APOBEC3B,APOBEC3D,APOBEC3F	8,2,2	PC,PC,PC
C15orf48	6	C15orf48,MIR147B	6,1	PC,miRNA
CD8B	16	CD8B,CD8B2	16,2	PC,PC
EPCAM	8	EPCAM,LOC132386	8,1	PC,PG
HMGA1	7	HMGA1,HMGA1P1,HMGA1P2, HMGA1P3,HMGA1P6	7,1,3,2,2	PC,PG,PG,PG, PG
KRT14	6	KRT14,KRT16P6	6,1	PC,PG
KRT8	16	KRT8,KRT8P33,KRT8P32,KRT8P22, KRT8P3,KRT8P42,KRT8P45,KRT8P17, KRT8P11,KRT8P2,KRT8P36,KRT8P37, KRT8P30	8,2,1,1,3, 1,2,1,1,2, 1,1,1	PC,PG,PG,PG, PG,PG,PG,PG, PG,PG,PG,PG, PG
MYLK	11	MYLK,MYLKP1	11,1	PC,PG
MYO5B	8	MYO5B,MYO5BP1,MYO5BP2, LOC102724787	8,1,1,1	PC,PG,PG,PG
POLR2J3	12	SPDYE2,SPDYE7P,SPDYE21,SPDYE16, SPDYE5,SPDYE18,SPDYE2B,POLR2J, POLR2J3,POLR2J2-UPK3BL1,POLR2J2, POLR2J3-UPK3BL2,POLR2J4	1,1,1,1, 1,1,1,3, 5,2,3,4,3, 2	PC,PG,PC,PC, PC,PC,PC,PC, PC,PC,lncRNA, lncRNA,PG
SCD	8	SCD,SCDP1	8,2	PC,PG
SERHL2	8	SERHL2,SERHL	8,7	PC,PG
SMS	8	SMS,LOC646347,LOC100419037, LOC642590	8,3,1,1	PC,PG,PG,PG
TOMM7	10	TOMM7,LOC107986346	3,2	PC,PC
TPD52	8	TPD52,TPD52-MRPS28	8,5	PC,PC
TPSAB1	2	TPSAB1,TPSB2,TPSD1	2,2,1	PC,PC,PC
TRAF4	9	TRAF4,LOC100420513	9,1	PC,PG
TUBB2B	8	TUBB2B,TUBB2BP1	8,1	PC,PG
VOPP1	11	VOPP1,LOC729086	10,8	PC,PG
VWF	11	VWF,VWFP1	8,1	PC,PG

Supplementary Table 3. OPT output of genes with predicted off-target binding based on perfect
sequence homology in CHESS. This table shows the 23 genes whose probes in the 10x Genomics Xenium
v1 Human Breast Gene Expression Panel that exhibit predicted off-target probe binding, where each off-
target alignment involves a perfect 40bp match to the probe sequence. The final column shows the gene
types, in order, of each of the off-target genes shown in column 3. Off-target alignments between
CCPG1 probes and *DNAAF1-CCPG1* were excluded. Abbreviations: PC = protein-coding; PG =
pseudogene; miRNA = microRNA.

Perfect match		GENCODE Basic	GENCODE Comprehensive	RefSeq	CHESS
Number of transcripts		158,339	387,944	181,627	158,363
All genes	Off-target probes	128	126	44	73
	Affected genes	37	38	14	23
Pseudogenes removed	Off-target probes	101	107	44	35
	Affected genes	29	29	15	11
Protein-coding genes only	Off-target probes	26	28	26	26
	Affected genes	11	11	10	9

944
945 **Supplementary Table 4.** The number of off-target probes and affected genes (from the set of 313 genes
946 in the Xenium panel) found when looking for perfect matches between probe sequences and transcripts
947 in four different reference annotations: GENCODE basic, GENCODE comprehensive, RefSeq, and CHESS.
948 Off-target alignments between *CCPG1* probes and *DNAAF1-CCPG1* were excluded.
949

Gene	GENCODE	RefSeq	CHESS
ADH1B	✓	✓	✓
AKR1C1	✓	✓	✓
APOBEC3A	✓	✓	✓
APOBEC3B	✓	✓	✓
CD8B	✓	✓	✓
POLR2J3	✓	✓	✓
TPSAB1	✓	✓	✓
AQP1	✓	x	x
C1QA	✓	x	x
CD79B	✓	x	x
CEACAM6	✓	x	x
S100A4	x	✓	x
TOMM7	x	✓	✓
TPD52	x	✓	✓

Supplementary Table 5. Union set of protein-coding genes that OPT predicts to be affected by off-target binding, across three different reference annotations: GENCODE basic, RefSeq, and CHESS.

Target gene	Number of probes	Predicted binding genes	Number of probes aligned	Gene types – GENCODE (v47)
ACTG2	14	*ACTG2,POTEE,ACTB,POTEJ,ACTBP16, ACTA1,POTEI,POTEM,ACTG1P14, ACTG1P12,POTEF,POTEKP,ACTBP15, ENSG00000241838,ACTG1P15, ENSG00000230080,ACTBP1,ACTG1P19*	14,1,1,1, 1,1,1, 1,1, 1,1,1, 1,1, 1,1,1,1	PC,PC,PC,PC,PG,PC, PC,PC,PG,PG, PC,PG,PG,PG,PG, PG,PG,PG
APOC1	2	*APOC1,APOC1P1*	2,1	PC, PG
CD83	8	*CD83,CD83P1*	8,2	PC,PG
CEACAM8	13	*CEACAM8,CEACAMP5,LIPE-AS1, CEACAMP8*	8,1,1,1	PC,PG,lncRNA,PG
ENAH	8	*ENAH,ENAHP1*	8,3	PC,PG
FBLIM1	9	*FBLIM1,FBLIM1P1*	9,1	PC,PG
FCGR3A	8	*FCGR3A,FCGR3B,ENSG00000289768*	8.4.1	PC,PC,PC
FLNB	14	*FLNB, ENSG00000263343*	14,1	PC,PG
KLRC1	8	*KLRC1,KLRC2,KLRC3,KLRC4, ENSG00000255641,KLRC4-KLRK1*	8,1,1,1,1	PC,PC,PC,PC,PC, PC
KRT16	5	*KRT16,ENSG00000290978 ,KRT16P3, KRT16P2,ENSG00000290657*	5,2,2,1,1	PC,lncRNA, lncRNA,PG, lncRNA
KRT6B	8	*KRT6B,KRT6C*	8,1	PC,PC
KRT7	5	*KRT7,KRT75,KRT8,KRT8P26,KRT8P20,KRT8P13,KRT8P7,KRT8P12,KRT8P3,KRT5, KRT8P21,KRT8P51,KRT79,KRT8P14 ,KRT8P33,KRT8P35,KRT8P17, KRT6C,KRT8P18,KRT8P10,KRT8P11, KRT8P50,KRT8P25,KRT8P32,KRT6B, KRT8P36,KRT126P,ENSG00000293192, KRT8P46,KRT8P39,KRT8P49*	5,1,1,1,1,1, 1, 1,1,1,1,1,1, 1, 1,1,1,1,1,1, 1, 1,1,1,1,1,1, 1,1,1,1	PC,PC,PC,PG,PG, PG,PG,PG,PG,PC, PG,PG,PC,PG,PG, PG,PG,PC,PG,PG, PG,PG,PG,PG,PC, PG,PG,lncRNA, PG,PG,PG
OCIAD2	8	*OCIAD2,ENSG00000253760, ENSG00000294977*	8,1,1	PC,PG,lncRNA
REXO4	8	*REXO4,ENSG00000259696*	8,1	PC,PG
SCGB2A1	6	*SCGB2A1,SCGB2A2*	6,1	PC,PC
SVIL	8	*SVIL,SVIL2P,ENSG00000290952*	8,1,1	PC,PG,lncRNA
ZEB2	21	*ZEB2,ENSG00000254198*	21,1	PC,PG

ZNF562	10	ZNF562,ENSG00000291231,ZNF561, ZNF812P,ENSG00000226810, ENSG00000267490	10,1, 1,1, 1,1	PC,lncRNA,PC,PG, PG,PG

Supplementary Table 6. 18 additional genes that were identified to exhibit potential off-target probe binding when allowing for a 10bp error margin on either side of the binding site when using GENCODE v47. Abbreviations: PC = protein-coding; PG = pseudogene; lncRNA = long non-coding RNA.

Target gene	Number of probes	Predicted binding genes	Number of additional probes aligned	Gene types – GENCODE (v47)
AKR1C1	9	AKR1C6P	1	PG
APOBEC3A	10	APOBEC3G	1	PC
CEACAM6	8	CEACAM1,CEACAM8	1,1	PC,PC
HMGA1	7	HMGA1P6	2	PG
KRT14	6	ENSG00000290978,KRT16,KRT16P2	1,1,1	lncRNA,PC,PG
KRT8	16	KRT8P8,KRT8P9,KRT8P44,KRT8P31, KRT8P46,KRT8P48,KRT8P6, ENSG00000235286,KRT8P43,KRT8P47, KRT8P13,KRT8P14,KRT8P50,KRT8P5, KRT8P35,KRT8P7,KRT8P15,KRT8P1, KRT8P19,GFAP,KRT80,KRT8P41, KRT8P34,KRT90P,KRT8P4,KRT8P49, KRT8P40,ENSG00000293192,KRT8P12, KRT8P21	1,1,1,1,1, 1,2,1,1,1, 1,1,1,1,1,2, 1,2,1,2,2,2, 1,1,2,2,2,2, 1,1,1	PG,PG,PG,PG, PG,PG,PG,PG, PG,PG,PG,PG, PG,PG,PG,PG, PG,PG,PG,PC, PC,PG,PG,PG, PG,PG,PG, lncRNA,PG,PG
LDHB	8	LDHBP1,LDHBP2	2,4	PG,PG
MYO5B	8	MYO5BP3	2	PG
SMS	8	ENSG00000251342	2	PG
TUBB2B	8	TUBB2A	1	PC

1009
Supplementary Table 7. 10 genes that were previously predicted to be affected by off-target binding
based on perfect matching now show additional predicted off-target interactions when a 10bp error
margin is allowed on either side of the binding site when using GENCODE v47. This effect is observed
either through the accumulation of new probes with predicted off-target binding or via the identification
of additional predicted off-target genes per probe. Abbreviations: PC = protein-coding; PG =
pseudogene; lncRNA = long non-coding RNA.

Target gene	Number of probes	Predicted binding genes	Number of additional probes aligned	Gene types – GENCODE (v47)
ANG	8	ANG,RNASE4,ENSG00000259171	8,1,1	PC,PC,PC
AOC3	8	AOC3,AOC4P	8,3	PC,PG
CCL3	6	CCL3, CCL3L3	6,2	PC,PG
CCNB2	8	CCNB2,CCNB2P1	7,1	PC,PG
CD209	11	CD209,ENSG00000288669,CLEC4M	10,2,2	PC,PC,PC
CD24	12	CD24,CD24P4	12,2	PC,PG
CD68	9	CD68,ENSG00000264772	9,8	PC,lncRNA
CLEC4M	9	CLEC4M,CD209	9,2	PC,PC
CSH1	5	CSH1,CSHL1,CSH2	5,2,5	PC,PC,PC
CSH2	8	CSH2,CSH1,GH2,GH1,ENSG00000285947,CSHL1	8,8,2,2,2,3	PC,PC,PC,PC,PC,PC
CSHL1	9	CSHL1,CSH1,CSH2,GH2,ENSG00000285947,GH1	9,3,3,2,1,2	PC,PC,PC,PC,PC,PC
CSPG4	8	CSPG4,ENSG00000290664,CSPG4P13,CSPG4P5,CSPG4P10,CSPG4P11,ENSG00000290947,ENSG00000290690,CSPG4P12,ENSG00000291211,CSPG4P4Y,CSPG4P3Y	8,1,5,2,2,2,1,2,2,1,1,1	PC,lncRNA,PG,PG,PG,PG,lncRNA,lncRNA,PG,lncRNA,PG,PG
CYP26A1	8	CYP26A1,ENSG00000285846	5,5	PC,lncRNA
EPCAM	8	EPCAM,ENSG00000225356	5,1	PC,PG
ERVFRD-1	8	ERVFRD-1,ENSG00000258619	8,1	PC,PG
ERVV-1	8	ERVV-1,ERVV-2	8,5	PC,PC
ERVV-2	8	ERVV-2,ERVV-1	8,1	PC,PC
ERVW-1	8	ERVW-1,RBM41	8,4	PC,PC
FCGR2B	8	FCGR2B,FCGR2A,ENSG00000289273,FCGR2C	7,2,1,3	PC,PC,PG,PC
FCGR3A	8	FCGR3A,FCGR3B,ENSG00000289768	8,4,1	PC,PC,PG
FLT1	11	FLT1,FLT1P1	10,2	PC,PG
HLA-A	8	HLA-H,HLA-A,HLA-L,HLA-B,HLA-J,HLA-C,ENSG00000290574,HLA-G,HLA-W,HLA-E	7,3,1,1,1,1,1,1,1,1	PC,PG,PG,PC,PG,PC,lncRNA,PC,PG,PC
HLA-C	8	HLA-C,HLA-B,HLA-L,HLA-J,HLA-H,HLA-A,ENSG00000290574	8,2,1,1,1,1,1	PC,PC,PG,PG,PG,PC,lncRNA
HLA-G	8	HLA-G,HLA-E	7,1	PC,PC
IFI27	5	IFI27,IFI27L1,IFI27L2	5,1,1	PC,PC,PC
IL6ST	10	IL6ST,IL6STP1,ENSG00000305843	10,2,1	PC,PG,lncRNA
IRF6	8	IRF6,ENSG00000289700	8,8	PC,PC

JAG2	8	JAG2,ENSG00000257622	8,3	PC,lncRNA
KIR2DL1	8	KIR2DL1,KIR2DP1,KIR3DL1,KIR2DL3	7,1,1,2	PC,PG,PC,PC
KLRC1	8	KLRC1,KLRC2,KLRC4,KLRC3,ENSG00000255641,KLRC4-KLRK1	8,1,1,1,1,1	PC,PC,PC,PC,PC,PC
KRT7	7	KRT7,KRT8P25,KRT8P14,KRT8P7,KRT6C,KRT8P50,KRT8,KRT8P39,KRT8P3,KRT75,KRT8P20,KRT8P18,KRT8P21,KRT126P,KRT8P12,KRT8P10,KRT8P32,KRT5,KRT8P33,ENSG00000293192,KRT8P13,KRT8P49,KRT8P26,KRT8P35,KRT6B,KRT8P51,KRT8P36,KRT79,KRT8P17,KRT8P11,KRT8P46,KRT8P1,KRT8P9,KRT8P31,KRT8P5,KRT8P41,KRT8P45,KRT8P23,KRT8P47,KRT8P43,KRT78,KRT6A,KRT3,KRT8P24,KRT8P22,KRT7-AS	3,1,2,1,1,1,2,1,2,2,2,2,1,1,2,1,2,1,2,2,1,2,1,2,1,1,2,2,2,1,1,1,1,1,1,1,1,1,1,1,1,1,1	
KRT8	10	KRT8,ENSG00000293192,KRT8P21,KRT8P12,KRT8P32,KRT8P48,KRT8P36,KRT8P3,ENSG00000235286,KRT8P19,KRT8P47,KRT8P1,KRT8P11,KRT8P31,KRT8P7,KRT8P2,KRT8P35,KRT8P13,KRT8P22,KRT8P15,KRT8P17,KRT8P43,KRT8P14,KRT8P42,KRT8P50,KRT8P6,KRT8P5,KRT8P9,KRT8P45	10,1,1,1,3,1,2,3,2,1,2,1,2,1,1,1,2,2,1,1,1,1,1,1,1,1,1,1,1	PC,lncRNA,PG,PG,PG,PG,PG,PG,PG,PG,PG,PG,PG,PG,PG,PG,PG,PG,PG,PG,PG,PG,PG,PG,PG,PG,PG,PG,PG
LAIR2	6	LAIR2,CDH23	5,1	PC,PC
LGALS9	8	LGALS9,ENSG00000266728,LGALS9DP,LGALS9B,LGALS9C	7,3,3,2,2	PC,PC,PG,PC,PC
NOS2	8	NOS2,NOS2P2,ENSG00000297633,NOS2P3,NOS2P1	8,1,1,1,1	PC,PG,lncRNA,PG,PG
PAGE4	6	PAGE4,ENSG00000235349,XAGE3,ENSG00000290520,ENSG00000169164	6,1,1,1,1	PC,PG,PC,lncRNA,PG
PCNA	8	PCNA,PCNAP3	8,2	PC,PG
PLAC8	11	PLAC8,ENSG00000256804	11,1	PC,PG
PPBP	8	PPBP,ENSG00000288796	8,4	PC,PC
PRG2	4	PRG2,ENSG00000254979	4,1	PC,PC
PSG1	8	PSG1,PSG10P,PSG5,PSG2,ENSG00000291235,PSG6,PSG7,PSG11,PSG4,PSG8,PSG3,PSG9	8,4,3,3,4,2,3,5,2,3,4,1	PC,PG,PC,PC,lncRNA,PC,PC,PC,PC,PC,PC,PC
PSG2	8	PSG2,PSG9,PSG5,PSG8,PSG4,PSG6,PSG3,PSG1,PSG7,PSG10P,PSG11,ENSG00000291235	8,3,4,5,2,2,4,4,3,2,3,1	PC,PC,PC,PC,PC,PC,PC,PC,PC,PG,PC,lncRNA
RPS4Y1	8	RPS4Y1,XGY2,ENSG00000213318	8,3,1	PC,PG,PG
SLC25A15	8	SLC25A15,SLC25A15P3,ENSG00000299646	7,1,1	PC,PG,lncRNA
SOX7	8	SOX7,PINX1	8,4	PC,PC
TEAD4	8	TEAD4,TEAD3	5,1	PC,PC

TPSAB1	8	*TPSAB1,TPSB2,TPSD1*	7,6,3	PC,PC,PC
VWF	8	*VWF,VWFP1*	8,1	PC,PG
XIST	15	*XIST,LNX2BP*	15,3	lncRNA,PG

1034
1035 **Supplementary Table 8**. OPT output for 49 genes with predicted off-target binding in the HuBMAP
1036 placenta custom probe panel, generated using GENCODE v47 and allowing a 10bp mismatch on either
1037 end of each probe. The final column lists the gene types, in order, corresponding to the off-target genes
1038 shown in column 3. Abbreviations: PC = protein-coding; PG = pseudogene; precursor_RNA = precursor
1039 RNA; misc_RNA = miscellaneous RNA; ncRNA = non-coding RNA.
1040

1041

1042

1043

1044

1045

1046

1047

1048

1049

1050

1051

1052

1053

1054

1055

1056

1057

Target gene	Number of probes	Predicted binding genes	Number of additional probes aligned	Gene types – GENCODE (v47)
ACSM2A	8	ACSM2A,ACSM2B	6,4	PC,PC
ACSM2B	8	ACSM2B,ACSM2A	8,4	PC,PC
ATRNL1	8	ATRNL1,ENSG00000285582	7,1	PC,lncRNA
CD68	7	CD68,ENSG00000264772	7,6	PC,lncRNA
CEACAM1	8	CEACAM1,CEACAM3,CEACAM8,CEACAM5	8,1,1,1	PC,PC,PC,PC
CFH	8	CFH,ENSG00000289697	7,7	PC,PC
CP	8	CP,CPP	8,3	PC,PG
CSPG4	8	CSPG4,CSPG4P11,ENSG00000290690,ENSG00000291211,CSPG4P12,CSPG4P13,CSPG4P10,CSPG4P5,ENSG00000290947,ENSG00000290664,CSPG4P4Y,CSPG4P3Y	8,2,2,1,2,5,2,2,1,1,1,1	PC,PG,lncRNA,lncRNA,PG,PG,PG,PG,lncRNA,lncRNA,PG,PG
CUBN	8	CUBN,CUBNP2	8,1	PC,PG
CYP2F1	8	CYP2F1,ENSG00000293114,CYP2F2P	6,2,3	PC,lncRNA,PG
EPCAM	8	EPCAM,ENSG00000225356	6,1	PC,PG
FCGR3A	8	FCGR3A,FCGR3B,ENSG00000289768	8,4,1	PC,PC,PC
FCGR3B	12	FCGR3B,FCGR3A,ENSG00000289768	10,7,4	PC,PC,PC
FLT1	17	FLT1,FLT1P1	16,4	PC,PG
HLA-DQA1	8	HLA-DQA1,HLA-DQA2	7,2	PC,PC
LDB3	13	LDB3,ENSG00000289258	11,7	PC,PC
PARP8	15	PARP8,LINC00869,LINC01138,LINC00623,ENSG00000290569,ENSG00000223804,LINC01145,ENSG00000274265,ENSG00000284964,ENSG00000285062	14,2,2,2,2,2,2,2,1,1	PC,lncRNA,lncRNA,lncRNA,lncRNA,PG,lncRNA,lncRNA,PG,PG
PLD5	8	PLD5,ENSG00000226113,PLD5P1	8,2,2	PC,PG,PC
REG1A	8	REG1A,ENSG00000290771,REG1CP,REG1B	7,2,2,1	PC,lncRNA,PG,PC
SERPINB3	8	SERPINB3,SERPINB4	8,3	PC,PC
SLAMF1	11	SLAMF1,ENSG00000228863	10,1	PC,lncRNA
SVIL	8	SVIL,ENSG00000290952,SVIL2P	8,1,1	PC,lncRNA,PG
TPSB2	1	TPSB2,TPSD1,TPSAB1	1,1,1	PC,PC,PC
VIM	8	VIM,ENSG00000234961	7,1	PC,lncRNA

1058

Supplementary Table 9. OPT output for 24 genes with predicted off-target binding in the HuBMAP multi
custom probe panel, generated using GENCODE v47 and allowing a 10bp mismatch on either end of each
probe. The final column lists the gene types, in order, corresponding to the off-target genes shown in
column 3. Abbreviations: PC = protein-coding; PG = pseudogene; precursor_RNA = precursor RNA;
misc_RNA = miscellaneous RNA; ncRNA = non-coding RNA.

ZIP FILE

Supplementary Table 10. OPT results for all publicly available 10x Genomics probe sets. Results include

all possible RNA species and using a pad length (-pl) of 10.

1067

1068

1069   **Supplementary Note: Differences between 3 Xenium breast gene panels and the impact on**

1070   **interpretation of results**

1071

In the previous iteration of this manuscript, we used the publicly available Xenium v1 Human

Breast Gene Expression Panel probe sequences from the 10x Genomics website (prior to April

10, 2025) and assumed it was used to generate the Xenium breast cancer data from Janesick et

al. based on the paper's methodological descriptions. 10x Genomics has since noted on their

public website as well as through private communications that this publicly available file

erroneously included extra probe sequences that are not used. Further, the Xenium breast cancer

data presented in Janesick et al. uses a human breast gene panel with probe sequences that

represent an earlier iteration of the commercially available Xenium v1 Human Breast Gene

Expression Panel. In particular, compared to the commercially available panel, the Janesick et al.

panel has 24 genes with one or more probes modified in the final panel, along with the probe

sequences targeting the 33 custom genes added by Janesick et al. This impacts our interpretation

of observed results to provide stronger evidence of putative imperfect sequence homology based

off-target probe binding.

1085

For clarity, we refer to the three probesets as:

- Probeset A: the previously analyzed publicly available Xenium v1 Human Breast Gene

Expression Panel probe sequences (prior to April 10, 2025)

- Probeset B: the Xenium probe sequences used in the Janesick et al. panel

- Probeset C: the currently commercial pre-designed Xenium v1 Human Breast Gene

Expression Panel (after April 10, 2025)

1092

To provide a specific example, in the previous iteration of this manuscript, OPT identified a probe

sequence targeting *TUBB2B* that exhibited perfect sequence homology with *TUBB2A*:

`>ENSG00000137285|TUBB2B|1dec8c0`

`GTTCATGATGCGGTCTGGGTACTCTTCCCGGATCTTGCTG`

Analysis with orthogonal spatial and single-cell transcriptome profiling technologies suggested

the *TUBB2B* gene expression pattern observed in the Xenium breast cancer data from Janesick

et al. represented an aggregation of both *TUBB2B* and *TUBB2A* consistent with off-target binding.

Given our assumption at the time that the Xenium breast cancer data had been generated using

Probeset A, we therefore believed this perfect homology probe was responsible for the observed

off-target signal.

1103

However, we now understand that the Xenium breast cancer data from Janesick et al. were

generated using Probeset B. Probeset B no longer contains this specific probe for *TUBB2B* with

perfect sequence homology with *TUBB2A*. However, the observed *TUBB2B* gene expression

pattern still represents an aggregation of both *TUBB2B* and *TUBB2A* based on analysis with

orthogonal spatial and single-cell transcriptome profiling technologies. Although Probeset B no

longer contains this specific probe for *TUBB2B* with perfect sequence homology with *TUBB2A*, it

contains one probe with imperfect sequence homology with *TUBB2A* (=30X1=6X3):

`>ENSG00000137285|TUBB2B|ed52e1c`

`TTGTCAATGCAGTAGGTTTCATCTGTGTTTTCCACCAGCT`

The observed comparative gene expression patterns therefore provide newfound strong support

for putative imperfect sequence homology off-target binding.

1115	

While we do not have Xenium breast cancer data generated using Probeset C, this *TUBB2B* probe

is also present in Probeset C:

`>TUBB2B (ENSG00000137285) | 3 | 5`

`TTGTCAATGCAGTAGGTTTCATCTGTGTTTTCCACCAGCT`

This suggests that off-target effects may still impact commercially available pre-designed panels

such as Probeset C (Supplementary Table 10).

1122