1　Archaeological Bolivian maize genomes suggest Inca cultural expansion augmented
2　maize diversity in South America
3

4　Huan Chen[1,2,4], Amy Baetsen-Young[3], Addie Thompson[2,3], Brad Day[1,2,3], Thelma
5　Madzima[4], Sally Wasef[5], Claudia Rivera Casanovas[6], William Lovis[7,*], Gabriel Wrobel[7,*]
6
[1]Department of Genetics and Genome Sciences, Michigan State University, East
Lansing, Michigan 48824, USA. [2]Graduate Program in Molecular Plant Sciences,
Michigan State University, East Lansing, Michigan 48824, USA. [3]Department of Plant,
Soil and Microbial Sciences, Michigan State University, East Lansing, MI 48824, USA.
[4]Department of Plant Biology, Michigan State University, East Lansing, Michigan 48824,
USA. [5]Genomics Research Centre, Queensland University of Technology, 60 Musk Ave,
Kelvin Grove, QLD, 4059, Australia. [6]Universidad Mayor de San Andres, Avenida Villazón
1995, Monoblock Central, La Paz, Bolivia. [7]Department of Anthropology, Michigan State
University, East Lansing, MI 48824, USA.
16
17　*corresponding authors: William Lovis, Gabriel Wrobel
18
Email: lovis@msu.edu and wrobelg@msu.edu

Author Contributions: HC, ABY, AT, and BD designed and conceived the experiments.
HC, ABY, AT, BD, TM, SW, CRC, WL, and GW analyzed the data and wrote the
manuscript. All authors commented and agreed on the manuscript before submission.

Competing Interest Statement: The authors declare no competing interests.

Classification: Biological Sciences and Social Sciences.

Keywords: archaeological maize, Inca culture, agriculture, domestication

Abstract

Previous archaeological and anthropological studies have demonstrated the myriad of
ways that cultural and political systems shape access to food and food preferences.
However, few studies have conducted a biocultural analysis linking specific
genotypic/phenotypic traits as evidence of cultural selection in ancient contexts. Here, we
provide insight into this topic through ancient genome data from Bolivia dating to ~500
BP, included as an offering with the mummified remains of a young girl, including 16
archaeological maize samples spanning at least 5,000 years of evolution, and 226
modern maize samples. Our phylogenetic analysis showed that the archaeological
Bolivian maize (aBM) has the closest genetic distance to the archaeological maize from
ancient Peru, which in turn shared the most similarities with archaeological Peruvian
maize. With the increased presence of the Inca state in the central Andes and its
interactions with local agricultural traditions, genetic diversity of maize increased. The
ovule development process was selected from modern maize and was compared to

archaeological maize; where it indicates the breeding programs aimed at enhancing seed
quality and yield in modern maize. Our study provides insights into the complex biocultural
role that Inca Empire expansion, including its economic, symbolic and religious cultural
practices, may have had in driving the expansion of maize diversity in South America.

Significance Statement

Maize has been improved through selective breeding delivering desirable horticultural,
agronomic, and cultural characteristics, related to environment, nutrition, etc. Inca culture
was closely associated with maize diversity, but there is little evidence to show how and
why culture influenced that diversity and modern maize biogeographic patterns in South
America. During expansion of the Inca Empire in the central Andes, maize diversity was
enhanced through selection for specific traits associated with reproduction. We show
ovule development related to improving seed quality and yield was selected in modern
maize as compared to archaeological maize. We hypothesize that this may have
contributed to maize becoming a staple under the Inca, as happened previously with
expanding Tiwanaku interactions and influence during the Middle Horizon.

Introduction

Modern maize (*Zea mays* ssp*. mays L*.) is one of the most important and productive global
food crops[1]. Prior to European contact in the Americas, maize had been adapted for a
wide range of environments. Its versatility is clear from its extensive geographic
distribution today, ranging from 58º north latitude to 40º south latitude, and growing below
sea level as well as 3,600 meters above sea level in the Andes[2]. In Bolivia there is
empirical evidence that small maize varieties can grow at altitudes of 3,800-3,900 meters
above sea level (masl) in some areas of the Lake Titicaca Basin[3], as well as in other
sheltered areas of Oruro and Potosí, although with lower levels of production[4, 5]Studies
of archaeological samples and modern molecular data indicate that maize originally
evolved in Mexico from the flowering species of grass, teosinte (*Zea mays* ssp*.*
parviglumus) ~9,000 years before the present (cal. BP) [6, 7]and spread into South America
by ~7,000 cal BP[8, 9] from the lowland and Eastern North America by ~2500 cal BP[10].
During this domestication process there was hybridization between teosinte and
mexicana (*Zea mays* ssp. *mexicana*)[11]. The dispersal of maize to the rest of the world
followed European colonization of the Americas in the 15th and 16th centuries AD[2, 12]. The
evolution and diversity of maize has been shaped by both cultural and natural selection
in a wide range of environmental and social contexts. For example, evidence has shown
that cultural preferences were a factor in maintaining otherwise undesirable traits[13, 14].
Maize seeds were selected and retained by farmers for use in subsequent growing
years to maintain or enhance specific morphological traits, such as color and texture,
based on changing cultural preferences, sometimes at the cost of reduced cultivability
and/or nutrition[15].

Politics and culture have had a strong influence on crop planting and food preferences in
ancient (and modern) civilizations. Indeed, in the Andes, previous research showed that

under the Inca empire, maize was a sacred crop[16], symbolic of political power, and tightly
associated with high-status or elite social groups[6, 17]. It was also an important element in
numerous cultural events and rituals[18]. From an agriculture-economic point-of-view,
planting a crop on the most suitable land takes priority if a farmer wishes to maximize the
output-to-input ratio. However, in certain civilizations, including Andean groups, maize
was not a crop for which yield was considered a deciding factor. For example, in Andean
highland communities, maize was not the most essential food crop during pre-Inca
times[19], not only because maize was ill-adapted to arid highland conditions and its
cultivation was risky during the short growing season, but also because planting maize
did not yield a return on investment. In the Andes during the Spanish conquest, the
chronicler Cobo observed that, "Indians dropped all that they were doing in order to
produce a little maize, even though it cost them more than it was worth". It seemed to him
that work in the fields was "…one of the major forms of recreation and festivals…", and
he was impressed by the "extraordinary fondness for farming"[20, 21]. This did not, however,
deter Andean farmers from planting an abundance of maize. In fact, while it was a
desirable food, it was more so a symbol of empire[17] and a ritual-feast crop, since its first
introduction to the region around 3000 BC[22].
103
Given the availability of archaeological maize samples, as well as the association of
maize with cultural rituals of an historically powerful civilization and its antecedents, we
were curious as to how – if at all – maize diversity was associated with Inca cultural
influence and modern maize biogeographic patterns in South America. The current study
focuses on maize samples from an offering associated with an elite stone tomb burial
context from south of La Paz, Plurinational State of Bolivia, and dated ca. 1470 cal AD.
We sequenced two archaeological Bolivian Maize (aBM) kernels with six libraries, using
short read sequencing. In parallel, we used [14]C Accelerator Mass Spectrometer (AMS)
radiocarbon dating to determine the approximate age of the samples. Phylogenetic
analysis of aBM, 16 archaeological and 226 modern maize samples (*Zea mays* ssp. *mays*
L.), 88 modern samples of Teosinte *(Zea mays* ssp. *parviglumis)*, 79 modern samples of
Mexicana teosinte *(Zea mays* ssp. *mexicana)* and 1 *Tripsacum* genotype revealed
genetic relationships among themselves (Table S1; Dataset S1). Finally, the unique
single nucleotide polymorphisms (SNPs) in aBM infer that the selection imposed to
enhance reproductive success in marginal highland growing environments resulted in
increased maize diversity. Based on the data presented herein, we conclude that a
combination of Inca culture, the social and political needs of local ethnic groups, and
mitigation of certain natural environmental factors may have contributed to the specific
preference for, and diversification of maize in at least this part of South America. As such,
our conclusions offer insight into the possible pathways that influenced maize genetic
diversity, thus providing a vital resource to investigate further the factors that drove
selection for maize domestication and diversity at the cusp of the cultural influence of the
Inca empire.
127
128
129 **Results**
130
Archaeological Bolivian maize sample (aBM)

La Paz is a major city located in the highlands of west-central modern Bolivia, which is
16°30′S, 68°09′W, 3,650 masl. Dry conditions in this region and the adjacent Andes
Mountain range have aided in the preservation of desiccated organic material. Our maize
samples were found within a woven camelid fiber pouch that was one of many offerings
associated with the mummified remains of a child reported as being found in a *chullpa*
(stone tower tomb) south of La Paz, Plurinational State of Bolivia around 1890 (Fig. 1).
Although it is impossible to know the exact location of the chullpa or the origin of the
individual, the assemblage was acquired by the U.S. Consul to Chile and was sent as a
gift to the Michigan State University Museum, where she was curated until 2019.
Following a series of recent analyses, she has since been voluntarily repatriated to
Bolivia[23] Given the purported archaeological context of the mummy and her abundant
offerings recorded in the MSU Museum intake records, it has been proposed that she
might represent an elite member of society. However, the nature of the coarse fabrics,
simple design motifs, and restricted color palette of her textiles would suggest a more
common individual, perhaps with an affiliation with a local high-status group. The vast
majority of the funerary towers that exist in a radius of 60 km south of La Paz city are
made of mudbricks. To our knowledge, the only exception is an Inca "cushioned" (i.e.,
rounded edges) stone and mudbrick tower located in Uypaca, in the nearby Achocalla
Valley, 8 km southwest of the city center[24]. Because of its characteristics, this tomb most
likely belonged to a local (possibly non-Inca) high-status group and was part of a more
extensive funerary site. Although it is impossible to know the exact provenience of the
mummy, it would not be unreasonable to suppose that she originated from this or another
similar tomb in the Achocalla Valley. Regardless, in the decades following her death,
community members periodically visited the *chullpa* and left offerings. These included the
pouch full of well-preserved maize from which we collected and analyzed samples from
two kernels. A single Accelerator Mass Spectrometer (AMS) date was obtained from one
kernel, yielding a radiocarbon age of 400±26 BP (D-AMS-027148; maize kernel;
corrected for fractionation, no $\delta^{13}C$ reported). For the date 400±26 BP the two possible
calibrated age ranges are 1440-1520 cal AD (p=0.823) and 1588-1621 cal AD (p=0.177),
with a median age of cal AD 1474. (Calibrated at 2σ with Calib 8.2 [IntCal 20]). Two
associated maize kernels from the same pouch were subjected to molecular analysis.
163
Genomic base modification patterns associated with ancient DNA (aDNA), including
cytosine deamination damage patterns, have been identified. Observed position-specific
substitutions at the ends of the sequence reads (e.g., C→T/G→A[25-29] (Fig. S1a), as well
as an increase in adenine (A) and guanine (G) residues at the 5′ end, concomitant with a
reduction of C and T residues[29-31] (Fig. S1b). Both features are indicative of postmortem
deamination patterns. From this, we evaluated all substitution types, error rate (Fig. S1c)
and read length (Fig. S2), including read length range[32] and high error rate types, which
are associated with aDNA genome data[33]. In total, these data are consistent with the
radiocarbon results, supporting that 1) with the maize is in fact consistent with an
archaeological biological sample, and 2) with based on the geographic origin of the
sample, the kernels potentially descended from the period of the last Inca empire before
the Spaniard Francisco Pizarro's conquest of the Inca capital, Cuzco, in A.D. 1533[34].
176

Our study focused on identifying the location from which aBM originally came,
establishing and explaining patterns of genetic variability of maize, with a specific focus
on maize strains that are related to our current aBM samples. We were especially
interested in exploring how these patterns of variability found in both archaeological and
modern samples reflect the influence of cultural systems of behavior on both the spread
of variants across the ancient landscape and selection for phenotypic traits by local
populations. In particular, it appears that the Inca empire had a large effect on this process
of genetic differentiation.
185
Genetic Identity and Genetic Relatedness
After Tupac Inca Yupanqui became the emperor of the Inca Empire in the 15th century,
the territory of the Inca empire was greatly expanded across much of western South
America, including the Qollasuyu, a broad territory located in what is now western Bolivia,
southern Perú, northern Chile, and non-highland Argentina. Different ethnic groups
populated this territory forming polities that were confederated at the time of Inca arrival.
Several Aymara Pakaje groups inhabited the highlands and the inter-Andean valleys
south of La Paz, from where the maize samples derive. Because maize was more of a
sacred than subsistence crop of the Inca and, to a degree symbolic of Inca imperial
control, the extent of its distribution as a crop may have been related to the extension of
social and political hegemony over newly acquired territories. Given the provenience of
the aBM and its age, it is very possible it was introduced into western highland Bolivia
from the Inca core area – modern Peru. To test this hypothesis, we first evaluated the
genetic relatedness of archaeological and contemporary modern maize samples using a
genome sequence approach to assist in identifying the origin(s) of aBM and the
relatedness to other maize from different locations. Our admixture f3-statistics test results
suggest aBM is not admixed. Still, f3 outgroup with *Tripsacum* suggests that aBM is
closely related to two archaeological lowland Peruvian maize (Z65 and Z61) among all
other ancient maize and modern maize (Fig. 2). These results indicate that the closest
relative of aBM is the archaeological maize that came from Peru. In addition, to predict
the geographic location of aBM, we used spatial auto-correlations in genetic data, which
allowed us to compare it to a set of archaeological maize samples of known geographic
origin[35]. Our results indicate the potential geographic location that aBM may have
originally come from (locator.py –seed 549657840) and the predicted location is inside of
Peru and very close to archaeological maize Z61 (Fig. 3; R2(x)=1.0, R2(y)=1.0). This
suggests that aBM might not only be genetically related to the archaeological maize from
ancient Peru, but also in the possible geographic location. This is interesting, as
historically, evidence from recordings of *capacocha* rituals showed that they were able to
source their food from the stations connecting high-altitude ritual sites and other regions
of the empire[34]. Indeed, data generated herein show that our aBM has a strong
relationship with archaeological Peruvian maize in terms of both genetic identity and
geographic location[36]. This finding is also consistent with the above previous research
revealing that one of the maize groups from Peru during Inca peak times spread into
newly conquered territories, then slowly diverged to fit the local environment[37] following
Inca cultural influence[20]. After this, diverged groups were introduced outside of Inca
conquered territories.
222

To investigate the genomic data from the aBM in relation to other archaeological and modern maize, principal component analysis (PCA) was used. As expected, the results showed that all the archaeological maize clustered with our aBM in the same cluster, which supports the authenticity of aBM as archaeological maize (Fig. 4). SM3, 5, and 10 were excluded due to their limited data to meet requirements for the PCA analysis. Unexpectedly, there is one modern Brazil maize in the ancient maize cluster that has overlap with archaeological Brazil maize Z6, which indicates that certain genetic traits from ancient Brazil maize might have been preserved in modern Brazil maize populations. There is also one modern tropicalis maize very close to archaeological Peruvian maize Z65. Considering the geographic origin of Z65 and its lowland characters, it is not surprising that modern tropicalis maize shares some level of genome sequence similarity. Following the establishment of an important maize culture center in Peru, maize was domesticated and diverged into different groups, which then spread along with Inca culture[38]. We found that the Central American cluster has overlap with both North and South America, which supports the distribution route of maize[39-41]. This finding also is consistent with previous research revealing that the original maize came from teosinte in Mexico, then dispersed northward into North America and southward into South America[6, 7, 9, 42, 43]. We also found that a few types of maize from Europe have a much closer distance to the archaeological maize cluster compared to other modern maize, which indicates maize from Europe might expectedly share certain traits or evolutionary characteristics with ancient maize. It is also consistent with the historical fact that maize spread to Europe from multiple areas and time periods after Christopher Columbus's late 15th century voyages to the Americas[12]. But as shown, maize also has diversity inside the European maize cluster. It is possible that European farmers and merchants may have favored different phenotypic traits, and the subsequent spread of specific varieties followed the new global geopolitical maps of the Colonial era[44, 45].

Taken together, these results support our hypothesis that aBM was introduced into western highland Bolivia from the Inca core area – modern Peru, and aBM may possibly reveal a history of adaptation to tropical conditions following cultural influence of the Inca empire[43, 46]. After this, maize, both as a primary food and as a symbol of power, was dispersed from Peru throughout the Inca empire, in the end enhancing maize diversity in South America.

Selection for aBM under Inca Culture Influence

After establishing the identity of aBM and its relatedness to other regional samples, we explored how aBM was shaped by human behavior, and specifically how cultural selection within the Inca empire influenced maize diversity in South America. To address this question, we examined the evolutionary relationship between aBM, its wild ancestor, and extant landraces. We constructed a Neighbour-Joining (NJ) and maximum-likelihood (ML) phylogenetic trees using the common genome-wide polymorphisms shared among aBM and modern maize samples[47]. Our data included 226 samples of modern maize, 88 modern samples of Teosinte (*Zea mays* ssp. *parviglumis*), 79 modern samples of Mexicana teosinte (*Zea mays* ssp. *mexicana*), and 1 *Tripsacum dactyloides* as an outgroup, totaling 28,588 genome-wide SNPs. As shown in Fig. S3a and Fig. S3b, both phylogenetic trees indicate that aBM lies outside the diversity of modern maize yet

clusters within a specific clade alongside one modern tropicalis maize. aBM tends to have
long branches compare to tropicalis maize, which can be explained by adaption for
specific local environment by time. This suggests that aBM shares a closer genetic
relationship with modern tropicalis maize while being genetically distinct from other
modern maize. Considering modern tropicalis maize is adapted to tropical climates
exhibiting traits such as drought tolerance, heat resistance, and the ability to thrive in low-
fertility soils, it is plausible that aBM might share these stress response characteristics
with tropicalis maize[48, 49].
277
To investigate the selection preference on aBM under Inca cultural influence, we
compared the allele frequency between aBM and other ancient maize samples. We
identified 18,668 SNPs unique to aBM. Notably, two genes - Zm00001eb123120 and
Zm00001eb178600-associated with traits "internode length below ear" and "nodes above
ear" traits[50]contain several unique SNPs in aBM (Dataset S2). This suggested that height
related traits in aBM may have been selected under Inca cultural practices. Additionally,
variations in the 5 or 3 untranslated regions UTR) can significantly affect gene expression.
We found 83 target genes harboring unique aBM SNPs in their UTR regions (Dataset
S3). Gene Ontology (GO) results showed that 3 of these genes are involved in
glycosyltransferase activity, while serval of others is involved in defense response to
biotic and abiotic stress and starch biosynthetic processes (Dataset S4). Combining all
this information, we infer that trait associated with plant development and stress response
processes might have been selected in maize under Inca cultural influence, although we
could not determine whether the selection was positive or negative.
292
Finally, to understand positive selection in modern maize compared to ancient maize, we
investigated the selective sweep using cross-population extended haplotype
homozygosity (XP-EHH) method. Following the same method of previous research[51], the
cutoff metric for significant signatures of selection was set at a value of 6 for 5 kb windows
across the genome. We only identified one target gene (Fig. 5): Zm00001eb105060,
whose biological process functions include plant ovule development, chromosome
segregation and asparaginyl-tRNA aminoacylation. Ovule development plays a crucial
role in improving maize breeding strategies via higher seed production and hybrid vigor[52].
The positive selection of this gene in modern maize suggests that characteristics aiming
for higher seed production were selected during maize diversification after aBM. This
finding may relate to breeding efforts to improve seed yield in modern maize compared
to ancient varieties.
305
In conclusion, the aBM included in the funerary accoutrements of a young elite Andean
girl belongs to the family of ancient Peruvian maize groups, and likely was introduced to
the Aymara region following Inca expansion in the late 15th century. Following its dispersal
as a crop across the entire Inca empire, maize was further modified by regional
populations. Some genomic changes reflect selection to mitigate local environmental
conditions, including response to stress. Thus, patterns of regional diversity in pre-
Contact South America maize reflect complex biocultural processes. These data may also
inform subsequent studies about Colonial dispersals and the origin of global maize
varieties[12].

Discussion

Previous archaeological and anthropological studies have demonstrated that Inca cultural
and political systems shaped access to and preferences for food. In our study we explored
the evidence for the contribution of Inca culture to maize diversity in South America by
analyzing 16 archaeological maize samples spanning at least 5,000 years of evolution,
and 226 modern maize samples. Recognizing their shared central Andean origins, we
propose that the archaeological Bolivian maize (aBM) sample came from ancient
Peruvian archaeological maize. After aBM was cultivated in the Andean highland, traits
adapting to climate change and environmental variability may have been selected by Inca
culture. For modern maize, traits related to improved seed production through ovule
development were selected compared to ancient maize. While our findings are consistent
with both genetic and archaeological evidence, they raise several questions. Notably,
what specific traits were selected in aBM, and how this selection occurred under Inca
cultural practices is unclear, especially given our limited aBM sample size and the lack of
the same genotype in archaeological maize under no Inca culture influence.

Early evidence has shown that cultural preferences were a factor in maintaining certain
aesthetically desirable traits[13, 14], such as color and texture, even at the cost of reduced
cultivability or nutrition[15] by farmers. This suggests it is possible that culture played an
important role in maize evolution and diversity. To better understand how exactly Inca
culture influenced maize diversity, we explored the possible origin of aBM and its closest
genetic relationships to ancient maize. We identified that aBM may have originated from
ancient Peru, and archaeological maize from Peru has the closest genetic relationship
with aBM, indicating that aBM is a product of Incan cultural dissemination. Following the
expansion of the Inca civilization, maize was introduced in new territories to demonstrate
the power of the Inca empire[20]. This might lead to the selection of specific traits in maize
due to cultural preferences, although this may have conflicted with natural selection
pressures.

We further investigated which traits were possibly selected under Incan culture by
analyzing unique SNPs in aBM. The results show that traits related to plant development,
and stress response were selected. These traits can aid in resisting drought and frost in
the relatively harsh conditions of the Andean highlands. The selected traits could facilitate
fermentation for the creation of *chicha* to meet the consumption requirement of elite
lifeways[17]. However, we are uncertain if these traits were directly selected under Inca
culture due to the limited aBM sample size, which prevented us from identifying positive
selection directly between aBM and other archaeological maize groups.

Climate change is expected to increasingly impact maize seed production in agricultural
areas worldwide. Traditional breeding systems have increased tolerance to biotic and
abiotic stress and contributed to yield in crops[53, 54]. In this study, we compared modern
maize to archaeological maize across the entire Western Hemisphere and found that the

ovule development traits were selected for breeding to improving maize seed production during its evolution.

Our results not only highlight the influence of Inca culture on maize diversity in South America but also point to the adaptive characteristics of maize in improving maize seed production during evolution. This provides a potential source for maize diversity studies and breeding programs. Most importantly, this work suggests that cultural influence plays a very important role in aiding maize diversity, alongside the traits selected by early farmers.

Materials and Methods

Plant material. The two archaeological Bolivian maize (aBM) kernels were collected prior to 1890 AD from the area south of modern La Paz, Bolivia, and later curated at the Michigan State University Museum until their repatriation in 2022[55]. AMS assay dates provided a range from 401 to 582 cal. BP, with a median age of 1474 cal AD. In addition, data from 16 archaeological samples collected from previous published research were also used, including two from Chile dating from 750 to 1,100 BP, two from Brazil dating from 510 to 690 BP, three from Peru dating from 600 to 1,000 BP, two from Argentina, dating from 980 to 1040 BP and from 70 to 130 BP, three from Honduras dating from 1,740 to 2,300 BP, and three from Mexico dating to 5,310 BP (Table S1)[36, 56].

aDNA extraction. Archaeological maize kernels were processed in a clean laboratory environment at Michigan State University. Whole kernels were crushed in a mortar and pestle, to which 600 uL of fresh PTB (*N*-phenacylthiazolium bromide) buffer (0.4 mg/mL proteinase K, T2.5 mM PTB, 50 nM dithiothreitol) was added. The sample was then added to 1260 uL of lysis buffer. An additional 600 uL of PTB buffer was used to rinse the mortar and added to the lysis buffer. The individually ground maize kernels were rotated overnight at 37 °C. The remaining DNA extraction followed Swarts et al., 2017[41].

Library construction, amplification, and sequence. DNA extracts were converted to Illumina-compatible paired-end DNA libraries in a clean setting using SMARTer ThruPLEX DNA-seq from Takara Bio and amplified with 7 polymerase chain reactions (PCR) cycles. The generated libraries were cleaned to remove remaining adaptors with AxyPrep Mag beads (Fisher Scientific) and analyzed on a 2100 Bioanalyzer (Agilent Technologies) to check for quality. The libraries were indexed to facilitate pooling on the NovaSeq SP at the University of Illinois (Urbana-Champaign).

Geographic location map. The geographical coordinates of each sample were obtained from previously published studies, as indicated (Table S1). Utilizing latitude and longitude information, the samples were plotted on a geographic map, resulting in the successful mapping of 17 ancient samples. Geographic data visualization was carried out in R using the ggplot2, sf, leaflet, rnaturalearth, rnaturalearthdata, ggspatial, and rnaturalearthhires packages.

Genome analysis. For the aBM sequence samples data with six libraries, we combined them into a single composite sample, referred to as aBMComb. This was achieved by mapping each individual aBM sequence sample to the B73 V5 soft-masked maize genome and subsequently removing duplicate reads before merging them. We merged the sequence sample to obtain higher coverage for the analysis was needed for this study.

Subsequently, we used MapDamage v.2.2.1[57] verified cytosine deamination profiles consistent with ancient DNA, and estimated the misincorporation frequencies. Based on these results we hard-masked 5′ thymine and 3′ adenine residues within 5nt of the two ends, where deaminations was most concentrated.

Data for 16 ancient samples with specific geographic location information (maize landraces) obtained from previously published work[36, 56, 58] by downloading sequencing from Sequence Read Archive (SRA) (https://www.ncbi.nlm.nih.gov/sra/)[59]. To ensure consistency and accuracy, we downloaded the raw sequence (SRA format) for each sample and converted it into FASTQ format using fastq-dump.

For the archaeological maize samples, adapters were removed and paired reads were merged using AdapterRemoval[60] with parameters --minquality 20 --minlength 30. All 5′ thymine and 3′ adenine residues within 5nt of the two ends were hard-masked, where deamination was most concentrated. Reads were then mapped to soft-masked B73 v5 reference genome using BWA v0.7.17 with disabled seed (-l 1024 -o 0 -E 3) and a quality control threshold (-q 20) based on the recommended parameter[61] to improve ancient DNA mapping. Duplicated reads were removed by GATK tools Picard (https://gatk.broadinstitute.org/hc/en-us). The coverage of each nucleotide position was estimated by bam-readcount (https://github.com/genome/bam-readcount) (Table S2), and the overall coverage of the assembled genome sequence was measured using Qualimap v2.2.1[62].

Variant Calling. We used SequenceTools pileupCaller (v 1.5.4.0; https://github.com/stschiff/sequenceTools) to generate pseudohaploid calls for 17 archaeological maize samples. The commands executed were: samtools mpileup -R -B -q20 -Q20 & pileupCaller -- sampleNames <ancient sample name> randomHaploid --singleStrandMode. This process yielded 16,815,782 single nucleotide polymorphisms (SNPs).

We future filtered the SNPs using estimated information from bam-readcount. SNPs supported by at least two reads were retained for each individual archaeological maize sample. We then combined the filtered SNPs from each sample, keeping only those SNPs that were present in at least one sample. This resulted in a final dataset of 2,808,917 SNPs for the 17 archaeological maize samples.

For all subsequent analyses, the ancient genotype data was merged with the genotype dataset for 394 samples from Grzybowski et al, ("Feb 03, 2023 version files" in Dryad repository: https://doi.org/10.5061/dryad.bnzs7h4f1). The merged dataset included

449   2,635,047 SNPs from chromosomes 1-10 that were common in both archaeological and
450   modern maize samples, which were used for downstream analysis.
451
452   ***f3 statistic and phylogenetic analyses***. We created a genotype file by combining 17
453   archaeological maize and publicly available SNP dataset form Grzybowski et al, resulting
454   in a total of 17-archaeological and 226-modern maize, 88-modern samples of Teosinte
455   *(Zea mays* ssp*. parviglumis)*, 79-modern samples of *Mexicana teosinte (Zea mays* ssp.
456   *mexicana)*, and 1-*Tripsacum* genotypes.
457
458   Next, we used plink v1.9[63] to prune for linkage disequilibrium with the parameters --indep-
459   pairwise 50 5 0.1, following previous research[45], to prepare the dataset for Principal
460   Component Analysis (PCA). The final results were plotted in R using the ggplot2
461   package[64].
462
463   We calculated the *f3* statistic[65, 66] using the admixr package[67] , which is part of the
464   package of ADMIXTURE software suite[68], to detect evidence of admixture involving aBM.
465   The outgroup f3 statistics was used to measure how closely aBM and each test group are
466   related to each other. The final results were plotted using Python.
467
A total of 28,588 SNPs present in both aBM and all modern maize, out of a total of
2,626,087 SNPs, was used to build the phylogenetic tree. A maximum likelihood (ML)
phylogenetic tree was constructed using SNPhylo[69] with parameter: -v snps.vcf -r -o
tripsacum -P tree -b 1000; A Neighbour-Joining (NJ) phylogenetic tree was constructed
using VCF2PopTree[70] with the "number of differences" model. Both ML and NJ
phylogenies trees were inferred and visualized using FigTree v1.4.4
(http://tree.bio.ed.ac.uk/software/figtree/).
475
476   **Geographical location prediction**. To predict the possible geographical location of aBM,
477   we applied a deep neural network method[35] using only 17 ancient maize samples, as all
478   the archaeological maize clustered in the same cluster. A total of 2,800,190 SNPs were
479   used to predict the possible geographical location of aBM. The command used was:
480   locator.py --vcf SNP_file --sample_data sample_data.txt –out name.
481
Infer selection on aBM. Due to the lack of archaeological maize with the same genotype
as aBM grows outside the influence of Inca culture, it was impossible to directly identify
the selection that occurred on aBM under Inca cultural influence. To determine if any of
our SNPs are related to known traits, we utilized existing information on SNP-trait
associations[50]. To achieve potential unique selection on aBM, we calculated the allele
frequency for each SNPs between aBM and other ancient maize, resulting in allele
frequency data for 49,896 SNPs. Of these,18,668 SNPs were unique to aBM. We
consulted SNPVersity 2.0 from MaizeGDB[47, 71] and found that 84 of these unique SNPs
were classified as "3_prime_UTR_variant" or "5_prime_UTR_variant". These SNPs
corresponded to 83 target genes, for which we retrieved Gone Ontology (GO) terms.
492
493   **Identification of selection sweeps.** To detect genomic signatures of recent selective
494   sweeps in modern maize compared to ancient maize groups, SNPs that present in at

least two archaeological samples and modern samples were used. We applied the cross-population extended haplotype homozygosity (XP-EHH)[72] method. A commonly used threshold $-\log_{10}(p) = 6$ was set as the cutoff threshold for identifying the signal of significant selection. The regions with $-\log10(p) >= 6$ were designated as candidate sweeps for candidate genes located within 5 KB of these regions identified, following the approach used in previous research[51]. MaizeMine from MaizeGDB[73] was used to obtain putative gene function information.

Data availability

The authors declare that all data supporting the findings of this study are included in the manuscript and its supplementary files or are available from the corresponding author upon request. Source data are provided with this paper. The genome sequence data of ancient Bolivian Maize has been deposited in the NCBI database under BioProject accession PRJNA886637. Custom scripts and single nucleotide polymorphism (SNP) calls are available on Dryad (DOI: https://doi.org/10.5061/dryad.w6m905qtd).

Acknowledgements

We are grateful to all participants who were involved in this project. We are thankful for the support of the Michigan State University Cloud Fellowship for computational skills training. This work was supported in part through computational resources and services provided by the Institute for Cyber-Enabled Research at Michigan State University. We thank Jennifer Wai, University of Chicago, for her help with DNA extraction and library preparation. We also thank Emily Josephs, Michigan State University, and Logan Kistler, Smithsonian Institution of National Museum of Natural History, for their help with the DNA analysis. Thanks to José Capriles Flores for his insightful reading and comments. The MSU Museum provided maize samples and images for documentation and analysis prior to repatriation to Bolivia. Research in the laboratory of Brad Day was supported by the MSU Research Foundation.

References

1. Ranum, P., Pena-Rosas, J.P. & Garcia-Casal, M.N. Global maize production, utilization, and consumption. *Ann N Y Acad Sci* **1312**, 105-112 (2014).
2. Bonavia, D. *Maize : origin, domestication, and its role in the development of culture*. (Cambridge University Press, Cambridge, NY; 2013).
3. Chavez, S.J.a.R.G.T. Early Maize on the Copacabana Peninsula: Implications for the Archaeology Introduction. *Histories of Maize*, pp. 415-428 (2006).
4. Hastorf, C., Whitehead, W., Bruno, M. & Wright, M. The Movements of Maize into Middle Horizon Tiwanaku, Bolivia Introduction. *Histories of Maize*, pp. 429-448. (2006).
5. Suca Gómez, A. Caracterizacion de granos de maiz andino(Zea mays L.) de colectas de germoplasma en la Facultad de Agronomia -UMSA. *Trabajo dirigido para optar por el título de licenciado en Ingeniería Agronómica.* (2011).
6. Doebley, J. The genetics of maize evolution. *Annu Rev Genet* **38**, 37-59 (2004).

7. Matsuoka, Y. *et al.* A single domestication for maize shown by multilocus microsatellite
genotyping. *Proc Natl Acad Sci U S A* **99**, 6080-6084 (2002).
8. Grobman, A. *et al.* Preceramic maize from Paredones and Huaca Prieta, Peru. *Proc Natl
Acad Sci U S A* **109**, 1755-1759 (2012).
9. Lombardo, U. *et al.* Early Holocene crop cultivation and landscape modification in
Amazonia. *Nature* **581**, 190-193 (2020).
10. Hart, J.P. & Lovis, W.A. Reevaluating What We Know About the Histories of Maize in
Northeastern North America: A Review of Current Evidence. *Journal of Archaeological
Research* **21**, 175-216 (2013).
11. Hufford, M.B. *et al.* The genomic signature of crop-wild introgression in maize. *PLoS
Genet* **9**, e1003477 (2013).
12. Rebourg, C. *et al.* Maize introduction into Europe: the history reviewed in the light of
molecular data. *Theor Appl Genet* **106**, 895-903 (2003).
13. Bellon, M.R. The dynamics of crop infraspecific diversity: A conceptual framework at the
farmer level. *Econ Bot* **50**, 26-39 (1996).
14. Knapp, G. Changing fortunes: Biodiversity and peasant livelihood in the Peruvian Andes.
Geogr Rev **87**, 568-570 (1997).
15. Louette, D. & Smale, M. Farmers' seed selection practices and traditional maize varieties
in Cuzalapa, Mexico. *Euphytica* **113**, 25-41 (2000).
16. Mintz, S.W. *Sweetness and power : the place of sugar in modern history*. (Viking, New
York, N.Y.; 1985).
17. Bray, T.L. Inka pottery as culinary equipment: Food, fasting, and gender in imperial state
design. *Lat Am Antiq* **14**, 3-28 (2003).
18. Ceruti, M.C. Frozen Mummies from Andean Mountaintop Shrines: Bioarchaeology and
Ethnohistory of Inca Human Sacrifice. *Biomed Res Int* **2015**, 439428 (2015).
19. Burger, R.L. & Vandermerwe, N.J. Maize and the Origin of Highland Chavin Civilization
- an Isotopic Perspective. *Am Anthropol* **92**, 85-95 (1990).
20. Doutriaux, M. Power, ideology and ritual: the practice of agriculture in the Inca Empire.
Pap Kroeber Anthropol Soc, 91-108 (2001).
21. Cobo, B. & Hamilton, R. *Inca religion and customs*, Edn. 1st. (University of Texas Press,
Austin; 1990).
22. Pearsall, D.M. Phytolith analysis of archeological soils: evidence for maize cultivation in
formative Ecuador. *Science* **199**, 177-178 (1978).
23. Lovis, W.A., J. M. Capriles & D. Trigo Rodríguez La repatriación de una momia andina
del Museo de la Universidad Estatal de Michigan al Estado Plurinacional de Bolivia. *Textos
Antropológicos* **21(1):23-41** (2023).
24. Michel López, M. Prospección arqueológica del valle de Achocalla. *XIV Reunión Anual de
Etnología* **Tomo 1: 209-222 (2001)** (2001).
25. Handt, O., Krings, M., Ward, R.H. & Paabo, S. The retrieval of ancient human DNA
sequences. *Am J Hum Genet* **59**, 368-376 (1996).
26. Krings, M. *et al.* Neandertal DNA sequences and the origin of modern humans. *Cell* **90**,
19-30 (1997).
27. Hansen, A., Willerslev, E., Wiuf, C., Mourier, T. & Arctander, P. Statistical evidence for
miscoding lesions in ancient DNA templates. *Mol Biol Evol* **18**, 262-265 (2001).
28. Gilbert, M.T. *et al.* Characterization of genetic miscoding lesions caused by postmortem
damage. *Am J Hum Genet* **72**, 48-61 (2003).

29.   Dabney, J., Meyer, M. & Paabo, S. Ancient DNA damage. *Cold Spring Harb Perspect Biol*
5 (2013).
30.   Briggs, A.W. *et al.* Patterns of damage in genomic DNA sequences from a Neandertal.
Proc Natl Acad Sci U S A **104**, 14616-14621 (2007).
31.   Stiller, M. *et al.* Patterns of nucleotide misincorporations during enzymatic amplification
and direct large-scale sequencing of ancient DNA. *Proc Natl Acad Sci U S A* **103**, 13578-
13584 (2006).
32.   Paabo, S. Ancient DNA: extraction, characterization, molecular cloning, and enzymatic
amplification. *Proc Natl Acad Sci U S A* **86**, 1939-1943 (1989).
33.   Prufer, K. *et al.* Computational challenges in the analysis of ancient DNA. *Genome Biol*
11, R47 (2010).
34.   Wilson, A.S. *et al.* Stable isotope and DNA evidence for ritual sequences in Inca child
sacrifice. *Proc Natl Acad Sci U S A* **104**, 16456-16461 (2007).
35.   Battey, C.J., Ralph, P.L. & Kern, A.D. Predicting geographic location from genetic
variation with deep neural networks. *Elife* **9** (2020).
36.   Kistler, L. *et al.* Multiproxy evidence highlights a complex evolutionary legacy of maize
in South America. *Science* **362**, 1309-1313 (2018).
37.   Vallebueno-Estrada, M. *et al.* Correction: Domestication and lowland adaptation of coastal
preceramic maize from Paredones, Peru. *Elife* **12** (2023).
38.   Cook, O.F. Peru as a center of domestication: Tracing the Origin of Civilization Through
the Domesticated Plants. *Journal of Heredity* **16**, 95-110 (1925).
39.   Wang, L. *et al.* Molecular Parallelism Underlies Convergent Highland Adaptation of Maize
Landraces. *Mol Biol Evol* **38**, 3567-3580 (2021).
40.   Dillehay, T.D. *et al.* Monte Verde: seaweed, food, medicine, and the peopling of South
America. *Science* **320**, 784-786 (2008).
41.   Swarts, K. *et al.* Genomic estimation of complex traits reveals ancient maize adaptation to
temperate North America. *Science* **357**, 512-515 (2017).
42.   van Heerwaarden, J. *et al.* Genetic signals of origin, spread, and introgression in a large
sample of maize landraces. *Proc Natl Acad Sci U S A* **108**, 1088-1092 (2011).
43.   Mercer, K., Martinez-Vasquez, A. & Perales, H.R. Asymmetrical local adaptation of maize
landraces along an altitudinal gradient. *Evol Appl* **1**, 489-500 (2008).
44.   Haberer, G. *et al.* European maize genomes highlight intraspecies variation in repeat and
gene content. *Nat Genet* **52**, 950-957 (2020).
45.   Tenaillon, M.I. & Charcosset, A. A European perspective on maize history. *C R Biol* **334**,
221-228 (2011).
46.   Cerda-Hurtado, I.M. *et al.* Climatic adaptation and ecological descriptors of wild beans
from Mexico. *Ecol Evol* **8**, 6492-6504 (2018).
47.   Grzybowski, M.W. *et al.* A common resequencing-based genetic marker data set for global
maize diversity. *Plant J* **113**, 1109-1121 (2023).
48.   Chakradhar, T., Hindu, V. & Reddy, P.S. Genomic-based-breeding tools for tropical maize
improvement. *Genetica* **145**, 525-539 (2017).
49.   Menkir, A. *et al.* Approaches and progress in breeding drought-tolerant maize hybrids for
tropical lowlands in west and central Africa. *Plant Genome* **17**, e20437 (2024).
50.   Wallace, J.G. *et al.* Association mapping across numerous traits reveals patterns of
functional variation in maize. *PLoS Genet* **10**, e1004845 (2014).

51. Brodie, A., Azaria, J.R. & Ofran, Y. How far from the SNP may the causative genes be? *Nucleic Acids Res* **44**, 6046-6054 (2016).
52. Yang, X. & Tucker, M.R. Establishing a regulatory blueprint for ovule number and function during plant development. *Curr Opin Plant Biol* **63**, 102095 (2021).
53. Galluzzi, G., Seyoum, A., Halewood, M., Lopez Noriega, I. & Welch, E.W. The Role of Genetic Resources in Breeding for Climate Change: The Case of Public Breeding Programmes in Eighteen Developing Countries. *Plants (Basel)* **9** (2020).
54. Kim, K.H. & Lee, B.M. Effects of Climate Change and Drought Tolerance on Maize Growth. *Plants (Basel)* **12** (2023).
55. Lovis, W.A. A Mummy Returns Home. *MSU Today, MSU 360, Faculty Voice* (2019).
56. Kistler, L. *et al.* Archaeological Central American maize genomes suggest ancient gene flow from South America. *Proc Natl Acad Sci U S A* **117**, 33124-33129 (2020).
57. Jonsson, H., Ginolhac, A., Schubert, M., Johnson, P.L. & Orlando, L. mapDamage2.0: fast approximate Bayesian estimates of ancient DNA damage parameters. *Bioinformatics* **29**, 1682-1684 (2013).
58. Vallebueno-Estrada, M. *et al.* The earliest maize from San Marcos Tehuacan is a partial domesticate with genomic evidence of inbreeding. *Proc Natl Acad Sci U S A* **113**, 14151-14156 (2016).
59. Chia, J.M. *et al.* Maize HapMap2 identifies extant variation from a genome in flux. *Nat Genet* **44**, 803-807 (2012).
60. Lindgreen, S. AdapterRemoval: easy cleaning of next-generation sequencing reads. *BMC Res Notes* **5**, 337 (2012).
61. Schubert, M. *et al.* Improving ancient DNA read mapping against modern reference genomes. *BMC Genomics* **13**, 178 (2012).
62. Okonechnikov, K., Conesa, A. & Garcia-Alcalde, F. Qualimap 2: advanced multi-sample quality control for high-throughput sequencing data. *Bioinformatics* **32**, 292-294 (2016).
63. Purcell, S. *et al.* PLINK: a tool set for whole-genome association and population-based linkage analyses. *Am J Hum Genet* **81**, 559-575 (2007).
64. Wickham, H. *Ggplot2 : elegant graphics for data analysis*. (Springer, New York; 2009).
65. Green, R.E. *et al.* A draft sequence of the Neandertal genome. *Science* **328**, 710-722 (2010).
66. Durand, E.Y., Patterson, N., Reich, D. & Slatkin, M. Testing for Ancient Admixture between Closely Related Populations. *Molecular Biology and Evolution* **28**, 2239-2252 (2011).
67. Petr, M., Vernot, B. & Kelso, J. admixr-R package for reproducible analyses using ADMIXTOOLS. *Bioinformatics* **35**, 3194-3195 (2019).
68. Alexander, D.H. & Lange, K. Enhancements to the ADMIXTURE algorithm for individual ancestry estimation. *BMC Bioinformatics* **12**, 246 (2011).
69. Lee, T.H., Guo, H., Wang, X., Kim, C. & Paterson, A.H. SNPhylo: a pipeline to construct a phylogenetic tree from huge SNP data. *BMC Genomics* **15**, 162 (2014).
70. Subramanian, S., Ramasamy, U. & Chen, D. VCF2PopTree: a client-side software to construct population phylogeny from genome-wide SNPs. *PeerJ* **7**, e8213 (2019).
71. Schnable, P.S. *et al.* The B73 maize genome: complexity, diversity, and dynamics. *Science* **326**, 1112-1115 (2009).
72. Sabeti, P.C. *et al.* Genome-wide detection and characterization of positive selection in human populations. *Nature* **449**, 913-918 (2007).

73.   Lawrence, C.J., Dong, Q., Polacco, M.L., Seigfried, T.E. & Brendel, V. MaizeGDB, the
679   community database for maize genetics and genomics. *Nucleic Acids Res* **32**, D393-397
680   (2004).

681
682

Figures and Tables

Fig.1. Location and appearance of archaeological Bolivian maize (aBM). a Map of maize sample collection and location information. b Photographs of kernels showing morphological characteristics.

Fig. 2. Admixture f3-statistics. In the form f3 (aBM; Test, tripsacum) where Test represents archaeological and modern samples.

Fig. 3. Predicted geographic location of archaeological Bolivian maize (aBM). The blue point is the predicted locations of aBM based on the SNPs from all archaeological maize. The pink points are the archaeological samples.

Fig. 4. Principal Component Analysis (PCA) of nuclear genomic data from archaeological and modern maize. Samples with greater affinity to one another cluster closer together. Samples with the same color and shape indicate they came from the same country.

Fig. 5. Genome-wide distribution of ancient maize specific selective sweeps. a Signal of selection of modern maize compared to archaeological maize population. Each point indicates a SNP. The blue horizontal line shows the genome-wide significance level ($p=1 \times 10^{-6}$).

Fig. S1. Ancient DNA (aDNA) damage pattern of archaeological Bolivian maize (aBM). a Cytosine deamination damage patterns for the combined bam file of six aBM sequence samples and individual six aBM sequence samples. The position specific substitutions from the 5' end (red) and the 3' end (green) of a read. The red line corresponds to C to T substitutions, the green line corresponds to G to A substitutions, and the blue line represents other types of substitutions. b Ancient DNA damage profile. The four upper plots show the base frequencies inside and outside of a read, where the open grey box corresponds to a read. The two lower plots show the position specific substitutions from the 5' end and the 3' end of a read. The red line corresponds to C to T substitutions, the blue line corresponds to G to A substitutions, and the fade line represents other types of substitutions.

Fig. S2. Archaeological Bolivian maize (aBM) DNA read length. a Read length distribution of combined aBM bam files prior to hard-masked 5′ thymine and 3′ adenine residues within 5nt of both ends.

Fig. S3. Phylogenetic trees of archaeological Bolivian maize (aBM). a Neighbour-Joining tree with VCF2Pop. b Maximum likelihood (ML) tree with SNPhylo. Parv (*Zea*

mays ssp. *parviglumis*), Zmex (*Zea mays* ssp. *mexicana*), Europe (modern Europe), SweetC (model sweet corn), Ztrol (modern Tropical maize), mMexico (modern Mexico), mParaguay (modern Paraguay), mBrazil (modern Brazil), mPeru (modern Peru), mEcuador (modern Ecuador), mGuatemala (modern Guatemala), and mColombia (modern Colombia).

Fig. S4. Q-Q plot for genome-wide selection in modern maize compared to archaeological maize. The Q-Q plot between modern and archaeological maize. Each point indicates a value from expected and observed. Red line represents the expected distribution of the p-value, while the blue trend represents the observed distribution.

Table S1. Geographical information of the archaeological maize samples used in this study.

Table S2: Paleogenomic characterization of archaeological Bolivian maize sequence samples with six libraries.

Table S3. The percentage of genomic sites covered at variable depths in the archaeological Bolivian maize (aBM) sample (aBMComb_rm5nt.rmDR.bam).