Green cities and the risk for vector-borne disease transmission for humans and animals: a scoping review

Mathilde Mercat[1†] and Colombine Bartholomée[1†], Florence Fournet[1], Magdalena Alcover Amengual[2], Maria Bourquia[3], Emilie Bouhsira[4], Anthony Cornel[5], Xavier Fernandez Cassi[2], Didier Fontenille[1], Adolfo Ibáñez-Justicia[6], Renaud Marti[7], Nicolas Moiroux[1], El Hadji Niang[8], Woutrina Smith[9], Jeroen Spitzen[6], Tessa M. Visser[10], Constantianus J.M. Koenraadt[10] and Frederic Simard*[1]

[1] MIVEGEC, Univ. Montpellier, IRD, CNRS – Montpellier, France
[2] Laboratory of Parasitology, Department of Biology, Healthcare and Environment, Faculty of Pharmacy and Food Sciences, University of Barcelona – Barcelona, Spain
[3] Unité de Parasitologie et Maladies Parasitaires, Département de Pathologie et Santé publique vétérinaires, Institut Agronomique et Vétérinaire Hassan II – Rabbat, Morocco
[4] InTheres, Univ. Toulouse, INRAE, Ecole Nationale Vétériniare de Toulouse – Toulouse, France
[5] Mosquito Control Research Laboratory, Department of Entomology & Nematology, University of California at Davis – Parlier, California, USA
[6] Centre for Monitoring of Vectors, Netherlands Institute for Vectors, Invasive plants and Plant Health, Netherlands Food and Consumer Product Safety Authority – Wageningen, the Netherlands
[7] TETIS, Univ. Montpellier, AgroParisTech, CIRAD, CNRS, INRAE – Montpellier, France
[8] Laboratoire d'Ecologie Vectorielle et Parasitaire, Faculty of Sciences and Techniques, Univ. Cheikh Anta Diop – Dakar, Senegal
[9] One Health Institute, School of Veterinary Medicine, University of California – Davis, California, USA
[10] Laboratory of Entomology, Wageningen University & Research, – Wageningen, the Netherlands

[†] Mathilde Mercat and Colombine Bartholomée contributed equally to this work.

*Corresponding author
Correspondence: frederic.simard@ird.fr

ABSTRACT
Background: Greening cities is a nature-based strategy to sustainable urban development that integrates natural elements like plants or water bodies, to mitigate climate change impacts and enhance human well-being. However, urban green infrastructures (UGIs) can influence the distribution of disease vectors, potentially affecting vector-borne diseases (VBDs). UGIs may provide new suitable environments for urban vectors, while also creating opportunities to mitigate VBD risks through

predation, competition, and dilution effects. This article examined the relationships
between UGIs, vectors, and associated pathogens, impacting both human and animal
health, highlighting knowledge gaps and identifying research priorities to support VBD
risk mitigation measures and to guide smart urban planning and design.
Methods: A systematic literature search was conducted following PRISMA guidelines
in three databases (Pubmed, Scopus, Web of Science). Selected articles involved (i)
any aspect of a urban vector system, (ii) in UGIs, and (iii) statistically analyzing the
effects of UGIs on VBD risk. Methods employed to characterize UGIs and VBDs were
described and the identified impacts were summarized by vector group.
Results: Among the 99 articles reviewed, most addressed mosquito-pathogen
systems (67) and tick-pathogen systems (29), with studies often confined to a single
city or several cities within the same country and focused on one vector group. Urban
vegetation generally appeared to heighten the risk of tick-borne diseases. In contrast,
the influence of UGIs on the risk of mosquito-borne diseases varied depending on the
vector system and on the environmental and climatic context. The diversity of
indicators used to assess UGIs and VBD risks may affect the observed impact on VBD
risk.
Conclusion: Given the increasing popularity of urban greening, it is crucial to
investigate its potential implications for public health, and thereby urban planning
decisions. However, the lack of standardized protocols complicates the accurate
assessment of the effects of UGIs on the risk for VBD emergence and transmission
and consequently, on potential mitigation measures.
63
64 *Keywords:* mosquitoes, ticks, sand flies, chigger mites, urban green infrastructure, city
65

Introduction

Urbanization is an ongoing global phenomenon that brings both opportunities and challenges. By 2050, approximately 70% of the world population will live in urban areas (UN Habitat, 2022). Urbanization offers opportunities to reduce poverty and inequality by increasing access to education and healthcare (United Nations, 2020). However, the rapid growth of cities, often poorly controlled, has led to social and economic inequalities, as well as health issues, primarily affecting the most vulnerable populations (Kuddus et al., 2020). City development drives environmental disruptions by modifying land use. As urban areas are responsible for most of the world's carbon emissions, increasing urbanization could further exacerbate pollution (UN Habitat, 2022). Urban landscapes are increasingly vulnerable to extreme events such as flooding or urban heat islands, negatively impacting human health (Heaviside et al., 2017). Climate change is expected to intensify both the frequency and severity of these events (Vautard et al., 2020). Additionally, urbanization induces habitat loss and fragmentation, leading to decreased biodiversity (E. L. Jones & Leather, 2012). Urban landscapes are a mosaic of highly heterogeneous environments (Breuste et al., 2008) that form complex interfaces connecting natural, rural, peri-urban, and urban ecosystems (Honnen & Monaghan, 2017). These interfaces increase interactions between humans, livestock, and wildlife, thereby increasing the risk of zoonotic pathogen emergence and transmission (Hassell et al., 2017; Pearce et al., 2018). High human density in urban areas and global connectivity through trade and travel further facilitate the introduction and spread of arthropod vectors and the pathogen agents they carry (Weaver & Reisen, 2010).

Greening cities is recognized as a key intervention to "make cities inclusive, safe, resilient, and sustainable," as outlined by the United Nations (UN) Sustainable Development Goals (United Nations, 2022). This concept involves integrating vegetation within urban environments through various initiatives, such as creating blue spaces (e.g, lakes, rivers, ponds) (Hyseni et al., 2021), preserving parks, planting trees, and establishing green corridors, which are defined as interconnected landscape elements that help to protect the environment (Aly & Amer, 2010). Urban green spaces mitigate environmental challenges and contribute to various ecosystem services. It reduces heat island effects (Gunawardena et al., 2017; Marando et al., 2022), alleviates flooding (Bai et al., 2018), and reduces air pollution by capturing carbon dioxide (Baró et al., 2014; Mexia et al., 2018). Moreover, urban green spaces restore and maintain city biodiversity (Archibald et al., 2017; Vergnes et al., 2012). Substantial evidence highlights the positive impact of urban greening on human well-being and health, including stress reduction, increased opportunities for physical activities that can lower the risk of non-communicable diseases like hypertension and obesity, and a decrease in respiratory diseases (Van Den Bosch & Ode Sang, 2017). These benefits align with recommendations from the recently established One Health High Level Expert Panel (OHHLEP), which promotes a holistic approach to health for people, animals, and the environment (WHO, 2021).

While Urban Green Infrastructures (UGIs; see Boxes 1 and 2) provide ecological and health benefits, they can also create new habitats within cities that favour the establishment and spread of arthropod vectors of human and animal pathogens. This raises the risk of Vector-Borne Disease (VBD, see Box 1) emergence and transmission in urban areas (Fournet et al., 2024). Hematophagous (blood-feeding) arthropods are key players in transmitting pathogens than can impact human and animal health. Many VBDs are on the rise worldwide, with their arthropod vectors spreading beyond native ranges due to climate change, globalization, and the intensification of international trade and travel (Cuthbert et al., 2023). Arthropod vectors such as mosquitoes and ticks are particularly able to adapt to new environments, often acting as invasive species due to their specific ecological, biological, and genetic traits (Renault et al., 2024). As such, UGIs may inadvertently increase habitat suitability for these disease vectors in urban areas, thereby increasing human-vector contact in these densely populated environments. On the other hand, UGIs may also support the establishment of predators and competitors of urban arthropod vectors, facilitating novel trophic networks that could reduce vector populations and mitigate VBD

risk (Prihandi & Nurvianto, 2022). Actually, increased biodiversity in UGIs can lead to shifts in vector
population dynamics, host availability, and/or the presence of biological control agents, potentially
resulting in dilution effects for pathogens (Keesing & Ostfeld, 2021).
This scoping review examined how UGIs and the process of urban greening influence VBD
risks for humans and animals. The objectives were to: (i) assess the influence of different UGIs on
the presence and abundance of vectors (e.g., mosquitoes, ticks, sand flies, and chigger mites) and
VBD dynamics (e.g., dengue fever, West Nile fever, malaria, Lyme disease, leishmaniasis, and
scrub typhus); (ii) identify UGIs characteristics such as connectivity, surface area, proximity to
residential areas, and plants species that most affect vector populations and VBD dynamics; and
(iii) compare the effects of UGIs on the same vector species across different locations, as well as
within the same location when possible. Moreover, we identified critical knowledge gaps that
require further research to inform VBD mitigation measures and guide the smart design of UGIs
aiming to minimize VBDs risks while fostering greener cities. This work will help city planners and
policymakers in understanding the VBD risks linked to urban greening, tailored to each city's
epidemiological context, enabling them to implement landscape management strategies in a
thoughtful and informed way.

Box 1. General definitions

135 **Vector:** an invertebrate, typically an arthropod, that transmits an infectious agent (Duvallet et
136 al., 2018; Wilson et al., 2017) from one host (human or animal) to another (Blanc & Gutiérrez,
137 2015).
138
Vector Borne Disease (VBD): human or animal diseases caused by parasites, viruses, and
bacteria that are transmitted by vectors (WHO, 2020).
Risk: According to the United Nations Office for Disaster Risk Reduction (UNDRR) (UNDRR,
2019), a risk is a "potential loss of life, injury or destroyed assets which could occur to a system,
society or a community, determined probabilistically as a function of hazard, exposure,
vulnerability, and capacity." The risk may be health, economic, ecological, social or political. It can
be summarized as risk = hazard x exposure x vulnerability.
VBD risk: According to the UNDRR disaster risk definition (UNDRR, 2019), VBD risk can be
defined as a hazard linked to an infected vector x exposure to this vector x vulnerability (of the
vertebrate host population).
Urban green infrastructure (UGI): L. Jones et al. (2022) introduced the term "Urban Green
Infrastructure" to categorize various green and blue spaces within urban environments. Their
typology resulted in nine categories, including gardens, parks, amenity areas, other public spaces
(e.g., cemeteries), linear features, constructed green infrastructures (e.g., green roofs or green
walls), and three additional categories for urban water spaces. This review used the overarching
term "urban green infrastructure" more broadly to also include vegetation in neglected urban
spaces, such as wastelands and vacant lots (Yang et al., 2019), urban agriculture farms and urban
forests (see Box 2).

Box 2. Classification of UGIs

160
161 We developed a classification system to describe urban green infrastructures (UGIs) as follows:

162
163 • **Urban green space typology:** This typology is based on the framework of L. Jones et al.
164   (2022), which classifies urban green spaces according to their use, land cover, ownership,
165   geometry, and management. For example, if an UGI is described as an 'urban park' or as

the connectivity between urban green spaces, it will be included in the class 'Urban green space typology'. This classification allows detailed descriptions including:

- o **Use:** urban parks, cemeteries, zoos, urban farms, urban agriculture, etc.
- o **Ownership:** residential areas, publicly-owned parks, etc.
- o **Geometry:** green roofs, walls, etc.

Key characteristics of urban green spaces can be described by:

- o **Shape and size:** This refers to the structural characteristics of urban green spaces, including overall shape, perimeter, and surface area.
- o **Connectivity:** This term denotes the interconnectedness of green spaces or their connection to peri-urban environments.
- o **Relevant features:** This refers to outstanding areas within urban green spaces, such as trails and edges.
- o **Surrounding land cover:** This refers to the type of vegetation or soil present near urban green spaces.
- o **Management:** This refers to maintenance approaches to the vegetation within the urban green space, such as grass-cutting or watering schedules.
- o **Distance to urban green spaces:** Distance of sampling points to urban green areas.

- **Urban vegetation description:** This description focuses on the detailed characterization of urban vegetation including land cover, vegetation indexes, and specific vegetation traits such as height, tree density, diversity, etc. It provides a comprehensive description of the vegetation around the sampling points across different studies. For example, vegetation described using Normalized Difference Vegetative Index (NDVI) within a 200-meter buffer around a mosquito trap will be included in this category. Key components include:
  - o **Type of vegetation:** This describes vegetation based on its land cover type, including grass, deciduous forest, etc.
  - o **Litter composition:** This describes the composition of organic litter found within the vegetation, such as leaves and twigs.
  - o **Vegetation index:** This index allows for an alternative vegetation description through remote sensing indices, tree cover percentage, or overall vegetation cover. The NDVI, is a remote sensing index used to determine the presence of vegetation. It ranges from -1 to 1, and the higher it is, the denser the vegetation (Martinez & Labib, 2023).
  - o **Vegetation characteristics:** This encompasses many aspects of vegetation traits, including height, diversity, shade, size of tree holes, tree density, etc.
  - o **Plant species:** This involves a thorough characterization of vegetation, with a focus on the presence and attributes of specific plant species studied within the area.
  - o **Distance to urban vegetation:** Distance of sampling points to urban vegetation.

Methods

The 'Preferred Reporting Items for Systematic Reviews and Meta-Analyses' (PRISMA) guidelines (Page et al., 2021) were used to conduct a scoping review examining the links between UGIs, vectors, and VBD risk (Box 1). Key methodological steps are outlined in Figure 1 and detailed below. The PRISMA check-list is available in Additional file 1: Text S1.

Search strategy

A comprehensive search was conducted in October 2022 using three online databases: PubMed, Scopus, and Web of Science. The search strategy combined key concepts related to

"urban," "greening," "vector," and "vector-borne disease." All possible associated keywords for each key concept were included. A proximity operator was used when available to ensure that "greening" keywords referred specifically to urban areas (Additional file 2: Text S2). The query targeted titles, abstracts, and keywords, and searches were conducted using the CADIMA web tool (Kohl et al., 2018). All articles published up to October 6, 2022 (with no specified start date) were considered. Given that urban greening is a dynamic and relatively new management approach, relevant articles were expected to be limited. As a proxy for "greening cities", all articles assessing any component of VBD risk related to any aspect of UGIs were included.

Article selection

After the systematic database search, duplicates were automatically removed using CADIMA tool. The remaining articles underwent an initial systematic screening based on titles and abstracts, using the following inclusion criteria: articles (i) must be original research published in peer-reviewed journals in English; (ii) must address any aspect of the vector system in relation to human or animal health (e.g., vector species, pathogens, or hosts); and (iii) must focus on at least one UGI (e.g., urban park, green/blue corridor, vegetation index or similar). Subsequently, a full-text screening was conducted to further refine the selection. In this step, articles were included if they: (i) referred specifically to urban areas; (ii) examined any aspect of UGIs (e.g., urban parks, vegetation cover indices, or tree species); and (iii) statistically analyzed the effect of different UGIs (e.g., multiple parks, varying vegetation cover, or different tree species) on VBD risk including hazard, exposure, and/or vulnerability. Commentaries, editorial reviews, opinion papers, and articles from grey literature were excluded, as well as those focusing on VBDs in urban environments without reference to green spaces. Two of the authors (MM and CB) independently reviewed both screenings (titles and abstracts, and full-text articles) for inclusion and extracted articles data, with any disagreement resolved by consensus

Data extraction and description

From each article, the following metadata was extracted: location (continent, country, city), year of publication, year of sampling, vectors involved (i.e., groups of vectors (mosquitoes, ticks, sand flies, chigger mites), including genus, species, and life stage), as well as the disease and host of interest, when available. The methodology used for vector sampling was also documented, including the type of trap and frequency of collection, as well as the diagnostic tests employed to detect pathogens. The risk indicators from the studies were categorized into three groups: **"vector data"**, for indicators addressing vector presence, abundance, or density; **"vector-borne pathogen data"**, for indicators related to vector infection by pathogens or host disease prevalence/incidence; and **"vector and vector-borne pathogen data"**, for studies encompassing both vector and pathogen aspects. Metadata related to the description of UGIs were extracted as well, including the type of methodology used (e.g., general classification based on usage, remote sensing, or field description), the variables monitored (e.g., land-use classification or Normalized Difference Vegetation Index (NDVI)), and the buffer size used around sampling points, when available. The variables used to describe UGIs were classified into two categories: (1) urban green space typology and (2) urban vegetation description (Box 2).

Finally, the statistical framework used to analyze the effect of UGIs on VBD risk was also extracted from each article. All combinations of VBD risk and UGI variables (*VBD risk variable ~ UGI variable*) tested statistically were collected. For each interaction, the effect of UGI on VBD risk was classified into four categories: (i) harmful, when a variable describing the UGI increases VBD risk, (ii) beneficial, when a variable describing the UGI decreases VBD risk, (iii) variable when the effect of the UGI variable on VBD risk varies across sampling sites, independently of site characteristics, or (iv) no association when there was no significant relationship between the UGI variable and VBD risk. The data extracted from all of the articles can be found in Additional file 3: Dataset S1. All data and additional files are available online (Bartholomée & Mercat, 2025).

Results

Overview

The literature search yielded 1558 articles, of which 99 were included in the final selection after the different screening steps (Figure 1). Two-thirds of the articles focused on mosquitoes and mosquito-borne diseases or pathogens (67.7%, 67/99). One third addressed ticks and tick-borne diseases or pathogens (29.3%, 29/99). The remaining 3 articles examined chigger mites and sand flies. The main results of each article are detailed in Additional file 4: Table S1. Ninety percent of the articles featured studies conducted in a single city and country. However, studies in six articles investigated multiple cities within the same country (Levytska et al., 2021; Overzier et al., 2013; Pedrosa et al., 2020; Schorn, et al., 2011a; Schorn, et al., 2011b; Yadouléton et al., 2010) , and three reported on studies carried out in cities of different countries (Carbó-Ramírez et al., 2017; Georganos et al., 2020; Talbot et al., 2021). The geographic distribution of studies varied (Figure 2A). Most mosquito-related articles originated from North America (37.3%, 25/67) and South America (28.4%, 19/67), whereas more than 70% of tick-related articles originated from Europe (72.4%, 21/29). The articles were published between 1999 and 2022, with a notable increase in mosquito-related articles after 2007 and tick-related articles after 2010 (Figure 2B).

Identification of studies via databases and registers

Identification

Date of search 5/09/22
Records identified from:
Databases (n = 2535)
  (Scopus n=775,
  Web of Science n=791,
  PubMed n=969)

Records removed before screening:
Duplicate records removed by :
Cadima automatic tools (n=972)
Authors (n=5)

Screening

Records screened on
title/abstract
(n = 1558)

Records excluded on title/abstract
(n = 997)

Records screened on full text
(n = 561)

Reason of exclusion on full text (n=463)
-    Not primary research
-    Not accessible
-    Not related to vector borne diseases
-    Not related to urban area
-    Not related to greening cities
-    No comparison between urban green spaces versus other spaces
-    No statistical analysis

Inclusion

Studies included in review
(n = 98 +1 = 99)

1 article added by the authors

281   **Figure 1** - PRISMA flow diagram for systematic review. It shows the number of
282   article identified through cross-checking references .

Figure 2 - Overview of the articles included. (A) Global distribution of articles (here a study refers to the investigation of a vector in a particular city, at a particular time, by a particular research team. For example, a single article may include two studies if the research was carried out in two different cities). An interactive map with the references of the studies (Bartholomée & Mercat, 2024) is available at

The selected articles were categorized based on their focus: those examining vectors alone
were classified under the 'vector data' category, while those studying pathogens within hosts or
vectors were classified under the 'vector-borne pathogen data' category (Table 1 and Figure 2C).
Among the 67 articles on mosquito-pathogen systems, 43 concentrated on mosquito data and 25
on mosquito-borne pathogens data. Only three articles addressed both mosquito and mosquito-
borne pathogens data: yellow fever virus and *Haemagogus leucocelaenus* and *Haemagogus*
janthinomys/capricornii (Abreu et al., 2022), dengue virus and *Aedes aegypti* (Francisco et al.,
2021), and malaria and *Anopheles gambiae* s.l. (Yadouléton et al., 2010). The most studied
mosquito genera were *Aedes* (35 articles), followed by *Culex* (13 articles) and *Anopheles* (three
articles). Associated mosquito-borne diseases included West Nile fever (10 articles), dengue fever
(six articles), and malaria (four articles). Of the 29 articles on tick pathogen systems, 19
investigated the effects of UGIs on tick vectors, 20 examined the effects of UGIs on tick-borne
pathogens, and 10 addressed both categories. Ticks from the *Ixodes* genus were studied the most
(18 articles), with a focus on *Ixodes ricinus* (14 articles). The most extensively studied pathogen
was the *Borrelia* spp. bacterium (11 articles) and small rodents were the most frequently studied
host group (seven articles).

Table 1: Classification of articles in accordance with the vector-pathogen system
and the UGIs under investigation

Group of vector-pathogen system and UGI description			Number
MOSQUITO-PATHOGEN SYSTEMS			67
Mosquito data			43
	Stage	Adult	25
		Immature	19
		Egg	8
	Species	*Aedes*	35
		Aedes aegypti	14
		Aedes albopictus	17
		Aedes hendersoni	1
		Aedes japonicas	1
		Aedes mediovittatus	1
		Aedes triseriatus	1
		Aedes vexans	1
		Culex	13
		Culex antillummagnorum	1
		Culex dolosus	1
		Culex pipiens	4
		Culex quinquefasciatus	6
		Culex restuans	3
		Culex salinarius	1
		Anopheles	3
		Anopheles gambiae s.l.	1
Mosquito-borne pathogen data			25
	Arbovirus		22
		Chikungunya virus – CHIKV	1
		Dengue virus – DENV	6
		Ross River virus – RRV	1
		Saint Louis Encephalitis virus - SLEV	1
		West Nile virus – WNV	10
		Yellow fever virus – YFV	1
	Avian haemosporidian parasites – *Avian haemosporidia*		1
	Malaria parasites – *Plasmodium* spp.		4
Mosquito and mosquito-borne pathogen data			3
	Dengue – *Aedes aegypti, Aedes albopictus*		1

	Malaria – *Anopheles gambiae* s.l.			1
	Yellow fever – *Haemagogus leucocelaenus, Haemagogus janthinomys/capricornii*			1
Pathogen diversity				2
	Avian haemosporidian parasites			1
	West Nile virus			1

TICK-PATHOGEN SYSTEMS				29
Tick data				19
	Stage	Adult		13
		Immature		18
	Species	*Amblyomma*		4
		Amblyomma americanum		1
		Dermacentor		2
		Dermacentor marginatus		1
		Haemaphysalis		5
		Haemaphysalis longicornis		1
		Ixodes		18
		Ixodes apronophorus		1
		Ixodes inopinatus		1
		Ixodes persulcatus		1
		Ixodes ricinus		13
		Ixodes scapularis		2
		Ixodes trianguliceps		1
		Rhipicephalus	*Rhipicephalus turanicus*	1
Tick-borne pathogen data				20
	Anaplasmosis – *Anaplasma phagocytophilum*			5
	Ehrlichiosis – *Ehrlichia* spp.			1
	Lyme-*Borrelia* spp.			11
	Piroplasmosis/Babesiosis – *Babesia* spp.			2
	Rickettsiosis - *Rickettsia* spp.			4
	Multiple tick-borne pathogens			4
Tick and Tick-borne pathogen data				10
	Anaplasmosis – *Ixodes ricinus*			2
	Lyme – *Ixodes ricinus*			7
	Lyme – *Ixodes scapularis*			1
	Rickettsiosis – *Ixodes ricinus*			1

OTHER VECTOR-PATHOGEN SYSTEMS				3
Sand flies				1
	Sand flies data	*Lutzomyia*	*Lutzomyia longipalpis*	1
Chigger mites				2
	Chigger mites-borne pathogen data	Human scrub typhus		1

TYPE OF UGI DESCRIPTION				
Urban green space typology				64
	Mosquito-pathogen systems			37
	Tick-pathogen systems			23
	Other vector-pathogen systems			3
Description of urban vegetation				58
	Mosquito-pathogen systems			44
	Tick-pathogen systems			11
	Other vector-pathogen systems			1

UGIs were described using the urban green space typology or the urban vegetation around the
sampling points (Table 1, Box 2). Both approaches were commonly used for mosquito-pathogen
systems (Figure 3), appearing in 44 and 37 articles out of 67, respectively. In contrast, for tick-
borne pathogen systems articles (Figure 4), the urban green space typology was the dominant
approach used in 21 out of 29 articles. Below, the impact of UGIs on mosquito-pathogen systems,
then on tick-pathogen systems and finally on other vector-pathogen systems is examined.

Figure 3 - Impact of UGIs on mosquito-pathogen systems. (A) Impact of UGIs on mosquito populations. (B) Impact of UGIs on mosquito-borne pathogen (prevalence of infection in host or vector). Numbers in brackets indicate the total number of variables tested and the total number of articles from which these variables were extracted. Colors represent the proportion of variables in all studies that increased risk (harmful effect, red), decreased risk (beneficial effect, blue), varied (variable effect, yellow), or showed no association (grey) with risk.

Figure 4 - Impact of UGIs on tick-pathogen systems. (A) Impact of UGIs on tick populations. (B) Impact of UGIs on tick-borne pathogen (prevalence of infection in host or vector). Numbers in brackets indicate the total number of variables tested and the total number of articles from which these variables were extracted. Colors represent the proportion of variables in all studies that increased risk (harmful effect, red), decreased risk (beneficial effect, blue), varied (variable effect, yellow), or showed no association (grey) with risk.

Mosquito-pathogen system

For each mosquito genus, we first addressed the effects of UGIs on mosquito populations, followed by their influence on mosquito-borne pathogens, and we concluded with overarching insights into the impact of UGIs on the specific mosquito-pathogen system relationship.

Effects on Aedes-pathogen system

Effects on *Aedes* populations

Of the 35 articles studying the genus *Aedes*, 17 focused on *Aedes albopictus* and 14 on *Aedes aegypti* (Table 1). *Aedes vexans*, *Aedes mediovittatus*, *Aedes japonicus*, *Aedes triseriatus* and *Aedes hendersoni* each appeared in a single article. In this section, we mainly describe the results for the two major disease vectors, *Ae. aegypti* and *Ae. albopictus*.

UGIs appeared to be unfavorable habitats for *Ae. aegypti* (Figure 5A). The impact of UGIs on *Ae. aegypti* was mainly assessed using urban vegetation description indicators rather than urban green space typology indicators (Figure 5A), with most studies focusing on immature stages (78.6%, 11/14). Consistently, higher numbers of *Ae. aegypti* immature stages and adults have been recorded in densely urbanized areas compared to UGIs in several locations in Brazil. This trend has been observed using urban vegetation description indicators, such as in Rio de Janeiro (Honório et al., 2009) and in São Paulo (Arduino et al., 2020), and with urban green spaces typology indicators, such as in Mariana, Ouro Preto (Pedrosa et al., 2020) and São Paulo (Lorenz et al., 2020).

Using urban vegetation description indicators, locations with a high percentage of vegetation coverage, like urban forests or urban grass in Singapore (Sun et al., 2021) or arboreal vegetation in Argentina (Benitez et al., 2020) exhibited low mosquito abundance. Moreover, in New Orleans (United States of America, USA) (de Jesús Crespo & Rogers, 2021) cemeteries with a higher heat index supported higher mosquito populations than cemeteries with lower heat index. Reduced temperature in vegetated areas may explain why *Ae. aegypti* was less commonly found in areas with more vegetation cover (de Jesús Crespo & Rogers, 2021).

Using urban green space typology indicators, *Ae. aegypti* oviposition rates were observed to be higher (i) near houses compared to deeper forested areas in Rio de Janeiro (Lourenço-de-Oliveira et al., 2004), and (ii) in the peripheral part of the park with less vegetation near urban areas compared to the densely vegetated inner sections of the park (Heinisch et al., 2019). Also, other countries, including the USA (Puerto Rico) (Cox et al., 2007) and Vietnam (Huynh & Minakawa, 2022), have shown similar results. Breeding sites could be associated with residential areas, where artificial breeding containers like jars or planters were common (Huynh & Minakawa, 2022).

Despite the general negative association between *Ae. aegypti* and UGIs, the mosquito still thrived in areas with vegetation near breeding sites, even in densely populated environments. For instance, using the urban vegetation description, Estallo et al. (2018) observed in Córdoba (Argentina) that vegetation adjacent to breeding sites was positively associated with the likelihood of these sites supporting mosquito larvae. In the same city, Andreo et al. (2021) found that the distance to vegetated areas could have a variable impact on the presence of breeding sites depending on the month, from decreasing their presence in November to increasing it in March. A study by Landau & van Leeuwen (2012) found that *Ae. aegypti* adult abundance in Tucson (USA) increased with the percentage of medium to tall trees within 10 to 50 meters of trap locations. No association was observed with herbaceous or shrub vegetation (Landau & van Leeuwen, 2012). Using the urban green space typology, Sun et al. (2021) observed that adult *Ae. aegypti* were more abundant in areas in Singapore with buildings and managed vegetation than in zones with dense forest cover. In contrast to *Ae. aegypti,* UGIs generally supported *Ae. albopictus* populations, although the effects varied depending on the specific relationship between mosquito stage and UGI studied, using urban vegetation description and urban green space typology indicators (Figure 5B).

387

 .

389
Figure 5 - Impact of UGIs on *Aedes*-pathogen systems. (A) Impact of UGIs on
Aedes aegypti populations. (B) Impact of UGIs on *Aedes albopictus* populations.
(C) Impact of UGIs on Dengue virus dynamics (prevalence of infection in host or
vector). Numbers in brackets indicate the total number of variables tested and the total number
of articles from which these variables were extracted. Colors represent the proportion of variables
in all studies that increased risk (harmful effect, red), decreased risk (beneficial effect, blue), varied
(variable effect, yellow), or showed no association (grey) with risk.

Aedes albopictus eggs and larvae were more abundant in urban green spaces near residential
areas than in densely urbanized zones. This trend has been observed in Brazilian cities using
urban green space typology indicators, such as in Mariana and Ouro Preto (Pedrosa et al., 2020)
and São Paulo (Heinisch et al., 2019; Wilke et al., 2017), as well as using urban vegetation
description indicators, such as in Rio de Janeiro (Honório et al., 2009; Lourenço-de-Oliveira et al.,
2004) and São Paulo (Arduino et al., 2020). Urban green spaces can support more immature
mosquito stages than more urbanized spaces. For instance, in Miami (USA), urban farms hosted
more *Ae. albopictus* larvae compared to densely built areas (Wilke et al., 2020). Similarly, in
Greensboro (USA) urban forests and urban parks yielded more *Ae. albopictus* eggs than
campuses (Schwarz et al., 2020). Urban vegetation significantly supported the presence of
breeding sites. Examples include the percentage of urban vegetation linked to egg abundance in
Italy (Cianci et al., 2015) and the vegetation shade associated with breeding sites in Vietnam
(Huynh & Minakawa, 2022). Francisco et al. (2021) also found a positive correlation between egg
abundance and the percentage of urban vegetation in the Philippines. Conversely, a study in New
Orleans (de Jesús Crespo & Rogers, 2021) reported no association between the presence of
vegetation next to breeding sites and the presence of larvae. Positive correlations between tree
cover and mosquito breeding may be explained by specific factors. Tree holes, for example, serve
as breeding sites for *Aedes* spp. mosquitoes, such as those found in *Delonix regia* and *Magnifera
indica* trees in Bangladesh (Chowdhury et al., 2014), or, for *Ae. albopictus* eggs in tree canopies
in Greensboro (Schwarz et al., 2020). Trees can also create a favorable microclimate for mosquito
development by mitigating urban heat island effects, reducing temperatures, and providing shelter
(de Jesús Crespo & Rogers, 2021). Some breeding sites, like water-filled flower vases, were
associated with gardening practices (Che Dom et al., 2016).

Regarding the abundance of *Ae. albopictus* adults, a similar trend emerged when using urban green space typology or urban vegetation description indicators. In Vietnam, urban parks hosted higher adult abundance than residential zones (Huynh & Minakawa, 2022). As observed in Brazil (Pedrosa et al., 2020), green spaces near residential zones also supported higher mosquito populations than in densely urbanized areas. In Malaysia's Klang Valley, urban farms had higher adult mosquito abundance than peri-urban farms (Lee et al., 2020). In Hong Kong (China), no differences in adult mosquito abundance were found between sites with green or bare roofs (Wong & Jim, 2018). Using the urban vegetation description, adult *Ae. albopictus* populations were generally more abundant in forested areas than in grassy areas, as observed in Singapore (Sun et al., 2021). Dense or poorly maintained vegetation can reduce the abundance of adults (Hendy et al., 2020; Sun et al., 2021). In Rome (Italy), Manica et al. (2016) found that adults preferred small urban green spaces, with a positive correlation between abundance and vegetation within 20 meters of the sampling trap and a negative correlation with vegetation at 300 meters. A potential explanation for the positive correlation between urban vegetation and *Ae. albopictus* adults is the role of certain plant species. Waxleaf Privet (*Ligustrum quihoui*) has been found to promote mosquito survival, while Glossy Abelia (*Abelia grandiflora*) enhanced female fecundity (Tian, 2019). However, in Manaus (Brazil), Hendy et al. (2020) reported no association between tree cover and adult abundance. The specific and context-dependent vegetation characteristics, such as the presence of certain plant species discussed here, played a critical role in mosquito abundance.

Articles addressing both *Ae. albopictus* and *Ae. aegypti* in UGIs where both species co-exist confirmed the contrasting effects of UGIs on these two mosquito species as presented in the section above (Arduino et al., 2020; de Jesús Crespo & Rogers, 2021; Heinisch et al., 2019; Honório et al., 2009; Huynh & Minakawa, 2022; Pedrosa et al., 2020; Sun et al., 2021). However, some articles found similar trends for both mosquito species distribution in UGIs. The abundance of immature stages of both species was positively correlated with the presence of vegetation adjacent to breeding sites, as observed in Brazil (Honório et al., 2009), Bangladesh (Paul et al., 2018), Malaysia (Che Dom et al., 2016), and the Philippines (Francisco et al., 2021). Moreover, a study from Rio de Janeiro observed a similar trend for both species, showing a decrease in the abundance of their immature stages with increasing distance to the peri-urban forest (Lourenço-de-Oliveira et al., 2004). In addition, a multi-country study from South America reported that the abundance of both species was positively correlated with decorative vegetation and the proximity of urban vegetation to residential areas (Talbot et al., 2021). In Bangladesh, *Aedes* spp. *(Ae. albopictus* and *Ae. aegypti*) breeding site positivity was associated with vegetation located near breeding sites (Paul et al., 2018).

Some studies have also explored the effects of UGIs on other *Aedes* species. For instance, *Aedes mediovittatus* larvae were more abundant in low-density residential areas, with over 20% vegetation cover, compared to forest patches and densely urbanized residential areas in Puerto Rico (Cox et al., 2007). In New Haven (USA), more *Aedes vexans* adults were found in moderately vegetated residential areas than in highly or un-vegetated areas (Brown et al., 2008). For *Aedes japonicus*, adult abundance showed no difference between green and bare roofs in Hong Kong (Wong & Jim, 2018). *Aedes triseriatus* and *Aedes hendersoni* eggs were more abundant in urban forests than on urban campuses in Greensboro (Schwarz et al., 2020).

Effects on dengue virus dynamics

Dengue is an arboviral disease in which the virus (DENV) is transmitted primarily by *Aedes* mosquitoes, particularly *Ae. aegypti* and *Ae. albopictus*. Six articles examined the relationship between dengue incidence and UGIs (Table 1), revealing varying effects across locations and periods (Figure 5C). Only one of them used urban green space typology indicators and observed that the impact of urban green spaces on dengue incidence in humans varied by city in China, with effects ranging from no association to increased risk (Chen et al., 2020). Using urban vegetation indicators, the same authors found that dengue incidence was positively correlated with forest coverage in two Chinese cities (Chen et al., 2020). Three other studies reported a negative

correlation between UGIs and dengue incidence. First, Zellweger et al. (2017) reported that
vegetation cover was either negatively correlated or not correlated with human cases, depending
on the epidemic (2008-2009 versus 2012-2013) in Nouméa (New Caledonia). In São Paulo, Araujo
et al. (2015) found higher human dengue incidence in areas with low to moderate vegetation cover
compared to dense vegetation zones, a pattern also observed in Manila (the Philippines), where
vegetation was negatively associated with dengue incidence (Francisco et al., 2021). The last two
studies used remote sensing indices like NDVI, the weekly Enhanced Vegetation Index (EVI), and
the Normalized Difference Water Index (NDWI) to assess the impact of vegetation on human
dengue and reported inconsistent correlations, with no clear trend emerging (Ogashawara et al.,
2019; Troyo et al., 2009).
484
Effects on Chikungunya virus dynamics
Chikungunya virus (CHIKV) is primarily transmitted by *Aedes* mosquitoes, particularly *Ae.*
aegypti and *Ae. albopictus*. To date, only one study explored the link between UGIs and CHIKV
dynamics and showed that, in Rio de Janeiro, where *Ae. aegypti* serves as the vector, a higher
proportion of green parks was associated with reduced CHIKV incidence in humans (Freitas et al.,
2021).
491
Concluding remarks on *Aedes*–pathogen systems
The impact of UGIs on *Aedes* mosquito populations differed between the two major arbovirus
vector species: *Ae. aegypti* was more abundant in densely urbanized areas with less vegetation
than in vegetated areas, while *Ae. albopictus* tended to be supported by UGIs, with variations
depending on mosquito life stages and the type of UGI studied. Results showed no consistent
trends regarding the effects of UGIs on dengue dynamics. This inconsistency may stem from
DENV being transmitted by both *Ae. aegypti* and *Ae. albopictus*, which responded differently to
UGIs. Identifying the primary vector in a given context is essential to assess dengue risk effectively.
Notably, Francisco et al. (2021) in Manila was the only study that examined the influence of UGIs
on both *Aedes* abundance and dengue incidence in humans. Their findings revealed a positive
correlation between urban vegetation and mosquito eggs abundance but a negative correlation
with dengue incidence in humans, underscoring the challenge of estimating VBD risk based solely
on vector abundance.
505

506 *Effects on Culex-pathogen systems*
507
Effects on *Culex* populations
Of the 13 articles reporting on the effect of UGIs on *Culex* mosquitoes, six focused on *Culex*
quinquefasciatus (Cox et al., 2007; de Jesús Crespo & Rogers, 2021; Landau & van Leeuwen,
2012; Sallam et al., 2017; Wilke et al., 2020; Wong & Jim, 2018), four on *Culex pipiens* (Brown et
al., 2008; DeMets et al., 2020; Gardner et al., 2013; Trawinski & Mackay, 2010) and three on *Culex*
restuans (Brown et al., 2008; DeMets et al., 2020; Gardner et al., 2013) (Table 1, Figure 6A and
Figure 6B).

Figure 6 - Impact of UGIs on *Culex*-pathogen system. (A) Impact of UGIs on *Culex quinquefasciatus* populations. (B) Impact of UGIs on *Culex pipiens* and *Culex restuans* populations. (C) Impact of UGIs on West Nile virus dynamics (prevalence of infection in host or vector). Numbers in brackets indicate the total number of variables tested and the total number of articles from which these variables were extracted. Colors represent the proportion of variables in all studies that increased risk (harmful effect, red), decreased risk (beneficial effect, blue), varied (variable effect, yellow), or showed no association (grey) with risk.

Regarding *Cx. quinquefasciatus*, the diversity of urban green vegetation description indicators studied made it difficult to identify a clear trend, regardless of the type of indicator used to describe UGIs (Figure 6A). When considering urban green space typology, Cox et al. (2007) found that *Cx. quinquefasciatus* larvae were more abundant in low-density residential areas with over 20% vegetation cover, compared to forest patches and densely urbanized residential areas in Puerto Rico. Urban agriculture positively influenced the abundance of *Cx. quinquefasciatus* in Miami, where Wilke et al. (2020) reported higher numbers of adults and immature stages in two urban farms than in other urban areas, particularly in farms surrounded by densely urbanized zones. A specific study on green roofs in Hong Kong found that there were less *Cx. quinquefasciatus* males near green roofs than near bare roofs (Wong & Jim, 2018). Using urban vegetation description, *Cx. quinquefasciatus* was mainly positively associated with the percentage of medium and high tree height between 10 and 50 m around the sampling traps, compared to the percentage of herbaceous and shrub vegetation in Tuscon (USA) (Landau & van Leeuwen, 2012). In New Orleans, Sallam et al. (2017) observed higher vector-host contact in urban non-forest wetlands compared to forest wetlands which was positively associated with tree density. In another study in New Orleans,de Jesús Crespo & Rogers (2021) found that *Cx. quinquefasciatus* larvae were more often located in cemeteries with open canopies than in cemeteries with closed canopies.

The impact of UGIs on *Cx. pipiens* and *Cx. restuans* did not reveal any clear trend as well Figure 6B. Urban vegetation, measured indirectly by NDWI, was positively correlated with the abundance of *Cx. pipiens* and *Cx. restuans* in Toronto (Canada) (DeMets et al., 2020), but showed the opposite effect or no correlation in Amherst (USA) (Trawinski & Mackay, 2010).) DeMets et al. (2020) noted that NDVI was not associated with the abundance of *Cx. pipiens* and *Cx. restuans*. In Chicago (USA), non-flowering shrubs and specific tree species, such as oak and pear, favored the abundance of immature stages of *Cx. pipiens* and *Cx. restuans*, while flowering shrubs and

ash trees had the opposite effect (Gardner et al., 2013). Additionally, deciduous tree density
showed no impact on abundance. When considering urban green spaces within residential areas,
Cx. pipiens and *Cx. restuans* preferred moderately vegetated residential areas over residential
areas with sparse or dense vegetation, as observed in New Haven (USA) (Brown et al., 2008).
Some studies have also explored the effects of UGIs on other *Culex* species. *Culex salinarius*
females were more abundant in highly vegetated residential areas than in sparsely or moderately
vegetated areas in New Haven (Brown et al., 2008). In Rio de Janeiro, *Cx. dolosus* was more
commonly found deeper in urban forests than near residential areas (Lourenço-de-Oliveira et al.,
2004). Similarly, *Cx. antillummagnorum* in Puerto Rico was more abundant in urban forests and
low-density residential areas, with over 20% vegetation cover, than in densely urbanized
residential areas (Cox et al., 2007). Additionally, in Hanoi (Vietnam), the abundance of female *Cx.*
vishnui and *Cx. tritaeniorhynchus* was higher in urban rice fields than in urban forest areas (Ha et
al., 2021). A study in Cleveland (USA) examined *Culex* mosquitoes without identifying them to
species level (Yang et al., 2019). In this study, adult mosquito abundance was positively correlated
with grass and shrub cover, while tree canopy coverage had no significant effect. The same study
found that vegetation diversity, biomass, and the presence of vacant lots did not influence *Culex*
abundance.
565
Effects on West Nile virus dynamics
West Nile virus (WNV) is primarily transmitted by *Culex* spp. mosquitoes and circulates within
bird populations, particularly among corvids (crows, blue jays, etc.) (Chancey et al., 2015; Levine
et al., 2016, 2017; Liu et al., 2011). Humans and horses serve as dead-end hosts. Ten articles
examined the effects of UGIs on WNV dynamics, all of which were conducted in North America
(Table 1). The impact of UGIs on WNV dynamics was mainly explored using urban green space
typology indicators compared to urban vegetation indicators (Figure 6C).
Eight of the ten studies focused on the effect of UGIs on WNV infection in *Culex* spp (Levine
et al., 2016, 2017; Liu et al., 2008; Liu & Weng, 2012; Lockaby et al., 2016; Myer & Johnston,
2019; Poh et al., 2020; Talbot et al., 2019). One study, conducted in Indianapolis reported a
positive association between mosquito infection and UGI in the form of urban agriculture using
green space typology indicators (Liu et al., 2011). Urban agriculture was suggested to offer stable
temperatures and moisture conditions more suitable for mosquito breeding (Liu et al., 2008). Four
studies highlighted that UGIs may have no effect on mosquito infection rates (Figure 6C), including
three using urban green space typology indicators. In Ottawa (Canada), Talbot et al. (2019) found
no relationship between the proportion of urban park areas and WNV infection rates. In Atlanta
(USA), Levine et al. (2016, 2017) found that mosquito infection rates were generally similar across
urban green spaces (urban wooded parks or urban parks with water, urban forests, and urban
zoos), though infected mosquitoes were more prevalent in wooded parks than in zoos. The fourth
study, conducted in Los Angeles (USA), used urban vegetation indicators and observed no
association between mosquito infection rates and the urban vegetation (described using NDVI),
possibly due to the cross-correlation of NDVI with higher elevation, which is less conducive to
mosquito habitats (Liu & Weng, 2012). In contrast, three studies found a negative correlation
between UGIs and mosquito infection, all using urban vegetation indicators (Figure 6C). Lockaby
et al. (2016) found a negative correlation between the probability of mosquito infection and both
the percentage of pine forests—likely due to drier conditions—and the area of forest patches,
possibly explained by a dilution effect. According to the authors, large forest patches were
associated with high avian species diversity, which reduced the probability of contact between
infected *Culex* mosquitoes and competent host species. Another study in Atlanta by Myer &
Johnston (2019) highlighted a negative correlation between NDVI and mosquito infection, a trend
also observed in Dallas (USA) by Poh et al. (2020).
Five studies investigated the association between WNV incidence/seroprevalence in hosts and
UGIs, focusing on birds and humans (Figure 6C). One study in Northern Virginia (USA) found a
positive correlation between urban vegetation indicators and WVN incidence in corvids, with a
peak of infection in medium and high canopy density areas compared to impervious surfaces with

lower canopy density (Liu et al., 2011). In contrast, four studies identified a negative or no association between UGIs and WNV incidence (Figure 6C), three using urban green space typology indicators and one using urban vegetation indicators. In Atlanta, WNV seroprevalence was higher in passerine birds in residential areas than in more vegetated areas and lower in urban forests compared to parks with wooded and paved areas (Levine et al., 2016, 2017). In Cleveland, WNV incidence in humans was lower in urban green spaces compared to residential areas (LaBeaud et al., 2008). Finally in Ottawa, no association was found between the proportion of green parks and human WN case incidence (Talbot et al., 2019).

Two studies addressed WNV genetic diversity in *Culex* spp. and bird hosts. In mosquitoes, the genetic diversity of WNV decreased or remained unchanged within urban green spaces (Amore et al., 2010; Bertolotti et al., 2008), while in birds, WNV genetic diversity was higher in urban green spaces compared to residential areas (Bertolotti et al., 2008). According to the authors, this higher diversity in urban green spaces may be explained by the greater diversity of bird species associated with a higher availability of natural habitats in urban green spaces compared to residential areas.

Effects on other *Culex*-borne pathogens

A study in Argentina found that prevalence in humans of Saint Louis encephalitis virus, transmitted by *Cx. quinquefasciatus*, was lower in low-density residential areas than in densely populated regions (Vergara Cid et al., 2013). Similarly, in Brisbane (Australia), incidence of Ross River virus in humans, a virus transmitted by *Aedes vigilax* and *Culex annulirostris*, was positively associated with the percentage of riparian and bushland areas (Muhar et al., 2000).

Concluding remarks on *Culex*-pathogen systems

The diversity and specificity of UGI indicators made it challenging to discern clear trends regarding their impact on *Culex* mosquitoes. For WNV dynamics, findings indicated that UGIs often negatively affected mosquito infection rates, though their effects can vary significantly depending on urban and environmental contexts. Additionally, UGIs tended to reduce or have no significant effect on WNV incidence in birds as well as in human hosts compared to impervious residential areas. The dynamics of WNV transmission remains complex and influenced by various factors, including plasticity in the feeding behavior of mosquitoes, host availability, and diversity in the ability of bird host species to amplify and further transmit the virus to the vector mosquitoes (Levine et al., 2016, 2017).

Effects on Anopheles-pathogen systems

Effects on *Anopheles* populations

Only three of the selected articles focused on *Anopheles* spp. mosquitoes, all indicating a positive correlation between urban agriculture and the presence or abundance of *Anopheles* mosquitoes and mainly using urban green space typology indicators (Dongus et al., 2009; Matthys et al., 2010; Yadouléton et al., 2010). Dongus et al. (2009) reported a higher presence of *Anopheles* larvae in urban crop cultivation areas, such as sugarcane and leafy vegetable farms, and in urban agricultural zones compared to non-agricultural areas in Dar es Salaam (Tanzania). The presence of wells or irrigation systems was also positively correlated with the probability of presence of *Anopheles* larvae (Dongus et al., 2009). Similarly, Matthys et al. (2010) found *Anopheles* larvae most frequently in agricultural trenches over other irrigation types in urban farming in Abidjan (Côte d'Ivoire). In Benin, Yadouléton et al. (2010) reported higher human biting rates near vegetable farms compared to households located farther away from these agricultural zones across several cities.

Effects on *Plasmodium* dynamics

Malaria is a parasitic disease caused by *Plasmodium* spp., which is transmitted between human hosts by *Anopheles* spp. mosquitoes. Four of the 99 selected studies explored the dynamics of malaria related to UGIs (Georganos et al., 2020; Kabaria et al., 2016; Matthys et al., 2006; Yadouléton et al., 2010), including two that examined the relationship with urban green space typology indicators and two with urban vegetation indicators. In Benin, Yadouléton et al. (2010) found that *Anopheles* spp*.* infection rate was higher in households close to vegetable farms than in households farther away. Similarly, malaria prevalence in the human population of Côte d'Ivoire was higher in urban agriculture areas (Georganos et al., 2020), particularly within mixed crop systems compared to traditional smallholder rice plots (Matthys et al., 2006). The prevalence of malaria was also positively linked to riparian vegetation in Tanzania and Uganda (Georganos et al., 2020) and to the percentage of urban vegetation based on NDVI in Tanzania (Kabaria et al., 2016).

Concluding remarks on *Anopheles*-pathogen systems
These results showed a positive association between UGIs, specifically urban agriculture, for both *Anopheles* spp. abundance as well as malaria prevalence in humans, and in the vector mosquitoes.

Effects on Haemagogus and on yellow fever dynamics
In Rio de Janeiro, Abreu et al. (2022) studied the relationships between UGIs and the yellow fever vector system, which involves the vectors *Haemagogus leucocelaenus* and *Haemagogus capricornii*, as well as human and non-human primates as hosts. The study found no difference in vector abundance between urban forests, urban pastures, and densely urbanized areas, but the abundance of vectors was positively associated with the NDVI. Furthermore, the prevalence of yellow fever in humans and non-human primates was higher in urban forest fragments than in heavily urbanized areas. Both *Hg. leucocelaenus* and *Hg. capricornii* showed higher infection rates for yellow fever virus (YFV) in urban forest settings compared to other urban areas.

Effects on other mosquito-pathogen systems
Only four studies evaluated the influence of UGIs on the abundance of all mosquitoes collected. In São Paulo, Medeiros-Sousa et al. (2017) found a positive association between mosquito abundance and urban parks. Another study in Sinop (Brazil) reported higher abundance of adult mosquitoes in urban forested parks and in urban forest fragments than in urbanized areas (Vieira et al., 2020). In Hong Kong, the abundance of all adult species was lower on green roofs compared to bare roofs (Wong & Jim, 2016) and low-elevation gardens (Wong & Jim, 2017). One multi-site study indicated that, in both Mexico and Germany, there was no association between seroprevalence of avian haemosporidia in birds and urban forests (Carbó-Ramírez et al., 2017).

Concluding remarks on the impact of UGIs on mosquito-pathogen systems
Globally, the influence of UGIs on mosquito populations and mosquito-borne disease transmission remain inconclusive and context specific: UGIs have been associated with increased, decreased, or negligible effects on vector risk, showing no consistent trend across cities and vector systems. Further research is urgently needed to clarify the effects of UGIs on mosquito vector populations dynamics and associated mosquito-borne disease risk in cities (Figure 3, Additional file 5: Table S2, Additional file 6: Table S3).

Effects of UGIs on tick-pathogen systems

Effects on Ixodes-pathogen system

Effects on *Ixodes* populations

Across 19 articles on UGIs and tick populations, thirteen articles centered on *Ixodes ricinus* in
Europe and two on *Ixodes scapularis* in North America (Gregory et al., 2022; VanAcker et al.,
2019). Six of the thirteen articles observed a positive correlation between UGIs and *I. ricinus*
population density, with one using urban green space typology indicators and five using urban
vegetation indicators (Figure 7A). Higher tick densities may be linked to greater connectivity
between urban and peri-urban environments, allowing movement of the host, as observed in
Antwerp (Belgium) (Heylen et al., 2019). Using urban vegetation indicators, *I. ricinus* was more
abundant at woodland edges and within forests than in grasslands, as demonstrated in Salisbury
(United Kingdom, UK) (Hansford et al., 2017), London (UK) (Hansford et al., 2021), and Rome (Di
Luca et al., 2013), as well as in urban parks in Bath (UK) (Hansford et al., 2022). Additionally, *I.*
ricinus was more prevalent in deciduous or mixed forests than in coniferous forests, as noted in
Poznań (Poland) (Michalik et al., 2003). This habitat preference can be explained by the presence
and the abundance of small mammals (e.g., hedgehogs, *Apodemus* spp.) (Borşan et al., 2020;
Michalik et al., 2003) and larger mammals such as white-tailed deer (Hansford et al., 2017) which
serve as hosts for ticks.

In contrast, eight articles reported no consistent trend between UGIs and tick population density
(Figure 7A), with seven articles using urban green space typology indicators and one using urban
vegetation indicators. Studies conducted in Lyon (France) (Mathews-Martin et al., 2020), Helsinki
(Finland) (Junttila et al., 1999), Cluj-Napoca (Romania) (Borşan et al., 2020), and Warsaw
(Poland) (Welc-Falęciak et al., 2014) revealed differences in *I. ricinus* abundance between urban
green spaces, like urban parks, though these studies did not explain the observed variations. Other
articles conducted in St Petersburg (Russia) (Tretyakov, 2009) and in Olsztyn (Poland) (Kubiak et
al., 2019) found no significant differences in vector abundance across urban parks and urban
forests. In Budapest (Hungary), Hornok et al. (2014) observed that tick abundance could vary by
the month of sampling, sometimes being higher in cemeteries than in urban forests or parks or
showing no differences. Finally, a study in Bath revealed no differences in tick vector abundance
between woodland and grassland or between mixed and deciduous forests (Hansford et al., 2022).

In the two articles on *I. scapularis* in the city of New York (USA), urban green space connectivity
and host accessibility were important in determining tick populations density. Enhanced
connectivity, measured by park-to-park links (VanAcker et al., 2019), was correlated with increased
tick numbers. Permeable garden edges did not seem to raise the probability of tick
detection (Gregory et al., 2022). High canopy cover around parks (VanAcker et al., 2019) or
gardens (Gregory et al., 2022) was associated with increased tick counts, while lower tick numbers
were found in grasslands or bare soil.

Other *Ixodes* species, such as *I. persulcatus, I. trianguliceps,* and *I. apronophorus*, have been
observed in urban parks, such as in St. Petersburg (Tretyakov, 2009). In Hanover (Germany),
Hauck et al. (2020) reported that *I. ricinus* and *I. inopinatus* densities were higher in urban mixed
forests than in urban parks, with no difference between deciduous and mixed forests and between
mixed forests and urban parks.

739
Figure 7 - Impact of UGIs on *Ixodes*-pathogen systems. (A) Impact of UGIs on
Ixodes ricinus populations. (B) Impact of UGIs on Lyme bacteria *Borrelia* spp.
(prevalence of infection in host or vector). Numbers in brackets indicate the total number
of variables tested and the total number of articles from which these variables were extracted.
Colors represent the proportion of variables in all studies that increased risk (harmful effect, red),
decreased risk (beneficial effect, blue), varied (variable effect, yellow), or showed no association
(grey) with risk.
747

Effects on *Borrelia* spp. dynamics
Eleven articles focused on *Borrelia spp.* (Table 1)*,* the bacterial pathogens causing Lyme
disease, primarily transmitted by *Ixodes* ticks. The presence of infected ticks is closely linked to
the movement and presence of competent hosts, such as the striped field mouse (*Apodemus*
agrarius), a major reservoir for *Borrelia burgdorferi* (Gryczyńska et al., 2021). The effects of UGIs
on tick infection rates were primarily studied using urban vegetation indicators compared to urban
green space typology indicators (Figure 7B).
Five articles reported a positive correlation between UGIs and *Borrelia* dynamics, with two
relying on urban green space typology and three on urban vegetation indicators (Figure 7B). In
Budapest (Hungary), the *Ixodes* infection rates were higher in urban forests than in cemeteries
(Hornok et al., 2014). In New York city, VanAcker et al. (2019) demonstrated that higher
connectivity among urban green spaces—measured by a park's contribution to network
connectivity—increased tick infection rates. Patterns of the effects of urban vegetation indicators
on *Borrelia* infection closely aligned with the impact on *Ixodes* tick abundance. Infection rates were
higher in urban woodlands compared to grasslands in London (Hansford et al., 2021) and at
woodland edges compared to grasslands in Bath (Hansford et al., 2022). In New Castle county
(USA), *Rosa multiflora* presence in urban forests was associated to higher infection rates in ticks
(Adalsteinsson et al., 2018), as did woody debris compared to leaf litter.
Seven articles, including four using urban green space typology and three employing urban
vegetation indicators, found no link between UGIs and *Borrelia* dynamics (Figure 7B). Urban parks
were not significantly associated with *Borrelia* spp. infection rates in ticks in some European cities
including Prague (Czech Republic), Helsinki, Budapest and Olsztyn (Hornok et al., 2014; Junttila
et al., 1999; Kubiak et al., 2019; Kybicová et al., 2017). There was no significant difference in ticks'
infection rates between grasslands and woodlands in Salisbury (Hansford et al., 2017) and in Bath
(Hansford et al., 2022), or between coniferous and deciduous forests in Poznań (Michalik et al.,
2003).
774
775 Effects on other tick-borne pathogens dynamics

Four studies focused on the *Rickettsia/I. ricinus* vector-pathogen system in various cities in Germany cities (May & Strube, 2014; Overzier et al., 2013; Schorn, et al. 2011a) and Poland cities (Welc-Falęciak et al., 2014). They found no differences in ticks' infection rates within urban green spaces, without offering specific explanations for these results. Five studies investigated the relationship between UGIs and the *Anaplasma phagocytophilum/I. ricinus* system. Infection rates for *An. phagocytophilum* varied within and between German cities' green spaces (May & Strube, 2014; Schorn et al., 2011b). However, studies from Hungary and Poland (Hornok et al., 2014; Welc-Falęciak et al., 2014) found no significant differences between different urban green spaces. Infection rates were higher in ticks in urban parks than in peri-urban parks in Prague (Kybicová et al., 2017) and higher in urban parks than in cemeteries in Budapest (Hornok et al., 2014). One study explored the *Babesia/I. ricinus* system and reported no association between ticks' infection rates and urban green spaces in Germany (Schorn, et al. 2011a).

Two studies conducted in Ukraine (Levytska et al., 2021) and in Belgium (Heylen et al., 2019) examined multiple pathogens (*Rickettsia spp.*, *Babesia spp.*, *Bartonella spp.*, and *Anaplasma phagocytophilum*) collectively. The findings indicated that and infection rates with these tick-borne pathogens were not affected by urban green spaces or their connectivity.

Concluding remarks on *Ixodes*-pathogen systems

The abundance of *Ixodes* ticks was influenced by host movement, likely shaped by connectivity between urban green spaces and by local vegetation characteristics, with ticks exhibiting a preference for deciduous or mixed forests. Variations in *Ixodes* tick abundance across different urban green spaces were location-specific and depended on the time of year. *Borrelia* infection rates in ticks were strongly linked to the presence and movement of competent hosts, particularly in woodlands or connected urban green spaces, reflecting the influence of UGIs on *Ixodes* populations. However, the effect of urban green spaces on ticks' infection rate varied by location and specific characteristics, which may explain why some UGIs have minimal impact. For other tick-borne pathogens, infection rates in ticks did not exhibit any consistent pattern in relation to UGIs.

Effects on Amblyomma-pathogen system

Only one study focused on *Amblyomma americanum*. This tick was positively associated with urban green spaces, particularly with permeable garden edges and increased canopy cover around yards in New York city (Gregory et al., 2022). The presence of log and brush piles in the yard also favored the probability of tick presence, which is not influenced by the presence of low canopy and grass around the yard.

Two studies reported results on other *Amblyomma*–pathogen systems. In Oklahoma (USA), infection rates of *A. americanum* with *Rickettsia spp.* or *Ehrlichia spp.* were comparable between residential and non-residential areas (Small & Brennan, 2021). Additionally, Noden et al. (2022) found that the percentage of developed land surrounding sampling points did not affect infection rates of *A. americanum* with *Borrelia spp.*, *Rickettsia spp.*, or *Ehrlichia spp*. Altogether, these results highlighted that *Amblyomma* tick populations were positively associated with the presence of urban green spaces, without evidence of differences in pathogen infection rates.

Effects on other tick species populations

Only one study focused on *Haemaphysalis longicornis* in New York city, showing that its presence was positively associated with logs and brush piles in yards, while it was unaffected by the permeable edge or the surrounding land cover of sampled yards (Gregory et al., 2022). In Italy, *Dermacentor marginatus* and *Rhipicephalus turanicus* were more prevalent in open landscapes, including pastures and wheat fields, than in deciduous woodlands (Di Luca et al., 2013).

Four studies explored the relationship between UGIs and various tick species from different genera. In Romania, *I. ricinus* and *Haemaphysalis punctata* were found to be unaffected by vegetation cover (Borşan et al., 2020). In Singapore, Kwak et al. (2021) observed higher infestation rates of small mammals by ticks such as *Amblyomma helvolum*, *Dermacentor auratus*,

Haemaphysalis semermis, and *Ixodes granulatus* in old secondary forests compared to young secondary forests, with no influence from scrubland. In Oklahoma, Roselli et al. (2022) found that bird infestations by ticks, including *A. americanum*, *Amblyomma maculatum*, and *Haemaphysalis leporispalustris*, were not affected by the surrounding land cover of park sampling sites. In New York City, Hassett et al. (2022) reported that the densities of *I. scapularis*, *Amblyomma americanum*, and *Haemaphysalis longicornis* were negatively influenced by park management practices and were higher along trails than in park edges. Human-tick contact was also more frequent on trails than in natural open spaces. These results revealed no consistent trends across tick species, although park management practices were thought to reduce overall tick abundance.

Concluding remarks on tick-pathogen systems

We found no record of negative effects of UGIs on tick populations. UGIs allowed the presence of ticks in cities and had either no or a positive effect on tick populations and the transmission of tick-borne diseases (Figure 4, Additional file 7: Table S4, Additional file 8: Table S5). Tick abundance and the prevalence of infected ticks were primarily determined by the movement of competent hosts between urban green spaces and peri-urban environments, as well as by local vegetation characteristics.

Effects of UGIs on others vector-pathogen systems

Only one study assessed the impact of UGIs on sand fly populations. In Correntes (Argentina), abundance of *Lutzomyia longipalpis* was higher in areas with low NDVI and in presence of farm animals compared to areas with high NDVI (Berrozpe et al., 2017).

Two studies in Asia investigated chigger mite vectors (Trombiculidae), known carriers of *Orientia tsutsugamushi*, the bacterium responsible for scrub typhus. Park et al. (2015) reported that human scrub typhus prevalence was higher in urban parks than in residential areas in South Korea. Additionally, in Thailand, rodent (*Rattus rattus*) infestation rates were higher in peri-urban and large parks than in small parks in urban centers and were negatively associated with distance from open fields (Wulandhari et al., 2021). This suggested that environmental characteristics of urban parks may influence vector abundance and that proximity to open fields may increase vector populations (Wulandhari et al., 2021).

It is noteworthy that, due to the lack of study, no conclusive trend can be drawn concerning the relationships between UGIs and other vector-pathogen systems, including some of major public health concerns.

Discussion

Since the early 2000's, a growing body of research has emphasized the need to investigate the relationship between UGIs and the risk of VBD introduction and expansion. While urban green spaces may result in increased VBD emergence and transmission for city-dwellers, they also hold potential for VBD risk mitigation through careful UGI planning and management (Fournet et al., 2024). This review highlighted the diverse impacts of UGIs and urban vegetation features on arthropod vectors of human and animals diseases and associated VBD dynamics in cities. These impacts varied significantly depending on vector system and the local socio-ecological and epidemiological context.

The selection of databases and keywords may have resulted in the exclusion of certain articles. For instance, one study lacking the specified keywords was identified through cross-referencing. However, the geographic and thematic heterogeneity of available studies underscores the need for more comprehensive and systematic sampling and monitoring of interactions between UGIs, disease vectors and VBDs affecting humans, but also animals. This should be achieved through an integrative One Health approach, as promoted by the UN in June 2021, which aims to sustainably balance and optimise the health of people, animals and ecosystems (OHHLEP, 2022). Such an approach should consider local sociological, political, and ecological contexts and use

876 harmonized protocols and robust theoretical frameworks for data acquisition and analysis. This will
877 help build scientific evidence and promote stakeholders' engagement to mitigate VBD risk in green
878 cities. Below, we discuss key challenges and propose research avenues to advance this effort.
879

Data acquisition

Developing standard protocols for longitudinal monitoring of disease vector populations,
pathogen dynamics, and urban vegetation features is essential. These protocols would enable
local in-depth assessment of the risk of exposure to VBDs in green cities, and support comparative
studies across different UGIs, study sites, and eco-socio-epidemiological contexts.
First, the studies included in this review encompassed considerable variability in sampling
schemes and arthropod trapping methods, even among studies focusing on the same vector
species. These inconsistencies can limit the representativeness of findings due to the limited
number of spatial or temporal replicates (Amore et al., 2010; Carbó-Ramírez et al., 2017; de Jesús
Crespo & Rogers, 2021; Dongus et al., 2009; Francisco et al., 2021; Georganos et al., 2020; Ha
et al., 2021; Huynh & Minakawa, 2022; Matthys et al., 2006; Sallam et al., 2017; Wilke et al., 2017).
The diversity of indicators used to assess VBD risk further complicates comparisons between
studies on diseases and pathogens. For example, some studies relied on indicators based on
vector abundance. Yet these metrics are known to vary significantly depending on the
physiological stage of the vector sampled (eggs, larvae, adults) and are often uncorrelated with
one another or with disease transmission risk (Corriveau et al., 2003). For example, a recent study
in Leiden (the Netherlands) demonstrated that distribution of *Cx. pipiens* life stages differed across
urban green and grey spaces where ovipositioning rates (number of egg rafts) and larval life stages
were far more abundant in residential areas compared to adults being more abundant in parks
(Krol et al., 2024). Other studies relied on indicators such as disease incidence and prevalence in
humans, or vector infection rates, each representing distinct temporal outcomes of pathogen
transmission, therefore challenging their comparison. Human health data are often aggregated
spatially (Freitas et al., 2021) or tied to residential locations of patients (Muhar et al., 2000; Troyo
et al., 2009), which may not correspond to sites where the patient acquired infection through the
bite of a vector (Lilienfeld, 1983). This mismatch can distort within-city fine-scale risk assessments.
Moreover, the approach can be limited due to a low number of reported cases (Talbot et al., 2019)
or underestimation of cases (Zellweger et al., 2017). Detection and monitoring of pathogens in
animal hosts remain scarce despite their importance for comprehensive VBD risk assessment
(ECDC, 2023). Advancements in technology offer promising solutions for improving spatial and
temporal resolution in monitoring. For instance, 'smart' mosquito traps based on machine learning
approaches can identify and register mosquito counts and species (ECDC, 2023). Non-invasive
eDNA surveys (L'Ambert et al., 2023) and metabarcoding (Bohmann et al., 2022.) can enable
precise pathogen and vector detection, as well as vertebrate host diversity monitoring. Merging
DNA-metabarcoding with ecological network analysis should provide novel insights into vector-
pathogen-host interactions (see more below) and is a promising approach to understand and build
resilient terrestrial ecosystems (Evans et al., 2016). Tailoring these methods to assess VBD risks
in UGIs, will pave the way for integrated and harmonized (meta)data collection and analysis.
Second, there is a need for precise, consistent characterization of urban green spaces, and
associated vegetation. The diversity of definitions and descriptions used to characterize urban
landscapes, particularly UGIs, made it difficult to generalize findings across studies (Cianci et al.,
2015; Talbot et al., 2019). To address this, controlled vocabularies and ontologies are needed to
standardize both field and remote data acquisition. The typology proposed by L. Jones et al. (2022)
provided a clear and comprehensive framework for classifying UGIs. Further development of this
typology could significantly enhance the comparability of UGI studies. Aligning the spatial scale of
measurements with the specific impacts being monitored presents an additional challenge (Landau
& van Leeuwen, 2012; Manica et al., 2016). Longitudinal studies (Mackenstedt et al., 2015;
Tarsitano, 2006), for instance, are complicated by the continuous evolution of urban landscape
design and use over time. Importantly, many studies have shown that landscape factors

influencing VBD risks are context-dependent (Brown et al., 2008; Chen et al., 2020; Cianci et al.,
2015; Gardner et al., 2013; Landau & van Leeuwen, 2012; Sun et al., 2021; Yang et al., 2019).
This emphasizes that relying on a single index cannot determine the relationship between
vegetation and VBD risk. For example, using NDVI alone cannot differentiate between deciduous
and coniferous vegetation, which may have opposite effects depending on the vector system
studied (Lockaby et al., 2016). Moreover, the effects of similar UGIs may vary depending on
factors such as the year of observation (Zellweger et al., 2017), or the specific city being
investigated (Chen et al., 2020). As a result, some landscape indices and summary statistics used
in these studies may not adequately capture the complexity of data needed for a thorough
investigation. Notably, biotic interactions significantly influence vector arthropod fitness and the
dynamics of pathogen transmission within ecosystems. For example, vector trophic behavior can
shift with the seasons (Hedeen et al., 2016; Levine et al., 2017), and host-switching or niche-
switching throughout the seasons can lead to complex epidemiological outcomes (Lefèvre et al.,
2013; Lejal et al., 2021). Indeed, vegetation may impact mosquito population dynamics in multiple
ways. It may directly serve as resting site or sugar source for adults (Peach et al., 2019). Indirectly,
vegetation can alter abiotic and biotic factors by modifying microclimates and by providing habitats
for various animal species, including pathogen hosts, mosquito competitors, and predators
(Benitez et al., 2020; Cianci et al., 2015; Myer & Johnston, 2019). As detailed below, integrating
the full range of interactions mediated by arthropod disease vectors into an ecological network
(food web) approach can help identify relevant biotic and abiotic factors to help design composite
indicators for VBD risk assessment in UGIs.

Finally, the multifaceted nature of VBD transmission dynamics requires consideration of
additional drivers beyond biological and physical parameters associated with vector and pathogen
prevalence. In particular, incorporating socio-economic parameters into VBD risk assessments
has been proposed to bridge ecological and epidemiological findings. While this information is
crucial to assess human exposure to arthropod bites and, to a certain extent to the pathogens they
may transmit, quantifying host-vector contact both within and outside UGIs remains challenging.
Targeted Knowledge-Attitude-Practices (KAP) surveys and citizen science-based approaches
should be encouraged. Mobile applications such as Mosquito Alert (Južnič-Zonta et al., 2022),
Muggenradar (Kampen et al., 2015), iMoustique mobile application (Gautreau, 2024) or Citique
(Hassler et al., 2024) are promising tools to improve completeness and reliability of data
acquisition. Earlier work has shown that such data are extremely valuable for modelling, vector
management, and communication with the general public (Abourashed et al., 2021; Južnič-Zonta
et al., 2022; Kampen et al., 2015). In this context, standardized (minimal) metadata sets should be
developed for studies and case reports to ensure unbiased data assessment and interpretation,
while also facilitating data sharing in accordance with FAIR (Findability, Accessibility,
Interoperability and Reuse) principles (Wilkinson et al., 2016).

Theorical frameworks and models

As this work demonstrated, most studies on the impacts of UGIs on vectors and VBD dynamics
relied on univariate analyses of associations between (sometimes ill-defined) urban green metrics
and vector and/or pathogen presence or abundance. These analyses frequently failed to account
for key confounding factors, such as human density (Bertolotti et al., 2008; Sallam et al., 2017;
Vergara Cid et al., 2013), income levels (Chen et al., 2020; LaBeaud et al., 2008; Matthys et al.,
2010; Zellweger et al., 2017), or urbanization features like building age (Sun et al., 2021). To
address these limitations, various theoretical, statistical, and/or process-driven frameworks and
models are being developed to explore the relationships between landscape features (mainly
abiotic parameters) and population dynamics of arthropod vectors or VBD epidemiology. These
models range from global-scale analyses (Eberhard et al., 2020) to regional-scale studies
(Ducheyne et al., 2018). At a more local scale, deterministic weather-driven population dynamic
models based on generic frameworks (Cailly et al., 2012) are particularly valuable. These models
can map the environmental carrying capacity and identify hotspots for arthropod vector
abundance, such as *Ae. albopictus,* in tropical and temperate environments (Marti et al., 2022;

Tran et al., 2020). When tailored to urban green spaces, these process-based models can be
generalized further by incorporating additional variables, such as local weather parameters of
importance, landscape features and vector data.

Furthermore, hematophagous arthropod vectors are part of complex ecological interaction
networks. These organisms contribute to ecosystem functioning and stability in multiple ways, with
food webs top-down and bottom-up effects (Bellekom et al., 2021; Macfadyen et al., 2009; Ostfeld
et al., 2018). Arthropod vectors may serve as prey for insectivore species at higher trophic levels,
act as competitors with other species sharing similar habitats and resources, and have a major
top-down influence as vectors of pathogens to a wide range of hosts. In this context, developing
multilayer ecological networks (Pocock et al., 2012) appears promising to shed light on complex
and/or unexpected ecological interactions with direct relevance to vector populations and disease
dynamics (Bellekom et al., 2021; Ostfeld et al., 2018; Renault et al., 2024).

Once trained and validated with robust field data, predictive modeling and machine-learning
tools can help to forecast the impact of urban green planning on VBD risks (Wilke et al., 2023).
When combined with future urban (re)naturation scenarios and considering relevant Shared Socio-
economic Pathways (SSPs), these models can simulate entomological and epidemiological
outcomes resulting from changes in urban vegetation across various time horizons. Such
simulations are crucial for identifying high-risk spatial and temporal scenarios, enhancing
preparedness for VBD outbreak.

999 Stakeholder engagement and implications for urban planning

Stakeholder engagement is vital for the development of effective public policies. City planners
should adopt integrated, participatory approaches (Gulachenski et al., 2016; Landau & van
Leeuwen, 2012; Millins et al., 2017; Sprong et al., 2018; Syal, 2021) and embrace a One Health
strategy that considers social, ecological, economic, and technical dimensions (Ellwanger et al.,
2022; Koren & Butler, 2006). Such an integrated approach requires robust communication and
education campaigns targeting many stakeholders, including citizens, health professionals, urban
planners, technicians, entomologists, biologists, architects, botanists, urban ecologists, and
sociologists (Ellwanger et al., 2022). These efforts are important for fostering collaboration and
ensuring comprehensive VBD risk assessments (Wilke et al., 2023). Gaps in knowledge as well
as public health and research priorities may be different across the world, prompting the need for
tailored strategies. For example, in New York (USA), the primary concerns are Lyme disease
(Gregory et al., 2022; Hassett et al., 2022; VanAcker et al., 2019) and West Nile fever (Myer &
Johnston, 2019) whereas in Madrid (Spain), recent leishmaniasis outbreaks highlighted the need
for a better understanding of sandfly ecology in UGIs (Arce et al., 2013). In both cases, urban park
managers are in the frontline to inform VBD risk assessments and contribute to mitigation plans
through wildlife management. In Africa and in South America, basic research is being conducted
on mosquito-pathogen systems, but there is a critical lack of studies on ticks and tick-borne
diseases systems in urban ecosystems despite emerging concerns such as the spread of
Amblyomma cajennense and transmission of *Rickettsia rickettsii* in urban green spaces in Rio de
Janeiro (Szabó et al., 2013).

International knowledge exchange is essential, particularly between countries with long-term
experience of vector presence and countries that have been recently affected. For example, *Ae.*
albopictus is well-established in Italy but is now colonizing Belgium (Deblauwe et al., 2022).
Similarly, countries actively advancing urban greening efforts may share insights with those that
are just beginning. Most studies on urban greening have been conducted in Europe or America,
with relatively few in Africa and Asia. However, a recent study in Gabon investigated the influence
of urban green spaces on VBDs, focusing specifically on urban forest ecosystems (Obame-
Nkoghe et al., 2023). This highlights the growing interest on these issues in different regions and
the need for international cooperation.

In 2019, the Food and Agricultural Organization (FAO) published a report advocating for the
global expansion of urban agriculture to address future food shortages (Taguchi & Santini, 2019).
The report promoted the development of urban farms and agriculture, often associated with

ecopastoralism (urban grazing) involving livestocks. Such developments may contribute to the introduction of competent hosts such as cattle for pathogens like the Crimean-Congo Hemorrhagic Fever Virus (CCHFV), transmitted by the invasive tick *Hyalomma marginatum* (Vial et al., 2016). While urban agriculture improves food access and supports ecological urban green space management, it also presents city planners with the challenge of controlling vector populations in these rapidly changing environments.

Although the impact of UGIs on plant health was out of the scope of this review, the risk of introducing agricultural pests, such as the tree moth (Ciceoi et al., 2017), into cities and contaminating peri-urban forests and/or surrounding crops cannot be ignored and should be carefully assessed when introducing greenery into urban area. This is particularly important when introducing exotic plant specimens that can serve as hosts and reservoirs for devastating pathogens or their vectors (Atkins et al., 2021).

Lastly, city planners should consider the impact of global warming in their risk assessments. We did not identify articles that explicitly consider global warming as a contributing factor to VBD risks associated with urban green spaces. However, climate warming can significantly influence the spatial and temporal distribution of vectors. For instance, the tick *Hyalomma marginatum* is expanding its range into southern France (Bah et al., 2022). Similarly, for *Ae. albopictus*, future projections—though to be interpreted cautiously—identified potential hotspots in the southern United Kingdom, Germany, and Benelux, while southern Spain and Portugal are predicted to become less suitable due to drought and hot summers (Caminade et al., 2012).

Conclusion

In the context of climate change that our world is facing today and given the dynamics of increasing urbanization, renaturing cities appears essential. However, this can lead to the emergence of vector-borne diseases affecting humans, animals, and plants. This risk must be considered in urban policies to enable prevention. The increase in the diversity of vectors, pathogens, and hosts could contribute to a dilution effect that would reduce the risk, but this must be assessed to provide solid foundation for comprehensive risk assessment and informed urban planning. In response to these concerns, only a close partnership between scientists, policymakers, urban planners, and citizens can enable the sustainable development of cities, thereby improving human health and well-being.

Acknowledgements

We would like to thank members of the VECT-OH consortium including Eline Ampt, Roser Fisa Saladrigas, Rosina Girones Llop, Gregory Lanzaro, Sonja Marie Neumeister and Xavier Roca Geronès for useful discussions during the review process.

Funding

This work was part of the VECT-OH initiative and the V2MOC project, funded by the University of Montpellier and RIVOC Défi Clé of Région Occitanie to FS and FF, respectively. CB received a PhD fellowship from the RIVOC Défi Clé of Région Occitanie.

Conflict of interest disclosure

The authors declare that they comply with the PCI rule of having no financial conflicts of interest in relation to the content of the article. The authors declare the following non-financial conflict of interest.

Authors' contributions

FS was the principal investigator of the VECTOH Project and coordinated the manuscript draft with FF. MM and CB conducted the literature review, analyzed the articles and drafted the original version of the manuscript. NM conceptualized the data extraction and description, including the interactive map with references. All co-authors contributed to revision of the manuscript, and approved the final version of the manuscript for publication. All authors read and approved the final manuscript

Data, scripts, code, and supplementary information availability

References

Abourashed, A., Doornekamp, L., Escartin, S., Koenraadt, C. J. M., Schrama, M., Wagener, M., Bartumeus, F., & Van Gorp, E. C. M. (2021). The potential role of school citizen science programs in infectious disease surveillance: a critical review. *International Journal of Environmental Research and Public Health*, *18*(13), 7019. https://doi.org/10.3390/ijerph18137019

Abreu, F. V. S. D., De Andreazzi, C. S., Neves, M. S. A. S., Meneguete, P. S., Ribeiro, M. S., Dias, C. M. G., De Albuquerque Motta, M., Barcellos, C., Romão, A. R., Magalhães, M. D. A. F. M., & Lourenço-de-Oliveira, R. (2022). Ecological and environmental factors affecting transmission of sylvatic yellow fever in the 2017–2019 outbreak in the Atlantic Forest, Brazil. *Parasites & Vectors*, *15*(1), 23. https://doi.org/10.1186/s13071-021-05143-0

Adalsteinsson, S. A., Shriver, W. G., Hojgaard, A., Bowman, J. L., Brisson, D., D'Amico, V., & Buler, J. J. (2018). Multiflora rose invasion amplifies prevalence of Lyme disease pathogen, but not necessarily Lyme disease risk. *Parasites & Vectors*, *11*(1), 54. https://doi.org/10.1186/s13071-018-2623-0

Aly, S. S. A., & Amer, M. S. E. (2010). *Green Corridors as a response for nature: Greening Alexandria city by creating a green infrastructure network*. 101–117. https://doi.org/10.2495/DN100101

Amore, G., Bertolotti, L., Hamer, G. L., Kitron, U. D., Walker, E. D., Ruiz, M. O., Brawn, J. D., & Goldberg, T. L. (2010). Multi-year evolutionary dynamics of West Nile virus in suburban Chicago, USA, 2005–2007. *Philosophical Transactions of the Royal Society B: Biological Sciences*, *365*(1548), 1871–1878. https://doi.org/10.1098/rstb.2010.0054

Andreo, V., Cuervo, P. F., Porcasi, X., Lopez, L., Guzman, C., & Scavuzzo, C. M. (2021). Towards a workflow for operational mapping of *Aedes aegypti* at urban scale based on remote sensing. *Remote Sensing Applications: Society and Environment*, *23*, 100554. https://doi.org/10.1016/j.rsase.2021.100554

Araujo, R. V., Albertini, M. R., Costa-da-Silva, A. L., Suesdek, L., Franceschi, N. C. S., Bastos, N. M., Katz, G., Cardoso, V. A., Castro, B. C., Capurro, M. L., & Allegro, V. L. A. C. (2015). São Paulo urban heat islands have a higher incidence of dengue than other urban areas. *The Brazilian Journal of Infectious Diseases*, *19*(2), 146–155. https://doi.org/10.1016/j.bjid.2014.10.004

Arce, A., Estirado, A., Ordobas, M., Sevilla, S., García, N., Moratilla, L., de la Fuente, S., Martínez, A. M., Pérez, A. M., Aránguez, E., Iriso, A., Sevillano, O., Bernal, J., & Vilas, F. (2013). Re-emergence of leishmaniasis in Spain: Community outbreak in Madrid, Spain, 2009 to 2012. *Eurosurveillance*, *18*(30). https://doi.org/10.2807/1560-7917.ES2013.18.30.20546

Archibald, C. L., McKinney, M., Mustin, K., Shanahan, D. F., & Possingham, H. P. (2017). Assessing the impact of revegetation and weed control on urban sensitive bird species. *Ecology and Evolution*, *7*(12), 4200–4208. https://doi.org/10.1002/ece3.2960

Arduino, M. de B., Mucci, L. F., Santos, L. M. dos, & Soares, M. F. de S. (2020). Importance of microenvironment to arbovirus vector distribution in an urban area, São Paulo, Brazil. *Revista Da Sociedade Brasileira de Medicina Tropical*, *53*, e20190504. https://doi.org/10.1590/0037-8682-0504-2019

Atkins, D. H., Davis, T. S., & Stewart, J. E. (2021). Probability of occurrence and phenology of pine wilt disease transmission by insect vectors in the Rocky Mountains. *Ecological Solutions and Evidence*, *2*(1), e12044. https://doi.org/10.1002/2688-8319.12044

Bah, M. T., Grosbois, V., Stachurski, F., Muñoz, F., Duhayon, M., Rakotoarivony, I., Appelgren, A., Calloix, C., Noguera, L., Mouillaud, T., Andary, C., Lancelot, R., Huber, K., Garros, C., Leblond, A., & Vial, L. (2022). The Crimean-Congo haemorrhagic fever tick vector *Hyalomma marginatum* in the south of France: Modelling its distribution and determination of factors influencing its establishment in a newly invaded area. *Transboundary and Emerging Diseases*, *69*(5). https://doi.org/10.1111/tbed.14578

Bai, T., Mayer, A. L., Shuster, W. D., & Tian, G. (2018). The hydrologic role of urban green space in mitigating flooding (Luohe, China). *Sustainability*, *10*(10), 3584. https://doi.org/10.3390/su10103584

Baró, F., Chaparro, L., Gómez-Baggethun, E., Langemeyer, J., Nowak, D. J., & Terradas, J. (2014). Contribution of ecosystem services to air quality and climate change mitigation policies: the case of urban forests in Barcelona, Spain. *AMBIO*, *43*(4), 466–479. https://doi.org/10.1007/s13280-014-0507-x

Bartholomée, C., & Mercat, M. (2024). *Interactive map of studies included in the Scoping Review on the influence of Urban Green Infrastructures on Vector-Borne Disease risk*. 10.5281/zenodo.15096123, https://zenodo.org/records/15096123

Bartholomée, C., & Mercat, M. (2025). Green cities and the risk for vector-borne disease transmission for humans and animals: a scoping review : additional files [Data set]. Zenodo. https://doi.org/10.5281/zenodo.15165981

Bellekom, B., Hackett, T. D., & Lewis, O. T. (2021). A network perspective on the vectoring of human disease. *Trends in Parasitology*, *37*(5), 391–400. https://doi.org/10.1016/j.pt.2020.12.001

Benitez, E. M., Ludueña-Almeida, F., Frías-Céspedes, M., Almirón, W. R., & Estallo, E. L. (2020). Could land cover influence *Aedes aegypti* mosquito populations? *Medical and Veterinary Entomology*, *34*(2), 138–144. https://doi.org/10.1111/mve.12422

Berrozpe, P., Lamattina, D., Santini, M. S., Araujo, A. V., Utgés, M. E., & Salomón, O. D. (2017). Environmental suitability for *Lutzomyia longipalpis* in a subtropical city with a recently established visceral leishmaniasis transmission cycle, Argentina. *Memórias Do Instituto Oswaldo Cruz*, *112*(10), 674–680. https://doi.org/10.1590/0074-02760170056

Bertolotti, L., Kitron, U. D., Walker, E. D., Ruiz, M. O., Brawn, J. D., Loss, S. R., Hamer, G. L., & Goldberg, T. L. (2008). Fine-scale genetic variation and evolution of West Nile Virus in a transmission "hot spot" in suburban Chicago, USA. *Virology*, *374*(2), 381–389. https://doi.org/10.1016/j.virol.2007.12.040

Blanc, S., & Gutiérrez, S. (2015). The specifics of vector transmission of arboviruses of vertebrates and plants. *Current Opinion in Virology*, *15*, 27–33. https://doi.org/10.1016/j.coviro.2015.07.003

Bohmann, K., Elbrecht, V., Carøe, C., Bista, I., Leese, F., Bunce, M., Yu, D. W., Seymour, M., Dumbrell, A. J., & Creer, S. (2022). *Strategies for sample labelling and library preparation in DNA metabarcoding studies*.

Borşan, S.-D., Toma-Naic, A., Péter, Á., Sándor, A. D., Peştean, C., & Mihalca, A.-D. (2020). Impact of abiotic factors, habitat type and urban wildlife on the ecology of hard ticks (Acari: Ixodidae) in urban and peri-urban habitats. *Parasites & Vectors*, *13*(1), 476. https://doi.org/10.1186/s13071-020-04352-3

Breuste, J., Niemelä, J., & Snep, R. P. H. (2008). Applying landscape ecological principles in urban environments. *Landscape Ecology*, *23*(10), 1139–1142. https://doi.org/10.1007/s10980-008-9273-0

Brown, H., Diuk-Wasser, M., Andreadis, T., & Fish, D. (2008). Remotely-sensed vegetation indices identify mosquito clusters of west nile virus vectors in an urban Landscape in the Northeastern United States. *Vector-Borne and Zoonotic Diseases*, *8*(2), 197–206. https://doi.org/10.1089/vbz.2007.0154

Cailly, P., Tran, A., Balenghien, T., L'Ambert, G., Toty, C., & Ezanno, P. (2012). A climate-driven abundance model to assess mosquito control strategies. *Ecological Modelling*, *227*, 7–17. https://doi.org/10.1016/j.ecolmodel.2011.10.027

Caminade, C., Medlock, J. M., Ducheyne, E., McIntyre, K. M., Leach, S., Baylis, M., & Morse, A. P. (2012). Suitability of European climate for the Asian tiger mosquito *Aedes albopictus*: Recent trends and future scenarios. *Journal of The Royal Society Interface*, *9*(75), 2708–2717. https://doi.org/10.1098/rsif.2012.0138

Carbó-Ramírez, P., Zuria, I., Schaefer, H. M., & Santiago-Alarcon, D. (2017). Avian haemosporidians at three environmentally contrasting urban greenspaces. *Journal of Urban Ecology*, *3*(1). https://doi.org/10.1093/jue/juw011

Chancey, C., Grinev, A., Volkova, E., & Rios, M. (2015). The global ecology and epidemiology of West Nile Virus. *BioMed Research International*, *2015*, 1–20. https://doi.org/10.1155/2015/376230

Che Dom, N., Faiz Madzlan, M., Nadira Yusoff, S. N., Hassan Ahmad, A., Ismail, R., & Nazrina Camalxaman, S. (2016). Profile distribution of juvenile *Aedes* species in an urban area of Malaysia. *Transactions of The Royal Society of Tropical Medicine and Hygiene*, *110*(4), 237–245. https://doi.org/10.1093/trstmh/trw015

Chen, Y., Yang, Z., Jing, Q., Huang, J., Guo, C., Yang, K., Chen, A., & Lu, J. (2020). Effects of natural and socioeconomic factors on dengue transmission in two cities of China from 2006 to 2017. *Science of The Total Environment*, *724*, 138200. https://doi.org/10.1016/j.scitotenv.2020.138200

Chowdhury, R., Chowdhury, V., Faria, S., Huda, M. M., Laila, R., Maheswary, N. P., & Dash, A. P. (2014). How dengue vector *Aedes albopictus* (Diptera: Culicidae) survive during the dry season in Dhaka City, Banglades. *J Vector Borne Dis*.

Cianci, D., Hartemink, N., Zeimes, C. B., Vanwambeke, S. O., Ienco, A., & Caputo, B. (2015). High resolution spatial analysis of habitat preference of *Aedes albopictus* (Diptera: Culicidae) in an urban environment. *Journal of Medical Entomology*, *52*(3), 329–335. https://doi.org/10.1093/jme/tjv026

Ciceoi, R., Gutue, C., Gutue, M., & Roşca, I. (n.d.). *Current status of pests associated with urban vegetation in Bucharest Area*.

Corriveau, R., Philippon, B., Yebakima, A., & Coleman, H. (with Institut de recherche pour le développement). (2003). *La dengue dans les départements français d'Amérique: Comment optimiser la lutte contre cette maladie ?* Institut de recherche pour le développement.

Cox, J., Grillet, M. E., Ramos, O. M., Amador, M., & Barrera, R. (2007). Habitat segregation of dengue vectors along an urban environmental gradient. *The American Journal of Tropical Medicine and Hygiene*, *76*(5), 820–826. https://doi.org/10.4269/ajtmh.2007.76.820

Cuthbert, R. N., Darriet, F., Chabrerie, O., Lenoir, J., Courchamp, F., Claeys, C., Robert, V., Jourdain, F., Ulmer, R., Diagne, C., Ayala, D., Simard, F., Morand, S., & Renault, D. (2023). Invasive hematophagous arthropods and associated diseases in a changing world. *Parasites & Vectors*, *16*(1), 291. https://doi.org/10.1186/s13071-023-05887-x

de Jesús Crespo, R., & Rogers, R. E. (2021). Habitat segregation patterns of container breeding mosquitos: the role of urban heat islands, vegetation cover, and income disparity in cemeteries of New Orleans. *International Journal of Environmental Research and Public Health*, *19*(1), 245. https://doi.org/10.3390/ijerph19010245

Deblauwe, I., De Wolf, K., De Witte, J., Schneider, A., Verlé, I., Vanslembrouck, A., Smitz, N., Demeulemeester, J., Van Loo, T., Dekoninck, W., Krit, M., Madder, M., Müller, R., & Van Bortel,

W. (2022). From a long-distance threat to the invasion front: A review of the invasive *Aedes*
mosquito species in Belgium between 2007 and 2020. *Parasites & Vectors*, *15*(1), 206.
https://doi.org/10.1186/s13071-022-05303-w

DeMets, S., Ziemann, A. K., Manore, C., & Russell, C. (2020). Too big, too small, or just right? The
influence of multispectral image size on mosquito population predictions in the greater Toronto
area. In D. W. Messinger & M. Velez-Reyes (Eds.), *Algorithms, Technologies, and Applications*
for Multispectral and Hyperspectral Imagery XXVI (p. 39). SPIE.
https://doi.org/10.1117/12.2558128

Di Luca, M., Toma, L., Bianchi, R., Quarchioni, E., Marini, L., Mancini, F., Ciervo, A., & Khoury, C.
(2013). Seasonal dynamics of tick species in an urban park of Rome. *Ticks and Tick-Borne*
Diseases, *4*(6), 513–517. https://doi.org/10.1016/j.ttbdis.2013.06.008

Dongus, S., Nyika, D., Kannady, K., Mtasiwa, D., Mshinda, H., Gosoniu, L., Drescher, A. W.,
Fillinger, U., Tanner, M., Killeen, G. F., & Castro, M. C. (2009). Urban agriculture and Anopheles
habitats in Dar es Salaam, Tanzania. *Geospatial Health*, *3*(2), 189.
https://doi.org/10.4081/gh.2009.220

Ducheyne, E., Tran Minh, N. N., Haddad, N., Bryssinckx, W., Buliva, E., Simard, F., Malik, M. R.,
Charlier, J., De Waele, V., Mahmoud, O., Mukhtar, M., Bouattour, A., Hussain, A., Hendrickx,
G., & Roiz, D. (2018). Current and future distribution of *Aedes aegypti* and *Aedes albopictus*
(Diptera: Culicidae) in WHO Eastern Mediterranean Region. *International Journal of Health*
Geographics, *17*(1), 4. https://doi.org/10.1186/s12942-018-0125-0

Duvallet, G., Boulanger, N., & Robert, V. (2018). Arthropods. In *Skin and Arthropod Vectors* (pp.
29–54). Elsevier. https://doi.org/10.1016/B978-0-12-811436-0.00002-2

Eberhard, F. E., Cunze, S., Kochmann, J., & Klimpel, S. (2020). Modelling the climatic suitability
of Chagas disease vectors on a global scale. *eLife*, *9*, e52072.
https://doi.org/10.7554/eLife.52072

Ellwanger, J. H., Byrne, L. B., & Chies, J. A. B. (2022). Examining the paradox of urban disease
ecology by linking the perspectives of Urban One Health and Ecology with Cities. *Urban*
Ecosystems, *25*(6), 1735–1744. https://doi.org/10.1007/s11252-022-01260-5

Estallo, E. L., Sangermano, F., Grech, M., Ludueña-Almeida, F., Frías-Cespedes, M., Ainete, M.,
Almirón, W., & Livdahl, T. (2018). Modelling the distribution of the vector *Aedes aegypti* in a
central Argentine city. *Medical and Veterinary Entomology*, *32*(4), 451–461.
https://doi.org/10.1111/mve.12323

European Food Safety Authority (EFSA) & European Centre for Disease Prevention and Control
(ECDC). (2023). The European Union One Health 2022 Zoonoses Report. *EFSA Journal*,
21(12). https://doi.org/10.2903/j.efsa.2023.8442

Evans, D. M., Kitson, J. J. N., Lunt, D. H., & Straw, N. A. (2016). Merging DNA metabarcoding and
ecological network analysis to understand and build resilient terrestrial ecosystems. *Functional*
Ecology.

Fournet, F., Simard, F., & Fontenille, D. (2024). Green cities and vector-borne diseases: Emerging
concerns and opportunities. *Eurosurveillance*, *29*(10). https://doi.org/10.2807/1560-
7917.ES.2024.29.10.2300548

Francisco, M. E., Carvajal, T. M., Ryo, M., Nukazawa, K., Amalin, D. M., & Watanabe, K. (2021).
Dengue disease dynamics are modulated by the combined influences of precipitation and
landscape: A machine learning approach. *Science of The Total Environment*, *792*, 148406.
https://doi.org/10.1016/j.scitotenv.2021.148406

Freitas, L. P., Schmidt, A. M., Cossich, W., Cruz, O. G., & Carvalho, M. S. (2021). Spatio-temporal
modelling of the first Chikungunya epidemic in an intra-urban setting: The role of socioeconomic
status, environment and temperature. *PLOS Neglected Tropical Diseases*, *15*(6), e0009537.
https://doi.org/10.1371/journal.pntd.0009537

Gardner, A. M., Anderson, T. K., Hamer, G. L., Johnson, D. E., Varela, K. E., Walker, E. D., &
Ruiz, M. O. (2013). Terrestrial vegetation and aquatic chemistry influence larval mosquito
abundance in catch basins, Chicago, USA. *Parasites & Vectors*, *6*(1), 9.
https://doi.org/10.1186/1756-3305-6-9

Gautreau, Y. (2024). *iMoustiques: Une application pour signaler les moustiques, EID Atlantique*. Vigilance - Moustiques. https://vigilance-moustiques.com/imoustiques-une-application-mobile-pour-signaler-les-moustiques-tigres/

Georganos, S., Brousse, O., Dujardin, S., Linard, C., Casey, D., Milliones, M., Parmentier, B., Van Lipzig, N. P. M., Demuzere, M., Grippa, T., Vanhuysse, S., Mboga, N., Andreo, V., Snow, R. W., & Lennert, M. (2020). Modelling and mapping the intra-urban spatial distribution of *Plasmodium falciparum* parasite rate using very-high-resolution satellite derived indicators. *International Journal of Health Geographics*, *19*(1), 38. https://doi.org/10.1186/s12942-020-00232-2

Gregory, N., Fernandez, M. P., & Diuk-Wasser, M. (2022). Risk of tick-borne pathogen spillover into urban yards in New York City. *Parasites & Vectors*, *15*(1), 288. https://doi.org/10.1186/s13071-022-05416-2

Gryczyńska, A., Sokół, M., Gortat, T., & Kowalec, M. (2021). *Borrelia miyamotoi* infection in *Apodemus spp.* Mice populating an urban habitat (Warsaw, Poland). *International Journal for Parasitology: Parasites and Wildlife*, *14*, 138–140. https://doi.org/10.1016/j.ijppaw.2021.01.009

Gulachenski, A., Ghersi, B., Lesen, A., & Blum, M. (2016). Abandonment, ecological assembly and public health risks in counter-urbanizing cities. *Sustainability*, *8*(5), 491. https://doi.org/10.3390/su8050491

Gunawardena, K. R., Wells, M. J., & Kershaw, T. (2017). Utilising green and bluespace to mitigate urban heat island intensity. *Science of The Total Environment*, *584–585*, 1040–1055. https://doi.org/10.1016/j.scitotenv.2017.01.158

Ha, T. V., Kim, W., Nguyen-Tien, T., Lindahl, J., Nguyen-Viet, H., Thi, N. Q., Nguyen, H. V., Unger, F., & Lee, H. S. (2021). Spatial distribution of *Culex* mosquito abundance and associated risk factors in Hanoi, Vietnam. *PLOS Neglected Tropical Diseases*, *15*(6), e0009497. https://doi.org/10.1371/journal.pntd.0009497

Hansford, K. M., Fonville, M., Gillingham, E. L., Coipan, E. C., Pietzsch, M. E., Krawczyk, A. I., Vaux, A. G. C., Cull, B., Sprong, H., & Medlock, J. M. (2017). Ticks and *Borrelia* in urban and peri-urban green space habitats in a city in southern England. *Ticks and Tick-Borne Diseases*, *8*(3), 353–361. https://doi.org/10.1016/j.ttbdis.2016.12.009

Hansford, K. M., McGinley, L., Wilkinson, S., Gillingham, E. L., Cull, B., Gandy, S., Carter, D. P., Vaux, A. G. C., Richards, S., Hayes, A., & Medlock, J. M. (2021). *Ixodes ricinus* and *Borrelia burgdorferi* sensu lato in the Royal Parks of London, UK. *Experimental and Applied Acarology*, *84*(3), 593–606. https://doi.org/10.1007/s10493-021-00633-3

Hansford, K. M., Wheeler, B. W., Tschirren, B., & Medlock, J. M. (2022). Questing *Ixodes ricinus* ticks and *Borrelia* spp. in urban green space across Europe: A review. *Zoonoses and Public Health*, *69*(3), 153–166. https://doi.org/10.1111/zph.12913

Hassell, J. M., Begon, M., Ward, M. J., & Fèvre, E. M. (2017). Urbanization and disease emergence: dynamics at the wildlife–livestock–human interface. *Trends in Ecology & Evolution*, *32*(1), 55–67. https://doi.org/10.1016/j.tree.2016.09.012

Hassett, E., Diuk-Wasser, M., Harrington, L., & Fernandez, P. (2022). Integrating tick density and park visitor behaviors to assess the risk of tick exposure in urban parks on Staten Island, New York. *BMC Public Health*, *22*(1), 1602. https://doi.org/10.1186/s12889-022-13989-x

Hassler, L., Durand, J., & Frey-Klett, P. (2024). *CiTIQUE-Tracker, INRAE, SK8*. https://www.citique.fr/

Hauck, D., Springer, A., Chitimia-Dobler, L., & Strube, C. (2020). Two-year monitoring of tick abundance and influencing factors in an urban area (city of Hanover, Germany). *Ticks and Tick-Borne Diseases*, *11*(5), 101464. https://doi.org/10.1016/j.ttbdis.2020.101464

Heaviside, C., Macintyre, H., & Vardoulakis, S. (2017). The Urban Heat Island: Implications for Health in a Changing Environment. *Current Environmental Health Reports*, *4*(3), 296–305. https://doi.org/10.1007/s40572-017-0150-3

Hedeen, D. L., Bisanzio, D., McMillan, J. R., Brosi, B. J., Mead, D. G., Hamer, G. L., Hedeen, M. W., Kitron, U. D., & Levine, R. S. (2016). Supersuppression: reservoir competency and timing

of mosquito host shifts combine to reduce spillover of West Nile Virus. *The American Journal of Tropical Medicine and Hygiene*, *95*(5), 1174–1184. https://doi.org/10.4269/ajtmh.15-0809

Heinisch, M. R. S., Diaz-Quijano, F. A., Chiaravalloti-Neto, F., Menezes Pancetti, F. G., Rocha Coelho, R., Dos Santos Andrade, P., Urbinatti, P. R., De Almeida, R. M. M. S., & Lima-Camara, T. N. (2019). Seasonal and spatial distribution of *Aedes aegypti* and *Aedes albopictus* in a municipal urban park in São Paulo, SP, Brazil. *Acta Tropica*, *189*, 104–113. https://doi.org/10.1016/j.actatropica.2018.09.011

Hendy, A., Hernandez-Acosta, E., Chaves, B. A., Fé, N. F., Valério, D., Mendonça, C., Lacerda, M. V. G. de, Buenemann, M., Vasilakis, N., & Hanley, K. A. (2020). Into the woods: Changes in mosquito community composition and presence of key vectors at increasing distances from the urban edge in urban forest parks in Manaus, Brazil. *Acta Tropica*, *206*, 105441. https://doi.org/10.1016/j.actatropica.2020.105441

Heylen, D., Lasters, R., Adriaensen, F., Fonville, M., Sprong, H., & Matthysen, E. (2019). Ticks and tick-borne diseases in the city: Role of landscape connectivity and green space characteristics in a metropolitan area. *Science of The Total Environment*, *670*, 941–949. https://doi.org/10.1016/j.scitotenv.2019.03.235

Honnen, A.-C., & Monaghan, M. T. (2017). City-dwellers and country folks: lack of population differentiation along an urban–rural gradient in the mosquito *Culex pipiens* (Diptera: Culicidae). *Journal of Insect Science*, *17*(5). https://doi.org/10.1093/jisesa/iex086

Honório, N. A., Castro, M. G., Barros, F. S. M. D., Magalhães, M. D. A. F. M., & Sabroza, P. C. (2009). The spatial distribution of *Aedes aegypti* and *Aedes albopictus* in a transition zone, Rio de Janeiro, Brazil. *Cadernos de Saúde Pública*, *25*(6), 1203–1214. https://doi.org/10.1590/S0102-311X2009000600003

Hornok, S., Meli, M. L., Gönczi, E., Halász, E., Takács, N., Farkas, R., & Hofmann-Lehmann, R. (2014). Occurrence of ticks and prevalence of *Anaplasma phagocytophilum* and *Borrelia burgdorferi* s.l. in three types of urban biotopes: Forests, parks and cemeteries. *Ticks and Tick-Borne Diseases*, *5*(6), 785–789. https://doi.org/10.1016/j.ttbdis.2014.05.010

Huynh, T. T. T., & Minakawa, N. (2022). A comparative study of dengue virus vectors in major parks and adjacent residential areas in Ho Chi Minh City, Vietnam. *PLOS Neglected Tropical Diseases*, *16*(1), e0010119. https://doi.org/10.1371/journal.pntd.0010119

Hyseni, C., Heino, J., Bini, L. M., Bjelke, U., & Johansson, F. (2021). The importance of blue and green landscape connectivity for biodiversity in urban ponds. *Basic and Applied Ecology*, *57*, 129–145. https://doi.org/10.1016/j.baae.2021.10.004

Jones, E. L., & Leather, S. R. (2012). Invertebrates in urban areas: A review. *European Journal of Entomology*, *109*(4), 463–478. https://doi.org/10.14411/eje.2012.060

Jones, L., Anderson, S., Læssøe, J., Banzhaf, E., Jensen, A., Bird, D. N., Miller, J., Hutchins, M. G., Yang, J., Garrett, J., Taylor, T., Wheeler, B. W., Lovell, R., Fletcher, D., Qu, Y., Vieno, M., & Zandersen, M. (2022). A typology for urban Green Infrastructure to guide multifunctional planning of nature-based solutions. *Nature-Based Solutions*, *2*, 100041. https://doi.org/10.1016/j.nbsj.2022.100041

Junttila, J., Peltomaa, M., Soini, H., Marjamäki, M., & Viljanen, M. K. (1999). Prevalence of *Borrelia burgdorferi* in *Ixodes ricinus* ticks in urban recreational areas of Helsinki. *Journal of Clinical Microbiology*, *37*(5), 1361–1365. https://doi.org/10.1128/JCM.37.5.1361-1365.1999

Južnič-Zonta, Ž., Sanpera-Calbet, I., Eritja, R., Palmer, J. R. B., Escobar, A., Garriga, J., Oltra, A., Richter-Boix, A., Schaffner, F., Della Torre, A., Miranda, M. Á., Koopmans, M., Barzon, L., Bartumeus Ferre, F., Mosquito Alert Digital Entomology Network, & Mosquito Alert Community. (2022). Mosquito alert: Leveraging citizen science to create a GBIF mosquito occurrence dataset. *GigaByte (Hong Kong, China)*, *2022*, gigabyte54. https://doi.org/10.46471/gigabyte.54

Kabaria, C. W., Molteni, F., Mandike, R., Chacky, F., Noor, A. M., Snow, R. W., & Linard, C. (2016). Mapping intra-urban malaria risk using high resolution satellite imagery: A case study of Dar es Salaam. *International Journal of Health Geographics*, *15*(1), 26. https://doi.org/10.1186/s12942-016-0051-y

Kampen, H., Medlock, J. M., Vaux, A., Koenraadt, C., Van Vliet, A., Bartumeus, F., Oltra, A., Sousa, C. A., Chouin, S., & Werner, D. (2015). Approaches to passive mosquito surveillance in the EU. *Parasites & Vectors*, *8*(1), 9. https://doi.org/10.1186/s13071-014-0604-5

Keesing, F., & Ostfeld, R. S. (2021). Dilution effects in disease ecology. *Ecology Letters*, *24*(11), 2490–2505. https://doi.org/10.1111/ele.13875

Kohl, C., McIntosh, E. J., Unger, S., Haddaway, N. R., Kecke, S., Schiemann, J., & Wilhelm, R. (2018). Online tools supporting the conduct and reporting of systematic reviews and systematic maps: A case study on CADIMA and review of existing tools. *Environmental Evidence*, *7*(1), 8. https://doi.org/10.1186/s13750-018-0115-5

Koren, H. S., & Butler, C. D. (2006). *The interconnection between the built environment ecology and health*.

Krol, L., Langezaal, M., Budidarma, L., Wassenaar, D., Didaskalou, E. A., Trimbos, K., Dellar, M., Van Bodegom, P. M., Geerling, G. W., & Schrama, M. (2024). Distribution of *Culex pipiens* life stages across urban green and grey spaces in Leiden, The Netherlands. *Parasites & Vectors*, *17*(1), 37. https://doi.org/10.1186/s13071-024-06120-z

Kubiak, K., Dziekońska-Rynko, J., Szymańska, H., Kubiak, D., Dmitryjuk, M., & Dzika, E. (2019). Questing *Ixodes ricinus* ticks (Acari, Ixodidae) as a vector of *Borrelia burgdorferi* sensu lato and *Borrelia miyamotoi* in an urban area of north-eastern Poland. *Experimental and Applied Acarology*, *78*(1), 113–126. https://doi.org/10.1007/s10493-019-00379-z

Kuddus, M. A., Tynan, E., & McBryde, E. (2020). Urbanization: A problem for the rich and the poor? *Public Health Reviews*, *41*(1), 1. https://doi.org/10.1186/s40985-019-0116-0

Kwak, M. L., Neves, E. S., Borthwick, S. A., Smith, G. J. D., Meier, R., & Mendenhall, I. H. (2021). Habitat impacts the abundance and network structure within tick (Acari: Ixodidae) communities on tropical small mammals. *Ticks and Tick-Borne Diseases*, *12*(3), 101654. https://doi.org/10.1016/j.ttbdis.2021.101654

Kybicová, K., Baštová, K., & Malý, M. (2017). Detection of *Borrelia burgdorferi* sensu lato and *Anaplasma phagocytophilum* in questing ticks *Ixodes ricinus* from the Czech Republic. *Ticks and Tick-Borne Diseases*, *8*(4), 483–487. https://doi.org/10.1016/j.ttbdis.2017.02.007

LaBeaud, A. D., Gorman, A.-M., Koonce, J., Kippes, C., McLeod, J., Lynch, J., Gallagher, T., King, C. H., & Mandalakas, A. M. (2008). Rapid GIS-based profiling of West Nile virus transmission: Defining environmental factors associated with an urbansuburban outbreak in Northeast Ohio, USA. *Geospatial Health*, *2*(2), 215. https://doi.org/10.4081/gh.2008.245

L'Ambert, G., Gendrot, M., Briolant, S., Nguyen, A., Pages, S., Bosio, L., Palomo, V., Gomez, N., Benoit, N., Savini, H., Pradines, B., Durand, A., Leparc, I., Grard, G., & Fontaine, A. (2023). *Analysis of trapped mosquito excreta as a non-invasive method to reveal biodiversity and arbovirus circulation*.

Landau, K. I., & van Leeuwen, W. J. D. (2012). Fine scale spatial urban land cover factors associated with adult mosquito abundance and risk in Tucson, Arizona. *Journal of Vector Ecology*, *37*(2), 407–418. https://doi.org/10.1111/j.1948-7134.2012.00245.x

Lee, J. M., Wasserman, R. J., Gan, J. Y., Wilson, R. F., Rahman, S., & Yek, S. H. (2020). Human Activities Attract Harmful Mosquitoes in a Tropical Urban Landscape. *EcoHealth*, *17*(1), 52–63. https://doi.org/10.1007/s10393-019-01457-9

Lefèvre, T., Vantaux, A., Dabiré, K. R., Mouline, K., & Cohuet, A. (2013). Non-genetic determinants of mosquito competence for malaria parasites. *PLoS Pathogens*, *9*(6), e1003365. https://doi.org/10.1371/journal.ppat.1003365

Lejal, E., Chiquet, J., Aubert, J., Robin, S., Estrada-Peña, A., Rue, O., Midoux, C., Mariadassou, M., Bailly, X., Cougoul, A., Gasqui, P., Cosson, J. F., Chalvet-Monfray, K., Vayssier-Taussat, M., & Pollet, T. (2021). Temporal patterns in *Ixodes ricinus* microbial communities: An insight into tick-borne microbe interactions. *Microbiome*, *9*(1), 153. https://doi.org/10.1186/s40168-021-01051-8

Levine, R. S., Hedeen, D. L., Bisanzio, D., McMillan, J. R., Brosi, B. J., Mead, D. G., Hamer, G. L., Hedeen, M. W., & Kitron, U. D. (2016). Supersuppression: reservoir competency and timing of

mosquito host shifts combine to reduce spillover of West Nile Virus. *The American Journal of Tropical Medicine and Hygiene*, *95*(5), 1174–1184. https://doi.org/10.4269/ajtmh.15-0809

Levine, R. S., Hedeen, D. L., Hedeen, M. W., Hamer, G. L., Mead, D. G., & Kitron, U. D. (2017). Avian species diversity and transmission of West Nile virus in Atlanta, Georgia. *Parasites & Vectors*, *10*(1), 62. https://doi.org/10.1186/s13071-017-1999-6

Levytska, V. A., Mushinsky, A. B., Zubrikova, D., Blanarova, L., Długosz, E., Vichova, B., Slivinska, K. A., Gajewski, Z., Gizinski, S., Liu, S., Zhou, L., & Rogovskyy, A. S. (2021). Detection of pathogens in ixodid ticks collected from animals and vegetation in five regions of Ukraine. *Ticks and Tick-Borne Diseases*, *12*(1), 101586. https://doi.org/10.1016/j.ttbdis.2020.101586

Lilienfeld, A. M. (n.d.). *Practical Limitations of Epidemiologic Methods*.

Liu, H., & Weng, Q. (2012). Enhancing temporal resolution of satellite imagery for public health studies: A case study of West Nile Virus outbreak in Los Angeles in 2007. *Remote Sensing of Environment*, *117*, 57–71. https://doi.org/10.1016/j.rse.2011.06.023

Liu, H., Weng, Q., & Gaines, D. (2008). Spatio-temporal analysis of the relationship between WNV dissemination and environmental variables in Indianapolis, USA. *International Journal of Health Geographics*, *7*(1), 66. https://doi.org/10.1186/1476-072X-7-66

Liu, H., Weng, Q., & Gaines, D. (2011). Geographic incidence of human West Nile virus in northern Virginia, USA, in relation to incidence in birds and variations in urban environment. *Science of The Total Environment*, *409*(20), 4235–4241. https://doi.org/10.1016/j.scitotenv.2011.07.012

Lockaby, G., Noori, N., Morse, W., Zipperer, W., Kalin, L., Governo, R., Sawant, R., & Ricker, M. (2016). Climatic, ecological, and socioeconomic factors associated with West Nile virus incidence in Atlanta, Georgia, U.S.A. *Journal of Vector Ecology*, *41*(2), 232–243. https://doi.org/10.1111/jvec.12218

Lorenz, C., Castro, M. C., Trindade, P. M. P., Nogueira, M. L., De Oliveira Lage, M., Quintanilha, J. A., Parra, M. C., Dibo, M. R., Fávaro, E. A., Guirado, M. M., & Chiaravalloti-Neto, F. (2020). Predicting *Aedes aegypti* infestation using landscape and thermal features. *Scientific Reports*, *10*(1), 21688. https://doi.org/10.1038/s41598-020-78755-8

Lourenço-de-Oliveira, R., Castro, M. G., Braks, M. A. H., & Lounibos, L. P. (2004). The invasion of urban forest by dengue vectors in Rio de Janeiro. *Journal of Vector Ecology*.

Macfadyen, S., Gibson, R., Polaszek, A., Morris, R. J., Craze, P. G., Planqué, R., Symondson, W. O. C., & Memmott, J. (2009). Do differences in food web structure between organic and conventional farms affect the ecosystem service of pest control? *Ecology Letters*, *12*(3), 229–238. https://doi.org/10.1111/j.1461-0248.2008.01279.x

Mackenstedt, U., Jenkins, D., & Romig, T. (2015). The role of wildlife in the transmission of parasitic zoonoses in peri-urban and urban areas. *International Journal for Parasitology: Parasites and Wildlife*, *4*(1), 71–79. https://doi.org/10.1016/j.ijppaw.2015.01.006

Manica, M., Filipponi, F., D'Alessandro, A., Screti, A., Neteler, M., Rosà, R., Solimini, A., della Torre, A., & Caputo, B. (2016). Spatial and temporal hot spots of *Aedes albopictus* abundance inside and outside a South European Metropolitan Area. *PLOS Neglected Tropical Diseases*, *10*(6), e0004758. https://doi.org/10.1371/journal.pntd.0004758

Marando, F., Heris, M. P., Zulian, G., Udías, A., Mentaschi, L., Chrysoulakis, N., Parastatidis, D., & Maes, J. (2022). Urban heat island mitigation by green infrastructure in European Functional Urban Areas. *Sustainable Cities and Society*, *77*, 103564. https://doi.org/10.1016/j.scs.2021.103564

Marti, R., Castets, M., Demarchi, M., Catry, T., Besnard, G., Chouin, S., Clément, C., Esteve-Moussion, I., Etienne, M., Foussadier, R., Habchi-Hanriot, N., Jouanthoua, F., L'Ambert, G., Godal, A., & Tran, A. (2022, 13/05). *ARBOCARTO: an operational spatial modeling tool to predict Aedes albopictus dynamics and the impact of vector control interventions (OC 4.1)*. 5th International Workshop on Aedes albopictus, Montpellier.

Martinez, A. D. L. I., & Labib, S. M. (2023). Demystifying normalized difference vegetation index (NDVI) for greenness exposure assessments and policy interventions in urban greening. *Environmental Research*, *220*, 115155. https://doi.org/10.1016/j.envres.2022.115155

Mathews-Martin, L., Namèche, M., Vourc'h, G., Gasser, S., Lebert, I., Poux, V., Barry, S., Bord,
S., Jachacz, J., Chalvet-Monfray, K., Bourdoiseau, G., Pamies, S., Sepúlveda, D., Chambon-
Rouvier, S., & René-Martellet, M. (2020). Questing tick abundance in urban and peri-urban
parks in the French city of Lyon. *Parasites & Vectors*, *13*(1), 576.
https://doi.org/10.1186/s13071-020-04451-1
Matthys, B., Koudou, B. G., N'Goran, E. K., Vounatsou, P., Gosoniu, L., Koné, M., Gissé, G., &
Utzinger, J. (2010). Spatial dispersion and characterisation of mosquito breeding habitats in
urban vegetable-production areas of Abidjan, Côte d'Ivoire. *Annals of Tropical Medicine &
Parasitology*, *104*(8), 649–666. https://doi.org/10.1179/136485910X12851868780108
Matthys, B., N'Goran, E. K., Gosoniu, L., Cissé, G., Tschannen, A. B., Becket, E. G., Vounatsou,
P., Raso, G., Utzinger, J., & Tanner, M. (2006). Urban farming and malaria risk factors in a
medium-sized town in Côte d'Ivoire. *The American Journal of Tropical Medicine and Hygiene*,
75(6), 1223–1231. https://doi.org/10.4269/ajtmh.2006.75.1223
May, K., & Strube, C. (2014). Prevalence of Rickettsiales (*Anaplasma phagocytophilum* and
Rickettsia spp.) in hard ticks (*Ixodes ricinus)* in the city of Hamburg, Germany. *Parasitology
Research*, *113*(6), 2169–2175. https://doi.org/10.1007/s00436-014-3869-x
Medeiros-Sousa, A. R., Fernandes, A., Ceretti-Junior, W., Wilke, A. B. B., & Marrelli, M. T. (2017).
Mosquitoes in urban green spaces: Using an island biogeographic approach to identify drivers
of species richness and composition. *Scientific Reports*, *7*(1), 17826.
https://doi.org/10.1038/s41598-017-18208-x
Mexia, T., Vieira, J., Príncipe, A., Anjos, A., Silva, P., Lopes, N., Freitas, C., Santos-Reis, M.,
Correia, O., Branquinho, C., & Pinho, P. (2018). Ecosystem services: Urban parks under a
magnifying glass. *Environmental Research*, *160*, 469–478.
https://doi.org/10.1016/j.envres.2017.10.023
Michalik, J., Hofman, T., Buczek, A., Skoracki, M., & Sikora, B. (2003). *Borrelia burgdorferi* s.l. in
Ixodes ricinus (Acari: Ixodidae) ticks collected from vegetation and small rodents in recreational
areas of the city of Poznań. *Journal of Medical Entomology*, *40*(5), 690–697.
https://doi.org/10.1603/0022-2585-40.5.690
Millins, C., Gilbert, L., Medlock, J., Hansford, K., Thompson, D. B., & Biek, R. (2017). Effects of
conservation management of landscapes and vertebrate communities on Lyme borreliosis risk
in the United Kingdom. *Philosophical Transactions of the Royal Society B: Biological Sciences*,
372(1722), 20160123. https://doi.org/10.1098/rstb.2016.0123
Muhar, A., Dale, P. E. R., Thalib, L., & Arito, E. (2000). The spatial distribution of ross river virus
infections in Brisbane: Significance of residential location and relationships with vegetation
types. *Environmental Health and Preventive Medicine*, *4*(4), 184–189.
https://doi.org/10.1007/BF02931256
Myer, M. H., & Johnston, J. M. (2019). Spatiotemporal Bayesian modeling of West Nile virus:
Identifying risk of infection in mosquitoes with local-scale predictors. *Science of The Total
Environment*, *650*, 2818–2829. https://doi.org/10.1016/j.scitotenv.2018.09.397
Noden, B. H., Roselli, M. A., & Loss, S. R. (2022). Effect of urbanization on presence, abundance,
and coinfection of bacteria and protozoa in ticks in the US Great Plains. *Journal of Medical
Entomology*, *59*(3), 957–968. https://doi.org/10.1093/jme/tjab228
Obame-Nkoghe, J., Makanga, B. K., Zongo, S. B., Koumba, A. A., Komba, P., Longo-Pendy, N.-
M., Mounioko, F., Akone-Ella, R., Nkoghe-Nkoghe, L. C., Ngangue-Salamba, M.-F., Yangari,
P., Aboughe-Angone, S., Fournet, F., Kengne, P., & Paupy, C. (2023). Urban green spaces
and vector-borne disease risk in africa: the case of an unclean forested park in Libreville
(Gabon, Central Africa). *International Journal of Environmental Research and Public Health*,
20(10), 5774. https://doi.org/10.3390/ijerph20105774
Ogashawara, I., Li, L., & Moreno-Madriñán, M. J. (2019). Spatial-temporal assessment of
environmental factors related to dengue outbreaks in São Paulo, Brazil. *GeoHealth*, *3*(8), 202–
217. https://doi.org/10.1029/2019GH000186
One Health High-Level Expert Panel (OHHLEP), Adisasmito, W. B., Almuhairi, S., Behravesh, C.
B., Bilivogui, P., Bukachi, S. A., Casas, N., Cediel Becerra, N., Charron, D. F., Chaudhary, A.,

Ciacci Zanella, J. R., Cunningham, A. A., Dar, O., Debnath, N., Dungu, B., Farag, E., Gao, G. F., Hayman, D. T. S., Khaitsa, M., … Zhou, L. (2022). One Health: A new definition for a sustainable and healthy future. *PLOS Pathogens*, *18*(6), e1010537. https://doi.org/10.1371/journal.ppat.1010537

Ostfeld, R. S., Levi, T., Keesing, F., Oggenfuss, K., & Canham, C. D. (2018). Tick-borne disease risk in a forest food web. *Ecology*, *99*(7), 1562–1573. https://doi.org/10.1002/ecy.2386

Overzier, E., Pfister, K., Thiel, C., Herb, I., Mahling, M., & Silaghi, C. (2013). Diversity of *Babesia* and *Rickettsia* species in questing *Ixodes ricinus*: a longitudinal study in urban, pasture, and natural habitats. *Vector-Borne and Zoonotic Diseases*, *13*(8), 559–564. https://doi.org/10.1089/vbz.2012.1278

Page, M. J., McKenzie, J. E., Bossuyt, P. M., Boutron, I., Hoffmann, T. C., Mulrow, C. D., Shamseer, L., Tetzlaff, J. M., Akl, E. A., Brennan, S. E., Chou, R., Glanville, J., Grimshaw, J. M., Hróbjartsson, A., Lalu, M. M., Li, T., Loder, E. W., Mayo-Wilson, E., McDonald, S., … Moher, D. (2021). The PRISMA 2020 statement: An updated guideline for reporting systematic reviews. *Systematic Reviews*, *10*(1), 89. https://doi.org/10.1186/s13643-021-01626-4

Park, S.-W., Ha, N.-Y., Ryu, B., Bang, J. H., Song, H., Kim, Y., Kim, G., Oh, M., Cho, N.-H., & Lee, J. (2015). Urbanization of Scrub Typhus Disease in South Korea. *PLOS Neglected Tropical Diseases*, *9*(5), e0003814. https://doi.org/10.1371/journal.pntd.0003814

Paul, K. K., Dhar-Chowdhury, P., Haque, C. E., Al-Amin, H. M., Goswami, D. R., Kafi, M. A. H., Drebot, M. A., Lindsay, L. R., Ahsan, G. U., & Brooks, W. A. (2018). Risk factors for the presence of dengue vector mosquitoes, and determinants of their prevalence and larval site selection in Dhaka, Bangladesh. *PLOS ONE*, *13*(6), e0199457. https://doi.org/10.1371/journal.pone.0199457

Peach, D. A. H., Ko, E., Blake, A. J., & Gries, G. (2019). Ultraviolet inflorescence cues enhance attractiveness of inflorescence odour to *Culex pipiens* mosquitoes. *PLOS ONE*, *14*(6), e0217484. https://doi.org/10.1371/journal.pone.0217484

Pearce, J. C., Learoyd, T. P., Langendorf, B. J., & Logan, J. G. (2018). Japanese encephalitis: The vectors, ecology and potential for expansion. *Journal of Travel Medicine*, *25*(Suppl_1), S16–S26. https://doi.org/10.1093/jtm/tay009

Pedrosa, M. C., Borges, M. A. Z., Eiras, Á. E., Caldas, S., Cecílio, A. B., Brito, M. F., & Ribeiro, S. P. (2020). Invasion of tropical montane cities by *Aedes aegypti* and *Aedes albopictus* (Diptera: Culicidae) depends on continuous warm winters and suitable urban biotopes. *Journal of Medical Entomology*, tjaa135. https://doi.org/10.1093/jme/tjaa135

Pocock, S. J., Ariti, C. A., Collier, T. J., & Wang, D. (2012). The win ratio: A new approach to the analysis of composite endpoints in clinical trials based on clinical priorities. *European Heart Journal*, *33*(2), 176–182. https://doi.org/10.1093/eurheartj/ehr352

Poh, K. C., Medeiros, M. C. I., & Hamer, G. L. (2020). Landscape and demographic determinants of Culex infection with West Nile virus during the 2012 epidemic in Dallas County, TX. *Spatial and Spatio-Temporal Epidemiology*, *33*, 100336. https://doi.org/10.1016/j.sste.2020.100336

Prihandi, D. R., & Nurvianto, S. (2022). The role of urban green space design to support bird community in the urban ecosystem. *Biodiversitas Journal of Biological Diversity*, *23*(4). https://doi.org/10.13057/biodiv/d230449

Renault, D., Derocles, S. A. P., Haubrock, P., Simard, F., Cuthbert, R. N., & Valiente-Moro, C. (2024). Biological, ecological and trophic features of invasive mosquitoes and other hematophagous arthropods: What makes them successful? *Biological Invasions*, *26*(1), 33–55. https://doi.org/10.1007/s10530-023-03158-5

Roselli, M. A., Noden, B. H., & Loss, S. R. (2022). Tick infestation of birds across a gradient of urbanization intensity in the United States Great Plains. *Urban Ecosystems*, *25*(2), 379–391. https://doi.org/10.1007/s11252-021-01160-0

Sallam, M., Michaels, S., Riegel, C., Pereira, R., Zipperer, W., Lockaby, B., & Koehler, P. (2017). Spatio-Temporal Distribution of Vector-Host Contact (VHC) Ratios and Ecological Niche Modeling of the West Nile Virus Mosquito Vector, Culex quinquefasciatus, in the City of New

Orleans, LA, USA. *International Journal of Environmental Research and Public Health*, *14*(8), 892. https://doi.org/10.3390/ijerph14080892

Schorn, S., Pfister, K., Reulen, H., Mahling, M., Manitz, J., Thiel, C., & Silaghi, C. (2011). Prevalence of *Anaplasma phagocytophilum* in *Ixodes ricinus* in Bavarian public parks, Germany. *Ticks and Tick-Borne Diseases*, *2*(4), 196–203. https://doi.org/10.1016/j.ttbdis.2011.09.009

Schorn, S., Pfister, K., Reulen, H., Mahling, M., & Silaghi, C. (2011). Occurrence of *Babesia* spp., *Rickettsia spp*. and *Bartonella* spp. in *Ixodes ricinus* in Bavarian public parks, Germany. *Parasites & Vectors*, *4*(1), 135. https://doi.org/10.1186/1756-3305-4-135

Schwarz, M., Byrd, B. D., Marayati, B. F., Blum, P. W., Wells, M. B., Greene, A. D., Taylor, M., & Wasserberg, G. (2020). Horizontal distribution affects the vertical distribution of native and invasive container-inhabiting *Aedes* mosquitoes within an urban landscape. *Journal of Vector Ecology*, *45*(1), 16–24. https://doi.org/10.1111/jvec.12369

Small, M., & Brennan, R. E. (2021). Detection of *Rickettsia amblyommatis* and *Ehrlichia chaffeensis* in *Amblyomma americanum* inhabiting two urban parks in Oklahoma. *Vector-Borne and Zoonotic Diseases*, *21*(5), 385–387. https://doi.org/10.1089/vbz.2020.2755

Sprong, H., Azagi, T., Hoornstra, D., Nijhof, A. M., Knorr, S., Baarsma, M. E., & Hovius, J. W. (2018). Control of Lyme borreliosis and other *Ixodes ricinus*-borne diseases. *Parasites & Vectors*, *11*(1), 145. https://doi.org/10.1186/s13071-018-2744-5

Sun, H., Dickens, B. L., Richards, D., Ong, J., Rajarethinam, J., Hassim, M. E. E., Lim, J. T., Carrasco, L. R., Aik, J., Yap, G., Cook, A. R., & Ng, L. C. (2021). Spatio-temporal analysis of the main dengue vector populations in Singapore. *Parasites & Vectors*, *14*(1), 41. https://doi.org/10.1186/s13071-020-04554-9

Syal, S. (2021). Learning from pandemics: Applying resilience thinking to identify priorities for planning urban settlements. *Journal of Urban Management*, *10*(3), 205–217. https://doi.org/10.1016/j.jum.2021.05.004

Szabó, M. P. J., Pinter, A., & Labruna, M. B. (2013). Ecology, biology and distribution of spotted-fever tick vectors in Brazil. *Frontiers in Cellular and Infection Microbiology*, *3*. https://doi.org/10.3389/fcimb.2013.00027

Taguchi, M., & Santini, G. (n.d.). *Urban agriculture in the Global North & South: A perspective from FAO*.

Talbot, B., Ardis, M., & Kulkarni, M. A. (2019). Influence of demography, land use, and urban form on West Nile Virus Risk and Human West Nile Virus incidence in Ottawa, Canada. *Vector-Borne and Zoonotic Diseases*, *19*(7), 533–539. https://doi.org/10.1089/vbz.2018.2366

Talbot, B., Sander, B., Cevallos, V., González, C., Benítez, D., Carissimo, C., Carrasquilla Ferro, M. C., Gauto, N., Litwiñiuk, S., López, K., Ortiz, M. I., Ponce, P., Villota, S. D., Zelaya, F., Espinel, M., Wu, J., Miretti, M., & Kulkarni, M. A. (2021). Determinants of *Aedes* mosquito density as an indicator of arbovirus transmission risk in three sites affected by co-circulation of globally spreading arboviruses in Colombia, Ecuador and Argentina. *Parasites & Vectors*, *14*(1), 482. https://doi.org/10.1186/s13071-021-04984-z

Tarsitano, E. (2006). Interaction between the environment and animals in urban settings: integrated and participatory planning. *Environmental Management*, *38*(5), 799–809. https://doi.org/10.1007/s00267-005-0148-8

Tian. (2019). Effect of common ornamental plants on the survivorship and fecundity of the *Aedes albopictus* (Diptera: Culicidae). *Florida Entomologist*, *102*(1), 36. https://doi.org/10.1653/024.102.0106

Tran, A., Mangeas, M., Demarchi, M., Roux, E., Degenne, P., Haramboure, M., Le Goff, G., Damiens, D., Gouagna, L.-C., Herbreteau, V., & Dehecq, J.-S. (2020). Complementarity of empirical and process-based approaches to modelling mosquito population dynamics with *Aedes albopictus* as an example—Application to the development of an operational mapping tool of vector populations. *PLOS ONE*, *15*(1), e0227407. https://doi.org/10.1371/journal.pone.0227407

Trawinski, P. R., & Mackay, D. S. (2010). Identification of environmental covariates of West Nile Virus Vector mosquito population abundance. *Vector-Borne and Zoonotic Diseases*, *10*(5), 515–526. https://doi.org/10.1089/vbz.2008.0063

Tretyakov, K. (2009). Small mammals and their parasites (ixodid ticks) in urban forests and parks of St. Petersburg. *Estonian Journal of Ecology*, *58*(3), 232. https://doi.org/10.3176/eco.2009.3.08

Troyo, A., Fuller, D. O., Calderón-Arguedas, O., Solano, M. E., & Beier, J. C. (2009). Urban structure and dengue incidence in Puntarenas, Costa Rica. *Singapore Journal of Tropical Geography*, *30*(2), 265–282. https://doi.org/10.1111/j.1467-9493.2009.00367.x

UN Habitat. (2022). *Envisaging the Future of Cities*. UN Habitat.

UNDRR. (2019). *Global Assessment Report on Disaster Risk Reduction, Geneva, Switzerland, United Nations Office for Disaster Risk Reduction (UNDRR)*.

United Nations. (2020, February 21). *Urbanization: Expanding opportunities, but deeper divides | UN DESA | United Nations Department of Economic and Social Affairs*. https://www.un.org/development/desa/en/news/social/urbanization-expanding-opportunities-but-deeper-divides.html

United Nations. (2022). *The Sustainable Development Goals Report* (p. 68).

Van Den Bosch, M., & Ode Sang, Å. (2017). Urban natural environments as nature-based solutions for improved public health – A systematic review of reviews. *Environmental Research*, *158*, 373–384. https://doi.org/10.1016/j.envres.2017.05.040

VanAcker, M. C., Little, E. A. H., Molaei, G., Bajwa, W. I., & Diuk-Wasser, M. A. (2019). Enhancement of risk for lyme disease by landscape connectivity, New York, New York, USA. *Emerging Infectious Diseases*, *25*(6), 1136–1143. https://doi.org/10.3201/eid2506.181741

Vautard, R., Van Aalst, M., Boucher, O., Drouin, A., Haustein, K., Kreienkamp, F., Van Oldenborgh, G. J., Otto, F. E. L., Ribes, A., Robin, Y., Schneider, M., Soubeyroux, J.-M., Stott, P., Seneviratne, S. I., Vogel, M. M., & Wehner, M. (2020). Human contribution to the record-breaking June and July 2019 heatwaves in Western Europe. *Environmental Research Letters*, *15*(9), 094077. https://doi.org/10.1088/1748-9326/aba3d4

Vergara Cid, C., Estallo, E. L., Almirón, W. R., Contigiani, M. S., & Spinsanti, L. I. (2013). Landscape determinants of Saint Louis encephalitis human infections in Córdoba city, Argentina during 2010. *Acta Tropica*, *125*(3), 303–308. https://doi.org/10.1016/j.actatropica.2012.12.005

Vergnes, A., Viol, I. L., & Clergeau, P. (2012). Green corridors in urban landscapes affect the arthropod communities of domestic gardens. *Biological Conservation*, *145*(1), 171–178. https://doi.org/10.1016/j.biocon.2011.11.002

Vial, L., Stachurski, F., Leblond, A., Huber, K., Vourc'h, G., René-Martellet, M., Desjardins, I., Balança, G., Grosbois, V., Pradier, S., Gély, M., Appelgren, A., & Estrada-Peña, A. (2016). Strong evidence for the presence of the tick *Hyalomma marginatum* Koch, 1844 in southern continental France. *Ticks and Tick-Borne Diseases*, *7*(6), 1162–1167. https://doi.org/10.1016/j.ttbdis.2016.08.002

Vieira, C. J. da S. P., Thies, S. F., da Silva, D. J. F., Kubiszeski, J. R., Barreto, E. S., Monteiro, H. A. de O., Mondini, A., São Bernardo, C. S., & Bronzoni, R. V. de M. (2020). Ecological aspects of potential arbovirus vectors (Diptera: Culicidae) in an urban landscape of Southern Amazon, Brazil. *Acta Tropica*, *202*, 105276. https://doi.org/10.1016/j.actatropica.2019.105276

Weaver, S. C., & Reisen, W. K. (2010). Present and future arboviral threats. *Antiviral Research*, *85*(2), 328–345. https://doi.org/10.1016/j.antiviral.2009.10.008

Welc-Falęciak, R., Kowalec, M., Karbowiak, G., Bajer, A., Behnke, J. M., & Siński, E. (2014). Rickettsiaceae and Anaplasmataceae infections in *Ixodes ricinus* ticks from urban and natural forested areas of Poland. *Parasites & Vectors*, *7*(1), 121. https://doi.org/10.1186/1756-3305-7-121

WHO. (2021). *One health*. Retrieved 31 March 2025, from https://www.who.int/health-topics/one-health

WHO. (2020). *Vector-borne diseases*. https://www.who.int/news-room/fact-sheets/detail/vector-
borne-diseases

Wilke, A. B. B., Carvajal, A., Vasquez, C., Petrie, W. D., & Beier, J. C. (2020). Urban farms in
Miami-Dade county, Florida have favorable environments for vector mosquitoes. *PLOS ONE*,
15(4), e0230825. https://doi.org/10.1371/journal.pone.0230825

Wilke, A. B. B., Chang, N.-B., Townsend, J., Benelli, G., & Ajelli, M. (2023). Unintended effects of
urban policies on the risk of arbovirus transmission. *Trends in Parasitology*, *39*(12), 1001–1003.
https://doi.org/10.1016/j.pt.2023.08.012

Wilke, A. B. B., Medeiros-Sousa, A. R., Ceretti-Junior, W., & Marrelli, M. T. (2017). Mosquito
populations dynamics associated with climate variations. *Acta Tropica*, *166*, 343–350.
https://doi.org/10.1016/j.actatropica.2016.10.025

Wilkinson, M. D., Dumontier, M., Aalbersberg, Ij. J., Appleton, G., Axton, M., Baak, A., Blomberg,
N., Boiten, J.-W., Da Silva Santos, L. B., Bourne, P. E., Bouwman, J., Brookes, A. J., Clark, T.,
Crosas, M., Dillo, I., Dumon, O., Edmunds, S., Evelo, C. T., Finkers, R., … Mons, B. (2016).
The FAIR Guiding Principles for scientific data management and stewardship. *Scientific Data*,
3(1), 160018. https://doi.org/10.1038/sdata.2016.18

Wilson, A. J., Morgan, E. R., Booth, M., Norman, R., Perkins, S. E., Hauffe, H. C., Mideo, N.,
Antonovics, J., McCallum, H., & Fenton, A. (2017). What is a vector? *Philosophical
Transactions of the Royal Society B: Biological Sciences*, *372*(1719), 20160085.
https://doi.org/10.1098/rstb.2016.0085

Wong, G. K. L., & Jim, C. Y. (2016). Do vegetated rooftops attract more mosquitoes? Monitoring
disease vector abundance on urban green roofs. *Science of The Total Environment*, *573*, 222–
232. https://doi.org/10.1016/j.scitotenv.2016.08.102

Wong, G. K. L., & Jim, C. Y. (2017). Urban-microclimate effect on vector mosquito abundance of
tropical green roofs. *Building and Environment*, *112*, 63–76.
https://doi.org/10.1016/j.buildenv.2016.11.028

Wong, G. K. L., & Jim, C. Y. (2018). Abundance of urban male mosquitoes by green infrastructure
types: Implications for landscape design and vector management. *Landscape Ecology*, *33*(3),
475–489. https://doi.org/10.1007/s10980-018-0616-1

Wulandhari, S. A., Paladsing, Y., Saesim, W., Charoennitiwat, V., Sonthayanon, P., Kumlert, R.,
Morand, S., Sumruayphol, S., & Chaisiri, K. (2021). High prevalence and low diversity of chigger
infestation in small mammals found in Bᴀɴɢᴋᴏᴋ Mᴇᴛʀᴏᴘᴏʟɪᴛᴀɴ parks. *Medical and Veterinary
Entomology*, *35*(4), 534–546. https://doi.org/10.1111/mve.12531

Yadouléton, A., N'Guessan, R., Allagbé, H., Asidi, A., Boko, M., Osse, R., Padonou, G., Kindé, G.,
& Akogbéto, M. (2010). The impact of the expansion of urban vegetable farming on malaria
transmission in major cities of Benin. *Parasites & Vectors*, *3*(1), 118.
https://doi.org/10.1186/1756-3305-3-118

Yang, L., Turo, K. J., Riley, C. B., Inocente, E. A., Tian, J., Hoekstra, N. C., Piermarini, P. M., &
Gardiner, M. M. (2019). Can urban greening increase vector abundance in cities? The impact
of mowing, local vegetation, and landscape composition on adult mosquito populations. *Urban
Ecosystems*, *22*(5), 827–839. https://doi.org/10.1007/s11252-019-00857-7

Zellweger, R. M., Cano, J., Mangeas, M., Taglioni, F., Mercier, A., Despinoy, M., Menkès, C. E.,
Dupont-Rouzeyrol, M., Nikolay, B., & Teurlai, M. (2017). Socioeconomic and environmental
determinants of dengue transmission in an urban setting: An ecological study in Nouméa, New
Caledonia. *PLOS Neglected Tropical Diseases*, *11*(4), e0005471.
https://doi.org/10.1371/journal.pntd.0005471

1743
1744

Identification of studies via databases and registers

Identification

Date of search 5/09/22
Records identified from:
Databases (n = 2535)
(Scopus n=775,
Web of Science n=791,
PubMed n=969)

Records removed before screening:
Duplicate records removed by :
Cadima automatic tools (n=972)
Authors (n=5)

Screening

Records screened on
title/abstract
(n = 1558)

Records excluded on title/abstract
(n = 997)

Records screened on full text
(n = 561)

Reason of exclusion on full text (n=463)
- Not primary research
- Not accessible
- Not related to vector borne diseases
- Not related to urban area
- Not related to greening cities
- No comparison between urban green spaces versus other spaces
- No statistical analysis

Inclusion

Studies included in review
(n = 98 +1 = 99)

1 article added by the authors

A. Mosquito population

Urban green space typology
(68 variables/24 articles)

Urban vegetation description
(122 variables/24 articles)

0.0   0.5   1.0

B. Mosquito-borne

Urban green space typology
(44 variables/12 articles)

Urban vegetation description
(33 variables/17 articles)

0.0   0.5   1.0

Impact of UGIs

Harmful
Variable
Beneficial
No association

A. *Aedes aegypti*

Urban green space typology
(8 variables/6 articles)

Urban vegetation description
(26 variables/9 articles)

B. *Aedes albopictus*

Urban green space typology
(11 variables/9 articles)

Urban vegetation description
(26 variables/11 articles)

Impact of UGIs

- Harmful
- Variable
- Beneficial
- No association

C. Dengue virus

Urban green space typology
(16 variables/1 article)

Urban vegetation description
(15 variables/6 articles)

A *Culex quinquefasciatus*

B *Culex pipiens* and *Culex restuans*

C. West Nile Virus

Impact of UGIs

- Harmful
- Variable
- Beneficial
- No association

A *Ixodes ricinus*

Urban green space typology
(17 variables/8 articles)

Urban vegetation description
(26 variables/5 articles)

B. Lyme bacteria *Borrelia spp.*

Urban green space typology
(6 variables/4 articles)

Urban vegetation description
(9 variables/4 articles)

Impact of UGIs

- Harmful
- Variable
- Beneficial
- No association