No evidence of reactive avoidance of baboons (*Papio ursinus and Papio anubis*) to the presence of predators

N. van Rooyen[1,2,*], L. Thel[3,*], J. A. Venter[3], H. Fritz[1,2], F. Prugnolle[1,2,#], V. Rougeron[1,2,#]

Affiliations

[1] International Research Laboratory, REHABS, CNRS-NMU-UCBL, George Campus, Nelson Mandela University, Madiba drive, 6529 George, South Africa

[2] Sustainability Research Unit, George Campus, Nelson Mandela University, George Campus, Madiba drive, 6529 George, South Africa

[3] Department of Conservation Management, Faculty of Science, Nelson Mandela University, George Campus, George, South Africa

* Co-first authors: N. van Rooyen, L. Thel; [#] Co-supervised the work: F. Prugnolle, V. Rougeron

Corresponding authors: Rougeron Virginie, International Research Laboratory, REHABS, CNRS-NMU-UCBL, Sustainability Research Unit, George Campus, Nelson Mandela University, George, South Africa, rougeron.virginie@gmail.com/virginie.rougeron@cnrs.fr; Nicholas van Rooyen, Department of conservation management, faculty of science, Nelson Mandela University George Campus, Nelson Mandela University, George, South Africa, vanrooyennicholas@gmail.com.

Abstract

Predators exert strong selective pressure on prey species, shaping their behavioral adaptations. Prey species use proactive responses, such as site selection and the adjustment of daily activity patterns to anticipate and avoid predation exposure. In contrast, reactive responses, including fleeing, referential signaling and retaliation, occur after encountering a predator to mitigate the immediate predation risk. The presence of a predator in an area can also generate longer-term responses such as reactive avoidance, defined here as the temporary avoidance of the area following the passage of a predator. Such a long-term reactive response remains understudied in primates. To investigate reactive avoidance, we analyzed an extensive camera trap dataset of more than 3,000 detections of both chacma (*Papio ursinus*) and olive (*Papio anubis*) baboons, including almost 7,000 detections of their main predators, lions (*Panthera leo*), leopards (*Panthera pardus*) and spotted hyenas (*Crocuta crocuta*), in four savanna sites (Madikwe Game Reserve, Tswalu Kalahari Private Wildlife Reserve and the Associated Private Nature Reserves, South Africa; Serengeti National Park, Tanzania). We examined whether baboons display species-specific reactive avoidance towards predators up to 72 hours after the detection of a predator using randomization tests. We found no significant decrease in baboon presence (defined as the number of baboon detections) 0 to 24 hours, 24 to 48 hours and 48 to 72 hours after the detection of any predator species. These results suggest that baboons may not display reactive avoidance and rely on alternative predator-avoidance strategies, such as proactive avoidance or aggressive retaliation, to mitigate predation risk.

Keywords: anti-predator behavior, camera trap, leopard, lion, predator-prey interactions, primate, randomization test, spotted hyena, tropical ecosystem.

Introduction

Predation exerts a strong selective pressure on prey species due to its impact on individual fitness and survival (Humphries & Driver, 1967; Harvell, 1990; Krebs & Davies, 2009). Prey species have evolved a diverse range of behavioral responses to enhance their survival as they encounter fluctuating predation risk in their environment (Lima & Dill, 1990; Kie, 1999). Behavioural adaptations are inherently flexible and allow prey to modify the nature and intensity of their responses based on their spatial and temporal perception of risk in the surrounding environment (i.e., the landscape of fear, Laundré et al., 2001; Creel & Winnie, 2005; Palmer et al., 2022), and can be classified into proactive and reactive responses (Broekhuis et al., 2013; Courbin et al., 2016; Creel, 2018; Palmer & Packer, 2021; Say-Sallaz et al., 2023).

Proactive responses are behavioral adjustments made in anticipation of predation risk, based on prior knowledge of predator locations and preferred habitats (Creel et al & Christianson, 2008; Broekhuis et al., 2013; Creel et al., 2014). These adjustments, which result in the interaction with resource and refuge availability in the landscape (Willems & Russell, 2009; Suscke et al., 2021), operate over broad spatial and temporal scales such as home ranges and seasonal cycles (Droge et al., 2016). For example, orangutans (*Pongo spp.*) proactively select nest sites that provide greater camouflage, making it more difficult for predators to detect both them and their young (van Casteren, 2012). Hamadryas baboons (*Papio hamadryas*) make use of cliff faces (Schreier & Swedell, 2012) as sleeping and refuge sites to limit predator accessibility to the group and increase predator detectability (Anderson, 1999; Bidner et al., 2018). Chacma baboons (*Papio ursinus*) use refuge areas on the edge of urban environments, spending less than 1% of their time within the urban area, as a strategy to avoid human presence (Fehlmann et al., 2017). They also increase vigilance when away from roosting sites or when isolated from group members, enhancing threat detection in

vulnerable situations (Fischer et al., 2001). However, when foraging, the item size and

complexity of the foraging task might reduce the level of vigilance of the individual (e.g.,

reduced vigilance when processing small food items requiring more attention, Allan et al.,

2024).

Reactive responses, in contrast, are immediate reactions triggered by the detection of

a predator (Broekhuis et al., 2013). These responses occur on smaller spatial and temporal

scales such as foraging sites and across diel activity periods (Droge et al., 2016), driven by

real-time threats. As a short term reactive response, white-faced capuchins (*Cebus*

capucinus), vervet monkeys (*Chlorocebus pygerythrus*) and spectral tarsiers (*Tarsius tarsier*)

commonly use referential signaling to alert other group members after the detection of a

predator (Seyfarth et al., 1980; Fichtel et al., 2005; Gursky, 2007). Bonobos (*Pan paniscus*),

colobine monkeys (*Colobus guereza*) and baboons (*Papio spp*.) seek refuge in elevated areas

(e.g., tree canopies and cliffs) to reduce their vulnerability to terrestrial predators (Cowlishaw,

1994; Von Hippel, 1998; Isbell et al., 2018; Druelle et al., 2020). Chacma baboons also

demonstrate aggressive predator mobbing when facing a predator, where the group

collectively confronts and harasses the predator to deter the attack (Bailey, 1993; Cowlishaw,

1994; Jooste et al., 2013; Isbell et al., 2018). In addition to these short-term responses, some

taxa also display long-term reactive avoidance, leaving the disturbed area for several hours to

several days, as frequently observed in ungulates (e.g., in Burchell's zebra (*Equus quagga*),

Courbin et al., 2016; in red deer (*Cervus elaphus*), Chassagneux et al., 2020). However, it

remains unclear whether primates actively move away from sites visited recently by predators

for extended periods of time (Allan et al., 2024; Hammond et al., 2025).

Baboons present an ideal study species to explore primate anti-predator behavior. As a

prey species, baboons encounter a diverse range of predators due to their broad geographic

distribution (Busse, 1980; Cowlishaw, 1994; Jooste et al., 2013; Isbell et al., 2018), such as

lions (*Panthera leo*), leopards (*Panthera pardus*) and spotted hyenas (*Crocuta crocuta*) (Busse, 1980; Jooste et al., 2013; Dröge et al., 2017; Isbell et al., 2018). Although baboons are usually not the main target for these predators (Busse, 1980; Jooste et al., 2013; Dröge et al., 2017; Isbell et al., 2018), they nevertheless contribute to significant baboon mortality (Busse, 1980; Cowlishaw, 1994). Leopards and lions are both ambush predator species, lions hunting baboons during the day, whereas leopards hunt during both diurnal and nocturnal periods, often targeting baboons at their roosting sites at night (Busse, 1980; Bidner et al., 2018; Isbell et al., 2018). Spotted hyenas are cursorial predators (Trinkel, 2010; Wentworth et al., 2011), which mostly target baboons when the opportunity arises as opposed to actively hunting them (Busse 1980; Cowlishaw, 1994; Trinkel, 2010; Wentworth et al., 2011). These different predation strategies are known to trigger different anti-predator responses in ungulate prey species (Berger, 1979; Palmer et al., 2021). Although species-specific responses among primates have been overlooked, there is emerging evidence of similar anti-predator responses occurring within this taxon (Gursky, 2007; Seiler et al., 2013; Adams & Kitchen, 2020). Suire et al. (2023) found that male olive baboons (*Papio anubis*) tend to position themselves at the periphery of the group to protect more vulnerable group members such as females and juveniles from cursorial predators.

So far, most studies on baboon-predator interactions have relied primarily on direct field observations (e.g., Busse, 1980; Hill & Weingrill, 2007; Allan et al., 2021). While these methods offer detailed insights into short-term reactive responses, they may influence animal behavior due to the presence of human observers (predator shelter hypothesis, Shannon et al., 2014). Additionally, they are often difficult to implement over extended periods or under specific conditions such as nighttime, limiting their usability to explore long-term reactive responses (Busse, 1980; Nowak et al., 2014; Dill & Frid, 2020; LaBarge et al., 2022; Allan et al., 2024). In contrast, camera traps are a non-invasive, uninterrupted monitoring alternative

(Newey et al., 2015), which proved to reliably detect large carnivores (Meek et al., 2016) as

well as primates (Boyer-Ontl & Pruetz, 2014; Bersacola et al., 2022). Camera traps have

largely been used to study predator-prey interactions (Smith et al., 2020), particularly space

use (e.g., Cusack et al., 2017; Keim et al., 2019) and activity pattern overlaps (e.g., Linkie &

Ridout, 2011; Foster et al., 2013). The widespread adoption of camera traps in recent decades

has led to the development of long-term monitoring programs across extensive areas

(Swanson et al., 2015; Blount et al., 2021), providing a new source of data for the study of

predator-prey interactions (e.g., Hammond et al., 2025). Although promising, are these non-

targeted, large-scale datasets adequate for examining long-term primate reactive responses to

predators?

In this study, we took advantage of the largest camera trap-based monitoring project

in Africa, Snapshot Safari (Swanson et al., 2015; Pardo et al., 2021) and analyzed 10,000

camera trap detections from three sites in South Africa and one site in Tanzania to test

whether baboons exhibit species-specific reactive avoidance in response to the presence of

predators. We tested three hypotheses related to the interaction between baboons and predator

characteristics: H1) we expected baboon presence to decrease significantly 0 to 24 hours

following the detection of spotted hyena, a terrestrial cursorial predator. Spotted hyenas only

opportunistically hunt baboons (Busse, 1980; Cowlishaw, 1994; Trinkel, 2010; Wentworth et

al., 2011) and are highly mobile across the landscape (Kolowski et al., 2007). As semi-

arboreal primates, baboons predominantly forage in proximity to trees or cliffs (Schreier &

Swedell, 2012; Fehlmann et al., 2017) that serve as potential refuges in the event of a

predation attempt. We therefore predicted that baboons should be weakly influenced by the

occurrence of spotted hyena, returning to the disturbed area shortly after the detection of

spotted hyena. H2) we expected baboon presence to decrease significantly 0 to 48 hours

following the detection of leopard, a semi-arboreal ambush predator. Leopards can pursue

baboons into trees, which may prompt the entire group to relocate for a longer period (Busse,

1980; Trinkel, 2010; Wentworth et al., 2011; Isbell et al., 2018). However, leopards are also

highly mobile and may not remain in the same area for long periods (Martins and Harris,

2013; Hubel et al., 2018). We therefore predicted that baboons should be more strongly

affected by leopard than spotted hyena presence, leading to their displacement from the

disturbed area with a gradual return following the detection of leopard. H3) we expected

baboon presence to decrease significantly 0 to 72 hours following the detection of lion, a

terrestrial ambush predator. Although baboons can evade lion predation by retreating to

elevated refuges, lions can exhibit limited daily displacement and may remain in the same

area for extended periods (Elliot et al., 2014), potentially forcing baboons to relocate for a

prolonged duration to maintain their normal activity patterns (particularly for foraging) in

safer conditions. We therefore predicted that baboons should be most strongly affected by

lion presence, resulting in the longest avoidance period compared to other predators, with a

slow return following the detection of lion.

Methods

Study site selection and description

In this study, we exploited data from the ongoing camera trap monitoring project *Snapshot*

Safari, which spans multiple sites and countries (Tanzania, data publicly available at Dryad:

http://dx.doi.org/10.5061/dryad.5pt92, Swanson et al., 2015; South Africa, Pardo et al.,

2021). Of the 32 sites surveyed, we included sites where all three target predators were

detected along with baboons. The South African study sites included Madikwe Game Reserve

(MAD, 24°49'S-26°13'E; Figure 1A), Tswalu Kalahari Private Wildlife Reserve (TSW,

27°14'S-22°23'E; Figure 1B), and the Associated Private Nature Reserves (APN, 24°20'S-

31°20'E; Figure 1C), inhabited by chacma baboons (Stone et al., 2013). The Tanzanian study

site was Serengeti National Park (SER, 2°19'S-34°49'E; Figure 1D), occupied by olive

baboons (Zinner et al., 2011). All four study sites were located within the savanna biome,

characterized by open grasslands interspersed with isolated trees (Sankaran et al., 2005).

MAD is a 750 km² fenced game reserve located in the Northwest Province, on the

border of South Africa and Botswana, within the central bushveld bioregion (Szott et al.,

2020). The landscape is characterized by ridges and thorny woodlands dominated by acacia

species (Szott et al., 2020). TSW, located in the Northern Cape Province, is a 1,020 km²

fenced reserve in the Eastern Kalahari bioregion, with portions extending into the Kalahari

dunefields. Elevation ranges from 1,020 m a.s.l. in lower-lying areas to 1,586 m a.s.l. at the

highest point (Davis et al., 2010). The vegetation consists of discontinuous annual and

perennial grasses interspersed with deep-rooted trees, such as the camel thorn (*Vachellia*

erioloba, Beaumont & Bednarik, 2015). APN is a collective of five interconnected reserves

(Timbavati, Thornybush, Klaserie, Balule, and Umbabat Private Nature Reserves) covering

1,800 km² in the mopaneveld bioregion of Limpopo and Mpumalanga provinces. It is open to

Kruger National Park (Bedetti et al., 2020; Rutherford et al., 2006). The APN vegetation is

highly variable, with isolated trees in a continuous grass understory interspersed with patches

of dense woody vegetation (Scholes & Archer, 1997). The SER study site corresponds to an

open sub-section of 1,125 km^2 inside the core of Serengeti National Park, Tanzania, part of

the East African savanna biome. It is covered by grasslands interspersed with scattered trees,

with steeper slopes to the North-West (Swanson et al., 2015).

Camera trap data

Data for MAD, TSW, and APN were collected between June 2017 and November 2019 (29

months), while data for SER were collected between June 2010 and May 2013 (35 months).

As is common in large-scale monitoring programs, camera traps (n = 357) from different

brands (Cuddeback's Professional Series x-change, white flash in South Africa, and

Scoutguard (SG565) incandescent cameras in SER) were deployed. Camera trap placement

followed a systematic layout in MAD, TSW, and SER, where camera traps were positioned at

the center of a 5 km² grid pattern. Camera trap placement in the APN followed a random

allocation layout (*sensu* Meek et al., 2014), based on habitat and terrain structure. We

removed five camera traps from the MAD dataset that were positioned specifically near

waterholes, as predator-prey interactions may likely differ at waterholes due to the trade-off

between the need for scarce resources (i.e., water) and predator avoidance (Edwards et al.,

2016). In the present study, we used a total of 346 camera traps (nMAD = 40; nTSW = 31;

nAPN = 56; nSER = 219) for our analyses (see below for data processing and selection

steps).

The camera traps were enclosed in steel casings and fixed at a height of

approximately 50 cm above the ground (Swanson et al., 2015; Pardo et al., 2021). To

minimize glare caused by the rising and setting sun, the camera traps were oriented to face as

far North or South as possible. Cameras were also directed based on vegetation cover,

avoiding areas of high grass density to reduce the probability of obtaining blank pictures and

maximizing animal detection. The sensitivity of each camera trap was set to a medium level

(50-75%) to minimize false triggers caused by windblown vegetation. No lures were used to

attract any species to the camera trap sites. Each camera trap was programmed to take three

pictures for every motion detection during the day and one at night (see Pardo et al., 2021 for

a more detailed description of the protocol in the South African sites; and Swanson et al.,

2015 for the Tanzanian site). To minimize the impact of potential variability in detection

probabilities due to differences between protocols and sites (Cusack et al., 2017; Herrera et

al., 2021), we analyzed our four sites separately.

Species were identified in each capture event (i.e., any single picture or consecutive

series of pictures initiated by an animal-trigger, regardless of the number of individuals

present in the picture(s), Meek et al., 2014) using two methods: (i) for SER, the citizen

science program Zooniverse (www.zooniverse.org) which uses classifications from the

general public (consensus after 10 matching classifications of species or species combination;

see details in Swanson et al., 2015); (ii) for the three South African sites, the online program

TrapTagger (www.wildeyeconservation.org), which was used to manually record animal

species as identified by trained laboratory technicians. After species classification, we

obtained a total of n = 1,625 detections of chacma baboons, n = 1,531 detections of olive

baboons, n = 6,929 detections of spotted hyenas, n = 449 detections of leopards and n = 3,886

detections of lions.

As our objective was to evaluate the presence of baboons up to 72 hours after (and

symmetrically, before) the detection of a predator, we exclude any predator detection that

occurred within the 72 hours following the camera's deployment and within 72 hours before

the end of the camera's operational lifespan. To ensure consistency, we applied a 30-minute

time-to-independence filter to predator detections, retaining the central detection within a

cluster of consecutive detections as the main detection for a given predator detection. This

threshold aligns with previous studies that showed that increasing the independence interval

has minimal impact on observed activity patterns of large carnivores (Linkie & Ridout, 2011;

Searle et al., 2021; Tjaden☐McClement et al., 2025). Retaining the central detection instead

of the first detection as usually conducted in the literature ensures symmetry when testing for

baboon response to a predator before and after the predator detection in our study design. Our

final dataset was thus composed of n = 1,584 detections of chacma baboons, n = 1,458

detections of olive baboons, n = 5,071 detections of spotted hyenas, n = 355 detections of

leopards and n = 1,532 detections of lions (Table 1).

Statistical analyses

To assess whether baboon presence after a predator detection was lower than expected by

chance, we used a randomization test to compare the number of baboon detections after the

predator detection to the ones obtained under the assumption of non-avoidance of predators

(Niedballa et al., 2019; Zhang & Zhao, 2023). We calculated the total number of baboon

detections following predator detections in each site, for each predator species independently.

This was done for three consecutive 24-hour time-blocks after predator detection: 0 to 24

hours after the predator detection, 24 to 48 hours after the predator detection and 48 to 72

after the predator detection. Extending the observation period beyond 72 hours after predator

detection was avoided due to the difficulty in associating the prey species presence or

absence with the original predator detection (Courbin et al., 2016; Swinkels et al., 2023). We

further analyzed baboon detections before the predation detection to establish a baseline for

their presence, with the expectation of no significant difference in the number of baboon

detections. This allowed us to determine the strength of the predator effect while accounting

for natural fluctuations in baboon activity. The analysis of baboon detections before a

predator detection was also conducted in three 24-hour time-blocks over a 72-hour period.

Predator species have been reported to sometimes follow each other (e.g., for

kleptoparasitism, Cusack et al., 2017). In our dataset, < 5% of the days when a predator was

detected at a camera trap site were also characterized by the detection of a second predator

species (0.7% in MAD, 0.0% in TSW, 2.5% in APN, 3.6% in SER). We thus considered that

there was no significant risk of interpreting baboon reactive avoidance due to a predator

species when examining the effect of another species.

To obtain the distribution of baboon detections under the null hypothesis (i.e., non-

avoidance of predators), we randomized the dates while preserving the original timing to

maintain natural diel activity patterns of each predator detection 1,000 times in each camera

trap-roll (i.e., the continuous working time of a camera trap between two services,

approximately three months) detecting both baboons and the predator of interest. We then

estimated, for each randomized data set, the total number of baboon detections in each time-

block. We excluded any camera trap with a lifespan < 30 consecutive days to avoid restricting

the randomization process. Predator detections captured by camera traps with a shorter

lifespan could only be randomized within a limited time window, which would produce

unreliable results.

For each 24-hour time-block, we evaluated whether the observed number of baboon

detections deviated significantly from the assumption of non-avoidance of predators. To do

so, we calculated the significance value (p) for each time-block as the proportion of iterations

in our randomization procedure in which the number of baboon detections was inferior

(superior, respectively) or equal to the observed number of baboon detections after (before,

respectively) the predator detection. All data processing and analyses were performed using

the R software (version 4.5.1; R Core Development Team, 2025).

Validation of the randomization approach

To validate our randomization approach, we tested it on Burchell's zebras (*Equus quagga).*

Burchell's zebras are known to display reactive avoidance in response to lions for up to 24

hours after an encounter (Thaker et al., 2011; Courbin et al., 2016; Say-Sallaz et al., 2023),

therefore we expected to find a significant response in the 0 to 24 hour time-block, after lion

detection.

APN had the highest number of lion detections among the South African sites ($n_{APN} =$

188, $n_{MAD} = 17$, $n_{TSW} = 51$; sample sizes after data selection process). Although SER had

higher numbers of both lions ($n_{SER} = 1,276$) and Burchell's zebras ($n_{SER} = 60,212$), analysing

these data revealed too computationally intensive using our randomization procedure.

Additionally, the number of Burchell's zebras (n_{APN} = 2,047) was of the same magnitude as

the number of baboons in SER, which represents our largest dataset in terms of baboon

detections (n_{SER} = 1,458), making it a suitable comparison point to assess the effect of sample

size on the results.

To assess the effect of sample size on the analyses, we performed 100 random subsets

of Burchell's zebra detections in APN to match the number of baboon detections (n_{APN} =

500). We thus created a down sampled data set, for which we conducted the randomization

analysis for each of the 100 iterations and calculated the frequency of significant versus non-

significant values.

Results

Method validation: reactive avoidance in Burchell's zebras

As expected, Burchell's zebras showed significant reactive avoidance 0 to 24 hours after lion

detection in APN (p = 0.024, Figure 2), providing support for the soundness of our

randomization approach. We did not find significant reactive avoidance 24 to 48 hours (p =

0.176) and 48 to 72 hours (p = 0.495) after lion detection (Table 2). After taking a subset of

Burchell's zebra detections to match the number of baboon detections in APN (n = 500), we

could no longer detect this response 0 to 24 hours following lion detection (*p (mean ± SD)* =

0.182 ± 0.103, Supplementary Figure 1).

Reactive avoidance in baboons

Our results did not show any significant reactive avoidance in baboons in response to the

detection of spotted hyenas, across any time-block after predator detection (0 to 24, 24 to 48,

or 48 to 72 hours) in any of the three study sites in South Africa (all p > 0.05, Table 2, Figure

3). Similarly, there was no evidence of baboons displaying reactive avoidance following the

detection of lions or leopards in any South African site and across any time-block after their

detection (all $p > 0.05$, Table 2, Figure 3). However, in APN, baboon presence increased

significantly 48 to 24 hours before the detection of lions ($p < 0.001$) and 24 to 0 hours before

the detection of spotted hyenas ($p < 0.007$, Table 2, Figure 3). In TSW, baboon presence

increased significantly 24 to 0 hours before the detection of lions ($p = 0.033$, Table 2, Figure

3).

In SER, where the number of baboon detections exceeded those at all South African sites

(SER detection rate approximately 4.21 times higher than MAD, 2.92 times higher than APN

and 1.98 times higher than TSW, Table 1), and was close to the number of Burchell's zebra

detections in APN, we still did not detect significant reactive avoidance of baboons in

response to any predator tested and across all time-blocks after predator detection (all $p >$

0.05, Table 2, Figure 3). Consistent with observations in APN, baboon detections in SER

were significantly higher than expected under the null hypothesis 24 to 0 hours before the

detection of spotted hyenas ($p < 0.008$, Table 2, Figure 3). Additionally, baboon detections in

SER were significantly higher than expected under the null hypothesis 24 to 0 hours before

the detection of leopards ($p = 0.012$, Table 2, Figure 3).

Discussion

Using camera trap data from a large-scale monitoring program in four different

savanna sites in South Africa and Tanzania, we found no evidence that either chacma or olive

baboons display reactive avoidance for as long as 72 hours after the detection of any of their

main predators (lions, leopards and spotted hyenas). Contrary to our expectations, there was

no detectable difference in the duration of the reactive avoidance response of baboons

according to predator characteristics (i.e., predator hunting strategy, predator daily mobility, predator specialization in baboon hunting) at such a time scale.

The absence of reactive avoidance at a one-to-three-day timescale suggests that baboons might rely on short-term rather than long-term anti-predator responses. Baboons usually forage in areas with tall trees which can serve as refuge against terrestrial predators such as lions and spotted hyenas (Cowlishaw, 1997). They also exhibit effective sentinel behaviors, one individual monitoring for threats while the rest of the group forages (Horrocks et al., 1986; Bezerra et al., 2008), which leads to the early detection of predation attempts. The danger is rapidly signaled to conspecifics via contact barks (Bailey, 1993; Cowlishaw, 1994; Jooste et al., 2013; Isbell et al., 2018), which allows for their prompt retreat. Baboons may thus remain in the vicinity of a predator, and retreat into refuges temporarily only in the instance of a predation attempt, rather than completely avoiding a potentially resource rich area for several days when frequented by a predator. Similarly, Hammond et al. (2025) found no evidence of predator-occupied area avoidance by chacma baboons and suggest that they might instead display short-term reactive avoidance. The modulation of reactive anti-predator responses by environmental variables such as the proximity of refuge areas reducing the vulnerability to predators has also been documented in other species (e.g., in woodchucks (*Marmota monax*), Kramer & Bonenfant, 1997; in European hares (*Lepus europaeus*), Weterings et al., 2016).

Additionally, baboons are notorious for their aggressive retaliation toward predators, particularly leopards (Buse, 1980; Bailey, 1993; Cowlishaw, 1994; Hayward et al., 2006: Jooste et al., 2013; Isbell et al., 2018). A review by Cowlishaw (1994) showed that, when baboons retaliate, they successfully deter predation in 93% of cases, and four out of 11 observed predation attempts resulted in the death of the attacking leopard. This behavior is predominantly exhibited by male in both chacma (Cowlishaw, 1994; Isbell et al., 2018) and

olive (MacCormick et al., 2012) baboons, owing to their larger teeth and body size (Virgadamo et al., 1972; Galbany et al., 2015; Johnson, 2003). Similarly, a large number of primate species, mainly males, display mobbing and counter-attack behavior as an anti-predator response (Schaik et al., 2022). Such a response could decrease predation attempts by lowering the predator's chance of successful capture and thus, their interest in the species (Lima & Dill, 1990; Cowlishaw, 1994; Alberts, 2018; Isbell et al., 2018). Therefore, predators could be perceived as less of a threat by baboons which would consequently be less inclined to avoid the area, even if frequented by predators.

Surprisingly, we detected evidence suggesting an increased presence of baboons before a predator detection across all study sites (except MAD). Although these results might be artifacts due to small sample sizes, they might also be related to the fact that baboons use the ground more frequently in the absence of predators as it is perceived as safer (Hammond et al., 2022; Hammond et al., 2025). This could lead to an increased capture rate of the same individuals during their terrestrial displacements in the monitored area. Although most primates primarily forage in trees, they may adopt more terrestrial foraging behaviors during periods when food resources are less abundant and require supplementation with grasses or other ground-based food items (e.g., Li, 2007), particularly baboons.

The ability to detect spatio-temporal avoidance confidently when applying a randomization procedure to camera trap data depends on the strength of the avoidance behavior and the number of detections of both species, the latter needing to be as high as 100 detections per species in the case of weak avoidance (Niedballa et al. 2019). In our method validation, we found that Burchell's zebras showed reactive avoidance toward lions within the 0 to 24 hour time-block after a lion detection, consistent with previous findings in this species (Thaker et al., 2011; Courbin et al., 2016; Say-Sallaz et al., 2023) and thus supporting the validity of our approach. However, after the number of detections of Burchell's zebras

was sub-sampled to match the highest number of baboon detections, we no longer detected

reactive avoidance. Sample sizes being relatively low in our four study sites (particularly in

MAD and TSW), we might lack statistical power to detect significant reactive avoidance in

baboons. To completely rule out this possibility, more camera traps may be necessary to

obtain better detection rates of both prey and predators. Si et al. (2014) and Evans et al.

(2019) showed that when studying coyotes, adding one camera trap at each capture site

increases the detection probability by four. Additionally, baboons being semi-arboreal, a

mixed design of terrestrial and arboreal camera traps could be adopted to cover all three

spatial dimensions of baboon movement and increase their detection rate (Bersacola et al.,

2022). Although large-scale camera-trap based monitoring programs present a novel

opportunity for preliminary explorations of predator-prey interactions, optimizing camera

trap density and placement appears essential to improve detection rates, particularly for low

density species with complex habitat use such as primates.

Conclusion

This study contributes to broadening our understanding of predator-prey interactions

in two primate species. The potential absence of reactive avoidance in chacma and olive

baboons raises important questions about the variability of their anti-predator strategies, not

only in savanna biomes but also across other environments in response to different predators

(Zinner et al., 2011; Stone et al., 2013). This study further highlights the influence of

detection rates on the detectability of behavioral responses when using camera traps, as

illustrated by the loss of significance of reactive avoidance in Burchell's zebras after the

reduction of sample size. These limitations related to camera trap density and placement

further emphasize the challenges of accurately quantifying prey-predator interactions among

416   semi-arboreal and low-density species using systematic and random layouts, common in

417   large-scale monitoring projects.

Authors' contributions

Conceptualization: VR, FP – Methodology: NVR, LT, VR, FP – Data acquisition: HF, JV – Formal analysis: NVR, LT, VR, FP – Investigation: NVR, VR, FP, LT – Writing: Original draft: NVR, VR, FP, LT – Writing: Review & Editing: NVR, LT, VR, FP, HF, JV – Supervision: VR, FP – Funding acquisition: VR, JV

Acknowledgments

The authors thank the CNRS for funding this project. They also thank the Wildlife Ecology Lab, Nelson Mandela University – George campus, CNRS, IRL REHABS, and the Snapshot Safari project for the provision and processing of the data. Funding for Snapshot Safari was provided by the National Research Foundation and the Foundational Biodiversity Information Program (Grant number FBIP170720256205). The authors thank the editor in chief and an anonymous reviewer for insightful comments on previous drafts.

Inclusion and diversity statement

The authors are committed to fostering an environment of open dialogue, respect, and cultural inclusion. Our collaboration between African and European researchers reflects our dedication to international scientific exchange and diverse perspectives. Within our research team and laboratory, we actively promote a supportive and inclusive environment. Our first author lives with temporal lobe epilepsy, a condition that affects both memory and language. We acknowledge the unique challenges this presents in research and take meaningful steps to ensure that our team remains understanding and accommodating to this disability. We firmly oppose all forms of discrimination based on sex, gender, race, or any other identity marker. Our partnership is grounded in fairness and merit, and we evaluate all contributions based on the quality and impact of the work, not on the individual's background or identity.

Data availability statement

445   Pictures from the Snapshot South Africa program can be requested at Snapshot Safari upon

446   reasonable request. Pictures from the Snapshot Serengeti program are available from the

447   Labeled Image Library of Alexandria – Biology and Conservation:

448   http://lila.science/datasets/snapshot-serengeti. All classification data and metadata are

449   publicly available at Dryad: http://dx.doi.org/10.5061/dryad.5pt92.

Code availability statement

452   The code to analyze and reproduce this study has been deposited in Zenodo and is available

453   online at https://zenodo.org/records/15018760.

References

Adams, D. B., & Kitchen, D. M. (2020). Model vs. playback experiments: The impact of sensory mode on predator☐specific escape responses in saki monkeys. Ethology, 126(5), 563–575. https://doi.org/10.1111/eth.13008

Alberts, S. C. (2019). Social influences on survival and reproduction: Insights from a long☐term study of wild baboons. Journal of Animal Ecology, 88(1), 47–66. https://doi.org/10.1111/1365-2656.12887

Allan, A. T. L., Bailey, A. L., & Hill, R. A. (2021). Consistency in the flight and visual orientation distances of habituated chacma baboons after an observed leopard predation. Do flight initiation distance methods always measure perceived predation risk? Ecology and Evolution, 11(21), 15404–15416. https://doi.org/10.1002/ece3.8237

Allan, A. T. L., LaBarge, L. R., Bailey, A. L., Jones, B., Mason, Z., Pinfield, T., Schröder, F., Whitaker, A., White, A. F., Wilkinson, H., & Hill, R. A. (2024). Behavioural compatibility, not fear, best predicts the looking patterns of chacma baboons. Communications Biology, 7(1), 980. https://doi.org/10.1038/s42003-024-06657-w

Anderson, J. R. (1998). Sleep, sleeping sites, and sleep-related activities: Awakening to their significance. American Journal of Primatology, 46(1), 63–75. https://doi.org/10.1002/(SICI)1098-2345(1998)46:1<63::AID-AJP5>3.0.CO;2-T

Bailey, T. N. (1993). 9. Social Organization of Leopards. In The African Leopard (pp. 240–297). Columbia University Press. https://doi.org/10.7312/bail90198-010

Beaumont, R. A. R.☐; P. B., & Bednarik, R. G. (2015). Concerning a cupule sequence on the edge of the Kalahari Desert in South Africa. 32(2), 163–177. https://doi.org/https://doi.org/10.69978/rar.v32i2.136

Bedetti, A., Greyling, C., Paul, B., Blondeau, J., Clark, A., Malin, H., Horne, J., Makukule, R., Wilmot, J., Eggeling, T., Kern, J., & Henley, M. (2020). System for Elephant Ear-pattern Knowledge (SEEK) to identify individual African elephants. Pachyderm, 61, 63–77. https://doi.org/10.69649/pachyderm.v61i.65

Berger, J. (1979). "Predator Harassment" as a Defensive Strategy in Ungulates. American Midland Naturalist, 102(1), 197. https://doi.org/10.2307/2425087

Bersacola, E., Hill, C. M., Nijman, V., & Hockings, K. J. (2022). Examining primate community occurrence patterns in agroforest landscapes using arboreal and terrestrial camera traps. Landscape Ecology, 37(12), 3103–3121. https://doi.org/10.1007/s10980-022-01524-7

Bezerra, B. M., & Souto, A. (2008). Structure and Usage of the Vocal Repertoire of *Callithrix jacchus*. International Journal of Primatology, 29(3), 671–701. https://doi.org/10.1007/s10764-008-9250-0

Bidner, L. R., Matsumoto□Oda, A., & Isbell, L. A. (2018). The role of sleeping sites in the predator□prey dynamics of leopards and olive baboons. American Journal of Primatology, 80(12). https://doi.org/10.1002/ajp.22932

Blount, J. D., Chynoweth, M. W., Green, A. M., & Şekercioğlu, Ç. H. (2021). Review: COVID-19 highlights the importance of camera traps for wildlife conservation research and management. Biological Conservation, 256, 108984. https://doi.org/10.1016/j.biocon.2021.108984

Boyer-Ontl, K. M., & Pruetz, J. D. (2014). Giving the Forest Eyes: The Benefits of Using Camera Traps to Study Unhabituated Chimpanzees (*Pan troglodytes verus*) in Southeastern Senegal. International Journal of Primatology, 35(5), 881–894. https://doi.org/10.1007/s10764-014-9783-3

Broekhuis, F., Cozzi, G., Valeix, M., McNutt, J. W., & Macdonald, D. W. (2013). Risk

avoidance in sympatric large carnivores: reactive or predictive? Journal of Animal

Ecology, 82(5), 1098–1105. https://doi.org/10.1111/1365-2656.12077

Busse, C. D. (1980). Leopard and Lion predation upon Chacma Baboons living in the

Moremi Wildlife Reserve. Botswana Society, 12, 15–21.

Chassagneux, A., Calenge, C., Marchand, P., Richard, E., Guillaumat, E., Baubet, E., & Saïd,

S. (2020). Should I stay or should I go? Determinants of immediate and delayed

movement responses of female red deer (*Cervus elaphus*) to drive hunts. PLOS ONE,

15(3), e0228865. https://doi.org/10.1371/journal.pone.0228865

Courbin, N., Loveridge, A. J., Macdonald, D. W., Fritz, H., Valeix, M., Makuwe, E. T., &

Chamaillé□Jammes, S. (2016). Reactive responses of zebras to lion encounters shape

their predator–prey space game at large scale. Oikos, 125(6), 829–838.

https://doi.org/10.1111/oik.02555

Cowlishaw, G. (1994). Vulnerability To Predation in Baboon Populations. Behaviour, 131(3–

4), 293–304. https://doi.org/10.1163/156853994X00488

Cowlishaw, G. (1997). Refuge use and predation risk in a desert baboon population. Animal

Behaviour, 54(2), 241–253. https://doi.org/10.1006/anbe.1996.0466

Creel, S. (2018). The control of risk hypothesis: reactive vs. proactive antipredator responses

and stress□mediated vs. food□mediated costs of response. Ecology Letters, 21(7),

947–956. https://doi.org/10.1111/ele.12975

Creel, S., & Christianson, D. (2008). Relationships between direct predation and risk effects.

Trends   in   Ecology   &   Evolution,   23(4),   194–201.

https://doi.org/10.1016/j.tree.2007.12.004

Creel, S., Schuette, P., & Christianson, D. (2014). Effects of predation risk on group size,

vigilance, and foraging behavior in an African ungulate community. Behavioral

Ecology, 25(4), 773–784. https://doi.org/10.1093/beheco/aru050

Creel, S., & Winnie, J. A. (2005). Responses of elk herd size to fine-scale spatial and

temporal variation in the risk of predation by wolves. Animal Behaviour, 69(5), 1181–

1189. https://doi.org/10.1016/j.anbehav.2004.07.022

Cusack, J. J., Dickman, A. J., Kalyahe, M., Rowcliffe, J. M., Carbone, C., MacDonald, D. W.,

& Coulson, T. (2017). Revealing kleptoparasitic and predatory tendencies in an African

mammal community using camera traps: a comparison of spatiotemporal approaches.

Oikos, 126(6), 812–822. https://doi.org/10.1111/oik.03403

Davis, A. L. V., Scholtz, C. H., Kryger, U., Deschodt, C. M., & Strümpher, W. P. (2010).

Dung Beetle Assemblage Structure in Tswalu Kalahari Reserve: Responses to a Mosaic

of Landscape Types, Vegetation Communities, and Dung Types. Environmental

Entomology, 39(3), 811–820. https://doi.org/10.1603/EN09256

Dill, L. M., & Frid, A. (2020). Behaviourally mediated biases in transect surveys: a predation

risk sensitivity approach. Canadian Journal of Zoology, 98(11), 697–704.

https://doi.org/10.1139/cjz-2020-0039

Dröge, E., Creel, S., Becker, M. S., & M'soka, J. (2017). Spatial and temporal avoidance of

risk within a large carnivore guild. Ecology and Evolution, 7(1), 189–199.

https://doi.org/10.1002/ece3.2616

Druelle, F., Aerts, P., Ngawolo, J. C. B., & Narat, V. (2020). Impressive Arboreal Gap-

Crossing Behaviors in Wild Bonobos, (*Pan paniscus*). International Journal of

Primatology, 41(1), 129–140. https://doi.org/10.1007/s10764-020-00140-z

Edwards, S., Gange, A. C., & Wiesel, I. (2016). An oasis in the desert: The potential of water

sources as camera trap sites in arid environments for surveying a carnivore guild.

Journal of Arid Environments, 124, 304–309.

https://doi.org/10.1016/j.jaridenv.2015.09.009

Elliot, N. B., Cushman, S. A., Loveridge, A. J., Mtare, G., & Macdonald, D. W. (2014).

Movements vary according to dispersal stage, group size, and rainfall: the case of the

African lion. Ecology, 95(10), 2860–2869. https://doi.org/10.1890/13-1793.1

Evans, B. E., Mosby, C. E., & Mortelliti, A. (2019). Assessing arrays of multiple trail cameras

to detect North American mammals. PLOS ONE, 14(6), e0217543.

https://doi.org/10.1371/journal.pone.0217543

Fehlmann, G., O'Riain, M. J., Kerr-Smith, C., Hailes, S., Luckman, A., Shepard, E. L. C., &

King, A. J. (2017). Extreme behavioural shifts by baboons exploiting risky, resource-

rich, human-modified environments. Scientific Reports, 7(1), 15057.

https://doi.org/10.1038/s41598-017-14871-2

Fichtel, C., Perry, S., & Gros-Louis, J. (2005). Alarm calls of white-faced capuchin monkeys:

an acoustic analysis. Animal Behaviour, 70(1), 165–176.

https://doi.org/10.1016/j.anbehav.2004.09.020

Fischer, J., Metz, M., Cheney, D. L., & Seyfarth, R. M. (2001). Baboon responses to graded

bark variants. Animal Behaviour, 61(5), 925–931.

https://doi.org/10.1006/anbe.2000.1687

Foster, V. C., Sarmento, P., Sollmann, R., Tôrres, N., Jácomo, A. T. A., Negrões, N., Fonseca,

C., & Silveira, L. (2013). Jaguar and Puma Activity Patterns and Predator‐Prey

Interactions in Four Brazilian Biomes. Biotropica, 45(3), 373–379.

https://doi.org/10.1111/btp.12021

Galbany, J., Tung, J., Altmann, J., & Alberts, S. C. (2015). Canine Length in Wild Male Baboons: Maturation, Aging and Social Dominance Rank. PLOS ONE, 10(5), e0126415. https://doi.org/10.1371/journal.pone.0126415

Gursky, S. L. (2007). The Response of Spectral Tarsiers Toward Avian and Terrestrial Predators. In Primate Anti-Predator Strategies (pp. 241–252). Springer US. https://doi.org/10.1007/978-0-387-34810-0_11

Hammond, P., Gaynor, K., Easter, T., Biro, D., & Carvalho, S. (2025). Landscape□Scale Effects of Season and Predation Risk on the Terrestrial Behavior of Chacma Baboons (*Papio ursinus*). American Journal of Biological Anthropology, 186(4). https://doi.org/10.1002/ajpa.70052

Hammond, P., Lewis□Bevan, L., Biro, D., & Carvalho, S. (2022). Risk perception and terrestriality in primates: A quasi□experiment through habituation of chacma baboons (*Papio ursinus*) in Gorongosa National Park, Mozambique. American Journal of Biological Anthropology, 179(1), 48–59. https://doi.org/10.1002/ajpa.24567

Harvell, C. D. (1990). The Ecology and Evolution of Inducible Defenses. The Quarterly Review of Biology, 65(3), 323–340. https://doi.org/10.1086/416841

Hayward, M. W., Henschel, P., O'Brien, J., Hofmeyr, M., Balme, G., & Kerley, G. I. H. (2006). Prey preferences of the leopard (*Panthera pardus*). Journal of Zoology, 270(2), 298–313. https://doi.org/10.1111/j.1469-7998.2006.00139.x

Herrera, D. J., Moore, S. M., Herrmann, V., McShea, W. J., & Cove, M. V. (2021). A shot in the dark: White and infrared LED flash camera traps yield similar detection probabilities for common urban mammal species. *Hystrix*, the Italian Journal of Mammalogy, 32(1), 72–75. https://doi.org/10.4404/hystrix-00429-2021

Hill, R. A., & Weingrill, T. (2007). Predation Risk and Habitat Use in Chacma Baboons

(*Papio hamadryas ursinus*). In Primate Anti-Predator Strategies (pp. 339–354).

Springer US. https://doi.org/10.1007/978-0-387-34810-0_16

Horrocks, J. A., & Hunte, W. (1986). Sentinel behaviour in vervet monkeys: Who sees whom

first? Animal Behaviour, 34(5), 1566–1568. https://doi.org/10.1016/S0003-

3472(86)80226-3

Hubel, T. Y., Golabek, K. A., Rafiq, K., McNutt, J. W., & Wilson, A. M. (2018). Movement

patterns and athletic performance of leopards in the Okavango Delta. Proceedings of

the Royal Society B: Biological Sciences, 285(1877), 20172622.

https://doi.org/10.1098/rspb.2017.2622

Humphries, D. A., & Driver, P. M. (1967). Erratic Display as a Device against Predators.

Science, 156(3783), 1767–1768. https://doi.org/10.1126/science.156.3783.1767

Isbell, L. A., Bidner, L. R., Van Cleave, E. K., Matsumoto-Oda, A., & Crofoot, M. C. (2018).

GPS-identified vulnerabilities of savannah-woodland primates to leopard predation and

their implications for early hominins. Journal of Human Evolution, 118, 1–13.

https://doi.org/10.1016/j.jhevol.2018.02.003

Johnson, S. E. (2003). Life history and the competitive environment: trajectories of growth,

maturation, and reproductive output among Chacma baboons. American Journal of

Physical Anthropology, 120(1), 83–98. https://doi.org/10.1002/ajpa.10139

Jooste, E., Pitman, R. T., van Hoven, W., & Swanepoel, L. H. (2013). Unusually High

Predation on Chacma Baboons (*Papio ursinus*) by Female Leopards (*Panthera pardus*)

in the Waterberg Mountains, South Africa. Folia Primatologica, 83(3–6), 353–360.

https://doi.org/10.1159/000339644

Keim, J. L., Lele, S. R., DeWitt, P. D., Fitzpatrick, J. J., & Jenni, N. S. (2019). Estimating the
intensity of use by interacting predators and prey using camera traps. Journal of Animal
Ecology, 88(5), 690–701. https://doi.org/10.1111/1365-2656.12960

Kie, J. G. (1999). Optimal Foraging and Risk of Predation: Effects on Behavior and Social
Structure in Ungulates. Journal of Mammalogy, 80(4), 1114–1129.
https://doi.org/10.2307/1383163

Kolowski, J. M., Katan, D., Theis, K. R., & Holekamp, K. E. (2007). Daily Patterns of
Activity in the Spotted Hyena. Journal of Mammalogy, 88(4), 1017–1028.
https://doi.org/10.1644/06-MAMM-A-143R.1

Kramer, D. L., & Bonenfant, M. (1997). Direction of predator approach and the decision to
flee to a refuge. Animal Behaviour, 54(2), 289–295.
https://doi.org/10.1006/anbe.1996.0360

Krebs, J. R., & Davies, N. B. (Eds.). (2009). *Behavioural ecology: an evolutionary approach*.
John Wiley & Sons.

LaBarge, L. R., Allan, A. T. L., Berman, C. M., Hill, R. A., & Margulis, S. W. (2022).
Cortisol metabolites vary with environmental conditions, predation risk, and human
shields in a wild primate, Cercopithecus albogularis. Hormones and Behavior, 145,
105237. https://doi.org/10.1016/j.yhbeh.2022.105237

Laundré, J. W., Hernández, L., & Altendorf, K. B. (2001). Wolves, elk, and bison:
reestablishing the "landscape of fear" in Yellowstone National Park, U.S.A. Canadian
Journal of Zoology, 79(8), 1401–1409. https://doi.org/10.1139/z01-094

Li, Y. (2007). Terrestriality and tree stratum use in a group of Sichuan snub-nosed monkeys.
Primates, 48(3), 197–207. https://doi.org/10.1007/s10329-006-0035-9

Lima, S. L., & Dill, L. M. (1990). Behavioral decisions made under the risk of predation: a review and prospectus. Canadian Journal of Zoology, 68(4), 619–640. https://doi.org/10.1139/z90-092

Linkie, M., & Ridout, M. S. (2011). Assessing tiger–prey interactions in Sumatran rainforests. Journal of Zoology, 284(3), 224–229. https://doi.org/10.1111/j.1469-7998.2011.00801.x

MacCormick, H. A., MacNulty, D. R., Bosacker, A. L., Lehman, C., Bailey, A., Anthony Collins, D., & Packer, C. (2012). Male and female aggression: lessons from sex, rank, age, and injury in olive baboons. Behavioral Ecology, 23(3), 684–691. https://doi.org/10.1093/beheco/ars021

Martins, Q., & Harris, S. (2013). Movement, activity and hunting behaviour of leopards in the Cederberg mountains, South Africa. African Journal of Ecology, 51(4), 571–579. https://doi.org/10.1111/aje.12068

Meek, P. D., Ballard, G., Claridge, A., Kays, R., Moseby, K., O'Brien, T., O'Connell, A., Sanderson, J., Swann, D. E., Tobler, M., & Townsend, S. (2014). Recommended guiding principles for reporting on camera trapping research. Biodiversity and Conservation, 23(9), 2321–2343. https://doi.org/10.1007/s10531-014-0712-8

Meek, P. D., Ballard, G., Fleming, P., & Falzon, G. (2016). Are we getting the full picture? Animal responses to camera traps and implications for predator studies. Ecology and Evolution, 6(10), 3216–3225. https://doi.org/10.1002/ece3.2111

Newey, S., Davidson, P., Nazir, S., Fairhurst, G., Verdicchio, F., Irvine, R. J., & van der Wal, R. (2015). Limitations of recreational camera traps for wildlife management and conservation research: A practitioner's perspective. Ambio, 44, 624–635. https://doi.org/10.1007/s13280-015-0713-1

Niedballa, J., Wilting, A., Sollmann, R., Hofer, H., & Courtiol, A. (2019). Assessing analytical methods for detecting spatiotemporal interactions between species from camera trapping data. Remote Sensing in Ecology and Conservation, 5(3), 272–285. https://doi.org/10.1002/rse2.107

Nowak, K., le Roux, A., Richards, S. A., Scheijen, C. P. J., & Hill, R. A. (2014). Human observers impact habituated samango monkeys' perceived landscape of fear. Behavioral Ecology, 25(5), 1199–1204. https://doi.org/10.1093/beheco/aru110

Palmer, M. S., Gaynor, K. M., Becker, J. A., Abraham, J. O., Mumma, M. A., & Pringle, R. M. (2022). Dynamic landscapes of fear: understanding spatiotemporal risk. Trends in Ecology & Evolution, 37(10), 911–925. https://doi.org/10.1016/j.tree.2022.06.007

Palmer, M. S., & Packer, C. (2021). Reactive anti-predator behavioral strategy shaped by predator characteristics. PLOS ONE, 16(8), e0256147. https://doi.org/10.1371/journal.pone.0256147

Pardo, L. E., Bombaci, S., Huebner, S. E., Somers, M. J., Fritz, H., Downs, C., Guthmann, A., Hetem, R. S., Keith, M., le Roux, A., Mgqatsa, N., Packer, C., Palmer, M. S., Parker, D. M., Peel, M., Slotow, R., Maartin Strauss, W., Swanepoel, L., Tambling, C., … Venter, J. A. (2021). Snapshot Safari: A large-scale collaborative to monitor Africa's remarkable biodiversity. South African Journal of Science, 117(1–2). https://doi.org/10.17159/SAJS.2021/8134

Rutherford, M. C., Mucina, L., Lötter, M. C., Bredenkamp, G. J., Smit, J. H. L., Scott-Shaw, C. R., Hoare, D. B., Goodman, P. S., Bezuidenhout, H., Scott, L., Ellis, F., Powrie, L., Siebert, F., Mostert, T. H., Henning, B. J., Ventner, C. E., Camp, K. G. T., Siebert, S., Matthews, W., & Hurter, J. (2006). Savanna Biome. In L. Mucina & M. C. Rutherford (Eds.), The Vegetation of South Africa, Lesotho and Swaziland. SANBI.

Sankaran, M., Hanan, N. P., Scholes, R. J., Ratnam, J., Augustine, D. J., Cade, B. S.,

Gignoux, J., Higgins, S. I., Le Roux, X., Ludwig, F., Ardo, J., Banyikwa, F., Bronn, A.,

Bucini, G., Caylor, K. K., Coughenour, M. B., Diouf, A., Ekaya, W., Feral, C. J., …

Zambatis, N. (2005). Determinants of woody cover in African savannas. Nature,

438(7069), 846–849. https://doi.org/10.1038/nature04070

Say-Sallaz, E., Chamaillé-Jammes, S., Périquet, S., Loveridge, A. J., Macdonald, D. W.,

Antonio, A., Fritz, H., & Valeix, M. (2023). Large carnivore dangerousness affects the

reactive spatial response of prey. Animal Behaviour, 202, 149–162.

https://doi.org/10.1016/j.anbehav.2023.05.014

Scholes, R. J., & Archer, S. R. (1997). Tree-Grass Interactions in Savannas. Annual Review

of Ecology and Systematics, 28(1), 517–544.

https://doi.org/10.1146/annurev.ecolsys.28.1.517

Schreier, A. L., & Swedell, L. (2012). Ecology and sociality in a multilevel society:

Ecological determinants of spatial cohesion in hamadryas baboons. American Journal

of Physical Anthropology, 148(4), 580–588. https://doi.org/10.1002/ajpa.22076

Searle, C. E., Smit, J. B., Cusack, J. J., Strampelli, P., Grau, A., Mkuburo, L., Macdonald, D.

W., Loveridge, A. J., & Dickman, A. J. (2021). Temporal partitioning and

spatiotemporal avoidance among large carnivores in a human-impacted African

landscape. PLOS ONE, 16(9), e0256876. https://doi.org/10.1371/journal.pone.0256876

Seiler, M., Schwitzer, C., & Holderied, M. (2013). Anti-predator behaviour of Sahamalaza

sportive lemurs, (*Lepilemur sahamalazensis*), at diurnal sleeping sites. Contributions to

Zoology, 82(3), 131-S1. https://doi.org/10.1163/18759866-08203003

Seyfarth, R. M., Cheney, D. L., & Marler, P. (1980). Vervet monkey alarm calls: Semantic

communication in a free-ranging primate. Animal Behaviour, 28(4), 1070–1094.

https://doi.org/10.1016/S0003-3472(80)80097-2

Shannon, G., Cordes, L. S., Hardy, A. R., Angeloni, L. M., & Crooks, K. R. (2014).

Behavioral Responses Associated with a Human-Mediated Predator Shelter. PLoS

ONE, 9(4), e94630. https://doi.org/10.1371/journal.pone.0094630

Si, X., Kays, R., & Ding, P. (2014). How long is enough to detect terrestrial animals?

Estimating the minimum trapping effort on camera traps. PeerJ, 2, e374.

https://doi.org/10.7717/peerj.374

Smith, J. A., Suraci, J. P., Hunter, J. S., Gaynor, K. M., Keller, C. B., Palmer, M. S., Atkins, J.

L., Castañeda, I., Cherry, M. J., Garvey, P. M., Huebner, S. E., Morin, D. J., Teckentrup,

L., Weterings, M. J. A., & Beaudrot, L. (2020). Zooming in on mechanistic predator–

prey ecology: Integrating camera traps with experimental methods to reveal the drivers

of ecological interactions. Journal of Animal Ecology, 89(9), 1997–2012.

https://doi.org/10.1111/1365-2656.13264

Stone, O. M. L., Laffan, S. W., Curnoe, D., & Herries, A. I. R. (2013). The Spatial

Distribution of Chacma Baboon (*Papio ursinus*) Habitat Based on an Environmental

Envelope Model. International Journal of Primatology, 34(2), 407–422.

https://doi.org/10.1007/s10764-013-9669-9

Suscke, P., Presotto, A., & Izar, P. (2021). The role of hunting on Sapajus xanthosternos"</i>

landscape of fear in the Atlantic Forest, Brazil. American Journal of Primatology, 83(5).

https://doi.org/10.1002/ajp.23243

Swanson, A., Kosmala, M., Lintott, C., Simpson, R., Smith, A., & Packer, C. (2015).

Snapshot Serengeti, high-frequency annotated camera trap images of 40 mammalian

species in an African savanna. Scientific Data, 2(1), 150026.
https://doi.org/10.1038/sdata.2015.26

Swinkels, C., van der Wal, J. E. M., Stinn, C., Monteza-Moreno, C. M., & Jansen, P. A.
(2023). Prey tracking and predator avoidance in a Neotropical moist forest: a camera-
trapping approach. Journal of Mammalogy, 104(1), 137–145.
https://doi.org/10.1093/jmammal/gyac091

Szott, I. D., Pretorius, Y., Ganswindt, A., & Koyama, N. F. (2020). Physiological stress
response of African elephants to wildlife tourism in Madikwe Game Reserve, South
Africa. Wildlife Research, 47(1), 34. https://doi.org/10.1071/WR19045

Thaker, M., Vanak, A. T., Owen, C. R., Ogden, M. B., Niemann, S. M., & Slotow, R. (2011).
Minimizing predation risk in a landscape of multiple predators: Effects on the spatial
distribution of African ungulates. Ecology, 92(2), 398–407. https://doi.org/10.1890/10-
0126.1

Tjaden□McClement, K., Gharajehdaghipour, T., Shores, C., White, S., Steenweg, R.,
Bourbonnais, M., Konanz, Z., & Burton, A. C. (2025). Mixed evidence for
disturbance□mediated apparent competition for declining caribou in western British
Columbia, Canada. The Journal of Wildlife Management, 89(6).
https://doi.org/10.1002/jwmg.70040

Trinkel, M. (2010). Prey selection and prey preferences of spotted hyenas Crocuta crocuta in
the Etosha National Park, Namibia. Ecological Research, 25(2), 413–417.
https://doi.org/10.1007/s11284-009-0669-3

van Casteren, A., Sellers, W. I., Thorpe, S. K. S., Coward, S., Crompton, R. H., Myatt, J. P.,
& Ennos, A. R. (2012). Nest-building orangutans demonstrate engineering know-how

to produce safe, comfortable beds. Proceedings of the National Academy of Sciences, 109(18), 6873–6877. https://doi.org/10.1073/pnas.1200902109

van Schaik, C. P., Bshary, R., Wagner, G., & Cunha, F. (2022). Male anti predation services in primates as costly signalling? A comparative analysis and review. Ethology, 128(1), 1–14. https://doi.org/10.1111/eth.13233

Virgadamo, P., Hodosh, M., Povar, M., & Shklar, G. (1972). The Dentition of *Papio anubis*. Journal of Dental Research, 51(5), 1338–1345. https://doi.org/10.1177/00220345720510051501

Von Hippel, F. A. (1998). Use of sleeping trees by black and white Colobus monkeys (*Colobus guereza*) in the Kakamega Forest, Kenya. American Journal of Primatology, 45(3), 281–290. https://doi.org/10.1002/(SICI)1098-2345(1998)45:3<281::AID-AJP4>3.0.CO;2-S

Wentworth, J. C., Tambling, C. J., & Kerley, G. I. H. (2011). Evidence for prey selection by spotted hyaena in the Eastern Cape, South Africa. Acta Theriologica, 56(4), 389–392. https://doi.org/10.1007/s13364-011-0033-1

Weterings, M. J. A., Zaccaroni, M., van der Koore, N., Zijlstra, L. M., Kuipers, H. J., van Langevelde, F., & van Wieren, S. E. (2016). Strong reactive movement response of the medium-sized European hare to elevated predation risk in short vegetation. Animal Behaviour, 115, 107–114. https://doi.org/10.1016/j.anbehav.2016.03.011

Willems, E. P., & Hill, R. A. (2009). Predator specific landscapes of fear and resource distribution: effects on spatial range use. Ecology, 90(2), 546–555. https://doi.org/10.1890/08-0765.1

Zhang, Y., & Zhao, Q. (2023). What is a Randomization Test? Journal of the American

Statistical   Association,   118(544),   2928–2942.

https://doi.org/10.1080/01621459.2023.2199814

Zinner, D., Buba, U., Nash, S., & Roos, C. (2011). Pan-African Voyagers: The

Phylogeography of Baboons. In Primates of Gashaka (pp. 319–358). Springer New

York. https://doi.org/10.1007/978-1-4419-7403-7_7

Tables

Table 1. Number of detections for the species of interest: chacma baboon (*Papio ursinus*), olive baboon (*Papio anubis*), Burchell's zebra (*Equus quagga*), spotted hyena (*Crocuta crocuta*), leopard (*Panthera pardus*), and lion (*Panthera leo*), in Associated Private Nature Reserves (APN), Madikwe Game Reserve (MAD), Tswalu Kalahari Private Wildlife Reserve (TSW) (South Africa, data collected between 2017 and 2019), and Serengeti National Park (SER) (Tanzania, data collected between 2011 and 2013). A 30-minute time-to-independence filter was applied to predator detections (see Methods section for details).

Species	Scientific name	Number of detections in MAD	Number of detections in TSW	Number of detections in APN	Number of detections in SER
Lion	*Panthera leo*	17	51	188	1,276
Leopard	*Panthera pardus*	28	8	159	160
Spotted hyena	*Crocuta crocuta*	110	9	1,570	3,382
Chacma baboon	*Papio ursinus*	346	738	500	0
Olive baboon	*Papio anubis*	0	0	0	1,458
Burchell's zebra	*Equus quagga*	3,602	1,394	2,047	60,212

Table 2. Significance values (*p*) calculated by comparing the observed number of detections

of chacma baboon (*Papio ursinus*) and olive baboon (*Papio anubis*) before and after predator

detections: spotted hyena (*Crocuta crocuta*), leopard (*Panthera pardus*), and lion (*Panthera*

leo), to the distribution of number of detections produced by randomizing predator detections

under the assumption of non-avoidance, across six consecutive 24-hour time-blocks relative

to the predator detection (72 to 48 hours before, 48 to 24 hours before, 24 to 0 hours before, 0

to 24 hours after, 24 to 48 hours after, 48 to 72 hours after) in the four study sites: Associated

Private Nature Reserves (APN), Madikwe Game Reserve (MAD), Tswalu Kalahari Private

Wildlife Reserve (TSW) (South Africa, data collected between 2017 and 2019), and

Serengeti National Park (SER) (Tanzania, data collected between 2011 and 2013).

Study site	Prey species	Predator species	-72 to -48 hours	-48 to -24 hours	-24 to 0 hours	0 to 24 hours	24 to 48 hours	48 to 72 hours
MAD	Chacma baboon	Lion	0.503	1.000	1.000	0.900	0.469	0.452
		Leopard	1.000	0.105	1.000	0.808	0.802	0.496
		Spotted hyena	0.877	1.000	0.904	0.742	0.251	0.107
TSW	Chacma baboon	Lion	0.451	0.288	0.033 *	0.925	0.664	0.354
		Leopard	1.000	1.000	0.149	0.792	0.647	0.773
		Spotted hyena	1.000	1.000	1.000	0.912	0.714	0.722
APN	Chacma baboon	Lion	0.671	0.001 *	1.000	0.526	0.251	0.373
		Leopard	0.547	0.739	0.597	0.573	0.114	0.600
		Spotted hyena	0.886	0.072	0.007 *	0.225	0.498	0.144
SER	Olive Baboon	Lion	0.869	0.781	0.160	0.098	0.460	0.679
		Leopard	0.551	0.434	0.012 *	0.599	0.141	0.788
		Spotted hyena	0.574	0.317	0.008 *	0.984	0.400	1.000

804   * Significance to the level alpha = 0.05.

 Figure captions

Figure 1. Map of the four study sites in South Africa (data collected between 2017 and 2019)

and Tanzania (data collected between 2011 and 2013). A) Madikwe Private Game Reserve

(MAD; n = 40 camera traps); B) Serengeti National Park (SER; n = 219 camera traps); C)

Tswalu Kalahari Game Reserve (TSW; n = 31 camera traps); D) Associated Private Nature

Reserves (APN; n = 56 camera traps). The green boundaries represent the borders of the

study areas, the red circles represent the camera trap locations, with their assigned code

printed above. Chacma baboon (*Papio ursinus*) occurs in the three South African sites, olive

baboon (*Papio anubis*) occurs in the Tanzanian site.

Figure 2. Distribution of Burchell's zebra (*Equus quagga*) detections obtained from the

randomized datasets (n = 1,000) before and after lion (*Panthera leo*) detection in Associated

Private Nature Reserves (APN), South Africa (data collected between 2017 and 2019). Each

panel corresponds to one of six consecutive 24-hour time-blocks relative to the predator

detection: 72 to 48 hours before, 48 to 24 hours before, 24 to 0 hours before, 0 to 24 hours

after, 24 to 48 hours after, 48 to 72 hours after. The red vertical lines represent the observed

number of Burchell's zebra detections for each time-block. The shaded area corresponds to

the density of randomization trials, the area under the curve thus being equal to 1. *:

significance to the level alpha = 0.05.

Figure 3. Distribution of baboon (*Papio ursinus* and *Papio anubis*) detections obtained from

the randomized datasets (n = 1,000) before and after predator detection: spotted hyena

(*Crocuta crocuta*), leopard (*Panther pardus*), and lion (*Panthera leo*), across the four study

sites: Associated Private Nature Reserves (APN), Madikwe Game Reserve (MAD), Tswalu

Kalahari Private Wildlife Reserve (TSW) (South Africa, data collected between 2017 and

2019), and Serengeti National Park (SER) (Tanzania, data collected between 2011 and 2013).

Each panel corresponds to one of the six consecutive 24-hour time-blocks relative to the

predator detection: 72 to 48 hours before, 48 to 24 hours before, 24 to 0 hours before, 0 to 24

hours after, 24 to 48 hours after, 48 to 72 hours after. The red vertical line represents the

observed number of baboon detections for each site/predator/time-block combination. The

shaded area corresponds to the density of randomization trials, the area under the curve thus

being equal to 1. *: significance to the level alpha = 0.05.

Figures

Figure 1

Figure 2

Figure 3

Supplementary material

Supplementary Figure 1. Distribution of significance values (*p*) from the randomization test

assessing the effect of lions (*Panthera leo*) on Burchell's zebra (*Equus quagga*) detections in

Associated Private Nature Reserves (APN), South Africa (data collected between 2017 and

2019) within the 0 to 24 hours following lion detection. The test was conducted on randomly

down-sampled Burchell's zebra detections datasets (n = 100). The red dashed line indicates

the significance value threshold of alpha = 0.05, marking the significance level for detecting

deviations from the assumption of non-avoidance.

43