# Absence of reactive temporal response of baboons to the occurrence of predators.

N. van Rooyen[1,2, *], L. Thel[3, *], J.A. Venter[3], H. Fritz[1,2], F. Prugnolle[1,2,#], V. Rougeron[1,2,#]

**Affiliations**

[1] International Research Laboratory, REHABS, CNRS-NMU-UCBL, George Campus, Nelson Mandela University, Madiba drive, 6529 George, South Africa

[2] Sustainability Research Unit, George Campus, Nelson Mandela University, George Campus, Madiba drive, 6529 George, South Africa

[3] Department of Conservation Management, Faculty of Science, Nelson Mandela University, George Campus, George, South Africa

* **Co-first authors:** N. van Rooyen[1,2, *], L. Thel[3, *]

# **Co-supervised the work:** F. Prugnolle[1,2,] V. Rougeron[1,2,#]

**Corresponding authors:**

Rougeron Virginie, International Research Laboratory, REHABS, CNRS-NMU-UCBL, Sustainability Research Unit, George Campus, Nelson Mandela University, George, South Africa, rougeron.virginie@gmail.com/virginie.rougeron@cnrs.fr;

19   Nicholas van Rooyen, Department of conservation management, faculty of science,

20   Nelson Mandela University George Campus, Nelson Mandela University, George,

21   South Africa, s216653207@mandela.ac.za

# Abstract

Predation exerts a strong selective pressure on prey species, shaping their behavioral adaptations. While proactive strategies such as habitat selection and increased vigilance allow prey to anticipate threats, reactive responses, including immediate flight or retaliation, occur on direct detection of a predator. Baboons (*Papio spp.*) encounter various predators across their extensive range, but their behavioral responses to these threats remain poorly understood. Using camera trap data from four savanna sites—three in South Africa (Madikwe Game Reserve, Tswalu Kalahari Private Wildlife Reserve, and the Associated Private Nature Reserves) and one in Tanzania (Serengeti National Park)—this study examined whether chacma (*P. ursinus*) and olive (*P. anubis*) baboons exhibit reactive temporal avoidance following the occurrence of their main predators, lions (*Panthera leo*), leopards (*Panthera pardus*), and spotted hyenas (*Crocuta crocuta*). Using a randomization test, we analysed baboon presence patterns within three 24-hour time blocks before and after the occurrence of a predator. There was no significant decrease in baboon presence following a predator event, suggesting the absence of reactive temporal avoidance in baboons. These results suggest that baboons may rely on alternative predator-avoidance strategies, such as early detection, sentinel behavior, or aggressive retaliation. This study highlights the importance of sample size in detecting behavioral responses and highlights the possible limitations of camera traps in capturing arboreal escape strategies.

**Keywords:** predator-prey interactions, *Papio spp.*, randomization test, anti-predator behavior, camera trap

## Introduction

Predation exerts a strong selective pressure on prey species due to its impact on individual fitness and survival (Humphries & Driver, 1967; Harvell, 1990; Krebs & Davies, 2009). In response, prey species have evolved a diverse range of behavioural responses to enhance their survival as they encounter fluctuating predation risk in their environment (Lima & Dill, 1990; Kie, 1999). Behavioural adaptations are inherently flexible and allow prey to modify the nature and intensity of their responses based on their spatial and temporal perception of risk in the surrounding environment (i.e., the landscape of fear, Laundré et al., 2001; Creel & Winnie, 2005; Palmer et al., 2022).

Behavioral adaptations can be classified into proactive and reactive responses (Broekhuis et al., 2013; Courbin et al., 2016; Creel, 2018; Palmer et al., 2021; Say-Sallaz et al., 2023). Proactive responses are behavioral strategies prey adopt before encountering a predator to reduce predation risk by anticipating threats rather than reacting after a predator appears (Creel et al., 2008; Broekhuis et al., 2013; Creel et al., 2014). These adjustments operate over broad spatial and temporal scales, such as home ranges or seasonal cycles (Droge et al., 2016). For example, Burchell's zebra (*Equus quagga burchellii*) regularly move away from previous lion (*Panthera leo*) encounter sites, such as water holes, to reduce the risk of predation (Thaker et al., 2011; Courbin et al., 2016; Say-Sallaz et al., 2023). Reactive responses, in contrast, are immediate reactions triggered by the direct detection of a predator (Broekhuis et al., 2013). These responses occur on smaller spatial and temporal scales, such as foraging sites and across diel activity periods (Droge et al., 2016), driven by real-time threats. These responses include fleeing the

area, alerting surrounding group members, or engaging in aggressive retaliation towards the threat (Cowlishaw, 1994; Lingle & Pellis, 2002; Isbell et al., 2018). For example, white-tailed deer (*Odocoileus virginianus*) flee upon detecting coyotes (*Canis latrans*), whereas mule deer (*Odocoileus hemionus*) may choose to actively retaliate (Lingle & Pellis, 2002).

Non-human primates (NHPs), as a prey species, have evolved different strategies in response to predation pressure. Some species such as the hamadryas baboons (*Papio hamadryas*), anticipate by forming large coalitions across troops at roosting sites to enhance group protection (Schreier & Swedell, 2012). In white-faced capuchins (*Cebus capucinus*), individuals increase vigilance near the ground, as they are more exposed to terrestrial predators (Campos & Fedigan 2014). Regarding reactive responses, many species use alarm-call systems as a response to the occurrence of a predator, such as white-faced capuchin and vervet monkeys (*Chlorocebus pygerythrus*) (Seyfarth et al., 1980; Fichtel et al., 2005). These calls help individuals select an appropriate response, such as fleeing, hiding, attacking the predator, or even attempting to rescue captured group members (Perry et al., 2003; Meno et al., 2013). Baboons (*Papio spp.),* bonobos (*Pan paniscus*) and colobine monkeys (*Colobus guereza*) seek refuge in elevated areas as a reactive response, relying on tree canopies and/or cliffs as escape routes while foraging on the ground, to reduce their vulnerability to terrestrial predators (Cowlishaw, 1994; Von Hippel, 1998; Isbell et al., 2018; Druelle et al., 2020).

Among NHP species, baboons encounter a diverse range of predators due to their broad geographic distribution (Buse, 1980; Cowlishaw, 1994; Jooste et al., 2013; Isbell et al., 2018). In southern, eastern and central Africa, their predators include large carnivores such as lions, leopards (*Panthera pardus*) and spotted

hyenas (*Crocuta crocuta*) (Busse, 1980; Jooste et al., 2013; Dröge et al., 2017; Isbell et al., 2018). Although these predators infrequently prey on baboons (Busse, 1980; Jooste et al., 2013; Dröge et al., 2017; Isbell et al., 2018), they nevertheless contribute to baboon mortality (Busse, 1980; Cowlishaw, 1994). Leopard and lion are both ambush predator species, but their hunting strategies differ as lions have been observed hunting baboons during the day, whereas leopards hunt both day and night, often targeting baboons at their roosting sites (Busse, 1980; Isbell et al., 2018). Spotted hyenas are cursorial predators (Trinkel, 2010; Wentworth et al., 2011) and although the evidence suggests that they do kill and consume baboons as prey, they may only target baboons when the opportunity arises as opposed to actively hunting them (Buse 1980; Cowlishaw, 1994; Trinkel, 2010; Wentworth et al., 2011).

These different predation strategies necessitate different vigilance and escape responses, leading baboons to employ both proactive and reactive responses to mitigate predation risk. Proactively, baboons increase vigilance when away from roost sites or when isolated from group members, enhancing threat detection in vulnerable situations (Fischer et al., 2001). Males also position themselves at the troop's periphery, acting as a guardian for more vulnerable troop members (Surie et al., 2023). Reactively, baboons demonstrate adaptability through aggressive predator mobbing, where they collectively confront and harass predators to deter attacks (Bailey, 1993; Cowlishaw, 1994; Jooste et al., 2013; Isbell et al., 2018). However, it remains unclear whether baboons actively move away from predator encounter sites as part of their antipredator strategies (Allan et al., 2021).

Most studies on baboon-predator interactions have primarily used direct in-field observations (Busse, 1980; Hill et al., 2007; Allan et al., 2021). While these methods provide valuable insights, they can be invasive, costly, and may influence

animal behavior due to the observer's presence (Busse, 1980; Basille et al., 2015;

Latombe et al., 2014). Camera traps offer a non-invasive, cost-effective alternative

for long-term monitoring across large areas, while minimising human disturbance

(Newey et al., 2015).

This study analyses camera trap data from three study sites in South Africa

and one in Tanzania, to test whether olive baboons (*Papio anubis*) and chacma

baboons (*Papio ursinus*) exhibit reactive temporal avoidance in response to

predators. Specifically, we test two hypotheses: baboons' presence decreases

following the occurrence of a predator (H1). The intensity of this response varies

based on the predator's hunting strategy (H2): we predicted a greater decrease in

baboons' presence after the occurrence of an ambush predator (lions and leopards)

than after a cursorial predator (spotted hyenas). As semi-arboreal animals, baboons

can escape cursorial predators by seeking refuge in elevated areas such as trees or

cliffs, however leopards may continue their pursuit into a tree, thereby forcing the

baboon troop to move out of the area. (Busse, 1980; Trinkel, 2010; Wentworth et al.,

2011; Isbell et al., 2018).

## Materials and methods

*Study site selection and description*

This study utilized data from the ongoing camera trap monitoring project

*Snapshot Safari*, which spans multiple countries (Swanson et al., 2015; Pardo et al.,

2021). We used data from three sites in South Africa (Pardo et al., 2021) and one

site in Tanzania (Swanson et al., 2015, data publicly available at Dryad:

). The focus was on two baboon species:

chacma baboons, primarily distributed in southern Africa, comprising South Africa,

Botswana, Zimbabwe and Namibia (Stone et al., 2013), and olive baboons, which

occupy the northern regions of central Africa, ranging from Mali to Ethiopia, Kenya,

and north-western Tanzania (Zinner et al., 2011).

Of the 32 sites available from the *Snapshot Safari* project, those included in

this study were selected based on data availability. Sites were selected only if all

three target predators (lions, leopards, and spotted hyenas) were present along with

baboons. After such selection, the southern African study sites included Madikwe

Game Reserve (MAD, 24°49'S - 26°13'E; Figure 1A), Tswalu Kalahari Private

Wildlife Reserve (TSW, 27°14'S - 22°23'E; Figure 1B), and the Associated Private

Nature Reserves (APNR, 24°20'S - 31°20'E; Figure 1C). Serengeti National Park

(SER, 2°19'S - 34°49'E; Figure 1D), in Tanzania, was the sole study site for olive

baboons. All four study sites were located within the savanna biome, characterised

by open grasslands interspersed with trees (Sankaran et al., 2005).

MAD is a 750 km² fenced game reserve located in the Northwest Province, on

the border of South Africa and Botswana, within the central bushveld bioregion of the

savanna biome (Szott et al., 2020). The landscape is characterized by ridges and

thorny woodlands dominated by acacia species (Szott et al., 2020). TSW, located in

the Northern Cape Province, is a 1,020 km² fenced reserve in the Eastern Kalahari

bioregion, with portions extending into the Kalahari Dunefields. Elevations range

from 1,020 m a.s.l. in lower-lying areas to 1,586 m a.s.l. at the highest point (Davis et

al., 2010). The vegetation consists of discontinuous annual and perennial grasses

interspersed with deep-rooted trees, such as the camel thorn (*Vachellia erioloba*)

(Beaumont & Bednarik, 2015). APNR is a collective of five interconnected reserves

(Timbavati, Thornybush, Klaserie, Balule, and Umbabat Private Nature Reserves)

covering 1,800 km² in the Mopaneveld bioregion of Limpopo and Mpumalanga

provinces. It is open to Kruger National Park (Bedetti et al., 2020; Rutherford et al.,

2006). The APNR vegetation is highly variable, with isolated tree canopies and a

continuous grass understory (Scholes & Archer, 1997). SER, located in Tanzania,

spans 25,000 km² and is part of the East African savanna biome, characterized by

vast grasslands interspersed with scattered trees (Swanson, 2015).

*Camera trap data*

The data for MAD, TSW, and APNR were collected between June 2017 and

November 2019 (29 months), while data for SER were collected between June 2010

and May 2013 (35 months). Camera trap placement followed a systematic layout in

MAD, TSW, and SER, where camera traps were positioned at the center of a 5 km²

grid pattern, whereas camera trap placement in the APNR followed a random

allocation layout (sensu Meek et al., 2014), based on habitat and geography (see

Pardo et al., 2021 for a more detailed description of the camera-trap network in the

South African sites; and Swanson et al., 2015 for the Tanzanian site). A total of 355

camera traps (Cuddeback's Professional Series x-change, white flash in South

Africa, and Scoutguard (SG565) incandescent cameras in SER) were deployed

across the four study sites.

In MAD, TSW and APNR, each camera trap was directed with the infrared

trigger facing towards game paths where animal presence had been confidently

identified through the presence of scats and/or spoor. In SER, cameras were

directed based on vegetation cover, avoiding areas of high grass density to avoid

blank images (i.e. a frame or sequence of frames that do not contain animals). The

camera traps were enclosed in steel casings and fixed at a height of 0.5-1 m above the ground (Swanson et al., 2015; Pardo et al., 2021). To minimize glare caused by the rising and setting sun, the camera traps were oriented to face as far North or South as possible. Each camera trap was programmed to take three images for every motion detection during the day and one at night (Pardo et al., 2021; Swanson et al., 2015). The sensitivity of each camera trap was set to a medium level (50-75%) to minimize false triggers caused by windblown vegetation. No lures were used to attract any species to the camera trap sites.

Capture events (i.e. any single image or consecutive series of images initiated by an animal-trigger, regardless of the number of individuals present in the image) were classified using two methods: (i) the citizen science program Zooniverse (www.zooniverse.org; Swanson et al., 2015), which utilizes the general public to identify animal species in camera trap images, was used exclusively for data collected from SER; (ii) for the three South African sites, species identification was performed by trained laboratory technicians using TrapTagger (www.wildeyeconservation.org).

Five camera traps were removed from the MAD dataset that were positioned specifically near waterholes, as predator-prey interactions may likely differ at waterholes due to the trade-off between the need for scarce resources (i.e. water) and predator avoidance (Edwards et al., 2016). The present study used a total of 350 camera traps, distributed as follows: ($n_{MAD}$ = 41; $n_{TSW}$ = 28; $n_{APNR}$ = 56; $n_{SER}$ = 225). After removing duplicate capture events, the final number of capture events for each of the species of interest was: chacma baboons (n = 1,694), olive baboons (n = 1,556), lions (n = 4,554), leopards (n = 442), and spotted hyenas (n = 7,410).

*Statistical analyses*

To assess whether baboon occurrences after a predator event were lower than expected by chance, we used a randomisation test to compare the observed presence of baboons after the predator event to the ones obtained under the assumption of non-avoidance behavior (Niedballa et al., 2019; Zhang & Zhao, 2023). We calculated the total number of baboon occurrences following all predator events, for each predator species independently. This was done for three consecutive 24h time blocks after each predator event: 0-24 after the predator event, 24-48 hours after the predator event and 48-72 after the predator event. Extending the observation period beyond 72 hours post predator event was avoided due to the difficulty in associating the prey species occurrence with the original predator capture event (Courbin et al., 2016; Swinkels et al., 2023). We further analyzed baboon occurrences before the predation event to establish a baseline of their presence and assess whether any observed decline was linked to the event. This allows us to determine the strength of the predation effect while accounting for natural fluctuations in baboon activity. The analysis of baboon occurrence before a predator event was also conducted in 24-hour intervals over a 72-hour period.

We excluded any predator events that occurred within the 72 hours following the camera's deployment and within 72 hours of the end of the camera's operational lifespan to prevent missing data in any pre- or post-encounter time blocks. In addition, we applied a 30-minute time-to-independence filter exclusively to predator capture events, retaining the central capture event within a cluster of consecutive detections as the main event for a given predator. This threshold aligns with previous studies that showed that increasing the independence interval has minimal impact on observed activity patterns of large carnivores (Meek et al., 2014; Peral et al., 2022).

Retaining the central event instead of the first event as usually done in the literature,

ensures symmetry when testing for predator-avoidance before and after the predator

event in our study design.

To obtain the distribution of baboon occurrences under the null hypothesis

(non-avoidance of predators), we randomised the dates of each predator capture

event 1,000 times (while preserving their original timing to maintain natural diel

activity patterns) and estimated, for each randomised data set, the total number of

baboon occurrences in each time block. We excluded any camera trap observation

period with a lifespan of < 30 consecutive days to avoid restricting the randomization

process. Predator events captured by cameras with a shorter lifespan can only be

randomized within a limited date window, which would produce unreliable results.

For each 24-hour time block, we evaluated whether baboon occurrences

deviated significantly from a distribution under the assumption of non-avoidance of

predators. To do so, we calculated the *P-value* for each time block as the proportion

of iterations in our randomization procedure in which the number of baboons was

superior or equal to the observed number of baboons before the predator event, and

inferior or equal to the observed number of baboons after the predator event. All data

processing and analyses were performed using R software (version 4.4.0; R Core

Development Team, 2022).

*Validation of the randomization approach*

To validate the randomization approach, we tested our method on Burchell's

zebras in APN. Burchell's zebras are known to display reactive temporal avoidance

behavior in response to lions for up to 24 hours after an encounter (Thaker et al.,

2011; Courbin et al., 2016; Say-Sallaz et al., 2023), therefore we expected to find a

significant response in the 0–24-hour time block, after lion capture events.

APN had the highest number of lion capture events among the South African

sites ($n_{APN}$ = 197, $n_{MAD}$ = 39, $n_{TSW}$ = 87). Although SER had higher numbers of both

lions ($n_{SER}$ = 4,266) and Burchell's zebra ($n_{SER}$ = 70,577), analysing these data

revealed too computationally intensive using our randomisation procedure.

Additionally, the number of Burchell's zebras ($n_{APN}$ = 2,267) was of the same

magnitude as the number of baboons in SER, which represents our largest dataset

in terms of baboon capture events ($n_{SER}$ = 1,556), making it a suitable comparison

point.

To assess the effect of sample size on the analyses, we performed 50

random subsets of Burchell's zebra capture events in APN to match the number of

baboon capture events ($n_{APN}$ = 500; see Table 1). This created a down sampled data

set, for which we conducted the randomization analysis for each iteration and

calculated the frequency of significant versus non-significant p-values.

**Results**

*Method validation: predator-avoidance in Burchell's zebras*

Burchell's zebras showed significant reactive temporal avoidance 0–24 hours

after the occurrence of lion in APNR (*P-value* = 0.018, Figure 2), providing support

for the soundness of our randomisation approach. However, after taking a subset of

Burchell's zebra capture events to match the number of chacma baboon capture

events in APNR, we could no longer detect this response 0-24 hours following a lion

occurrence (mean *P-value* = 0.160, supplementary figure 1). Despite this absence of

significance, a general trend of avoidance remained observable in this species

(Figure 2).

*Predator-avoidance in baboons*

Obtained results did not show any significant temporal avoidance in baboons

in response to the occurrence of our selected cursorial predator, spotted hyena,

across any time block (0–24, 24–48, or 48–72 hours) in any of the three study sites

in South Africa (all *P-values* > 0.05, Figure 3, supplementary table 1). Similarly, there

was no evidence of baboons displaying reactive temporal avoidance following the

occurrence of any of the two ambush predators tested here, namely lions and

leopards (all *P-values* > 0.05, Figure 3, supplementary table 1). However, in APNR,

baboon presence increased significantly 48–24 hours before the occurrence of lion

(*P-value* = 0.001) and 24–0 hours before the occurrence of spotted hyena (*P-value* =

0.003, Figure 3).

In SER, where the number of baboon capture events exceeded those at all

South African sites (SER 3.11 times higher than APNR, 4.01 times higher than MAD

and 2.11 times higher than TSW, Table 1), and was close to the number of

Burchell's zebra capture events in APNR, we still did not detect significant reactive

temporal avoidance of baboons in response to any predators tested (all *P-values* >

0.05, Figure 3, supplementary Table 1). Consistent with observations in APNR,

baboon occurrences in SER were significantly higher than expected under the

assumption of non-avoidance of the predator 24–0 hours before the occurrence of

lion (*P-value* = 0.043, Figure 3). Additionally, baboon occurrences in SER increased

significantly 24-0 hours before the occurrence of leopard (*P-value* = 0.019) but

showed no significant increase before spotted hyena occurrences (Figure 3).

## Discussion

Our results suggest that chacma and olive baboons do not exhibit any

significant reactive temporal avoidance toward any predator (ambush or cursorial) in

any of the four study sites (Figure 3), thus invalidating both baboons' presence

decreases following the occurrence of a predator H1 and the intensity of this

response varies based on the predator's hunting strategy H2. In contrast, Burchell's

zebra showed reactive temporal avoidance toward lions within the 0−24-hour time

block following the occurrence of lion (Figure 2), as expected given previous

observations (Thaker et al., 2011; Courbin et al., 2016; Say-Sallaz et al., 2023). After

reducing zebra capture events (to match the highest number of baboon capture

events), this significant decrease was no longer observable, suggesting that sample

size may impact our ability to detect avoidance response. Although we cannot rule

out the problem of statistical power for baboons, the non-significant results in SER

(in which the number of baboon occurrences are close to that of zebras in our

validation test) suggest the absence of reactive avoidance behavior in olive baboons,

and probably chacma baboons.

The absence of reactive temporal avoidance in baboons may be explained by

their reliance on alternative predator avoidance strategies. Lima and Dill (1990)

proposed that the probability of prey survival increases with the prey's ability to

successfully detect and effectively avoid or interrupt the threat as early as possible.

By doing so, prey might be able to prevent the predation attempt and reduce their

energetic costs of being displaced from a favorable area. Baboons are highly social primates that exhibit sentinel behavior, where one individual monitors for threats while the rest of the troop forages (Horrocks et al., 1986; Bezerra et al., 2008) and use contact barks to signal the presence of a predator (Bailey, 1993; Cowlishaw, 1994b; Jooste et al., 2013; Isbell et al., 2018). This altruistic threat detection mechanism may reduce energy expenditure among kin and be favoured over the forced displacement of the entire troop as described in meerkats (*Suricata suricatta*) (Santema & Clutton-Brock, 2013; Rauber et al., 2018).

Additionally, moving away from the area at risk does not necessarily imply horizontal displacement for arboreal species. Baboon troops are known for using refuge areas, such as trees or cliffs, employing a vertical escape strategy (Cowlishaw 1997a; Fischer et al. 2001; Isbell et al. 2018). This strategy protects them from terrestrial predators such as spotted hyenas and lions, which are less adapted to climbing (Buse 1980; Cowlishaw 1994). Refuge-based avoidance behaviors have been documented in other species worldwide, including the Nubian ibex (*Capra nubiana*), mule deer (*Odocoileus hemionus*) and klipspringer (*Oreotragus oreotragus*) (Hayward et al. 2006; Hochman & Kotler 2007; Druce et al. 2009; Iribarren & Kotler 2012). Similarly to these species that focus their foraging efforts on areas closer to cliffs, rocky outcrops, or higher ground, baboons might favor foraging in areas with sufficient tall trees as an escape route, a mechanism difficult to capture with the camera-trap setup used in this study.

When interaction with a predator is inevitable, baboons have been described to respond with aggressive retaliation (Buse, 1980; Hayward et al., 2006; Bailey, 1993; Cowlishaw, 1994; Jooste et al., 2013; Isbell et al., 2018). A review by Cowlishaw (1994) analysed predation success rates on baboons across 14 sites in

multiple countries. This study showed that leopards were twice as successful as lions and spotted hyenas when preying on baboons. However, when baboons retaliated, they successfully deterred predation in 93% of cases, and in four of 11 observed predation attempts, baboons killed the attacking leopard. This behavior is predominantly exhibited by male baboons in both chacma (Cowlishaw, 1994; Isbell et al., 2018) and olive (MacCormick et al., 2012) baboons, owing to their larger teeth and body size (Virgadamo et al., 1972; Galbany et al., 2015; Johnson, 2003). Such a response could decrease predation attempts rate by lowering the predator's chance of successful capture and their interest in the species (Lima & Dill, 1990; Cowlishaw, 1994; Alberts, 2018; Isbell et al., 2018). In response to these effective anticipatory anti-predator strategies, predators may simply not actively hunt baboons and focus their efforts on more cost-effective prey such as ungulates. Therefore, the predators would then be viewed as less of a threat by baboons which would consequently be less inclined to leave the area after the arrival of a predator.

Despite these results, we cannot completely rule out the possibility that we lacked sufficient statistical power to detect reactive temporal avoidance in our South African sites. To capture the true prey and predator densities, more cameras may be necessary, as is shown by Si et al. (2014) and Evans et al. (2019) with coyotes, whereby increasing the number of cameras by one at each capture site, the probability of detecting the species was four times higher. In addition, it has been shown that the ability to detect spatio-temporal avoidance with camera trap data depends on both interaction strength and sample size (Niedballa et al. 2019). This suggests that detection remains constrained by species-specific activity patterns and the number of observations, reinforcing the need for specific survey design. Optimizing camera density and placement seems essential to mitigate biased

detection, particularly for species with low detection probabilities or complex habitat use.

## Conclusion

This study contributes to broadening our understanding of predator-prey dynamics in two non-human primates' species. The potential absence of reactive temporal avoidance in chacma and olive baboons raises important questions about the variability of their anti-predator strategies, not only in savannah biomes but also across other environments in response to different predators (Zinner et al., 2011; Stone et al., 2013). Interspecific interactions are complex, and species have a wide array of behavioral responses at their disposal to avoid interactions, such as partitioning space or resources instead of time (Morales-Castilla et al., 2015). This study highlights the potential influence of sample size on the detectability of behavioral responses when using camera traps, as illustrated by the loss of significance of reactive temporal avoidance in Burchell's zebra after reduction of the sample size. Limitations in camera trap detection range (i.e. camera traps density and vertical detection capabilities) further emphasize the challenges of accurately quantifying prey-predator interactions among arboreal and low-density species. Future research should prioritize increasing sample sizes and optimizing camera trap configurations to enhance detection probabilities.

**Authors' contributions**

Conceptualization: VR and FP - Methodology: NVR, VR, FP, LT - Data acquisition: HF, JV - Formal analysis: NVR, VR, FP, LT - Investigation: NVR, VR, FP, LT - Writing: Original draft: NVR, VR, FP, LT - Writing: NVR, VR, FP, LT - Review & Editing : NVR, VR, FP, LT, HF, JV - Supervision :  VR, FP - Funding acquisition : VR, JV

**Acknowledgments**

The authors thank the CNRS for funding this project. They also thank the Wildlife Ecology Lab, Nelson Mandela University's George campus, CNRS, IRL REHABS, and the Snapshot Safari project for the provision and processing of the data. Funding for SnaphotSafari was provided by the National Research Foundation and the Foundational Biodiversity Information Programme (Grant number FBIP170720256205).

**Data availability statement**

Pictures from the Snapshot South Africa program can be requested at SnapShot Safari upon reasonable request. Pictures from the Snapshot Serengeti program are available from the Labelled Image Library of Alexandria – Biology and Conservation: <http://lila.science/datasets/snapshot-serengeti>. All classification data and metadata are publicly available at Dryad: http://dx.doi.org/10.5061/dryad.5pt92.

**Code availability**

The code to analyze and reproduce this study has been deposited in Zenodo and is available online at https://zenodo.org/records/15018760

## References

**References**

Alberts, S. C. (2019). Social influences on survival and reproduction: Insights from a

long-term study of wild baboons. *Journal of Animal Ecology*, *88*(1), 47–66.

https://doi.org/10.1111/1365-2656.12887

Allan, A. T. L., Bailey, A. L., & Hill, R. A. (2021). Consistency in the flight and visual

orientation distances of habituated chacma baboons after an observed leopard

predation. Do flight initiation distance methods always measure perceived predation

risk? *Ecology and Evolution*, *11*(21), 15404–15416.

https://doi.org/10.1002/ece3.8237

Bailey, T. N. (1993). 9. Social Organization of Leopards. In *The African Leopard* (pp.

240–297). Columbia University Press. https://doi.org/10.7312/bail90198-010

Basille, M., Fortin, D., Dussault, C., Bastille-Rousseau, G., Ouellet, J.-P., & Courtois,

R. (2015). Plastic response of fearful prey to the spatiotemporal dynamics of

predator distribution. *Ecology*, *96*(10), 2622–2631. https://doi.org/10.1890/14-1706.1

Beaumont, R. A. R.; P. B., & Bednarik, R. G. (2015). Keywords: Cupule-Age

estimate-Palaeoenvironment-Tswalu Kalahari Reserve-South Africa concerning a

cupule sequence on the edge of the kalahari desert in South Africa (Vol. 32, Issue 2).

Bedetti, A., Greyling, C., Paul, B., Blondeau, J., Clark, A., Malin, H., Horne, J.,

Makukule, R., Wilmot, J., Eggeling, T., Kern, J., & Henley, M. (2020). System for

Elephant Ear-pattern Knowledge (SEEK) to identify individual African elephants.

*Pachyderm*, *61*, 63–77. https://doi.org/10.69649/pachyderm.v61i.65

Belsky, A. J. (1985). Long-Term Vegetation Monitoring in the Serengeti National

Park, Tanzania. *The Journal of Applied Ecology*, *22*(2), 449.

https://doi.org/10.2307/2403177

Bezerra, B. M., & Souto, A. (2008). Structure and Usage of the Vocal Repertoire of

(*Callithrix jacchus)*. *International Journal of Primatology*, *29*(3), 671–701.

https://doi.org/10.1007/s10764-008-9250-0

Broekhuis, F., Cozzi, G., Valeix, M., McNutt, J. W., & Macdonald, D. W. (2013). Risk

avoidance in sympatric large carnivores: reactive or predictive? *Journal of Animal*

*Ecology*, *82*(5), 1098–1105. https://doi.org/10.1111/1365-2656.12077

Busse, C. D. (1980). *Leopard and Lion predation upon Chacma baboons living in the*

*Moremi Wildlife Reserve*.

Campos, F. A., & Fedigan, L. M. (2014). Spatial ecology of perceived predation risk

and vigilance behavior in white-faced capuchins. *Behavioral Ecology*, *25*(3), 477–

486. https://doi.org/10.1093/beheco/aru005

Charles-Dominique, T., Staver, A. C., Midgley, G. F., & Bond, W. J. (2015). Functional

differentiation of biomes in an African savanna/forest mosaic. *South African Journal*

*of Botany*, *101*, 82–90. https://doi.org/10.1016/j.sajb.2015.05.005

Coleman, B. T. (2013). Spatial and temporal determinants of samango monkey

(*Cercopithecus mitis erythrarchus)* resource acquisition and predation avoidance

behaviour [ Doctoral theses]. Durham University.

Courbin, N., Loveridge, A. J., Macdonald, D. W., Fritz, H., Valeix, M., Makuwe, E. T.,

& Chamaillé-Jammes, S. (2016). Reactive responses of zebras to lion encounters

shape their predator–prey space game at large scale. *Oikos*, *125*(6), 829–838.

https://doi.org/10.1111/oik.02555

Cowlishaw, G. (1994). Vulnerability To Predation in Baboon Populations. *Behaviour*,

*131*(3–4), 293–304. https://doi.org/10.1163/156853994X00488

Creel, S. (2018). The control of risk hypothesis: reactive vs. proactive antipredator

responses and stress-mediated vs. food-mediated costs of response. *Ecology*

*Letters*, *21*(7), 947–956. https://doi.org/10.1111/ele.12975

Creel, S., & Christianson, D. (2008). Relationships between direct predation and risk

effects.    *Trends    in    Ecology    &    Evolution*,    *23*(4),    194–201.

https://doi.org/10.1016/j.tree.2007.12.004

Creel, S., Schuette, P., & Christianson, D. (2014). Effects of predation risk on group

size, vigilance, and foraging behavior in an African ungulate community. *Behavioral*

*Ecology*, *25*(4), 773–784. https://doi.org/10.1093/beheco/aru050

Creel, S., & Winnie, J. A. (2005). Responses of elk herd size to fine-scale spatial and

temporal variation in the risk of predation by wolves. *Animal Behaviour*, *69*(5), 1181–

1189. https://doi.org/10.1016/j.anbehav.2004.07.022

Davis, A. L. V., Scholtz, C. H., Kryger, U., Deschodt, C. M., & Strümpher, W. P.

(2010). Dung Beetle Assemblage Structure in Tswalu Kalahari Reserve: Responses

to a Mosaic of Landscape Types, Vegetation Communities, and Dung Types.

*Environmental Entomology*, *39*(3), 811–820. https://doi.org/10.1603/EN09256

Dröge, E., Creel, S., Becker, M. S., & M'soka, J. (2017). Spatial and temporal

avoidance of risk within a large carnivore guild. *Ecology and Evolution*, *7*(1), 189–

199. https://doi.org/10.1002/ece3.2616

Druce, D. J., brown, J. S., Kerley, G. I. H., Kotler, B. P., Mackey, R. L., & Slotow, R.

(2009). Spatial and temporal scaling in habitat utilization by klipspringers

(Oreotragus *oreotragus*) determined using giving-up densities. *Austral Ecology*,

*34*(5), 577–587. https://doi.org/10.1111/j.1442-9993.2009.01963.x

Druelle, F., Aerts, P., Ngawolo, J. C. B., & Narat, V. (2020). Impressive Arboreal Gap-

Crossing Behaviors in Wild Bonobos, (*Pan paniscus*). *International Journal of*

*Primatology*, *41*(1), 129–140. https://doi.org/10.1007/s10764-020-00140-z

Eberle, M., & Kappeler, P. M. (2008). Mutualism, reciprocity, or kin selection?

Cooperative rescue of a conspecific from a boa in a nocturnal solitary forager the

gray mouse lemur. *American Journal of Primatology*, *70*(4), 410–414.

https://doi.org/10.1002/ajp.20496

Edwards, S., Gange, A. C., & Wiesel, I. (2016). An oasis in the desert: The potential

of water sources as camera trap sites in arid environments for surveying a carnivore

guild. *Journal of Arid Environments*, *124*, 304–309.

https://doi.org/10.1016/j.jaridenv.2015.09.009

Evans, B. E., Mosby, C. E., & Mortelliti, A. (2019). Assessing arrays of multiple trail

cameras to detect North American mammals. *PLOS ONE*, *14*(6), e0217543.

https://doi.org/10.1371/journal.pone.0217543

Fichtel, C., Perry, S., & Gros-Louis, J. (2005). Alarm calls of white-faced capuchin

monkeys: an acoustic analysis. *Animal Behaviour*, *70*(1), 165–176.

https://doi.org/10.1016/j.anbehav.2004.09.020

Fischer, J., Metz, M., Cheney, D. L., & Seyfarth, R. M. (2001). Baboon responses to

graded bark variants. *Animal Behaviour*, *61*(5), 925–931.

https://doi.org/10.1006/anbe.2000.1687

Galbany, J., Tung, J., Altmann, J., & Alberts, S. C. (2015). Canine Length in Wild

Male Baboons: Maturation, Aging and Social Dominance Rank. *PLOS ONE*, *10*(5),

e0126415. https://doi.org/10.1371/journal.pone.0126415

Hamilton, W. J., & Busse, C. (1982). Social dominance and predatory behavior of

chacma baboons. *Journal of Human Evolution*, *11*(7), 567–573.

https://doi.org/10.1016/S0047-2484(82)80003-1

Harvell, C. D. (1990). The Ecology and Evolution of Inducible Defenses. *The*

*Quarterly Review of Biology*, *65*(3), 323–340. https://doi.org/10.1086/416841

Hayward, M. W., Henschel, P., O'Brien, J., Hofmeyr, M., Balme, G., & Kerley, G. I. H.

(2006). Prey preferences of the leopard (*Panthera pardus*). *Journal of Zoology*,

*270*(2), 298–313. https://doi.org/10.1111/j.1469-7998.2006.00139.x

Hill, R. A., & Weingrill, T. (2007). Predation Risk and Habitat Use in Chacma

Baboons (*Papio hamadryas ursinus*). In *Primate Anti-Predator Strategies* (pp. 339–

354). Springer US. https://doi.org/10.1007/978-0-387-34810-0_16

Hochman, V., & Kotler, B. P. (2007). Patch use, apprehension, and vigilance

behavior of Nubian Ibex under perceived risk of predation. *Behavioral Ecology*,

*18*(2), 368–374. https://doi.org/10.1093/beheco/arl087

Horrocks, J. A., & Hunte, W. (1986). Sentinel behaviour in vervet monkeys: Who

sees whom first? *Animal Behaviour*, *34*(5), 1566–1568.

https://doi.org/10.1016/S0003-3472(86)80226-3

Humphries, D. A., & Driver, P. M. (1967). Erratic Display as a Device against

Predators. *Science*, *156*(3783), 1767–1768.

https://doi.org/10.1126/science.156.3783.1767

Iribarren, C., & Kotler, B. P. (2012). Foraging patterns of habitat use reveal

landscape of fear of Nubian ibex (*Capra nubiana). Wildlife Biology*, *18*(2), 194–201.

https://doi.org/10.2981/11-041

Isbell, L. A., Bidner, L. R., Van Cleave, E. K., Matsumoto-Oda, A., & Crofoot, M. C.

(2018). GPS-identified vulnerabilities of savannah-woodland primates to leopard

predation and their implications for early hominins. *Journal of Human Evolution*, *118*,

1–13. https://doi.org/10.1016/j.jhevol.2018.02.003

Jared, C., Antoniazzi, M. M., Jordão, A. E. C., Silva, J. R. M. C., Greven, H., &

Rodrigues, M. T. (2009). Parotoid macro glands in toad (*Rhinella jimi*): Their structure

and functioning in passive defence. *Toxicon*, *54*(3), 197–207.

https://doi.org/10.1016/j.toxicon.2009.03.029

Jooste, E., Pitman, R. T., van Hoven, W., & Swanepoel, L. H. (2013). Unusually High

Predation on Chacma Baboons (*Papio ursinus*) by Female Leopards (Panthera

pardus) in the Waterberg Mountains, South Africa. *Folia Primatologica*, *83*(3–6),

353–360. https://doi.org/10.1159/000339644

Karpanty, S. M., & Wright, P. C. (2007). Predation on Lemurs in the Rainforest of

Madagascar by Multiple Predator Species: Observations and Experiments. In

*Primate Anti-Predator Strategies* (pp. 77–99). Springer US.

https://doi.org/10.1007/978-0-387-34810-0_4

Kerbs, John. R., & Davies, Nicholas. B. (2009). *Behavioural ecology: an evolutionary*

*approach*. John Wiley & Sons.

Kie, J. G. (1999). Optimal Foraging and Risk of Predation: Effects on Behavior and

Social Structure in Ungulates. *Journal of Mammalogy*, *80*(4), 1114–1129.

https://doi.org/10.2307/1383163

Latombe, G., Fortin, D., & Parrott, L. (2014). Spatio-temporal dynamics in the

response of woodland caribou and moose to the passage of grey wolf. *Journal of*

*Animal Ecology*, *83*(1), 185–198. https://doi.org/10.1111/1365-2656.12108

Laundré, J. W., Hernández, L., & Altendorf, K. B. (2001). Wolves, elk, and bison:

reestablishing the "landscape of fear" in Yellowstone National Park, U.S.A. *Canadian*

*Journal of Zoology*, *79*(8), 1401–1409. https://doi.org/10.1139/z01-094

Lima, S. L., & Dill, L. M. (1990). Behavioral decisions made under the risk of

predation: a review and prospectus. *Canadian Journal of Zoology*, *68*(4), 619–640.

https://doi.org/10.1139/z90-092

Lingle, S., & Pellis, S. (2002). Fight or flight? Antipredator behavior and the

escalation of coyote encounters with deer. *Oecologia*, *131*(1), 154–164.

https://doi.org/10.1007/s00442-001-0858-4

MacCormick, H. A., MacNulty, D. R., Bosacker, A. L., Lehman, C., Bailey, A., Anthony

Collins, D., & Packer, C. (2012). Male and female aggression: lessons from sex,

rank, age, and injury in olive baboons. *Behavioral Ecology*, *23*(3), 684–691.

https://doi.org/10.1093/beheco/ars021

Morales-Castilla, I., Matias, M. G., Gravel, D., & Araújo, M. B. (2015). Inferring biotic

interactions from proxies. In Trends in Ecology and Evolution (Vol. 30, Issue 6, pp.

347–356). Elsevier Ltd. https://doi.org/10.1016/j.tree.2015.03.014

Meek, P. D., Ballard, G., Claridge, A., Kays, R., Moseby, K., O'Brien, T., O'Connell,

A., Sanderson, J., Swann, D. E., Tobler, M., & Townsend, S. (2014). Recommended

guiding principles for reporting on camera trapping research. In *Biodiversity and*

*Conservation* (Vol. 23, Issue 9, pp. 2321–2343). Kluwer Academic Publishers.

https://doi.org/10.1007/s10531-014-0712-8

Meno, W., Coss, R. G., & Perry, S. (2013). Development of Snake-Directed Antipredator Behavior by Wild White-faced Capuchin Monkeys: I. Snake-Species Discrimination. *American Journal of Primatology*, *75*(3), 281–291. https://doi.org/10.1002/ajp.22106

Mirza, R. S., Mathis, A., & Chivers, D. P. (2006). Does Temporal Variation in Predation Risk Influence the Intensity of Antipredator Responses? A Test of the Risk Allocation Hypothesis. *Ethology*, *112*(1), 44–51. https://doi.org/10.1111/j.1439-0310.2006.01111.x

Newey, S., Davidson, P., Nazir, S., Fairhurst, G., Verdicchio, F., Irvine, R. J., & van der Wal, R. (2015). Limitations of recreational camera traps for wildlife management and conservation research: A practitioner's perspective. *Ambio*, *44*, 624–635. https://doi.org/10.1007/s13280-015-0713-1

Niedballa, J., Wilting, A., Sollmann, R., Hofer, H., & Courtiol, A. (2019). Assessing analytical methods for detecting spatiotemporal interactions between species from camera trapping data. Remote Sensing in Ecology and Conservation, 5(3), 272–285. https://doi.org/10.1002/rse2.107

Northfield, T. D., Barton, B. T., & Schmitz, O. J. (2017). A spatial theory for emergent multiple predator–prey interactions in food webs. *Ecology and Evolution*, *7*(17), 6935–6948. https://doi.org/10.1002/ece3.3250

Palmer, M. S., & Packer, C. (2021). Reactive anti-predator behavioral strategy shaped by predator characteristics. *PLOS ONE*, *16*(8), e0256147. https://doi.org/10.1371/journal.pone.0256147

Pardo, L. E., Bombaci, S., Huebner, S. E., Somers, M. J., Fritz, H., Downs, C., Guthmann, A., Hetem, R. S., Keith, M., le Roux, A., Mgqatsa, N., Packer, C., Palmer,

M. S., Parker, D. M., Peel, M., Slotow, R., Maartin Strauss, W., Swanepoel, L., Tambling, C., … Venter, J. A. (2021). Snapshot Safari: A large-scale collaborative to monitor Africa's remarkable biodiversity. *South African Journal of Science*, *117*(1–2). https://doi.org/10.17159/SAJS.2021/8134

Peral, C., Landman, M., & Kerley, G. I. H. (2022). The inappropriate use of time-to-independence biases estimates of activity patterns of free-ranging mammals derived from camera traps. *Ecology and Evolution*, *12*(10). https://doi.org/10.1002/ece3.9408

Perry, S., Manson, J. H., Dower, G., & Wikberg, E. (2003). White-Faced Capuchins Cooperate to Rescue a Groupmate from a Boa constrictor. *Folia Primatologica*, *74*(2), 109–111. https://doi.org/10.1159/000070008

Prates, I., Antoniazzi, M. M., Sciani, J. M., Pimenta, D. C., Toledo, L. F., Haddad, C. F. B., & Jared, C. (2012). Skin glands, poison, and mimicry in dendrobatid and leptodactylid amphibians. *Journal of Morphology*, *273*(3), 279–290. https://doi.org/10.1002/jmor.11021

Rauber, R., & Manser, M. B. (2018). Experience of the signaller explains the use of social versus personal information in the context of sentinel behaviour in meerkats. Scientific Reports, 8(1), 11506. https://doi.org/10.1038/s41598-018-29678-y

Rutherford, M. C., Mucina, L., Lötter, M. C., Bredenkamp, G. J., Smit, J. H. L., Scott-Shaw, C. R., Hoare, D. B., Goodman, P. S., Bezuidenhout, H., Scott, L., Ellis, F., Powrie, L., Siebert, F., Mostert, T. H., Henning, B. J., Ventner, C. E., Camp, K. G. T., Siebert, S., Matthews, W., & Hurter, J. (2006). Savanna Biome. In *The Vegetation of South Africa, Lesotho and Swaziland* (pp. 440–529).

Sankaran, M., Hanan, N. P., Scholes, R. J., Ratnam, J., Augustine, D. J., Cade, B. S., Gignoux, J., Higgins, S. I., Le Roux, X., Ludwig, F., Ardo, J., Banyikwa, F., Bronn,

A., Bucini, G., Caylor, K. K., Coughenour, M. B., Diouf, A., Ekaya, W., Feral, C. J., …

Zambatis, N. (2005). Determinants of woody cover in African savannas. *Nature*,

*438*(7069), 846–849. https://doi.org/10.1038/nature04070

Santema, P., & Clutton-Brock, T. (2013). Meerkat helpers increase sentinel behaviour

and bipedal vigilance in the presence of pups. *Animal Behaviour*, *85*(3), 655–661.

https://doi.org/10.1016/j.anbehav.2012.12.029

Say-Sallaz, E., Chamaillé-Jammes, S., Périquet, S., Loveridge, A. J., Macdonald, D.

W., Antonio, A., Fritz, H., & Valeix, M. (2023). Large carnivore dangerousness affects

the reactive spatial response of prey. *Animal Behaviour*, *202*, 149–162.

https://doi.org/10.1016/j.anbehav.2023.05.014

Scholes, R. J., & Archer, S. R. (1997). Tree-Grass Interactions in Savannas. *Annual*

*Review of Ecology and Systematics*, *28*(1), 517–544.

https://doi.org/10.1146/annurev.ecolsys.28.1.517

Schreier, A. L., & Swedell, L. (2012). Ecology and sociality in a multilevel society:

Ecological determinants of spatial cohesion in hamadryas baboons. *American*

*Journal of Physical Anthropology*, *148*(4), 580–588.

https://doi.org/10.1002/ajpa.22076

Seyfarth, R. M., Cheney, D. L., & Marler, P. (1980). Vervet monkey alarm calls:

Semantic communication in a free-ranging primate. *Animal Behaviour*, *28*(4), 1070–

1094. https://doi.org/10.1016/S0003-3472(80)80097-2

Si, X., Kays, R., & Ding, P. (2014). How long is enough to detect terrestrial animals?

Estimating the minimum trapping effort on camera traps. *PeerJ*, 2, e374.

https://doi.org/10.7717/peerj.374

Stankowich, T. (2012). Armed and dangerous: predicting the presence and function

of defensive weaponry in mammals. *Adaptive Behavior*, *20*(1), 32–43.

https://doi.org/10.1177/1059712311426798

Stankowich, T., & Campbell, L. A. (2016). Living in the danger zone: Exposure to

predators and the evolution of spines and body armor in mammals. *Evolution*, *70*(7),

1501–1511. https://doi.org/10.1111/evo.12961

Stone, O. M. L., Laffan, S. W., Curnoe, D., & Herries, A. I. R. (2013). The Spatial

Distribution of Chacma Baboon (*Papio ursinus*) Habitat Based on an Environmental

Envelope Model. *International Journal of Primatology*, *34*(2), 407–422.

https://doi.org/10.1007/s10764-013-9669-9

Suire, A., Kunita, I., Harel, R., Crofoot, M., Mutinda, M., Kamau, M., Hassel, J. M.,

Murray, S., Kawamura, S., & Matsumoto-Oda, A. (2023). Estimating individual

exposure to predation risk in group-living baboons, *Papio anubis*. *PLOS ONE*,

*18*(11), e0287357. https://doi.org/10.1371/journal.pone.0287357

Swanson, A. B. (2014). *Living with lions: Spatiotemporal aspects of coexistence in*

*savanna carnivores*.

Swanson, A., Kosmala, M., Lintott, C., Simpson, R., Smith, A., & Packer, C. (2015).

Snapshot Serengeti, high-frequency annotated camera trap images of 40

mammalian species in an African savanna. *Scientific Data*, *2*(1), 150026.

https://doi.org/10.1038/sdata.2015.26

Swinkels, C., van der Wal, J. E. M., Stinn, C., Monteza-Moreno, C. M., & Jansen, P.

A. (2023). Prey tracking and predator avoidance in a Neotropical moist forest: a

camera-trapping approach. *Journal of Mammalogy*, *104*(1), 137–145.

https://doi.org/10.1093/jmammal/gyac091

Szott, I. D., Pretorius, Y., Ganswindt, A., & Koyama, N. F. (2020). Physiological stress
response of African elephants to wildlife tourism in Madikwe Game Reserve, South
Africa. *Wildlife Research*, *47*(1), 34. https://doi.org/10.1071/WR19045

Thaker, M., Vanak, A. T., Owen, C. R., Ogden, M. B., Niemann, S. M., & Slotow, R.
(2011). Minimizing predation risk in a landscape of multiple predators: Effects on the
spatial distribution of African ungulates. *Ecology*, *92*(2), 398–407.
https://doi.org/10.1890/10-0126.1

Trapanese, C., Meunier, H., & Masi, S. (2019). What, where and when: spatial
foraging decisions in primates. *Biological Reviews*, *94*(2), 483–502.
https://doi.org/10.1111/brv.12462

Trinkel, M. (2010). Prey selection and prey preferences of spotted hyenas *Crocuta*
*crocuta* in the Etosha National Park, Namibia. *Ecological Research*, *25*(2), 413–417.
https://doi.org/10.1007/s11284-009-0669-3

Virgadamo, P., Hodosh, M., Povar, M., & Shklar, G. (1972). The Dentition of *Papio*
*anubis*. *Journal of Dental Research*, *51*(5), 1338–1345.
https://doi.org/10.1177/00220345720510051501

Von Hippel, F. A. (1998). Use of sleeping trees by black and white Colobus monkeys
(*Colobus guereza*) in the Kakamega Forest, Kenya. *American Journal of*
*Primatology*, *45*(3), 281–290. https://doi.org/10.1002/(SICI)1098-
2345(1998)45:3<281:AID-AJP4>3.0.CO;2-S

Wentworth, J. C., Tambling, C. J., & Kerley, G. I. H. (2011). Evidence for prey
selection by spotted hyaena in the Eastern Cape, South Africa. *Acta Theriologica*,
*56*(4), 389–392. https://doi.org/10.1007/s13364-011-0033-1

Zhang, Y., & Zhao, Q. (2023). What is a Randomization Test? *Journal of the*

*American Statistical Association*, *118*(544), 2928–2942.

https://doi.org/10.1080/01621459.2023.2199814

Zinner, D., Buba, U., Nash, S., & Roos, C. (2011). Pan-African Voyagers: The

Phylogeography of Baboons. In *Primates of Gashaka* (pp. 319–358). Springer New

York. https://doi.org/10.1007/978-1-4419-7403-7_7

 **Tables**

712 Table 1. Number of capture events for the species of interest in Associated Private

713 Nature Reserves (APNR), Madikwe Game Reserve (MAD), Tswalu Kalahari Private

714 Wildlife Reserve (TSW) (South Africa, 2017-2019), and Serengeti National Park

715 (SER, Tanzania, 2011-2013). A 30-minute independence interval was applied on the

716 predator detections (see Material and methods section for more details).

| Species | Scientific name | No. of capture events in MAD | No. of capture events in TSW | No. of capture events in APNR | No. of capture events in SER |
|---|---|---|---|---|---|
| Lion | *Panthera leo* | 39 | 87 | 197 | 4,266 |
| Leopard | *Panthera pardus* | 43 | 8 | 168 | 228 |
| Spotted hyena | *Crocuta crocuta* | 187 | 11 | 1,676 | 5,303 |
| Chacma baboon | *Papio ursinus* | 388 | 806 | 500 | 0 |
| Olive baboon | *Papio anubis* | 0 | 0 | 0 | 1,556 |
| Burchell's zebra | *Equus quagga* | 5,099 | 828 | 2,267 | 70,577 |

**Supplementary table 1.**

722 P-values calculated by comparing the observed occurrences of chacma baboons

723 (*Papio ursinus*) and olive baboons (*Papio anubis*) after a predator event (lion,

724 leopard, spotted hyena) to the distribution of occurrences produced by randomizing

725 the predator event under the assumption of non-avoidance, across six-time blocks

726 (see material and methods section for more details).

| Site | Prey species | Predator species | (-72 to -48 hr) | (-48 to -24 hr) | (-24 to 0 hr) | (0 to 24 hr) | (24 to 48 hr) | (48 to 72 hr) |
|---|---|---|---|---|---|---|---|---|
| MAD | Chacma baboon | Lion | 0,158 | 1 | 1 | 0,894 | 0,46 | 0,459 |
| | | Leopard | 1 | 0,105 | 1 | 0,807 | 0,813 | 0,509 |
| | | Spotted hyena | 0,899 | 1 | 0,91 | 0,753 | 0,235 | 0,085 |
| TSW | Chacma baboon | Lion | 0,451 | 0,32 | 0,042 (***) | 0,924 | 0.686 | 0,348 |
| | | Leopard | 1 | 1 | 0,118 | 0,804 | 0,661 | 0,791 |
| | | Spotted hyena | 1 | 1 | 1 | 0,94 | 0,765 | 0,724 |
| APNR | Chacma baboon | Lion | 0,645 | 0,001 (***) | 1 | 0,503 | 0,241 | 0,415 |
| | | Leopard | 0,555 | 0,714 | 0,536 | 0,563 | 0,123 | 0,606 |
| | | Spotted hyena | 0,921 | 0,064 | 0,003 (***) | 0,202 | 0,47 | 0,152 |
| SER | Olive Baboon | Lion | 0,935 | 0,219 | 0,043 (***) | 0,589 | 0,507 | 0,87 |
| | | Leopard | 0,714 | 0,613 | 0,019 (***) | 0,386 | 0,071 | 0,636 |
| | | Spotted hyena | 0.637 | 0.378 | 0.257 | 0.997 | 0.266 | 0.847 |

 **Figure legends**

**Figure 1. Map of the four study sites in South Africa and Tanzania.** A) Madikwe

Private Game Reserve (MAD; n = 41 camera traps); B) Tswalu Kalahari Game

Reserve (TSW; n = 28 camera traps) and C) Associated Private Nature Reserves

(APNR; n = 56 camera traps) in South Africa. D) Serengeti National Park (SER; n =

225 camera traps) in Tanzania. The green boundaries represent the borders of the

study areas, the red circles represent the camera trap locations, with their assigned

code printed above. *P. ursinus* occurs in all three South African sites, and *P. anubis*

occurs only in Serengeti National Park, Tanzania.

**Figure 1. Map of the four study sites in South Africa and Tanzania.** A) Madikwe

Private Game Reserve (MAD; n = 41 camera traps); B) The Serengeti National Park

(SER; n = 225 camera traps) in Tanzania; C) Tswalu Kalahari Game Reserve (TSW;

n = 28 camera traps) and D) Associated Private Nature Reserves (APNR; n = 56

camera traps) in South Africa. The green boundaries represent the borders of the

study areas, the red circles represent the camera trap locations, with their assigned

code printed above. Pictograms are included to show the presence of *P. ursinus*

(indicated for all three South African sites) and *P. anubis* (indicated only for the

Serengeti National Park in Tanzania).

**Figure 2. Distribution of Burchell's zebra occurrences obtained from the**

**randomized dataset (n = 1,000) in APNR (Associated Private Nature Reserves)**

**before and after a lion event.** Each panel corresponds to one of six consecutive

24-hour time blocks relative to the predator's occurrence: [-72 to -48 hours before], [-

48 to -24 hours before], [-24 to 0 hours before], [0–24 hours after], [24–48 hours

after], and [48–72 hours after]. The black vertical lines represent the observed

number of zebra occurrences for each time period. "*" illustrates significant *P-values* (calculated as the proportion of randomized datasets in which zebra occurrences were equal to or lower than the observed value).

**Figure 3. Distribution of baboon occurrences obtained from the randomized dataset (n = 1,000) across different sites and predator category combinations, with each panel representing a distinct time period before and after the predator event.** Each panel corresponds to one of the six consecutive 24-hour time blocks relative to the predator's event: [-72 to -48 hours before], [-48 to -24 hours before], [-24 to 0 hours before], [0–24 hours after], [24–48 hours after], and [48–72 hours after]. The red vertical lines represent the observed number of baboon occurrences for each site-predator-time period combination. "*" illustrates significant *P-values* (calculated as the proportion of randomized datasets in which baboon occurrences were equal to or lower than the observed value). Colors indicate the predator species: blue for spotted hyena; green for leopard; and red for lion. Sites abbreviations: TSW: Tswalu Kalahari Private Wildlife Reserve; SER: Serengeti National Park; MAD: Madikwe Game Reserve; APNR: Associated Private Nature Reserves.

**Supplementary Figure legend.**

**Supplementary Figure 1. Distribution of *P-values* from the randomization test assessing the effect of lions on zebra occurrence in APNR (Associated Private Nature Reserves) within the 0–24 hours following a lion event.** The test was conducted on randomly down sampled zebra occurrence datasets (n = 50). The red dashed line indicates the *P-value* threshold of 0.05, marking the significance level for detecting deviations from random expectations.

780 **Figure 1**

782 **Figure 2**

 **Figure 3**

789
790 **Supplementary Figure 1.**