Neural Tracking of the Maternal Voice in the Infant Brain

Sarah Jessen[1,3], Martin Orf[2,3], Jonas Obleser[2,3]

[1] Institute of Medical Psychology, University of Lübeck, 23562 Lübeck, Germany

[2] Department of Psychology, University of Lübeck, 23562 Lübeck, Germany

[3] Center of Brain, Behavior, and Metabolism, University of Lübeck, 23562 Lübeck, Germany

Corresponding Author:

Sarah Jessen

Institute of Medical Psychology
University of Lübeck
Ratzeburger Allee 160
23562 Lübeck
Germany

sarah.jessen@uni-luebeck.de

Tel.:(+49) (0)451 3101 7449

Number of Pages: 21
Number of Tables: 0
Number of Figures: 5
Number of words (abstract): 174
Number of words (introduction): 645
Number of words (discussion): 1471

Conflict of interest
The authors declare no competing financial interests.

Acknowledgements
The authors are grateful to the families for their participation. We further would like to thank the German Research Foundation (DFG) for funding to SJ (JE781/3-1) and to JO (JO352/2-2). Generative Artificial Intelligence (DeepL by DeepL; chatGTP by openAI) was used for proofreading certain paragraphs, for the content of which the authors retain full responsibility.

Abstract

Infants preferentially process familiar social signals, but the neural mechanisms underlying continuous processing of maternal speech remain unclear. Using EEG-based neural encoding models based on temporal response functions, we investigated how 7-month-old infants track maternal vs. unfamiliar speech and whether this affects simultaneous face processing. Infants showed stronger neural tracking of their mother's voice, independent of acoustic properties, suggesting an early neural signature of voice familiarity. Face tracking responses differed depending on the voice infants heard. When listening to a stranger's voice, face-tracking accuracy at central electrodes increased with occipital face tracking, suggesting heightened attentional engagement. However, we found no evidence for differential processing of happy vs. fearful faces, contrasting previous findings on early emotion discrimination. Our results reveal interactive effects of voice familiarity on multimodal processing in infancy: while maternal speech enhances neural tracking, it may also alter how other social cues, such as faces, are processed. The findings suggest that early auditory experiences shape how infants allocate cognitive resources to social stimuli, emphasizing the need to consider cross-modal influences in early development.

Significance statement

How infants continuously process familiar social signals and how this familiarity shapes perception has remains unknown. Here we demonstrate that infants' brains preferentially track maternal speech over unfamiliar voices, highlighting the early tuning of the auditory system to socially relevant signals. Moreover, the maternal voice modulates the neural encoding of concurrently presented faces—altering the interplay between early visual and attentional processing—without eliciting emotion-specific differences. These findings underscore the role of caregiver signals in shaping multisensory integration during early development.

Keywords:

Neural tracking, EEG, infant, voice perception, face perception, voice familiarity, social development

Introduction

Infants are born into a social environment and, within their first year, develop a range of abilities which allow them to interact with their environment with increasing competence.

One fundamental aspect is the ability to recognize familiar people, most importantly their primary caregivers. A number of studies suggest that newborns quickly learn to recognize their mother's face (Sai, 2005) and voice (DeCasper & Fifer, 1980), and in the first months of life show a preference for such maternal signals (DeCasper & Fifer, 1980; Sai, 2005). Furthermore, social signals like speech elicit differential brain activation when originating from the mother versus an unfamiliar person (Dehaene-Lambertz et al., 2010; Naoi et al., 2012).

While these studies suggest a preferential orientation towards the maternal voice, it is unclear (a) whether infants preferentially process maternal speech over a longer duration and (b) whether this affects other social processes. To investigate these questions, we examined neural tracking of a continuous stream of maternal vs. unfamiliar speech. Neural tracking reflects the neural representation of a continuous signal, and is increasingly used to investigate infant speech processing (Jessen et al., 2019; Kalashnikova et al., 2018; Menn et al., 2022; Ortiz Barajas et al., 2021).

In particular, several studies have investigated which features of an auditory signal infants track most effectively. For example, infants show stronger tracking of infant-directed versus adult-directed speech (Kalashnikova et al., 2018; Menn et al., 2022) and intelligible versus unintelligible speech (Florea et al., 2024). Furthermore, infants' neural tracking reflects their attunement to their native language; while newborns show equal tracking of familiar and unfamiliar language input, by 6 months, they show weaker amplitude but not phase tracking of their native language (Ortiz Barajas et al., 2021). Auditory tracking in infants is also found for variants of speech input, such as songs (Nguyen et al., 2023) and nursery rhymes (Attaheri et al., 2022), where increased tracking can be found depending on song type (Nguyen et al., 2023). In summary, most previous studies compare infants' tracking of speech signals with different acoustic properties (infant-directed speech vs. adult-directed speech, familiar vs. unfamiliar language, play song vs. lullaby).

However, maternal and unfamiliar speech do not systematically differ in acoustic properties; the same speech signal can be familiar to one infant and unfamiliar to the next. Hence, potential differences in neural tracking cannot arise from low-level stimulus properties but must arise from familiarity with the voice as such, or from increased attention due to personal significance. While in adults, voice familiarity indeed enhances neural tracking (Yahav et al., 2024), it is unknown whether similar effects can be observed in infancy.

In addition to investigating the differential processing of maternal compared to an unfamiliar voice, we were interested in whether hearing the mother's voice influences unrelated social processes, as seen with other signals of maternal presence. Recent work for instance shows that maternal odor influences face categorization (Leleu et al., 2020) and emotional face processing (Jessen, 2020). However, social odor is a more subtle social signal, and often not consciously processed (Loos et al., 2025). Hence, hearing the mother's voice might be more attention grabbing, potentially

120 interfering with the processing of other social signals. For instance, it has been
121 observed in adults that hearing a task-irrelevant voice in the background impairs face
122 memory (Bell et al., 2019).
123
To investigate these questions, we studied 7-month-old infants in two sessions. In one,
they listened to a recording of their mother reading a children's story; in the other, they
heard another infant's mother reading the same story. While listening, infants viewed
photos of happy and fearful facial expressions. We analyzed event-related potentials
(ERPs) linked to attention (Nc; Guy et al., 2016; Xie et al., 2019) and face processing
(N290, P400; Guy et al., 2016; Xie et al., 2019), as well as neural tracking of faces and
voices, which allowed us to directly link voice and face tracking.
131

Methods

Pre-registration
Rationale, sampling plan, and major analyses were pre-registered prior to data acquisition at https://aspredicted.org/x4xc-rwbt.pdf. In what follows we will refer to this and indicate which analyses deviated from the pre-registered analyses.

Participants
A total of 39 7-month-old infants were invited to participate in the study. Five additional infants participated only in the first appointment and were therefore excluded from further analysis. The exact composition of the analysis sample differs for the two analysis tracks we will present (see below).

Infants were recruited via the maternity unit of the local hospital (Universitätsklinikum Schleswig-Holstein). All infants were born full-term (at least 37 weeks gestational age) with a birthweight of at least 2500 g, and had no known neurological deficits or visual or hearing impairments. The study was approved by the local ethics committee, conducted according to the Declaration of Helsinki, and parents provided written informed consent prior to the infant's participation.

For the evoked-potentials averaging (ERP) analysis, the final sample included N = 25 infants (age at first appointment: 214 ± 7 days (mean ± standard deviation), age at second appointment: 223 ± 8 days; 13 boys, 12 girls). The remaining 14 infants were excluded from the analysis because they did not contribute at least 10 trials per Emotion condition at one of the appointments (n= 9); had an overall mean Nc amplitude more than 2 standard deviations from the mean (n=3); or continued to cry at one of the appointments (n=2).

For the encoding-model analysis using multivariate temporal response functions (mTRF), the final sample included N = 30 7-month-old infants (15 boys, 15 girls, age at first appointment: 214 ± 7 days, age at second appointment: 221 ± 7 days). The remaining 9 infants were excluded because they did not contribute at least 100 artifact-free data segments at one of the appointments (n=6); continued to cry at one of the appointments (n=2); or because the trigger for the sound onset was not recorded (n=1). All but one infant included in the ERP analysis were also included in the mTRF analysis.

We aimed for a sample size of N = 25 in the ERP analysis based on our pre-registration. A sample size of N = 25 allowed us to detect effects of at least medium size (f = .25) in a 2 x 2 within-subjects ANOVA with a typical type-I error rate of .05 with a power of .8 (assuming a correlation of .5 of the repeated measures, G*Power 3.1), which seemed sufficient given the large effect observed in Jessen (2020) for the interaction between *Emotion* and *Familiarity* in a very similar paradigm (f = .39).

Stimuli
As visual stimuli, we used photographs of Caucasian female faces displaying happy and fearful facial expressions from the FACES database ((Ebner et al., 2010), ID 54, 63, 85, 90, 115, 173). Faces were cropped to an oval shape so that as little of the hair as possible was visible.

As auditory stimuli, we used an approximately one-minute-long recording of a
children's story, read and recorded by each infant's mother. The story was identical for
all participants and the recording was made at the first appointment prior to the actual
experiment. The mother was given a printout of the story and instructed to read the
story "as if you were reading it to your child". No other restrictions or instructions were
given regarding speaking style or speed. The recording was made via a standard
headset using Audacity and normalized to –23 RMS dBFS (dB full scale) after recording.
Each audio recording was used twice in the study: Once as a familiar stimulus for the
mother's own infant and once as an unfamiliar stimulus for a different infant.
191

Experimental Design
To investigate the impact of a familiar voice on the processing of emotional faces, we
followed a 2 x 2 design with the factors *Emotion* (happy, fearful) and *Voice* (familiar,
unfamiliar), in which infants heard a voice over loudspeakers while simultaneously
viewing photographs of faces presented on a screen.
Infants were invited to two appointments, ideally within two weeks. At one
appointment, infants would hear their mother's voice, while at the other appointment,
they would hear the unfamiliar voice of a different infant's mother. The order of
Familiarity condition was counterbalanced across infants.
Otherwise, the experimental design was identical for both appointments. A maximum
of 216 pictures were presented (18 per emotion and actress) arranged in blocks of 24
pictures (12 happy, 12 fearful), which were played consecutively without interruption.
Pictures were presented in a pseudorandomized order, ensuring that no picture was
repeated more than once. Each picture was presented for 800 ms, preceded by a
fixation cross shown for 300 ms and followed by an intertrial interval jittered between
700 and 1100 ms. The audio recording (familiar or unfamiliar voice, depending on the
appointment) was played without interruption in a loop throughout the entire
experiment in parallel to the visual presentation.
210

212 ***Figure 1. Experimental Set-up and Design****. Infants were seated in a car seat in front of a*
213 *screen with speakers on both sides of the screen. While the recording of the mother or the*
214 *stranger reading the same story was played continuously in the background, infants viewed*
215 *photographs of happy and fearful faces. Each recording was used twice in the study; once as a*
216 *familiar voice for the mother's own child and once as a stranger's voice for a different child.*
217

Procedure
Prior to the first appointment, parents were sent a set of questionnaires which they
were asked to complete at home and bring along to the lab appointment (EPDS (Cox et
al., 1987), IBQ-R (Gartstein & Rothbart, 2003; Vonderlin et al., 2012), and a lab-internal
questionnaire assessing demographic information as well as feeding and sleeping
routines of the infant). Questionnaire data are not considered further in the present
analysis.

Upon arrival at the lab, parents and infants were familiarized with the new environment
and parents were again explained the procedure and had the opportunity to ask any
remaining questions, before signing the informed consent form. The EEG recording
was prepared while the infant was sitting on their parent's lap. An elastic cap
(BrainCap, Easycap GmbH) containing 27 AgAgCl electrodes according to the
international 10-20 system was used for recording. The EEG signal was recorded at a
sampling rate of 500 Hz using a BrainAmp amplifier and the software
BrainVisionRecorder (both BrainProducts).
During the experiment, the infant was seated in an age-appropriate car seat (Maxi Cosi)
placed on the floor approximately 70 cm in front of a 24-inch monitor on which the
visual stimuli were presented. The face height was approximately 28 cm. Auditory
stimuli were presented via loudspeakers positioned left and right of the monitor set at
the same volume for all participants. Matlab (The MathWorks, Inc., Natick, MA) and the
toolbox Psychtoolbox (Brainard, 1997) were used to run the experiment. The parents
remained in the room with the infant but were instructed to maintain a distance of at
least 1 m behind the infant.
The infant was observed and videotaped throughout the experiment by a small camera
mounted on top of the monitor. If necessary, the experimenter could interrupt the
experiment to play short video clips containing colorful moving shapes and ringtones
to redirect the infant's attention to the screen. The experiment continued until all 216
trials had been presented (approximately 8 minutes) or the infant became too fussy to
continue the experiment.
247
<u>Data analysis</u>
EEG data were analyzed using Matlab 2023b (The MathWorks, Inc., Natick, MA), using
custom-made scripts was well as the toolbox FieldTrip (Oostenveld et al., 2011).
Jamovi (version 2.6.26) was used for the correlation analysis and mixed models for the
ERP results using the GAMLj package (version 2.4.0).
As preregistered, we analyzed the data focusing on differences in the ERP response to
emotional faces, neural tracking of familiar vs. unfamiliar voices, as well as the relation
between both measures. Exploratorily and in addition to the preregistered analyses,
we also included the responses to emotional faces in the encoding model.
257
<u>EEG preprocessing</u>
In a first step, data were offline re-referenced to the average reference; filtered using a
1- Hz highpass and 40-Hz lowpass filter; and segmented in 1-second epochs. Epochs
in which the standard deviation exceeded 100 μV in a sliding window of 200 ms at any
electrode were removed from further analysis. On the remaining data, we computed
an independent component analysis (ICA) to identify artifactual components. On
average, 5 $\pm$ 2 components (range: 0-9) were removed based on visual inspection of
time courses and topographies of the components.
266
For the ERP analysis, data were then filtered between 0.2 and 20 Hz, re-referenced to
the linked mastoids; and epoched into trials ranging from 200 ms before to 800 ms after
the picture onset. Any trials in which the standard deviation exceeded 80 μV in a sliding
window of 200 ms at any electrode were removed, and the data were inspected visually
for any remaining artifacts. Finally, trials in which the infants did not attend to the
screen based on the video recording were removed from further analysis.
Infants who contributed fewer than 10 trials per condition were excluded from further
analysis (n=9). Infants in the final sample contributed an average of 37 $\pm$ 19 trials per
condition (mother-happy: 34 $\pm$ 18; mother-fearful: 33 $\pm$ 16; stranger-happy: 42 $\pm$ 22;

stranger-fearful: 41 ± 20; since infants contributed more trials when they heard a
stranger's as opposed to their mother's voice [F(1,24) = 3.99, p=.057], we exploratorily
included number of trials as a covariate in the GLM, which had no effect on the
observed results).
280
281 For the neural encoding-model analysis, data were filtered between 1 and 10 Hz, but
282 no further visual inspection was carried out (Fiedler et al., 2019; Jessen et al., 2019; Orf
283 et al., 2023).
284
Encoding model analysis
A temporal response function (TRF) serves as a simplified model of brain activity,
illustrating how the brain processes a given stimulus feature to generate the observed
EEG signal, assuming a linear filter mechanism. To compute the TRF, we utilized a
multiple linear regression approach (Crosse et al., 2016). To predict the recorded EEG
response, we trained a forward model per EEG channel using the envelopes of the
mothers and stranger streams (e.g., Fiedler et al., 2019). In this framework, we
examined delays between envelope changes and brain responses of between -200 and
800 ms.
To account for EEG variance associated with the processing of presented faces and
their corresponding evoked responses, we incorporated triggers for face presentations
as regressors using stick functions. A stick function is a modeling vector in which non-
event times are denoted by zeros, while events, such as the onset of a face, are marked
with a value of one.
To prevent overfitting, we used ridge regression to estimate the TRF and determined the
optimal ridge parameter through leave-one-out cross-validation for each participant.
We predefined a range of ridge values, calculated a separate model for each value, and
averaged across trials to predict the neural response for each test set. The ridge
parameter with the lowest mean squared error (MSE) was selected as the optimal
value specific to each subject. One model was trained using predictor variables for the
envelopes of mother and stranger streams, as well as stick functions for presented
faces (Orf et al., 2023). These were modelled jointly (same regressor matrix) using the
same regularization.
Neural speech tracking quantifies the representation of a single stream within the EEG
signal by using TRFs to predict the EEG response. The neural tracking metric (r) was
computed by correlating the predicted EEG response with the actual EEG signal using
Pearson correlation. A leave-one-out cross-validation approach was used to predict
the EEG signal. To assess neural tracking accuracy over TRF time lags, we used a sliding
time window (48 ms, 6 samples, 24 ms overlap), resulting in a time-resolved measure
of neural tracking (Fiedler et al., 2019; Hausfeld et al., 2018; Kraus et al., 2021;
O'Sullivan et al., 2015).
316
ERP analysis
For the ERP analysis, we computed single-subject averages for all four conditions
(mother-happy, mother-fearful, stranger-happy, stranger-fearful).
We computed the Nc response as the mean value in a time-window of 400–800 ms
after picture onset averaged across F3, Fz, F4, C3, Cz, C4, and the N290 and P400 as
the mean value in a time-window of 200–300 ms respectively 300–500 ms after picture
onset averaged across O1 and O2. These time-windows and electrodes correspond to
those preregistered and are based on prior studies (Jessen, 2020; Quadrelli et al.,
2019). Exploratorily, we also computed the Nc in a time-window from 350 to 600 ms,

as inspection of the average across all conditions revealed an Nc peak in this earlier
time-window.
Participants whose Nc amplitude was more than 2 standard deviations from the
average across all conditions between 400 and 800 ms were excluded from further
analysis (n=3).
331
<u>Statistical Analysis</u>
We statistically evaluated both, conventional ERP results and encoding model
outcomes, using linear mixed-effects models. These models account for the
hierarchical structure in multi-condition, within-subject designs by combining fixed
effects and random effects. Predictors were centered and scaled as needed. Random
intercepts were included where appropriate to account for inter-individual differences.
338
339 The mTRFs were analyzed in Matlab using the fitlme function and the following model:
340
341   `Audio_EncodingAccuracy ~ 1 + VoiceFamiliarity + Session +`
342   `Faces_EncodingAccuracy_central + BaselinePower + (1 | SubjectID)`
343
where *VoiceFamiliarity* was the voice heard (mother or stranger). We additionally
included *Session* (1 or 2) to account for potential order effects and *SubjectID* to
account for interindividual variance. Furthermore, *BaselinePower* (log-transformed)
served as a measure of noise in the infant data, while
Faces_EncodingAccuracy_central at central electrodes (C3, Cz, C4) was included to
control for the impact of the visual stimuli (faces viewed by the infant) on encoding
accuracy.
351
352 We also computed the following model to the assess impact of voice encoding on
353 central face processing:
354
355   `Face_EncodingAccuracy_central ~ 1 + VoiceFamiliarity + Emotion +`
356   `Audio_EncodingAccuracy + Face_EncodingAccuracy_occipital + Session`
357   `+ (1 | SubjectID)`
358
359 Here, we additionally included *Emotion*, accounting for the emotion expressed in the
360 picture seen (happiness or fear) and face encoding at occipital electrodes (O1, O2) as
361 *Face_EncodingAccuracy_occipital* to account for visual processing. Including
362 BaselinePower in the model did not further improve the model fit.
363
Not shown in the pseudo-formula above are all two-way interaction terms of the
regressors of interest, which we included in the model as well. A random slope for the
significant *Face_EncodingAccuracy_occipital*VoiceFamiliarity* effect improved this
model further (likelihood ratio test <.001) and was accordingly included in the final
model, although the conclusions from this model do not depend on this modelling
choice.
370
371 Furthermore, we followed this up with an alternative but similarly set up model, where
372 the roles of the variables *Face_EncodingAccuracy_central* and
373 *Face_EncodingAccuracy_occipital* were swapped to model face encoding at occipital
374 electrodes specifically.
375
376 We analyzed the ERP data in Jamovi using the following model for each ERP component
377 separately:

```
ERP amplitude ~ 1 + VoiceFamiliarity *Emotion + VoiceFamiliarity +
Emotion + Session + (1 | SubjectID)
```

Models additionally including *Trialnumber* (reflecting the number of trials in each condition) and/or *Breastfeeding* (whether the infant was still breastfed or not) yielded the same results as reported below.

Finally, to link our two EEG analysis approaches and for completeness, we also report a simple Pearson correlation between *Audio_EncodingAccuracy* and the ensuing difference in Nc response to happy and fearful faces.
For the pre-registered hypotheses (see below) and substantive null effects of interest (i.e., the absence of effects of Emotion in face encoding models), we report Bayes Factors in favor of the null effect derived from the Jamovi model output using Eq. 6 in Faulkenberry (2018) (see also Wagenmakers, 2007).

Data availability
Upon publication, data and anaylsis scripts will be made available at http://osf.

Acoustic properties of voice stimuli

Figure 2. Acoustic properties of voice signals. *As regressor for auditory neural tracking, we used the voice envelope, as shown in the left panel. The envelope was extracted from both the mother and stranger condition audio signal, and the peak frequency was identified from the spectrum using the Fast Fourier Transform (FFT). Since the same recordings were used once as a stranger and once as a familiar (mother's) voice, they showed very similar acoustic properties, as can be seen in the right panel. Histograms and Kernel Density Estimates (KDE) with a bandwidth of 0.5 were used to visualize the distribution of peak frequencies across subjects for both conditions. (Note that the two stimulus distributions are not identical as not all infants could be included in the final sample, so that for a small number of infants the recording was only included in one of the conditions).*

Results

<u>Infants' neural speech tracking is sensitive to voice familiarity</u>
The temporal response function showed clear signatures of the infant brain neurally tracking both, the mother's as well as the stranger's voice. The frontocentral bilateral distribution (Figure 3A) of this neural tracking is expected and well in line with the adult neural tracking literature, broadly commensurate with bilateral auditory (i.e., perislyvian cortical) origins.

However, tracking of the mother's voice was consistently stronger than tracking of a stranger's voice (Figure 3), despite both having comparable acoustic properties (Figure 2). This is highlighted in Figure 3C, which compares the encoding accuracy within the time window with the highest TRF and time-shifted neural tracking (200–300 ms). A statistically significant enhancement in neural tracking for the mother's voice compared to the stranger's voice is evident (b = -0.004, SE = 0.001, t(84.8) = -4.72, p < .001, for full results see Supp Material, Table SM1).

<u>Infant encoding models of faces are sensitive to voice familiarity</u>
Furthermore, we clearly observed neural tracking of faces at both, central and occipital electrodes (i.e., face encoding accuracies greater than zero; Figure 4). This neural face-encoding, however, was modulated by the concurrent presence or absence of the mother's familiar voice.

When infants heard a stranger as opposed to their mother, central electrodes showed an enhanced face encoding (Effect of *VoiceFamiliarity* on *Central face tracking*; b=-0.005, SE = 0.002, t(80.5) = -3.06, p=.003, for full results see Supp Material, Table SM2). This can also be expressed as a relatively diminished encoding of (unknown) faces when the mother's voice is present (Figure 4A).

Note that this central face tracking was also partially accounted for by simultaneous encoding accuracy from occipital electrodes (*occipital face tracking*; b=0.011, SE = 0.0023, t(22.9) = 4.52, p<.001). This lends plausibility to the overall results of the forward encoding model (which establishes encoding accuracies per electrode): It confirms that infants who showed stronger tracking at occipital electrodes also showed stronger tracking at central electrodes.

Intruigingly, the data also revealed an interaction between *VoiceFamiliarity* and *occipital face tracking*, as shown in Figure 4B (interaction: b= -0.003, SE = 0.0015, t(35.8) = -2.27, p=.029). Only when infants heard a stranger's voice and occipital face tracking was accordingly high, did infants showed increased central-electrode face tracking. When infants instead heard their mother, occipital face tracking did not show this 'gating effect' on central face tracking accuracy.

When, for completeness, modelling the face encoding response at occipital electrodes in the same time window, the generally positive association with face trackign at central electrodes replicates (b = 0.014, SE = 0.0027, t(18.5) = 5.03, p<.001, for full results see Supp Material, Table SM3), also reflected in a simple and expected bivariate correlation between central and occipital face tracking (r=.51, p<.001). However, in contrast to the pattern observed at central electrodes, we did not find the interaction pattern with *VoiceFamiliarity*. We will return to this in the Discussion section, but this pattern of results is indirect evidence for a plausible, directed flow of neural information, where brain areas captured by occipital-electrode encoding

463 accuracy hierarchically gate or precede information contained in more fronto-central
464 brain areas that constitute the Nc-like face encoding accuracies at central electrodes.
465
<u>Absence of facial emotion effects</u>
We did not find an effect of *Emotion* expressed in the face stimuli on the temporal
response function or the resulting encoding accuracy at either central or occipital
electrodes (both Bayes Factors >10 in favour of the null hypothesis). Neither of the
Emotion differences discernable in some time windows and electrodes did surpass
conventional levels of statistical significance. This speaks to a generally reduced
impact of *Emotion* on infants' neural responses in this paradigm with a spoken-
language backing track, compared to many previous infant studies on face processing
without such auditory input.

Figure 3. Temporal response functions (TRFs) and neural speech tracking. A. *TRF weights*
are averaged across infants and channels of interest (Cz, C3, C4). The shaded areas show the
standard error for each time lag across infants. The inset displays the topographical
distribution of TRF weights for a time window between 200-300 ms (gray bar). Blue colors
indicate TRFs for the mother's voice, while orange colors indicate TRFs for the stranger's voice.
B. *Neural tracking unfolds over time lags (-200 to 800 ms). Solid lines show the averaged*
neural tracking (r) across subjects and channels of interest (topographic map). Shaded areas
represent the standard error for each time lag across infants. ***C.*** *Unadjusted neural tracking (r)*
refers to the encoding accuracy (200-300 ms), calculated by multiplying the estimated TRFs by
the presented audio envelope. The line plot shows single-infants data (N = 30), averaged over
channels of interest (Cz, C3, C4). Connecting lines between dots indicate the same subject.
Blue dots represent tracking of the mother's voice, while orange dots represent tracking of the
stranger's voice. The inset shows adjusted neural tracking based on predictive model results,
including covariates such as baseline power and face tracking.
490

A Visual mTRF at central and occipital electrodes

B Visual tracking accuracy at occipital and central electrodes

491
Figure 4. Temporal response function and neural face tracking. A. *Clear tracking was found*
at central (C3, Cz, C4) and occipital (O1, O2) electrodes, which was lower when infants heard
their mother (left panel; blue/green) compared to a stranger (right panel; red/ orange). We did
not find an impact of face emotion, suggesting that happy (green/red) and fearful faces
(blue/orange) were tracked to a similar extent. Faces were encoded at both, central and
occipital electrodes, as can be seen in the topographic representations (350 - 500 ms). **B.**
Encoding at central electrodes was modulated by both, voice identity (blue = mother, orange =
stranger) and occipital encoding, such that higher encoding at occipital electrodes was
associated with higher encoding at central electrodes specifically when a stranger's voice was
heard. As can be seen from the topographic representations, differences where focused on
*central and occipital electrodes as can be expected for face processing. [*p<.05]*

 Infant evoked responses to faces do not yield effects of voice familiarity or emotion
content
The purported superior sensitivity of the encoding-model approach is underscored
when looking at the conventional ERP analysis to faces in these data. We did not find
significant main effects of *VoiceFamiliarity* or *Emotion*, nor an interaction between the
two on any of the ERP components investigated (all p>. 1, all BF10 < .35; see Figure 5
for ERP response at central and occipital electrodes and Supp Material Table SM4–7
for the full statistical results).

The difference in Nc response to fearful and happy faces, as obtained in the
conventional ERP analysis and often reported in infant ERP studies, was mildly
positively correlated with the strength of auditory tracking of the ongoing voice, as
derived from the encoding model (r=.24, p=.10, Supp Material Table SM8).

Figure 5. Conventional evoked-response analysis to faces. *Shown are ERP responses at central (A) and occipital electrodes (B) to the presentation of happy (lighter colors) and fearful faces (darker colors) while infants listened to either their mother's voice (left panel, blue/green) or a stranger's voice (right panel, orange/red). Topographic representations show the average for time-windows analyzed for ERP responses (A: Nc response in a time-window of 400 to 800 ms; B: N290 in a time-window of 200 to 300 ms and P400 in a time-window of 300 to 500 ms. Dots indicate the electrodes averaged (A: F3, Fz, F4, C3, Cz, C4; B: O1, O2). No significant differences were observed.*

Discussion

Can we trace infants' neural processing of familiar versus unfamiliar voices and its potential impact on the processing emotional faces? In doing so, we have leveraged neural encoding models of the infant EEG and have contrasted them with conventional evoked-response analyses (ERP).

Infants showed a stronger neural tracking of their mother's compared to a stranger's voice. Furthermore, infants' face processing was affected by the concurrently heard voice; infants hearing their mother showed a stronger increase in central face tracking with increasing occipital tracking. While infants thus successfully encode faces, we found no difference in processing between fearful and happy faces.

Neural encoding of maternal versus unfamiliar voices
The robust tracking of voice information and the enhanced tracking of maternal over an unfamiliar voices adds to a growing body of literature investigating the continuous tuning of the developing brain to relevant auditory social signals (Jessen et al., 2019; Kalashnikova et al., 2018; Menn et al., 2022; Nguyen et al., 2023).

For the first time, we show that enhanced tracking occurs not only for auditory signals that are either specifically tailored to infants (e.g., infant-directed speech or nursery rhymes, (Kalashnikova et al., 2018; Menn et al., 2022; Nguyen et al., 2023)) or native language input (Ortiz Barajas et al., 2021), but also for voices carrying distinct signals for an individual infant. This enables the infant to attend to personally relevant social cues, facilitating bonding and social learning.

One explanation for stronger tracking of maternal voice is better predictability due to the high familiarity. Such an explanation would be comparable to the mechanisms observed in the case of infant-directed speech or lullabies, both of which are characterized by a high degree of rhythmicity, large modulations in pitch, and simplified structure, all features that increase predictability and thereby facilitate processing (Kalashnikova et al., 2018). For maternal speech, increased predictability is more subtle and individualized, resulting from the idiosyncrasies of each speaker. A similar effect is seen in adults, where speaker familiarity improves comprehension for hearing-impaired listeners (Souza et al., 2013) and even brief familiarization benefits listeners with intact hearing (Nygaard & Pisoni, 1998).

While such an explanation is primarily based on perceptual facilitation, another explanation could be derived from attention-related processes. Since maternal signals are highly salient and of particular importance to the infant, infants allocate more attentional resources to maternal signals (DeCasper & Fifer, 1980; Rigato et al., 2023). As has been extensively studied in adults, increased attention allocation leads to increased neural tracking (e.g., Fiedler et al., 2019; Mesgarani & Chang, 2012; Orf et al., 2023; Reetzke et al., 2021), which is likely to be the case also in infancy.

Neural encoding of faces is influenced by voice familiarity
While an increasing number of studies investigate auditory tracking, visual tracking is less commonly investigated. However, initial studies suggest neural tracking also of continuous visual signals (Jessen et al., 2019; Ki et al., 2021). In our study, we did not investigate the tracking of a continuous visual signal but of discrete visual events optimized for an ERP design. Nevertheless, we observed clear visual tracking at both

central and occipital electrodes, corresponding in both, topography and latency, to well-known infant ERP components typically observed in face processing (occipital N290-P400 and central Nc; Conte et al., 2020; Xie et al., 2019). This observation is confirmed by the more traditional ERP analysis. Thus, infants successfully encode faces even when listening to unrelated voice information in parallel, and such a processing is observed not only in the ERP response but also in neural tracking.

Moreover, the encoding of faces is not independent of voice processing. Voice information modulated the relation between occipital and central encoding; when hearing an unfamiliar voice, stronger occipital encoding was linked to stronger central encoding. Here, using mTRFs to analyze face processing offers a distinct advantage as it allows us to directly control for simultaneous auditory tracking in the same model.

While purely speculative at this point, these data might suggest that an unfamiliar voice causes infants to attend more closely to other types of social information. Conversely, maternal presence reduces the infants' need to attend to unfamiliar faces (see Jessen, 2020, for a qualitatively similar effect of maternal odor).

Specifically, the interaction of maternal-voice presence and occipital encoding accuracy on the simultaneous central, Nc-like face encoding accuracy suggests that this process depends on an interplay between sensory and attentional processes. Only infants who show a better occipital encoding also show an increased effect on central, attention-related processes (i.e., a "gating effect"). Note that while encoding at occipital and central electrodes does correlate and there is certainly a link between the two processes, they share only $r^2 = 26\%$ of variance and are not identical.

While linear models per se can rarely be interpreted causally, this interaction pattern in explaining central-electrode face encoding was intriguingly non-reversible: When instead modelling occipital-electrode encoding accuracy of faces, putatively further 'upstream' in a cortical processing hierarchy, no interaction of (central) face encoding with voice familiarity was found. We take these results as putative evidence that the impact of voice familiarity on more 'downstream', Nc-like central face tracking processes is 'gated by' (i.e., statistically dependent on) the relative strength of a more 'upstream' occipital encoding of these faces. The result clearly calls for a more elaborate neuroimaging approach to sensory–occipital vs attentional–frontocentral processes in the infant brain.

In sum, the analyses of neural tracking revealed that infants track faces effectively even while hearing an unrelated voice, and voice familiarity influences how faces are processed

Absence of emotion effect in facial processing
Contrary to our prediction, we found no evidence that maternal voice influenced emotion processing, nor did we find evidence for emotion processing per se. This is surprising, given the vast body of literature showing that 7-month-old infants reliably discriminate emotions at the neural level (e.g. Aran et al., 2023; Bowman et al., 2022; Leppanen et al., 2007; Peltola et al., 2009; Xie et al., 2019), and our own previous work, which found robust evidence for emotion discrimination in a very similar set-up using the same visual stimuli (Jessen, 2020). At this point, we can only speculate about possible explanations.

Note that the absence of an emotion effect for infants exposed to maternal signals
aligns with our previous findings, where infants showed evidence of emotion
discrimination only when exposed to social signals from a stranger (Jessen, 2020).
Descriptively, this is also observed in the present study at later occipital responses,
which show emotion-related differences when the infant is hearing a stranger's voice
but not when they are hearing their mother (in both ERP and mTRF).
644
However, the question remains as to why no significant difference was observed. While
infants at this age typically show neural emotion differentiation, not all studies report
emotion differentiation in all three ERP components. For example, both Aran et al.
(2023) and Xie et al. (2019) only found a differential processing of happy and fearful
faces on the N290 but not on the P400 or Nc, while Leppanen et al. (2007) report no
difference on the N290 but rather the P400 and Nc. Hence, while emotion processing
in infancy has been extensively studied, it is still unclear under which conditions which
neural processes are affected by affective content.
653
One possibility is that while background voices did not disrupt face processing, they
may have hindered detailed extraction of emotional cues. To our knowledge, no EEG
study so far has investigated emotional face processing with concurrent, unrelated
voice signals. Infants might for instance have been distracted by attending to the voice;
an indicator of such an effect might be the fact that infants contributed more trials
when listening to a stranger versus their mother (although this effect was only
marginally significant). Other potential indicators, such as an interaction between
auditory encoding accuracy and emotion or a correlation between auditory encoding
accuracy and difference in Nc amplitude were not observed. On the contrary, on a
descriptive level, higher auditory encoding accuracy was associated with a higher Nc
amplitude for fearful compared to happy faces ($r = .24$, $p = .1$), which would suggest an
improved discrimination with increasing encoding accuracy.
Future studies are clearly necessarily to further explore this issue and examine the
robustness of infants' emotion processing under conditions of potential distraction.
668
669
Conclusion
Infants are exposed to a wide variety of social signals in their daily lives. We
investigated the interplay between two different types of signals, voice and faces,
providing unrelated information. Infants showed a stronger tracking of their mother's
than of a stranger's voice and this tracking impacted how well unrelated faces were
encoded. Surprisingly, we found no evidence for more sophisticated face processing,
such as emotion discrimination. Our results further highlight the potential of using
neural tracking to uncover the complex relationship between different types of social
processes in early development.
679

680 **References**
681
682

Aran, O., Garcia, S. E., Hankin, B. L., Hyde, D. C., & Davis, E. P. (2023). Signatures
of emotional face processing measured by event-related potentials in 7-
month-old infants. *Developmental Psychobiology*, *65*(2), e22361.
https://doi.org/10.1002/dev.22361
Attaheri, A., Choisdealbha, A. N., Di Liberto, G. M., Rocha, S., Brusini, P., Mead,
N., Olawole-Scott, H., Boutris, P., Gibbon, S., Williams, I., Grey, C.,
Flanagan, S., & Goswami, U. (2022). Delta- and theta-band cortical
tracking and phase-amplitude coupling to sung speech by infants.
Neuroimage, *247*, 118698.
https://doi.org/10.1016/j.neuroimage.2021.118698
Bell, R., Mieth, L., Roer, J. P., & Buchner, A. (2019). Effects of Auditory Distraction
on Face Memory. *Sci Rep*, *9*(1), 10185. https://doi.org/10.1038/s41598-
019-46641-7
Bowman, L. C., McCormick, S. A., Kane-Grade, F., Xie, W., Bosquet Enlow, M.,
& Nelson, C. A. (2022). Infants' neural responses to emotional faces are
related to maternal anxiety. *J Child Psychol Psychiatry*, *63*(2), 152-164.
https://doi.org/10.1111/jcpp.13429
Brainard, D. H. (1997). The Psychophysics Toolbox. *Spat Vis*, *10*(4), 433-436.
https://www.ncbi.nlm.nih.gov/pubmed/9176952
Conte, S., Richards, J. E., Guy, M. W., Xie, W., & Roberts, J. E. (2020). Face-
sensitive brain responses in the first year of life. *Neuroimage*, *211*,
116602. https://doi.org/10.1016/j.neuroimage.2020.116602
Cox, J. L., Holden, J. M., & Sagovsky, R. (1987). Detection of postnatal
depression. Development of the 10-item Edinburgh Postnatal
Depression Scale. *Br J Psychiatry*, *150*, 782-786.
https://doi.org/10.1192/bjp.150.6.782
Crosse, M. J., Di Liberto, G. M., Bednar, A., & Lalor, E. C. (2016). The Multivariate
Temporal Response Function (mTRF) Toolbox: A MATLAB Toolbox for
Relating Neural Signals to Continuous Stimuli. *Front Hum Neurosci*, *10*,
604. https://doi.org/10.3389/fnhum.2016.00604
DeCasper, A. J., & Fifer, W. P. (1980). Of Human Bonding - Newborns Prefer Their
Mothers Voices. *Science*, *208*(4448), 1174-1176. https://doi.org/DOI
10.1126/science.7375928
Dehaene-Lambertz, G., Montavont, A., Jobert, A., Allirol, L., Dubois, J., Hertz-
Pannier, L., & Dehaene, S. (2010). Language or music, mother or Mozart?
Structural and environmental influences on infants' language networks.
Brain Lang, *114*(2), 53-65. https://doi.org/10.1016/j.bandl.2009.09.003
Ebner, N. C., Riediger, M., & Lindenberger, U. (2010). FACES--a database of
facial expressions in young, middle-aged, and older women and men:
development and validation. *Behav Res Methods*, *42*(1), 351-362.
https://doi.org/10.3758/BRM.42.1.351
Faulkenberry, T. J. (2018). Computing Bayes factors to measure evidence from
experiments: An extension of the BIC approximation. *Biometrical*
Letters, *55*(1), 31-43. https://doi.org/10.2478/bile-2018-0003

Fiedler, L., Wostmann, M., Herbst, S. K., & Obleser, J. (2019). Late cortical
tracking of ignored speech facilitates neural selectivity in acoustically
challenging conditions. *Neuroimage*, *186*, 33-42.
https://doi.org/10.1016/j.neuroimage.2018.10.057

Florea, C., Reimann, M., Schmidt, F., Preiss, J., Angerer, M., Ameen, M., Heib,
D., Roehm, D., & Schabus, M. (2024). Neural speech tracking in
newborns: prenatal learning and contributing factors. *bioRxiv*.
https://doi.org/https://doi.org/10.1101/2024.03.18.585222

Gartstein, M. A., & Rothbart, M. K. (2003). Studying infant temperament via the
Revised Infant Behavior Questionnaire. *Infant Behavior & Development*,
26(1), 64-86. https://doi.org/Pii S0163-6383(02)00169-8
Doi 10.1016/S0163-6383(02)00169-8

Guy, M. W., Zieber, N., & Richards, J. E. (2016). The Cortical Development of
Specialized Face Processing in Infancy. *Child Dev*, *87*(5), 1581-1600.
https://doi.org/10.1111/cdev.12543

Hausfeld, L., Riecke, L., Valente, G., & Formisano, E. (2018). Cortical tracking of
multiple streams outside the focus of attention in naturalistic auditory
scenes. *Neuroimage*, *181*, 617-626.
https://doi.org/10.1016/j.neuroimage.2018.07.052

Jessen, S. (2020). Maternal odor reduces the neural response to fearful faces in
human infants. *Dev Cogn Neurosci*, *45*, 100858.
https://doi.org/10.1016/j.dcn.2020.100858

Jessen, S., Fiedler, L., Munte, T. F., & Obleser, J. (2019). Quantifying the
individual auditory and visual brain response in 7-month-old infants
watching a brief cartoon movie. *Neuroimage*, *202*, 116060.
https://doi.org/10.1016/j.neuroimage.2019.116060

Kalashnikova, M., Peter, V., Di Liberto, G. M., Lalor, E. C., & Burnham, D. (2018).
Infant-directed speech facilitates seven-month-old infants' cortical
tracking of speech. *Sci Rep*, *8*(1), 13745.
https://doi.org/10.1038/s41598-018-32150-6

Ki, J. J., Dmochowski, J. P., Touryan, J., & Parra, L. C. (2021). Neural responses to
natural visual motion are spatially selective across the visual field, with
selectivity differing across brain areas and task. *Eur J Neurosci*, *54*(10),
7609-7625. https://doi.org/10.1111/ejn.15503

Kraus, F., Tune, S., Ruhe, A., Obleser, J., & Wostmann, M. (2021). Unilateral
Acoustic Degradation Delays Attentional Separation of Competing
Speech. *Trends Hear*, *25*, 23312165211013242.
https://doi.org/10.1177/23312165211013242

Leleu, A., Rekow, D., Poncet, F., Schaal, B., Durand, K., Rossion, B., & Baudouin,
J. Y. (2020). Maternal odor shapes rapid face categorization in the infant
brain. *Dev Sci*, *23*(2), e12877. https://doi.org/10.1111/desc.12877

Leppanen, J. M., Moulson, M. C., Vogel-Farley, V. K., & Nelson, C. A. (2007). An
ERP study of emotional face processing in the adult and infant brain.
Child Dev, *78*(1), 232-245. https://doi.org/10.1111/j.1467-
8624.2007.00994.x

Loos, H. M., Schaal, B., Pause, B. M., Smeets, M. A. M., Ferdenzi, C., Roberts, S.
C., de Groot, J., Lubke, K. T., Croy, I., Freiherr, J., Bensafi, M., Hummel, T.,

& Havlicek, J. (2025). Past, Present, and Future of Human Chemical
Communication Research. *Perspect Psychol Sci*, *20*(1), 20-44.
https://doi.org/10.1177/17456916231188147

Menn, K. H., Michel, C., Meyer, L., Hoehl, S., & Mannel, C. (2022). Natural infant-
directed speech facilitates neural tracking of prosody. *Neuroimage*, *251*,
118991. https://doi.org/10.1016/j.neuroimage.2022.118991

Mesgarani, N., & Chang, E. F. (2012). Selective cortical representation of
attended speaker in multi-talker speech perception. *Nature*, *485*(7397),
233-236. https://doi.org/10.1038/nature11020

Naoi, N., Minagawa-Kawai, Y., Kobayashi, A., Takeuchi, K., Nakamura, K.,
Yamamoto, J., & Kojima, S. (2012). Cerebral responses to infant-directed
speech and the effect of talker familiarity. *Neuroimage*, *59*(2), 1735-
1744. https://doi.org/10.1016/j.neuroimage.2011.07.093

Nguyen, T., Reisner, S., Lueger, A., Wass, S. V., Hoehl, S., & Markova, G. (2023).
Sing to me, baby: Infants show neural tracking and rhythmic movements
to live and dynamic maternal singing. *Dev Cogn Neurosci*, *64*, 101313.
https://doi.org/10.1016/j.dcn.2023.101313

Nygaard, L. C., & Pisoni, D. B. (1998). Talker-specific learning in speech
perception. *Percept Psychophys*, *60*(3), 355-376.
https://doi.org/10.3758/bf03206860

O'Sullivan, J. A., Power, A. J., Mesgarani, N., Rajaram, S., Foxe, J. J., Shinn-
Cunningham, B. G., Slaney, M., Shamma, S. A., & Lalor, E. C. (2015).
Attentional Selection in a Cocktail Party Environment Can Be Decoded
from Single-Trial EEG. *Cereb Cortex*, *25*(7), 1697-1706.
https://doi.org/10.1093/cercor/bht355

Oostenveld, R., Fries, P., Maris, E., & Schoffelen, J. M. (2011). FieldTrip: Open
Source Software for Advanced Analysis of MEG, EEG, and Invasive
Electrophysiological Data. *Computational Intelligence and*
Neuroscience, *2011*. https://doi.org/Artn 156869
10.1155/2011/156869

Orf, M., Wostmann, M., Hannemann, R., & Obleser, J. (2023). Target
enhancement but not distractor suppression in auditory neural tracking
during continuous speech. *iScience*, *26*(6), 106849.
https://doi.org/10.1016/j.isci.2023.106849

Ortiz Barajas, M. C., Guevara, R., & Gervain, J. (2021). The origins and
development of speech envelope tracking during the first months of life.
Dev Cogn Neurosci, *48*, 100915.
https://doi.org/10.1016/j.dcn.2021.100915

Peltola, M. J., Leppanen, J. M., Maki, S., & Hietanen, J. K. (2009). Emergence of
enhanced attention to fearful faces between 5 and 7 months of age. *Soc*
Cogn Affect Neurosci, *4*(2), 134-142.
https://doi.org/10.1093/scan/nsn046

Quadrelli, E., Conte, S., Cassia, V. M., & Turati, C. (2019). Emotion in motion:
Facial dynamics affect infants' neural processing of emotions.
Developmental Psychobiology, *61*(6), 843-858.
https://doi.org/10.1002/dev.21860

Reetzke, R., Gnanateja, G. N., & Chandrasekaran, B. (2021). Neural tracking of
the speech envelope is differentially modulated by attention and
language experience. *Brain Lang*, *213*, 104891.
https://doi.org/10.1016/j.bandl.2020.104891

Rigato, S., Stets, M., Charalambous, S., Dvergsdal, H., & Holmboe, K. (2023).
Infant visual preference for the mother's face and longitudinal
associations with emotional reactivity in the first year of life. *Sci Rep*,
13(1), 10263. https://doi.org/10.1038/s41598-023-37448-8

Sai, F. Z. (2005). The role of the mother's voice in developing mother's face
preference: Evidence for intermodal perception at birth. *Infant and Child*
Development, *14*(1), 29-50. https://doi.org/10.1002/icd.376

Souza, P., Gehani, N., Wright, R., & McCloy, D. (2013). The advantage of knowing
the talker. *J Am Acad Audiol*, *24*(8), 689-700.
https://doi.org/10.3766/jaaa.24.8.6

Vonderlin, E., Ropeter, A., & Pauen, S. (2012). Erfassung des fruehkindlichen
Temperaments mit dem Infant Behavior Questionnaire Revised.
Zeitschrift fuer Kinder- und Jugendpsychiatrie und Psychotherapie,
40(5), 307-314.

Wagenmakers, E. J. (2007). A practical solution to the pervasive problems of p
values. *Psychon Bull Rev*, *14*(5), 779-804.
https://doi.org/10.3758/bf03194105

Xie, W., McCormick, S. A., Westerlund, A., Bowman, L. C., & Nelson, C. A.
(2019). Neural correlates of facial emotion processing in infancy. *Dev*
Sci, *22*(3), e12758. https://doi.org/10.1111/desc.12758

Yahav, P. H.-s., Sharaabi, A., & Golumbic, E. Z. (2024). The effect of voice
familiarity on attention to speech in a cocktail party scenario. *Cereb*
Cortex, *34*, 1-16.
https://doi.org/https://doi.org/10.1093/cercor/bhad475