Title: Effects of urbanization on the microbial fungal communities from water, sediment, and

amphibian hosts across a rural-to-urban waterway in Worcester, Massachusetts

Sara J. Wheeler[a], Manning DelCogliano[a], Brennan Hare[a], Alexander J. Bradshaw[a], Madison R. Hincher[a],

Jasper P. Carleton[a], Philip J. Bergmann[a], Nathan Ahlgren[a], Javier F. Tabima[a,1]

a. Biology Department, Clark University. Worcester, Massachusetts.

1. Corresponding author (jtabima@clarku.edu)

ABSTRACT:

In a world of increasingly urbanized environments, it is critical to understand the impact of urbanization

on microbial communities, including fungal communities, as a measure of ecosystem health, and to

document how these environments are changing. Aquatic environments, in particular, can be highly

sensitive to urbanization with removal of local habitat and inputs of wastewater and contaminants, leading

to displacement or extinction of natural flora, fauna, and funga. Aquatic microbial communities,

especially fungal communities, are extremely complex and far less characterized compared to their macro

counterparts. Here, we characterized the fungal microbiome, i.e. "mycobiome", across multiple habitats—

water, sediment, and frog fecal matter from three species (American bullfrog (*Lithobates catesbeianus*),

Green frog (*Lithobates clamitans*), and Pickerel frog (*Lithobates palustris*)) — within the Tatnuck Brook

waterway in Worcester, MA. The Tatnuck Brook waterway is a connected group of streams, ponds, and

lakes that transition from protected areas at its headwaters, below the city's drinking water reservoir, to

highly urbanized regions within Worcester. Using metabarcode sequencing, we found that water,

sediment, and frog gut habitats harbor distinct fungal communities, but all exhibit parallel shifts in

diversity along an urbanization gradient. In particular, we identified fungal taxa from the amphibian

mycobiome and environmental samples that exhibited a range of sensitivities to urbanization. These taxa,

including *Basidiobolus*, *Cladosporium*, and *Lemonniera* in fecal and sediment samples, as well as

Candida, which was found in all habitats, are potential indicators of shifts in aquatic ecosystems due to

urbanization. Their responses to urban stressors may serve as a baseline for further studies of aquatic

fungal communities.

IMPORTANCE:

As cities grow, they can change nearby rivers, streams, and ponds in ways that affect plants, animals, and

even tiny microbes. Fungi are an important part of these ecosystems because they help break down

materials and can affect the health of animals like frogs. However, we still know very little about how

fungi respond to urban environments. In this study, we looked at fungi living in water, sediment, and frog

guts along a range of places from natural to highly developed areas in Worcester, Massachusetts. We

found that some fungi were more common in cleaner, less developed sites, while others were more

common in urban ones. These patterns suggest that fungi could be useful indicators of environmental

health. Our work helps scientists better understand how urbanization affects nature and offers a new way

to monitor and protect freshwater ecosystems.

KEYWORDS

Urbanization, Fungal communities, Aquatic environments, Amphibian symbionts.

INTRODUCTION

Urbanization alters biodiversity and ecosystem function, affecting both macroscopic and microscopic life. Freshwater biodiversity is crucial for maintaining water quality and nutrient cycling, but urbanization degrades these systems by increasing pollution, habitat fragmentation, and impervious surface areas (ISA), which intensify runoff and pollutant loads (1–4). Additionally, urban discharge introduces novel microbial taxa, altering community composition and potentially reshaping ecosystem functions (5). ISAs, such as buildings and pavement, contribute to environmental degradation by intensifying heat island effects and increasing runoff elevating sediment loads (6). Stormwater and wastewater further introduce excess nutrients, heavy metals, and pharmaceuticals, degrading water quality (5, 7).

Urbanization negatively impacts waterway-associated organisms, including amphibians, which serve as environmental indicators due to their sensitivity to habitat changes. Habitat loss is a major driver of global amphibian declines (8, 9), while pollution and habitat modification also disrupt microbial communities, including those associated with amphibians (5, 10). As amphibians persist in varying degrees of degraded environments, shifts in their microbiomes serve as signals of ecosystem health before extreme habitat loss occurs. While the effects of urbanization on amphibians have been documented (11, 12), urban-driven changes also extend to microbial communities, which is comparatively less well studied. Despite the generally cryptic nature of the members of a microbial community, their composition is essential to overall ecosystem function.

Microorganisms play essential roles in nutrient and energy cycling and exhibit rapid growth, making them highly sensitive to environmental changes (10, 13, 14). As a result, microbial taxa can serve as potential biological indicators of environmental and host health. Microbiomes are the community of microbes comprised by viruses, bacteria, archaea and microscopic eukaryotes. The microbiomes of gastrointestinal tracts are particularly critical for host physiology, immune function, and behavior, influencing the response to environmental shifts (14).

Despite their ecological importance, fungi have been historically overlooked in microbiome

studies, even though they play key roles in host metabolism, carbohydrate degradation, pathogen

resistance, and immune regulation (14–16). Understanding fungal communities (the "mycobiome") is

essential, as host-fungal interactions can support amphibian health or contribute to population declines,

especially in urban environments where microbial diversity is reshaped by environmental pressures (14).

Microbiome research provides valuable insights into amphibian conservation by revealing how

urbanization affects microbial communities, host health, and ecosystem stability [12–14]. While most

urbanization studies on microbial communities have focused on bacteria (2, 6, 17, 18), recent research

highlights fungi and the mycobiome as potential environmental health indicators(19–21). However, prior

microbiome studies typically examine urban impacts on only a single habitat type, such as water,

sediment, or host-associated communities (22–24).

Here, we studied the shifting biodiversity and ecology of the mycobiome along an urbanization

gradient in Worcester, MA, across water, sediment, and frog-associated microbiomes. Worcester is the

second most populous city in New England (USA) and has a long history as a manufacturing hub, with

high levels of human activity and pollution from goods production factories. Across the city, there are

numerous points of contacts with these historical production centers that directly connect with many

waterways that converge into the Blackstone river (25). The northernmost parts of Worcester have lower

population density and larger areas of protected land, as well as freshwater reservoirs. The Tatnuck Brook

watershed presents a gradient of rural, more protected areas in the north to increasingly developed areas

to the south, making it ideal to study the impact of urbanization on freshwater ecosystems.

We tested the hypothesis that increasing urbanization alters fungal community structure in the

waterway, leading to shifts in diversity across habitats. Specifically, we predicted that urbanization would

drive changes in community composition, with certain fungal taxa increasing in abundance in highly

urbanized sites while others decline, reflecting environmental filtering along the urbanization gradient.

We focused on accomplishing two main objectives: 1) To elucidate the differences of the fungal

microbial communities in freshwater ecosystems across the microbiomes of sediment, water and frog

gastrointestinal tracts, and 2) to characterize potential changes in taxonomic and functional fungal

microbial diversity of these communities associated with a rural to urban gradient.

METHODS

Study Site Selection

We collected samples from three main habitats: fecal pellets from three frog species, sediment cores, and water samples across the Tatnuck Brook watershed system (Figure 1). Ten sites spanning approximately 10 km from north to south were sampled. Sites were a mix of ponds and streams that varied strongly in their exposure to human activity. Percent impervious surface area (PISA) was calculated for each site to determine urbanization levels using QGIS V 3.30.1 (26). The impervious surface area (ISA) layer, created by data collected in 2005, was extracted from publicly available MassGIS data (https://www.mass.gov/info-details/massgis-data-impervious-surface-2005). Urbanization levels were classified by the PISA value for each site, characterized as low (<10%), medium (10–30%), and high (>30%) urbanization categories as determined by Hincher et al.(27).

Three frog species were collected: *Lithobates catesbeianus* (American bullfrog)*, Lithobates clamitans* (Green frog)*,* and *Lithobates palustris* (Pickerel frog). These are common local frog species and of no conservation concern and thus were selected for non-destructive capture and fecal pellet isolation. All amphibians were captured at sites authorized by the Massachusetts Division of Fisheries and Wildlife (Permit 16.23SCRA). The capture-release protocol was approved by Clark University's Institutional Animal Care and Use Committee (IACUC, Protocol 037R). Frogs were temporarily housed in sterile collection bags until defecation occurred, after which they were released at their original capture site. Fecal samples were then collected using a sterile scoopula. The fecal sample was covered with 20% glycerol and stored at -20 °C until DNA extraction.

Sediment core samples were taken at stream sites using a Large Universal Corer (Aquatic Research Instruments, Idaho, USA) to a depth of ~5 cm at four locations randomly chosen at each site and kept separate. After sampling, each core was homogenized and ~50 ml duplicate subsamples were stored separately of each core at -80 °C until DNA extraction.

Water samples were collected using acid-washed 4 L cubitainers, rinsed using water downstream

of the collection point. Water samples were placed on ice until filtered later that day. Water samples were

filtered by peristaltic pump by drawing water through a 25 µm Nitex mesh covering the intake opening of

the peristaltic tubing and then passing through a 47 mm 3 µm polycarbonate filter (Cat # PCT3047100,

Sterlitech, USA) within a stainless-steel filter holder to collect eukaryotic cells. The 3 µm filters were

then aseptically removed and stored in tubes kept at -80 °C until DNA extraction.

Environmental DNA Extraction

Fecal DNA was extracted from frog fecal samples and sediment samples using the QiAmp

PowerSoil Pro DNA Kit (Cat # 47021, Qiagen, Maryland, USA). For water DNA extraction, a phenol

chloroform protocol was used to extract DNA from filters. Modifications to the protocols can be found in

the Supplementary Materials. DNA concentrations of all samples were determined using a Qubit 4

Fluorometer with a dsDNA High Sensitivity kit (ThermoFisher Scientific, Massachusetts, USA). The

purity was measured with a Nanodrop One Microvolume UV-Vis Spectrophotometer (ThermoFisher

Scientific, Massachusetts, USA).

ITS1 Metabarcode sequencing and analysis

The ITS1 region was amplified using the standardized primer set in Bellemain et al. (2010) to

conduct metabarcode community analysis of fungal communities. Amplification, library preparation, and

2X250 PE sequencing was done by NovoGene Inc. (Sacramento, CA) using on an Illumuna

NovaSeq6000 sequencer. Illumina adapter sequences were removed, and sequences were demultiplexed

by NovoGene. The resulting raw reads were quality trimmed with cutadapt (v3.4) (28). Downstream

processing of trimmed reads was performed using DADA2 (v1.16) (29), and taxonomy was assigned to

resulting amplicon sequence variants (ASVs) using the general Fungi UNITE database (version 9.0) with

a naïve Bayesian classifier and a minimum bootstrap confidence threshold of 80% (30).

Fungal Community Analysis

All data analyses were conducted in R (4.4.2) (31) and RStudio (32). Phyloseq (V 1.44.0) (33)

was used for microbiome data processing such as deduplicating reads, defining ASVs and generating

counts per ASV. Relative abundances were calculated for downstream analysis. Rarefaction curves were

generated with the function rarecurve from vegan (v. 2.6) (34). Reads were not rarified for the subsequent

analyzes as this type of data stratification tends to remove rare, but important taxa (35).

For alpha-diversity analyzes, Shannon and Simpson diversity indices were computed with the

diversity function in vegan (34).   Analysis of Variance (ANOVA) was performed to analyze differences

in alpha diversity, using Shannon or Simpson's indices as response variable, and habitat and urbanization

levels as the explanatory variable. Assumptions for ANOVA were checked using Q-Q plots for normality

and Bartlett's test for homogeneity of variance. Post hoc analyses for significant ANOVA results were

conducted using Tukey's HSD tests. Boxplots were generated using ggplot2 (36).

Beta diversity was compared using permutational multivariate ANOVA (PERMANOVA) using

the adonis2 function in the vegan package (37) with 1,000 permutations. Nonmetric multidimensional

scaling ordination (NMDS) plots were generated using Bray-Curtis dissimilarity values. For the

PERMANOVA, the Bray-Curtis dissimilarity values were used as the response variables, and the habitat

or urbanization levels as the categorical explanatory variables. For the frog fecal samples, we also

included the frog host species and the interaction of urbanization level and host species as the categorical

variables. Post hoc pairwise PERMANOVA analyses were conducted to determine which specific

variables exhibited significant community differences from each other using the pairwiseAdonis package

(v. 0.4.1) (38) with Benjamini-Hochberg (BH) p-value correction for multiple test comparisons (39). The

top 20 most abundant taxa per habitat were ranked by mean relative abundance.

Differential abundance analyses were conducted using DESeq2 (40) to identify taxa with

significant changes in absolute abundance per source across urbanization levels. Variance-stabilizing

transformation (VST) was applied to normalize count data. Differentially abundant taxa between

urbanization categories were identified using the DESeq function, with a significance threshold of

adjusted p-value < 0.05. Log2 fold changes were computed, where an ASV with log2FC $\geq$ 5 or $\leq$ -5

indicates a significant presence in one condition compared to the other.

Fungal trophic guild abundances

Fungal ecological guild information was extracted using FUNGuild (v. 1.1) (41) and its

associated database, with ASVs without annotation being removed from further analysis. After

assignment of guild to each ASV, analysis targeted ASVs assigned to only one trophic guild with a

confidence ranking of 'probable' or 'highly probable.' Significant differences in the absolute abundance

of each trophic mode were analyzed using a Kruskal-Wallis test in R, after assumptions for ANOVA were

checked and violated (Q-Q plots for normality and Bartlett's test for homogeneity of variance) as well as

a permutation ANOVA using the aovp function of the lmPerm R package using 5,000 permutations (42)

and pairwise permutation tests in the pairwisePermutationTest function from the rcompanion R package

using 10,000 permutations and the false discovery rate (FDR) correction (43). For each guild (Saprotroph,

pathogen and symbiont), the abundance was used as the response variable and the urbanization category

as the explanatory variable) for both Kruskal-Wallis test and permutation ANOVA.

RESULTS

<u>**Sampling and Meta-amplicon Sequencing**</u>

Water (planktonic), stream sediment, and frog fecal samples were collected from the Tatnuck

Brook waterway, spanning a north-to-south urbanization gradient in Worcester (Fig. 1 (Table 1). The

northernmost sites—Kinney Woods (5.1%) and Tatnuck Brook (9.0%)—were classified as low

urbanization. The next four sites had medium urbanization (17.2–25.6%), while the five most southern,

downstream sites exceeded 45% PISA, with Holy Cross (56.7%) and Beaver Brook (56.6%) showing the

highest urbanization levels (Table 1).. This 10 km gradient reflects a transition from rural to developed

regions.

A total of 129 microbiome samples were collected, with 112 yielding high-quality sequencing

data (fecal n=60, sediment n=30, water n=22) (Supp. Table 1). These samples generated 25,608,648

paired-end raw reads, resulting in 121,368 unique amplicon sequence variants (ASVs) spanning 20 phyla,

80 classes, 214 orders, 596 families, 1,740 genera, and 2,879 species (Table 2). Rarefaction curves

plateaued, confirming sufficient sequencing depth to capture taxonomic diversity (Supplementary Figure

1).

Alpha and Beta diversity of Fungal Communities Across Freshwater Habitats

Alpha and beta diversity analyses revealed significant differences in fungal communities across habitats

and urbanization levels. Using Shannon's diversity index, sediment samples were significantly more

diverse than fecal and water samples (ANOVA, $F = 37.95$, $p<0.001$; Tukey's post hoc, $p<0.001$), while

fecal and water samples did not differ (Tukey's post hoc, $p = 0.132$) (Figure 2). In contrast, there were no

significant differences across habitats for Simpson's diversity index (ANOVA, $F = 1.39$, $p=0.253$),

suggesting similar dominance patterns among sources (Supplementary Figure 2a). Beta diversity analysis

(PERMANOVA) confirmed distinct microbial communities across habitats ($F = 5.79$, $p = 0.001$, $R^2 =$

0.095$), with pairwise comparisons showing significant differences between all groups ($p = 0.001$).

Urbanization significantly influenced fungal community composition ($F = 3.32$, $p = 0.001$, $R^2 = 0.050$),

though habitat type explained more variance, indicating a stronger role in shaping community structure.

Fecal samples were dominated by *Basidiobolus* and *Cladosporium*, while sediments were dominated by

Lemonniera (Supplementary Figure 3). *Only Candida* was consistently found in all habitats

(Supplementary Table 2 and Supplementary Figure 3).

Effects of Urbanization on Fungal Communities

Fecal Fungal Communities

To control for the effect of habitat, samples were analyzed separately by habitat type to further test for relationships between community diversity and urbanization. In frog fecal samples, urbanization did not significantly impact alpha diversity using either Shannon's diversity ($F = 1.804$, $p = 0.174$) or Simpson's diversity ($F = 0.01$, $p = 0.994$) . There was also no effect of frog species (Shannon: $F = 0.502$, $p = 0.608$, Simpson: $F = 0.546$, $p = 0.608$) or the interaction of frog species and PISA categories (Shannon: $F = 0.507$, $p = 0.605$, Simpson: $F = 0.585$, $p = 0.560$).

However, although there were no differences in alpha diversity, there were significant differences in fungal community composition between frog species (F=1.29, p=0.020, R^2=0.038) (Supplementary Figure 2b) and urbanization levels (F=3.88, p=0.001, R^2=0.116) (Figure 3a), with the latter having a stronger effect. The interaction between host species and urbanization was not significant (F=1.091, p=0.169, R^2=0.032). Pairwise PERMANOVA tests confirmed significant community composition differences across urbanization levels (p=0.001 for all comparisons). Notably, *Basidiobolus* and *Candida* were abundant in low urbanization, while *Botrytis*, *Alternaria*, and *Parasarocladium* were more common in highly urbanized sites (Supplementary Table 2). Differential abundance analysis identified 125 taxa showing significant differences between low and high urbanization, with 82 favoring low and 43 favoring high urbanization (Table 3, Supplementary File 1).

Sediment Fungal Communities

Sediment samples exhibited no significant difference in Shannon's ($F = 2.16$, $p = 0.132$) or Simpson's ($F = 1.60$, $p = 0.217$) diversity indices across urbanization levels. However, beta diversity was significantly influenced by urbanization (PERMANOVA, $F = 2.32$, $p = 0.001$), in this case explaining 13% of the observed variation (Figure 3b). Post hoc pairwise PERMANOVA confirmed significant differences across all urbanization levels as well (p=0.001 for all comparisons). Low urbanization sites had unique genera like *Apodus* and *Podospora*, while high urbanization sites were characterized by *Neoascochyta* and *Candida* (Supplementary Table 2). Out of 33 taxa that were differentially abundant

between low and high urbanization, ten and twenty-three were more common at low and high

urbanization, respectively (Table 3, Supplementary File 2)

Planktonic Fungal Communities

As with the other habitats, planktonic fungal communities showed no significant change in

Shannon's ($F = 0.24$, $p = 0.787$) or Simpson's diversity indices ($F = 0.32$, $p = 0.729$) with urbanization.

Nevertheless, community composition was significantly structured by urbanization (PERMANOVA, $F =$

1.33, $p = 0.003$, $R^2 = 0.159$) (Figure 3c). 16% of sample variance was explained by urbanization, again

notably lower than for frog fecal mycobiomes. There was overlap in the clustering of medium and high

urbanization samples, but PERMANOVA shows a significant difference in community composition

between these groups ($F = 1.528$, $p = 0.006$, $R^2 = 0.105$). There were only two low-urbanization water

samples, so we were unable to test for significant differences in community composition of these sites

from other urbanization sites.

The top 20 genera analysis showed that low and medium urbanization sites had greater

abundances of *Basidiobolus*, *Clavispora*, and *Zygophlyctis*, while high urbanization sites featured

Aureobasidium and *Metarhizium* (Supplementary Table 2). We identified 95 differentially abundant taxa

between low and high urbanization sites. Chytridiomycota ASV's represented the most differentially

abundant taxa, with 45 ASV's represented in low urbanization, in comparison to 13 at higher urbanization

(Table 3, Supplementary File 3).

Fungal Trophic Mode Analysis

A total of 35,854 fungal ASVs were classified into pathotrophs, saprotrophs, symbiotrophs, and

unknowns, with 58% of the ASVs labeled as unknown due to multiple assignments or low confidence

rankings (Figure 4). While most Kruskal-Wallis and permutation-based ANOVA tests did not detect

significant differences in trophic mode abundance across urbanization levels (Supplementary Tables 3

and 4), the saprotroph community in water samples showed a significant effect of urbanization in the

global permutation ANOVA ($p = 0.036$). However, post hoc pairwise permutation tests did not reveal

significant differences between specific urbanization categories after false discovery rate (FDR)

correction, despite a marginal trend between high and medium sites and low and medium size (adjusted p

= 0.16 for both paiwise comparisons). These findings suggest that while saprotrophic communities in

water are influenced by urbanization at a broad level, the directional differences between individual

urbanization categories are subtle and may reflect gradual community shifts rather than discrete stepwise

transitions. Such shifts may still carry ecological implications for nutrient cycling and organic matter

processing in urban-impacted aquatic systems.

DISCUSSION

Urbanization is reshaping ecosystems, altering both macroscopic biodiversity and microbial communities. While urban impacts on bacterial and archaeal communities have been studied, fungal communities remain underexplored. Primary, previous research has often focused on a single habitat, be it environmental (44–46) or host associated (47–50) and less frequently in aquatic environments. This study examines three interconnected microbial environments, including both environmental and host derived sources. Our results show that while alpha diversity remained stable, beta diversity shifted significantly across urbanization level, indicating taxonomic turnover rather than biodiversity loss. This aligns with previous studies showing that urbanization reshapes microbial composition without reducing species richness (51–53).

There were some notable differences in strength of correlation of urbanization with community composition. Urbanization influenced fungal community composition across habitats, but habitat type was a stronger predictor (9.5% vs. 6%) when considering all samples together. This is not surprising because habitat type is a strong driver of prokaryotic community similarity (54). When considering each habitat separately, urbanization explained similar proportions of variance in community composition (12%, 13%, and 16% for fecal, sediment, and water habitats respectively). Within fecal samples, host species identity explained only 4% of variance, suggesting that urbanization exerts a stronger influence on microbial turnover than interspecies differences. These findings indicate urbanization has broad-sweeping impacts of altering fungal communities across various components of the aquatic ecosystem.

The observed shifts in fungal community composition suggest that urban environments act as ecological filters, favoring taxa adapted to pollutants, altered nutrient dynamics, and habitat disturbances (22–24, 47, 55, 56). However, the specific mechanisms driving these shifts remain unclear. Future metagenomic studies could identify key genes and pathways involved. Additionally, the implications for amphibian health and water quality require further investigation. By analyzing fungal taxa across freshwater habitats, our findings highlight how urbanization shapes microbial diversity and underscore

the need for research into the functional roles of urban-adapted fungi in host-microbe interactions,

biogeochemical cycling, and ecosystem stability.

Differences in Fungal Community Responses Across Sources

The three primary habitats—sediment, water, and host associated fecal matter—demonstrate

distinct community compositions influenced by urbanization, reflecting the diverse ecological niches

within the watershed. Sediment, with its high fungal diversity, supports saprotrophic fungi crucial for

organic matter breakdown, likely due to nutrient-rich microenvironments that favor fungal growth and

metabolic diversity [40, 41, 42]. The species richness in sediment may also stem from its role as a

microbial sink, capturing inputs from water and detritus, and supporting taxa that interact with organic

substrates.

The planktonic communities exhibited fewer taxa but showed notable resilience to urbanization-

induced changes, particularly in community composition. The persistence of certain genera, like

Clavispora and *Aureobasidium*, across urbanization levels in water suggests their functional adaptation to

the aquatic environment, highlighting their ecological relevance as nutrient recyclers and decomposers in

freshwater systems (60). The resilience observed may be attributed to the natural flow dynamics in water

bodies, which might buffer against certain urban pollutants (61, 62).

Frog fecal samples showed the greatest compositional shifts with urbanization. *Basidiobolus*, a

genus linked to amphibian gut health, was abundant in low urbanization sites but absent in high

urbanization samples, suggesting sensitivity to urban stressors. Previous studies found *Basidiobolus* to be

a dominant gut fungus, representing 84% of gut fungi across various salamander species, 29% of gut

fungi in various species of frogs (63), and comprising 60% of gut fungi in slimy salamanders (*Plethodon*

spp.) (64). Despite its prevalence in amphibians, in this study it was abundant only in low urbanization

sites. This suggests *Basidiobolus* may be a useful bioindicator for habitat quality and amphibian-

microbiome interactions.

Novel discoveries in fungal ecology

The *Rozellomycota* were highly represented across all sample types and mostly in fecal and

sediment samples. They are typically cryptic or parasitic, so they are often linked to other fungi or protists

in soil. The presence of *Rozellomycota* in both amphibian fecal pellets and sediment expands their known

habitat range. Their detection in aquatic environments suggests potential roles in organic matter

decomposition, symbiosis, or host-microbe interactions that warrant further study.

Additionally, this study revealed the presence of *Basidiobolus* within the planktonic water

communities, marking one of the first records of this genus in freshwater systems. This genus is typically

associated with amphibian gastrointestinal tracts or terrestrial environments such as leaf litter, so its

presence in water possibly suggests environmental transmission pathways that extend beyond known host

associations and a broader ecological versatility (65–67). These findings suggest fungal adaptability in

urban-impacted freshwater systems, with urbanization potentially facilitating the movement of terrestrial

taxa into aquatic and host-associated environments. The presence of *Rozellomycota* in novel habitats

highlights the need for further studies using multi-marker approaches to clarify their ecological roles,

diversity, and contributions to nutrient cycling and ecosystem function.

Implications of Trophic Mode Shifts in Urbanized Environments

Trophic mode analysis provides insights into how urbanization may influence the ecological roles

of fungal communities. While no significant differences in mode were found with urbanization level, this

apparent stability may reflect ecological resilience rather than a lack of response. The consistency in

saprotrophs, pathotrophs, and symbiotrophs suggests that key ecosystem functions are maintained despite

shifts in community composition. This points to possible functional redundancy within fungal

communities, allowing them to sustain roles like decomposition and symbiosis even under urban stress.

Other trends, such as saprotrophic fungi in water samples exhibited a significant effect of urbanization in

the global permutation ANOVA (p = 0.036), although follow-up pairwise permutation tests did not yield

statistically significant differences between specific urbanization categories after FDR correction. This

pattern suggests a broad but gradual shift in saprotrophic composition in response to increasing urban

impact, potentially linked to altered nutrient regimes or detrital inputs in urban waterways. In parallel, we

observed non-significant but consistent trends of increasing pathotroph abundance in sediment and

planktonic habitats with higher urbanization, echoing patterns reported in other studies (68, 69). This shift

could impact amphibians and aquatic ecosystems by increasing pathogen loads, contributing to

biodiversity loss. Increased urban organic waste may support more opportunistic fungi, while reduced

plant diversity often seen with urbanization likely limits symbiotic fungi. The heightened pathotroph

abundance in urbanized sediments and water raises concerns about potential amphibian health risks, as

pathogenic fungi, such as chytrids, are known to have devastating effects on amphibian populations (70).

However, the high proportion of unclassified ASVs underscores the need for further metagenomic

research to clarify functional shifts in fungal communities along urban gradients.

Urbanization as a Filter for Resilient or Opportunistic Fungal Taxa

The observed shifts in fungal communities align with the concept of biotic homogenization,

where urbanization filters out sensitive taxa and selects for those resilient to anthropogenic pressures.

This effect, where resilient taxa dominate urban ecosystems, highlights a trend toward microbial

communities dominated by opportunistic or anthropophilic species (71–73). Such homogenization can

reduce functional diversity within ecosystems, potentially diminishing the resilience of microbial

communities to future environmental changes (71–73).

Conservation and Restoration Implications

The distinct fungal community compositions across urbanization levels and sources in this study

highlight the need for conservation and restoration efforts tailored to urbanized freshwater systems. The

high fungal diversity observed in sediment, especially in low urbanization sites, suggests that sediment

habitats act as refuges for microbial biodiversity. Preserving and restoring sediment-rich areas along

urban watersheds could help maintain fungal diversity and associated ecosystem functions, such as

nutrient cycling and organic matter decomposition.

These results also imply that some fungal taxa, particularly those in frog GI tracts, may be more

sensitive to urban stressors and can be bioindicators for assessing environmental health in freshwater

systems (74, 75). For instance, the absence of *Basidiobolus* in high urbanization fecal samples suggests it

could serve as a sensitive indicator of habitat quality. Conservation strategies could include monitoring

this genus across watersheds to identify areas of significant environmental degradation and inform

mitigation actions.

Conclusions

This study serves as an initial step toward understanding the effects of urbanization on fungal

communities in freshwater ecosystems, and the interconnected nature they form across different trophic

levels. Here we find consistent patterns of impact across three inter-connected microbial habitats in a

freshwater ecosystem: stable alpha diversity but significant shifts in beta diversity associated with

urbanization. Future research should expand upon these findings by integrating shotgun metagenomics to

determine specific metabolic pathways affected by urbanization. Additionally, studies tracking temporal

fungal communities could inform the responses of these communities to increasing urban development.

This interconnectivity of freshwater systems renders it critical to study environmental microbial

communities in tandem with host microbial communities since they serve as source pools for host

microbiomes. Therefore, sediment and planktonic microbial communities in freshwater systems are also

of interest as they not only perform vital ecosystem services but also contribute to the assembly of

microbial communities in amphibians.

Acknowledgements

Special thanks to Dr. Anne Devan Song for her field expertise and help with our collection of amphibians.

This project was funded by the Clark University's Academic Innovation Fund.

Conflicts of Interest

The authors report no conflicts of interest

Data availability

All raw sequencing data has been deposited in the Short Read Archive (SRA) under Bioproject number

PRJNA1244802. All supplementary data, including all supplementary files, tables, figures, R objects and

code can be downloaded from GitHub at

(https://github.com/TabimaLab/Fungal_Watershed_Mycobiome).

REFERENCES

1. Uttara S, Bhuvandas N, Aggarwal V. 2012. IMPACTS OF URBANIZATION ON ENVIRONMENT. Applied Sciences 2.

2. Hosen JD, Febria CM, Crump BC, Palmer MA. 2017. Watershed Urbanization Linked to Differences in Stream Bacterial Community Composition. Front Microbiol 8:1452.

3. Abrego N, Crosier B, Somervuo P, Ivanova N, Abrahamyan A, Abdi A, Hämäläinen K, Junninen K, Maunula M, Purhonen J, Ovaskainen O. 2020. Fungal communities decline with urbanization— more in air than in soil. ISME J 14:2806–2815.

4. Chen Y, Martinez A, Cleavenger S, Rudolph J, Barberán A. 2021. Changes in Soil Microbial Communities across an Urbanization Gradient: A Local-Scale Temporal Study in the Arid Southwestern USA. Microorganisms 9:1470.

5. Roberto AA, Van Gray JB, Leff LG. 2018. Sediment bacteria in an urban stream: Spatiotemporal patterns in community composition. Water Research 134:353–369.

6. Simonin M, Voss KA, Hassett BA, Rocca JD, Wang S, Bier RL, Violin CR, Wright JP, Bernhardt ES. 2019. In search of microbial indicator taxa: shifts in stream bacterial communities along an urbanization gradient. Environmental Microbiology 21:3653–3668.

7. Paul MJ, Meyer JL. 2001. Streams in the Urban Landscape. Annual review of Ecology and Systematics 32:333–365.

8. Scheffers BR, Paszkowski CA. 2012. The effects of urbanization on North American amphibian species: Identifying new directions for urban conservation. Urban Ecosyst 15:133–147.

9. Riley SPD, Busteed GT, Kats LB, Vandergon TL, Lee LFS, Dagit RG, Kerby JL, Fisher RN, Sauvajot RM. 2005. Effects of Urbanization on the Distribution and Abundance of Amphibians and Invasive Species in Southern California Streams. Conservation Biology 19:1894–1907.

10. Astudillo-García C, Hermans SM, Stevenson B, Buckley HL, Lear G. 2019. Microbial assemblages and bioindicators as proxies for ecosystem health status: potential and limitations. Appl Microbiol Biotechnol 103:6407–6421.

430   11.   Effects of Urbanization on Occupancy of Stream Salamanders.

431   12.   Yang L, Zhao S, Liu S. 2022. A global analysis of urbanization effects on amphibian richness:
432   Patterns and drivers. Global Environmental Change 73:102476.

433   13.   Karimi B, Maron PA, Chemidlin-Prevost Boure N, Bernard N, Gilbert D, Ranjard L. 2017.
434   Microbial diversity and ecological networks as indicators of environmental quality. Environ Chem
435   Lett 15:265–281.

436   14.   Simonin M, Voss KA, Hassett BA, Rocca JD, Wang S-Y, Bier RL, Violin CR, Wright JP, Bernhardt
437   ES. 2019. In search of microbial indicator taxa: shifts in stream bacterial communities along an
438   urbanization gradient. Environmental Microbiology 21:3653–3668.

439   15.   Kearns PJ, Fischer S, Fernández-Beaskoetxea S, Gabor CR, Bosch J, Bowen JL, Tlusty MF,
440   Woodhams DC. 2017. Fight Fungi with Fungi: Antifungal Properties of the Amphibian Mycobiome.
441   Front Microbiol 8.

442   16.   Wagg C, Schlaeppi K, Banerjee S, Kuramae EE, Van Der Heijden MGA. 2019. Fungal-bacterial
443   diversity and microbiome complexity predict ecosystem functioning. Nat Commun 10:4841.

444   17.   Wang S-Y, Sudduth EB, Wallenstein MD, Wright JP, Bernhardt ES. 2011. Watershed Urbanization
445   Alters the Composition and Function of Stream Bacterial Communities. PLoS ONE 6:e22972.

446   18.   Belt KT, Hohn C, Gbakima A, Higgins JA. 2007. Identification of culturable stream water bacteria
447   from urban, agricultural, and forested watersheds using 16S rRNA gene sequencing. Journal of
448   Water and Health 5:395–406.

449   19.   Kearns PJ, Fischer S, Fernández-Beaskoetxea S, Gabor CR, Bosch J, Bowen JL, Tlusty MF,
450   Woodhams DC. 2017. Fight Fungi with Fungi: Antifungal Properties of the Amphibian Mycobiome.
451   Front Microbiol 8:2494.

452   20.   Sam Q, Chang M, Chai L. 2017. The Fungal Mycobiome and Its Interaction with Gut Bacteria in
453   the Host. IJMS 18:330.

454   21.   Richard ML, Sokol H. 2019. The gut mycobiota: insights into analysis, environmental interactions
455   and role in gastrointestinal diseases. Nat Rev Gastroenterol Hepatol https://doi.org/10.1038/s41575-
456   019-0121-2.

457   22.   Numberger D, Zoccarato L, Woodhouse J, Ganzert L, Sauer S, Márquez JRG, Domisch S, Grossart
458   H-P, Greenwood AD. 2022. Urbanization promotes specific bacteria in freshwater microbiomes
459   including potential pathogens. Science of The Total Environment 845:157321.

460   23.   Li G, Cao L, Li X, Zhao H, Chang M, Lu D, Zhang L, Li M, Qin T, Jin H, Li X, Wang J, Li Y. 2023.
461   Effects of urbanization on fungal communities and their functions in the sediment of the Haihe
462   River. J Soils Sediments 23:4069–4080.

24. Mora J, Olson ,Matthew, Rocks ,Sara S., and Zahn G. 2025. Watershed urbanization alters aquatic
plant mycobiomes through the loss of rare taxa. Mycologia 117:235–245.

25. Sinha DR. 2010. The environmental Kuznets curve hypothesis and legacy pollution: a geohistorical
analysis of the environmental consequences of industrialization in Worcester, Massachusetts (USA).
The Industrial Geogrpaher 7:1–18.

26. QGIS Development Team. QGIS Geographic Information System. QGIS Association.
https://www.qgis.org.

27. Hincher MR, Carleton JP, Wheeler SJ, DelCogliano M, Mathis K, Tabima JF. Ubiquity and diversity
of Basidiobolus across amphibian species inhabiting an urbanization gradient. Mycologia 0:1–13.

28. Martin M. 2011. Cutadapt removes adapter sequences from high-throughput sequencing reads.
EMBnet j 17:10.

29. Callahan BJ, McMurdie PJ, Rosen MJ, Han AW, Johnson AJA, Holmes SP. 2016. DADA2: High-
resolution sample inference from Illumina amplicon data. Nat Methods 13:581–583.

30. Nilsson RH, Larsson K-H, Taylor AFS, Bengtsson-Palme J, Jeppesen TS, Schigel D, Kennedy P,
Picard K, Glöckner FO, Tedersoo L, Saar I, Kõljalg U, Abarenkov K. 2019. The UNITE database
for molecular identification of fungi: handling dark taxa and parallel taxonomic classifications.
Nucleic Acids Research 47:D259–D264.

31. R Core Team. 2023. R: A Language and Environment for Statistical Computing. R Foundation for
Statistical Computing, Vienna, Austria. https://www.R-project.org/.

32. Posit team. 2024. RStudio: Integrated Development Environment for R. Posit Software, PBC,
Boston, MA. http://www.posit.co/.

33. McMurdie PJ, Holmes S. 2013. phyloseq: An R Package for Reproducible Interactive Analysis and
Graphics of Microbiome Census Data. PLoS ONE 8:e61217.

34. Oksanen J, Simpson GL, Blanchet FG, Kindt R, Legendre P, Minchin PR, O'Hara RB, Solymos P,
Stevens MHH, Szoecs E, Wagner H, Barbour M, Bedward M, Bolker B, Borcard D, Carvalho G,
Chirico M, Caceres MD, Durand S, Evangelista HBA, FitzJohn R, Friendly M, Furneaux B,
Hannigan G, Hill MO, Lahti L, McGlinn D, Ouellette M-H, Cunha ER, Smith T, Stier A, Braak
CJFT, Weedon J, Borman T. 2025. vegan: Community Ecology Package (2.6-10).

35. McMurdie PJ, Holmes S. 2014. Waste Not, Want Not: Why Rarefying Microbiome Data Is
Inadmissible. PLOS Computational Biology 10:e1003531.

36. Wickham H ggplot2. 2011. WIREs Computational Stats 3. 180–185.

37.   Anderson MJ. 2017. Permutational Multivariate Analysis of Variance (PERMANOVA), p. 1–15. *In*
495   Balakrishnan, N, Colton, T, Everitt, B, Piegorsch, W, Ruggeri, F, Teugels, JL (eds.), Wiley StatsRef:
496   Statistics Reference Online, 1st ed. Wiley.

497   38.   Arbizu PM. 2017. pairwiseAdonis: Pairwise Multilevel Comparison using Adonis.
498   https://github.com/pmartinezarbizu/pairwiseAdonis.

499   39.   Benjamini Y, Hochberg Y. 1995. Controlling the False Discovery Rate: A Practical and Powerful
500   Approach to Multiple Testing. Journal of the Royal Statistical Society: Series B (Methodological)
501   57:289–300.

502   40.   Love MI, Huber W, Anders S. 2014. Moderated estimation of fold change and dispersion for RNA-
503   seq data with DESeq2. Genome Biol 15:550.

504   41.   Nguyen NH, Song Z, Bates ST, Branco S, Tedersoo L, Menke J, Schilling JS, Kennedy PG. 2016.
505   FUNGuild: An open annotation tool for parsing fungal community datasets by ecological guild.
506   Fungal Ecology 20:241–248.

507   42.   Wheeler B, Torchiano M. 2025. lmPerm: Permutation Tests for Linear Models.
508   https://github.com/mtorchiano/lmperm.

509   43.   Mangiafico SS. 2025. rcompanion: Functions to Support Extension Education Program Evaluation.
510   Rutgers Cooperative Extension, New Brunswick, New Jersey. https://CRAN.R-
511   project.org/package=rcompanion/.

512   44.   Ortiz-Vera MP, Olchanheski LR, da Silva EG, de Lima FR, Martinez LR del PR, Sato MIZ, Jaffé R,
513   Alves R, Ichiwaki S, Padilla G, Araújo WL. 2018. Influence of water quality on diversity and
514   composition of fungal communities in a tropical river. Sci Rep 8:14799.

515   45.   Fukasawa Y, Matsukura K. 2021. Decay stages of wood and associated fungal communities
516   characterise diversity–decomposition relationships. Sci Rep 11:8972.

517   46.   Tedersoo L, Bahram M, Põlme S, Kõljalg U, Yorou NS, Wijesundera R, Ruiz LV, Vasco-Palacios
518   AM, Thu PQ, Suija A, Smith ME, Sharp C, Saluveer E, Saitta A, Rosas M, Riit T, Ratkowsky D,
519   Pritsch K, Põldmaa K, Piepenbring M, Phosri C, Peterson M, Parts K, Pärtel K, Otsing E, Nouhra E,
520   Njouonkou AL, Nilsson RH, Morgado LN, Mayor J, May TW, Majuakim L, Lodge DJ, Lee SS,
521   Larsson K-H, Kohout P, Hosaka K, Hiiesalu I, Henkel TW, Harend H, Guo L, Greslebin A, Grelet
522   G, Geml J, Gates G, Dunstan W, Dunk C, Drenkhan R, Dearnaley J, De Kesel A, Dang T, Chen X,
523   Buegger F, Brearley FQ, Bonito G, Anslan S, Abell S, Abarenkov K. 2014. Global diversity and
524   geography of soil fungi. Science 346:1256688.

525   47.   García-Sánchez JC, Arredondo-Centeno J, Segovia-Ramírez MG, Tenorio Olvera AM, Parra-Olea
526   G, Vredenburg VT, Rovito SM. 2023. Factors Influencing Bacterial and Fungal Skin Communities
527   of Montane Salamanders of Central Mexico. Microb Ecol 86:670–686.

528   48.   Zhu L, Wu Q, Dai J, Zhang S, Wei F. 2011. Evidence of cellulose metabolism by the giant panda gut
529   microbiome. Proceedings of the National Academy of Sciences 108:17714–17719.

530   49.   Weinstein SB, Stephens WZ, Greenhalgh R, Round JL, Dearing MD. 2022. Wild herbivorous
531   mammals (genus Neotoma) host a diverse but transient assemblage of fungi. Symbiosis 87:45–58.

532   50.   Bradshaw AJ, Autumn KC, Rickart EA, Dentinger BTM. 2022. On the origin of feces: Fungal
533   diversity, distribution, and conservation implications from feces of small mammals. Environmental
534   DNA 4:608–626.

535   51.   Ortiz-Vera MP, Olchanheski LR, da Silva EG, de Lima FR, Martinez LR del PR, Sato MIZ, Jaffé R,
536   Alves R, Ichiwaki S, Padilla G, Araújo WL. 2018. Influence of water quality on diversity and
537   composition of fungal communities in a tropical river. Sci Rep 8:14799.

538   52.   Tai X, Li R, Zhang B, Yu H, Kong X, Bai Z, Deng Y, Jia L, Jin D. 2020. Pollution Gradients Altered
539   the Bacterial Community Composition and Stochastic Process of Rural Polluted Ponds.
540   Microorganisms 8:311.

541   53.   Mora J, Olson ,Matthew, Rocks ,Sara S., and Zahn G. 2025. Watershed urbanization alters aquatic
542   plant mycobiomes through the loss of rare taxa. Mycologia 117:235–245.

543   54.   Thompson LR, Sanders JG, McDonald D, Amir A, Ladau J, Locey KJ, Prill RJ, Tripathi A, Gibbons
544   SM, Ackermann G, Navas-Molina JA, Janssen S, Kopylova E, Vázquez-Baeza Y, González A,
545   Morton JT, Mirarab S, Zech Xu Z, Jiang L, Haroon MF, Kanbar J, Zhu Q, Jin Song S, Kosciolek T,
546   Bokulich NA, Lefler J, Brislawn CJ, Humphrey G, Owens SM, Hampton-Marcell J, Berg-Lyons D,
547   McKenzie V, Fierer N, Fuhrman JA, Clauset A, Stevens RL, Shade A, Pollard KS, Goodwin KD,
548   Jansson JK, Gilbert JA, Knight R, The Earth Microbiome Project Consortium, Rivera JLA, Al-
549   Moosawi L, Alverdy J, Amato KR, Andras J, Angenent LT, Antonopoulos DA, Apprill A, Armitage
550   D, Ballantine K, Bárta J, Baum JK, Berry A, Bhatnagar A, Bhatnagar M, Biddle JF, Bittner L,
551   Boldgiv B, Bottos E, Boyer DM, Braun J, Brazelton W, Brearley FQ, Campbell AH, Caporaso JG,
552   Cardona C, Carroll J, Cary SC, Casper BB, Charles TC, Chu H, Claar DC, Clark RG, Clayton JB,
553   Clemente JC, Cochran A, Coleman ML, Collins G, Colwell RR, Contreras M, Crary BB, Creer S,
554   Cristol DA, Crump BC, Cui D, Daly SE, Davalos L, Dawson RD, Defazio J, Delsuc F, Dionisi HM,
555   Dominguez-Bello MG, Dowell R, Dubinsky EA, Dunn PO, Ercolini D, Espinoza RE, Ezenwa V,
556   Fenner N, Findlay HS, Fleming ID, Fogliano V, Forsman A, Freeman C, Friedman ES, Galindo G,
557   Garcia L, Garcia-Amado MA, Garshelis D, Gasser RB, Gerdts G, Gibson MK, Gifford I, Gill RT,
558   Giray T, Gittel A, Golyshin P, Gong D, Grossart H-P, Guyton K, Haig S-J, Hale V, Hall RS, Hallam
559   SJ, Handley KM, Hasan NA, Haydon SR, Hickman JE, Hidalgo G, Hofmockel KS, Hooker J, Hulth
560   S, Hultman J, Hyde E, Ibáñez-Álamo JD, Jastrow JD, Jex AR, Johnson LS, Johnston ER, Joseph S,
561   Jurburg SD, Jurelevicius D, Karlsson A, Karlsson R, Kauppinen S, Kellogg CTE, Kennedy SJ,
562   Kerkhof LJ, King GM, Kling GW, Koehler AV, Krezalek M, Kueneman J, Lamendella R, Landon
563   EM, Lane-deGraaf K, LaRoche J, Larsen P, Laverock B, Lax S, Lentino M, Levin II, Liancourt P,
564   Liang W, Linz AM, Lipson DA, Liu Y, Lladser ME, Lozada M, Spirito CM, MacCormack WP,
565   MacRae-Crerar A, Magris M, Martín-Platero AM, Martín-Vivaldi M, Martínez LM, Martínez-
566   Bueno M, Marzinelli EM, Mason OU, Mayer GD, McDevitt-Irwin JM, McDonald JE, McGuire KL,

McMahon KD, McMinds R, Medina M, Mendelson JR, Metcalf JL, Meyer F, Michelangeli F, Miller K, Mills DA, Minich J, Mocali S, Moitinho-Silva L, Moore A, Morgan-Kiss RM, Munroe P, Myrold D, Neufeld JD, Ni Y, Nicol GW, Nielsen S, Nissimov JI, Niu K, Nolan MJ, Noyce K, O'Brien SL, Okamoto N, Orlando L, Castellano YO, Osuolale O, Oswald W, Parnell J, Peralta-Sánchez JM, Petraitis P, Pfister C, Pilon-Smits E, Piombino P, Pointing SB, Pollock FJ, Potter C, Prithiviraj B, Quince C, Rani A, Ranjan R, Rao S, Rees AP, Richardson M, Riebesell U, Robinson C, Rockne KJ, Rodriguezl SM, Rohwer F, Roundstone W, Safran RJ, Sangwan N, Sanz V, Schrenk M, Schrenzel MD, Scott NM, Seger RL, Seguin-Orlando A, Seldin L, Seyler LM, Shakhsheer B, Sheets GM, Shen C, Shi Y, Shin H, Shogan BD, Shutler D, Siegel J, Simmons S, Sjöling S, Smith DP, Soler JJ, Sperling M, Steinberg PD, Stephens B, Stevens MA, Taghavi S, Tai V, Tait K, Tan CL, Taș N, Taylor DL, Thomas T, Timling I, Turner BL, Urich T, Ursell LK, Van Der Lelie D, Van Treuren W, Van Zwieten L, Vargas-Robles D, Thurber RV, Vitaglione P, Walker DA, Walters WA, Wang S, Wang T, Weaver T, Webster NS, Wehrle B, Weisenhorn P, Weiss S, Werner JJ, West K, Whitehead A, Whitehead SR, Whittingham LA, Willerslev E, Williams AE, Wood SA, Woodhams DC, Yang Y, Zaneveld J, Zarraonaindia I, Zhang Q, Zhao H. 2017. A communal catalogue reveals Earth's multiscale microbial diversity. Nature 551:457–463.

55. Galitskaya P, Biktasheva L, Blagodatsky S, Selivanovskaya S. 2021. Response of bacterial and fungal communities to high petroleum pollution in different soils. Sci Rep 11:164.

56. Tai X, Li R, Zhang B, Yu H, Kong X, Bai Z, Deng Y, Jia L, Jin D. 2020. Pollution Gradients Altered the Bacterial Community Composition and Stochastic Process of Rural Polluted Ponds. Microorganisms 8:311.

57. Bridge P, Spooner B. 2002. Soil fungi: diversity and detection, p. 147–154. *In* Powlson, DS, Bateman, GL, Davies, KG, Gaunt, JL, Hirsch, PR (eds.), Interactions in the Root Environment: An Integrated Approach. Springer Netherlands, Dordrecht.

58. Wang Y, Liu J, Wang J, Gao G, Bartlam MG. 2015. Distribution and diversity of fungi in freshwater sediments on a river catchment scale. Front Microbiol 6.

59. Hobbie EA, Macko SA, Shugart HH. 1999. Insights into nitrogen and carbon dynamics of ectomycorrhizal and saprotrophic fungi from isotopic evidence. Oecologia 118:353.

60. Grossart H-P, Van Den Wyngaert S, Kagami M, Wurzbacher C, Cunliffe M, Rojas-Jimenez K. 2019. Fungi in aquatic ecosystems. Nat Rev Microbiol 17:339–354.

61. Chilton D, Hamilton DP, Nagelkerken I, Cook P, Hipsey MR, Reid R, Sheaves M, Waltham NJ, Brookes J. 2021. Environmental Flow Requirements of Estuaries: Providing Resilience to Current and Future Climate and Direct Anthropogenic Changes. Front Environ Sci 9.

62. Utz RM, Hopkins KG, Beesley L, Booth DB, Hawley RJ, Baker ME, Freeman MC, L. Jones K. 2016. Ecological resistance in urban streams: the role of natural and legacy attributes. Freshwater Science 35:380–397.

63. Vargas-Gastélum L, Romer AS, Ghotbi M, Dallas JW, Alexander NR, Moe KC, McPhail KL, Neuhaus GF, Shadmani L, Spatafora JW, Stajich JE, Tabima JF, Walker DM. 2024. Herptile gut

microbiomes: a natural system to study multi-kingdom interactions between filamentous fungi and bacteria. mSphere e00475-23.

64. Walker DM, Hill AJ, Albecker MA, McCoy MW, Grisnik M, Romer A, Grajal-Puche A, Camp C, Kelehear C, Wooten J, Rheubert J, Graham SP. 2020. Variation in the Slimy Salamander (Plethodon spp.) Skin and Gut-Microbial Assemblages Is Explained by Geographic Distance and Host Affinity. Microb Ecol 79:985–997.

65. Srinivasan MC, Thirumalachar MJ. 1967. Studies on Basidiobolus species from India with discussion on some of the characters used in the speciation of the genus. Mycopathologia et Mycologia Applicata 33:56–64.

66. Manning RJ, Waters SD, Callaghan AA. 2007. Saprotrophy of Conidiobolus and Basidiobolus in leaf litter. Mycological Research 111:1437–1449.

67. Nickerson MA, Hutchison JA. 1971. The Distribution of the Fungus Basidiobolus ranarum Eidam in Fish, Amphibians and Reptiles. American Midland Naturalist 86:500.

68. Pagani DM, Ventura SPR, Vu D, Mendes-Pereira T, Ribeiro Tomé LM, Carvalho DS de, Costa-Rezende DH, Kato RB, García GJY, Geml J, Robert V, They NH, Brenig B, Azevedo V, Scroferneker ML, Valente P, Góes-Neto A. 2023. Unveiling Fungal Community Structure along Different Levels of Anthropic Disturbance in a South American Subtropical Lagoon. 9. Journal of Fungi 9:890.

69. Jian Y, Lin J, Mu C, Wang Y, He Z, Chen G, Ding W. 2025. Short-Term Effects of Three Tree Species on Soil Physicochemical Properties and Microbial Communities During Land-Use Change from Farmland to Forests. 2. Forests 16:362.

70. Longcore JE, Pessier AP, Nichols DK. 1999. Batrachochytrium dendrobatidis gen. et sp. nov., a chytrid pathogenic to amphibians. Mycologia 91:219–227.

71. McKinney ML. 2006. Urbanization as a major cause of biotic homogenization. Biological Conservation 127:247–260.

72. Lokatis S, Jeschke JM. 2022. Urban biotic homogenization: Approaches and knowledge gaps. Ecological Applications 32:e2703.

73. Sidemo-Holm W, Ekroos J, Reina García S, Söderström B, Hedblom M. 2022. Urbanization causes biotic homogenization of woodland bird communities at multiple spatial scales. Global Change Biology 28:6152–6164.

74. Bai Y, Wang Q, Liao K, Jian Z, Zhao C, Qu J. 2018. Fungal Community as a Bioindicator to Reflect Anthropogenic Activities in a River Ecosystem. Front Microbiol 9:3152.

75. Warnasuriya SD, Udayanga D, Manamgoda DS, Biles C. 2023. Fungi as environmental bioindicators. Science of The Total Environment 892:164583.

640   **Fig 1.**

642   Map of the sampling sites along the Tatnuck Brook waterway in Worcester, Massachusetts. The

643   y-axis represents longitude, and the x-axis represents latitude. Circles depict the location

644   sampling sites, color-coded by urbanization level, and corresponding site names are listed in the

645   legend.

656
657 **Fig. 2**

Diversity and community composition of microfungi associated with amphibian fecal pellets,

sediment, and water samples. (a) Shannon diversity index of microfungal communities across

three habitats: fecal pellets (Represented by a Green Frog), sediment (Represented by a soil

sample in a tube), and water (Represented by a cubitainer). Letters above bars indicate statistical

similarity or differences, with shared letters representing no significant difference (e.g., "a"). (b)

Non-metric multidimensional scaling (NMDS) plot illustrating differences in microfungal

community composition among the three habitats. Each point represents a sample, and its

position reflects similarity in community composition. Axes represent the first two dimensions of

ordination space (NMDS1 and NMDS2). Ellipses represent 95% confidence intervals for the

multivariate t-distribution of the ordinated samples. All illustrations were generated in Adobe

Illustrator.

Fig. 3

Effects of urbanization on microfungal community composition in amphibian fecal pellets,

sediment, and water samples. NMDS plots show the ordination of microfungal communities with

respect to different levels of urbanization (High, Medium, Low) for samples segregated by

habitat type: a) frog fecal samples b) stream sediment samples, and c) planktonic water samples.

For each figure the axes represent the first two dimensions of ordination space (NMDS1 and

NMDS2). Each point represents a sample, which are color coded by urbanization level. Ellipses

represent 95% confidence intervals for the multivariate t-distribution of the ordinated samples.

Only two low-urbanization water samples were generated (c) so an ellipse could not be drawn

for that sample subset.

Fig. 4

Proportional representation of fungal trophic modes across fecal, sediment, and water samples

under varying urbanization levels. The bar plots illustrate the relative abundance of fungal

trophic modes (pathotrophs, saprotrophs, and symbiotrophs) and unknowns according to the

number of ASVs assigned to each in samples collected from fecal pellets, sediment, and water.

Urbanization levels are compared within each sample type, highlighting shifts in trophic mode

proportions in response to urbanization gradients.

TABLES

Table 1. Classification of sampling sites in the Tatnuck Brook watershed based on percent impervious

surface area (PISA).

Site Name	Site ID	Percent Standard Impervious Surface Area (PISA)	Urbanization Category
Kinney Woods	KP	5.1%	Low
Tatnuck Brook	PRS, CP2	9.0%	Low
Cook Pond	Cook	17.2%	Medium
Patch Reservoir	PR	22.0%	Medium
Mill Pond	MP	25.6%	Medium
Coes Pond	Coes, CR2, CR3	24.7%	Medium
Beaver Brook	BB1	56.6%	High
	BB2	48.1%	High
	BB3	52%	High
Cemetery	MR1	46.2%	High
Holy Cross	MB	56.7%	High
Middle River	MR, MR2	51.8%	High

Table 2. Number of fungal ASVs assigned to each phylum across fecal, water, and sediment habitats. The

table presents the number of ASVs associated with each fungal phylum (identified using the UNITE

database) across the three sample habitats. Notably, *Ascomycota* and *Basidiomycota* dominate all habitats,

while less common phyla such as *Rozellomycota*, *Chytridiomycota*, and *Mortierellomycota* show variable

representation. The presence of rare phyla, including *Neocallimastigomycota* and *Entomophthoromycota*,

highlights the diversity and complexity of fungal communities in these ecosystems

Phyla (UNITE)	Fecal	Water	Sediment
Aphelidiomycota	15	9	15
Ascomycota	6313	1532	6580
Basidiobolomycota	130	35	48
Basidiomycota	2087	364	3812
Blastocladiomycota	14	9	53
Chytridiomycota	944	943	3506
Entomophthoromycota	1	0	5
Glomeromycota	11	7	42
GS01	18	6	16
Kickxellomycota	12	0	11
Monoblepharomycota	8	16	24
Mortierellomycota	189	24	192
Mucoromycota	342	8	129
Neocallimastigomycota	0	0	5
Olpidiomycota	29	13	30
Rozellomycota	602	189	771
Zoopagomycota	9	6	8

706 **Table 3.** Fungal genera significantly enriched in low or high urbanization samples across different

707 habitats, based on DESeq2 analysis. In most cases, genera were detected in one level but not the other, but

708 otherwise the log2-fold difference is listed in parentheses after the genus name.

Habitat	More abundant in low or high urbanization samples	Genera
Fecal	High	*Mucor, Basidiobolus, Penicillium, Dichostereum, Dothiora, Apiotrichum, Aureobasidium, Taphrina, Coniella*
	Low	*Paraphaeosphaeria, Colletotrichum, Xenooacremonium, Leucosporidium, Pseudocercospora, Cadophora, Cytospora, Comoclathris, Cystofilobasidium, Acremonium*
Sediment	High	*Hymenoscyphus, Linnemannia, Saitozyma (4.7), Ganoderma (4.1)*
	Low	*Pseudorhypophila, Cirrenalia, Melanomma, Babjevia, Nectria, Conocybe, Pseudophialocephala, Stagonosporopsis, Helicodendron, Epicoccum*
Water	High	*Hebeloma, Zygophlyctis, Basidiobolus, Pseudeurotium, Rhizoclosmatium*
	Low	*Cyberlindnera, Betamyces, Fusicolla*

LEGEND

🟡 **Low Urbanization**
1. Kinney Woods
2. Tatnuck Brook

🔴 **Medium Urbanization**
3. Cook Pond
4. Patch Reservoir
5. Mill Pond
6. Coes Pond

🟣 **High Urbanization**
7. Beaver Brook
8. Cemetery
9. Holy Cross
10. Middle River

A. Fecal   **B. Sediment**   **C. Water**

Stress: 0.224

Stress: 0.204

Stress: 0.118

Urbanization — High — Medium — Low