Title: Impacts on the microbial fungal communities across water, sediment, and amphibian hosts

across an urbanization gradient waterway in Worcester, Massachusetts

Sara J. Wheeler[a], Manning DelCogliano[a], Brennan Hare[a], Alexander J. Bradshaw[a], Madison R. Hincher[a],

Jasper P. Carleton[a], Philip Bergmann[a], Nathan Ahlgren[a], Javier F. Tabima[a,1]

a. Biology Department, Clark University. Worcester, Massachusetts.

1. Corresponding author (jtabima@clarku.edu)

8
ABSTRACT:

In a world of increasingly urbanized environments, it is critical to understand the impact on microbial

communities, including fungal communities, as a measure of ecosystem health, and to document how

these environments are changing. Aquatic environments, in particular, can be highly sensitive to

urbanization with removal of local habitat and inputs of wastewater and contaminants, leading to a

displacement or extinction of natural flora, fauna, and funga. Aquatic microbial communities, especially

fungal communities, are extremely complex and far less characterized compared to their macro

counterparts. Here, we characterized the fungal microbiome, i.e. "mycobiome", across multiple habitats—

water, sediment, and amphibian fecal matter from three species (American bullfrog, Green frog and

Pickerel frog) — within the Tatnuck Brook waterway in Worcester, MA. The Tatnuck Brook waterway is

a connected group of streams, ponds, and lakes that transition from protected areas at its headwaters,

below the city's drinking water reservoir, to highly urbanized regions within Worcester. Using

metabarcode sequencing, we found that water, sediment, and host habitats harbor distinct fungal

communities, but all exhibit parallel shifts in diversity along an urbanization gradient. Furthermore, we

identified fungal taxa from the amphibian mycobiome and environmental samples that exhibited a range

of sensitivities to urbanization. These taxa, including *Basidiobolus*, *Cladosporium*, and *Lemonniera* in

fecal and sediment samples, as well as *Candida*, which was found across all habitats, are potential

indicators of shifts in aquatic ecosystems due to urbanization. Their responses to urban stressors may

serve as a baseline for further studies of aquatic fungal communities.

KEYWORDS

Urbanization, Fungal communities, Aquatic environments, Amphibian symbionts.

INTRODUCTION

Urbanization alters biodiversity and ecosystem function, affecting both macroscopic and microscopic life. Freshwater biodiversity is crucial for maintaining water quality and nutrient cycling, but urbanization degrades these systems by increasing pollution, habitat fragmentation, and impervious surface areas (ISA), which intensify runoff and pollutant loads (Uttara et al. 2012; Hosen et al. 2017; Abrego et al. 2020; Chen et al. 2021). Additionally, urban discharge introduces novel microbial taxa, altering community composition and potentially reshaping ecosystem functions (Roberto et al. 2018). ISAs, such as buildings and pavement, contribute to environmental degradation by intensifying heat island effects and increasing runoff elevating sediment loads (Simonin et al. 2019a). Stormwater and wastewater further introduce excess nutrients, heavy metals, and pharmaceuticals, degrading water quality (Paul and Meyer 2001; Roberto et al. 2018).

Urbanization negatively impacts waterway-associated organisms, including amphibians, which serve as environmental indicators due to their sensitivity to habitat changes. Habitat loss is a major driver of global amphibian declines (Riley et al. 2005; Scheffers and Paszkowski 2012), while pollution and habitat modification also disrupt microbial communities, including those associated with amphibians (Roberto et al. 2018; Astudillo-García et al. 2019). As amphibians persist in varying degrees of degraded environments, shifts in their microbiomes serve as signals for ecosystem health before extreme habitat loss occurs. While the effects of urbanization on amphibians have been documented (Roberto et al. 2018; Astudillo-García et al. 2019) urban-driven changes also extend to microbial communities, which is comparatively less well studied. Despite the generally cryptic nature of the members of a microbial community, their composition is essential to overall ecosystem function.

Microorganisms play essential roles in nutrient and energy cycling and exhibit rapid growth, making them highly sensitive to environmental changes (Karimi et al. 2017; Astudillo-García et al. 2019; Simonin et al. 2019b). As a result, microbial taxa can serve as potential as biological indicators of environmental and host health. Microbiomes are the community of microbes comprised by viruses, bacteria, archaea and microscopic eukaryotes. The microbiomes of gastrointestinal tracts are particularly

critical for host physiology, immune function, and behavior, influencing the response to environmental

shifts (Simonin et al. 2019b).

Despite their ecological importance, fungi have been historically overlooked in microbiome

studies, even though they play key roles in host metabolism, carbohydrate degradation, pathogen

resistance, and immune regulation (Kearns et al. 2017a; Simonin et al. 2019b; Wagg et al. 2019).

Understanding fungal communities (the "mycobiome") is essential, as host-fungal interactions can

support amphibian health or contribute to population declines, especially in urban environments where

microbial diversity is reshaped by environmental pressures (Simonin et al. 2019b). Microbiome research

provides valuable insights into amphibian conservation by revealing how urbanization affects microbial

communities, host health, and ecosystem stability [12–14]. While most urbanization studies on microbial

communities have focused on bacteria (Belt et al. 2007; Wang et al. 2011; Hosen et al. 2017; Simonin et

al. 2019a), recent research highlights fungi and the mycobiome as potential environmental health

indicators(Kearns et al. 2017b; Sam et al. 2017; Richard and Sokol 2019). However, prior microbiome

studies typically examine urban impacts on only a single habitat type, such as water, sediment, or host-

associated communities (Mora et al. 2025).

Here, we studied the shifting biodiversity and ecology of the mycobiome along an urbanization

gradient in Worcester, MA, across water, sediment, and frog-associated microbiomes. Worcester is the

second most populous city in New England (USA) and has a long history as a manufacturing hub, with

high levels of human activity and pollution by goods production factories. Across the city, there are

numerous points of contacts with this historical production centers that directly connect with many

waterways converge into the Blackstone river (Sinha 2010). The northernmost parts of Worcester have

lower population density and larger areas of protected land, as well as freshwater reservoirs. The Tatnuck

Brook watershed in Worcester presents a gradient of rural, more protected areas in the north to

increasingly developed areas to the south, making it ideal to study the impact of urbanization on

freshwater ecosystems.

We tested the hypothesis that increasing urbanization alters fungal community structure in the

waterway, leading to shifts in diversity across habitats. Specifically, we predicted that urbanization would

drive changes in community composition, with certain fungal taxa increasing in abundance in highly

urbanized sites while others decline, reflecting environmental filtering along the urbanization gradient.

We focused on accomplishing two main objectives: 1) To elucidate the differences of the fungal

microbial communities in freshwater ecosystems across the microbiomes of sediment, water and frog

gastrointestinal tracts, and 2) to characterize potential changes in taxonomic and functional fungal

microbial diversity of these communities associated with a rural to urban gradient.

METHODS

Study Site Selection

We collected samples from three main habitats: fecal pellets from three frog species, sediment cores, and water samples across the Tatnuck Brook watershed system (Figure 1). Ten sites spanning approximately 10 km from north to south were sampled. Sites were a mix of ponds and streams that varied strongly in their exposure to human activity. Percent impervious surface area (PISA) was calculated for each site to determine Urbanization levels using QGIS V 3.30.1 (QGIS Development Team). The impervious surface area (ISA) layer, created by data collected in 2005, was extracted from publicly available MassGIS data (https://www.mass.gov/info-details/massgis-data-impervious-surface-2005). Urbanization levels were classified by the PISA value for each site, characterized as low (<10%), medium (10–30%), and high (>30%) urbanization categories as determined by Hincher et al.(Hincher et al.).

Three frog species were collected: *Lithobates catesbeianus* (American bullfrog)*, Lithobates clamitans* (Green frog)*,* and *Lithobates palustris* (Pickerel frog). These are common local frog species and of no conservation concern and thus were selected for non-destructive capture and fecal pellet isolation. All amphibians were captured at sites authorized by the Massachusetts Division of Fisheries and Wildlife (Permit 16.23SCRA). The capture-release protocol was approved by Clark University's Institutional Animal Care and Use Committee (IACUC , Protocol 037R). Frogs were temporarily housed in sterile collection bags until defecation occurred, after which they were released at their original capture site. Fecal samples were then collected using a sterile scoopula. The fecal sample was covered with 20% glycerol and stored at -20 °C until DNA extraction.

Sediment core samples were taken at stream sites using a Large Universal Corer (Aquatic Research Instruments, Idaho, USA) to a depth of ~5 cm at four locations randomly chosen at each site. After sampling, each core was homogenized and ~50 ml duplicate subsamples were stored separately of each core at -80 °C until DNA extraction.

Water samples were collected using acid-washed 4 L cubitainers, rinsed using water downstream

of the collection point. Water samples were placed on ice until filtered later that day. Water samples were

filtered by peristaltic pump by drawing water through a 25 µm Nitex mesh covering the intake opening of

the peristaltic tubing and then passing through a 47 mm 3 µm polycarbonate filter (Cat #XX4404700,

Millipore Sigma, USA) within a stainless-steel filter holder to collect eukaryotic cells. The 3 µm filters

were then aseptically removed and stored in falcon tubes kept at -80 °C until DNA extraction.

Environmental DNA Extraction

Fecal DNA was extracted from frog fecal samples and sediment samples using the QiAmp

PowerSoil Pro DNA Kit (Cat # 47021, Qiagen, Maryland, USA). For water DNA extraction, a phenol

chloroform protocol was used to extract DNA from filters. Modifications to the protocols can be found in

the Supplementary Materials. DNA concentrations of all samples were determined using a Qubit 4

Fluorometer with a dsDNA High Sensitivity kit (ThermoFisher Scientific, Massachusetts, USA). The

purity was measured with a Nanodrop One Microvolume UV-Vis Spectrophotometer (ThermoFisher

Scientific, Massachusetts, USA).

ITS1 Metabarcode sequencing and analysis

The ITS1 region was amplified using the standardized primer set in (Bellemain et al. 2010) to

conduct metabarcode community analysis of fungal communities. Amplification, library preparation, and

2X250 PE sequencing was done by NovoGene Inc. (Sacramento, CA) using on an Illumuna

NovaSeq6000 sequencer. Illumina adapter sequences were removed, and sequences were demultiplexed

by NovoGene. The resulting raw reads were quality trimmed with cutadapt (v3.4) (Martin 2011).

Downstream processing of trimmed reads was performed using DADA2 (v1.16) (Callahan et al. 2016),

and taxonomy was assigned to resulting amplicon sequence variants (ASVs) using the general Fungi

UNITE database (version 9.0) with a naïve Bayesian classifier and a minimum bootstrap confidence

threshold of 80% (Nilsson et al. 2019).

Fungal Community Analysis

All data analyses were conducted in R (4.4.2) (R Core Team 2023) and RStudio (Posit team

2024). Phyloseq (V 1.44.0) (McMurdie and Holmes 2013) was used for microbiome data processing such

as deduplicating reads, defining ASV's and generating counts per ASV. Relative abundances were

calculated for downstream analysis. Rarefaction curves were generated with the function rarecurve from

vegan (v. 2.6) (Oksanen et al. 2025). Reads were not rarified for the subsequent analyzes as this type of

data stratification tends to remove rare, but important taxa (McMurdie and Holmes 2014).

For alpha-diversity analyzes, Shannon and Simpson diversity indices were computed with the

diversity function in the vegan (Oksanen et al. 2025).   Analysis of Variance (ANOVA) were performed

to analyze differences in alpha diversity, using Shannon or Simpson's indices as response variable, and

source and urbanization levels as the explanatory variable. Assumptions for ANOVA were checked using

Q-Q plots for normality and Bartlett's test for homogeneity of variance. Post hoc analyses for significant

ANOVA results were conducted using Tukey's HSD tests. Boxplots were generated using ggplot2

(Wickham 2011).

Beta diversity was compared using permutational multivariate ANOVA (PERMANOVA) using

the adonis2 function in the vegan package (Anderson 2017) with 1,000 permutations. Nonmetric

multidimensional scaling ordination (NMDS) plots were generated using Bray-Curtis dissimilarity

values. For the PERMANOVA, the Bray-Curtis dissimilarity values were used as the response variables,

and the habitat or urbanization levels as the categorical explanatory variables. For the frog fecal samples,

we also included the frog host species and the interaction of urbanization level and host species as the

categorical variables. Posthoc pairwise PERMANOVA analyses were conducted to determine which

specific variables exhibited significant community differences from each other using the pairwiseAdonis

package (v. 0.4.1) (Arbizu 2017) with Benjamini-Hochberg (BH) p-value correction for multiple test

comparisons (Benjamini and Hochberg 1995). The top 20 most abundant taxa per habitat were ranked by

mean relative abundance.

Furthermore, differential abundance analyses were conducted using DESeq2 (Love et al. 2014) to

identify taxa with significant changes in abundance per source across urbanization levels. Variance-

stabilizing transformation (VST) was applied to normalize count data. Differentially abundant taxa

between urbanization categories were identified using the DESeq function, with a significance threshold

of adjusted p-value < 0.05. Log2 fold changes were computed, where an ASV with $log2FC \geq 5$ or ≤ -5

indicates a significant presence in one condition compared to the other.

Fungal trophic guild abundances

ASVs that had no annotation available were removed from downstream analyses and Fungal

ecological guild information was extracted using FUNGuild (v. 1.1) (Nguyen et al. 2016) and its

associated database. After assignment of guild to each ASV, analysis targeted ASVs assigned to only one

trophic guild with a confidence ranking of 'probable' or 'highly probable.' Significant differences in the

abundance of each trophic mode were analyzed using a Kruskal-Wallis test in R, after assumptions for

ANOVA were checked and violated (Q-Q plots for normality and Bartlett's test for homogeneity of

variance). For each guild (Saprotroph, pathogen and symbiont), the abundance was used as the response

variable and the urbanization category as the explanatory variable).

RESULTS

<u>**Sampling and Meta-amplicon Sequencing**</u>

Water (planktonic), stream sediment, and frog fecal samples were collected from the Tatnuck

Brook waterway, spanning a north-to-south urbanization gradient in Worcester (Fig. 1). Urbanization

levels were categorized as low (<10%), medium (10–30%), and high (>30%) based on PISA values

(Table 1). The northernmost sites—Kinney Woods (5.1%) and Tatnuck Brook (9.0%)—were classified as

low urbanization. The next four sites had medium urbanization (17.2–25.6%), while the five most

southern, downstream sites exceeded 45% PISA, with Holy Cross (56.7%) and Beaver Brook (56.6%)

showing the highest urbanization levels. This 10 km gradient reflects a transition from rural to developed

regions.

A total of 129 microbiome samples were collected, with 112 yielding high-quality sequencing

data (fecal n=60, sediment n=30, water n=22) (Supp. Table 1). These samples generated 25,608,648

paired-end raw reads, resulting in 121,368 unique amplicon sequence variants (ASVs) spanning 20 phyla,

80 classes, 214 orders, 596 families, 1,740 genera, and 2,879 species (Table 2). Rarefaction curves

plateaued, confirming sufficient sequencing depth to capture taxonomic diversity (Supplementary Figure

1).

Alpha and Beta diversity of Fungal Communities Across Freshwater Habitats

Alpha and beta diversity analyses revealed significant differences in fungal communities across habitats

and urbanization levels. Shannon's diversity index showed that sediment samples were significantly more

diverse than fecal and water samples (ANOVA, F = 37.95, p<0.001; Tukey's post hoc, p<0.001), while

fecal and water samples did not differ (Tukey's post hoc, p = 0.132). In contrast, Simpson's diversity

index showed no significant differences across habitats (ANOVA, F = 1.39, p=0.253), suggesting similar

dominance patterns among sources. Beta diversity analysis (PERMANOVA) confirmed distinct microbial

communities across habitats (F = 5.79, p = 0.001, R² = 0.095), with pairwise comparisons showing

significant differences between all groups (p = 0.001). Across all habitats, urbanization significantly

influenced fungal community composition (F = 3.920, p = 0.001, R² = 0.117), though habitat type

explained more variance, indicating a stronger role in shaping community structure. Fecal samples were

dominated by *Basidiobolus* and *Cladosporium*, while sediments were dominated by *Lemonniera*

(Supplemental Figure 3). *Only Candida* was consistently found across all habitats (Supplementary Figure

3).

Effects of Urbanization on Fungal Communities

Fecal Fungal Communities

To control for the effect of habitat, samples were analyzed separately by habitat type to look for

further assess patterns in community diversity according to urbanization. In frog fecal samples,

urbanization did not significantly impact alpha diversity using either Shannon's diversity ($F = 1.804$, $p =$

0.174) or Simpson's diversity ($F = 0.01$, $p = 0.994$) (Figure 1 and Supplementary Figure 2). Neither did

frog species (Shannon: $F = 0.502$, $p = 0.608$, Simpson: $F = 0.546$, $p = 0.608$) nor the frog species and

urbanization interaction (Shannon: $F = 0.507$, $p = 0.605$, Simpson: $F = 0.585$, $p = 0.560$).

PERMANOVA results indicate significant differences in fungal communities between frog

species (F=1.29, p=0.020, R²=0.038) (Supplementary Figure 2B) and urbanization levels (F=3.88,

p=0.001, R²=0.116) (Figure 3a). However, urbanization had a stronger effect on beta diversity, while the

interaction between host species and urbanization was not significant (F=1.091, p=0.169, R²=0.032). The

model explains 12% of the variance due to urbanization, 4% due to host species, and only 3% from their

interaction. Pairwise PERMANOVA tests confirmed significant community composition differences

across urbanization levels (p=0.001 for all comparisons). Notably, *Basidiobolus* and *Candida* were

abundant in low urbanization, while *Botrytis*, *Alternaria*, and *Parasarocladium* were more common in

highly urbanized sites (Supplementary Table 2, Supplementary Figure 3). Differential abundance analysis

identified 125 taxa shifting significantly between low and high urbanization, with 82 favoring low and 43

favoring high urbanization (Table 3, Supplementary File 1).

Sediment Fungal Communities

Sediment samples exhibited no significant changes in Shannon's ($F = 2.16$, $p = 0.132$) or

Simpson's ($F = 1.60$, $p = 0.217$) diversity indices across urbanization levels. However, beta diversity was

significantly influenced by urbanization (PERMANOVA, $F = 2.32$, $p = 0.001$), in this case explaining

13% of the observed variation (Figure 3b Pairwise PERMANOVA shows significant differences across

urbanization levels as well (p=0.001 for all comparisons). Low urbanization sites had unique genera like

Apodus and *Podospora*, while high urbanization sites were characterized by *Neoascochyta* and *Candida*

(Supplementary Table 2). Out of 33 taxa that were differentially abundant between low and high

urbanization, ten and twenty-three were more common at low and high urbanization, respectively (Table

3, Supplementary File 2)

Planktonic Fungal Communities

As with the other habitats, planktonic fungal communities showed no significant change in

Shannon's ($F = 0.24$, $p = 0.787$) or Simpson's diversity indices ($F = 0.32$, $p = 0.729$) with urbanization.

Nevertheless, PERMANOVA results indicated that community composition was significantly structured

by urbanization (PERMANOVA, $F = 1.33$, $p = 0.003$, R2 = 0.159) (Figure 3c). A16% of sample variance

was explained by urbanization, again notably lower than for frog fecal mycobiomes. There was overlap in

the clustering of medium and high urbanization samples, but PERMANOVA shows a significant

difference in community composition between these groups (F = 1.528, p = 0.006, R2 = 0.105). There

were only two low-urbanization water samples, so we were unable to test for significant differences in

community composition of these sites from other urbanization sites.

The top 20 genera analysis showed that low and medium urbanization sites had greater

abundances of *Basidiobolus*, *Clavispora*, and *Zygophlyctis*, while high urbanization sites featured

Aureobasidium and *Metarhizium* (Supplementary Table 2). We identified 95 differential abundant taxa

between low and high urbanization sites. Chytridiomycota ASV's represented the most differentially

abundant taxa, with 45 ASV's represented in low urbanization, in comparison to 13 at higher urbanization

(Table 3, Supplementary File 3).

Fungal Trophic Mode Analysis

The 35,854 ASVs were classified into pathotrophs, saprotrophs, symbiotrophs, and unknowns, with most

ASVs labeled as unknown due to multiple assignments or low confidence rankings (Figure 4). Relative

abundances showed saprotrophs as the dominant trophic mode across all habitats and urbanization levels.

While no significant differences were detected in trophic mode abundance across urbanization levels

(Supplementary Table 2), the data indicate a trend of increasing pathotroph abundance in highly

urbanized sediment and water samples. This suggests a potential shift in community composition that

may have implications for ecosystem health and host resilience in urban environments.

DISCUSSION

Urbanization is reshaping ecosystems, altering both macroscopic biodiversity and microbial communities. While urban impacts on bacterial and archaeal communities have been studied, fungal communities remain underexplored. Primary, previous research has often focused on a single habitat, be it environmental (Tedersoo et al. 2014; Ortiz-Vera et al. 2018) or host associated(García-Sánchez et al. 2023), less frequent in difficult to sample aquatic environments. This study examines three interconnected microbial environments, including both environmental and host derived sources. Our results show that while alpha diversity remained stable, beta diversity shifted significantly across habitats, indicating taxonomic turnover rather than biodiversity loss. This aligns with previous studies showing urbanization reshapes microbial composition without reducing species richness (Abrego et al. 2020).

There were some notable differences in strength of correlation of urbanization with community composition. Urbanization influenced fungal community composition across habitats, but habitat type was a stronger predictor. PERMANOVA results showed that urbanization explained 12% of variance in fecal communities, 13% in sediment, and 16% in water, while habitat type accounted for significantly more variation. Within fecal samples, host species identity explained only 4% of variance, suggesting that urbanization exerts a stronger influence on microbial turnover than interspecies differences. These findings indicate that while urbanization alters fungal communities, it primarily drives taxonomic shifts rather than biodiversity loss. This is not surprising because habitat type is a strong driver of prokaryotic community similarity (Thompson et al. 2017)

The observed shifts in fungal community composition suggest that urban environments act as ecological filters, favoring taxa adapted to pollutants, altered nutrient dynamics, and habitat disturbances [42]. However, the specific mechanisms driving these shifts remain unclear. Future metagenomic studies could identify key genes and pathways involved. Additionally, the implications for amphibian health and water quality require further investigation. By analyzing fungal taxa across freshwater habitats, our findings highlight how urbanization shapes microbial diversity and underscore the need for research into

290   the functional roles of urban-adapted fungi in host-microbe interactions, biogeochemical cycling, and

291   ecosystem stability.

292   **Differences in Fungal Community Responses Across Sources**

The three primary habitats—sediment, water, and host associated fecal matter—demonstrate distinct

community compositions influenced by urbanization, reflecting the diverse ecological niches within the

watershed. Sediment, with its high fungal diversity, supports saprotrophic fungi crucial for organic matter

breakdown, likely due to nutrient-rich microenvironments that favor fungal growth and metabolic

diversity. The species richness in sediment may also stem from its role as a microbial sink, capturing

inputs from water and detritus, and supporting taxa that interact with organic substrates.

299   The planktonic communities exhibited fewer taxa but showed notable resilience to urbanization-

300   induced changes, particularly in community composition. The persistence of certain genera, like

301   *Clavispora* and *Aureobasidium*, across urbanization levels in water suggests their functional adaptation to

302   the aquatic environment, highlighting their ecological relevance as nutrient recyclers and decomposers in

303   freshwater systems (Grossart et al. 2019). The resilience observed may be attributed to the natural flow

304   dynamics in water bodies, which might buffer against certain urban pollutants.

305   Frog fecal samples showed the greatest compositional shifts with urbanization. *Basidiobolus*, a

genus linked to amphibian gut health, was abundant in low urbanization sites but absent in high

urbanization samples, suggesting sensitivity to urban stressors. Previous studies found *Basidiobolus* to be

a dominant gut fungus, representing 84.4% of salamander gut fungi and 29% of frog gut fungi (Vargas-

Gastélum et al. 2024) and comprising 60% of gut fungi in slimy salamanders (*Plethodon* spp.) (Walker et

al. 2020). Despite its prevalence in amphibians, our study found it abundant only in low urbanization

sites. This suggests *Basidiobolus* may be a useful bioindicator for habitat quality and amphibian-

microbiome interactions

Novel discoveries in fungal ecology

314   **T**he *Rozellomycota* were highly represented across all sample types and mostly in fecal and

sediment samples. Typically, cryptic or parasitic, they are often linked to other fungi or protists in soil.

The presence of *Rozellomycota* in both amphibian fecal pellets and sediment expands their known habitat

range. Their detection in aquatic environments suggests potential roles in organic matter decomposition,

symbiosis, or host-microbe interactions that warrant further study.

Additionally, this study revealed the unexpected presence of *Basidiobolus* within the planktonic

water communities, marking one of the first records of this genus in freshwater systems. Typically

associated with amphibian gastrointestinal tracts or terrestrial environments such as leaf litter,

Basidiobolus in water points to possible environmental transmission pathways that extend beyond known

host associations, suggesting a broader ecological versatility.

These findings suggest fungal adaptability in urban-impacted freshwater systems, with

urbanization potentially facilitating the movement of terrestrial taxa into aquatic and host-associated

environments. The presence of *Rozellomycota* in novel habitats highlights the need for further studies

using multi-marker approaches to clarify their ecological roles, diversity, and contributions to nutrient

cycling and ecosystem function.

Implications of Trophic Mode Shifts in Urbanized Environments

Trophic mode analysis provides insights into how urbanization may influence the ecological roles of

fungal communities. While no significant statistical differences were found, an increase in pathotroph

abundance with urbanization in sediment and from medium to high urbanization in planktonic habitats

suggests that urban ecosystems may favor pathogenic fungi, a trend reported in previous studies (Pagani

et al. 2023; Jian et al. 2025). This shift could impact amphibians and aquatic ecosystems by increasing

pathogen loads, contributing to biodiversity loss. Urban organic waste may support opportunistic fungi,

while reduced plant diversity limits symbiotic fungi. The heightened pathotroph abundance, particularly

in urbanized sediment and water, raises concerns about potential amphibian health risks, as pathogenic

fungi, such as chytrids, are known to have devastating effects on amphibian populations (Longcore et al.

1999). However, the high proportion of unclassified ASVs underscores the need for further metagenomic

research to clarify functional shifts in fungal communities along urban gradients.

Urbanization as a Filter for Resilient or Opportunistic Fungal Taxa

The observed shifts in fungal communities align with the concept of biotic homogenization, where urbanization filters out sensitive taxa and selects for those resilient to anthropogenic pressures. This effect, where resilient taxa dominate urban ecosystems, highlights a trend toward microbial communities dominated by opportunistic or anthropophilic species (McKinney 2006; Lokatis and Jeschke 2022; Sidemo-Holm et al. 2022). Such homogenization can reduce functional diversity within ecosystems, potentially diminishing the resilience of microbial communities to future environmental changes.

Conservation and Restoration Implications

The distinct fungal community compositions across urbanization levels and sources in this study highlight the need for conservation and restoration efforts tailored to urbanized freshwater systems. The high fungal diversity observed in sediment, especially in low urbanization sites, suggests that sediment habitats act as refuges for microbial biodiversity. Preserving and restoring sediment-rich areas along urban watersheds could help maintain fungal diversity and associated ecosystem functions, such as nutrient cycling and organic matter decomposition.

The results also imply that some fungal taxa, particularly those in frog GI tracts, may be more sensitive to urban stressors and can be bioindicators for assessing environmental health in freshwater systems (Bai et al. 2018; Warnasuriya et al. 2023). For instance, the absence of *Basidiobolus* in high urbanization fecal samples suggests it could serve as a sensitive indicator of habitat quality. Conservation strategies could include monitoring this genus across watersheds to identify areas of significant environmental degradation and inform mitigation actions.

Conclusions

This study serves as an initial step toward understanding the effects of urbanization on fungal communities in freshwater ecosystems, and the interconnected nature they form across different trophic levels. Here we find consistent patterns of impact across three inter-connected microbial habitats: stable alpha diversity but significant shifts in beta diversity associated with urbanization in a freshwater

ecosystem. Future research should expand upon these findings by integrating shotgun metagenomics to

determine specific metabolic pathways affected by urbanization. Additionally, studies tracking temporal

fungal communities could information on the variability of these communities due to urban development.

This interconnectivity of freshwater systems renders it critical to study environmental microbial

communities in tandem with host microbial communities since they serve as source pools for host

microbiomes. Therefore, sediment and planktonic microbial communities in freshwater systems are also

of interest as they not only perform vital ecosystem services but also contribute to the assembly of

microbial communities in amphibians.

REFERENCES

Abrego N, Crosier B, Somervuo P, Ivanova N, Abrahamyan A, Abdi A, Hämäläinen K, Junninen K,
Maunula M, Purhonen J, Ovaskainen O. 2020. Fungal communities decline with urbanization—more in
air than in soil. The ISME Journal 14:2806–2815.

Anderson MJ. 2017. Permutational Multivariate Analysis of Variance (PERMANOVA). In: N.
Balakrishnan, T. Colton, B. Everitt, W. Piegorsch, F. Ruggeri, and J.L. Teugels, eds. Wiley StatsRef:
Statistics Reference Online. Wiley. p. 1–15.

Arbizu PM. 2017. pairwiseAdonis: Pairwise Multilevel Comparison using Adonis.

Astudillo-García C, Hermans SM, Stevenson B, Buckley HL, Lear G. 2019. Microbial assemblages and
bioindicators as proxies for ecosystem health status: potential and limitations. Appl Microbiol Biotechnol
103:6407–6421.

Bai Y, Wang Q, Liao K, Jian Z, Zhao C, Qu J. 2018. Fungal Community as a Bioindicator to Reflect
Anthropogenic Activities in a River Ecosystem. Frontiers in Microbiology 9:3152.

Belt KT, Hohn C, Gbakima A, Higgins JA. 2007. Identification of culturable stream water bacteria from
urban, agricultural, and forested watersheds using 16S rRNA gene sequencing. Journal of Water and
Health 5:395–406.

Benjamini Y, Hochberg Y. 1995. Controlling the False Discovery Rate: A Practical and Powerful
Approach to Multiple Testing. Journal of the Royal Statistical Society: Series B (Methodological)
57:289–300.

Bridge P, Spooner B. 2002. Soil fungi: diversity and detection. In: D.S. Powlson, G.L. Bateman, K.G.
Davies, J.L. Gaunt, and P.R. Hirsch, eds. Interactions in the Root Environment: An Integrated Approach.
Dordrecht: Springer Netherlands. p. 147–154.

Callahan BJ, McMurdie PJ, Rosen MJ, Han AW, Johnson AJA, Holmes SP. 2016. DADA2: High-
resolution sample inference from Illumina amplicon data. Nature Methods 13:581–583.

Chen Y, Martinez A, Cleavenger S, Rudolph J, Barberán A. 2021. Changes in Soil Microbial Communities across an Urbanization Gradient: A Local-Scale Temporal Study in the Arid Southwestern USA. Microorganisms 9:1470.

García-Sánchez JC, Arredondo-Centeno J, Segovia-Ramírez MG, Tenorio Olvera AM, Parra-Olea G, Vredenburg VT, Rovito SM. 2023. Factors Influencing Bacterial and Fungal Skin Communities of Montane Salamanders of Central Mexico. Microbial Ecology 86:670–686.

Grossart H-P, Van Den Wyngaert S, Kagami M, Wurzbacher C, Cunliffe M, Rojas-Jimenez K. 2019. Fungi in aquatic ecosystems. Nature Reviews Microbiology 17:339–354.

Hincher MR, Carleton JP, Wheeler SJ, DelCogliano M, Mathis K, Tabima JF. Ubiquity and diversity of Basidiobolus across amphibian species inhabiting an urbanization gradient. Mycologia 0:1–13.

Hobbie EA, Macko SA, Shugart HH. 1999. Insights into nitrogen and carbon dynamics of ectomycorrhizal and saprotrophic fungi from isotopic evidence. Oecologia 118:353.

Hosen JD, Febria CM, Crump BC, Palmer MA. 2017. Watershed Urbanization Linked to Differences in Stream Bacterial Community Composition. Frontiers in Microbiology 8:1452.

Jian Y, Lin J, Mu C, Wang Y, He Z, Chen G, Ding W. 2025. Short-Term Effects of Three Tree Species on Soil Physicochemical Properties and Microbial Communities During Land-Use Change from Farmland to Forests. Forests 16:362.

Karimi B, Maron PA, Chemidlin-Prevost Boure N, Bernard N, Gilbert D, Ranjard L. 2017. Microbial diversity and ecological networks as indicators of environmental quality. Environmental Chemistry Letters 15:265–281.

Kearns PJ, Fischer S, Fernández-Beaskoetxea S, Gabor CR, Bosch J, Bowen JL, Tlusty MF, Woodhams DC. 2017a. Fight Fungi with Fungi: Antifungal Properties of the Amphibian Mycobiome. Frontiers in Microbiology 8.

Kearns PJ, Fischer S, Fernández-Beaskoetxea S, Gabor CR, Bosch J, Bowen JL, Tlusty MF, Woodhams DC. 2017b. Fight Fungi with Fungi: Antifungal Properties of the Amphibian Mycobiome. Frontiers in Microbiology 8:2494.

Lokatis S, Jeschke JM. 2022. Urban biotic homogenization: Approaches and knowledge gaps. Ecological Applications 32:e2703.

Longcore JE, Pessier AP, Nichols DK. 1999. Batrachochytrium dendrobatidis gen. et sp. nov., a chytrid pathogenic to amphibians. Mycologia 91:219–227.

Love MI, Huber W, Anders S. 2014. Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. Genome Biol 15:550.

Martin M. 2011. Cutadapt removes adapter sequences from high-throughput sequencing reads. EMBnetJournal 17:10.

McKinney ML. 2006. Urbanization as a major cause of biotic homogenization. Biological Conservation 127:247–260.

McMurdie PJ, Holmes S. 2013. phyloseq: An R Package for Reproducible Interactive Analysis and
Graphics of Microbiome Census Data. PLoS ONE 8:e61217.

McMurdie PJ, Holmes S. 2014. Waste Not, Want Not: Why Rarefying Microbiome Data Is Inadmissible.
PLOS Computational Biology 10:e1003531.

Mora J, Olson ,Matthew, Rocks ,Sara S., Zahn G and. 2025. Watershed urbanization alters aquatic plant
mycobiomes through the loss of rare taxa. Mycologia 117:235–245.

Nguyen NH, Song Z, Bates ST, Branco S, Tedersoo L, Menke J, Schilling JS, Kennedy PG. 2016.
FUNGuild: An open annotation tool for parsing fungal community datasets by ecological guild. Fungal
Ecology 20:241–248.

Nilsson RH, Larsson K-H, Taylor AFS, Bengtsson-Palme J, Jeppesen TS, Schigel D, Kennedy P, Picard
K, Glöckner FO, Tedersoo L, Saar I, Kõljalg U, Abarenkov K. 2019. The UNITE database for molecular
identification of fungi: handling dark taxa and parallel taxonomic classifications. Nucleic Acids Research
47:D259–D264.

Oksanen J, Simpson GL, Blanchet FG, Kindt R, Legendre P, Minchin PR, O'Hara RB, Solymos P,
Stevens MHH, Szoecs E, Wagner H, Barbour M, Bedward M, Bolker B, Borcard D, Carvalho G, Chirico
M, Caceres MD, Durand S, Evangelista HBA, FitzJohn R, Friendly M, Furneaux B, Hannigan G, Hill
MO, Lahti L, McGlinn D, Ouellette M-H, Cunha ER, et al. 2025. vegan: Community Ecology Package.

Ortiz-Vera MP, Olchanheski LR, Silva EG da, Lima FR de, Martinez LR del PR, Sato MIZ, Jaffé R, Alves
R, Ichiwaki S, Padilla G, Araújo WL. 2018. Influence of water quality on diversity and composition of
fungal communities in a tropical river. Scientific Reports 8:14799.

Pagani DM, Ventura SPR, Vu D, Mendes-Pereira T, Ribeiro Tomé LM, Carvalho DS de, Costa-Rezende
DH, Kato RB, García GJY, Geml J, Robert V, They NH, Brenig B, Azevedo V, Scroferneker ML, Valente
P, Góes-Neto A. 2023. Unveiling Fungal Community Structure along Different Levels of Anthropic
Disturbance in a South American Subtropical Lagoon. Journal of Fungi 9:890.

Paul MJ, Meyer JL. 2001. Streams in the Urban Landscape. Annual Review of Ecology and Systematics
32:333–365.

Posit team. 2024. RStudio: Integrated Development Environment for R. Boston, MA: Posit Software,
PBC.

QGIS Development Team. QGIS Geographic Information System. QGIS Association.

R Core Team. 2023. R: A Language and Environment for Statistical Computing. Vienna, Austria: R
Foundation for Statistical Computing.

Richard ML, Sokol H. 2019. The gut mycobiota: insights into analysis, environmental interactions and
role in gastrointestinal diseases. Nature Reviews Gastroenterology & Hepatology.

Riley SPD, Busteed GT, Kats LB, Vandergon TL, Lee LFS, Dagit RG, Kerby JL, Fisher RN, Sauvajot
RM. 2005. Effects of Urbanization on the Distribution and Abundance of Amphibians and Invasive
Species in Southern California Streams. Conservation Biology 19:1894–1907.

Roberto AA, Van Gray JB, Leff LG. 2018. Sediment bacteria in an urban stream: Spatiotemporal patterns
in community composition. Water Research 134:353–369.

Sam Q, Chang M, Chai L. 2017. The Fungal Mycobiome and Its Interaction with Gut Bacteria in the
Host. International Journal of Molecular Sciences 18:330.

Scheffers BR, Paszkowski CA. 2012. The effects of urbanization on North American amphibian species:
Identifying new directions for urban conservation. Urban Ecosyst 15:133–147.

Sidemo-Holm W, Ekroos J, Reina García S, Söderström B, Hedblom M. 2022. Urbanization causes biotic
homogenization of woodland bird communities at multiple spatial scales. Global Change Biology
28:6152–6164.

Simonin M, Voss KA, Hassett BA, Rocca JD, Wang S, Bier RL, Violin CR, Wright JP, Bernhardt ES.
2019a. In search of microbial indicator taxa: shifts in stream bacterial communities along an urbanization
gradient. Environmental Microbiology 21:3653–3668.

Simonin M, Voss KA, Hassett BA, Rocca JD, Wang S-Y, Bier RL, Violin CR, Wright JP, Bernhardt ES.
2019b. In search of microbial indicator taxa: shifts in stream bacterial communities along an urbanization
gradient. Environmental Microbiology 21:3653–3668.

Sinha DR. 2010. The environmental Kuznets curve hypothesis and legacy pollution: a geohistorical
analysis of the environmental consequences of industrialization in Worcester, Massachusetts (USA). The
Industrial Geogrpaher 7:1–18.

Tedersoo L, Bahram M, Põlme S, Kõljalg U, Yorou NS, Wijesundera R, Ruiz LV, Vasco-Palacios AM,
Thu PQ, Suija A, Smith ME, Sharp C, Saluveer E, Saitta A, Rosas M, Riit T, Ratkowsky D, Pritsch K,
Põldmaa K, Piepenbring M, Phosri C, Peterson M, Parts K, Pärtel K, Otsing E, Nouhra E, Njouonkou AL,
Nilsson RH, Morgado LN, et al. 2014. Global diversity and geography of soil fungi. Science
346:1256688.

Thompson LR, Sanders JG, McDonald D, Amir A, Ladau J, Locey KJ, Prill RJ, Tripathi A, Gibbons SM,
Ackermann G, Navas-Molina JA, Janssen S, Kopylova E, Vázquez-Baeza Y, González A, Morton JT,
Mirarab S, Zech Xu Z, Jiang L, Haroon MF, Kanbar J, Zhu Q, Jin Song S, Kosciolek T, Bokulich NA,
Lefler J, Brislawn CJ, Humphrey G, Owens SM, et al. 2017. A communal catalogue reveals Earth's
multiscale microbial diversity. Nature 551:457–463.

Uttara S, Bhuvandas N, Aggarwal V. 2012. IMPACTS OF URBANIZATION ON ENVIRONMENT.
Applied Sciences 2.

Vargas-Gastélum L, Romer AS, Ghotbi M, Dallas JW, Alexander NR, Moe KC, McPhail KL, Neuhaus
GF, Shadmani L, Spatafora JW, Stajich JE, Tabima JF, Walker DM. 2024. Herptile gut microbiomes: a
natural system to study multi-kingdom interactions between filamentous fungi and bacteria. mSphere
e00475-23.

Wagg C, Schlaeppi K, Banerjee S, Kuramae EE, Van Der Heijden MGA. 2019. Fungal-bacterial diversity
and microbiome complexity predict ecosystem functioning. Nat Commun 10:4841.

Walker DM, Hill AJ, Albecker MA, McCoy MW, Grisnik M, Romer A, Grajal-Puche A, Camp C,
Kelehear C, Wooten J, Rheubert J, Graham SP. 2020. Variation in the Slimy Salamander (Plethodon spp.)

Skin and Gut-Microbial Assemblages Is Explained by Geographic Distance and Host Affinity. Microbial Ecology 79:985–997.

Wang S-Y, Sudduth EB, Wallenstein MD, Wright JP, Bernhardt ES. 2011. Watershed Urbanization Alters the Composition and Function of Stream Bacterial Communities. PLoS ONE 6:e22972.

Wang Y, Liu J, Wang J, Gao G, Bartlam MG. 2015. Distribution and diversity of fungi in freshwater sediments on a river catchment scale. Frontiers in Microbiology 6.

Warnasuriya SD, Udayanga D, Manamgoda DS, Biles C. 2023. Fungi as environmental bioindicators. Science of The Total Environment 892:164583.

Wickham H ggplot2. 2011. WIREs Computational Stats 3. 180–185.

Conflicts of Interest

The authors report no conflicts of interest

Data availability

All raw sequencing data has been deposited in the Short Read Archive (SRA) under Bioproject number

PRJNA1244802. All supplementary data, including all supplementary files, tables, figures, R objects

and code can be downloaded from GitHub at

(https://github.com/TabimaLab/Fungal_Watershed_Mycobiome).

529 **FIGURES**

530 **Fig 1.**

531
532 Map of the sampling sites along the Tatnuck Brook waterway in Worcester, Massachusetts. The

533 y-axis represents longitude, and the x-axis represents latitude. Each colored ID represents a

534 distinct site, and its name is shown within the black label.

536   **Fig. 2**

Diversity and community composition of microfungi associated with amphibian fecal pellets,

sediment, and water samples. (a) Shannon diversity index of microfungal communities across

three habitats: fecal pellets (Represented by a Green Frog), sediment (Represented by a soil

corer), and water (Represented by a 4L cubitainer). Letters above bars indicate statistical

similarity or differences, with shared letters representing no significant difference (e.g., "a"). (b)

Non-metric multidimensional scaling (NMDS) plot illustrating differences in microfungal

community composition among the three habitats. Each point represents a sample, and its

position reflects similarity in community composition. Axes represent the first two dimensions of

ordination space (NMDS1 and NMDS2)

Fig. 3

Effects of urbanization on microfungal community composition in amphibian fecal pellets,

sediment, and water samples. NMDS plots show the ordination of microfungal communities with

respect to different levels of urbanization (High, Medium, Low) for samples segregated by

habitat type: a) frog fecal samples b) stream sediment samples, and c) planktonic water samples.

For each figure the axes represent the first two dimensions of ordination space (NMDS1 and

NMDS2). Each point represents a sample, which are color coded by urbanization level. Ellipses

represent 95% of the data. Only two low-urbanization water samples were generated (c) so an

ellipse could not be drawn for that sample subset.

Fig. 4

Proportional representation of fungal trophic modes across fecal, sediment, and water samples

under varying urbanization levels. The bar plots illustrate the relative abundance of fungal

trophic modes (pathotrophs, saprotrophs, and symbiotrophs) and unknowns according to the

number of ASVs assigned to each in samples collected from fecal pellets, sediment, and water.

Urbanization levels are compared within each sample type, highlighting shifts in trophic mode

proportions in response to urbanization gradients.

TABLES

Table 1. Urbanization classification of Worcester watershed sampling sites by percent impervious surface

area.

Site	Site ID	Percent Standard Impervious Surface Area (PISA)	PISA Classification
Kinney Woods	KP	5.1%	Low
Tatnuck Brook	PRS, CP2	9.0%	Low
Cook Pond	Cook	17.2%	Medium
Patch Reservoir	PR	22.0%	Medium
Mill Pond	MP	25.6%	Medium
Coes Pond	Coes, CR2, CR3	24.7%	Medium
Beaver Brook	BB1	56.6%	High
	BB2	48.1%	High
	BB3	52%	High
Cemetery	MR1	46.2%	High
Holy Cross	MB	56.7%	High
Middle River	MR, MR2	51.8%	High

Table 2. Distribution of Amplicon Sequence Variants (ASVs) across fungal phyla in fecal, water, and

sediment habitats. The table presents the number of ASVs associated with each fungal phylum (identified

using the UNITE database) across the three sample habitats. Notably, *Ascomycota* and *Basidiomycota*

dominate all habitats, while less common phyla such as *Rozellomycota*, *Chytridiomycota*, and

Mortierellomycota show variable representation. The presence of rare phyla, including

Neocallimastigomycota and *Entomophthoromycota*, highlights the diversity and complexity of fungal

communities in these ecosystems

Phyla (UNITE)	Fecal	Water	Sediment
Aphelidiomycota	15	9	15
Ascomycota	6313	1532	6580
Basidiobolomycota	130	35	48
Basidiomycota	2087	364	3812
Blastocladiomycota	14	9	53
Chytridiomycota	944	943	3506
Entomophthoromycota	1	0	5
Glomeromycota	11	7	42
GS01	18	6	16
Kickxellomycota	12	0	11
Monoblepharomycota	8	16	24
Mortierellomycota	189	24	192
Mucoromycota	342	8	129
Neocallimastigomycota	0	0	5
Olpidiomycota	29	13	30
Rozellomycota	602	189	771
Zoopagomycota	9	6	8

583
Table 3. Differentially abundant genera across habitats and urbanization levels. The table

displays genera that showed significant differences in abundance between habitat types (fecal, sediment,

and water) and urbanization categories (low vs. high). Numbers in parentheses indicate the fold-change of

the most abundant amplicon sequence variant (ASV) within each genus, representing how many times

more abundant the ASV is in one condition compared to another. Values of log2FC ≥ 5 or ≤ -5 indicates a

presence in one condition compared to absence in the other, highlighting taxa that are sensitive to

urbanization-driven environmental changes.

Fecal		Sediment		Water	
Low Urbanization	**High Urbanization**	**Low Urbanization**	**High Urbanization**	**Low Urbanization**	**High Urbanization**
Mucor (29.66)	*Paraphaeosphaeria (26.53)*	*Hymenoscyphus (29.99)*	*Pseudorhypophila (24.14)*	*Hebeloma (27.36)*	*Cyberlindnera (26.56)*
Basidiobolus (28.89)	*Colletotrichum (25.78)*	*Linnemannia (28.13)*	*Cirrenalia (22.18)*	*Zygophlyctis (27.09)*	*Betamyces (22.82)*
Penicillium (27.78)	*Xenooacremonium (25.39)*	*Saitozyma (4.73)*	*Melanomma (21.08)*	*Basidiobolus (25.95)*	*Fusicolla (19.95)*
Dichostereum (27.02)	*Leucosporidium (24.62)*	*Ganoderma (4.05)*	*Babjevia (20.34)*	*Pseudeurotium (25.58)*	
Dothiora (26.99)	*Pseudocercospora (23.88)*		*Nectria (20.31)*	*Rhizoclosmatium (25.47)*	
Apiotrichum (26.47)	*Cadophora (23.62)*		*Conocybe (20.10)*		
Aureobasidium (26.35)	*Cytospora (22.18)*		*Pseudophialocephala (16.95)*		
Taphrina (25.99)	*Comoclathris (11.57)*		*Stagonosporopsis (7.39)*		
Coniella (10.74)	*Cystofilobasidium (10.33)*		*Helicodendron (7.38)*		
	Acremonium (10.16)		*Epicoccum (7.35)*		

LEGEND

🟡 **Low Urbanization**
  1. Kinney Woods
  2. Tatnuck Brook

🔴 **Medium Urbanization**
  3. Cook Pond
  4. Patch Reservoir
  5. Mill Pond
  6. Kinney Woods

🟣 **High Urbanization**
  7. Beaver Brook
  8. Cemetery
  9. Holy Cross
  10. Middle River

A. Fecal   **B. Sediment**   **C. Water**

Stress: 0.224

Stress: 0.204

Stress: 0.118

Urbanization — High — Medium — Low