Metagenomics reveals fibre fermentation and AMR pathways in red grouse (_Lagopus_

scotica) microbiota

Anum Ali Ahmad[1], Kathy Fletcher[2], Nicholas Hesford[2], Laura Glendinning[1]*

[1] The Roslin Institute, The University of Edinburgh, Easter Bush Campus, Midlothian,

Edinburgh, EH25 9RG, UK.

[2] The Game and Wildlife Conservation Trust, Hopetoun Estates Office, Home Farm,

Hopetoun, South Queensferry, EH30 9SL, UK.

*Corresponding author: Laura.glendinning@roslin.ed.ac.uk

Abstract:

Background: The avian caecal microbiota plays a vital role in host nutrition, enabling non-digestible, fibrous material to be converted into compounds that can be absorbed and used as an energy source by the host. The diet of adult red grouse (*Lagopus scotica*) is dominated by heather (*Calluna vulgaris*), which is particularly high in fibre. It is therefore likely that the caecal microbiota plays a key role in enabling grouse to thrive on this diet. In this study, we present the first characterisation of the caecal microbiota of red grouse using modern sequencing methods.

Results: We performed metagenomic sequencing on caecal content samples from fifteen red grouse from three upland estates in Scotland. From this data, we constructed and characterised twelve high-quality, species-level metagenome assembled genomes (MAGs). Eleven of these MAGs could not be assigned a taxonomic label at species level, indicating that they may be novel species. MAGs belonged to diverse taxa (5 phyla) and several encoded genes and pathways for the digestion of fibres, including cellulose, hemi-cellulose, xylooligosaccharides and pectin. Several MAGs also contained antimicrobial resistance genes, predominantly related to vancomycin resistance.

Conclusions: This study is the first to reconstruct commensal microbial genomes from red grouse. The caeca contain diverse, often novel, microbial taxa capable of fermenting various fibres, potentially aiding in the digestion of the red grouse's high-fibre diet. Further research is necessary to explore how these bacteria support red grouse nutrition and health.

Keywords: Microbiota; avian; metagenomics; gut; caeca; caecum; grouse; MAG; fibre; AMR.

Background:

The avian gut microbiota plays a vital role in maintaining host health and contributing to host nutrition (1). While the vast majority of avian microbiota studies have been conducted in farmed species, there are an increasing number of studies examining the microbiota of wild bird populations. These studies have shown that various environmental and genetic factors are important influencers of the composition and function of the gut microbiota (2).

An important challenge when studying wild species is acquiring samples that are truly representative of the gut microbiota. Studies that use wild birds are often limited to collecting non-invasive samples, such as faeces and caecal swabs. However, these samples do not quantitatively represent the microbiota of the large and small intestine (3, 4). Obtaining samples directly from the gastrointestinal section of interest is therefore advantageous, but is often not possible as it requires killing the bird. Game bird species offer a potential avenue for studying the microbiota in "wild" or minimally-managed avian populations, where gastrointestinal samples can be obtained in an ethical manner. Game birds are regularly shot during hunts, or to collect samples to monitor population health. The gastrointestinal tracts from these birds are usually discarded and can therefore be seen as a waste product of this industry. This allows researchers to access gastrointestinal samples without requiring additional animal sacrifice.

Diet has been reported as the most important factor affecting the wild avian microbiota (5). Most game bird species consume plants as a large component of their diet. For wild avian species that consume a predominantly or partially plant-based diet, it is likely that the gut microbiota plays an important role in enabling the host to extract energy from plant fibres. Members of the gut microbiota can ferment fibre into short chain fatty acids that can be used as an energy source by the host (6). In avians the vast majority of fibre fermentation occurs in the caeca (7), blind pouches located near the end of the gastrointestinal tract. In general, herbivorous avian species have longer caeca than non-herbivorous species (8).

Red grouse (*Lagopus scotica*) represent a particularly interesting game bird species in which

to study microbiota fibre fermentation. The main component of the adult red grouse diet is

heather (*Calluna vulgaris*), which is high in fibre (25%) and low in protein (7%) (9). Red

grouse have long caeca (~75 cm), which increase in size when they are supplied with a

higher fibre diet (10). From feed digestibility studies, there is some evidence that the red

grouse microbiota is able to digest a wide variety of fibres, including those that are generally

considered more resistant to fermentation (e.g. lignin) (9).

There is also increasing interest in studying antimicrobial resistance in wild-living animal

populations through a one health lens (11). As game birds are frequently consumed by

humans as food, the presence of AMR in these birds is of interest from this perspective.

Previous studies have identified AMR in gut-associated microbes in UK pheasants and

partridges (12, 13), but this has not been examined in red grouse.

The gut microbiota has been studied in several grouse species using 16S rRNA gene

sequencing and/or metagenomic sequencing, including in capercaillies (14, 15), greater

sage grouse (16-18), wild prairie grouse (19) and ptarmigans (20-22). However, while

bacteria have been cultured from the gastrointestinal tracts of red grouse (23, 24), their

microbiota has not been characterised using these modern sequencing methods. In this

study, we perform metagenomic sequencing analysis of red grouse caecal content samples.

We construct twelve high-quality bacterial metagenome assembled genomes, and

characterise their functional potential to metabolise fibre and their AMR potential.

Methods:

<u>Sample collection and processing</u>

Samples were collected from red grouse killed during legal game bird shoots within the open season in three upland estates in Scotland. Samples were collected in the 2023 Scottish red grouse (*Lagopus scotica*) shooting season (August-October), at three distinct areas of moorland managed for recreational hunting of red grouse (Latitude and longitude: 55.75 - 3.01 (Samples RS1-5), 55.98 -3.33 (Samples RB1-5), and 57.02 -4.12 (Samples CA1-CA5)). Samples were collected from five birds per area. The intestines were removed from birds and frozen on site using a standard domestic freezer within 24 hours. Samples were then transported to the Roslin Institute within one month of sample collection, and stored at -80°C until DNA extraction. DNA was extracted from caecal contents using the QIAamp PowerFecal Pro DNA Kit according to the manufacturer's instructions, using 250-300 mg of caecal contents. DNA quality and quantity were assessed using a NanoDrop spectrophotometer and a Qubit Fluorometer with a Qubit dsDNA Quantification Assay Kit (broad range). Shotgun metagenomic library preparation and sequencing was performed by Novogene, using an Illumina NovaSeq 6000 producing 150 bp paired-end reads. Samples RS1 and RS4 were produced on two separate sequencing runs in order to produce sufficient data quantity, then fastq files from the same sample were concatenated prior to analysis.

<u>Bioinformatic analysis</u>

Sequences were checked for quality using FastQC (v.0.12.1) (25) and MultiQC (v.1.6) (26). Quality control was performed using fastp (v.0.24) (27), with the options "--detect_adapter_for_pe -l 30 -q 20 --trim_poly_g --trim_poly_x". The host genome and feed genome were removed from adaptor trimmed read files. There is currently no RefSeq or GenBank genome available for *Lagopus scotica*, therefore the *Lagopus muta* (rock ptarmigan) genome was downloaded from RefSeq (GCF_023343835.1) and used instead of

a host genome. Heather (*Calluna vulgari*) is the main food source for adult red grouse (28);

therefore the *Calluna vulgari* genome was downloaded from GenBank (GCA_964145215.1).

Sample reads were mapped to host and feed reference genomes using BWA-MEM

(v.0.7.18) (29), followed by SAMtools (v.1.21, samtools fastq -f 12) (30) to select reads

where both paired-end reads were unmapped. The unmapped reads were taxonomically

classified using Kraken2 (v.2.3.1) (31) with a custom database built using *Lagopus muta* and

bacterial, fungal, archaeal, viral, plant, plasmid, and protozoal RefSeq genomes (built on 23rd

June 2025). Bracken (v.3.0.1) (32) was then used to estimate their relative abundances

(options: -k 35 -l 150). Single-sample assemblies and coassemblies were constructed using

Megahit (v.1.2.9) (33).

CoverM (v.0.7) (34) was used to create depth files, which were used as input for genome

binning by MetaBAT 2 (v.2.17) (35). Contamination and completeness of bins were

estimated using CheckM2 (v.1.0.2) (36). dRep (v.3.5.0) (37) was used to dereplicate

genome bins into MAGs, with minimum completeness set at 80% and maximum

contamination at 10%. Genomes were dereplicated at 95% ANI for species-level MAGs, and

99% ANI for strain-level MAGs. Viruses were identified from megahit assemblies using

VirSorter2 (v.2.2.4: database downloaded 3rd February 2025) (38) (options: --include-groups

dsDNAphage,ssDNA --min-length 5000 --min-score 0.5 -j 28 all).

GTDB-Tk (v.2.4.0) (39) was used to assign taxonomy to MAGs (database downloaded 24th

January 2025). METABOLIC was used to assign metabolic function to MAGs (v.4.0) (40).

Antimicrobial resistance (AMR) genes present in MAGs and megahit assemblies were

identified using RGI main (v. 6.0.3) (41) with the Comprehensive Antibiotic Resistance

Database (database downloaded 17th March 2025). Loose hits were included, and the --

low_quality option was used for megahit assemblies. The abundance of MAGs within

samples was calculated using CoverM (options: -p bwa-mem -m mean relative_abundance

trimmed_mean covered_bases variance length reads_per_base count rpkm tpm --min-read-

aligned-percent 75 --min-read-percent-identity 95 --min-covered-fraction 0). Graphs and data

summaries were produced in R using the packages reshape2 (v.1.4.4) (42), cowplot

(v.1.1.3) (43), dplyr (v.1.1.4) (44), ggplot2 (v.3.5.1) (45) and tidyverse (v.2.0.0) (46). MAG

taxonomic trees were constructed using PhyloPhlAn (v.3.1.1) (47) (options: --diversity high -f

supermatrix_aa.cfg --subsample phylophlan –fast). Taxonomic trees were midpoint rerooted

and visualised using figtree (v.1.4.3) (48).

Results:

<u>Sequence quality</u>

We performed metagenomic sequencing on caecal content samples from fifteen Scottish red

grouse. After removal of poor-quality sequences, and trimming of adaptors, polyAs and

polyGs, there were on average 102,686,250 ± 24,122,717 reads per sample. After removal

of host and feed sequences the average number of reads per sample dropped by 89%, to

10,902,109 ± 7,733,350. This was predominantly due to the removal of host reads, indicating

that our samples had a large proportion of host contamination. From single sample

assemblies and a coassembly, we constructed 161 putative genome bins. Quality-filtering

and dereplication resulted in twelve high-quality (<10% contamination, >80% complete)

species-level MAGs (**Table 1**). We also constructed strain-level MAGs, but as we found only

one more MAG than at species-level we decided to restrict our further analyses to species

level. Eighteen high quality viral genomes were identified (**Table S1**), two of which were

proviruses.

Table 1: **Genome stats for species-level MAGs**

MAG	GTDB taxonomy	Complete ness (%)	Contamin ation (%)	Contig N50	Genome size (bp)	Total coding sequences	GC (%)
metabat 2_RB1.3	d__Bacteria;p __Bacillota_A ;c__Clostridia ;o__Oscillospi rales;f__Acut alibacteracea	100	0.17	167,26 2	1,021,94 3	930	33

	e;g__CASPLG01;s__						
metabat2_RB3.3	d__Bacteria;p__Bacillota_A;c__Clostridia;o__Christensenellales;f__Borkfalkiaceae;g__Coproplasma;s__	97.91	0	54,893	1,424,430	1,267	49
metabat2_RS5.3	d__Bacteria;p__Campylobacterota;c__Campylobacteria;o__Campylobacterales;f__Helicobacteraceae;g__Helicobacter_B;s__	99.78	0.01	27,466	1,541,840	1,592	37
metabat2_coassembly.1	d__Bacteria;p__Bacillota_A;c__Clostridia;o__Lachnospirales;f__CA	98.34	0.27	28,654	1,333,034	1,241	28

	G-274;g__;s__						
metabat 2_coass embly.10	d__Bacteria;p __Verrucomic robiota;c__Ve rrucomicrobia e;o__Opitutal es;f__LL51;g __JAIDVZ01; s__	83.77	0.37	8,043	804,428	898	32
metabat 2_coass embly.14	d__Bacteria;p __Bacillota_A ;c__Clostridia ;o__Christens enellales;f__ Borkfalkiacea e;g__Copropl asma;s__	88.08	0.05	6,612	1,044,10 0	1,098	52
metabat 2_coass embly.16	d__Bacteria;p __Actinomyc etota;c__Acti nomycetes;o_ _Actinomycet ales;f__Actin omycetaceae; g__Scrofimicr	96.84	0	30,426	1,474,56 2	1,248	50

	obium;s__						
metabat2_coassembly.28	d__Bacteria;p__Bacillota_A;c__Clostridia;o__Lachnospirales;f__Lachnospiraceae;g__Eubacterium_I;s__	100	0.02	13,892	2,419,852	2,339	48
metabat2_coassembly.36	d__Bacteria;p__Bacillota_A;c__Clostridia;o__Oscillospirales;f__Oscillospiraceae;g__;s__	100	0.06	20,806	2,734,688	2,460	57
metabat2_coassembly.50	d__Bacteria;p__Actinomycetota;c__Coriobacteriia;o__Coriobacteriales;f__UMGS124;g__JAHZGC01;s__	92.48	3.21	11,155	2,307,052	1994	68

metabat2_coassembly.53	d__Bacteria;p__Bacteroidota;c__Bacteroidia;o__Bacteroidales;f__Bacteroidaceae;g__Prevotella;s__	81.06	1.36	8,397	2,145,104	1,915	56
metabat2_coassembly.7	d__Bacteria;p__Bacillota_A;c__Clostridia;o__Oscillospirales;f__Ruminococcaceae;g__RGIG1955;s__RGIG1955 sp017400345	97.55	0.04	73,819	871,746	975	35

Taxonomy of MAGs

All MAGs belonged to bacteria (**Figure 1**), with the most common phylum being Bacillota_A

(n=7), all of the members of which belonged to the class Clostridia. Two members of this

phylum were identified as members of the genus *Coproplasma,* within the family

Christensenellales. The remaining MAGs in this phylum belonged to Lachnospirales (one

identified to family CAG-274 and one identified to the genus *Eubacterium*), and

Oscillospirales (one identified to family Oscillospiraceae, one identified to genus *CASPLG01*,

and one identified as *RGIG1955 sp017400345*). This latter MAG was the only one from our

dataset to be identified to species level by GTDB-Tk. Two MAGs were identified as members

of the phylum Actinomycetota, and were assigned to the genera *Scrofimicrobium* and

JAHZGC01. One MAG belonged to the phylum Bacteroidota (genus *Prevotella*), one

belonged to the phylum Campylobacterota (genus *Helicobacter_B*) and one belonged to the

phylum Verrucomicrobiota (genus *JAIDVZ01*).

Figure 1: Taxonomic tree of species-level, high-quality metagenome assembled

genomes constructed from red grouse caecal metagenomic data. Tip labels represent

the lowest taxonomy assigned to each MAG by GTDB-Tk.

For most samples, the majority of reads did not map to MAGs (69.5 ± 14.3% relative

abundance) (**Figure 2**). By far the most abundant MAG was metabat2_RS5.3 (genus

Helicobacter_B) (26.3 ± 15.4%), followed by metabat2_RB1.3 (genus *CASPLG01*) (1.9 ±

1.3%) and metabat2_RB3.3 (genus *Coproplasma*) (1.4 ± 1.9%). Some samples contained

particularly high levels of *Helicobacter_B*, including RB2 (64.5%), RB4 (46.7%) and RB5

(44.0%).

Figure 2: Relative abundance of species-level, high-quality metagenome assembled

genomes in red grouse caecal samples. Taxonomies represent the lowest taxonomy

assigned to each MAG by GTDB-Tk.

<u>Metabolism of MAGs</u>

Carbohydrate active enzymes (CAZymes) play a key role in carbohydrate fermentation by

the microbiota, including the fermentation of fibrous compounds. Overall, 40 different

CAZymes were identified in our MAGs. The most common CAZyme (18 genes) was GH13,

followed by GH3 (16 genes), GH25 (11 genes), GH2 (9 genes), GH43 (8 genes), GH18, (7

genes), GH77 (7 genes), GH23 (5 genes), GH28 (5 genes). All other CAZymes had less

than 5 genes present in our MAGs (**Table S2**). MAG metabat2_coassembly.53 (*Prevotella*)

contained substantially more CAZyme genes than any other MAG: 55 genes belonging to 27

CAZyme families. Metabat2_coassembly.28 (*Eubacterium*) and metabat2_coassembly.36

(Oscillospiraceae) also contained a large number of CAZyme genes: 25 genes from 17

CAZyme families and 20 genes from 13 families respectively. The three most abundant

MAGs in our samples contained far fewer CAZyme genes: metabat2_RS5.3

(*Helicobacter_B*) contained 3 genes from 2 families, metabat2_RB1.3 (*CASPLG01*)

contained 7 genes from 4 families and metabat2_RB3.3 (*Coproplasma*) contained 9 genes

from 5 families.

Several MAGs were predicted to be able to metabolise complex carbohydrates commonly

found in plants. Cellulose degradation via beta-glucosidase was present in five of our MAGs,

including the taxa *Coproplasma*, *Eubacterium*, Oscillospiracea and *Prevotella*. Cellulose

degradation via cellobiosidase or cellulase was not present in our dataset. Hemicellulose-

debranching enzymes were also present amongst our MAGs, including arabinosidase

(*Prevotella*), beta-glucuronidase (*Coproplasma* and *JAHZGC01*) and alpha-L-rhamnosidase

(Oscillospiraceae and *Prevotella*). One endohemicellulase gene (alpha-D-xyloside

xylohydrolase) was also identified in our *Prevotella* MAG. Beta-xylosidase (targeting

xylooligosaccharides) was also only identified in this *Prevotella* MAG.

Heather is known to contain various phenolic compounds. Enzymes for converting phenol

into benzoyl-CoA were found in two MAGs (Oscillospiraceae and *Helicobacter_B*). Although

starch is present in low concentrations in heather, amylolytic enzymes involved in starch

digestion were also present in two MAGs: alpha-amylase in the *Prevotella* MAG and

pullulanase in the *JAIDVZ01* MAG. Only two genes were identified related to sulphur

metabolism, namely sulphite reduction in the *Eubacterium* MAG, and sulphur oxidation in the

CAG-274.

<u>Antimicrobial resistance genes</u>

We next investigated whether our MAGs contained antimicrobial resistance (AMR) genes

(**Table S3**). Whilst we do not expect any of the sampled red grouse to have been directly

treated with antibiotics, AMR genes occur naturally in many bacteria, even without antibiotic

exposure. The RGI "strict" algorithm is used to detect previously unknown variants of known

AMR genes. Twenty-nine AMR genes were detected in our MAGs using this algorithm

(**Table 2**), the vast majority of which encoded resistance to vancomycin (27 genes in 9

MAGs). A metronidazole resistance gene was detected in metabat2_RB3.3 (*Coproplasma*)

and a tetracycline gene in metabat2_RS5.3 (*Helicobacter*).

The RGI "loose" algorithm aims to identify more distant homologs of AMR genes. While this

method is more prone to false positives, we decided to cautiously employ it as members of

the grouse microbiota are likely to be poorly represented in the CARD database. This

method identified 13,159 potential AMR gene homologs amongst our MAGs, with the top

three targeted antibiotics as oxacillin, tetracycline and vancomycin. The MAGs with the most

putative AMR gene homologs were metabat2_coassembly.36 (n=1972, Oscillospiraceae),

metabat2_coassembly.28 (n=1882, *Eubacterium*), and metabat2_coassembly.50 (n=1525,

JAHZGC01).

Table 2: AMR genes present in MAGs (strict RGI hits)

MAG	Taxonomy	Antibiotic	Gene	% Identity	Resistance Mechanism	AMR Gene Family
metabat2_coassembly.1	f__CAG-274	vancomycin; teicoplanin	vanW	30.04	antibiotic target alteration	vanW; glycopeptide resistance gene cluster
metabat2_coassembly.1	f__CAG-274	vancomycin	vanG	41.88	antibiotic target alteration	glycopeptide resistance gene cluster; Van ligase

metabat2_coassembly.1	f__CAG-274	vancomycin	vanT	36.04	antibiotic target alteration	glycopeptide resistance gene cluster; vanT
metabat2_coassembly.14	g__Coproplasma	vancomycin; teicoplanin	vanW	30.75	antibiotic target alteration	vanW; glycopeptide resistance gene cluster
metabat2_coassembly.28	g__Eubacterium	vancomycin; teicoplanin	vanW	32.82	antibiotic target alteration	vanW; glycopeptide resistance gene cluster
metabat2_coassembly.28	g__Eubacterium	vancomycin	vanT	38.24	antibiotic target alteration	glycopeptide resistance gene cluster; vanT
metabat2_coassembly.36	f__Oscillospiraceae	vancomycin; teicoplanin	vanW	39.9	antibiotic target alteration	vanW; glycopeptide de

						resistance gene cluster
metabat2_coassembly.36	f__Oscillospiraceae	vancomycin	vanT	35.22	antibiotic target alteration	glycopeptide resistance gene cluster; vanT
metabat2_coassembly.36	f__Oscillospiraceae	vancomycin	vanY	61.62	antibiotic target alteration	vanY; glycopeptide resistance gene cluster
metabat2_coassembly.36	f__Oscillospiraceae	vancomycin; teicoplanin	vanW	39.59	antibiotic target alteration	vanW; glycopeptide resistance gene cluster
metabat2_coassembly.36	f__Oscillospiraceae	vancomycin	vanG	39.94	antibiotic target alteration	glycopeptide resistance gene cluster; Van ligase

metabat2_coassembly.36	f__Oscillospiraceae	vancomycin	vanT	48.37	antibiotic target alteration	glycopeptide resistance gene cluster; vanT
metabat2_coassembly.36	f__Oscillospiraceae	vancomycin	vanG	53.78	antibiotic target alteration	glycopeptide resistance gene cluster; Van ligase
metabat2_coassembly.36	f__Oscillospiraceae	vancomycin	vanW	56.25	antibiotic target alteration	vanW; glycopeptide resistance gene cluster
metabat2_coassembly.50	g__JAHZGC01	vancomycin	vanXY	41.85	antibiotic target alteration	glycopeptide resistance gene cluster; vanXY
metabat2_coassembly.50	g__JAHZGC01	vancomycin	vanG	40.51	antibiotic target alteration	glycopeptide resistance

						gene cluster; Van ligase
metabat2_coassembly.50	g__JAHZGC01	vancomycin	vanT	40.51	antibiotic target alteration	glycopeptide resistance gene cluster; vanT
metabat2_coassembly.50	g__JAHZGC01	metronidazole	nimJ	49.07	antibiotic inactivation	nitroimidazole reductase
metabat2_coassembly.50	g__JAHZGC01	vancomycin	vanH	43.71	antibiotic target alteration	vanH; glycopeptide resistance gene cluster
metabat2_coassembly.53	g__Prevotella	vancomycin	vanT	36.98	antibiotic target alteration	glycopeptide resistance gene cluster; vanT
metabat2_coassembly.7	s__RGIG1955 sp017400	vancomycin	vanY	37.74	antibiotic target alteration	vanY; glycopeptide de

						resistance gene cluster
metabat2_coassembly.7	s__RGIG1955 sp017400345	vancomycin	vanG	61.21	antibiotic target alteration	glycopeptide resistance gene cluster; Van ligase
metabat2_coassembly.7	s__RGIG1955 sp017400345	vancomycin	vanXY	53.66	antibiotic target alteration	glycopeptide resistance gene cluster; vanXY
metabat2_coassembly.7	s__RGIG1955 sp017400345	vancomycin	vanT	53.12	antibiotic target alteration	glycopeptide resistance gene cluster; vanT
metabat2_RB1.3	g__CASPLG01	vancomycin	vanT	37.67	antibiotic target alteration	glycopeptide resistance gene cluster; vanT

metabat2_RB1.3	g__CASPLG01	vancomycin	vanG	41.13	antibiotic target alteration	glycopeptide resistance gene cluster; Van ligase
metabat2_RB1.3	g__CASPLG01	vancomycin	vanY	36.21	antibiotic target alteration	vanY; glycopeptide resistance gene cluster
metabat2_RB3.3	g__Coproplasma	vancomycin; teicoplanin	vanW	28.37	antibiotic target alteration	vanW; glycopeptide resistance gene cluster
metabat2_RS5.3	g__Helicobacter	Tetracycline	adeF	43.68	antibiotic efflux	resistance-nodulation-cell division (RND) antibiotic efflux pump

Taxonomic classification of reads

As MAGs normally only capture the most abundant bacterial and archaeal taxa in samples,

we used Kraken2/Bracken to taxonomically classify our metagenomic reads and

characterise taxonomies that may not have been captured by MAGs (**Figure 3**). On average,

85.0 ± 1.63% metagenomic reads remained unclassified in our data. The classified reads

were assigned to 4 domains, 82 phyla, and 2,632 genera across all samples. At the domain

level, the reads were dominated by bacteria (56.3 ± 12.8%, relative abundance) and

Eukaryota (43.2 ± 12.7%) across all samples **(Figure 3A)**. Streptophyta (40.3 ± 12.0%) and

Campylobacterota (27.7 ± 13.4 %) were dominant phyla **(Figure 3B)**. Other minor phyla

included Pseudomonadota (9.27 ± 3.01%), Bacillota (8.87 ± 3.25%), Bacteroidota (4.41 ±

2.84%), and Actinomycetota (3.49 ± 2.87%). At the genus level, *Helicobacter* (18.8 ± 9.42%)

and *Campylobacter* (6.21 ± 2.89%) were highly abundant across all samples **(Figure 3C)**.

Other minor genera included *Cryptomeria* (2.28 ± 0.57%), *Escherichia* (2.18 ± 2.35%),

Solanum (1.88 ± 0.69%), and *Cucumis* (1.14 ± 0.52%).

253 **Figure 3: Barplot showing Kracken2/Bracken classified metagenomic reads at the**

254 **domain (A), phylum (B), and genus (C) levels. The top 10 highly abundant taxa are**

255 **displayed, while the remaining taxa are grouped as "Other".**

Discussion:

The red grouse represents a particularly interesting species in which to study the microbiota,

due to its reliance on a single, high-fibre plant species (heather - *Calluna vulgaris*) for the

majority of its diet (9). Avians do not produce fibre fermenting enzymes, and therefore in

order to acquire nutrition from fibre they are reliant on their gut microbiota to ferment fibre

into compounds that they can absorb and use as an energy source (e.g. short chain fatty

acids) (6). In this study, we examined the gut microbiota of red grouse for the first time using

metagenomics to construct metagenome assembled genomes. Unlike metabarcoding

methods (e.g. 16S rRNA gene sequencing), metagenomics allows us to not only study the

taxa that are present in a community, but also provides functional information on their

metabolic and virulence potential.

Taxonomic profiling using Kraken2/Bracken revealed that the majority of our metagenomic

reads remained unclassified. This is likely due to the fact that the gut microbiota of red

grouse has not previously been examined using metagenomic sequencing methods, so red

grouse-specific members of the gut microbiota will not currently be included in genomic

databases. Similar results have been reported for other wild bird species (49-51),

demonstrating the understudied nature of the microbiota of these species. A large proportion

of our reads originated from plants, likely from the bird's diet. Due to our quality control

methods these reads are unlikely to have impacted our microbiota analysis, but these results

highlight the possibility of using gut metagenomics to study the diets of wild birds. The

results from our Kraken2/Bracken analysis should be interpreted cautiously. As members of

the red grouse microbiota are unlikely to be represented in the RefSeq database, they are

unlikely to be present in the Kraken2 database used in our analysis. Misassignment of

taxonomies is therefore likely, particularly at lower taxonomic levels.

We constructed MAGs to gain a deeper understanding of the microbial community

composition and its functional potential, and to begin to address the lack of red grouse

microbiota representatives in public sequence databases. In this study, we identified twelve

high-quality, species-level MAGs. Of our twelve MAGs, eleven could not be assigned to a

species by GDTB-Tk. The GTDB is constructed from high-quality genomes obtained from

RefSeq and GenBank, and the release used in this study contains 596,859 microbial

genomes in 113,104 species clusters, including genomes constructed from metagenomic

data. If genomes cannot be assigned a taxonomy by this tool it is therefore a good indication

that they are likely to be novel. Only one MAG was assigned a species label, RGIG1955

sp017400345 within the Ruminococcaceae family. This is a poorly characterised species,

which has previously only been identified in metagenomics data from ruminants (52, 53). A

MAG from the genus *Helicobacter_B* was the most abundant MAG in our samples,

comprising on average 26% of the microbiota. Kraken2/Bracken analysis also identified this

genus to be highly abundant in our metagenomic reads. *Helicobacter* is a highly diverse

genus, which while commonly associated with gastrointestinal disease (e.g. *H.pylori* infection

in humans) is also a common member of the gut microbiota in many species, including

grouse (16, 21). In ptarmigan intestines, Bjørnsen *et al.* found similar average abundances

of *Helicobacter* as observed in our study (24% ± 3) (54).

The majority of our MAGs were identified as Clostridia, which is a commonly identified taxon

in the guts of other grouse species (16, 17, 20). Of these MAGs, most are from poorly

described taxa identified from metagenomic data: the genus *Coproplasma* has been

identified in the chicken gut (55), the genus *CASPLG01* has been identified in mice (56-58),

and the family CAG-274 has been identified in metagenomic datasets from diverse species

(59-61). In contrast, the diverse genus *Eubacterium* is relatively well studied, and has been

proposed as a potential probiotic in humans due to the ability of some species to produce

butyrate (62). Of the two MAGs assigned to the phylum Actinomycetota, one belonged to the

genus *JAHZGC01* which has only one representative in the GTDB, obtained from human

metagenomic data (63). The other MAG belonged to the genus *Scrofimicrobium*, which has

been isolated from the gastrointestinal tracts of diverse species (64, 65).Only one MAG was

identified as belonging to the phylum Verrucomicrobiota, a member of the genus *JAIDVZ01*,

which is poorly characterised and has only been identified in metagenomic data (66, 67).

Conversely, the only MAG identified in the phylum Bacteroidota belonged to the very well

characterised genus *Prevotella*. This genus is known to play a key role in the fermentation of

fibre in various herbivores (68) and is commonly found in the microbiota of avians, including

grouse species (16, 17).

It is therefore unsurprising that our *Prevotella* MAG contained the largest quantity of

CAZyme genes amongst our MAGs, and was predicted to degrade fibrous compounds

including cellulose, hemi-celluloses and xylooligosaccharides. However, several MAGs from

less well characterised taxa were also found to encode for diverse CAZyme genes and to be

capable of degrading fibres, highlighting the potential role of poorly characterised taxa in

fibre fermentation in the grouse caeca. Several CAZymes families with xylooligosaccharide,

pectin and hemicellulose degrading activity (e.g. GH36, GH42, GH73) were identified both in

our dataset and in the microbiota of other grouse species (14, 20). Various phenolic

compounds are also found in heather, and have been shown to affect fermentation in the

rumen microbiota (69). Pathways for the degradation of phenols were found in both our

MAGs and other grouse species (17, 20), as were pathways for starch degradation (20).

These findings indicate some degree of shared function in the grouse gut microbiota, in

relation to digestion of plant-derived components of the diet.

Our MAGs also contained several antimicrobial resistance genes. It is likely that many of the

hits found using the "loose" algorithm from RGI are false-positives, due to their high number.

Using the strict RGI algorithm, the most commonly targeted antibiotic by AMR genes was

vancomycin. This included the gene families: vanW, Van ligase, vanT, vanY, vanXY, and

vanH. These genes are present in a wide range of taxa bacterial taxa (70) and have also

been identified in samples from wild western capercaillies (14), but not amongst wild prairie

grouse (19). Many of these resistance genes are not well characterised, with limited

information on their molecular structures (70). vanT is found within resistance cassettes and

encodes a membrane-bound serine racemase, while vanH encodes a dehydrogenase that

converts pyruvate to d-lactate. The mechanism of action of vanW is currently unknown. vanY

and vanXY are not required for vancomycin resistance but may contribute to increased

resistance. A resistance-nodulation-cell division antibiotic efflux pump, predicted to target

tetracycline was found in our *Helicobacter* MAG. Similar efflux pumps have been found in

other species of *Helicobacter* (71). A nitroimidazole reductase gene was also identified in our

JAHZGC01 MAG. Similar genes can be found in a wide variety of bacterial taxa (72), but to

our knowledge this is the first time this gene has been identified in this taxon. The ecological

and one health relevance of these AMR genes is currently unknown, and is worth further

investigation.

The main weakness of our study is the relatively small number of MAGs which we were able

to construct in comparison to studies in other wild avians (14, 51, 73). However, it is worth

noting that the MAG quality cut-offs used for these studies were often substantially lower

than in our study (e.g. minimum 50% completeness). We chose to use a higher cut-off of

80% completeness as we were predominantly interested in predicting the metabolism of

these bacterial species, and lower genome completeness can have a significant effect on

functional inference (74). The small number of taxa observed in our samples may be truly

representative of the microbiota, with the community being dominated by a small number of

more abundant bacteria. In general, the more abundant a species is in a sample then the

more likely it will be able to be constructed into a high-quality MAG. The high abundance of

Helicobacter in our samples may have led to other species not being present in a high

enough relative abundance to be constructed into MAGs. The low number of MAGs may

also be due to reduced sequencing depth, due to the large amount of reads that belonged to

host (89%).This percentage is similar to that found in saliva, throat, and vaginal swabs (75),

but is unusual for the caeca or large intestine. One possible methodological explanation for

this high degree of contamination could be how samples were stored. In order to enhance

ease of sampling in the field and prevent spoiling of the carcass, intestines were initially

stored whole, then defrosted and the contents removed for DNA extraction. This may have

led to increased lysis of host cells, and thereby increased host DNA in our samples.

Conclusion:

This is the first study to reconstruct microbial genomes from the caeca of red grouse. The

red grouse caeca harbours diverse microbial taxa, many of which are novel. These taxa

have the potential to ferment various forms of fibre, thereby contributing to the digestion of

the red grouse's high fibre diet. Further studies are needed to understand how these

bacteria contribute to grouse nutrition and health.

List of abbreviations:

AMR: Antimicrobial resistance

ANI: Average nucleotide identity

Bp: Base pairs

CAZymes Carbohydrate active enzymes

GH: Glycoside hydrolases

GTDB: Genome taxonomy database

MAG: Metagenome assembled genome

rRNA: ribosomal RNA

Declarations

Ethics approval and consent to participate:

Red grouse carcasses were privately owned by the estates on which the birds were shot. The Game Wildlife and Conservation Trust accessed and collected the gastrointestinal samples for this study. Informed consent was obtained from the owners to use these samples in our study. Field studies were conducted in accordance with local legislation. In the United Kingdom the birds listed in Schedule 2 of the Wildlife and Countryside Act 1981 (which include red grouse) are exempt from the general provision of protection. Any person is able to kill or 'take' a species listed outside of the close season for that bird. Therefore, red grouse can be legally taken between 12th August and 10th December within the United Kingdom. Red grouse were not killed for the purpose of providing samples for this study. Samples were collected from red grouse killed during legal game bird shoots within the open season in Scotland, from gastrointestinal samples that would normally be discarded as waste material. Therefore, approval for this study from an Animal Welfare and Ethical Review Body was not required.

Consent for publication:

Not applicable

Availability of data and materials:

The datasets generated and analysed during the current study (fastq files, assemblies, genome bins and metagenome assembled genomes) are available in the European Nucleotide Archive under the project number PRJEB85541.

Competing interests:

The authors declare that they have no competing interests.

Funding:

This work was supported by the Biotechnology and Biological Sciences Research Council

Institute through Strategic Program Programme Grant funding (BBS/E/RL/230001A). Laura

Glendinning is supported by a University of Edinburgh Chancellor's Fellowship. As a UK

registered charity, the Game and Wildlife Conservation Trust relies mainly on fundraising

from the private sector. For the purpose of open access, the author has applied a Creative

Commons Attribution (CC BY) licence to any Author Accepted Manuscript version arising

from this submission.

Authors' contributions:

The authors confirm contribution to the paper as follows: study conception and design - LG;

data collection - AA, NH and KF; analysis and interpretation of results - AA and LG; writing –

original draft - LG; writing – editing and review - AA, LG, NH and KF. All authors reviewed

the results and approved the final version of the manuscript.

Acknowledgements:

We would like to acknowledge the upland estate staff who provided the red grouse samples.

References:

1.   Sun F, Chen J, Liu K, Tang M, Yang Y. The avian gut microbiota: Diversity, influencing factors, and future directions. Front Microbiol. 2022;**13**: 934272 doi: 10.3389/fmicb.2022.934272

2.   Bodawatta KH, Hird SM, Grond K, Poulsen M, Jønsson KA. Avian gut microbiomes taking flight. Trends Microbiol. 2022;**30**: 268-80 doi: 10.1016/j.tim.2021.07.003

3.   Videvall E, Strandh M, Engelbrecht A, Cloete S, Cornwallis CK. Measuring the gut microbiome in birds: Comparison of faecal and cloacal sampling. Mol Ecol Resour. 2018;**18**: 424-34 doi: 10.1111/1755-0998.12744

4.   Stanley D, Geier MS, Chen H, Hughes RJ, Moore RJ. Comparison of fecal and cecal microbiotas reveals qualitative similarities but quantitative differences. BMC Microbiol. 2015;**15**: 51 doi: 10.1186/s12866-015-0388-6

5.   Matheen MIA, R. GM, and Dudaniec RY. Dominant factors shaping the gut microbiota of wild birds. Emu. 2022;**122**: 255-68 doi: 10.1080/01584197.2022.2114088

6.   Mahmood T, Guo Y. Dietary fiber and chicken microbiome interaction: Where will it lead to? Anim Nutr. 2020;**6**: 1-8 doi: 10.1016/j.aninu.2019.11.004

7.   Józefiak D, Rutkowski A, Martin SA. Carbohydrate fermentation in the avian ceca: A review. Anim Feed Sci Technol. 2004;**113**: 1-15 doi: 10.1016/j.anifeedsci.2003.09.007

8.   Hunt A, Al-Nakkash L, Lee AH, Smith HF. Phylogeny and herbivory are related to avian cecal size. Sci Rep. 2019;**9**: 4243 doi: 10.1038/s41598-019-40822-0

9.   Moss R, Parkinson JA. The digestion of heather (*Calluna vulgaris*) by red grouse (*Lagopus lagopus scoticus*). Br J Nutr. 1972;**27**: 285-98 doi: 10.1079/BJN19720093

10.   Moss R, and Trenholm IB. Food intake, digestibility and gut size in red grouse. Br Poult Sci. 1987;**28**: 81-9 doi: 10.1080/00071668708416939

11.   Larsson DGJ, Gaze WH, Laxminarayan R, Topp E. AMR, one health and the environment. Nat Microbiol. 2023;**8**: 754-5 doi: 10.1038/s41564-023-01351-9

12.   Bain C, Rathor G, Jones BP, Hassan MM, Papke P, et al. β-Lactam resistance genes

present in UK pheasants and red-legged partridges. Vet Rec. 2023;**192**: e2540 doi:

https://doi.org/10.1002/vetr.2540

13.   Chintoan-Uta C, Seguino A, editors. Antimicrobial resistance of *Campylobacter* spp.

in wild game pheasants (*Phasianus colchicus*) in Scotland. Microbiology Society Annual

Conference; 2018.

14.   Rasmussen JA, Chua PYS. Genome-resolving metagenomics reveals wild western

capercaillies (*Tetrao urogallus*) as avian hosts for antibiotic-resistance bacteria and their

interactions with the gut-virome community. Microbiol Res. 2023;**271**: 127372 doi:

10.1016/j.micres.2023.127372

15.   Wienemann T, Schmitt-Wagner D, Meuser K, Segelbacher G, Schink B, et al. The

bacterial microbiota in the ceca of Capercaillie (*Tetrao urogallus*) differs between wild and

captive birds. Syst Appl Microbiol. 2011;**34**: 542-51 doi: 10.1016/j.syapm.2011.06.003

16.   Drovetski SV, O'Mahoney MJV, Matterson KO, Schmidt BK, Graves GR. Distinct

microbiotas of anatomical gut regions display idiosyncratic seasonal variation in an avian

folivore. Anim microbiome. 2019;**1**: 2 doi: 10.1186/s42523-019-0002-6

17.   Kohl KD, Connelly JW, Dearing MD, Forbey JS. Microbial detoxification in the gut of

a specialist avian herbivore, the Greater Sage-Grouse. FEMS Microbiol Lett. 2016;**363**:

fnw144 doi: 10.1093/femsle/fnw144

18.   Vaasjo E, Stothart MR, Black SR, Poissant J, Whiteside DP. The impact of

management on the fecal microbiome of endangered greater sage-grouse (*Centrocercus*

urophasianus) in a zoo-based conservation program. Conserv Physiol. 2024;**12**: coae052

doi: 10.1093/conphys/coae052

19.   Drovetski SV, Schmidt BK, Lai JE, Gross MS, Hladik ML, et al. Exposure to crop

production alters cecal prokaryotic microbiota, inflates virulome and resistome in wild prairie

grouse. Environ Pollut. 2022;**306**: 119418 doi: 10.1016/j.envpol.2022.119418

20.   Salgado-Flores A, Tveit AT, Wright A-D, Pope PB, Sundset MA. Characterization of

the cecum microbiome from wild and captive rock ptarmigans indigenous to Arctic Norway.

PLOS ONE. 2019;**14**: e0213503 doi: 10.1371/journal.pone.0213503

21.   Durnova E, Karmanova E, Sorokina T, Mayorova K, Aksenov A. Bacterial diversity

and composition in the internal organs of Taiga Bean Goose, Greater White-Fronted Goose

and Willow Ptarmigan as a new tools in the Arctic biomonitoring system. Diversity. 2025;**17**:

101 doi: 10.3390/d17020101

22.   Ingvaldsen EW, Østnes JE, Kleven O, Davey M, Fossøy F, Nilsen EB. Fecal DNA

metabarcoding reveals seasonal and annual variation in willow ptarmigan diet. R Soc Open

Sci. 2024;**11**: 231518 doi: 10.1098/rsos.231518

23.   Dunn JC, Clegg SR. High prevalence of multi-drug resistant bacteria in faecal

samples from UK passerine birds. bioRxiv. 2024: 2024.07.24.604896 doi:

10.1101/2024.07.24.604896

24.   Fantham HB. Observations on the parasitic protozoa of the Red Grouse (*Lagopus*

scoticus), with a note on the Grouse Fly. Proc Zool Soc Lond. 1910;**80**: 692-708 doi:

10.1111/j.1096-3642.1910.tb01911.x

25.   FastQC: A quality control analysis tool for high throughput sequencing data.

https://github.com/s-andrews/FastQC.

26.   Ewels P, Magnusson M, Lundin S, Käller M. MultiQC: Summarize analysis results for

multiple tools and samples in a single report. Bioinformatics. 2016;**32**: 3047-8 doi:

10.1093/bioinformatics/btw354

27.   Chen S, Zhou Y, Chen Y, Gu J. fastp: An ultra-fast all-in-one FASTQ preprocessor.

Bioinformatics. 2018;**34**: i884-i90 doi: 10.1093/bioinformatics/bty560

28.   Eastman DS, Jenkins D. Comparative food habits of red grouse in northeast

Scotland, using fecal analysis. J Wildl Manag. 1970;**34**: 612-20 doi: 10.2307/3798870

29.   Li H. Aligning sequence reads, clone sequences and assembly contigs with BWA-

MEM. arXiv 2013;**1303**: 3997 doi: 10.48550/arXiv.1303.3997

30.   Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, et al. The Sequence

Alignment/Map format and SAMtools. Bioinformatics. 2009;**25**: 2078-9 doi:

10.1093/bioinformatics/btp352

31.   Wood DE, Lu J, Langmead B. Improved metagenomic analysis with Kraken 2.

Genome Biol. 2019;**20**: 257 doi: 10.1186/s13059-019-1891-0

32.   Lu J, Breitwieser FP, Thielen P, Salzberg SL. Bracken: Estimating species

abundance in metagenomics data. PeerJ Comput Sci. 2017;**3**: e104 doi: 10.7717/peerj-

cs.104

33.   Li D, Liu C-M, Luo R, Sadakane K, Lam T-W. MEGAHIT: An ultra-fast single-node

solution for large and complex metagenomics assembly via succinct de Bruijn graph.

Bioinformatics. 2015;**31**: 1674-6 doi: 10.1093/bioinformatics/btv033

34.   Aroney ST, Newell RJ, Nissen JN, Camargo AP, Tyson GW, Woodcroft BJ. CoverM:

Read alignment statistics for metagenomics. arXiv. 2025;**preprint**: 2501.11217 doi:

2501.11217

35.   Kang DD, Li F, Kirton E, Thomas A, Egan R, et al. MetaBAT 2: An adaptive binning

algorithm for robust and efficient genome reconstruction from metagenome assemblies.

PeerJ. 2019;**7**: e7359 doi: 10.7717/peerj.7359

36.   Chklovski A, Parks DH, Woodcroft BJ, Tyson GW. CheckM2: A rapid, scalable and

accurate tool for assessing microbial genome quality using machine learning. Nat Methods.

2023;**20**: 1203-12 doi: 10.1038/s41592-023-01940-w

37.   Olm MR, Brown CT, Brooks B, Banfield JF. dRep: A tool for fast and accurate

genomic comparisons that enables improved genome recovery from metagenomes through

de-replication. 2017;**11**: 2864-8 doi: 10.1038/ismej.2017.126

38.   Guo J, Bolduc B, Zayed AA, Varsani A, Dominguez-Huerta G, et al. VirSorter2: A

multi-classifier, expert-guided approach to detect diverse DNA and RNA viruses.

Microbiome. 2021;**9**: 37 doi: 10.1186/s40168-020-00990-y

39.   Chaumeil P-A, Mussig AJ, Hugenholtz P, Parks DH. GTDB-Tk: A toolkit to classify

genomes with the Genome Taxonomy Database. Bioinformatics. 2019;**36**: 1925-7 doi:

10.1093/bioinformatics/btz848

40.   Zhou Z, Tran PQ, Breister AM, Liu Y, Kieft K, et al. METABOLIC: High-throughput

profiling of microbial genomes for functional traits, metabolism, biogeochemistry, and

community-scale functional networks. Microbiome. 2022;**10**: 33 doi: 10.1186/s40168-021-

01213-8

41.   Alcock BP, Huynh W, Chalil R, Smith KW, Raphenya Amogelang R, et al. CARD

2023: Expanded curation, support for machine learning, and resistome prediction at the

Comprehensive Antibiotic Resistance Database. Nucleic Acids Res. 2022;**51**: D690-D9 doi:

10.1093/nar/gkac920

42.   Wickham H. Reshaping data with the reshape package. J Stat Softw. 2007;**21**: 1 - 20

doi: 10.18637/jss.v021.i12

43.   Wilke CO. cowplot: Streamlined plot theme and plot annotations for 'ggplot2'. 2024.

44.   Wickham H, François R, Henry L, Müller K, Vaughan D. dplyr: A grammar of data

manipulation. 2023.

45.   Wickham H. ggplot2: Elegant graphics for data analysis: Springer-Verlag New York;

2016.

46.   Wickham H, Averick M, Bryan J, Chang W, McGowan LDA, et al. Welcome to the

Tidyverse. J Open Source Softw. 2019;**4**: 1686 doi: 10.21105/joss.01686

47.   Asnicar F, Thomas AM, Beghini F, Mengoni C, Manara S, et al. Precise phylogenetic

analysis of microbial isolates and genomes from metagenomes using PhyloPhlAn 3.0. Nat

Commun. 2020;**11**: 2500 doi: 10.1038/s41467-020-16366-7

48.   Figtree. https://github.com/rambaut/figtree.

49.   Pallen MJ, Ponsero AJ, Telatin A, Moss CJ, Baker D, et al. Faecal metagenomes of

great tits and blue tits provide insights into host, diet, pathogens and microbial biodiversity.

Access Microbiol. 2025;**7**:   doi: 10.1099/acmi.0.000910.v3

50.   Liu J, Li XY, Song WT, Zeng XX, Li H, et al. Metagenomic analysis of environmental

samples from wildlife rescue station at Poyang Lake, China. Biomed Environ Sci. 2023;**36**:

595-603 doi: 10.3967/bes2023.087

51.   Wang Y, Zhai J, Tang B, Dong Y, Sun S, et al. Metagenomic comparison of gut

communities between wild and captive Himalayan griffons. Front vet sci. 2024;**11**: 1403932

doi: 10.3389/fvets.2024.1403932

52.   Mitchell AL, Scheremetjew M, Denise H, Potter S, Tarkowska A, et al. EBI

Metagenomics in 2017: enriching the analysis of microbial communities, from sequence

reads to assemblies. Nucleic Acids Res. 2017;**46**: D726-D35 doi: 10.1093/nar/gkx967

53.   Xie F, Jin W, Si H, Yuan Y, Tao Y, et al. An integrated gene catalog and over 10,000

metagenome-assembled genomes from the gastrointestinal microbiome of ruminants.

Microbiome. 2021;**9**: 137 doi: 10.1186/s40168-021-01078-x

54.   Bjørnsen MB, Valerón NR, Vásquez DP, Velasco EM, Hansen AJ, Hauptmann AL.

Microbiota in the ptarmigan intestine—An Inuit delicacy and its potential in popular cuisine.

PLOS ONE. 2024;**19**: e0305317 doi: 10.1371/journal.pone.0305317

55.   Gilroy R, Ravi A, Getino M, Pursley I, Horton DL, et al. Extensive microbial diversity

within the chicken gut microbiome revealed by metagenomics and culture. PeerJ. 2021;**9**:

e10941 doi: 10.7717/peerj.10941

56.   Beresford-Jones BS, Forster SC, Stares MD, Notley G, Viciani E, et al. The Mouse

Gastrointestinal Bacteria Catalogue enables translation between the mouse and human gut

microbiotas via functional mapping. Cell Host Microbe. 2022;**30**: 124-38.e8 doi:

10.1016/j.chom.2021.12.003

57.   Rosshart SP, Herz J, Vassallo BG, Hunter A, Wall MK, et al. Laboratory mice born to

wild mice have natural microbiota and model human immune responses. Science. 2019;**365**:

eaaw4361 doi: doi:10.1126/science.aaw4361

58.   Lesker TR, Durairaj AC, Gálvez EJC, Lagkouvardos I, Baines JF, et al. An integrated

metagenome catalog reveals new insights into the murine gut microbiome. Cell Rep.

2020;**30**: 2909-22.e6 doi: 10.1016/j.celrep.2020.02.036

59. Riva A, Sahin E, Volpedo G, Petretto A, Lavarello C, et al. Identification of an

epilepsy-linked gut microbiota signature in a pediatric rat model of acquired epilepsy.

Neurobiol Dis. 2024;**194**: 106469 doi: 10.1016/j.nbd.2024.106469

60. Zheng J, Sun Q, Zhang M, Liu C, Su Q, et al. Noninvasive, microbiome-based

diagnosis of inflammatory bowel disease. Nat Med. 2024;**30**: 3555-67 doi: 10.1038/s41591-

024-03280-4

61. Blyton MDJ, Soo RM, Hugenholtz P, Moore BD. Maternal inheritance of the koala gut

microbiome and its compositional and functional maturation during juvenile development.

Environ Microbiol. 2022;**24**: 475-93 doi: 10.1111/1462-2920.15858

62. Mukherjee A, Lordan C, Ross RP, Cotter PD. Gut microbes from the phylogenetically

diverse genus *Eubacterium* and their various contributions to gut health. Gut Microbes.

2020;**12**: 1802866 doi: 10.1080/19490976.2020.1802866

63. Kim CY, Lee M, Yang S, Kim K, Yong D, et al. Human reference gut microbiome

catalog including newly assembled genomes from under-represented Asian metagenomes.

Genome Med. 2021;**13**: 134 doi: 10.1186/s13073-021-00950-7

64. Wylensek D, Hitch TCA, Riedel T, Afrizal A, Kumar N, et al. A collection of bacterial

isolates from the pig intestine reveals functional and taxonomic diversity. Nat Commun.

2020;**11**: 6389 doi: 10.1038/s41467-020-19929-w

65. IJdema F, De Smet J, Crauwels S, Lievens B, Van Campenhout L. Meta-analysis of

larvae of the black soldier fly (*Hermetia illucens*) microbiota based on 16S rRNA gene

amplicon sequencing. FEMS Microbiol Ecol. 2022;**98**: fiac094 doi: 10.1093/femsec/fiac094

66. Salgado JFM, Hervé V, Vera MAG, Tokuda G, Brune A. Unveiling lignocellulolytic

potential: a genomic exploration of bacterial lineages within the termite gut. Microbiome.

2024;**12**: 201 doi: 10.1186/s40168-024-01917-7

67. Bowerman KL, Knowles SCL, Bradley JE, Baltrūnaitė L, Lynch MDJ, et al. Effects of

laboratory domestication on the rodent gut microbiome. ISME Commun. 2021;**1**: 49 doi:

10.1038/s43705-021-00053-9

68.   Betancur-Murillo CL, Aguilar-Marín SB, Jovel J. *Prevotella*: A key player in ruminal

metabolism. 2022;**11**: 1 doi: 10.3390/microorganisms11010001

69.   Murray AH, Iason GR, Stewart C. Effect of simple phenolic compounds of heather

(*Calluna vulgaris*) on rumen microbial activity *in vitro*. J Chem Ecol. 1996;**22**: 1493-504 doi:

10.1007/bf02027727

70.   Stogios PJ, Savchenko A. Molecular mechanisms of vancomycin resistance. Protein

Sci. 2020;**29**: 654-69 doi: 10.1002/pro.3819

71.   Li J, Li X-Z. Antimicrobial resistance and drug efflux pumps in *Helicobacter*. In: Li X-

Z, Elkins CA, Zgurskaya HI, editors. Efflux-mediated antimicrobial resistance in bacteria:

Mechanisms, regulation and clinical implications: Springer International Publishing; 2016. p.

489-513.

72.   Alauzet C, Lozniewski A, Marchandin H. Metronidazole resistance and nim genes in

anaerobes: A review. Anaerobe. 2019;**55**: 40-53 doi: 10.1016/j.anaerobe.2018.10.004

73.   Tang B, Wang Y, Dong Y, Cui Q, Zeng Z, et al. The catalog of microbial genes and

metagenome-assembled genomes from the gut microbiomes of five typical crow species on

the Qinghai–Tibetan Plateau. Microorganisms. 2024;**12**: 2033 doi:

10.3390/microorganisms12102033

74.   Eisenhofer R, Odriozola I, Alberdi A. Impact of microbial genome completeness on

metagenomic functional inference. ISME Commun. 2023;**3**: 12 doi: 10.1038/s43705-023-

00221-z

75.   Gao Y, Luo H, Lyu H, Yang H, Yousuf S, et al. Benchmarking short-read

metagenomics tools for removing host contamination. GigaScience. 2025;**14**: giaf004 doi:

10.1093/gigascience/giaf004