Theory on the accurate estimation of Michaelis-Menten enzyme kinetic parameters from steady state and progress curve datasets

Rajamanickam Murugan

Department of Biotechnology, Indian Institute of Technology Madras
Chennai, India. Email: rmurugan@gmail.com

ABSTRACT

We show that neither pure steady state nor pure progress curve analysis yield reliable estimate of enzyme K_M values since these methods are valid only at specific timescales and also use different type of datasets. All the currently proposed validity conditions of these methods assume *a priori* knowledge on K_M. Hence, there is no way to check whether the obtained K_M from a given dataset is a reliable estimate or not. Here we propose an integrated approach in which the same time course dataset will be analysed both in the progress curve as well as steady state perspectives at different reaction timescales across replications and substrate concentrations. Our theory shows that there exists an optimum reaction time at which the error in the estimation of K_M using various progress curve and steady state methods show the least possible value so that the coefficient of variation of the median K_M values obtained across various methods attains a minimum. Using detailed stochastic simulations, we confirm that the K_M value obtained with minimum coefficient of variation across various methods is actually the reliable estimate that is close to the original K_M value. We further show that using multiple nonlinear regression methods, the type of inhibition viz. competitive, uncompetitive and mixed can be accurately classified apart from obtaining the accurate inhibitor constants and IC_{50} values from Dixon type average velocity steady state datasets.

KEY WORDS

Michaelis-Menten enzyme kinetics, K_M, enzyme kinetic parameters, accurate estimation, nonlinear least square fitting.

Introduction

Enzymes catalyse several crucial biochemical reactions especially at physiologically relevant timescales and environmental conditions [1-3]. The mechanistic view of single substrate enzyme catalysis can be well described by the Michaelis-Menten scheme (MMS) [4, 5]. In this scheme (**Fig. 1A**), the enzyme first binds the substrate to form an enzyme-substrate complex which reduces the activation energy barrier of substrate to product conversion. Subsequently, the substrate will be converted into the respective product and released. The MMS scheme is characterized by three different rate constants viz. diffusion-controlled bimolecular forward k_1 (1/mol/lit/second), unimolecular reverse k_{-1} (1/second) and product formation rate constants k_2 (1/second). These rate constants along with the initial enzyme level (e_0) are generally summarized by two important enzyme kinetic parameters viz. $K_M = (k_{-1} + k_2) / k_1$ (mol/lit) and $v_{max} = k_2\, e_0$ (mol/lit/s). Here k_2 is also termed as k_{cat} and the ratio $k_E = k_{cat} / K_M$ is defined as the catalytic efficiency of the enzyme. When $k_2 \gg k_{-1}$, then k_E reaches the diffusion-controlled rate limit k_1.

Exact closed form analytical solution to the nonlinear set of ODEs corresponding to MMS scheme is not known though the integral solution can be expanded in terms of ordinary [6] and singular perturbation series [7] or over slow manifolds [8, 9]. Singular perturbation expansions always yield inner and outer solutions corresponding to the pre-steady state and post-steady state regimes which should be combined via proper matching at the temporal boundary layer [7, 10-15] to obtain the complete solution. Whereas, ordinary perturbation series can give the complete solution that is valid for both pre- and post-steady state regimes [6]. Obtaining (K_M, k_{cat}, v_{max}) from the steady state and progress curve experimental datasets is essential to characterize an enzyme. Since complete closed form analytical solution to MMS is unknown, one needs to rely on the widely used standard quasi steady state approximation (sQSSA) of the reaction velocity $v = dp/dt$ as $v = v_{max}\, s / (K_M + s)$ that is valid only when $k_2 \ll (k_1\, s_0)$ [16]. Here p represents the product concentration. When $e_0 \ll s_0$, then one can use the stationary reactant assumption to further simplify as $v \cong v_0 = v_{max}\, s_0 / (K_M + s_0)$ which connects K_M, v_{max} and initial substrate concentration s_0. Nonlinear least square fitting of the data on (v, s_0) over original hyperbolic function or linearization techniques such as Lineweaver-Burk on ($1/v$, $1/s_0$) and Eadie-Hofstee on (v, v/s_0) [17] can be used to obtain the estimates of kinetic parameters [18, 19]. Several statistical analysis procedures were also proposed [20, 21] to compute the error on these parameter estimates.

Enzymes of pathogenic organisms are important targets for drug molecules to control various diseases. Accurate estimation of various MMS parameters especially K_M along with K_I of a competitive inhibitor is essential to compute the IC50 values of drug molecules against a given target enzyme [22-25]. In steady state experiments, a fixed amount of enzyme will be incubated with a series of different substrate concentrations (s_0) for fixed amount of time (t_r) at constant temperature and pH conditions from which the *average product formation velocity* ($\bar{v}$) will be calculated by the ratio = total product formed during incubation time / total incubation time (t_r). Generally, t_r will be fixed approximately at 1 % depletion of s_0 however without any theoretical basis. Under sQSSA conditions with $e_0 \ll s_0$, it will be approximately assumed that the system evolves with constant reaction velocity $v_0 = v_{max}\, s_0 / (K_M + s_0)$ after a short pre-steady state timescale $t_c \ll t_r$ so that the total product formed over time t_r will be approximately equal to ($p = v_0\, t_r$) and therefore one sets the average

reaction velocity $\bar{v} \cong \bar{v}_{sQSSA} = (p/t_r) \cong v_0$. This equation is being used widely across enormous volumes of literature to estimate K_M and v_{max} from the dataset on ($\bar{v}$, s_0) obtained from steady state experiments. To estimate the inhibition constants K_I, this average velocity experiments will be conducted at different inhibitor concentrations i_0, and the data on ($\bar{v}$, s_0, i_0) will be used to classify the type of inhibition (i.e. competitive, uncompetitive or mixed type) and estimate v_{max}, K_M and K_I using Dixon plots [26] or sequential or simultaneous nonlinear least square fitting procedures [27, 28]. Using these parameters IC_{50} value of the inhibitory drug will be computed using Cheng-Prusoff equation [23]. Here $\bar{v}$ is the average velocity = product formed in a given total reaction time / total reaction time.

Detailed studies proposed the essential and sufficient conditions for the validity of the approximation $\bar{v} \cong v_0$ viz. (a) $k_2 \ll (k_1 s_0)$ [16] (b) $e_0 \ll s_0$ [29, 30] (c) ($e_0 / (s_0 + K_M)$) $\ll 1$ [29, 30]. However, except (b), conditions (a) and (c) strictly require *a priori* knowledge on k_1, k_2 and K_M. Otherwise, one needs to set very high values of s_0 along with condition (b) so that all the conditions (a-c) are satisfied. Although the inequality (b) is possible under *in vitro* laboratory conditions, there are several situations such as single molecule enzyme kinetics [31] and other *in vivo* conditions where one cannot manipulate the ratio of substrate to enzyme concentrations much. Setting high s_0 along with infinitesimal $e_0 \ll s_0$ would eventually drives $\bar{v}$ towards infinitesimal values below the background noise level which results in noisy experimental data on ($\bar{v}$, s_0) that leads to enormous amount of error in the estimation of K_M. Several recent studies [32-35] reported this issue and also put forth alternate methods such as total QSSA (tQSSA) to obtain accurate K_M values from such experimental datasets.

The source datasets for the estimation of K_M will be either time dependent progress curve type [36] or steady state average velocity type. Here progress curve data on (p, s, t) contains information on both pre- as well as post steady state dynamics that in turn depends on the timescale of experimental data collection. Whereas, steady state experimental setups generally capture the details of only the post-steady state regime. Progress curve methods can yield accurate estimates of K_M only when the data collection done at those timescale regimes with maximum change in the curvature of the kinetic trajectory [37]. Further, the role of total reaction time t_r is not considered anywhere in the sQSSA calculations i.e., $\bar{v}_{sQSSA} \cong v_0$ though the final results strongly depend on t_r. In this context, integrated form of sQSSA (isQSSA) is more superior than the general sQSSA since it also considers t_r.

Since the conditions of validity of various methods like sQSSA, isQSSA and progress curve analysis are different from each other, K_M values of the same enzyme obtained from each of these methods under identical conditions using the same source dataset will not be consistent. Further, absence of *a priori* knowledge on K_M and other kinetic parameters will eventually introduces uncertainty in setting up the appropriate experimental conditions to minimize the error in the estimation of these parameters using sQSSA methods with stationary reactant assumption or progress curve methods. Apart from the sequential and simultaneous nonlinear fitting, analysis of Dixon type inhibition datasets on ($\bar{v}$, s_0, i_0) warrants multiple nonlinear regression methods (MNLR) to obtain reliable estimates of IC_{50} of drug molecules. In this article, we develop an integrated theoretical and computational approach

which can produce consistent estimate of v_{max}, K_M, K_I and IC50 upon considering all these uncertainty aspects.

Theoretical Methods
Single substrate Michaelis-Menten kinetics
The nonlinear coupled differential rate equations corresponding to MM **Scheme A** shown in **Fig. 1** can be written as follows [16].

$$\frac{de}{dt} = -k_1 es + (k_{-1} + k_2)x \,. \tag{1.1}$$

$$\frac{ds}{dt} = -k_1 es + k_{-1}x. \tag{1.2}$$

$$\frac{dx}{dt} = k_1 es - (k_{-1} + k_2)x \,. \tag{1.3}$$

$$\frac{dp}{dt} = v = k_2 x \,. \tag{1.4}$$

Here k_1 (1/mol/lit/second), k_{-1} (1/second) and k_2 (1/second) are the rate constants corresponding to bimolecular enzyme-substrate complex formation, dissociation and product formation respectively. Further, (*e*, *s*, *x*, *p*) (mol/lit) are the concentrations of enzyme, substrate, enzyme-substrate complex, product respectively and v (mol/lit/second) is the reaction velocity. The initial conditions at $t \to 0$ are $(e, s, x, v, p) = (e_0, s_0, 0, 0, 0)$ and the reaction trajectory ends at $(e, s, x, v, p) = (e_0, 0, 0, 0, s_0)$ as $t \to \infty$. The dynamical variables (*e*, *s*, *x*, *p*) obey the mass conservation laws viz. $s = (s_0 - x - p), e = (e_0 - x)$ where one can also replace $x = v/k_2$. Here the required enzyme kinetic parameters are defined as $K_M = (k_{-1} + k_2)/k_1$ and $v_{max} = k_2 e_0$. From the definition of K_M, one can conclude that accurate estimation of K_M requires detailed data points representing both pre- and post-steady state dynamics. To simplify the system of **Eqs. 1,** we first introduce the following set of scaling transformations [16].

$$\tau = k_2 t; \ X = \frac{x}{e_0}; \ P = \frac{p}{s_0}; \ S = \frac{s}{s_0}; \ E = \frac{e}{e_0}; \ V = \frac{dP}{d\tau} \tag{2}$$

In **Eqs. 2**, the variables (e, s, x, p) are normalized so that $0 \le (E, S, X, P) \le 1$ with the mass conservations laws $S = 1 - \varepsilon X + P$, $E + X = 1$. We define $V = \varepsilon X$ so that $V + P + S = 1$ where $\varepsilon = \left(\frac{e_0}{s_0}\right)$. **Eqs. 1** can be rescaled in multiple ways at different variable spaces by the following set of dimensionless nonlinear ODEs [16].

$$\eta \frac{d^2 P}{d\tau^2} + \alpha \frac{dP}{d\tau} - \varepsilon(1 - P) = \frac{dP}{d\tau}\left(\frac{dP}{d\tau} + P\right). \tag{3.1}$$

$$\eta \frac{dV}{d\tau} + \alpha V - \varepsilon(1 - P) = V(V + P); \ \frac{dP}{d\tau} = V. \tag{3.2}$$

$$\eta \frac{dV}{d\tau} + \alpha V - \varepsilon(V + S) = V(1 - S); \ \eta \frac{dS}{d\tau} = (-(\varepsilon - V)S + \kappa V). \tag{3.3}$$

$$(-(\varepsilon - V)S + \kappa V)\frac{dV}{dS} + \alpha V - \varepsilon(V + S) = V(1 - S). \tag{3.4}$$

$$\eta V \frac{dV}{dP} + \alpha V - \varepsilon(1 - P) = V(V + P). \tag{3.5}$$

$$\frac{dP}{dS} = -\frac{\eta(1 - P - S)}{(\varepsilon - (1 - P - S))S - \kappa(1 - P - S)} \,. \tag{3.6}$$

Here **Eq. 3.1** describes the MMS dynamics in (P, τ) space, **Eqs. 3.2** in (V, P, τ), **Eqs. 3.3** (V, S, τ), **Eq. 3.4** in (V, S), **Eq. 3.5** in (V, P) and **Eq. 3.6** in (P, S) spaces. Various dimensionless parameters in **Eqs. 3** are defined as follows.

$$\varepsilon = e_0/s_0 \; ; \; \eta = k_2/k_1 s_0 \; ; \; \kappa = k_{-1}/k_1 s_0 \; ; \; \mu = \eta + \kappa = K_M/s_0 \; ; \; \alpha = 1 + \varepsilon + \mu \; . \qquad [4]$$

When $\eta \to 0$, then **Eq. 3.4** results in sQSSA with the velocity v_{sQSSA} in the original (v, s) space. Similarly, **Eq. 3.5** results in tQSSA with the velocity v_{tQSSA} in the (v, p) space which can be written in terms of the original dynamical variables as follows.

$$v_{sQSSA} \cong \frac{v_{max}S}{K_M+s} \; ; \; v_{tQSSA} \cong \frac{1}{2s_0}\Big(K_M + e_0 + s_0 - p - $$
$$s_0\sqrt{e_0^2 + 2e_0(p + K_M - s_0) + (p - K_M - s_0)^2}\Big). \qquad [5]$$

We denote **Eqs. 5** as η-approximations since they are valid only when $\eta \to 0$. The main flaw of these sQSSA as well as tQSSA methods is that the left-hand sides of **Eqs. 5** are still time dependent quantities along with time dependent s and p on the right-hand sides. That is to say, v_{sQSSA} and v_{tQSSA} are tangents of progress curve in (p, t) space at a given time point and not the time averaged velocities as used in most of the steady state experiments. To derive equations which are useful to fit steady state experimental data, one needs to further set $\varepsilon \to 0$ in **Eqs. 5**. This leads to the stationary reactant assumption $s \cong s_0 - p \cong s_0$ that results in the constant velocity as $v_{sQSSA} \cong v_{tQSSA} \cong v_0 = \frac{v_{max}s_0}{K_M+s_0}$ since $lim_{\varepsilon \to 0} v_{tQSSA} \cong v_{sQSSA}$. In this scenario, one can conduct the average steady state velocity experiments only within the reaction timescales in which these conditions are valid.

Upon setting $\varepsilon \to 0$ in **Eq. 3.6**, one finds that $\frac{dP}{dS} \cong \frac{\eta}{S+\kappa}$ in the pre-steady state regime of (P, S) space. Solving this ODE for the initial condition P = 0 for S = 1, we obtain $P \cong \eta ln\left(\frac{S+\kappa}{1+\kappa}\right)$. Since $S \cong 1$ when $(\eta, \varepsilon) \to (0,0)$, one finds that $P \cong 0$ and $V \cong 1 - S$ in the pre-steady state regime ($p \cong 0$ and $v \cong k_2(s_0 - s)$ in terms of original variables). Since $s \cong s_0$ when $s_0 \gg e_0$, one finds that $v \cong 0$ for infinitesimal values of k_2 near the steady state regime. The appropriate working reaction timescales for the average steady state velocity experiments can be derived as follows. Using the transformation rule $P = \varepsilon\eta^2 U$, **Eq. 3.1** can be rewritten as the following ordinary perturbation equation with the scaling parameter $\phi = \eta\varepsilon$ as follows.

$$\frac{d^2U}{d\tau^2} + \frac{\alpha}{\eta}\frac{dU}{d\tau} + \frac{\varepsilon}{\eta}U - \frac{1}{\eta^3} = \phi\frac{dU}{d\tau}\left(\frac{dU}{d\tau} + U\right) ; U(0) = 0; \; U'(0) = 0. \qquad [6]$$

When $\varepsilon \to 0$, then one arrives at the following approximations from **Eqs. 6** using the definition of dimensionless velocity $V = \varepsilon\eta^2 F$.

$$\frac{d^2U}{d\tau^2} + \frac{(1+\mu)}{\eta}\frac{dU}{d\tau} - \frac{1}{\eta^3} \cong 0; \; \frac{dU}{d\tau} = F; \; \frac{dF}{d\tau} + \frac{(1+\mu)}{\eta}F - \frac{1}{\eta^3} \cong 0; F(0) = 0 \; . \qquad [7]$$

We denote **Eqs. 7** as ε-approximations. Solving **Eqs. 7** for (U, F) in terms of τ and then reverting back to (P, V) and subsequently transforming back in to the original variables (p, v, t), one arrives at the following ε-approximations.

$$p_\varepsilon \cong (v_0 t - v_\varepsilon t_c) \cong v_0(t - t_c); \quad v_\varepsilon \cong v_0\left(1 - exp\left(-\frac{t}{t_c}\right)\right); \quad t_c \cong \frac{1}{k_1(K_M + s_0)}. \qquad [8]$$

Here $v_0 = \frac{v_{max}s_0}{K_M + s_0}$ is the sQSSA with stationary reactant assumption i.e., $s \cong s_0$. When $t \gg t_c$, then one finds that $p_\varepsilon \cong v_0(t - t_c)$. From **Eqs. 8** one can also obtain $s_\varepsilon \cong s_0 - v_0 t - v_\varepsilon(1/k_2 - t_c)$ which follows from the mass conservation law $s_\varepsilon \cong s_0 - p_\varepsilon - v_\varepsilon/k_2$. The subscript ε in p_ε and s_ε corresponds to the ε-approximations. The pre-steady state timescale in **Eq. 8** is t_c. When $t \gg t_c$, then one finds that $s_\varepsilon \cong \left(s_0 - v_0(1/k_2 - t_c)\right) - v_0 t$. When $t = t_c$, then the steady state concentration of substrate can be approximated as $s_{\varepsilon c} \cong s_0(1 - e_0/(K_M + s_0))$. Clearly, the stationary reactant assumption [38] $s_{\varepsilon c} \cong s_0$ will be true only when $(e_0/(K_M + s_0)) \ll 1$ which will be eventually true when $\varepsilon \to 0$. When the reaction time t_r is such that $t_r < t_c$, then $p_\varepsilon \cong 0$ in the pre-steady state regime (infinitesimal quantity especially for $\varepsilon \to 0$). When $t_r \gg t_c$, then one finds the linear product evolution regime with reaction time t_r as $p_\varepsilon \cong v_0(t_r - t_c)$ along with the constant reaction velocity $v_\varepsilon \cong v_0$. However, this linear dependence of p_ε on reaction timescale will break down towards the end of reaction since the substrate concentration starts to decline towards zero beyond this timescale t_m. This approximately occurs when $t_r \geq t_m \cong \frac{s_0}{v_0} = \left(\frac{K_M + s_0}{k_2 e_0}\right)$. Since the product concentration evolves linearly with time as $v_0 t$ in the post-steady state regime, we find that $t_m \cong \frac{s_0}{v_0}$. These timescale behaviors are demonstrated using numerically computed integral trajectories of **Eqs. 1** in **Fig. 1B** and **1C**. Clearly, one finds that $t_m \propto s_0$, $lim_{s_0 \to \infty} t_c \to 0$, $lim_{s_0 \to \infty} t_m \to \infty$ and $lim_{s_0 \to \infty} v \to v_{max}$. Further, one can define the timescale separation ratio χ as follows.

$$\chi = \frac{t_m}{t_c} = \frac{k_1(K_M + s_0)^2}{k_2 e_0} = \frac{(1+\mu)^2}{\eta\varepsilon} = \frac{(1+\mu)^2}{\phi}. \qquad [9]$$

Here χ is inversely proportional to the perturbation parameter ϕ. The successfulness of sQSSA methods strongly depends on extent of timescale separation that is decided by the condition $\chi \gg 1$ [16, 30] which also means that $\phi = \eta\varepsilon \ll (1 + \mu)^2$. The linear dependence of product formation on time will be valid only when the reaction time falls in the range $t_c \ll t_r < t_m$ where one can conduct average reaction velocity experiments using $\bar{v} \cong \bar{v}_{sQSSA} = v_0$. However, fitting of steady state data on ($\bar{v}$, s₀) with v_0 will be biased towards v_max over K_M since there is not enough datapoints from the pre-steady state regime. When $t_r < t_m$, then one can also use **Eqs. 8** to conduct average velocity experiments in the following form.

$$\bar{v}_\varepsilon \cong \frac{p_\varepsilon}{t_r} \cong \frac{v_{max}s_0}{K_M + s_0}\left(1 - \frac{t_c}{t_r}\left(1 - exp\left(-\frac{t_r}{t_c}\right)\right)\right) = v_0\left(1 - \frac{t_c}{t_r}\left(1 - exp\left(-\frac{t_r}{t_c}\right)\right)\right). \qquad [10]$$

Here p_ε is the total product formed at the end of reaction time t_r as in **Eq. 8** and $\bar{v}_\varepsilon$ is the average velocity obtained from the steady state experiments. Unlike $\bar{v}_{sQSSA}$, $\bar{v}_\varepsilon$ also considers the nonlinearity of the pre-steady state regime. The conditions for the validity of **Eq. 10** are

$\varepsilon \to 0$ and $t_r < t_m$. When $t_m > t_r \gg t_c$, then $\bar{v}_\varepsilon \to \bar{v}_{sQSSA} \cong v_0$. Here, $\bar{v}_{sQSSA} \cong v_0$ directly connects the average reaction velocity with the total substrate concentration s_0. Clearly, the conditions for the validity of the widely used $\bar{v}_{sQSSA} \cong v_0$ are viz. $(\eta, \varepsilon) \to (0,0)$ and $t_c \ll t_r < t_m$. Nonlinear least square fitting of the data on $(\bar{v}, s_0)$ for a given reaction time t_r with **Eq. 10** can directly yield the parameters (K_M, v_{max} and k_1). However, such fitting procedures favour accurate estimation of v_{max} over (K_M, k_1) since they mainly rely on the datapoints from the pre-steady state timescale regime which is actually a transient one compared to the post-steady state timescale. To overcome this issue, we will fix k_1 in the definition of $\bar{v}_\varepsilon$ with the median k_1 obtained from the progress curve methods in the later sections. To minimize the error in the estimation of K_M one needs to set t_r such that it is much lesser than the maximum reaction time t_m corresponding to the lowest s_0 value used in the entire experimental setup. When $\eta \to 0$, then one finds as in **Eqs. 5** that $V \cong \frac{\varepsilon S}{\mu + S}$. When $\varepsilon \to 0$, then one finds that $\frac{dP}{d\tau} \cong -\frac{dS}{d\tau}$ in the post-steady state regime and therefore one obtains $-\frac{dS}{d\tau} \cong \frac{\varepsilon S}{\mu + S}$ (or $\frac{dP}{d\tau} \cong \frac{\varepsilon(1-P)}{\mu + (1-P)}$). Upon solving this nonlinear first order ODE with the initial condition S = 1 at $\tau = 0$, and then reverting back to the original dynamical variables, one finds the following implicit solution.

$$\bar{v}_{isQSSA} = \frac{s_0 - s}{t_r} \cong \frac{p}{t_r} \cong v_{max} + \frac{K_M}{t_r} ln\left(\frac{s}{s_0}\right) \cong v_{max} + \frac{K_M}{t_r} ln\left(\frac{s_0 - p}{s_0}\right). \qquad [11]$$

Eqs. 10 is another form of integrated rate equation [35] that directly connects the average reaction velocity $\bar{v}_{isQSSA}$ with the enzyme kinetic parameters and reaction time t_r. Data on $(\bar{v}, p, s, s_0)$ at a given reaction time t_r can be directly used to perform linear least square fitting with **Eq. 11** i.e., $\bar{v}_{isQSSA}$ versus $\frac{1}{t_r} ln\left(\frac{s_0 - p}{s_0}\right)$ (type I) or $\frac{1}{t_r} ln\left(\frac{s}{s_0}\right)$ (type II) to obtain K_M and v_{max}. In the later sections, we will show that type II form gives more accurate estimate of K_M than type I. The conditions for the validity of **Eq. 11** are $(\eta, \varepsilon) \to (0,0)$ and $t_c < t_r < t_m$. One can also integrate $v_{sQSSA} \cong -\frac{ds}{dt} \cong \frac{v_{max}s}{K_M + s}$ given in **Eqs. 5** under the conditions $\varepsilon \to 0$ for the initial condition s = s_0 at t = 0 and derive the following expression for the average reaction velocity.

$$\bar{v}_{csQSSA} \cong \frac{p_{csQSSA}}{t_r} = \frac{1}{t_r}\left(s_0 - s_{csQSSA} - \frac{v_{csQSSA}}{k_2}\right); S_{csQSSA} \cong K_M W\left(\frac{s_0}{K_M} exp\left(\frac{s_0 - v_{max}t_r}{K_M}\right)\right). [12]$$

In this equation, we have used the mass conservation law $p = (s_0 - s - v/k_2)$ and $v_{csQSSA} = -\frac{ds_{csQSSA}}{dt} = \frac{v_{max}s_{csQSSA}}{K_M + s_{csQSSA}}$. When $\left(\frac{e_0}{K_M + s_{csQSSA}}\right) \ll 1$, then one finds from **Eqs. 12** that $\bar{v}_{csQSSA} \cong \frac{1}{t_r}\left(s_0 - s_{csQSSA}\right)$ where $p_{csQSSA} \cong \left(s_0 - s_{csQSSA}\right)$ which is an alternate form on **Eq. 11**. In **Eqs. 12**, W(x) is the Lambert W function [39, 40] which is the solution of y exp(y) = x for y. The conditions of validity of **Eqs. 12** is similar to **Eqs. 11**. We use **Eq. 11** for comparison purposes since **Eq. 11** is simple and computationally cheaper than **Eq. 12**. K_M and v_{max} can be obtained from the steady state datasets using any one of the expressions ($\bar{v}_{sQSSA}, \bar{v}_{isQSSA}, \bar{v}_\varepsilon, \bar{v}_{csQSSA}$) under appropriate validity conditions and timescales. However, except the condition on ε, all the others conditions (on η, t_c, t_m) require *a priori* knowledge on k_1, k_2 and K_M which are generally not available for unknown enzyme systems. In this scenario, one can perform direct non-linear least square fitting of the time course kinetic data on (p, t) or (s, t) with the system of nonlinear ODEs given in **Eqs. 1** to obtain k_1, k_{-1} and k_2. However, accuracy of such fitting procedure strongly depends on the detailed data points on both pre- and post-steady state

regimes that warrants progress curve data collection on sub-millisecond timescales apart from the steady state timepoint.

Michaelis-Menten kinetics with competitive inhibition

When $(\varepsilon_I, \eta_I, \eta_S, \varepsilon_S) \to (0,0,0,0)$ and $\delta = \frac{v_{max}(K_{MI}+e_0)}{u_{max}(K_{MS}+e_0)} \cong 1$, then earlier detailed theoretical studies revealed [22] the following approximations for the average velocities of the fully competitive inhibition **Scheme B** described in **Fig. 1**.

$$\bar{v} \cong v_0\left(1 - \frac{t_{cs}}{t_r}\left(1 - exp\left(-\frac{t_r}{t_{cs}}\right)\right)\right); \; t_{cs} \cong \frac{\left(1+\frac{i_0}{K_{MI}}\right)}{k_1\left(s_0+K_{MS}\left(1+\frac{i_0}{K_{MI}}\right)\right)} \quad\quad [13.1]$$

$$\bar{u} \cong u_0\left(1 - \frac{t_{ci}}{t_r}\left(1 - exp\left(-\frac{t_r}{t_{ci}}\right)\right)\right); \; t_{ci} \cong \frac{\lambda\left(1+\frac{s_0}{K_{Ms}}\right)}{k_i\left(i_0+K_{MI}\left(1+\frac{s_0}{K_{MS}}\right)\right)} \quad\quad [13.2]$$

In fully competitive inhibition, same enzyme will catalyse the conversion of two different substrates (s, i) in to their respective products (p, q) with reaction velocities (v = dp/dt, u = dq/dt). In these equations, various parameters as defined as follows.

$$K_{MS} = \frac{k_2+k_{-1}}{k_1}; K_{MI} = \frac{k_{-i}+k_3}{k_i}; \; v_{max} = k_2e_0; \; u_{max} = k_3e_0 \; . \quad\quad [13.2]$$

$$\eta_I = \frac{k_3}{k_i i_0}; \; \varepsilon_I = \frac{e_0}{i_0}; \eta_S = \frac{k_2}{k_1 s_0}; \; \varepsilon_S = \frac{e_0}{s_0}; \lambda = \frac{k_3}{k_2} = \frac{u_{max}}{v_{max}} \quad\quad [13.3]$$

When the total reaction time is such that $t_r \gg t_{cs}$ and $t_r \gg t_{ci}$ where (t_{cs}, t_{ci}) are the pre-steady state timescales, then **Eqs. 13.1-2** reduce to the following well-known steady state velocity equations.

$$\bar{v} \cong v_0 = \frac{v_{max}s_0}{s_0+K_{MS}\left(1+\frac{i_0}{K_{MI}}\right)}; \; \bar{u} \cong u_0 = \frac{u_{max}i_0}{i_0+K_{MI}\left(1+\frac{s_0}{K_{MS}}\right)} \quad\quad [13.4]$$

When $\delta \ll 1$ or $\delta \gg 1$, the validity of **Eqs. 13.4** will break down since the fully competitive inhibition system can exhibit multiple steady states [22] under such scenarios. Similarly for pure partial competitive inhibition given **Scheme C** of **Fig. 1**, one can derive the following average velocity expression under the conditions $(\varepsilon_S, \chi_I, \varepsilon_I) \to (0,0,0)$ where $\chi_I = \frac{k_2}{k_i i_0}$.

$$\bar{v} \cong \frac{v_{max}s_0}{s_0+K_{MS}\left(1+\frac{i_0}{K_I}\right)}; \; K_I = \frac{k_{-i}}{k_i} \quad\quad [14]$$

Similarly, for the pure uncompetitive inhibition given in Scheme C **of Fig. 1** one can derive the following average velocity expression.

$$\bar{v} \cong \frac{v_{max}s_0}{s_0\left(1+\frac{i_0}{K_Z}\right)+K_{MS}}; \; K_Z = \frac{k_{-z}}{k_z} \quad\quad [15]$$

Finally, for the mixed competitive inhibition where both competitive and uncompetitive inhibition scenarios operate as given in Scheme C of **Fig. 1**, one can derive the following combined average velocity expression.

$$\bar{v} \cong \frac{v_{max}s_0}{s_0\left(1+\frac{i_0}{K_Z}\right)+K_{MS}\left(1+\frac{i_0}{K_I}\right)} = f(s_0, i_0 | v_{max}, K_{MS}, K_Z, K_I) \qquad [16.1]$$

IC_{50} is the inhibitor concentration which is required to decrease 50% of the inhibitor free enzyme activity which can be straightforwardly calculated [23] as follows for the generalized mixed inhibition scenario by solving the following equation for i_0.

$$\frac{f(s_0,0|v_{max},K_{MS},K_Z,K_I)-f(s_0,i_0|v_{max},K_{MS},K_Z,K_I)}{f(s_0,0|v_{max},K_{MS},K_Z,K_I)} = \frac{1}{2}; \ IC_{50} = \frac{K_Z K_I(s_0+K_{MS})}{K_Z K_{MS}+K_I s_0} \qquad [16.2]$$

When the inhibition is non-competitive, then the inhibitor binds at the allosteric site of enzyme that is different from the substrate binding active-site. In such scenario, the average steady state velocity equation becomes as follows.

$$\bar{v} \cong \frac{v_{max}s_0}{\left(1+\frac{i_0}{K_I}\right)(s_0+K_{MS})} \qquad [17]$$

Eq. 17 is similar to **Eq. 16.1** with $K_Z = K_I$ and for non-competitive inhibition one finds that $IC_{50} = K_I$. For pure competitive inhibition one finds that $\lim_{K_Z\to\infty} IC_{50} = K_I(1 + s_0/K_{MS})$ and for pure uncompetitive inhibition one finds that $\lim_{K_I\to\infty} IC_{50} = K_Z(1 + K_{MS}/s_0)$.

Numerical Simulation

The stochastic version of nonlinear ODEs given in **Eqs. 1.1-1.4** can be numerically integrated using the following Euler type scheme on chemical Langevin equations [41, 42].

$$x_{n+1} = x_n + (k_1(e_0 - x_n)(s_0 - x_n - p_n) - (k_{-1} + k_2)x_n)\Delta t + \Gamma_{x,t}. \qquad [18.1]$$

$$\Gamma_{x,t} = \rho\xi_{p,t}\sqrt{(k_1(e_0 - x_n)(s_0 - x_n - p_n) - (k_{-1} + k_2)x_n)\Delta t}. \qquad [18.2]$$

$$p_{n+1} = p_n + k_2 x_n \Delta t + \rho\sqrt{k_2 x_n \Delta t}\xi_{p,t}. \qquad [18.3]$$

$$e_n = (e_0 - x_n); s_n = (s_0 - x_n - p_n); \langle\xi_{p,t}\rangle = 0; \langle\xi_{x,t}\rangle = 0. \qquad [18.4]$$

$$\langle\xi_{p,t}\xi_{p,t'}\rangle = \delta(t - t'); \langle\xi_{x,t}\xi_{x,t'}\rangle = \delta(t - t'); \langle\xi_{x,t}\xi_{p,t'}\rangle = 0. \qquad [18.5]$$

Here the initial conditions are $x = 0$, $s = s_0$, $e = e_0$, and $p = 0$ at $t = 0$, $\xi_{q,t}$ where q = (x, p) are delta correlated gaussian white noise terms and ρ is the additional noise control parameter. The time increment parameter Δt will be adjusted to capture both pre- as well as post-steady state dynamics. Setting $\rho = 0$, results in the deterministic trajectory of the MMS scheme for the given set of parameters and initial conditions. To check the consistency among the data fitting methods ($\bar{v}_{sQSSA}, \bar{v}_{isQSSA}, \bar{v}_\varepsilon$), several stochastic trajectories of **Eqs. 1** were generated with fixed t_r and e_0 and different s_0 values. Stochastic simulated datasets are posted at (DOI: 10.5281/zenodo.15094623) and the sample code implementations of ML algorithm are given in the **Supporting Materials**. To obtain the kinetic parameters k_1, k_{-2} and k_2 from the progress curve data with initial s_0 and e_0, we used nonlinear least square fitting [43] using Marquardt-Levenberg (ML) algorithm [44-47]. The computational workflow of ML scheme starts with

minimizing the function $L = \sum_{i=1}^{n}(f(t_i|r) - y_i)^2$ where $f(t|r)$ is the numerical solution to **Eqs. 1** with the parameter set $r = [k_1, k_{-1}, k_2]$, and y is the observed dataset where $(f, y) = (p, s)$. Upon expanding L in Taylor series around $r_u = [k_1^u, k_{-1}^u, k_2^u]$ and then truncating with the first order terms, one finds the following approximation.

$$L \cong \sum_{i=1}^{n}\left(\Delta y_i + \frac{\partial f(t_i|r_u)}{\partial k_1}(k_1^u - k_1) + \frac{\partial f(t_i|r_u)}{\partial k_1}(k_{-1}^u - k_{-1}) + \frac{\partial f(t_i|r_u)}{\partial k_1}(k_2^u - k_2)\right)^2. \quad [19]$$

In this equation $\Delta y_i = f(t_i|r_u) - y_i$. The conditions for the minimum with respect to r will be $\frac{\partial L}{\partial k_1} = 0$; $\frac{\partial L}{\partial k_{-1}} = 0$; $\frac{\partial L}{\partial k_2} = 0$ which can be explicitly written as follows.

$$\sum_{i=1}^{n}\left(\Delta y_i + \frac{\partial f(t_i|r_u)}{\partial k_1}(k_1^u - k_1) + \frac{\partial f(t_i|r_u)}{\partial k_{-1}}(k_{-1}^u - k_{-1}) + \frac{\partial f(t_i|r_u)}{\partial k_2}(k_2^u - k_2)\right)\frac{\partial f(t_i|r_u)}{\partial k_1} = 0. \quad [20.1]$$

$$\sum_{i=1}^{n}\left(\Delta y_i + \frac{\partial f(t_i|r_u)}{\partial k_1}(k_1^u - k_1) + \frac{\partial f(t_i|r_u)}{\partial k_{-1}}(k_{-1}^u - k_{-1}) + \frac{\partial f(t_i|r_u)}{\partial k_2}(k_2^u - k_2)\right)\frac{\partial f(t_i|r_u)}{\partial k_{-1}} = 0. \quad [20.2]$$

$$\sum_{i=1}^{n}\left(\Delta y_i + \frac{\partial f(t_i|r_u)}{\partial k_1}(k_1^u - k_1) + \frac{\partial f(t_i|r_u)}{\partial k_{-1}}(k_{-1}^u - k_{-1}) + \frac{\partial f(t_i|r_u)}{\partial k_2}(k_2^u - k_2)\right)\frac{\partial f(t_i|r_u)}{\partial k_2} = 0. \quad [20.3]$$

Eqs. 15.1-3 can be written in a matrix form [43] as $J^T\Delta y + (J^TJ)\Delta r = 0$. Definition of various terms in this matrix equation are given below.

$$J = \begin{bmatrix} \frac{\partial f(t_1|r_u)}{\partial k_1} & \frac{\partial f(t_1|r_u)}{\partial k_{-1}} & \frac{\partial f(t_1|r_u)}{\partial k_2} \\ \vdots & \vdots & \vdots \\ \frac{\partial f(t_n|r_u)}{\partial k_1} & \frac{\partial f(t_n|r_u)}{\partial k_{-1}} & \frac{\partial f(t_n|r_u)}{\partial k_2} \end{bmatrix}; \quad \Delta r = \begin{bmatrix} k_1^u - k_1 \\ k_{-1}^u - k_{-1} \\ k_2^u - k_2 \end{bmatrix}; \quad \Delta y = \begin{bmatrix} f(t_1|r_u) - y_1 \\ \vdots \\ f(t_n|r_u) - y_n \end{bmatrix}. \quad [21]$$

We compute the partial derivatives with respect to parameters $[k_1, k_{-1}, k_2]$ at time point t_i as follows where $f(t|r)$ is the numerical integrated solution to **Eqs. 1** with the parameter set $r = [k_1, k_{-1}, k_2]$.

$$\frac{\partial f(t_i|r_u)}{\partial k_1} = \frac{\partial f(t_i|[k_1^u, k_{-1}^u, k_2^u])}{\partial k_1} \cong \frac{f(t_i|[k_1^u+\alpha, k_{-1}^u, k_2^u]) - f(t_i|[k_1^u, k_{-1}^u, k_2^u])}{\alpha}. \qquad [22.1]$$

$$\frac{\partial f(t_i|r_u)}{\partial k_{-1}} = \frac{\partial f(t_i|[k_1^u, k_{-1}^u, k_2^u])}{\partial k_{-1}} \cong \frac{f(t_i|[k_1^u, k_{-1}^u+\alpha, k_2^u]) - f(t_i|[k_1^u, k_{-1}^u, k_2^u])}{\alpha}. \qquad [22.2]$$

$$\frac{\partial f(t_i|r_u)}{\partial k_2} = \frac{\partial f(t_i|[k_1^u, k_{-1}^u, k_2^u])}{\partial k_2} \cong \frac{f(t_i|[k_1^u, k_{-1}^u, k_2^u+\alpha]) - f(t_i|[k_1^u, k_{-1}^u, k_2^u])}{\alpha}. \qquad [22.3]$$

We set α to the lowest possible value depending on the computational accuracy requirement. In **Eqs. 16**, J is the Jacobian matrix and explicitly one finds the following.

$$J^T\Delta y = \begin{bmatrix} \sum_{i=1}^{n}\left(\Delta y_i \frac{\partial f(t_i|r_u)}{\partial k_1}\right) \\ \sum_{i=1}^{n}\left(\Delta y_i \frac{\partial f(t_i|r_u)}{\partial k_{-1}}\right) \\ \sum_{i=1}^{n}\left(\Delta y_i \frac{\partial f(t_i|r_u)}{\partial k_2}\right) \end{bmatrix}. \qquad [23]$$

$$J^T J = \begin{bmatrix} \sum_{i=1}^n \left(\frac{\partial f(t_i|r_u)}{\partial k_1} \frac{\partial f(t_i|r_u)}{\partial k_1} \right) & \sum_{i=1}^n \left(\frac{\partial f(t_i|r_u)}{\partial k_{-1}} \frac{\partial f(t_i|r_u)}{\partial k_1} \right) & \sum_{i=1}^n \left(\frac{\partial f(t_i|r_u)}{\partial k_2} \frac{\partial f(t_i|r_u)}{\partial k_1} \right) \\ \sum_{i=1}^n \left(\frac{\partial f(t_i|r_u)}{\partial k_1} \frac{\partial f(t_i|r_u)}{\partial k_{-1}} \right) & \sum_{i=1}^n \left(\frac{\partial f(t_i|r_u)}{\partial k_{-1}} \frac{\partial f(t_i|r_u)}{\partial k_{-1}} \right) & \sum_{i=1}^n \left(\frac{\partial f(t_i|r_u)}{\partial k_2} \frac{\partial f(t_i|r_u)}{\partial k_{-1}} \right) \\ \sum_{i=1}^n \left(\frac{\partial f(t_i|r_u)}{\partial k_1} \frac{\partial f(t_i|r_u)}{\partial k_2} \right) & \sum_{i=1}^n \left(\frac{\partial f(t_i|r_u)}{\partial k_{-1}} \frac{\partial f(t_i|r_u)}{\partial k_2} \right) & \sum_{i=1}^n \left(\frac{\partial f(t_i|r_u)}{\partial k_2} \frac{\partial f(t_i|r_u)}{\partial k_2} \right) \end{bmatrix} . \quad [24]$$

From **Eqs. 15-19**, one obtains the following Marquardt-Levenberg (ML) iterative scheme where λ is the ML parameter and $\lambda = 0$ represents the Newton's method [43].

$$r_{u+1} = r_u + \left(J^T J + \lambda \operatorname{diag}(J^T J) \right)^{-1} J^T \Delta y . \qquad [25]$$

The reduced sum of squares of deviation will be computed at the end of u^{th} iteration as $Z_u = \sum_{i=1}^n (y_{u,i} - y_i)^2$ where $y_{u,i} = f(t_i|r_u)$ is the predicted data point values for r_u which will be obtained by numerical integration of **Eqs. 1** using Euler type scheme given in **Eqs. 13** with ρ = 0. Here λ should be adjusted depending on changing pattern of Z_u over iterations. When $Z_u < Z_{u+1}$, then we set $\lambda \to \lambda \delta$. When $Z_u > Z_{u+1}$ then we set $\lambda \to \lambda/\delta$. Here the δ is the ML tuning parameter [44]. When $|Z_u - Z_{u+1}| < \theta$ where θ is the required tolerance limit, then we exit from iteration and the final parameters $\hat{r}$ will be obtained. For NLS fit of various datasets we used $\lambda = 10^{-2}$, $\delta = 2$ and $\theta = 10^{-6}$. the final sum of squares error (SSE) and the covariance matrix will be calculated as follows.

$$SSE = \sum_{i=1}^n (\hat{y}_i - y_i)^2 \; ; \; \hat{\sigma}^2 = \frac{SSE}{n-3}; \; Cov(\hat{r}) = (J^T J)^{-1} \hat{\sigma}^2. \qquad [26]$$

Upon computing the total sum of squares (SST), one can compute the final nonlinear regression coefficient R^2 as follows.

$$SST = \sum_{i=1}^n y_i^2 - \frac{1}{n} \left(\sum_{i=1}^n y_i \right)^2; \; R^2 = 1 - \frac{SSE}{SST} \qquad [27]$$

Here $\hat{y}_i = f(t_i|\hat{r})$ where $\hat{r} = [\hat{k}_1, \hat{k}_{-1}, \hat{k}_2]$ is the final fit parameter values. We use the final SSE and R^2 to compare the goodness of fit across trajectories with different s_0 and replications. Similar workflow was used to obtain K_M and v_{max} using $\bar{v}_\varepsilon$ as given in **Eq. 10** with the median of k_1 obtained from the progress curve analysis across different s_0 over several replications outlined in **Eqs. 14-21**. Linear least square fitting with **Eq. 11** [43] for $\bar{v}_{isQSSA}$ yields the required parameter estimates along with the errors.

Multiple nonlinear regression for competitive inhibition
In this section, we will outline the multiple nonlinear regression fitting procedure for the Dixon type steady state average velocity inhibition dataset on $(\bar{v}, s_0, i_0)$ over **Eqs. 13-17**. For example, let us consider the case of pure competitive inhibition i.e., $\bar{v} \cong f(s_0, i_0|K_{MS}, K_I, v_{max})$ as given in **Eq. 14**. For simplicity we use the notation $(a, b, c) = (K_{MS}, K_I, v_{max})$ for the parameters and $(s_{0,i}, i_{0,k}) = (x_i, y_k)$ for the concentrations of substrate and inhibitor. We denote the substrate levels $s_{0,i}$ where i = 1 to n and inhibitor levels $i_{0,k}$ where k = 1 to m and the corresponding average velocities as $\bar{v}_{i,k}$ = product formed in the reaction time t_r / total

reaction time t_r. With these settings one can define the chi-square function to be minimized as follows.

$$L = \sum_{i=1}^{n} \sum_{k=1}^{m} \left(\bar{v}_{i,k} - f(x_i, y_k | a, b, c) \right)^2 . \tag{28}$$

Here L needs to be minimized with respect to the parameters (a, b, c) along with two different independent random variables. In this equation, $\Delta \bar{v}_{i,k} = \bar{v}_{i,k} - f_{i,k}$ where $f_{i,k} = f(x_i, y_k | a_0, b_0, c_0)$ is the fit function. As in case of single variable nonlinear fitting procedure, we expand L in terms of Taylor's series around (a_0, b_0, c_0) and truncate the series with the first order term as follows.

$$\frac{\partial L}{\partial a} \cong -2 \sum_{i=1}^{n} \sum_{k=1}^{m} \left(\Delta v_{i,k} - \frac{\partial f_{i,k}}{\partial a}(a - a_0) - \frac{\partial f_{i,k}}{\partial b}(b - b_0) - \frac{\partial f_{i,k}}{\partial c}(c - c_0) \right) \frac{\partial f_{i,k}}{\partial a} . \tag{29.1}$$

$$\frac{\partial L}{\partial b} \cong -2 \sum_{i=1}^{n} \sum_{k=1}^{m} \left(\Delta v_{i,k} - \frac{\partial f_{i,k}}{\partial a}(a - a_0) - \frac{\partial f_{i,k}}{\partial b}(b - b_0) - \frac{\partial f_{i,k}}{\partial c}(c - c_0) \right) \frac{\partial f_{i,k}}{\partial b} . \tag{29.2}$$

$$\frac{\partial L}{\partial c} \cong -2 \sum_{i=1}^{n} \sum_{k=1}^{m} \left(\Delta v_{i,k} - \frac{\partial f_{i,k}}{\partial a}(a - a_0) - \frac{\partial f_{i,k}}{\partial b}(b - b_0) - \frac{\partial f_{i,k}}{\partial c}(c - c_0) \right) \frac{\partial f_{i,k}}{\partial c} . \tag{29.3}$$

Eqs. 24 can be summarized in the following matrix form.

$$\sum_{k=1}^{m} \left(J_k^T \Delta \bar{v}_k \right) - \left(\sum_{k=1}^{m} \left(J_k^T J_k \right) \right) \begin{bmatrix} (a - a_0) \\ (b - b_0) \\ (c - c_0) \end{bmatrix} = 0 \quad . \tag{30}$$

Various terms in **Eq. 25** are defined as follows.

$$J_k = \begin{bmatrix} \frac{\partial f_{1,k}}{\partial a} & \frac{\partial f_{1,k}}{\partial b} & \frac{\partial f_{1,k}}{\partial c} \\ \vdots & \vdots & \vdots \\ \frac{\partial f_{n,k}}{\partial a} & \frac{\partial f_{n,k}}{\partial b} & \frac{\partial f_{n,k}}{\partial c} \end{bmatrix} \quad . \tag{31}$$

$$\Delta \bar{v}_k = \begin{bmatrix} \bar{v}_{1,k} - f_{1,k} \\ \vdots \\ \bar{v}_{n,k} - f_{n,k} \end{bmatrix} ; J_k^T \Delta \bar{v}_k = \begin{bmatrix} \sum_{i=1}^{n} \left(\Delta \bar{v}_{i,k} \frac{\partial f_{i,k}}{\partial a} \right) \\ \sum_{i=1}^{n} \left(\Delta \bar{v}_{i,k} \frac{\partial f_{i,k}}{\partial b} \right) \\ \sum_{i=1}^{n} \left(\Delta \bar{v}_{i,k} \frac{\partial f_{i,k}}{\partial c} \right) \end{bmatrix} ; r = \begin{bmatrix} a \\ b \\ c \end{bmatrix} ; r_0 = \begin{bmatrix} a_0 \\ b_0 \\ c_0 \end{bmatrix} . \tag{32}$$

From **Eqs. 25**, one finds the Newton's iterative scheme for the multiple nonlinear fitting procedure to minimize L as follows.

$$\begin{bmatrix} a \\ b \\ c \end{bmatrix} = \begin{bmatrix} a_0 \\ b_0 \\ c_0 \end{bmatrix} + \left(\sum_{k=1}^{m} J_k^T J_k \right)^{-1} \left(\sum_{k=1}^{m} J_k^T \Delta \bar{v}_k \right) . \tag{33}$$

Clearly, before the inversion of $J^T J$ matrix, one needs to sum over the index k. Various partial derivative terms can be numerically computed as follows. This idea can be extended to any number of independent random variables apart from $(s_0$ and $i_0)$.

$$\frac{\partial f_{i,k}}{\partial a} = \frac{\partial f(x_i, y_k | a_0, b_0, c_0)}{\partial a} \cong \left[\frac{f(x_i, y_k | a + \Delta a, b, c) - f(x_i, y_k | a, b, c)}{\Delta a} \right]_{r=r_0} . \tag{34.1}$$

$$\frac{\partial f_{i,k}}{\partial b} = \frac{\partial f(x_i,y_k|a_0,b_0,c_0)}{\partial b} \cong \left[\frac{f(x_i,y_k|a,b+\Delta b,c)-f(x_i,y_k|a,b,c)}{\Delta b}\right]_{r=r_0} \quad . \tag{34.2}$$

$$\frac{\partial f_{i,k}}{\partial c} = \frac{\partial f(x_i,y_k|a_0,b_0,c_0)}{\partial c} \cong \left[\frac{f(x_i,y_k|a,b,c+\Delta c)-f(x_i,y_k|a,b,c)}{\Delta c}\right]_{r=r_0} \quad . \tag{34.3}$$

As in case of single variable NLS fitting described in **Eqs. 20**, Marquardt-Levenberg algorithm can be embedded in the **Eq. 26** to speed up the convergence. Sample datasets for competitive inhibitions schemes described by **Eqs. 13-17** were generated by adding gaussian white noise terms to the Dixon type steady state velocity datasets with appropriate noise control parameter ρ as $\bar{v} \to \bar{v} + \rho\xi$ where ξ is the random number drawn from the standard normal population N (0, 1).

Results and Discussion

There are several issues and limitations associated with the currently available steady state as well as progress curve methods to estimate the K_M and v_{max} of single substrate MM enzymes. First let us consider sQSSA with stationary reactant assumption ($\bar{v}_{sQSSA}$) which is valid only when $t_c \ll t_r < t_m$ and $(\eta, \varepsilon) \to (0,0)$. Since K_M values are not generally available for the unknown enzyme systems, to achieve these criteria, one needs to set higher values for s_0 than e_0 which will lead to high background fluctuations in the measured average reaction velocities. For example, let $\bar{v}_0 = (\bar{v}_{0,1}, \bar{v}_{0,2})$ correspond to respectively $s_0 = (s_{0,1}, s_{0,2})$. When $s_0 \to \infty$, then one finds the following.

$$lim_{s_0 \to \infty}|\bar{v}_{0,2} - \bar{v}_{0,1}| = lim_{s_0 \to \infty} v_{max}\left|\frac{s_{0,2}}{K_M+s_{0,2}} - \frac{s_{0,1}}{K_M+s_{0,1}}\right| \to 0. \tag{35}$$

Eq. 23 leads to high noise level in the observed average reaction velocities that in turn results in high level of uncertainty in the estimation of K_M. Similar effects are prevalent in the progress curve analysis using especially the trajectories on (p, t) space since $p \cong 0$ in the pre-steady state regime that leads to high level of background fluctuations in the measured values of p. This in turn results in high uncertainty in the estimation of K_M using the data on (p, t) which significantly depends on the pre-steady state dynamics. Clearly, the data on (p, t) can be used only in the steady state methods which mainly work well in the post-steady state regime. Detailed NLS fitting analysis of the data on (p, t) and (s, t) spaces in the presence as well as absence of noise suggests the following.

a) Progress curve methods using Marquardt-Levenberg algorithm outlined in the methods section works very well over data on (s, t) as well as (p, t) in the absence of noise i.e. $\rho = 0$ in **Eqs. 13** as demonstrated in **Figs. 2A** and **2B**. NLS fitting over the progress curve data on (p, t) as well as (s, t) can recover the original K_M and v_{max} with minimal error.

b) When $\rho > 0$, then the NLS fit procedure work better with the data on (s, t) than the data on (p, t) as demonstrated in **Figs. 2C-F**. Particularly, fluctuations in the trajectories of (p, t) in the pre-steady state regime is demonstrated in **Fig. 2C**. NLS fitting methods can capture k_1 and k_2 very well and the estimate on k_{-1} is prone to severe uncertainties especially when $k_{-1} \ll (k_1, k_2)$ as demonstrated with sample trajectories from **Figs. 2C** and **2D** in **Figs. 2E** and **2F**. This in turn leads to significant amount error in the estimation of K_M although v_{max} is not much affected by this issue.

c) The distribution of NLS fit parameters obtained at different noise levels are shown in **Figs. 3** for both (s, t) and (p, t) time course datasets. These results clearly suggest that K_M values obtained from the NLS fit of time course dataset on (p, t) space are unreliable and prone to significant fluctuations in the presence of high noise levels. When the noise level increases, then the distribution of fit parameters corresponding to (p, t) space dataset exhibits bimodality as shown in (**Figs. 3D**, **3F**). Whereas, NLS fit over time course dataset on (s, t) space can yield reliable median estimates of K_M even at high noise levels. Particularly, monomodal type distribution of fit values of K_M and v_{max} are observed for the dataset on (s, t) space as demonstrated in **Figs. 3A**, **3C**, **3E**.

d) Remarkably both time course datasets on (s, t) and (p, t) spaces can yield reliable NLS fit median estimates of v_{max} across various noise levels as demonstrated in **Fig. 3**. This observation is reasonable since v_{max} mostly relies on the post-steady state dynamics and therefore the fluctuations in the pre steady state regime of (p, t) space will not influence the error in the estimation of v_{max}.

These observations suggest that neither progress curve analysis alone nor steady state methods alone can provide consistent estimates of K_M. Multiple ways of analyzing the same dataset of (p, t) and (s, t) is clearly required to check for the consistency across various steady state and progress curve methods and obtain reliable estimates. In this context, one can use the median value of k_1 obtained from progress curve methods into the NLS fit on $\bar{v}_\varepsilon$ as given in **Eq. 10** to obtain consistent estimates of K_M and v_{max} across progress curve analysis and steady state methods. We consider the following methods I, II, III and IV using the same main time course datasets on (s, p, t). For Methods II, III and IV, sub-dataset on average reaction velocity versus s_0 at different t_r will be generated from the main dataset. Here the average velocity is defined as (total product formed at the end of t_r) / t_r where t_r will be iterated approximately from the steady state timescale t_c to t_m. Various NLS fit methods I, II, III and IV are summarized as follows.

a) **Method I** is the direct NLS fitting of the main progress curve dataset on (s, t) with fixed e_0 and different s_0 across several replications with the original rate equations **Eqs. 1.1-1.4** using Marquardt-Levenberg algorithm as outlined in the methods section. K_M values will be calculated from the fit values of (k_1, k_{-1}, k_2) and the median of (k_1, k_{-1}, k_2, K_M) values will also be obtained.

b) In **Method II**, the parameter k_1 in the definition of $\bar{v}_\varepsilon$ of **Eqs. 10** will be fixed with the median of k_1 values obtained from Method I and the sub-dataset on ($\bar{v}$, s_0) computed from the main time course dataset on (p, t) at different t_r will be used for NLS fitting with fixed k_1 to obtain K_M and v_{max}.

c) In **Method III**, $\bar{v}_{isQSSA}$ as defined in **Eq. 11** will be used to fit the sub-dataset on ($\bar{v}$, p, s_0) computed from the main dataset on (p, t) at different t_r and s_0. One can also use $\bar{v}_{csQSSA}$ that is computationally costlier than $\bar{v}_{isQSSA}$.

d) In **Method IV** that is widely used across various literature, $\bar{v}_{sQSSA} \cong v_0$ will be used to fit the sub-dataset on ($\bar{v}$, s_0) computed from the main dataset on (p, t) at different reaction time t_r and s_0.

In Methods II, III and IV, the K_M value corresponding to t_r with minimum sum of square error (SSE) as given in **Eq. 21** will be computed as demonstrated in **Fig. 4**. There exists a critical reaction time t_r at which the sum of square error in the estimation of K_M attains the minimum which is clearly observed in the absence of noise in the dataset ($\rho = 0$, **Fig. 4A**). Remarkably, **Fig. 4B** suggest that the error in the estimation of K_M strongly depends on the reaction time t_r in steady state Methods II, III and IV. In the presence of noise in the dataset, the median of SSE corresponding to K_M decreases monotonically and the fit parameter value converges to the original K_M as t_r is increased towards t_m. The speed of convergence of fit K_M towards the original K_M is much faster in Method II than in Methods III and IV as demonstrated in **Fig. 4B**. Sum of square errors of various approximations can be defined as $SSE = \sum_{i=0}^{n}\left(\bar{v}_{t_i} - \bar{v}_{G,t_i}\right)^2$ where $t_n = t_r$ and $\bar{v}_{t_i}$ is the numerically simulated average velocity at time $t = t_i$, and $\bar{v}_{G,t_i}$ are the approximations where the subscripts G = sQSSA, csQSSA, ε, isQSSA (I, II) associated with $\bar{v}_{sQSSA}, \bar{v}_{csQSSA}, \bar{v}_{\varepsilon}, \bar{v}_{isQSSA}$. The computed SSE seems to be strongly dependent on t_r and range of iteration of s_0 as demonstrated in **Fig. 4C** and **4D**. Clearly, SSE of $\bar{v}_{\varepsilon}$ is minimum near t_c and the SSEs corresponding to $(\bar{v}_{isQSSA}(I, II), \bar{v}_{csQSSA})$ attain minimum approximately near t_m. The timescale t_Q at which the SSE of $\bar{v}_{sQSSA}$ attains minimum can be computed by solving $\bar{v}_{\varepsilon} - \bar{v}_{csQSSA} = 0$ for t as shown in **Fig. 4E**. This is the critical time point at which both pre- and post-steady state approximations ($\bar{v}_{\varepsilon}, \bar{v}_{csQSSA}$) start to deviate from the original simulated average reaction velocity values.

When $t_r = t_Q$, then the reaction velocity will be $\bar{v} \cong v_0$ leading to minimum amount of error in the approximation $\bar{v}_{sQSSA}$. Remarkably, all the fitting methods ($\bar{v}_{sQSSA}, \bar{v}_{isQSSA}, \bar{v}_{\varepsilon}, \bar{v}_{csQSSA}$) approximately perform well near t_Q, at which the variation of the fit parameters across these methods will be at minimum. The entire procedure will be repeated with different e_0 until obtaining consistent median estimates of K_M and v_{max} with minimal threshold variation across methods I to IV. We consider the coefficient of variation (CV = standard deviation / mean) of the median K_M values obtained from methods I-IV as the *reliability index*. **Figs. 4C**, **4D** and **4E** suggest that the reliability index will be at minimum when the reaction time is set as $t_r = t_Q$. In the absence of *a priori* knowledge on enzyme kinetic parameters, one can use the reliability index to obtain best fit values of K_M which occurs near the optimum reaction time t_Q. Upon obtaining such consistent median estimates of K_M corresponding to a given threshold *reliability index*, all the K_M values obtained from the same dataset using different methods will be then pooled and the overall median of K_M and v_{max} will be computed as depicted in **Fig. 5**.

The distributions of various fit parameters obtained from Methods I to IV over several replications are shown in **Fig. 6** along with the overall pooled distribution of K_M and v_{max}. The distribution of fit parameters (k_1, k_{-1}, k_2) show skewness towards lower values (**Figs. 6A1-3**). Particularly, the distribution of k_{-1} exhibits a bimodal type with zero spike. As result, the distribution of K_M also exhibits spike near zero and v_{max} exhibits skewed distribution (**Figs. 6B1-2**). Since the median of k_1 obtained from Method I is used for the NLS fitting calculations in Method II, the distribution of K_M values obtained from Method II also exhibits a spike near zero as shown in **Fig. 6B-3**. Interestingly, both K_M and v_{max} obtained from Method III exhibit symmetric type distribution (**Figs. 6C-1**, **6D-1**). However, the distribution of K_M values obtained using Method III span towards negative values (**Fig. 6C-1**). The distribution of K_M values obtained from Method IV also exhibit a spike near zero (**Fig. 6C-2**). Clearly, the inconsistency

among the K_M values obtained from different methods originates mainly from the type of distribution of fit parameter in the presence of noise.

Efficient progress curve fitting also seems to be dependent on the detailed data collection around the transition regions in (s, t) and (p, t) spaces with maximum change in curvature [37]. Clearly, there exists at least three different timescale regimes in the trajectory on (p, t) space with significant change in curvature viz. exponential increasing phase in the pre-steady state regime, linear increasing phase in the steady state region and reaction-ending phase. Similar timescale regimes also exist in (s, t) space. Significant change in the curvature occurs at the interface of these timescale regimes approximately near t_c and t_m. Accurate values of these interface timescales can be calculated from s_ε and s_{csQSSA} for the pre- and post-steady state regimes respectively. The approximate curvature ζ_t associated with s_ε of **Eq. 8** can be defined as follows [48].

$$\zeta_t \cong \left(\frac{d^2 s_\varepsilon}{dt^2}\right)\bigg/\left(1 + \left(\frac{ds_\varepsilon}{dt}\right)^2\right)^{3/2} \; ; \; s_\varepsilon \cong s_0 - p_\varepsilon - v_\varepsilon/k_2 \qquad [36]$$

Here p_ε and v_ε are defined as in **Eqs. 8**. Upon solving $\frac{d\zeta_t}{dt} = 0$ for t, one finds the time scale associated with the maximum change in curvature at the interface of pre-steady to steady steady-state transition t_R as follows.

$$t_R \cong -t_c ln\left(\frac{t_c k_2}{4v_0(t_c k_2 - 1)}\left(v_0 - \sqrt{9v_0^2 + 8}\right)\right); \; v_0 = \frac{v_{max}s_0}{K_M + s_0} \qquad [37]$$

Similarly noting the definition of s_{csQSSA} as given in **Eqs. 12**, the approximate curvature ψ_t in the steady state to post-steady state transition regime of (s, t) space can be defined as follows.

$$\psi_t \cong \left(\frac{d^2 s_{csQSSA}}{dt^2}\right)\bigg/\left(1 + \left(\frac{ds_{csQSSA}}{dt}\right)^2\right)^{3/2} \qquad [38]$$

Upon solving $\frac{d\psi_t}{dt} = 0$ for t, one finds the timescale associated with the maximum change in curvature at the interface of steady state to reaction ending regime t_U as follows.

$$t_U \cong \frac{1}{v_{max}}\left|s_0 - K_M\left(h + ln\left(\frac{K_M h}{s_0}\right)\right)\right|; \; h = \left(\sqrt{8v_{max}^2 + 9} - 1\right)\big/4(v_{max}^2 + 1) \qquad [39]$$

Since the enzyme system evolves with constant velocity v_0 in the timescale region $t_R < t < t_U$, progress curve fitting methods can yield reliable estimate of K_M only when the experimental data collection is done around the timescales (t_R, t_U) with maximum change in the curvature of (s, t) trajectory. Whereas, steady state methods work very well in the constant velocity regime with minimal change in the curvature of the trajectory of (s, t). Change in curvature of the trajectory in (s, t) space with respect to time is shown in **Fig. 7A** along with the approximations given in **Eqs. 25-27** which are in good agreement with the numerical simulation results. Further, **Eqs. 25** and **27** suggest that (t_R, t_U) will be close to (t_c, t_m) and for practical purposes one can use (t_c, t_m) for checking the validity conditions of various model fitting methods. The simulated average reaction velocity (p / t) along with the approximations

$(\bar{v}_{sQSSA}, \bar{v}_{\varepsilon}, \bar{v}_{isQSSA} \ or \ \bar{v}_{csQSSA})$ are shown in **Fig. 7B**. These suggest that the approximation $\bar{v}_{\varepsilon}$ works well in the pre-steady to steady state regime and $\bar{v}_{isQSSA} \ or \ \bar{v}_{csQSSA}$ work well in the post steady state regime and $\bar{v}_{sQSSA}$ work only in narrow range of timescales with constant reaction velocity. Clearly, the reaction time t_r plays important role in deciding the reliability index among Methods I to IV. The error in the estimation of K_M using $\bar{v}_{\varepsilon}$ proportionally increases when t_r increases well beyond the steady state. Conversely, inclusion of data points from the pre-steady state regime increases the error in $\bar{v}_{isQSSA} \ or \ \bar{v}_{csQSSA}$. These means that the error in the estimation of KM using the widely used $\bar{v}_{sQSSA}$ will be always higher than the error associated with $(\bar{v}_{\varepsilon}, \bar{v}_{isQSSA} \ or \ \bar{v}_{csQSSA})$. Reliability index can be used as a check point to optimize the appropriate t_r to obtain accurate estimate of K_M values from progress curve and various steady state methods.

Software packages such as global kinetic explorer and dynafit [49-51] were developed to fit the data obtained from progress curve and steady state experiments. Several comparative studies were done earlier over steady state and progress curve methods to optimize the analysis workflow towards minimizing the estimation error [52, 53]. Selective removal of data points [54] over certain progress curve regime were also proposed. However, all these methods are solely either progress curve or steady state type and there is no way to check the consistency of fit parameters across various methods since each method requires different type of input dataset. Accurate estimation of kinetic parameters depends on strict impose of validity conditions of each method as given in **Table 1**. Unfortunately, these validity conditions require *a priori* knowledge on various enzyme kinetic parameters which are not generally available for unknown systems. Since the conditions of validity for each method is different, a global analysis workflow with a common input dataset is required to obtain consistent estimates. In this context, our proposed computational workflow depicted in **Fig. 5** can yield reliable estimates of K_M from the common time course dataset at different initial substrate concentrations s_0 and replications. The reliability index which is the coefficient of variation of median K_M values obtained from Methods I-IV can act as a check point against unreliable estimates of K_M. When the computed reliability index is more than the given threshold value, then one needs to iterate the reaction time until obtaining the desired threshold level. Clearly, the proposed methods do not require *a priori* knowledge on K_M values unlike the currently available sQSSA methods.

Identification of the correct inhibition mode using the experimental dataset on $(\bar{v}, s_0, i_0)$ viz. competitive, uncompetitive, noncompetitive and mixed is essential to design drug molecules for the target enzyme and obtain their IC$_{50}$ values for the comparison purposes. Dixon type graphical methods [26] used to identify the type of inhibition are not accurate in the presence of noise in the dataset on $(\bar{v}, s_0, i_0)$. It will be difficult to visually identify and measure the intersection points in the Dixon plot. Sequential and simultaneous nonlinear regression methods [27, 28] were proposed to overcome such issues. In this context, we have developed the multiple nonlinear regression (MNLS) method to evaluate various inhibition parameters directly by fitting dataset on $(\bar{v}, s_0, i_0)$ over the respective inhibition model equations as shown in **Fig. 8**. Using MNLS method, one can directly obtain the parameters fit values (K_M, K_I, K_Z, v_{max}) along with their standard errors from the Dixon type datasets via minimizing the overall sum of square deviations as defined in **Eq. 28**. Fit parameter values for sample datasets in the presence and absence of noise are given in **Tables 2** and **3**. MNLS method can recover the exact parameter values in the absence of noise in the datasets as demonstrated in **Figs. 8A**,

8C, **8E**, **8F** and **Table 2**. Further, an algorithm based on the sum of square error (SSE) values of various multiple nonlinear fit functions and standard errors of fit parameters is also developed as shown in **Fig. 9**. In this algorithm, the given enzyme inhibition dataset on ($\bar{v}$, s_0, i_0) will be simultaneously fitted with competitive, uncompetitive, noncompetitive and mixed inhibition model equations as given in **Eqs. 14-17**. The overall sum of square errors computed with respect to each of these model functions were compared and the model function showing the lowest overall SSE will be chosen as the best fit model. When the chosen model is a mixed-competitive inhibition model, then further investigations will be carried out as follows to avoid ambiguities arising due to the generalized nature of mixed inhibition model equation.

a. When $K_I = K_Z$, then it will be interpreted as the noncompetitive inhibition model. This scenario occurs when one tries to fit noninhibition dataset over mixed inhibition model function.
b. When the parameter error corresponding to K_I or K_Z is higher than the next lowest SSE, then the next lowest SSE model will be the best fit model.
c. When $K_I \gg K_Z$, then it will be interpreted as the uncompetitive inhibition model. When the dataset on uncompetitive inhibition is fitted with mixed inhibition model equation, then one observes $K_I \gg K_Z$ in the fit results as shown in **Tables 2** and **3**.
d. When $K_I \ll K_Z$, then it will be interpreted as the competitive inhibition model. When the dataset on competitive inhibition is fitted with mixed inhibition model equation, then one observes $K_I \ll K_Z$ in the fit results as shown in **Table 2** and **3**.

Performance of the inhibition type classification algorithm in the presence of noise is shown in **Fig. 10**. Clearly, our inhibition type classification algorithm works very well especially at low noise levels. Further, fitting of competitive inhibition dataset may be wrongly shown as mixed inhibition type at moderate noise levels as demonstrated in **Fig. 10A**. At high noise levels, uncompetitive inhibition dataset may be wrongly shown as (**Fig. 10B**) noncompetitive inhibition and noncompetitive inhibition dataset may be wrongly detected as mixed, competitive and uncompetitive inhibition (**Fig. 10C**). These results clearly suggests that repeated data collection and MNLS analysis is required to obtain consistent classification of inhibition schemes and the respective parameters and IC_{50}.

Conclusion

Accurate estimation of enzyme kinetic parameters such as K_M and v_{max} from the experimental datasets is critical to characterize any enzyme. The validity of the widely used standard quasi steady state method over decades relies on the *a priori* information of K_M and other kinetic parameters which are generally not available for newly identified enzyme systems. Progress curve methods were proposed to alleviate such issues. However, nonlinear least square fitting of the time course dataset with the original differential rate equations works well only in the timescale regimes with maximum change in the curvature of the trajectory of substrate evolution. Identifying these timescales requires the knowledge on enzyme kinetic parameters. Though experimental trial and error methods can be used to identify such timescales, they are generally cost ineffective.

Further, progress curve methods work better with the time course data on substrate than the time course data on product since the product concentration is close to zero in the pre-steady

state regime. Since steady state and progress curve analysis use different types of datasets in the current analysis set up, the estimated K_M values will be inconsistent among these methods. This leads to the dilemma of identifying correct K_M values among various steady state and progress curve analyses. Here we have shown that the error in the estimation of K_M using steady state methods is strongly dependent on the total reaction time. We have further shown that the steady state methods work well over the reaction timescales with minimum change in the curvature of the trajectory of product evolution with constant reaction velocity. Considering these theoretical factors, we have developed an integrated method of analysis which comprises of both progress curve and steady state analysis using the same time course datasets that also consider the total reaction time. We considered three different methods of steady state analyses that involve fitting of the average reaction velocity with the total substrate concertation. We have defined the reliability index which is the coefficient of variation of the median K_M values obtained from progress curve and other steady state fitting methods using the same time course dataset at different substrate concentrations and replications. We further demonstrate that reliable estimate of K_M can be obtained by iterating the total reaction time to minimize the reliability index across time course and steady state curve fitting methods.

We have developed a multiple nonlinear fitting procedure-based algorithm to classify the average velocity, total substrate and inhibitor concentrations dataset over competitive, uncompetitive, noncompetitive and mixed types. This method works very well at low to medium noise levels. However, at high noise levels the competitive inhibition dataset may be wrongly shown as mixed inhibition type, uncompetitive inhibition dataset may be wrongly shown as noncompetitive inhibition and noncompetitive inhibition dataset may be wrongly detected as mixed, competitive and uncompetitive inhibition types. Repeated data collection and multiple nonlinear fitting analysis is required to obtain consistent estimates of inhibition parameters and IC_{50}.

Data availability statement
The datasets generated during and/or analysed during the current study are available in the Zenodo. DOI: 10.5281/zenodo.15094623

References

1. Alberts B. Molecular biology of the cell. New York: Garland Science; 2002.
2. Stryer L. Biochemistry. New York: W.H. Freeman; 1988.
3. Voet D, Voet JG. Biochemistry. New York: J. Wiley & Sons; 1995.
4. Michaelis L, Menten ML. {Die kinetik der invertinwirkung}. Biochem Z. 1913;49(333-369):352. doi: citeulike-article-id:5936552.
5. Briggs GE, Haldane JBS. A Note on the Kinetics of Enzyme Action. Biochemical Journal. 1925;19(2):338-9. PubMed PMID: PMC1259181.
6. Murugan R. Solution to Michaelis–Menten enzyme kinetic equation via undetermined gauge functions: Resolving the nonlinearity of Lineweaver–Burk plot. The Journal of chemical physics. 2002;117(9):4178-83.
7. Murray JD. Mathematical biology. New York: Springer; 2002.
8. Fraser SJ. Slow manifold for a bimolecular association mechanism. J Chem Phys. 2004;120(7):3075-85. doi: 10.1063/1.1634555. PubMed PMID: 15268460.
9. Roussel MR, Fraser SJ. Invariant manifold methods for metabolic model reduction. Chaos. 2001;11(1):196-206. doi: 10.1063/1.1349891. PubMed PMID: 12779453.
10. Dingee JW, Anton AB. A new perturbation solution to the Michaelis-Menten problem. AIChE Journal. 2008;54(5):1344-57. doi: 10.1002/aic.11461.
11. Vogt D. A simplified perturbation solution of Michaelis-Menten kinetics equations in a "total" framework. Journal of Mathematical Chemistry. 2013;51(9):2257-63. doi: 10.1007/s10910-013-0221-0.
12. Dell'Acqua G, Bersani AM. A perturbation solution of Michaelis–Menten kinetics in a "total" framework. Journal of Mathematical Chemistry. 2012;50(5):1136-48. doi: 10.1007/s10910-011-9957-6.
13. Seshadri MS, Fritzsch G. The time evolution of sequential enzyme reactions: a singular perturbation approach. J Theor Biol. 1981;93(1):197-205. PubMed PMID: 7334820.
14. Segel LA, Slemrod M. The Quasi-Steady-State Assumption: A Case Study in Perturbation. SIAM Review. 1989;31(3):446-77. doi: 10.1137/1031091.
15. Dell'Acqua G, Bersani AM. On the appropriate use of asymptotic expansions in enzyme kinetics. Journal of Mathematical Chemistry. 2014;52(10):2475-81. doi: 10.1007/s10910-014-0403-4.
16. Murugan R. Theory on the rate equation of Michaelis–Menten type single-substrate enzyme catalyzed reactions. Journal of Mathematical Chemistry. 2018;56(2):508-56. doi: 10.1007/s10910-017-0791-3.
17. Hofstee BH. Non-inverted versus inverted plots in enzyme kinetics. Nature. 1959;184:1296-8. doi: 10.1038/1841296b0. PubMed PMID: 14402470.
18. Atkins GL, Nimmo IA. A comparison of seven methods for fitting the Michaelis-Menten equation. Biochem J. 1975;149(3):775-7. PubMed PMID: 1201002; PubMed Central PMCID: PMCPMC1165686.
19. Lineweaver H, Burk D. The Determination of Enzyme Dissociation Constants. Journal of the American Chemical Society. 1934;56(3):658-66. doi: 10.1021/ja01318a036.
20. Wilkinson GN. Statistical estimations in enzyme kinetics. Biochem J. 1961;80(2):324-32. doi: 10.1042/bj0800324. PubMed PMID: 13785321; PubMed Central PMCID: PMCPMC1244002.
21. Fernley HN. Statistical estimations in enzyme kinetics. The integrated Michaelis equation. Eur J Biochem. 1974;43(2):377-8. doi: 10.1111/j.1432-1033.1974.tb03423.x. PubMed PMID: 4838986.

22. Murugan R. Theory on the rate equations of Michaelis-Menten type enzyme kinetics with competitive inhibition. PLOS ONE. 2024;19(7):e0302679. doi: 10.1371/journal.pone.0302679.

23. Cheng Y, Prusoff WH. Relationship between the inhibition constant (K1) and the concentration of inhibitor which causes 50 per cent inhibition (I50) of an enzymatic reaction. Biochem Pharmacol. 1973;22(23):3099-108. doi: 10.1016/0006-2952(73)90196-2. PubMed PMID: 4202581.

24. Yadav J, Maldonato BJ, Roesner JM, Vergara AG, Paragas EM, Aliwarga T, et al. Enzyme-mediated drug-drug interactions: a review of in vivo and in vitro methodologies, regulatory guidance, and translation to the clinic. Drug Metab Rev. 2024:1-33. Epub 20240726. doi: 10.1080/03602532.2024.2381021. PubMed PMID: 39057923.

25. Jang HJ, Song YM, Jeon JS, Yun HY, Kim SK, Kim JK. Optimizing enzyme inhibition analysis: precise estimation with a single inhibitor concentration. Nat Commun. 2025;16(1):5217. Epub 20250605. doi: 10.1038/s41467-025-60468-z. PubMed PMID: 40473596; PubMed Central PMCID: PMCPMC12141731.

26. Dixon M. The determination of enzyme inhibitor constants. Biochem J. 1953;55(1):170-1. doi: 10.1042/bj0550170. PubMed PMID: 13093635; PubMed Central PMCID: PMCPMC1269152.

27. Ernest CS, 2nd, Hooker AC, Karlsson MO. Methodological comparison of in vitro binding parameter estimation: sequential vs. simultaneous non-linear regression. Pharm Res. 2010;27(5):866-77. Epub 20100311. doi: 10.1007/s11095-010-0082-1. PubMed PMID: 20221897.

28. Kakkar T, Boxenbaum H, Mayersohn M. Estimation of Ki in a competitive enzyme-inhibition model: comparisons among three methods of data analysis. Drug Metab Dispos. 1999;27(6):756-62. PubMed PMID: 10348808.

29. Segel LA. On the validity of the steady state assumption of enzyme kinetics. Bull Math Biol. 1988;50(6):579-93. PubMed PMID: 3219446.

30. Borghans JM, De Boer RJ, Segel LA. Extending the quasi-steady state approximation by changing variables. Bulletin of Mathematical Biology. 1996;58(1):43-63. doi: http://dx.doi.org/10.1016/0092-8240(95)00306-1.

31. Grima R, Walter NG, Schnell S. Single-molecule enzymology a la Michaelis-Menten. FEBS J. 2014;281(2):518-30. doi: 10.1111/febs.12663. PubMed PMID: 24289171.

32. Choi IY, Lee SP, Kim SG, Gruetter R. In vivo measurements of brain glucose transport using the reversible Michaelis-Menten model and simultaneous measurements of cerebral blood flow changes during hypoglycemia. J Cereb Blood Flow Metab. 2001;21(6):653-63. doi: 10.1097/00004647-200106000-00003. PubMed PMID: 11488534.

33. Lee BY, Kwon KI, Kim MS, Baek IH. Michaelis-Menten elimination kinetics of etanercept, rheumatoid arthritis biologics, after intravenous and subcutaneous administration in rats. Eur J Drug Metab Pharmacokinet. 2016;41(4):433-9. doi: 10.1007/s13318-015-0270-9. PubMed PMID: 25725773.

34. Choi B, Rempala GA, Kim JK. Beyond the Michaelis-Menten equation: Accurate and efficient estimation of enzyme kinetic parameters. Scientific Reports. 2017;7(1):17018. doi: 10.1038/s41598-017-17072-z.

35. Frère JM, Verlaine O, Matagne A. The measurement of true initial rates is not always absolutely necessary to estimate enzyme kinetic parameters. Sci Rep-Uk. 2023;13(1). doi: ARTN 15053
10.1038/s41598-023-41805-y. PubMed PMID: WOS:001168651100052.

36.   Duggleby RG, Clarke RB. Experimental designs for estimating the parameters of the Michaelis-Menten equation from progress curves of enzyme-catalyzed reactions. Biochim Biophys Acta. 1991;1080(3):231-6. PubMed PMID: 1954231.

37.   Stroberg W, Schnell S. On the estimation errors of KM and V from time-course experiments using the Michaelis-Menten equation. Biophys Chem. 2016;219:17-27. doi: 10.1016/j.bpc.2016.09.004. PubMed PMID: 27677118.

38.   Hanson SM, Schnell S. Reactant stationary approximation in enzyme kinetics. J Phys Chem A. 2008;112(37):8654-8. doi: 10.1021/jp8026226. PubMed PMID: 18714952.

39.   Corless RM, Gonnet GH, Hare DEG, Jeffrey DJ, Knuth DE. On the LambertW function. Advances in Computational Mathematics. 1996;5(1):329-59. doi: 10.1007/BF02124750.

40.   Abramowitz M, Stegun IA. Handbook of mathematical functions, with formulas, graphs, and mathematical tables. New York: Dover Publications; 1965.

41.   Gardiner CW. Handbook of stochastic methods for physics, chemistry, and the natural sciences. Berlin; New York: Springer-Verlag; 1985.

42.   Risken H. The Fokker-Planck equation : methods of solution and applications. Berlin; New York: Springer-Verlag; 1989.

43.   Press WH. Numerical recipes in C++ : the art of scientific computing. 2nd ed. Cambridge, UK: Cambridge University Press; 2002.

44.   Marquardt DW. An Algorithm for Least-Squares Estimation of Nonlinear Parameters. Journal of the Society for Industrial and Applied Mathematics. 1963;11(2):431-41. doi: 10.1137/0111030.

45.   Levenberg K. A method for the solution of certain non-linear problems in least squares. Quart Appl Math. 1944;2:164--8. doi: doi:10.1090/qam/10666.

46.   Zimmerle CT, Frieden C. Analysis of progress curves by simulations generated by numerical integration. Biochem J. 1989;258(2):381-7. doi: 10.1042/bj2580381. PubMed PMID: 2705989; PubMed Central PMCID: PMCPMC1138373.

47.   Kuzmic P. Program DYNAFIT for the analysis of enzyme kinetic data: application to HIV proteinase. Anal Biochem. 1996;237(2):260-73. doi: 10.1006/abio.1996.0238. PubMed PMID: 8660575.

48.   Courant R, John F. Introduction to calculus and analysis. Repr. of the 1989 ed. Berlin: Springer; 1989.

49.   Johnson KA. Fitting enzyme kinetic data with KinTek Global Kinetic Explorer. Methods Enzymol. 2009;467:601-26. doi: 10.1016/S0076-6879(09)67023-3. PubMed PMID: 19897109.

50.   Johnson KA, Simpson ZB, Blom T. Global kinetic explorer: a new computer program for dynamic simulation and fitting of kinetic data. Anal Biochem. 2009;387(1):20-9. Epub 20081225. doi: 10.1016/j.ab.2008.12.024. PubMed PMID: 19154726.

51.   Mak DA, Dunn S, Coombes D, Carere CR, Allison JR, Nock V, et al. Enzyme Kinetics Analysis: An online tool for analyzing enzyme initial rate data and teaching enzyme kinetics. Biochemistry and Molecular Biology Education. 2024;52(3):348-58. doi: https://doi.org/10.1002/bmb.21823.

52.   Cho YS, Lim HS. Comparison of various estimation methods for the parameters of Michaelis-Menten equation based on in vitro elimination kinetic simulation data. Transl Clin Pharmacol. 2018;26(1):39-47. Epub 20180316. doi: 10.12793/tcp.2018.26.1.39. PubMed PMID: 32055546; PubMed Central PMCID: PMCPMC6989224.

53.	Marasović M, Marasović T, Miloš M. Robust Nonlinear Regression in Enzyme Kinetic Parameters Estimation. Journal of Chemistry. 2017;2017(1):6560983. doi: https://doi.org/10.1155/2017/6560983.

54.	Petric B, Golicnik M, Bavec A. The Removal of Time-Concentration Data Points from Progress Curves Improves the Determination of K(m): The Example of Paraoxonase 1. Molecules. 2022;27(4). Epub 20220215. doi: 10.3390/molecules27041306. PubMed PMID: 35209091; PubMed Central PMCID: PMCPMC8874660.

FIGURE 1. Dynamics of product formation (p, μM) in Michaelis-Menten scheme of single substrate enzyme catalyzed reactions. Here $t_c \cong 1/\big(k_1(K_M + s_0)\big)$, $t_m \cong ((K_M + s_0)/k_2 e_0)$

are the approximate pre-steady state and reaction-end timescales (t, seconds) and numerical integration settings from **Eqs. 1** are k_1 = 0.1 $\mu M^{-1} s^{-1}$, k_{-1} = 0.01 s^{-1} and k_2 = 0.2 s^{-1}, with K_M = 2.1 μM. **A**. s_0 = 100 μM and e_0 was iterated in (0.1, 1, 10, 100) μM. **B**. e_0 = 100 μM and s_0 was iterated in (1, 10, 100-1000, with increment of 100) μM along the arrow. Scheme B describes the fully competitive inhibition and Scheme C describes various partial inhibition schemes viz. competitive (k_z = 0), uncompetitive (k_i = 0) and mixed types. In fully competitive inhibition both substrate and inhibitor will be converted into their respective products by the same enzyme. Whereas, in case of partial competitive inhibition the enzyme-inhibitor will be a dead-end complex.

FIGURE 2. Progress curve analysis of data on (p, t) and (s, t). Common simulation settings for the numerical integration scheme in **Eqs. 13** are k_1 = 5.1 $\mu M^{-1} s^{-1}$, k_{-1} = 0.5 s^{-1}, k_2 = 1.1 s^{-1}, Δt = 10^{-3} s, e_0 = 0.85 μM, s_0 = 50 μM so that K_M = 0.313725 μM, $t_c \approx 0.0039$ s, $t_m \approx 118.38$ s, total simulation time = 10 s and v_{max} = 0.935 μM s^{-1}. **A**. Simulation of **Eqs. 13** with ρ = 0 and NLS fit of the simulation trajectory with **Eqs. 1** using ML algorithm yielded the parameters k_1 = 5.1 ± 1.25 x 10^{-5}, k_{-1} = 0.499 ± 7 x 10^{-6}, k_2 = 1.1 ± 4.5 x 10^{-8} for the data on (p, t) with K_M = 0.314 at 95% confidence level. **B**. NLS fit using ML algorithm over **Eqs. 1** yielded k_1 = 5.1 ± 6 x 10^{-6}, k-1 = 0.499 ± 9 x 10^{-6}, k_2 = 1.1 ± 4.5 x 10^{-8} for data on (s, t) generated using ρ = 0 in **Eqs. 13** with K_M = 0.313639 at 95% confidence level. **C**. stochastic trajectories (1000 numbers) of **Eqs. 13** with ρ = 10^{-2} demonstrating the fluctuations in the pre-steady state regime of (p, t) space. **D**. stochastic trajectories of **Eqs. 13** with ρ = 10^{-2} demonstrating less fluctuations in the pre-steady state regime of (s, t) space. **E**. NLS fit of sample (p, t) trajectory with ρ = 10^{-2} yielded k_1 = 2.29 ± 0.09, k_{-1} = 1.53 ± 0.1, k_2 = 1.12 ± 2 x 10^{-3} with K_M = 1.18 and v_{max} = 0.95 at 95% confidence level. **F**. NLS fit of sample (s, t) trajectory with ρ = 10^{-2} over **Eqs. 1** yielded k_1 = 5.1 ± 0.07, k_{-1} = 1.3 ± 0.1, k_2 = 1.1 ± 5 x 10^{-4} with K_M = 0.47 and v_{max} = 0.94 at 95% confidence level.

FIGURE 3. Distribution of nonlinear least squares (NLS) fit parameters across various noise levels. Common simulation settings for the numerical integration scheme in **Eqs. 13** are k_1 = 5.1 $\mu M^{-1} s^{-1}$, k_{-1} = 0.5 s^{-1}, k_2 = 1.1 s^{-1}, Δt = 10^{-3} s, total simulation time = 10 s, e_0 = 0.85 μM so that K_M = 0.313725 μM and v_{max} = 0.935 $\mu M\ s^{-1}$, s_0 = 50 μM for which one finds that $t_c \approx 0.0039$ s and $t_m \approx 118.38$ s. NLS fit was done using Marquardt-Levenberg algorithm as outlined in the methods section. In A, C and E, NLS fittings were done using data on (s, t). Whereas, in B, D

and F the fittings were done using data on (p, t). Upon obtaining k_1, k_{-1} and k_2, K_M and v_{max} will be calculated. Noise levels in **Eqs. 13** were set as follows. **A**, **B**. $\rho = 10^{-3}$. **C**, **D**. $\rho = 10^{-2}$. **E**, **F**. $\rho = 10^{-1}$. The median values of K_M and v_{max} were computed across 10^3 replications. **A**. the median values of $(k_1, k_{-1}, k_2, K_M, v_{max})$ = (5.1005, 0.5009, 1.100, 0.3140, 0.9350). **B**. the median values of $(k_1, k_{-1}, k_2, K_M, v_{max})$ = (4.9188, 0.3849, 1.1001, 0.3163, 0.9351). **C**. the median values of $(k_1, k_{-1}, k_2, K_M, v_{max})$ = (5.2031, 0.5499, 1.1021, 0.3197, 0.9368). **D**. the median values of $(k_1, k_{-1}, k_2, K_M, v_{max})$ = (4.7469, 1.1193, 1.0981, 1.5888, 0.9334). **E**. the median values of $(k_1, k_{-1}, k_2, K_M, v_{max})$ = (6.5060, 0.4776, 1.1348, 0.3727, 0.9646). **F**. the median values of $(k_1, k_{-1}, k_2, K_M, v_{max})$ = (16.0345, 0.7332, 1.1380, 3.1860, 0.9673).

FIGURE 4. Variation of the fitted K_M values with respect to reaction time t_r across different fitting methods. Common simulation settings for the numerical integration scheme in **Eqs. 13** are $k_1 = 5.1\ \mu M^{-1} s^{-1}$, $k_{-1} = 0.5\ s^{-1}$, $k_2 = 1.1\ s^{-1}$, $\Delta t = 10^{-3}\ s$, total simulation time = 10 s, $e_0 = 0.85$ μM so that $K_M = 0.313725\ \mu M$ and $v_{max} = 0.935\ \mu M\ s^{-1}$, s_0 was iterated from 50 to 250 μM with interval of 25 μM. For $s_0 = 50\ \mu M$ one finds that $t_c \approx 0.0039$ s and $t_m \approx 118.38$ s. The reaction time t_r was iterated from 0.1 to 10 s. The sub datasets on ($\bar{v}$, p, s_0) at different t_r were generated from the main progress curve dataset on (p, t) and used to fit **Eqs. 10** (ϵ-approximation), **Eq. 11** (isQSSA) and v_0 (sQSSA) and the median values of K_M were obtained across 10^3 replications. **A.** $\rho = 0$ in **Eqs. 13**. **B.** $\rho = 10^{-2}$ in **Eqs. 13**. **C.** $\Delta t = 10^{-5}$ s, s_0 was iterated from 50 to 250 μM with interval of 25 μM. For p_{csQSSA} / t_r, s_0 was iterated from 100 to 200 μM with interval of 10 μM. **D.** For all the steady state methods, $\Delta t = 10^{-5}$ s, s_0 was iterated from 100 to 200 μM with interval of 10 μM. **E.** $s_0 = 100\ \mu M$ and $\Delta t = 10^{-5}$ s for which $t_Q \approx 10.49$ s.

FIGURE 5. Computational algorithm to obtain consistent estimate of K_M from the progress curve dataset. Here, the progress curve data on (p, s, t) will be the main input for Methods I, II, III and IV. Here t_{max} is the maximum data collection time. Method I is the direct nonlinear least square fitting of the dataset on (s, t) with fixed e_0 and different s_0 across replications with the original **Eqs. 1.1-1.4** using Marquardt-Levenberg algorithm as outlined in the methods section. The parameter k_1 in $\bar{v}_\varepsilon$ of **Eqs. 10** of Method II will be fixed with the median of k_1 values obtained from Method I. For Methods II-IV, sub-dataset on average velocity ((total product formed at the end of t_r) / t_r, where t_r will be iterated from t_c towards t_m) versus s_0 at different t_r will be generated from the main dataset. NLS fittings over ($\bar{v}$, p, s_0) were done using models corresponding to Methods II-IV at different t_r. The K_M value corresponding to t_r with minimum sum of square error (SSE) will be considered. Upon obtaining consistency on the estimates of K_M across Methods I-IV corresponding to the given threshold *reliability index*, all the K_M values obtained from the same dataset using different methods will be then pooled and the overall median K_M will be computed. Here *reliability index* is the coefficient of variation of median K_M values obtained from Methods I-IV.

FIGURE 6. Distribution of fit parameters across various methods I-IV. Common simulation settings for the numerical integration scheme in **Eqs. 13** are $k_1 = 5.1\ \mu M^{-1}s^{-1}$, $k_{-1} = 0.5\ s^{-1}$, $k_2 = 1.1\ s^{-1}$, $\Delta t = 10^{-3}$ s, total simulation time = 10 s, $e_0 = 0.85\ \mu M$ so that $K_M = 0.313725\ \mu M$ and $v_{max} = 0.935\ \mu M\ s^{-1}$, s_0 was iterated from 50 to 250 μM with interval of 25 μM. For $s_0 = 50\ \mu M$ one finds that $t_c \approx 0.0039$ s and $t_m \approx 118.38$ s. The median values corresponding to progress curve fit over 10^3 trajectories on (s, t) dataset were $(k_1, k_{-1}, k_2, K_M, v_{max}) = (5.164, 0.587, 1.102, 0.331, 0.937)$. **A1-3, B1-2**. NLS fit method I using Marquardt-Levenberg algorithm. Upon obtaining k_1, k_{-1} and k_2, K_M and v_{max} will be calculated as in **B1, B2**. **B3, C3**. Method II. Here the obtained median values from datasets on (p, t) were $(K_M, v_{max}) = (0.338, 0.935)$. **C1, D1**. Method III. Here the obtained median values from datasets on (p, t) were $(K_M, v_{max}) = (0.332, 0.935)$. **C2, D2**. Method IV. Here the obtained median values from datasets on (p, t) were $(K_M, v_{max}) = (0.366, 0.935)$. **D3**. Overall distribution of pooled K_M values across all the methods I-IV with overall median of $K_M = 0.3418\ \mu M$ which correspond to the coefficient of variation of median K_M values obtained from Methods I, II, III and IV as 0.048.

FIGURE 7. Change in the curvature of trajectory on (s, t) space and average reaction velocity. Common simulation settings for the numerical integration scheme in **Eqs. 13** are k_1 = 5.1 μM^{-1} s^{-1}, k_{-1} = 0.5 s^{-1}, k_2 = 1.1 s^{-1}, Δt = 0.5 x 10^{-4} s, total simulation time = 1000 s, noise level set as ρ = 0, e_0 = 0.85 μM so that K_M = 0.313725 μM and v_{max} = 0.935 μM s^{-1}, and s_0 = 100 μM. For these settings, one finds that $t_c \approx 0.002$ s, $t_m \approx 91.2$ s and $t_Q \approx 10.49$ s. **A**. The transition timescales (t_R, t_U) at which maximum change in the curvature occurs can be calculated using **Eqs. 25-27** as t_R = 0.0126 s and t_U = 92 s. The time dependent curvatures associated with the trajectory in (s, t) space ζ_t and ψ_t corresponding to the pre- and post-steady state regimes respectively are defined as in **Eqs. 24-27**. Progress curve Method I works very well in the regions with maximum change in the curvature. Whereas, steady state methods II, III and IV work very well in the regions with minimal change in the curvature. **B**. Change in the average velocity (p / t) with time. Method II works well from the pre-steady to steady state regime. Whereas, Method III works well in the steady to post-steady state regime. Method IV works well only in a narrow timescale regime with constant average velocity.

FIGURE 8. Multiple nonlinear regression fitting of Dixon type enzyme inhibition dataset on ($\bar{v}$, s_0, i_0) in the presence and absence of noise. A, C, E and G are in the absence of noise. B, D, F and H are in the presence of noise with strength $\rho = 10^{-3}$. The preset values of parameters are K_M

= 2.80952 μM, K_I = 0.354839 μM, K_Z = 2.12903 μM and v_{max} = 4.62 μM/s. s_0 was iterated from 10 to 100 μM with an increment of 10 μM and i_0 was iterated in 10 to 110 μM with an increment of 10 μM. **Eqs. 14-17** were used to generate datasets for (1) competitive, (2) uncompetitive, (3) noncompetitive, (4) mixed-competitive modes of enzyme inhibition and the multiple nonlinear fitting were done as outline in the methods section. The fit results are summarized **Tables 2** and **3**. **A**, **B**. Competitive inhibition data fitted with (1)-(4) modes of inhibition. **C**, **D**. Uncompetitive inhibition data fitted with (1)-(4) modes of inhibition. **E**, **F**. Noncompetitive inhibition data fitted with (1)-(4) modes of inhibition. **G**, **H**. Mixed-competitive inhibition data fitted with (1)-(4) modes of inhibition.

FIGURE 9. Classification algorithm for the type of inhibition from Dixon type dataset. Here the given dataset on ($\bar{v}$, s_0, i_0) was fitted with competitive, uncompetitive, noncompetitive and mixed inhibition models and the sum of square errors were compared and model showing lowest overall SSE will be chosen. If the model is a mixed-competitive inhibition model, then

the following investigations should be carried out. (1) when $K_I = K_Z$, then it will the noncompetitive inhibition model. (2) when the parameter error corresponding to K_I or K_Z is higher than the next lowest SSE, then the next lowest SSE model will be the best fit model. (3) when $K_I \gg K_Z$, then it will be the uncompetitive inhibition model. (4) when $K_I \ll K_Z$, then it will be the competitive inhibition model.

FIGURE 10. Performance of the inhibition type classification algorithm. The preset values of parameters are K_M = 2.80952 μM, K_I = 0.354839 μM, K_Z = 2.12903 μM and v_{max} = 4.62 μM/s. s_0 was iterated from 10 to 100 μM with an increment of 10 μM and i_0 was iterated in 10 to 110 μM with an increment of 10 μM. **Eqs. 14-17** were used to generate datasets for (1) competitive, (2) uncompetitive, (3) noncompetitive, (4) mixed-competitive modes of enzyme inhibition and the multiple nonlinear fitting were done as outlined in the methods section. Gaussian white noise with zero mean and unit variance was added through the control parameter ρ that was iterated from 10^{-6} to 1 as $\bar{v} \rightarrow \bar{v} + \rho\xi$ where ξ is the random number drawn from the standard normal population N (0, 1). Fraction fit results were computed over 1000 randomly generated datasets. **A.** Competitive inhibition datasets were fitted with model equations corresponding to (1)-(4) modes of inhibition. **B.** Uncompetitive inhibition datasets were fitted with model equations corresponding to (1)-(4) modes of inhibition. **C.** Noncompetitive inhibition datasets were fitted with model equations corresponding to (1)-(4) modes of inhibition. **D.** Mixed-competitive inhibition datasets were fitted with the model equations corresponding to (1)-(4) modes of inhibition.

TABLE 1. Conditions of validity of various fitting methods to obtain K$_M$ and v$_{max}$

Description	Definition	Validity conditions
Progress curve analysis	Time course dataset on (s, p, t).	timescale interval should capture both pre- and post-steady state regimes. (Method I)
$\bar{v}_{sQSSA}$	$\bar{v}_{sQSSA} \cong v_{max}s_0/(K_M + s_0)$	$t_c \ll t_r < t_m, (\eta, \varepsilon) \to (0,0)$, $(e_0/(K_M + s_0)) \ll 1$, Method IV
$\bar{v}_{isQSSA}$	$\bar{v}_{isQSSA} \cong v_{max} + \frac{K_M}{t_r} ln\left(\frac{s_0-p}{s_0}\right)$ (I) $\bar{v}_{isQSSA} \cong v_{max} + \frac{K_M}{t_r} ln\left(\frac{s}{s_0}\right)$ (II)	$t_c < t_r < t_m, (\eta, \varepsilon) \to (0,0)$ Method III.
$\bar{v}_{csQSSA}$	$\bar{v}_{csQSSA} \cong \frac{1}{t_r}\left(s_0 - s_{csQSSA}\right)$ s_{csQSSA} $\cong K_M W\left(\frac{s_0}{K_M} exp\left(\frac{s_0 - v_{max}t_r}{K_M}\right)\right)$	$t_c < t_r < t_m, (\eta, \varepsilon) \to (0,0)$ Method III. Alternate form of $\bar{v}_{isQSSA}$.
$\bar{v}_\varepsilon$	$\bar{v}_\varepsilon \cong v_0(1 - t_c h(t_r)/t_r)$ $v_0 = v_{max}s_0/(K_M + s_0)$ $t_c \cong 1/k_1(K_M + s_0)$ $h(t_r) = \left(1 - exp(-t_r/t_c)\right)$	$t_r < t_m, \varepsilon \to 0$ Method II. median of k_1 obtained from Method I will be substituted in t_c.

TABLE 2. Multiple nonlinear regression of enzyme inhibition data in the absence of noise

Parameter fit	Competitive	Uncompetitive	Non competitive	Mixed
Competitive inhibition dataset				
K$_M$	2.81 ± 1.4 x 10^{-10}	709019 ± 0.3	141.9 ± 7.2	2.81 ± 5.4 x 10^{-9}
K$_I$	0.355 ± 1.7 x 10^{-11}	-	9.48 ± 0.4	0.355 ± 7 x 10^{-10}
K$_Z$	-	0.00124 ± 3.6 x 10^{-10}	-	278930 ± 15
v$_{max}$	4.62 ± 8.3 x 10^{-12}	35824.9± 1.4 x 10^{-10}	13.21 ± 0.47	4.62 ± 6.1 x 10^{-9}
SSE	9.64496e-25	0.0220512	0.00144751	1.08116e-11
Uncompetitive inhibition dataset				
K$_M$	0.003 ± 1.6 x 10^{-9}	2.80952 ± 8.8 x 10^{-10}	0.36 ± 0.03	2.81 ± 7.5 x 10^{-7}
K$_I$	0.0005 ± 2.6 x 10^{-10}	-	2.3 ± 0.029	118254 ± 103
K$_Z$	-	2.12903 ± 1.2 x 10^{-9}	-	2.13 ± 8.8 x 10^{-9}
v$_{max}$	1.4 ± 5 x 10^{-7}	4.62 ± 2.3 x 10^{-10}	4.32072 ± 0.047	4.62 ± 1.8 x 10^{-8}
SSE	0.0159053	2.29059e-23	1.59847e-06	2.80365e-14
Noncompetitive inhibition dataset				
K$_M$	0.00184 ± 6 x 10^{-5}	135276 ± 382	2.81 ± 4.1 x 10^{-10}	2.81 ± 8.8 x 10^{-9}
K$_I$	0.000189 ± 6 x 10^{-6}	-	0.35 ± 2.4 x 10^{-10}	0.35 ± 1 x 10^{-9}
K$_Z$	-	0.00021 ± 4.3 x 10^{-8}	-	0.35 ± 9 x 10^{-11}
v$_{max}$	0.33 ± 0.007	7454.34 ± 1.4	4.62 ± 3 x 10^{-9}	4.62 ± 1.2 x 10^{-9}
SSE	0.000445374	2.17735e-06	4.79886e-24	2.81224e-25
Mixed inhibition dataset				
K$_M$	8.4 x 10^{-5} ± 2 x 10^{-3}	383529 ± 287	14.985 ± 0.14	2.81 ± 2 x 10^{-9}
K$_I$	7.8 x 10^{-6} ± 10^{-4}	-	1.7 ± 0.04	0.35 ± 3 x 10^{-10}
K$_Z$	-	0.0004 ± 7 x 10^{-8}	-	2.13 ± 1 x 10^{-10}
v$_{max}$	1.58 ± 2.39	20448 ± 3	5.625 ± 0.1	4.62 ± 2 x 10^{-9}
SSE	0.0038	0.000377	1.63 x 10^{-6}	2 x 10^{-24}

TABLE 3. Multiple nonlinear regression of enzyme inhibition data in the presence of noise

Parameter fit	Competitive	Uncompetitive	Non competitive	Mixed
Competitive inhibition dataset				
K_M	2.87 ± 0.14	642519 ± 0.06	142.1 ± 16.7	2.73 ± 3.9
K_I	0.3 ± 0.02	-	9.5 ± 0.6	1.17 ± 31
K_Z	-	$0.0014 ± 8.6 \times 10^{-11}$	-	5.6e+06 ± 0
v_{max}	4.62 ± 0.009	$32449 ± 2 \times 10^{-3}$	13.3 ± 12.2	4.70554± 3.9
SSE	1.063e-06	0.0220808	0.00145037	0.830483
Uncompetitive inhibition dataset				
K_M	$0.0031 ± 3 \times 10^{-7}$	2.81 ± 0.19	0.37 ± 0.04	2.67 ± 0.2
K_I	$0.0005 ± 5 \times 10^{-8}$	-	2.28 ± 0.04	243.82 ± 388
K_Z	-	2.12 ± 0.03	-	2.12532 ± 0.004
v_{max}	$1.4 ± 9 \times 10^{-5}$	4.66 ± 0.05	4.34 ± 0.06	4.62 ± 0.008
SSE	0.0159049	1.06653e-06	2.477e-06	1.07078e-06
Noncompetitive inhibition dataset				
K_M	$0.002 ± 8.3 \times 10^{-6}$	104307 ± 409	2.78 ± 0.19	0.0023 ± 0.001
K_I	$0.0002 ± 9 \times 10^{-7}$	-	0.35 ± 0.11	0.0003 ± 0.0001
K_Z	-	$0.00026 ± 7 \times 10^{-8}$	-	0.38 ± 0.02
v_{max}	$0.33 ± 9 \times 10^{-4}$	5928.66 ± 1.58	4.67 ± 1.43	4.314 ± 0.17
SSE	0.00044	3.51923e-06	1.01839e-06	1.01705e-06
Mixed inhibition dataset				
K_M	$6.2e-05 ± 1 \times 10^{-9}$	329020 ± 42	14.95 ± 0.19	1.89 ± 1.3
K_I	$5.7e-06 ± 1 \times 10^{-10}$	-	1.7 ± 0.05	0.24 ± 0.16
K_Z	-	$0.00045 ± 1 \times 10^{-8}$	-	2.16 ± 0.06
v_{max}	$1.58 ± 2 \times 10^{-5}$	17591 ± 0.45	5.63 ± 0.14	4.55 ± 0.12
SSE	0.00381337	0.000379777	2.76054e-06	8.3677e-07

Theory on the accurate estimation of Michaelis-Menten enzyme kinetic parameters from steady state and progress curve datasets

SUPPORTING MATERIALS

Datasets for the global analysis

FIGURES 2C, 2D
sampleptr.txt and samplestr.txt
Simulation settings **(Eqs. 13**) are k_1 = 5.1 µM^{-1}s^{-1}, k_{-1} = 0.5 s^{-1}, k_2 = 1.1 s^{-1}, Δt = 10^{-3} s, e_0 = 0.85 µM, s_0 = 50 µM and ρ = 10^{-2}. Totally there are 1000 replications. First column is the time in seconds, and subsequent columns are concentrations of product (sampleptr.txt) and substrate (samplestr.txt) respectively.

FIGURE 3
Simulation settings (Eqs. 13) are k_1 = 5.1 µM^{-1}s^{-1}, k_{-1} = 0.5 s^{-1}, k_2 = 1.1 s^{-1}, Δt = 10^{-3} s, e_0 = 0.85 µM and s_0 = 50 µM. Totally there are 1000 replications. First column is the time in seconds, and subsequent columns are replications of concentrations of product (sampleptr.txt) and substrate (samplestr.txt) respectively.

sampledat_ptpm1.txt, sampledat_ptpm2.txt, sampledat_ptpm3.txt respectively with ρ = 10^{-1}, 10^{-2} and 10^{-3} for product evolution.
sampledat_stpm1.txt, sampledat_stpm2.txt, sampledat_stpm3.txt respectively with ρ = 10^{-1}, 10^{-2} and 10^{-3} for substrate evolution.

FIGURES 2E, 2F, 4 and 6
Datasets were generated by numerical integration of **Eqs. 13** as given in main text with the following settings k_1 = 5.1 µM^{-1}s^{-1}, k_{-1} = 0.5 s^{-1}, k_2 = 1.1 s^{-1}, ρ = 10^{-2}, Δt = 10^{-3} s, e_0 = 0.85 µM, s_0 was iterated from 50 to 250 with interval 25 µM. Totally 1000 replications were generated for the global analysis as in **Fig. 5** of the main text. The data files are named as "sample_enzkin_**w**t_noise_data_R_**N**.dat" where N is the replication number taking values from 0 to 999 and w = s or p depending on the trajectory of substrate or product. The first column is time and columns 2 to 10 represents the data corresponding to different initial substrate concentrations s_0 viz. 50, 75, 100…250 µM. Time course sample datasets on the trajectories of substrate depletion were using in direct nonlinear least square fit with **Eqs. 1** to obtain rate constants $[k_1, k_{-1}, k_2]$ using Marquardt-Levenberg algorithm as depicted in methods section. Using these fit values K_M and v_{max} were calculated as $K_M = (k_{-1} + k_2)/k_1$ and $v_{max} = k_2 e_0$. Analysis over 1000 such time course datasets, one obtains 1000 such values of $[k_1, k_{-1}, k_2, K_M, v_{max}]$ which were used

to construct the histogram as depicted in **Figs. 6**. Trajectories on the product evolution were used for the steady state analysis by computing average velocities (p / t) at a given reaction time across various substate concentrations and replications.

Fig. 2E: "sample_enzkin_**p**t_noise_data_R_**0**.dat", first and second columns were used as time course data (s_0 = 50 µM) to NLS fit of (p, t).

Fig. 2F: "sample_enzkin_**s**t_noise_data_R_**0**.dat", first and second columns were used as time course data (s_0 = 50 µM) to NLS fit of (s, t).

Sample MATLAB code for NLS fit of time course (p, t) data to Eqs. 1 of main text

```matlab
function out = enzkinfitnlsp(tdat,pdat,ez,sz,alpha)
outn = [tdat pdat];
maxt = outn(length(outn(:,1)),1);
delt = tdat(2,1)-tdat(1,1);
gx = gca;
gx.FontSize = 20;
gx.LineWidth = 2;
v = outn(:,2);
semilogx(outn(:,1),v,'LineWidth',2); hold on;
dik1 = @(k1,k_1,k2,dpram,ez,sz,maxt,delt) ((enzkinpt(k1+dpram,k_1,k2,ez,sz,delt,maxt)-
enzkinpt(k1,k_1,k2,ez,sz,delt,maxt))/dpram);
dik_1 = @(k1,k_1,k2,dpram,ez,sz,maxt,delt) ((enzkinpt(k1,k_1+dpram,k2,ez,sz,delt,maxt)-
enzkinpt(k1,k_1,k2,ez,sz,delt,maxt))/dpram);
dik2 = @(k1,k_1,k2,dpram,ez,sz,maxt,delt) ((enzkinpt(k1,k_1,k2+dpram,ez,sz,delt,maxt)-
enzkinpt(k1,k_1,k2,ez,sz,delt,maxt))/dpram);
lambda = 1e-1; lambda_l = 1e-8; lambda_h = 1e8; tune_p = 5; dpram = 1e-4; nite = 200000; stat = 1;
dof = length(outn(:,1))-3;
param = zeros(3,1); param(1:3,1) = [0.1 1 0.1]; tol = 1e-12;
out =
struct('alpha',alpha,'params',0,'total_iterations',0,'initial_lambda',0,'RMSE',0,'final_lambda',0,'tun
e_param',0,'tpvalues',0,'errDOF',0,'totDOF',0,'regDOF',0,'MSE',0,'nparam',0,'tstat',0,'covmat',0,'SE_p
arams',0,'H0_params',0,'SST',0,'SSR',0,'SSE',0,'fcal',0,'H0_f',0,'fpvalue',0,'R_square',0,'R_square_ad
j',0);
out.initial_lambda = lambda;
out.errDOF = dof;
out.regDOF = 2;
out.totDOF = length(outn(:,1))-1;
out.nparam = 3;
for w=1:nite
    if stat == 1
        jmat = [dik1(param(1,1),param(2,1),param(3,1),dpram,ez,sz,maxt,delt)';
dik_1(param(1,1),param(2,1),param(3,1),dpram,ez,sz,maxt,delt)';
dik2(param(1,1),param(2,1),param(3,1),dpram,ez,sz,maxt,delt)']';
        jtjm = jmat'*jmat;
        dy = v - enzkinpt(param(1,1),param(2,1),param(3,1),ez,sz,delt,maxt);
        if w == 1, chia = sum(dy.^2); end
    end
    for i=1:length(jtjm), jtjm(i,i) = jtjm(i,i) + jtjm(i,i) * lambda; end
    ijtjm = jtjm^-1;
    dyn = v - enzkinpt(param(1,1),param(2,1),param(3,1),ez,sz,delt,maxt);
    nparam = param + ijtjm * jmat'*dyn;
    dym = v - enzkinpt(nparam(1,1),nparam(2,1),nparam(3,1),ez,sz,delt,maxt);
    chib = sum(dym.^2);
    if abs(chia-chib) < tol || w == nite
        param = nparam;
        out.params = param;
        out.final_lambda = lambda;
        out.tune_param = tune_p;
        SSE = chib; SST = sum(v.^2) - sum(v)^2/length(v);
        out.SSE = SSE; out.SST = SST; out.SSR = out.SST-out.SSE;
        out.fcal = (out.SSR/out.regDOF)/(out.SSE/out.errDOF);
        out.fpvalue = fcdf(out.fcal,out.regDOF,out.errDOF,'upper');
        if out.fpvalue < alpha
            out.H0_f = 1;
        else
```

```matlab
                out.H0_f = 0;
            end
        SSEr = SSE/dof; RMSE = sqrt(SSEr);
        rsquare = 1 - SSE/SST;
        out.R_square = rsquare;
        rsqadj = 1-((length(v)-1)/dof) * (1-rsquare);
        out.R_square_adj = rsqadj;
        cov_mat = ijtjm * SSEr;
        out.covmat = cov_mat;
        param_err = zeros(3,2);
        out.MSE = out.SSE/out.errDOF;
        param_err(:,1) = param - (abs(tinv(alpha/2,dof)) * sqrt(diag(cov_mat)));
        param_err(:,2) = param + (abs(tinv(alpha/2,dof)) * sqrt(diag(cov_mat)));
        out.tstat = param.*sqrt(diag(cov_mat)).^-1;
        out.tpvalues = zeros(3,1); out.H0_params = zeros(3,1);
        for ip=1:3
            out.tpvalues(ip,1) = tpval(out.tstat(ip,1),out.errDOF);
            if out.tpvalues(ip,1) < alpha
                out.H0_params(ip,1) = 1;
            else
                out.H0_params(ip,1) = 0;
            end
        end
        SET = (abs(tinv(alpha/2,dof)) * sqrt(diag(cov_mat)));
        out.SE_params = param_err;
        disp(['param (k1, k_1, k2), ' 'param_low, ' 'param_high, ' 'Std. Err']);
        disp([param param_err SET]);
        disp(['total iterations, ' 'SSE, ' 'RMSE, ' 'rsquare, ' 'rsqadj, ' 'lambda']);
        disp([w SSE RMSE rsquare rsqadj lambda]);
        out.total_iterations = w;
        out.RMSE = RMSE;
        break;
        end
    if chib < chia
        if lambda/tune_p >= lambda_l, lambda = lambda/tune_p; end
        param = nparam;
        chia = chib;
        stat = 1;
    else
        if lambda * tune_p <= lambda_h, lambda = lambda * tune_p; end
        stat = 0;
    end
end
% get the NLS fit dataset
qpt = enzkinpt(param(1,1),param(2,1),param(3,1),ez,sz,delt,maxt);
semilogx(outn(:,1),qpt,'LineWidth',2,'MarkerSize',10);
hold off;
legend('Simulation','NLS fit');
legend('boxoff');
xlabel('t'); ylabel('p');
pt = gca; pt.FontSize = 20;
xscale('log');
yscale('log');
end

function out = enzkinpt(k1,k_1,k2,ez,sz,delt,tot)
nt = ceil(tot/delt);
out = zeros(nt,1);
t = 0; pt = 0; xt = 0; i = 1;
while t <= tot
    out(i,1) = pt;
    xt = xt + delt * (k1*(ez-xt)*(sz-xt-pt)-(k_1+k2)*xt);
    pt = pt + delt * k2 * xt;
    t = t + delt;
    i = i + 1;
end
end
```

Sample MATLAB code for NLS fit of time course (s, t) data to Eqs. 1 of main text

```matlab
function out = enzkinfitnlss(tdat,sdat,ez,sz,alpha)
outn = [tdat sdat];
maxt = outn(length(outn(:,1)),1);
delt = tdat(2,1)-tdat(1,1);
gx = gca;
gx.FontSize = 20;
gx.LineWidth = 2;
v = outn(:,2);
semilogx(outn(:,1),v,'LineWidth',2); hold on;
dik1 = @(k1,k_1,k2,dpram,ez,sz,maxt,delt) ((enzkinst(k1+dpram,k_1,k2,ez,sz,delt,maxt)-
enzkinst(k1,k_1,k2,ez,sz,delt,maxt))/dpram);
dik_1 = @(k1,k_1,k2,dpram,ez,sz,maxt,delt) ((enzkinst(k1,k_1+dpram,k2,ez,sz,delt,maxt)-
enzkinst(k1,k_1,k2,ez,sz,delt,maxt))/dpram);
dik2 = @(k1,k_1,k2,dpram,ez,sz,maxt,delt) ((enzkinst(k1,k_1,k2+dpram,ez,sz,delt,maxt)-
enzkinst(k1,k_1,k2,ez,sz,delt,maxt))/dpram);
lambda = 1e-1; lambda_l = 1e-8; lambda_h = 1e8; tune_p = 5; dpram = 1e-4; nite = 200000; stat = 1;
dof = length(outn(:,1))-3;
param = zeros(3,1); param(1:3,1) = [0.1 1 0.1]; tol = 1e-12;
out =
struct('alpha',alpha,'params',0,'total_iterations',0,'initial_lambda',0,'RMSE',0,'final_lambda',0,'tun
e_param',0,'tpvalues',0,'errDOF',0,'totDOF',0,'regDOF',0,'MSE',0,'nparam',0,'tstat',0,'covmat',0,'SE_p
arams',0,'H0_params',0,'SST',0,'SSR',0,'SSE',0,'fcal',0,'H0_f',0,'fpvalue',0,'R_square',0,'R_square_ad
j',0);
out.initial_lambda = lambda;
out.errDOF = dof;
out.regDOF = 2;
out.totDOF = length(outn(:,1))-1;
out.nparam = 3;
for w=1:nite
    if stat == 1
        jmat = [dik1(param(1,1),param(2,1),param(3,1),dpram,ez,sz,maxt,delt)';
dik_1(param(1,1),param(2,1),param(3,1),dpram,ez,sz,maxt,delt)';
dik2(param(1,1),param(2,1),param(3,1),dpram,ez,sz,maxt,delt)']';
        jtjm = jmat'*jmat;
        dy = v - enzkinpt(param(1,1),param(2,1),param(3,1),ez,sz,delt,maxt);
        if w == 1, chia = sum(dy.^2); end
    end
    for i=1:length(jtjm), jtjm(i,i) = jtjm(i,i) + jtjm(i,i) * lambda; end
    ijtjm = jtjm^-1;
    dyn = v - enzkinst(param(1,1),param(2,1),param(3,1),ez,sz,delt,maxt);
    nparam = param + ijtjm * jmat'*dyn;
    dym = v - enzkinst(nparam(1,1),nparam(2,1),nparam(3,1),ez,sz,delt,maxt);
    chib = sum(dym.^2);
    if abs(chia-chib) < tol || w == nite
        param = nparam;
        out.params = param;
        out.final_lambda = lambda;
        out.tune_param = tune_p;
        SSE = chib; SST = sum(v.^2) - sum(v)^2/length(v);
        out.SSE = SSE; out.SST = SST; out.SSR = out.SST-out.SSE;
        out.fcal = (out.SSR/out.regDOF)/(out.SSE/out.errDOF);
        out.fpvalue = fcdf(out.fcal,out.regDOF,out.errDOF,'upper');
        if out.fpvalue < alpha
            out.H0_f = 1;
        else
            out.H0_f = 0;
        end
        SSEr = SSE/dof; RMSE = sqrt(SSEr);
        rsquare = 1 - SSE/SST;
        out.R_square = rsquare;
        rsqadj = 1-((length(v)-1)/dof) * (1-rsquare);
        out.R_square_adj = rsqadj;
        cov_mat = ijtjm * SSEr;
        out.covmat = cov_mat;
        param_err = zeros(3,2);
        out.MSE = out.SSE/out.errDOF;
        param_err(:,1) = param - (abs(tinv(alpha/2,dof)) * sqrt(diag(cov_mat)));
        param_err(:,2) = param + (abs(tinv(alpha/2,dof)) * sqrt(diag(cov_mat)));
        out.tstat = param.*sqrt(diag(cov_mat)).^-1;
        out.tpvalues = zeros(3,1); out.H0_params = zeros(3,1);
        for ip=1:3
```

```matlab
            out.tpvalues(ip,1) = tpval(out.tstat(ip,1),out.errDOF);
            if out.tpvalues(ip,1) < alpha
                out.H0_params(ip,1) = 1;
            else
                out.H0_params(ip,1) = 0;
            end
        end
        SET = (abs(tinv(alpha/2,dof)) * sqrt(diag(cov_mat)));
        out.SE_params = param_err;
        disp(['param (k1, k_1, k2), ' 'param_low, ' 'param_high, ' 'Std. Err']);
        disp([param param_err SET]);
        disp(['total iterations, '  'SSE, ' 'RMSE, ' 'rsquare, ' 'rsqadj, ' 'lambda']);
        disp([w SSE RMSE rsquare rsqadj lambda]);
        out.total_iterations = w;
        out.RMSE = RMSE;
        break;
    end
    if chib < chia
        if lambda/tune_p >= lambda_l, lambda = lambda/tune_p; end
        param = nparam;
        chia = chib;
        stat = 1;
    else
        if lambda * tune_p <= lambda_h, lambda = lambda * tune_p; end
        stat = 0;
    end
end
% get the NLS fit dataset
qpt = enzkinst(param(1,1),param(2,1),param(3,1),ez,sz,delt,maxt);
semilogx(outn(:,1),qpt,'LineWidth',2,'MarkerSize',10);
hold off;
legend('Simulation','NLS fit');
legend('boxoff');
xlabel('t'); ylabel('p');
pt = gca; pt.FontSize = 20;
xscale('log');
yscale('log');
end

function out = enzkinst(k1,k_1,k2,ez,sz,delt,tot)
nt = ceil(tot/delt);
out = zeros(nt,1);
t = 0; pt = 0; xt = 0; i = 1;
while t <= tot
    out(i,1) = sz-xt-pt;
    xt = xt + delt * (k1*(ez-xt)*(sz-xt-pt)-(k_1+k2)*xt);
    pt = pt + delt * k2 * xt;
    t = t + delt;
    i = i + 1;
end
end
```