A lipidomic exploration of the effects of high-intensity interval exercise in healthy men after metformin intake

René Neuhaus[1†], Thomas Meikopoulos[2,3†], Ana Gradillas[1], Carolina González-Riaño[1], Georgios Theodoridis[2], Stefanos Nikolaidis[4], Vassilis Mougios[4], Coral Barbas[1*], Helen Gika[6,3*] and Alma Villaseñor[1,5*]

[1] Centro de Metabolómica y Bioanálisis (CEMBIO), Facultad de Farmacia, Universidad San Pablo-CEU, CEU Universities, Urbanización Montepríncipe, 28660, Boadilla del Monte, Spain.

[2] Department of Chemistry, Aristotle University of Thessaloniki, Greece.

[3] Biomic AUTh, Center for Interdisciplinary Research and Innovation (CIRI-AUTH), Balkan Thessaloniki, Greece

[4] Laboratory of Evaluation of Human Biological Performance, School of Physical Education and Sport Science at Thessaloniki, Aristotle University of Thessaloniki, Greece.

[5] Departamento de Ciencias Médicas Básicas, Instituto de Medicina Molecular Aplicada (IMMA) Nemesio Díez, Facultad de Medicina, Universidad San Pablo-CEU, CEU Universities, Boadilla del Monte, Spain.

[6] Department of Medicine, Aristotle University of Thessaloniki, Greece.

[†]These authors have contributed equally

[*]Corresponding authors

Corresponding authors:

Coral Barbas

CEMBIO, Centro de Metabolómica y Bioanálisis, Facultad de Farmacia, Universidad San Pablo CEU. Avda. Montepríncipe s/n, 28668 Boadilla del Monte, Madrid, SPAIN.

c.barbas@ceu.es

Helen Gika

Department of Medicine and Biomic AUTh, Aristotle University of Thessaloniki, Greece

gkikae@auth.gr

Alma Villaseñor

CEMBIO, Centro de Metabolómica y Bioanálisis, Facultad de Farmacia, Universidad San Pablo CEU. Avda. Montepríncipe s/n, 28668 Boadilla del Monte, Madrid, SPAIN.

alma.villasenor@ceu.es

ABSTRACT

We have previously found that high-intensity interval exercise (HIIE) affected metformin pharmacokinetics, causing higher maximal plasma concentration compared with rest. In this scenario, changes in individual lipids could play an important role. The prolonged responses of the lipidome to HIIE have not been explored.

Thus, this study aimed to explore differences in the plasma lipidomic profiles between HIIE and rest, both under metformin treatment.

Nine healthy males participated in two sessions where they received 1,000 mg of metformin. In session A, they performed HIIE at an average intensity of 67% of maximum heart rate for a total duration of 76 min, whereas in session B they rested. Plasma was collected before taking metformin and during each session (in total 14 time points spanning 12 h). Samples were analysed through lipidomics using mass spectrometry. Paired Wilcoxon tests between sessions were applied for statistics.

We found several variations in the lipid profiles due to HIIE, which persisted until 4 h post-exercise. The main discriminant lipid classes were fatty acids, acyl carnitines, glycerophosphocholins, sphingomyelins, and triglycerides. These changes were followed in time up to 12 h, showing the effect of the meals taken during the session. We hypothesize the changes are a synergic effect of HIIE and metformin in the lipidome with the effect of HIIE being the predominant.

These findings provide important insights into the dynamic and complex physiological response of humans to intensive exercise under metformin intake.

Keywords: LC-MS, HIIE, acyl carnitines, fatty acids, triglycerides.

Word count: 3,077

INTRODUCTION

Maintaining adequate levels of physical activity is essential for good health. Physical activity is widely recommended to prevent and treat obesity-related complications such as type 2 diabetes mellitus (T2DM)(1). Recent studies have shown that high-intensity interval exercise (HIIE) which includes alternating short periods of intense exercise with brief recovery periods, is effective in improving cardiometabolic markers in patients with insulin resistance and T2DM(2–4). Physical exercise is known to produce large changes in the concentrations of multiple metabolites(5), including lipids(6). Among the latter, ceramides (Cer), diglycerides (DG), and acyl carnitines (CAR) have been shown to activate intracellular signalling cascades and regulate metabolic pathways(7–11). It has been described that physical exercise induces lipid alterations such as an increased utilization of triglycerides (TG) that in exchange increase the concentration of fatty acids (FA) and CAR(12). Other studies have addressed the regulation of the TG in the skeletal muscle upon exercise(13,14), The overall regulation of circulating lipid species to HIIE is poorly understood. To cope with this challenge, lipidomic analysis employing high throughput techniques, such as liquid chromatography coupled to mass spectrometry (LC-MS) can be used, as it allows the acquisition of a lipid profile consisting of numerous lipid species(2–4).

Metformin is a first line antihyperglycemic drug for T2DM treatment(15). As a first step in the exploration of the relationship between metformin administration and HIIE, we have previously shown that HIIE during the absorption phase of metformin affected metformin pharmacokinetics, causing higher maximal plasma concentration (C_{max}) compared with rest without compromising glucose homeostasis in healthy men(16). In this scenario, changes in the plasma concentrations of individual lipids could play an important role in the response to HIIE. However, the way that the plasma lipidome changes, over consecutive hours in a controlled environment of food intake and either HIIE or rest, has not been explored.

Thus, the aim of this work was to explore the changes in the plasma lipidome of healthy individuals for 12 h after metformin intake followed by either HIIE or rest.

To our knowledge this is the first study reporting HIIE effect on the blood lipidome of healthy individuals over a long period of time covering exercise session and post-exercise, using a comprehensive, untargeted lipidomic approach.

RESULTS

To study the effect of exercise on the lipidome, we collected plasma samples from nine healthy men that participated in two sessions that lasted 12 h, one including 76 min of HIIE, the other one resting **(Figure 1)**. In both sessions the participants took 1,000 mg of metformin at the beginning. For practical reasons, from all collected samples, a subgroup was analyzed via reverse phase ultra high-performance liquid chromatography coupled to mass spectrometry with an electrospray ionization source and a quadrupole–time of flight mass analyzer (RP-UHPLC-ESI-QTOF-MS).

Figure 1.

Study design.

Outline of the two sessions showing food intake, metformin intake, blood sampling, and exercise. Blood
samples in red means samples were analysed using lipidomics.

The samples collected at the same time point in both sessions (T0-T5) were analyzed and
compared (**Figure S1**). The untargeted lipidomic analysis yielded a total of 1,303 and 299
chemical features in ESI(+) and ESI(−) modes, respectively. The unsupervised model of these
features using PCA showed tight clustering of QC samples in the plots for each ionization mode
(**Figure S2A, B**). After lipid annotation from both ionization modes and removal of duplicates, a
single data matrix was obtained consisting of 247 unique lipids (**Table S1**). The PCA model of
these data continued to show clustering of QC samples (**Figure S2C**) indicating good quality of
the data and that the variation shown is due to inter-individual variation of lipids in the plasma of
the participants. Interestingly, PCA models based on the samples alone from all-time points did
not reveal any discrimination between rest and exercise sessions (**Figure S2D-F**). Thus, further
statistical analysis was applied to get a deeper insight into the data. No sample outliers were
detected with the profile of 247 annotated lipids (**Figure S2F**).

Lipidomic changes after HIIE involve a complex network of different lipid classes.

To study the effects of HIIE on the plasma lipidome, the time points T0-T5 were considered
separately and lipidome was compared under rest and exercise conditions using the Wilcoxon
test. Specific focus was set on a time frame up to 6 hours (**Figure S1**), where early effects might
be reflected by HIIE. Univariate statistical analysis showed several lipid differentiations (**Figure
2A**, **Tables S2-S5**).

Figure 2.

Significantly different lipid species between exercise and resting session under metformin intake.

A) Stack bar plot of significantly different lipids (p < 0.05, pFDR< 0.2) per lipid species at different time
points after Wilcoxon test between exercise and resting session for healthy subjects taking metformin.
Abbreviations: acyl carnitines (CAR), fatty acids (FA), lysoglycerophosphocholines (LPC), glycerol-
phosphocholines (PC), ether-glycerophosphocholines (PC-O/P), lysoglycerophosphoethanolamines (LPE),
sphingomyelins (SM), ceramides (Cer), hexosylceramides (HexCer), cholesteryl esters (CE), triglycerides
(TG). **B)** Venn diagram showing shared lipids between T3 (after finishing HIIE), T4 (1 h post-HIIE) and
T5 (4 h post-HIIE), the intensity of the grey color corresponds to the amount of shared lipids. Abbreviations:
high-intensity interval exercise (HIIE).

Starting with T0 (1.5 h after breakfast) it was found that 34 lipids had significantly different levels
between the two sessions (p < 0.05, pFDR < 0.2). These lipids belong to TG,
glycerophosphocholines (PC), lysoglycerophosphocholines (LPC), and
lysoglycerophosphoethanolamines (LPE). Their connections were presented using lipid networks
by Linex as shown in **Figure S3A**. As can be seen, most of the TGs, PCs, and LPCs were
connected within their classes, with TGs (e.g. TG(44:1), TG(42:1), and TG(42:0) exhibiting the
highest fold change in exercise vs rest. All these lipids were higher in the exercise session before
HIIE mostly probable due to a food intake effect (**Table S2**).

These lipid differences found at T0 seem to fade out at T1 and T2, which correspond to prior and
40 minutes into exercise, respectively. Interestingly, it was after 76 minutes of HIIE, at T3, that
52 lipid species (**Table S3**) emerged as statistically different from resting conditions ($p < 0.05$,
pFDR < 0.2). The main lipid classes of these species were sphingomyelins (SM), PC, TG, and
FA, followed by CAR and Cer (**Figure 2A**). The lipid network suggested again that most of the
lipids were connected within their classes and that acetylcarnitine [CAR(2:0)] and palmitoleic
acid [FA(16:1)] were the lipids with the highest fold change (**Figure 3A**). Almost in all cases,
except for LPE(22:6), increased levels of these lipids were observed in the plasma of the exercise
group (**Table S3**).
After almost one hour of finishing HIIE at T4 only 15 significant lipid species were significantly
different, in almost all cases increased with the exception of LPC(16:0), compared with resting
conditions ($p < 0.05$, pFDR < 0.2). These included CAR(2:0) and members of the FA and PC
classes (**Figure S3B**, **Table S4**). The lipid network shown in **Figure S3B** revealed the importance
of CAR(2:0) and FAs such as, FA(20:1), FA(20:2), and FA(18:2). Moving away from the
completion of the exercise session, three hours later (at T5, six hours from beginning of the
session and after banana intake), 64 lipids were found increased in the exercise group (**Table S5**).
Again, the lipid classes that mainly found differentiated were PC, SM, and TG (**Figure 2A**). Lipid
network pointed out additional lipid classes, such as FA and CAR, which showed high fold
changes (**Figure 3B**). A Venn diagram comparing the significant lipid changes between T3 (after
finishing HIIE), T4 (1 h post-HIIE) and T5 (4 h post-HIIE) is shown in **Figure 2B**. In this figure,
we observed that T3 and T5 shared around 50% of the significant lipids showing the strong effect
of HIIE on the lipidome even after 4 h post-exercise.
To understand better the changes observed, the levels of some of the most altered lipid species
due to exercise such as CAR(2:0), FA(16:1), FA(20:1), FA(20:2), FA(20:3), and FA(20:4) were
plotted for the two sessions (**Figure 4**). It can be observed that CAR(2:0) increased dramatically
after starting HIIE (T2) and continued increasing, reaching its highest level right after finishing
exercise (T3). One hour afterwards (T4), a decrease was observed, nevertheless, even 4h after
the termination of HIIE (T5) the level is still higher than in resting conditions.
In the case of the FA, a rise in their level was also observed, which however continued to increase
after the completion of HIIE, reaching a maximum at T4, 1 h after HIIE. Similarly to CAR(2:0),
most FAs did not return to the levels at resting conditions at T5.
Other cases of altered lipids, such as PC(16:0_18:0), LPE(22:6), SM(d40:1), Cer(d18:1/22:0),
and some TGs; TG(56:2), and TG(42:0), exhibited diverse trajectories indicating mainly an
increased effect around the exercise time window (T1-T3) that tends to get back at rest levels later
on (**Figure S4**). PC(16:0_18:0), SM(d40:1), Cer(d18:1/22:0), TG(56:2), and TG(42:0) increased
upon HIIE (to a lower extent compared to CAR and FA) whereas LPE(22:6) showed a decreasing
trend.

Figure 3.

Lipid networks of the statistically significant lipids at T3 and T5.

Lipid network at **A)** T3 (2:05 h) just after finishing HIIE and **B)** T5 (6:00 h) 4 hours after finishing HIIE

and after a snack. The size of the dots represents the fold change for individuals at exercise session against

resting conditions. The color of the dots represents the lipid class. All lipid names are depicted as the

composition ("_"), to see the *sn*-positions ("/") see Tables S3 and S5. Lipids with the same sum composition

are represented as one lipid. Abbreviations: acyl carnitines (CAR), fatty acids (FA), lysoglycerophospho-

cholines (LPC), glycerophosphocholines (PC), ether-glycerophospho-cholines (PC-O/P), lysoglycero-

phosphoethanolamines (LPE), sphingomyelins (SM), ceramides (Cer), hexosylceramides (HexCer),

cholesteryl esters (CE), triglycerides (TG).

Figure 4.

Lipid trajectories for lipids with high fold changes due to HIIE from T0 (0 h) to T5 (6 h).

Abundances in exercise and resting sessions of CAR(2:0), FA(16:1), FA(20:1), FA(20:2), FA(20:3) and

FA(20:4). Values are represented as mean of normalized area ± SEM, significant differences were found

using Wilcoxon test; * p< 0.05; ** p< 0.01.

Trajectories of exercise-induced lipid changes over 12 hours show interaction with meals

To further examine the trend of lipids most affected by exercise beyond 6 h (T5), all 14 sampling points of the exercise session, up to 12 h (T9), were plotted (**Figure 5**). These graphs show that CAR(2:0) continued decreasing, reaching baseline (that is, the pre-exercise values) at 12 h, regardless of the food intake in between. In the case of the FA, the downward trajectory starting after the completion of exercise already described above, was interrupted by the meal taken at 6.5 h, after which the FA concentrations dropped again (**Figure 5**). This pattern was more pronounced with FA(16:1) but less with FA(20:1), FA(20:2), FA(20:3), and FA(20:4).

The 12 h trajectories of the lipid species that were affected less by exercise are presented in **Figure S5**. In these cases of lipids, like PC(16:0_18:0), LPE(22:6), SM(d40:1), and Cer(d18:1/22:0) the levels after 6 h did not change considerably and no impact of food intake was detected. Regarding the altered TGs, it was found that TG(56:2) strongly increased after the meal, peaking at 10 h, whereas TG(42:0) increased to a lower extent and only transiently after the meal, after which it continued dropping.

Overall, the trends of all annotated lipids in relation to exercise can be seen in the heatmap which shows the average intensities at each of the 14 time points from 0 to 12 h (**Figure 6**). It can be clearly seen that plasma CAR, FA, SM, some specific TG (saturated and unsaturated), and PC increased during and after HIIE (T2 and T3), whereas LPC decreased. From the increased lipids, TG and PC decreased 30 min post-exercise, whereas CAR and FA remained elevated until 2 hours after HIIE (T4b-T4c). It can be also noted that, after the meal (T6 and on), mainly PC, LPC, and TG (especially polyunsaturated TG) increased.

Figure 5.

Lipid trajectories for lipids with high fold changes due to HIIE from T0 (0 h) to T9 (12 h).

Abundances in exercise and resting sessions of CAR(2:0), FA(16:1), FA(20:1), FA(20:2), FA(20:3) and FA(20:4). Values are represented as mean of normalized area ± SEM, significant differences were found using Wilcoxon test; * p< 0.05; ** p< 0.01.

235   **Figure 6.**

236
237   **Trends of lipid classes.**
238   Heatmap of average abundances of all annotated lipids at all time points (12 h) in exercise session.
239   Abbreviations: acyl carnitines (CAR), fatty acids (FA), sphingomyelins (SM), triglycerides (TG),
240   lysoglycerophosphocholines (LPC), glycerophosphocholines (PC).

DISCUSSION

It has been reported that HIIE interventions may play a role in improving long-term
cardiovascular health(17) via favourable changes in the blood lipid content such as decreasing
low-density lipoprotein (LDL) and total cholesterol (TC) and increasing high-density lipoprotein
(HDL), following high-intensity exercise(18). Increased exercise intensity correlates with high
levels of catecholamines, cortisol, and growth hormone which have a lipolytic effect and in
combination to the additional increase in fat oxidation that occurs, it can positively affect levels
of circulating lipids(19). The lipolytic effect occurs as breakdown of TG in adipose tissue releases
free fatty acids (FFA) into the bloodstream. FFA serve as an energy source for muscles and other

tissues but their oxidation is suppressed during high-intensity exercise (above ~75% VO₂ max) as

a shift toward carbohydrate metabolism occurs(20). In parallel, muscle contraction increases

lipoprotein lipase (LPL) activity, which leads to declined levels of circulating TG(21).

Even a single HIIE session can lead to immediate TG reduction, temporary increase in HDL, and

sustained fat oxidation post-exercise(22–25). Increased fat oxidation is sustained post-HIIE and

it is shown that FFA remain elevated for a period as the body continues to rely on fat metabolism

for recovery and decreased levels of TG may persist for hours after exercise due to enhanced

clearance by LPL(26). However, the findings are diverse as the effect on lipid metabolism is

related to the intensity of the exercise and the variations in the HIIE protocol applied.

Up to now, scarce data exist on the effect of HIIE by comprehensive lipidomic analysis. Studies

are very limited on the effect of HIIE on the plasma lipidome of healthy individuals and have

been performed only for comparative purposes to other type of exercise after repetitive sessions

over a period of time(27), either in combination with diet(28) or in comparison to a disease

group(3). In most of the cases, studies performed of the effect of HIIE present data limited to

typical blood lipid parameters like cholesterol, LDL, HDL and TG(29,30). In these cases, positive

impact on lipid metabolism has been shown. However, there is no further evidence on the effect

of HIIE on diverse lipid species. In this study, by performing untargeted lipidomic analysis we

were able to capture information for 247 unique lipid species from various classes.

Our findings show that significant changes occur in different lipid species during and after HIIE,

suggesting complex correlations networks. CAR and FA species showed to be mostly affected

with a steep increase in their levels, while other classes of lipids such as SM, PC, LPE, Cer and

TG also showed alterations.

The applied methodology captures comprehensively the impact of HIIE on the plasma lipidome,

thus changes in lipid metabolism during and after HIIE over a period of 12 h in healthy individuals

are shown for the first time.

In one of the very few relevant lipidomic studies where a wide range of lipid classes was assessed

in serum with the aim to compare a 6-week moderate-intensity continuous training group with

sprint interval training(31), alterations in various classes of lipids have also been observed.

SM(d18:1/22:5), SM(d18:1/22:6), MG(18:2), LPE(18:1), LPE(20:1), PC(O-16:0/16:0), and

PI(16:1/18:1) were found increased, while most PC, Cer and hexosylceramides (HexCer) were

decreased. Certainly, that study focuses on different training protocols, however, its findings via

a holistic approach indicate the diversity of lipids altered as a result of exercise.

In our study we have observed acute increase of acetyl carnitine, CAR(2:0), which has been

reported in several studies, suggesting that exercise leads to increased carnitine release into the

plasma, especially when glycogen stores are depleted, as the body shifts more towards fat

metabolism. The rapid increase in acetyl-CoA production and the role of carnitine as acetyl

buffering in skeletal muscle metabolism led to CAR(2:0) build up in muscle which then is released
into the bloodstream(32).
Significant FA rise observed in the study is also in agreement with lipolysis activation and break
down of stored TGs expected to occur in the training session(33). SM and ceramides found
elevated are known to be involved in stress signalling and muscle adaptation, whereas PC are
major component of cell membranes that may play a role in lipid remodeling to maintain
membrane integrity during HIIE and muscle cells stress(34,35). Lipid mobilization and sharp
increase in energy demand during HIIE can also explain the observed increase of some TG.
These observations are the reflection of complex metabolic perturbations due to a strong factor
such as intense exercise, while metformin intake should not be disregarded.
Although earlier, based on studies measuring basic lipid characteristics e.g. TG, cholesterol and
HDL, LDL, it was considered that acute metformin intake did not affect circulating lipid
concentrations(36), more recent studies applying comprehensive lipidomic analysis approaches
have indicated that metformin can alter plasma lipid composition, including decrease in saturated
fatty acids, associated with insulin resistance, increase in polyunsaturated fatty acids and altered
ceramide levels. Dahabiyeh *et al*(37) have recently provided evidence by applying a mass
spectrometry shotgun lipidomic approach that a single dose of 500 mg metformin induced acute
alterations in the plasma lipidome of healthy subjects. Thirty-three lipids were significantly
increased and 192 decreased in plasma in response to acute metformin intake. The altered lipids
are mainly involved in arachidonic acid metabolism, steroid hormone biosynthesis, and
glycerophospholipid metabolism. It was also shown that metformin had the most impactful effect
on lipidome at its Cmax. Furthermore, several lipids displayed a similar change in their levels
compared to metformin, among which are monoacylglycerols; a larger number of lipids acted in
an opposed manner to metformin levels, including fatty acyls (fatty esters, fatty acids, and
eicosanoids), sterol lipids, glycerolipids, and glycerophospholipids (PC,
glycerophosphoinositols)(37). In an earlier study from the same group(38), several lipids were
revealed with opposed trends to metformin patterns, mainly involving arachidonic and linoleic
acid metabolism and including glycerophospholipids (such as lysoglycerophosphates (LPA) and
LPC). It was concluded that the pleiotropic effect of metformin probably is associated with its
effect in lipid metabolism and lipid signalling pathways(38). Hence, we could hypothesize that,
although to a smaller extent than exercise, metformin intake could have played a role on the lipid
dysregulation observed in this study and that a combinatorial effect in lipid metabolism might
have been captured with the HIIE being the predominant. Taking into the account that in one of
our previous studies in healthy male individuals, we observed that the levels of metformin in
plasma were higher during HIIE compared to resting conditions(16), an increased impact might
be expected.

It could also be mentioned that a similar trend with that described by Dahabiyeh *et al*(37)
regarding the similar trend of altered lipids with the metformin levels along time have been noted
in our study. CAR(2:0) maxima coincided with that of metformin and they seem to follow similar
trajectories over time (**Figure S6A**). A similar curve trend of CAR found in our study was also
noted by Dahabiyeh *et al*(37). On the other side, TG(56:2) shows an opposite trend to that of
metformin overtime (**Figure S6B**).

These findings could be of high relevance for pathologies such T2DM, where metformin is
administered and routine exercise is recommended. Recently HIIE has been popularized among
nonathletes for improving cardiovascular health(39), weight loss(40), and insulin sensitivity(41).
Providing deeper insight on the biochemical modulations triggered by high intensity exercise
could provide further knowledge to enhance its beneficial impact in health, as the complex
metabolic effects that are induced are yet not fully understood.

Definitely, further investigation is required to enlighten the biochemical processes behind the
current findings and to evaluate their importance to improve the contribution of intensive exercise
in the battle against metabolic diseases such as obesity and diabetes. To our knowledge, this is
the first study that monitored the effect of HIIE on the plasma lipidome under metformin for a
period of 12 h. A limitation of the study is the small number of participants, which was primarily
due to its complex and demanding design. Additionally, as based on previous evidence, acute
intake of a therapeutic dose of metformin had probably modest effect on plasma lipids, therefore
the interaction between metformin and HIIE effect is difficult to interpret. Further studies with
larger sample sizes and a control group without metformin intake are needed to confirm or refute
our findings.

All in all, this study helps to understand better the highly dynamic lipid alterations in blood by
HIIE after metformin intake. The main discriminant lipid classes were fatty acids, acyl carnitines,
glycerophosphocholins, sphingomyelins, and triglycerides. These lipid classes continued to be
significant up to 4 h after finishing HIIE.

MATERIALS AND METHODS
Study design
This work is part of a wider study examining the effect of exercise on the pharmacokinetics of
metformin(16). In this, nine healthy, non-smoking male subjects participated in two sessions, A
and B, each lasting 12 h, performed in random counterbalanced order with one or two weeks in
between. They were not taking any medication and abstained from drinking alcohol and coffee
for 72 and 12 h, respectively, before each experimental session, as well as during the sessions.
Schematically, the study design is presented in **Figure 1**.

In both sessions, the participants received 1,000 mg of metformin (Glucophage®, immediate-release tablets) at around 10 am, 1-1.5 hours after receiving a light, low-fat, high-carbohydrate breakfast. In session A, they performed a HIIE protocol with an average intensity of 67% of maximum heart rate (HRmax) and a total duration of 76 min. HIIE was performed 45 min after taking metformin and consisted of 5 min of warm-up by walking at 4-5 km/h, followed by 11 sets of 6 min, each consisting of 3 min of walking or running at 90% of HRmax and 3 min of walking at 4-5 km/h. Between sets 6 and 7, a 5-min rest was included. The participants then rested throughout the remainder of the day. In session B, they rested throughout the whole day.

In both sessions, the participants consumed one or two bananas at 3.5 h after taking metformin and a regular serving of spaghetti at 6.5 h. The study was in accordance with the Helsinki Declaration and was approved by the Research Ethics Committee of the School of Physical Education and Sport Science at Thessaloniki, Aristotle University of Thessaloniki, following the Regulation of Principles and Operation of the Research Ethics Committee of the Aristotle University of Thessaloniki and the applicable legislation (approval number 113/2022).

Sample collection and processing

Fourteen venous blood samples were collected in EDTA tubes in each session at the Laboratory of Evaluation of Human Biological Performance as follows: before taking metformin (0 h, T0) and after taking metformin at 40 min (T1), 1 h 25 min (T2), 2 h 5 min (T3), 2 h 30 min (T3a), 3 h (T4), 3 h 30 min (T4a), 4 h (T4b), 4 h 30 min (T4c), 6 h (T5), 7 h (T6), 8 h (T7), 10 h (T8) and 12 h (T9) (**Figure 1**). Plasma was collected by centrifugation and stored at –80 ºC until shipment to CEMBIO, where it was kept at –80 ºC until analysis. For practical reasons, only samples taken at T0, T1, T2, T3, T4, and T5 of session B were subjected to lipidomic analysis, whereas all samples taken during session A were analyzed.

Reagents and chemicals

For all aqueous solutions, reverse-osmosed ultrapure water was obtained from a Milli-Qplus185 system (Millipore, Billerica, MA, USA). LC-MS-grade isopropanol, methanol, and acetonitrile were purchased from Fisher Scientific (Pittsburgh, PA, USA). Methyl *tert*-butyl ether (MTBE) (HPLC grade, ≥ 99.8%), ammonium fluoride (ACS reagent, ≥ 98%), and palmitic acid-d31 were purchased from Sigma-Aldrich (Steinheim, Germany). Analytical grade ammonia solution (28%, GPR RECTAPUR®) and glacial acetic acid (AnalaR® NORMAPUR®) were obtained from VWR Chemicals (Radnor, PA, USA). Sphinganine (d17:0) primary standard and LightSPLASH® LIPIDOMIX® Quantitative Mass Spec Primary Standard were obtained from Avanti Polar Lipids (Alabaster, AL, USA).

Sample preparation for lipidomics

Lipidomics analysis was carried out following previously published methodology (42). Plasma samples were thawed on ice for approximately 1 h and vortex-mixed for 2 min. Then, 50 μL of each sample were transferred to an Eppendorf vial on ice. Subsequently, 175 μL of ice-cold methanol (stored at −20 °C), containing 2.3 ppm of sphinganine (d17:0) and 4.6 ppm of palmitic acid-d31 as internal standards, were added for deproteinization. After shaking the sample for 1 min, 175 μL of MTBE and 10 μL of LightSPLASH LIPIDOMIX Quantitative Mass Spec Primary Standard mix were added. Samples were vortex-mixed for 30 min and centrifuged for 15 min at 15 °C and 16,100 x *g*. Finally, 100 μL of supernatants were added to chromatography vials, equipped with 300-μL inserts (one for each ionization mode, positive and negative), centrifuged for 5 min at 15 °C and 2,000 x *g*, and then injected into the LC-MS system (**Figure S7A**).

In addition, 7 types of quality control samples (QC) were prepared to control the quality of the analysis and to facilitate lipid annotations. First, a "QCpool" sample was prepared by mixing 5 μL of each of the 180 analyzed samples. Then, the rest of the QC samples were used for iterative MS/MS experiments, a "QC0 (A+B)" sample was prepared by mixing 15 μL of samples taken at T0 of both sessions (18 in total). Additionally, a "QC1 (A)" sample was prepared by mixing 25μL of samples collected at T1 of session A (9 in total). Similarly, "QC5 (A)" (T5 of session A), "QC1 (B)" (T1 of session B), "QC5 (B)" (T5 of session B), and "QC9 (B)" (T9 of session B) samples were prepared.

Analysis in Liquid chromatography coupled to QTOF mass spectrometry analysis (LC-QTOF-MS)

Samples were analyzed using an Agilent 1290 Infinity II UHPLC system coupled to an Agilent 6545 quadrupole time-of-flight (QTOF) mass spectrometer (Agilent, Santa Clara, CA, USA). An Agilent InfinityLab Poroshell 120 EC-C18 (3.0 × 100 mm, 2.7 μm) column, protected by a compatible guard column (Agilent InfinityLab Poroshell 120 EC-C18, 3.0 × 5 mm, 2.7 μm), was employed for the analysis. The column temperature was set at 50°C, while the autosampler's temperature was set at 15 °C to preserve compound stability and prevent lipid precipitation. Data were acquired in both ionization modes in separate runs. For positive ionization (ESI+), the injection volume was set at 0.5 μL, while, for negative ionization (ESI-), at 1 μL. Elution was performed with a solvent system consisting of (A) 10 mM ammonium acetate, 0.2 mM ammonium fluoride in water/methanol (9:1, v/v) and (B) 10 mM ammonium acetate, 0.2 mM ammonium fluoride in acetonitrile/methanol/isopropanol (2:3:5, v/v/v). The elution program applied was 70% B at 0–1 min, linear gradient to 86% B until 3.5 min, held until 10 min, linear gradient to 100% B until 11 min, and held until 17 min. Then, the system returned to the initial

condition to achieve column equilibration for a total of 2 min. Thus, the total analysis time was
19 min.

The QTOF mass spectrometer, equipped with a dual atmospheric jet stream electrospray
ionization (ESI) ion source, was configured with the following parameters: 150 V fragmentor, 65
V skimmer, 3500 V capillary voltage, 750 V octopole radio frequency voltage, 10 L/min nebulizer
gas flow, 200 °C gas temperature, 50 psi nebulizer gas pressure, 12 L/min sheath gas flow, and
300 °C sheath gas temperature. In full scan mode, operated from 50 to 1.800 m/z with a scan rate
of 3 spectra/s, data were collected for both ionization modes. A solution of two reference mass
compounds was used throughout the analysis: purine ($C_5H_4N_4$) at m/z 121.0509 for the positive
and m/z 119.0363 for the negative ionization modes; and HP-0921 ($C_{18}H_{18}O_6N_3P_3F_{24}$) at m/z
922.0098 for the positive and m/z 980.0163 (HP-0921 + acetate) for the negative ionization
modes. These masses were continuously infused into the system to provide constant mass
correction.

At the end of the analytical run, the 7 types of QC were used to perform iterative runs of tandem
mass spectrometry (MS/MS, 5 for each type) in both ionization modes. They were analyzed with
MS and MS/MS scan rates of 3 spectra/s, 40-1,700 m/z mass window, narrow (~ 1.3 amu) MS/MS
isolation width, 3 precursors per cycle, 5,000 counts, and 0.001% of MS/MS threshold. Five
iterative-MS/MS runs for each QC sample were set at a collision energy of 20 eV, and the
subsequent five runs for each QC sample were performed at 40 eV. To avoid inclusion in the
iterative-MS/MS data, reference masses and contaminants detected in blank samples were
excluded from the analysis.

Data treatment

The Agilent MassHunter Profinder B.10.0.2 software was used to reprocess the data collected
after the LC-MS analyses in both positive and negative ion modes, yielding 1,303 and 299
features, respectively. The datasets were extracted using the Batch Recursive Feature Extraction
(RFE) algorithm integrated into the software. The selected adducts were the following: [M+H]+,
[M+Na]+, [M+K]+, and [M+NH$_4$]+ in ESI(+); [M-H]−, [M+CH$_3$COO]−, and [M+Cl]− in ESI(−).
The neutral loss of water was also considered for both ion modes.

Prior to statistical analysis, quality assurance (QA) was accomplished by data filtration and
normalization. Missing values were estimated using the k-Nearest Neighbor algorithm(43) in
MATLAB (2018, MathWorks, Netik, MA, USA). Then, the features were filtered based on the
coefficient of variation of the QC samples, with a cutoff threshold of 30%. As the number of
samples was high, data were normalized using the quality control samples and support vector
regression (QC-SVRC) algorithm(44) in MATLAB. Finally, to ensure the quality of the obtained
data and detect sample patterns and possible outliers, we applied principal component analysis
(PCA) models using SIMCA 16 (Sartorius, Göttingen, Germany).

Lipid annotation

The workflow for lipid annotation consisted of three steps (**Figure S7B**): First, the raw LC-MS/MS data obtained were imported into the Lipid Annotator software (Agilent). The software created a fragmentation-based MS/MS library containing the *m/z* and matching retention time (RT) of all precursors recognized as lipids. The Lipid Annotator method was set as follows: ion species $[M+H]^+$, $[M+Na]^+$, and $[M+NH_4]^+$ for ESI(+); and $[M-H]^-$ and $[M+CH_3COO]^-$ for ESI(−). Then, for both ion modes, the Q-score was set at ≥ 50; all lipid classes were selected, mass deviation was established as ≤ 20 ppm, fragment score threshold was fixed as ≥ 30, and total score was set at ≥ 60.

Second, raw LC-MS/MS data were reimported into the open-source MS-DIAL software(45) in order to achieve an additional lipid identification with the aim of combining the information from both software to confirm lipid annotations and/or obtain new ones. Like Lipid Annotator, MS-DIAL can build a fragmentation-based MS/MS library comprising the *m/z* of all precursors identified as lipids by the software and the corresponding RT. The same ion species used with Lipid Annotator were used with MS-DIAL.

Finally, manual MS/MS spectral inspection was carried out using the Agilent MassHunter Qualitative software (version 10.0), comparing the fragmentation patterns of the displayed duplicates from both software. To ensure a correct annotation, the co-elution of possible adducts for the specific lipid subclass was examined(42). The final list of putatively annotated lipids was further verified and corrected with an in-house LC-MS database of human plasma lipids(46).

Statistical analysis

To examine the effects of HIIE, the signals of the putatively annotated lipids were compared between samples taken at the same time point in sessions A and B using the Wilcoxon test, performed in MATLAB. To control for type I error due to multiple comparisons, false discovery rate (FDR) was calculated by using the Benjamini-Hochberg correction. Statistical significance was set at 95% ($p < 0.05$) with a corrected p-value (pFDR) < 0.2 for paired analysis and exploratory studies(47,48).

The MetaboAnalyst online tool v6.0 (https://www.metaboanalyst.ca) was used to produce heat maps with hierarchical clustering(49). For lipid networks, the web-tool LINEX Lipid Network Explorer (version 1) was used (50). GraphPad Prism v9.5.1 (GraphPad Software, San Diego, CA, USA) was used to represent plots with stock bars and lipid trajectories.

Acknowledgments:

Authors would like to thank Santiago Angulo for his advice on statistics.

Author contributions: CRediT:

H.G. and **V.M.**: Conceptualization, Funding acquisition, Methodology, Supervision, and Writing
– original draft. **C.B.**, **C.G-R.**, and **G.T.** Funding acquisition, Methodology, Conceptualization,
and Supervision. **S.N.**, and **V. M.** Methodology, Investigation, and Writing – original draft. **T.M.**
and **C.G-R.** Formal analysis, Data curation, and Investigation. **R.N.**, **A.V.**, and **A.G.** Data
curation, Investigation, Methodology, and Writing – original draft. **A.V.**, and **A.G.** Supervision.
All authors: Writing – review and editing.

Competing Interests Statement: The authors declare no conflict of interest.

Sources of funding for the research reported in the article: This work was supported by
the HORIZON-WIDERA-2021 Project 101079370, BiACEM: BiOMIC_AUTh, Center of
Excellence in Metabolomics research.
This work was supported by the Ministry of Science and Innovation of Spain (MCIN)
MCIN/AEI/10.13039/501100011033, by ERDF A way of making Europe, grant number
PID2021-122490NB-I00. A.V. acknowledges financial support from "Ayudas Puente curso
24/25" from San Pablo CEU University and grant number PEJ-2023-AI/SAL-GL-27622 from the
Community of Madrid for hiring a research assistant.

Availability of data and materials: All data generated and analyzed during this study are
included in this article and its supplementary information files (Supplementary Tables). The raw
files of lipidomics generated in this study have been deposited in the Metabolomics Workbench
under Study ID ST003807 [http://dx.doi.org/10.21228/M8VR88].

REFERENCES

1. Szapary PO, Bloedon LT, Foster GD. Physical activity and its effects on lipids. Curr Cardiol Rep. noviembre de 2003;5(6):488-93.

2. Liu J xin, Zhu L, Li P jun, Li N, Xu Y bing. Effectiveness of high-intensity interval training on glycemic control and cardiorespiratory fitness in patients with type 2 diabetes: a systematic review and meta-analysis. Aging Clin Exp Res. mayo de 2019;31(5):575-93.

3. Bravo GM, Paramasivam P, Bellissimo GF, Jacquez Q, Zheng H, Amorim F, et al. High-Intensity Interval Training Decreases Circulating HMGB1 in Individuals with Insulin Resistance; Plasma Lipidomics Identifies Associated Cardiometabolic Benefits [Internet]. Molecular Biology; 2024 [citado 10 de marzo de 2025]. Disponible en: http://biorxiv.org/lookup/doi/10.1101/2024.08.21.608998

4. De Nardi AT, Tolves T, Lenzi TL, Signori LU, Silva AMVD. High-intensity interval training versus continuous training on physiological and metabolic variables in prediabetes and type 2 diabetes: A meta-analysis. Diabetes Research and Clinical Practice. marzo de 2018;137:149-59.

5. Belhaj MR, Lawler NG, Hoffman NJ. Metabolomics and Lipidomics: Expanding the Molecular Landscape of Exercise Biology. Metabolites. 7 de marzo de 2021;11(3):151.

6. Hoene M, Li J, Li Y, Runge H, Zhao X, Häring HU, et al. Muscle and liver-specific alterations in lipid and acylcarnitine metabolism after a single bout of exercise in mice. Sci Rep. 26 de febrero de 2016;6(1):22218.

7. Powell DJ, Turban S, Gray A, Hajduch E, Hundal HS. Intracellular ceramide synthesis and protein kinase Cζ activation play an essential role in palmitate-induced insulin resistance in rat L6 skeletal muscle cells. Biochemical Journal. 1 de septiembre de 2004;382(2):619-29.

8. Itani SI, Ruderman NB, Schmieder F, Boden G. Lipid-Induced Insulin Resistance in Human Muscle Is Associated With Changes in Diacylglycerol, Protein Kinase C, and IκB-α. Diabetes. 1 de julio de 2002;51(7):2005-11.

9. Kusminski CM, Shetty S, Orci L, Unger RH, Scherer PE. Diabetes and apoptosis: lipotoxity. Apoptosis. diciembre de 2009;14(12):1484-95.

10. Koves TR, Ussher JR, Noland RC, Slentz D, Mosedale M, Ilkayeva O, et al. Mitochondrial Overload and Incomplete Fatty Acid Oxidation Contribute to Skeletal Muscle Insulin Resistance. Cell Metabolism. enero de 2008;7(1):45-56.

11. Amati F, Dubé JJ, Alvarez-Carnero E, Edreira MM, Chomentowski P, Coen PM, et al. Skeletal Muscle Triglycerides, Diacylglycerols, and Ceramides in Insulin Resistance. Diabetes. 1 de octubre de 2011;60(10):2588-97.

12. Schenk S, Horowitz JF. Acute exercise increases triglyceride synthesis in skeletal muscle and prevents fatty acid–induced insulin resistance. J Clin Invest. 1 de junio de 2007;117(6):1690-8.

13. Hargreaves M, Spriet LL. Skeletal muscle energy metabolism during exercise. Nat Metab. 3 de agosto de 2020;2(9):817-28.

14. Lundsgaard AM, Fritzen AM, Kiens B. Molecular Regulation of Fatty Acid Oxidation in Skeletal Muscle during Aerobic Exercise. Trends in Endocrinology & Metabolism. enero de 2018;29(1):18-30.

15. Home PD. An overview of insulin therapy for the non-specialist. Diabetes Obes Metab. 4
de marzo de 2025;

16. Nikolaidis S, Virgiliou C, Vekiou M, Skari A, Kechagidou A, Gika H, et al. Effect of
exercise on key pharmacokinetic parameters related to metformin absorption in healthy
humans: A pilot study. Scandinavian Med Sci Sports. mayo de 2020;30(5):858-64.

17. The Effects of HIIT Versus MICT and Sedentary Controls on Blood Lipid Concentrations
in Nondiabetic Overweight and Obese Young Adults: A Meta-analysis. Int J Exerc Sci
[Internet]. 2023 [citado 10 de marzo de 2025];16(3). Disponible en:
https://digitalcommons.wku.edu/ijes/vol16/iss3/8/

18. O'Donovan G, Owen A, Bird SR, Kearney EM, Nevill AM, Jones DW, et al. Changes in
cardiorespiratory fitness and coronary heart disease risk factors following 24 wk of
moderate- or high-intensity exercise of equal energy cost. Journal of Applied Physiology.
mayo de 2005;98(5):1619-25.

19. Soria M, Ansón M, Lou-Bonafonte JM, Andrés-Otero MJ, Puente JJ, Escanero J. Fat
Oxidation Rate as a Function of Plasma Lipid and Hormone Response in Endurance
Athletes. Journal of Strength and Conditioning Research. enero de 2020;34(1):104-13.

20. Zhu JY, Guo L. Exercise-regulated lipolysis: Its role and mechanism in health and diseases.
Journal of Advanced Research. noviembre de 2024;S2090123224005502.

21. Zhu JY, Guo L. Exercise-regulated lipolysis: Its role and mechanism in health and diseases.
Journal of Advanced Research. noviembre de 2024;S2090123224005502.

22. Ferguson MA, Alderson NL, Trost SG, Essig DA, Burke JR, Durstine JL. Effects of four
different single exercise sessions on lipids, lipoproteins, and lipoprotein lipase. Journal of
Applied Physiology. 1 de septiembre de 1998;85(3):1169-74.

23. Atakan MM, Li Y, Koşar ŞN, Turnagöl HH, Yan X. Evidence-Based Effects of High-
Intensity Interval Training on Exercise Capacity and Health: A Review with Historical
Perspective. IJERPH. 5 de julio de 2021;18(13):7201.

24. Lee MC, Chung YC, Thenaka PC, Wang YW, Lin YL, Kan NW. Effects of different HIIT
protocols on exercise performance, metabolic adaptation, and fat loss in middle-aged and
older adults with overweight. Int J Med Sci. 2024;21(9):1689-700.

25. Bellou E, Magkos F, Kouka T, Bouchalaki E, Sklaveniti D, Maraki M, et al. Effect of high-
intensity interval exercise on basal triglyceride metabolism in non-obese men. Appl Physiol
Nutr Metab. agosto de 2013;38(8):823-9.

26. Warren A, Howden EJ, Williams AD, Fell JW, Johnson NA. Postexercise Fat Oxidation:
Effect of Exercise Duration, Intensity, and Modality. International Journal of Sport
Nutrition and Exercise Metabolism. diciembre de 2009;19(6):607-23.

27. Su W, Liu J, Wang A, Zhang H, Sun Y, Yan Z, et al. Distinct lipidomic profiles but similar
improvements in aerobic capacity following sprint interval training versus moderate-
intensity continuous training in male adolescents. Front Physiol. 30 de enero de
2025;16:1475391.

28. Cipryan L, Kosek V, García CJ, Dostal T, Bechynska K, Hajslova J, et al. A lipidomic and
metabolomic signature of a very low-carbohydrate high-fat diet and high-intensity interval
training: an additional analysis of a randomized controlled clinical trial. Metabolomics. 23
de diciembre de 2023;20(1):10.

29. Zaer Ghodsi N, University of Urmia, Urmia, Iran, Zolfaghari MR, University of Urmia,
Urmia, Iran, Fattah A, School of Medicine, Tabriz University of Medical Sciences, Tabriz,
Iran. The Impact of High Intensity Interval Training On Lipid Profile, Inflammatory
Markers and Anthropometric Parameters in Inactive Women. mljgoums. 1 de marzo de
2016;10(1):56-60.

30. Cavalli NP, De Mello MB, Righi NC, Schuch FB, Signori LU, Da Silva AMV. Effects of
high-intensity interval training and its different protocols on lipid profile and glycaemic
control in type 2 diabetes: A meta-analysis. Journal of Sports Sciences. 16 de febrero de
2024;42(4):333-49.

31. Su W, Liu J, Wang A, Zhang H, Sun Y, Yan Z, et al. Distinct lipidomic profiles but similar
improvements in aerobic capacity following sprint interval training versus moderate-
intensity continuous training in male adolescents. Front Physiol. 30 de enero de
2025;16:1475391.

32. Klepochová R, Niess F, Meyerspeer M, Slukova D, Just I, Trattnig S, et al. Correlation
between skeletal muscle acetylcarnitine and phosphocreatine metabolism during
submaximal exercise and recovery: interleaved 1H/31P MRS 7 T study. Sci Rep. 8 de
febrero de 2024;14(1):3254.

33. Lundsgaard AM, Fritzen AM, Kiens B. The Importance of Fatty Acids as Nutrients during
Post-Exercise Recovery. Nutrients. 21 de enero de 2020;12(2):280.

34. Tan-Chen S, Guitton J, Bourron O, Le Stunff H, Hajduch E. Sphingolipid Metabolism and
Signaling in Skeletal Muscle: From Physiology to Physiopathology. Front Endocrinol. 7 de
agosto de 2020;11:491.

35. Helge JW, Dobrzyn A, Saltin B, Gorski J. Exercise and training effects on ceramide
metabolism in human skeletal muscle. Experimental Physiology. enero de 2004;89(1):119-
27.

36. Widén EIM, Eriksson JG, Groop LC. Metformin Normalizes Nonoxidative Glucose
Metabolism in Insulin-Resistant Normoglycemic First-Degree Relatives of Patients With
NIDDM. Diabetes. 1 de marzo de 1992;41(3):354-8.

37. Dahabiyeh LA, Mujammami M, AlMalki RH, Arafat T, Benabdelkamel H, Alfadda AA,
et al. Lipids Alterations Associated with Metformin in Healthy Subjects: An Investigation
Using Mass Spectrometry Shotgun Approach. IJMS. 29 de septiembre de
2022;23(19):11478.

38. Dahabiyeh LA, Mujammami M, Arafat T, Benabdelkamel H, Alfadda AA, Abdel Rahman
AM. A Metabolic Pattern in Healthy Subjects Given a Single Dose of Metformin: A
Metabolomics Approach. Front Pharmacol. 15 de julio de 2021;12:705932.

39. Martin-Smith R, Cox A, Buchan DS, Baker JS, Grace F, Sculthorpe N. High Intensity
Interval Training (HIIT) Improves Cardiorespiratory Fitness (CRF) in Healthy, Overweight
and Obese Adolescents: A Systematic Review and Meta-Analysis of Controlled Studies.
IJERPH. 24 de abril de 2020;17(8):2955.

40. Wewege M, Van Den Berg R, Ward RE, Keech A. The effects of high-intensity interval
training vs. moderate-intensity continuous training on body composition in overweight and
obese adults: a systematic review and meta-analysis. Obesity Reviews. junio de
2017;18(6):635-46.

41. Ryan BJ, Schleh MW, Ahn C, Ludzki AC, Gillen JB, Varshney P, et al. Moderate-Intensity
Exercise and High-Intensity Interval Training Affect Insulin Sensitivity Similarly in Obese
Adults. The Journal of Clinical Endocrinology & Metabolism. 1 de agosto de
2020;105(8):e2941-59.

42. Martínez S, Fernández-García M, Londoño-Osorio S, Barbas C, Gradillas A. Highly
reliable LC-MS lipidomics database for efficient human plasma profiling based on NIST
SRM 1950. Journal of Lipid Research. noviembre de 2024;65(11):100671.

43. Armitage EG, Godzien J, Alonso-Herranz V, López-Gonzálvez Á, Barbas C. Missing value
imputation strategies for metabolomics data. Electrophoresis. diciembre de
2015;36(24):3050-60.

44. Kuligowski J, Sánchez-Illana Á, Sanjuán-Herráez D, Vento M, Quintás G. Intra-batch
effect correction in liquid chromatography-mass spectrometry using quality control samples
and support vector regression (QC-SVRC). Analyst. 2015;140(22):7810-7.

45. Tsugawa H, Cajka T, Kind T, Ma Y, Higgins B, Ikeda K, et al. MS-DIAL: data-
independent MS/MS deconvolution for comprehensive metabolome analysis. Nat Methods.
junio de 2015;12(6):523-6.

46. Martínez S, Fernández-García M, Londoño-Osorio S, Barbas C, Gradillas A. Highly
reliable LC-MS lipidomics database for efficient human plasma profiling based on NIST
SRM 1950. Journal of Lipid Research. noviembre de 2024;65(11):100671.

47. Chen M, Zhou K, Chen X, Qiao S, Hu Y, Xu B, et al. Metabolomic Analysis Reveals
Metabolic Changes Caused by Bisphenol A in Rats. Toxicological Sciences. 1 de abril de
2014;138(2):256-67.

48. Fiehn O, Garvey WT, Newman JW, Lok KH, Hoppel CL, Adams SH. Plasma Metabolomic
Profiles Reflective of Glucose Homeostasis in Non-Diabetic and Type 2 Diabetic Obese
African-American Women. Gimble JM, editor. PLoS ONE. 10 de diciembre de
2010;5(12):e15234.

49. Pang Z, Lu Y, Zhou G, Hui F, Xu L, Viau C, et al. MetaboAnalyst 6.0: towards a unified
platform for metabolomics data processing, analysis and interpretation. Nucleic Acids
Research. 5 de julio de 2024;52(W1):W398-406.

50. Köhler N, Rose TD, Falk L, Pauling JK. Investigating Global Lipidome Alterations with
the Lipid Network Explorer. Metabolites. 28 de julio de 2021;11(8):488.