Nutritional state and predator exposure jointly mediate stress resilience and behavioural flexibility

Stefan Fischer[1,2], Katharina Hirschenhauser[3], Barbara Taborsky[4], Leonida Fusani[1,2], Virginie Canoine[2] & Sabine Tebbich[2]

[1]Konrad Lorenz Institute of Ethology, Department of Interdisciplinary Life Sciences, University of Veterinary Medicine Vienna, Savoyenstrasse 1, 1160 Vienna, Austria

[2]Department of Behavioral and Cognitive Biology, University of Vienna, Djerassiplatz 1, 1030 Vienna, Austria

[3]University College for Education of Upper Austria (PH OÖ), Kaplanhofstraße 40, 4020 Linz, Austria

[4]Division of Behavioural Ecology, University of Bern, Wohlenstrasse 50a, CH-3032 Hinterkappelen, Switzerland

Correspondence and requests for materials should be addressed to S.F. (email:Stefan.fischer@vetmeduni.ac.at)

ORCIDs:

0000-0001-8811-7518 (Stefan Fischer)

0000-0001-8710-9022 (Katharina Hirschenhauser)

0000-0003-1690-8155 (Barbara Taborsky)

0000-0001-8900-796X (Leonida Fusani)

0000-0003-2272-0634 (Virginie Canoine)

0000-0003-3971-2139 (Sabine Tebbich)

Key words:

Neolamprologus pulcher; stress; HPI-axis; wear-and-tear; cortisol; allostatic load

Abstract

Organisms must adapt to changing environments, and behavioural flexibility plays an important role in adaptation within an individual's lifetime. Stress resilience influences flexibility, as short-term stress can enhance attention and memory, while prolonged stress may impair cognitive function. Nutritional state also plays a role, with overweight individuals often suffering greater consequences of chronic stress. The allostatic load model predicts that these individuals, under prolonged stress, remain in a chronically activated state, potentially reducing flexibility.

Using a social cichlid, we manipulated body condition and exposed individuals to occasional or frequent predator presentations. We then assessed stress responses and behavioural flexibility via a reversal learning test. Contrary to predictions, fish in a high nutritional state exposed to frequent predators recovered rapidly from stress and exhibited enhanced flexibility. Our results provide direct evidence that nutritional state interacts with behavioural flexibility, revealing the costs of coping with repeated stressors.

We highlight a link between brief stress responses, characterized by rapid cortisol recovery, and improved flexibility. This association only emerged in well-nourished individuals facing frequent stressors, suggesting energy reserves buffer the costs of repeated stress. Thus, resilience benefits behavioural flexibility, but it is also energetically costly, as a reduced diet hindered adaptation to environmental change.

Introduction

Animals have to constantly cope with and adapt to stressors in the environment such as the presence

of predators or changing environmental conditions [1]. Stress resilience, i.e. how rapid, efficient, and

complete the recovery of the stress response after a perturbation has occurred [sensu 2], is a major

component of behavioural and physiological adaptations for coping with extreme conditions. The

hypothalamic-pituitary-adrenal/interrenal (HPA/HPI) axis plays a central role in mediating

physiological stress resilience [3]. This hormonal cascade induces the release of glucocorticoids (GCs),

such as cortisol or corticosterone, which help mobilising energy to cope with the stressor [4].

Stressors disrupt the homeostasis of an organism, and the concept of allostasis refers to the process

by which the body reestablishes homeostasis through physiological or behavioural changes [5-7].

Allostasis is costly and organisms accumulate allostatic load when exposed to stressors [8, 9]. An

effective termination of the stress response is therefore essential to prevent the detrimental effects

of prolonged exposure to stress hormones such as GCs [5]. Thus, acute stress responses are

characterised by a short activation of the HPA/HPI axis with the result of increased GC secretion from

the adrenal, followed by a swift decrease. If the stressors occur repeatedly, but only occasionally,

there is sufficient time for the organism to quickly return to homeostasis, thus keeping the cost of

allostasis low. An acute stress response is beneficial, because it effectively activates physiological and

behavioural processes to cope with the stressor. Such acute stress responses are adaptive and

important for survival with positive effects on health and cognition [10-12].

In contrast, prolonged stress responses are characterised by a long activation of the HPA/HPI axis with

persistently elevated GC concentrations [13]. This persistent accumulation of allostatic load is

energetically costly and may lead to 'wear and tear' of the organism's body [8]. If stressors occur

frequently and the organisms cannot recover, these wear and tear costs accumulate and may

contribute to a range of health problems and impair cognition [14].

Behavioural flexibility is the capacity to adapt behaviour in response to changing environmental

conditions [15-17]. Arguably, cognition is among the most important determinants of behavioural

flexibility because learned behaviour can be reversed, allowing for significant adaptability [18]. While

exposure to acute stressors may enhance behavioural flexibility [19-21], prolonged or frequent

stressors impair behavioural flexibility, possibly as a result of persistently elevated stress responses

[22].

The type and the duration of the stressor as well as the energetic reserves of the organism determine

whether a stressor elicits an acute or prolonged stress response [5, 6, 23]. Previous studies proposed

that overnutrition results in the accumulation of excessive baseline energy reserves that fuel

prolonged allostasis; therefore the assumption was that obese organisms would be more susceptible

to the effects of prolonged stress responses than lean organisms [5, 6]. The consequences can be

severe. For instance, in humans chronic stress has been associated to a heightened risk for obesity,

which in turn results in negative health consequences such as depression and an increased risk of

inflammations [24-26].

A resilient organism is able to quickly overcome the allostatic load induced by a stressor and returns

to homeostasis. Resilience is affected by several factors including the energetic reserves and life-

history characteristics of the species [2]. In humans, resilient stress response patterns reveal

important health considerations, e. g. poor stress recovery correlates with poor cardiovascular status

[27] and obesity has been linked to lower stress responsiveness and impaired cognition [28, 29].

Characterising the specific factors that determine resilience in humans is a medically important

question. However, identifying these important and diverse factors is extremely difficult, particularly

in humans, due to the limits of experimental manipulations [30]. Thus, more research about the

interaction between nutrition, stress responses and cognition is needed, particularly in animal

models. So far, studies in laboratory rodents and birds showed rather mixed results [31-34] with

positive, negative, or no relationships between body condition and glucocorticoid responsiveness. We

are not aware of any previous studies that have experimentally investigated the influence of

nutritional state on the recovery pattern of the stress response and the resulting consequences for

behavioural flexibility

Fish and in particular cichlids are excellent models to further develop the research fields of cognition

and stress. The physiological mechanisms to cope with stressors are evolutionary conserved among

vertebrates [35] and cichlids display a broad repertoire of socio-cognitive abilities [36] including

behavioural flexibility [37, 38]. Standard procedures for experimental stress exposure are available

and individuals can be easily tagged to obtain individual characteristics (e.g. body condition) [39, 40].

Combined with detailed physiological measurements, this allows using both functional and

mechanistic approaches in experiments with cichlids. *Neolamprologus pulcher* is a highly social

cichlid, in which numerous previous studies have demonstrated the significance of the stress axis and

stress reactivity for its social behaviour and cognition [35, 38, 41-44]. Thus, these fish are a well

suited model to test how the frequency of stressors and nutritional status influence stress response

patterns and behavioural flexibility.

Here we performed an experiment using *N. pulcher* as study organisms to test the following

hypothesis (1) The frequency of stressors and the nutritional state changes individual cortisol

response patterns which ultimately influences behavioural flexibility. To test this prediction, we

manipulated the nutritional status of individuals while exposing them to different frequencies of

recurrent stressors in the form of occasional or frequent predator presentations. At the end of the

experiment, we assessed individual stress response patterns to a novel stressor and behavioural

flexibility using a reversal learning test paradigm [37]. Based on the allostatic load model we

formulated two specific predictions: (1) Frequent exposure to stressors and overnutrition separately

induce prolonged allostatic states including persistently prolonged high cortisol concentrations and

impaired behavioural flexibility, whereas an occasional exposure to stressors and a lower nutritional

status induce brief allostatic states. (2) When these two factors are combined, overweight individuals

are particularly susceptible to the negative effects of frequent exposure to stress, compared to lean

individuals.

Methods

<u>Subjects and housing</u>

The experiment was conducted at the Konrad Lorenz Institute of Ethology, University of Veterinary

Medicine, Vienna, Austria, under the licence (ETK-101/06/2019). All experimental tanks were

equipped with a 2 cm layer of sand, a biological filter, and a heater. The water temperature was kept

at 27 ± 1°C with a light-dark regime of 13:11 h and a dimmed-light phase of 10 min.

The study species was the cooperatively breeding *N. pulcher*, which is endemic to Lake Tanganyika. It

lives in stable social groups always including a dominant breeding pair and related or unrelated

helpers [45-48]. As an ecologically relevant stress stimulus, we used the piscivorous cichlid

Lepidiolamprologus elongatus, which coexists with *N. pulcher* [49]. *L. elongatus* preys on *N. pulcher* of

all sizes except for the eggs and is one of its most significant predators [50].

<u>Experimental design</u>

For this experiment we used 126 (63 male and 63 female) adult *N. pulcher* randomly selected from

the institute`s stock population. The experiment was run in three separate blocks with 42 (21

females) fish each that were housed in sex separated groups of 3-7 individuals (see below and

Supplementary Fig. S1). A block design with three blocks was chosen for logistical reasons and we

controlled for potential temporal effects by always including all six treatments in each experimental

block. To test the predictions outlined in the introduction, 30 groups of fish were created in total and

each group received one of the following six treatments, resulting in five groups and a total of 21 test

fish per treatment (Fig. 1 and Supplementary Fig. S2): (1) increased diet, no predators (IN); (2)

increased diet, occasional predators [IO]; (3) increased diet, frequent predators [IF]; (4) reduced diet,

no predators [RN]; (5) reduced diet, occasional predators [RO]; and (6) reduced diet, frequent

predators [RF]. At the end of the treatments, we randomly selected 96 test fish from each group and

assessed either their stress response patterns to a novel stressor (1 or 2 per group, Fig. S1) or their

behavioural flexibility (1, 2 or 4 per group, Fig. S1). From the remaining fish (N=30, 1 per group) we

collected the entire brains for a separate study that will be presented elsewhere. The experiment

consisted of four phases: (1) the feeding regime, (2) predator presentations, (3) reversal learning task,

and (4) hormone sampling. Each block of four phases lasted nine weeks (Fig. 1). Throughout the

experiment we constantly monitored for signs of aggression between group members, and in cases

where overt aggression was observed, we placed a transparent divider with holes in the tank for the

remainder of the experiment to physically separate fish from other group members without

interfering with visual or olfactory contact.

Experimental setup

Prior to the start of the experiment, we measured the standard length of the test fish (from the tip of

the snout to the end of the caudal peduncle [SL], to the nearest 0.1 cm) and body mass (BM, to the

nearest 0.1 g). To obtain individual growth rates (see Supplementary Figs. S6, S7) and to avoid testing

the same fish repeatedly, we used visible implant elastomer (VIE, Northwest Marine Technology, Inc.)

tags to individually mark the fish [40]. In block 1, all test fish were divided into six groups of seven,

separated by sex (three female and three male groups) and placed in 160L tanks. In blocks 2 and 3,

test fish showed more signs of aggression prior to the start of the experiment, and we decided, for

ethical reasons, to separate the fish into 12 groups of three (six male and six female groups) and 12

groups of four (six male and six female groups) again separated by sex (see Supplementary Fig. S1).

Feeding regime

Each group was assigned to one of two feeding regimes: (1) an increased (2) or a reduced diet. Fish

were fed with commercial JBL Stick M for cichlids® once a day six times per week. The increased and

reduced diet consisted of either 1.5 or 0.5 sticks per fish respectively (please see Supplementary

Methods S1 for results of a pilot experiment which confirms that the two diets induced fast and long-

lasting changes in body condition). We chose JBL Stick M for cichlids® as food because of two

reasons: (1) Individual sticks within a package are uniform and have nearly the same weight (0.045 ±

0.002g [mean ± SEM], N=10), which enabled us to create the two different food regimes by varying

the number of sticks provided to the groups. (2) The sticks float on the water surface for a prolonged

time and this assured that all group members received an equal amount of food.

In each block the experiment started with the assignment of each group to its respective feeding

regime and fish were fed the diet until the start of the hormone sampling phase (phase 4). During

feeding we always placed two transparent dividers inside the tanks to assure that all fish within the

same group received similar amounts of food. The dividers were arranged in a way to split up larger

groups of fish into individual fish or small groups of 2-3 individuals depending on the initial group

size. Particular attention was paid to avoid the repeated formation of small feeding groups that

consisted of the same individuals. Dividers were left in place until most of the food was consumed

and removed thereafter.

<u>Predator presentations</u>

After fish had received their respective dietary treatments for three weeks, each group was assigned

to one of three predator presentations. In blocks 2 and 3, two groups were assigned to the same

predator presentations resulting in the same number of fish per treatment overall. All predator

presentations consisted of a 5 min video playback of either (1) a recording of an empty tank, without

any predators visible, or (2) a recording of an attacking predator. To manipulate the frequency of

predator exposure, we presented the recording of the attacking predator either three times per week

(occasional exposure) or three times per day (frequent exposure).

We presented the video playbacks using a 25.7cm [10.1⬛⬛] display using a laptop (Lenovo Ideapad

D330 81MD-0006 Tablet Top). Videos were pre-recorded and validated to elicit similar responses as

live predators [39]. When presenting a video of an attacking predator, we randomly selected one of

the four available videos and randomly placed the screen either left or right next to the experimental

tank. To ensure that fish perceive the predators also using olfactory cues, we transferred 20ml of

predator water to create a more naturalistic experience of an approaching predator. Predator water

was always derived from one of two tanks (400L) containing five adult *L. elongatus* each. To avoid

that test fish habituated to the predator presentations we spread out the presentations semi-

randomly throughout the entire day and extended it with additional standardized stress

manipulations (see Supplementary methods S2). To control for potential disturbances caused by

adding predator water during the presentation of predator recordings, we also added 20ml of water

from tanks containing groups of *N. pulcher* not used in the experiment during the presentation of

control recordings.

<u>Reversal learning task</u>

We used a well-established reversal learning task to measure behavioural flexibility [37, 38]. For this,

fish first had to learn a visual discrimination task and subsequently perform the reverse of the

previously learned reward contingency. From each treatment group, we randomly selected 10 fish to

be tested individually (in total N=60 test fish [Block 1: N=22; Block 2,3: N=18]). The task was

separated in three phases: (1) pre-training and habituation phase, (2) training and colour

discrimination phase and (3) reversal learning phase. One fish died before the start of the training

phase reducing the overall sample size to N=59.

(1) Pre-training and habituation phase: During the last two weeks of the predator presentations, we

conducted the pre-training phase to habituate the fish to the test apparatus. The test apparatus

consisted of an experimental dark grey slate placed on the sandy bottom of the tank. The slate

contained 16 wells arranged in 4 columns, which could be covered with coloured discs of 15 mm

diameter. During the normal feeding schedules (as described above), when the groups were

separated into smaller feeding groups, we also placed an experimental slate in each separation. We

hid a small amount of food (crushed JBL Stick M for cichlids, according to their dietary treatments) in

some of the wells which were either uncovered or only partially covered with a coloured disc

(irrespective of the colour). During the two-week period we consecutively increased the coverage of

the wells by the coloured discs (25%, 50%, 75%, 100%) [see 51].

(2) Training phase: After the two-week pre-training and habituation phase we randomly selected fish

from each group for the reversal learning task. Each fish was weighed and measured as described

above and transported to an experimental training tank (30 x 30 x 50 cm). Each training tank was

equipped with a 2cm layer of sand, a biological filter, a heater, and a half clay flowerpot as shelter.

Each training tank was divided into a home compartment (2/3 of the tank) and an experimental

compartment (1/3 of the tank). The two compartments were separated by an opaque barrier

attached to strings on pulleys, allowing the observer to lift the barrier remotely while hiding behind a

curtain to minimize disturbance of the fish during the procedure. The training tanks were placed side

by side to allow the fish to interact socially, ensuring the welfare of the experimental fish and

encouraging their participation during the tests. During tests we placed opaque barriers between the

training tanks which avoided that fish in neighbouring tanks influenced or eavesdropped on the

behaviour of the focal fish. Each fish was allowed to acclimate overnight to the new environment

before the training phase started. Human observers were hidden behind a curtain and observations

were done blind to the experimental treatments. All observations were recorded using a web cam

placed on a tripod in front of the curtain which was connected via a USB cable to a laptop at the

other side of the curtain. This set-up allowed us to record and directly observe each test without the

need to lift the curtain and potentially disturb the fish.

As soon as the fish had habituated to the novel experimental tank, we started to train them to access

food from the feeding slate. We placed a feeding slate in the experimental compartment, removed

the barrier to the home compartment and allowed the fish to access the experimental compartment

and the food on the feeding slate. In the first training trial, a small green and a blue plastic disc were

placed again next to one of the holes containing a reward (a red mosquito larvae). On each successive

trial, the discs were placed so that they covered more and more of the holes, until the discs

completely covered the holes with the rewards, and the fish had to learn to move the disc by pushing

it away to reach the reward. We always performed six trials per day for a maximum of 4 days or until

fish successfully moved a disc to reach a reward that was entirely covered. Once they successfully

moved a disc (N=56) or repeatedly entered the experimental compartment to eat the reward (N=2),

fish proceeded to the acquisition phase on the next day. One fish did not proceed to the acquisition

phase because it never entered the experimental compartment to eat the reward from the slate

during the habituation phase.

(3) Colour discrimination in the acquisition phase: In this phase, fish (N=58) were given the choice

between two coloured discs (blue and green), one with an accessible reward and one with no reward

underneath. To prevent fish from uncovering the non-rewarded hole, we attached a pin below the

unrewarded disc, which made it hard for the test fish to push the disc away and uncover the well. We

placed a few drops of water which had previously contained mosquito larvae but not the actual

larvae, into the hole covered by the unrewarded disc. This ensured that fish used visual cues (i.e. the

colour of the discs) and not any other cues (i.e. the smell of the reward) to discriminate between the

rewarded and unrewarded discs. The rewarded colour was randomly assigned so that half of the fish

always found the reward underneath the blue disc, while the other half of the fish found the reward

underneath the green disc. We always placed one disc on the left and the other disc on the right side

of the slate on a randomly chosen well. On the next trial, the disc previously placed on the right side

was placed on the left side on a randomly selected well and vice versa, and we continued to switch

sides on each subsequent trial.

After the feeding slate was prepared, the observer moved behind the curtain and lifted the barrier

between the experimental and the home compartment using a string. The angle of the video camera

was positioned in a way that allowed us to test 3 fish at the same time. After the barrier was lifted,

each fish was allowed to explore the experimental compartment for a maximum of 3 minutes. In a

successful trial, the fish first uncovered the rewarded disc and accessed the reward. Immediately

afterwards the observer closed the barrier and the fish moved back into the home compartment

waiting for the next trial to start. In an unsuccessful trial, the fish either first touched the unrewarded

disc or did not touch any discs until the end of the trial. If the fish first touched the unrewarded disc

or did not touch any discs, the observer did not close the barrier allowing the fish more time to move

the rewarded disc and to uncover the reward. If the fish did not move the correct disc within an

additional 2-minutes, the observer closed the barrier and moved the fish back into the home

compartment. Afterwards, the observer moved the rewarded disc to reveal the reward underneath

and opened the barrier until the fish consumed the reward or for a maximum of 2 min. If the fish did

not consume the reward within these additional 2 minutes, the observer closed the barrier again and

the fish moved back into the home compartment. The uneaten reward was transferred from the

feeding slate into the home compartment and the fish could consume the reward while the

preparations for the next trial took place. This ensured that all fish consumed the same amount of

food during each testing day. This procedure was repeated for six consecutive trials per day until the

fish reached the learning criterion. We combined each set of 12 consecutive trials to one "block" of

trials. As we did 6 trials per day, a block lasted at least for two days. To reach the learning criterion,

the fish had to first choose the rewarded disc in 10 out of 12 trials (i.e., 80% of correct choices per

block) with a maximum of one unsuccessful choice per day. If fish did not reach the learning criterion

at the end of a block of 12 trials, the fish entered the next block of 12 trials. This procedure was

repeated until the fish either reached the learning criterion or had passed five blocks (60 trials)

without reaching the criterion. Once a fish reached the learning criterion in the acquisition phase, we

started the reversal phase on the next day. Two fish did not reach the learning criterion and did not

proceed to the reversal phase, reducing the sample size to N=56 for the next phase.

(3) Reversal learning phase. In this phase we reversed the rewarded and unrewarded colour discs, so

that the previously unrewarded disc became the rewarded one and vice versa. Otherwise, all

procedures and recordings were done as described for the acquisition phase. The reversal phase

ended when a fish had reached the learning criterion or after five blocks. All fish tested in the reversal

phase reached the learning criterion.

<u>Hormone sampling</u>

After the reversal learning task, we selected 1-2 fish from each group (N=36 fish, Block 1: N=11; Block

2: N=13: Block 3: N=12, Supplementary Fig. S1) that had received the food and predator treatments

but not the reversal learning task. At the same time all predator presentations stopped, and all

groups were transferred to an intermediate food diet (1 stick of JBL Stick M for cichlids® per fish.) The

apparatus (see below) enabled the collection of water samples from two fish at the same time. We

extended the static holding water method [52] with a continuous water flow that allows the repeated

water sampling without further disturbing the fish [41]. The collections for each pair of fish took

place over four consecutive days which allowed us to quantify cortisol concentrations (as ng

cortisol/L) immediately after a new stressor (i.e. catching and handling of the fish on day 1) as well as

cortisol concentrations after fish were habituated to this stressor, i.e. the same catching and handling

procedures on day 4 (see below). The involved procedures, including catching, measuring, weighing,

and placing the fish in a novel container are perceived as stressful by the experimental animals [41,

53]. However, another study showed that *N. pulcher* habituate to this procedure after three days,

which allows the collection of cortisol concentrations of fish before and after habituation [41] and

thus serves, as a reliable baseline measure.

On the first hormone sampling day we caught two fish and measured their body sizes and body

masses. We then placed each of the two fish individually in a sampling container filled with 500ml of

water and collected four 500ml water samples over 2h. All water samples were collected at the same

time in the afternoon (after 14:00h) to control for diurnal variation of cortisol [41]. On two

consecutive days, we caught the same two fish and transferred them to the sampling containers filled

with 500ml of water for 2h, but did not collect any water samples. This ensured that fish habituated

to the catching and handling procedures of the hormone sampling. On the fourth day, we repeated

the same procedures, including measuring and weighing the fish and collected another set of four

water samples for 2h (see Supplementary Fig. S3). In block 1, due to logistical constraints, we only

collected 2 water samples per day, instead of 8, and we decided to always collect the 3rd and 4th water

samples on the 1st and 4th day respectively. All other procedures were the same for all three blocks.

This resulted in a total of 240 water samples.

<u>Hormone measurement</u>

Apparatus. To repeatedly and minimal invasively sample individual cortisol levels, we used a

modification of the holding-water method previously used and validated for *N. pulcher* [41]. Exact

details of the apparatus can be found in the Supplementary Methods S3. In short, we used a water

flow-through system to collect four water samples per fish throughout a 2h sampling period.

Cortisol measures. Prior analysing cortisol levels, an extraction was performed as following: Water

samples were defrosted overnight and filtered in the morning of the next day (paper filter 1/2595

grade, diameter 320 mm, Whatman, Sigma-Aldrich, Switzerland) to remove solid particles, and

lipophilic compounds were extracted using solid phase extraction (SPE). After the hormone samples

were loaded onto a cartridge (SPE column Isolute C18 (EC), 500 mg/6mL, Biotage, Sweden), the

cartridges were stored at - 20 ⬚C until further processing at the University of Vienna.

Conditioned cartridges were defrosted, then placed on a vacuum manifold (Biotage VacMaster 20)

and eluted with 5ml ethyl acetate (HPLC -grade 99.9%). Ethyl acetate proved to be a very satisfactory

solvent for the elution of cortisol (mainly free CORT) from solid phase extraction cartridges [54].

Collected cortisol samples were dried down with N_2 at 45 °C, resuspend with 750ul assay buffer,

vortexed and placed overnight in a fridge. Free cortisol concentrations were quantified using a

commercial enzymatic immunoassay for cortisol (ENZO Life Sciences Cat # ADI 900-071).

Reconstituted samples were brought to RT on a shaker for 30 min and then further processed

following the manufacture's instruction. All samples were analysed in duplicates. The sensitivity of

the assay was 24.7 pg/ml, intra-assay CV% was ≤ 5% and inter assay CV% was below 13% using a

control sample. Sample concentrations were corrected for dilution. Of the 36 sampled test fish, we

were not able to reliably measure cortisol from 2 fish in any of the collected water samples, reducing

the overall sample size to 34 fish.

<u>Statistical analysis</u>

For statistical analyses we used R 4.3.3 [55] and the packages lme4 [56], and afex [57]. We used linear

mixed effects models (LMM) and generalized linear mixed effects models (GLMM). To obtain p-values,

we used either the mixed() function in the package 'afex' with a Satterthwaite's approximation for

degrees of freedom, or the drop1() function to do a likelihood ratio test using an F statistics for LMM

or an ⬚2 statistics for GLMM. Interactions and covariates were stepwise removed if non-significant

[58] and reported p-values refer to the final model without non-significant interactions. An overview

of all full models is provided in Table S3. To account for familiarity of fish kept in the same group tanks

and for potential variations between blocks we always included group tank ID and block ID as random

factors.

Based on our predictions and the rationale outlined in the introduction, we aimed to investigate the

influence of nutritional state and not feeding regime *per se* on stress reactivity and behavioural

flexibility. To compare fish with high and low body condition, we classified them as 'high nutritional

state' or 'low nutritional state' depending on whether their body condition was higher or lower than

the mean body condition of all fish tested in this study (3.02 ± 0.03 [mean ± SE]). We verified

nevertheless that the feeding regimes had the desired effect, and found that fish on the increased

diet gained more weight and grew more throughout the experiment than fish on the reduced diet

(see Supplementary Table S4, Figs. S6, S7). Consequently, the majority (71%) of fish on the increased

diet had above-average body condition and were classified as 'high nutritional state' whereas the

majority (74%) of fish on the reduced diet had below average body condition and were classified as

'low nutritional state'.

To analyse stress response patterns, we first classified each individual as a brief or prolonged

responder based on their cortisol pattern in response to the hormone sampling on the first day. Brief

responders had shortly elevated cortisol peaks whereas prolonged responders had persistently

elevated cortisol concentrations within the 2h sampling period. We used a GLMM with a binomial

error distribution to analyse whether the probability to display a brief stress response is influenced by

body condition or predator exposures.

To estimate the duration of individual stress response patterns, we calculated an index of stress

recovery on the first day of sampling. This index was calculated as the difference between the

individual peak cortisol concentration and the final value after 2h (at sampling point 4). Thus, a higher

recovery value indicates a stress response pattern characterized by a short cortisol peak with a rapid

decrease of cortisol within 2h. A lower recovery value indicates a stress response pattern

characterised by prolonged periods of high cortisol levels with a slow decrease within 2h. A recovery

value of zero indicates a stress response pattern characterised by prolonged periods of high cortisol

levels with no decrease within 2h. To assess whether the duration of the stress response is

determined by the body condition of the fish, we used a LMM with stress recovery as the dependent

variable and the actual body condition as a covariate. We only conducted this analysis after

confirming that fish exposed to frequent predators were more likely to show a brief stress response

pattern. This result, although not significant, was clearly driven by fish with high body condition (see

results). Therefore, to increase the resolution of the post-hoc analyses, we used the stress recovery

indices and individual body condition, and focused on the experimental group where the majority of

fish showed a brief stress response (i.e. the fish exposed to frequent predators).

To analyse behavioural flexibility, we used the number of trials until fish reached the learning

criterion in the reversal learning phase as the dependent variable and included body condition ('high',

'low'), predator exposure ('no predators', frequent predators', 'occasional predators') as fixed factors

in an LMM. To assess whether the treatments influenced the accuracy of fish during the reversal

phase we used the proportion of wrong choices over all trials in this phase ('error rate') [59] as the

depend variable and included body condition and predator exposure as fixed effects in a GLMM with

a binomial error distribution. To assess whether the treatments influenced the fish's performance

during the acquisition phase we used the number of trials and the error rate in a LMM and GLMM

respectively with body condition and predator exposure as fixed effects.

Results

<u>(1) Do nutritional state and the frequency of stressors determine stress recovery patterns?</u>

Contrary to the predictions, the majority of fish exposed to frequent predators had a brief stress response pattern when exposed to a novel stressor, which was not the case for fish exposed to occasional or no predators (Table 1 factor: 'Predator exposure [χ2=8.2, p=0.02], for orthogonal contrasts see Table S5, Fig. 2). In addition, we found that all fish in a high nutritional state that experienced frequent predators showed a brief stress response pattern (see Fig. 2, N=4 out of 4). To test whether body condition determines stress recovery patterns we performed a post-hoc analysis that used the stress recovery index of fish frequently exposed to predators. Although nutritional status did not predict whether fish were classified as brief or prolonged responders (Table 1 factor: 'Body condition' [χ2=1.93, p=0.17]), the post-hoc analysis revealed that the magnitude of the stress response, in terms of the individual stress recovery indices, was clearly influenced by body condition (Table 2 factor: 'Body condition' [$F_{1,9}$=8.78, p=0.02]). Fish with a higher body condition had a more rapid recovery pattern than fish with a lower body condition (see Fig. S8). The frequency of predator exposures explained stress response patterns when fish were exposed to a new stressor on day 1 (see Table 1), but not when fish were potentially habituated to the hormone sampling procedures on day 4 (Table S6 factor 'Predator exposure' [χ2=1.48, p=0.48]).

<u>(2) Do nutritional status and the frequency of stressors determine behavioural flexibility?</u>

In the reversal learning test, fish took on average 27.2 ± 1.5 and 30.6 ± 1.1 (mean ± SEM) trials to reach the learning criterion in the acquisition and reversal phase, respectively. Two fish did not reach the learning criterion in the acquisition phase and were excluded from the analysis. Behavioural flexibility, in terms of the number of trials until fish reached the learning criterion in the reversal phase, varied depending on the frequency of predator exposures and the nutritional status of the fish (Table 3a factor 'Body condition x Predator exposure' [χ2=5.51, p=0.06]). Contrary to the prediction that frequent stressors and overnutrition would impair behavioural flexibility, orthogonal post-hoc comparisons revealed that fish with high body condition reached the learning criterion faster when exposed to frequent predators compared to fish exposed to occasional predators (Table S7c factor 'Occasional predators vs. frequent predators (high body condition)' [estimate: -0.16, z=-3.4, p<0.01, Fig. 3). Predator exposure did not influence behavioural flexibility of fish with a low body condition (see Table S7d,e). Neither predator exposure nor nutritional status influenced the number of trials required to reach the learning criterion in the acquisition phase (Table 3b factors 'Body condition' [x^2=0.05, p=0.83], 'Predator exposure' [x^2=0.14, p=0.93]). Similarly, the experimental manipulations

did not affect the error rates of the fish during the acquisition phase (Table S8a) and the reversal

learning phase (Table S8b).

Discussion

Based on the allostatic load model we predicted that overnutrition and frequent exposure to

stressors induce prolonged allostatic states including persistently prolonged stress response patterns

and impaired behavioural flexibility. Contrary to this prediction, high body condition fish exposed to

frequent predators showed brief stress response patterns when exposed to a new stressor. In

addition, high body condition fish exposed to frequent predators also showed greater behavioural

flexibility. Therefore, the results do not support our initial hypothesis. An alternative interpretation is

that our treatments were not sufficient to induce the conditions on which the hypotheses were

based: (1) Although fish on the increased diet gained more weight and had more energy resources

than fish on the reduced diet, they were not obese, with body masses within the adult range [60],

and therefore their condition did not represent overnutrition [6]; (2) The predator presentations

clearly induced stress responses in the test fish, however, the frequent predator encounters may not

have been enough to induce severe chronic stress. In any case, our results highlight a potential

association between brief stress response patterns, characterised by a quick termination of cortisol

concentrations, and improved behavioural flexibility measured as reversal learning speed. This

association only became apparent when individuals with high body condition, i. e. with more energy

reserves available, were exposed to frequent stressors.

Glucocorticoid receptors (GR) which are expressed in the brain play an important role in terminating

the stress response [61, 62]. A previous study that pharmacologically blocked GRs in *N. pulcher*

showed that GR blockade reduced behavioural flexibility in a detour task, highlighting a potential link

between stress resilience and the ability to cope with changing conditions. Although cortisol was not

measured directly, GR-blocked fish also showed more fear-related behaviours, a possible indication

that blocking GRs may have led to an accumulation of cortisol due to the reduced ability to terminate

the stress response [63]. Based on the results of the current study, we can conclude that while

resilience, defined as an acute stress response, is strongly associated with improved behavioural

flexibility, it also comes with associated costs that well-fed individuals are better able to overcome.

A prolonged phase of allostasis is the result of a chronic exposure to stressors with high energetic

costs and consequences for health and fitness [8], however, the determination of chronic stress load

particularly under natural conditions, depends on species-specific factors [64]. We did not find

support for the prediction that frequent predator exposures and overnutrition will lead to prolonged

allostasis. This prediction was derived from the classic version of the allostatic load model which

introduced the concept of allostasis as maintaining homeostastis through change [6]. An extension of

the classic version, the reactive scope model, predicts that animals with more energy resources will

be better able to cope with repeatedly occurring stressors [8]. This might explain the result that fish

with higher body condition had shorter stress response patterns and an improved behavioural

flexibility when exposed to frequent predators. Possibly, the higher energetic resources enabled them

to express an adequate response to a new stressor, i.e. a brief stress response pattern.

Interestingly, fish that had both a high body condition and were exposed frequently to predators also

showed more behavioural flexibility. This suggests that the ability to exhibit a stress response pattern

with a brief cortisol peak followed by a timely recovery, is associated with higher behavioural

flexibility. Stress influences cognition in a complex way that is dependent on the type and duration of

the stressor, the individual physiological response type, and the cognitive task being studied [10, 65].

Generally, short term stressors are considered to have positive effects on cognitive abilities required

to overcome the stressor, whereas long-term or chronic stress is detrimental [65, 66]. Our results

provide a valuable contribution to the fields of stress and cognition by highlighting the importance of

the nutritional status in determining stress response and behavioural flexibility, i.e. the potential

benefits of brief stress response patterns.

There are two mutually exclusive explanations as to why fish adapted to the increased frequency of

the stressor with beneficial effects on physiology and flexibility. First, fish exposed to frequent

predator presentations might have been habituated to the re-occurring stressor leading to shorter

stress responses when exposed to a new stressor and with beneficial effects for behavioural

flexibility. We took great care to avoid habituation of fish to the predator presentations by varying the

timing of the exposures and adding additional stressors such as chasing with a net or creating a

moving shadow. Yet, the fish potentially may have developed habituation simply because they were

exposed more frequently to the predator presentations. Second, the brief stress responses and

improved behavioural flexibility could have been an adaptive response to cope with the frequent

predator presentations, which would support the idea of preparative effects of glucocorticoid actions

[4, 67, 68]. The 'preparative hypothesis' suggests that prior glucocorticoid actions modulate the

response to a subsequent stressor in a mediating or suppressive way [4]. Although not yet tested

experimentally, this hypothesis has been used to explain seasonal GC variation. According to the

preparative hypothesis both, baseline and stress induced GC levels are higher at challenging times of

the year [68, 69]. Yet, stressful periods do not necessarily occur at fixed periods of the year and

perceived stress may depend on the social status or demands of the individual [70, 71]. An interesting

extension of the preparative hypothesis suggests that GC actions i) prepare to better cope with a

subsequent stressor and ii) increase the threshold when certain stimuli are perceived as a stressor

[72]. The observed interaction between body condition and adaptive stress response patterns when

exposed to predators is in line with this extension of the hypothesis because the available energy

stores affect the ability of organisms to cope with stressors [73].

In conclusion, our results show a potential relationship between stress resilience and behavioural

flexibility. Groups of fish that terminated stress responses sooner showed improved behavioural

flexibility. This relationship was only seen in groups of fish exposed to frequent predator exposures,

which simulated a higher frequency of stressful events. Experiencing more frequent stressful events

thus might have prepared fish to better cope with subsequent stressors, particularly in those with

greater energy reserves. These results are also consistent with the concept of 'wear and tear', which

introduces the idea that frequent exposure to stressors is costly [8] and animals in a better

nutritional condition should be better able to cope with repeated stressors due to their higher

energy reserves compared to animals in a poor nutritional status. Our results provide direct evidence

for the costs of coping with repeated stressors as the manipulation of the nutritional condition

interacted with behavioural flexibility. This work is advancing our understanding of the way how

animals respond to environmental change.

Acknowledgments

We would like to thank Claudia Martina and Noémie Liesch for their help during the learning task,

Arne Jungwirth for his help with the predator presentations, Martina Krakhofer for help with animal

husbandry and Roland Sasse for logistical support. We are particularly grateful to Stefan Graf, whose

invaluable support and dedication at all stages of the project was crucial to its success. This study was

funded by the Vienna Science and Technology Fund (CS18-042). The text of this manuscript has been

improved with the assistance of AI tools, such as ChatGPT and DeepL, to enhance clarity and

language quality.

Literature

[1] MacLeod, K.J., English, S., Ruuskanen, S.K. & Taborsky, B. 2023 Stress in the social context: a
behavioural and eco-evolutionary perspective. *J. Exp. Biol.* **226**. (doi: 10.1242/jeb.245829).

[2] Wingfield, J.C. 2013 The comparative biology of environmental stress: behavioural endocrinology
and variation in ability to cope with novel, changing environments. *Anim. Behav.* **85**, 1127-1133. (doi:
10.1016/j.anbehav.2013.02.018).

[3] Creel, S., Dantzer, B., Goymann, W. & Rubenstein, D.R. 2013 The ecology of stress: effects of the
social environment. *Funct. Ecol.* **27**, 66-80. (doi: 10.1111/j.1365-2435.2012.02029.x).

[4] Sapolsky, R.M., Romero, L.M. & Munck, A.U. 2000 How do glucocorticoids influence stress
responses? Integrating permissive, suppressive, stimulatory, and preparative actions. *Endocr. Rev.* **21**,
55-89. (doi: 10.1210/er.21.1.55).

[5] McEwen, B.S. 2005 Stressed or stressed out: what is the difference? *J. Psychiatr. Neurosci.* **30**, 315-
318.

[6] McEwen, B.S. & Wingfield, J.C. 2003 The concept of allostasis in biology and biomedicine. *Horm.*
Behav. **43**, 2-15. (doi: 10.1016/s0018-506x(02)00024-7).

[7] McEwen, B.S. & Wingfield, J.C. 2010 What is in a name? Integrating homeostasis, allostasis and
stress. *Horm Behav* **57**, 105-111. (doi: 10.1016/j.yhbeh.2009.09.011).

[8] Romero, L.M., Dickens, M.J. & Cyr, N.E. 2009 The reactive scope model - a new model integrating
homeostasis, allostasis, and stress. *Horm. Behav.* **55**, 375-389. (doi: 10.1016/j.yhbeh.2008.12.009).

[9] Wright, J., Buch, K., Beattie, U.K., Gormally, B.M.G., Romero, L.M. & Fefferman, N. 2023 A
mathematical representation of the reactive scope model. *Journal of Mathematical Biology* **87**, 51.
(doi: 10.1007/s00285-023-01983-9).

[10] Sandi, C. & Pinelo-Nava, M.T. 2007 Stress and memory: behavioral effects and neurobiological
mechanisms. *Neural Plasticity* **2007**, 2007: 78970. (doi: 10.1155/2007/78970).

[11] Schwabe, L., Joels, M., Roozendaal, B., Wolf, O.T. & Oitzl, M.S. 2012 Stress effects on memory: an
update and integration. *Neurosci. Biobehav. Rev.* **36**, 1740-1749. (doi:
10.1016/j.neubiorev.2011.07.002).

[12] de Kloet, E.R., Joels, M. & Holsboer, F. 2005 Stress and the brain: from adaptation to disease.
Nat. Rev. Neurosci. **6**, 463-475. (doi: 10.1038/nrn1683).

[13] Dickens, M.J. & Romero, L.M. 2013 A consensus endocrine profile for chronically stressed wild
animals does not exist. *Gen Comp Endocrinol* **191**, 177-189. (doi: 10.1016/j.ygcen.2013.06.014).

[14] D'Amico, D., Amestoy, M.E. & Fiocco, A.J. 2020 The association between allostatic load and
cognitive function: a systematic and meta-analytic review. *Psychoneuroendocrino.* **121**, 104849. (doi:
10.1016/j.psyneuen.2020.104849).

[15] Diamond, A. 2013 Executive functions. In *Annual Review of Psychology, Vol 64* (ed. S.T. Fiske), pp.
135-168. Palo Alto, Annual Reviews.

[16] Sol, D., Timmermans, S. & Lefebvre, L. 2002 Behavioural flexibility and invasion success in birds.
Anim. Behav. **63**, 495-502. (doi: 10.1006/anbe.2001.1953).

[17] Burkart, J.M., Schubiger, M.N. & van Schaik, C.P. 2017 The evolution of general intelligence.
Behav. Brain Sci. **40**, e195. (doi: 10.1017/s0140525x16000959).

[18] Tebbich, S., Sterelny, K. & Teschke, I. 2010 The tale of the finch: adaptive radiation and
behavioural flexibility. *Philos. Trans. R. Soc. B-Biol. Sci.* **365**, 1099-1109. (doi:
doi.org/10.1098/rstb.2009.0291).

[19] Bryce, C.A. & Howland, J.G. 2015 Stress facilitates late reversal learning using a touchscreen-
based visual discrimination procedure in male Long Evans rats. *Behav. Brain Res.* **278**, 21-28. (doi:
10.1016/j.bbr.2014.09.027).

[20] Graybeal, C., Feyder, M., Schulman, E., Saksida, L.M., Bussey, T.J., Brigman, J.L. & Holmes, A. 2011
Paradoxical reversal learning enhancement by stress or prefrontal cortical damage: rescue with BDNF.
Nat. Neurosci. **14**, 1507-1509. (doi: 10.1038/nn.2954).

[21] Thai, C.A., Zhang, Y. & Howland, J.G. 2013 Effects of acute restraint stress on set-shifting and
reversal learning in male rats. *Cogn. Affect. Behav. Neurosci.* **13**, 164-173. (doi: 10.3758/s13415-012-
0124-8).

[22] George, S.A., Rodriguez-Santiago, M., Riley, J., Abelson, J.L., Floresco, S.B. & Liberzon, I. 2015
Alterations in cognitive flexibility in a rat model of post-traumatic stress disorder. *Behav. Brain Res.*
286, 256-264. (doi: 10.1016/j.bbr.2015.02.051).

[23] Canoine, V., Hayden, T.J., Kevin, R. & Goymann, W. 2002 The stress response of european
stonechats depends on the type of stressor. *Behaviour* **139**, 1303-1311. (doi:
10.1163/156853902321104172).

[24] Benson, S., Arck, P.C., Tan, S., Mann, K., Hahn, S., Janssen, O.E., Schedlowski, M. & Elsenbruch, S.
2009 Effects of obesity on neuroendocrine, cardiovascular, and immune cell responses to acute
psychosocial stress in premenopausal women. *Psychoneuroendocrino.* **34**, 181-189. (doi:
10.1016/j.psyneuen.2008.08.019).

[25] Petry, N.M., Barry, D., Pietrzak, R.H. & Wagner, J.A. 2008 Overweight and obesity are associated
with psychiatric disorders: results from the National Epidemiologic Survey on Alcohol and Related
Conditions. *Psychosom Med* **70**, 288-297. (doi: 10.1097/PSY.0b013e3181651651).

[26] Kühnel, A., Hagenberg, J., Knauer-Arloth, J., Ködel, M., Czisch, M., Sämann, P.G., Binder, E.B. &
Kroemer, N.B. 2023 Stress-induced brain responses are associated with BMI in women. *Commun Biol*
6, 1031. (doi: 10.1038/s42003-023-05396-8).

[27] Chida, Y. & Steptoe, A. 2010 Greater cardiovascular responses to laboratory mental stress are
associated with poor subsequent cardiovascular risk status: a meta-analysis of prospective evidence.
Hypertension **55**, 1026-1032. (doi: 10.1161/hypertensionaha.109.146621).

[28] Mujica-Parodi, L.R., Renelique, R. & Taylor, M.K. 2009 Higher body fat percentage is associated
with increased cortisol reactivity and impaired cognitive resilience in response to acute emotional
stress. *Int J Obes (Lond)* **33**, 157-165. (doi: 10.1038/ijo.2008.218).

[29] Jones, A., McMillan, M.R., Jones, R.W., Kowalik, G.T., Steeden, J.A., Deanfield, J.E., Pruessner, J.C.,
Taylor, A.M. & Muthurangu, V. 2012 Adiposity is associated with blunted cardiovascular,
neuroendocrine and cognitive responses to acute mental stress. *Plos ONE* **7**, e39143. (doi:
10.1371/journal.pone.0039143).

[30] Chbeir, S. & Carrión, V. 2023 Resilience by design: How nature, nurture, environment, and
microbiome mitigate stress and allostatic load. *World J Psychiatry* **13**, 144-159. (doi:
10.5498/wjp.v13.i5.144).

[31] Kitaysky, A.S., Piatt, J.F., Wingfield, J.C. & Romano, M. 1999 The adrenocortical stress-response of
Black-legged Kittiwake chicks in relation to dietary restrictions. *Journal of Comparative Physiology B*
169, 303-310. (doi: 10.1007/s003600050225).

[32] Spencer, S.J. & Tilbrook, A. 2009 Neonatal overfeeding alters adult anxiety and stress
responsiveness. *Psychoneuroendocrino.* **34**, 1133-1143. (doi: 10.1016/j.psyneuen.2009.02.013).

[33] Lynn, S.E., Hunt, K.E. & Wingfield, J.C. 2003 Ecological factors affecting the adrenocortical
response to stress in chestnut-collared and McCown's longspurs (*Calcarius ornatus, Calcarius*
mccownii). *Physiol Biochem Zool* **76**, 566-576. (doi: 10.1086/375435).

[34] Schultner, J., Kitaysky, A.S., Welcker, J. & Hatch, S. 2013 Fat or lean: adjustment of endogenous
energy stores to predictable and unpredictable changes in allostatic load. *Funct. Ecol.* **27**, 45-55. (doi:
10.1111/j.1365-2435.2012.02058.x).

[35] Nyman, C., Fischer, S., Aubin-Horth, N. & Taborsky, B. 2018 Evolutionary conserved neural
signature of early life stress affects animal social competence. *Proc. R. Soc. B-Biol. Sci.* **285**, 20172344.
(doi: 10.1098/rspb.2017.2344).

[36] Jordan, A., Taborsky, B. & Taborsky, M. 2021 Cichlids as a Model System for Studying Social
Behaviour and Evolution. In *The Behavior, Ecology and Evolution of Cichlid Fishes* (eds. M.E. Abate &
D.L.G. Noakes), pp. 587-635. Dordrecht, Springer Netherlands.

[37] Fischer, S., Balshine, S., Hadolt, M.C. & Schaedelin, F.C. 2021 Siblings matter: Family
heterogeneity improves associative learning later in life. *Ethology* **127**, 897-907. (doi:
10.1111/eth.13196).

[38] Reyes-Contreras, M. & Taborsky, B. 2022 Stress axis programming generates long-term effects on
cognitive abilities in a cooperative breeder. *Proc. R. Soc. B-Biol. Sci.* **289**, 20220117. (doi:
10.1098/rspb.2022.0117).

[39] Watve, M., Prati, S. & Taborsky, B. 2019 Simulating more realistic predation threat using attack
playbacks. *PeerJ* **7**, 16. (doi: 10.7717/peerj.8149).

[40] Jungwirth, A., Balzarini, V., Zottl, M., Salzmann, A., Taborsky, M. & Frommen, J.G. 2019 Long-term
individual marking of small freshwater fish: the utility of Visual Implant Elastomer tags. *Behav. Ecol.*
Sociobiol. **73**. (doi: 10.1007/s00265-019-2659-y).

[41] Antunes, D.F., Reyes-Contreras, M., Glauser, G. & Taborsky, B. 2021 Early social experience has
life-long effects on baseline but not stress-induced cortisol levels in a cooperatively breeding fish.
Horm. Behav. **128**, 104910. (doi: 10.1016/j.yhbeh.2020.104910).

[42] Nyman, C., Fischer, S., Aubin-Horth, N. & Taborsky, B. 2017 Effect of the early social environment
on behavioural and genomic responses to a social challenge in a cooperatively breeding vertebrate.
Mol. Ecol. **26**, 3186-3203. (doi: 10.1111/mec.14113).

[43] Reyes-Contreras, M., Glauser, G., Rennison, D.J. & Taborsky, B. 2019 Early-life manipulation of
cortisol and its receptor alters stress axis programming and social competence. *Philos. Trans. R. Soc.*
B-Biol. Sci. **374**, 20180119. (doi: 10.1098/rstb.2018.0119).

[44] Taborsky, B., Tschirren, L., Meunier, C. & Aubin-Horth, N. 2013 Stable reprogramming of brain
transcription profiles by the early social environment in a cooperatively breeding fish. *Proceedings of*
the Royal Society of London Series B-Biological Sciences **280**, 20122605. (doi:
10.1098/rspb.2012.2605).

[45] Balshine, S., Leach, B., Neat, F., Reid, H., Taborsky, M. & Werner, N. 2001 Correlates of group size
in a cooperatively breeding cichlid fish (*Neolamprologus pulcher*). *Behav. Ecol. Sociobiol.* **50**, 134-140.
(doi: 10.1007/s002650100343).

[46] Heg, D., Brouwer, L., Bachar, Z. & Taborsky, M. 2005 Large group size yields group stability in the
cooperatively breeding cichlid *Neolamprologus pulcher*. *Behaviour* **142**, 1615-1641. (doi:
10.1163/156853905774831891).

[47] Taborsky, M. 1984 Broodcare helpers in the cichlid fish *Lamprologus brichardi*: their costs and
benefits. *Anim. Behav.* **32**, 1236-1252. (doi: 10.1016/s0003-3472(84)80241-9).

[48] Taborsky, M. 1985 Breeder-helper conflict in a cichlid fish with broodcare helpers: an
experimental analysis. *Behaviour* **95**, 45-75. (doi: 10.1163/156853985x00046).

[49] Ochi, H. & Yanagisawa, Y. 1998 Commensalism between cichlid fishes through differential
tolerance of guarding parents toward intruders. *J. Fish Biol.* **52**, 985-996. (doi: 10.1111/j.1095-
8649.1998.tb00598.x).

[50] Groenewoud, F., Frommen, J.G., Josi, D., Tanaka, H., Jungwirth, A. & Taborsky, M. 2016 Predation
risk drives social complexity in cooperative breeders. *Proc. Natl. Acad. Sci. U.S.A.*, 4104–4109. (doi:
10.1073/pnas.1524178113).

[51] Buechel, S.D., Boussard, A., Kotrschal, A., van der Bijl, W. & Kolm, N. 2018 Brain size affects
performance in a reversal-learning test. *Proc. R. Soc. B-Biol. Sci.* **285**. (doi: 10.1098/rspb.2017.2031).

[52] Scott, A.P., Hirschenhauser, K., Bender, N., Oliveira, R., Earley, R.L., Sebire, M., Ellis, T., Pavlidis,
M., Hubbard, P.C., Huertas, M., et al. 2008 Non-invasive measurement of steroids in fish-holding
water: important considerations when applying the procedure to behaviour studies. *Behaviour* **145**,
1307-1328. (doi: 10.1163/156853908785765854).

[53] Wong, S.C., Dykstra, M., Campbell, J.M. & Earley, R.L. 2008 Measuring water-borne cortisol in
convict cichlids (*Amatitlania nigrofasciata*): is the procedure a stressor? *Behaviour* **145**, 1283-1305.
(doi: 10.1163/156853908785765863).

[54] Scott, A.P. & Ellis, T. 2007 Measurement of fish steroids in water--a review. *Gen Comp Endocrinol*
153, 392-400. (doi: 10.1016/j.ygcen.2006.11.006).

[55] R Core Team. 2024 R: a language and environment for statistical computing.

[56] Bates, D., Maechler, M., Bolker, B.M. & Walker, S. 2015 Fitting linear mixed-effects models using
{lme4}. *J. Stat. Softw.* **67**, 1-48. (doi: 10.18637/jss.v067.i01).

[57] Singmann, H., Bolker, B., Westfall, J., Aust, F. & Ben-Shachar, M.S. 2021 afex: analysis of factorial
experiments. https://CRAN.R-project.org/package=afex

[58] Engqvist, L. 2005 The mistreatment of covariate interaction terms in linear model analyses of
behavioural and evolutionary ecology studies. *Anim. Behav.* **70**, 967-971. (doi:
10.1016/j.anbehav.2005.01.016).

699 [59] Bannier, F., Tebbich, S. & Taborsky, B. 2017 Early experience affects learning performance and
700 neophobia in a cooperatively breeding cichlid. *Ethology* **123**, 712-723. (doi: 10.1111/eth.12646).

701 [60] Hirschenhauser, K., Taborsky, M., Oliveira, T., Canario, A.V.M. & Oliveira, R.F. 2004 A test of the
702 'challenge hypothesis' in cichlid fish: simulated partner and territory intruder experiments. *Anim.*
703 *Behav.* **68**, 741-750. (doi: 10.1016/j.anbehav.2003.12.015).

704 [61] de Kloet, E.R. 2014 From receptor balance to rational glucocorticoid therapy. *Endocrinology* **155**,
705 2754-2769. (doi: 10.1210/en.2014-1048).

706 [62] Joels, M., Karst, H., DeRijk, R. & de Kloet, E.R. 2008 The coming out of the brain mineralocorticoid
707 receptor. *Trends Neurosci.* **31**, 1-7. (doi: 10.1016/j.tins.2007.10.005).

708 [63] Fischer, S., Ferlinc, Z., Hirschenhauser, K., Taborsky, B., Fusani, L. & Tebbich, S. 2024 Does the
709 stress axis mediate behavioural flexibility in a social cichlid, *Neolamprologus pulcher*? *Physiol. Behav.*
710 **287**, 114694. (doi: 10.1016/j.physbeh.2024.114694).

711 [64] Boonstra, R. 2013 Reality as the leading cause of stress: rethinking the impact of chronic stress in
712 nature. *Funct. Ecol.* **27**, 11-23. (doi: 10.1111/1365-2435.12008).

713 [65] Giovanniello, J., Bravo-Rivera, C., Rosenkranz, A. & Matthew Lattal, K. 2023 Stress, associative
714 learning, and decision-making. *Neurobiol Learn Mem* **204**, 107812. (doi: 10.1016/j.nlm.2023.107812).

715 [66] Roozendaal, B. 2002 Stress and memory: opposing effects of glucocorticoids on memory
716 consolidation and memory retrieval. *Neurobiol Learn Mem* **78**, 578-595. (doi:
717 10.1006/nlme.2002.4080).

718 [67] Harris, B.N. 2020 Stress hypothesis overload: 131 hypotheses exploring the role of stress in
719 tradeoffs, transitions, and health. *Gen. Comp. Endocrinol.* **288**, 113355. (doi:
720 10.1016/j.ygcen.2019.113355).

721 [68] Lattin, C.R. & Romero, L.M. 2013 Seasonal variation in corticosterone receptor binding in brain,
722 hippocampus, and gonads in house sparrows (*Passer domesticus*). *The Auk* **130**, 591-598. (doi:
723 10.1525/auk.2013.13043).

724 [69] Bauer, C.M., Hayes, L.D., Ebensperger, L.A. & Romero, L.M. 2014 Seasonal variation in the degu
725 (*Octodon degus*) endocrine stress response. *Gen. Comp. Endocrinol.* **197**, 26-32. (doi:
726 10.1016/j.ygcen.2013.11.025).

727 [70] Kotrschal, K., Van Staaden, M.J. & Huber, R. 1998 Fish brains: evolution and environmental
728 relationships. *Reviews in Fish Biology and Fisheries* **8**, 373-408. (doi: 10.1023/a:1008839605380).

729 [71] Kotrschal, K., Hirschenhauser, K. & Möstl, E. 1998 The relationship between social stress and
730 dominance is seasonal in greylag geese. *Anim. Behav.* **55**, 171-176. (doi: 10.1006/anbe.1997.0597).

731 [72] Vera, F., Zenuto, R. & Antenucci, C.D. 2017 Expanding the actions of cortisol and corticosterone in
732 wild vertebrates: a necessary step to overcome the emerging challenges. *Gen Comp Endocrinol* **246**,
733 337-353. (doi: 10.1016/j.ygcen.2017.01.010).

734 [73] Landys, M.M., Ramenofsky, M. & Wingfield, J.C. 2006 Actions of glucocorticoids at a seasonal
735 baseline as compared to stress-related levels in the regulation of periodic life processes. *Gen Comp*
736 *Endocrinol* **148**, 132-149. (doi: 10.1016/j.ygcen.2006.02.013).

Factors	Estimate ± SE	X^2-value	*P*-value
Intercept	-0.47 ± 0.75	-	-
Nutritional state	-1.37 ± 0.89	2.71	0.09
Predator exposure	-*	6	**0.04**

*For estimates please see orthogonal contrasts in Table S3

Table 1: Probability of fish showing acute stress response to a novel stressor. Results are shown from
a generalized linear mixed effect model with a binomial error distribution. Predator exposure
significantly influenced the probability to show an acute stress response which was irrespective of the
nutritional state. Fish were classified as acute or prolonged responders depending on their cortisol
concentration profile for two hours on the first water collection day (see methods for more details).
In short acute responders decreased cortisol concentrations within two hours whereas prolonged
responders did not. Experimental treatments consisted of the independent manipulations of the
subject's nutritional state and predator exposure. Nutritional state was included as a two-level factor
(high, low) and the estimate is shown as difference to the reference level ('high'). Predator exposure
was included as a three-level factor ('no predator', 'occasional predators', frequent predators') and
estimates are shown as orthogonal contrasts in Table S3. To obtain p-values a likelihood ration test
was used to compare the models with and without the factor of interest. N=34 fish from 20 groups
and three blocks. p<0.05 is highlighted in bold.

Factors	Estimate ± SE	Num. D.F.	Den. D.F	F-value	*P*-value
Intercept	-28.27 ± 6.86	-	-	-	-
Body condition	9.31 ± 2.35	1	4.78	15.64	**0.01**

Table 2: The intensity of the stress reactivity after experiencing a novel stressor. Results are shown
from a linear model with a gaussian error distribution. To obtain normally distributed residuals the
dependent variable was log transformed. Fish body condition positively correlated with higher stress
reactivity. For each fish cortisol concentrations were measured every 30 minutes (four samples) for
two hours on the first water collection day (for more details see methods). The intensity of the stress
reactivity was calculated as the difference between the sample showing peak cortisol concentrations
and sample four (2h after novel stressor) and a positive value indicates a faster decrease in cortisol
concentrations. Fish were exposed to frequent predator encounters and body condition was included
as a continuous variable. To obtain p-values a likelihood ration test was used to compare the models
with and without the factor of interest. N=34 fish from 20 tanks and 3 blocks. p<0.05 are highlighted
in bold. N=11 fish from seven groups and three blocks. p<0.05 is highlighted in bold

Factors	Estimate ± SE	X^2-value	*P*-value
(a) Number of trials until learning criterion in the reversal phase			
Intercept	3.57 ± 0.07	-	-
Nutritional state	-0.1 ± 0.09	1.45	0.22
Predator exposure	-*	6.23	**0.03**
Body condition x Predator exposure	-*	5.51	*0.05*
(b) Number of trials until learning criterion in the acquisition phase			
Intercept	3.29 ± 0.12	-	-
Nutritional state	0.02 ± 0.12	0.04	0.84
Predator exposure	-	0.23	0.99
Occasional predators	$-6e^{-6}$ ± 0.14	-	-
Frequent predators	0.02 ± 0.14	-	-

*For estimates please see orthogonal contrasts in Table S5

Table 3: The number of trials until fish reached the learning criterion in the (a) reversal and (b)
acquisition phase of the learning task. Results are presented from generalized linear mixed effect
models with (a) poisson and (b) negative binomial error distributions. Nutritional state and predator
exposure interactively influenced the number of trials until fish reached the learning criterion in the
reversal but not in the acquisition phase. During the acquisition phase fish had to discriminate
between two colours to access a food reward. During the reversal phase the reward consistency was
reversed, and fish were again tested until they associated the new colour with the food reward.
Experimental treatments consisted of the independent manipulations of the subject's nutritional
state and predator exposure. Nutritional state was included as a two-level factor (high, low) and the
estimate is shown as difference to the reference level ('high'). Predator exposure was included as a
three-level factor ('no predator', 'occasional predators', frequent predators') and estimates in (a) are
shown as orthogonal contrasts in Table S5 and in (b) as difference to the reference level 'no
predators'. To obtain p-values a likelihood ration test was used to compare the models with and
without the factor of interest. N=34 fish from 20 groups and three blocks. p<0.05 is highlighted in
bold and p<0.1 is italicised.

Figure 1

Fig. 1: Timeline of the experiment. The experiment was done in three blocks (not shown) and each
block lasted nine weeks which were separated into four phases. In the first phase, fish were kept
either on an increased or reduced diet for three weeks. In the second phase, while continuing the
feeding regimes, fish were exposed to three different predator exposures: (1) a control exposure
without any predators, (2) a occasional or (3) a frequent exposure. In the third phase, we assessed
behavioural flexibility using a reversal learning task. In the fourth phase, we collected water samples
after exposure to a new stressor to measure stress response patterns in terms of cortisol
concentrations.

Figure 2

Fig. 2: Proportion (mean ± SE) of fish (N=34) showing brief stress response pattern within two hours
on the first water sample collection day. Fish were classified as brief or prolonged responders
depending on their cortisol concentration profile (see methods for more details). In short brief
responders decreased cortisol concentrations within two hours whereas prolonged responders did
not. Fish were classified based on their body condition (bc) into: 'low body condition' (red bars, fish
with a bc lower than the average) or 'high body condition' (blue bars, fish with a bc higher than the
average). Groups were exposed to three predator exposures treatments that included the
presentation of a control condition without any predators present ('no predators'), or the
presentations of predators with varying frequencies ('occasional predators' or 'frequent predators').
Fish exposed to the frequent predator treatment were more likely to show a brief stress response
pattern compared to fish exposed to occasional predators. For statistical analysis see Table 1 and
Table S3.

Figure 3

Fig. 3: The number of trials (log mean ± SE) until fish reached the learning criterion in the reversal
phase of the learning task. Fish were separated into two groups depending on their body condition:
low body condition (red bars, lower than the average) or high body condition (blue bars, higher than
the average). Groups were exposed to three predator exposure treatments that included the
presentation of a control condition without any predators present ('no predators'), or the
presentations of predators with varying frequencies ('occasional predators' or 'frequent predators').
Fish with a high body condition that were exposed to the frequent predator treatment reached the
learning criterion faster than fish with a high body condition that were exposed to the occasional
predator treatment. For statistical analysis see Table 3 and Table S5