1 **Cortical motor activity modulates respiration and reduces apnoea in**

2 **neonates**

5 Coen S. Zandvoort[1], Fatima Usman[1], Shellie Robinson[1], Odunayo Fatunla[1], Eleri

6 Adams[2], Kyle T.S. Pattinson[3,4], Simon F. Farmer[5,6], Caroline Hartley[1,*]

[1] | Department of Paediatrics, University of Oxford, Oxford, United Kingdom

[2] | Newborn Care Unit, John Radcliffe Hospital, Oxford University Hospitals NHS
Foundation Trust, Oxford, United Kingdom

[3] | Nuffield Department of Clinical Neurosciences, University of Oxford, Oxford, UK.

[4] | Nuffield Department of Anaesthetics, Oxford University Hospitals NHS Foundation
Trust, Oxford, UK.

[5] | Department of Neurology, National Hospital for Neurology and Neurosurgery,
London, United Kingdom

[6] | Department of Clinical and Human Neuroscience, UCL Institute of Neurology,
London, United Kingdom

[*] | For correspondence: caroline.hartley@paediatrics.ox.ac.uk

Abstract

Respiration is governed by a widespread network of cortical and subcortical structures.

This complex communication between the brain and lungs is altered in pathological

conditions. Apnoea – the cessation of respiration – is a common condition in infants,

particularly those born prematurely. Apnoea in infants is believed to relate to

immaturity of brainstem respiratory centres; involvement of the cortex in respiration in

infants has yet to be explored. We investigated if there was any evidence for cortical

coupling with respiration in newborn humans and whether it relates to apnoea. Using

simultaneous electroencephalography (EEG) and impedance pneumography we

investigated interactions between cortical and respiratory activity (known as cortico-

respiratory coupling) using phase-amplitude coupling. We show that cortico-

respiratory coupling is present in premature and term newborns (104 recordings from

68 infants; 34.5 ± 2.6 weeks post-menstrual age), identifying an interplay between

breathing phase and EEG amplitude. We further shed light on the biological meaning

by revealing that the strongest coupling occurs during inspiration and that cortical

activity precedes respiration, with coupling strongest over frontocentral regions. Whilst

our study was limited in spatial resolution, and determining causality is challenging,

we believe these findings support the notion that the cortico-respiratory coupling

observed here constitutes communication between cortical motor areas and lung

effectors. Moreover, we show that cortico-respiratory coupling is negatively correlated

with the rate of apnoea, revealing novel insight into this common and potentially life-

threatening neonatal pathology.

Introduction

Apnoea – the cessation of respiration – is a potentially life-threatening event[1], affecting more than 50% of preterm infants and some term infants[2]. Apnoeic episodes can result in physiological instability including oxygen desaturation and bradycardia[1], are associated with suppression of brain activity[3], and have been linked to poorer neurodevelopmental outcomes[4]. Apnoea of prematurity is caused by immaturity of the lungs, chest wall, and nervous system[5]. Immaturity of the brainstem plays a significant role in apnoea[1]; however, the relationship with other respiratory-linked brain regions is less clear. In adults with congenital central hypoventilation syndrome[6] or obstructive sleep apnoea[7], a respiratory-linked increase in cortical motor activity suggests that the motor cortex plays an important role in maintaining autonomous respiration, with the authors postulating that cortico-respiratory drive whilst participants are awake may prevent the hypoventilation/apnoea observed in these patients whilst they are asleep.

Recently, it was proposed that communication between the cortex and lungs, known as cortico-respiratory coupling, can be identified and quantified through phase-amplitude coupling[8,9]. This functional coupling arises when the amplitude of cortical activity is modulated by the respiration phase, or *vice versa*. Phase-amplitude coupling is typically quantified using non-invasive recordings capturing respiratory and neural activity (e.g., magneto- or electroencephalography [EEG]). Studies have demonstrated that the respiratory phase at low frequencies (<1 Hz) relates to higher-frequency oscillations in the cortex[9,10]. Phase-amplitude coupling therefore allows us to quantify the interplay between respiratory and cortical activity. In adults, cortico-respiratory coupling can be observed in a wide network of subcortical and cortical regions including motor areas[9]. Moreover, cortico-respiratory coupling is aberrated in pathological conditions[11].

Here, using phase-amplitude coupling, we investigated whether cortical areas are coupled with respiration in infants (Figure 1). We hypothesised that cortico-respiratory coupling occurs in newborns and that the strength of cortico-respiratory coupling is negatively associated with apnoea rate (in line with the suggestions made from studies of adults with congenital central hypoventilation syndrome[6] and obstructive sleep apnoea[7]). To this end, we first examined whether cortico-respiratory coupling exists in both premature and term infants. To investigate if the coupling might relate to central motor drive, we further studied the spatial localisation, timing

83   throughout the respiratory cycle and coupling directionality. Finally, we examined the
84   relationship between cortico-respiratory coupling and apnoea rate.

86
87   *Figure 1. Schematic of the data acquisition setup (left panel) and main findings (right panel).*
88   *Each recording consisted of cortical activity (using electroencephalography) and respiratory activity*
89   *(using impedance pneumography).*

Results and Discussion

Respiratory and cortical activity exhibit phase-amplitude coupling in infants

We analysed simultaneous 8-channel resting-state EEG (to measure cortical activity)
and impedance pneumography (IP; to measure respiration) in 68 infants (28-42 weeks
postmenstrual age [PMA] at time of recording) on 104 occasions. We first assessed
whether and how EEG and IP are interrelated using cross-frequency coupling[9,12,13]. In
line with findings in adults[9], we expected that the amplitude of oscillatory brain activity
is modulated as a function of the respiratory cycle. That is, we expected to detect an
EEG component that systematically varied with timing within the respiratory cycle. To

this end, we identified individual breaths (and thus respiratory phase) from the IP

signal[14]. Through averaging the EEG (time-locked to the IP signal) across breaths, we

observed that EEG amplitude was modulated in the delta- (0.5-4 Hz) and theta-band

(4-8 Hz) frequencies depending on the respiratory phase (Figure 2A-B). This analysis

can be seen as a first indication that the EEG amplitude in specific frequency ranges

changes systematically with the respiratory phase.

107
Figure 2. Cortico-respiratory coupling in infants*. A) Electroencephalographic (EEG) amplitude*
modulation depending on the respiratory cycle and frequency. EEG amplitude is averaged over breaths,
recordings and channels FCz and Cz. The blue graph below is the impedance pneumographic signal
time-locked from breath to breath and averaged over breaths and recordings. Grey and white bars
*below indicate expiration (Ex; in grey) and inspiration (In; in white). **B)** The time-resolved amplitude*
modulations averaged within four frequency bands (delta: 0.5-4 Hz, theta: 4-8 Hz, alpha: 8-13 Hz, beta:
13-25 Hz), focusing on channels FCz and Cz again. Shaded light-green area is the standard error
*across recordings after pooling over channels. **C)** Spatial map of coherence-based phase-amplitude*
coupling (PAC) between the respiratory phase (ωrespiratory phase) and EEG amplitude (ωEEG amplitude) – see
*Figure S1 for the pan-spectral PAC estimates. Spectra are the mean over all recordings. **D)** Spatial*
map for the corresponding (uncorrected) statistical significance relative to surrogate PAC obtained
through epoch shuffling (reported on a logarithmic scale). For all panels, data included is from 68 infants
(28-42 weeks postmenstrual age [PMA] at time of recording) on 104 recording occasions. See Table 1
for further clinical and demographic characteristics.

We then statistically confirmed this association between EEG and respiration

using phase-amplitude coupling (PAC). Using cross-frequency coherence, we

observed significant PAC between the phase at the mean respiratory frequency and

the EEG amplitude in delta and theta frequency bands (Figure 2C-D). PAC was

strongest at central electrodes (channels FCz and Cz; Figure 2C-D; *p*<0.001 false

discovery rate [FDR] corrected). In line with the results presented in Figure 2A, we did

not find significant PAC at higher EEG frequencies such as the alpha (8-13 Hz) and

beta (13-25 Hz) frequency bands (Figure S1).

These findings confirm that cortico-respiratory coupling occurs in infants. Whilst

we are the first to show that cortico-respiratory coupling already exists in early life,

adult studies have revealed widespread brain-respiratory networks covering both

cortical and subcortical areas, as well as pan-spectral components of oscillatory brain

activity (2-150 Hz)[9]. These networks have been linked to a diverse range of perceptual,

cognitive and motor processes[8,15-17]. In these studies, respiratory rhythm has been

found to relate to task performance (e.g., better performances are found during a

certain respiratory phase) as well as neural oscillations in the brain[11]. For example,

during an isometric motor task, the strength of cortico-muscular communication at

beta-band frequencies cyclically changes with the respiratory phase[16]. Moreover, the

probability of initiating goal-directed movements relates to cortical motor activity

(measured as the readiness potential amplitude) and respiratory phase[17]. These

relationships between task performance, respiration and cortical activity are well-

established and replicated in a variety of animal and human adult studies,

demonstrating the importance of cortico-respiratory coupling.

Cortico-respiratory coupling is strongest during inspiration and may be driven by

cortical activity

To further shed light on the functional meaning of cortico-respiratory coupling in

neonates, we next examined the timing and directionality of the coupling. To study

whether coupling strength fluctuates throughout the respiratory cycle, we focused on

the EEG channels with strongest coupling (FCz and Cz), and frequencies where

statistically significant coupling occurs (Figure 2D; $p<0.001$ FDR-corrected). We

confirmed that PAC changes in a time-dependent manner, observing the strongest

PAC during the inspiratory phase (Figure 3A-B). Such within-breath modulation is

evident for both delta- and theta-frequencies (Figure 3A-B). This indicates that the

strength of cortico-respiratory coupling changes throughout the respiratory cycle.

During the inspiratory phase, neural activation from the central nervous system is

required to drive respiratory motoneurons causing contraction of respiratory muscles.

Given the phasic PAC modulation as seen in Figure 3A-B, this suggests that cortical

areas centred around the vertex may play an important role in the neural activation of
respiratory muscles.

164
Figure 3. Timing and directionality of the cortico-respiratory coupling. A) *Phase-amplitude*
coupling modulation (estimated as cross-frequency coherence) throughout the respiratory cycle.
Graphs are colour-coded for the delta- and theta-frequency bands (dark- and light-blue colours
respectively), which are the EEG's primary frequency bands for which we observed statistically
significant PAC (Figure 2C-D). Shaded regions indicate standard error over recordings. PAC was
estimated for EEG channels FCz and Cz. Grey waves at the top and grey bar at the bottom indicate the
phase of the respiratory cycle. Coloured dots indicate points that are statistically significant (alpha level
was set to 0.001 with false discovery rate correction). ***B)*** *Corresponding (uncorrected) statistical*
significance analogous to the coloured dots at the top of panel A. ***C)*** *Spatial map of the cross-frequency*
phase-slope index between respiratory phase ($\omega_{respiratory\ phase}$) and EEG amplitude ($\omega_{EEG\ amplitude}$) (see
Figure S2 for the pan-spectral estimates) and ***D)*** *its (uncorrected) statistical significance (reported on a*
logarithmic scale). The spectra show the mean over all recordings. For all panels, data included is from
68 infants (28-42 weeks postmenstrual age [PMA] at time of recording) on 104 recording occasions.
See Table 1 for further clinical and demographic characteristics.

Standard coherence (i.e., our PAC measure) only quantifies coupling in a
bidirectional manner and cannot directly show any directionality, meaning that it is not
possible to distinguish whether cortical activity precedes or follows respiratory activity.
To investigate whether cortical activity precedes respiration or *vice versa*, we
employed a cross-frequency version of the phase-slope index[13]. This revealed that
delta- and theta-band EEG amplitude leads respiratory activity (Figure 3C). We
observed the strongest phase-slope index around central areas, in particular for
channel FCz (Figure 3D; *p*<0.05 FDR-corrected). Moreover, in line with the PAC
analysis (Figure 2), phase-slope index was not statistically significant for alpha- and

beta-band frequencies (Figure S2). However, caution is needed in the interpretation
of these results as signal processing techniques such as the phase-slope index
estimate directionality but do not confirm causality. Rather, they show a statistical
relationship which can be influenced by a multitude of factors (e.g., signal-to-noise
ratio and preprocessing steps). Nevertheless, the results suggest that cortical activity
may precede respiration in newborns. Future work is needed to confirm this
association by, for example, employing other metrics to estimate directionality, such
as the time-lagged cross-correlation and Granger causality and through direct
mechanistic studies.

Considering the cortico-respiratory coupling's spatial location, its timing in the
respiratory cycle and its directionality of action, combined with previous work in adults
and animal models, leads us to suggest that this coupling observed may represent the
respiratory drive from higher-order motor centres, primarily encompassing the cortical
motor areas. It is important to note that the number of electrodes in our montage (with
only 8 recording electrodes) is limited, and so source localisation was not possible;
higher-density recordings are warranted to confirm whether the motor cortex plays a
role. Structurally, the motor cortex connects to respiratory muscles both directly and
indirectly[18]. This allows for functional control over respiratory muscles (comprising
diaphragm and intercostal muscles) via the corticospinal[19] and reticulospinal tracts[20].
In cat preparations, transection of the ventrolateral part of the spinal cord (which
essentially blocks the brainstem input to respiratory motoneurons) does not prevent
cortical stimulation from eliciting responses in the respiratory intercostal muscle
motoneurons, signifying the cortico-spinal drive[20]. Such motor-evoked potentials can
also be elicited by stimulating the cortical motor areas of humans using transcranial
magnetic stimulation[21,22].

Whilst we believe that it is possible that our coupling reflects cortical motor drive,
there are several limitations to consider and further mechanistic studies are warranted.
One might also argue that cortico-respiratory coupling could reflect changes in blood
flow or oxygen saturation which induces 'coupling', where blood flow/oxygen
saturation acts as a confounding factor essentially modulating respiratory and brain
activity in a cycle-dependent manner. It is, however, unlikely that this is the case for
our results given the focal spatial localisation around central electrodes and narrow
spectral bandwidth comprising EEG frequencies between 1-8 Hz. Moreover, the

amplitude modulation of blood flow reveals a different time-frequency representation

over the respiratory cycle (Figure S3) when compared to cortical activity (Figure 2A).

To further our understanding of whether the coupling in infants relates directly

to central motor drive, future work should examine potential coupling in the cortico-

muscular domain. Electrical activity of the diaphragm (Edi) in infants can be recorded

using an Edi catheter[23], as is used in ventilators with neurally adjusted ventilatory

assist[24]. Correlating this electromyographic (EMG) activity from the diaphragm with

brain activity (e.g., EEG) will enable cortico-muscular assessments that can quantify

whether oscillatory components in the EEG and diaphragmatic EMG are phase-locked.

It is of particular interest whether linear coherence can be observed in the frequency

range of 1-8 Hz, which is expected from our PAC findings presented in Figure 2.

Interestingly, such coherence analysis has been conducted at the muscular level in

rodents. Invasively recording activity from the diaphragm and intercostal muscles

revealed strong inter-muscular coherence in the lower frequencies between 1 and 8

Hz[25]. Future work should also establish to what extent cortical motor activity is related

to other respiratory functionalities such as altering respiratory rate. In the current

investigation, we focused on how cortical amplitude directly related to the respiratory

phase but phase-amplitude coupling of cortical activity has also been revealed to play

a role in adjusting respiratory rate[26] (i.e., respiratory rate rather than individual breaths).

Cortico-respiratory coupling relates to apnoea in infants

Having identified significant cortico-respiratory coupling in infants, we then tested our

hypothesis that the strength of the cortico-respiratory coupling is negatively associated

with apnoea. Apnoea of prematurity can lead to periods of physiological instability

when apnoea is accompanied by bradycardia and desaturation[5]. These episodes may

have long-term neurological effects[4] and, hence, it is important to better understand

the pathophysiology. In addition to pulmonary immaturity, apnoea of prematurity has

been primarily attributed to immaturity and weakened neural drive from the respiratory

centres in the brainstem. These respiratory centres are principally located in the

medulla oblongata and pons within the brainstem, which ensure the periodical

activation of lung muscles to enable respiration. However, as our results suggest that

further central input to respiration may be provided by higher-order brain areas such

as the cortical motor areas, investigating whether there is an association between
cortico-respiratory coupling and apnoea in infants is warranted.

To test our hypothesis, apnoea rate was estimated over a time window of up to
24 hours (infants have their vital signs continuously monitored at the neonatal care
unit), whereas cortico-respiratory coupling was assessed on the same day from EEG
recordings with an average duration of 1.7 ± 0.6 hours. Choosing different time
windows ensured that any relationship between apnoea rate and cortico-respiratory
coupling was not the mere result of the number of breaths included in the analysis.
For instance, when the same respiratory signal is used in both analyses, a higher
apnoea rate would mean that fewer windows are used for the PAC estimation which
could result in an increased magnitude. We selected PAC pooled over EEG channels
Cz and FCz since their coupling magnitudes were the highest.

We found support for our hypothesis that apnoea and cortico-respiratory-
coupling are related. PAC was negatively associated with apnoea rate (t(98):-2.37,
p:0.02, ρ:-0.23; Figure 4A; this relationship was also identified when focusing on
individual channels [Figure S4] and preterm infants [≤36 weeks, Figure S5]). This
suggests that cortico-respiratory coupling may relate to the likelihood of infant's
experiencing apnoeas. As caffeine is given to infants for the treatment of apnoea of
prematurity, we also conducted a stratified analysis according to whether infants were
receiving caffeine at the time of the EEG recording or not (Figure S6). Conducting a
stratified analysis substantially reduces sample size, nevertheless, there was a strong
trend towards a relationship between apnoea rate and PAC in the infants receiving
caffeine, who are the infants most likely to experience apnoea (Figure S6B).

Overall, the negative association between apnoea rate and PAC provides
further support that our measures of cortico-respiratory coupling (Figures 2-3) are
related to central motor drive. It is plausible that in infants cortical motor activity assists
the brainstem's central pattern generators (CPGs) to establish increased control over
the respiratory muscles, which in our analysis, is reflected as an association with
apnoea rate. The motor cortex can exert coordination over the CPGs through its
downstream projections to the thalamus[27], which, in turn is connected to the
preBötzinger complex (the structure that primarily hosts the CPGs[28]). Nevertheless, it
is important to recognise that a limitation of this analysis is that correlation does not

imply causation, and future mechanistic studies are required to determine whether and
how cortico-respiratory coupling plays a role in reducing apnoea in infants.

288
Figure 4. Cortico-respiratory coupling dependency on apnoea rate and postmenstrual age. A)
Relationship between apnoea rate and statistically significant phase-amplitude coupling (PAC; defined
as coherence encompassing delta- and theta-band frequencies). Each blue dot is the data of an
individual recording (data of 68 infants were included from 104 test occasions). Black linear graph is
the best fit of the linear mixed-effects model (fixed factors: coupling averaged over statistically
significant samples from channels FCz and Cz [see Figure S4 for channel-specific relationships], data
length to determine apnoea rate, mode of ventilation, and postmenstrual age; random factor: infant).
Panel title contains the test statistic of the predictor's regression slope (t), its significance (p), and partial
correlation coefficient (ρ). See Figures S5-S7 for data according to PMA and whether or not the infant
received caffeine at the time of recording. ***B-C)*** *PAC spectra between the respiratory phase ($\omega_{respiratory}$*
$_{phase}$) and EEG amplitude ($\omega_{EEG\ amplitude}$) for ***B)*** *preterm (58 recordings; 33.0-36.86 weeks), and* ***C)*** *very*
preterm (28 recordings; 28.0-32.86 weeks) infants. Both spectra are averaged over recordings and EEG
channels FCz and Cz.

In theory, the strength of cortico-respiratory coupling and its inverse relationship
with apnoeas could reflect a shared relationship with PMA, where both coupling
strength and apnoea rate change depending on PMA. The apnoea rate is expected to
decrease with increased PMA across a broad age range[29]. It is also plausible that the
strength of cortico-respiratory coupling (i.e., PAC magnitude) may change with PMA,
with younger (i.e., preterm) infants potentially having weaker coupling. However, in

our dataset, we observe statistically significant PAC in preterm infants (33^{+0}-36^{+6}

weeks^{+days}; Figure 4B), and very preterm infants (28^{+0}-32^{+6} weeks^{+days}; Figure 4C).

Moreover, PAC and apnoea rate were not significantly associated with PMA (Figure

S7A-B, linear mixed-effects analysis). Similarly, PAC and apnoea rate were not

significantly related to postnatal age (PNA; defined as the age in weeks from birth;

Figure S7C-D). Whilst future studies should investigate the relationship between

cortico-respiratory coupling and age across an even broader age range, the lack of

significant associations here suggests that the relationship we observed between

apnoea rate and cortico-respiratory coupling is not determined by age.

It is known that most apnoeas in infants occur during active sleep[6,30] and delta-

and theta-band frequencies in EEG are strongly related to sleep state[31]. A limitation of

our study is that we did not record the sleep state of the infant. We speculate that

cortico-respiratory coupling may change depending on sleep state, giving rise to more

episodes of apnoea during certain sleep stages. Indeed, adult studies have suggested

that cortical respiratory drive may occur predominantly during wakefulness[6,32]. Future

work should study the effects of sleep stages on cortico-respiratory coupling to shed

further light on the role of cortical motor drive for respiration in infants.

In summary, we demonstrate the existence of cortico-respiratory coupling in

newborn infants. We believe that this coupling may be related to central motor drive

due to its spatial location (predominantly over frontocentral EEG channels), timing

(during inspiration) and directionality (with the cortex driving respiration). Consistent

with our hypothesis, we found a significant relationship between apnoea rate and

coupling strength, with infants with stronger cortico-respiratory coupling less likely to

experience apnoea. The limitations of our study need to be considered, and in

particular, directionality of the cortico-respiratory coupling, improved spatial

localisation, and a direct mechanistic link between cortico-respiratory coupling and

apnoea rate, should be investigated in future work. Nevertheless, our results have far-

reaching implications for our understanding of apnoea of prematurity as they suggest

that cortical motor areas play a pivotal role in apnoea in infants.

Data and code availability

•   Due to ethical restrictions, the raw data that support the findings of this study

are available on reasonable request. Requests can be made directly to the corresponding author (caroline.hartley@paediatrics.ox.ac.uk) or directed to the Paediatric Neuroimaging Group (University of Oxford) through the institutional email: neonatal.research@ouh.nhs.uk. Researchers will be required to agree to destroy the data within the time limits stipulated in the ethically approved study protocols. Upon agreement, data will be provided within 1 month.

- Summary-level data (before regression model adjustments of Figure 4), and channel- and recording-specific PAC spectra are provided at https://gitlab.com/paediatric_neuroimaging/cortico_respiratory_coupling.

- The codes for all analyses are available at: https://gitlab.com/paediatric_neuroimaging/cortico_respiratory_coupling. The algorithm for inter-breath interval detection is online at: https://gitlab.com/paediatric_neuroimaging/identify_ibi_from_ip.

- Any additional information required to reanalyse the data reported in this paper is available from the lead contact upon request.

Acknowledgements

This work is funded by the Wellcome Trust and Royal Society through a Sir Henry Dale Fellowship awarded to Caroline Hartley (grant number: 213486/Z/18/Z). Fatima Usman is funded by the Commonwealth Scholarship Commission. Odunayo Fatunla is funded by the Clarendon Fund in partnership with the Lincoln College Kingsgate Graduate Scholarship. We would like to thank members of the Paediatric Neuroimaging Group (University of Oxford) for help with data collection, and the babies and their parents who took part in this research.

Author contributions

Conceptualization: CSZ and CH; **Data curation:** FU, SR, OF, and CH; **Formal analysis:** CSZ; **Funding acquisition:** CH; **Investigation:** FU, SR, and OF; **Methodology:** CSZ; **Project administration:** CH; **Software:** CSZ; **Resources:** EA, KTSP and SFF; **Supervision:** CH and EA; **Visualization:** CSZ; **Writing – original draft:** CSZ; **Writing – review & editing:** all authors

Declaration of interests

The authors declare no competing interests.

Material & Methods

Participants and study design

This study was conducted at the Newborn Care Unit of the John Radcliffe Hospital,
Oxford University Hospitals NHS Foundation Trust, Oxford, UK. The National
Research Ethics Service approved the study (references: 19/LO/1085; 12/SC/0447).
Eligible families were approached and given verbal and written details about the study.
Parents provided written consent to take part in the study. The study conformed to the
standards set by the Declaration of Helsinki and Good Clinical Practice.

Clinically stable infants aged between 28- and 42-weeks postmenstrual age
(PMA) at the time of the test occasion were eligible to be included in the study. Infants
were excluded if they had intraventricular haemorrhage grades III or IV, hypoxic
ischaemic encephalopathy, congenital malformations, there was a known history of
maternal substance misuse during pregnancy, or at the time of the study, they were
receiving mechanical ventilation, opioid analgesics, or antibiotics (as a surrogate
marker of infection). These conditions are known to alter respiratory and brain activity.
We enrolled 74 infants who were recorded on 123 separate occasions (range: 1-8
recordings per infant). Fifty-one infants were included in one or two test occasions; 23
infants took part in the ongoing *Breathing and Brain Development* study where they
are studied approximately once a week whilst in the Newborn Care Unit (included on
3±2 [median ± interquartile range] occasions). Demographic details of all infants can
be found in Table 1. Parts of this dataset have been reported earlier in Zandvoort et
al. [33] to address a different research question (this study investigated the development
of sensory-evoked potentials, which were also recorded in these infants; it did not
explore respiration).

Factors	
Age	
Postmenstrual age at recording (weeks, mean±SD)	34.5 ± 2.6
Gestational age at birth (weeks, mean±SD)	31.1 ± 3.4
Postnatal age at recording (weeks, mean±SD)	3.4 ± 2.2

Birthweight (*g*, mean±SD*)	1,726 ± 816
Sex	
Females (n)	27
Males (n)	47
Mode of delivery	
Normal vaginal delivery (n)	16
Vaginal assisted (ventouse/forceps/kiwi) (n)	4
Elective Caesarean section (n)	7
Emergency Caesarean section (n)	47
Apgar scores	
Apgar at 1 minute (median [IQR])	8 [3]
Apgar at 5 minutes (median [IQR])	10 [2]
Apgar at 10 minutes (median [IQR])	10 [0]
Respiratory distress syndrome	
Ongoing condition (n)	39
Past treated (n)	48
No (n)	36
Persistent pulmonary hypertension	
Past treated (n)	5
No (n)	118
Infection	
None (n)	48
Treated (n)	75
Respiratory support	
High flow therapy (n)	25
Low flow therapy (n)	16
Spontaneous ventilation (n)	82
Caffeine received on the day of the EEG recording	
Yes (n)	59
No (n)	64

401 **Table 1. Infant demographics.** *Reported values are number of babies or recordings, or mean and*
402 *standard deviation [SD]/median and interquartile range [IQR] – as indicated. Values are reported over*
403 *recordings for ages (postmenstrual, gestational, and postnatal), respiratory distress syndrome,*
404 *persistent pulmonary hypertension, infection and respiratory support. Birthweight, sex, mode of delivery,*
405 *and Apgar scores are provided over infants.*

Data recordings

During each test occasion, approximately one to two hours of simultaneous electroencephalography (EEG) and vital signs (respiration, heart rate and oxygen saturation) were recorded. We continuously recorded the vital signs throughout their hospital stay of the infants who took part in the *Breathing and Brain Development* study.

Vital signs were recorded for each infant using Phillips IntelliVue MX800 or MX750 monitors. These data were continuously downloaded from the vital signs monitor using an electronic data capture software (iXtrend, ixitos, Germany). Both electrocardiograph (ECG; sampling frequency: 250 Hz) and impedance pneumography (IP; sampling frequency: 62.5 Hz) were recorded with three electrodes placed on the infant's chest. Peripheral photoplethysmography was recorded at 125 Hz from a probe placed on the infant's foot or hand.

EEG was recorded from DC to 400 Hz using a SynAmps RT 64-channel headbox and amplifiers and CURRYscan7 neuroimaging suite (Compumedics Neuroscan) at a sampling rate of 2 kHz. Eight electrodes were placed at Cz, CPz, C3, C4, Oz, FCz, T3, and T4. The ground electrode was placed at FPz and the reference electrode at Fz. The infant's scalp was cleaned with NuPrep gel (D.O. Weaver and Co., Aurora, USA) to achieve impedances of 5-10 kΩ. A conductive paste (Elefix EEG paste, Nihon Kohden, Tokyo, Japan) was used to attach disposable Ag/AgCl cup EEG electrodes (Ambu Neuroline, Ballerup, Denmark) to the infant's scalp. EEG was recorded for an average of 1.7 ± 0.7 hours (mean ± SD). ECG was co-registered with our EEG recordings using a single electrode (Ambu Neuroline, Ballerup, Denmark) placed on the infant's chest (referenced to Fz).

To synchronise the vital signs and EEG recordings, we registered approximately 30 annotations simultaneously on both recordings using a bespoke triggering device (the PiNe box)[34]. Annotations were used to mark resting state as well as stimulus-evoked activity (these responses were used in other studies to investigate infant brain maturity – see Zandvoort *et al.* [33] – but were used here only to synchronise the EEG and vital signs recordings). The PiNe box enables synchronisation with an average precision of 105 ± 14 ms (mean ± SD)[34]. Annotation timings were matched

between both vital signs and EEG recordings. Next, to further improve synchronisation,

we utilised the ECG signals which were recorded on both the vital signs and EEG

recording devices. To do so, ECG epochs were created from -10 to 10 seconds around

each annotation. We aligned the ECG signals around each of these annotation epochs

using cross correlations, providing a temporal delay between vital signs and EEG[34].

Temporal synchronisation between vital signs and EEG recordings was achieved by

taking the median of the time differences across all annotations.

From the total of 123 recordings, we excluded 19 recordings due to (a) missing

IP recordings (5 recordings), (b) missing annotations (9 recordings), and (c)

inaccuracies in EEG-IP syncing (5 recordings; standard deviations over annotations

exceeding 1 sec for annotation alignment or 0.1 sec during ECG alignment). Hence,

data comprising 104 recordings from 68 infants were used in the analysis.

Breath and apnoea identification

IP signals were processed to identify breaths (including apnoeas). To this end, we

used an algorithm validated to detect inter-breath intervals in infants[14]. Briefly, the

algorithm attenuates noise introduced by movements and cardiac activity by filtering

the signal. An adaptive amplitude threshold then identifies individual breaths. This

threshold was set at 0.4 times the standard deviation of the signal over the preceding

15 breaths. Finally, a support vector machine algorithm was applied to all long (here,

15 seconds or greater) inter-breath intervals to separate episodes of apnoea

(cessation of respiration) and periods of low amplitude erroneously identified as

apnoeas due to noise or shallow breathing. The inter-breath intervals marked as

noise/shallow breathing were discarded from further analysis. The model was trained

and validated on IP signals acquired during a clinical trial[35], which was a completely

independent data sample to the one used in the current study.

Phase-amplitude coupling

In line with Osipova *et al.* [12], we defined phase-amplitude coupling (PAC) as cross-

frequency coherence between a low-frequency phase and high-frequency amplitude.

Note that the term "phase-amplitude coupling" is used interchangeably in the literature

to mean the coupling principle of the statistical relationship between phases and
amplitudes as well as the actual measure to detect this. We reserve the abbreviation
for the latter. For our PAC measure, this meant that coherence was computed between
the filtered IP signal and instantaneous amplitude of the EEG signal at some higher
frequency (i.e., the second-order moment between the IP phase and EEG amplitude).
More formally, in our case PAC was defined as:

$$C(f_1, f_2) = \frac{|\sum_{n=1}^{N} X(f_1) * Y(f_1, f_2)^*|}{\sqrt{\sum_{n=1}^{N}|X(f_1)|^2 * \sum_{n=1}^{N}|Y(f_1, f_2)|^2}}$$ Eq. 1

with X being the IP signal at frequency f_1, Y the instantaneous EEG amplitude, and $*$
the complex conjugate. This instantaneous amplitude was obtained using the
magnitude of the Hilbert transform after narrowly filtering the data at frequency f_2. This
component is often filtered with a bandwidth that relates to the low-frequency
component f_1. Following earlier recommendations, the filter bandwidth of the high-
frequency component (at f_2) was equal to a single time the frequency of the low-
frequency component f_1[36] (e.g., when f_1 equals 5 Hz the filter bandwidth of f_2 is set to
5 Hz).

Data were segmented into 4-sec epochs (windows) (50% overlap), which
meant that the frequency resolution was equal to 0.25 Hz (allowing to capture low
delta-band frequencies), which comprised around four breaths. To focus on regular
respiration, we discarded epochs which contained fewer than three detected breaths.
An epoch was also removed from further analysis when its standard deviation
exceeded 3 times the mean or median of the standard deviation of all epochs (which
was done separately for the EEG and IP). Finally, we rejected any epoch containing
an absolute value greater than 10 times the mean or median of the standard deviation
of all epochs. Before cross-spectrum computation, EEG and IP epochs were
convolved with a wavelet constructed from a Hanning window.

Changes in PAC throughout the respiratory cycle

To investigate how cortico-respiratory coupling changes throughout the respiratory
cycle, we estimated PAC at 50 equally spaced timepoints within the respiratory cycle.
To this end, we time-locked both EEG and IP to each inter-breath interval as identified
by the algorithm of Adjei *et al.* [14]. At each respiratory cycle phase, cross- and auto-
spectra were obtained to estimate PAC over all breaths within a recording. We limited
the analysis to EEG channels FCz and Cz, because they exhibited the highest PAC
magnitude (Figure 2), as well as statistically significant frequency-frequency samples
relative to the surrogate PAC (see Figure 2D for the *p*-values of this statistical
comparison). PAC was computed within the different frequency bands (delta: 0.5-4
Hz; and theta: 4-8 Hz) to see whether its modulation was specific for each frequency
band. The same preprocessing steps apply as for PAC computation described earlier
apart from the fact that data were segmented into 2/3 sec windows (so that temporal
changes would appear within the respiratory cycle).

Directionality of phase-amplitude coupling

To assess whether the cortical activity was preceding or trailing relative to the
respiratory activity, we employed a cross-frequency version of the phase-slope index.
Briefly, this metric estimates the phase lags over narrow frequency ranges and
quantifies whether they increase or decrease over this range by estimating the phase-
frequency slopes. Increasing or decreasing slopes signify changing time lags
depending on the frequency of the oscillation and can therefore be indicative of
whether a signal leads or trails another signal. We used the phase-slope index (Ψ) as
proposed by Jiang *et al.* [13], which was defined as:

$$\Psi(f_1, f_2) = imag\left(\sum_{f_2 - \frac{\beta}{2}}^{f_2 + \frac{\beta}{2}} \tilde{C}^*(f_1, f_2)\, \tilde{C}(f_1, f_2 + \Delta f)\right) \qquad \text{Eq. 2}$$

with $\tilde{C}$ being the coherency (i.e., the normalised cross-spectrum) of C defined in Eq. 1.
Δf is the frequency resolution and β the bandwidth used to estimate the phase slope.

In line with previous recommendations by Jiang *et al.* [13], we set β to 1 Hz (i.e., four

times the frequency resolution).

Surrogate analysis

To assess the statistical significance of PAC (i.e., coherence) and phase-slope index

at each of the EEG channels, we performed a surrogate analysis under the null

hypothesis that the means of the true and surrogate PAC (and phase-slope index)

were equal. The surrogate analysis was performed by randomly permuting the epoch

(4-s windows used for PAC calculation – see previous sections) order of the EEG

amplitude while leaving the respiration epoch order unchanged (i.e., respiration

segment/epoch S1 was paired with an EEG segment/epoch Sj, j ≠ 1). Importantly, no

temporal samples were shuffled within epochs. Thus, the full within-epoch temporal

structure, including autocorrelation and spectral profile (auto-spectra), was preserved

for both signals. This permutation destroys trial-wise cross-signal phase alignment

(and therefore coherence) while retaining the intrinsic dynamics of each signal. We

obtained a surrogate spectrum for each recording. Next, the true and surrogate

spectra were statistically compared on a sample-by-sample basis using dependent *t*-

tests[37] (using the FieldTrip toolbox – version 20210325). This comparison was

performed over all recordings. Statistical significance was defined using the false

discovery rate correction (with an alpha level set at 0.05 [phase-slope index] or 0.001

[coherence]).

Statistical analysis of relationships between cortico-respiratory coupling and apnoea

We tested our main hypothesis that PAC is negatively associated with apnoea rate.

Vital signs were continuously monitored for the infants that took part in the *Breathing*

and Brain Development study, which allowed us to analyse their vital signs in longer

time windows than the EEG. We here selected up to 24 hours of vital signs from the

day of the EEG recording (mean ± standard deviation: 9.5 ± 9.9 hours) and expressed

this as a rate per hour. We then constructed linear mixed-effects models to study the

relationships between PAC and apnoea rate. We included the coherence magnitude

as a fixed factor, which was averaged over the statistically significant samples from

channels FCz and Cz (see Figure S4 for the channel-specific relationships). As fixed

effects, we included (1) data length of the respiratory signal, (2) mode of ventilation

('Self-ventilating', 'Low flow', 'High flow'; Figure S8), and (3) PMA (Figure S7). Infant

ID was included as a random effect acting on the intercept. Data length, mode of

ventilation and PMA were included as confounding factors. P-values (for the

regression coefficient of PAC) and partial correlation coefficients were reported from

these models. The normality assumption was verified with Q-Q plots. We also tested

the linearity assumption (see Supplementary text). For visualisation (Figure 4), we

used the plotAdjustedResponse function in MATLAB to adjust for the fixed factors

(calculating a separate model without the random effect of an infant). Regression lines

are visualised by calculating the line-of-best-fit on the adjusted data.

References

1. Martin, R.J., and Abu-Shaweesh, J.M. (2005). Control of breathing and neonatal apnea. Neonatology *87*, 288-295.
2. Finer, N.N., Higgins, R., Kattwinkel, J., and Martin, R.J. (2006). Summary proceedings from the apnea-of-prematurity group. Pediatrics *117*, S47-S51.
3. Zandvoort, C.S., Dereymaeker, A., Baxter, L., Jansen, K., Naulaers, G., de Vos, M., and Hartley, C. (2024). Apnoea suppresses brain activity in infants. Imaging Neuroscience *2*, 1-14.
4. Williamson, M., Poorun, R., and Hartley, C. (2021). Apnoea of prematurity and neurodevelopmental outcomes: Current understanding and future prospects for research. Frontiers in Pediatrics *9*, 755677.
5. Martin, R.J., Abu-Shaweesh, J.M., and Baird, T.M. (2004). Apnoea of prematurity. Paediatric Respiratory Reviews *5*, S377-S382.
6. Tremoureux, L., Raux, M., Hudson, A.L., Ranohavimparany, A., Straus, C., and Similowski, T. (2014). Does the supplementary motor area keep patients with Ondine's curse syndrome breathing while awake? PLoS One *9*, e84534.
7. Launois, C., Attali, V., Georges, M., Raux, M., Morawiec, E., Rivals, I., Arnulf, I., and Similowski, T. (2015). Cortical drive to breathe during wakefulness in patients with obstructive sleep apnea syndrome. Sleep *38*, 1743-1749.
8. Kluger, D.S., Balestrieri, E., Busch, N.A., and Gross, J. (2021). Respiration aligns perception with neural excitability. Elife *10*, e70907.
9. Kluger, D.S., and Gross, J. (2021). Respiration modulates oscillatory neural network activity at rest. PLoS biology *19*, e3001457.
10. Tort, A.B., Brankačk, J., and Draguhn, A. (2018). Respiration-entrained brain rhythms are global but often overlooked. Trends in neurosciences *41*, 186-197.
11. Brændholt, M., Kluger, D.S., Varga, S., Heck, D.H., Gross, J., and Allen, M.G. (2023). Breathing in waves: Understanding respiratory-brain coupling as a gradient of predictive oscillations. Neuroscience & Biobehavioral Reviews *152*, 105262.
12. Osipova, D., Hermes, D., and Jensen, O. (2008). Gamma power is phase-locked to posterior alpha activity. PloS one *3*, e3990.
13. Jiang, H., Bahramisharif, A., van Gerven, M.A., and Jensen, O. (2015). Measuring directionality between neuronal oscillations of different frequencies. Neuroimage *118*, 359-367.
14. Adjei, T., Purdy, R., Jorge, J., Adams, E., Buckle, M., Fry, R.E., Green, G., Patel, C., Rogers, R., and Slater, R. (2021). New method to measure interbreath intervals in infants for the assessment of apnoea and respiration. BMJ Open Respiratory Research *8*, e001042.
15. Perl, O., Ravia, A., Rubinson, M., Eisen, A., Soroka, T., Mor, N., Secundo, L., and Sobel, N. (2019). Human non-olfactory cognition phase-locked with inhalation. Nature human behaviour *3*, 501-512.
16. Kluger, D.S., and Gross, J. (2020). Depth and phase of respiration modulate cortico-muscular communication. Neuroimage *222*, 117272.
17. Park, H.-D., Barnoud, C., Trang, H., Kannape, O.A., Schaller, K., and Blanke, O. (2020). Breathing is coupled with voluntary action and the cortical readiness potential. Nature communications *11*, 289.

18. Urfy, M.Z., and Suarez, J.I. (2014). Breathing and the nervous system.
Handbook of clinical neurology *119*, 241-250.
19. Evans, K.C., Shea, S.A., and Saykin, A.J. (1999). Functional MRI localisation
of central nervous system regions associated with volitional inspiration in
humans. The Journal of Physiology *520*, 383-392.
20. Aminoff, M., and Sears, T. (1971). Spinal integration of segmental, cortical and
breathing inputs to thoracic respiratory motoneurones. The Journal of
Physiology *215*, 557-575.
21. Gandevia, S., and Rothwell, J. (1987). Activation of the human diaphragm from
the motor cortex. The Journal of physiology *384*, 109-118.
22. Welch, J.F., Argento, P.J., Mitchell, G.S., and Fox, E.J. (2020). Reliability of
diaphragmatic motor-evoked potentials induced by transcranial magnetic
stimulation. Journal of Applied Physiology *129*, 1393-1404.
23. Essouri, S., Baudin, F., Mortamet, G., Beck, J., Jouvet, P., and Emeriaud, G.
(2019). Relationship between diaphragmatic electrical activity and esophageal
pressure monitoring in children. Pediatric Critical Care Medicine *20*, e319-e325.
24. Fang, S.-J., Chen, C.-C., Liao, D.-L., and Chung, M.-Y. (2023). Neurally
adjusted ventilatory assist in infants: a review article. Pediatrics & Neonatology
64, 5-11.
25. Rice, A., Fuglevand, A.J., Laine, C.M., and Fregosi, R.F. (2011).
Synchronization of presynaptic input to motor units of tongue, inspiratory
intercostal, and diaphragm muscles. Journal of neurophysiology *105*, 2330-
2336.
26. Hammer, M., Schwale, C., Brankačk, J., Draguhn, A., and Tort, A.B. (2021).
Theta-gamma coupling during REM sleep depends on breathing rate. Sleep *44*,
zsab189.
27. Mckay, L.C., Evans, K.C., Frackowiak, R.S., and Corfield, D.R. (2003). Neural
correlates of voluntary breathing in humans. Journal of applied physiology *95*,
1170-1178.
28. Del Negro, C.A., Funk, G.D., and Feldman, J.L. (2018). Breathing matters.
Nature Reviews Neuroscience *19*, 351-367.
29. Henderson-Smart, D. (1981). The effect of gestational age on the incidence and
duration of recurrent apnoea in newborn babies. Journal of Paediatrics and
Child Health *17*, 273-276.
30. Lehtonen, L., and Martin, R.J. (2004). Ontogeny of sleep and awake states in
relation to breathing in preterm infants. In 3. (Elsevier), pp. 229-238.
31. Jenni, O.G., Borbély, A.A., and Achermann, P. (2004). Development of the
nocturnal sleep electroencephalogram in human infants. American journal of
physiology-regulatory, integrative and comparative physiology *286*, R528-
R538.
32. Dubois, M., Chenivesse, C., Raux, M., Morales-Robles, A., Nierat, M.-C.,
Garcia, G., Navarro-Sune, X., Chavez, M., Martinerie, J., and Similowski, T.
(2016). Neurophysiological evidence for a cortical contribution to the
wakefulness-related drive to breathe explaining hypocapnia-resistant
ventilation in humans. Journal of Neuroscience *36*, 10673-10682.
33. Zandvoort, C.S., van der Vaart, M., Robinson, S., Usman, F., Mellado, G.S.,
Fry, R.E., Worley, A., Adams, E., Slater, R., and Baxter, L. (2024). Sensory
event-related potential morphology predicts age in premature infants. Clinical
Neurophysiology *157*, 61-72.

34. Worley, A., Fabrizi, L., Boyd, S., and Slater, R. (2012). Multi-modal pain
measurements in infants. J Neurosci Methods *205*, 252-257.
35. Hartley, C., Moultrie, F., Hoskin, A., Green, G., Monk, V., Bell, J.L., King, A.R.,
Buckle, M., van der Vaart, M., and Gursul, D. (2018). Analgesic efficacy and
safety of morphine in the Procedural Pain in Premature Infants (Poppi) study:
randomised placebo-controlled trial. The Lancet *392*, 2595-2605.
36. Zandvoort, C.S., and Nolte, G. (2021). Defining the filter parameters for phase-
amplitude coupling from a bispectral point of view. Journal of Neuroscience
Methods *350*, 109032.
37. Maris, E., and Oostenveld, R. (2007). Nonparametric statistical testing of EEG-
and MEG-data. J Neurosci Methods *164*, 177-190.
673
674