RNA Editors Sculpt the Transcriptome During Terminal Erythropoiesis

Areum Han[1,2], Alena Yermalovich[3,4], Mohamad Ali T. Najia[1,4,5,6], Daniel S. Pearson[7], Yuko Fujiwara[10,11], Michael Bolgov[1], Caroline Kubaczka[6], Trista E. North[6,8], Vanessa Lundin[9], Stuart Orkin[10, 11], George Q. Daley[1,2,6*]

[1]Division of Hematology/Oncology, Boston Children's Hospital, Boston, MA, USA. [2]Department of Biological Chemistry and Molecular Pharmacology, Harvard Medical School, Boston, MA, USA. [3]Department of Medical Oncology, Dana-Farber Cancer Institute, Boston, MA, USA. [4]Broad Institute of Harvard and MIT, Cambridge, MA, USA. [5]Harvard-MIT Division of Health Sciences and Technology, Massachusetts Institute of Technology, Cambridge, MA, USA. [6]Stem Cell Program, Boston Children's Hospital, Boston, MA, USA. [7]Department of Pathology, Massachusetts General Hospital, Boston, MA, USA. [8]Developmental and Regenerative Biology Program, Harvard Medical School, Boston, MA, USA. [9]Center for Hematology and Regenerative Medicine, Department of Medicine Huddinge, Karolinska Institutet, Huddinge, Sweden. [10]Dana-Farber/Boston Children's Hospital Cancer and Blood Disorder Center, Department of Pediatrics, Harvard Medical School, Boston, MA, USA. [11]Howard Hughes Medical Institute, Harvard Medical School, Boston, MA, USA.

*Corresponding author: george.daley@childrens.harvard.edu

ABSTRACT

Selective RNA degradation during terminal erythropoiesis results in a globin-rich transcriptome in mature erythrocytes, but the specific RNA decay pathways remain unknown. We found that deficiency of the terminal uridylyl transferase enzyme Zcchc6 and the 3'-5' exoribonuclease Dis3l2 in mouse models led to fetal and perinatal reticulocytosis, an accumulation of RNA-rich precursors of terminal erythroid cells, suggesting their crucial roles in terminal red cell differentiation. Notably, knockout embryos exhibited persistent high-level expression of *Hbb-bh1* globin, the ortholog of human fetal γ-globin. Perturbation of the Zcchc6-Dis3l2 pathway in mice engineered to express the human β-globin locus likewise increased γ-globin levels in fetal erythroid cells, suggesting that globin switching entails post-transcriptional mechanisms of mRNA destabilization in addition to transcriptional down-regulation. We cultured human hematopoietic stem and progenitor cells (HSPCs), performed CRISPR/Cas9-mediated knockout of ZCCHC6 and DIS3L2, and observed accumulation of RNA and elevated γ-globin levels in terminal erythroid cells. Our findings reveal a conserved role for the

37 ZCCHC6/DIS3L2 RNA editors in terminal erythropoiesis and demonstrate a post-
38 transcriptional mechanism for γ-globin gene switching, advancing research into *in vitro*
39 erythrocyte generation and γ-globin stabilization to ameliorate hemoglobinopathies.

INTRODUCTION

Terminal differentiation of red blood cells (RBCs)[1,2] entails the loss of key cellular
components such as the nucleus, mitochondria, and ribosomes[3–7]. After enucleation,
orthochromatic erythroid cells become reticulocytes, which then undergo extensive RNA
degradation to develop into mature erythrocytes, primarily composed of globins[8].
Previous examination of RNA half-life suggests that active decay of non-erythrocyte
transcripts contributes to this globin-rich, low-entropy transcriptome[9,10], in addition to
global proteome remodeling through protein tagging by ubiquitination[11].

The Terminal Uridylyl Transferase ZCCHC6 (TUTase 7) has recently been
implicated in regulating the degradation of various RNAs during cellular transitions, rather
than in steady-state conditions[12–20]. This re-sculpting of the transcriptome occurs in
transcriptionally compromised or quiescent contexts, such as when cells are treated with
the transcription inhibitor Actinomycin D[21] or during the maternal-to-zygotic transition
(MZT)[22]. During MZT, a sharp decline in maternal transcripts accelerates the emergence
of the zygotic transcriptome. This coordinated transcriptome remodeling is comparable to
the terminal stages of erythropoiesis.

Embryonic, fetal, and adult hemoglobins are sequentially expressed during mouse
and human erythroid ontogeny[23]. In humans, during the switch from fetal globin (HbF;
α2γ2) to adult globin (α2β2), γ-globins are replaced by adult β-globins. Conversely, faulty
adult β-globin with pathogenic mutations can be rescued by reactivating γ-globin
expression. Indeed, elevated γ-globin levels in adults have been shown to ameliorate
symptoms of sickle cell disease[24,25]. Due to this therapeutic potential, the transcriptional
regulation of γ-globin has been extensively studied[26,27], while its post-transcriptional
regulation remains less explored[28,29].

Prior studies have shown that the ZCCHC6 adds uridines to the 3' end of
target RNAs, which serves as a recognition signal for the 3' to 5' exonuclease DIS3L2,
leading to the degradation of target RNAs[12,18,30,31]. *Zcchc6*, uniquely recognized for its

RNA-tagging capability, is upregulated in the peripheral blood of mice exposed to chronic

hypoxia and subsequently downregulated during normoxic recovery [32], suggesting that

increased levels of ZCCHC6 might be necessary to process the higher number of

reticulocytes generated during erythropoietic stress. ZCCHC6 appears to be a rate-

limiting factor in RNA editing, as it is slower than nuclease activity. Based on all the above,

we posited that the ZCCHC6-DIS3L2 axis might regulate RNA fate during terminal

erythropoiesis, accelerating RNA degradation and potentially influencing globin switching.

In this study, we explored the role of the RNA editor-exonuclease axis during terminal

erythropoiesis using murine models and in vitro erythroid differentiation of human blood

progenitors. Our findings revealed that loss of this axis leads to RNA accumulation and

upregulation of fetal globin expression, resulting in a reversed globin switching pattern

during terminal erythropoiesis. This highlights a novel, conserved post-transcriptional

gene regulation pathway during late erythropoiesis.

RESULTS

Whole-body *Zcchc6/Dis3l2* KO mice exhibit reticulocytosis and anemia.

We began our investigation by analyzing the expression profiles of Zcchc6 and

Dis3L2 throughout erythropoietic ontogeny, examining blood from knockout (KO)

models across embryonic stages to adulthood. Analysis of a public dataset

demonstrated that the expression of *Zcchc6* is upregulated as erythroid cells

differentiate into the terminal stages of fetal and adult definitive erythropoiesis[33,34]

(Extended Data Fig. 1, upper panel), suggesting that Zcchc6 plays an important role in

terminal erythropoiesis. In contrast, *Dis3l2* did not show high expression in the terminal

stages (Extended Data Fig. 1, lower panel). As an exonuclease, Dis3L2 may not be a

limiting factor in RNA regulation.

We generated a whole-body *Dis3l2* KO model (Extended Data Fig. 2), but the

Dis3l2 KOs exhibited postnatal lethality. Adult *Zcchc6* KOs survived and demonstrated

an increase in reticulocytes compared to wild-type (WT) controls, suggesting a role for

Zcchc6 during the transition from reticulocytes to mature RBCs (Extended Data Fig. 3c).

Other hematological parameters in the *Zcchc6* KO were mostly comparable to those of

the WT in young adult stages (Extended Data Fig. 3 a&b). Zcchc6 KO animals exhibited

increased neutrophil counts compared to heterozygous (HET) animals (Extended Data

Fig. 3b).

At embryonic day (E) 18.5, cells are undergoing RNA degradation as they

differentiate from reticulocytes to erythrocytes, and both KOs are viable. Therefore, we

investigated animals at this age and observed embryonic anemia and reticulocytosis in

both KO models. In detail, we collected peripheral blood and spleen cells from E18.5

embryos, and performed histology, flow cytometry, and hemogram analysis (Fig.1a).

Examination of blood smears from *Zcchc6* and *Dis3l2* KO embryos showed

polychromasia and increased RNA retention in red cells, as revealed by staining with

methylene blue, indicating reticulocytosis (Fig.1b). We quantified reticulocytosis using

flow cytometry, pre-gating erythroid (TER119+) cells and staining for the CD71

transferrin receptor and the nucleic acid dye Thiazole Orange (TO) [35,36] (Fig.1c). WTs

displayed expected patterns of CD71 loss and RNA (TO signal) loss, while KOs showed

higher levels of RNA and a significantly higher $CD71^{low}TO^{high}$ population relative to

HETs and WTs (Fig.1c). These data demonstrate that Zcchc6 and Dis3l2 impact RNA

degradation during terminal erythropoiesis. We also profiled erythroid RNA dynamics in

the embryonic spleen, an erythropoietic organ at E18.5[37]. Embryonic spleen

harbors nucleated erythroid cells, and we observed a large drop in TO signal in the

$CD71^{high}$ population due to enucleation[35] (Extended Data Fig.4a). However, we did not

observe differences in enucleation ratio across genotypes. On the other hand, in the

later stage of erythroid cell differentiation ($TER119^{+}CD71^{low}$), the $CD71^{low}TO^{high}$

population was elevated in KOs compared to HETs and WTs (Extended Data Fig.4a),

again indicating inefficient RNA degradation in KOs. In both peripheral blood and

spleen, this accumulation of the $CD71^{low}TO^{high}$ population was genotype-dependent on

Zcchc6 (Extended Data Fig.4a, right graph). This result suggests that cellular levels of

Zcchc6 determine the extent of RNA degradation.

Both KOs at E18.5 had reduced RBC numbers, decreased hematocrit (HCT) and

increased Mean Corpuscular Hemoglobin (MCH) compared to HETs and WTs (Fig.1d).

Zcchc6 KO also showed increased Red Cell Distribution Width (RDW); a trend observed

in *Dis3l2* KO that failed to reach statistical significance in WT. For *Dis3l2* KOs, we

observed a significant increase in Mean Corpuscular Volume (MCV) and lower

hemoglobin concentration (Hb), a trend also observed in *Zcchc6* KOs but without statistical significance. Both KOs were anemic from a reduced number of RBCs, and both showed lymphocytosis (Extended Data Fig. 4b). *Zcchc6* KO generally showed more marked differences in the leukocyte panel with an increased absolute number and percentage of monocytes (MO), a decreased percentage of basophils (BA), and increased number in neutrophils (NE). These interesting observations could be further examined in future studies.

Conditional *Zcchc6/Dis3l2* KO mice phenocopy whole-body KOs.

To examine erythrocyte-specific phenotypes, we utilized animals in which the *Zcchc6/Dis3l2* alleles were flanked by LoxP sites (floxed) to generate conditional KOs of these genes by crossing to strains expressing *EpoR-Cre* [38] (Fig. 2a & Extended Data Fig. 2). Conditional KO in erythroid cells (TER119+) was confirmed by western blot (Extended Data Fig. 2b). Indeed, cKOs phenocopied the reticulocytosis observed in whole-body KOs (Fig. 2b), indicating that this phenotype is cell-autonomous. Both cKOs exhibited a statistically significant increase in the CD71^{low}TO^{high} population in peripheral blood (Fig. 2b) and spleen (Extended Data Fig. 5a) compared to *EpoR-Cre*-carrying and cWT animals. Hematological readings in both cKOs showed consistent reduction in RBC numbers (Fig. 2c). cKOs again showed a trend of increased absolute WBC numbers, specifically in the absolute number of lymphocytes (Extended Data Fig. 5b). To demonstrate that the observed reticulocytosis reflected failed red cell maturation and not hemolytic anemia, we confirmed preserved levels of haptoglobin and the absence of elevated LDH and bilirubin levels [39] (Extended Data Fig. 6). Taken together, these results indicate that the observed reticulocytosis arises from an RNA degradation defect intrinsic to the erythroid cell during terminal differentiation.

Normal terminal erythropoiesis in KO mice lacking *Lin28a/b* and *Zcchc11*.

An enzymatic axis regulating microRNA biogenesis has been described whereby the RNA binding proteins LIN28A/B dock to the loop of the microRNA precursor *pre-let-7*, which then recruits ZCCHC6 or the paralog ZCCHC11, resulting in an inhibitory tri-complex[18]. The assembled tri-complex facilitates oligouridylation, and the

oligouridylated RNA species then become substrates for DIS3L2[30] (Extended Data Fig.

7a). However, neither *Lin28a* nor *Lin28b* KOs exhibited reticulocytosis (Extended Data

Fig. 7b). Paralogs often perform similar functions and manifest redundancy. Among

seven TUTases in humans, ZCCHC6 and ZCCHC11 are the two largest TUTases with

CCHC and C2H2-type zinc finger domains[18]. Previous studies have shown that

ZCCHC6 and ZCCHC11 have redundant RNA editing activity [21,31,40]. Importantly, while

Zcchc11 heterozygotes exhibited slight reticulocytosis, this was not observed in

Zcchc11 KOs, suggesting that Zcchc11 may provide at best partial redundancy

(Extended Data Fig. 7b). The *Zcchc6* and *Zcchc11* quad-flox system also showed that

significant reticulocytosis was observed only in animals with *Zcchc6* cKO (homozygous

floxed alleles of *Zcchc6* driven by *EpoR-Cre)*, but not in *Zcchc11* cKO (Extended Data

Fig. 8).

Reticulocytosis showed postnatal attenuation.

Dis3l2 whole-body KO mice die shortly after birth, whereas *Dis3l2* cKOs under

EpoR-Cre survive postnatally, allowing us to investigate adult peripheral blood. Both

Zcchc6 KO and *Dis3l2* cKOs showed significant reticulocytosis at Postnatal day (P) 2

and P21 (Extended Data Fig. 9). Interestingly, the degree of reticulocytosis was

attenuated with age. We postulate there may be differential RNA regulation in the bone

marrow and adult peripheral blood compared to the embryonic spleen and blood.

Characterization of accumulated RNA in *Zcchc6* and *Dis3l2* KO mice.

Next, we investigated the nature of RNA accumulated in *Zcchc6* and *Dis3l2* KOs

at E18.5 utilizing RNA-seq (Fig. 3). The estimated expression levels in *Zcchc6* and

Dis3l2 KOs clustered by KO genotype (Extended Data Fig. 10b). In PCA analysis, the

Dis3l2 KO differed more from the WT than did the *Zcchc6* KO (Fig. 3b), consistent with

more severe RNA accumulation in *Dis3l2* KO than in *Zcchc6* KO. Differential gene

expression analysis also showed that *Dis3l2* KOs had more differentially expressed

transcripts than did the *Zcchc6* KOs (Fig. 3c). Both *Zcchc6* and *Dis3l2* KOs had more

over-represented transcripts (3.0% and 9.6%) than under-represented transcripts (1.3%

and 5.3%). Downregulated transcripts lacked any significant gene ontology

categories, suggesting a lack of target specificity for the population of affected mRNAs.

On the other hand, upregulated genes for each KO demonstrated some gene ontology

term enrichment. "Cytoplasmic translation" and "ribonucleoprotein complex

biogenesis" were the most enriched terms in *Zcchc6* KO, whereas enriched terms for

Dis3l2 KO were "regulation of the mRNA metabolic process", "proteasomal protein

catabolic process", "NIK/NF-kappaB signaling", "generation of precursor metabolites

and energy", and "ribonucleoprotein complex subunit organization" (Extended Data Fig.

10c). When the differentially expressed genes from both KOs were overlaid (Fig. 3d),

we found that most genes (75% of the total) were aligned in the same regulatory

direction, with the majority being accumulated in both KOs. This included the non-

coding RNA *H19*, a known target of Dis3l2, identified in a previous nephron progenitor

study [41]. "Cytoplasmic translation" and "ribonucleoprotein complex regulation" were

gene ontology terms common to both KOs (Fig. 3e). We further validated the top

upregulated and downregulated genes by performing qPCRs on cKO samples (Fig. 3f).

The direction of regulation was consistent, with several genes significantly upregulated

in both cKOs: *Hbb-bh1*, *Ccnd3*, *Gnb2*, *Hspa8*, and *Alad*. The *Hbb-bh1* transcript was

more upregulated in the *Dis3l2* KO than in the *Zcchc6* KO, consistent with our

observation of greater RNA accumulation in the *Dis3l2* KO. *Hbb-bh1* is the

embryonic/fetal globin, which is unusual to observe at our dissection time point of

E18.5, as globin switching has already occurred in wild type mice by this point [42,43].

Upregulation of *γ*-globin in *Zcchc6* and *Dis3l2* KOs in Humanized Mice.

Hbb-bh1 is the ortholog of the human fetal γ-globin (gamma; *HBG1/HBG2*).

Research has established that patients with pathogenic sickle cell mutations in adult β-

globin show improved symptomology following reactivation of fetal γ-globin[25,44]. Thus,

we aimed to test whether *Zcchc6/Dis3l2* deficiency would upregulate the human γ-

globin gene. We crossed our *Zcchc6* and *Dis3l2* KOs with a humanized β-YAC model

mouse harboring the human β-globin gene cluster as a yeast artificial chromosome

transgene [45] (Fig. 4a) and validated the new model through genomic and protein

analysis (Extended Data Fig.11a, b). The human *γ*-globin was significantly

upregulated in erythroid cells at E18.5 in the *Zcchc6* and *Dis3l2* KOs (Fig. 4b). These

humanized β-YAC mice also preserved human globin switching in the mouse, enabling

the profiling of the dynamics of all β-like globins[45]. We examined erythroid cells at an

earlier time point (E14.5 fetal liver) and observed up-regulation of human fetal γ-

globins and down-regulation of adult *β*-globin relative to WT (Fig. 4c). At E18.5, this

pattern of reverse switching persisted in the peripheral blood, albeit to a lesser degree

(Fig. 4d), and was independent of maturation differences between KO and WT mice

(Extended Data Fig. 11 c&d). Together, these results highlight that ZCCHC6/DIS3L2

upregulates human fetal γ-globin in a conserved manner.

Profiling ZCCHC/DIS3L2 during human *in vitro* erythropoiesis.

Following our mouse studies, we adopted a human *in vitro* system for

erythropoiesis that enables a single, synchronized wave of differentiation, as

described[46], to allow detailed molecular studies. We performed *in vitro* erythropoiesis

with human umbilical cord blood-derived CD34+ cells and profiled protein expression

and globin mRNA expression (Extended Data Fig. 12). Flow cytometry profiles of

differentiating erythroid precursors from cord blood mirrored that of native erythroid

ontogeny, with expression of CD71, followed by glycophorin A, and ultimately

incomplete terminal loss of CD71 compared to adult peripheral blood (Extended Data

Fig. 12b). Immunoblotting showed increasing pan-hemoglobin levels through D20,

followed by decreased levels at D23 when some cell lysis occurs. DIS3L2 was highly

expressed starting on D9 and thereafter decreased, while ZCCHC6 was highly

expressed on D12 and D16, and decreased thereafter (Extended Data Fig. 12c).

Expression of mRNA for HBE (*ε*) and HBG1/2 (*γ*) increased until terminal stages when

levels dropped in the setting of cell lysis (Extended Data Fig. 12d). The expression of

HBB (β-globin) increased significantly until D16, after which it declined. Histological view

at D12 and D16 revealed that erythroid cells become extensively hemoglobinized and

exhibit highly condensed or absent nuclei (Extended Data Fig. 12e).

CRISPR KOs during *in vitro* erythropoiesis phenocopy KO mice.

Next, we used CRISPR/Cas9 to generate KOs by introducing ribonucleoprotein

complexes targeting ZCCHC6, DIS3L2, and the AAVS1 locus as a control (Fig. 5a) [47,48].

We conducted genomic analysis at D12 of differentiation, selected samples with over
60% frameshift indels, and confirmed significantly decreased protein levels (Extended
Data Fig. 13a&b). KO of the RNA editing machinery did not impact maturation profiles,
as reflected by flow cytometry for CD71 and GlyA (Extended Data Fig. 13c). However,
we observed significant accumulation of RNA, as reflected by staining with TO (Fig. 5b).
In both ZCCHC6 and DIS3L2 KOs, fetal γ-globin was upregulated while adult β-globin
was relatively downregulated (Fig. 5c). However, the effect size was greater for KO of
ZCCHC6 relative to DIS3L2. Taken together, these data demonstrate RNA
accumulation in CRISPR knockout (KO) cells, which is consistent with the KO mice
study, and a reversed globin-switching pattern in these cells, aligning with the
humanized murine model study.
265

***In vitro* erythroid cells from PB with CRISPR KOs and HU treatment.**
Reactivation or elevation of the expression of human fetal γ-globin by genetic or
pharmacologic means is an effective therapeutic strategy in sickle cell disease and
transfusion-dependent thalassemia [49]. Hydroxyurea (HU) treatment is known to
elevate γ-globin levels in the peripheral blood of patients with hemoglobinopathies [50,51].
While a direct mechanistic connection between HU and ZCCHC6 has not been
described, a previous study revealed that HU-induced DNA replication arrest triggers
the rapid degradation of histone mRNA, mediated by uridylation from ZCCHC6 [52].
In this experiment, we aim to test whether the genetic deficiency of
ZCCHC6/DIS3L2 affects PB (Peripheral Blood) HPSCs, which better reflect a clinical
setting compared to CB HPSCs due to their low baseline levels of fetal hemoglobin. The
PB CD34+ cells were subjected to CRISPR knockout (KO) of the RNA editing
machinery and differentiated after day 9 in the absence or presence of HU (Fig. 6a) [53,54].
Consistently, accumulation of reticulocytes (GlyA+CD71^{low}TO^{high} cells), elevation of γ-
globin levels, and reversed switching patterns were noted in ZCCHC6 CRISPR KOs
(Fig. 6b, c, d), while DIS3L2 CRISPR KOs showed a similar trend with a marginal p-
value for the first two features. Upon DIS3L2 CRISPR KOs, adult β-globin was also
upregulated, resulting in our inability to observe reverse-switching. Genomic analysis at
day 12 demonstrated similar frame-shift editing rates in both PB and CB CD34+ cells.

No differential impact on erythroid maturation profiles were observed upon CRISPR

KOs and HU treatment (Extended Data Fig. 14a & b).

When compared with HU treatment, KO of ZCCHC6 alone showed better

upregulation of γ-globin levels and reverse switching between γ and β globins than HU

treatment (Fig. 6c &d). Furthermore, HU treatment did not augment the impact, likely

because the cells reached the maximum capacity to carry fetal globin. Taken together,

these results highlight that ZCCHC6 is a consistent regulator in both CB and PB-derived

cells. Our data implicating ZCCHC6 as a regulator of γ-globin mRNA stabilization

suggests that ZCCHC6 may be a target worth testing further as a therapeutic strategy

for enhancing γ-globin expression in hemoglobinopathies.

DISCUSSION

During terminal differentiation, erythroid cells dramatically discard cellular

components, including the nucleus, mitochondria, and ribosomes[3–7]. Nuclear

condensation and enucleation entail a loss of transcription[1,3,4], implicating post-

transcriptional regulatory mechanisms as prominent drivers of transcriptome shedding.

The resulting transcriptome, with its low-entropy, globin-rich composition in terminal

erythroid cells, is thought to be achieved by the active decay of RNAs that are

nonessential for mature RBC function[9,10]. However, the exact molecular pathways

driving this process have not been elucidated. In this study, we discovered that the

ZCCHC6-DIS3L2 RNA editing pathway plays a role in RNA shedding during terminal

erythropoiesis, with a prominent target for degradation being fetal (γ-) hemoglobin. Our

finding complements the previous discovery of global proteome shedding through

ubiquitination[11], as both mechanisms involve post-translational and post-transcriptional

tagging, respectively, to target molecules for degradation. Our finding begs the question

of how β-globin mRNA escapes this shedding process. Previous studies have shown

that pyrimidine-rich cis-elements in the 3' UTR, along with associated trans-elements

such as αCP and nucleolin, stabilize β-globin mRNAs [55,56]. This RNP complex may

outcompete ZCCHC6 or form an untaggable substrate for the ZCCHC6-DIS3L2

pathway.

Hereditary Persistence of Fetal Hemoglobin (HPFH), a benign genetic condition

manifesting as incomplete globin switching, mediates significant amelioration of

symptoms when inherited along with the genetic mutation in Sickle Cell Disease

(SCD)[25]. This experiment of nature has inspired a therapeutic strategy to reactivate fetal

γ-globin via genetic perturbation of the γ-globin repressor BCL11A, resulting in

transcriptional reactivation and considerable clinical improvement for SCD

patients[44,49,57,58]. Reactivation of γ-globin by genetic editing has achieved proof of

concept for treating hemoglobinopathies, marking a seminal translational application of

gene editing technology [44,49]. However, it is challenging to deliver genetic therapy for

SCD patients in under-resourced areas of the globe, where SCD is most prevalent [59,60].

This challenge arises from the scarcity of bone marrow transplantation centers and

trained healthcare providers[59,60], and by the high costs of genetic therapies[61]. The

development of small molecule inhibitors that can reactivate γ-globin expression,

including inhibition of BCL11A, remains an appealing alternative that could be more

readily distributed in under-resourced locales[62–66]. However, achieving therapeutic

levels of γ-globin reactivation with a single drug has, to date, proven challenging due to

the suboptimal potency of monotherapy and heterogenous pathological symptoms[65].

The successful treatment may therefore require the use of multiple pharmacological

agents[65].

Here, we report a novel post-transcriptional mechanism whereby γ-globin mRNA

is subject to active degradation. This discovery raises the possibility that therapeutic

strategies to stabilize γ-globin mRNA could synergize with pharmacologic strategies that

reactivate γ-globin mRNA expression to ameliorate the symptoms and deleterious

health consequences of hemoglobinopathies. Recent studies have reported small

molecule activators of γ-globin expression[65,66], as well as inhibitors of ZCCHC6/11

which have been developed as anti-cancer agents[67,68]. In the latter case, inhibitors

reduced the uridylation of target RNAs, including *pre-let-7*[67], and demonstrated

antiproliferative activity in focadhesin (FOCAD)-deleted cancer[68]. Testing existing

ZCCHC6 inhibitors or developing novel inhibitors using assays with γ-globin RNA as a

substrate in erythroid cells could offer an accessible pharmacological option.

While we observed delayed globin switching, prolonged γ-globin mRNA half-life,

reticulocytosis and anemia in the embryonic circulation of *Zcchc6* and *Dis3l2* KO mice,

these phenotypes were attenuated in post-weanling mice, when fetal erythropoiesis is

fully extinguished in favor of adult patterns of red cell development. Resculpting of the

terminal transcriptome during adult murine erythropoiesis most likely entails an additive

or distinct regulatory mechanism. In our *in vitro* studies of human erythropoiesis from

adult peripheral blood, we observed modest stabilization of γ-globin mRNA by ZCCHC6

knock-out, suggesting that the ZCCHC6 machinery is active in adult erythropoiesis.

Strategies to reactivate and stabilize γ-globin in adult red blood cells remains to be

further optimized.

Akin to human β-globins, human α-globin mRNAs are selectively stabilized

during terminal erythropoiesis, by a mechanism that entails binding of the αCP complex

at polypyrimidine-rich sites within the 3' UTR [69]. In $\alpha^{Constant\ Spring\ (CS)}$ thalassemia,

mutation of the 3' UTR disrupts αCP complex formation, leading to accelerated mRNA

decay, characterized by shortened poly(A) tails [70]. Upon *Zcchc6* KO (but not in *Dis3l2*

KO), RNA-seq results show that the embryonic α-like ζ-globin gene (*Hba-x*) was

significantly upregulated in terminal erythroid cells at E18.5. It is therefore likely that ζ-

globin and γ-globin share the RNA editing machinery that regulates their mRNA stability

post-transcriptionally, most notably ZCCHC6, warranting future studies to investigate

the consequences of ZCCHC6 KO in $\alpha^{Constant\ Spring\ (CS)}$ thalassemia.

METHODS

Animal Models

All experiments were performed with previously reported KO models for *Lin28a*

and *Lin28b*, as well as KO and flox lines for *Zcchc6* and *Zcchc11*[13,71,72]. Whole body and

conditional *Dis3l2* KO mice were generated in-house as follows: Targeted mESCs

(Dis3l2[tm1a(EUCOMM)Hmgu]) were obtained from Helmholtz Zentrum München via the

European Conditional Mouse Mutagenesis Program (EUCOMM)[73]. The gene-targeted

ESCs were injected into chimeric blastocysts. Once injected, they were implanted into

pseudopregnant Balb/c females and mutant mice were established. To ascertain

germline transmission, we crossed chimeras to WT mice, screened the offspring, and

obtained several Dis3l2 transgenic animals with germline transmission, after which the

strain was expanded. Constitutive knockout of Dis3l2 was generated by crossing

females carrying *Vasa-Cre* allele (Jackson laboratory, stock #006954) to *Dis3l2*

transgenic males. The cross resulted in the constitutive deletion of Dis3l2 in all tissues

by virtue of *Cre* expression in oocytes[74]. Conditional *Dis3l2* flox line was generated by

crossing *Dis3l2* transgenic mice to Actin-Flippase (Flp) first (B6.Cg-

Tg(ACTFLPe)9205Dym/J, the Jackson Laboratory Strain #005703) to remove Lac-Z

reporter cassette, followed by *Cre* breeding.

The *Zcchc6* and *Dis3l2* flox animals were crossed with previously published

EpoR-Cre mice to generate an erythroid-specific conditional KO mouse [38]. Humanized

beta-globin YAC mice were bred with *Zcchc6/Dis3l2* KO mice [45]. Embryonic and adult

ear tissue genotyping was performed either manually or automatically (using

Transnetyx) to confirm the genotypes of KO, flox, and humanized lines. Conditional

Zcchc6 and *Dis3l2* KOs in the newly generated murine models were validated by

western blots (1:1000 for both antibodies, α-ZCCHC6; Protein Tech 25196-1-AP and α-

DIS3l2; NBP1-84740) on the sorted erythroid cells (PI-Hoechst-TER119+ cells). Animal

experiments were performed by protocols approved by the Institutional Animal Care and

Use Committee at Boston Children's Hospital.

Murine Blood/Tissue Collection and Analysis

Embryonic blood was collected from embryos harvested via Caesarian section.

Blood of postnatal and adult mice was harvested through tail clipping or retro-orbital

bleeding. During time-series analysis across various developmental ages, we bled each

animal only once to avoid reticulocytosis secondary to anemia. Blood was collected

using heparinized capillary tubes into EDTA-coated collection tubes, then subsequently

processed using a HEMAVET machine and Fluorescence-Activated Single Cell Sorting

(FACS) profiling.

One ul of blood was diluted into 50 ul of a buffered saline citrate glucose solution

(BSCG) and mixed with equal parts of staining buffer (PBS + 2% serum + 1X pen/strep)

containing a 1:100 dilution of each antibody: TER119 (BD Pharmingen 557909), CD71

(BD Clone: C2), Thiazole Orange (Sigma-Aldrich-390062), Hoechst 33342 (Invitrogen

R37165), and Propidium Iodide (BD Pharmingen 51-66211E). All acquisition was

performed on an LSR/FORTESSA cytometer, and the results were analyzed using

FlowJo and Prism software. Spleen and liver cells were dissociated in ice-cold PBS,

filtered, and then subjected to FACS analysis using the same staining panel.

For hemolytic marker assays, we collected plasma and serum samples,

incubating the latter at room temperature for 30 minutes to allow for coagulation. We

performed centrifugation at 1,500x g for 10 minutes at 4°C and collected plasma and

serum samples for downstream analysis (western blots and blood biochemistry assays).

The plasma was then mixed and heated in a protein preparation solution (RIPA buffer

with protease and phosphatase inhibitors, and DTT). The protein samples were

loaded onto Bis-Tris gels, and the transferred membranes were probed with primary

antibodies (1:500 and 1:1000 for each antibody respectively: α-Haptoglobin; Life

Technologies # 16665-1-AP and α-IgG; ab190475). LDH (Abcam no. ab102526) and

bilirubin (Sigma Aldrich # MAK126) assays were performed according to the

manufacturers' instructions. The LDH samples were harvested from heparin-coated

collection tubes instead of the EDTA-coated tubes.

Blood Histology

Blood smears were prepared on frosted slides and stained with May-Grünwald

and Giemsa solutions (Sigma-Aldrich; MG500 and GS500). The slides were placed in

the May-Grünwald solution for two minutes, then were subsequently incubated in a

diluted Giemsa solution (1:25 with milliQ H_2O) for 12 minutes and washed with

autoclaved Milli-Q H_2O. Reticulocytes were identified using methylene blue staining

protocol (Sigma-Aldrich; R4132). The stained slides were coated (VECTASHIELD) and

examined under a microscope.

Transcriptomic (RNA-seq) and Transcript Analysis (RT-qPCRs)

Enucleated erythroid cells (PI$^-$Hoechst$^-$TER119$^+$ cells) were sorted directly into

concentrated TRIzol (TRIzol™ LS Reagent). Total RNA was isolated using a column

assay with DNase treatment (Direct-zol MicroPrep, ZYMO). The quantity and quality of

RNA were evaluated using a nanodrop machine and RNA screen tape. We constructed

an RNA-seq library with high-quality RNA (Both 260/280 and 260/230 over 1.7 with

RNA integrity number >7) after ribosomal RNA depletion. For regular gene expression

analysis, adapter-trimmed reads from the sequencer were mapped to the mouse

genome, quantified, and analyzed using seq data analysis tools (Cutadapt, Bowtie,

TopHat, HTSeq, R, and edgeR). For RT-qPCRs, cDNA was reverse transcribed from

the isolated RNA (using the Maxima First Strand cDNA Synthesis Kit, Thermo Fisher

Scientific). Then, quantitative PCR (qPCR) was performed using either SYBR Green or

TaqMan assays (on a QuantStudio machine, Applied Biosystems).

Human *In vitro* Erythropoiesis from Hematopoietic Stem/Progenitor Cells

Hematopoietic stem and progenitor (CD34+ HSPCs) cells derived from umbilical

cord blood or adult peripheral blood were purchased (mixed donor vials,

ALLCELLS/STEMCELL technologies) and subjected to the *in vitro* RBC differentiation

protocol described in the previous study [46]. Genetic KOs were generated in CD34+

HSPCs via nucleofection of CRISPR/Cas9 ribonucleoproteins (RNPs)[48]. First, CD34+

cells were thawed dropwise and cultured for two days before nucleofection in SFEM II

media (StemCell Technologies, 09605) supplemented with 100 ng/mL SCF, TPO,

FLT3L, and IL-6. The CD34+ HSPCs were nucleofected with Cas9 RNPs (125 pmol of

sgRNA complexed with 105 pmol of Cas9 protein) using a Lonza 4D-Nucleofector. Cells

were subsequently cultured for recovery and expansion, followed by the initiation of *in*

vitro differentiation assays, and then harvested for genomic DNA extraction using a lysis

buffer. The gRNA-targeted genomic loci (~200 bp flanking regions of the expected Cas9

cut site) were PCR-amplified, gel-purified, and sanger-sequenced. The frameshifting

Indel frequencies for each gRNA were quantified using TIDE analysis (http://tide.nki.nl).

Hydroxyurea (Sigma) was purchased, freshly resuspended in ultra-pure water each

time, and added to the cell cultures. For downstream qPCR analysis, cells were washed

twice with ice-cold PBS, lysed in TRIzol, and their cDNAs were subsequently subjected

to qPCRs. For Western blot and FACS analysis, we used the same method described in

the murine sample analysis, except for the use of specific human antibodies:

Hemoglobin β/γ/δ/ε Antibody (sc-390668) for western blot, and anti-Human CD71

(Fisher Scientific, BDB555537) and GlyA (CD235a, Beckman Coulter, A71564) for
FACS analysis.
470
Statistical Analysis
Data are expressed as mean ± SD. For hematological profile analysis, adjusted
p-values were calculated from a linear mixed-effects model adjusted for sex (fixed
effect) and litter information (random effect), followed by post-hoc multiple pairwise
comparisons based on Tukey's Honest Significant Difference (HSD) adjustment. For all
other analyses, p-values were calculated using t-tests (* <0.05, ** <0.01, *** <0.001, ****
<0.0001).
478

479   **Data availability**
480   The resulting RNA-seq data is available on the Gene Expression Omnibus (GEO)
481   database (GSE151184).
482

483   **Code availability**
484   All scripts and codes are available upon request.
485

REFERENCES

1.   Dzierzak, E. & Philipsen, S. Erythropoiesis : Development and Differentiation. (2013).
2.   Palis, J. Primitive and definitive erythropoiesis in mammals. *Front Physiol* **5 JAN**, 1–9
(2014).
3.   Keerthivasan, G., Wickrema, A. & Crispino, J. D. Erythroblast enucleation. *Stem Cells Int*
2011, (2011).
4.   Menon, V. & Ghaffari, S. Erythroid enucleation: a gateway into a "bloody" world. *Exp*
Hematol **95**, 13–22 (2021).
5.   Moras, M., Lefevre, S. D. & Ostuni, M. A. From erythroblasts to mature red blood cells:
Organelle clearance in mammals. *Front Physiol* **8**, 1–9 (2017).
6.   Kundu, M. *et al.* Ulk1 plays a critical role in the autophagic clearance of mitochondria and
ribosomes during reticulocyte maturation. *Blood* **112**, 1493–1502 (2008).
7.   Schweers, R. L. *et al.* NIX is required for programmed mitochondrial clearance during
reticulocyte maturation. *Proc Natl Acad Sci U S A* **104**, 19500–19505 (2007).
8.   Ovchynnikova, E., Aglialoro, F., von Lindern, M. & van den Akker, E. The shape shifting
story of reticulocyte maturation. *Front Physiol* **9**, 1–14 (2018).
9.   Bastos, R. N. & Aviv, H. Theoretical analysis of a model for globin messenger RNA
accumulation during erythropoiesis. *J Mol Biol* **110**, 205–218 (1977).

10.   Bastos, R. N., Volloch, Z. & Aviv, H. Messenger RNA population analysis during
erythroid differentiation: A kinetical approach. *J Mol Biol* **110**, 191–203 (1977).

11.   Nguyen, A. T. *et al.* UBE2O remodels the proteome during terminal erythroid
differentiation. *Science* **357**, (2017).

12.   Thomas, M. P. *et al.* Apoptosis Triggers Specific, Rapid, and Global mRNA Decay with
3' Uridylated Intermediates Degraded by DIS3L2. *Cell Rep* **11**, 1079–1089 (2015).

13.   Morgan, M. *et al.* MRNA 3′ uridylation and poly(A) tail length sculpt the mammalian
maternal transcriptome. *Nature* **548**, 347–351 (2017).

14.   Kozlowski, E. *et al.* The RNA uridyltransferase Zcchc6 is expressed in macrophages and
impacts innate immune responses. *PLoS One* **12**, 1–16 (2017).

15.   Ansari, M. Y. *et al.* Genetic Inactivation of ZCCHC6 Suppresses Interleukin-6 Expression
and Reduces the Severity of Experimental Osteoarthritis in Mice. *Arthritis Rheumatol* **71**,
583–593 (2019).

16.   Lin, C. C. *et al.* Terminal uridyltransferase 7 regulates TLR4-triggered inflammation by
controlling Regnase-1 mRNA uridylation and degradation. *Nat Commun* **12**, (2021).

17.   Gupta, A. *et al.* TUT4/7-mediated uridylation of a coronavirus subgenomic RNAs delays
viral replication. *Commun Biol* **6**, 1–9 (2023).

18.   Lee, M., Kim, B. & Kim, V. N. Emerging roles of RNA modification: M6A and U-Tail.
Cell **158**, 980–987 (2014).

19.   Lackey, P. E., Welch, J. D. & Marzluff, W. F. TUT7 catalyzes the uridylation of the 3′
end for rapid degradation of histone mRNA. *Rna* **22**, 1673–1688 (2016).

20.   Le Pen, J. *et al.* Terminal uridylyltransferases target RNA viruses as part of the innate
immune system. *Nat Struct Mol Biol* **25**, 778–786 (2018).

21.   Lim, J. *et al.* Uridylation by TUT4 and TUT7 Marks mRNA for Degradation. *Cell* **159**,
1365–1376 (2014).

22.   Chang, H. *et al.* Terminal Uridylyltransferases Execute Programmed Clearance of
Maternal Transcriptome in Vertebrate Embryos. *Mol Cell* **70**, 72-82.e7 (2018).

23.   Barbarani, G., Labedz, A., Stucchi, S., Abbiati, A. & Ronchi, A. E. Physiological and
Aberrant γ-Globin Transcription During Development. *Front Cell Dev Biol* **9**, 1–10
(2021).

24.   Wienert, B., Martyn, G. E., Funnell, A. P. W., Quinlan, K. G. R. & Crossley, M. Wake-up
Sleepy Gene: Reactivating Fetal Globin for β-Hemoglobinopathies. *Trends in Genetics*
34, 927–940 (2018).

25.   Akinsheye, I. *et al.* Fetal hemoglobin in sickle cell anemia. *Blood* **118**, 19–27 (2011).

26.   Blobel, G. A. & Khandros, E. Elevating fetal hemoglobin - recently discovered regulators
and mechanisms. *Blood Journal* (2024) doi:10.1182/blood.2023022190.

27.   Lu, H. Y., Orkin, S. H. & Sankaran, V. G. Fetal Hemoglobin Regulation in Beta-
Thalassemia. *Hematol Oncol Clin North Am* **37**, 301–312 (2023).

28.   Elagooz, R. *et al.* PUM1 mediates the posttranscriptional regulation of human fetal
hemoglobin. *Blood Adv* **6**, 6016–6022 (2022).

29.   Wakabayashi, A. *et al.* Identification and characterization of RBM12 as a novel regulator
of fetal hemoglobin expression. *Blood Adv* **6**, 5956–5968 (2022).

30.   Faehnle, C. R., Walleshauser, J. & Joshua-Tor, L. Mechanism of Dis3l2 substrate
recognition in the Lin28-let-7 pathway. *Nature* **514**, 252–256 (2014).

31.   Pirouz, M., Du, P., Munafò, M. & Gregory, R. I. Dis3l2-Mediated Decay Is a Quality
Control Pathway for Noncoding RNAs. *Cell Rep* **16**, 1861–1873 (2016).

32. Mosqueira, M., Willmann, G., Zeiger, U. & Khurana, T. S. Expression profiling reveals
novel hypoxic biomarkers in peripheral blood of adult mice exposed to chronic hypoxia.
PLoS One **7**, 1–10 (2012).
33. Palis, J. Ontogeny of erythropoiesis. *Curr Opin Hematol* **15**, 155–161 (2008).
34. Kingsley, P. D. *et al.* Ontogeny of erythroid gene expression. *Blood* **121**, 5–13 (2013).
35. Lee, L. G., Chen, C. H. & Chiu, L. A. Thiazole orange: a new dye for reticulocyte
analysis. *Cytometry* **7**, 508–517 (1986).
36. Malleret, B. *et al.* Plasmodium vivax: Restricted tropism and rapid remodeling of CD71-
positive reticulocytes. *Blood* **125**, 1314–1324 (2015).
37. Yumine, A., Fraser, S. T. & Sugiyama, D. Regulation of the embryonic erythropoietic
niche: A future perspective. *Blood Res* **52**, 10–17 (2017).
38. Heinrich, A. C., Pelanda, R. & Klingmüller, U. A mouse model for visualization and
conditional mutations in the erythroid lineage. *Blood* **104**, 659–666 (2004).
39. Barcellini, W. & Fattizzo, B. Clinical Applications of Hemolytic Markers in the
Differential Diagnosis and Management of Hemolytic Anemia. *Dis Markers* **2015**, (2015).
40. Thornton, J. E., Chang, H.-M., Piskounova, E. & Gregory, R. I. Lin28-mediated control of
let-7 microRNA expression by alternative TUTases Zcchc11 (TUT4) and Zcchc6 (TUT7).
RNA vol. 18 1875–1885 Preprint at https://doi.org/10.1261/rna.034538.112 (2012).
41. Hunter, R. W. *et al.* Loss of Dis3l2 partially phenocopies perlman syndrome in mice and
results in upregulation of Igf2 in nephron progenitor cells. *Genes Dev* **32**, 903–908 (2018).
42. Sankaran, V. G., Xu, J. & Orkin, S. H. Advances in the understanding of haemoglobin
switching: Review. *Br J Haematol* **149**, 181–194 (2010).
43. McGrath, K. E. *et al.* A transient definitive erythroid lineage with unique regulation of the
β-globin locus in the mammalian embryo. *Blood* **117**, 4600–4608 (2011).
44. Orkin, S. H. & Bauer, D. E. Emerging Genetic Therapy for Sickle Cell Disease. *Annu Rev*
Med **70**, 257–271 (2019).
45. Porcu, S. *et al.* The human β globin locus introduced by YAC transfer exhibits a specific
and reproducible pattern of developmental regulation in transgenic mice. *Blood* **90**, 4602–
4609 (1997).
46. Huang, N. J. *et al.* Genetically engineered red cells expressing single domain camelid
antibodies confer long-Term protection against botulinum neurotoxin. *Nat Commun* **8**, 1–
13 (2017).
47. Bak, R. O., Dever, D. P. & Porteus, M. H. CRISPR/Cas9 genome editing in human
hematopoietic stem cells. *Nat Protoc* **13**, 358–376 (2018).
48. Sanson, K. R. *et al.* Optimized libraries for CRISPR-Cas9 genetic screens with multiple
modalities. *Nat Commun* **9**, 1–15 (2018).
49. Daley, G. Q. Welcoming the Era of Gene Editing in Medicine. *N Engl J Med* **390**, 1642–
1645 (2024).
50. Rodgers, G. P., Dover, G. J., Noguchi, C. T., Schechter, A. N. & Nienhuis, A. W.
Hematologic responses of patients with sickle cell disease to treatment with hydroxyurea.
N Engl J Med **322**, 1037–1045 (1990).
51. Platt, O. S. *et al.* Hydroxyurea enhances fetal hemoglobin production in sickle cell
anemia. *J Clin Invest* **74**, 652–656 (1984).
52. Holmquist, C. E., He, W., Meganck, R. M. & Marzluff, W. F. Erratum: Knockouts of
TUT7 and 3′hExo show that they cooperate in histone mRNA maintenance and

degradation (RNA (2022) 28 (1519–1533) DOI: 10.1261/rna.079233.122). *Rna* **29**, 862
(2023).

53. Zhu, X. *et al.* Hydroxyurea differentially modulates activator and repressors of γ-globin
gene in erythroblasts of responsive and non-responsive patients with sickle cell disease in
correlation with Index of Hydroxyurea Responsiveness. *Haematologica* **102**, 1995–2004
(2017).

54. Cokic, V. P. *et al.* Hydroxyurea induces fetal hemoglobin by the nitric oxide–dependent
activation of soluble guanylyl cyclase. *Journal of Clinical Investigation* **111**, 231–239
(2003).

55. Jiang, Y., Xu, X.-S. & Russell, J. E. A Nucleolin-Binding 3′ Untranslated Region Element
Stabilizes β-Globin mRNA In Vivo. *Mol Cell Biol* **26**, 2419–2429 (2006).

56. Yu, J. & Russell, J. E. Structural and Functional Analysis of an mRNP Complex That
Mediates the High Stability of Human β-Globin mRNA. *Mol Cell Biol* **21**, 5879–5888
(2001).

57. Xu, J. *et al.* Correction of sickle cell disease in adult mice by interference with fetal
hemoglobin silencing. *Science (1979)* **334**, 993–996 (2011).

58. Sankaran, V. G. *et al.* Developmental and species-divergent globin switching are driven
by BCL11A. *Nature* **460**, 1093–1097 (2009).

59. Kumar, A. & Bhattacharya, S. Sickle cell disease: a comparative perspective on global
and national initiatives. *Frontiers in Hematology* **3**, 1–6 (2024).

60. John, T. D., Namazzi, R., Chirande, L. & Tubman, V. N. Global perspectives on cellular
therapy for children with sickle cell disease. *Curr Opin Hematol* **29**, 275–280 (2022).

61. Basu, A. *et al.* Sickle Cell Disease in the United States : **177**, 155–164 (2024).

62. Zheng, G. *et al.* A tetramer of BCL11A is required for stable protein production and fetal
hemoglobin silencing. *Science* **386**, 1010–1018 (2024).

63. Saunthararajah, Y. Targeting sickle cell disease root-cause pathophysiology with small
molecules. *Haematologica* **104**, 1720–1730 (2019).

64. Takase, S. *et al.* A specific G9a inhibitor unveils BGLT3 lncRNA as a universal mediator
of chemically induced fetal globin gene expression. *Nat Commun* **14**, 1–18 (2023).

65. Telen, M. J., Malik, P. & Vercellotti, G. M. Therapeutic strategies for sickle cell disease:
towards a multi-agent approach. *Nat Rev Drug Discov* **18**, 139–158 (2019).

66. Ting, P. Y. *et al.* A molecular glue degrader of the WIZ transcription factor for fetal
hemoglobin induction. *Science* **385**, 91–99 (2024).

67. Lin, S. & Gregory, R. I. Identification of small molecule inhibitors of Zcchc11 TUTase
activity. *RNA Biol* **12**, 792–800 (2015).

68. Triboulet, R. *et al.* Targeting the synthetic lethal relationship between FOCAD and TUT7
represents a potential therapeutic opportunity for TUT4/7 small molecule inhibitors in
cancer. *Mol Cancer Ther* 1779–1788 (2024) doi:10.1158/1535-7163.mct-24-0176.

69. Russell, J. E., Morales, J. & Liebhaber, S. A. *The Role of MRNA Stability in the Control of*
Globin Gene Expression. Progress in nucleic acid research and molecular biology vol. 57
(Elsevier Masson SAS, 1997).

70. Morales, J., Russell, J. E. & Liebhaber, S. A. Destabilization of human α-globin mRNA
by translation anti-termination is controlled during erythroid differentiation and is
paralleled by phased shortening of the poly(A) tail. *Journal of Biological Chemistry* **272**,
6607–6613 (1997).

71.   Osborne, J. K. *et al.* Lin28 paralogs regulate lung branching morphogenesis. *Cell Rep* **36**,
641   109408 (2021).
642   72.   Yermalovich, A. V. *et al.* Lin28 and let-7 regulate the timing of cessation of murine
643   nephrogenesis. *Nat Commun* **10**, (2019).
644   73.   Skarnes, W. C. *et al.* A conditional knockout resource for the genome-wide study of
645   mouse gene function. *Nature* **474**, 337–344 (2011).
646   74.   Gallardo, T., Shirley, L., John, G. B. & Castrillon, D. H. Generation of a germ cell-
647   specific mouse transgenic Cre line, Vasa-Cre. *Genesis (United States)* **45**, 413–417
648   (2007).
649   75.   Kingsley, P. D. *et al.* Ontogeny of erythroid gene expression. *Blood* **121**, 5–13 (2013).
650

Acknowledgements

The authors would like to thank Dr. Ursula Klingmüller for granting permission to use the *EpoR-cre* mouse model; Dr. Susanna Porcu for permission to use the Human *β-Globin YAC* mouse model; Dr. Dónal O'Carroll for sharing the *Zcchc6* and *Zcchc11* whole-body and conditional knockout mouse models; Dr. Vijay Sankaran for providing the globin primer lists and engaging in scientific discussions; current and former members of the Daley, Thorsten, and North labs for valuable discussions and technical assistance; the Harvard Medical School Biopolymers Facility for sequencing; and the Research Computing Group for providing the O2 High-Performance Compute Cluster. This work was supported by PCTC (Progenitor Cell Translational Consortium; NHLBI UO1-HL134812) and T32 grants from the National Heart, Lung, and Blood Institute (NHLBI), U54DK110805: A Center of Molecular Hematology from National Institute of Diabetes and Digestive and Kidney Diseases (NIDDK), as well as an early career grant from the National Blood Foundation.

Author contributions

A.H. conceived the research question, performed experiments, and wrote the manuscript. A.Y. designed and conducted the Dis3l2 mice experiments. M.A.T.N. performed CRISPR/Cas9 knockouts and analyzed genome editing data. D.P. designed and established the ZCCHC6/11 mice study. Y.F. performed the blastocyst injections for the Dis3l2 study and guided the EpoR-cre and YAC mice studies. M.B. conducted mouse genotyping and RNA experiments. C.K. designed and provided guidance on FACS profiling. T.N. advised on the study. V.L. set up the human *in vitro* erythropoiesis

674   study and revised the manuscript. S.O. designed the EpoR-cre and YAC mice studies

675   and advised on the study. G.Q.D. conceived the idea, supervised the study, and wrote

676   the manuscript.

a

Mouse Embryo Dissection → Blood/Tissue Collection → Analysis

Zcchc6 WT vs. KO
Dis3l2 WT vs. KO

- HISTOLOGY
- FACS
- HEMOGRAM

b

	Zcchc6	*Dis3l2*
WT		
KO		

c E18.5 PB TER119+ cells

TO / CD71

CD71^lowTO^high / CD71^low (%)

d

p-values (<0.05 *, <0.01 **, <0.001***)

Parameter	Unit	*Zcchc6*			p-value		*Dis3l2*			p-value	
		WT	HET	KO	WT vs. KO	HET vs. KO	WT	HET	KO	WT vs. KO	HET vs.KO
RBC	M/µl	3.06 ± 0.26	3.05 ± 0.14	2.73 ± 0.30	**	**	3.38 ± 0.34	3.49 ± 0.32	2.44 ± 0.34	***	***
HCT	%	44.18 ± 4.21	44.14 ± 3.46	40.00 ± 3.59	*	*	42.88 ± 4.17	44.31 ± 4.24	34.52 ± 2.93	***	***
MCH	pg	33.61 ± 1.87	33.68 ± 2.20	36.30 ± 2.09	**	**	32.42 ± 1.46	32.78 ± 1.57	35.25 ± 1.72	**	**
MCHC	g/dL	23.25 ± 0.98	23.30 ± 1.21	24.75 ± 1.31		*	25.62 ± 2.00	25.81 ± 0.92	25.62 ± 1.05		
MCV	fL	144.55 ± 4.43	144.62 ± 6.51	146.90 ± 9.88			126.98 ± 5.74	127.10 ± 6.46	137.85 ± 7.7	**	***
RDW	%	19.56 ± 0.53	19.06 ± 0.66	19.89 ± 0.32	*	***	17.82 ± 0.33	17.91 ± 0.81	18.82 ± 0.79		**
Hb	g/dL	10.26 ± 0.95	10.28 ± 0.91	9.90 ± 1.05			10.96 ± 1.15	11.44 ± 1.23	8.65 ± 1.02	***	***

Fig. 1. Analysis of Whole-Body *Zcchc6/Dis3l2* Knockout Mice.
(a) The overall scheme of the embryonic blood studies on *Zcchc6* and *Dis3l2* knockout (KO) mice.
(b) Histology of peripheral blood smears stained by May-Grünwald-Giemsa (left) and methylene blue (right).
Scale bar = *10um*.
(c, left figure) Flow cytometry profiles of erythroid cells from peripheral blood (PB).
Representative profiles of each KO and littermate WT were displayed.
(c, right graphs) Quantification of reticulocyte population per genotype (mean ± SD). Multiple litters were used.
(d) Hematological profiles of the peripheral blood (mean ± SD).
HCT; Hematocrit, MCH; Mean corpuscular hemoglobin, MCHC; Mean corpuscular hemoglobin concentration,
MCV; Mean corpuscular volume, RDW; Red cell Distribution Width.

b

E18.5 PB TER119+ cells

c

p-values (<0.05 *, <0.01 **, <0.001***)

Parameter	Unit	Zcchc6			p-value		Dis3l2			p-value	
		EpoR-Cre	cWT	cKO	cWT vs. cKO	EpoR-Cre vs. cKO	EpoR-Cre	cWT	cKO	cWT vs. cKO	EpoR-Cre vs. cKO
RBC	M/µl	3.43 ± 0.22	3.45 ± 0.10	3.19 ± 0.14	*	*	3.70 ± 0.56	3.64 ± 0.54	3.49 ± 0.20	*	*
HCT	%	52.00 ± 1.66	51.93 ± 2.03	48.13 ± 4.21			49.00 ± 10.17	48.86 ± 7.66	47.29 ± 4.76		*
MCH	pg	34.28 ± 0.85	34.18 ± 1.98	36.52 ± 1.04	*	*	33.08 ± 2.13	32.71 ± 4.25	33.64 ± 3.09		
MCHC	g/dL	22.58 ± 0.82	22.68 ± 1.31	24.20 ± 0.75			25.13 ± 0.33	24.48 ± 3.68	24.86 ± 2.13		
MCV	fL	151.78 ± 6.82	150.63 ± 4.08	150.85 ± 8.27			131.73 ± 9.68	134.08 ± 8.55	135.50 ± 8.15		
RDW	%	19.38 ± 1.13	18.73 ± 0.43	19.7 ± 0.76			19.58 ± 1.88	19.50 ± 0.79	19.49 ± 1.32		
Hb	g/dL	11.75 ± 0.47	11.78 ± 0.36	11.63 ± 0.77			12.30 ± 2.47	11.74 ± 1.16	11.74 ± 1.42		

Fig. 2. Analysis of Conditional *Zcchc6/Dis3l2* Knockout Mice.
(a) Breeding scheme to generate erythroid-specific, conditional KO mice.
(b, left figures) Representative FACS profiles from peripheral blood of embryos
  carrying only the *EpoR-Cre* allele (top lane), only the flox allele (cWT, middle lane),
  and both alleles; erythroid-specific conditional KOs (cKO, bottom lane).
(b, right graphs) Quantification of reticulocyte (CD71^{low}TO^{high}) population per genotype.
(c) Hematological profile results from the PB of conditional KOs (erythrocyte panel).

Fig. 3. RNA Profiling.

(a) RNA-seq experimental scheme from embryonic erythroid cells.
(b) Principal component analysis of global transcriptome from *Zcchc6* and *Dis3l2* KOs and corresponding WTs.
(c) Differentially expressed genes in each KO model.
(d) Overlap analysis of differentially expressed genes in each KO model.
(e) Gene ontology analysis of up-regulated genes from both KOs.
(f) Validation of top hits using qPCRs with samples from cKOs.

Fig. 4. Humanized Murine Model Study.
(a) Breeding scheme for crossing *Zcchc6/Dis3l2* KO mice with humanized globin YAC mice.
(b) Fold change in normalized *γ-globin* expression (Ct values for *γ-globin* transcripts were first normalized to those of *r18S* transcripts, and the fold change was then calculated relative to the value of littermate WTs).
(c-d) Human *β-like* globin profiles on sorted erythroid cells from E14.5 fetal liver and E18.5 PB of humanized mouse embryos using qPCRs. For each embryonic age and mouse model, embryos from two litters were used.

a

b

c

Fig. 5. Human *In Vitro* Erythropoiesis Study Using CRISPR Knockout.
(a) Scheme of the CRISPR/Cas9 knockout followed by *in vitro* erythropoiesis.
(b) Representative profiles of TO and CD71 within erythroid (GlyA+) cells and the corresponding graph
with statistical assessment.
(c) Globin profiles upon CRISPR/Cas9 knockouts.

Fig. 6. *In Vitro* Erythropoiesis Study with CRISPR KO and HU(Hydroxyurea) Treatment.
(a) Scheme of the CRISPR/Cas9 knockout and HU treatment followed by *in vitro* erythropoiesis.
(b) Summary graph and statistical assessment of CD71^low^TO^high^ population upon CRISPR/Cas9 KOs.
(c) Human *γ-globin* mRNA expression upon CRISPR/Cas9 knockouts and HU treatment.
(d) Globin profiles upon CRISPR/Cas9 KOs and HU treatment.

Extended Data Fig. 1

Zcchc6

P : **P**roerythroblast
B : **B**asophilic Erythroblast
O : Poly**o**rthochromatic Erythroblast
R : **R**eticulocyte

Dis3l2

Extended Data Fig. 1.
Expression Profiles of Zcchc6 and Dis3l2 mRNAs During Erythropoiesis (from the Erythron Database).
Expression Levels of Zcchc6 and Dis3l2 mRNAs During Mouse Primitive and Definitive
(fetal liver and adult bone marrow) Erythropoiesis.

Extended Data Fig. 2

a

b

Extended Data Fig. 2. Allele Map of the Dis3l2 Knockout Model and Conditional Knockout Validation.
(a) Allele Map of the *Dis3l2* Knockout Models.
(b) Western Blots of Erythroid Cells from Embryos Carrying only *floxed* (cWT)
and both *floxed* and *EpoR-Cre* (cKO) Alleles for Validation.

Extended Data Fig. 3

a

Parameter	Unit	Zcchc6			p-value	
		WT	HET	KO	WT vs. KO	HET vs. KO
RBC	M/µl	9.09 ± 0.39	9.03 ± 0.29	9.05 ± 0.39		
HCT	%	43.28 ± 1.52	43.35 ± 1.71	43.99 ± 1.66		
MCH	pg	14.52 ± 0.73	14.66 ± 0.69	15.15 ± 0.76		
MCHC	g/dL	30.48 ± 0.96	30.55 ± 1.17	31.15 ± 1.24		
MCV	fL	47.65 ± 1.50	48.04 ± 1.44	48.61 ± 1.26		
RDW	%	16.49 ± 0.44	16.74 ± 0.60	16.60 ± 0.47		
Hb	g/dL	13.18 ± 0.57	13.23 ± 0.54	13.70 ± 0.79		

b

Parameter	Unit	Zcchc6			p-value	
		WT	HET	KO	WT vs. KO	HET vs.KO
Leukocytes:						
WBC	K/µl	9.65 ± 1.25	9.56 ± 1.57	10.29 ± 1.41		
LY	K/µl	6.15 ± 0.89	6.17 ± 1.29	6.38 ± 1.04		
NE	K/µl	2.86 ± 0.45	2.67 ± 0.48	3.13 ± 0.52		*
MO	K/µl	0.49 ± 0.20	0.64 ± 0.18	0.67 ± 0.15		
EO	K/µl	0.13 ± 0.08	0.07 ± 0.04	0.09 ± 0.05		
BA	K/µl	0.02 ± 0.01	0.01 ± 0.01	0.01 ± 0.00		
LY	%	63.71 ± 3.99	64.20 ± 5.30	61.88 ± 4.04		
NE	%	29.74 ± 3.29	28.25 ± 4.55	30.54 ± 3.55		*
MO	%	4.96 ± 1.50	6.66 ± 1.50	6.63 ± 1.77		
EO	%	1.40 ± 0.85	0.78 ± 0.50	0.86 ± 0.38		
BA	%	0.19 ± 0.14	0.12 ± 0.10	0.10 ± 0.04		
Thrombocytes:						
PLT	K/µl	870.15 ± 83.11	923.91 ± 120.69	884.38 ± 125.59		
MPV	fL	4.60 ± 0.12	4.63 ± 0.14	4.66 ± 0.15		

c

Extended Data Fig. 3. Analysis of Whole-Body *Zcchc6* Knockout Young Adult Mice (6–7 Weeks).
(a) Hematological Profiles of Peripheral Blood from Retro-Orbital Bleeding (mean ± SD).
  HCT; Hematocrit, MCH; Mean corpuscular hemoglobin, MCHC; Mean corpuscular hemoglobin concentration,
  MCV; Mean corpuscular volume, RDW; Red cell Distribution Width. Multiple litters were used.
(b) Leukocytes and Thrombocyte panels.
  LY; lymphocytes, NE; neutrophils, MO; monocytes, EO; eosinophils, BA; basophils,
  PLT; Platelet or thrombocyte count, MPV; Mean Platelet Volume.
(c) Quantification of reticulocyte (TO+) population per genotype (mean ± SD).

Extended Data Fig. 4

a

E18.5 Spleen TER119+ cells

b

Parameter	Unit	Zcchc6			p-value		Dis3l2			p-value	
		WT	HET	KO	WT vs. KO	HET vs.KO	WT	HET	KO	WT vs. KO	HET vs. KO
Leukocytes:											
WBC	K/µl	7.10 ± 1.39	8.96 ± 3.72	21.13 ± 8.52	***	***	3.79 ± 1.22	4.42 ± 1.08	5.53 ± 1.36	*	
LY	K/µl	5.85 ± 1.25	7.60 ± 3.50	18.61 ± 7.52	***	***	1.65 ± 0.62	1.96 ± 0.49	2.77 ± 1.10	*	*
NE	K/µl	0.96 ± 0.14	0.98 ± 0.31	1.66 ± 0.74	**	**	1.58 ± 0.57	1.82 ± 0.57	2.00 ± 0.73		
MO	K/µl	0.17 ± 0.08	0.22 ± 0.07	0.71 ± 0.29	***	***	0.12 ± 0.10	0.12 ± 0.09	0.15 ± 0.09		
EO	K/µl	0.10 ± 0.04	0.11 ± 0.05	0.12 ± 0.04			0.34 ± 0.11	0.43 ± 0.18	0.46 ± 0.10		
BA	K/µl	0.04 ± 0.01	0.04 ± 0.02	0.05 ± 0.03			0.10 ± 0.05	0.10 ± 0.05	0.16 ± 0.09		*
LY	%	82.10 ± 2.33	83.63 ± 5.61	87.87 ± 1.61			43.52 ± 9.67	44.52 ± 7.34	49.51 ± 11.79		
NE	%	13.75 ± 2.22	11.93 ± 4.92	7.92 ± 1.44	*		41.88 ± 9.86	40.86 ± 6.63	36.35 ± 9.34		
MO	%	2.24 ± 0.92	2.60 ± 0.63	3.38 ± 0.63	*		2.73 ± 1.84	2.75 ± 2.05	2.51 ± 1.23		
EO	%	1.37 ± 0.44	1.37 ± 0.61	0.60 ± 0.21		*	9.37 ± 1.14	9.64 ± 3.13	8.58 ± 2.26		
BA	%	0.54 ± 0.06	0.47 ± 0.19	0.22 ± 0.09	**	**	2.51 ± 1.15	2.23 ± 1.16	3.05 ± 1.26		
Thrombocytes:											
PLT	K/µl	496.75 ± 49.31	517.67 ± 35.27	519.13 ± 24.93			425.60 ± 71.20	464.71 ± 86.20	411.64 ± 66.27		
MPV	fL	6.31 ± 0.17	6.42 ± 0.15	6.50 ± 0.11			6.32 ± 0.18	6.10 ± 0.27	6.13 ± 0.22		

Extended Data Fig. 4. Additional Analysis of Whole-Body *Zcchc6/Dis3l2* Knockout Mice.
(a, left figures) Flow cytometry profiles of erythroid cells from spleens.
Representative profiles of each KO mouse and WT littermates were displayed.
(a, right graphs) Quantification of reticulocyte population per genotype.
(b) Hematological profiles in the PB of whole-body KOs (Leukocytes and Thrombocyte panels).
LY; lymphocytes, NE; neutrophils, MO; monocytes, EO; eosinophils, BA; basophils,
PLT; Platelet or thrombocyte count, MPV; Mean Platelet Volume.

Extended Data Fig. 5

a

E18.5 Spleen TER119+ cells

b

p-values (<0.05 *, <0.01 **, <0.001***)

Parameter	Unit	Zcchc6 EpoR-Cre	cWT	cKO	p-value cWT vs. cKO	EpoR-Cre vs. cKO	Dis3l2 EpoR-Cre	cWT	cKO	p-value cWT vs. cKO	EpoR-Cre vs. cKO
Leukocytes:											
WBC	K/µl	15.60 ± 5.12	16.54 ± 6.04	30.89 ± 11.89	0.0686	*	17.14 ± 15.10	13.37 ± 7.68	26.49 ± 16.83	***	*
LY	K/µl	14.28 ± 5.13	15.24 ± 5.72	27.93 ± 11.59	0.0913	*	15.49 ± 14.10	12.25 ± 7.40	24.19 ± 15.79	***	*
NE	K/µl	0.92 ± 0.21	0.86 ± 0.20	1.86 ± 0.89			1.18 ± 0.49	0.88 ± 0.25	1.49 ± 0.52	**	
MO	K/µl	0.32 ± 0.13	0.36 ± 0.22	0.91 ± 0.43		*	0.40 ± 0.55	0.17 ± 0.17	0.73 ± 0.66	**	
EO	K/µl	0.06 ± 0.03	0.06 ± 0.03	0.16 ± 0.08			0.06 ± 0.04	0.06 ± 0.03	0.04 ± 0.02		
BA	K/µl	0.03 ± 0.01	0.02 ± 0.01	0.03 ± 0.01			0.02 ± 0.01	0.02 ± 0.01	0.02 ± 0.01		
LY	%	90.50 ± 4.55	91.88 ± 1.40	89.39 ± 4.33			87.57 ± 7.37	88.59 ± 7.74	90.05 ± 3.11		
NE	%	6.85 ± 4.25	5.58 ± 1.66	7.21 ± 4.31			10.20 ± 7.96	9.72 ± 7.94	7.41 ± 3.66	*	
MO	%	1.99 ± 0.27	2.02 ± 0.70	2.77 ± 1.08			1.74 ± 0.93	0.94 ± 0.64	2.21 ± 1.09	***	
EO	%	0.47 ± 0.38	0.36 ± 0.20	0.51 ± 0.13			0.38 ± 0.18	0.56 ± 0.34	0.21 ± 0.10	*	
BA	%	0.20 ± 0.15	0.17 ± 0.11	0.11 ± 0.08			0.12 ± 0.06	0.17 ± 0.11	0.11 ± 0.11		
Thrombocytes:											
PLT	K/µl	590.75 ± 38.23	579.50 ± 16.09	609.67 ± 48.13			610.5 ± 133.57	609.11 ± 94.52	569.50 ± 57.43		
MPV	fL	6.00 ± 0.14	6.10 ± 0.18	6.02 ± 0.13			5.98 ± 0.15	6.12 ± 0.16	6.06 ± 0.28		

Extended Data Fig. 5. Additional Analysis of Conditional *Zcchc6/Dis3l2* KO.
(a) Flow cytometry profiles of erythroid cells from the spleen.
Representative profiles and reticulocyte quantification of embryos carrying
only the *EpoR-Cre* allele, only the *floxed* allele; cWTs, and both alleles; cKOs.
(b) Hematological profiles in the PB of conditional KOs. (Leukocytes and Thrombocyte panels).
LY; lymphocytes, NE; neutrophils, MO; monocytes, EO; eosinophils, BA; basophils,
PLT; Platelet or thrombocyte count, MPV; Mean Platelet Volume.

Extended Data Fig. 6

a

b

Extended Data Fig. 6. Analysis of Hemolytic Anemia Markers.
(a) Western blots of Haptoglobin in cKOs and cWTs.
(b) LDH and Bilirubin assay results from serum and plasma of cKO mice.

Extended Data Fig. 7

a

b

E18.5 PB TER119+

Lin28a

Lin28b

Zcchc11

Extended Data Fig. 7. Characterizing the Impact of Knockouts in Pathway-Related Genes.
(a) Previously published model for the RNA decay machinery (LIN28-ZCCHC6/11-DIS3L2)
on the microRNA regulation.
(b) FACS profiles and population quantification for each whole-body KO model
(*Lin28a*, *Lin28b*, and *Zcchc11* sequentially).

Extended Data Fig. 8

Extended Data Fig. 8. Analysis of *Quad-floxed* Conditional Knockouts.
(a) *Quad-floxed* conditional KO breeding scheme for *Zcchc6* and its paralog *Zcchc11*.
(b) Representative FACS profiles of the *quad-floxed* embryos and quantification of reticulocytes.

Extended Data Fig. 9

Extended Data Fig. 9. Postnatal Profiling of *Zcchc6* KO and *Dis3l2* cKO.
(a) Postnatal PB FACS profiling and quantification of whole body *Zcchc6* KO model
(b) and *Dis3l2* cKO model.

Extended Data Fig. 10

a

Genotype	Litter	Total Read No.	Mapped Read No.	Mapping Rate
Zcchc6 WT	L3	78,543,026	36,284,397	46.2%
Zcchc6 WT	L4	90,445,954	51,073,608	56.5%
Zcchc6 WT	L5	59,412,646	30,752,512	51.8%
Zcchc6 KO	L3	81,304,604	37,445,722	46.1%
Zcchc6 KO	L4	79,495,636	44,989,077	56.6%
Zcchc6 KO	L5	74,278,610	37,835,979	50.9%
Dis3l2 WT	L1	68,767,710	36,391,345	52.9%
Dis3l2 WT	L2	72,958,170	35,684,609	48.9%
Dis3l2 WT	L2	64,899,346	32,206,562	49.6%
Dis3l2 KO	L1	76,216,868	36,395,571	47.8%
Dis3l2 KO	L2	70,791,404	27,438,581	38.8%
Dis3l2 KO	L2	75,760,250	35,673,350	47.1%

b

c

Enriched Gene Ontology (Biological Process)

Up-regulated upon *Zcchc6* KO

Up-regulated upon *Dis3l2* KO

Extended Data Fig. 10. RNA-seq Data Analysis.
(a) Depth and mapping rates of bulk RNA-seq experiments.
(b) Hierarchical clustering of samples based on the expression of *Zcchc6* and *Dis3l2*.
(c) Enriched gene ontology terms and their statistical significances for each KO model.

Extended Data Fig. 11

Extended Data Fig. 11. Analysis of Humanized Mice Upon *Zcchc6/Dis3l2* KO.
(a) Genomic PCR analysis of the human β-YAC allele in offspring.
(b) Validation of preserved *Zcchc6/Dis3l2* KO in the crossed humanized mice using western blots.
(c & d) FACS profiles of erythroid maturation from E14.5 fetal liver and E18.5 PB of subject mice.

Extended Data Fig. 12

Extended Data Fig. 12. Human *In Vitro* Erythropoiesis

(a) Scheme of *in vitro* erythropoiesis from human cord blood-derived hematopoietic stem/progenitor cells
   (HSPCs; CD34+ cells).

(b) Time-series FACS profiles to follow erythroid maturation status
   (Adult human PB served as a positive control).

(c) Time-series protein expression profiles of ZCCHC6, DIS3L2, and hemoglobin
   during the *in vitro* erythropoiesis by western blots.

(d) RNA expression profiles of β-like globins using qPCRs during the in vitro erythropoiesis.

(e) Histology of cytospined cells (stained by May-Grünwald Giemsa method).

Extended Data Fig. 13

Extended Data Fig. 13. Analysis of CRISPR KO in Differentiated Erythroid Cells from CB.
(a) Genomic editing and **(b)** Protein profiles to access efficiency of CRISPR/Cas9-mediated KOs.
(c) Summary graph of erythroid maturation status by FACS (CD71 and GlyA) in CRISPR/Cas9 KOs.

Extended Data Fig. 14

a

b

Extended Data Fig. 14.
Analysis of CRISPR KO in Differentiated Erythroid Cells from PB Following HU treatment.
Summary graph of erythroid maturation status by FACS (CD71 and GlyA) in **(a)** CRISPR/Cas9 KOs
(b) CRISPR/Cas9 KOs & HU treatment

SUPPLEMENTAL TABLES

Supplemental Table 1

qPCR Primers for RNA-seq Validation

Category	Gene	Source	Forward primer seq	Reverse primer seq	Reference
Up	*Hbb-bh1*	Literature			McGrath *et al*. (PubMed ID:21378272)
Up	*Zfpm1*	Commercial			OriGene;MP219016
Up	*Anapc5*	Commercial			OriGene;MP200648
Up	*Pcbp2*	Commercial			OriGene;MP221925
Up	*H19*	Literature	AAGAGCTCGGACTGGAGACT	GACCACACCTGTCATCCTCG	Giovarelli *et al*. (PubMed ID:25385579)
Up	*Ahcyl1*	Commercial			OriGene;MP214966
Up	*Ccnd3*	Commercial			OriGene;MP221497
Up	*Gnb2*	Commercial			OriGene;MP205834
Up	*Hspa8*	Commercial			OriGene;MP205962
Up	*Ddx3x*	Literature	TGGAAATAGTCGCTGGTGTG	GGAGGACAGTTGTTGCCTGT	Lennox *et al*. (PubMed ID:32135084)
Up	*Donson*	Commercial			OriGene;MP203922
Up	*Slc38a5*	Commercial			OriGene;MP215857
Up	*Alad*	Commercial			OriGene;MP200605
Down	*Fam117a*	Commercial			OriGene;MP214105
Down	*Ell2*	Literature	CGCTGGAGACTTACCAGAGC	CATTGAAGGGATCATTTTTGG	Pascal *et al*. (PubMed ID:28870994)
Down	*Dnmt1*	Literature	CCCATGCATAGGTTCACTTCCTTC	TGGCTTCGTCGTAACTCTCTACCT	Noh *et al*. (PubMed ID:15671176)
Down	*Ube2o*	Commercial			OriGene;MP217952

Supplemental Table 2

qPCR Primers for β-like globin profiling

Description	Gene	Forward primer seq	Reverse primer seq
Mus musculus *β* globin	*εy*	TGGCCTGTGGAGTAAGGTCAA	GAAGCAGAGGACAAGTTCCCA
	βh1	TGGACAACCTCAAGGAGACC	ACCTCTGGGGTGAATTCCTT
	βmajor/minor	TTTAACGATGGCCTGAATCACTT	CAGCACAATCACGATCATATTGC
Homo sapiens *β* globin	*ε (HBE)*	GCAAGAAGGTGCTGACTTCC	ACCATCACGTTACCCAGGAG
	γ (HBG)	TGGATGATCTCAAGGGCAC	TCAGTGGTATCTGGAGGACA
	β (HBB)	CTGAGGAGAAGTCTGCCGTTA	AGCATCAGGAGTGGACAGAT

Supplemental Table 3

Guide RNA (gRNA) Sequences for CRISPR/Cas9 Knockout (KO)

Name	sgRNA sequence
gAAVS1	GGACGCACCATTCTCACAAA
gDIS3L2 No.1	TTCCCCAGAGCATAGCAGCG
gDIS3L2 No.2	AATACTAACAGAGTATGGCG
gZCCHC6 No.1	TTGACTGTTGATCAGAAACG
gZCCHC6 No.2	TGAGTCTGTACTTTCAACGA