**Title:** CIT tumor lines: A novel series of immunogenic squamous cell skin carcinoma cell lines

derived from chemical carcinogenesis

**Authors**

Miho Tanaka[1,*], Robert Letchworth[2,3,*], Alina A Barnes[2,3], Lotus Lum[1], Savannah Hughes[2], Grant

Schlauderaff[2], Piyush Chaudhary[2,3], Kenneth M Ng[1], Daphne Superville[1], Corinna Martinez

Luna[1], Maria Gonzalez[1], Allie Grossmann[4], Melissa Q Reeves[2,3]

**Affiliations**

[1]Department of Microbiology and Immunology, University of California, San Francisco

[2]Huntsman Cancer Institute, University of Utah

[3]Department of Pathology, University of Utah

[4]Earle A. Chiles Research Institute, Providence Cancer Institute of Oregon, Portland, OR

[*]These authors contributed equally.

Correspondence: melissa.reeves@hci.utah.edu

**Keywords**

Squamous cell skin carcinoma, tumor immunology, tumor neoantigen, immune checkpoint

inhibitor, mouse models of cancer

**Abstract**

Immunotherapy is now widely used to treat cancer, but its efficacy in many cancer types

remains modest. To overcome current barriers, preclinical mouse models that faithfully

recapitulate the diversity of cancer types, tumor genetics, mutation burdens, and neoantigen

patterns of human tumors are essential. Currently, there are relatively few transplantable

murine models of squamous cell carcinomas (SCC). Here we describe a novel series of 11 skin

SCC tumor lines, the Carcinogen-Induced Tumor (CIT) lines, syngeneic to the FVB strain. The CIT

lines were established from skin carcinomas induced by DMBA and TPA treatment, and harbor

genetic drivers and overall tumor mutational burdens that recapitulate those found across

multiple human SCCs. Each CIT line gives rise to tumors with a consistent immune infiltration

pattern, ranging from T cell-rich "hot" tumors to T cell-poor "cold" tumors. Hot CIT lines exhibit

partial responses to treatment with immune checkpoint inhibitors, and we have identified two

neoantigens present in an immunotherapy-responsive CIT line. The CIT lines thus provide a

valuable new series of preclinical models for studying anti-tumor immune responses and

developing strategies to improve immunotherapy efficacy in SCCs.

**Introduction**

Over the past decade, immunotherapy has become a new pillar of cancer treatment[1,2]. While

immunotherapy holds immense potential to cure even advanced disease, this promise has only

been realized in a handful of cancer types[1]. Murine models play a vital role in basic and

translational cancer research, providing a means of conducting mechanistic experiments to

dissect cancer and immune cell biology and their interactions with therapy. Transplantable

tumor models, in which cancer cell lines are implanted into immunocompetent syngeneic hosts,

play a particularly important role in preclinical research, as the reliability of their growth

kinetics makes them a workhorse for testing the efficacy of new therapies or combinations.

However, a large proportion of murine tumor immunology studies to date have relied on a

small number of widely-used transplantable tumor models, such as MC38 colorectal cancer and

B16 melanoma. Recent years have also seen a rise in the use of genetically engineered mouse

models (GEMMs) of lung adenocarcinoma[3–6], pancreatic cancer[7,8], and melanoma[9] in

immunotherapy studies.

However, there remain significant gaps in mouse models for studying the anti-tumor

immune response. Many barriers to the efficacy of immunotherapy are cancer type-specific,

and there is a need for mouse models that accurately recapitulate the unique features of each

disease. Additionally, the majority of mouse models currently used for immunotherapy studies

are syngeneic to C57BL/6 mice, which are known to have a Th1-biased immune response[10].

While this is likely beneficial for tumor control—and C57BL/6 mice are widely known for their

resistance to tumorigenesis[11,12]—it is not necessarily representative of the diversity of human

immune responses. While no single mouse model or mouse strain can represent the full

spectrum of human immunity, increasing the diversity of mouse strains used to model anti-

tumor immune responses is essential to capture human diversity.

Here we describe the Carcinogen-Induced Tumor (CIT) cell line panel, a novel series of

murine squamous cell skin carcinoma (skin SCC) cell lines that help to address many of these

gaps. Skin SCC is one of the most common cancer types in the United States[13,14]. While early

stage disease is frequently curable by surgery, immune checkpoint inhibitors (ICIs) including

anti-PD-1 are FDA-approved for patients who have locally advanced, recurrent or metastatic

skin SCC[15]. ICI treatment of skin SCC patients yields response rates of 50% in locally advanced

disease[16,17] and drops as low as 35% in metastatic disease[16]. While this makes skin SCC a

disease type that is considered to respond well to ICIs, there is nonetheless a substantial

proportion of patients with advanced skin SCC who currently do not benefit from

immunotherapy. Additionally, SCCs at other anatomical sites—which share similar genetic

drivers to skin SCC[13,18–20]—exhibit even more limited responses to ICI, with head and neck SCC

patients exhibiting a 15-18% response rate[21,22], and esophageal SCC patients exhibiting 14-28%

response rates[23].

SCCs are highly underrepresented among murine models of cancer and have few cell

line models available, with the MOC series of three oral SCC lines[24] being the main SCC models

to be widely used. To address this gap, we have established the CIT cell lines. The CIT lines are

derived from DMBA/TPA-induced skin SCCs, and they are syngeneic to the FVB strain, which—in

contrast to C57BL/6—is known to exhibit Th2-biased immune responses[10,25] and to be readily

susceptible to tumorigenesis. The CIT cell lines recapitulate both the mutational profiles and

antigen burdens of human SCCs. Although all the CIT lines are derived from skin SCCs, they

harbor alterations in driver genes that are relevant to many SCC types, including *Hras* and

*Trp53*. This is consistent with our previous work showing DMBA/TPA-induced murine skin SCCs

are driven by the same genetic programs as human lung, head and neck, and esophageal

SCCs[26]. As a result of being chemically induced in immune-competent hosts, the CIT lines

further carry endogenous neoantigens that have survived immune selection during tumor

progression of the original carcinoma. Furthermore, tumors arising from the CIT lines exhibit a

range of reproducible patterns of immune infiltration—from high to low T cell infiltrate—along

with a corresponding range of responses to immunotherapy.

The CIT cell line series thus offers a panel of SCC tumor models that exhibit a range of

mutational burdens, immune infiltrates, and neoantigen expression, while being driven by SCC-

relevant genomic drivers and exhibiting histopathologic features characteristic of SCCs. These

properties make the CIT lines a valuable set of preclinical models for investigating tumor

biology and responses to therapy, particularly immunotherapy, in SCCs.

**Results**

**CIT lines display a range of reproducible immune infiltration phenotypes.**

The CIT cell lines were derived from skin carcinomas that arose by topically treating *K5-*

*CreERT2-Confetti* FVB/N mice with a single dose of 7,12-dimethylbenzanthracene (DMBA)

followed by 12-O-tetradecanoylphorbol-13-acetate (TPA) twice weekly for 20 weeks, using a

well-established skin carcinogenesis protocol[27,28] (**Fig. 1A**). In the DMBA/TPA model, benign

papillomas arise after about 6 weeks of TPA treatment, and approximately 10% of these

progress to malignant carcinomas. Primary carcinomas were resected when they were

approximately 1cm in longest diameter, digested into a single cell suspension, and grown under

standard tissue culture conditions (DMEM with 10% FBS, antibiotics, and antimycotics). Each

cell line was established in tissue culture for >10 passages. Implantation of $1.25 \times 10^5$ tumor cells

into the subcutaneous flank of FVB mice gave rise to tumors with 100% penetrance, and we

subsequently profiled the histopathologic and immune characteristics of these tumors.

111  We carried out immune profiling of 11 CIT cell lines by flow cytometry, quantifying the

112 number of total T cells, CD4+ and CD8+ T cells in tumors from each CIT model **(Supp. Fig. 1A)**.

113 We found each CIT line gave rise to tumors with a consistent pattern of immune infiltration,

114 spanning the range of immune hot (highly infiltrated by T cells, e.g., CIT18) to immune cold

115 (poorly infiltrated by T cells, e.g., CIT9) (**Fig. 1B**). Interestingly, we observed a consistent bias in

116 the T cell compartment toward CD4+ T cells, about half of which were Tregs **(Supp. Fig. 1B)**. We

117 selected 5 CIT lines, including representation of both hot and cold models, for more extensive

118 immune profiling **(Supp. Fig. 1A)**. On the whole, we found the myeloid compartment of CIT

119 tumors was strongly biased toward macrophages (CD11b+MHCII+F4/80+), with relatively fewer

120 Ly6C+ and Ly6G+ myeloid cells (monocytes and neutrophils, respectively) (**Fig. 1C**). Macrophage

121 infiltration was highest in the two coldest (lowest T cell infiltrate) CIT models, CIT22 and CIT9.

122 Conversely, hot models CIT18 and CIT6 exhibited the highest B cell infiltration (**Fig. 1C**). There

123 was no clear trend between hot and cold tumors for Ly6G$^{hi}$Ly6C$^{mid}$ neutrophils, Ly6C$^{hi}$CD11b+

124 monocytes, or CD11c+MHCII+ dendritic cells.

125  Tumor histology for the same five selected CIT tumor models was reviewed by a board-

126 certified pathologist. Tumors from all CIT models were poorly differentiated, spindled, arranged

127 in fascicles, and highly mitotic (**Fig. 1D**). Some CIT18 tumors contained pleomorphic, rhabdoid,

128 or epithelioid malignant cell morphologies. Among the hot models, CIT18 and CIT17 tumors

129 contained brisk tumor-infiltrating lymphocyte (TIL) infiltration, while CIT6 tumors contained

130 focal clusters of TILs or TILs localized to the edge of the tumor. In comparison, tumors arising

131 from the cold lines CIT9 and CIT22 displayed very little TIL infiltration. CIT6, CIT17, CIT18 and

132 CIT22 tumors contained diffuse infiltration of polymorphonuclear (PMN) cells, while PMNs in

cold CIT9 tumors were largely restricted to the edge of the tumors. Altogether, we find that the

CIT cell lines give rise to tumors that recapitulate the histologic features of high grade,

sarcomatoid SCC, and each exhibit characteristic patterns of immune infiltration.

**Genomic profiles of CIT lines recapitulate human SCC.**

All 11 CIT cell lines were subjected to exome sequencing, to a median depth of 153X.

Consistent with previous sequencing of DMBA/TPA tumors by ourselves and others[26,29,30], CIT

cell lines were genomically characterized by *Hras* driver mutations, a substantial mutation

burden, and whole-chromosome copy number alterations. The mutation burden among the CIT

lines ranged from 46 to 256 non-synonymous mutations (**Fig. 2A, Table 1**). Ten of our CIT lines

exhibited an *Hras* Q61L mutation, while one, CIT16, exhibited an *Hras* G12V as well as a *Kras*

G13R mutation (**Fig. 2B**). Seven CIT lines displayed additional non-synonymous mutations in

driver genes known to be mutated in human cutaneous, head and neck, lung, and esophageal

SCC, including *Trp53* and *Fgfr3*. (**Fig. 2B**). The majority of total mutations were T>A

transversions, with a particular prevalence of CTN>A mutations (with N representing any

nucleotide) consistent with the known mutagenic signature of DMBA[26,29,30] (**Fig. 2C**). The CIT

lines also harbored whole-chromosome alterations, primarily amplifications, especially on

chromosomes 6 and 15 (**Fig. 2D**). Chromosome 6 amplifications are common in DMBA/TPA

carcinomas[26]. Chromosome 15 amplifications were also previously observed, but they were

more heavily represented in the CIT lines than in a previously sequenced cohort of primary

tumors[26], suggesting perhaps a selection in cell culture for lines with this amplification.

Interestingly, we saw no whole-chromosome amplifications of chromosome 1 and only one

amplification of chromosome 7. We previously reported that chromosome 1 and 7

amplifications are common in DMBA/TPA carcinomas with an epithelial-like "squamous"

morphology, but largely absent in DMBA/TPA carcinomas with a mesenchymal-like "spindle"

morphology [26]. The underrepresentation of these alterations is perhaps not surprising given

that the CIT lines primarily give rise to tumors with spindle cell morphology, but may again

suggest a selection bias of the cell lines we were able to establish in culture.

Given that mutation burden information and immune phenotypes were available for all

11 cell lines, we examined whether a relationship existed between these metrics. While two of

the tumors with the highest T cell infiltration, CIT18 and CIT6, exhibited two of the three

highest mutation burdens, we observed that tumor model with the second-highest T cell

infiltration, CIT16, had a relatively low mutation burden. CIT23 tumors, which had the second-

highest mutation burden, were by contrast very poorly infiltrated by T cells (**Fig. 1B, 2A**). Thus,

aligning with observations from patients, mutation burden alone was insufficient to predict the

immune response to any given tumor.

**Immunologically hot CIT lines respond to immunotherapy.**

Because our CIT lines exhibited moderate to high tumor mutation burdens, we

hypothesized they would carry neoantigens and would be sensitive to immune checkpoint

inhibitors (ICIs). We tested the efficacy of a combination ICI regimen—anti-PD-1 plus anti-CTLA-

4, administered as two doses 3 days apart—on CIT18, CIT6, and CIT9 tumors (**Fig. 3A**).

CIT18 and CIT6 tumors—both of which we had observed to be immunologically hot—

grew significantly slower than controls when treated with anti-PD-1 plus anti-CTLA-4 (hereafter,

"ICI") **(Fig. 3B)**. Conversely, immunologically cold CIT9 tumors displayed no difference in growth

between ICI-treated and control tumors **(Fig. 3B)**. We carried out immune profiling of ICI- and

control-treated tumors to interrogate how ICI remodeled the immune response in these

models. CIT18 tumors had the strongest response to ICIs, both in terms of the reduction in

growth and in the increase in tumor infiltrating T cells. Total T cells, CD8+ T cells and CD4+ T

cells were all more abundant in ICI-treated CIT18 tumors (**Fig. 3C**, **Supp. Fig. 2A**). CIT6 tumors,

which also showed a response to ICI, only exhibited an influx of CD8+ T cells following ICI, while

CIT9 tumors showed no increase in overall T cell infiltration. **(Fig. 3D-E**, **Supp. Fig. 2A)**.

ICI treatment also altered the activation profile of tumor-infiltrating T cells. Effective ICI

treatment in CIT18 and CIT6 tumors was accompanied by an increase in the percentage of CD8+

and CD4+ T cells which exhibited a CD44+CD62L- "effector-like" phenotype **(Fig. 3F-G**, **Supp.**

**Fig. 2B).** Notably, in hot tumors this increase in "effector-like" CD8+ T cells resulted in a skewing

of the CD8+ compartment away from a CD44+CD62L+ "central memory-like" phenotype, even

though the abundance of central memory-like CD8+ T cells as a percent of all immune cells was

unchanged  **(Supp. Fig. 2C)**. In contrast, ICI treatment in cold CIT9 tumors was unsuccessful at

shifting the phenotype of either CD8+ or CD4+ T cells **(Fig. 3H**, **Supp. Fig. 2B-C)**. We further

examined the expression of PD-1, which is present on both activated and exhausted T cells, and

Tim3 and TIGIT, both of which mark exhaustion, in CD8+ T cells. The proportion of tumor-

infiltrating CD8+ T cells expressing PD-1 increased after ICI treatment in all three tumor models

**(Fig. 3I-K)**. Interestingly, the "activated" PD-1+Tim3-TIGIT- T cell population was the primary

CD8+ population to increase in CIT18 tumors following ICI, and this corresponded to the best

tumor growth control **(Fig. 3I, Supp. Fig. 2D)**. By contrast in CIT6 tumors, which showed more

modest tumor growth control, the proportion of "activated" PD-1+Tim3-TIGIT- CD8+ T cells did

not increased as a fraction of PD-1+ CD8+ T cells **(Fig. 3J)**. Instead, there was an increase in the

proportion of PD-1+ CD8+ T cells with an exhausted (PD-1+Tim3+ or PD-1+Tim3+TIGIT+)

phenotype after ICI treatment **(Supp. Fig. 2D)**. Together, this data shows that CIT tumor models

exhibit a range of responses to immunotherapy, and that the degree of sensitivity appears to

be approximately correlated with the level of T cell infiltration in untreated tumors.

**CIT tumors display identifiable and immunotherapy-sensitive neoantigens.**

207            Given that improvement in the CD8+ T cell response corresponded to effective tumor

control following ICI, we next sought to identify the neoantigens recognized during this

response. We focused here on the CIT6 tumor model, which exhibited the intermediate

response to ICI among the three models tested and thus may be most representative of patient

tumors which are recognized by the host immune response but benefit only marginally from

current ICI strategies. We employed the NetH2pan algorithm[31] to predict the binding strength

between the H2-Dq and H2-Kq MHCI alleles carried by FVB/N mice and the neoepitopes

predicted to be generated by all non-synonymous mutations identified in our exome

sequencing analysis. We identified 27 top binders, and these were further filtered to prioritize

those for which the mutant sequence was predicted to exhibit better binding than the wild-

type sequence, and for which we could detect expression of the gene by RNAseq (**Fig. 4A-B**,

**Table 2)**.

219            Candidate neoantigens identified through this pipeline were subsequently screened by

*in vitro* IFNγ ELISPOT assay[32,33]. Whole splenocytes, CD8+ T cells, or CD4+ T cells from the spleen

of CIT6 tumor-bearing mice were incubated with synthetic peptides corresponding to each

candidate epitope sequence (**Fig. 4C**). We identified two peptides, CPAFLPCTV (resulting from a

Brinp3 T436P mutation) and VPPSSLLPL (resulting from a Ttll4 Q454L mutation), which

consistently elicited IFNγ production in sorted CD8+ T cells from CIT6-bearing mice in a peptide

concentration-dependent manner (**Fig. 4D-G**). These peptides elicited no response from sorted

CD4+ T cells (**Supp. Fig. 3A**). When the peptide assay was carried out with whole splenocytes

(no prior sorting of CD8+ or CD4+ T cells), the dose-dependent response to VPPSSLLPL (Ttll4

Q454L mutation) was still observed, but the response to CPAFLPCTV (Brinp3 T436P mutation)

was not detectable above background (**Supp. Fig. 3B**).

We next interrogated the effect of ICI therapy on the T cell response to VPPSSLLPL,

focusing on VPPSSLLPL because its response was more clearly detected in whole splenocytes

(**Supp. Fig. 3B**). CIT6 tumors were treated with three doses of anti-PD-1 and anti-CTLA-4

combination therapy (**Fig. 4H**), and an IFNγ ELISPOT assay was carried out on whole splenocytes

harvested 3 days after the final dose of ICI. Splenocytes from ICI-treated mice produced a

robust IFNγ response when incubated with VPPSSLLPL compared to no peptide **(Fig. 4I)**,

indicating that response to the neoantigen VPPSSLLPL is prominent during immunotherapy

response.

**Discussion**

We have established the CIT lines as a novel series of *Hras*-driven skin SCC tumor lines, and

here present comprehensive histologic, genomic, and immune characterization of these 11 cell

lines. The CIT lines carry genomic drivers relevant to human SCC, as well as a moderate-to-high

mutation burden[13,18,19]. We show that each CIT line exhibits a characteristic immune infiltrate,

with individual cell lines ranging from immune hot to immune cold. Immune hot CIT lines CIT6

and CIT18 exhibit partial responses to an immunotherapy regimen of anti-CTLA-4 and anti-PD-1.

We have also identified two neoantigens that elicit CD8+ T cell responses in the CIT6 cell line. Of

note, while numerous neoantigens have been identified in tumor models syngeneic to

C57BL/6[6,32–35], there has been a notable dearth of neoantigen identification in other mouse

strains including FVB. Antigen identification efforts in FVB mice to date have largely focused on

breast cancer models and identification of antigens arising from the MMTV-PyMT construct or

identification of tumor-associated antigens, rather than neoantigens arising from tumor

mutations[36]. The CIT lines, with the identification of two MHCI-restricted neoantigens in CIT6,

thus provide a valuable new series of models for immunotherapy research in squamous cell

carcinoma.

An additional feature of the CIT lines, having been derived from *K5CreERT2-Confetti*

mice, is that cells can be labeled with fluorescent tags following exposure to Cre

recombinase[28,37]. We have successfully labeled CIT lines *in vitro* using adenoviral Cre

recombinase. While the *K5CreERT2* allele will also allow for labeling of K5+ cells following

tamoxifen treatment, we have found low expression of K5 in these cell lines—as well as in

DMBA/TPA tumors following progression to the carcinoma stage[28] —making this approach less

effective. Introducing Cre and subsequently sorting cells enables the establishment of isogenic

cell lines with each GFP, YFP, RFP or CFP[37]; while introducing Cre to activate labeling without

sorting could allow for multiclonal lineage tracing.

Improving immunotherapy outcomes in SCC remains an unmet clinical need. The CIT

panel of SCC tumor lines provides an important new set of tools to better understand the

mechanisms behind immunotherapy resistance in SCC, and to preclinically model novel

approaches to overcoming ICI resistance and improving SCC patient outcomes.

**Materials and methods:**

Carcinogenesis and cell line generation

To induce tumors, *K5CreER-Confetti FVB/N* mice were treated with 25 µg DMBA followed by

TPA (200 µl of a $10^{-4}$ M solution in acetone) two times a week for 20 weeks. To generate cell

lines, carcinomas were resected at a size of >1 cm in longest diameter, finely chopped, digested

in DMEM containing Collagenase I and IV and DNase I for 45 minutes at 37 °C, washed with PBS,

plated in DMEM plus 10% heat-inactivated FBS and L-glutamine, with penicillin, streptomycin,

and amphotericin B, and passaged until cells stably grew in culture.

In vivo experiments and immune checkpoint inhibitor treatment

A total of $1.25 \times 10^5$ cells were injected subcutaneously into the dorsal flank of 8 to 16 week old

male and female FVB/N mice. For baseline flow cytometric immune profiling and histological

analysis (Fig. 1), tumors were harvested approximately 21 days post-engraftment, when tumors

measured ~1cm in diameter. For immune checkpoint inhibitor (ICI) experiments (Fig. 3), mice

were enrolled in treatment 14 days post-tumor engraftement, when tumors grew to 25-100

mm$^3$ in volume. Mice were treated with 200 µg anti-PD-1 (BioXCell Catalog# BE0146) and 200

µg anti-CTLA-4 (BioXCell Catalog# BE0131) or 200 µg isotype control (BioXCell Catalog#

BE0089). Mice received two doses 3 days apart. Mice were harvested when tumors were

~10mm in length, typically 17-21 days post-ICI enrollment and 31-35 days post-tumor

engraftment, or when an ethical endpoint was reached. All animal experiments were approved

by the by the University of Utah Office of Comparative Medicine (IACUC #22-02003) or by the

University of California San Francisco Laboratory Animal Resource Center (IACUC #AN187679).

This work complies with all the relevant ethical regulations regarding animal research.

<u>Flow cytometric immune profiling</u>

Harvested tumors were finely chopped, digested in DMEM containing Collagenase IV (Sigma-

Aldrich Catalog# 9001-12-1) and DNase I (Sigma-Aldrich Catalog# D5025-15KU) for 45 minutes

at 37 °C, and filtered through 70um filters. Cells were counted using a Countess III, and 5 million

cells were plated and stained with antibodies and Live/Dead NIR stain. Antibodies used for flow

cytometric analyses are listed in **Table 1**. Flow cytometry analysis was performed on a Cytek

Aurora.

<u>Histology</u>

Harvested tumors were fixed in formalin for 24 hours, placed in cassettes (VWR Catalog#

18000), submerged in 70% ethanol, and submitted to ARUP Laboratories Research Histology

Core Laboratory, where they were embedded in paraffin, cut into 4μm-thick slices, stained with

hematoxylin and eosin (H&E) and mounted on slides. The 4μm slides were then analyzed by a

board-certified pathologist, A.G..

<u>Cell line sequencing</u>

DNA was extracted from cultured cells from each cell line and from tails of the mice in which

each tumor originated using the Qiagen DNeasy Blood & Tissue DNA purification kit (Qiagen).

Whole exome sequencing was performed using the Agilent Sure Select Mouse Exon capture kit,

and sequenced to an average depth of 153X on an Illumina HiSeq X by MedGenome, Inc. in

Redwood City, CA. Reads were trimmed with cutadapt (v3.5)[38] and aligned to the mm10 mouse

genome using BWA (v0.7.17)[39]. Reads were deduplicated using MarkDuplicates and

recalibrated using BaseRecalibrator from GATK (v4.1)[40]. Mutations were called using MuTect2

(GATK v4.1)[40,41] and Strelka2 (v2.9.10)[42], using tails from the mice in which each tumor

developed as the matched normal. Mutation calls were filtered using the following criteria:

minimum read depth of 10 at the mutation position in both tumor and normal samples;

minimum of 4 reads supporting the mutation call; for Strelka only, minimum quality QSS score

of 25. Mutations that passed all filters with both callers were kept and were additionally filtered

to remove any germline SNP detected in the panel of normals. Final mutation calls were

annotated with Annovar[43]. Copy number calling was performed with CNVkit[44], using all tails as a

pooled normal. Exome sequencing data from CIT lines is available in SRA (accession number

PRJNA1129114).

<u>Neoantigen peptide prediction</u>

Mutations identified by exome sequencing were used to construct wild-type and mutant amino

acid sequences of all 7mer, 8mer, and 9mer peptides that included each identified single

nucleotide variant. The NetH2pan algorithm was used to predict binding affinity and percentile

rank for each wild-type and mutant peptide sequence. This data was filtered using the following

criteria: mutant peptide with a percentile rank <0.5, log affinity ratio >0 (affinity ratio defined as

ratio of predicted binding affinity of mutant sequence dividesd by wild-type sequence), and a

gene expression >300 RPM based on RNAseq of whole CIT6 and CIT9 tumors.

<u>ELISpot assays</u>

Whole spleens were harvested from FVB mice and processed by crushing through a 70μm mesh

filter, pre-wet with 1X PBS, using a syringe plunger and washed. Red blood cells were lysed for 5

minutes using ACK lysis buffer and splenocytes counted. Spleens from CIT6-bearing FVB mice

were harvested 2-4 weeks after tumor inoculation. CD8+ T cells were isolated from whole

spleens using EasySep$^{TM}$ Mouse CD8+ T Cell Isolation Kits  (STEMCELL Technologies Catalog#

19853) and CD4+ T cells were isolated from whole spleens using EasySep$^{TM}$ Mouse CD4+ T Cell

Isolation (STEMCELL Technologies Catalog# 19852).

Interferon-gamma (IFNγ) ELISpots were performed over a three-day assay.

Multiscreen®HTS 96-well filtration plate(s) were coated with IFNγ capture antibody (clone AN-

18, Catalog# 517902), diluted 1:500 in 1X PBS. Plates were wrapped in aluminum foil to equally

distribute temperature and incubated overnight, or at least 18 hours, at 4°C. The following day

plates were washed twice with 200μl RPMI-2 (RPMI with 2% FBS, 1% L-glutamate, 1%

streptococcus and penicillin antibiotic solution, and 0.09% 2-Mercaptoethanol (BME)) and

blocked for two hours with 200μl of RPMI-20 (RPMI with 20% FBS, 1% L-glutamate, 1%

streptococcus and penicillin antibiotic solution, and 0.09% 2-Mercaptoethanol (BME)) before

splenocytes or T cells were added. For assays where total splenocytes were used, 500,000

splenocytes were plated per well. For assays using CD8+ and CD4+ T cells, 20,000 cells per well

were co-cultured with 500,000 splenocytes from a non-tumor-bearing mouse to provide a

source of antigen presenting cells. Stimulation medium was prepared by diluting synthetic

peptide in RPMI-10 to desired peptide concentration(s). All synthetic peptides used for these

assays were ordered through GenScript. Peptides were reconstituted in DMSO or MilliQ water

according to solubility testing provided by the manufacturer. Reconstituted peptides were

aliquoted and frozen at -80°C. The negative control consisted of MilliQ water and RPMI-10.

Positive control contained 0.04% Phorbol 12-myristate 13-acetate (PMA) and 0.09% Ionomycin

in RPMI-10.

Plates were wrapped in aluminum foil and incubated undisturbed at 37°C and 5% $CO_2$

for 44-48 hours before developing using biotinylated anti-IFNγ detection antibody (clone R4-

6A2, Catalog# 505704), streptavidin-HRP enzyme conjugation (Thermofisher, Catalog# 405210),

and AEC (3-amino-9-ethyl-carbazole) substrate. AEC substrate was prepared by dissolving 20

mg of AEC in 2 ml of N, N Dimethylforamamide (Sigma-Aldrich, Catalog# 227056), diluting in 0.1

M Acetate Solution, adding hydrogen peroxide and filtering with a 0.45µM filter. Spots were

Imaged using the Agilent BioTek Cytation7 and analyzed using the Gen5 software.

**Data availability**

Exome sequencing of CIT lines is available in SRA (accession number PRJNA1129114).

**Acknowledgements**

This work was supported by the UCSF Program for Breakthrough Biomedical Research (PBBR)

Sandler Fellowship, US National Cancer Institute (NCI) grant R21CA264599, the Huntsman

Cancer Institute Cancer Center Support Grant P30CA040214, the American Cancer Society IRG-

21-131-01, Five For The Fight, The Cancer League, and the V Foundation (V2024-002). G.S. is

supported by the Institute of General Medical Sciences grant T32GM141848. We thank the

University of Utah Flow Cytometry Core, the Associated Regional and University Pathologists,

Inc. (ARUP Laboratories) Research Histology Core, the University of Utah Center for High

Performance Computing (CHPC), and the Preclinical Research Resource (PRR), in particular D.

Lum and W. Sun, and the UCSF Parnassus Flow CoLab. We also thank A. Lai for help establishing

the ELISPOT methods. The computational resources used for exome sequencing analysis were

partially funded by the NIH Shared Instrumentation Grant 1S10OD021644-01A1. We thank and

all members of the Reeves Lab for thoughtful input during the development of this story. Some

figures were created with Biorender.com.

**387     Author contributions**

M.T., R.L., L.L., A.B., and M.Q.R. designed the project and experiments. R.L., S.H., M.T., L.L. and

M.Q.R. led experiment implementation and data analysis. A.B. designed and carried out the

neoantigen identification experiments. S.H. and G.S. carried out the immune checkpoint

inhibitor experiments, with support from R.L. Board-certified pathologist A.G. provided

histology review. R.L., M.T. and M.Q.R. wrote the manuscript, with input from other co-authors.

P.C., K.N., D.S., C.M.L. and M.G. assisted with experiments.

**References**

1.    Ribas, A. & Wolchok, J. D. Cancer immunotherapy using checkpoint blockade. *Science*

**359**, 1350–1355 (2018).

2.    Chen, D. S. & Mellman, I. Elements of cancer immunity and the cancer-immune set point.

*Nature* **541**, 321–330 (2017).

3.    Schenkel, J. M. *et al.* Conventional type I dendritic cells maintain a reservoir of

proliferative tumor-antigen specific TCF-1+ CD8+ T cells in tumor-draining lymph nodes.

*Immunity* **54**, 2338-2353.e6 (2021).

4.    Burger, M. L. *et al.* Antigen dominance hierarchies shape TCF1+ progenitor CD8 T cell

phenotypes in tumors. *Cell* **184**, 4996-5014.e26 (2021).

5.    Nguyen, K. B. *et al.* Decoupled neoantigen cross-presentation by dendritic cells limits

anti-tumor immunity against tumors with heterogeneous neoantigen expression. *Elife*

**12**, e85263 (2023).

6.    Roerden, M. *et al.* Neoantigen architectures define immunogenicity and drive immune

evasion of tumors with heterogenous neoantigen expression. *J Immunother Cancer* **12**,

e010249 (2024).

7.    Li, J. *et al.* Tumor Cell-Intrinsic Factors Underlie Heterogeneity of Immune Cell Infiltration

and Response to Immunotherapy. *Immunity* **49**, 178-193.e7 (2018).

8.    Byrne, K. T. & Vonderheide, R. H. CD40 Stimulation Obviates Innate Sensors and Drives T

Cell Immunity in Cancer. *Cell Rep* **15**, 2719–2732 (2016).

9.    Wee, Y. *et al.* Tumour-intrinsic endomembrane trafficking by ARF6 shapes an

immunosuppressive microenvironment that drives melanomagenesis and response to

checkpoint blockade therapy. *Nat Commun* **15**, 6613 (2024).

10.   Sellers, R. S., Clifford, C. B., Treuting, P. M. & Brayton, C. Immunological variation

between inbred laboratory mouse strains: Points to consider in phenotyping genetically

immunomodified mice. *Vet Pathol* **49**, 32–43 (2012).

11.   Hennings, H., Shores, R., Mitchell, P., Spangler, E. F. & Yuspa, S. H. Induction of

papillomas with a high probability of conversion to malignancy. *Carcinogenesis* **6**, 1607–

10 (1985).

12.   Davie, S. A. *et al.* Effects of FVB/NJ and C57Bl/6J strain backgrounds on mammary tumor

phenotype in inducible nitric oxide synthase deficient mice. *Transgenic Res* **16**, 193–201

(2007).

13.   Chang, D. & Shain, A. H. The landscape of driver mutations in cutaneous squamous cell

carcinoma. *NPJ Genom Med* **6**, 61 (2021).

14.   Nehal, K. S. & Bichakjian, C. K. Update on Keratinocyte Carcinomas. *New England Journal*

*of Medicine* **379**, 363–374 (2018).

15.   Ansary, T. M., Hossain, M. R., Komine, M. & Ohtsuki, M. Immunotherapy for the

Treatment of Squamous Cell Carcinoma: Potential Benefits and Challenges. *International*

*Journal of Molecular Sciences* **23**, 8530 (2022).

16.   Hughes, B. G. M. *et al.* Pembrolizumab for locally advanced and recurrent/metastatic

cutaneous squamous cell carcinoma (KEYNOTE-629 study): an open-label,

nonrandomized, multicenter, phase II trial. *Annals of Oncology* **32**, 1276–1285 (2021).

17.    Migden, M. R. *et al.* PD-1 Blockade with Cemiplimab in Advanced Cutaneous Squamous-

439            Cell Carcinoma. *New England Journal of Medicine* **379**, 341–351 (2018).

440    18.    Pickering, C. R. *et al.* Mutational landscape of aggressive cutaneous squamous cell

441            carcinoma. *Clin Cancer Res* **20**, 6582–92 (2014).

442    19.    Stransky, N. *et al.* The mutational landscape of head and neck squamous cell carcinoma.

443            *Science* **333**, 1157–60 (2011).

444    20.    The Cancer Genome Atlas. Comprehensive genomic characterization of squamous cell

445            lung cancers. *Nature* **489**, 519–25 (2012).

446    21.    Cohen, E. E. W. *et al.* Pembrolizumab versus methotrexate, docetaxel, or cetuximab for

447            recurrent or metastatic head-and-neck squamous cell carcinoma (KEYNOTE-040): a

448            randomised, open-label, phase 3 study. *The Lancet* **393**, 156–167 (2019).

449    22.    Chow, L. Q. M. Head and Neck Cancer. *New England Journal of Medicine* **382**, 60–72

450            (2020).

451    23.    Hirano, H. & Kato, K. Systemic treatment of advanced esophageal squamous cell

452            carcinoma: Chemotherapy, molecular-targeting therapy and immunotherapy. *Jpn J Clin*

453            *Oncol* **49**, 412–420 (2019).

454    24.    Kono, M., Saito, S., Egloff, A. M., Allen, C. T. & Uppaluri, R. The mouse oral carcinoma

455            (MOC) model: A 10-year retrospective on model development and head and neck cancer

456            investigations. *Oral Oncol* **132**, 106012 (2022).

457    25.    Kim, E. M., Bae, Y. M., Choi, M. H. & Hong, S. T. Cyst formation, increased anti-

458            inflammatory cytokines and expression of chemokines support for Clonorchis sinensis

459            infection in FVB mice. *Parasitol Int* **61**, 124–129 (2012).

26.    McCreery, M. Q. *et al.* Evolution of metastasis revealed by mutational landscapes of

chemically induced skin cancers. *Nat Med* **21**, 1514–1520 (2015).

27.    Wong, C. E. *et al.* Inflammation and Hras signaling control epithelial-mesenchymal

transition during skin tumor progression. *Genes Dev* **27**, 670–82 (2013).

28.    Reeves, M. Q., Kandyba, E., Harris, S., Del Rosario, R. & Balmain, A. Multicolour lineage

tracing reveals clonal dynamics of squamous carcinoma evolution from initiation to

metastasis. *Nat Cell Biol* **20**, 699–709 (2018).

29.    Nassar, D., Latil, M., Boeckx, B., Lambrechts, D. & Blanpain, C. Genomic landscape of

carcinogen-induced and genetically induced mouse skin squamous cell carcinoma. *Nat*

*Med* **21**, 946–954 (2015).

30.    Li, Y. R. *et al.* Long Term Latency of Highly Mutated Cells in Normal Mouse Skin is

Reversed by Exposure to Tumor Promoters and Chronic Tissue Damage. *Cancer Discov*

(2025).

31.    DeVette, C. I. *et al.* NetH2pan: A computational tool to guide MHC peptide prediction on

murine tumors. *Cancer Immunol Res* **6**, 636–644 (2018).

32.    Castle, J. C. *et al.* Exploiting the mutanome for tumor vaccination. *Cancer Res* **72**, 1081–

1091 (2012).

33.    Alspach, E. *et al.* MHC-II neoantigens shape tumour immunity and response to

immunotherapy. *Nature* **574**, 696–701 (2019).

34.    Matsushita, H. *et al.* Cancer exome analysis reveals a T-cell-dependent mechanism of

cancer immunoediting. *Nature* **482**, 400–404 (2012).

35.    Yadav, M. *et al.* Predicting immunogenic tumour mutations by combining mass

spectrometry and exome sequencing. *Nature* **515**, 572–576 (2014).

36.    DeVette, C. I. *et al.* A pipeline for identification and validation of tumor-specific antigens

in a mouse model of metastatic breast cancer. *Oncoimmunology* **9**, 1685300 (2020).

37.    Tanaka, M. *et al.* Tumor cell heterogeneity drives spatial organization of the intratumoral

immune response. *J Exp Med* **222**, e20242282 (2025).

38.    Martin, M. Cutadapt removes adapter sequences from high-throughput sequencing

reads. *EMBnet J* **17**, 10 (2011).

39.    Li, H. & Durbin, R. Fast and accurate short read alignment with Burrows-Wheeler

transform. *Bioinformatics* **25**, 1754–60 (2009).

40.    Van der Auwera, G. A. & O'Connor, B. D. *Genomics in the Cloud: Using Docker, GATK, and*

*WDL in Terra*. (O'Reilly Media, 2020).

41.    Cibulskis, K. *et al.* Sensitive detection of somatic point mutations in impure and

heterogeneous cancer samples. *Nat Biotechnol* **31**, 213–9 (2013).

42.    Kim, S. *et al.* Strelka2: fast and accurate calling of germline and somatic variants. *Nat*

*Methods* **15**, 591–594 (2018).

43.    Wang, K., Li, M. & Hakonarson, H. ANNOVAR: functional annotation of genetic variants

from high-throughput sequencing data. *Nucleic Acids Res* **38**, e164 (2010).

44.    Talevich, E., Shain, A. H. & Bastian, B. C. CNVkit: Copy number detection and

viziualization for targeted sequencing using off-target reads. *PLoS Comput Biol* **12**,

e1004873 (2016).

# Figure 1

**Figure 1. Immune infiltration and histology profiles of CIT lines.**

**(A)** Schematic of CIT cell line generation. Tumors were induced with the DMBA/TPA

carcinogenesis model, and cell lines were established from malignant carcinomas. **(B)**

Quantification of total T cells, CD4+ T cells and CD8+ T cells by flow cytometry in each CIT line.

**(C)** Quantification of macrophages, neutrophils, monocytes, B cells and dendritic cells by flow

cytometry in 5 selected CIT lines. **(D)** Representative images of tumors from selected CIT lines

stained with hematoxylin and eosin (H&E) for histological analysis. Scale bar = 20 $\mu$m. Statistical

analysis in (B-C) by one-way ANOVA with Tukey post-hoc test. * $p < 0.05$, ** $p < 0.01$, *** $p <$

0.001.

# Figure 2

**Figure 2. Genetic profiling of CIT lines.**

**(A)** Number of non-synonymous mutations per cell line, based on exome sequencing. **(B)** Genes

mutated in CIT lines which are known to be commonly mutated in human SCC. **(C)**  Frequencies

of mutations observed in each of 96 possible trinucleotide contexts for all mutations in CIT cell

lines. Trinucleotide contexts, arranged on the x axis, are grouped by the base pair change of the

mutation. Highest peaks are observed at CTG>A and GTG>A. **(D)** CNV alterations detected in

each chromosome in CIT lines based on exome sequencing. Red denotes copy number gains,

blue denotes copy number losses.

# Figure 3

**Figure 3. Immunologically hot CIT lines are responsive to combined anti-PD-1 and anti-CTLA-4 immune checkpoint inhibitor (ICI) therapy.**

**(A)** Schematic of ICI treatment regimen on CIT tumor-bearing mice. **(B)** Average tumor volumes of immunologically hot CIT18 and CIT6 tumors and immunologically cold CIT9 tumors after combined ICI therapy, compared to tumors treated with isotype control antibodies. **(C-E)** Total T cell, CD4+ T cell, and CD8+ T cell infiltration of CIT18 (C), CIT6 (D) and CIT9 (E) tumors as a percentage of all CD45+ immune cells. **(F-H)** Abundance of effector subsets of CD8+ T cells defined by the combinatorial expressions of CD44 and CD62L in CIT18 (F), CIT6 (G) and CIT9 (H) tumors. **(I-K)** Abundance of CD8+ T cells expressing PD-1, and frequency of "activated" PD-1+Tim3-TIGIT- CD8+ T cells in CIT18 (I), CIT6 (J) and CIT9 (K) tumors. All statistical analyses performed with Welch's t-test. * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$, **** $p < 0.001$.

# Figure 4

**Figure 4. CIT lines express identifiable neoantigens whose cognate CD8+ T cells are responsive**

**to immunotherapy.**

**(A)** Selection criteria for CIT6 mutations predicted to generate neoantigens. **(B)** Predicted

peptide log affinity ratio (affinity ratio is defined as ratio between predicted binding affinity of

mutant versus wild-type sequence) and gene expression level of mutations predicted to be top

CIT6 candidate neoantigens, shown for H2-Dq and H2-Kq alleles. Affinity predictions were

generated using NetH2Pan algorithm. **(C)** Schematic of IFNγ ELISpot assay. For screening of

CD8+ T cell and CD4+ T cell responses, CD8+ or CD4+ T cells were isolated from the spleens of

tumor bearing mice and co-cultured with total splenoyctes from a healthy tumor-naïve mouse.

**(D)** Representative IFNγ ELISpot images of assay peformed with CD8+ T cells from the spleen of

CIT6-bearing mice cultured with 160ug/ml of Brinp3 peptide CPAFLPCTV (top row) and negative

control (no peptide, bottom row). **(E)** Quantification of average IFNγ ELISpot spot counts for

CD8+ T cells from the spleen of CIT6-bearing mice cultured with different concentrations of

Brinp3 peptide CPAFLPCTV. Each colored line represents data points from a different mouse.

**(F)** Representative IFNγ ELISpot images of assay peformed with CD8+ T cells from the spleen of

CIT6-bearing mice cultured with 160ug/ml of Ttll4 peptide VPPSSLLPL (top row) and negative

control (no peptide, bottom row). **(G)** Quantification of average IFNγ ELISpot spot counts for

CD8+ T cells from the spleen of CIT6-bearing mice cultured with different concentrations of

Ttll4 peptide VPPSSLLPL. **(H)** Schematic of immune checkpoint inhibitor (ICI) treatment regimen

and ELISpot analysis. **(I)** Average spot counts of ICI-treated splenocytes cultured with 80ug/ml

of Ttll4 peptide vs negative control. Statistical analysis in panels E, G, and I was done by paired

t-test. * $p < 0.05$, ** $p < 0.01$.

# Supplemental Figure 1

**A**

**B**

### Regulatory T cells (%CD4)

560 **Supp. Fig 1. Flow cytometry gating strategy and regulatory T cell infiltration in CIT tumors.**

561 **(A**) Flow cytometric gating strategy of main immune cell types. Immune analysis of a CIT6

562 tumor is shown as representative data. **(B)** Abundance of Foxp3+CD25+ regulatory T cells

563 (Tregs) as percentage of CD4+ T cells in CIT18, CIT6 and CIT9 tumors. Statistical analysis by One-

564 way ANOVA with Tukey post-hoc test.

# Supplemental Figure 2

**Supp. Fig. 2. Tumor-infiltrating T cell effector activity after treatment with combined anti-PD-**

**1 and anti-CTLA-4 immune checkpoint inhibitor (ICI) therapy.**

**(A)** Total T cell, CD4+ T cell, and CD8+ T cell infiltration of CIT18, CIT6 and CIT9 as percentage of

all live cells, including tumor and stromal cells. **(B)** Abundance of effector subsets of CD4+ T

cells defined by the combinatorial expressions of CD44 and CD62L in CIT18, CIT6 and CIT9. **(C)**

Infiltration of CD44+CD62L- "effector-like" CD8+ T cells and CD44+CD62L+ "central memory-

like" CD8+ T cells as percentage of all CD45+ immune cells. **(D)** Abundance of CD8+ PD-1+ T cells

expressing specific combinations of TIGIT and Tim3 exhaustion markers in CIT18, CIT6 and CIT9

tumors. All statistical analyses performed with Welch's t-test. * $p < 0.05$, ** $p < 0.01$, *** $p <$

0.001, **** $p < 0.001$.

# Supplemental Figure 3

**A**

**B**

**Supp. Fig. 3. CIT6 ELISpot analysis of unfractionated splenocytes and CD4+ splenocytes.**

**(A)** Spot counts from ELISpot assays performed with CD4+ T cells isolated from the spleens of

CIT6-bearing mice and cultured with splenocytes from tumor-naïve mice loaded with increasing

concentrations of Brinp3 and Ttll4 peptides. **(B)** Spot counts from ELISpot assays performed

with total splenocytes from CIT6-bearing mice cultured with increasing concentrations of

Brinp3 and Ttll4 peptides. All statistical analyses performed with paired t-test. * $p < 0.05$, ** $p <$

0.01, *** $p < 0.001$.

## Table 1: Number of Non-Synonymous Mutations per CIT Cell Line

| Cell Line | Non-Synonymous Mutations |
|-----------|--------------------------|
| CIT18 | 170 |
| CIT16 | 86 |
| CIT6 | 256 |
| CIT17 | 74 |
| CIT19 | 56 |
| CIT20 | 113 |
| CIT21 | 111 |
| CIT22 | 94 |
| CIT23 | 176 |
| CIT24 | 46 |
| CIT9 | 102 |

588 **Table 1: Number of non-synonymous mutations per CIT cell line.** Number of non-synonymous

589 mutations based on exome sequencing of CIT cell lines.

## Table 2: List of Filtered Peptide Candidates for CIT6 Antigens

| Gene | Amino Acid Change | Peptide | Allele | Percentile Rank |
|---|---|---|---|---|
| Slc39a10 | S286I | LPEHIGHEL | Dq | 0.0225 |
| Ttll4 | Q454L | VPPSSLLPL | Dq | 0.1367 |
| Brinp3 | T436P | CPAFLPCTV | Dq | 0.2119 |
| Cep76 | P158A | VPCACEADF | Dq | 0.4240 |
| Atp9a | I216F | EPNIDFHNF | Dq | 0.4662 |
| Fam83f | Q4L | VPALPTESL | Dq | 0.1392 |
| Emc1 | H307L | SHYALLHYL | Dq | 0.2106 |
| Cnot10 | A543P | VPLALGDNL | Dq | 0.0063 |
| Fam13b | D766H | QEKLARHL | Kq | 0.1266 |
| Setbp1 | N1160Y | HDYLSGLF | Kq | 0.2688 |
| Pcdh18 | M631I | NEENIFII | Kq | 0.1341 |
| Cpt1c | Q29H | HEIYLCAL | Kq | 0.3203 |
| Trip4 | Q423L | HQHQLRIL | Kq | 0.3234 |

592    **Table 2: List of filtered peptide candidates for CIT6 neoantigens.** List of all neoantigen peptide

593    candidates that met the criteria specified in Fig. 4A. Percentile ranked based on NetH2pan

594    algorithm.

| Target | Fluorophore | Clone | Dilution | Manufacturer | Catalog # |
|---|---|---|---|---|---|
| **Table 3**: List of Antibodies and Conjugated Fluorophores used for Flow Cytometry Analysis | | | | | |
| B220 | BUV661 | RA3-6B2 | 1:200 | BD Biosciences | 612972 |
| CD11b | BV711 | M1/70 | 1:100 | BioLegend | 101242 |
| CD11c | BV605 | N418 | 1:100 | BioLegend | 117334 |
| CD16/CD32 (FcR Block) | N/A | 2.4G2 | 1:167 | Tonbo | 70-0161-U500 |
| CD3e | BUV395 | 145-2C11 | 1:62.5 | BD Biosciences | 563565 |
| CD4 | BUV563 | GK1.5 | 1:200 | BD Biosciences | 612923 |
| CD44 | BV785 | 1M7 | 1:200 | BioLegend | 103059 |
| CD45 | AlexaFluor700 | 30-F11 | 1:400 | BioLegend | 103128 |
| CD62L | FITC | MEL-14 | 1:200 | Invitrogen | 11-0621-81 |
| CD8a | BV570 | 53-6.7 | 1:200 | BioLegend | 100739 |
| CD8a | BV785 | 53-6.7 | 1:200 | BioLegend | 100750 |
| F4/80 | APC | BM8 | 1:100 | BioLegend | 123116 |
| FoxP3 | PE-Cy7 | FJK-16s | 1:50 | eBioscience | 25-5773-82 |
| Live/Dead | NIR | N/A | 1:1000 | Invitrogen | L34976A |
| Ly6C | BV650 | HK1.4 | 1:200 | BioLegend | 128049 |
| Ly6G | PE-CF594 | 1A8 | 1:200 | BD Biosciences | 562700 |
| MHCII | PE-Cy7 | M5/114.15.2 | 1:400 | BioLegend | 107630 |
| PD-1 | BV605 | 29F.1A12 | 1:100 | BD Biosciences | 135220 |
| TIGIT | BV421 | 1G9 | 1:100 | BioLegend | 142111 |
| Tim3 | APC | RMT3-23 | 1:100 | BioLegend | 119705 |

597   **Table 3: List of antibodies used for flow cytometry analysis.** All antibodies, clone number, and

598   conjugated fluorophores for antibodies used for flow cytometry.

600