SYMPOSIUM

Rhagovelia uses interfacial run-and-tumble locomotion to improve prey capture in flowing environments

Ishant Tiwari,[*][1] Nithil Nagappan,[*][1] Jacob S. Harrison[1] and Saad Bhamla[1,†]

[1]School of Chemical and Biomolecular Engineering, Georgia Institute of Technology, Atlanta, Georgia, United States
[*]Corresponding author. saadb@chbe.gatech.edu

Abstract

Rhagovelia oriander is a neustonic insect native to the rivers and streams of North and South America, known for its distinctive skating movement on the water's surface. This movement resembles the correlated random-walk pattern seen in microorganisms such as *Escherichia coli*. Previous studies have primarily focused on limb adaptations and biomechanics, leaving the ecological significance inadequately addressed. We combine field observations with controlled laboratory experiments using a flow mill to investigate the dynamics of *R. oriander* under relevant flow conditions. Our findings indicate that this insect exhibits a two-dimensional run-and-tumble motion, often incorporating lateral tumbles following straight runs (run distance: 30.7 ± 9.7 mm). We find that this behavior is resilient to changes in flow speed. In-silico simulations of predator-prey interactions demonstrated that this locomotion method significantly enhances prey capture efficiency compared to linear movement, particularly when the prey is following a rapid flow field. Our results document run-and-tumble locomotion in a millimeter-scale organism, showcasing a unique example of convergent behavior across diverse taxonomic groups and providing valuable insights into locomotion ecology while serving as a source of inspiration for bioinspired robotics and environmental exploration algorithms.

Key words: Run-and-Tumble, Neustonic Insects, Water-striders, *Rhagovelia oriander*, Insect foraging

Introduction

Navigating complex environments efficiently for exploration and resource acquisition presents key challenges for mobile organisms. Various species from different taxa have developed unique foraging and search strategies Bell (2012), often striking a balance between directional persistence and exploratory randomness to enhance their chances of survival and reproduction. Strategies such as correlated random walks Kareiva and Shigesada (1983), Lévy flights Viswanathan et al. (1996); Pyke (2015), and area-restricted searches Othmer et al. (1988); Smith (1974) have been thoroughly studied in both terrestrial and aquatic organisms, showcasing how selective pressures tailor organismal behavior to specific ecological settings.

The run-and-tumble locomotion strategy effectively alternates between straight, directional runs and quick, random reorientations (tumbles). This behavior has been extensively researched in microorganisms such as *Escherichia coli*, which use run-and-tumble to navigate chemical gradients, thus improving their ability to efficiently find nutrients Berg and Brown (1972); Patteson et al. (2015). Similarly, the unicellular algae *Chlamydomonas reinhardtii* also employs this strategy by adjusting flagellar synchronization, allowing a shift between straight swimming and rapid reorientations, facilitating effective spatial exploration Polin et al. (2009).

Although there has been considerable research at the microbial level Berg and Brown (1972); Ishihara et al. (1983); Polin et al. (2009); Patteson et al. (2015); Gomez-Marin et al. (2011); Johnson et al. (2024), run-and-tumble behavior is still largely unexplored at the millimeter scale. *R. oriander*, commonly seen skimming over flowing streams in North and South America, exhibits a movement that resembles the previously described run-and-tumble behavior (Figure 1). *Rhagovelia* are a unique genus within the water strider family (Velidae). They excel in navigating fast-flowing streams, enabling them to inhabit areas that other

---

*These authors contributed equally to this work.

water striders do not, and possibly take advantage of new ecological niches. Prior studies have mainly focused on *R. oriander's* distinct morphological traits, including specialized hydrophilic fans on their middle legs, which enable swift turns through elastocapillary-driven morphing Santos et al. (2017); Ortega-Jimenez et al. (2024). These fans are believed to be the morphological innovation allowing them to thrive in demanding, high-flow environments; however, their unique locomotion behavior might also play a critical role in their success.

In this study, we combine empirical field observations with controlled tests in a custom-designed flow mill to explore how varying flow conditions influence the run-and-tumble behavior of *R. oriander*. We propose that *R. oriander* utilizes run-and-tumble motion to maneuver across the water's surface, with changes in its runs and tumbles expected to correlate with different flow rates. Furthermore, through mathematical modeling, we anticipate that this behavior will enhance food capture rates in simulated *R. oriander* predator-prey interactions.

Materials and methods

Animals

We collected *R. oriander* (Figure 1A) from the East Palisades Trail in Atlanta, GA, USA (33°53'18.8"N 84°26'21.4"W), its natural habitat in ponds and creeks (Figure 1B), and maintained them in a controlled laboratory environment. The insects were kept in a plastic enclosure measuring (12x24x12) inches, filled with water, and maintained at ($\sim20°C$). An aquarium filter and air stones were included in the water to ensure a steady flow. The enclosure was supplemented with aquatic plants, algae, and decomposing leaves to ensure that the environmental conditions mimicked their natural habitat. Insects were exposed to a circadian lighting cycle of 12 hours and were fed a diet of live springtails (*Folsomia candida*) daily. Since our study involved working with only invertebrates, it did not require an Institutional Animal Care and Use Committee (IACUC) approval.

Behavioral Testing

Initial studies were performed at the field site with a GoPro camera mounted on a tripod and a top-down view of a section of the flowing creek (Figure 1C). We selected the recording area by identifying a region with populations of *R. oriander* actively moving on the water's surface. A subset (n=1-3) of these individuals was enclosed within a mesh boundary that allowed them to remain in the field of view, without obstructing water flow. Characteristic tracks captured during the field recordings are illustrated in Figure 1 D. The movement trajectory of the *R. oriander* displays a unique pattern, characterized by the bug traveling in relatively straight lines, followed by sudden directional changes.

To study the locomotion of *R. oriander* in a controlled laboratory setting, we designed a custom flow mill (Figure 2A) consisting of a three-sided acrylic enclosure (9 x 9 x 2 inches) open at the top. Filtered deionized water was continuously pumped into one side and allowed to flow out over the opposite edge. Below this enclosure, a reservoir housed the pumps for water recirculation and contained an upward-facing light source that was diffused by the acrylic enclosure to create uniform illumination.

We verified the velocity fields within the setup using Particle Image Velocimetry (PIV), with lycopodium powder serving as tracer particles to visualize flow trajectories. Flow velocities were adjusted and confirmed at 8–10 mm/s (low flow), 29–31 mm/s (medium flow), and 34–36 mm/s (high flow) before introducing the test individuals to the flow mill. Previous studies Ortega-Jimenez et al. (2024) have reported observing *Rhagovelia* in natural flows as high as 1600 mm/s with significant turbulence levels (i.e., ratio between root-mean-square of flow velocity fluctuations to the mean velocity) as high as 65%.

For our experiment, we tested *R. oriander* moving on the surface of the flowing water arena under the three flow regimes (low flow, medium flow, and high flow). Five adult individuals were tested for each flow, resulting in 15 trials, 1 individual per trial. During experimentation, a GoPro Hero 9 was placed viewing the arena top-down and recording the insects at 240 fps 2K resolution for five minutes.

Data Analysis

Video recordings of trials were processed using Adobe$^©$ Premiere Pro. The video processing included removing fisheye lens distortion and cropping the video to our region of interest. We then used DeepLabCut, a neural network-based markerless tracking software Nath et al. (2019), to track two points on the animal: the anterior-most tip of the head and the posterior-most tip of the abdomen, throughout the trial. We extracted the coordinates of the head and tail points' motion and then took their instantaneous average to determine the approximate location of the body center. From the extracted positional data, we computed various metrics such as instantaneous speed, mean squared displacement curves, run-distance distributions, and turning angles. To reduce noise in our data, we excluded any low-confidence estimates from DeepLabCut as well as any points close to the boundary of the observation arena. The trimmed data yielded periods of trajectory data, which were used for downstream analysis. Exemplary data, along with rejection boundary margins and water flow directions, are visualized in Figure 2B.

Statistical Analysis

We performed various statistical tests to check for significant differences in distributions across test conditions and across populations. We observed three conditions (low flow, medium flow and high flow conditions) with n=5 individual replicates for each flow.

For linearly distributed metrics such as speed, run distance, and mean squared displacement (MSD) exponents, we conducted the following statistical analyses. We quantified the variability within each flow condition by computing the standard deviation (SD) of the time series for each individual replicate and then applied Levene's test Levene (1960) to evaluate differences in variance across flow conditions. We then computed the per-replicate medians and conducted non-parametric Kruskal–Wallis tests Kruskal and Wallis (1952) to assess whether the median values of these metrics differed between conditions. If a significant main effect emerged, we performed post-hoc pairwise comparisons using Bonferroni corrections to account for multiple testing. Lastly, to directly compare MSD scaling exponents at short versus long timescales within individuals, we employed the Wilcoxon signed-rank test Wilcoxon (1992), a non-parametric paired test appropriate for small sample sizes and non-normally distributed data. In

Fig. 1. (A) *R. oriander* is a millimeter scale member of the family *Veliidae*. (B) The natural habitat of *R. oriander* consists of shallow streams and creeks, where individuals are typically found near gently flowing water, along edges, and within small eddies or calm surface regions. (C) Environmental observation set up: 10"x10"x10" netted enclosure designed to be dropped over *R. oriander* in its natural habitat (D) Exemplar trajectories of *R. oriander* extracted from field videos show sequential linear motions and sharp turning behavior. (E) Run-and-tumble motion has been observed across a diverse set of taxa and lengthscales in nature, ranging from *E. coli* bacteria and *C. reinhardtii* at the micro-scale to *R. oriander* at the millimeter scale.

addition to reporting the *p*-values, we quantified effect sizes
using Cliff's delta Cliff (1993) (δ), providing a robust, non-
parametric measure of the magnitude of observed differences
between short- and long-time MSD exponents.
We applied the Hermans–Rasson Hermans and Rasson
(1985) test for each condition on the pooled heading data to
assess directional uniformity, which is sensitive to multimodal
deviations and well-suited for detecting non-uniform heading
distributions. Circular statistics were calculated using the
CircStat toolbox Berens (2009). Circular mean, variance and
skewness were calculated for each replicate in a flow condition
and the differences of these statistics across flow conditions
estimated using the Kruskal-Wallis test. Post-hoc tests were
only performed if there was significant difference. Together,

these analyses allowed us to evaluate both linear and directional
behavioral metrics as a function of flow condition, accounting
for variability within and across individuals.

Simulations

175 We simulate the insects' motion using a correlated random
176 walk model. The simulations were performed using MATLAB
177 programming language. The complete code for this simulation
178 is available on GitHub Tiwari et al. (2025).
179

Fig. 2. (A) Schematic of the laboratory observational setup of *R. oriander* locomotion. (B) Example snapshot of the experimental arena showing a superposition of insect trajectories. Dashed lines indicate the boundaries used for selecting trajectories included in subsequent analysis. Scale bar, 10 mm. (See SI video 1 for sample motion of *R. oriander*) (C) Instantaneous speed time series corresponding to the highlighted trajectory in panel B. Blue points indicate local speed peaks, which align with the blue markers on the trajectory shown in panel B. (D) Close-up trajectories of the mid-legs and body center as the *R. oriander* performs a sequence of "tumble–run–tumble" movements. Color gradients along each track represent the instantaneous speed of the corresponding limb. Violin plots showing the distribution of individual (E) median speeds and (F) median run distance across different flow conditions in the arena. Both the quantities were not found to be significantly different across flow conditions (Kruskal-Wallis test for Median speed p>0.1 and Median run distance p>0.4) White dot in the center indicates median of the N=5 replicates. Grey box shows the interquartile range. The whiskers are upper and lower adjacent values.

Fig. 3. (A) Mean squared displacement (MSD) as a function of lag time shows an early-time scaling exponent of approximately 2, indicative of ballistic motion, followed by a reduced exponent between 1 and 1.5 at later times, consistent with superdiffusive behavior. (B) Early-time exponents (α_{Early}) are significantly higher than late-time exponents (α_{Late}) across all flow conditions, indicating a shift in locomotor behavior at longer timescales. (Wilcoxon signed-rank tests; low flow: $p = 0.0625$, Cliff's $\delta = 0.84$; medium flow: $p = 0.0625$, Cliff's $\delta = 1.00$; high flow: $p = 0.0625$, Cliff's $\delta = 1.00$). White dot in the center indicates median of the N=5 replicates. Grey box shows the interquartile range. The whiskers are upper and lower adjacent values. (C) Polar histograms of *Rhagovelia oriander*'s turning angles show a strong preference for lateral left and right turns. (Herman-Rasson test for multimodality in circular distributions; low flow: $p < 0.0001$, $T = 9.34 \times 10^5$; medium flow: $p < 0.0001$, $T = 1.04 \times 10^6$; high flow: $p < 0.0001$, $T = 2.69 \times 10^7$) The angular distribution is consistent across all flow conditions when the N=5 replicate means of circular statistics were compared.(Kruskal-Wallis test applied: on mean vector length: $p = 0.932$; on circular variance: $p = 0.932$; on circular skewness: $p = 0.698$)

Results

Kinematics

We observe that *R. oriander* generates bursts of forward motion by actively rowing with its middle legs. Each stroke produces a sharp increase in speed (from $\sim$100mm/s to $\sim$300mm/s), marking the start of a propulsion phase. Immediately after, the insect enters a passive coasting phase during which it decelerates smoothly while maintaining a relatively straight trajectory. Following each run, *R. oriander* typically executes a sudden directional change or tumble. These tumble events occur at the end of the deceleration experienced during the insect's linear motion. The insect reorients its body, often by several tens of degrees during a tumble, in what appears to be a random direction. It then swiftly resumes motion by rowing with its mid legs to initiate a new run in the updated direction. This rapid ($\sim$5 rows per second), recurring pattern of propulsion followed by glide creates a characteristic speed-time series: sharp peaks corresponding to rowing events followed by gradual declines in speed. The profile presented in Figure 2C is consistent across individuals and flow conditions, reflecting the biomechanical rhythm underlying this insect's locomotion. Previous work Ortega-Jimenez et al. (2024)

have also reported that these insects are actively swimming in this fashion for most of their lifetimes. Figure 2D depicts a close-up visualization of this run-and-tumble behavior by overlaying the trajectory of the insect's center body along with the midleg segments, specifically the middle joints and terminal tarsi. The trails are color-coded by their respective instantaneous speed, revealing that peak body velocity occurs shortly after the tumble, as the insect initiates propulsion in the new direction. The figure indicates two distinct tumbles and one run, highlighting the sharp reorientation between runs and the biomechanical coordination involved in each transition. This visualization confirms that run-and-tumble behavior in *R. oriander* emerges from coordinated leg dynamics that alternate between propulsion and directional change.

R. oriander locomotion is independent of flow rate

To assess whether external fluid forces affect locomotor output, we quantified the distribution of insect speeds and run distances across three flow regimes. Figure 2E shows the distribution of median speeds for individual insects under low, medium, and high flow conditions. The violin plots reveal overlapping distributions, and statistical analysis using the Kruskal-Wallis

Fig. 4. (A, B) Sample turning angle distributions and simulated trajectories for $\Delta\phi = 0°, 90°,$ and $180°$, illustrating increasing angular reorientations. (C) Schematic of the simulation setup showing the vision triangle of the predator and prey drifting from left to right. (D) Heatmap of prey capture count as a function of Angular Separation ($\Delta\phi$) and prey speed (v_f), averaged over 10 simulations. Prey capture is maximized at intermediate $\Delta\phi$ values for slow prey, while faster prey favor sharper directional changes ($\Delta\phi \approx 130°$–$150°$), indicating an advantage of run-and-tumble-like strategies in dynamic environments.

test indicates that there is no significant difference between conditions ($p > 0.1$).

We similarly compared the distances covered during each run phase across flow conditions. As shown in Figure 2F, the distribution of run distances remains consistent, with a typical run length of approximately 40 mm across all tested flows. Once again, no significant differences were observed ($p > 0.4$), suggesting that *R. oriander* maintains its locomotor pattern regardless of ambient flow speed. These findings indicate that the insect's run-and-tumble strategy is robust against environmental variation within the tested flow range, enabling consistent movement performance across variable aquatic habitats.

R. oriander performs Run and Tumble locomotion

To quantitatively characterize the movement behavior of *R. oriander*, we calculate the mean squared displacement (MSD) as a function of time (Figure 3A). On a log-log scale, the MSD initially rises approximately t^2 at short timescales

($t < 0.25\,\mathrm{s}$), indicating ballistic motion. However, beyond 0.25 seconds, the MSD growth slows, transitioning into a superdiffusive regime, marked by exponents greater than 1 but less than 2. This shift represents a qualitative reduction in effective movement speed over longer durations. Importantly, the observed timescale for this transition closely corresponds to the average interval between consecutive rowing events (approximately 0.25 seconds).

To further validate this behavioral shift, we compare the MSD scaling exponents calculated at short and long timescales across all three flow conditions (Figure 3B). In each flow regime (low, medium, and high), the average MSD exponent consistently decreases from short to long timescales, indicating a steady transition from ballistic to super-diffusive behavior. Statistically significant support for this shift is shown by Wilcoxon signed-rank tests, nearing significance in every condition (low flow: $p = 0.0625$, Cliff's $\delta = 0.84$; medium flow: $p = 0.0625$, Cliff's $\delta = 1.00$; high flow: $p = 0.0625$, Cliff's $\delta = 1.00$). This evidence quantitatively confirms the dynamic locomotion patterns of the insect characterized by alternating run-and-tumble movement.

To confirm whether the observed slowdown involves directional changes characteristic of run-and-tumble behavior, we analyze the turning angles between consecutive runs (Figure 3C). A polar histogram (rose plot) of turning angles reveals a strong preference for sharp directional reorientations, with approximately equal probabilities of turning left or right. To quantitatively test this multimodal distribution, we apply the Hermans–Rasson test for circular uniformity. The test strongly rejects uniformity for pooled turning angle data under all flow conditions (low flow: $p < 0.0001$, $T = 9.34 \times 10^5$; medium flow: $p < 0.0001$, $T = 1.04 \times 10^6$; high flow: $p < 0.0001$, $T = 2.69 \times 10^7$), confirming a significant tendency for lateral directional changes. Furthermore, we test whether the turning angle distributions vary across flow regimes by calculating circular statistics (mean vector length R, circular variance, and circular skewness) per insect and applying the Kruskal–Wallis test. No significant differences emerge among flow conditions (mean vector length: $p = 0.932$; circular variance: $p = 0.932$; circular skewness: $p = 0.698$). Together, these analyses strengthen the characterization of $R.\ oriander$'s locomotion as consistent run-and-tumble behavior, defined by periodic runs punctuated by distinct lateral reorientations.

Run-and-tumble improves prey capture when prey are in motion

Having established that $R.\ oriander$ employs a run-and-tumble locomotion strategy, we now investigate the potential ecological advantages of this behavior. To address this, we develop a numerical simulation that mimics the insect's locomotion dynamics and prey-searching behavior.

The simulated $R.\ oriander$ moves at speeds drawn from a Gaussian distribution based on experimental observations and executes runs of approximately uniform duration. After each run, the simulated insect experiences directional changes modeled by a superposition of two circular Gaussian distributions with means separated by an adjustable parameter $\Delta\phi$ (Figure 4A,B). Specifically, a value of $\Delta\phi = 0°$ corresponds to predominantly straight movement, while $\Delta\phi = 180°$ indicates frequent sharp turns that are equally likely to be left or right, with some finite variance around $\pm90°$. Characteristic turning angle distributions and representative trajectories for $\Delta\phi = 0°, 90°$, and $180°$ are shown in Figure 4A,B.

In the simulation environment, prey particles passively drift from left to right within a square domain of side length L under periodic boundary conditions, moving at a constant horizontal speed v_f (Figure 4C). This prey speed v_f serves as a second adjustable parameter in our model. The simulated predator is equipped with a forward-directed "vision triangle," defined as an isosceles triangle with a vertex angle of 45 degrees and side lengths of 40 mm, reflecting the typical run length observed experimentally. Prey are considered captured when they enter this triangular visual region.

We explore the effects of turning sharpness $\Delta\phi$ (0–180 degrees) and prey speed v_f (0–200 mm/s) on prey capture success, quantifying the number of prey captured over fixed-duration simulations. Each parameter combination is evaluated across ten replicate simulations to ensure robust statistical characterization of model performance.

We visualize the combined effect of turning sharpness ($\Delta\phi$) and prey speed (v_f) on prey capture success using a two-dimensional heatmap (Figure 4D). The x-axis represents angular separation $\Delta\phi$, the y-axis corresponds to prey speed v_f, and the color encodes the total prey captures during the simulations. At low prey speeds, capture rates show a clear unimodal relationship with $\Delta\phi$, peaking around $\Delta\phi \approx 100°$. Notably, at these low speeds, extreme directional changes ($\Delta\phi \approx 180°$) result in the lowest prey capture rates, performing even worse than straight-line movements ($\Delta\phi \approx 0°$). As prey speed increases, however, this peak in prey capture shifts toward higher angular separations ($\Delta\phi \approx 130°$–$150°$), suggesting that more pronounced lateral turning becomes increasingly advantageous as prey move faster.

These results indicate that while moderate lateral turning optimizes prey capture when prey move slowly or remain nearly stationary, sharper directional changes become increasingly beneficial as prey speed increases. Therefore, the run-and-tumble strategy employed by $R.\ oriander$, characterized by frequent lateral turns, likely represents an adaptive response that maximizes prey encounters in environments with flowing substrates.

Discussion

In this study, we document and quantitatively characterize the run-and-tumble locomotion exhibited by $R.\ oriander$, an interface-dwelling insect navigating dynamic, fluid environments. Our observations demonstrate that $R.\ oriander$ employs repeated cycles of rapid directional propulsion (runs) interspersed with sharp lateral reorientations (tumbles), a behavioral pattern well-documented in microbial taxa such as $E.\ coli$ and $C.\ reinhardtii$, but seldom observed at macroscopic scales. Through controlled laboratory experiments and computational simulations, we reveal that this distinctive locomotion pattern provides significant ecological advantages by optimizing prey encounters, particularly when prey are found on a moving or flowing substrate.

The behavior of $R.\ oriander$ is a compelling example of behavioral convergence. Despite significant morphological and ecological differences, diverse taxa ranging from microscopic bacteria and algae to interfacial insects at millimeter scales independently evolve similar locomotion strategies to efficiently search for resources or more favorable environments. The consistent run-and-tumble movement observed across three orders of magnitude in length highlights its importance

as a search strategy, especially in dynamic or uncertain environments where continuous linear motion is less effective.

Run-and-tumble locomotion in *R. oriander* persisted across ambient flow conditions found in its natural environment. Our research shows that essential locomotion parameters, including speed and run length, remain largely unchanged across the ecologically relevant flow regimes we explored. This stability likely allows *R. oriander* to take advantage of different microhabitats near gentle stream edges and within small eddies or surface areas, areas where prey availability and fluid dynamics can vary unpredictably. Such resilience in behavioral strategies is important for organisms living in fluid environments marked by varying external conditions.

From a broader ecological perspective, the run-and-tumble strategy observed here serves as a fascinating evolutionary solution for effective spatial exploration and resource gathering. Previous theoretical and empirical research has recognized search patterns such as Lévy flights and correlated random walks as advantageous in specific scenarios. Our research contributes to this ongoing conversation by showcasing run-and-tumble locomotion in a neustonic environment, which is particularly suited to the dynamic and complex conditions faced by mobile organisms.

Beyond fundamental ecological insights, the behavioral mechanisms of run-and-tumble locomotion documented in *R. oriander* hold significant potential for biomimetic technological applications. Specifically, the observed locomotion patterns can inspire the design of search algorithms for autonomous surface robots. Such biomimetic algorithms could be particularly valuable in developing mobile swarms Rubenstein et al. (2014) for applications including oceanic plastic debris capture, oil spill mitigation, and search-and-rescue operations Murphy (2007). Robots operating at fluid interfaces must efficiently explore large spatial areas while adapting to complex surface flows and unpredictable targets, conditions analogous to those encountered by interfacial insects. Therefore, embedding principles derived from *R. oriander*'s run-and-tumble locomotion into algorithmic control frameworks could lead to enhanced operational efficiency, robustness, and adaptability of autonomous robotic systems.

In conclusion, we recognize several limitations within our study. Although laboratory experiments are systematic and controlled, they might not completely reflect the intricacies of natural habitats. Moreover, our computational model streamlines prey dynamics and predator sensory systems, presenting room to integrate more realistic ecological interactions and environmental variability in future simulations. Despite these limitations, our findings lay a strong groundwork for further investigation into the influence of environmental complexity on behavioral evolution and offer promising avenues for bioinspired innovations in robotics and computational algorithms.

Supplementary data

Supplementary data available at *ICB* online.

Competing interests

There is NO Competing Interest.

Author contributions statement

I.T., J.S.H. and S.B. conceptualized the research. I.T., N.N. and J.S.H. designed the experiments. N.N. conducted the experiments, for which I.T. performed the analysis. I.T. did the simulations. S.B. supervised the research. All authors contributed to writing, discussion, and revising the manuscript.

Acknowledgments

S.B. acknowledges funding support from NIH Grant R35GM142588; NSF Grants MCB-1817334; CMMI-2218382; CAREER IOS-1941933; PHY-2310691 and the Open Philanthropy Project. Text in this paper was revised using LLMs.

References

Bell, W. J. (2012). *Searching behaviour: the behavioural ecology of finding resources.* Springer Science & Business Media.

Berens, P. (2009). Circstat: a matlab toolbox for circular statistics. *Journal of statistical software*, 31:1–21.

Berg, H. C. and Brown, D. A. (1972). Chemotaxis in escherichia coli analysed by three-dimensional tracking. *nature*, 239(5374):500–504.

Cliff, N. (1993). Dominance statistics: Ordinal analyses to answer ordinal questions. *Psychological bulletin*, 114(3):494.

Gomez-Marin, A., Stephens, G. J., and Louis, M. (2011). Active sampling and decision making in drosophila chemotaxis. *Nature communications*, 2(1):441.

Hermans, M. and Rasson, J. (1985). A new sobolev test for uniformity on the circle. *Biometrika*, 72(3):698–702.

Ishihara, A., Segall, J. E., Block, S. M., and Berg, H. C. (1983). Coordination of flagella on filamentous cells of escherichia coli. *Journal of Bacteriology*, 155(1):228–237.

Johnson, S., Freedman, B., and Tang, J. X. (2024). Run-and-tumble kinematics of enterobacter sp. sm3. *Physical Review E*, 109(6):064402.

Kareiva, P. and Shigesada, N. (1983). Analyzing insect movement as a correlated random walk. *Oecologia*, 56:234–238.

Kruskal, W. H. and Wallis, W. A. (1952). Use of ranks in one-criterion variance analysis. *Journal of the American statistical Association*, 47(260):583–621.

Levene, H. (1960). Robust tests for equality of variances. *Contributions to probability and statistics*, pages 278–292.

Murphy, R. R. (2007). Search and rescue robotics. *Springer handbook of robotics*, pages 1151–1173.

Nath, T., Mathis, A., Chen, A. C., Patel, A., Bethge, M., and Mathis, M. W. (2019). Using deeplabcut for 3d markerless pose estimation across species and behaviors. *Nature protocols*, 14(7):2152–2176.

Ortega-Jimenez, V. M., Kim, D., Kumar, S., Kim, C., Koh, J.-S., and Bhamla, S. (2024). Ultrafast elastocapillary fans control agile maneuvering in ripple bugs and robots. *bioRxiv*, pages 2024–11.

Othmer, H. G., Dunbar, S. R., and Alt, W. (1988). Models of dispersal in biological systems. *Journal of mathematical biology*, 26(3):263–298.

Patteson, A. E., Gopinath, A., Goulian, M., and Arratia, P. E. (2015). Running and tumbling with e. coli in polymeric solutions. *Scientific reports*, 5(1):15761.

Polin, M., Tuval, I., Drescher, K., Gollub, J. P., and Goldstein, R. E. (2009). Chlamydomonas swims with two "gears" in a

eukaryotic version of run-and-tumble locomotion. *Science*, 325(5939):487–490.

Pyke, G. H. (2015). Understanding movements of organisms: it's time to abandon the lévy foraging hypothesis. *Methods in Ecology and Evolution*, 6(1):1–16.

Rubenstein, M., Cornejo, A., and Nagpal, R. (2014). Programmable self-assembly in a thousand-robot swarm. *Science*, 345(6198):795–799.

Santos, M. E., Le Bouquin, A., Crumière, A. J., and Khila, A. (2017). Taxon-restricted genes at the origin of a novel trait allowing access to a new environment. *Science*, 358(6361):386–390.

Smith, J. N. (1974). The food searching behaviour of two european thrushes. *Behaviour*, 48(1-4):276–301.

Tiwari, I., Nagappan, N., Harrison, J. S., and Bhamla, S. (2025). Rhagovelia run and tumble icb 2025 paper. `https://github.com/bhamlalab/Rhagovelia-run-and-tumble-ICB-2025`. Accessed: 2025-04-02.

Viswanathan, G. M., Afanasyev, V., Buldyrev, S. V., Murphy, E. J., Prince, P. A., and Stanley, H. E. (1996). Lévy flight search patterns of wandering albatrosses. *Nature*, 381(6581):413–415.

Wilcoxon, F. (1992). Individual comparisons by ranking methods. In *Breakthroughs in statistics: Methodology and distribution*, pages 196–202. Springer.