Title: Simultaneous knockout of VITAMIN C DEFECTIVE 2 and 3 exacerbates ascorbate

deficiency and light stress sensitivity

Running title: Functional and genetic relationship between VTC2 and VTC3

Authors: Yasuhiro Tanaka[1,2,†], Takanori Maruta[1,2,3,4,†,*], Koki Arima[3], Hiroki Nakayama[3], and

Takahiro Ishikawa[1,2,3,4]

Affiliations:

[1] Graduate School of Natural Science and Technology, Shimane University, 1060 Nishikawatsu,

Matsue, Shimane 690-8504, Japan

[2] Bioresource and Life Sciences, The United Graduate School of Agricultural Sciences, Tottori

University, 4-101 Koyama-Minami, Tottori, Tottori 680-8553, Japan

[3] Department of Life Sciences, Faculty of Life and Environmental Science, Shimane University,

1060 Nishikawatsu, Matsue, Shimane 690-8504, Japan

[4] Institute of Agricultural and Life Sciences, Academic Assembly, Shimane University, 1060

Nishikawatsu, Matsue, Shimane 690-8504, Japan

[†] These authors contributed equally to the article.

[*] Corresponding author

E-mail addresses: Yasuhiro Tanaka (yt.y.kk.s6@gmail.com), Takanori Maruta

(maruta@life.shimane-u.ac.jp), Koki Arima (kouchan506@gmail.com), Hiroki Nakayama

(umikihiroki@icloud.com), Takahiro Ishikawa (ishikawa@life.shimane-u.ac.jp)

The date of submission: April 4, 2025

Figures and Tables: Eight figures

Supplementary data: One supplementary figure and two supplementary tabels

The word count: 4,248 words

Highlight

VTC2 and VTC3 are crucial for activating ascorbate biosynthesis. Their simultaneous knockout further exacerbates ascorbate deficiency under low- and high-light conditions, causing severe photooxidative damage.

Abstract

Light-dependent activation of ascorbate biosynthesis is essential for accumulating ascorbate to mitigate photooxidative stress. The *VTC2* and *VTC3* genes play key roles in this process: *VTC2* encodes GDP-L-galactose phosphorylase (GGP), the rate-limiting enzyme in ascorbate biosynthesis, and *VTC3* encodes a putative protein kinase/phosphatase with an unknown function. Here, we investigated their functional and genetic relationship. In *vtc3* mutants, *VTC2* transcription and GGP activity were slightly enhanced, suggesting that VTC3 is not required for VTC2 expression. Additionally, the *vtc3* mutation had negligible effects on the transcriptome and the activity of enzymes involved in ascorbate redox cycle regulation, implying that VTC3 regulates ascorbate biosynthesis at a step other than VTC2. Under low-light conditions, ascorbate levels were lower in *vtc2* than in *vtc3* mutants, but *vtc2* retained the ability to increase ascorbate levels under high-light stress, unlike *vtc3*. The simultaneous knockout of VTC2 and VTC3 further reduced ascorbate levels compared to single mutants and severely impaired light stress-induced ascorbate accumulation, leading to pronounced photooxidative damage. These findings highlight the additive effects of VTC2 and VTC3 on ascorbate biosynthesis and stress tolerance. The *vtc2 vtc3* double mutants provide a valuable model for further elucidating the physiological roles of ascorbate in plants.

Keywords: Arabidopsis thaliana, ascorbate biosynthesis, GDP-L-galactose phosphorylase, high light, oxidative stress, vitamin C-defective

Abbreviations: APX, ascorbate peroxidase; ASC, ascorbate; DHA, dehydroascorbate; DHAR, dehydroascorbate reductase; GGP, GDP-L-galactose phosphorylase; GME, GDP-D-mannose-3',5'-epimerase; GMP, GDP-D-mannose pyrophosphorylase; GPP, L-galactose-1-phosphate phosphatase; GSH, reduced glutathione; L-GalDH, L-galactose dehydrogenase; L-GalLDH, L-galactono-1,4-lactone dehydrogenase; MDHA, monodehydroascorbate; MDAR, monodehydroascorbate reductase; NPQ, non-photochemical quenching; PMI, phosphomannose isomerase; PMM, phosphomannomutase; PP2C, protein phosphatase 2C; ROS, reactive oxygen species; uORF, upstream open reading frame; VTC, vitamin c defective

Introduction

In plants, reactive oxygen species (ROS) can be generated as byproducts of photosynthesis (Asada, 1999). While ROS are highly reactive and can damage cellular components at elevated levels, they also serve as critical signaling molecules in plant stress acclimation (Mittler *et al.*, 2022). Maintaining ROS homeostasis is essential for harnessing their signaling functions without compromising cellular integrity. This balance is achieved through robust antioxidant systems, among which ascorbate is one of the most abundant soluble antioxidants in plant leaves, present in all intracellular compartments at millimolar (mM) concentrations (Zechmann *et al.*, 2011; Smirnoff, 2018; Maruta *et al.*, 2024). Ascorbate plays a central role in scavenging ROS through both direct nonenzymatic reactions and enzymatic pathways mediated by ascorbate peroxidase (APX), an enzyme specific to eukaryotic photosynthetic organisms (Shigeoka *et al.*, 2002; Maruta *et al.*, 2016). Additionally, ascorbate serves as a substrate for violaxanthin de-epoxidase, an enzyme crucial for the xanthophyll cycle, which dissipates excess light energy as heat and limits ROS production during photosynthesis (Müller-Moulé *et al.*, 2004). Beyond its role in ROS management, ascorbate is involved in diverse biological processes, including photosynthetic electron transport, iron uptake, cell division, anthocyanin accumulation, and hormone biosynthesis (Smirnoff, 2018).

Several pathways for ascorbate biosynthesis have been proposed; however, the D-mannose/L-galactose (D-Man/L-Gal) pathway, also known as the Smirnoff–Wheeler pathway, is widely recognized as the primary route (Wheeler *et al.*, 1998; Smirnoff, 2018; Maruta, 2022). This notion is supported by substantial experimental evidence demonstrating its involvement in ascorbate biosynthesis across land plants and algae (Wheeler *et al.*, 1998; Conklin *et al.*, 1999; Vidal-Meireles *et al.*, 2017; Sodeyama *et al.*, 2021; Ishida *et al.*, 2024). The D-Man/L-Gal pathway synthesizes ascorbate from the hexose phosphate pool through eight enzymatic reactions catalyzed sequentially by phosphomannose isomerase (PMI), phosphomannomutase (PMM), GDP-D-mannose pyrophosphorylase (GMP), GDP-D-mannose-3',5'-epimerase (GME), GDP-L-galactose phosphorylase (GGP), L-galactose-1-phosphate phosphatase (GPP), L-galactose dehydrogenase (L-GalDH), and L-galactono-1,4-lactone dehydrogenase (L-GalLDH) (Smirnoff, 2018). The final step, catalyzed by L-GalLDH, occurs in the inner mitochondrial membrane, whereas all other enzymatic steps take place in the cytosol (Smirnoff, 2018). The significance of the D-Man/L-Gal pathway has been confirmed mainly through the characterization of Arabidopsis vitamin C-defective (*vtc*) mutants. The genes *VTC1*, *VTC2*, and *VTC4* encode GMP, GGP, and GPP, respectively (Conklin *et al.*, 1999, 2006; Jander *et al.*, 2002; Laing *et al.*, 2007; Linster *et al.*, 2007). Arabidopsis also possesses a minor isoform of GGP, encoded by *VTC5*. Notably, a double knockout mutant of

VTC2 and *VTC5* results in undetectable ascorbate levels and exhibits seedling lethality unless

supplemented with ascorbate or its precursors (Dowdle *et al.*, 2007).

Light is a potent stimulus for increasing the ascorbate pool size in plants. Leaf ascorbate

content rises in response to light intensity and exposure duration (Dowdle *et al.*, 2007; Yabuta *et al.*,

2007). The key enzyme regulating ascorbate content under light conditions is VTC2/GGP, whose

transcription is induced by light (Dowdle *et al.*, 2007; Yabuta *et al.*, 2007). Interestingly, the

light-dependent activation of *VTC2* transcription is suppressed by photosynthesis inhibitors,

suggesting a role for photosynthesis in the regulation of ascorbate biosynthesis (Yabuta *et al.*, 2007).

GGP activity is significantly enhanced under high-light stress, whereas the activities of other

biosynthetic enzymes remain unaffected (Dowdle *et al.*, 2007). Moreover, overexpression of VTC2

effectively enhances ascorbate pool sizes in various plant species (Bulley *et al.*, 2009, 2012;

Yoshimura *et al.*, 2014). These findings confirm that VTC2/GGP is the rate-limiting enzyme in the

D-Man/L-Gal pathway, a conclusion further supported by computational simulations (Fenech *et al.*,

2021). The regulation of VTC2 by light also occurs at the post-translational level. In the dark, GGP

activity is inhibited by a new class of blue-light receptors, PAS/LOV proteins, which directly

associate with GGP (Ogura *et al.*, 2008; Aarabi *et al.*, 2023; Bournonville *et al.*, 2023). Upon

exposure to blue light, PAS/LOV proteins are dissociated from GGP, promoting its enzymatic

activity. Thus, light-dependent activation of GGP plays a pivotal role in ascorbate biosynthesis.

Recent studies have proposed that the biosynthetic enzymes in the cytosol form a protein complex

(Fenech *et al.*, 2021), potentially contributing to the regulation of ascorbate biosynthesis.

Additionally, ascorbate itself tightly regulates its biosynthesis. High ascorbate levels repress VTC2

translation via an upstream open reading frame (uORF) in the 5′-untranslated region of the gene,

counterbalancing the light-induced activation of GGP (Laing *et al.*, 2015). Consequently, ascorbate

biosynthesis is finely tuned through transcriptional, translational, and post-translational regulation.

However, the precise mechanisms underlying light-dependent GGP activation remain to be

elucidated.

VTC3, the gene responsible for the *vtc3* mutant, encodes a unique protein containing both

serine/threonine protein kinase and protein phosphatase 2C (PP2C) domains (Conklin *et al.*, 2000,

2013). Two distinct *vtc3* alleles have been identified: *vtc3-1*, which carries a G202E missense

mutation in the kinase domain, and *vtc3-2*, which has a nonsense mutation in the PP2C domain

(Conklin *et al.*, 2013). Both alleles exhibit a moderate ascorbate-deficient phenotype, with

approximately 50% of wild-type ascorbate levels, a deficiency less severe than that observed in *vtc2*

mutants (20–30% of wild-type levels). VTC3 is highly conserved across land plants and algae

(Conklin *et al.*, 2013; Maruta *et al.*, 2024), and its loss also results in ascorbate deficiency in the

moss *Physcomitrella patens* (Sodeyama *et al.*, 2021). Notably, light-induced ascorbate accumulation

is significantly impaired in Arabidopsis *vtc3* mutants (Conklin *et al.*, 2013), suggesting that VTC3

plays a crucial role in the light regulation of VTC2/GGP. Since VTC3 has been reported to be a

chloroplast protein, it is unlikely to regulate ascorbate biosynthesis through direct

(de)phosphorylation of biosynthetic enzymes. The precise physiological function of VTC3 remains

unclear, and further studies are required to elucidate its role.

The size of the ascorbate pool is regulated not only by biosynthesis but also by its

oxidation, recycling, and degradation. Reduced ascorbate (ASC) undergoes oxidation through APX

or direct interaction with ROS, generating monodehydroascorbate (MDHA) (Asada, 1999; Smirnoff,

2018). MDHA radicals are disproportionated into ASC and dehydroascorbate (DHA). MDHA and

DHA are recycled back to ASC by monodehydroascorbate reductase (MDAR) and dehydroascorbate

reductase (DHAR), respectively (Foyer and Halliwell, 1977; Hossain and Asada, 1985). MDAR uses

NAD(P)H, while DHAR relies on reduced glutathione (GSH) as the electron donor. Additionally,

GSH can nonenzymatically reduce DHA to ASC (Winkler *et al.*, 1994; Terai *et al.*, 2020; Hamada *et*

al., 2023). However, DHA is highly unstable and undergoes irreversible degradation if not rapidly

reduced. These interconnected processes are essential for maintaining ascorbate homeostasis.

This study aims to achieve two primary objectives. First, we seek to elucidate the

potential involvement of VTC3 in the light-dependent regulation of VTC2/GGP. Considering that

vtc3 mutants exhibit impaired light-induced ascorbate accumulation (Conklin *et al.*, 2013), VTC3

may play a role in facilitating the activation of VTC2/GGP under light conditions. Second, we aim to

investigate the impact of simultaneously knocking out VTC2 and VTC3 on ascorbate metabolism

and stress tolerance. The *vtc2* mutant has significantly lower ascorbate levels than the *vtc3* mutant,

yet it retains the ability to increase ascorbate accumulation under light stress (Hamada *et al.*, 2023;

Terai *et al.*, 2020), likely due to the presence of VTC5, a minor isoform of VTC2. In contrast, *vtc3*

mutants exhibit defective light-induced ascorbate accumulation despite having higher basal

ascorbate levels than *vtc2* mutants (Conklin *et al.*, 2013). If both VTC2 and VTC3 are absent, will

ascorbate levels remain severely depleted even under light stress? By analyzing single and double

mutants of *vtc2* and *vtc3*, we aim to uncover how these genes collectively regulate ascorbate

homeostasis and influence plant tolerance to oxidative stress.

Materials and methods

Plant Materials and Growth Conditions

Arabidopsis thaliana ecotype Col-0 was used as the wild type. The T-DNA insertion lines *vtc2-4*

(SAIL_769_H05) and *vtc3-3* (SALK_095743) were obtained from the Arabidopsis Biological

Resource Center and used for further crossings. Unless otherwise specified, wild-type and mutant

seeds were sown in Jiffy-7 peat pellets and stratified in the dark at 4°C for 2–3 days. Plants were

grown in a growth chamber (KCLP-1000CCFL-6-8L-CO□, NK System) at 22°C under a 16-hour

photoperiod with low light intensity (40–60 µmol photons m^{-2} s^{-1}) and at 20°C in darkness for 8

hours. At three weeks of age, plants were exposed to moderate- or high-light stress (750 or 1,500

µmol photons m^{-2} s^{-1}) using an LED lighting unit (PFQ-600PT, NK System) without a preceding

dark period. The LED system minimized heat production, maintaining leaf temperature at

approximately 24–25°C. To ensure uniform light exposure, plants were placed on a turntable beneath

the LED panel. Fully expanded rosette leaves were collected from at least three plants and treated as

a single biological replicate. Rosette and lesion areas (percentage of the total rosette area) were

quantified using ImageJ software.

177   For plate-grown experiments (**Figures 3 and 4**), wild-type and mutant plants were

cultivated on half-strength Murashige and Skoog (MS) medium supplemented with 1% sucrose

under a 16-hour photoperiod at 100 µmol photons m^{-2} s^{-1} and 22°C, with an 8-hour dark period at

20°C (LH-240S, NK System). Plants were then subjected to moderate-light stress (650 µmol

photons m^{-2} s^{-1}) without an intermediate dark period using the same LED panel and turntable system.

Shoots from at least ten plants were pooled and treated as a single biological replicate.

Semi-quantitative and quantitative reverse transcription-polymerase chain reaction

Semi-quantitative reverse transcription-polymerase chain reaction (RT-PCR) was performed

according to Terai *et al*. (2020). Quantitative RT-PCR (qPCR) was performed as described by

Kameoka *et al*. (2021). All the primers used are listed in **Supplementary Table S1**.

Western blotting

To examine VTC2 protein expression, we attempted to generate two independent anti-VTC2 peptide

antibodies. The peptide sequences used were "NH₂-C+QNSSGNVNQKSNRT-COOH" and

"NH₂-C+KRKEDYEGASEDNA-COOH". The design and synthesis of the antigenic peptides, as

well as antibody production, were outsourced to Eurofins Genomics. In addition, we purchased a

commercial anti-GDP-L-galactose phosphorylase antibody (AS10 723, Agrisera). Using these

antibodies, we performed western blotting following the method described by Kameoka *et al*. (2021),

but we were unable to detect the VTC2 protein.

Measurements of Antioxidants and Enzyme Activity

Ascorbate content was quantified using an ultra-fast liquid chromatography (UFLC) system equipped with a C18 column, following the method of Hamada and Maruta (2024). Glutathione levels were measured according to Noctor *et al.* (2016) without modifications. Soluble and membrane-bound APX activity was assessed using the method described by Kameoka *et al.* (2021), while DHAR and MDAR activities were measured as outlined by Tanaka *et al.* (2021).

GGP activity was measured according to Hamada *et al.*, (2023). *Arabidopsis* leaves (0.2 g) frozen in liquid nitrogen were ground and homogenized in 1 mL of extraction buffer containing 50 mM Tris–HCl (pH 7.5), 10% glycerol, 2 mM DTT, 10 mM $MgCl_2$, 1 mM aminocaproic acid, 1 mM benzamidine, 1 mM phenylmethanesulfonylfluoride. After centrifugation ($15,300 \times g$) for 20 min at 4 °C, the supernatant was concentrated by ultrafiltration ($15,300 \times g$, 30 min, 4 °C) using an Amicon ultra centrifugal filter (10 kDa, NMWL) and used for the following assay. The extract (20 µL) was added to 80 µL of a reaction mixture containing 50 mM Tris–HCl (pH 7.5), 2 mM $MgCl_2$, 10 mM KCl, 1mM dithiothreitol, and 100 µM GDP-D-glucose with or without 5 mM sodium phosphate and incubated for 1 h at 26 °C. The reaction was stopped by boiling for 5 min, and the mixture was passed through 0.22 µm filters (Merck Millipore). The samples were injected into an ultra-fast liquid chromatography (UFLC) system (Prominence UFLC, Shimadzu) equipped with a C-18 column (LUNA C18(2), Column 150 x 4.6 nm, Shimadzu). The mobile phase consisted of 50 mM phosphate buffer (pH 6.7), 5 mM tetrabutylammonium dihydrogen phosphate, 5 mg L^{-1} EDTA, and 3.5% methanol, and the flow rate was 1 mL min^{-1}. GDP-D-glucose was detected at 254 nm using the SPD-20A UV-Vis detector (Shimadzu). Pi-independent degradation was subtracted from the Pi-dependent decrease in the substrate to calculate GDP-D-glucose degradation activity.

RNA-Seq Analysis

Two-week-old wild-type, *vtc2-4*, and *vtc3-3* seedlings, grown on half-strength MS medium supplemented with 1% sucrose under the same conditions described above, were incubated in darkness for two days, followed by a six-hour exposure to growth light (100 µmol photons m^{-2} s^{-1}). Total RNA was extracted from shoot tissues using RNAiso Plus (Takara, Japan) and purified with NucleoSpin RNA Plus (Takara). RNA was fragmented and converted to double-stranded DNA using the TruSeq RNA Sample Preparation Kit v2 (Illumina). Library quality and quantity were assessed using a Bioanalyzer (Agilent) and a KAPA Library Quantification Kit (KAPA Biosystems), respectively. Sequencing was performed using an Illumina MiSeq with the MiSeq Reagent Kit v3

(300 cycles; Illumina). RNA-seq data were obtained from triplicate biological samples (n = 3 per

group) and statistically analyzed. Read mapping, data processing, and statistical analyses were

conducted using the CLC Genomics Workbench (version 9.5.2). The gene expression dataset has

been deposited in the Gene Expression Omnibus under accession number GSE293768. Gene

ontology (GO) enrichment analysis was performed using agriGO (version 2; Tian *et al.*, 2017).

Chlorophyll Fluorescence Measurements

Chlorophyll fluorescence measurements were performed using a Closed FluorCam 800MF (Photon

Systems Instruments). Plants were dark-adapted for 30 minutes before measurements. Chlorophyll

fluorescence parameters, F_v/F_m and NPQ, were calculated according to Baker (2008) using the

following equations: $F_v/F_m = (F_m - F_o)/F_m$, and $NPQ = (F_m - F_m\square)/F_m\square$. NPQ was measured under

actinic light at an intensity of 90 µmol photons $m^{-2}\ s^{-1}$.

Accession Numbers

VTC2, At4g26850; VTC3, At4g39830.

Data Analysis

The statistical methods used are described in the respective figure legends (Tukey–Kramer,

Dunnett's test, or Student's *t*-test). Statistical calculations were performed using at least three

independent biological replicates (see figure legends for details).

Results

VTC3 is required for ascorbate accumulation under high-light stress

To evaluate the role of VTC3 in ascorbate accumulation under high-light conditions, we used a

T-DNA insertion mutant for the *VTC3* gene, referred to hereafter as *vtc3-3*. This mutant has been

previously characterized as a knockout line (Conklin *et al.*, 2013), lacking a full-length *VTC3*

transcript (see **Supplementary Figure S1**). Additionally, the *vtc2-4* mutant (Lim *et al.*, 2016) was

included in the experiments. Under low-light conditions, *vtc2-4* and *vtc3-3* mutants exhibited

approximately 26% and 45% of wild-type ascorbate levels, respectively (**Figure 1**), consistent with

previous reports (Conklin *et al.*, 2013; Lim *et al.*, 2016). Exposure to high-light stress induced a

time-dependent increase in ascorbate content in wild-type plants. In contrast, the *vtc3-3* mutant

displayed only a minimal and statistically insignificant increase in ascorbate levels. The ascorbate

pool size in *vtc2-4* remained significantly lower than in wild-type plants under both low-light and

high-light conditions. However, high-light stress markedly increased ascorbate levels in *vtc2-4*

(**Figure 1**), corroborating previous findings (Hamada *et al.*, 2023; Terai *et al*., 2020). The elevated

amounts of ascorbate after 48 hours of high-light exposure were approximately 5.5 µmol g^{-1} FW in

the wild type, 3.4 µmol g^{-1} FW in *vtc2-4*, and 0.7 µmol g^{-1} FW in *vtc3-3* (**Figure 1**). Neither the

vtc2-4 nor *vtc3-3* mutation affected the ascorbate redox state under either low- or high-light

conditions. These results demonstrate that VTC3 is essential for the accumulation of ascorbate in

response to high-light exposure.

Loss of VTC3 does not limit GDP-L-galactose phosphorylase activity

Since high-light stress induces ascorbate accumulation through GGP activation (Dowdle *et al.*, 2007),

we hypothesized that VTC3 might regulate GGP. To test this, we analyzed the transcript levels of

VTC2 and *VTC5* before and after 24 hours of high-light exposure. In *vtc2-4* mutants, *VTC2*

transcripts were nearly undetectable (**Figure 2**). In wild-type plants, the transcript levels of both

VTC2 and *VTC5* showed a modest increase in response to high-light stress. Intriguingly, these

increases were more pronounced in *vtc3-3* mutants. This upregulation of *VTC2* and *VTC5* in *vtc3-3*

mutants may represent a compensatory response to ascorbate deficiency. However, no similar

increase in *VTC5* expression was observed in *vtc2-4* mutants after high-light exposure (**Figure 2**),

possibly because *vtc2-4* mutants retained higher ascorbate levels than *vtc3-3* mutants under

high-light conditions (**Figure 1**).

To assess VTC2 protein levels, we obtained three anti-VTC2 antibodies targeting

different VTC2 peptides. Unfortunately, none of these antibodies successfully detected VTC2 via

western blotting, likely due to low antibody specificity or the very low abundance of VTC2 protein.

As an alternative, we measured GGP activity using a Pi-dependent GDP-D-glucose degradation

assay, a reliable indicator of endogenous GGP activity (Hamada *et al.*, 2023). In wild-type plants,

GGP activity increased in response to high-light exposure, whereas in *vtc2-4* mutants, activity was

nearly undetectable (**Figure 2**). Notably, the absence of VTC3 did not reduce GGP activity; rather, it

appeared to enhance it, consistent with the observed increase in *VTC2* and *VTC5* transcript levels in

vtc3-3 mutants. Contrary to our hypothesis, these results indicate that VTC3 is not essential for the

expression and activity of GGP.

Effects of VTC3 on ascorbate redox cycle regulation and transcriptome

Because of the largely unknown physiological function of VTC3, we hypothesized that it may

contribute to ascorbate redox cycling rather than biosynthesis. To test this, we examined the impact

of VTC3 on glutathione levels and the activities of APX, DHAR, and MDAR. Plants grown under
plate conditions were used for these experiments. Two-week-old wild-type, *vtc2-4*, and *vtc3-3* plants
on agar plates were exposed to moderate-light stress (650 µmol photons m^{-2} s^{-1}; ML^{650}) for 24
hours. This stress increased ascorbate levels in both wild-type and *vtc2-4* plants, but this response
was nearly absent in *vtc3-3* mutants, consistent with the data obtained from *in vivo* soil experiments
(**Figure 3**).

The size of the glutathione pool increased under ML^{650} stress in all genotypes, with no
significant differences observed among the wild type, *vtc2-4*, and *vtc3-3*. Additionally, APX, DHAR,
and MDAR activities were upregulated under these conditions, but neither *vtc2-4* nor *vtc3-3*
mutations influenced their activities under either low-light or stress conditions (**Figure 3**). These
findings suggest that VTC3 does not significantly affect ascorbate redox cycling.

To further investigate the function of VTC3, we performed transcriptome analysis. For
this experiment, a drastic light/dark shift was used instead of light stress to induce a stronger
response. Two-week-old wild-type, *vtc2-4*, and *vtc3-3* plants grown on agar plates under low-light
conditions were incubated in darkness for two days, followed by a 6-hour exposure to growth light
(100 µmol photons m^{-2} s^{-1}). Differentially expressed genes (false discovery rate < 0.05) were
identified from triplicate datasets by comparing the mutants to wild-type plants (**Figure 3**). In total,
over 200 genes were differentially expressed in *vtc2-4* mutants, whereas only 24 genes, including
VTC3 itself, were affected in *vtc3-3* mutants. Among these, seven genes were commonly affected in
both mutants. The relatively minor transcriptomic impact of VTC3 compared to VTC2 likely reflects
the weaker ascorbate-deficient phenotype of *vtc3-3*. Gene ontology (GO) enrichment analysis
showed that "defense response to pathogens" was the most enriched GO term among the 12 genes
upregulated in *vtc3* mutants (**Figure 3**). Furthermore, downregulated genes specific to *vtc3-3*
included cytochrome P450 superfamily protein (AT3G44970), transmembrane protein (AT5G01881),
mto1 responding down 1 (AT1G53480), and phosphoinositide 4-kinase gamma 1 (AT2G40850),
none of which are known to be directly associated with ascorbate metabolism. These results suggest
that VTC3 plays a more limited role in transcriptomic regulation than VTC2.

Additive effects of the* vtc2-4 *and* vtc3-3 *mutations on ascorbate pool size under low- and
high-light conditions
As shown in **Figure 1**, the absence of VTC2 severely restricted ascorbate levels under low-light
conditions, whereas the loss of VTC3 had a comparatively milder effect. However, while *vtc2-4*
mutants still exhibited a light-dependent increase in ascorbate pool size, this response was

significantly diminished in *vtc3-3* mutants. Given these distinct characteristics, we generated *vtc2-4*

vtc3-3 double mutants. Homozygous double mutants were identified by genotyping the F_2

population following crossing (**Supplementary Figure S1**). Compared to single mutants, the double

mutants exhibited the lowest ascorbate levels, retaining only approximately 51% of the ascorbate

content observed in *vtc2-4* single mutants, equivalent to about 14% of wild-type levels (**Figure 5**).

Notably, except for mutants entirely lacking detectable ascorbate (e.g., *vtc2 vtc5* double mutants),

vtc2-4 vtc3-3 double mutants displayed the lowest ascorbate levels reported among

ascorbate-deficient mutants. Despite these extremely low levels, their growth under low-light

conditions for three weeks was comparable to that of wild-type plants and single mutants (**Figure 5**),

suggesting that even 14% of wild-type ascorbate levels is sufficient to support Arabidopsis growth

under controlled laboratory conditions.

When exposed to high-light stress for 24 hours, *vtc2-4* and *vtc3-3* single mutants

exhibited photo-bleaching, primarily in older leaves (**Figure 6**). In contrast, the *vtc2-4 vtc3-3* double

mutants displayed a more severe photo-bleaching phenotype, extending to mature leaves, and

exhibited the most pronounced cell death, consistent with their extremely low ascorbate content.

Reflecting these phenotypes, the reduction in the maximum quantum yield of photosystem II (F_v/F_m)

under high-light stress was most pronounced in the double mutants (**Figure 6**), indicating severe

damage to photosynthesis.

Considering the heightened sensitivity of *vtc2-4 vtc3-3* mutants to high-light stress, we

measured ascorbate content in wild-type plants and mutants during the early stages of high-light

exposure (6 and 12 hours). Ascorbate levels in wild-type plants and *vtc2-4* mutants increased

approximately 1.5-fold by 12 hours after high-light exposure (**Figure 7**), but there was no statistical

difference. Furthermore, no significant increase in ascorbate levels was observed in *vtc3-3* or *vtc2-4*

vtc3-3 mutants. However, the early-stage ascorbate increase in wild-type plants and *vtc2-4* mutants

was relatively modest, making detailed comparisons between mutants challenging.

To further investigate the ascorbate accumulation response under different light

intensities, we conducted similar experiments under moderate-light conditions (750 µmol photons

m^{-2} s^{-1}; ML^{750}). Under ML^{750} stress, ascorbate accumulation was observed even in *vtc3-3* and

vtc2-4 vtc3-3 mutants; however, the increase was significantly smaller compared to *vtc2-4*, with the

double mutants exhibiting the lowest levels of stress-induced ascorbate accumulation (**Figure 7**).

These results demonstrate that the simultaneous knockout of VTC2 and VTC3 severely limits both

steady-state ascorbate levels and stress-induced ascorbate accumulation, thereby dramatically

increasing sensitivity to photooxidative stress.

Non-photochemical quenching in **vtc2-4 vtc3-3** *double mutants*

As ascorbate is implicated in the induction of NPQ (Müller-Moulé *et al.*, 2004), a photoprotective

mechanism that dissipates excess absorbed light energy as heat, we investigated how the severe

ascorbate deficiency in *vtc2-4 vtc3-3* double mutants affects NPQ. Plants were grown under

low-light conditions for three weeks before being subjected to high-light stress for 24 hours. Under

low-light conditions, NPQ induction was partially suppressed in all mutants compared to wild-type

plants, with statistically significant but relatively small differences (**Figure 8A and C**). Among the

single mutants, *vtc2-4* exhibited a greater reduction in NPQ induction compared to *vtc3-3*, consistent

with its lower ascorbate pool size. Although the *vtc2-4 vtc3-3* double mutants showed a trend toward

further reduced NPQ values, the difference was not statistically significant compared to *vtc2-4* single

mutants. Following high-light exposure, NPQ suppression was most pronounced in *vtc2-4 vtc3-3*

double mutants, significantly lower than in either *vtc2-4* or *vtc3-3* single mutants (**Figure 8B and D**).

Interestingly, while *vtc2-4* exhibited lower NPQ values than *vtc3-3* under low-light conditions

(**Figure 8 C**), this difference disappeared under high-light stress. (**Figure 8D**) These variations in

NPQ induction closely paralleled the differences in ascorbate pool sizes among the mutants. These

findings provide new insights into the relationship between NPQ induction and ascorbate levels,

highlighting the additive effects of *vtc2-4* and *vtc3-3* mutations on NPQ suppression under high-light

stress. The results suggest that ascorbate plays a critical role in NPQ induction under both low- and

high-light conditions and that VTC2 and VTC3 contribute to this regulation through their roles in

ascorbate accumulation.

Discussion

Among the *vtc* mutant series, *vtc3* is the only mutant with a mutation in a gene that does not encode

a biosynthetic enzyme (Conklin *et al.*, 2013). VTC3 is a unique protein containing both protein

kinase and phosphatase domains and is essential for ascorbate accumulation in response to

continuous exposure to growth light. Given that VTC2/GGP is the rate-limiting enzyme in ascorbate

biosynthesis and that its expression and activity are enhanced by light, we initially hypothesized that

VTC3 regulates ascorbate biosynthesis at the VTC2/GGP step. However, this hypothesis was

strongly refuted, as the loss of VTC3 did not reduce *VTC2/VTC5* transcription or GGP activity

(**Figure 2**). As previously reported, VTC2 translation is controlled by ascorbate via an uORF in the

5′-untranslated region of the *VTC2* gene (Laing *et al.*, 2015). Because we were unable to examine

VTC2/GGP expression at the protein level, we cannot entirely rule out the possibility that VTC3

influences VTC2 translation. However, if this were the case, the absence of VTC3 would likely lead

to reduced GGP activity, which was not observed (**Figure 2**). Thus, we conclude that VTC3 is not

required for the expression and activity of VTC2/GGP.

We also found that the loss of VTC3 did not affect glutathione profiles or the activities of

APX, DHAR, and MDAR (**Figure 3**). Since we did not examine the expression or activity of

individual isoforms of these enzymes, we cannot rule out the possibility that VTC3 affects ascorbate

redox cycling in specific compartments. However, recent studies have demonstrated that the

complementary actions of MDARs, DHARs, and glutathione maintain a robust ascorbate recycling

system in Arabidopsis (Maruta *et al.,* 2024). For instance, the loss of all three *DHAR* genes has a

negligible impact on ascorbate levels, even under stress conditions (Rahantaniaina *et al.*, 2017; Terai

et al., 2020). Considering the resilience of the ascorbate recycling system, it is unlikely that VTC3

plays a role in regulating ascorbate redox cycling. Another possible function of VTC3 could be the

inhibition of ascorbate degradation. However, this is also unlikely, as ascorbate degradation begins

with DHA, and ascorbate recycling capacity was unaffected in *vtc3* mutants.

Since VTC3 contains both protein kinase and phosphatase domains, it was also plausible

that it mediates light-dependent signaling pathways to enhance ascorbate pool size. However,

transcriptome analysis revealed that only a limited number of genes were specifically up or

downregulated in *vtc3-3* mutants, and none had known associations with ascorbate metabolism. GO

enrichment analysis identified "defense response to pathogens" as the most enriched term among the

12 genes whose expression was specifically enhanced in *vtc3* mutants (**Figure 4 and**

Supplementary Table S2). This finding aligns with previous observations in *vtc1* mutants, where

ascorbate deficiency was linked to enhanced pathogen responses (Pastori *et al.*, 2003). Thus, the

transcriptional activation of some defense-related genes in *vtc3-3* mutants is likely a secondary effect

of ascorbate deficiency rather than the direct involvement of VTC3 in defense signaling. In *vtc2-4*

mutants, 115 genes were upregulated (**Figure 4**), with GO terms related to "response to salicylic

acid" and "response to chitin" enriched (**Supplementary Table S2**), further suggesting a link

between ascorbate deficiency and pathogen responses. Taken together, these findings indicate that

the loss of VTC3 has minimal impact on VTC2/GGP, ascorbate redox cycling, and the transcriptome.

Given its significant effect on ascorbate pool size, we hypothesize that VTC3 regulates ascorbate

biosynthesis at steps other than VTC2/GGP.

Consistent with previous reports (Terai *et al.*, 2020; Hamada *et al.*, 2023), *vtc2-4* mutants

retained the ability to increase ascorbate accumulation in response to light stress (**Figure 1**). We

initially hypothesized that this ability might be due to the activation of residual VTC5. However,

VTC5 transcription showed little to no increase in *vtc2-4* mutants compared to the wild type, and

GGP activity was nearly undetectable in *vtc2-4* mutants even under high-light stress (**Figure 2**). This

could be due to GGP activity in *vtc2-4* being below the detection limit. Indeed, the simultaneous loss

of VTC2 and VTC5 results in ascorbate levels below the detection limit, indicating that VTC5

functions as an active enzyme in ascorbate biosynthesis (Dowdle *et al.*, 2007; Lim *et al.*, 2016).

Alternatively, because we used GDP-D-glucose instead of GDP-L-galactose (the physiological

substrate, which is commercially unavailable) to measure GGP activity, our assay may not fully

reflect endogenous GGP activity. Nevertheless, it is intriguing that ascorbate levels in *vtc2* mutants

can increase significantly without a measurable increase in GGP activity. If alternative biosynthetic

pathways are not considered, this suggests a mechanism that enhances the GGP response

independent of measurable GGP activity. The recently reported metabolon formation of ascorbate

biosynthetic enzymes may be involved (Fenech *et al.*, 2021), but further experimental validation is

required.

Another key objective of this study was to examine the impact of the simultaneous loss

of VTC2 and VTC3 on ascorbate pool size. As expected, *vtc2-4 vtc3-3* double mutants exhibited the

lowest ascorbate levels reported among viable ascorbate-deficient mutants, retaining only about 14%

of wild-type levels (**Figure 5**). Nevertheless, under low-light conditions, *vtc2-4 vtc3-3* double

mutants displayed a healthy phenotype similar to wild-type plants. The original *vtc2-1* mutant,

isolated via an ethyl methanesulfonate (EMS) mutagenesis screen, has been reported to exhibit

dwarfism under standard growth conditions, whereas the T-DNA insertion mutant *vtc2-4*, despite

having comparable ascorbate levels, does not exhibit dwarfism (Lim *et al*., 2016). It has been

suggested that the dwarfism observed in *vtc2-1* may be due to a mutation in another gene. The

relationship between ascorbate levels and growth must be carefully evaluated, as it is highly

influenced by growth light intensity. For instance, when grown under relatively higher light

conditions (200 µmol photons m^{-2} s^{-1}), both *vtc2-1* and *vtc2-4* exhibit similar degrees of dwarfism

(Plumb *et al.*, 2018). Considering this complexity, our data suggest that under artificial low-light

conditions, as little as 14% of wild-type ascorbate levels are sufficient for normal growth in

Arabidopsis. This finding provides the idea that the primary reason plants accumulate high

concentrations of ascorbate is to protect against oxidative stress rather than to fulfill its coenzyme

functions.

The loss of VTC3 significantly impaired high-light-induced ascorbate accumulation in

both wild-type and *vtc2-4* backgrounds (**Figure 7**). *vtc2-4 vtc3-3* double mutants maintained

extremely low ascorbate levels even under high-light conditions, leading to more severe

photooxidative damage than either single mutant (**Figure 6**). In terms of photobleaching severity,

vtc2-4 was more sensitive to high light than *vtc3-3*, and the sensitivity of the double mutant was

entirely consistent with its steady-state ascorbate pool size (**Figure 5**). NPQ values were statistically

significantly lower in all *vtc* mutants compared to wild-type plants under low-light conditions, with

vtc2-4 and the double mutant exhibiting greater reductions than *vtc3-3* (**Figure 8**). However, no

significant difference was observed between *vtc2-4* and the double mutant under low-light

conditions, suggesting that NPQ suppression due to ascorbate deficiency may reach a threshold at

~30% of wild-type ascorbate levels. Interestingly, this difference became more pronounced after

high-light exposure, paralleling the fact that *vtc2-4* mutants still exhibited light-induced ascorbate

accumulation, whereas this response was severely impaired in the double mutants (**Figure 7**). So far,

vtc1 and *vtc4* mutants have also been used to study ascorbate deficiency, but these mutants have

pleiotropic effects unrelated to ascorbate metabolism (Torabinejad *et al.*, 2009; Barth *et al.*, 2010). In

contrast, *vtc2-4* and *vtc3-3* mutants provide a more targeted approach. Since the *vtc2-4 vtc3-3* double

mutant retains the characteristics of both single mutants, comparative analysis of these mutants

represents a powerful tool for elucidating the physiological roles of ascorbate.

In conclusion, our findings indicate that VTC3 is not required for the expression or

activity of VTC2/GGP, ascorbate redox cycling (including ascorbate recycling), or the regulation of

the transcriptome. While the fundamental function of VTC3 remains unresolved, our study has

significantly narrowed down its possible roles. We propose that VTC3 regulates ascorbate

biosynthesis at a step other than VTC2, and future research will focus on validating this hypothesis.

Furthermore, this study revealed that the simultaneous knockout of VTC2 and VTC3 exacerbates

ascorbate deficiency and increases sensitivity to light stress. The newly generated *vtc2-4 vtc3-3*

double mutant retains only 14% of wild-type ascorbate levels. Given its distinct physiological

characteristics, the double mutant, in combination with the single mutants, is expected to serve as a

valuable tool for further dissecting the physiological functions of ascorbate in plants.

Figure legends

Figure 1 Ascorbate accumulation in *vtc3-3* under high light

Wild-type, *vtc2-4*, and *vtc3-3* plants were grown in soil under low-light (40–60 μmol photons m^{-2} s^{-}

^{1}) conditions for three weeks and then exposed to high-light stress (1,500 μmol photons m^{-2} s^{-1}) for

48 h. Fully expanded leaves were used for ascorbate measurements. (A) Total ascorbate content

(sum of ASC and DHA contents). (B) The amounts of ascorbate increase from before to after

high-light exposure (48 h). Data are presented as the mean ± SE of four biological replicates.

Different letters indicate significant differences ($P < 0.05$, one- or two-way ANOVA followed by

Tukey–Kramer test). ASC, reduced ascorbate; DHA, dehydroascorbate; FW, fresh weight.

Figure 2 Expression and activity of GGP in *vtc3-3* under high light

Wild-type, *vtc2-4*, and *vtc3-3* plants were grown in soil under low-light (40–60 μmol photons m^{-2} s^{-}

^{1}) conditions for three weeks and then exposed to high-light stress (1,500 μmol photons m^{-2} s^{-1}) for

24 h. Fully expanded leaves were used in subsequent experiments. (A, B) Transcript levels of *VTC2*

and *VTC5* were determined using qPCR analysis. (C) GGP activity was determined as

GDP-D-glucose degrading activity. Data are presented as the mean ± SE of three biological

replicates. (A, C) Significant differences between wild type and *vtc3-3* (Student's *t*-test): $^{*}P < 0.05$,

$^{**}P < 0.01$, versus the values of the wild-type plants. Because VTC2 transcript levels and GGP

activity were negligible in vtc2-4, statistical analysis was performed only between wild-type and

vtc3-3. (B) Different letters indicate significant differences between genotypes ($P < 0.05$, one-way

ANOVA followed by Tukey–Kramer test).

Figure 3 Antioxidant levels and enzyme activities related to ascorbate redox turnover in *vtc3-3*

under high light

Wild-type, *vtc2-4*, and *vtc3-3* plants were grown on MS plates under low-light (40–60 μmol photons

m^{-2} s^{-1}) conditions for two weeks and then exposed to moderate-light stress (650 μmol photons m^{-2} s

$^{-1}$) for 24 h. (A) Ascorbate oxidation by ascorbate peroxidase (APX) and its recycling through

monodehydroascorbate reductase (MDAR) and dehydroascorbate reductase (DHAR). (B) Total

ascorbate content (sum of ASC and DHA contents). (C) Total glutathione content (sum of GSH and

GSSG contents). (D) APX activity. (E) MDAR activity. (F) DHAR activity. Data are presented as the

mean ± SE of three or four biological replicates. Different letters indicate significant differences ($P <$

0.05, two-way ANOVA followed by Tukey–Kramer test). ASC, reduced ascorbate; DHA,

dehydroascorbate; FW, fresh weight; GSH, reduced glutathione; GSSG, oxidized glutathione.

Figure 4 Transcriptomic changes in *vtc2-4* and *vtc3-3*

Wild-type, *vtc2-4*, and *vtc3-3* plants were grown on MS plates under low-light (40–60 μmol photons

m^{-2} s^{-1}) conditions for two weeks and then incubated in darkness for two days (to decline ascorbate

pool size), followed by a 6-hour exposure to light (100 μmol photons m^{-2} s^{-1}). From the triplicate

data, we selected the genes whose expression was significantly up-regulated (false discovery rate <

0.05) in *vtc2-4* or *vtc3-3* compared to the wild-type plants. (A, B) The Venn diagrams show the

overlap between sets of genes upregulated (A) or downregulated (B) in *vtc2-4* and *vtc3-3*. (C) Gene

ontology enrichment analysis (biological process) of 12 genes whose expression was upregulated in

vtc3-3 compared to the wild-type plants. (C) Gene ontology (GO) terms enriched in genes whose

expression was specifically promoted in *vtc3-3*. BG, background; Ref, reference.

Figure 5 Phenotypes and ascorbate profiles of *vtc2-4 vtc3-3* double mutants

Wild-type, *vtc2-4*, *vtc3-3*, and *vtc2-4 vtc3-3* plants were grown in soil under low-light (40–60 μmol

photons m^{-2} s^{-1}) conditions for three weeks. Fully expanded leaves were used for ascorbate

measurements. (A) Representative phenotypes of plants. (B, C) Box plots illustrate the rosette area

(B) and fresh weights (C). Horizontal bars and cross marks within the box represent median and

mean values from 12 biological replicates, respectively, with boxes representing the 25th and 75th

percentiles. The outer error bars (whiskers) represent maximum and minimum values. There was no

significant difference between genotypes (one-way ANOVA). (D) Total ascorbate content (sum of

ASC and DHA contents). Data are presented as the mean ± SE of five biological replicates. Different

letters indicate significant differences ($P < 0.05$, one-way ANOVA followed by Tukey–Kramer test).

ASC, reduced ascorbate; DHA, dehydroascorbate; FW, fresh weight.

Figure 6 Sensitivity of *vtc2-4 vtc3-3* double mutants to high-light stress

Wild-type, *vtc2-4*, *vtc3-3*, and *vtc2-4 vtc3-3* plants were grown in soil under low-light (40–60 μmol

photons m^{-2} s^{-1}) conditions for three weeks and then exposed to high-light stress (1,500 μmol

photons m^{-2} s^{-1}) for 24 h. (A) Representative phenotypes of plants before and after high-light stress.

(B) Box plots illustrate lesion area (% of whole rosette area). The data were obtained from nine

plants per genotype in three independent experiments. As wild-type plants showed no lesions, lesion

area analysis was not performed. Horizontal bars and cross marks within the box represent median

and mean values from five biological replicates, respectively, with boxes representing the 25th and

75th percentiles. The outer error bars (whiskers) and diamonds represent the maximum and minimum

values, and an outlier, respectively. Different letters indicate significant differences ($P < 0.05$,

one-way ANOVA followed by Tukey–Kramer test). (C) The maximum quantum yield of

photosystem II (F_v/F_m) before and after high-light stress. Significant differences between genotypes

(one-way ANOVA followed by Tukey–Kramer test): [*]$P < 0.05$, [**]$P < 0.01$.

Figure 7 Ascorbate profiles in *vtc2-4 vtc3-3* under high- or moderate-light stress

Wild-type, *vtc2-4*, *vtc3-3*, and *vtc2-4 vtc3-3* plants were grown in soil under low-light (40–60 μmol

photons m^{-2} s^{-1}) conditions for three weeks and then exposed to high-light stress (1,500 µmol

photons m^{-2} s^{-1}) for 24 h (A, B), or to moderate-light stress (750 µmol photons m^{-2} s^{-1}) for 48 h (C,

D). Fully expanded leaves were used for ascorbate measurements. (A, C) Total ascorbate content

(sum of ASC and DHA contents). Data are presented as the mean ± SE of three or four biological

replicates. Significant differences between data before and after light stress for each genotype

(one-way-ANOVA with Dunnett's post hoc test for total content): $P^* < 0.05$, $^{**}P < 0.01$, compared to

the values of 0 h. (B, D) The amounts of ascorbate increase from before to after high- or

moderate-light exposure (24 or 48 h, respectively). Data are presented as the mean ± SE of three or

four biological replicates. Different letters indicate significant differences ($P < 0.05$, one-way

ANOVA followed by Tukey–Kramer test). ASC, reduced ascorbate; DHA, dehydroascorbate; FW,

fresh weight; n.s., not significant.

Figure 8 Non-photochemical quenching in *vtc2-4 vtc3-3* before and after high-light stress

Wild-type, *vtc2-4*, *vtc3-3*, and *vtc2-4 vtc3-3* plants were grown in soil under low-light (40–60 µmol

photons m^{-2} s^{-1}) conditions for three weeks and then exposed to high-light stress (1,500 µmol

photons m^{-2} s^{-1}) for 24 h. (A, B) Non-photochemical quenching (NPQ) induction upon the onset of

actinic light (90 µmol photons m-2 s-1) before (A) and after (B) stress. Data are presented as the

mean ± SE of three or four biological replicates. The red triangles indicate the data points used for

statistical analysis (C, D). (C, D) Box plots illustrate NPQ data before (C) and after (D) stress,

corresponding to the red triangles in (A) and (B), respectively. Horizontal bars and cross marks

within the box represent median and mean values from 12 biological replicates, respectively, with

boxes representing the 25th and 75th percentiles. The outer error bars (whiskers) represent

maximum and minimum values. Different letters indicate significant differences ($P < 0.05$, one-way

ANOVA followed by Tukey–Kramer test).

Supplementary Data

Supplementary Figure S1 Expression of *VTC2* and *VTC3* in *vtc* mutant lines

Supplementary Table S1 List of primers used

Supplementary Table S2 List of genes whose expression is altered in *vtc* mutants and gene

ontology enrichment analysis

Author Contributions

YT, TM, and TI: conceptualization; YT, TM, and TI: methodology; YT, and TM: formal analysis;

YT, TM, KA, HN, AH, and TI: investigation; YT, and TM: data curation; YT, and TM: writing -

original draft; KA, HN, AH, and TI: writing - review & editing; TM: visualization; TM, TI:

supervision; TM, TI: funding acquisition

Conflicts of Interest

The authors declare no conflict of interest.

Funding Statement

This work was conducted as an SDGs Research Project of Shimane University (T.M.) and supported

by JSPS Bilateral Program Number JPJSBP120232302 (T.M.) and JSPS KAKENHI Grant Number

24K01680 (T.I.).

Data availability

The data underlying this article are available in the article and its online supplementary material. The

raw and processed data of RNA-seq have been deposited in the Gene Expression Omnibus under

accession number GSE293768.

Acknowledgments

We would like to thank Hiromi Ueno, Reona Tanimura, and Natsumi Ito for their technical

assistance.

References

Aarabi F, Ghigi A, Wijesingha Ahchige M, Bulut M, Geigenberger P, Neuhaus HE,

Sampathkumar A, Alseekh S, Fernie AR. 2023. Genome-Wide Association Study unveils

ascorbate regulation by PAS/LOV PROTEIN during high light acclimation. Plant Physiology doi:

10.1093/plphys/kiad323.

Asada K. 1999. THE WATER-WATER CYCLE IN CHLOROPLASTS: Scavenging of active

oxygens and dissipation of excess photons. Annual Review of Plant Physiology and Plant molecular

biology **50**, 601–639.

Baker NR. 2008. Chlorophyll fluorescence: a probe of photosynthesis in vivo. Annual Review of

Plant Biology **59**, 89–113.

Barth C, Gouzd ZA, Steele HP, Imperio RM. 2010. A mutation in GDP-mannose

pyrophosphorylase causes conditional hypersensitivity to ammonium, resulting in Arabidopsis root

growth inhibition, altered ammonium metabolism, and hormone homeostasis. Journal of

experimental botany **61**, 379–394.

Bournonville C, Mori K, Deslous P, *et al.* 2023. Blue light promotes ascorbate synthesis by

deactivating the PAS/LOV photoreceptor that inhibits GDP-L-galactose phosphorylase. The Plant

Cell **35**, 2615–2634.

Bulley SM, Rassam M, Hoser D, Otto W, Schünemann N, Wright M, MacRae E, Gleave A,

Laing W. 2009. Gene expression studies in kiwifruit and gene over-expression in Arabidopsis

indicates that GDP-L-galactose guanyltransferase is a major control point of vitamin C biosynthesis.

Journal of Experimental Botany **60**, 765–778.

Bulley S, Wright M, Rommens C, *et al.* 2012. Enhancing ascorbate in fruits and tubers through

over-expression of the L-galactose pathway gene GDP-L-galactose phosphorylase. Plant

Biotechnology Journal **10**, 390–397.

Conklin PL, DePaolo D, Wintle B, Schatz C, Buckenmeyer G. 2013. Identification of

Arabidopsis VTC3 as a putative and unique dual function protein kinase::protein phosphatase

involved in the regulation of the ascorbic acid pool in plants. Journal of Experimental Botany **64**,

2793–2804.

Conklin PL, Gatzek S, Wheeler GL, Dowdle J, Raymond MJ, Rolinski S, Isupov M, Littlechild

JA, Smirnoff N. 2006. *Arabidopsis thaliana VTC4* encodes L-galactose-1-P phosphatase, a plant

ascorbic acid biosynthetic enzyme. The Journal of Biological Chemistry **281**, 15662–15670.

Conklin PL, Norris SR, Wheeler GL, Williams EH, Smirnoff N, Last RL. 1999. Genetic

evidence for the role of GDP-mannose in plant ascorbic acid (vitamin C) biosynthesis. Proceedings

of the National Academy of Sciences of the United States of America **96**, 4198–4203.

Conklin PL, Saracco SA, Norris SR, Last RL. 2000. Identification of ascorbic acid-deficient

Arabidopsis thaliana mutants. Genetics **154**, 847–856.

Dowdle J, Ishikawa T, Gatzek S, Rolinski S, Smirnoff N. 2007. Two genes in *Arabidopsis*

thaliana encoding GDP-L-galactose phosphorylase are required for ascorbate biosynthesis and

652 seedling viability. The Plant Journal **52**, 673–689.

653 **Fenech M, Amorim-Silva V, Esteban Del Valle A, Arnaud D, Ruiz-Lopez N, Castillo AG,**

654 **Smirnoff N, Botella MA**. 2021. The role of GDP-l-galactose phosphorylase in the control of

655 ascorbate biosynthesis. Plant Physiology **185**, 1574–1594.

656 **Foyer CH, Halliwell B**. 1977. Purification and properties of dehydroascorbate reductase from

657 spinach leaves. Phytochemistry **16**, 1347–1350.

658 **Hamada T, Maruta T**. 2024. Measurements of ascorbate and dehydroascorbate in plants using

659 high-performance liquid chromatography. Methods in Molecular Biology (Clifton, N.J.) **2798**,

660 131–139.

661 **Hamada A, Tanaka Y, Ishikawa T, Maruta T**. 2023. Chloroplast dehydroascorbate reductase and

662 glutathione cooperatively determine the capacity for ascorbate accumulation under photooxidative

663 stress conditions. The Plant Journal **114**, 68–82.

664 **Hossain MA, Asada K**. 1985. Monodehydroascorbate reductase from cucumber is a flavin adenine

665 dinucleotide enzyme. The Journal of Biological Chemistry **260**, 12920–12926.

666 **Ishida T, Tanaka Y, Maruta T, Ishikawa T**. 2024. The D-mannose/L-galactose pathway plays a

667 predominant role in ascorbate biosynthesis in the liverwort *Marchantia polymorpha* but is not

668 regulated by light and oxidative stress. The Plant Journal **117**, 805–817.

669 **Jander G, Norris SR, Rounsley SD, Bush DF, Levin IM, Last RL**. 2002. Arabidopsis map-based

670 cloning in the post-genome era. Plant Physiology **129**, 440–450.

671 **Kameoka T, Okayasu T, Kikuraku K, Ogawa T, Sawa Y, Yamamoto H, Ishikawa T, Maruta T**.

672 2021. Cooperation of chloroplast ascorbate peroxidases and proton gradient regulation 5 is critical

673 for protecting Arabidopsis plants from photo-oxidative stress. The Plant Journal **107**, 876–892.

674 **Laing WA, Martínez-Sánchez M, Wright MA,** *et al.* 2015. An upstream open reading frame is

675 essential for feedback regulation of ascorbate biosynthesis in Arabidopsis. The Plant Cell **27**,

676 772–786.

677 **Laing WA, Wright MA, Cooney J, Bulley SM**. 2007. The missing step of the L-galactose pathway

678 of ascorbate biosynthesis in plants, an L-galactose guanyltransferase, increases leaf ascorbate content.

Proceedings of the National Academy of Sciences of the United States of America **104**, 9534–9539.

Lim B, Smirnoff N, Cobbett CS, Golz JF. 2016. Ascorbate-deficient *vtc2* mutants in Arabidopsis

do not exhibit decreased growth. Frontiers in Plant Science **7**, 1025.

Linster CL, Gomez TA, Christensen KC, Adler LN, Young BD, Brenner C, Clarke SG. 2007.

Arabidopsis *VTC2* encodes a GDP-L-galactose phosphorylase, the last unknown enzyme in the

Smirnoff-Wheeler pathway to ascorbic acid in plants. The Journal of Biological Chemistry **282**,

18879–18885.

Maruta T. 2022. How does light facilitate vitamin C biosynthesis in leaves? Bioscience,

Biotechnology, and Biochemistry **86**, 1173–1182.

Maruta T, Sawa Y, Shigeoka S, Ishikawa T. 2016. Diversity and evolution of ascorbate peroxidase

functions in chloroplasts: More than just a classical antioxidant enzyme? Plant & Cell Physiology **57**,

1377–1386.

Maruta T, Tanaka Y, Yamamoto K, Ishida T, Hamada A, Ishikawa T. 2024. Evolutionary

insights into strategy shifts for the safe and effective accumulation of ascorbate in plants. Journal of

Experimental Botany **75**, 2664–2681.

Mittler R, Zandalinas SI, Fichman Y, Van Breusegem F. 2022. Reactive oxygen species

signalling in plant stress responses. Nature Reviews. Molecular Cell Biology **23**, 663–679.

Müller-Moulé P, Golan T, Niyogi KK. 2004. Ascorbate-deficient mutants of Arabidopsis grow in

high light despite chronic photooxidative stress. Plant Physiology **134**, 1163–1172.

Noctor G, Mhamdi A, Foyer CH. 2016. Oxidative stress and antioxidative systems: recipes for

successful data collection and interpretation. Plant, Cell & Environment **39**, 1140–1160.

Ogura Y, Komatsu A, Zikihara K, Nanjo T, Tokutomi S, Wada M, Kiyosue T. 2008. Blue light

diminishes interaction of PAS/LOV proteins, putative blue light receptors in *Arabidopsis thaliana*,

with their interacting partners. Journal of Plant Research **121**, 97–105.

Pastori GM, Kiddle G, Antoniw J, Bernard S, Veljovic-Jovanovic S, Verrier PJ, Noctor G,

Foyer CH. 2003. Leaf vitamin C contents modulate plant defense transcripts and regulate genes that

control development through hormone signaling. The Plant Cell **15**, 939–951.

Plumb W, Townsend AJ, Rasool B, Alomrani S, Razak N, Karpinska B, Ruban AV, Foyer CH.

2018. Ascorbate-mediated regulation of growth, photoprotection, and photoinhibition in *Arabidopsis*

thaliana. Journal of Experimental Botany **69**, 2823–2835.

Rahantaniaina M-S, Li S, Chatel-Innocenti G, Tuzet A, Issakidis-Bourguet E, Mhamdi A,

Noctor G. 2017. Cytosolic and chloroplastic DHARs cooperate in oxidative stress-driven activation

of the salicylic acid pathway. Plant Physiology **174**, 956–971.

Shigeoka S, Ishikawa T, Tamoi M, Miyagawa Y, Takeda T, Yabuta Y, Yoshimura K. 2002.

Regulation and function of ascorbate peroxidase isoenzymes. Journal of Experimental Botany **53**,

1305–1319.

Smirnoff N. 2018. Ascorbic acid metabolism and functions: A comparison of plants and mammals.

Free Radical Biology and Medicine **122**, 116–129.

Sodeyama T, Nishikawa H, Harai K, Takeshima D, Sawa Y, Maruta T, Ishikawa T. 2021. The

d-mannose/l-galactose pathway is the dominant ascorbate biosynthetic route in the moss

Physcomitrium patens. The Plant Journal **107**, 1724–1738.

Tanaka M, Takahashi R, Hamada A, Terai Y, Ogawa T, Sawa Y, Ishikawa T, Maruta T. 2021.

Distribution and functions of monodehydroascorbate reductases in plants: Comprehensive reverse

genetic analysis of *Arabidopsis thaliana* Enzymes. Antioxidants **10**.

Terai Y, Ueno H, Ogawa T, Sawa Y, Miyagi A, Kawai-Yamada M, Ishikawa T, Maruta T. 2020.

Dehydroascorbate reductases and glutathione set a threshold for high-light-induced ascorbate

accumulation. Plant Physiology **183**, 112–122.

Tian T, Liu Y, Yan H, You Q, Yi X, Du Z, Xu W, Su Z. 2017. agriGO v2.0: a GO analysis toolkit

for the agricultural community, 2017 update. Nucleic Acids Research **45**, W122–W129.

Torabinejad J, Donahue JL, Gunesekera BN, Allen-Daniels MJ, Gillaspy GE. 2009. VTC4 is a

bifunctional enzyme that affects myoinositol and ascorbate biosynthesis in plants. Plant Physiology

150, 951–961.

Vidal-Meireles A, Neupert J, Zsigmond L, Rosado-Souza L, Kovács L, Nagy V, Galambos A,

Fernie AR, Bock R, Tóth SZ. 2017. Regulation of ascorbate biosynthesis in green algae has

evolved to enable rapid stress-induced response via the *VTC2* gene encoding GDP-l-galactose

phosphorylase. The New Phytologist **214**, 668–681.

Wheeler GL, Jones MA, Smirnoff N. 1998. The biosynthetic pathway of vitamin C in higher
plants. Nature **393**, 365–369.

Winkler BS, Orselli SM, Rex TS. 1994. The redox couple between glutathione and ascorbic acid: a
chemical and physiological perspective. Free Radical Biology & Medicine **17**, 333–349.

Yabuta Y, Mieda T, Rapolu M, Nakamura A, Motoki T, Maruta T, Yoshimura K, Ishikawa T,
Shigeoka S. 2007. Light regulation of ascorbate biosynthesis is dependent on the photosynthetic
electron transport chain but independent of sugars in Arabidopsis. Journal of Experimental Botany
58, 2661–2671.

Yoshimura K, Nakane T, Kume S, Shiomi Y, Maruta T, Ishikawa T, Shigeoka S. 2014. Transient
expression analysis revealed the importance of VTC2 expression level in light/dark regulation of
ascorbate biosynthesis in Arabidopsis. Bioscience, Biotechnology, and Biochemistry **78**, 60–66.

Zechmann B, Stumpe M, Mauch F. 2011. Immunocytochemical determination of the subcellular
distribution of ascorbate in plants. Planta **233**, 1–12.

Tanaka & Maruta et al., Figure 1

Tanaka & Maruta et al., Figure 2

Tanaka & Maruta et al., Figure 3

A *vtc2*   *vtc3*

115   4   12

Upregulated genes vs wild type
(FDR *P* value < 0.05)

B *vtc2*   *vtc3*

88   3   5

Downregulated genes vs wild type
(FDR *P* value < 0.05)

C

GO term	Description	Number in input list	Number in BG/Ref	*p*-value	FDR
GO:0006952	defense response	7	1566	9.80E-07	1.90E-05
GO:0050832	defense response to fungus	5	504	1.30E-06	1.90E-05
GO:0006950	response to stress	9	3506	1.10E-06	1.90E-05
GO:0098542	defense response to other organism	6	936	1.00E-06	1.90E-05
GO:0009620	response to fungus	5	573	2.40E-06	2.80E-05
GO:0043207	response to external biotic stimulus	6	1212	4.60E-06	3.80E-05
GO:0051707	response to other organism	6	1210	4.50E-06	3.80E-05
GO:0009607	response to biotic stimulus	6	1253	5.50E-06	4.00E-05
GO:0009605	response to external stimulus	6	1582	2.10E-05	0.00014
GO:0051704	multi-organism process	6	1631	2.50E-05	0.00015

Tanaka & Maruta et al., Figure 4

Tanaka & Maruta et al., Figure 5

Tanaka & Maruta et al., Figure 6

Tanaka & Maruta et al., Figure 7

Tanaka & Maruta et al., Figure 8