1 *Title Page*

A transfer RNA methyltransferase with an unusual domain composition catalyzes 2′-*O*-methylation at position 6 in tRNA.

Teppei Matsuda[1], Ryota Yamagami[1]*, Aoi Ihara[2], Takeo Suzuki[3], Akira Hirata[2] and Hiroyuki Hori[1]*

1. Department of Materials Science and Biotechnology, Graduate School of Science and Engineering, Ehime University, 3 Bunkyo-cho, Matsuyama, Ehime 790-8577, Japan

2. Department of Natural Science, Graduate School of Technology, Industrial and Social Science, Tokushima University, 2-1 Minamijosanjima-cho, Tokushima, Tokushima 770-8506, Japan

3. Department of Medical Biochemistry, Graduate School of Medicine, University of the Ryukyus, 207 Uehara, Nishihara, Okinawa 903-0125, Japan.

* To whom correspondence should be addressed:

Hiroyuki Hori

Department of Materials Science and Biotechnology, Graduate School of Science and Engineering, Ehime University, 3 Bunkyo-cho, Matsuyama, Ehime 790-8577, Japan.

Phone: +81-89-927-8548

E-mail: hori.hiroyuki@outlook.jp, hori.hiroyuki.my@ehime-u.ac.jp

Ryota Yamagami

Department of Materials Science and Biotechnology, Graduate School of Science and Engineering, Ehime University, 3 Bunkyo-cho, Matsuyama, Ehime 790-8577, Japan.

Phone: +81-89-927-9919

E-mail: yamagami.ryota.bn@ehime-u.ac.jp

Running Title: tRNA methyltransferase that possesses an unprecedented domain composition.

Keywords: RNA modification, tRNA, archaea, TrmTS, THUMP, SPOUT

ABSTRACT

***Thermococcus kodakarensis* tRNA$^{\text{Trp}}$ contains 2′-*O*-methylcytidine at position 6 (Cm6).**

However, the tRNA methyltransferase responsible for the modification has not been

identified.   Using comparative genomics we predicted TK1257 as a candidate for the

modification.   Biochemical and mass spectrometry studies of purified recombinant

TK1257 gene product show it to possess a tRNA methyltransferase activity for Cm6

formation.   This protein has a highly unusual composition of domains, containing

N-terminal ferredoxin-like, SPOUT catalytic and THUMP domains.   Previous to this

study, all known THUMP-related tRNA methyltransferases were shown to contain a

Rossmann fold catalytic domain and the nucleosides they produced were

***N*2-methylguanosine and/or *N*2, *N*2-dimethylguanosine.   Therefore, our findings extend**

the knowledge of architecture of tRNA methyltransferases.   We named the TK1257 gene

product TrmTS and showed it can synthesize Am6 and Um6 as well as Cm6.   A *trmTS*

gene deletion strain showed slight growth retardation at high temperatures.   Site-directed

mutagenesis studies based on structural model revealed catalytically and structurally

important amino acid residues in TrmTS and identified a TrmTS-specific linker is

structurally essential.   We showed that TrmTS recognizes the 3′-CCA terminal region

and a stretch loop connected with at least two stems in RNA.   Finally, we constructed a

model of the binding between TrmTS and tRNA.

INTRODUCTION

To date, more than one hundred modified nucleosides have been discovered in tRNA (1-3). Of the modified nucleosides in tRNA, methylated nucleosides are the most abundant (4). Almost all methylated nucleosides in tRNA are synthesized by site-specific tRNA methyltransferases (4).

Although a few RNA methyltransferases use N5, N10-methylenetetrahydrofolate as a methyl group donor (5-7), the majority of RNA methyltransferases use S-adenosyl-L-methionine (SAM) (4). SAM-dependent methyltransferases can be classified into more than five classes by the structures of their catalytic domain (8). The majority of SAM-dependent tRNA methyltransferases are classified as the class I enzymes, which possess a Rossmann fold catalytic domain. The classes II, III and V enzymes methylate low molecular weight compounds or protein and do not methylate RNAs. The majority of class IV enzymes methylate RNA and possess the SpoU-TrmD (SPOUT) catalytic domain (9-13). SpoU is a classical name of TrmH (14, 15) and remains in the superfamily name. TrmD is the tRNA methyltransferase for synthesis of 1-methylguanosine at position 37 (m^{1}G37) (16, 17). In addition, TrmO (a SAM-dependent tRNA methyltransferase for N6-methylation in N6-threonylcarbamoyladenosine synthesis) exceptionally possesses a β-barrel catalytic domain (18).

The existence of the SPOUT RNA methyltransferase superfamily (class IV enzymes) was initially predicted by a bioinformatic study (9) and then confirmed by structural studies (19-25). The SPOUT catalytic domain of Class IV enzymes contains a trefoil-knot structure, in which its C-terminal polypeptide threads through a polypeptide loop. Three conserved amino acid sequence motifs (motifs 1, 2 and 3) have been proposed in the catalytic domain of SPOUT RNA methyltransferase superfamily (9). In the cases of TrmH and a subset of other SPOUT RNA

2′-*O*-methyltransferases such as TrmL, the catalytic arginine residue is contained in the motif 1

and conserved amino acid residues in the motif 2 and motif 3 contribute to formation of the

trefoil-knot structure and SAM-binding pocket (26-28).

The SPOUT RNA methyltransferases can be further classified by their products such as

2′-*O*-methylated nucleosides, m^{1}G, 1-methyladenosine (m^{1}A), and 1-methylpseudouridine

(m^{1}Ψ) (11). Within the SPOUT RNA 2′-*O*-methyltransferases, four tRNA

2′-*O*-methyltransferase families [TrmH (14, 15, 29), TrmL (30), TrmJ (31) and Trm56 (32, 33)

families] have been identified. TrmH catalyzes the methylation of 2′-OH of ribose of G18 and

produces 2′-*O*-methylguanosine at position 18 (Gm18) in tRNA (34-36). Likely, TrmL, TrmJ,

and Trm56 catalyze the 2′-*O*-methylation at positions 34, 32, and 56, respectively, and produces

2′-*O*-methylcytidine (Cm34) and 5-carboxylmethylaminomethyl-2′-*O*-methyluridine

(cnmm^{5}Um34), Cm32 and 2′-*O*-methyluridine (Um32), and Cm56, respectively (30, 32, 33, 37,

38).

In general, 2′-*O*-methylation in RNA tilts the equilibrium of ribose puckering from the

C2′-endo form to the C3′-endo form (39). Therefore, 2′-*O*-methylation stabilizes the local

structure of RNA. Furthermore, because 2′-*O*-methylation prevents cleavage by RNases, it

may prolong the half-life of RNA (40). In the case of thermophilic archaea, 2′-*O*-methylations

at multiple positions in tRNA coordinately stabilize the L-shaped tRNA structure (41). In fact,

thermophile-specific modified nucleosides often contain 2′-*O*-methylation, for example 1,

2′-*O*-dimethylinosine (m^{1}Im) and N^2, N^2, 2′-*O*-trimethylguanosine (m$^2_2$Gm) (42). Furthermore,

2′-*O*-methylations in tRNAs are involved in higher biological phenomena such as

stress-resistance or stress-response (43-46), immune response (47-53) and human diseases (54).

RNA modification enzymes often contain an RNA binding domain(s). The thiouridine

synthetase, methyltransferase and pseudouridine synthase (THUMP) domain is a typical RNA

binding domain (55) and has only been found in tRNA modification enzymes (56).

THUMP-related tRNA modification enzymes can be classified into five types, namely

4-thiouridine synthase (57-59), deaminase (60), methyltransferase (61-66), a partner protein of

acetyltransferase (67, 68) and pseudouridine synthase (69-72).   Although the THUMP domain

can fold autonomously, its affinity for tRNA is very low (73).   The crystal structure of *Bacillus*

anthracis TtuI (previous name ThiI (74); tRNA 4-thiouridine synthetase) revealed that the

THUMP domain is composed of α-helices and β-strands (75) as predicted by a bioinformatics

study (55).   Furthermore, in this report (75), a tRNA-binding model was constructed and the

THUMP domain of TtuI was placed near the CCA-terminus of substrate tRNA because CCA

terminus is essential for the sulfur-transfer reaction by TtuI (76).   THUMP domain recognizes

the 3′-end of RNA (*i.e.,* CCA terminus of tRNA).   This concept was established by numerous

biochemical and structural studies of THUMP-related tRNA modification enzymes (60, 63, 75,

77-81).   The N-terminal ferredoxin-like domain adjusts the angle and distance between the

THUMP and catalytic domains (79).   Furthermore, it should be mentioned that a

THUMP-related structure is contained in an accessory domain of human PUS10 (tRNA

pseudouridine synthase for Ψ54 and Ψ55 in tRNA) (82) and that the CCA terminus in tRNA is

not essential for pseudouridine modification by PUS10 (83).   All THUMP-related tRNA

methyltransferases discovered are tRNA N2-guanine methyltransferases, which produce

N2-methylguanosine (m^2G) and/or N2, N2-dimethylguanosine (m^2_2G) (56).

In 2019, we reported the nucleotide sequence of tRNA[Trp] from *Thermococcus kodakarensis*,

a hyper-thermophilic archaeon (Figure 1A, ref. 84).   This tRNA[Trp] contains Cm6.   However,

the tRNA methyltransferase responsible for the synthesis of Cm6 has not been identified yet.

In this study, we have found that a SPOUT 2′-*O*-methyltransferase with a THUMP domain

produces the Cm6 modification in tRNA.   This tRNA methyltransferase, which has a very

1 unusual domain composition, acts on the $2'$-O-methylation of aminoacyl-stem in tRNA.   In this

2 article, we report the identification of the gene encoding this tRNA methyltransferase, its

3 enzymatic properties, and predicted structure as well as the presence of conserved amino acid

4 residues in the catalytic domain and proposed tRNA-binding mode.

MATERIALS AND METHODS

Materials

[Methyl-^{3}H]-SAM (2.47 TBq/mmol) and [Methyl-^{14}C]-SAM (1.95 GBq/mmol) were purchased

from PerkinElmer.   Non-radioisotope-labeled SAM was obtained from Sigma.   DNA

oligonucleotides were obtained from Thermo Fisher Scientific.

Strains, media, and culture conditions

Thermococcus kodakarensis strains and plasmids are listed in Supplementary Table 1.

Thermococcus kodakarensis KUW1 strain was a kind gift from Dr. Tamotsu Kanai (Toyama

Prefectural University, Japan; ref. 85).   *Thermococcus kodakarensis* KUWA strain, which was

used for the construction of the TK1257 gene deletion (*ΔTK1257*) strain, is auxotrophic for

agmatine (86).   All *T. kodakarensis* strains were cultivated under anaerobic conditions at 85°C

or 93°C in nutrient-rich medium (MA-YT).   MA-YT medium (1 L) contains $0.8 \times$ Marine Art

SF1 reagent (Osaka Yakken Co. Ltd., Osaka, Japan), 5 g yeast extract (Y), and 5 g tryptophan

(T).   2 g elemental sulfur (S^0) or 5 g sodium pyruvate (Pyr) and 5 g Amycol #3-L (Mdx) was

supplied into 1 L MA-YT medium prior to culture.   Amycol #3-L (Nippon Starch Chemical,

Osaka, Japan) contains a mixture of malto-oligosaccharides with lengths from 1 to 12 glucose

units.   Growth characteristics of the KUW1 and the *ΔTK1257* strains were examined in

MA-YT-Pyr-Mdx medium.   Growth at 85°C and 93 °C was examined by monitoring the

optical density at 660 nm.   For all liquid media, resazurine (final concentration, 0.5 mg/L) was

supplemented as an oxygen indicator, and 5.0% Na$_2$S H$_2$O was added until the medium became

colorless.   For colony isolation, solid MA-YT media containing 1 g of gelrite and 0.4 g of

polysulfide per 0.1 L were used.   *Escherichia coli* DH5α strain was used for construction of

the gene disruption plasmids and was grown at 37 °C in LB medium containing ampicillin (final

concentration, 100 mg/L).   Unless otherwise described in this study, chemicals were purchased

from Nacalai Tesque (Kyoto, Japan) except for the gelrite, which was obtained from Wako Pure

Chemicals (Osaka, Japan),

Comparative genomic analysis

The genomes used in this study were obtained from the complete genomes available in the

BioProject of the NCBI database.   Protein sequence analyses were performed using BLASTP

with an inclusion threshold E-value of 1e-10.

Construction of TK1257 gene product expression vector in *Escherichia coli* cells

Thermococcus kodakarensis strain KUW1 genomic DNA was extracted from the cells with a

phenol-chloroform mixture.   TK1257 gene was amplified by polymerase chain reaction (PCR)

from the genomic DNA using the following primers: TK1257_F, 5'-GAA GGA GAT ATA CAT

ATG AAG TTT CTC GTC AAG ACT CAG AGG- 3'; TK1257_R, 5'-GAG CTC GAA TTC

GGA TCC TTA TCA GGA AGA GTC CTC GGC TTT TTC- 3'.   The underlined sequences

show the restriction enzyme NdeI and BamHI cleavage sites, respectively.   The amplified

TK1257 gene was cloned into a pET30a vector using NEBuilder HiFi DNA Assembly Master

Mix (New England Biolabs) in accordance with the manufacturer's instruction.

Expression of TK1257 gene product in *E. coli* cells

The TK1257 expression vector was introduced into *E. coli* BL21 (DE3) Rosetta 2 strain cells.

The transformants were cultivated in 250 ml of LB liquid medium containing 50 µg/ml of

kanamycin at 37ºC for 12 h, then added into 1 L of the same medium, and the culture was

continued at 37ºC.   When the optical density at 600 nm ($OD_{600\ nm}$) reached ~0.8,

isopropyl-ß-D-thiogalactopyranoside (IPTG) was added into the medium to a final

concentration of 1 mM and then the culture was continued at 37ºC for 4 h.   The cells were

collected by centrifugation at 10,000 x g at 4ºC for 20 min, frozen by liquid nitrogen, and stored

at -80ºC.

Purification of TK1257 gene product

0.7 g of wet cells were suspended in 10 mL buffer A [50 mM Tris-HCl (pH 7.6), 5 mM $MgCl_2$,

200 mM KCl, 6 mM 2-mercaptoethanol and 5% (v/v) glycerol] supplemented with 100 µL

protease inhibitor solution (Nacalai Tesque) and then disrupted with an ultra-sonic disruptor

(model VCX-500, Sonics and Materials. Inc) on ice for 30 min.   The supernatant fractions

were collected by centrifugation at 10,000 x g at 4ºC for 20 min.   The sample was incubated at

75ºC for 30 min and denatured proteins were removed by centrifugation at 10,000 x g at 4ºC for

20 min.   The supernatant fraction was loaded on to a HiTrap Q HP (5 ml, Cytiva) column.

The unbound proteins were washed out with 25 mL buffer A.   The bound proteins were eluted

by a linear gradient of KCl from 200 mM to 1000 mM in buffer A.   The fractions containing

TK1257 gene product were combined and then loaded onto a HiTrap Heparin HP column (5 mL,

Cytiva).   The unbound proteins were washed out with 25 mL buffer A.   TK1257 gene product

was eluted with a linear gradient of KCl from 200 mM to 1000 mM in 50 mL buffer A.   The

purity of the fractions was assessed by 10% SDS-polyacrylamide gel electrophoresis (PAGE).

The TK1257 gene product fractions were combined and then the buffer was changed to storage

buffer [50 mM Tris-HCl (pH 7.6), 5 mM $MgCl_2$, 500 mM KCl, 6 mM 2-mercaptoethanol, and

50% (v/v) glycerol] using a Vivaspin 15R centrifugal filter unit (code VS15RH01, 10,000

MWCO, Sartorius).   The purified TK1257 gene product was concentrated to 3.5 mg/mL in a

Vivaspin 15R centrifugal filter unit and stored at - 30ºC.

Preparation of tRNA transcripts

Transfer RNA transcripts were prepared by *in vitro* transcription using T7 RNA polymerase as reported (87). The sequences of oligonucleotides used for the construction of template DNA are given in Supplementary Table 2. The transcripts were purified on 10% polyacrylamide gels containing 7 M urea electrophoresis [10% PAGE (7 M urea)] (88).

Measurements of tRNA methyltransferase activity of TK1257 gene product

The time course experiment was performed as follows. 3 µM TK1257 gene product, 35 µM *T. kodakarensis* tRNA^{Trp} transcript and 100 µM ^{3}H-labeled SAM were incubated at 75ºC in 200 µL of buffer B [50 mM Tris-HCl (pH 7.6), 5 mM MgCl$_2$, 300 mM KCl and 6 mM 2-mercaptoethanol]. At 0, 1, 3, 5, 10, 20, 30 and 40 min time periods, 20 µL of the reaction mixture was spotted onto a Whatman 3MM filter. The filters were washed with 5% (w/v) trichloroacetic acid eight times at 4 ºC for 5 min and then dried. Methyl group incorporation was measured by a liquid scintillation counter. This experiment was independently replicated three times (n = 3). The data was fitted to the single-exponential equation (eq. 1)

$$f_a = A + Be^{-k_{obs} \cdot t} \qquad \text{(eq.1)}$$

where f_a is the methyl group incorporation activity [dpm] at a specific time, A is the maximum methyl group incorporation activity at completion, -B is the amplitude of the observable phase, k_{obs} is the observed first-order rate constant for the methyl group transfer reaction, and t is time.

Modified nucleotide analyses by two-dimensional thin layer chromatography

The reaction was performed at 75ºC in 200 µL of buffer B containing 3 µM TK1257 gene product, 35 µM *T. kodakarensis* tRNA^{Trp} transcript and 100 µM ^{14}C-labeled SAM. The RNA

was treated with phenol-chloroform and then precipitated by ethanol.   The tRNA was

dissolved in 5 µl of 50 mM ammonium acetate (pH 5.3) and digested with 2.5 units of nuclease

P1.   1 µl of standard nucleotide mixture (pA, pG, pC and pU, 0.05 A260 units each) was added

into the sample and then spotted onto a cellulose thin-layer plate (code 1.05565.0001, Merck).

The solvent systems were as follows: first dimension; isobutyric acid/concentrated

NH_4OH/water = 66/1/33 (v/v/v), second dimension; isopropyl alcohol/HCl/water = 70:15:15

(v/v/v) (89).   The standard nucleotides were marked using a pencil with irradiation of

ultra-violet light at 254 nm.   The autoradiogram of the thin-layer plate was obtained using

Typhoon FLA 7000 Laser Scanner (GE Healthcare).

Construction of *T. kodakarensis* Δ*trmTS* strain

To construct the gene disruption plasmid (pUTS) for the Δ*trmTS* strain, the *trmTS* (TK1257)

gene along with their 5′- and 3′-flanking regions (~1.0 kbp) were amplified from *T. kodakarensis*

KUW1 genomic DNA by PCR using the primer sets TS-F/TS-R.   Each fragment was inserted

into the linearized plasmid pUC19 at the 5′- and 3′ blunt ends of partial BamHI sites by the

in-fusion reaction.   Inverse PCR was conducted with the primer sets TS-delF/TS-delR to

remove the target gene (*trmTS*) from the resulting plasmid, and then the fragment was in-fused

with the fragment including the PpadD-*pdaD* (Tk0149) marker gene, which was amplified with

the primer set TS-AGF/TS-AGR from pTK02 plasmid (90).   The pUTS plasmid was used to

transform the *T. kodakarensis* KUWA strain, exhibiting agmatine auxotroph.   Cells grown in

MA-YT-S^0 medium containing 1.25 mg/mL agmatine at 85 °C for 10 h were harvested and

suspended in 200 µl of 0.8 × MA, and kept on ice for 30 min.   3 µg of the plasmid was gently

added into the suspended cells and then kept on ice for 1h.   The cells were cultivated in

MA-YT-S^0 without agmatine at 85 °C for 10 h.   100 µL of the cultures were spread onto

MA-YT solid medium without agmatine and then incubated at 85 °C for 24 h. Only cells with a phenotype exhibiting agmatine prototrophy by homologous recombination (Supplementary Figure S1) can grow in the absence of agmatine. Single colonies were picked up and then cultured in MA-YT-S^0 medium without agmatine at 85°C for 10 h. The candidate *ΔtrmTS* cells were harvested and suspended in distilled water. Genomic DNA was extracted from the cells by phenol-chloroform treatment. DNA sequencing of the recombination region was performed. The primers used for the disruption of *trmTS* gene are listed in Supplementary Table 3.

Western blotting

Customized rabbit anti-TrmTS serum was prepared by Kitayama Labes Co., LTD, Japan. The polyclonal antibody fraction was prepared using a MAbTrap Kit (GE Healthcare). Cell extracts from *T. kodakarensis* wild-type and *ΔtrmTS* strains were prepared as follows. Wet cells (0.1 g) were suspended in 100 µL of 2x SDS-PAGE loading buffer [100 mM Tris-HCl (pH 6.8), 200 mM dithiothreitol, 2.5% SDS, 0.2% bromophenol blue, and 20% (v/v) glycerol], disrupted by sonication and boiled for 5 min. 0.5 µL of the samples were loaded onto a 12.5% SDS-polyacrylamide gel. Half of the gel was stained with Coomassie Brilliant Blue for visualization and half of the gel was electroblotted onto a polyvinylidene difluoride membrane (code 10600029, 0.45 µm, GE Healthcare). TrmTS was detected using horse radish peroxidase (HRP)-conjugated Affinipure Goat Anti-Rabbit IgG (H+L) (code AB-2313567, Proteintech) as a secondary antibody and the chemiluminescence derived from HRP was detected with a Typhoon FLA 7000 Laser Scanner (GE Healthcare) in accordance with the manufacturer's instruction.

Purification of tRNA^{Trp} from *T. kodakarensis* wild-type and *ΔtrmTS* strains using a solid-phase DNA probe method

Total RNAs were prepared from *T. kodakarensis* wild-type and *ΔtrmTS* strains as described previously (91).　The tRNA fraction was further purified by 10% PAGE (7 M urea).　Transfer RNA^{Trp} molecules were purified from the tRNA mixtures using a solid-phase DNA probe method (91, 92).　The sequence of DNA probe was complimentary to U20-A36 in *T. kodakarensi*s tRNA^{Trp}: 5′-TGG AGC CCG CGA TGA TGG A-biotin-3′.

Mass spectrometry for RNA modification analysis

To generate RNase T1 fragments of RNA, 40 pmol of tRNA was digested in a 20 μL mixture containing 100 units of RNase T1 (Thermo Fisher Scientific) and 5 mM ammonium acetate (NH$_4$OAc, pH 5.3) at 37°C for 60 min.　To generate RNase A fragments of RNA, 40 pmol of tRNA was digested with 1 μg of RNase A (Thermo Fisher Scientific) and 1 unit of calf intestinal alkaline phosphatase (NEB), if required, in total 20 μL of 5 mM ammonium bicarbonate (AMBIC) at 37°C for 60 min.　The nucleases were removed by phenol-chloroform extraction as follows: Twenty μl of water-saturated phenol was added to the reaction mixture and vigorously vortexed for 10 s, and then 20 μl of chloroform were added and vortexed as well. After centrifugation at 15,000 ×g for 5 min at room temperature, the upper aqueous layer was transferred to a new tube and dried in vacuum.　The dried samples were dissolved with 10 μL of 70–80% acetonitrile (ACN) and analyzed on a liquid chromatography-mass spectrometry (LC-MS) system (Vanquish Flex UHPLC system and Orbitrap Exploris 240 (Thermo Fisher Scientific)).　The digests were separated on an iHILIC-Fusion column (PEEK, 100×2.1 mm, 3.5 μm, HILCON) with a guard column (PEEK, 20×2.1 mm, 5 μm, HILCON) in 5 mM NH$_4$OAc (pH 5.3) with a linear gradient of acetonitrile (80 to 50% ACN over 20 min) at 150

µl/min. Negative ion scanning ranged from 600 to 2000 *m/z*. Nucleoside measurement was

conducted according to the literature (93) with slight modifications. Twenty pmol of tRNA

was digested by the following 3-step reaction at 37°C for 1 h in each step: (1) 0.03 units of

nuclease P1 (Fujifilm Wako Pure Chemical) in 10 mM NH_4OAc (pH 5.3), (2) 0.04 units of

phosphodiesterase I (Worthington Biochemical) in 50 mM AMBIC and (3) 0.03 units of alkaline

phosphatase (*E. coli* C75, Nippon Gene) in 50 mM AMBIC. Enzymatic digests were separated

on a SunShell C18 column (150×2.1 mm, 2.6 µm, ChromaNik Technologies) with a guard

cartridge column RP (ChromaNik Technologies). The solvent system comprised 5 mM NH_4OAc

(pH 5.3) and ACN with a multi-step gradient (1–11% ACN from 0 to 10 min, 11–24% ACN

from 10 to 20 min, 24–90% ACN from 20 to 25 min, and 90% ACN from 25 to 35 min,

followed by 1% CAN equilibration for 10 min) at 100 µl/min. Positive ion scanning ranged

from 105 to 700 *m/z*. Mass spec data were analyzed using the Xcalibur Qual Browser (Thermo

Fisher Scientific).

Measurements of kinetic parameters of TrmTS for SAM and tRNA

The kinetic parameters for SAM and tRNA were determined by a custom-made Python program

(94), in which the data were fitted to the Michaelis-Menten equation. The following mixtures

were used for the measurement of kinetic parameters of the wild-type TrmTS (or TrmTS mutant

protein) for SAM: 3 µM TrmTS, 35 µM tRNA, various concentrations (0.25, 0.5, 1.0, 2.0, 4.0,

8.0, 16.0, 32.0, 64.0 µM) of SAM (a mixture of non-radioisotope and [3]H-labeled SAM;

molecular ratio 1000:1) in 25 µL of buffer B. In the case of measurement of kinetic

parameters for tRNA transcripts, 3 µM TrmTS (or TrmTS mutant protein), various

concentrations (0.25, 0.5, 1.0, 2.0, 4.0, 8.0, 16.0, 32.0 µM) of tRNA transcript, and 100 µM

SAM were mixed in 25 µL of buffer B. The reaction mixtures were incubated at 75°C for 10

min and then spotted onto Whatman 3MM filters.   Incorporation of ^{3}H-methyl groups was monitored by conventional filter assay.   In brief, the filters were washed in 50 mL 5% (w/v) trichloroacetic acid solution five times.   The filters were dipped into 50 mL 99.5% ethanol to remove water and then dried.   The ^{3}H-methyl group incorporation was measured by a liquid scintillation counter.   The filter assay was independently replicated three times (n = 3).

Alignment analysis

TrmTS homologs were searched using BLAST with default algorithm parameters except for Max target sequences (increased to 500).   To depict the distribution of conserved amino acids, sequence alignments were generated by ClustalW (95) and ESPript 3 (96) with sequences from eight selected organisms including *Thermococcus kodakarensis* with E-value less than 1e-20.

Site-directed mutagenesis and purification of mutant proteins

Site-directed mutagenesis was performed using reverse polymerase chain reaction with the primers listed in Supplementary Table 4.   These mutations were verified by DNA sequencing. All TrmTS mutant proteins were expressed and purified using the same methods for the wild-type TrmTS protein.

Measurements of CD-spectra

TrmTS wild-type and mutant proteins were dialyzed against CD buffer [50 mM Tris-HCl (pH 7.6), 5 mM $MgCl_2$, 500 mM KCl].   After dialysis, the protein concentrations were adjusted to 2 μM.   CD spectra were measured on a JASCO J-820 spectropolarimeter equipped with a JASCO PTC-423L thermo-controller at 25, 50 and 75ºC.   Cuvettes with a 1 mm path length

1 were used.   The spectra were recorded from 300 nm to 200 nm.   The scan speed was 50

2 nm/min.   The spectra shown in this report are the average of five scans.

Inhibition experiments with mutant tRNA^Trp transcripts

5 3 µM TrmTS, 35 µM the wild-type tRNA^{Trp} transcript, various concentrations (0, 10, 20, 30, 40,

6 50, 60, 70, 80 µM) of mutant tRNA^{Trp} transcript, 100 µM ^{3}H-labeled SAM in 25 µL of buffer B

7 were incubated at 75°C for 25 min and then the filter assay was performed as described in

8 "Measurements of tRNA methyltransferase activity of TK1257 gene product" section.   The

9 assay was independently replicated three times (n = 3).

Structural models

12 The structural model of *T. kodakarensis* TrmTS was constructed by AlphaFold 3 (97).   Electro

13 potential map was generated by APBS Electrostatics in PyMOL using default settings (-5.000

14 kBT/e ~ 5.000 kBT/e).   The conserved amino acid residues map was generated by Consurf-DB

15 using default settings (98).

Prediction of the secondary structure of truncated tRNA mutants

The secondary structure of truncated tRNA mutant transcripts (transcript 6, transcript 7, and

transcript   8)   were   predicted   using   the   RNAfold   Web   Server

(http://rna.tbi.univie.ac.at//cgi-bin/RNAWebSuite/RNAfold.cgi) with default parameters.

RESULTS

Search for a candidate gene using comparative genomics

Initially, we searched for a candidate gene for Cm6 formation in *T. kodakarensis* tRNA by

comparative genomics (Fig. 1B and C). 2,306 genes are encoded in the *T. kodakarensis*

genome (99). Fortunately, activities of tRNA modification enzymes in a cell extract of

Pyrococcus furiosus have been reported (100): when *Haloferax volcanii* tRNA$^{\text{Ile}}$ transcript was

used as a substrate, Am6 was produced by the cell extract of *P. furiosus*. The

2′-*O*-methyltransferase responsible for the Am6 formation in *P. furiosus* has not yet been

identified. If our target methyltransferase catalyzes the 2′-*O*-methylation of A6 as well as C6,

an ortholog of the target methyltransferase will be encoded in the *P. furiosus* genome.

Therefore, we predicted that the candidate gene is encoded in both *T. kodakarensis* and *P.*

furiosus genomes. In contrast, tRNAs from *H. volcanii*, *Sulfolobus acidocaldarius*, and

Thermoplasma acidophilum do not contain the Cm6 modification (101-105). We therefore

predicted that these archaea genomes do not contain the candidate gene. Using this analysis,

we were able to disregard a large fraction of the *T. kodakarensis* genome such that only 570

genes remained. Of the 570 genes, ten genes are annotated as RNA methyltransferase or

tRNA-binding protein genes and two genes are annotated as function unknown genes

(Supplementary Table 5). One gene (TK1257) product, which is composed of 355-amino acid,

is predicted to possess N-terminal ferredoxin-like, THUMP and SPOUT RNA methyltransferase

domains. As described in the Introduction, the THUMP domain recognizes the CCA terminus

of tRNA. Furthermore, the majority of the SPOUT RNA methyltransferase superfamily are

2′-*O*-methyltransferases (11). Moreover, the N-terminal ferredoxin-like domain often adjusts

the distance and angle between the THUMP and the Rossmann fold RNA methyltransferase

catalytic domains (79). Therefore, we considered the TK1257 gene to be the best candidate.

Recombinant TK1257 gene product possesses tRNA methyltransferase activity for Cm6

formation.

The TK1257 gene product was expressed in *Escherichia coli* cells and purified as shown in Fig.

2A.   We observed that TK1257 gene product aggregated during dialysis against buffer with

low concentrations of KCl (below 50 mM).   Initially, therefore, we investigated the optimum

KCl concentration for methylation by TK1257 gene product (Supplementary Fig. 1).   In this

experiment, we used *E. coli* tRNA mixtures because we assumed that methylation by the

TK1257 gene product may require other modifications of tRNA.   It should be mentioned that

E. coli tRNAs do not possess the Cm6 modification.   As shown in Supplementary Fig. 1, the

velocity of methyl-group incorporation into *E. coli* tRNA mixture was maximum at 300 mM

KCl.   Therefore, 300 mM KCl was added to the reaction mixtures in this study.   When

TK1257 gene product, *T. kodakarensis* tRNA$^{\text{Trp}}$ transcript, and ^{3}H-labeled SAM were incubated

at 75°C, ^{3}H-methyl groups were clearly incorporated into tRNA$^{\text{Trp}}$ transcript (Fig. 2B).   Thus,

this result shows that methylation by TK1257 gene product does not require any other

modifications in tRNA.   To address the modification position, we prepared three mutant

tRNA$^{\text{Trp}}$ transcripts in addition to the wild-type tRNA$^{\text{Trp}}$: the C6-G67 base pair was replaced by

A6-U67, U6-A67 or G6-C67 base pairs (Fig. 2C).   The mutant tRNA$^{\text{Trp}}$ A6-U67 and U6-A67

transcripts were methylated by TK1257 gene product as well as the wild-type tRNA$^{\text{Trp}}$ transcript

(Fig. 2D).   In contrast, the mutant tRNA$^{\text{Trp}}$ G6-C67 transcript was not methylated (Fig. 2D).

To confirm the modified nucleotides, these tRNA$^{\text{Trp}}$ transcripts were treated with TK1257 gene

product and ^{14}C-labeled SAM, and then digested with nuclease P1.   The resultant nucleotides

were separated by two-dimensional thin-layer chromatography and the autoradiograms of the

thin-layers were obtained (Fig. 2E).   As shown in Fig. 2E, the spot corresponding to ^{14}C-pCm

was clearly detected from the wild-type tRNA[Trp] sample.   Furthermore, [14]C-pAm and [14]C-pUm

were detected from the tRNA[Trp] A6-U67 and tRNA[Trp] U6-A67 mutant samples, respectively

(Fig. 2E).   In contrast, no spot was observed from the tRNA[Trp] G6-C67 mutant sample

(negative control) (Fig. 2E) consistent with the results in Fig. 2D.   Thus, these results clearly

show that the methylation position is the ribose at position 6 in tRNA.   Furthermore, the result

with tRNA[Trp] A6-U67 mutant transcript is in line with the observation in an early study (100):

when *H. volcanii* tRNA[Ile] transcript was used as a substrate, Am6 was formed by *P. furiosus* cell

extract.   The modification position was further checked by LC/MS analysis of methylated

tRNA[Trp] (Supplementary Fig. 2) and tRNA[Pro] (Supplementary Fig. 3) transcripts.   When the

ribose at position 6 in tRNA[Trp] transcript was methylated by TK1257 gene product, RNase A did

not cleave the phosphodiester bond between C6 and G7.   As a result, an RNA fragment,

5′-HO-GGGGGCmGUp-3′ (m/z 876.79) appeared (Supplementary Fig. 2).   Similarly, when

methylated tRNA[Pro] was digested with RNase T1, 5′-HO-CCmGp-3′ (m/z 986.13) appeared

(Supplementary Fig. 3).   Taking these experimental results together, we conclude that the

TK1257 gene product possesses a tRNA methyltransferase activity for Cm6 formation in

tRNA[Trp] and tRNA[Pro] transcripts.   Based on these results, hereafter we describe TK1257 gene

product as TrmTS (Transfer RNA methylation gene product with THUMP and SPOUT

domains).   To investigate the enzymatic properties of TrmTS, kinetic parameters of TrmTS and

tRNA[Trp] transcript were determined at 75°C.   Because the Km value of TrmTS for the

wild-type tRNA[Trp] transcript was determined to be 7.1 μM (Fig. 2F), we used 100 μM RNAs as

substrates in latter experiments.   Furthermore, kinetic analyses of mutant tRNA[Trp] A6-U67 and

U6-A67 transcripts reveal that these mutant tRNA[Trp] transcripts are efficiently methylated by

TrmTS.   It should be noted that all tRNA molecular species encoded in the *T. kodakarensis*

genome possess C6 or G6 (98).   Therefore, the Am6 and Um6 formations by TrmTS do not

occur in native tRNAs.   Moreover, the Km value of TrmTS for SAM was determined to be

20.3 μM (Fig. 2G).   This Km value is slightly large for a Km value of tRNA

methyltransferases for SAM: for example, the Km value of *T. thermophilus* TrmH for SAM is

10.0 μM (26).   Based on this experimental result, we used 200 μM SAM as the methyl group

donor in latter experiments.

The *ΔtrmTS* strain showed a slight growth retardation at 93°C.

To confirm that the *trmTS* gene is expressed in living cells, we constructed a *trmTS* gene

deletion (*ΔtrmTS*) strain (Supplementary Fig. 4).   To detect TrmTS, we prepared anti-TrmTS

polyclonal antibody fraction from an immunized rabbit.   Proteins in the cell-extracts of the

wild-type and *ΔtrmTS* strains were separated by 12.5% SDS-polyacrylamide gel electrophoresis

(PAGE) (Fig. 3A).   20 ng of purified TrmTS was used as a positive control (right lane in Fig.

3A): the band of TrmTS is not visible by Coomassie Brilliant Blue staining because only 20 ng

of protein was used.   The proteins were electro-blotted onto a polyvinylidene difluoride

membrane filter and then western blotting analysis was performed using the anti-TrmTS

polyclonal antibody fraction.   The band corresponding to the TrmTS was clearly observed in

the cell-extract of the wild-type strain.   In contrast, this band was not observed in the

cell-extract of the *ΔtrmTS* strain.   This result clearly shows that TrmTS is expressed in the

wild-type strain and this protein is not expressed in the *ΔtrmTS* strain.   Transfer RNA[Trp]

molecules were purified from the wild-type and *ΔtrmTS* strains by the solid-phase DNA probe

method (91, 92) (Fig. 3B).   The methylated cytidines in these tRNA[Trp] molecules were

analyzed by LC/MS (Fig. 3C).   As shown in Fig. 1A, *T. kodakarensis* tRNA[Trp] possesses two

m[5]C and four Cm modifications.   In the tRNA[Trp] from the wild-type strain, the amount of Cm

is larger than that of m[5]C (Fig. 3C left).   In contrast, in the tRNA[Trp] from the *ΔtrmTS* strain the

amount of Cm is smaller than that of m⁵C due to disappearance of Cm6 (Fig. 3C right). It

should be mentioned that the modification level of each Cm modification is not 100%.

Therefore, this data shows decrease only in the total Cm modification level. Taking these

experimental results together, we conclude that TrmTS is expressed in living *T. kodakarensis*

cells and methylates tRNA$^{\text{Trp}}$ *in vivo*. To address the phenotype of the *ΔtrmTS* strain, we

compared the growth curves of the wild-type and *ΔtrmTS* strains at 85 and 93°C. As shown in

Fig. 3D, the *ΔtrmTS* strain showed a slight growth retardation at 93°C: doubling times of the

wild-type and *ΔtrmTS* strains were 2.19 h and 2.58 h, respectively. When TrmTS was

expressed in the *ΔtrmTS* strain, the growth phenotype was recovered (Fig. 3D right panel, blue

line). Therefore, the presence of TrmTS may help the survival of *T. kodakarensis* at high

temperatures.

Identification of three conserved amino acid sequence motifs and amino acid residues

essential for enzymatic activity

Using the amino acid sequence of *T. kodakarensis* TrmTS as a query, we searched for

TrmTS-like proteins in other organisms. As shown in Fig. 4A, TrmTS-like proteins are

encoded in several thermophilic archaea and eubacteria. The amino acid sequence alignment

predicted that TrmTS contains N-terminal ferredoxin-like, THUMP and SPOUT domains.

Between the THUMP and SPOUT domains, a linker region exists. Also, we identified several

conserved amino acid residues in these TrmTS-like proteins (Fig. 4A). TrmH is known to

possess three conserved amino acid sequence motifs, namely motifs 1, 2 and 3 (9). The

catalytic center is an arginine residue in motif 1 (Arg41 in TrmH) (25-27). The conserved

amino acid residues in motif 2 and motif 3 contribute to the formations of the trefoil-knot

structure and SAM-binding pocket (25, 26). To investigate the catalytically and structurally

important residues in TrmTS, we first examined if TrmTS also possesses these motifs found in

TrmH. A multiple sequence alignment of the SPOUT domain sequences from TrmTS, TrmL,

TrmJ, Trm56, and TrmH revealed that motifs 1, 2, and 3, as characterized in TrmH, are not

clearly conserved in TrmTS (Supplementary Fig. 5A).

Next, we constructed a structural model of TrmTS using AlphaFold 3 (Fig. 4B).   The

parameters of the structural model are given in Supplementary Table 6.   The predicted template

modeling (pTM) value of monomer subunit is 0.78: in general, the pTM value more than 0.5

means that the model has been constructed with high precision.   Subsequently, the conserved

amino acid residues in TrmTS were mapped onto the structural model of the SPOUT domain

(Fig. 4C and Supplementary Fig. 5B).   For this purpose, we constructed a dimer structure

model   of   TrmTS   by   AlphaFold   3,   as   all   previously   characterized   SPOUT

2′-O-methyltransferases   are   known   to   form   a   dimer   structure   (Supplementary   Fig.   5C,

Supplementary Fig. 6, and ref. 11).   The ipTM value of the dimer model is 0.78, indicating a

confident   prediction.   The   location   of   dimer   interface   of   the   model   coincided   with   that

observed in known SPOUT 2′-O-methyltransferases (Supplementary Fig. 6A): the hydrophobic

areas   (area   size,   2635.6 $□^2$;   orange   area   in   the   SPOUT   catalytic   domain   of   TrmTS   in

Supplementary Fig. 6A) of monomer subunits assemble into the dimer structure.   Furthermore,

topological   diagrams   of   TrmTS   and   other   SPOUT   enzymes   showed   similar   structural

arrangements (Supplementary Fig. 6B). These findings strongly suggest that TrmTS forms a

dimer   structure   like   other   SPOUT   2′-O-methyltransferases.   To   determine   whether   TrmTS

forms   a   dimer   structure,   we   analyzed   the   purified   full-length   TrmTS   by   Superdex-200

gel-filtration   column   chromatography   (Supplementary   Fig.   7).   The   protein   was   eluted   in   the

fractions I-IV (Supplementary Fig. 7A and 7B).   The peak observed in the fraction IV

corresponds to the molecular size of TrmTS dimer: the elution points of molecular weight

markers are shown by arrows in Supplementary Fig. 7A.   In addition to the dimer peak, two

additional peaks were observed: one peak in fraction I, which eluted at the void volume, and

another peak in fractions II and III, corresponding to the molecular size of TrmTS tetramer.

Thus, the full-length TrmTS formed three peaks.   As shown in Supplementary Fig. 6A, the

THUMP domain of TrmTS possesses hydrophobic areas.   Therefore, in the absence of

substrate tRNA, purified TrmTS may form a tetramer and multimer structures in addition to a

dimer structure.   Furthermore, because the predicted dimer structure of TrmTS is not globular

(Supplementary Fig. 6A), the molecular weight of full-length TrmTS dimer is not correctly

measured by gel-filtration column chromatography.   To directly assess dimerization mediated

by the SPOUT catalytic domain, we prepared the SPOUT catalytic domain of TrmTS lacking

the N-terminal ferredoxin-like and THUMP domains (inset in Supplementary Fig. 7C).   When

the SPOUT catalytic domain of TrmTS was analyzed by Superdex-200 gel-filtration column

chromatography, the protein was eluted at the location of dimer (molecular weight, 48 kDa) as a

single peak (peak ii in Supplementary Fig. 7C).   Because the peaks i and iii do not contain any

protein (Supplementary Fig. 7D), this experimental result clearly shows that the SPOUT

catalytic domain of TrmTS forms a dimer.

Based on these findings, we performed single-point mutational analyses. We selected

several amino acid residues for mutation, based on structural comparison between TrmTS and

TrmH (Supplementary Fig. 5B and Supplementary Fig. 5C).   The mutant TrmTS proteins were

purified as shown in Fig. 5A.   The relative initial velocities of mutant proteins were assessed

(Fig.5B).   After these pilot experiments, we measured the CD spectra of both wild-type and

mutant proteins (Fig. 6A).   Furthermore, we determined the kinetic parameters of the mutant

proteins (Fig. 6B and C).   These experiments elucidated that mutations in several conserved

amino acid residues (e.g. Lys155, Arg185, Ser229 and Glu326, as shown in Fig. 6A) altered the

peak shape of CD spectra compared to the wild-type protein, suggesting that these residues are

structurally important.   The amino acid sequence alignment and structural comparison between

TrmTS and TrmH suggests that Arg193 is the catalytic residues.   Consistent with this, the

R193A mutant protein lost all enzymatic activity (Fig. 5B).   Furthermore, the CD spectra

indicated that the substitution of this Arginine with alanine does not affect the overall protein

structure (Fig. 6C). Thus, it is likely that the Arg193 residue plays a catalytic role rather than a

structural one. Next, we identified the serine residue conserved in TrmH.   In the case of TrmH,

Ser150 in the motif 3 in one subunit forms a hydrogen bond with the catalytic center (Arg41) in

another subunit and is essential for the methyl-transfer activity (26) (Supplementary Fig. 5C).

Initially, we assumed that Ser299 in TrmTS might be the serine residue that plays the same role

with that Ser in TrmH.   However, the Ser299A mutation did not influence the enzymatic

activity (Fig. 5B).   We checked the structural model again and found that Thr322 could move

near to the catalytic Arg193 residue because Thr322 is located in a polypeptide loop

(Supplementary Fig. 5C).   Therefore, we substituted Thr322 with serine or alanine.   As

expected, the T322A mutant protein completely lost enzymatic activity (Fig. 5B).   Furthermore,

the substitution of Thr322 by serine (T322S mutant) also abolished the enzymatic activity (Fig.

5B).   This result shows that the methyl group of Thr322 is essential for the enzymatic activity

of TrmTS.   Because this residue is conserved as serine in TrmH, the methyl group of Thr322

may possess a specific role in, for example, stabilization of trefoil-knot structure in the case of

TrmTS.   To address this issue, we measured the CD spectra of T322A and T322S mutant

proteins (Fig. 6A).   The CD spectra of these mutant proteins coincide with that of the

wild-type enzyme.   Thus, these results suggest that the mutation of Thr322 residue does not

cause the disruption of trefoil-knot structure.   One possibility is that the methyl group of

Thr322 is involved in SAM-binding (formation of SAM-binding pocket).   To clarify the role of

Thr322, structural study of the complex between TrmTS and SAM (or tRNA) is necessary. Next,

we identified the glutamic acid residue conserved in TrmH.   The structural model suggested

that Glu301 residue in TrmTS corresponds to the glutamic acid residue in the motif 2 in TrmH.

When the Glu301 residue was substituted by alanine, considerable enzymatic activity remained

(Fig. 6B).   This phenomenon was also reported in the mutant protein of Trm56 (27).

Although the glutamic acid residue in the motif 2 is important for folding of the catalytic

domain of TrmH (26), the glutamic acid residue may be not so important for the activities of

archaeal SPOUT 2'-*O*-methyltransferases including Trm56 and TrmTS.   The Glu326 residue in

TrmTS was expected to be the amino acid residue corresponding Asn150 residue in TrmH

which is essential for the TrmH activity (26).   When Glu326 was substituted by alanine, the

enzymatic activity of TrmTS was completely lost (Fig. 5B and Fig. 6C).   Thus, Glu326 residue

is one of the conserved amino acid residues in TrmTS that plays a catalytic role.   The P319

residue in TrmTS was expected to be the amino acid residue corresponding to the P143 residue

in TrmH. This proline residue is semi-conserved in TrmH and is replaced by glutamine in some

eukaryotic SPOUT RNA 2'-*O*-methyltransferases (for example, *Saccharomyces cerevisiae* Trm3

(105)).   We expressed the TrmTS P139A mutant protein in *E. coli* cells, however the expressed

protein was heavily degraded in *E. coli* cells.   Therefore, we could not obtain the mutant

protein.   This result suggests that the Pro319 residue in TrmTS is structurally important.   We

continued the alanine-substitution experiments and found that six amino acid residues (Arg193,

Ser229, Asp270, Thr322, Thr325 and Glu326) were essential for the methyl-transfer activity of

TrmTS (Fig. 6B).   Because Ser229 and Thr325 assemble around the catalytic center (Arg193),

these residues are probably involved in the catalytic mechanism and/or binding of substrate(s).

To clarify the roles of these residues, a crystal structure study of substrate (SAM or

tRNA)-bound forms is necessary: these residues probably move according to substrate-binding.

The TrmTS-specific linker region contains conserved lysine residues.

The multiple sequence alignment of TrmTS-like proteins showed that these proteins possess a

linker region between the THUMP and SPOUT domains (Fig. 4A).   This linker region is not

observed in other SPOUT RNA methyltransferases.   Thus, the conserved amino acid sequence

in the linker region is TrmTS-specific.   Two lysine residues (Lys155 and Lys160) are

conserved in all TrmTS-like proteins.   This linker locates near to the Asp270 residue of the

catalytic domain in another subunit (Fig. 7) and the ε-amino group of Lys160 can contact the

Asp270 residue.   Furthermore, the replacement of Asp270 by alanine causes the complete loss

of enzymatic activity.   To address the role of the linker region, we substituted Lys155 and

Lys160 residues with alanine.   In the case of TrmTS Lys155Ala mutant protein, the

methyl-transfer activity was considerably decreased (Fig. 5B).   The measurement of kinetic

parameters for SAM and tRNA^{Trp} transcript (Fig. 6B and C) revealed that the substitution of

Lys155 with alanine critically decreases the Vmax value.   Because the Lys155 residue is

located far from the catalytic pocket, the Lys155 residue may be involved in movement of

domains during the methylation process.   The expressed TrmTS Lys160Ala mutant protein was

heavily degraded in *E. coli* cells.   Therefore, we could not obtain purified TrmTS Lys160Ala

mutant protein.   This result suggests the importance of the linker region as a structural factor.

The recognition sites of TrmTS in tRNA

To clarify the binding-mode of TrmTS to tRNA, we prepared eight truncated tRNA mutant

transcripts (Fig. 8A and B).   These truncated tRNA mutant transcripts were incubated with

TrmTS and ^{14}C-labeled SAM, and then RNAs were analyzed by 10% PAGE (7 M urea) (Fig. 8C

left panels).   Autoradiograms of the same gels were obtained (Fig. 8C right panels).   When

the D-arm (transcript 2), the anticodon-arm (transcript 3) and the T-arm (transcript 4) were

individually deleted, the methyl group acceptance activities were retained (Fig. 8C).   Thus, the

L-shaped tRNA structure is not essential for methylation by TrmTS.   In contrast, when the

discriminator and CCA terminus (ACCA) were deleted (transcript 5), the methyl group

acceptance activity was completely lost (Fig. 8C lane 5).   Thus, the 3′-terminal region is

essential for methylation by TrmTS.   Furthermore, a deletion mutant of the anticodon- and

T-arms (transcript 6), a deletion mutant of the D- and T-arms (transcript 7) and a deletion mutant

of the D- and anticodon-arms were methylated by TrmTS (Fig.8C lanes 6-8).   In contrast, a

micro-helix RNA, which mimics the aminoacyl-stem, was not methylated by TrmTS at all (Fig.

8C lane 9).   To explore these results deeply, we predicted the secondary structure of these

truncated transcripts (Supplementary Fig. 8).   The transcript 6, in which the anticodon arm and

T-arm were removed, was predicted to have the acceptor stem and two stems forming a

three-way junction with a loop region (Supplementary Fig. 8A).   The transcript 7, in which the

D-arm and T-arm ware removed, was predicted to fold into a long stem and loop structure with

three short bulges (Supplementary Fig. 8B).   Finally, the transcript 8 in which D-arm and

anticodon arm were removed, was predicted to have an extended acceptor stem, a stretch loop,

and a shortened T-arm (Supplementary Fig. 8C).   Of these mutant transcripts, transcript 7

showed relatively low methyl group acceptance activity (Fig. 8C), suggesting that TrmTS

recognize a minimal substrate RNA having a stretch loop connected with at least two stems for

2′-O-methylation, and the L-shaped tRNA structure itself is not essential.

The THUMP domain of TrmTS is required for the methylation in the binding process.

For TrmH, a SPOUT tRNA 2′-O-methyltransferase, the tRNA-binding mode consists of at least

two processes, namely the initial binding and induced-fit processes (107, 108).   To clarify

whether the THUMP domain of TrmTS works in the binding process, we performed an

inhibition experiment.   Two tRNA^{Trp} mutant transcripts were prepared.   One mutant tRNA^{Trp}

transcript is a deletion mutant of the 3'-ACCA.   Another mutant tRNA^{Trp} transcript possesses a

G6-C67 base pair instead of the C6-G67 base pair.   Neither mutant tRNA^{Trp} transcripts were

methylated by TrmTS as shown in Fig. 2 and Fig. 8C.   We tested whether these transcripts act

as competitive inhibitors.   When the concentration of tRNA^{Trp} G6-C67 mutant transcript was

increased, the methyl-transfer to the wild-type tRNA^{Trp} transcript was clearly inhibited (blue in

Fig. 8D): the IC$_{50}$ of tRNA^{Trp} G6-C67 mutant transcript was around 30 µM.   Given that the Km

value of TrmTS for the wild-type tRNA^{Trp} transcript is 7.1 µM (Fig. 7C), this inhibitor possesses

a considerable affinity for TrmTS.   In contrast, when the concentration of tRNA^{Trp} (-ACCA)

was increased, the methylation speed of the wild-type tRNA^{Trp} was not decreased (red in Fig.

8D).   This result clearly shows that the CCA terminal region in tRNA is essential for the

binding of TrmTS.   Thus, the THUMP domain, which captures the CCA terminal region, is

required in the binding process.

Prediction of tRNA binding mode

Figure 9A shows the electro-charges of the TrmTS surface: the blue and red areas represent

positive and negative charges, respectively.   The amino acid residues in the positively charged

areas of the THUMP and SPOUT domain are highly conserved in TrmTS-like proteins (Fig. 9B).

We therefore predicted the tRNA binding mode of TrmTS based on the experimental results (Fig.

9C).   When a tRNA structure was modeled onto the TrmTS structure, both tRNA and TrmTS

structures were crushed.   This observation suggests that the methylation by TrmTS requires an

induced-fit process.   It should be mentioned that clashes between the enzyme and tRNA

structures is often observed in other docking models between SPOUT 2'-O-methyltransferases

and the substrate tRNAs (38, 109, 110).　Furthermore, in the case of TrmD (a member of

SPOUT tRNA methyltransferases), movements of the SPOUT catalytic and C-terminal RNA

binding domains according to SAM and tRNA bindings have been reported (111-113).　In the

binding process, the CCA terminal region is captured by the THUMP domain of TrmTS and

then the three-dimensional core of tRNA contacts the positively charged areas of TrmTS.

During the induced-fit process, movements of the linker region and THUMP domain may be

cased.　Finally, the ribose of C6 in tRNA is introduced into the catalytic pocket and accesses

the catalytic Arg193 residue.

DISCUSSION

In this study, we have discovered a new tRNA methyltransferase using comparative genomics.

This tRNA methyltransferase, TrmTS possesses highly unusual composition of domains,

namely N-terminal ferredoxin-like, THUMP and SPOUT catalytic domains.   Until this study,

all THUMP-related tRNA methyltransferases reported possessed a Rossmann fold catalytic

domain and nucleosides they produced were restricted to m^2G and m^2_2G (56).   Thus, our

findings extend the concept of architecture of tRNA methyltransferases.

Although simple amino acid sequence alignment could not find the three conserved

motifs in TrmH (Supplementary Fig. 5A), our mutagenesis study based on the structural

comparison between the predicted TrmTS structure and TrmH identified the conserved amino

acid residues which are important for the catalysis and structure. These results implies that the

catalytic   mechanism   of   TrmTS   is   probably   common   with   other   SPOUT   RNA

2′-*O*-methyltransferases.   However, one important residue is missing in TrmTS.   The

asparagine residue in the motif 1 of TrmH (Asn35 in *T. thermophilus* TrmH) is involved in the

release of product (S-adenosyl-L-homocysteine) (26).   However, the corresponding amino acid

residue is not found in the sequence of TrmTS.   Because Trm56 does not possess the

asparagine residue as well (27), these archaeal enzymes (TrmTS and Trm56) may possess a

different mechanism for the release of products.   To clarify the details of roles of amino acid

residues in TrmTS, a crystal structural study of the substrate-bound form of TrmTS is necessary.

We found a TrmTS-specific linker region between the THUMP and SPOUT catalytic

domains.   Two lysine residues (Lys155 and Lys160) are highly conserved in the linker region

of the TrmTS-like proteins.   Lys160 in one subunit can interact with Asp270 residue in another

subunit.   Our mutagenesis study suggests that the Lys160 residue is important for the structure

of TrmTS because the TrmTS Lys160Ala mutant protein was heavily degraded in *E. coli* cells.

Furthermore, the substitution of Lys155 with alanine decreases the Vmax value.   Because the Lys155 residue is located far from the catalytic pocket, Lys155 residue may be involved in the movement of domains during the methylation process.   Taking these results together, although the linker region between the THUMP and SPOUT domain is TrmTS-specific, this linker is essential for the activity of TrmTS.

TrmTS can form Am6 and Um6 as well as Cm6 in tRNA.   Thus, the guanine base at position 6 is a negative determinant for the methylation by TrmTS.   This enzymatic property is unique in tRNA 2'-O-methyltransferases.   For example, TrmH (36), Trm56 (114) and archaeal TrmJ (37) strictly recognize G18, C56 and C32, respectively, in tRNA.   Furthermore, TrmL recognizes only pyrimidine nucleosides at position 34 (115).   Moreover, bacterial TrmJ does not recognize the base at position 32 (38).   In the case of TrmTS, the 2-amino group of guanine base at position 6 in tRNA may cause steric hindrance around the catalytic pocket.

The THUMP-domain captures the CCA terminus.   This concept has been established by numerous biochemical and structural studies (60, 63, 66, 75-81).   In this study, we demonstrate that TrmTS, a novel THUMP-related tRNA methyltransferase, also requires the CCA terminal region for methylation.   Furthermore, the deletion of the CCA terminal region from tRNA[Trp] transcript causes the loss of affinity for TrmTS: the deletion mutant tRNA[Trp] transcript does not work as a competitive inhibitor.   Thus, our biochemical experiment shows that the interaction between the CCA terminus in tRNA and the THUMP domain of TrmTS is essential for the binding process.   Our hypothetical model in Fig. 9C suggests that the three-dimensional core of substrate tRNA contacts the positively charged areas of TrmTS and then an induced-fit process occurs.   To clarify the precise interaction between TrmTS and tRNA, the structural study of the complex between TrmTS and tRNA is necessary.

The physiological role of the Cm6 modification by TrmTS is unclear.   Because

TrmTS-like proteins are encoded in the genomes from thermophilic archaea and bacteria, we assumed that deletion of the *trmTS* gene might affect growth at high temperatures. However, the *ΔtrmTS* strain showed only a slight growth retardation at high temperatures. In *T. kodakarensis* tRNA^{Trp}, numerous modified nucleosides exist (Fig. 1). Therefore, the Cm6 modification may work with other modifications at high temperatures. This idea is in line with the previous report that initiator tRNA^{Met} from *Pyrodictium occultum*, a hyper-thermophilic archaeon, is stabilized by 2′-*O*-methylation at multiple positions (41). Furthermore, in the case of *Pseudomonas aeruginosa*, the Am32 modification by TrmJ confers oxidative-stress resistance (42). Therefore, there is a possibility that the Cm6 modification by TrmTS may function in a stressful environment. To clarify the physiological role of Cm6 modification, further study is required.

ACKNOWLEDGEMENTS

We thank the RI room staffs in the Advanced Research Support Center (Ehime University) for use of radioisotope compounds. Mass spectrometry experiments were conducted with the support of University of the Ryukyus, Research Facility Center.

DATA AVAILABILITY

All data used in this study are available from the corresponding authors upon request.

SUPPLEMENTARY DATA

Supporting information is avairable at NAR Online: Supplementary Figures 1-8, and Supplementary Tables 1-6.

FUNDING

This work was supported by Grant-in-Aid for Scientific Research from the JAPAN Society for the Promotion of Science (JSPS) KAKENHI (JP24K09352 to AH, and JP20H03211 and JP24K09381 to HH) and Funding from the Institute for Fermentation, Osaka, Grant Number G-2024-2-059 to RY.

CONFLICT OF INTERST

The authors declare no conflict of interest.

REFERENCES

1. Cappannini, A., Ray, A., Purta, E., Mukherjee, S., Boccaletto, P., Moafinejad, S. N., Lechner, A., Barchet, C., Klaholz, B. P., Stefaniak, F. and Bujnicki, J. M. (2023) MODOMICS: a database of RNA modifications and related information. 2023 update. *Nucleic Acids Res*. **52**, D239-D244.

2. Lorenz, C., Lünse, C. E. and Mörl, M. (2017) tRNA Modifications: Impact on Structure and Thermal Adaptation. *Biomolecules* **7**, pii: E35.

3. Väre, V. Y., Eruysal, E. R., Narendran, A., Sarachan, K. L. and Agris, P. F. (2017) Chemical and Conformational Diversity of Modified Nucleosides Affects tRNA Structure and Function. *Biomolecules*, **7**, pii E29.

4. Hori, H. (2014) Methylated nucleosides in tRNA and tRNA methyltransferases. *Front. Genet*. **5**, 144.

5. Urbonavicius, J., Skouloubris, S., Myllykallio, H. and Grosjean, H. (2005) Identification of a novel gene encoding a flavin-dependent tRNA:m^5U methyltransferase in bacteria--evolutionary implications. *Nucleic Acids Res*. **33**, 3955-3964.

6. Moukadiri, I., Prado, S., Piera, J., Velázquez-Campoy, A., Björk, G. R. and Armengod M. E. (2009) Evolutionarily conserved proteins MnmE and GidA catalyze the formation of two methyluridine derivatives at tRNA wobble positions. *Nucleic Acids Res*. **37**, 7177-7193.

7. Lartigue, C., Lebaudy, A., Blanchard, A., El Yacoubi, B., Rose, S., Grosjean, H., and Douthwaite, S. (2014) The flavoprotein Mcap0476 (RlmFO) catalyzes m^5U1939 modification in *Mycoplasma capricolum* 23S rRNA. *Nucleic Acids Res*. **42**, 8073-8082.

8. Schubert, H. G., Blumenthal, R. M. and Cheng, X. (2003) Many paths to methyltransfer: a chronicle of convergence. *Trends Biochem. Sci*. **28**, 329-32.

9. Anantharaman, V., Koonin, E. V. and Aravind, L. (2002) SPOUT: a class of methyltransferases that includes *spoU* and *trmD* RNA methylase superfamilies, and novel superfamilies of predicted prokaryotic RNA methylases. *J. Mol. Microbiol. Biotechnol*. **4,** 71-75.

10. Tkaczuk, K. L.*;* Dunin-Horkawicz, S.*;* Purta, E. *and* Bujnicki, J. M. *(2007)* Structural and evolutionary bioinformatics of the SPOUT superfamily of methyltransferases. *BMC Bioinformatics* **8***,* 73.

11. Hori, H. (2017) Transfer RNA methyltransferases with a SpoU-TrmD (SPOUT) fold and their modified nucleosides in tRNA. *Biomolecules* **7**, 23.

12. Krishnamohan, A. and Jackman, J. E. (2019) A Family Divided: Distinct Structural and Mechanistic Features of the SpoU-TrmD (SPOUT) Methyltransferase Superfamily. *Biochemistry* **58**, 336-345.

13. Strassler, S. E., Bowles, I. E., Dey, D., Jackman, J. E. and Conn, G. L. (2022) Tied up in knots: Untangling substrate recognition by the SPOUT methyltransferases. *J. Biol. Chem.* **298**, 102393.

14. Gustafsson, C., Reid, R., Greene, P. J. and Santi, D. V. (1996) Identification of new RNA modifying enzymes by iterative genome search using known modifying enzymes as probes. *Nucleic Acids Res.* **24**, 3756-3762.

15. Persson, B. C., Jäger, G. and Gustafsson, C. (1997) The spoU gene of *Escherichia coli*, the fourth gene of the spot operon, is essential for tRNA (Gm18) 2'-*O*-methyltransferase activity. *Nucleic Acid Res*. **25**, 3969-3973.

16. Byström, A.S. and Björk, G. R. (1982) Chromosomal location and cloning of the gene (trmD) responsible for the synthesis of tRNA (m^{1}G) methyltransferase in *Escherichia coli* K-12. *Mol. Gen. Genet*. **188**, 440-446.

17. Hou, Y. M., Matsubara, R., Takase, R., Masuda, I. and Sulkowska, J. I.. (2017) TrmD: A Methyl Transferase for tRNA Methylation With m^{1}G37. *Enzymes* **41**, 89-115.

18. Kimura, S., Miyauchi, K., Ikeuchi, Y., Thiaville, P. C., de Crécy-Lagard, V., and Suzuki, T. (2014) Discovery of the β-barrel-type RNA methyltransferase responsible for N6-methylation of N6-threonylcarbamoyladenosine in tRNAs. *Nucleic Acids Res.* **42**, 9350-9365.

19. Michel, G., Sauvé, V., Larocque, R., Li, Y., Matte, A. and Cygler, M. (2002) The structure of the RlmB 23S rRNA methyltransferase reveals a new methyltransferase fold with a unique knot. *Structure*, **10**, 1303-1315.

20. Nureki, O., Shirouzu, M., Hashimoto, K., Ishitani, R., Terada, T., Tamakoshi, M., Oshima, T., Chijimatsu, M., Takio, K., Vassylyev, D. G., Shibata, T., Inoue, Y., Kuramitsu, S. and Yokoyama, S. (2002) An enzyme with a deep trefoil knot for the active-site architecture. *Acta Cryst. Section D, Biol. Cryst.* **D58**, 1129-1137.

21. Lim, K., Zhang, H., Tempczyk, A., Krajewski, W., Bonander, N., Toedt, J., Howard, A., Eisenstein, E., and Herzberg, O. (2003) Structure of the YibK methyltransferase from *Haemophilus influenzae* (HI0766): a cofactor bound at a site formed by a knot. *Proteins* **51**, 56-67.

22. Ahn, H. J., Kim, H. W., Yoon, H. J., Lee, B., Suh, S. S., and Yang, J. K. (2003) Crystal structure of tRNA (m^{1}G37) methyltransferase: insights into tRNA recognition. *EMBO J.* **22**, 2593-2603.

23. Elkins, P. A., Watts, J. M., Zalacain, M., van Thiel, A., Vitazka, P. R., Redlak, M., Andraos-Selim, C., Rastinejad, F. and Holmes, W. M. (2003) Insights into catalysis by a knotted TrmD tRNA methyltransferase. *J. Mol. Biol.* **333**, 931-949.

24. Liu, J., Wang, W., Shin, D. H., Yokota, H., Kim, R. and Kim, S.-H. (2003) Crystal structure

of tRNA (m^{1}G37) methyltransferase from *Aquifex aeolicus* at 2.6 A resolution: a novel methyltransferase fold. *Proteins* **53**, 326-328.

25. Nureki, O, Watanabe, K., Fukai, S., Ishii, R., Endo, Y., Hori, H. and Yokoyama, S. (2004) Deep knot structure for construction of active site and cofactor binding site of tRNA modification enzyme. *Structure* **12**, 593-604.

26. Watanabe, K., Nureki, O., Fuakai, S., Ishii, R., Okamoto, H., Yokoyama, S., Endo, Y. and Hori, H. (2005) Roles of conserved amino acid sequence motifs in the SpoU (TrmH) RNA methyltransferase family. *J. Biol. Chem*. **280**, 10368-10377.

27. Kuratani, M., Bessho, Y., Nishimoto, M., Grosjean, H. and Yokoyama, S. (2008) Crystal structure and mutational study of a unique SpoU family archaeal methylase that forms 2′-*O*-methylcytidine at position 56 of tRNA. *J. Mol. Biol*. 375, 1064-1075.

28. Liu, R. J.; Zhou, M.; Fang, Z. P.; Wang, M.; Zhou, X. L.; Wang, E. D. (2013) The tRNA recognition mechanism of the minimalist SPOUT methyltransferase, TrmL. *Nucleic Acids Res*. **41**, 7828-7842.

29. Hori, H., Suzuki, T., Sugawara, K., Inoue, Y., Shibata, T., Kuramitsu S., Yokoyama, S., Oshima, T., and Watanabe, K. (2002) Identification and characterization of tRNA (Gm18) methyltransferase from *Thermus thermophilus* HB8: domain structure and conserved amino acid sequence motifs *Genes Cells,* **7**, 259-272.

30. Benítez-Páez, A., Villarroya, M., Douthwaite, S., Gabaldón, T. and Armengod, M. E. (2010) YibK is the 2′-*O*-methyltransferase TrmL that modifies the wobble nucleotide in *Escherichia coli* tRNA(Leu) isoacceptors. RNA **16**, 2131-2143.

31. Purta, E., van Vliet, F., Tkaczuk, K. L., Dunin-Horkawicz, S., Mori, H., Droogmans, L. and Bujnicki, J. M. (2006) The *yfhQ* gene of *Escherichia coli* encodes a tRNA:Cm32/Um32 methyltransferase. *BMC Mol. Biol. 7, 23*.

32. Clouet-d′Orval, B., Gaspin, C. and Mougin, A. (2005) Two different mechanisms for tRNA ribose methylation in Archaea: a short survey. *Biochimie* **87**, 889-895.

33. Renalier, M. H., Joseph, N., Gaspin, C., Thebault, P. and Mougin, A. (2005) The Cm56 tRNA modification in archaea is catalyzed either by a specific 2′-*O*-methylase or a C/D sRNP. RNA **11**, 1051-1063.

34. Kumagai, I., Watanabe, K. and Oshima, T. (1980) Thermally induced biosynthesis of 2′-O-methylguanosine in tRNA from an extreme thermophile, *Thermus thermophilus* HB27. *Proc. Natl. Acad. Sci. USA* **77**, 1922-1926.

35. Matsumoto, T., Nishikawa, K., Hori, H., Ohta, T., Miura, K., and Watanabe, K. (1990) Recognition sites of tRNA by a thermostable tRNA(guanosine-2′-)-methyltransferase from *Thermus thermophilus* HB27. *J. Biochem*. **107**, 331-338.

36. Hori, H., Yamazaki, N., Matsumoto, T., Watanabe, Y., Ueda, T., Nishikawa, K., Kumagai, I.

and Watanabe, K. (1998) Substrate recognition of tRNA (Guanosine-2′-)-methyltransferase from *Thermus thermophilus* HB27 *J. Biol. Chem.*, **273**, 25721-25727.

37. Somme, J., Van Laer, B., Roovers, M., Steyaert, J., Versées, W. and Droogmans, L. (2014) Characterization of two homologous 2′-*O*-methyltransferases showing different specificities for their tRNA substrates. RNA **20**, 1257-1271.

38. Liu, R. J., Long, T., Zhou, M., Zhou, X. L. and Wang, E. D. (2015) tRNA recognition by a bacterial tRNA Xm32 modification enzyme from the SPOUT methyltransferase superfamily *Nucleic Acids Res.* **43**, 7489-7503.

39. Kawai, G., Yamamoto, Y., Kamimura, T., Masegi, T., Sekine, M., Hata, T., Iimori, T., Watanabe, T., Miyazawa, T. and Yokoyama, S. (1992) Conformational rigidity of specific pyrimidine residues in tRNA arises from posttranscriptional modifications that enhance steric interaction between the base and the 2′-hydroxyl group. *Biochemistry*, **43**, 1040-1046.

40. Kumagai, I., Watanabe, K. and Oshima, T. (1982) A thermostable tRNA (guanosine-2′)-methyltransferase from *Thermus thermophilus* HB27 and the effect of ribose methylation on the conformational stability of tRNA. *J. Biol. Chem.* **257**, 7388-7395.

41. Ushida, C., Muramatsu, T., Mizushima, H., Ueda, T., Watanabe, K., Stetter, K. O., Crain, P. F., McCloskey, J. A. and Kuchino, Y. (1996) Structural feature of the initiator tRNA gene from Pyrodictium occultum and the thermal stability of its gene product, tRNA(imet). *Biochimie* **78**, 847-855.

42. Hori, H., Kawamura, T., Awai, T., Ochi, A., Yamagami, R., Tomikawa, C. and Hirata, A. (2018) Transfer RNA Modification Enzymes from Thermophiles and Their Modified Nucleosides in tRNA. *Microorganisms* **6**, 110.

43. Jaroensuk, J., Atichartpongkul, S., Chionh, Y. H., Wong, Y. H., Liew, C. W., McBee, M. E., Thongdee ,N., Prestwich, E. G., DeMott, M. S., Mongkolsuk, S., Dedon, P. C., Lescar, J. and Fuangthong, M. (2016) Methylation at position 32 of tRNA catalyzed by TrmJ alters oxidative stress response in *Pseudomonas aeruginosa*. *Nucleic Acids Res.* **44**, 10834-10848.

44. Han, L., Guy, M. P., Kon, Y. and Phizicky, E. M. (2018) Lack of 2′-O-methylation in the tRNA anticodon loop of two phylogenetically distant yeast species activates the general amino acid control pathway. *PLoS Genet.* **14**, e1007288.

45. Galvanin, A., Vogt, L. M., Grober, A., Freund, I., Ayadi, L., Bourguignon-Igel, V., Bessler, L., Jacob, D., Eigenbrod, T., Marchand, V., Dalpke, A., Helm, M., and Motorin, Y. (2020) Bacterial tRNA 2′-O-methylation is dynamically regulated under stress conditions and modulates innate immune response. *Nucleic Acids Res.* **48**, 12833-12844.

46. Endres, L., Rose, R. E., Doyle, F., Rahn, T., Lee, B., Seaman, J., McIntyre, W. D. and Fabris, D. (2020) 2′-O-ribose methylation of transfer RNA promotes recovery from oxidative stress in *Saccharomyces cerevisiae*. *PLoS One* **15**, e0229103.

47. Gehrig, S., Eberle, M. E., Botschen, F., Rimbach, K., Eberle, F., Eigenbrod, T., Kaiser, S., Holmes, W. M., Erdmann, V. A., Sprinzl, M., Bec, G., Keith, G., Dalpke, A. H. and Helm, M. (2012) Identification of modifications in microbial, native tRNA that suppress immunostimulatory activity. *J. Exp. Med.* **209**, 225-233.

48. Jöckel, S., Nees, G., Sommer, R., Zhao, Y., Cherkasov, D., Hori, H., Ehm, G., Schnare, M., Nain, M., Kaufmann, A. and Bauer, S. (2012) The 2′-O-methylation status of a single guanosine controls transfer RNA-mediated Toll-like receptor 7 activation or inhibition. *J. Exp. Med.* **209**, 235-241.

49. Rimbach, K., Kaiser, S., Helm, M., Dalpke, A. H. and Eigenbrod, T. (2015) 2′-O-Methylation within Bacterial RNA Acts as Suppressor of TLR7/TLR8 Activation in Human Innate Immune Cells. *J. Innate Immun.* **7**, 482-493.

50. Schmitt, F. C. F., Freund, I., Weigand, M. A., Helm, M., Dalpke, A. H. and Eigenbrod, T. (2017) Identification of an optimized 2′- *O*-methylated trinucleotide RNA motif inhibiting Toll-like receptors 7 and 8. *RNA* **23**, 1344-1351.

51. Keller, P., Freund, I., Marchand, V., Bec, G., Huang, R., Motorin, Y., Eigenbrod, T., Dalpke, A. and Helm, M. (2018) Double methylation of tRNA-U54 to 2′-O-methylthymidine (Tm) synergistically decreases immune response by Toll-like receptor 7. *Nucleic Acids Res.* **46**, 9764-9775.

52. Freund, I., Eigenbrod, T., Helm, M. and Dalpke, A. H. (2019) RNA Modifications Modulate Activation of Innate Toll-Like Receptors. *Genes* **10**, 92.

53. Freund, I., Buhl, D. K., Boutin, S., Kotter, A., Pichot, F., Marchand, V., Vierbuchen, T., Heine, H., Motorin, Y., Helm, M., Dalpke, A. H. and Eigenbrod, T. (2019) 2′-*O*-methylation within prokaryotic and eukaryotic tRNA inhibits innate immune activation by endosomal Toll-like receptors but does not affect recognition of whole organisms. *RNA* **25**, 869-880.

54. Chujo, T. and Tomizawa, K. (2021) Human transfer RNA modopathies: diseases caused by aberrations in transfer RNA modifications. *FEBS J.* **288**, 7096-7122.

55. Aravind, L. and Koonin, E. V. (2001) THUMP--a predicted RNA-binding domain shared by 4-thiouridine, pseudouridine synthases and RNA methylases. *Trends Biochem. Sci.* **26**, 215–217.

56. Hori, H. (2023) Transfer RNA Modification Enzymes with a Thiouridine Synthetase, Methyltransferase and Pseudouridine Synthase (THUMP) Domain and the Nucleosides They Produce in tRNA. *Genes* **14**, 382.

57. Mueller, E. G., Buck, C. J., Palenchar, P. M., Barnhart, L. E. and Paulson, J. L. (1998) Identification of a gene involved in the generation of 4-thiouridine in tRNA. *Nucleic Acids Res.* **26**, 2606–2610.

58. Liu, Y., Vinyard, D. J., Reesbeck, M. E., Suzuki, T., Manakongtreecheep, K., Holland, P. L.,

Brudvig, G. W. and Söll, D. (2016) A [3Fe-4S] cluster is required for tRNA thiolation in archaea and eukaryotes. *Proc. Natl. Acad. Sci. USA* **113**, 12703–12708.

59. He, N., Zhou, J., Bimai, O., Oltmanns, J., Ravanat, J. L., Velours, C., Schünemann, V., Fontecave, M. and Golinelli-Pimpaneau, B. (2022) A subclass of archaeal U8-tRNA sulfurases requires a [4Fe-4S] cluster for catalysis. *Nucleic Acids Res.* **50**, 12969–12978.

60. Randau, L., Stanley, B. J., Kohlway, A., Mechta, S., Xiong, Y. and Söll, D. (2009) A cytidine deaminase edits C to U in transfer RNAs in Archaea. *Science* **324**, 657–659.

61. Armengaud, J., Urbonavicius, J., Fernandez, B., Chaussinand, G., Bujnicki, J. M. and Grosjean, H. (2004) N2-methylation of guanosine at position 10 in tRNA is catalyzed by a THUMP domain-containing, S-adenosylmethionine-dependent methyltransferase, conserved in Archaea and Eukaryota. *J. Biol. Chem.* **279**, 37142–37152.

62. Purushothaman, S. K., Bujnicki, J. M., Grosjean, H. and Lapeyre, B. (2005) Trm11p and Trm112p are both required for the formation of 2-methylguanosine at position 10 in yeast tRNA. *Mol. Cell Biol.* **25**, 4359–4370.

63. Wang, C., Van Tran, N., Jactel, V., Guérineau, V. and Graille, M. (2020) Structural and functional insights into *Archaeoglobus fulgidus* m^{2}G10 tRNA methyltransferase Trm11 and its Trm112 activator. *Nucleic Acids Res.* **48**, 11068–11082.

64. Roovers, M., Oudjama, Y., Fislage, M., Bujnicki, J. M., Versées, W. and Droogmans, L. (2012) The open reading frame TTC1157 of *Thermus thermophilus* HB27 encodes the methyltransferase forming N^2-methylguanosine at position 6 in tRNA. *RNA* **18**, 815–824.

65. Menezes, S., Gaston, K. W., Krivos, K. L., Apolinario, E. E., Reich, N. O., Sowers, K. R., Limbach, P. A. and Perona, J. J. (2011) Formation of m^{2}G6 in *Methanocaldococcus jannaschii* tRNA catalyzed by the novel methyltransferase Trm14. *Nucleic Acids Res.* **39**, 7641–7655.

66. Yang, W. Q., Xiong, Q. P., Ge, J. Y., Li, H., Zhu, W. Y., Nie, Y., Lin, X., Lv, D., Li, J., Lin, H. and Liu, R. J. (2021) THUMPD3-TRMT112 is a m^{2}G methyltransferase working on a broad range of tRNA substrates. *Nucleic Acids Res.* **49**, 11900–11919.

67. Sharma, S., Langhendries, J. L., Watzinger, P., Kötter, P., Entian, K. D. and Lafontaine, D. L. (2015) Yeast Kre33 and human NAT10 are conserved 18S rRNA cytosine acetyltransferases that modify tRNAs assisted by the adaptor Tan1/THUMPD1. *Nucleic Acids Res.* **43**, 2242–2258.

68. Sas-Chen, A., Thomas, J. M., Matzov, D., Taoka, M., Nance, K. D., Nir, R., Bryson, K. M., Shachar, R., Liman, G. L. S., Burkhart, B. W., Gamage, S. T., Nobe, Y., Briney, C. A., Levy, M. J., Fuchs, R. T., Robb, G. B., Hartmann, J., Sharma, S., Lin, Q., Florens, L., Washburn, M. P., Isobe, T., Santangelo, T. J., Shalev-Benami, M., Meier, J. L. and Schwartz, S. (2020) Dynamic RNA acetylation revealed by quantitative cross-evolutionary mapping. *Nature* **583**,

638–643.

69. Roovers, M., Hale, C., Tricot, C., Terns, M. P., Terns, R. M., Grosjean, H. and Droogmans, L. (2006) Formation of the conserved pseudouridine at position 55 in archaeal tRNA. *Nucleic Acids Res*. **34**, 4293–4301.

70. Gurha, P. and Gupta, R. (2008) Archaeal Pus10 proteins can produce both pseudouridine 54 and 55 in tRNA. *RNA* **14**, 2521–2527.

71. Deogharia, M., Mukhopadhyay, S., Joardar, A. and Gupta, R. (2019) The human ortholog of archaeal Pus10 produces pseudouridine 54 in select tRNAs where its recognition sequence contains a modified residue. *RNA* **25**, 336–351.

72. Mukhopadhyay, S., Deogharia, M. and Gupta, R. (2021) Mammalian nuclear TRUB1, mitochondrial TRUB2, and cytoplasmic PUS10 produce conserved pseudouridine 55 in different sets of tRNA. *RNA* **27**, 66–79.

73. Gabant, G., Auxilien, S., Tuszynska, I., Locard, M., Gajda, M. J., Chaussinand, G., Fernandez, B., Dedieu, A., Grosjean, H., Golinelli-Pimpaneau, B., Bujnicki, J. M. and Armengaud, J. (2006) THUMP from archaeal tRNA:m^2_2G10 methyltransferase, a genuine autonomously folding domain. *Nucleic Acids Res.* **34**, 2483-2494.

74. He, N., Zhou, J., Bimai, O., Oltmanns, J., Ravanat, J. L., Velours, C., Schünemann, V., Fontecave M, Golinelli-Pimpaneau B. (2022) A subclass of archaeal U8-tRNA sulfurases requires a [4Fe-4S] cluster for catalysis. *Nucleic Acids Res.* **50**, 12969-12978.

75. Waterman, D. G., Ortiz-Lombardía, M., Fogg, M. J., Koonin, E. V. and Antson, A. A. (2006) Crystal structure of *Bacillus anthracis* ThiI, a tRNA-modifying enzyme containing the predicted RNA-binding THUMP domain. *J. Mol. Biol*. **356**, 97–110.

76. Lauhon, C.T., Erwin, W.M. and Ton, G.N. (2004) Substrate specificity for 4-thiouridine modification in *Escherichia coli*. Substrate specificity for 4-thiouridine modification in *Escherichia coli*. *J. Biol. Chem*. **279**, 23022–23029.

77. Neumann, P., Lakomek, K., Naumann, P. T., Erwin, W. M., Lauhon, C. T. and Ficner, R. (2014) Crystal structure of a 4-thiouridine synthetase-RNA complex reveals specificity of tRNA U8 modification. *Nucleic Acids Res*. **42**, 6673–6685.

78. Fislage, M., Roovers, M., Tuszynska, I., Bujnicki, J. M., Droogmans, L. and Versées, W. (2012) Crystal structures of the tRNA:m^2G6 methyltransferase Trm14/TrmN from two domains of life. *Nucleic Acids Res*. **40**, 5149–5161.

79. Hirata, A., Nishiyama, S., Tamura, T., Yamauchi, A. and Hori, H. (2016) Structural and functional analyses of the archaeal tRNA m^2G/m^2_2G10 methyltransferase aTrm11 provide mechanistic insights into site specificity of a tRNA methyltransferase that contains common RNA-binding modules. *Nucleic Acids Res*. **44**, 6377–6390.

80. Bourgeois, G., Marcoux, J., Saliou, J. M., Cianférani, S. and Graille, M. (2017) Activation

mode of the eukaryotic m^2G10 tRNA methyltransferase Trm11 by its partner protein Trm112. *Nucleic Acids Res*. **45**, 1971–1982.

81. Nishida, Y., Ohmori, S., Kakizono, R., Kawai, K., Namba, M., Okada, K., Yamagami, R., Hirata, A. and Hori, H. (2022) Required Elements in tRNA for Methylation by the Eukaryotic tRNA (Guanine-N^2-) Methyltransferase (Trm11-Trm112 Complex). *Int. J. Mol. Sci*. **23**, 4046.

82. McCleverty, C. J., Hornsby, M., Spraggon, G. and Kreusch, A. (2007) Crystal structure of human Pus10, a novel pseudouridine synthase. *J. Mol. Biol*. **373**, 1243–1254.

83. Deogharia, M., Mukhopadhyay, S., Joardar, A. and Gupta, R. (2019) The human ortholog of archaeal Pus10 produces pseudouridine 54 in select tRNAs where its recognition sequence contains a modified residue. *RNA* **25**, 336–351.

84. Hirata, A., Suzuki, T., Nagano, T., Fujii, D., Okamoto, M., Sora, M., Lowe, T. M., Kanai, T., Atomi, H., Suzuki, T. and Hori, H. (2019) Distinct Modified Nucleosides in tRNA^{Trp} from the Hyperthermophilic Archaeon *Thermococcus kodakarensis* and Requirement of tRNA m^2G10/m^2_2G10 Methyltransferase (Archaeal Trm11) for Survival at High Temperatures. *J. Bacteriol*. **201**, e00448–e00519.

85. Sato, T., Fukui, T., Atomi, H. and Imanaka, T. (2003) Targeted gene disruption by homologous recombination in the hyperthermophilic archaeon *Thermococcus kodakaraensis* KOD1. *J. Bacteriol*. **185**, 210–220.

86. Kawamura, T., Hirata, A., Ohno, S., Nomura, Y., Nagano, T., Nameki, T., Yokogawa, T. and Hori, H. (2016) Multisite-specific archaeosine tRNA-guanine transglycosylase (ArcTGT) from *Thermoplasma acidophilum*, a thermo-acidophilic archaeon. *Nucleic Acids Res*. **44**, 1894-1908.

87. Matsuda, T., Hori, H. and Yamagami, R. (2024) Rational design of oligonucleotides for enhanced in vitro transcription of small RNA. *RNA* **30**, 710-727.

88. Yamagami R, Hori H. (2023) Functional analysis of tRNA modification enzymes using mutational profiling. *Methods Enzymol*. 692:69-101.

89. Keith, G. (1995) Mobilities of modified ribonucleotides on two-dimensional cellulose thin-layer chromatography. *Biochimie* **77**, 142-144.

90. Nagaoka, E., Hidese, R., Imanaka, T. and Fujiwara, S. (2013) Importance and determinants of induction of cold-induced DEAD RNA helicase in the hyperthermophilic archaeon *Thermococcus kodakarensis*. *J Bacteriol* **195**, 3442-3450.

91. Kazayama, A., Yamagami, R., Yokogawa, T. and Hori, H. (2015) Improved solid-phase DNA probe method for tRNA purification: large scale preparation, and alteration of DNA fixation. *J. Biochem*. **157**, 411-418.

92. Yokogawa, T., Kitamura, Y., Nakamura, D., Ohno, S., and Nishikawa, K. (2010)

Optimization of the hybridization-based method for purification of thermostable tRNAs in the presence of tetraalkylammonium salts. *Nucleic Acids Res.* **38**, e89.

93. Takakura, M., Ishiguro, K., Akichika, S., Miyauchi, K. and Suzuki, T. (2019) Biogenesis and functions of aminocarboxypropyluridine in tRNA. *Nat. Commun*. **10**, 5542.

94. Fujita, S., Sugio, Y., Kawamura, T., Yamagami, R., Oka, N., Hirata, A., Yokogawa, T. and Hori, H. (2024) ArcS from *Thermococcus kodakarensis* transfers L-lysine to preQ$_0$ nucleoside derivatives as minimum substrate RNAs. *J. Biol. Chem.* **300**, 107505.

95. Thompson, J. D., Gibson, T. J. and Higgins, D. G. (2002) Multiple sequence alignment using ClustalW and ClustalX. *Curr. Protoc. Bioinformatics* Chapter 2, Unit 2.3. doi: 10.1002/0471250953.bi0203s00.

96. Robert, X. and Gouet, P. (2014) Deciphering key features in protein structures with the new ENDscript server. *Nucleic Acids Res.* **42**, W320-324.

97. Abramson, J., Adler, J., Dunger, J., Evans, R., Green, T., Pritzel, A., Ronneberger, O., Willmore, L., Ballard, A. J., Bambrick, J., Bodenstein, S. W., Evans, D. A., Hung, C. C., O′Neill, M., Reiman, D., Tunyasuvunakool, K., Wu, Z., Žemgulytė, A., Arvaniti, E., Beattie, C., Bertolli, O., Bridgland, A., Cherepanov, A., Congreve, M., Cowen-Rivers, A. I., Cowie, A., Figurnov, M., Fuchs, F. B., Gladman, H., Jain, R., Khan, Y. A., Low, C. M. R., Perlin, K., Potapenko, A., Savy, P., Singh, S., Stecula, A., Thillaisundaram, A., Tong, C., Yakneen, S., Zhong, E. D., Zielinski, M., Žídek, A., Bapst, V., Kohli, P., Jaderberg, M., Hassabis, D. and Jumper, J. M. (2024) Accurate structure prediction of biomolecular interactions with AlphaFold 3. *Nature* **630**, 493-500.

98. Ben Chorin, A., Masrati, G., Kessel, A., Narunsky, A., Sprinzak, J., Lahav, S., Ashkenazy, H. and Ben-Tal, N. (2020) ConSurf-DB: An accessible repository for the evolutionary conservation patterns of the majority of PDB proteins. *Protein Sci.* **29**, 258-267.

99. Fukui, T., Atomi, H., Kanai, T., Matsumi, R., Fujiwara, S., and Imanaka, T. (2005) Complete genome sequence of the hyperthermophilic archaeon *Thermococcus kodakaraensis* KOD1 and comparison with *Pyrococcus* genomes. *Genome Res* **15**: 352-363.

100. Constantinesco, F., Motorin, Y., and Grosjean, H. (1999) Transfer RNA modification enzymes from *Pyrococcus furiosus*: detection of the enzymatic activities in vitro. *Nucleic Acids Res.* **27**, 1308-1315.

101. Gupta, R. (1984) *Halobacterium volcanii* tRNAs. Identification of 41 tRNAs covering all amino acids, and the sequences of 33 class I tRNAs. *J. Biol. Chem.* **259**, 9461-9471.

102. Wolff, P., Villette, C., Zumsteg, J., Heintz, D., Antoine, L., Chane-Woon-Ming, B., Droogmans, L., Grosjean, H. and Westhof, E. (2020) Comparative patterns of modified nucleotides in individual tRNA species from a mesophilic and two thermophilic archaea. *RNA* **26**, 1957–1975.

103.   Kilpatrick, M.W. and Walker, R. T. (1981) The nucleotide sequence of the tRNAMMet from the archaebacterium *Thermoplasma acidophilum*. *Nucleic Acids Res*. **9**, 4387-4390.

104.   Walker, R. T. (1983) Mycoplasma evolution: a review of the use of ribosomal and transfer RNA nucleotide sequences in the determination of phylogenetic relationships. *Yale J. Biol. Med.* **56**, 367-372.

105.   Tomikawa, C., Ohira, T., Inoue, Y., Kawamura, T., Yamagishi, A., Suzuki, T., and Hori, H. (2013) Distinct tRNA modifications in the thermo-acidophilic archaeon, *Thermoplasma acidophilum*. *FEBS Lett.* **587**, 3575-3580.

106.   Cavaillé, J., Chetouani, F., and Bachellerie, J. P. (1999) The yeast Saccharomyces cerevisiae YDL112w ORF encodes the putative 2′-O-ribose methyltransferase catalyzing the formation of Gm18 in tRNAs. *RNA* **5**, 66-81.

107.   Ochi, A., Makabe, K., Kuwajima, K. and Hori, H. (2010) Flexible recognition of the tRNA G18 methylation target site by TrmH methyltransferase through first binding and induced fit processes. *J. Biol. Chem.* **285**, 9018-9029.

108.   Ochi, A., Makabe, K., Yamagami, R., Hirata, A., Sakaguchi, R., Hou, Y. M., Watanabe, K., Nureki, O., Kuwajima, K. and Hori, H. (2013) The catalytic domain of topological knot tRNA methyltransferase (TrmH) discriminates between substrate tRNA and nonsubstrate tRNA *via* an induced-fit process. *J. Biol. Chem.* **288**, 25562-25574.

109.   Watanabe, K., Nureki, O., Fukai, S., Endo, Y. and Hori, H. (2006) Functional categorization of the conserved basic amino acid residues in TrmH (tRNA(Gm18)methyltransferase) enzymes *J. Biol. Chem.* **281**, 34630-34639.

110.   Kohno, Y., Ito, A., Okamoto, A., Yamagami, R., Hirata, A. and Hori, H. (2023) Escherichia coli tRNA (Gm18) methyltransferase (TrmH) requires the correct localization of its methylation site (G18) in the D-loop for efficient methylation. *J. Biochem.* **175**, 43-56.

111.   Ito, T., Masuda, I., Yoshida, K., Goto-Ito, S., Sekine, S., Suh, S. W., Hou, Y. M. and Yokoyama, S. (2015) Structural basis for methyl-donor-dependent and sequence-specific binding to tRNA substrates by knotted methyltransferase TrmD. *Proc. Natl. Acad. Sci. USA* **112**, E4197-E4205.

112.   Christian, T., Sakaguchi, R., Perlinska, A. P., Lahoud, G., Ito, T., Taylor, E. A., Yokoyama, S., Sulkowska, J. I. and Hou, Y. M. (2016) Methyl transfer by substrate signaling from a knotted protein fold. *Nat. Struct. Mol. Biol.* **23**, 941-948.

113.   Jaroensuk, J., Wong, Y. H., Zhong, W., Liew, C. W., Maenpuen, S., Sahili, A. E., Atichartpongkul, S., Chionh, Y. H., Nah, Q., Thongdee, N., McBee, M. E., Prestwich, E. G., DeMott, M. S., Chaiyen, P., Mongkolsuk, S., Dedon, P. C., Lescar, J. and Fuangthong, M. (2019) Crystal structure and catalytic mechanism of the essential m^{1}G37 tRNA methyltransferase TrmD from *Pseudomonas aeruginosa*. *RNA* **25**, 1481-1496.

114.   Kawamura, T., Anraku, R., Hasegawa, T., Tomikawa, C. and Hori, H. (2014) Transfer RNA methyltransferases from *Thermoplasma acidophilum*, a thermoacidophilic archaeon. *Int. J. Mol. Sci.* **16**, 91-113.

115.   Zhou, M., Long, T., Fang, Z. P., Zhou, X. L., Liu, R. J. and Wang, E. D. (2015) Identification of determinants for tRNA substrate recognition by *Escherichia coli* C/U34 2′-*O*-methyltransferase. *RNA Biol.* **12**, 900-911.

 FIGURE LEGENDS

 Figure 1. Nucleotide sequence of *Thermococcus kodakarensis* tRNA$^{\text{Trp}}$ and scheme of

 comparative genomics. **A**, The secondary structure of *T. kodakarensis* tRNA$^{\text{Trp}}$ is depicted as

 a cloverleaf structure. The Cm6 is highlighted in red. **B**, The responsible gene for the Cm6

 modification in tRNA exists in *T. kodakarensis* and *P. furiosus* genomes was predicted by

 comparative genomics. *H. volcanii*, *S. acidocaldarius* and *T. acidophilum* do not possess the

 Cm6 modification in tRNA. The 570 genes were common in *T. kodakarensis* and *P. furiosus*.

 C, Scheme of comparative genomics is illustrated. An explanation of details is described in

 the main text.

 Figure 2. TK1257 gene product possesses a tRNA methyltransferase activity for

 2′-*O*-methylation at position 6. A, Purified TK1257 gene product (3 µg) was analyzed by 15%

 SDS-PAGE. The gel was stained with Coomassie Brilliant Blue. **B**, The methyl-transfer

 activity of TK1257 gene product was tested at 75ºC using tRNA$^{\text{Trp}}$ transcript and ^{3}H-labeled

 SAM. This experiment was independently replicated three times (n = 3). The error bars

 show the standard deviations. **C**, Three mutant tRNA$^{\text{Trp}}$ transcripts, in which the C6-G67 base

 pair was replaced by A6-U67, U6-A67 or G6-C67 base pair, were prepared. The mutation site

 is highlighted in red. **D**, The methylation speeds for the wild-type and mutant tRNA$^{\text{Trp}}$

 transcripts were compared. The methylation for the wild-type tRNA$^{\text{Trp}}$ is expressed as 1.00.

 This experiment was independently replicated three times (n = 3). The error bars show the

 standard deviations. **E**, ^{14}C-methylated nucleotides were analyzed by two-dimensional

 thin-layer chromatography. When the wild-type tRNA$^{\text{Trp}}$ transcript (WT) was used, ^{14}C-pCm

 was detected. In contrast, when the mutant tRNA$^{\text{Trp}}$ A6-U67 and U6-A67 transcripts were

 used, ^{14}C-pAm and ^{14}C-pUm were detected, respectively. Furthermore, when the mutant

tRNA^{Trp} G6-C67 transcript was used, ^{14}C-labeled nucleotide was not detected, consistent with

the result in panel D.   **F**, Kinetic parameters of TK1257 gene product for the wild-type and

mutant tRNA^{Trp} A6-U67 and U6-A67 transcripts were determined.   These transcripts had

comparable methyl group acceptance activities.   **G**, Kinetic parameters of TK1257 gene

product for SAM were determined.

Figure 3. TrmTS is expressed in *T. kodakarensis* wild-type strain and the *trmTS* gene

deletion (Δ*trmTS*) strain showed a slight growth retardation at 93°C.   **A,** The cell extracts

from wild-type and *ΔtrmTS* strains were analyzed by western-blotting.   Purified TrmTS (right

lane) was used as a positive control.   The gel was stained with Coomassie Brilliant Blue.

Because only 20 ng of TrmTS was used, the band of TrmTS was not visible by Coomassie

Brilliant Blue staining.   **B**, Native tRNA^{Trp} were purified from the wild-type and *ΔtrmTS*

strains.   The gel was stained with methylene blue.   **C**, Contents of methylated cytidines (m^{5}C

and Cm) in purified tRNA^{Trp} from the wild-type (upper) and *ΔtrmTS* (lower) strains were

analyzed by LC/MS.   The peak area of the m^{5}C in tRNA^{Trp} from the wild-type strain is

expressed as 100%.   **D**, Growth curves of the wild-type (black) and *ΔtrmTS* (red) strains are

compared at 85°C and 93°C.   At 93°C, a slight growth retardation of *ΔtrmTS* strain was

observed.   When TrmTS was expressed in the *ΔtrmTS* strain (blue), the growth speed

recovered.   This experiment was independently replicated three times (n = 3).   The error

bars show the standard deviations.

Figure 4. **Amnio acid sequence alignment of TrmTS-like proteins and SPOUT tRNA**

2′-*O*-methyltranserases and location of conserved amino acid residues.   **A**, The amino acid

sequence alignment of TrmTS-like proteins is depicted.   N-terminal ferredoxin-like (NFLD),

THUMP and SPOUT domains, and linker region are highlighted in gray, pale blue and purple,

and green, respectively.   Fifteen conserved amino acid residues were substituted by alanine

and/or other amino acid residues.   B, Subunit structural model of TrmTS was constructed by

AlphaFold 3.   A SAM molecule was modeled on the structure where the crystal structure of *T.*

thermophilus TrmH in complexed with SAM (PDB: 1V2X) was aligned with the TrmTS

structure.   The TrmH structure was then removed.   SAM is colored red.   The domains and

linker region are colored in the same as in panel A.   **C**, The mutation sites are mapped onto the

SPOUT catalytic domain and linker region.

Figure 5. **Purities of TrmTS mutant proteins and their activities.**   **A**, The wild-type and

mutant TrmTS proteins (2 μg each) were analyzed by 15% SDS-PAGE.   The gel was stained

with Coomassie Brilliant Blue.   **B**, Relative methylation speed of the wild-type and mutant

TrmTS proteins were compared using tRNA^{Trp} transcript and ^{3}H-SAM as substrates.   The

methylation speed of the wild-type TrmTS is expressed as 1.00.   This experiment was

independently replicated three times (n = 3).   The error bars show the standard deviations.

Figure 6.   CD spectra and kinetic parameters of TrmTS mutant proteins.   **A**, CD spectra

were measured at 25, 50 and 75ºC.   **B**, Kinetic parameters of mutant TrmTS proteins for SAM

and tRNA^{Trp} transcripts were measured.   **C**, Kinetic parameters of the wild-type and mutant

proteins are summarized.   "n. d." means that methyl-transfer activity was not detectable.

Figure 7.   Two conserved amino acid residues (Lys155 and Lys160) in the linker region.

A, The linker from one monomer contacts the SPOUT domain from the other monomer.

Asp270 residue of the catalytic domain in one subunit is able to contact with Lys160 residue of

the linker region in another subunit.   **B** and **C**, Kinetic parameters of TrmTS Lys155Ala mutant

protein for (B) SAM and (C) tRNA^{Trp} transcript were determined.

Figure 8.   Recognition sites in tRNA.   A, Eight mutant tRNA^{Trp} transcripts were prepared.

Transcript 1 is the wild-type tRNA^{Trp} transcript.   **B**, The features of mutant tRNA^{Trp} transcripts

are summarized.   **C**, The mutant tRNA^{Trp} transcripts were treated with TrmTS and 14C-SAM

and then analyzed by 10% PAGE (7 M urea).   The gel was stained with methylene blue (left).

The autoradiogram of the same gel was obtained (right).   The lane numbers correspond to the

transcript numbers in panel A and B.   **D**, Two mutant tRNA^{Trp} transcripts (-ACCA; red and

G6-C67; blue) which are not methylated by TrmTS, were used for the inhibition experiments.

Each mutant tRNA^{Trp} transcript (0 µM, 10 µM, 20 µM, 30 µM, 40 µM, 50 µM, 60 µM, 70 µM,

and 80 µM) was titrated to the TrmTS reaction in the presence of 35 µM wild-type tRNA^{Trp}. The

reaction was performed at 75 ºC for 10 min. The experiments were replicated three times (n=3).

The error bars show the standard deviations.

Figure 9. Hypothetical binding model between TrmTS and tRNA.   A, Positive (blue) and

negative (red) charged areas are mapped onto the surface of TrmTS dimer structure where +/– 5

kBT/e values were used for the visualization of the electrostatic potential map.   **B**, Conserved

and non-conserved amino acid regions are colored in magenta and dark green, respectively.   **C**,

A binding model between TrmTS and tRNA^{Trp} was predicted using AlphaFold3.   The THUMP

domain of TrmTS captures the CCA terminus of substrate tRNA and then the positively charged

area of the SPOUT domain interact with the three-dimensional core of tRNA.   Finally, the

modification site (ribose of C6) in tRNA is introduced into the catalytic pocket of SPOUT

domain and accesses the catalytic Arg193 residue.

Fig. 1

A

B

C

2306 genes in *T. kodakarensis*
↓
570 genes are common in
T. kodakarensis and *P. furiosus*
↓
10 genes annotated as
a tRNA methyltransferase gene
↓
1 gene product predicted as
a function-unknown
RNA methyltransferase
with THUMP and SPOUT domains

Fig. 2

Fig. 3

A
(kDa)

Marker | Wild-type cell extract | ΔtrmTS cell extract | TrmTS 20 ng | Wild-type cell extract | ΔtrmTS cell extract | TrmTS 20 ng

12.5% SDS-PAGE   Western blotting

B
Wild-type   ΔtrmTS
Total RNA | tRNA^Trp | tRNA mixture | tRNA^Trp

10% PAGE (7 M Urea)
Methylene Blue Staining

C
Wild-type tRNA^Trp   C + methyl m/z 258.11
m⁵C 100%   Cm 245%
Relative Abundance
0   Time (min)   40

ΔtrmTS tRNA^Trp   C + methyl m/z 258.11
m⁵C 100%   Cm 87 %
Relative Abundance
0   Time (min)   40

D
85°C
● : Wild-type
■ : ΔtrmTS
▲ : ΔtrmTS + trmTS
Optical density (660 nm)
Time [h]

93°C
● : Wild-type
■ : ΔtrmTS
▲ : ΔtrmTS + trmTS
Optical density (660 nm)
Time [h]

Fig. 4

Fig. 5

A

15％ SDS-PAGE

B

Fig. 6

A

Wild-type, K155A, R185A, G188A, R193A, S229A

S229C, S229T, Q234A, D270A, D270E, D270N

E301A, T322A, T322S, T325A, T325S, E326A

25°C
50°C
75°C

B

SAM: R185A, G188A, S229T, Q234A, E301A

tRNA^{Trp}: R185A, G188A, S229T, Q234A, E301A

C

Name	SAM			tRNA		
	Vmax [pmol·h^{-1}]	Km [μM]	Relative Vmax/Km	Vmax [pmol·h^{-1}]	Km [μM]	Relative Vmax/Km
WT	319	20	1	345	7.1	1
K155A	71	43	0.10	71	3.7	0.39
R185A	31	12	0.16	44	7.7	0.12
G188A	31	12	0.16	38	4.7	0.17
R193A	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.
S229A	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.
S229C	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.
S229T	48	19	0.16	63	11	0.12
Q234A	68	23	0.19	81	6.4	0.26
D270A	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.
D270E	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.
D270N	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.
E301A	62	20	0.19	83	14	0.12
T322A	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.
T322S	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.
T325A	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.
T325S	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.
E326A	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.

Fig. 7

A

B

SAM

Vmax: 71 pmol·h^{-1}
Km: 43 µM

(y-axis) Cm6 formation [pmol·h^{-1}]
(x-axis) SAM [µM]

C

tRNA^{Trp}

Vmax: 71 pmol·h^{-1}
Km: 3.7 µM

(y-axis) Cm6 formation [pmol·h^{-1}]
(x-axis) tRNA^{Trp} transcript [µM]

D270

K160'   K155'

Fig. 8

A

B

Number	Acceptor-stem	D-arm	Anticodon-arm	T-arm	Acceptor-stem	Feature
1	GGGGGCG UG	GUGUAGCCUGGUCCAUCAU	C GCGGGCUCCAGACCCGC	GGAC CGGGGUUCAAAUCCCCG	CGCCCCC A CCA	Full-length tRNA^Trp
2	GGGGGCG UG		C GCGGGCUCCAGACCCGC	GGAC CGGGGUUCAAAUCCCCG	CGCCCCC A CCA	Deletion of D-arm
3	GGGGGCG UG	GUGUAGCCUGGUCCAUCAU	C	GGAC CGGGGUUCAAAUCCCCG	CGCCCCC A CCA	Deletion of anticodon-arm
4	GGGGGCG UG	GUGUAGCCUGGUCCAUCAU	C GCGGGCUCCAGACCCGC	GGAC	CGCCCCC A CCA	Deletion of T-arm
5	GGGGGCG UG	GUGUAGCCUGGUCCAUCAU	C GCGGGCUCCAGACCCGC	GGAC CGGGGUUCAAAUCCCCG	CGCCCCC	Deletion of ACCA terminus
6	GGGGGCG UG	GUGUAGCCUGGUCCAUCAU	C	GGAC	CGCCCCC A CCA	Deletion of anticodon-arm & T-arm
7	GGGGGCG UG		C GCGGGCUCCAGACCCGC	GGAC	CGCCCCC A CCA	Deletion of D-arm & T-arm
8	GGGGGCG UG		C	GGAC CGGGGUUCAAAUCCCCG	CGCCCCC A CCA	Deletion of D-arm & anticodon-arm
9	GGGGGCG UG		C	GGAC	CGCCCCC A CCA	Deletion of D-arm & anticodon-arm & T-arm

C

1 2 3 4 5   1 2 3 4 5

10% PAGE (7 M Urea)   Autoradiogram

1 6 7 8 9   1 6 7 8 9

10% PAGE (7 M Urea)   Autoradiogram

D

Relative ³H-methyl group incorporation rate

Mutant tRNA^Trp transcript [μM]

tRNA^Trp (-ACCA)

tRNA^Trp C6G-G67C

Fig. 9