Potential title: A national assessment of waterbird hunting in coastal wetlands of Suriname,

South America

Author list: David S. Mizrahi[1*], Arie L. Spaans[2], Marianne J. Spaans-Scheen[2], Amelia R.

Cox[3,4], Benoit Laliberte[3], Eduardo Gallo-Cajiao[5, 6], Christian Roy[3]

[1] New Jersey Audubon
[2]Foundation Friends of Nature Conservation Suriname
[3]Canadian Wildlife Service, Environment and Climate Change Canada, Gatineau, Quebec,
Canada
[4] Current address: Northwest Territories Cumulative Impact Monitoring Program,
Government of the Northwest Territories, Yellowknife, Northwest Territories, Canada
[5]School of Marine and Environmental Affairs, University of Washington, Seattle,
Washington, USA
[6] Department of Human Dimensions of Natural Resources Management, Colorado State
University, Fort Collins, Colorado, USA
*Corresponding author: david.mizrahi@njaudubon.org

ORCID
David Mizrahi ID: https://orcid.org/0000-0003-4338-2443
Amelia Cox ID: https://orcid.org/0000-0002-4804-5115
Eduardo Gallo-Cajiao ID https://orcid.org/0000-0002-8607-2675
Christian Roy ID: https://orcid.org/0000-0002-5599-6234

Funding statement: The surveys conducted in 2006 received financial support from NVD

Nature Conservation Fund, BirdLife Netherlands. Funding for the survey conducted in 2016

was provided to New Jersey Audubon by US Fish and Wildlife Service, through a grant from

National Fish and Wildlife Foundation. The project received additional funding from Prince

Bernhard Culture Fund, Dierenrampenfonds Foundation, Van der Hucht De Beukelaar

Foundation, and Environment and Climate Change Canada.

Author contributions: The conception of the project was led by DSM and ALS. The

questionnaire was developed by ALS, MJSS, and DSM. Data collection was led by ALS and

MJSS. ARC and CR analyzed the data and prepared the tables and figures. All the authors

where invovled in the interpretation of the data. The first draft of the paper was written by

EGC and ARC. All authors approved the final version of the manuscript.

Acknowledgements: We are deeply grateful to all the hunters who participated in this study.

We also extend our sincere thanks to Marie Djosetro, Bryan Drakenstein, and Hesdy Esajas

for their valuable contributions at the outset of the fieldwork in 2006 and 2016, respectively.

Special thanks to Chanderbhan Dwarka, Amir Imraan, Astrid Moor, and Ashok Pherai for

their assistance in conducting the surveys, and to Brendan Gray for his support in compiling

the hunter database. F. Tremblay, two reviewers, and the editors also provided invaluable

comments that greatly helped to improve this manuscript.

Conflict of interest statement: The authors declare no conflicts of interest relevant to this

publication.

Ethic Statement: Informed consent was obtained from all participants before interviews

Abstract

The central northern coast of South America has extensive wetlands critical for waterbird conservation. While waterbird harvest occurs in the region, the impact on species population dynamics remains unclear. This study assesses waterbird hunting in the coastal wetlands of Suriname, addressing: (i) the extent of waterbird harvest, (ii) changes in harvest magnitude over time, and (iii) the motivations and methods used by hunters. We collected data via a national survey of licensed hunters in 2006 and 2016 using structured interviews. A Bayesian hierarchical model was used to analyze the data. We estimated harvest levels for 11 species and three groups (small herons, small shorebirds, large shorebirds). For most species, mean harvest per hunter significantly decreased from 2006 to 2016, except for blue-winged teal and migratory shorebirds. Most hunting was for non-commercial purposes (personal consumption and recreational). This is the first national assessment of waterbird hunting in Suriname. Harvest levels vary by species, and the sustainability of these levels remains uncertain. Managing hunting in Suriname requires addressing both legal and illegal hunting. Given Suriname's importance for waterbirds, particularly species like the scarlet ibis and migratory shorebirds, it should be a priority for conservation efforts.

Keywords: harvest management, scarlet ibis, shorebirds, Western Atlantic Flyway, western hemisphere, wildlife management

1. Introduction

Wetlands are biologically productive ecosystems that have sustained humans globally. The use of wetlands has included the harvesting of plants and animals, such as fish and waterbirds, with multiple purposes (e.g., cultural, subsistence, recreational, commercial). However, following increased pressure on the environment during the second phase of the Anthropocene (i.e., the Great Acceleration), the exploitation of some species may have contributed to population declines, and in some cases, extinction (Steffen et al. 2011; Benitez-Lopez et al. 2017; Cornford et al. 2023). Additionally, for long-distance migratory species, such as birds, harvest at specific locales can be compounded over large spatial scales and become unsustainable (e.g., Gallo-Cajiao et al. 2020). Consequently, it is necessary to evaluate harvest levels, their impacts on population dynamics, and the social and economic factors that drive them, in order to inform effective conservation and management policies for wetland-dependent species. Against this backdrop, there is a strong need to assess waterbird harvest worldwide, but particularly tropical regions, where people's livelihoods often rely on natural resources, governance capacity for wildlife management may be weak, and harvest monitoring is relatively limited (Amano et al. 2018; Brotherton et al. 2020).

The central northern coastal region of South America include extensive wetlands crucial for waterbird species conservation (Butler et al. 2001; Murray et al. 2019). These ecosystems comprise mangrove forests, brackish lagoons, freshwater swamps, and the most extensive mudflats of South America, a flat coastline, and is strongly influenced by the Guiana Current. Noteworthy, this region has a low human footprint with little intertidal wetland losses (Murray et al. 2022), so threats associated with land use change are generally not prevalent (Allan et al. 2023). These coastal wetlands serve as important non-breeding and stopover areas for migratory shorebirds (i.e., Aves: Scolopacidae, Charadriidae) along the Western Atlantic Flyway. Additionally, over 100 species of waterbirds, many of which are non-migratory, occur in these coastal wetlands, including ducks (Anatidae), flamingoes (Phoenicopteridae), and herons (Ardeidae). Along the central northern coast of South America, Suriname has the largest concentration of some waterbird species (Spaans 1978; Morrison & Ross 1989), including scarlet ibis (*Eudocimus ruber*), semipalmated sandpiper (*Calidris pusilla*), greater yellowlegs (*Tringa melanoleuca*), lesser yellowlegs (*Tringa flavipes*), and short-billed dowitcher (*Limnodromus griseus*). The importance of this region has been recognized through the designation of Important Bird Areas by BirdLife International, Ramsar wetlands of international importance, and sites of hemispheric

importance under the Western Hemisphere Shorebird Reserve Network (Ramsar Convention

Secretariat 1997; Western Hemisphere Shorebird Reserve Network 2024; BirdLife

International 2026).

Waterbird population trajectories in this region are generally unknown with some exceptions.

For instance, the central northern coast of South America is not surveyed as part of the

International Waterbird Census, hindering long-term monitoring of populations (Wetlands

International 2026). This data gap has also impeded assessments of protected area

effectiveness, which is essential for management (Wauchope et al. 2022). However, some

species that occur in this region are declining across their entire range, such as the scarlet ibis,

wood stork (*Mycteria americana*), and white-cheeked pintail (*Anas bahamensis*; BirdLife

International 2025). Similarly, several migratory shorebirds species (e.g., semipalmated

sandpiper, lesser yellowlegs, and short-billed dowitcher) have shown a significant decline of

over three decades along the central northern coast of South America (Morrison et al. 2012),

consistent with overall declines observed in North America (Smith et al. 2023).

Harvest, whether for commercial, recreational or subsistence purposes, can significantly

contribute to mortality rates and influence population dynamics of some waterbird species.

Hence, harvest assessments can play a crucial role in the development of sustainable harvest

management frameworks and to inform conservation objectives (Watts et al. 2015; Koneff et

al. 2017; Cox et al. 2024). Particularly, harvest assessments are important for shorebirds, as

they typically have slow life history traits, such as small clutch sizes and only one breeding

attempt per season, making them highly vulnerable to high harvest levels or even to

seemingly low harvest levels when populations are declining (Watts et al. 2015). For

migratory bird species, harvest assessments must also occur in multiple countries to account

for the cumulative effects of harvest to be estimated. Recent studies from Guyana (Andres et

al. 2022), French Guiana (Taylor 2017), and Brazil (Bosi de Almeida et al. 2018) show

widespread shorebird hunting in the central northern coast of South America. Beyond this

specific region, migratory shorebirds have also been hunted along the Western Atlantic

Flyway in parts of the Lesser Antilles (Watts & Turrin 2016; Gebauer 2018; Reed et al.

2018). In the United States and Canada, the extent of hunting remains unclear, as part of

subsistence hunting (but see Naves et al. 2019) and illegal harvest during recreational hunting

(Katzner et al. 2020).

In addition to evaluating harvest from an ecological standpoint, understanding the human

dimensions of hunting is crucial for informing conservation policy. These dimensions include

parameters such as demographics of resource users (e.g., age, gender, income), hunting's

contribution to livelihoods, and attitudes toward hunting and regulations (Vaske & Manfredo

2012; Eriksson et al. 2023). For instance, hunting of migratory shorebirds in Southeast Asia

can make a considerable contribution to household income (Zöckler et al. 2010; Gallo-Cajiao

et al. 2020). In Guyana, shorebird hunters are typically fishermen supplementing their income

when fishing conditions are poor (Andres et al. 2022). Understanding these parameters in the

context of the central northern coast of South America could help inform more effective and

equitable harvest management policies.

Here, we present a case study on a national assessment of waterbirds harvested in coastal

wetlands of Suriname. Our aim is to contribute to the evaluation of harvest sustainability of

waterbird species in Suriname within the context of central northern South America and the

Western Atlantic Flyway. Specifically, we ask: (i) what is the recent extent of waterbird

harvest in coastal wetlands of Suriname?; (ii) has magnitude of waterbird harvest changed

over time?; and, (iii) what are the motivations of waterbird hunters and, what harvest methods

they use?

2. Methods

2.1 Study area

Suriname, located on the central northern coast of South America, covers 163,800 km² and

has a 386 km coastline. It borders the Atlantic Ocean, Guyana, French Guiana, and Brazil.

The country has a tropical climate with two dry and two wet seasons. Major rivers include the

Courantyne, Coppename, Saramacca, Suriname, Commewijne, and Marowijne. Most of

Suriname's population (87%) lives in the coastal area, where the economy is primarily driven

by mining and agriculture (Food and Agriculture Organization 2024). The coastal zone,

mostly low and flat, comprises the lower and upper coastal plains, which are key waterbird

habitats (Spaans et al. 2018). Wetland ecosystems in the lower coastal plain include mudflats,

beaches, estuaries, mangroves, salt pans, coastal marshes, swamp forests, and shallow

brackish lagoons. The upper coastal plain includes herbaceous and peat swamps. Our study

focused on all coastal districts except Marowijne, where hunters were difficult to reach

(Figure 1).

Figure 1: Map of Suriname with district borders, showing coastal districts numbered: 1)

Nickerie, 2) Coronie, 3) Saramacca, 4) Wanica, 5) Paramaribo, 6) Commewijne, and 7)

Marowijne. Note: the Marowijne district and non-coastal districts (Para [purple], Brokopondo

[yellow] and Sipaliwini [green]) were not surveyed during this study.

The coastal zone of Suriname has various designations under different international

institutional frameworks, including one wetland of international importance under the Ramsar

Convention, three hemispherically important sites under the Western Hemisphere Shorebird

Reserve Network, and four Important Bird Areas under BirdLife International (Ramsar

Convention Secretariat 1997; Western Hemisphere Shorebird Reserve Network 2024;

BirdLife International 2026). Domestically, there are four Multiple-Use Management Areas

(MUMAS) and four nature reserves as well (Spaans et al. 2018). Some of these areas have

overlapping designations.

The Hunting Act 1954 and the corresponding implementing legislation (i.e. Hunting Decree

2002 as amended in 2009), determine the species of wildlife that may be hunted and when.

All bird species occurring in Suriname are fully protected, except those designated as 'game',

'cage birds', and 'predominantly harmful species'. Waterbird species without full protection

under Suriname regulations and considered in this paper are: black-bellied whistling-duck

(*Dendrocygna autumnalis*), muscovy duck (*Cairina moschata*), white-cheeked pintail, blue-

winged teal (*Spatula discors*), Neotropic cormorant (*Nannopterum brasilianum*), and anhinga

(*Anhinga anhinga*; Table 1).

Table 1: Species and species groups of waterbirds included in the hunter survey conducted in

six districts of Suriname in 2006 and 2016. International Union for Conservation of Nature's

(IUCN) Red List of Threatened Species information was sourced from the version 2022-2

(IUCN 2022).

Common Name	Scientific Name	IUCN Listing	Migratory	Hunting status according to Suriname's law
blue-winged teal	*Spatula discors*	LC	Yes	Game species
white-cheeked pintail	*Anas bahamensis*	LC	No	Game species
black-bellied whistling-duck	*Dendrocygna autumnalis*	LC	No	Game species
muscovy duck	*Cairina moschata*	LC	No	Game species
American flamingo	*Phoenicopterus ruber*	LC	No	Protected
cocoi heron	*Ardea cocoi*	LC	No	Protected
great egret	*Ardea alba*	LC	No	Protected
wood stork	*Mycteria americana*	LC	No	Protected
jabiru	*Jabiru mycteria*	LC	No	Protected
scarlet ibis	*Eudocimus ruber*	LC	No	Protected
roseate spoonbill	*Platalea ajaja*	LC	No	Protected
small herons (e.g., little blue heron, snowy egret, cattle egret, tricolored heron, black-crowned / yellow-crowned night-herons, boat-billed heron)	*Egretta caerulea, Egretta thula, Ardea ibis, Egretta tricolor, Nycticorax nycticorax, Nyctanassa violacea, Cochlearius cochlearius*	—	—	Protected
small shorebirds (e.g., semipalmated and least sandpiper, semipalmated	*Calidris pusilla, C. minutilla, Charadrius semipalmatus*	—	Yes	Protected

plover)				
large shorebirds (e.g., greater and lesser yellowlegs, short-billed dowitcher, willet, whimbrel, black-bellied plover)	*Tringa melanoleuca, T. flavipes, Limnodromus griseus, Tringa semipalmata, Numenius phaeopus, Pluvialis squatarola*	—	Yes	Protected

2.1 Study design and data collection

This is a cross-sectional and longitudinal single case study based on surveys conducted at two points in time that are ten years apart. Travel is difficult, particularly on unpaved roads during the rainy season in Suriname, so data were collected in April 2006 and 2016, during what is typically the short dry season, to improve our chances of reaching hunters in their communities.

At the time the surveys were conducted, no ethics approval was required in Suriname. The surveys were conducted in collaboration with the Suriname Forest Service/Nature Conservation Division, a department within the Ministry of Land Policy and Forest Management. Staff from the agency were actively involved in survey implementation.

In 2016, potential participants were identified from hunter licensing records provided by the District Commissioners in each of the included districts. Neither the District Commissioners nor their staff were informed of the identities of the selected participants. No minors were involved in the study, as individuals under the age of 18 are not permitted to hold a hunting license in Suriname. The participant pool consisted of those who could be reached and agreed to participate. Due to challenges in contacting hunters, we could not create a stratified sample proportional to the number of hunters in each district.

Prior to administering the survey, all participants were informed of the study's purpose and how their responses would be used. They were advised that submission of a completed survey would be considered as informed consent to participate. While participants were encouraged to answer all questions, they were also clearly informed that responding to any specific question was entirely voluntary. All surveys were administered in Dutch, the official language of Suriname. Survey responses were recorded on paper and later digitized. All responses remained anonymous; no names or identifying information were included on the

questionnaires. Because we did not record personal identification information, we could not
intentionally interview any hunters from the 2006 survey in 2016.

The survey questions in 2006 focused primarily on hunting frequency, magnitude and
purpose, and species or group of species of waterbirds harvested (Appendix 1). For some taxa,
species reporting was not possible, or at least not accurately as hunters do not distinguish
between species, so for these we present data for groups of species which share broad
morphological and ecological traits (Table 1). The 2016 survey used all the same questions
used in 2006 but included additional questions to improve our understanding of the hunter
community (e.g., age, family members, hunting purpose), hunting seasonality, and hunting
method (Appendix 2).

Survey respondents self-reported their harvest, a common method for assessing waterbird
hunting (Geissler 1990; Smith et al. 2022). However, this approach has potential limitations,
including recruitment bias and underreporting of mortality, such as wounded birds which may
die later (i.e., crippling losses). In 2006, we piloted group surveys with hunters, but found that
participants were conferring with each other, making the results unreliable. As a result, we
switched to individual one-on-one interviews at hunters' homes, continuing this method in
2016. If interviewers suspected dishonesty, they cross-checked answers with game wardens
familiar with the hunters' activities. Data deemed unreliable were excluded from analyses. All
data were kept anonymous and confidential, and participation was voluntary.

223 2.2 Data analysis

224 *2.2.1 Estimation of licensed hunters in 2016*

In 2006, there was no reliable centralized hunter license database, so the total number of
licensed hunters that year is unknown. For 2016, we used the Suriname hunter license
database, which tracks only license registrations. According to Suriname law, hunting licenses
are valid for up to five years, from January 1st to December 30th. We considered the total
number of licensed hunters in 2016 as the unique registrations between January 1, 2011, and
December 30, 2016. This method does not account for licenses that may have been suspended
before 2016, as the database lacked records on renewals or suspensions, and no corrective
measures were applied.

2.2.2 Estimation of hunting levels

We used a Bayesian hierarchical model to analyze the harvest survey data (Appendix 3). We

modelled harvest data for 11 waterbirds species and four taxa (i.e., small herons, small

shorebirds, large shorebirds, and all shorebirds). Because of the limited number of

respondents in each region, we pooled all the data together and did not consider variation in

hunting across regions/districts. In 2006, survey respondents were asked to provide estimates

for all shorebird species combined. In 2016, respondents were asked to distinguish between

harvest of large and small shorebird species. Therefore, the survey of small and large

shorebirds was set to zero for 2006 and we modeled the harvest of all shorebirds directly. For

2016, we instead modelled the harvest of small and large shorebirds and derived the harvest

of all shorebirds from those two categories.

During the interview process, hunters reported the number of birds they harvested that year

via 8 categories (either 0; 1; 2 – 10; 11 – 50; 51 – 100; 101 – 500; 501 – 1000; or > 1000

birds). We therefore modeled the reported harvest of each species *i* by each individual hunter

interviewed *j* via a categorical distribution:

$$Harvest_{i,j} \sim Categorical(\pi_{1:8_{j,t}}) \tag{1}$$

where $\pi_{1:8}$ represents the vector of probabilities that the hunter will report their harvest in that

category for a given species *j* in year *t*.

We assumed that the underlying distribution of harvest of each species would follow a

negative binomial distribution, which allows for over dispersion that we observed in harvest

data. We parametrized the negative binomial distribution via the mean and variance (Lindén

& Mäntyniemi 2011) and modelled the mean harvest $(\mu_{j,t})$ for a given species *j* in year *t* such

as:

$$log(\mu_{j,t}) = \eta_t + \varepsilon_{j,t} \tag{2}$$
$$\varepsilon_{j,t} \sim Normal(0, \sigma_t) \tag{3}$$

where η_t is the average harvest across all species on the log scale and a $\varepsilon_{t,j}$ is species specific

derivation from this mean in this given year. The species specific derivations are drawn from

a normal distribution with year specific standard distribution. This hierarchical formulation

allowed us to share information across species (i.e. 'borrowing the strength' from the

ensemble) but to also consider the change in harvest level across years. We then parametrized

the variance ($\sigma^2_{j,t}$) of the model in function of the mean such as:

$$\sigma^2_{j,t} = \mu_{j,t} + \theta\mu_{j,t}^2 \tag{4}$$

where θ is the overdispersion parameter for the quadratic mean–variance relationship of the

classic negative binomial regression (Lindén & Mäntyniemi 2011). We used moment

matching to link estimates of the mean and variance to the species-specific and year specific

probability $\left(p_{j,t} = \frac{\mu_{j,t}}{\sigma^2_{j,t}}\right)$ and size parameter $\left(r_{j,t} = \frac{\mu_{j,t}^2}{(\sigma^2_{j,t} - \mu_{j,t})}\right)$ and then linked the

probability of the harvest categories with the cumulative probability distribution (CDF;

$F(k, p_{j,t}, r_j)$) of the negative binomial distribution. The first category of harvest represents the

probability that harvest for a given species each year is exactly zero and can be derived

directly from the CDF:

$$\pi_{1_{j,t}} = F(0, p_{j,t}, r_j) \tag{5}$$

269   The probability of the harvest for categories 2 to 7 can be derived by subtracting the CDF at

270   the lower bound of the category from the CDF at the upper bound of category. For example,

271   for the probability that harvest is between 2 to 10 birds (i.e., Category 3) for a given species in

272   a given year can be estimated via:

$$\pi_{3_{j,t}} = F(10, p_{j,t}, r_j) - F(2, p_{j,t}, r_j) \tag{6}$$

273   The last harvest category ($\pi_8; > 1000\ birds$) is the probability that the harvest is larger than

274   the lower bound of the category:

$$\pi_{8_{j,t}} = 1 - F(1000, p_{j,t}, r_j) \tag{7}$$

Hunters did not always complete the survey and did not report their harvest for all the species.

In those cases, we used a data augmentation scheme and let the model assign a harvest

category for the hunter. Because of the hierarchical formulation of the model, the category of

harvest assigned to this hunter by the model will tend to be close to the most frequent

category of harvest observed for that species during that year.

Finally, we derived the total annual harvest in 2016 by multiplying the number of registered

hunters with the mean annual harvest ($\mu_{j,2016}$) for a given species j. Since the Suriname

hunter license database was incomplete for 2006, we were not able to estimate the number of

registered hunters and total annual harvest in 2006.

We used R version 4.2.0 for data processing (R Core Team 2020) and JAGS version 4.3.0

(Plummer 2003), called via the R package jagsUI (Kellner 2021) for analyses. We used non-

informative priors for all parameters and ran three chains with randomized initial start values

for 69,500 iterations, using the first 2,000 iterations as burn in and thinning the chains to save

every 12th iteration, with 1,000 iterations of adaption yielding 45,000 posterior samples for

each parameter. We assessed model convergence visually and by making sure that the

Gelman-Rubin statistic ($\hat{R}$) was less or equal than 1.1 (Gelman et al. 2013). We also used a

posterior predictive check to evaluate the performance of the model. All results are reported

as the median with 90% Bayesian Credible Intervals, calculated as high-density intervals

using *bayestestR* package (Makowski et al. 2019)

294 3. Results

295 3.1 Hunters in Suriname

Surveys were conducted in six districts in 2006 and five districts in 2016, with the highest

number of participants from Nickerie, Saramacca, and Commewijne (Table 2). Additionally,

we were unable to recruit the same number of hunters from each district over the two time

periods. Paramaribo District, despite having the largest population of licensed hunters, had the

fewest participants (Table 2).

Table 2: Numbers of hunters interviewed about their harvest of waterbirds per district in
Suriname in 2006 and 2016.

District	2006	2016
Nickerie	115	186
Coronie	8	21
Saramacca	100	29
Wanica	3	11
Paramaribo	3	0
Commewijne	47	42
Total	**276**	**289**

In 2006, 276 licensed hunters answered the survey (Table 2) but we excluded 45 of them from

the analysis as the answers were deemed to be unreliable. Of the remaining 231 surveys, only

69% (n=159) of the hunters reported their harvest of all species included in the survey.

Hunters completed the survey and reported their harvest most frequently for muscovy duck

(91%), black-bellied whistling-duck (91%), white-cheeked pintail (90%), while response rates

were lower for jabiru (*Jabiru mycteria*; 82%) and American flamingo (*Phoenicopterus ruber*;

81%).

In 2016, a total of 289 male hunters answered the survey (Table 2) out of the 4727 hunters

that were registered in the Suriname hunter license database. The answers to 16 surveys were

deemed to be unreliable so we kept 273 surveys for the analysis. The ages of the respondents

ranged from 19 to 85 years, with a mean age of 49 (SD = 11). Of the 273 surveys 99%

(n=270) of the hunters reported their harvest for all species included in the survey. Of the

surveyed hunters, 190 hunters indicated that they did not harvest any birds, 72 hunters

reported harvesting 5 species or less, 8 hunters harvested between 6 and 10 species, and 3

hunters harvested more than 10 species.

In addition to the 11 species and four taxa that we modelled, hunters also reported the harvest

of Neotropic cormorant, anhinga, brown pelican (*Pelecanus occidentalis*), black skimmer

(*Rynchops niger*), and undefined Laridae taxa (i.e., *gulls and terns*). However, because the

harvest levels for these species were too low for statistical modeling, we excluded them from

the analysis.

3.2   Estimated waterbirds harvest in Suriname

3.2.1   *Average harvest for 2006 and 2016*

In both 2006 and 2016, mean harvest per hunter of all shorebirds was higher than for any

other species or group (Figure 2). Results from the 2016 survey indicate that hunters

harvested on average more small shorebirds than large shorebirds. After shorebirds, the most

harvested species in 2006 were black-bellied whistling-duck, scarlet ibis, and cocoi heron; the

jabiru and American flamingo were the least harvested species (Figure 2). In 2016, the most

harvested species outside of shorebirds was blue-winged teal followed by black-bellied

whistling-duck, while American flamingo and the jabiru were still the two least harvested

species (Figure 2).

Figure 2: Mean birds harvested in a year (μ) per hunter based on surveys of hunters in
Suriname in 2006 and 2016. Data points represent the median, with error bars indicating the
90% high-density intervals. American Ornithological Society species codes are presented in
Table 3. Note that in 2006, hunters reported all shorebirds species combined, but in 2016
large and small shorebirds were distinguished.

3.2.2. Has hunting of waterbird species changed over time?

For most waterbird species, mean harvest per hunter declined between 2006 and 2016 (Figure

2). The exceptions were the blue-winged teal and the total migratory shorebirds (i.e., small

and large combined), for which harvest did not decrease significantly between 2006 and 2016.

In particular, wood stork, roseate spoonbill (*Platalea ajaja*), and jabiru harvest declined below

one bird per hunter on average in 2016. Hunters surveyed in 2016 generally agreed that their

overall harvest had declined. Among respondents who answered this question and hunted in

both 2006 and 2016, most reported hunting less in 2016 (77%), while a small minority (7%)

reported hunting more than in 2006. When asked why they had changed their hunting

practices, out of 66 responses provided the most common response was that there were fewer

birds available for harvest than there were in the past (29%). Some individuals also cited the

increase in hunting regulations and increased enforcement (12%) and associated costs (6%).

3.2.3. Total derived harvest 2016

Shorebirds were the most harvested group in general in 2016 with a total harvest of

approximately 174 thousand birds (90% BCI = 94 − 276 thousand; Table 3). Small shorebirds

(107 [90% BCI = 37 − 202] thousand) were harvested nearly twice as much as large

shorebirds (59 [90% BCI = 23 − 109] thousand). Ducks were the next group of species with

the most harvest, with the blue-winged teal having the highest total harvest (47 [90% BCI =

25 − 76] thousand), followed by herons, egrets and ibis, with the scarlet ibis (19 [90% BCI = 8

− 36] thousand; Table 3) having the highest total harvest. Finally, the wood stork, roseate

spoonbill, American flamingo, and jaribu were less frequently harvested species with only the

wood stork (1230 [90% BCI = 413 − 2408]; Table 3) harvest exceeding a thousand birds.

Table 3: Estimated total waterbird harvest by all registered hunters in Suriname in 2016.
Estimates are presented as medians with 90% Bayesian Credible Intervals (BCI).

Family/Species	Species code	2016 Annual harvest (90% BCI)
Anatidae		
black-bellied whistling-duck (*Dendrocygna autumnalis*)	BBWD	31824 (16955 – 50389)
blue-winged teal (*Spatula discors*)	BWTE	46974 (24505 – 76414)
muscovy duck (*Cairina moschata*)	MUDU	6075 (3191 – 9713)
white-cheeked pintail (*Anas bahamensis*)	WCHP	18520 (8987 – 30855)
Phoenicopteridae		
American flamingo (*Phoenicopterus ruber*)	AMFL	512 (81 – 1505)
Ardeidae		
cocoi heron (*Ardea cocoi*)	COHE	5129 (2115 – 9566)
great egret	GREG	7631 (2470 – 16363)

(*Ardea alba*)		
Small herons	SMHE	13960 (3769 – 32142)
Ciconiidae		
jabiru	JABI	
(*Jabiru mycteria*)		92 (10 – 241)
wood stork	WOST	
(*Mycteria americana*)		1230 (413 – 2408)
Threskiornithidae		
roseate spoonbill	ROSP	
(*Platalea ajaja*)		865 (186 – 2262)
scarlet ibis	SCIB	
(*Eudocimus ruber*)		19124 (7667 – 36053)
Charadriidae,		
Recurvirostridae,		
Scolopacidae		
Large shorebirds	Large	58915 (23218 – 109019)
Small shorebirds	Small	107042 (36968 – 202281)
Total shorebirds	Total	173700 (93970 – 276450)

3.2.4. What are the motivations of waterbird hunters and what hunting methods do they use?

Across both survey years (2006 and 2016), 63% of respondents travelled to their hunting

location with the intention of engaging in another activity, generally fishing or recreation.

Based on data from 2016 only, nearly all hunters who reported their hunting method (N=90)

reported hunting primarily with a firearm (99%), while a small proportion reported using a net

(1 %). In 2016, hunters who reported their primary purpose for hunting (N=90) indicated that

it was predominantly non-commercial in purpose, which includes personal consumption and

recreational activities. Though there is overlap between these two motivations, the majority of

hunters reported hunting for the former (70%) and only a small proportion for the latter

(13%). Additionally, 13% of the hunters reported selling the birds inferring a commercial

purpose.

4. Discussion

Harvest of waterbird species is pervasive across coastal wetlands of Suriname, has declined

over time, and is primarily practiced for personal consumption. To the best of our knowledge,

this is the first national assessment of waterbird harvest from Suriname, part of a region of

critical importance to shorebird conservation in central northern South America and along the

Western Atlantic Flyway. While the importance of waterbird harvest in Suriname has been

internationally recognized for many years (Ottema & Spaans 2008; AFSI Harvest Working

Group 2017, 2020), no structured assessment has been previously conducted. Waterbird

harvest occurs along the entire coast of Suriname and includes species across a wide range of

taxonomic groups. Amongst all waterbirds hunted in Suriname, the scarlet ibis and migratory

shorebirds are of particular conservation concern due to higher hunting levels, declining

populations, and slow life history traits (Olmos & Silva e Silva 2001; Watts et al. 2015;

BirdLife International 2016; Smith et al. 2023).

In Suriname, waterbirds are primarily harvested using firearms, with only a small proportion

using alternative methods, such as nets and choking wires. This pattern is similar to French

Guiana (Taylor 2017) but contrasts with neighboring Guyana and Brazil, where choking

wires, fishing nets and night lighting are the most common methods (Bosi de Almeida et al.

2018; Andres et al. 2022). In line with broader understanding of wetland use by humans,

waterbird harvest did not happen in isolation in our study but rather alongside other resource

uses, particularly fishing.

Based on hunters' self-reported motivations, waterbird harvest in Suriname was primarily

non-commercial and driven by recreational interest and personal consumption; the latter was

the most frequently cited motivation in our study. This pattern contrasts with Guyana, where

commercial harvest of migratory shorebirds occurs at a relatively larger scale although the

overall harvest is less than in Suriname (Andres et al. 2022). While the limited

commercialization reported in Suriname is encouraging, cross-border transport of waterbirds

from western Suriname to markets in Guyana has been documented and warrants further

investigation. Beyond subsistence, some of the harvest done for personal consumption may

have cultural dimensions, as suggested by the consumption of birds such as the scarlet ibis at

social celebrations (Ottema & Spaans 2008), a practice noted elsewhere in the region as being

driven by the bird's status as a traditional delicacy (BirdsCaribbean 2017).

Species hunted represented a wide range of taxonomic groups and harvest levels varied

between species, a pattern consistent with tropical wetlands elsewhere (Iskandar & Karlina

2004; Ramachandran et al. 2017; Bosi de Almeida et al. 2018; Deniau et al. 2022). Ducks

were among the most heavily harvested groups,likely reflecting both their high availability

and hunter preferences for larger-bodied species. Of all the duck species hunted, only the

blue-winged teal is considered migratory. Harvest levels of blue-winged teal from Suriname

were modest compared to North America, representing about 1.68% of the combined harvest

estimated for the US and Canada during the same year (Raftovich et al. 2018). The scarlet ibis

was also among the species with the highest levels of hunting, and although it is hunted

elsewhere in the tropics, the levels reported here are unusually high compared to recent

reports from other regions (Deniau et al. 2022). Harvest of other species of waterbirds such as

the American flamingo, the wood stork, the jabiru, and the roseate spoonbill, were relatively

low and in line with level reported for other areas such as the Sahel (Deniau et al. 2022).

Current estimates of migratory shorebird harvest in Suriname exceed those available for any

other country along the central northern coast of South America and the Lesser Antilles. We

estimated that ≈ 174 thousand migratory shorebirds were harvested in 2016, but this is a

conservative estimate as it does not account for crippling loss, under reporting because of

illegality or unlicensed hunters. While empirical data on crippling rates for shorebirds, and

particularly in South America, are virtually non-existent, estimates from other migratory game

birds provide valuable context. For species such as doves and waterfowl, crippling losses

typically range from 10% to over 25% and may be influenced by various hunting factors,

including the type of ammunition used (i.e., steel shot vs lead shot; Schulz et al. 2013; Ellis &

Miller 2022; Moreno-Zárate et al. 2023). Nevertheless, our estimate for shorebird harvest in

Suriname greatly exceeds combined estimates reported for Guyana, the Lesser Antilles, and

Brazil (Bosi de Almeida et al. 2018; Andres et al. 2022; AFSI Harvest Working Group 2020).

Our findings are limited by the qualitative, self-reported nature of the survey data, including

recall bias, rounding, or intentional misreporting. Broad harvest categories and assumptions

about the underlying distribution may affect precision but are unlikely to alter the overall

magnitude of our estimates. Misclassification near range category boundaries and limited

information on extreme values may affect estimation of the underlying distribution, and the

assumption of a negative binomial distribution may not fully capture its shape, potentially

biasing mean and total harvest estimates. This may affect precision but are unlikely to alter

the overall magnitude of our estimates. We also assumed surveyed hunters were

representative of all registered hunters. Future surveys could improve accuracy through

narrower harvest categories and a stratified sampling.

The sustainability of waterbird harvest in Suriname remains largely unknown, but inferences

can be drawn from ecological theory and species life-history traits. The population-level

effects of harvest range from compensatory to additive, or even super-additive mortality,

depending on hunting intensity and species life history (Anderson & Burnham 1976).

Additive mortality occurs when harvest reduces survival, while compensatory mortality arises

when harvest does not affect survival due to density dependence or targeting individuals that

would have died naturally (Sandercock et al. 2011; Cooch et al. 2014). Generally, species

with large clutch sizes, high natural mortality, and large populations experiencing density

dependence can generally sustain higher harvest rates than species with small clutch sizes,

high annual survival, and small populations (e.g., most shorebird species; Nichols et al. 1984;

Cooch et al. 2014). The timing of harvest relative to natural mortality is also important, with

hunting timed during or after periods of high natural mortality more likely to be additive

(Kokko 2001; Blomberg 2015). For many species, migration represents a period of elevated

natural mortality particularly for juveniles (Lok et al. 2015; Rushing et al. 2017; Sergio et al.

2019).

While duck hunting may pose relatively lower conservation concern, uncertainty around

population status of resident ducks in Suriname suggests a precautionary approach is

warranted. The scarlet ibis population is declining across its range (Sainz-Borgo 2025), and

given its slow life history traits (i.e., small clutch size, delayed sexual maturity; Olmos &

Silva e Silva 2001), current harvest levels may be unsustainable. A more reliable assessment

of the scarlet ibis population could encourage better adherence to regulations, as hunters may

use uncertainty as a justification to dismiss the need for the law (i.e., neutralization theory;

Eliason 1999). Compliance promotion campaigns could help, but poachers often comply only

in the presence of law enforcement (Jacoby 2014; von Essen et al. 2014). Any harvest

management strategies and compliance promotion efforts will need to consider broader socio-

economic contexts such as local wetland use, livelihoods, and governance capacity to be

effective (Ostrom 2009; Sutherland et al. 2014).

For migratory shorebirds, species-level harvest estimates are lacking, but concern is

particularly high for the semipalmated sandpiper, a Near Threatened species (BirdLife

International 2024) that likely accounts for a large percentage of the small shorebirds

harvested in Suriname given its prevalence in the region (Morrison & Ross 1989). Shorebird

harvest in Suriname remains high, and up to 44% of the flyway-wide sustainable hunting

levels could already be taken from neighboring Guyana (Watts et al. 2015; Andres et al.

2022).

Most species in our study are protected and the harvest of flamingoes, herons, storks, ibises,

and shorebirds in our surveys were illegal under Suriname's Hunting Act (1954). The

observed declines for the harvest of flamingoes, herons, storks, and ibises are encouraging.

These declines are also concurrent with an increase in compliance promotion and enforcement

of harvest regulations, which likely contributed to the reduction in take. Shorebirds harvest

level remains problematic. The status of shorebirds as protected species also complicates the

engagement of hunters over species identification and creates challenges for the assessment of

shorebird harvest sustainability (Solomon et al. 2015). This situation is not unique to

Suriname however and has been observed elsewhere such as in Guyana (Andres et al. 2022)

and the East Asian-Australasian Flyway (Gallo-Cajiao et al. 2020).

Furthermore, a patchwork approach to regulating hunting of migratory shorebirds can blur its

full effects until data are aggregated at a full life-cycle scale (Gallo-Cajiao et al. 2020).

Regulatory frameworks for shorebird harvest in particular are variable across the north coast

of South America (Watts & Turrin 2016). Shorebirds are totally protected from harvest in

Brazil, legally harvested for sport in French Guiana, and entirely unregulated in Guyana,

including commercial hunting. This regulatory heterogeneity creates opportunities for

laundering, whereby shorebirds illegally harvested in one jurisdiction may be sold in others

where regulations are weak or absent. Although international agreements such as the

Convention on International Trade in Endangered Species of Wild Fauna and Flora (CITES)

and the Convention on Migratory Species (CMS) are relevant, their effectiveness is limited in

this context. Suriname, for example, is a party to CITES but not CMS, while migratory

shorebird species are listed under CMS but not CITES.

Addressing these challenges will require a coordinated, multi-jurisdictional approach

grounded in improved, species-level harvest assessments and population monitoring, allowing

harvest levels to be explicitly linked to population trends (Cox et al. 2025). Equally important

is a better understanding of the human dimensions of shorebird harvest in the Western

Atlantic Flyway, including equity, livelihood dependence, and perceptions of regulations and

conservation measures, as policies lacking user acceptance are unlikely to succeed. Both lines

of research on harvest effects and human dimensions will need to be further integrated into

the policy process balancing conservation and human needs. More specifically, outputs from

this research agenda could inform institutional arrangements already in place that have

incorporated harvest mitigation as a priority, such as the Atlantic Flyway Shorebird Initiative

(AFSI Harvest Working Group 2017, 2020).

5.   Conclusion

510   Our study contributes to the understanding of wetland use more generally and to the

511   consumptive use of waterbirds more specifically, both in the tropics as well as in Suriname

512   within the context of central northern South America and the Western Atlantic Flyway. In

513   doing so, we discovered that harvest could be happening at unsustainable levels both for

resident and migratory species, supporting global analyses on the importance of

overharvesting as a threat to biodiversity (Maxwell et al. 2016). Illegally hunted species are

typically those that are of greatest conservation concern. The issue of waterbird harvest, and

shorebirds in particular, has been recognized for decades in the central northern coast of South

America and the Lesser Antilles (Hutt, 1991). However, structured and systematic

assessments have only recently become available, highlighting the challenges involved in

addressing this issue. More dedicated resources to monitor the status of populations and the

level of harvest will greatly help develop more effective regulatory frameworks for waterbird

conservation in the region. However, while enhancing existing regulatory frameworks is

important, it is crucial to also prioritize efforts in promoting compliance, engaging with local

communities, and developing alternative livelihoods. Given the importance of Suriname for

waterbirds in the Americas, developing a cost-effective approach to Suriname's harvest

management of species with international conservation concern, such as the scarlet ibis and

migratory shorebirds, should remain a priority for the international community.

6. References

AFSI Harvest Working Group. (2017). *Achieving a sustainable shorebird harvest in the Caribbean and northern South America, progress report, 2011–2017* [Unpublished report]. U.S. Fish & Wildlife Service, Migratory Bird Program. https://www.shorebirdplan.org/wp-content/uploads/2018/08/AFSI-Shorebird-Hunting-Progress-Report-Sep-2017.pdf

AFSI Harvest Working Group. (2020). *Actions for the Atlantic Flyway Shorebird Initiative's Shorebird Harvest Working Group 2020–2025*. U.S. Fish & Wildlife Service, Migratory Bird Program. https://www.shorebirdplan.org/wp-content/uploads/2020/09/AFSI-Shorebird-Harvest-Action-Plan-2020_25-April-2020.pdf

Allan, H., Levin, N., & Kark, S. (2023). Quantifying and mapping the human footprint across Earth's coastal areas. *Ocean & Coastal Management*, 236, 106476. https://doi.org/10.1016/j.ocecoaman.2022.106476

Amano, T., Székely, T., Sandel, B., Nagy, S., Mundkur, T., Langendoen, T., Blanco, D., Soykan, C. U., & Sutherland, W. J. (2018). Successful conservation of global waterbird populations depends on effective governance. *Nature*, 553, 199–202. https://doi.org/10.1038/nature25139

Anderson, D. R., & Burnham, K. P. (1976). *Population ecology of the mallard: VI. The effect of exploitation on survival*. Resource Publication 128, Federal Government Series, U.S. Fish and Wildlife Service, Washington, D.C., USA. http://pubs.er.usgs.gov/publication/rp128

Andres, B. A., Moore, L., Cox, A. R., Frei, B., & Roy, C. (2022). A preliminary assessment of shorebird harvest in coastal Guyana. *Wader Study*, 129, 39–47. https://doi.org/10.18194/ws.00262

Benitez-Lopez, A., Alkemade, R., Schipper, A., Ingram, D., Verweij, P., Eikelboom, J., & Huijbregts, M. (2017). The impact of hunting on tropical mammal and bird populations. *Science*, 356, 180–183. https://doi.org/10.1126/science.aaj1891

BirdsCaribbean. (2017). *Scarlet Ibis: A national symbol under siege*. https://www.birdscaribbean.org/2017/08/scarlet-ibis-a-national-symbol-under-siege/

BirdLife International. (2016). S*pecies factsheet: Scarlet Ibis Eudocimus ruber.* https://datazone.birdlife.org/species/factsheet/scarlet-ibis-eudocimus-ruber

BirdLife International. (2024). *Species factsheet: Semipalmated Sandpiper Calidris pusilla.* https://datazone.birdlife.org/species/factsheet/semipalmated-sandpiper-calidris-pusilla

BirdLife International. (2025). *IUCN Red List for birds*. https://datazone.birdlife.org

BirdLife International. (2026). *Country factsheet: Suriname*. Retrieved January 3, 2026, from https://datazone.birdlife.org/country/factsheet/suriname

Blomberg, E. J. (2015). The influence of harvest timing on greater sage-grouse survival: A cautionary perspective. *The Journal of Wildlife Management*, 79, 695–703. https://doi.org/10.1002/jwmg.877

Bosi de Almeida, J., Gallon de Aguiar, A., Damasceno, J. P., & Carvalho, M. R. (2018). *Assessment of shorebird hunting in northern Brazil* [Unpublished report]. Sociedade para a Conservação das Aves do Brasil.

Brotherton, S., Joyce, C. B., & Scharlemann, J. P. (2020). Global offtake of wild animals from wetlands: critical issues for fish and birds. *Hydrobiologia*, 847, 1631–1649. https://doi.org/10.1007/s10750-020-04222-2

Butler, R. W., Davidson, N. C., & Morrison, R. G. (2001). Global-scale shorebird distribution in relation to productivity of near-shore ocean waters. *Waterbirds*, 24, 224–232. https://doi.org/10.2307/1522030

Cooch, E. G., Guillemain, M., Boomer, G. S., Lebreton, J.-D., & Nichols, J. D. (2014). The effects of harvest on waterfowl populations. *Wildfowl*, Special Issue 4, 220–276.

Cornford, R., Spooner, F., McRae, L., Purvis, A., & Freeman, R. (2023). Ongoing over-exploitation and delayed responses to environmental change highlight the urgency for action to promote vertebrate recoveries by 2030. *Proceedings of the Royal Society B*, 290, 20230464. https://doi.org/10.1098/rspb.2023.0464

Cox, A. R., Roy, C., Hanson, A., & Robertson, G. J. (2024). Canadian murre harvest management in the face of uncertainty: a potential biological removal approach. *The Journal of Wildlife Management*, 88, e22573. https://doi.org/10.1002/jwmg.22573

Cox, A. R., Gallo-Cajiao, E., Tremblay, F., Rateau, F., Uroy, K., Laliberté, B., Euphrosine, D., Lavanne, T., & Roy, C. (2025). *Legal harvest of shorebirds and resident game birds on a Caribbean island: A Martinique case study* [Preprint]. bioRxiv. https://doi.org/10.1101/2025.06.17.660260

Deniau, C. P., Mathevet, R., Gautier, D., Besnard, A., Cornu, G., & Le Bel, S. (2022). Flyways to hell? An empirical assessment of Palearctic migratory waterbird harvest practices in key wetlands of Sahel-sub-Saharan Africa. *Biological Conservation*, 265, 109419. https://doi.org/10.1016/j.biocon.2021.109419

Eliason, S. L. (1999). The illegal taking of wildlife: Toward a theoretical understanding of
poaching. *Human Dimensions of Wildlife*, 4, 27–39.

Ellis, M. B., & Miller, C. A. (2022). The effect of a ban on the use of lead ammunition for
waterfowl hunting on duck and goose crippling rates in Illinois. *Wildlife Biology*,
2022(2), Article e01001. https://doi.org/10.1002/wlb3.01001

Eriksson, L., Månsson, J., Liljebäck, N., Sandström, C., Johansson, M., Eklund, A., &
Elmberg, J. (2023). The importance of structural, situational, and psychological factors
for involving hunters in the adaptive flyway management of geese. *Scientific Reports*,
13, 7112. https://doi.org/10.1038/s41598-023-34316-x

Food and Agriculture Organization of the United Nations. (2024). *Country profile –*
Suriname: Economy, agriculture and food security. FAO Knowledge Repository.
https://openknowledge.fao.org/handle/20.500.14283/ca0427en

Gallo-Cajiao, E., Morrison, T. H., Woodworth, B. K., Lees, A. C., Szabo, J. K., & Fuller, R.
A. (2020). Extent and potential impact of hunting on migratory shorebirds in the Asia-
Pacific. *Biological Conservation*, 246, 108582.
https://doi.org/10.1016/j.biocon.2020.108582

Gebauer, M. (2018). *Historic and current shorebird hunting and management structure in*
Trinidad and Tobago [Unpublished report]. Environment and Climate Change
Canada, Canadian Wildlife Service.

Geissler, P. (1990). Estimation of confidence intervals for federal waterfowl harvest surveys.
Journal of Wildlife Management, 54, 201–205.

Gelman, A., Carlin, J. B., Stern, H. S., Dunson, D. B., Vehtari, A., & Rubin, D. B. (2013).
Bayesian data analysis (3rd ed.). CRC Press/Chapman & Hall.

Hutt, M. B. (1991). Shooting of migrating shorebirds in Barbados. In T. Salathé (Ed.),
Conserving migratory birds (pp. 77–91). International Council for Bird Preservation.

Iskandar, S., & Karlina, E. (2004). Kajian pemanfaatan jenis burung air di pantai utara
Indramayu, Jawa Barat. *Buletin Plasma Nutfah*, 10, 43–48.

International Union for Conservation of Nature (IUCN). (2022). *The IUCN Red List of*
Threatened Species. Version 2022-2. https://www.iucnredlist.org/

Jacoby, K. (2014). *Crimes against nature: Squatters, poachers, thieves, and the hidden*
history of American conservation. University of California Press.

Katzner, T., Nelson, D. M., Diffendorfer, J. E., Duerr, A. E., Campbell, S. P., Leslie, D., &
Yee, J. L. (2020). Illegal killing of nongame wildlife and recreational shooting in
conservation areas. *Conservation Science and Practice*, 2, e279.
https://doi.org/10.1111/csp2.279

Kellner, K. (2021). *jagsUI: A wrapper around 'rjags' to streamline 'JAGS' analyses* [R
package]. https://CRAN.R-project.org/package=jagsUI

Kokko, H. (2001). Optimal and suboptimal use of compensatory responses to harvesting:
Timing of hunting as an example. *Wildlife Biology*, 7, 141–150.
https://doi.org/10.2981/wlb.2001.018

Koneff, M. D., Zimmerman, G. S., Dwyer, C. P., Fleming, K. K., Padding, P. I., Devers, P.
K., Johnson, F. A., Runge, M. C., & Roberts, A. J. (2017). Evaluation of harvest and
information needs for North American sea ducks. *PLoS One*, 12, e0175411.
https://doi.org/10.1371/journal.pone.0175411

Lindén, A., & Mäntyniemi, S. (2011). Using the negative binomial distribution to model
overdispersion in ecological count data. *Ecology*, 92, 414–1421.
https://doi.org/10.1890/10-1081.1

Lok, T., Overdijk, O., & Piersma, T. (2015). The cost of migration: spoonbills suffer higher
mortality during trans-Saharan spring migrations only. *Biology Letters*, 11, 20140944.
https://doi.org/10.1098/rsbl.2014.0944

Makowski, D., Ben-Shachar, M. S., & Lüdecke, D. (2019). bayestestR: Describing effects and
their uncertainty, existence and significance within the Bayesian framework. *Journal*
of Open Source Software, 4, 1541. https://doi.org/10.21105/joss.01541

Sainz-Borgo, C. (2025). Scarlet Ibis (*Eudocimus ruber*), version 2.0. In *Birds of the World* (N.
D. Sly, Editor). Cornell Lab of Ornithology, Ithaca, NY, USA.
https://doi.org/10.2173/bow.scaibi.02

Maxwell, S. L., Fuller, R. A., Brooks, T. M., & Watson, J. E. (2016). Biodiversity: The
ravages of guns, nets and bulldozers. *Nature*, 536, 143–145.
https://doi.org/10.1038/536143a

Moreno-Zárate, L., Arroyo, B., Sardà-Palomera, F., Rocha, G., Bota, G., & Peach, W. J.
(2023). Age ratio, crippling losses and factors affecting daily hunting bags of
European turtle-dove in Spain: Implications for sustainable harvest management of a
declining migratory species. *Science of The Total Environment*, 868, Article 161192.
https://doi.org/10.1016/j.scitotenv.2022.161192

Morrison, R. G., Mizrahi, D. S., Ross, R. K., Ottema, O. H., de Pracontal, N., & Narine, A.
(2012). Dramatic declines of Semipalmated Sandpipers on their major wintering areas
in the Guianas, northern South America. *Waterbirds*, 35, 120–134.
https://doi.org/10.1675/063.035.0112

Morrison, R. G., & Ross, R. K. (1989). Atlas of Nearctic shorebirds on the coast of South
America. Canadian Wildlife Service Special Publication, Ottawa, Ontario, Canada.

Murray, N. J., Phinn, S. R., DeWitt, M., Ferrari, R., Johnston, R., Lyons, M. B., Clinton, N.,
Thau, D., & Fuller, R. A. (2019). The global distribution and trajectory of tidal flats.
Nature, 565, 222–225. https://doi.org/10.1038/s41586-018-0805-8

Murray, N. J., Worthington, T. A., Fuller, R. A., Bi, J. J., Choi, C. Y., Clinton, N., & Lyons,
M. B. (2022). High-resolution mapping of losses and gains of Earth's tidal wetlands.
Science, 376, 744–749. https://doi.org/10.1126/science.abm9583

Naves, L. C., Keating, J. M., Tibbitts, T. L., & Ruthrauff, D. R. (2019). Shorebird subsistence
harvest and indigenous knowledge in Alaska: Informing harvest management and
engaging users in shorebird conservation. *The Condor*, 121, duz023.
https://doi.org/10.1093/condor/duz023

Nichols, J., Conroy, M., Anderson, D., & Burnham, K. (1984). Compensatory mortality in
waterfowl populations: a review of the evidence and implications for research and
management. *Transactions of the North American Wildlife and Natural Resources*
Conference, 49, 535–554.

Olmos, F., & Silva e Silva, R. (2001). Breeding biology and nest characteristics of the Scarlet
Ibis in Southeastern Brazil. *Waterbirds*, 24, 58–67. https://doi.org/10.2307/1522244

Ottema, O. H., & Spaans, A. L. (2008). Challenges and advances in shorebird conservation in
the Guianas, with a focus on Suriname. *Ornitología Neotropical*, 19, 339–346.

Ostrom, E. (2009). A general framework for analyzing sustainability of social-ecological systems. *Science*, 325(5939), 419-422.

Plummer, M. (2003). *JAGS: A program for analysis of Bayesian graphical models using Gibbs sampling* [Conference paper]. 3rd International Workshop on Distributed Statistical Computing.

R Core Team. (2020). *R: A language and environment for statistical computing*. R Foundation for Statistical Computing. https://www.R-project.org/

Raftovich, R., Chandler, S., & Fleming, K. (2018). *Migratory bird hunting activity and harvest during the 2016–17 and 2017–18 hunting seasons*. U.S. Fish and Wildlife Service.

Ramachandran, R., Kumar, A., Gopi Sundar, K. S., & Bhalla, R. S. (2017). Hunting or habitat? Drivers of waterbird abundance and community structure in agricultural wetlands of southern India. *Ambio*, 46, 613–620. https://doi.org/10.1007/s13280-016-0861-4

Ramsar Convention Secretariat. (1997). *Information sheet on Ramsar wetlands (RIS): Coppenamemonding Nature Reserve*. Ramsar Sites Information Service. https://rsis.ramsar.org/ris/304

Reed, E. T., Kardynal, K. J., Horrocks, J. A., & Hobson, K. A. (2018). Shorebird hunting in Barbados: Using stable isotopes to link the harvest at a migratory stopover site with sources of production. *The Condor*, 120, 357–370. https://doi.org/10.1650/CONDOR-17-134.1

Rushing, C. S., Hostetler, J. A., Sillett, T. S., Marra, P. P., Rotenberg, J. A., & Ryder, T. B. (2017). Spatial and temporal drivers of avian population dynamics across the annual cycle. *Ecology*, 98, 2837–2850. https://doi.org/10.1002/ecy.1983

Sandercock, B. K., Nilsen, E. B., Brøseth, H., & Pedersen, H. C. (2011). Is hunting mortality additive or compensatory to natural mortality? Effects of experimental harvest on the survival and cause-specific mortality of willow ptarmigan. *Journal of Animal Ecology*, 80, 244–258. https://doi.org/10.1111/j.1365-2656.2010.01769.x

Schulz, J. H., Bonnot, T. W., Millspaugh, J. J., & Mong, T. W. (2013). Harvest and crippling rates of mourning doves in Missouri. *Wildlife Society Bulletin*, 37, 287–292. https://doi.org/10.1002/wsb.249

Sergio, F., Tavecchia, G., Tanferna, A., Blas, J., Blanco, G., & Hiraldo, F. (2019). When and where mortality occurs throughout the annual cycle changes with age in a migratory bird: Individual vs population implications. *Scientific Reports*, 9, 17352. https://doi.org/10.1038/s41598-019-53531-z

Smith, A. C., Villeneuve, T., & Gendron, M. (2022). Hierarchical Bayesian integrated model for estimating migratory bird harvest in Canada. *The Journal of Wildlife Management*, 86, e22160. https://doi.org/10.1002/jwmg.22160

Smith, P. A., Smith, A. C., Andres, B., Francis, C. M., Harrington, B., Friis, C., Morrison, R. G., Paquet, J., Winn, B., & Brown, S. (2023). Accelerating declines of North America's shorebirds signal the need for urgent conservation action. *Ornithological Applications*, 125, duad003. https://doi.org/10.1093/ornithapp/duad003

Solomon, J. N., Gavin, M. C., & Gore, M. L. (2015). Detecting and understanding non-compliance with conservation rules. *Biological Conservation*, 189, 1–4. https://doi.org/10.1016/j.biocon.2015.04.029

Spaans, A. L. (1978). Status and numerical fluctuations of some North American waders
along the Surinam coast. *Wilson Bulletin*, 90, 60–83.

Spaans, A. L., Ottema, O. H., & Ribot, J. H. J. M. (2018). F*ield guide to the birds of*
Suriname (2nd ed.). Koninklijke Brill.

Steffen, W., Grinevald, J., Crutzen, P., & McNeill, J. (2011). The Anthropocene: From global
change to planetary stewardship. *Ambio*, 40, 739–761. https://doi.org/10.1007/s13280-
011-0185-x

Sutherland, W. J., Gardner, T. A., Haider, L. J., & Dicks, L. V. (2014). How can local and
traditional knowledge be effectively incorporated into international assessments?
Oryx, 48(1), 1-2.

Taylor, A. R. (2017). *Hunter surveys in the Mana rice fields, French Guiana* [Final report].
Department of Geography and Environmental Studies, University of Alaska
Anchorage.

Vaske, J. J., & Manfredo, M. J. (2012). Social psychological considerations in wildlife
management. In D. J. Decker, S. J. Riley, & W. F. Siemer (Eds.), *Human dimensions*
of wildlife management (pp. 43–57). JHU Press.

von Essen, E., Hansen, H. P., Nordström Källström, H., Peterson, M. N., & Peterson, T. R.
(2014). Deconstructing the poaching phenomenon: a review of typologies for
understanding illegal hunting. *British Journal of Criminology*, 54, 632–651.
https://doi.org/10.1093/bjc/azu022

Watts, B. D., Reed, E. T., & Turrin, C. (2015). Estimating sustainable mortality limits for
shorebirds using the Western Atlantic Flyway. *Wader Study*, 122, 37–53.
https://doi.org/10.18194/ws.00002

Watts, B. D., & Turrin, C. (2016). Assessing hunting policies for migratory shorebirds
throughout the Western Hemisphere. *Wader Study*, 123, 6–15.
https://doi.org/10.18194/ws.00023

Wauchope, H. S., Jones, J. P., Geldmann, J., Simmons, B. I., Amano, T., Blanco, D. E.,
Fuller, R. A., Johnston, A., Langendoen, T., Mundkur, T., & Nagy, S. (2022).
Protected areas have a mixed impact on waterbirds, but management helps. *Nature*,
605, 103–107. https://doi.org/10.1038/s41586-022-04617-0

Wetlands International. (2026). Waterbird Populations Portal. Retrieved January 1, 2026,
from https://wpp.wetlands.org

Western Hemisphere Shorebird Reserve Network. (2024). *WHSRN site profile: Suriname*
coastal wetlands. https://whsrn.org/whsrn-sites/

Zöckler, C., Hla, T. H., Clark, N., Syroechkovskiy, E., Yakushev, N., Daengphayon, S., &
Robinson, R. (2010). Hunting in Myanmar is probably the main cause of the decline of
the Spoon-billed Sandpiper Calidris pygmeus. *Wader Study Group Bulletin*, 117, 1–8.