Correlated evolution of the neck, head and forelimb across the theropod-bird transition

Ryan D. Marek[1*] and Ryan N. Felice[1,2,3]

[1]Centre for Integrative Anatomy, Department of Cell and Developmental Biology, University College London, London, UK

[2]Natural History Museum, London, UK

[3]Department of Genetics, Evolution and Environment, University College London, London, UK

[*]Corresponding author

Correspondence (email): r.marek@ucl.ac.uk

Correspondence (Twitter): @ryndmrk (Ryan D. Marek), @rfelice (Ryan N. Felice)

Acknowledgements: We would like to thank the following museum curatorial staff for their help in organizing and hosting trips to their institutions: Dr. Matthew Carrano and Amanda Millhouse at the Smithsonian Museum of Natural History, Kevin Seymour at the Royal Ontario Museum, Dr. Matt Lamanna and Amy Henrici at the Carnegie Museum of Natural History, Carrie Levitt-Bussian and Dr. Randy Irmis at the Natural History Museum of Utah, Carl Mehling, Ruth O'Leary, Suzann Goldberg and Prof. Jin Meng at the American Museum of Natural History. We would also like to thank the Leverhulme Trust (Research Project Grant award number: 182676) and the Bogue Fellowship scheme at University College London for funding this work.

Abstract

Powered flight has required birds to undergo numerous dramatic and coordinated
evolutionary responses across the entire body, yet studies are limited to a small number of
traits and often exclude a critical component of the vertebrate skeleton – the vertebrae. The
neck is a critical region of the avian spine as it operates in tandem with the head as a
surrogate forelimb across a diverse array of behaviours. However, the drivers of cervical
vertebral evolution remain poorly understood. Here, we model shifts in adaptive optima and
evolutionary rates of the neck, forelimb and head of extinct dinosaurs and extant birds to test
if these modules co-evolved. We observe a co-occurrence of adaptive optima shifts for neck
and forelimb proportion at the base of Avialae – to vertebrae adapted for stability and a
forelimb better adapted for flight. These patterns are due to shifts in neck and forelimb
allometry and suggest that heterochrony is an important factor in avian neck and forelimb
evolution. Further, we find lower rates of both neck and forelimb evolution in birds compared
to their non-avian theropod ancestors. The coordinated evolutionary response of the neck
and forelimb is a derived feature of Avialae that initially evolved to stabilise in-flight vision.
This axio-appendicular co-evolution has contributed to avian macroevolutionary dynamics by
facilitating the evolution of a novel locomotory mode without sacrificing the grasping
capability needed to directly interact with a huge diversity of environments.

Introduction

Powered flight is a key behavioural innovation, one that has allowed multiple clades across
the tree of life to thrive in entirely new niches (1–4). Birds are the exemplars of this
innovation as flight has allowed them to become the most speciose clade of extant
Tetrapoda (1). Such ecological diversification is thanks to an extraordinary transformation
from their iconic non-avian dinosaur antecedents that involved, among other morphological
and neurosensory transformations, sustained miniaturization (5), a decoupling of fore- and
hindlimb modules (6–8) and numerous changes to skull morphology and integration patterns
(9–12). Research into the evolution of avian powered flight is focused on cranial (10,11) and
appendicular adaptations (7,8,13), with little focus on other functional and anatomical
modules, nor on the coordinated response to flight between these modules (although studies
that consider multiple anatomical modules are starting to appear, see references 15–18). A
key component of the skeleton that is rarely discussed in this context is the neck. In extant
avians the neck is an important element of the axial skeleton that positions the head to
partake in a wide variety of behaviours – from preening to tool use to even tripedal
locomotion (18–23).

Over the course of theropod evolution the neck appears to have undergone a morphological
shift, from a robust musculoskeletal system primarily adapted for supporting a large head
and assisting in hypercarnivory (24–26), to an elongated and gracile configuration that
supports a myriad of key behaviours (27–29). This transformation is theorized to be
correlated with shifts in other major components of the prethoracic skeleton such as
decreases in head size and the loss of a grasping-capable forelimb as powered flight
evolved (16,29–32). However, studies into early avian evolution often consider
morphological systems in isolation and exclude the axial skeleton entirely (7,8,10,11,13). As
such, the timings and mechanisms that govern the evolution of this critical portion of the
avian skeleton remain unknown. One such mechanism that is responsible the evolution of
morphological adaptations is heterochrony, which is defined as the change in timing and
rates of developmental processes (33,34). Heterochronic shifts to a shorter development
period result in paedomorphosis, whereby the descendent clade evolves to resemble the
juvenile form of the ancestral clade. Paedomorphosis is an important mechanism as it is

credited as the mechanism behind many key avian innovations such as a small body size
(5,35,36) and a relatively enlarged orbit and neurocranium (37). However, the contribution of
heterochrony to the evolution of avian postcranial morphology, especially axial morphology,
is understudied.

The potential correlated evolution of the neck, head and forelimb across the dinosaur-bird
transition is theorized to result in the neck functioning as a 'surrogate forelimb' in extant
birds, allowing the functional burden of environmental manipulation to be shifted to the neck
from the forelimb (29–31). Evidence for the correlated evolution of head, neck and forelimb
is present in extant avians as significant morphological integration exists between all three of
these modules in many groups of birds (32). This pattern is exemplified in arboreal and
aquatic birds, as integration promotes shifts in the rate of neck morphological evolution (32).
Thus, the integration of the neck, head and forelimb is a keystone innovation of birds as it
has facilitated a coordinated response of the prethoracic skeleton when entering new niches,
allowing birds to successfully interact with new ecosystems (29,30,32). Understanding how
and when this correlation between the head, neck and forelimb evolved in avians and their
dinosaurian ancestors is thus a critical step in further understanding how birds came to
succeed in modern ecosystems.

There are numerous examples of separate anatomical modules co-evolving during the
evolution of birds (6–8,14,15,17,38) in a mosaic fashion (whereby separate anatomical
modules display distinct evolutionary modes and timings (38)) and these are key features of
avian macroevolution (6,8,10,14). Across the theropod-bird transition, the pelvic and pectoral
locomotor modules decoupled and this promoted the co-evolution of each locomotor module
with other anatomical modules (6–8). Coordinated evolution of the forelimb and tail gave rise
to the flight control system exhibited by extant avians, and by emphasizing different
combinations of the wing, hindlimb and tail, the diversity of avian locomotor diversity
increased (6,7). Recent work has shown that in extant avians this mosaic pattern of
modularity also includes the head and neck, as both affiliate with the forelimb to allow birds
to succeed in arboreal and aquatic environments (32). This paper aims to build on this
recent work and uncover how the potential head-neck-forelimb integration evolved in birds,
and when this feature arose in avian evolutionary history. By understanding this we will gain
a more complete insight into how integration and modularity promote the evolution of key
innovations across the broader avian bauplan in response to the evolution of powered flight.

The correlated evolution between the head and the neck may have arisen in birds due to the
reduction in head mass along the avian stem lineage (31,32). Head mass is a known
constraint on neck morphology across many terrestrial vertebrates as the weight of the head
must be supported by the neck at all times (39–42). As relative head size is thought to
decrease as theropod diet shifts away from hypercarnivory along the avian stem lineage
(43,44), it is predicted that neck morphology would be released from the constraint of head
size (29,31). We hypothesise that this reduction in head size could be the cause of this
head-neck integration (32). Neck-forelimb integration may have arisen alongside the loss of
the forelimb's grasping capability as powered flight evolved in early avialans (29–32). It has
been proposed the neck of birds evolved due to a morphofunctional compensation that was
driven by heterochrony between the loss of hand functions of theropods as the forelimb was
integrated into the avian wing and as the kinematic functions of the avian beak were
enhanced (30). This 'surrogate forelimb' hypothesis is supported our recent work on extant
avians (31,32), however has yet to be formally tested by assessing the tempo and
directionality of neck and forelimb evolution across the theropod-bird transition

Here we investigate how and when the unique avian neck evolved by exploring how the
interplay between the evolution of the neck and other regions of the body influenced the
macroevolutionary dynamics of birds and their theropod ancestors, as well as exploring the
heterochronic patterns of the neck, head and forelimb. We use Bayesian Ornstein
Uhlenbeck (OU) modelling approaches (45–47) to model shifts in adaptive optima as well as
shifts in evolutionary rates of head, neck and forelimb allometry across 222 species of avian
and non-avian theropods. We predict that concurrent shifts in the adaptive regimes of neck
morphology and forelimb morphology, to shorter centrum lengths and a more distally-
dominated forelimb respectively, would support the surrogate forelimb hypothesis.
Conversely, if neck evolution is primarily driven by a relaxing of head mass constraint as
head size decreases along the avian stem lineage we predict that a shift to a lower head
mass around the base of Maniraptora would be immediately followed by a shift in adaptive
regimes of neck morphology, as well as an increase in rates of neck morphological evolution.
We predict that both of these hypothesis will involve paedomorphic shifts in neck, forelimb
and head allometry. Since decreases body size are an important facet of early avialan
evolution due to the extensive miniaturization along the avian stem lineage we explore the
degree to which the observed patterns in neck, head and forelimb morphology were related
to allometric effects. We use this methodology to specifically answer whether the evolution
of the avian cervical spine was correlated with the loss of a grasping-capable forelimb and if
neck evolution was constrained by large head size in non-avian theropods. By investigating
the correlated evolution of multiple anatomical modules this study aims to unearth how a key
innovation, powered flight, led to an evolutionary cascade of adaptations across the
prethoracic skeleton that negated the negative consequences of evolving a grasping-
incapable wing.

Methods

Specimen information & data collection

We collected length measurements for the cervical centra, forelimb bones (humerus, radius,
ulna, longest metacarpal element) and skull of 222 species (224 specimens) of avian and
non-avian theropods (Table S1). 112 of these species were extant avians and the remaining
110 were extinct avialan and non-avialan theropods (Table S1). We measured 20 of the 110
species of theropods directly from surface scans collected with an Artec Leo 3D surface
scanner from multiple North American museums (Table S1). We then measured the
remaining 90 specimens from the literature (Table S1). Finally, we took avian skeletal
measurements from the dataset of Marek & Felice 2023. We assessed centrum length along
the ventral midline of the centrum across all specimens (27), and then calculated centrum
length for extant avians using the interlandmark distance between appropriate centrum
landmarks used in Marek & Felice 2023 (see Fig S1 of ref. 33). We chose length of forelimb
elements and relative forelimb proportion (brachial index) as they capture the major axes of
forelimb morphological variation (17) and correlate with flight capability in extant avians (48–
51). Centrum length was chosen as it correlates with measures of intervertebral flexibility
(52,53) (although see ref. 28)).

Phylogenetic Hypotheses

We assembled an informal supertree using a recent hypothesis of the phylogenetic
relationship of birds based on 63k intergenic coding regions (54) as a backbone. We then
grafted a time-calibrated supertree of non-neornithine birds (55) on to this backbone using
the 'tree.merger' function in the 'RRphylo' R package (56). This crown and stem bird
supertree was extended by grafting on a second supertree that spanned the entire Mesozoic
theropod phylogeny (including early diverging avialans) (57). We added theropod and early

avialan taxa that were present in our dataset but missing from the supertree at this stage by
grafting individual taxa from a composite tree of Dinosauria that was pruned to include just
theropods (35). We time calibrated the trees from Wang & Zhou (57) and Benson et al. (35)
prior to grafting the backbone. This was achieved by using the tip dates bracketed by the
first and last appearance dates of the geological stages in which each taxon was collected.
We calibrated the respective trees using these first and last appearance dates using the
'minimum branch length ('mbl', one million years) methods within the 'timePaleoPhy' function
of the 'paleotree' R package (58). These time calibration methods were similar to those used
in the original publications (57). We adapted procedures from Cooney et al. (59) that
combined the backbone of major avian relationships from (54) with the fine-scale species
relationships from a maximum clade credibility tree generated from www.birdtree.org (60) to
produce a phylogeny that contained all the extant avian species. This final tree of 10,199
species was pruned to match the 222 species in our dataset.

Evolutionary shifts in the adaptive optima of neck, forelimb and head morphology

We estimated adaptive shifts in cervical, forelimb and head morphology using a Bayesian
multi-regime Ornstein Uhlenbeck (OU) modelling approach in the R package 'bayou' (45). A
phylogenetic generalized least squares (PGLS) regression indicated that the lengths of the
cervical centra, forelimb and head and forelimb proportion metrics all displayed a significant
relationship ($p < 0.001$) with body mass. We therefore created custom bayou models that
included body mass a predictor variable to test for shifts in the allometric scaling of cervical,
pectoral and cranial morphology. We used these custom models to test specific biological
hypotheses. First, we tested whether adaptive shifts to a more distally-dominated forelimb
were concurrent with shifts in neck morphology at the base of Avialae, this would allow us to
understand whether the evolution of the avian neck was correlated with the loss of a
grasping-capable forelimb. Second, we tested whether adaptive shifts to a shorter relative
head length were concurrent with shifts in neck morphology to understand if neck evolution
was constrained at all by large head size in non-avian theropods. We used the
'shiftSummaries' and 'plotShiftSummaries' bayou functions to determine if adaptive shifts
were heterochronic by observing if a shift in y-intercept (trait morphology) occurred between
the ancestral and descendent regimes that were identified by the above bayou models.

We selected priors for the constraint parameter (α), evolutionary rate (σ^2), and phenotypic
optimum (θ) to ensure acceptance ratios remained between 0.2 and 0.4 for each parameter.
We set the prior for the body mass regression coefficient (β_{BM}) to 0.7, drawing on previous
studies of avian neck allometry (31). Once the priors were set, we ran five independent
MCMC chains and assessed convergence for each chain using the Gelman and Rubin's R
statistic (61,62). We used the combined chains resulting from this procedure to assess
allometric shifts in adaptive optima of each measurement of cervical, forelimb and head
morphology. For each individual dataset (first/mid/last centrum length,
humerus/radius/ulna/MC2 length and head length) five MCMC chains of 1,000,000
generations were run independently and then combined, with the first 30% of each chain
were removed as burn-in. We performed model selection by examining marginal likelihoods
and Bayes Factors of each model (Tables S2, S3). In each case (aside for MC2), the highest
likelihood model allowed for separate intercepts and global slopes where shifts have an
equal probability across branches and can only occur once per branch.

Evolutionary shifts in the rate of trait evolution of the neck, forelimb and head

We used fossil BAMM v2.5.0 (46,47) to model the variation and potential shifts in the rate of
morphological evolution of neck, forelimb and head morphology. We modelled rates using a
MCMC chain with 90,000,000 generations for each trait analysed. We set model priors

automatically using the 'setBAMMpriors' function in the BAMMtools R package. We sampled
parameters every 10000 iterations, the first 10% of which were deleted as burn-in. We
analysed the effective sample size of the log-likelihood and number of shift events of each
chains in order to assess convergence and found that all chains in this study successfully
converged (effective sample size, ESS, > 200). We accounted for changes in body mass by
running a second round of MCMC chains that used the residuals from phylogenetic GLS
regressions of each morphological trait versus body mass. We carried out these regressions
using the 'gls' function of the R package 'nlme'.

Assessing the influence of body mass on shifts in neck, head and forelimb
morphology

We calculated the phylogenetic means for head, neck and forelimb morphology, as well as
body mass for each of the regimes ('descendent' grade) found in the bayou analysis and
compared them to the mean of the ancestral grade following procedures from Smaers et al.
(63–65) We used this procedure to assess the extent to which body mass influenced the
recovered patterns of neck, head and forelimb morphology across the shifts recovered in the
bayou analysis. The ratio of the difference in ancestral-to-descendent mean trait
(neck/head/forelimb) morphology to the difference in ancestral-to-descendant mean body
mass is an indication of the proportionality of ancestral-to-descendant change in mean trait
morphology to mean body mass. We set the expected proportionality of this ratio as the
scaling coefficient of the trait-to-body relationship of the ancestral grade. We then used the
upper bound of the 95% confidence interval of the ancestral grade scaling coefficient to infer
that the change in mean trait morphology from ancestral-to-descendant grade is
characterized by more or less change in the trait than mean body mass. This process was
inversed to evaluate the mass-to-trait scaling relationship (63,64).

Results

Evolutionary shifts in adaptive allometric optima of neck, forelimb and head
morphology

There is strong support for a singular regime shift in the adaptive allometric optima of
vertebral morphology on the branch leading to Avialae in all three cervical vertebral regions
studied (posterior probability = 0.70-0.82, Figs 1, S1). Across all three studied vertebral
regions, avialans and non-avian theropod dinosaurs occupied distinct adaptive optima, with
avialans displaying shorter cervical vertebrae relative to non-avian theropods (Figs 1, S1).
The aforementioned shift in vertebral morphology is associated with a downshift in the
allometric scaling of centrum length (Figs 1, S1) meaning that centrum length is lower than
expected for a given body mass in avialans when compared to non-avialan theropods. The
slopes of these scaling relationship does not change across any of the observed regime
shifts.

Our models of humerus, radius, ulna and MC2 length scaling recovered an adaptive
optimum regime shift in the branch leading to Apodiformes (posterior probability = 0.44-0.95,
Fig 2). In each case, these shifts are to shorter bone lengths (Fig S2). This represents a shift
to a distinct forelimb phenotypic trait optima in Apodiformes (hummingbirds and allies). The
low number of Apodiformes present in this study makes this shift difficult to visualise on a
phenogram (Fig 2), however the shift is well supported with high values of posterior
probability ($PP_{humerus}$ = 0.96, PP_{radius} = 0.67, PP_{ulna} = 0.70, PP_{MC2} = 0.45) and low standard
error values ($SE_{humerus}$ = 0.002, SE_{radius} = 0.003 , SE_{ulna} = 0.003, SE_{MC2} = 0.002) across all
forelimb elements (Fig S3). The shifts to lower individual forelimb element length in
Apodiformes are associated with a downshift in allometric scaling relationship (Fig 2, S2). In

the radius and ulna these downshifts are also associated with an increase in the slope of the
allometric scaling relationship (Fig S2). The humerus displays three additional regime shifts
in allometric adaptive optima: one on the branch leading to Psittacopasseres (i.e., parrots
and perching birds; bayou posterior probability = 0.52), one within Ornithomimosauria on the
branch leading to Ornithomimidae and Deinocheridae (bayou posterior probability = 0.63,Fig
2a) and one on the branch leading to *Ornitholestes hermanni* and Compsognathidae (bayou
posterior probability = 0.34, Fig 2a). In the case of Psittacopasseres, this shift is towards a
shorter humerus and this clade appears to form a distinct phenotypic optima compared to
the rest of the species studied. The shift leading to Ornithomimidae and Deinocheridae is
characterized by an increase in humerus length, but does not appear to form a distinct
adaptive zone. These additional shifts are associated with a downshift in relative humerus
length in Psittacopasseres and Compsognithidae+*Ornitholestes* and an upshift within
Ornithomimosauria meaning that for a given humeri are relatively longer per a given body
mass in Ornithomimidae+Deinocheiridae than compared to their ancestors.

There are multiple regime shifts in the adaptive optima of forelimb proportions (Fig 2). There
is a regime shift in brachial index representing a shift in phenotypic optima to relatively
longer proximal portion of the forelimb on the branch leading to Ceratosauria (bayou
posterior probability = 0.81). There are two further regime shifts in adaptive optima in BI: one
on the branch leading to Avialae (bayou posterior probability = 0.58) and one on the branch
leading to Sphenisciformes (bayou posterior probability = 0.66, Fig 3). Avialans and non-
avian theropods appear to occupy separate adaptive optima in BI, with avialans exhibiting
lower BI scores than non-avian theropods Fig 3). Avialans display a downshift in the
allometric scaling of BI relative to non-avialan theropods (Fig 3). Sphenisciformes are the
exception to this rule and display much higher values for BI than any other avialan (extinct or
extant), and fall well within the non-avian theropod BI adaptive peak (Fig 3). This shift is
associated with an upshift in the allometric scaling of brachial index when compared to
sphenisciform ancestors (Fig 3).

Two regime shifts in adaptive optima were observed for head length: one on the branch
leading to *Mei long* (an early Cretaceous troodontid) and the other along the branch leading
to *Parus atricapillus* (black capped chickadee). These shifts correspond to a decrease in BI,
however neither form a new adaptive peak.

Shifts in the evolutionary rate of neck, forelimb and head morphology

Rates of cervical morphological evolution were generally lower in avialans than in non-avian
theropod dinosaurs (Fig 1). Shifts to decreased rates of centrum length evolution were
observed on the branch leading to Ornithuromorpha for the middle cervical vertebrae
(BAMM marginal shift probability = 0.35), and on the branch leading to Aves for the last
cervical vertebrae (BAMM marginal shift probability = 0.49). The first cervical vertebrae
displayed no shifts to lower rates of evolution. Shifts to higher rates of centrum length
evolution are frequently observed on branches leading to clades of herbivorous non-avian
theropods for the first and middle cervical vertebrae (Therizinosauria – first/middle: BAMM
marginal shift probabilities = 0.93/0.83, Alvarezsauroidea – first/middle: BAMM marginal shift
probabilities = 0.78/0.69, Ornithomimidae – first/middle: BAMM marginal shift probabilities =
0.55/0.44, Oviraptorosauria – first: BAMM marginal shift probability = 0.56). Additional shifts
to higher rates of centrum length evolution can be observed at the base of Bohaiornithidae
(first/middle: BAMM marginal shift probabilities = 0.61/0.79) and Emberizoidea (first/last:
BAMM marginal shift probabilities = 0.34/0.51). None of the above shifts directly correlate to
the recovered shifts in adaptive optima, however the two shifts to lower rates of centrum
length evolution for the middle and last cervical vertebrae are found along the avian stem

lineage, crown-ward of the avialan base. The support for the best shift configurations of
centrum length evolution across all cervical vertebrae was low (<4.1% of sampled posterior
distributions).

Similar to the patterns observed for centrum length rate evolution, shifts to higher rates of
morphological evolution can be observed on the branches leading to herbivorous theropods
across all individual forelimb elements studied (Alvarezsauroidea: BAMM marginal shift
probability = 0.45-0.73, Oviraptorosauria: BAMM marginal probability = 0.35-0.93, Fig 2).
Alvarezsauroidea displays the fastest rates of humerus, radius and MC2 evolution. An
additional shift to higher rates of morphological evolution can be observed on the branch
leading to Therizinosauria for MC2 length (BAMM marginal probability = 0.55, Fig 2). Rate
shifts across all studied individual forelimb elements did not match the shifts found for their
adaptive optima (Fig 2). Support for these 'best' configurations of rate shifts was low across
all individual forelimb elements (all represented <1% of the sampled posterior distribution.

A shift to a lower rate of brachial index evolution is observed on the branch leading to
Maniraptora (exclusive of Alvarezsauroidea, BAMM marginal shift probability = 0.18, Fig 3)
and this shift precedes the shift in brachial index adaptive optima. Shifts to faster rates of
brachial index evolution are reported along the branches leading to Anchiornithidae (BAMM
marginal shift probability = 0.67) and Longipteryigidae (BAMM marginal shift probability =
0.83), Fig 3.

There appears to be little support for any rate shifts in head length across theropod and bird
evolution, with the best shift configuration (accounting for 0.1% of the posterior distribution)
displaying no shifts in rate. In three of the top 10 most credible rate shift configurations a
shift to lower rates of head length evolution is observed on the branch leading to Avialae and
these three shift configurations account for 1.23% of the posterior distribution. These rate
shifts disappear after body mass is taken into account.

Influence of allometry on shifts in neck and forelimb morphology

Ancestor-to-descendent grade comparisons indicate that the shift to a lower centrum length
in Avialans that is observed across all neck vertebrae appears to be driven by decreases to
centrum length outpacing changes to body mass (Tables S4, S5). The avialan ancestral
grade has a mean centrum length of 3.29, 3.55 and 2.97 (log scale) and mean body mass of
1.20, 1.36 and 1.34 (log scale) for the first, middle and last cervical vertebrae respectively.
The avialan descendent grade has a mean centrum length of 1.56, 1.67 and 1.52 (log scale)
and mean body mass of -0.60, -0.59 and -0.58 (log scale) for the first, middle and last
cervical vertebrae respectively. The difference in mean centrum length from the avialan
ancestral grade to the avialan grade is -1.73, -1.89 and -1.45, whilst the difference in body
mass is -1.80, -1.95 and -1.92 the for the first, middle and last cervical vertebrae
respectively. The resultant $\frac{\Delta centrum\ length}{\Delta body\ mass}$ ratios across all cervical vertebrae (0.97, 0.97,
0.75) are higher (by 0.07, 0.31 and 0.01) that the upper bounds of the centrum-to-mass
expectations. The $\frac{\Delta body\ mass}{\Delta centrum\ length}$ ratios across all cervical vertebrae (1.04, 1.03 and 1.33)
are lower (by 0.41, 0.35 and 0.01) that the upper bounds of the mass-to-centrum
expectations.

A shift to a more distally-dominated forelimb at the base of Avialae is observed for brachial
index (Tables S4, S5). This shift appears to be driven by changes to body mass outpacing
the changes in forelimb proportions, however there is weak support for the opposite pattern
(Tables S4, S5). The avialan ancestral grade has a mean BI of 0.21 (log scale) and a mean
body mass of 1.23 (log scale). The avialan descendent grade has a mean BI of -0.06 (log

scale) and a mean body mass of -0.60 (log scale). The difference in BI from the avialan
ancestral grade to the avialan grade is -0.28, whilst the difference in body mass is -1.84. The
resultant $\frac{\Delta body\ mass}{\Delta BI}$ ratio for BI is higher (by 2.78) than the upper bounds of the mass-to-BI
expectations. The $\frac{\Delta BI}{\Delta body\ mass}$ ratio for BI is only slightly higher (by 0.01) than the upper
bounds of the BI-to-mass expectation.

Discussion

We observe a concurrent shift in the adaptive optima of neck morphology (Fig 1) and
forelimb proportion at the base of Avialae (Fig 3) to shorter centrum lengths in the former
and to a relative increase in the distal portion of the latter. We also observe a shift in the
allometric scaling of neck morphology and forelimb proportions (Figs 1, 3). Interestingly, we
find no significant shifts in either adaptive optima or evolutionary rate of the length of the
head (here used as a proxy for head size) across non-avian theropods or birds. These
results show the neck and forelimb morphology of birds and non-avian dinosaurs display
distinct evolutionary patterns as these two groups occupy separate phenotypic optima in
both the neck and forelimb. Avians also exhibit slower rates of morphological evolution (Figs
1, 2) across the head, neck and forelimbs, with significant rate shifts in forelimb proportion
and proximal neck morphology can occurring at the base of Maniraptora and within
Ornithuromorpha respectively.

We find no support for the hypothesis that the loss of hypercarnivory in theropods released a
constraint on neck evolution as no shifts in head length exist that are concurrent with (or
directly proceed or precede) shifts in neck morphology at the base of Avialae. We also find
very few shifts in relative head size across Theropoda more widely. Large hypercarnivorous
clades are emblematic of Theropoda and the lack of significant shifts to larger relative head
sizes is surprising given that hypercarniverous taxa have been shown to possess a greater
selective pressure to maintain a larger relative head size (66). Phylogeny is a strong
predictor theropod skull shape, more so than functional performance (67), therefore the lack
of head length shifts may be a product of phylogenetic constraint. The lack of significant
shifts in head size adaptive optima may also result from the measure of head size included
in this study, head length. We used this metric of head size to gain access to a wider
selection of theropod specimens, as relatively few are represented by a three-dimensionally
preserved skull, which is necessary for volumetric reconstructions. Despite this surprising
lack of adaptive optima shifts, our recovered trends in the rates of head morphological
evolution match prior work and suggest that birds have a lower rate of skull evolution
compared to non-avian theropods. Although head size and neck morphology may not co-
evolve, recent evidence suggests that the shape of the skull (specifically the occiput) and
neck morphology are integrated in extant birds due to their shared function and underlying
developmental mechanisms (68), and this represents an exciting future area of research.

Our results echo the established pattern that the sustained miniaturization that occurred
along the avian stem lineage heavily influenced the evolution of the Avialan appendicular
skeleton (5,35,57,69) as changes to body mass outpace shifts to a more distally-dominated
forelimb. Whereas forelimb evolution may be linked to selection towards a smaller body size,
the shift in morphology of cervical vertebrae in basal Avialae is not, as the change in centrum
length across this shift outpaces the change in body mass (Tables S2, S3). Despite not
resulting in any significant shifts in head size, carnivory is still frequently associated with
specific neck morphologies in both non-avian dinosaurs and extant avians (21,25,29,31) to
provide stability and increased retraction forces to allow flesh to be ripped from prey items.
Thus, we hypothesize that shifts in cervical morphology towards new adaptive zones is

driven by the diets of basal avialans shifting away from carnivory (70,71) rather than by
changes in body mass.

A shift to a more distally-dominated forelimb at the base of Avialae and to lower rates of
forelimb evolution at the base of Maniraptora is supported by previous research (13,57,69).
The overall proportions of the avian forelimb govern its functional capabilities. For example,
an ulna that is longer than the humerus is a structural requirement of avian volant flight (72–
74). This structural requirement of powered flight has been suggested to restrict the forelimb
morphology of early avialans and this restriction results in decelerated rates of forelimb
morphological evolution (57). This shift in forelimb proportion is associated with a concurrent
shift in vertebral morphology (centrum length), and this association has important functional
consequences for the evolution of powered flight. The shift to a shorter centrum indicates an
increase in cervical intervertebral stability (53,75,76), and this shift is also likely due to the
structural demands imposed by the evolution of flight. The neck of extant avians plays a
pivotal role in counteracting the effect of each wingbeat in order to stabilize vision during
even the most complex of aerial manoeuvres (77–79). This vision stabilisation function acts
as a constraint on extant avian neck morphology as soaring birds display higher levels of
morphological differentiation than their continual flapping counterparts (31). Whereas the
vestibulo-ocular reflex is also a critical component of avian gaze stabilisation (80), the
biomechanics of the avian neck (and hence the morphology of the neck vertebrae
themselves) allow for a stable vision by passively attenuating body oscillations produced
from each wingbeat (78). Gaze stabilization is critical to successful locomotion in any animal
(77,78,81) and is inferred here to act as a constraint on the evolution of neck morphology in
the early evolution of birds.

It is this constraint of providing stable vision during flight that may be responsible for the
decelerated rates of neck evolution that are observed throughout Avialae when compared to
non-avian theropods. Previous work has noted that extant avian cervical morphology is
generalized, with morphological adaptations only occurring to accommodate extreme neck
kinematics such as those present in carnivory (31). With both early diverging avialans and
extant avians showing slower rates of neck morphological evolution we suggest that the
generalized morphology of the extant avian neck is a product of the constraining effect of
gaze stabilization during flight. This is exemplified in the rate shift configuration of the last
cervical vertebrae, whereby a shift to a much lower rate of evolution can be observed at the
base of Aves (Fig. 1). The last cervical vertebrae is the most critical portion of the neck with
regards to gaze stabilisation as it functions as a fulcrum for the entire cervical spine (78) thus
the constraining effects of maintaining stable vision during flight should be, and are,
observed in this portion of the neck. In addition, we suggest that transitions in neck
morphology and forelimb proportion are not driven by shifts in the rate of morphological
evolution (and vice versa) as shifts in rate and adaptive optima appear decoupled across
Theropoda. The underlying drivers of evolutionary rate shifts in head size versus neck and
forelimb morphology are disparate. Rate shifts in both neck and forelimb morphology are
consistently found on the branches leading to herbivorous theropods (Alvarezsauroidea,
Ornithomimidae, Therizinosauria and Oviraptorosauria (70)). This suggests that whereas
phylogeny constrains head size rate evolution, shifts to elevated rates of neck and forelimb
evolution appear adaptive and are associated with dramatic ecological shifts away from
carnivory.

Heterochrony is a key mechanism of evolutionary change across vertebrates and has been
found to be a vital component of dinosaur evolution (37,82–85). By analysing the changes to
the intercept and slope of the allometric coefficients associated with the adaptive optima
shifts of neck and forelimb morphology (Figs 1, 2, S1, S2) we can assess potential

mechanisms behind the evolution of the avian neck and wing (86). Allometric shifts in
adaptive optima to lower relative centrum length (Figs 1, S1) and to a lower relative brachial
index are associated (Fig 3) with decreased y-intercepts and no changes to the slopes of
these scaling relationships. This suggests heterochronic shifts that delay the onset of growth
of these features (postdisplacement) are responsible for shifts in neck morphology and wing
proportion at the base of Avialae (33,34,86). Postdisplacement is a paedomorphic process,
one that results in the adult of the descendent clade (birds) resembling the juvenile stages of
the ancestral clade (non-avialan theropods). Paedomorphosis is responsible for the
evolution of key avian cranial characteristics (35,37) and this current work suggests that
paedomorphosis is a vital mechanism behind the axial and appendicular skeleton also. Our
results emphasise the importance of paedomorphosis in the emergence of avian powered
flight as it provides a provides the functional adaptations (the wing) as well as the processing
systems (neuroanatomical and vision stabilisation) required for life on the wing.

With concurrent shifts in adaptive optima we suggest that neck and forelimb evolution are
correlated in Avialae. The grasping capability of the theropod forelimb was lost during the
evolution of powered flight (72) and it has been hypothesized that the neck subsequently
evolved a surrogate forelimb functionality to provide birds with a means of interacting with
their environment without a grasping-capable forelimb (29–32). Some evidence for this
hypothesis exists, as morphological integration between the neck and forelimb is significant
in extant Aves and serves to influence the evolutionary variance of the neck (32). Neck-
forelimb integration in extant avians is a keystone innovation that can allow birds to explore
entirely new niches by utilizing the neck instead of the forelimb to interact successfully within
these niches (32). The concurrent shifts in neck and forelimb evolution displayed here
provide further support for the surrogate forelimb hypothesis and suggest that neck-forelimb
evolution is correlated during the evolution of avian powered flight (29,31). Powered flight is
a synapomorphy of birds and has contributed to their abundant diversity within modern
ecosystems (1). This correlated evolution of the neck and forelimb to new adaptive zones
represents a key innovation that has enabled birds to evolve this defining behavioral trait
without losing the functionality of the forelimb.

Mosaic evolution is a core concept of avian macroevolution (6,8). This mosaicism implies
that traits are organised into modules that share genetic and developmental origins. Due to
these shared origins, as well as their potential shared function, traits within these modules
will display a correlated response to selection. The neck and forelimb of extant birds display
a coordinated evolutionary response in the form of integration, and this may have resulted
from the contribution of migratory neck muscle precursor cells to forelimb muscle
development (40,76,87). These developmental pathways facilitating this integration were
likely present in basal avialans as they are shared across Tetrapoda (40,76) therefore the
concurrent shifts in forelimb and neck evolution at the base of Avialae represent the
beginning of this neck-forelimb integration. Our finding that postdisplacement is responsible
for both the shifts to relative shorter centra and to a more distally-dominated forelimb provide
further evidence that the neck and forelimb of avialans experience similar developmental
pathways. Body mass is an important determinant of the evolution of new modules within
organisms (6,15,88,89) and larger body sizes impose constraints on the evolvability of new
modules within species. The miniaturization that occurred along the Avialan stem lineage
may have eased these constraints and allowed for the evolution of new modules such as the
one observed between the neck and forelimb. Coordinated responses of anatomical
modules is a key feature of avian mosaicism and elaborations of different combinations of
locomotor modules (forelimb, hindlimb and tail) have allowed birds to occupy a diverse range
of niches by evolving a disparate array of locomotor behaviours (6,8,90). This study builds
on recent work (31,32) to establish that avian mosaicism is not limited to locomotor modules

alone (6,7). The coordinated response of neck and forelimb morphological evolution appears to be a derived feature of Avialans, initially providing them with stable vision as flight first evolved and then negating the negative consequences of losing a grasping capable forelimbs (29,31,32). In turn, this facilitated the ability for birds to interact with a multitude of different environments, and has contributed to the great taxonomic, morphological, and ecological diversity observed today.

References

1. Brusatte SL, O'Connor JK, Jarvis ED. The Origin and Diversification of Birds. Current Biology. 2015 Oct 5;25(19):R888–98.

2. Gill FB. Ornithology. Macmillan; 1995.

3. Dudley R, Pass G. Wings and powered flight: Core novelties in insect evolution. Arthropod Structure & Development. 2018 Jul 1;47(4):319–21.

4. Anderson SC, Ruxton GD. The evolution of flight in bats: a novel hypothesis. Mammal Review. 2020;50(4):426–39.

5. Lee MSY, Cau A, Naish D, Dyke GJ. Sustained miniaturization and anatomical innovation in the dinosaurian ancestors of birds. Science. 2014 Aug;345(6196):562–6.

6. Gatesy SM, Dial KP. Locomotor modules and the origin of avian flight. Evolution. 1996 Feb;50(1):331–40.

7. Eliason CM, Proffitt JV, Clarke JA. Early diversification of avian limb morphology and the role of modularity in the locomotor evolution of crown birds. Evolution. 2023 Feb 1;77(2):342–54.

8. Clarke JA, Middleton KM. Mosaicism, Modules, and the Evolution of Birds: Results from a Bayesian Approach to the Study of Morphological Evolution Using Discrete Character Data. Systematic Biology. 2008 Apr 1;57(2):185–201.

9. Torres CR, Norell MA, Clarke JA. Bird neurocranial and body mass evolution across the end-Cretaceous mass extinction: The avian brain shape left other dinosaurs behind. Science Advances. 2021 Jul 30;7(31):eabg7099.

10. Felice RN, Goswami A. Developmental origins of mosaic evolution in the avian cranium. Proceedings of the National Academy of Sciences. 2018 Jan 16;115(3):555–60.

11. Felice RN, Watanabe A, Cuff AR, Hanson M, Bhullar BAS, Rayfield ER, et al. Decelerated dinosaur skull evolution with the origin of birds. PLOS Biology. 2020 Aug 18;18(8):e3000801.

12. Chiappe LM, Navalón G, Martinelli AG, Carvalho I de S, Miloni Santucci R, Wu YH, et al. Cretaceous bird from Brazil informs the evolution of the avian skull and brain. Nature. 2024 Nov;635(8038):376–81.

13. Benson RBJ, Choiniere JN. Rates of dinosaur limb evolution provide evidence for exceptional radiation in Mesozoic birds. Proceedings of the Royal Society B: Biological Sciences. 2013 Oct 7;280(1768):20131780.

14. Navalón G, Bjarnason A, Griffiths E, Benson RBJ. Environmental signal in the evolutionary diversification of bird skeletons. Nature. 2022 Nov;611(7935):306–11.

15. Orkney A, Hedrick BP. Small body size is associated with increased evolutionary lability
of wing skeleton proportions in birds. Nat Commun. 2024 May 28;15(1):4208.

16. Böhmer C, Plateau O, Cornette R, Abourachid A. Correlated evolution of neck length
and leg length in birds. Royal Society Open Science. 2019 May 8;6(5):181588.

17. Orkney A, Bjarnason A, Tronrud BC, Benson RBJ. Patterns of skeletal integration in
birds reveal that adaptation of element shapes enables coordinated evolution between
anatomical modules. Nat Ecol Evol. 2021 Sep;5(9):1250–8.

18. Boas JEV. Biologisch-anatomische Studien über den Hals der Vögel. København: A.F.
Høst & søn; 1929. 122, [4] p.

19. Zweers G, Bout R, Heidweiller J. Motor Organization of the Avian Head-Neck System. In:
Davies MNO, Green PR, editors. Perception and Motor Control in Birds: An Ecological
Approach. 1994: 201-221

20. van der Leeuw AHJ, Bout RG, Zweers GA. Control of the Cranio-Cervical System During
Feeding in Birds. American Zoologist. 2001 Dec 1;41(6):1352–63.

21. Böhmer C, Prevoteau J, Duriez O, Abourachid A. Gulper, ripper and scrapper: anatomy
of the neck in three species of vultures. Journal of Anatomy. 2020;236(4):701–23.

22. Young MW, Dickinson E, Flaim ND, Granatosky MC. Overcoming a 'forbidden
phenotype': the parrot's head supports, propels and powers tripedal locomotion.
Proceedings of the Royal Society B: Biological Sciences. 2022 May
18;289(1975):20220245.

23. Young MW, Webster C, Tanis D, Schurr AF, Hanna CS, Lynch SK, et al. What does
climbing mean exactly? Assessing spatiotemporal gait characteristics of inclined
locomotion in parrots. J Comp Physiol A. 201.1:19-3

24. Snively E, Russell AP. Functional morphology of neck musculature in the
Tyrannosauridae (Dinosauria, Theropoda) as determined via a hierarchical inferential
approach. Zoological Journal of the Linnean Society. 2007 Dec 1;151(4):759–808.

25. Snively E, Russell AP, Powell GL, Theodor JM, Ryan MJ. The role of the neck in the
feeding behaviour of the Tyrannosauridae: inference based on kinematics and muscle
function of extant avians. Journal of Zoology. 2014;292(4):290–303.

26. Snively E, Cotton JR, Ridgely R, Witmer LM. Multibody dynamics model of head and
neck function in Allosaurus (Dinosauria, Theropoda). 2013;

27. Kambic RE, Biewener AA, Pierce SE. Experimental determination of three-dimensional
cervical joint mobility in the avian neck. Frontiers in Zoology. 2017 Jul 24;14(1):37.

28. Böhmer C, Rauhut OWM, Wörheide G. Correlation between Hox code and vertebral
morphology in archosaurs. Proceedings of the Royal Society B: Biological Sciences.
2015 Jul 7;282(1810):20150077.

29. Marek RD. A surrogate forelimb: Evolution, function and development of the avian
cervical spine. Journal of Morphology. 2023;284(10):e21638.

30. Bhullar BAS, Hanson M, Fabbri M, Pritchard A, Bever GS, Hoffman E. How to Make a
Bird Skull: Major Transitions in the Evolution of the Avian Cranium, Paedomorphosis,

and the Beak as a Surrogate Hand. Integrative and Comparative Biology. 2016 Sep
1;56(3):389–403.

31. Marek RD, Falkingham PL, Benson RBJ, Gardiner JD, Maddox TW, Bates KT.
Evolutionary versatility of the avian neck. Proceedings of the Royal Society B: Biological
Sciences. 2021 Mar 3;288(1946):20203150.

32. Marek RD, Felice RN. The neck as a keystone structure in avian macroevolution and
mosaicism. BMC Biology. 2023 Oct 13;21(1):216.

33. Klingenberg CP, Spence JR. Heterochrony and Allometry: Lessons from the Water
Strider Genus Limnoporus. Evolution. 1993;47(6):1834–53.

34. McNamara KJ. Heterochrony: the Evolution of Development. Evo Edu Outreach. 2012
Jun 1;5(2):203–18.

35. Benson RBJ, Campione NE, Carrano MT, Mannion PD, Sullivan C, Upchurch P, et al.
Rates of Dinosaur Body Mass Evolution Indicate 170 Million Years of Sustained
Ecological Innovation on the Avian Stem Lineage. PLOS Biology. 2014 May
6;12(5):e1001853.

36. Dececchi TA, Larsson HCE. Body and limb size dissociation at the origin of birds:
Uncoupling allometric constraints across a macroevolutionary transition. Evolution. 2013
Sep 1;67(9):2741–52.

37. Bhullar BAS, Marugán-Lobón J, Racimo F, Bever GS, Rowe TB, Norell MA, et al. Birds
have paedomorphic dinosaur skulls. Nature. 2012 Jul;487(7406):223–6.

38. de Beer GR. Archaeopteryx and evolution. Adv Sci. 1954;11(12):160.

39. Arnold P, Esteve-Altava B, Fischer MS. Musculoskeletal networks reveal topological
disparity in mammalian neck evolution. BMC Evol Biol. 2017 Dec;17(1):251.

40. Arnold P. Evolution of the Mammalian Neck from Developmental, Morpho-Functional,
and Paleontological Perspectives. J Mammal Evol. 2021 Jun;28(2):173–83.

41. Cardini A, Polly PD. Larger mammals have longer faces because of size-related
constraints on skull form. Nat Commun. 2013 Sep 18;4(1):2458.

42. Cardini A, Polly D, Dawson R, Milne N. Why the Long Face? Kangaroos and Wallabies
Follow the Same 'Rule' of Cranial Evolutionary Allometry (CREA) as Placentals. Evol
Biol. 2015 Jun 1;42(2):169–76.

43. Rayfield EJ. Structural performance of tetanuran theropod skulls, with emphasis on the
Megalosauridae, Spinosauridae and Carcharodontosauridae.

44. Van Valkenburgh B. Trophic diversity in past and present guilds of large predatory
mammals. Paleobiology. 1988;14(2):155–73.

45. Uyeda JC, Harmon LJ. A Novel Bayesian Method for Inferring and Interpreting the
Dynamics of Adaptive Landscapes from Phylogenetic Comparative Data. Systematic
Biology. 2014 Nov 1;63(6):902–18.

46. Rabosky DL, Santini F, Eastman J, Smith SA, Sidlauskas B, Chang J, et al. Rates of
speciation and morphological evolution are correlated across the largest vertebrate
radiation. Nat Commun. 2013 Jun 6;4(1):1958.

47. Rabosky DL. Automatic Detection of Key Innovations, Rate Shifts, and Diversity-
Dependence on Phylogenetic Trees. PLOS ONE. 2014 Feb 26;9(2):e89543.

48. Wagner GP, Zhang J. The pleiotropic structure of the genotype–phenotype map: the
evolvability of complex organisms. Nat Rev Genet. 2011 Mar;12(3):204–13.

49. Wang X, Clarke JA. Phylogeny and forelimb disparity in waterbirds. Evolution. 2014 Oct
1;68(10):2847–60.

50. Nudds RL, Dyke GJ, Rayner JMV. Forelimb proportions and the evolutionary radiation of
Neornithes. Proceedings of the Royal Society of London Series B: Biological Sciences.
2004 Aug 7;271(suppl_5):S324–7.

51. Karoullas C, Nudds RL. The link between avian brachial index, flight capability and the
neornithine evolutionary radiation. J Morphol. 2021 Nov;282(11):1698–707.

52. Pierce SE, Clack JA, Hutchinson JR. Comparative axial morphology in pinnipeds and its
correlation with aquatic locomotory behaviour. Journal of Anatomy. 2011;219(4):502–14.

53. Koob TJ, Long JH Jr. The Vertebrate Body Axis: Evolution and Mechanical Function1.
American Zoologist. 2000 Feb 1;40(1):1–018.

54. Stiller J, Feng S, Chowdhury AA, Rivas-González I, Duchêne DA, Fang Q, et al.
Complexity of avian evolution revealed by family-level genomes. Nature. 2024 Apr 1;1–
3.

55. Yu Y, Zhang C, Xu X. Deep time diversity and the early radiations of birds. Proceedings
of the National Academy of Sciences. 2021 Mar 9;118(10):e2019865118.

56. Castiglione S, Serio C, Mondanaro A, Melchionna M, Raia P. Fast production of large,
time-calibrated, informal supertrees with tree.merger. Palaeontology. 2022;65(1):e12588.

57. Wang M, Zhou Z. Low morphological disparity and decelerated rate of limb size
evolution close to the origin of birds. Nat Ecol Evol. 2023 Aug;7(8):1257–66.

58. Bapst DW. paleotree: an R package for paleontological and phylogenetic analyses of
evolution. Methods in Ecology and Evolution. 2012;3(5):803–7.

59. Cooney CR, Bright JA, Capp EJ, Chira AM, Hughes EC, Moody CJ, et al. Mega-
evolutionary dynamics of the adaptive radiation of birds. Nature. 2017;542(7641):344–7.

60. Jetz W, Thomas GH, Joy JB, Hartmann K, Mooers AO. The global diversity of birds in
space and time. Nature. 2012 Nov;491(7424):444–8.

61. Gelman A, Rubin DB. Inference from Iterative Simulation Using Multiple Sequences.
Statistical Science. 1992;7(4):457–72.

62. Brooks SP, Gelman A. General methods for monitoring convergence of iterative
simulations. Journal of computational and graphical statistics. 1998;7(4):434–55.

63. Smaers JB, Rothman RS, Hudson DR, Balanoff AM, Beatty B, Dechmann DKN, et al.
The evolution of mammalian brain size. Science Advances. 2021 Apr
28;7(18):eabe2101.

64. Ksepka DT, Balanoff AM, Smith NA, Bever GS, Bhullar BAS, Bourdon E, et al. Tempo
and Pattern of Avian Brain Size Evolution. Current Biology. 2020 Jun 8;30(11):2026-
2036.e3.

65. Smaers JB, Rohlf FJ. Testing species' deviation from allometric predictions using the
phylogenetic regression. Evolution. 2016 May 1;70(5):1145–9.

66. Maher AE, Burin G, Cox PG, Maddox TW, Maidment SCR, Cooper N, et al. Body size,
shape and ecology in tetrapods. Nat Commun. 2022 Jul 27;13(1):4340.

67. Brusatte SL, Sakamoto M, Montanari S, Harcourt Smith WEH. The evolution of cranial
form and function in theropod dinosaurs: insights from geometric morphometrics. Journal
of Evolutionary Biology. 2012 Feb 1;25(2):365–77.

68. Marugán-Lobón J, Nebreda SM. Craniocervical morphological integration in birds.
Geobios. 2025.

69. Brusatte SL, Lloyd GT, Wang SC, Norell MA. Gradual Assembly of Avian Body Plan
Culminated in Rapid Rates of Evolution across the Dinosaur-Bird Transition. Current
Biology. 2014 Oct 20;24(20):2386–92.

70. Zanno LE, Makovicky PJ. Herbivorous ecomorphology and specialization patterns in
theropod dinosaur evolution. Proceedings of the National Academy of Sciences. 2011
Jan 4;108(1):232–7.

71. Ma W, Pittman M, Butler RJ, Lautenschlager S. Macroevolutionary trends in theropod
dinosaur feeding mechanics. Current Biology. 2022 Feb 7;32(3):677-686.e3.

72. Middleton KM, Gatesy SM. Theropod forelimb design and evolution. Zoological Journal
of the Linnean Society. 2000 Feb 1;128(2):149–87.

73. Nudds RL, Dyke GJ, Rayner JMV. Avian brachial index and wing kinematics: putting
movement back into bones. Journal of Zoology. 2007;272(2):218–26.

74. Sullivan C, Hone DWE, Xu X, Zhang F. The asymmetry of the carpal joint and the
evolution of wing folding in maniraptoran theropod dinosaurs. Proceedings of the Royal
Society B: Biological Sciences. 2010 Mar 3;277(1690):2027–33.

75. Functional morphology of indrid lumbar vertebrae - Shapiro - 1995 - American Journal of
Physical Anthropology. 98.3:323-342.

76. Buchholtz EA, Bailin HG, Laves SA, Yang JT, Chan MY, Drozd LE. Fixed cervical count
and the origin of the mammalian diaphragm. Evolution & Development. 2012;14(5):399–
411.

77. Kress D, Bokhorst E van, Lentink D. How Lovebirds Maneuver Rapidly Using Super-Fast
Head Saccades and Image Feature Stabilization. PLOS ONE. 2015 Jun
24;10(6):e0129287.

78. Pete AE, Kress D, Dimitrov MA, Lentink D. The role of passive avian head stabilization in
flapping flight. Journal of The Royal Society Interface. 2015 Sep 6;12(110):20150508.

79. Quinn D, Kress D, Chang E, Stein A, Wegrzynski M, Lentink D. How lovebirds maneuver
through lateral gusts with minimal visual information. Proceedings of the National
Academy of Sciences. 2019 Jul 23;116(30):15033–41.

80. Theunissen LM, Troje NF. Head Stabilization in the Pigeon: Role of Vision to Correct for
Translational and Rotational Disturbances. Front Neurosci. 11:551.

81. Egelhaaf M, Böddeker N, Kern R, Kurtz R, Lindemann JP. Spatial vision in insects is
facilitated by shaping the dynamics of visual input through behavioral action. Front
Neural Circuit. 6:108.

82. Long J, McNamara K. Heterochrony: the key to dinosaur evolution. The Dinofest
International (Wolberg, DL, Stump, E & Rosenberg, GD, eds). 1997;113–23.

83. Erickson GM, Makovicky PJ, Currie PJ, Norell MA, Yerby SA, Brochu CA. Gigantism and
comparative life-history parameters of tyrannosaurid dinosaurs. Nature. 2004
Aug;430(7001):772–5.

84. Guenther MF. Influence of Sequence Heterochrony on Hadrosaurid Dinosaur Postcranial
Development. The Anatomical Record. 2009;292(9):1427–41.

85. Canale JI, Novas FE, Salgado L, Coria RA. Cranial ontogenetic variation in Mapusaurus
roseae (Dinosauria: Theropoda) and the probable role of heterochrony in
carcharodontosaurid evolution. Paläontol Z. 2015 Dec 1;89(4):983–93.

86. McKinney ML, McNamara KJ. Heterochrony: the evolution of ontogeny. Springer
Science & Business Media; 2013.

87. Greer JJ, Allan DW, Martin-Caraballo M, Lemke RP. An overview of phrenic nerve and
diaphragm muscle development in the perinatal rat. Journal of Applied Physiology. 1999
Mar;86(3):779–86.

88. Felice RN, Randau M, Goswami A. A fly in a tube: Macroevolutionary expectations for
integrated phenotypes. Evolution. 2018 Dec 1;72(12):2580–94.

89. Evans KM, Larouche O, Watson SJ, Farina S, Habegger ML, Friedman M. Integration
drives rapid phenotypic evolution in flatfishes. Proceedings of the National Academy of
Sciences. 2021 May 4;118(18):e2101330118.

90. Abourachid A, Höfling E. The legs: a key to bird evolutionary success. J Ornithol. 2012
Aug 1;153(1):193–8.

Figures

-Fig 1: Shifts in adaptive optima (yellow triangles) and evolutionary rates (red circles) of neck
morphology across non-avian theropods and avialans (including extant birds). A-C)
Phylogenetic trees of non-avian theropods and avialans depicting the evolutionary rates of
neck vertebral morphology (warmer branch colours denote faster rates of morphological
evolution) for the first (A), middle (B) and last (C) cervical vertebrae. D) Traitgrams depicting
the mode of phenotypic optimum evolution for centrum length (the measurement of which is
demonstrated in the bottom left of the traitgram) evolution of the last cervical vertebrae. E)
Density chart for the phenotypic optima of centrum length for non-avialan theropods (purple)
and Avialae (yellow). F) Allometric scaling relationships of centrum length for non-avialan

theropods (purple) and Avialae (yellow). Silhouettes are taken from PhyloPic.org and
correspond to representative members of Alvarezsauroidea (*Shuvuuia deserti*),
Oviraptorosauria (*Oviraptor philoceratops*, credit: Ivan Iofrida), Emberizoidea (*Emberiza sp.*),
Ornithomimidae (*Struthiomimus altus*), Therizinosauria (*Nothronychus mckinleyi*, credit:
Scott Hartman), Bohaiornithidae (*Bohaiornis guoi)*, Aves (*Ciconia ciconia*, credit: L. Shyamal)
and Neotheropoda (*Ceratosaurus nasicornis*, credit: Scott Hartman).

-Fig 2: Shifts in adaptive optima (yellow triangles) and evolutionary rates (red circles) of
individual forelimb bone lengths across non-avian theropods and avialans (including extant
birds). A-D) Phylogenetic trees of non-avian theropods and avialans depicting the
evolutionary rates for individual forelimb bone length (warmer branch colours denote faster
rates of morphological evolution for the humerus (A), radius (B), ulna (C), and longest
metacarpal/carpometacarpus (D). Forelimb silhouettes of *Archaeopteryx* (adapted from
Middleton & Gatesy 2000) with individual bones highlighted in yellow. Sillhouettes of non-
avian theropods and avialans are taken from PhyloPic.org and correspond to representative
members of Compsognithidae (*Sinosauropteryx prima*, credit: Maija Karala), Ornithomimidae
(*Struthiomimus altus*), Alvarezsauroidea (*Shuvuuia deserti*), Oviraptorosauria (*Oviraptor
philoceratops*, credit: Ivan Iofrida), Anchiornithidae (*Aurornis xui*, credit: Gareth Monger),
Apodiformes (Trochilidae), Therizinosauria (*Nothronychus mckinleyi,* credit: Scott Hartman)

-Fig 3: Shifts in adaptive optima (triangles) and evolutionary rates (red circles) of brachial
index across non-avian theropods and avialans (including extant birds). A) Phylogenetic
trees of non-avian theropods and avialans depicting the evolutionary rates of brachial index
morphology (warmer branch colours denote faster rates of morphological evolution) for
brachial index. Forelimb silhouette depicts the brachial index as the ratio of humerus (yellow)
to radius-ulna (purple) length. B) Traitgram depicting the mode of phenotypic optimum
evolution for brachial index (changes in brachial index are depicted at the extremes of the Y-
axis) evolution of the last cervical vertebrae. C) Allometric scaling relationships of brachial
index across non-avialan theropods and Avialae. D) Density chart for the phenotypic optima
of brachial index across non-avian theropods and avialans. Triangle colours in A and line
colours in B-D denote regimes of non-avian theropods and avialans where a shift in brachial
index adaptive optima has occurred (yellow: Ceratosauria, purple: Neotheropoda, Green:
Maniraptora (sans Alvarezsauroidea, blue: Sphenisciformes). Silhouettes are taken from
PhyloPic.org and correspond to representative members of Ceratosauria (*Ceratosaurus
nasicornis*, Scott Hartman), Maniraptora (*Deinonychus antirrhopus*, credit: Emily
Willoughby), Anchiornithidae (*Aurornis xui*, credit Gareth Monger), Longipteryigidae
(*Iberomesornis romerali*, credit: Durbed/T. Michael Keesey), Sphenisciformes (*Aptenodytes
patagonicus*).

-Fig S1: a) tree with adaptive optima shift regimes for the first cervical vertebrae, F) tree with
adaptive optima shift regimes for the middle cervical vertebrae. B, G) Phylogenetic trees
depicting adaptive optima regimes. C, H) Allometric scaling relationships of centrum length
across the first (C) and middle (H) cervical vertebrae. D, I) Density chart displaying
phenotypic optima of the first (D) and middle (I) cervical vertebrae. E, J) Density chart for
assigned slope values of body mass in the associated Bayesian OU model. Across all
charts, purple denotes non-avialan theropod dinosaurs and yellow denotes members of
Avialae. Sillhouttes are taken from PhyloPic.org and correspond to representative members
of Neotheropoda (*Ceratosaurus nasicornis*, credit: Scott Hartman), Maniraptora
(*Deinonychus antirrhopus*, credit: Emily Willoughby), Aves (*Ciconia ciconia*, credit: L.
Shyamal).

-Fig S2: Adaptive optima regimes (A, C, I, K) and their associated allometric scaling
relationships (B, D, J, L), phenotypic optima density charts (E, G, M, O) and body mass
slope density charts for humerus length (A, B, E, F), radius length (C, D, G, H), ulna length
(I, J, M, N) and longest metacarpal length (K, L, O, P).