A Regulatory Axis for Tonotopic MYO7A Expression in Cochlear Hair Cells

Sihan Li [1], Sujin Jun [6, 7, 9, 10, 12], Ye-Ri Kim [8, 11], Ayathi Gogineni [1], Franklin Lee [1], Chul Hoon Kim [13], Un-Kyung Kim [7, 9, 10, 12], Anthony W. Peng [6], Jung-Bum Shin[1 ,3, 4, 5, #]

Departments of Neuroscience[1], Biochemistry and Molecular Genetics[2], Cell Biology[3], Membrane and Cell Physiology[4], Otolaryngology[5], School of Medicine, University of Virginia, Charlottesville, VA 22903, USA.

[6]Department of Physiology and Biophysics, University of Colorado Anschutz Medical Campus, Aurora, CO.

Department of Biology[7], Immunology[8], Advanced Bio-Resource Research Center[9], School of Life Sciences[10], School of Medicine[11], BK21 plus KNU Creative BioResearch Group[12], Kyungpook National University, Daegu, Republic of Korea.

[13]Department of Pharmacology, BK21 PLUS Project for Medical Science, Brain Research Institute, Yonsei University College of Medicine, 03722, Seoul, Korea.

[#] Corresponding authors: Jung-Bum Shin, email: js2ee@virginia.edu

Classification: Biological Sciences, Neuroscience.

Keywords: Auditory hair cells; transcriptional enhancer; hair cell development; MYO7A; SIX2.

Competing interest statement

The Authors declare no competing interests

1. Significance

Cochlear hair cells rely on the molecular motor MYO7A for mechanosensory function. Our previous study revealed that Myo7a isoforms are differentially expressed in auditory hair cells, however, the mechanisms regulating this isoform-specific expression remain unclear. In this study, we identify a cis-regulatory element, *EnhancerA*, that governs MYO7A expression in a tonotopically graded manner. Furthermore, we demonstrate that the transcription factor SIX2 plays a role in regulation of MYO7A, and is essential for hair cell maintenance. These findings establish a mechanistic link between cochlear position and protein isoform diversity, highlighting how hair cells adapt to frequency-specific mechanical demands. Importantly, the tonotopic regulatory function of *EnhancerA* also offers new avenues for targeted gene therapy in auditory disorders.

2. Abstract

Myo7a, a gene mutated in Usher syndrome and non-syndromic deafness, encodes an unconventional myosin essential for hair cell function. Our previous work revealed that cochlear hair cells express distinct *Myo7a* isoforms with unique spatial and cell type-specific patterns. The canonical isoform (*Myo7a-C*) and a novel isoform (*Myo7a-N*) are co-expressed in outer hair cells (OHCs) but exhibit opposing tonotopic gradients, while inner hair cells (IHCs) primarily express *Myo7a-C*. These isoforms arise from distinct transcriptional start sites, indicating separate regulatory inputs. Here, we identify an intronic cis-regulatory element, *EnhancerA*, essential for tonotopically graded *Myo7a* expression. *EnhancerA* deletion reduces MYO7A protein levels in a tonotopically varied manner, disrupts hair bundle morphogenesis, alters OHC mechanotransduction, and leads to hair cell degeneration and hearing loss. We further identify SIX2, a tonotopically expressed transcription factor that may interact with *EnhancerA* to regulate *Myo7a-N* in OHCs. These findings define a cis-trans regulatory axis critical for isoform-specific *Myo7a* expression and cochlear function.

3. Introduction

Sensory hair cells of the inner ear rely on the unconventional myosin motor protein MYO7A for proper development[1–4], maintenance of hair bundles[5,6], and mechanotransduction[7]. Loss-of-function mutations in the *Myo7a* gene cause both syndromic and non-syndromic forms of hereditary hearing loss. Notably, mutations in *MYO7A* underlie Usher syndrome type 1B, a disorder characterized by congenital deafness, vestibular dysfunction, and progressive vision loss[1,2,8]. In non-syndromic cases, *Myo7a* mutations account for a significant proportion of autosomal recessive deafness[9,10]. Given its central role in auditory function, MYO7A has emerged as a key target for inner

ear gene therapy.

Previous studies have identified several *cis*-regulatory elements near the *Myo7a*

locus[11,12], including a proximal promoter located upstream of the canonical transcriptional

start site[7,11]. This promoter has been incorporated into gene therapy vectors and can drive

robust expression in inner hair cells (IHCs). However, its activity in outer hair cells (OHCs)

is weak or inconsistent, limiting its effectiveness for pan-hair cell targeting[13,14]. Regulatory

sequences derived from non-mammalian species, such as conserved elements in

zebrafish[12], have demonstrated hair cell-specific activity in transgenic models, but their

correspondence to mammalian regulatory mechanisms and isoforms remains poorly
understood.

Our previous work showed that cochlear hair cells express multiple *Myo7a* isoforms,
including the genomically annotated canonical isoform (*Myo7a-C*), the short isoform
(*Myo7a-S*), and more recently, a novel isoform (Myo7a-N) using long-read RNA

sequencing from cochlear cDNA[15]. Hair cells lacking both *Myo7a-C* and *Myo7a-N* exhibit

near complete loss of MYO7A levels at P5 and severe hair bundle disorganization with

profound hearing loss by P20[15], indicating that *Myo7a-S* is minimally expressed and that

Myo7a-C and *Myo7a-N* are the predominant MYO7A isoforms in cochlear hair cells.
Notably, *Myo7a-C* and *Myo7a-N* exhibit distinct cell type-specific and tonotopic
distributions. *Myo7a-C* and *Myo7a-N* are both expressed in OHCs, but exhibit opposing
gradients along the cochlear tonotopic axis: *Myo7a-N* expression increases toward the
cochlear base, while *Myo7a-C* is enriched toward the apex. In contrast, *Myo7a-C* is

predominant in IHCs across all tonotopic regions [10,15].

Myo7a isoforms arise from separate transcriptional start sites[16,17], suggesting
differential promoter usage and distinct trans-regulatory controls. Here, we identify a
previously uncharacterized intronic enhancer, *EnhancerA*, that is essential for
tonotopically graded expression of *Myo7a* isoforms in both IHCs and OHCs. Deletion of
EnhancerA leads to a progressive loss of MYO7A protein toward the cochlear base,
impaired hair bundle morphogenesis, altered mechanotransduction, hair cell degeneration,
and hearing loss. We further identify SIX2, a transcription factor with tonotopically graded
expression, as a potential trans-regulator that may interact with *EnhancerA* to selectively
control *Myo7a-N* expression in OHCs. Loss of SIX2 does not produce immediate or severe
phenotypes, but in aging mice, reduces *Myo7a-N* expression, leads to disorganization of
OHCs, and causes progressive auditory dysfunction. Together, these findings define a
previously unrecognized *cis-trans* regulatory axis that governs spatial and isoform-specific

expression of *Myo7a* in the cochlea and highlight new molecular mechanisms essential for
hair cell function and therapeutic intervention.
101

3. Results

3.1. *EnhancerA* is required for tonotopically graded expression of *Myo7a*

We previously showed that *Myo7a* is expressed in multiple isoforms within the
cochlea, each exhibiting distinct tonotopic expression patterns[16,17]. Importantly, these
isoforms are generated with different transcriptional start sites (**Fig. 1A**), suggesting that
their regulation may involve distinct combinations of *cis-* and *trans*-acting elements.
Previous studies have identified multiple enhancers and promoters of *Myo7a*, yet none

have been shown to support broad and robust expression across all cochlear hair cells[13,14].

We therefore hypothesized that additional regulatory elements may contribute to the
spatially patterned and cell type-specific expression of *Myo7a*.
112

To identify such elements, we reanalyzed publicly available ATAC-seq datasets from

purified cochlear hair cells and supporting cells[18]. This analysis revealed a prominent, hair

cell-specific peak of chromatin accessibility located within the third intron of the *Myo7a*
gene, approximately 400 bp in length (**Fig. 1A,** genomic location provided in the figure
legend). This peak represents the most accessible chromatin region within a 1Mb window
surrounding *Myo7a*, consistent with a potential regulatory function. We designated this
element *EnhancerA*.
120

To assess its functional role *in vivo*, we generated a mouse line harboring a 399 bp
deletion targeting *EnhancerA* (Δ*EnhancerA* mice). Quantitative PCR analysis of cochlear
cDNA from postnatal day 0 (P0) Δ*EnhancerA* mice revealed significantly reduced
expression of all known *Myo7a* isoforms relative to wild-type (WT) controls, including
Myo7a-C and *Myo7a-N*, and *Myo7a-S*. This result indicates that *EnhancerA* is essential
for robust *Myo7a* transcription but not selective to *Myo7a* isoforms (**Fig. 1B**).
127

Strikingly, immunofluorescence analysis of MYO7A protein levels in Δ*EnhancerA* mice
at P7 revealed a pronounced and spatially patterned reduction in both IHCs and OHCs
(**Fig. 1C,** quantified for IHCs and OHCs in **Fig. 1D, E**). In the basal cochlea, MYO7A
protein was nearly undetectable, indicating a severe transcriptional or post-transcriptional
deficit. In contrast, the apical cochlear region retained robust MYO7A expression, though
levels were reduced by approximately 50% compared to WT controls. Notably, reduced
MYO7A levels are observed in both OHCs and IHCs, consistent with our qPCR results
showing that all MYO7A isoforms are affected by *EnhancerA* deletion. MYO7A expression
is broadly comparable across tonotopic regions in WT controls. The progressive
base-to-apex gradient of MYO7A loss observed upon *EnhancerA* deletion suggests that
EnhancerA functions as a key cis-regulatory element that supports robust, spatially biased
MYO7A expression along the cochlear axis. Importantly, this regulatory effect appears to

act without isoform selectivity.

3.2. *EnhancerA* deletion impairs hair bundle development

To evaluate the developmental consequences of *EnhancerA* deletion, we performed
scanning electron microscopy (SEM) on Δ*EnhancerA* cochleae at P7, a stage when hair
bundle maturation is largely complete. In the most basal cochlear region ("hook" region),
where MYO7A loss was the most severe, we observed marked disorganization of hair
bundles in both IHCs and OHCs (**Fig. 2A, B**). Defects included the near-complete
absence of the shortest row of stereocilia, a reduced number of middle and tallest row
stereocilia, and disorganized stereocilia clusters. The clustering phenotype suggests
impaired formation or maintenance of interstereociliary links, a process dependent on
MYO7A function[19–21].

In contrast, hair cells in more apical regions, where residual MYO7A protein remained,
displayed largely normal bundle morphology resembling that of WT controls (**Fig. 2A**).
These findings indicate that *EnhancerA* is required for proper stereocilia development in a
spatially graded manner: based on the observation of impaired hair bundles in the
Δ*EnhancerA* "hook" region, it is likely that the reduced level of MYO7A in this region is
insufficient to support proper hair bundle development or organization. In contrast, in the
middle and apical regions, MYO7A levels are reduced to a lesser extent, allowing hair
bundles to maintain a WT-like morphology.

3.3. Hair cell loss and hearing loss in Δ*EnhancerA* mice

To determine whether these structural defects lead to auditory dysfunction, we
measured auditory brainstem responses (ABRs) and distortion product otoacoustic
emissions (DPOAEs). At 3 weeks of age, Δ*EnhancerA* mice exhibited mild to moderate
hearing loss, with significant threshold shifts at mid-to-high frequencies (**Fig. 2C**),
consistent with the basal-to-apical gradient of MYO7A depletion. Lower frequencies also
showed a statistically insignificant trend toward elevated ABR thresholds, likely reflecting
that the residual MYO7A levels are insufficient to fully support optimal hair cell function.
Distortion product otoacoustic emissions (DPOAEs) were largely unaffected at this stage.
However, by 8 weeks, Δ*EnhancerA* mice exhibited profound hearing loss across all
frequencies, accompanied by the near-complete loss of DPOAEs (**Fig. 2C**).

To further explore the cellular basis of this rapid progression, we performed
immunofluorescence analysis of MYO7A and hair cell morphology at 8 weeks. By this
stage, MYO7A expression was nearly absent throughout the cochlea in Δ*EnhancerA* mice
(**Fig. 2D**), and was accompanied by extensive degeneration of hair cells in the basal turn,
with partial loss observed in the middle and apical turn (**Fig. 2D**). These results suggest
that *EnhancerA* is not only essential for the early developmental expression of *Myo7a*, but
also becomes increasingly critical for maintaining *Myo7a* expression and hair cell survival
as the cochlea matures.

183
Together, these findings demonstrate that *EnhancerA* plays a critical role in regulating
MYO7A expression in a tonotopically graded manner. Its deletion disrupts hair bundle
morphogenesis, compromises hair cell maintenance, and impairs long-term auditory
function, initially affecting high-frequency regions and progressively extending toward
lower-frequency areas with age. Notably, the highest ABR frequency tested was 32 kHz,
which corresponds to the mid-basal region of the cochlea. Therefore, functional deficits in
the more basal, higher-frequency region may be underestimated in our measurements.
191

3.4. Deletion of *EnhancerA* causes reduced resting open probability of MET channels
in a tonotopically graded manner

Our previous work demonstrated that isoform-specific deletion of *Myo7a-C* results in
approximately 80% reduction of MYO7A protein in IHCs, leading to decreased resting
open probability and delayed activation kinetics of MET currents[17]. These findings
established a role for MYO7A-C in maintaining resting tension on the MET complex in
IHCs. However, the contribution of MYO7A to MET function in OHCs was not
characterized in our previous study. The *ΔEnhancerA* mouse model presents an
opportunity to address this gap. Unlike the isoform-specific KO, *ΔEnhancerA* mice exhibit
reduced expression of all known *Myo7a* isoforms in both IHCs and OHCs. Furthermore,
MYO7A loss occurs in a tonotopically graded manner, allowing us to correlate spatial
differences in protein levels with MET channel properties, independent of isoform bias.
204

To assess MET function, we recorded transduction currents from OHCs in the basal
and apical cochlea of *ΔEnhancerA* homozygous mice ($EA^{-/-}$) at postnatal day 4-5 (P4-5),
using fluid jet stimulation and high-speed imaging to monitor bundle deflections.
Considering that the "hook" region of *ΔEnhancerA* mice displays disorganized hair
bundles, we first confirmed that stereocilia in the basal region preceding the "hook" region
exhibit WT-like development, thereby avoiding potential confounds arising from disrupted
hair bundle morphology (**Supplementary Fig. 1A**). All recordings were restricted to
regions with intact stereocilia morphology from the mid-basal to apical sections of the
cochlea.
214

Apical OHCs, where MYO7A expression is reduced by only 50%, exhibited no
significant differences in resting open probability, peak current, or adaptation compared to
heterozygous controls ($EA^{+/-}$) (**Fig. 3A, B, E, F, Supplementary Fig. 2B**). However, in
basal OHCs, where MYO7A levels are strongly reduced but bundles are still intact,
sinusoidal stimulation revealed a significant decrease in the resting open probability of
MET channels, while peak current amplitudes remained unchanged (**Fig. 3C, D**). Similarly,
step-like stimuli confirmed a reduction in resting open probability, with no detectable
changes in current amplitude or adaptation kinetics at P4-P5 (**Fig. 3E, G; Supplementary**
Fig. 2B). Collectively, these results confirm our previous study that MYO7A levels are
critical for maintaining resting tension on the MET apparatus[17].
225

3.5. SIX2 is tonotopically expressed in the cochlea

While *EnhancerA* confers a tonotopically graded expression pattern on *Myo7a* expression, its deletion disrupts both *Myo7a-C* and *Myo7a-N* isoforms, indicating that additional mechanisms are involved in mediating isoform-specific regulation. To identify potential trans-regulatory factors, we queried the Unbind2021[22] and ReMap2022[23] ChIP-seq databases and discovered a prominent SIX2 binding site centrally located within *EnhancerA*[24,25] (**Fig. 4A, B**). SIX2 is a homeodomain transcription factor known for its roles in kidney[26–29], craniofacial skeleton development[30]. Although haploinsufficiency of SIX2 has been associated with mild high-frequency hearing loss in humans[31], its function in auditory and vestibular hair cells remains largely uncharacterized.

First, we investigated the expression pattern of SIX2 in the mouse cochlea. Previously, it has been reported that a widely used commercially available SIX2 antibody also interacts with SIX1[24]. To circumvent this, we utilized the *Six2::cre-egfp* mouse line and visualized EGFP expression as a proxy for SIX2 activity. At P5, we observed strong SIX2 driven EGFP signal in OHCs and a subset of supporting cells within the greater epithelial ridge (GER), with comparatively weaker expression in IHCs (**Fig. 4C–E**). Notably, *Six2::cre* activity displayed a tonotopic gradient, highest in the basal cochlear turn and tapering toward the apex, mirroring the distribution of *Myo7a-N* isoform expression. This parallel pattern suggests a potential regulatory role for SIX2 in modulating *Myo7a-N* expression in the cochlea.

To confirm endogenous SIX2 expression and define its subcellular localization, we generated a *Six2-HA* knock-in (KI) mouse line, tagging the C-terminus of the endogenous gene with an HA epitope. The HA tag did not grossly affect SIX2 function, since *Six2-HA* homozygous mice developed normally, while SIX2 deletion is known to cause perinatal lethality[26]. Immunostaining of these cochleae revealed a similar tonotopic pattern of SIX2-HA signal as observed in *Six2::cre-egfp* mice, with strong nuclear expression in OHCs and GER cells in the basal region, decreasing toward the apex (**Fig. 4F–H**). Consistent with *Six2::cre* activity, SIX2-HA expression was low in IHCs, supporting the idea that SIX2 acts as an OHC-specific transcription factor within the hair cell population. The nuclear enrichment of the HA signal further suggests that SIX2 is transcriptionally active in these cells.

To further characterize the expression pattern of SIX2 at different developmental stages in the cochlea, we extended our SIX2-HA immunostaining analyses to P1 and P7. At P1, SIX2 expression was detected in OHCs of the basal and middle turns, while minimal signal was observed in the apical turn (**Fig. 4I, J**). By P7, SIX2 expression in

264 OHCs increased across all cochlear regions and became relatively uniform (**Fig. 4I, J**),
265 indicating a loss of the initial tonotopic gradient. In contrast, SIX2 expression remained low
266 in IHCs at both P1 and P7. These results suggest that SIX2 expression initiates at the
267 basal end of the cochlea around P1 and subsequently expands to all cochlear regions by
268 P7.
269
270 To determine whether SIX2 transcripts are expressed at embryonic and adult stages,
271 we performed qPCR to assess SIX2 transcript levels at E17, P0, and P20 (**Fig. 4K**). SIX2
272 transcript levels showed low expression between E17 and P0. In contrast, SIX2
273 expression was significantly increased at P20, when hair cells are fully matured,
274 suggesting a role for SIX2 in postnatal and adult hair cells.
275
276 Taken together, these findings identify SIX2 as a transcription factor that is
277 preferentially expressed in OHCs relative to IHCs and whose expression is
278 developmentally regulated, initiating around P0 with a transient tonotopic pattern that
279 becomes uniform across the cochlea by P7.
280

3.6. Loss of SIX2 does not disrupt hair cell development but impairs hair cell
maintenance
Based on the expression pattern of SIX2 in the cochlea, particularly its enrichment in
OHCs, we hypothesized that SIX2 may play a critical role in OHC development or
maintenance. We first tested whether SIX2 is required for OHC development. Because
global deletion of SIX2 results in early postnatal lethality[26], we generated a hair
cell-specific SIX2 conditional knockout by crossing *Gfi1-cre* mice with *Six2$^{flox/flox}$* mice
(*Gfi1$^{cre/-}$*; *Six2$^{flox/flox}$*, hereafter referred to as, *Gfi1-Six2-cKO*). In this model, SIX2 is
specifically deleted in hair cells during early embryonic stages, likely preceding the onset
of endogenous SIX2 expression.
291
We first examined hair cell development at P7 using immunofluorescence microscopy.
In the basal turn of the cochlea, no loss of IHCs or OHCs was observed in Gfi1-Six2-cKO
mice, and OHCs maintained the normal three-row organization (**Fig. 5A, B**). Notably,
MYO7A immunofluorescence signals in both basal IHCs and OHCs were comparable to
those in control mice (**Fig. 5B**). Given that SIX2 is highly enriched in OHCs, we further
examined hair bundle morphology in basal OHCs using SEM. This analysis revealed no
disorganization of hair bundles, stereocilia degeneration, or other overt morphological
abnormalities (**Fig. 5C**). Based on these results, we conclude that SIX2 is not required for
early hair cell development, consistent with the postnatal-onset of its expression in OHCs.
301
Based on the expression of SIX2 that starts around P0 and progressively increases
into adult stages, we hypothesized that SIX2 contributes to hair cell maintenance. Since
Gfi1-cre causes hearing loss in adult mice due to haploinsufficiency[32], we chose
Myo15a-cre to investigate the role of SIX2 in mature hair cells (*Myo15a$^{cre/+}$*; *Six2$^{flox/flox}$* mice

hereafter referred to as *Myo15a-Six2-cKO*). Because *Myo15a-Cre* expression begins

during the early postnatal period [33], a few days after the onset of SIX2 expression, this

strategy enables selective deletion of SIX2 in hair cells after initial hair cell differentiation

and development. At 8 weeks of age, *Myo15a-Six2-cKO* mice showed no apparent loss of

hair cells (**Fig. 5D**). However, by 16 weeks, we observed progressive degeneration of

OHCs, starting in the basal turn of the cochlea (**Fig. 5E-F**).

Because no OHC loss was observed in *Myo15a-Six2-cKO* mice at 8 weeks of age,

whereas near-complete OHC loss was evident in the basal turn by 16 weeks, we sought to

identify an intermediate stage to examine OHC hair bundle morphology at the basal turn

and further assess the effects of SIX2 deletion on hair bundles. We first confirmed that

10-week, *Myo15a-Six2-cKO* mice exhibit increased OHC loss by quantifying the length of

the organ of Corti with complete OHC loss, which was significantly longer compared with

controls (~800 µm in cKO vs. ~250 µm in control) (**Fig. 5G-H**). In addition, surviving OHCs

located approximately 800 µm from the basal end displayed degeneration of the third row

of stereocilia (**Fig. 5I**).

OHC degeneration was accompanied by progressive hearing loss, with threshold

shifts first emerging at mid-to-high frequencies by 8 weeks and expanding to all

frequencies by 16 weeks (**Fig. 6J–K**). These findings suggest that SIX2 is required for the

maintenance of hair cell integrity post-development, and its deletion leads to hair cell loss,

hair bundle degeneration and hearing impairment.

3.7. SIX2 regulates expression of *Myo7a*

Lastly, given that SIX2 is highly enriched in OHCs and exhibits a tonotopic expression

pattern during early postnatal stages that correlates with *Myo7a-N*, we further tested the

hypothesis that SIX2 regulates the expression of MYO7A protein and *Myo7a* transcripts.

At 10 weeks of age, we observed a significant reduction of MYO7A levels in surviving

OHCs in the basal turn of *Myo15a-Six2-cKO* mice. In addition to the basal turn, OHCs in

the middle turn also showed a significant decrease in MYO7A signal, whereas the apical

turn was not affected (**Fig. 6A, B**). Notably, we also observed an unexpectedly low

expression level of MYO7A in IHCs. However, this low expression was also found in

regular WT controls, indicating that the decreased MYO7A signal in IHCs is not caused by

the *Six2$^{fl/fl}$* genotype but instead reflects a normal effect of aging (**Fig. 6A**). Overall, this

tonotopically restricted reduction of MYO7A correlates with the high-frequency hearing

loss observed in *Myo15a-Six2-cKO* mice, suggesting that SIX2 regulates MYO7A

expression preferentially in the high-frequency regions of the cochlea at 10 weeks of age.

We next analyzed *Myo7a* isoform expression using qPCR. We chose to perform this

at postnatal day 20 (P20), reasoning that transcript changes will precede protein loss, and

any transcript changes at this stage likely reflect regulatory effects rather than cell loss.

Notably, expression of total *Myo7a* and the *Myo7a-N* isoform was significantly reduced in

SIX2-deficient mice, whereas the *Myo7a-C* isoform was not significantly affected (**Fig. 6C**).

These results suggest that SIX2 regulates *Myo7a* expression and preferentially influences
the expression of MYO7A-N.
351
352 Together, these data reveal that SIX2 plays a crucial role in maintaining gene
353 expression in mature hair cells, including the regulation of specific *Myo7a* isoforms, and
354 that its loss compromises hair cell survival and auditory function.
355

4. Discussion

***EnhancerA* orchestrates tonotopic *Myo7a* expression in cochlear hair cells**
Cochlear hair cells express two *Myo7a* isoforms, *Myo7a-C* and *Myo7a-N*, with
complementary tonotopic and cell type-specific expression[16,17]. While IHCs
predominantly express *Myo7a-C*, OHCs co-express both isoforms in opposing spatial
gradients. Furthermore, these two isoforms exhibit distinct ATPase activity, suggesting
their roles in regulating MET properties. Thus, understanding how this expression
diversity is regulated is expected to be important for decoding their distinct functional
roles.
365
Here, we identify *EnhancerA*, a conserved intronic *cis*-regulatory element, as
essential for maintaining robust *Myo7a* expression in both IHCs and OHCs. Deletion of
EnhancerA during early postnatal development causes a progressive, tonotopically
graded reduction in MYO7A transcript and protein level, culminating in near-complete
loss by 8 weeks. This suggests that *EnhancerA* activity becomes increasingly critical as
chromatin landscapes mature.
372
Importantly, *EnhancerA* deletion impacts both isoforms, indicating that it serves as a
shared enhancer of isoform-selective promoters. The widely used canonical promoter
upstream of *Myo7a-C* is highly active in IHCs but weak in OHCs[7,13,14], suggesting that
Myo7a-N is driven by an alternative promoter. A previous study in zebrafish identified
two regulatory 'blocks' upstream of the *Myo7a* gene that drive hair cell–specific
expression of reporter genes[12]. Our recent findings suggest that one of these blocks
corresponds to the first exon of a previously uncharacterized *Myo7a* isoform we named
Myo7a-N[15], while the other is located just upstream, consistent with the location of a
putative *Myo7a-N* promoter. In the future, it will be important to investigate how
EnhancerA interacts with these isoform-specific promoters to regulate MYO7A
expression in a cell type- and isoform-specific manner.
384
MYO7A tunes MET gating
It has been proposed that MYO7A acts as a motor protein that provides tension to the
MET complex at UTLDs[34–36]. However, due to the disruption of hair bundle development in

MYO7A depletion models, it has been challenging to isolate its specific role in MET function.
Previously, we employed the *Myo7a-ΔC* model to examine the effects of severe MYO7A

depletion specifically in IHCs[7]. In this model, reduction of MYO7A at UTLDs led to

decreased resting P_o and slowed onset of MET currents, supporting the hypothesis that
MYO7A contributes to tensioning the MET complex.
393
In the current study, the *ΔEnhancerA* model serves as a tool to test MYO7A function,
as it reduces expression of all MYO7A isoforms in both IHCs and OHCs. Notably, the
tonotopic gradient of MYO7A depletion in this model enables investigation of MET
function across a range of MYO7A expression levels. At P4-5, MYO7A reduction is most
severe in basal OHCs, where we observed a significant decrease in MET resting P_o,
while peak current amplitude and adaptation kinetics remained unaffected. In contrast,
apical OHCs, which exhibited only mild MYO7A depletion, showed no measurable
changes in MET properties. Together, these results reinforce our previous findings that

MYO7A depletion leads to a reduction in tip-link tension[7].

403
Notably, a recent study reported that MYO7A loss in mature hair cells (P20–P30)
led to reduced bundle stiffness and diminished peak MET currents, without affecting

resting Po[5]. The contrast with our findings at P4-5 may be due to developmental

stage-specific functions of MYO7A. During early postnatal stages, MYO7A may play a
more active role in generating tip-link tension and supporting MET complex assembly,
whereas in mature cells, its function may shift toward structural stabilization. Collectively,
these results underscore the dynamic, isoform- and stage-dependent contributions of
MYO7A to hair cell MET that warrants further research to gain a more complete picture.
412
Other regulatory mechanisms that drive *Myo7a* expression
We propose that *EnhancerA* functions as a shared scaffold for transcriptional
activation in both IHCs and OHCs, with isoform specificity determined by the local
presence or absence of trans-acting factors. Using publicly available ChIP-seq datasets,

we identified a SIX2 binding site within *EnhancerA*[24]. Our qPCR and

immunofluorescence analyses demonstrate that conditional deletion of SIX2 leads to a
reduction in MYO7A expression, with the MYO7A-N isoform being more strongly
affected than other isoforms. These results suggest a potential regulatory role for SIX2
in isoform-specific control of MYO7A expression: in OHCs, particularly in basal regions
where SIX2 expression is highest, SIX2 may cooperate with *EnhancerA* to drive
Myo7a-N expression via a potential *Myo7a-N* promoter, thereby establishing an
increase in *Myo7a-N* levels toward the cochlear base. In contrast, in IHCs, which
express little or no SIX2, *EnhancerA* activity may preferentially be coupled with the
canonical promoter of *Myo7a-C* via other transcription factors. This combinatorial
mechanism, where a broadly active enhancer integrates input from spatially restricted

transcription factors, offers a potential explanation for the observed isoform gradients.
429
While our data implicate SIX2 in regulating MYO7A expression, its precise
contribution remains unclear, as we did not test direct binding of SIX2 to *EnhancerA*.
Meanwhile, the residual MYO7A expression observed in *ΔEnhancerA* mice suggests
the existence of additional regulatory mechanisms governing *Myo7a* expression.
Notably, a recent parallel study independently identified the *EnhancerA* region (referred
to as E3 in their study), along with several additional *cis*-regulatory elements within the
Myo7a locus[37]. In their study, E3 drives reporter gene expression only in combination
with other *Myo7a* enhancers, suggesting that *Myo7a* expression is regulated by a more
complex cooperative *cis/trans*-regulatory element network.
439
SIX2 is dispensable for hair cell development but required for maintenance
SIX family transcription factors are known to be critical for cell proliferation and
differentiation, which is important for kidney development[27–29,38]. SIX1 and SIX2 share
high similarity in amino-acid sequence, structural domains and DNA-binding motif. Both
have been implicated in auditory system development, with SIX1 being extensively
characterized for its essential role in early inner ear development[39–43]. A recent human
genetic study associates SIX2 haploinsufficiency with mid-high frequency hearing loss[31],
however, the molecular function of SIX2 in the cochlea has been less defined.
448
In this study, we demonstrated that SIX2 is not required for hair cell development.
However, SIX2 contributes to the maintenance of mature hair cells, as hair cell-specific
deletion using the *Myo15-cre* mouse leads to progressive loss of OHCs beginning in the
basal region of the cochlea by 10 weeks of age, accompanied by hair bundle
degeneration and high-frequency hearing loss. Deletion of SIX2 reduces overall Myo7a
expression, with the Myo7a-N isoform showing a more pronounced reduction, while
Myo7a-C exhibits only a modest, statistically insignificant decrease. Notably, it is an
intriguing observation that postnatal deletion of SIX2 results in a relatively mild
phenotype, despite the strong enrichment of SIX2 expression in OHCs, suggesting
redundancy and compensation by other transcription factors. As a future direction, it will
be important to identify downstream targets of SIX2 using RNA sequencing. In addition,
because SIX2 is also expressed in the GER and other supporting cell types, it will be
informative to delete SIX2 using an inner ear sensory epithelium-specific Cre driver and
perform single-cell RNA sequencing to characterize its downstream targets in non-hair
cell populations.
464
As mentioned above, SIX2 likely operates within a wider transcriptional network
involving other regulators such as ATOH1[44], POU4F3[45], and GFI1[46]. While other

transcription factors initiate hair cell differentiation, SIX2 may refine OHC identity by
regulating genes that are required for hair cell maturation and maintenance. Meanwhile,
EYA1/2 are known transcription co-factors that interact with SIX2 for gene expression,

and EYA2 showed tonotopic hair-cell specific expression[47]. Interactions among these

transcription factors are likely essential for shaping MYO7A isoform diversity and for
establishing the tonotopic architecture of cochlear hair cells.
473
Implications for isoform-specific gene therapy
Our findings have potential translational implications, given the compact size of
EnhancerA and its unique spatiotemporal regulatory pattern. Meanwhile, *EnhancerA*
and SIX2-responsive elements could be leveraged to design synthetic promoters that
enable targeted transgene expression in hair cells, or even regionally along the cochlear
axis. This spatiotemporal control would be particularly valuable for treating late-onset,
genetic forms of deafness, in which precise regulation of expression timing and location
is critical for therapeutic efficacy.
482
483   In summary, this study identifies *EnhancerA* and SIX2 as regulators of MYO7A
484 isoform expression in the cochlea, linking transcriptional control to regional and cell
485 type-specific functional specialization. These findings deepen our understanding of
486 cochlear gene regulation and lay the groundwork for targeted, isoform-aware therapies
487 for sensorineural hearing loss.
488

5. Figures

Figure 1: EnhancerA is required for tonotopically graded expression of Myo7a

A. Schematic indicating the transcriptional start sites of *Myo7a* isoforms. ATAC-seq tracks are
derived from a publicly available dataset. *EnhancerA* is an around 400 bp open chromatin
region that locates in the third intron of *Myo7a-N*, which is indicated with red bracket
(GRCm38/mm10: chr7: 98105422-98105821). This region is specifically knocked out in
ΔEnhancerA mice without affecting exons. Three major isoforms of Myo7a with distinct
transcriptional start sites are demonstrated in the "*Myo7a*-isoform by 5' RACE". Basewise
conservation shows multiple alignments of 35 vertebrate species as measurements of
evolutionary conservation. **B**. Normalized expression level of *Myo7a* isoforms measured by
qPCR on P0 cochlear cDNA. *Myo7a* expression level is normalized to *Gapdh* and then
log-transformed. (WT vs. *ΔEnhancerA* mice t-test p-values of different isoforms: *pan-Myo7a*:
1.30e-4, *Myo7a-C*=2.60e-3, *Myo7a-N*=1.32e-07, *Myo7a-S*=5.59e-05. Number of Cochlea:
WT=8, *ΔEnhancerA*=8. Pipetting repetition=5 per group). Boxplots show medians, 25th, and
75th percentiles as box limits and minima and maxima as whiskers. **C**. Immunofluorescence
images of whole mount *ΔEnhancerA* cochlea at P7. Compared to WT cochlea, *ΔEnhancerA*
mice showed tonotopic reduction of MYO7A immunoreactivity, with a stronger reduction in the
basal region and less reduction at the apical region. (Scale bar: left: 100 µm, right 10 µm). **D,**
E. Quantification of MYO7A immunoreactivity in IHCs and OHC at P7. MYO7A signals were
normalized to MYO6. (Normalization method: MYO7A signal in the cell body is divided by
F-actin signal in the cell body. T-test on the average normalized MYO7A immunofluorescence
signal of each cochlea. Number of cochlea: WT: 4; *ΔEnhancerA*: 4. T-test p-value between
WT and *ΔEnhancerA*: IHC: base: 5.63e-3; middle: 1.10e-3; apex: 1.43e-2. OHC: base:
1.77e-3; middle: 8.17e-3; apex: 2.89e-2).

Figure 2: **Hair cell degeneration and hearing loss of *ΔEnhancerA* mice.**

A. Scanning electron microscopy (SEM) images of WT and *ΔEnhancerA* cochleae at P7.
ΔEnhancerA hair cells display disorganized hair bundles near the base of the cochlea. (Scale
bar: 10 μm). **B**. Higher-magnification SEM images of WT and *ΔEnhancerA* cochleae at P7
reveal detailed structural abnormalities in *ΔEnhancerA* hair bundles. (Scale bar: 1 μm). **C**.
ABR thresholds and DPOAE response of WT and *ΔEnhancerA* at 3 weeks and 8 weeks of
age. *ΔEnhancerA* mice exhibit mild hearing loss at 3 weeks, which progresses to profound
hearing loss by 8 weeks. (ABR ANOVA p-value: 3-week=1.25e-08, 8-week<2e-16. Number of
animals: WT: 3-week=8, 8-week=9; *ΔEnhancerA*: 3-week=8, 8-week=5. DPOAE p-value:
3-week=0.0060, 8-week=1.12e-11. Number of animals: WT: 3-week=8, 8-week=10;
ΔEnhancerA: 3-week=7, 8-week=5). Line-plots show means of the ABR or DPOAE response.
Standard error is presented as error bars. **D**. Immunofluorescence images of WT and
ΔEnhancerA mice at 8 weeks. MYO7A expression is nearly absent across all cochlear
regions in *ΔEnhancerA* mice, accompanied by extensive hair cell loss. (Scale bar: 10 μm.
Asterisk labels the remaining OHCs at basal regions of *ΔEnhancerA* mice. Arrowheads label
the IHCs loss at basal area. White dashed box highlights the OHC loss in middle and apical
turn of cochlea).

Figure 3: **Reduction of resting open probability of *ΔEnhancerA* OHC.**

A. Evaluation of the MET current of apical OHCs from $EA^{+/-}$ (black) and $EA^{-/-}$ (pink) at P4-P5.
MET currents are evoked by sinusoidal stimulation. **B**. Summary plots of the peak current and
P_o observed from apical OHCs. Number of cells (animals): $EA^{+/-}$ 8 (8), $EA^{-/-}$ 7 (7). **C**.
Evaluation of the MET current of basal OHCs from $EA^{+/-}$ and $EA^{-/-}$ at P4-P5. MET currents are
evoked by sinusoidal stimulation. **D**. Summary plots of the peak current and P_o observed from
basal OHCs. ** $p \leq 0.01$. Number of cells (animals): $EA^{+/-}$ 13 (13), Basal $EA^{-/-}$ 10 (10). **E**.
Comparison of MET currents in apical and basal turns of $EA^{+/-}$ and $EA^{-/-}$ mice measured using
step-like force stimuli. Hair bundle displacement traces are shown at the top, and evoked
peak MET currents are shown at the bottom. **F, G.** Summary plots of the peak current and
resting open probability for apical (**F**) and basal (**G**) turn outer hair cells. * $p \leq 0.05$. Number of
cells (animals): Apical $EA^{+/-}$ = 8 (8), Basal $EA^{+/-}$ = 13 (13), Apical $EA^{-/-}$ = 7 (7), Basal $EA^{-/-}$ =
10(10). (For panel B, D, F and G, whiskers of the plot represent mean ± SD).

Figure 4: **Tonotopic expression of SIX2 in cochlea.**

A. The diagram demonstrating the SIX2 binding site in the *EnhancerA* tested by published ChIP-seq data.   **B**. Human SIX2 binding motif reported by JASPAR database. The DNA sequence of SIX2 binding site in the *EnhancerA* is provided below the binding motif. **C**. Whole mount immunofluorescence image of *Six2::cre-egfp* cochlea at P5. GFP is tonotopically expressed in the cochlea, with a higher expression level at base and decrease toward apex. (Scale bar: 100 µm). **D**. Immunofluorescence image of *Six2::cre-egfp* cochlea at base, middle and apex region. GFP was observed in the OHCs and GER cells. (Scale bar: 10 µm. "I": IHCs; "O": OHCs; "GER": greater epithelial ridge cells). **E**. immunofluorescence image of *Six2::cre-egfp* hair cell regions at base, middle and apex region. (Scale bar: 10 µm). **F**. Whole mount immunofluorescence image of *Six2-HA* cochlea mice at P7. Similar with *Six2::cre-egfp* cochlea, SIX2 is also observed to be expressed tonotopically. (Scale bar: 100 µm). **G**. immunofluorescence image of *Six2-HA* cochlea at base, middle and apex region. HA-tag signals were observed in the OHCs and GER cells. Oncomodulin (OCM) is stained as OHC marker (Scale bar: 10 µm). **H**. immunofluorescence image of *Six2-HA* hair cell regions at base, middle and apex region. (Scale bar: 10 µm). **I**. Representative SIX2-HA immunostaining images at P0 and P7. Scale bar = 10 µm. **J**. Quantification of SIX2-HA fluorescence signal at P0 and P7. At P0, SIX2 is tonotopically expressed in OHCs, with the highest levels in the basal turn of the cochlea. SIX2 expression is significantly increased across all cochlear regions at P7. In contrast, SIX2 expression in IHCs remains low at both P0 and P7. (Normalization method: SIX2-HA signals are collected in the cell nuclei and then normalized to DAPI. Number of cochleae: P0 = 8; P7 = 6. T-test p-values: P0: IHC: base vs. middle = 0.060; middle vs. apex = 0.62; OHC: base vs. middle = 5.00e-3; middle vs. apex = 5.58e-6. P7: IHC: base vs. middle = 0.36; middle vs. apex = 0.45; OHC: base vs. middle = 0.28; middle vs. apex = 0.033. **K**. Transcript levels of Six2 measured at E17, P0, and P20. (Normalization method: Cq value of SIX2 is first normalized to Cq value of Gapdh and then log-transformed. p-values: E17 vs. P0 = 0.66; P0 vs. P20 = 9.75e-3).

Figure 5: **SIX2 is important for hair cell maintenance.**

A. Immunofluorescence images of $Six2^{fl/fl}$ mice and $Gfi^{cre/+}; Six2^{fl/fl}$ mice at P7. (Scale bar: 10

μm). **B**. Quantification of MYO7A level in the cell body of basal turn. No significant MYO7A

reduction was observed (Normalization method: MYO7A signal in the cell body is divided by

F-actin signal in the cell body. Number of animals: $Six2^{fl/fl}$: 4; $Gfi^{cre/+}; Six2^{fl/fl}$: 4. T-test p-value

of $Six2^{fl/fl}$ vs. $Gfi^{cre/+}; Six2^{fl/fl}$: IHC: 0.66, OHC: 0.87). **C**. SEM images of $Six2^{fl/fl}$ and $Gfi^{cre/+};$

$Six2^{fl/fl}$ OHCs at basal turn at P7. $Gfi^{cre/+}; Six2^{fl/fl}$ mice do not show an obvious difference

between control OHCs. (Scale bar: 1 μm). **D**. Representative image of 8-week control

($Myo15a^{cre/+}$) and $Myo15a^{cre/+}; Six2^{fl/fl}$ cochlea at middle and basal turns. (Scale bar: 10 μm). **E**.

Representative image of 16-week control and $Myo15a^{cre/+}; Six2^{fl/fl}$ cochlea at middle and

basal turns. $Myo15a^{cre/+}$ induced conditional deletion of $Six2$ results in OHC degeneration

starting from basal turn. (Scale bar: 10 μm). **F**. Quantification of OHCs of control and

$Myo15a^{cre/+}; Six2^{fl/fl}$ cochlea at middle and basal turns at 16-week. (Number of animals:

$Myo15a^{cre/+}$: 3, $Myo15a^{cre/+}; Six2^{fl/fl}$ = 3. T-test p-value: base = 1.85e-4; middle = 0.39.). **G**.

SEM images of $Six2^{fl/fl}$ and $Myo15a^{cre/+}; Six2^{fl/fl}$ cochleae at 10-week. Images were taken at

approximately 800 μm from the basal end of the cochlea. (Scale bar: 10 μm). **H**.

Quantification of the length of the cochlear region exhibiting complete OHC loss, measured

from the beginning of basal end. (Number of animals: $Six2^{fl/fl}$: 5; $Myo15a^{cre/+}; Six2^{fl/fl}$: 5.

Average length: $Six2^{fl/fl}$: 240.02 μm; $Myo15a^{cre/+}; Six2^{fl/fl}$: 791.80 μm. T-test p-value: 2.41e-05).

I. SEM images of $Six2^{fl/fl}$ and $Myo15a^{cre/+}; Six2^{fl/fl}$ OHCs in the basal turn at 10 weeks of age.

The third row of stereocilia is highlighted in light blue. Degenerating stereocilia is marked by

yellow arrows. (Scale bar: 1 μm). **J**. ABR test of $Myo15a^{cre/+}; Six2^{fl/fl}$ mice at 8 weeks. Mild

ABR threshold shift was observed in the high frequencies tested. (ANOVA p-value: 3.23e-06.

Number of animals: $Six2^{fl/fl}$ = 4, $Myo15a^{cre/+}; Six2^{fl/fl}$ = 10). **K**. ABR test of $Myo15a^{cre/+}; Six2^{fl/fl}$

mice at 16 weeks. ABR threshold shift was observed in all of frequencies tested. (ANOVA

p-value: 7.85e-15. Number of animals: $Six2^{fl/fl}$ = 6, $Myo15a^{cre/+}; Six2^{fl/fl}$ = 10).

Figure 6: SIX2 regulates MYO7A expression.

A. Representative OHC image of 10-week WT (C57BL/6), $Myo15a^{cre/+}$; $Six2^{fl/fl}$ and $Six2^{fl/fl}$ cochlea at middle and basal turns. (Scale bar: 10 µm). **B**. Quantification of MYO7A level in the cell body of OHCs at 10-week. MYO7A is significantly reduced in basal and middle turn of the cochlea (Normalization method: MYO7A signal in the cell body is divided by F-actin signal in the cell body. Number of animals: $Six2^{fl/fl}$: 5; $Myo15a^{cre/+}$; $Six2^{fl/fl}$: 5. T-test p-value of $Six2^{fl/fl}$ vs. $Myo15a^{cre/+}$; $Six2^{fl/fl}$: base: 0.039, middle: 0.030, apex: 0.80). **C**. Normalized expression level of $Myo7a$ isoforms of $Myo15a^{cre/+}$; $Six2^{fl/fl}$ and control $Myo15a^{cre/+}$ cochlear cDNA at P20. $Myo7a$ expression level is normalized to $Gapdh$ and then log-transformed. (T-test p-value: $pan\text{-}Myo7a$ = 1.95e-4; $Myo7a\text{-}C$ = 7.04e-2; $Myo7a\text{-}N$ = 2.66e-3. Technical replication: 5 per group).

Supplementary Figure 1: A. Immunofluorescence images of WT and *ΔEnhancerA* mice at

P5 in the basal region of the cochlea. Hair bundle morphology in *ΔEnhancerA* mice is

comparable to that of WT mice. (Scale bar: 1 µm). **B**. Summary plots of the adaptation for

apical OHCs, basal OHCs and basal IHCs. Slow adaptation was measured at the 50% point

of the total peak current. Whiskers of the plot represent for Mean ± SD. Number of cells

(animals): Apical $EA^{+/-}$ = 8 (8), Basal $EA^{+/-}$ 13 (13), Apical $EA^{-/-}$ 7 (7), Basal $EA^{-/-}$ 10(10).

7. Methods:

Animal care and handling

The care and use of animals for all experiments described conformed to NIH guidelines. Experimental mice were housed in a 12:12 hour light:dark cycle with free access to chow and water in standard laboratory cages in a temperature and humidity-controlled vivarium. The protocols for the care and use of animals were approved by the Institutional Animal Care and Use Committees at the University of Virginia, the University of Colorado Denver, and the National Institute on Deafness and Other Communication Disorders. All the institutions mentioned above are accredited by the American Association for the Accreditation of Laboratory Animal Care. C57BL/6J (Bl6, from Jackson Laboratory, ME, USA) mice and sibling mice served as control mice for this study. Neonatal mouse pups (P0–P5) were sacrificed by rapid decapitation. Mature mice were euthanized by CO_2 asphyxiation followed by cervical dislocation.

Generation of Mouse strains for this study

For CRISPR/Cas mediated generation of the mouse models, we used the online tool CRISPR (http://crispor.tefor.net/crispor.py) to select suitable target sequences to ablate *Myo7a* isoforms. Single-guide (sg)RNAs are purchased from IDT (Integrated DNA Technologies, IA, USA). To produce genetically engineered mice, fertilized eggs were coinjected with Cas9 protein (PNA Bio, 50 ng/µl) and the sgRNA (30 ng/µl). Two-cell stage embryos were implanted on the following day into the oviducts of pseudopregnant ICR female mice (Envigo). Genotyping was performed by PCR amplification of the region of interest. Target sequences, sgRNAs, repair templates, and genotyping primers are listed below:

ΔEnhancerA **mice:**

To ensure the large deletion of *EnhancerA*, a pair of guide RNA surrounding the *EnhancerA* was injected.

sgRNA1: 5'-ACAACAGCGACUUCCCUGCAUGG-3', sgRNA2: 5'-CCTACGTTCCTAG CCAGTCTCGC-3'. Genotyping primers: Forward: 5'-ACCATGACCCCATAAGGTTC-3', W T reverse: 5'-GGGATGAATCATCACTCCTGC-3', Mutant Reverse: 5'-AGTGTTCTCCCGA GAGCCTTTC-3'.

Six2-HA **mouse:**

sgRNA: 5'-AGAGCTCTGTTCGCCTCCGG-3'. Repair template: GCGGCGCGGACCC ACTGCAGCATCACCACAGCCTGCAGGACTCCATACTCAACCCCATGTCGGCCAACCTTG TGGACCTGGGCTCCTACCCATACGATGTTCCAGATTACGCTTAGAGCTCTGTTCGCCTC CGGGCGCTTGCTTTCCGTGCTTGATGAAGCGGTGTGGGAGGGCGATAGATTCTGGAGT CTGTGCCGCGT. Genotyping primers: Forward: 5'-TCCATACTCAACCCCATGTCGG-3', Reverse: 5'-CCAAAGGATACCGAGCAGACCA-3'.

*Gfi1-cre***:**

*Gfi1^tm1(cre)Gan^* (*Gfi1-cre*) mouse line is Dr. Lin Gan at the University of Rochester. A targeting construct was used to replace the coding sequence of *Gfi1* from the translation initiation codon through exon 5 with the coding sequence of cre recombinase. Genotyping primers: CRE-Forward: 5'-GGGATAACGGACCAGTTG-3', Reverse: 5'-GCCCAAATGTTG CTGGATAGT-3'. *Gfi1-cre* WT allele genotyping primer: Forward: 5'-GGGATAACGGACCAGTTG-3', Reverse: 5'-CCGAGGGGCGTTAGGATA-3'

(More information:
https://www.informatics.jax.org/allele/MGI:4430258#:~:text=Mutation%20details%3A%20A%
20targeting%20construct,coding%20sequence%20of%20cre%20recombinase.)
***Myo15a-Cre*:**
To generate a *Myo15a-Cre* knockin (KI) mouse line, CRISPR/Cas9-mediated geno
me engineering was used to insert a "Cre-rBG (rat beta-globin intron) polyA" cassette
upstream of the translational start codon (ATG) in exon 2 of the mouse *Myo15* gene
(NCBI Reference Sequence: NM_010862.2), which is located on chromosome 11 and
contains 66 exons (Ensembl Transcript: ENSMUST00000071880). The cassette was pl
aced under the control of the endogenous *Myo15a* promoter to ensure physiologically
relevant Cre expression. Homology arms (~2–3 kb) flanking the insertion site were am
plified by high-fidelity PCR using BAC clone RP23-329M3 as a template and assemble
d with the Cre-rBG cassette into the targeting vector. For genome editing, Cas9 protei
n, guide RNAs (gRNAs), and the targeting vector were co-injected into fertilized C57B
L/6J one-cell embryos. Two gRNAs were designed to flank the insertion site with mini
mal predicted off-target activity: gRNA-A1 (GAGGGCCACCATGGCGGATG) matching th
e forward strand and gRNA-B1 (TTTCTTCTCCTCATCCGCCA) matching the reverse st
rand. Injected embryos were transferred into pseudopregnant females, and resulting pu
ps were genotyped by PCR using genomic DNA from tail biopsies. Correct integration
of the cassette was confirmed by Sanger sequencing across the homology arm–casset
te junctions. The mouse line was generated by Cyagen Biosciences (Santa Clara, CA),
and correctly targeted founders were bred with C57BL/6J animals to establish a stabl
e *Myo15a-Cre* line.
Genotyping primers: CRE-Forward: 5'-GGGATAACGGACCAGTTG-3', Reverse:
5'-GCCCAAATGTTGCTGGATAGT-3'. *Myo15a-Cre* WT allele genotyping primer: Forward:
5'-GGTCTCCAAGCAGTGTCTCCAA-3', Reverse: 5'- CTTGAGGCTCCGTTTGGGTTTC -3'
Six2::cre-egfp mice is purchased from The Jackson Laboratory (Strain #:009606), and
Six2^{flox/flox} are generated by previous studies from other labs. The genotyping methods are

provided by their corresponding reference[48,49].

Quantitative polymerase chain reaction (qPCR)
qPCR primers were designed to detect either *pan-Myo7a* transcripts or *Myo7a*
isoform-specific transcripts to evaluate their expression levels. Total RNA was extracted using
TRIzol reagent (15596026, ThermoFisher Scientific). Reverse transcription was performed
using the SuperScript™ IV Reverse Transcription system (18090010, ThermoFisher
Scientific). qPCR reactions were conducted using iTaq Universal SYBR Green Supermix
(1725121, Bio-Rad, CA, USA) on a CFX Opus 384 Real-Time PCR System (12011319,
Bio-Rad) for fluorescence measurement. Following analysis and plotting is done by R
(version 4.4.1).
Primers: *pan-Myo7a*: forward: 5'-CCTTCTCATTCGCTACCGGG-3', reverse: 5'-CCGG
TTGTTGCGTTTCATGT-3'; *Myo7a-C*: forward: 5'-CCGGTTGTTGCGTTTCATGT-3', revers
e: 5'-CCGACTCCCCGCTGATAATAC-3' ; *Myo7a-N*: forward: 5'-GTGCGAAACTTGAACCA
AACG-3', reverse: 5'-CCGACTCCCCGCTGATAATAC-3' ; *Myo7a-S*: forward: 5'-ACAACA

GCGACUUCCCUGCAUGG-3', reverse: 5'-CCGACTCCCCGCTGATAATAC-3' ; *Atp8b1*: fo

rward: 5'-ACGGCCGGTGGTCTTACATA-3', reverse: 5'-GCTTGTCACTCACGTCCTGAT-3'

; *Tmc1*: forward: 5'-GTGGGCCCTTCAGTGGTAAA-3', reverse: 5'-TTTCTGGCCCTTGG

CAGTAG-3' ; *Tmc2*: forward: 5'-GCCGGAAGGATCCTGCTAAG-3', reverse: 5'-CAAGGA

AGTGGACCTGGGTT-3'.

Immunofluorescence

Inner ear organs were fixed in 3% paraformaldehyde (PFA, Electron Microscopy

Sciences, PA) immediately after dissection for 20 min. Samples were washed three times

with phosphate-buffered saline (GIBCO® PBS, Thermo Fisher Scientific, Waltham, MA) for

5 min each. After blocking for 2 h with blocking buffer (1% bovine serum albumin, 3%

normal donkey serum, and 0.2% saponin in PBS), tissues were incubated in blocking buffer

containing primary antibody at 4 °C overnight. The following antibodies were used in this

study: rabbit polyclonal Myosin-VIIa antibody (catalog#: 25-6790, Proteus Biosciences Inc,

Ramona, CA. 1:100), mouse monoclonal Myosin-VI antibody (A-9, Santa Cruz, 1:100), rabbit

anti-HA antibody (C29F4, Cell Signaling Technologies, Catalog #: #3724, Danvers, MA).

Fluorescence imaging was performed using a Leica Stellaris 5 Confocal Microscope Platform

or Zeiss LSM880 with AiryScan.

Hearing tests in mice

Mice were anesthetized with a single intraperitoneal injection of 100 mg/kg ketamine

hydrochloride (Fort Dodge Animal Health) and 10 mg/kg xylazine hydrochloride (Lloyd

Laboratories). ABR and DPOAE were performed in a sound-attenuating cubicle

(Med-Associates, product number: ENV-022MD-WF), and mice were kept on a Deltaphase

isothermal heating pad (Braintree Scientific) to maintain body temperature.

ABRs of *ΔEnhancerA* mice and their BL6 controls were conducted on the equipment

purchased from Intelligent Hearing Systems (Miami, Fl). Recordings were captured by

subdermal needle electrodes (FE-7; Grass Technologies). The non-inverting electrode was

placed at the vertex of the midline, the inverting electrode over the mastoid of the right ear,

and the ground electrode on the upper thigh. Stimulus tones (pure tones) were presented at a

rate of 21.1/s through a high-frequency transducer (Intelligent Hearing Systems). Responses

were filtered at 300–3000 Hz, and threshold levels were determined from 1024 stimulus

presentations at 8, 11.3, 16, 22.4, and 32 kHz. Stimulus intensity was decreased in

5–10 dB steps until a response waveform could no longer be identified. Stimulus intensity

was then increased in 5 dB steps until a waveform could again be identified. If a waveform

could not be identified at the maximum output of the transducer, a value of 5 dB was added

to the maximum output as the threshold.

DPOAEs of the same group of WT and *Myo7a-ΔC* mice were recorded. While under

anesthesia for ABR testing, DPOAE was recorded using SmartOAE ver. 5.20 (Intelligent

Hearing Systems). A range of pure tones from 8 to 32 kHz (16 sweeps) was used to obtain

the DPOAE for the right ear. DPOAE recordings were made for $f2$ frequencies from 8.8 to

35.3 kHz using a paradigm set as follows: $L1 = 65$ dB, $L2 = 55$ dB SPL, and

$f_2/f_1 = 1.22$.

ABRs of *Myo15a::cre; Six2^{fl/fl}* mice and its corresponding control are conducted on RZ6

Multi-I/O Processor (Tucker Davis, FL 32615 USA) with Medusa4Z ABR Amplifier.

Responses are averaged by 512 stimulus sweeps presentations at 5, 10, 16, 24, 32, 40 and

48 kHz. Same ABR threshold determination strategies mentioned above are applied.

Whole-cell voltage-clamp electrophysiology recordings

Electrophysiological recordings

P4-P5 *ΔEnhancerA* homozygous and heterozygous mice were used for whole-cell patch

clamp experiments. Recordings were collected on the first or second row of outer hair cells

using an Axon 200B amplifier or a Multiclamp 700B (Molecular Devices). Initially, a patch

pipette filled with extracellular solution was used to clean the epithelium, and 2-3 adjacent

cells near the target cell were removed. Subsequently, a patch pipette filled with intracellular

solution was used to perform the whole-cell patch clamp on the cell. Apical perfusion was

continuously maintained until a gigaseal was achieved. After breakthrough of the patch

membrane to enter whole-cell configuration, a minimum waiting time of 5 minutes was

allowed for the intracellular solution to equilibrate. A fluid jet, using thin-walled borosilicate

pipettes (World Precision Instrument), was then applied to deliver step-like force stimuli to the

hair bundle. Experiments were performed at 18-22°C. Whole-cell currents were filtered at 10

kHz and sampled at 50 kHz using USB-6356 or USB6366 (National Instruments) controlled

by jClamp (SciSoft Company). All experiments used -80 mV holding potential unless

otherwise noted and did not account for the liquid junction potential.

Hair-bundle stimulation and motion recording

For the electrophysiological recordings, hair bundles were stimulated with a custom

three-dimensional—printed fluid jet driven by a piezoelectric disc bender (27 mm, 4.6 kHz;

Murata Electronics or PUI Audio). Thin-wall borosilicate pipettes were pulled to tip diameters

of 5 to 15 μm, filled with extracellular solution, and mounted in the fluid-jet stimulator. The

piezoelectric disc bender was driven by waveforms generated using jClamp. Stimulus

waveforms were filtered using an eight-pole Bessel filter at 1 kHz (L8L 90PF, Frequency

Devices Inc.) and variably attenuated (PA5, Tucker Davis) before being sent to a

high-voltage/high-current Crawford amplifier to drive the piezoelectric disc bender. During

stimulations, videos were taken of the hair-bundle motion with high-speed imaging at 10,000

frames per second using a Phantom Miro 320s or Veo 410L camera (Vision Research) when

illuminated with a TLED+ centered at 530 nm (Sutter Instruments). Videos were saved for

each stimulation and analyzed offline. Hair bundle displacement was determined as

described previously[50–52]. Briefly, hair bundle position was extracted using a Gaussian fit to a

high-pass–filtered hair-bundle image for a given vertical row of pixels in the image.

Analysis

All data were analyzed offline using jClamp (SciSoft), Matlab (MathWorks), Excel

(Microsoft), and Prism 7 (GraphPad). Figures were generated using Matlab, Prism 7, and

Adobe Illustrator.

Electrophysiological data analysis: The resting open probability of MET channels

(defined as P_{open}, or P_o) was determined from sine wave stimulation of the hair bundle and

calculated using Eq. (2), where I_{max} is the current elicited during maximum positive
stimulation, I_{leak} is the current remaining during maximum negative stimulation, and $I_{resting}$ is
the resting mechanosensitive current (defined as the current in the absence of stimulation,
subtracted by I_{leak}). We assumed we could observe a P_o of 0% and 100% during the
maximum negative and positive stimulations, respectively,

798
$$P_0 = \frac{I_{resting}}{I_{max} - I_{leak}} \quad (2)$$

799   For mechanical stimulus steps, adaptation time constant fits were obtained at ~50% peak
800 current using a double exponential equation

801
$$y = y_0 + A_1 e^{-\frac{x-x_0}{\tau_1}} + A_2 e^{-\frac{x-x_0}{\tau_2}} \quad (1)$$

802 where τ_1 and τ_2 are the decay constants, and A_1 and A_2 are the respective amplitudes;
803 the hair-bundle displacement was also fit with the same equation starting at a time after the
804 force stimulus plateaued, which was 0.5 ms after stimulus onset. All fits were automated in
805 MATLAB.
806
Scanning electron microscopy
Adult mice were euthanized by CO2 asphyxiation before intracardiac perfusion with 2.5%
glutaraldehyde (Electron Microscopy Sciences, Hatfield, PA) and 2% PFA. The otic capsule
was dissected and incubated in postfixation buffer at 4 °C overnight (2.5% glutaraldehyde, in
0.1 M cacodylate buffer, with 3 mM CaCl2). For neonatal mouse pups, the samples were
dissected and treated with postfixation buffer immediately. The otic capsules from adult mice
were incubated for two weeks in 4.13% EDTA for decalcification and then further dissected to
expose the organ of Corti. Samples underwent the OTOTO procedure and were dehydrated
using gradient ethanol and critical point drying. After sputter coating with platinum, the
samples were imaged on Zeiss Sigma VP HD field emission SEM using the in-lens
secondary electron detector.
818

6. Reference:

1. Hasson, T., Heintzelman, M. B., Santos-Sacchi, J., Corey, D. P. & Mooseker, M. S. Expression in cochlea and retina of myosin VIIa, the gene product defective in Usher syndrome type IB. *Proc. Natl. Acad. Sci. U. S. A.* **92**, 9815–9819 (1995).

2. Well, D. *et al.* Defective myosin VIIA gene responsible for Usher syndrome type IB. *Nature* **374**, 60–61 (1995).

3. Self, T. *et al.* Shaker-1 mutations reveal roles for myosin VIIA in both development and function of cochlear hair cells. *Development* **125**, 557–566 (1998).

4. Gibson, F. *et al.* A type VII myosin encoded by the mouse deafness gene shaker-1. *Nature* vol. 374 62–64 Preprint at https://doi.org/10.1038/374062a0 (1995).

5. Anna Underhill, Samuel Webb, Fiorella C Grandi, Jing-Yi Jeng, Jacques B de Monvel, Baptiste Plion, Adam J Carlton, Ana E Amariutei, Niovi Voulgari, Francesca De Faveri, Federico Ceriani, Mirna Mustapha, Stuart L Johnson, Saaid Safieddine, Corné J Kros, W. M. MYO7A is required for the functional integrity of the mechanoelectrical transduction complex in hair cells of the adult cochlea. *Proceedings of the National Academy of Sciences* **120**, (2025).

6. Corns, L. F. *et al.* Mechanotransduction is required for establishing and maintaining mature inner hair cells and regulating efferent innervation. *Nature Communications 2018 9:1* **9**, 1–15 (2018).

7. Li, S. *et al.* Myosin-VIIa is expressed in multiple isoforms and essential for tensioning the hair cell mechanotransduction complex. *Nat. Commun.* 1–15 (2020) doi:10.1038/s41467-020-15936-z.

8. Weil, D. *et al.* Human myosin VIIA responsible for the Usher 1B syndrome: A predicted membrane-associated motor protein expressed in developing sensory epithelia. *Proc. Natl. Acad. Sci. U. S. A.* **93**, 3232–3237 (1996).

9. Liu, X.-Z. *et al.* Autosomal dominant non-syndromic deafness caused by a mutation in the myosin VIIA gene. *Nat. Genet.* **17**, 268–269 (1997).

10. Liu, X. *et al.* Mutations in the myosin VIIA gene cause non-syndromic recessive deafness. *Nat. Genet.* 188–190 (1997) doi:10.1038/ng0697-188.

11. Boëda, B., Weil, D. & Petit, C. A specific promoter of the sensory cells of the inner ear defined by transgenesis. *Hum. Mol. Genet.* **10**, 1581–1589 (2001).

12. Ernest, S. & Rosa, F. M. A genomic region encompassing a newly identified exon provides enhancing activity sufficient for normal myo7aa expression in zebrafish

852 sensory hair cells. *Dev. Neurobiol.* **75**, 961–983 (2015).

13. Kawashima, Y. *et al.* Mechanotransduction in mouse inner ear hair cells requires transmembrane channel – like genes. *Journal of Clinical Investigation* **121**, 4796–4809 (2011).

14. Park, S., Jang, S., Lee, J.-H., Kim, S.-H. & Ryoo, Z. Y. Attempted rescue of circling mice by hair cell-specific expression (Myo7a promoter) of tmie transgene. *Med. Weter.* **73**, 399–403 (2017).

15. Sihan Li, Jinho Park, Tobey M. Phan, Edward H. Egelman, Jonathan E. Bird, J.-B. S. Tonotopic Specialization of MYO7A Isoforms in Auditory Hair Cells. *bioRxiv* **6**, (2025).

16. Sihan Li, Jinho Park, Tobey M. Phan, Edward H. Egelman, Jonathan E. Bird, J.-B. S. Tonotopic Specialization of MYO7A Isoforms in Auditory Hair Cells. *bioRxiv* **6**, (2025).

17. Li, S. *et al.* Myosin-VIIa is expressed in multiple isoforms and essential for tensioning the hair cell mechanotransduction complex. *Nat. Commun.* 1–15 (2020) doi:10.1038/s41467-020-15936-z.

18. Jen, H. I. *et al.* Transcriptomic and epigenetic regulation of hair cell regeneration in the mouse utricle and its potentiation by Atoh1. *Elife* **8**, 1–27 (2019).

19. Self, T. *et al.* Shaker-1 mutations reveal roles for myosin VIIA in both development and function of cochlear hair cells. *Development* **125**, 557–566 (1998).

20. Yu, I.-M. *et al.* Myosin 7 and its adaptors link cadherins to actin. *Nat. Commun.* **8**, 15864 (2017).

21. Michalski, N. *et al.* Molecular characterization of the ankle-link complex in cochlear hair cells and its role in the hair bundle functioning. *Journal of Neuroscience* **27**, 6478–6488 (2007).

22. Puig, R. R., Boddie, P., Khan, A., Castro-Mondragon, J. A. & Mathelier, A. UniBind: maps of high-confidence direct TF-DNA interactions across nine species. *BMC Genomics* **22**, 1–17 (2021).

23. Hammal, F., De Langen, P., Bergon, A., Lopez, F. & Ballester, B. ReMap 2022: A database of Human, Mouse, Drosophila and Arabidopsis regulatory regions from an integrative analysis of DNA-binding sequencing experiments. *Nucleic Acids Res.* **50**, D316–D325 (2022).

24. O'Brien, L. L. *et al. Transcriptional Regulatory Control of Mammalian Nephron Progenitors Revealed by Multi-Factor Cistromic Analysis and Genetic Studies. PLoS Genetics* vol. 14 (2018).

885   25.   Rauluseviciute, I. *et al.* JASPAR 2024: 20thãnniversary of the open-access database
886   of transcription factor binding profiles. *Nucleic Acids Res.* **52**, D174–D182 (2024).

887   26.   Self, M. *et al.* Six2 is required for suppression of nephrogenesis and progenitor
888   renewal in the developing kidney. *EMBO Journal* **25**, 5214–5228 (2006).

889   27.   Weber, S. *et al.* SIX2 and BMP4 mutations associate with anomalous kidney
890   development. *Journal of the American Society of Nephrology* **19**, 891–903 (2008).

891   28.   Li, J. *et al.* Chromatin remodelers interact with Eya1 and Six2 to target enhancers to
892   control nephron progenitor cell maintenance. *Journal of the American Society of*
893   *Nephrology* **32**, 2815–2833 (2021).

894   29.   Xu, J. *et al.* Eya1 interacts with Six2 and Myc to regulate expansion of the nephron
895   progenitor pool during nephrogenesis. *Dev. Cell* **31**, 434–447 (2014).

896   30.   Liu, Z. *et al.* Crucial and Overlapping Roles of Six1 and Six2 in Craniofacial
897   Development. *J. Dent. Res.* **98**, 572–579 (2019).

898   31.   Guan, J. *et al.* SIX2 haploinsufficiency causes conductive hearing loss with ptosis in
899   humans. *J. Hum. Genet.* **61**, 917–922 (2016).

900   32.   Matern, M. *et al.* Gfi1Cre mice have early onset progressive hearing loss and induce
901   recombination in numerous inner ear non-hair cells. *Sci. Rep.* **7**, 1–13 (2017).

902   33.   Caberlotto, E. *et al.* Usher type 1G protein sans is a critical component of the tip-link
903   complex, a structure controlling actin polymerization in stereocilia. *Proc. Natl. Acad.*
904   *Sci. U. S. A.* **108**, 5825–5830 (2011).

905   34.   Kros, C. J. *et al.* Reduced climbing and increased slipping adaptation in cochlear hair
906   cells of mice with Myo7a mutations. *Nat. Neurosci.* **5**, 41–47 (2002).

907   35.   Grati, M. & Kachar, B. Myosin VIIa and sans localization at stereocilia upper tip-link
908   density implicates these Usher syndrome proteins in mechanotransduction.
909   *Proceedings of the National Academy of Sciences* **108**, 11476–11481 (2011).

910   36.   Sato, O. *et al.* Human myosin VIIa is a very slow processive motor protein on various
911   cellular actin structures. *Journal of Biological Chemistry* **292**, 10950–10960 (2017).

912   37.   Zhao, S. *et al.* Deciphering enhancers of hearing loss genes for efficient and targeted
913   gene therapy of hereditary deafness. *Neuron* **113**, 1579-1596.e5 (2025).

914   38.   Self, M. *et al.* Six2 is required for suppression of nephrogenesis and progenitor
915   renewal in the developing kidney. *EMBO Journal* **25**, 5214–5228 (2006).

39.   Wong, E. Y. M., Ahmed, M. & Xu, P. X. EYA1-SIX1 complex in neurosensory cell fate

induction in the mammalian inner ear. *Hear. Res.* **297**, 13–19 (2013).

40.　Xu, J. *et al.* Chromatin remodelers and lineage-specific factors interact to target
enhancers to establish proneurosensory fate within otic ectoderm. *Proc. Natl. Acad.*
Sci. U. S. A. **118**, (2021).

41.　Ahmed, M. *et al.* Eya1-Six1 Interaction Is Sufficient to Induce Hair Cell Fate in the
Cochlea by Activating Atoh1 Expression in Cooperation with Sox2. *Dev. Cell* **22**,
377–390 (2012).

42.　Zhang, T., Xu, J., Maire, P. & Xu, P. X. Six1 is essential for differentiation and
patterning of the mammalian auditory sensory epithelium. *PLoS Genet.* **13**, 1–23
(2017).

43.　Weiming Zheng, Li Huang, Zhu-Bo Wei, Derek Silvius, Bihui Tang, P.-X. X. The role of
Six1 in mammalian auditory system development. *Development*
https://doi.org/10.1242/dev.00628 (2003) doi:10.1242/dev.00628.

44.　N A Bermingham, B A Hassan, S D Price, M A Vollrath, N Ben-Arie, R A Eatock, H J
Bellen, A Lysakowski, H. Y. Z. Math1: an essential gene for the generation of inner ear
hair cells. **284**, 1837–1842 (1999).

45.　Hertzano, R. *et al.* Transcription profiling of inner ears from Pou4f3ddl/ddl identifies
Gfi1 as a target of the Pou4f3 deafness gene. *Hum. Mol. Genet.* **13**, 2143–2153
(2004).

46.　Wallis, D. *et al.* The zinc finger transcription factor Gfi1, implicated in
lymphomagenesis, is require for inner ear hair cell differentiation and survival.
Development **130**, 221–232 (2003).

47.　Zhang, T., Xu, J. & Xu, P. X. Eya2 expression during mouse embryonic development
revealed by Eya2lacZ knockin reporter and homozygous mice show mild hearing loss.
Developmental Dynamics **250**, 1450–1462 (2021).

48.　Yang, H., Xie, X., Deng, M., Chen, X. & Gan, L. Generation and characterization of
Atoh1-Cre knock-in mouse line. *Genesis* **48**, 407–413 (2010).

49.　Li, C., Liu, H., Hu, Y. C., Lan, Y. & Jiang, R. Generation and characterization of Six2
conditional mice. *Genesis* **58**, (2020).

50.　Caprara, G. A., Mecca, A. A. & Peng, A. W. Decades-old model of slow adaptation in
sensory hair cells is not supported in mammals. *Sci. Adv.* **6**, 1–13 (2020).

51.　Caprara, G. A., Mecca, A. A., Wang, Y., Ricci, A. J. & Peng, A. W. Hair bundle
stimulation mode modifies manifestations of mechanotransduction adaptation. *Journal*

of Neuroscience **39**, 9098–9106 (2019).

52.   Peng, A. W. *et al.* Fluid Jet Stimulation of Auditory Hair Bundles Reveal Spatial
Non-uniformities and Two Viscoelastic-Like Mechanisms. *Front. Cell Dev. Biol.* **9**, 1–15
(2021).

Acknowledgement:

We thank Wenhao Xu and Daniel Grigsby at Genetically Engineered Murine Model Core (University of Virginia, RRID:SCR_025473) for generating mouse models used in this study. We thank Sijie Hao from Advanced Microscopy Facility (University of Virginia, RRID: SCR_018736) for the support of SEM imaging. We thank Ruliang Jiang (Cincinnati Children's Hospital Medical Center) for providing *Six2*$^{flox/flox}$ mice. We thank Maria Luisa S. Sequeira Lopez (University of Virginia) for providing the *Six2::cre-egfp* mice.

Funding source: J.B.S was supported by NIH grant RO1DC018842. Zeiss SEM of Advanced Microscopy Facility was supported by NIH SIG grant 1S10OD011966-01A1. The Titan Krios of Molecular Electron Microscopy Core was purchased with NIH grant SIG S10-RR025067. Y.R.K was supported by the National Research Foundation of Korea (NRF) grant funded by the Korea government (MSIT) (RS-2025-24535196).

Author contributions
Conceptualization: S.L., A. W.P., J.B. Shin; Methodology: S.L., S.J., Y.R.K., C.K., A.W.P., J.B. Shin; Experiments: S.L., S.J., Y.R.K., A.G., F.L.; Writing—review & editing: S.L., S.J., A. W.P., J.B. Shin; Supervision: U.K., A. W.P., J.B. Shin.

A

Myo7a isoforms from 5' RACE

Myo7a-C — Start codon

Myo7a-N — Start codon

Myo7a-S — Start codon

Basewise Conservation 3.296 –

−3.94 –

EnhancerA

ATAC-seq
Cochlea Hair cell
Supporting cell

1kb

B

Norm. Expression level by qPCR (%)

● WT
● ΔEnhancerA

*** *** *** ***

Pan-Myo7a   Myo7a-C   Myo7a-N   Myo7a-S

C

MYO7A   WT   ΔEnhancerA

WT

ΔEnhancerA

Apex

Middle

Base

D

IHC

* ** **

Apex   Middle   Base

Norm. MYO7A immunoreactivity

E

OHC

● WT
● ΔEnhancerA

* ** **

Apex   Middle   Base

Norm. MYO7A immunoreactivity

A *Myo7a* isoforms from 5' RACE: *Myo7a-C*, *Myo7a-N*. ATG. *EnhancerA*. 1kb. ATG. ATAC-seq track from Segil Lab. Cochlea Hair cell. SIX2 binding site by ChIP-seq ⟶. 10bp.

B SIX2 binding motif. ATGCAACCTGACCC. SIX2 binding site in EA.

C *Six2::cre-egfp*. GFP. Apex. Middle. Base.

D GFP / DAPI / MYO7A / Merge. Apex. Middle. Base. O. I. GER.

E DAPI/GFP. Apex. Middle. Base. O. I.

F *Six2-HA*. SIX2-HA. Middle. Apex. Base.

G SIX2-HA / DAPI / OCM / Merge. Apex. Middle. Base. O. I. GER.

H DAPI/SIX2-HA/OCM. Apex. Middle. Base. O. I.

I DAPI/SIX2-HA. P1. P7. Apex. Middle. Base. O. I.

J OHC. IHC. Normalized SIX2-HA IF Signal. Base Middle Apex. P1. P7. n.s. ** *** n.s. * n.s. n.s. n.s. n.s.

K Normalized SIX2 Transcript Level. E17 P0 P20. n.s. ***

A
F-actin   MYO7A   Merge

Six2fl/fl

Gficre/+ Six2fl/fl

B

IHC   OHC

Normalized MYO7A intensity

n.s.   n.s.

Six2fl/fl   *Gficre/+ Six2fl/fl*   *Six2fl/fl*   *Gficre/+ Six2fl/fl*

C

Six2fl/fl

Gficre/+ Six2fl/fl

D 8-week
MYO7A   F-Actin   Merge

Middle

Myo15acre/+

Myo15acre/+ Six2fl/fl

Base

Myo15acre/+

Myo15acre/+ Six2fl/fl

E 16-week
MYO7A   F-Actin   Merge

Middle

Myo15acre/+

Myo15acre/+ Six2fl/fl

Base

Myo15acre/+

Myo15acre/+ Six2fl/fl

F

Cell Survival (%)

***   n.s.

Base   Middle
Region

■ *Myo15acre/+*
■ *Myo15acre/+ Six2fl/fl*

G Base ~800µm Region

Six2fl/fl

Myo15acre/+ Six2fl/fl

H

Region of Complete OHC loss (µm)

***

Six2fl/fl   *Myo15acre/+ Six2fl/fl*

I OHC

Six2fl/fl

Myo15acre/+ Six2fl/fl

J 8-week

ABR threshold (dB)

n.s. n.s. n.s. n.s.   **   *   n.s.

5   10   16   24   32   40   48
Frequency (kHz)

■ *Myo15acre/+*
■ *Myo15acre/+ Six2fl/fl*

K 16-week

ABR threshold (dB)

**   **   **   ***   ***   ††   *↑
                                    *

5   10   16   24   32   40   48
Frequency (kHz)

■ *Myo15acre/+*
■ *Myo15acre/+ Six2fl/fl*

A

WT (C57BL/6)   *Six2^(fl/fl)*   *Myo15a^(cre/+) Six2^(fl/fl)*

F-actin MYO7A

MYO7A

B

Norm. MYO7A fluorescence signal

Base   Middle   Apex

*   *   n.s.

■ *Six2^(fl/fl)*
■ *Myo15a^(cre/+) Six2^(fl/fl)*

C

Norm. Expression by qPCR

pan *Myo7a*   *Myo7a-C*   *Myo7a-N*

***   n.s.   ***

■ *Six2^(fl/fl)*
■ *Myo15a^(cre/+) Six2^(fl/fl)*