Lipid metabolism of hepatocyte-like cells supports intestinal tumor growth by promoting tracheogenesis

Huang, K.*[1], Miao, T.*[1], Dantas, E.[3,4], Han, M.[1,2], Hu, Y.[1,2], Wang, K.[1], Sanford, J.[3,4], Goncalves, M.[3,4], and Perrimon, N[1,2]

1 Department of Genetics, Blavatnik Institute, Harvard Medical School, Harvard University, Boston, MA 02115, USA
2 Howard Hughes Medical Institute, Boston, MA, USA
3 Division of Endocrinology, Department of Medicine, Weill Cornell Medicine, New York, NY 10065, USA.
4 Meyer Cancer Center, Weill Cornell Medicine, New York, NY 10065, USA.

^Correspondence:
Kerui Huang: Kerui_Huang@hms.harvard.edu
Norbert Perrimon: perrimon@genetics.med.harvard.edu

*Authors contribute equally

Abstract

Tumors require metabolic adaptations to support their rapid growth, but how they influence lipid metabolism in distant tissues remains poorly understood. Here, we uncover a novel mechanism by which gut tumors in adult flies reprogram lipid metabolism in distal hepatocyte-like cells, known as oenocytes, to promote tracheal development and tumor growth. We show that tumors secrete a PDGF/VEGF-like factor, Pvf1, that activates the TORC1-Hnf4 signaling pathway in oenocytes. This activation enhances the production of specific lipids, including very long-chain fatty acids and wax esters, that are required for tracheal growth surrounding the gut tumor. Importantly, reducing expression in oenocytes of either the transcription factor *Hnf4,* or the elongase mElo that generates very long chain fatty acid suppresses tumor growth, tracheogenesis, and associated organ wasting/cachexia-like phenotypes, while extending lifespan. We further demonstrate that this regulatory pathway is conserved in mammals, as VEGF-A stimulates lipid metabolism gene expression in human hepatocytes, and lung tumor-bearing mice show increased hepatic expression of *Hnf4* and the lipid elongation gene *Elovl7*. Our findings reveal a previously unrecognized tumor-host interaction where tumors non-autonomously reprogram distal lipid metabolism to support their growth. This study not only identifies a

novel non-autonomous role of the TORC1-Hnf4 axis in lipid-mediated tumor progression

but also highlights potential targets for therapeutic intervention in cancer-associated

metabolic disorders.

Introduction

Tumor growth demands a growing supply of nutrients and oxygen. The growing

tumor constantly shapes its microenvironment to favor fast growth and expansion, through

secreted factors and extracellular matrix. The composition of the tumor microenvironment

(TME) varies between tumor types, but the hallmarks include increased blood vessels,

immune cells, stromal cells, and extracellular matrix [1]. To overcome the increasingly

hypoxic and acidic environment, TME promotes angiogenesis to restore oxygen and

nutrient supply. In addition to modulating the local TME, malignant cells can exert systemic

effects on distant organs—including the liver, skeletal muscle, and brain—to reshape the

host's metabolic pathways, ultimately promoting further tumor progression [2-6]. Liver is a

metabolic center for amino acids, carbohydrates, and lipid. A key liver-specific metabolic

function is the complete urea cycle (UC), which eliminates excess ammonia by converting

into urea, a less toxic waste. Previous studies have identified evidence of decreased UC

activity in the livers of 4T1 breast cancer bearing mice, and the plasma of children with

cancer [5], demonstrating an existing interaction between tumors and the liver. Delineating

molecular pathways involving tumor-host interaction can provide novel therapeutic targets.

Lipids play a critical role in cancer progression, by serving as building blocks, energy

sources and signaling molecules. It has been observed that tumor cells reprogram lipid

metabolism to enhance growth, characterized by increased lipid synthesis, lipid uptake,

fatty acid oxidation (FAO) and lipid storage [7]. Tumor cells also actively reshape lipid

metabolism in cells from the TME through secreted signaling molecules and drive TME to

favor tumor growth. For example, increased lipid uptake and FAO in regulatory T cells

(Tregs), tumor-associated macrophages (TAMs) and myeloid-derived suppressor cells

(MDSCs), promotes their immunosuppressive function [8-10]. However, whether tumor

cells also reshape lipid metabolism in distal organs, such as liver, which is a metabolic

center for lipid transport, is not known.

To better understand the reciprocal communication between liver and tumor cells,

we utilize *Drosophila* which has emerged as a simple yet powerful model to study tumor-

host interactions [11]. The fly midgut, oenocytes, fat body and malpighian tubules are

functionally equivalent to human digestive tract, hepatocytes, adipose tissue and kidney,

respectively [12]. Oenocytes, which are hepatocyte-like cells, are important for mobilizing

stored lipids during starvation and interact closely with metabolic tissues, such as the fat

body to regulate lipid uptake and reprocessing [13, 14]. In addition to regulating lipid

metabolism under nutritional deprivation, oenocytes are the major site for synthesis of

cuticular hydrocarbons in pterygote insects [15-17]. Adult oenocytes synthesize

hydrocarbons using fatty acyl-CoA precursors via reduction to fatty aldehyde and oxidative

decarbonylation [18, 19]. Many fatty acid synthases, elongases and desaturases expressed

in adult oenocytes produce various types of fatty acyl-CoA substrates that are processed to

form a complex blend of cuticular hydrocarbons [20-23]. Hepatocyte nuclear factor 4

(Hnf4), an oenocyte enriched transcription factor, plays an important role in regulating the

conversion of lipids to very long chain fatty acid, which are important substrates for

hydrocarbon synthesis [23]. Oenocytes are the major site for lipid transport and

processing, reminiscent of human hepatocytes. However, whether and how oenocytes

regulate tumor growth is unknown.

The *Drosophila* trachea is a network of oxygen transporting tubes, the functional

equivalent of mammalian blood vessels [24]. Trachea is composed of terminal tracheal

cells (TTCs), which are highly plastic cells, and the terminal branches (TBs) or tracheoles

which deliver oxygen to the tissues. TTCs are oxygen sensing cells, and hypoxic conditions

promote the formation of TBs through hypoxia-inducible factor-1α (HIF-1α; Similar (Sima)

in *Drosophila*), similar to the role of HIF-1α in mammalian angiogenesis. Trachea resides in

proximity of many tissues, including the midgut. Trachea penetrate the visceral muscles to

access the intestinal epithelium to supply it with oxygen [25]. The intestinal epithelium is

apicobasally polarized, composed of intestinal stem cells (ISCs) and their progeny.

Damage induced by enteric infection, oxidative agents and tumor growth can increase

tracheogenesis on the midgut, which is necessary for efficient damage-induced ISC-

mediated regeneration [26, 27]. Additionally, tumor-induced tracheogenesis in *Drosophila*

is reminiscent of cancer-induced angiogenesis in mammals [26].

Using a Drosophila gut tumor model, we uncover a novel mechanism through which

gut tumors reprogram lipid metabolism in distal oenocytes to promote both tracheal

development and tumor growth. We find that tumors secrete a PDGF/VEGF-like factor,

Pvf1, which activates the TORC1-Hnf4 signaling pathway in oenocytes. This activation

boosts the production of specific lipids, including very long-chain fatty acids and wax

esters, essential for tracheal growth around the tumor. Notably, reducing the activity of

Hnf4 and mElo, an elongase involved in generating very long-chain fatty acids, in oenocytes

suppress tumor growth, tracheal development, and cachexia-like phenotypes, while also

extending lifespan. Additionally, we demonstrate that this regulatory pathway is conserved

in mammals, as VEGF-A stimulates lipid metabolism gene expression in human

hepatocytes, and tumor-bearing mice exhibit increased hepatic expression of lipid

elongation genes. Our findings highlight a previously unrecognized tumor-host interaction,

where tumors non-autonomously reprogram distal lipid metabolism to facilitate their

growth.

Results

Oenocyte lipid metabolism regulated by Hnf4 controls tumor proliferation

To investigate how tumor growth rewires lipid metabolism of the whole fly, we

expressed an active form of *Yorkie (yki^{3SA}),* in adult gut stem cells, using the intestinal stem

cells (ISCs)/enteroblast driver (*Esg-Gal4* or *Esg-LexA*), hereafter referred to as Yki flies [28].

Esg-Gal4 or *Esg-LexA* was combined with the *tubulin* promoter-driven temperature-

sensitive *GAL80 (tubP-GAL80^{TS}),* to control the temporal expression of Gal4 or LexA. Next,

we conducted untargeted lipidomic profiling using an LC-MS/MS-based platform of whole-

body samples of adult female flies (*Esg-LexA > LexAop-yki^{3SA}*) expressing *yki^{3SA}* for 3 days,

together with control flies (*Esg > +*). The complete *Drosophila* genotype information can be

found in Supplementary Table 1. Female flies are used in the analysis unless otherwise

noted, because they show more significant and consistent tumor-associated cachexia

phenotypes. We detected 2471 lipid species in total representing 45 lipid classes

(Supplementary Table 2). The lipidomic analysis showed an increase in wax ester (WE),

monogalactosyldiacylglycerol (MGDG), phosphatidylglycerol (PG), phosphatidylcholine

(PC), and ceramide (Cer) level in Yki flies (Figure 1A). We also observed a slight reduction of

triglycerides (TG) levels in Yki flies at early stage, in agreement with previous findings [29].

WEs are major components of cuticular lipids, which cover nearly all parts of terrestrial

arthropods [30] to restrict water loss, and WE levels are significantly increased in Yki flies

(Figure 1B), yet the significance of this regulation is unknown. As the biosynthesis of

cuticular lipids mainly occurs in *Drosophila* oenocytes and Hnf4 is a main regulator of this

process [23], we tested whether oenocyte Hnf4 regulates the level of WEs in wild type flies.

WEs were reduced when *Hnf4* was knocked down in wild type adult oenocytes using the

PromE(800)-Gal4 Gene-switch driver (hereafter referred to as PromE^{gs}) [14], where the

driver is activated upon mifepristone feeding (RU) (Figure 1C). Next, to test whether Hnf4

regulates WE synthesis in Yki flies, we used the LexA-LexAop system to induce Yki tumors

in the gut and the Gal4/UAS system combining GAL80 in oenocytes (*PromE-GAL4,Tub-*

GAL80^{TS}, referred to as *PromE*) to knockdown *Hnf4*. Knockdown of *Hnf4* in oenocytes

decreased Yki-induced WE levels (Figure 1D), indicating that *Hnf4* expression in oenocytes

is required for the increase of WE level in Yki flies.

As a nuclear receptor, the localization of Hnf4 is highly enriched in oenocyte nuclei

compared to the fat body [23, 31]. To visualize the localization of Hnf4, we tagged the

endogenous *Hnf4* gene with 127D01, a small epitope tag that can be detected with a

specific nanobody [32, 33], using an homology directed repair pathway CRISPR-Cas9

targeted insertion method [32]. To validate the Hnf4-127D01 line, we knocked down *Hnf4* in

oenocytes (Figure S1A), which significantly reduced its nuclear localization (Figure S1B).

Hnf4-127D01 in wild type oenocytes showed high level of nuclear localization, consistent

with previous findings (Figure 1E) [31]. Interestingly, increased nuclear localization of Hnf4-

127D01 was observed in the oenocytes of Yki flies (Figure 1E-1F). Altogether, these data

suggest that oenocytes in Yki flies have increased Hnf4 activity. To test whether oenocyte

Hnf4 signaling is required for tumor growth, we measured ISC-mitosis using phospho-

histone 3 (pH3+) staining. Inhibition of *Hnf4* in oenocytes of Yki flies *(Esg-LexA>yki³ˢᴬ;*

PromE>Hnf4-i) for 3 days showed a decrease in pH3+ staining compared to controls (*Esg>*

UAS-yki³ˢᴬ), and a slower growth rate for tumor (Figure 1G). Importantly, knockdown of *Hnf4*

alone in wild type animals does not affect pH3+ number compared to control (Figure 1G),

indicating that the role of Hnf4 in Yki flies relates to the aberrant regulation of Hnf4 in Yki

flies. In wild type flies, knocking down *Hnf4* in oenocytes for 7 days resulted in a slight

reduction in ovary size (Figure S1C-D), suggesting long-term reduction of *Hnf4* is

detrimental to the ovary development. Further, oenocyte *Hnf4* knockdown in Yki flies for 3

days could rescue the level of both ovary wasting, bloating and induced wet weight (Figure

1H-1I, Figure S1E), indicating that aberrant Hnf4 activity in oenocytes contribute to tumor

growth and associated cachexia phenotypes.

Our result suggests that Hnf4 transcriptional activity is induced in Yki oenocytes. To

validate this, we measured whether the expression of Hnf4 targets in Yki oenocytes also

increased. Published studies identify that Hnf4 target genes are involved in beta-oxidation

and very long chain fatty acid (VLCFA) biosynthesis in oenocytes [23, 34], we tested

whether VLCFA biosynthesis genes (*Hnf4, mElo, CG30008, CG16904, FarO, ACC, HDAC,*

TER, Cyp4g1, Cpr), as well as genes involved in beta-oxidation (*Whd, Scu, Yip2* and

CG7461) were dysregulated in Yki flies. *CG30008, CG16904, FarO, ACC, HDAC, TER,*

Cyp4g1, Cpr, Whd, Scu, Yip2 showed no difference between control and Yki flies (Figure

S1F). However, we observed a significant increase in *mElo* (Figure 1K), which encodes a

methyl-branched cuticular hydrocarbon elongase [21] and a minor increase for *Hnf4 and*

CG7461 (Figure S1F).

mElo has recently been shown to produce long chain methyl-branched cuticular

hydrocarbons in oenocytes that help prevent desiccation [21], a key function of *Hnf4* in

oenocytes [23]. Additionally, *mElo* expression is highly enriched in wild type adult

oenocytes according to snRNAseq data (Figure 1L). Inhibition of *Hnf4* reduced *mElo*

expression, and overexpression of *Hnf4* led to increased levels of *mElo* (Figure S1G-H),

consistent with a previous report that mElo is a transcriptional target of *Hnf4* [23]. Further*,*

mElo is highly induced in the oenocytes of Yki flies and this induction is blocked when *Hnf4*

is knocked down (Figure 1K).

In wild type flies, reduction of *mElo* in oenocytes did not affect pH3+ number in the

gut (Figure 1M). To test the function of oenocyte *mElo* during tumor growth, we reduced the

level of oenocyte *mElo* in Yki flies and observed fewer pH3+ in the gut, suggesting that *mElo*

contributes to Yki tumor growth (Figure 1M). We validated the effectiveness of *mElo*

knockdown in the oenocytes (Figure S1I). Unlike *Hnf4,* reducing *mElo* expression in wild

type oenocytes did not cause steatosis and a reduction of ovary size, suggesting that other

targets of *Hnf4* regulate oenocytes steatosis and ovary size (Figure 1N and Figure S1J-S1K).

Consistently, *mElo* knockdown did not rescue ovary wasting in Yki flies (Figure S1L).

To assess whether oenocyte lipid metabolism affects the viability of Yki flies, we

inhibited *Hnf4* in oenocytes and observed a significant lifespan extension (Figure 1O).

Inhibition of mElo in oenocytes of Yki flies also significantly increased lifespan (Figure 1P).

Finally, *mElo* has a sex specific function in regulating different types of hydrocarbons [21].

Therefore, we tested whether *mElo* also regulates Yki viability in male flies. Strikingly*,*

inhibition of *mElo* also significantly increased the lifespan in Yki males (Figure S1M).

Oenocyte Hnf4 activity is regulated by TORC1

We next sought to determine the factors regulating Hnf4 level in oenocytes. Previous

studies have shown that inhibiting the TORC1 pathway in oenocytes leads to steatosis [35],

a phenotype that mimics reduced Hnf4 expression in these cells [23], suggesting that the

TORC1 pathway positively regulates Hnf4 transcription in oenocytes. The Tuberous

Sclerosis Complex (TSC) is known to negatively regulate TORC1 by inhibiting RAS homolog

enriched in brain (Rheb) activity [36]. Using a previously published snRNAseq dataset [35],

we found that overexpression of Tsc1 together with Tsc2 (TSC1,2) in oenocytes, which

inhibits TORC1, resulted in reduced Hnf4 expression specifically in oenocytes (Figure 2A),

but not in muscle nor fat body tissues, suggesting that that TORC1 positively regulates Hnf4

level in oenocytes.

Next, we examined the protein level of endogenously tagged Hnf4-127D01 following

TSC1,2 over-expression and observed a significant reduction in Hnf4-127D01 levels in

oenocytes, particularly in the nucleus as Hnf4 is a transcription factor (Figure 2B-C). Given

that Hnf4 regulates hydrocarbon synthesis in oenocytes and its knockdown increases the

susceptibility to dry starvation [23], we overexpressed *TSC1,2* in oenocytes and found that

it led to an increased sensitivity of flies to dry starvation (Figure S2A). Additionally, as Hnf4

regulates the expression of genes involved in very-long-chain fatty acid (VLCFA) synthesis

[23], we analyzed the expression of VLCFA synthesis genes in snRNAseq data from control

(*PromE>EGFP*) versus TSC1,2-overexprerssing oenocytes (*PromE>TSC1,2*). Using uniform

Manifold Approximation and Projection (UMAP), which identifies cell types in mixed

populations based on snRNAseq data, we found that VLCFA synthesis genes were highly

enriched in oenocytes (Figure 2D). The ModuleScore, calculated by averaging the

expression of the VLCFA synthesis pathway genes and subtracting the expression of a

randomly selected gene set expression, decreased in oenocytes upon *TSC1,2*

overexpression (Figure 2D).

Inhibition of TORC1 through overexpressing *TSC1,2,* or overexpressing Proline-rich

Akt substrate 40 kDa (*PRAS40),* or by feeding flies with rapamycin, all induced steatosis in

oenocytes (Figure 2E and S2B-C), as observed following *Hnf4* knockdown in oenocytes

(Figure 1N). However, simultaneous overexpression of *Hnf4* with *TSC1*,2 prevented the

development of steatosis (Figure 2E), indicating that TORC1 regulates oenocyte lipid

metabolism via Hnf4. Lastly, *TSC1,2* overexpression also reduced the expression of *mElo*

and *FASN2,* which are Hnf4 target genes, and overexpressing *Hnf4* simultaneously blocked

this decrease (Figure 2F-G). Together, these findings suggest that Hnf4 acts downstream of

TORC1 to regulate lipid metabolism in oenocytes.

Next, we also tested the effects of oenocyte *TSC1,2* in tumor growth. Consistent

with *Hnf4* and *mElo* knockdown, *TSC1,2* overexpression in oenocytes reduced tumor

induced WE levels (Figure 2H). We also observed an increased level of TORC1 activity, as

measured by phospho-4EBP (p4EBP) staining in oenocytes of Yki flies (Figure 2I-J).

Additionally, overexpressing *TSC1,2* blocked *mElo* expression in oenocytes of Yki flies

(Figure 2K), Yki-induced pH3+ cells in the gut (Figure 2L), as well as Yki-induced ovary

wasting and bloating (Figure 2M and S2D). Altogether, these results indicate that TORC1

regulate Hnf4 transcription and protein level in oenocytes and its activation is required for

tumor growth.

ISCs secreted Pvf1 regulates Hnf4 through PvR signaling in oenocytes and affects

tumor associated phenotypes

TORC1 activity is regulated through various pathways, one of which includes

PDGF/VEGF signaling. Studies have shown that PvR activation can stimulate the TOR

pathway [35, 37, 38]. Consistent with this, muscle-secreted Pvf1 can regulate TORC1

activity in the oenocytes [35], suggesting that the activation of the TORC1-Hnf4 axis in

oenocytes of Yki flies may occur through PvR activation. Yki-expressing ISCs are capable of

secreting higher levels of the Pvf1 ligand, which activates PvR signaling in surrounding

tissues [39]. However, *Pvf1* is expressed at low levels in oenocytes and is not significantly

induced in Yki flies (Figure S3A-C). To test whether ISC secreted Pvf1 can induce oenocyte

Hnf4 levels, we overexpressed *Pvf1* specifically in adult ISCs (*Esg>Pvf1*) and measured

Hnf4-127D01 nuclear level in oenocytes. *Pvf1* overexpression in ISCs increased nuclear

Hnf4-127D01 levels (Figure 3A-B), suggesting an increased transcriptional activity as

measured by increased level of *mElo* expression (Figure 3C). To test whether ISC-derived

Pvf1 activates PvR in oenocytes, we overexpressed an active form of PvR in oenocytes

which was associated with increased nuclear Hnf4-127D01 level, as well as increased level

of *mElo* expression (Figure 3D-F). To determine whether tumor secreted Pvf1 non-

autonomously activates Hnf4 level in oenocytes, we knocked down *Pvf1* expression in the

ISCs of Yki flies (*Esg>yki³ˢᴬ; Pvf1-i*) and observed its effect on Hnf4-127D01 nuclear level.

Esg>yki³ˢᴬ; Pvf1-i significantly reduced the level of Hnf4-127D01 nuclear localization

compared to *Yki* flies (*Esg>yki³ˢᴬ; Luc-i)* (Figure 3G-H). Consistently, *mElo* expression was

reduced in *Esg>yki³ˢᴬ; Pvf1-i* flies compared to Yki flies (Figure 3I). We also tested whether

Pvf1 reduction in Yki flies would affect tumor development, like Hnf4 inhibition in

oenocytes. As reported previously [39], inhibition of *Pvf1* in the tumor gut rescues both

bloating and ovary wasting (Figure 3J-K). Gut tumors in *Esg>yki³ˢᴬ;Pvf1-i* flies exhibited

slower growth at early stage, as shown by reduced number of pH3+ cells (Figure 3L).

Finally, to demonstrate that Pvf1 secreted from Yki tumors directly controls *mElo*

expression through PvR, we used the dual Gal4 and LexA systems to reduce *PvR* level in

oenocytes while inducing Yki in ISCs. *Esg-LexA>yki³ˢᴬ;PromE>PvR-i* animals showed

reduced *mElo* expression (Figure 3M), decreased the level of bloating and ovary wasting,

reduced the number of pH3+ cells in gut tumors (Figure 3N-P), and extended lifespan,

compared to Yki flies (Figure 3Q). Altogether, these data suggests that PDGF/VEGF induced

by Yki tumor non-autonomously increases Hnf4 activity in oenocytes through PvR.

Circulating lipids target trachea but not ISCs to promote trachea growth

To identify the lipid species induced by oenocytes in Yki flies, we performed

lipidomic analysis on the fly hemolymph of control flies, *Esg-LexA> yki³ˢᴬ* and *Esg-LexA>*

yki³ˢᴬ; PromE>Hnf4-i flies (Figure 4A, Supplementary Table 3). Among the lipid species that

are induced by Yki and decreased by *Esg-LexA > yki³ˢᴬ; PromE>Hnf4-i*, the majority are

phospholipids including phosphotidyl choline (PC), phosphatidylglycerol (PG) and

phosphatidylethanolamine (PE), triglycerides (TGs) and acylcarnitines (AcCas). More

specifically, AcCa 24:1 and AcCa 24:2, which are long-chain acylcarnitines (C$\geq$12)

secreted by mitochondria when metabolism is inhibited [40], were elevated in Yki flies

while reduced by *Hnf4* knockdown (Figure 4B-C). Circulating PC 36:4, and PE 32:5

phospholipids and TG(18:3_16:0_16:1), TG(18:2_16:0_18:0) triglycerides were induced in

Yki flies and rescued by *Hnf4* knockdown (Figure 4D-G). Interestingly, PC 36:4 is elevated in

the serum and plasma samples from breast cancer patients [41] and TG(18:2_16:0_18:0) is

induced in bladder cancer patients [42].

PC and PE are the most abundant phospholipids of cellular membranes and play

important role in lipoprotein metabolism. In insects, lipid transport in circulation relies on

lipophorins, which are the major lipoproteins. The increased circulation of PC 36:4 and PE

32:5 in Yki flies suggests an increase in lipoproteins that transport lipids from oenocytes

towards trachea or the midgut.

The transport of lipids, such as phospholipids from oenocytes, to the trachea or

ISCs to promote tumor growth has not been reported [26]. Thus, we inhibited Hnf4 activity

in oenocytes by overexpressing *TSC1,2*, knocking down or overexpressing *Hnf4*, and

observed no significant effect on the number of pH3+ cells in the midgut (Figure 4H). Since

95% of the hemolymph lipids are transported through lipophorin (Lpp) in flies [43], and that

low-density lipoprotein (LDL) receptor homologues LpR1 and LpR2 promote Lpp

internalization [44], we reduced LpR1 or LpR2 level in the ISCs. However, this had no

impact on the trachea area, skeleton length, and only a minor increase in the number of

branches in the *LpR1* knockdown condition (Figure S4A-C). Reducing LpR1 and LpR2

activities in the ISCs also did not affect the proliferation (Figure S4D). To further examine

the role of LpR1 or LpR2 in ISC tumors, we reduced LpR1 and LpR2 levels while

overexpressing Yki (*Esg> yki³ˢᴬ;LpR1-i and Esg> yki³ˢᴬ;LpR2-i*), and found that these

manipulations did not impair Yki-induced tracheal growth (Figure 4I-K) or prevent Yki-

induced ISC proliferation (Figure 4L). These results suggest that ISC proliferation does not

rely on circulating lipids in the hemolymph.

We further tested whether disrupting *LpR1* and *LpR2* activities in trachea could

blunt tracheal growth. Knocking down *LpR1* using a temperature sensitive trachea-specific

driver (*btl-GAL4, UAS-srcGFP, Tub-GAL80ᵀˢ,* referred to as *Btl*) reduced tracheal number of

branches, tube area and skeleton length compared to control flies (*Btl>+*), whereas

knocking down *LpR2* in trachea did not have a significant effect (Figure 4M-O). Reducing

LpR1 and LpR2 activities in the trachea did not affect ISC proliferation (Figure S4E). This

suggests that trachea growth, but not ISC growth, requires lipids generated from oenocytes

transported through lipophorins.

Oenocyte Hnf4 and mElo regulates tracheogenesis in the midgut

Next, we aimed to understand whether oenocyte lipid metabolism controls trachea

growth in wild type guts. Consistent with Hnf4 knockdown, oenocyte specific *mElo*

reduction and *TSC1,2* overexpression showed reduced level of whole body WE in non-Yki

flies (Figure 5A-B). WEs are essential parts of the outer layer of the trachea [45-48], and

waterproof (*wat*), which is involved in biosynthesis of wax, has been implicated in hydrophobic tracheal coating in *Drosophila* embryos [48]. Knocking down *Hnf4* did not affect the transepithelial barrier mediated by septate junctions in the midgut trachea on the midgut, as revealed by red dextran injection experiment (Figure 5C). However, reduction of oenocyte *Hnf4* altered the morphology of trachea associated with the midgut (R2-R4 regions) (Figure 5D). Specifically, we observed a reduction in the number of branches and total skeleton length, normalized to the area of the gut surface (Figure 5E-G). Consistently, we also observed a reduction in branch numbers, tube area and skeleton length when *mElo* was knocked down in oenocytes of wild type flies (Figure 5H-J). The effect of oenocyte Hnf4 on trachea number is not specific to the midgut, as we also observed a reduction of trachea total tube area in the ovary of wild type flies (Figure S5A-D). Overexpressing *Hnf4* or *mElo* is sufficient to induce total tracheal tube area and skeleton length in the midguts of wild type flies (Figure 5K-P). Finally, to eliminate the effect of RU feeding on trachea morphology, we fed control (*PromE>ctrl*) flies and observed no difference in trachea morphology (Figure S5E-G).

Hnf4-mElo axis in oenocytes mediates lipid metabolism and tracheogenesis induced by Yki tumors

Tumor growth in the adult *Drosophila* midgut has been previously shown to promote tracheole density [26]. Similarly, we observed an increase in the number of trachea branches, total tube area, and total tube length in the surrounding tissue of the Yki midgut (Figure S6A-D). This led us to investigate whether oenocyte lipid metabolism mediated by *Hnf4-mElo* axis is crucial for trachea growth in the Yki midgut. Our lipidomic analysis showed reduced whole body WE level when *Hnf4* is knocked down, or when *TSC1,2* is overexpressed in the oenocytes of Yki flies (Figure S6E).

To explore whether perturbations of lipid metabolism in oenocytes could block Yki-induced tracheogenesis in the midgut, we manipulated lipid metabolism by overexpressing TSC1,2 or knocking down Hnf4 or mElo. Using the LexA/LexAop and Gal4/UAS systems, we found that overexpressing TSC1,2 or reducing mElo significantly decreased the number of

tracheal branches, total tube area, and total tube length in the Yki midgut. Reducing Hnf4

levels, on the other hand, only decreased the total tube area (Figure 6A-C).

In addition, to verify whether midgut secreted Pvf1 regulates tracheal growth, we

knocked down *Pvf1* in ISC in addition to Yki overexpression. Reduction of *Pvf1* blocked Yki

induced number of branches, tube area and skeleton length (Figure 6D-F), indicating that

oenocyte lipid metabolism is important for trachea growth induced by Yki.

Regulation of hepatocyte lipid metabolism by pro-angiogenic factors in tumor models

Using the *Drosophila* Yki tumor model, we found that the tumor secreted

PDGF/VEGF-like factor Pvf1 can non-autonomously regulate Hnf4 activity in hepatocyte-

like cells to promote the production of VLCFA or related lipids. To test whether this

mechanism is conserved in mammals, we incubated HepG2 cells with vascular

endothelial growth factor A (VEGF-A), which is orthologous to Pvf1 in flies and a potent pro-

angiogenic growth factor expressed in many types of tumors. We measure the expression

level of ELOVL fatty acid elongase 1 – 7 (*ELOVL1-7*). Incubating HepG2 with VEGF-A for 24

hours can significantly increase the level of *ELOVL1, 2, 4, 6* and *7*, suggesting a conserved

regulation of PDGF/VEGF on lipid metabolism in hepatocytes (Figure 7A-E). In addition, we

collected liver samples from *KrasG12D/+;Lkb1f/f* (KL) mice, in which tumors were

specifically induced in the lungs, and analyzed the expression levels of *Hnf4a* and *Elovl1-7*

in the hepatocytes. *Hnf4a* and *Elovl7* showed higher expression level in the tumor bearing

mice, compared to control, suggesting that tumor growth in the lung increased the level of

genes involved in lipid metabolism in distal hepatocyte tissues in the KL mouse model

(Figure 7F-G). Altogether, these data showed that a conserved mechanism of PDGF/VEGF-

like factor in regulating VLCFA metabolism in hepatocytes through Hnf4.

Discussion

We reveal a novel mechanism by which gut tumors in Drosophila reprogram lipid

metabolism in oenocytes to promote tracheal development and tumor growth. Our

findings demonstrate that tumors secrete a PDGF/VEGF-like factor, Pvf1, which activates

TORC1 activity, triggering the expression of mElo through Hnf4 in oenocytes. This TORC1-

Hnf4 signaling pathway enhances the production of specific lipids, such as very long-chain

fatty acids and wax esters, which stimulate tracheal growth, a crucial process for tumor

expansion. We also show that reducing Hnf4 or mElo activity in oenocytes suppresses

tumor growth, tracheogenesis, and tumor-induced cachexia, while extending lifespan.

Additionally, we highlight the conservation of this regulatory pathway in mammals, where

VEGF-A stimulates lipid metabolism gene expression in human hepatocytes. Furthermore,

lung tumor-bearing mice exhibit increased Hnf4 and Elovl7 expression in peripheral

hepatocytes.

Pvf1, a tumor-secreted factor, has previously been linked to tumor-induced muscle

and fat body wasting, as well as renal dysfunction in *Drosophila* [39, 49, 50]. In this study,

we uncover a novel role for tumor-secreted Pvf1 in modulating oenocyte lipid metabolism

through TORC1 and Hnf4, which regulates the expression of the oenocyte-enriched gene

mElo (Figure 3). While Hnf4 activity is known to be induced by starvation and during the

developmental transition from pupae to larvae [23, 34], its upstream regulators remain

poorly understood. In MEF cells, the Hnf4 binding motif is highly enriched in the promoters

of TORC1-induced genes [51], suggesting Hnf4α mediates the transcriptional activities of

TORC1. Here, we demonstrate that TORC1 positively regulates Hnf4 protein levels in the

oenocyte nucleus through transcriptional regulation (Figure 2A). However, the precise

mechanisms controlling *Hnf4* transcription downstream of TORC1 remain unclear. REPTOR

and REPTOR-BP are likely key players in this process, as they are the major effectors of the

transcriptional response induced by TORC1 inhibition [52].

Our study identified a novel mode of communication between oenocytes and the

gut. Oenocytes produce lipid substrates that support gut tracheal growth, a process crucial

for tumor expansion and potentially ISC growth. Similar to tumor, bacterial infections and

epithelial damage also trigger ISC proliferation [53-55], suggesting that oenocyte-secreted

lipids may also support stress-induced ISC expansion. We found that lipids from oenocytes

are acquired by the tracheal cells instead of ISCs, as knockdowns of *LpR1* and *LpR2* in ISCs

have no effects in tumor growth (Figure 5I-L). In contrast, reducing *LpR1* in tracheal cells

was sufficient to decrease tracheal development (Figure 5M-O). This may be due to the

trachea's direct contact with hemolymph, allowing it to actively acquire lipids, while the

midgut is covered by visceral muscle and basement membrane [56, 57].

The mechanism by which VLCFA metabolism regulates tracheogenesis remains

unclear, though it may involve the induction of Reactive Oxygen Species (ROS) in the

trachea. ROS has been implicated in the activation of HIF-1α/Sima, and that VLCFAs is

known to induce mitochondrial dysfunction and ROS production [58-60]. Increased ROS

levels, through treatments such as paraquat, H_2O_2 or bacterial infection, have been shown

to induce tracheogenesis in the midgut [26]. Furthermore, manipulating ROS level in the

trachea alone can trigger tracheal branching and ISC proliferation [26]. Interestingly,

overexpressing *Hnf4* alone in oenocytes could promotes tracheogenesis, but does not

significantly induce ISC proliferation (Figure 4H). This difference may be attributed to

variations in driver strength, with tracheal development occurring prior to ISC proliferation.

We observed that VEGF-A stimulates lipid metabolism gene expression in human

hepatocytes, and lung tumor-bearing mice showed increased expression of *Hnf4* and

Elovl7 in peripheral hepatocytes. This work provides new insights into tumor-host

interactions, challenging the conventional view that tumor-associated metabolic changes

are merely passive processes and instead demonstrating their role in promoting tumor

growth. Specifically, inhibiting *Hnf4* and *mElo* in oenocytes slows tumor growth, halts

tumor-induced cachexia, and extends the lifespan of flies (Figures 1N, 1P, and S1M). Yki

activation increased levels of AcCas, PCs, and TAGs in the hemolymph, and inhibiting *Hnf4*

in oenocytes reversed these lipid changes induced by Yki (Figure 4A). This study identifies

the TORC1-Hnf4-Elovl7 axis and tumor-induced lipid metabolism as potential therapeutic

targets for managing tumor growth or as biomarkers for early tumor detection.

Hnf4 is a master regulator of multiple metabolic pathways in *Drosophila* and

inhibiting it could have broad effects on the host. For instance, knocking down *Hnf4* in non-

tumor-bearing flies led to increased steatosis and smaller ovaries (Figures 1N, S1C-D).

These wide-ranging effects may limit the efficacy of Hnf4 inhibition as an anti-tumor

strategy. In contrast, mElo, which is more specifically involved in lipid elongation, presents

a more targeted inhibition with lower toxicity and a greater extension of lifespan (Figures 1P,

S1M). Our study also suggests a potential role for elongases in tumor-induced

angiogenesis, as overexpression of ELOVL7 has been linked to prostate cancer and is

required for cancer growth [61]. While mElo is exclusively expressed in oenocytes in flies

(Figure 1L), its orthologue, ELOVL7, is widely expressed in multiple tissues, including the

prostate, kidney, brain, small intestine, and thyroid [62]. It is plausible that ELOVL7 is

upregulated in various tissues in cancer patients, indicating additional tissue targets for

anti-angiogenic therapies.

In summary, our data reveal a novel paradigm in which growing tumors actively

acquire substrates for tracheogenesis by secreting a PDGF/VEGF-like factor that activates

the TORC1-Hnf4 axis in distal oenocytes, inducing the expression of mElo. This promotes

the synthesis of VLCFAs, which are transported to the trachea to further support tumor

growth (Figure 7H).

Materials and Methods

Drosophila stocks and husbandry

All experiments were conducted using female flies due to their more consistent

bloating phenotype and larger oenocyte, which facilitated dissection. For temperature-

sensitive experiments, fly crosses were set up and maintained under a 12:12 hour

light:dark cycle at 18°C. Crosses were discarded after 5 days to maintain consistent

population density. Following eclosion, adult flies were collected and kept at 18°C for 2–3

days. GAL80^{TS} driver activation was achieved by incubating the flies at 29°C for the

durations specified in each experiment. For experiments involving the GeneSwitch driver,

fly crosses were maintained at 25°C under a 12:12 hour light:dark cycle and discarded after

3 days. Adult progenies were collected and fed on food containing 200 μM RU486

(Mifepristone, Cayman Chemicals, 10006317), prepared by dissolving the compound in

95% ethanol before adding it to standard fly food. Flies were fed on RU486-containing food

for 7 days.

Esg^{ts} (*esg-Gal4, UAS-GFP, tub-GAL80^{TS}*), *Esg-LexA* and *LexAop-yki^{3SA}-GFP, Attp40 UAS-*

Luciferase are from the Perrimon lab stock collection.

The following strains were obtained from the Bloomington *Drosophila* Stock Center

(BL), Vienna Drosophila Resource Center (V) and flyORF (F): *UAS-Hnf4 RNAi* (BL# 29375),

UAS-mElo RNAi (BL# 44510), *UAS-yki^{3SA}* (BL#28817), *UAS-Pvf1-3xHA* (F002862), *UAS-*

PvR[A] (BL#58496), *UAS-Pvf1 RNAi* (V#102699), *UAS-PvR RNAi* (V#43459), *UAS-Luciferase*

RNAi (BL#31603), Oe^{ts} (*Desat1-Gal4, tubP-Gal80*) (BL# 65406), *UAS-LpR1 RNAi* (BL#

50737), *UAS-LpR2 RNAi* (BL# 54461).

UAS-Hnf4 is a gift from Dr. Carl Thummel [31]; *UAS-TSC1, TSC2* is a gift from Dr.

Christen K Mirth [63] and UAS-PRAS40 was a gift from Dr. Aurelio Teleman [64], Btl^{TS} (*Btl-*

Gal4, UAS-srcGFP) and *QF6 >QUAS-mtdTomato* was a kind gift from Dr. Chrysoula Pitsouli

[26]. PromE^{GS}(*PromE-GeneSwitch-Gal4*) was a gift from Dr. Heinrich Jasper [65]. *UAS-*

CG18609 and *mFas-Gal4* were gifts from Dr. Henry Chung [66, 67]. *UAS-PRAS40* is a gift

from Dr. Aurelio A Teleman [52].

Hnf4-127D01 was generated through CRISPR-mediated tagging of endogenous Hnf4

with 127D01-tag as described previously [32]. Briefly, we cloned the sequence 1 kb

upstream and 1 kb downstream of the stop codon into the donor vector pScarlessHD-C-

3x127D01-DsRed (Addgene#171578). A sgRNA plasmid pCFD3-Hnf4-sgRNA that target the

seed sequence (CCAGAGACTGGTTACTAGAAGAA) near the stop codon of Hnf4 was injected

into yw;nos-cas9/CyO embryos together with the donor plasmid. Positive transformants

with red eye fluorescent eyes were outcrossed and successful Kis were confirms by PCR (F:

GTACAACCGGAGCGAGGGTA; R: CAAAATGGTTCAAGGCATAACA) and sequencing.

Subsequently, 3xP3dsRed was excised using piggyBac transposase (BL#8285).

Untargeted lipidomic analysis

Lipidomics and metabolomics of fly samples was performed following the Folch

method (Folch et al., 1957). 15 adult flies, or hemolymph from 200-300 adult flies were

used for lipidomics and metabolomics. Solid samples were homogenized in a 1 mL Dounce

homogenizer (TIGHT) on ice with 0.6 mL chloroform and 0.3 mL methanol (2:1 chloroform-

methanol mixture). Homogenized samples were transferred to a 15 mL glass tube with a

Teflon cap, and an additional 0.6 mL chloroform and 0.3 mL methanol were added to make

the final volume approximately 1.8 mL. Hemolymph was transferred to a 15 mL glass tube

containing 1.2 mL chloroform and 0.6 mL methanol (2:1 chloroform-methanol mixture).

Samples were vortexed briefly and incubated at room temperature on a rotator for 30–45

minutes. Following incubation, 0.2 volumes (0.36 mL) of HPLC-grade deionized water

(HPLC dH2O) were added. The mixture was vortexed three times, followed by

centrifugation at 1000 × g for 10 minutes at 4°.

The upper aqueous phase, containing polar metabolites, was carefully collected

using a glass pipette, while avoiding the middle phase. Metabolites were transferred to

microcentrifuge tubes, evaporated under vacuum using a SpeedVac rotary evaporator

overnight, and stored at -80°C. Polar metabolites were resuspended in 20 µL of HPLC-grade

water and then injected and analyzed using a hybrid 6500 QTRAP triple quadrupole mass

spectrometer (AB/SCIEX) connected to a Prominence UFLC HPLC system (Shimadzu). The

lower phase, containing non-polar lipids, was transferred to a clean 1.85 mL glass test tube

using a glass pipette. The phase was dried under nitrogen gas for approximately 30

minutes, and dried lipids were stored at -80°C to prevent oxidation. Non-polar lipids were

resuspended in 35 µL of HPLC-grade 50% methanol/50% isopropyl alcohol (IPA) and then

injected and analyzed using untargeted high-resolution LC-MS/MS on a Thermo QExactive

Plus Orbitrap mass spectrometer. LipidSearch version 4.2 (Thermo Scientific) was used to

identify lipid molecules, with the quantification of ion intensity by measuring the area size

of identified peaks.

The full names of the lipid species can be found in Supplementary Table 4. For

lipidomic quantification using whole fly bodies, the same number of flies was used for lipid

extraction. Lipid species were normalized to the total reads in each sample. For

hemolymph lipidomic analysis, lipid species were normalized to both the volume of

hemolymph extracted and the total reads in each sample.

RNA extraction Quantitative RT-PCR

Adult female fly oenocytes were dissected in cold 1xPBS before RNA extraction. For oenocyte dissection, the fat body was removed through liposuction and then oenocytes were removed from the cuticle using a fine-tip glass needle. Tissue lysis, RNA extraction, and cDNA synthesis were performed using Cells-to-CT Kit (Thermo Scientific, #44-029-53). For female adult whole body mRNA extraction, four flies per biological replicate were collected. Total RNA was extracted using Direct-zol RNA MicroPrep kit (Zymo Research, #R2060). mRNA was extracted from HepG2 cells using Directo-zol RNA MicroPrep kit (Zymo Research, #R2060). 30-50mg of mice livers were dissected and total RNA was extracted using Qiagen RNeasy Mini kit (Qiagen #74104). All the cDNAs were synthesized using the iScript cDNA synthesis kit (Bio-Rad, #1708896) in a 40ul reaction mixture using 1000ng total RNA. Quantitative real-time RT-PCR (RT-qPCR) assays were performed using iQ SYBR Green Supermix (Bio-Rad, #1708880) on a CFX96 Real-Time PCR Detection system (Bio-Rad). Two to four independent biological replicates were performed with two technical replicates. The mRNA abundance of each gene was normalized to the expression of *RpL32* or *aTub84B* for fly samples, Gapdh for mouse samples and ACTB for HepG2 samples. Primer sequences are listed in the Supplementary Table 5.

Fly imaging

Drosophila ovaries were dissected in cold PBS. ovaries were imaged immediately in cold PBS using a ZEISS Axiozoom V16 fluorescence microscope. Flies were anesthetized on CO_2 pad and imaged using ZEISS Axiozoom V16 fluorescence microscope. The size of the ovary and abdomen is measured using Fiji (version: 2.14.0/1.54f).

Oil Red O staining

The method was as previously described [13]. Flies were dissected in PBS and fat bodies were removed with liposuction. Specimens were fixed in 4% paraformaldehyde for 20 minutes. Specimens were rinsed twice with distilled water. Oil Red O (O0625-, Sigma-Aldrich) was dissolved in isopropanol to make 0.1% Oil Red O stock. A working solution was prepared by combining 3 mL of 0.1% Oil Red O in isopropanol with 2 mL of distilled

water. The solution was freshly prepared and filtered through a 0.2-µm syringe filter before

use. Specimens were incubated for 25 minutes in Oil Red O stain, rinsed briefly with PBS

and mounted in Vector Shield for imaging. Oil Red O images were taken using Olympus

VS200 Slide Scanner, equipped with color camera (2448x2048 pixels, 3.45µm).

Immunostaining and image analysis

For immunostaining using the nanobodies, fly abdomens were dissected in PBS at

room temperature with fat bodies removed and fixed in 4% formaldehyde in PBS for 30

minutes. Specimens were washed three times in PBST (0.1% Triton-X-100), 5 minutes each.

The specimens were then incubated in 5% NGS (normal goat serum in PBST) for 1 hour.

Nb127D01-ALFA, 0.2mg/ml was used at a dilution of 1:500, incubating the samples for 1

hour and a half with mild shaking. Followed by 0.1% PBST washes, 5 minutes each wash.

AlexaFluor 594 anti-Alpaca (Jackson Immuno Research, 128-585-232) was diluted 1:400 in

PBS-TX + 5% NGS and incubate with the specimens at room temperature with mild shaking

for 1 hours. After washes and DAPI staining, tissues were mounted in Vectashield and

imaged with Olympus VS200 Slide Scanner or confocal microscope Zeiss Axio Observer Z1.

For regular immunostainings, fly tissues were dissected in PBS at room temperature.

Tissues were fixed in 4% paraformaldehyde in PBS for 30 minutes. Specimens were washed

three times in 0.1% PBST and incubated in 5% NDS solution for 1 hour. Tissues were then

incubated with primary antibodies diluted in PBST and blocking solution overnight at 4°.

Then the tissues were washed three times with PBST at room temperature. Secondary

antibodies were applied for 1 hour. Excess antibodies were washed in PBST, and tissues

were incubated with DAPI (Invitrogen, D1306) dilute in PBST for 5 minutes. Tissues were

then mounted in Vectashield (Vector Laboratories Inc, H-1000). The following primary

antibodies were used: rabbit-anti-pH3 (1:500, Cell Signaling, 9701), rabbit-anti-p4E-BP

(1:800, Cell Signaling, 2855S). Secondary antibodies against mouse, rabbit, chicken or

alpaca conjugated to Alexa Fluor 488 and 594 (Jackson Immuno Research) were used at

1:800 for anti-p-4EBP, 1:1000 for anti-pH3 and 1:400 for Nb127D01-ALFA.

Lifespan analysis

For survival analysis, flies were collected within 24 hours post-eclosion (20 females and 3 males per vial) and maintained on standard laboratory fly food at 18°C, with 60% humidity, and a 12-hour light/dark cycle. After two days, mature and mated females were transferred to 29°C to induce gene expression. To maintain vial cleanliness, flies were transferred to fresh vials every two days, and the number of dead flies was recorded daily.

snRNAseq analysis

Data and plots were generated from the Fly Cell Atlas, generated using the 10X platform, "Stringent All" dataset. Other snRNAseq dataset was analyzed as described in [35], where a resolution of 0.4 was chosen as the clustering parameter. For the Seurat Module Score analysis, Seurat AddModuleScore function was used to calculate the average expression of a group of genes in Very Long Chain Fatty Acid Synthesis pathway in each cell (for details of the list, see Supplementary Table 6), then subtracted by a randomly picked gene set expression.

Rapamycin feeding

Rapamycin was dissolved in DMSO to make 2mM rapamycin stock. Rapamycin stock was further diluted to 200μM in ethanol and evenly distributed on to food surface. A small piece of Kimwipe was used to remove excess liquids on the food. DMSO dissolved in ethanol was used as control. Flies were transferred to new food every two days. Flies were fed for 6-7 days before dissection.

Dextran injection

Using a fine glass needle, ~100nl of 50 mg/ml 10,000 MW TR-dextran (Thermo Fisher Scientific, D-1817) diluted in water was injected in the abdomen of adult females. 6 hours after injection, abdomen and thoraces were separated and whole abdomen were fixed in

4% paraformaldehyde for 30 minutes with mild shaking. After fixation, midguts were

dissected and washed briefly with PBS and mounted in Vectashield. Midguts were imaged

using a single point laser scanning confocal, Olympus IX83 to measure the trachea

leakage. A Z-stack was taken, and maximum projections were used to generate final image

for analysis.

Trachea imaging

Generating fly strains that incorporate both a trachea reporter and other genetic

modifications is challenging. Therefore, the following method, as described in [26], was

used to quantify the trachea. Flies were dissected in cold PBS, and intestines were

carefully extracted to maintain their original structure. Tissues were then fixed in 4%

paraformaldehyde for 20 minutes, mounted in Vectashield, and imaged within 3 days.

Whole intestine images were captured using an Olympus VS200 Slide Scanner equipped

with a color camera. Only the upper layer of the tissue, in contact with the coverslip, was

analyzed.

Trachea analysis

To automate trachea measurement, we used the previously developed software

AutoTube [68]. Only the topmost Z-stack, which provided the clearest image of the trachea,

was analyzed. Images corresponding to the R2-R4 regions of the gut were cropped,

avoiding areas with the major trunk of the trachea to prevent skewing the results. Images

were thresholded on a white background and analyzed with AutoTube using the following

settings:

- Input Color Channel: Green

- Intensity adjustment: Autocontrast

- Illumination correction: Illumination size set to 51

- Noise correction: BM3D

- Tube detection: Finer tubes with a tube size of 3

- Small regions removed: 0.01% of Ima

- Hole filling: Hole size set to 1

- Threshold type: MultiOtsu

- Short ramifications removed: Spur length set to 15

- Spatial branch merging: Spatial distance set to 10

The analysis provided results for "Number of Branches (μm)," "Total Tube Area (μm)," and

"Total Skeleton Length (μm)," which were normalized to the corresponding intestinal area

and were multiplied by an arbitrary number 1000000. Results from the R2-R4 regions were

averaged for each gut.

Animal Care and Tumor Induction

$Kras^{G12D}/+;Lkb1^{f/f}$ mice, previously described [3, 69], were backcrossed to the FVB

strain. Male mice were maintained on a 12-hour light/dark cycle at an ambient temperature

of 22°C, with ad libitum access to rodent chow (PicoLab Rodent 5053; Lab Diet, 3.43

kcal/g) and drinking water. Tumors were induced in adult male and female mice (12 to 20

weeks old) via intranasal administration of 75 μL PBS containing 2.5×10^7 pfu of adenovirus

SMV-CRE (Ad5CMV-Cre) purchased from the University of Iowa Gene Transfer Vector Core

(Iowa City, IA). All animal experiments were conducted with the approval of the Institutional

Animal Care and Use Committee (IACUC) at Weill Cornell Medicine (WCM) under protocol

number 2013-0116.

Mice tissue collection

664   Male mice were euthanized with CO_2, followed by blood collection via cardiac

665   puncture. Liver, gonadal adipose tissue, and skeletal muscles were dissected and flash-

666   frozen in liquid nitrogen and stored at -80°. The control mice were a mix of fasted and fed

667   mice. "Fasted" mice were fasted for 12 hours and "fed" mice were fasted for 12 hours and

668   then allowed free access to food for 4 hours before tissue collection.

VEGFA incubation

HepG2 cells were cultured in HyClone Dulbecco's Modified Eagle Medium (DMEM) with high glucose (SH30022.01): with L-glutamine; without sodium pyruvate. Additional 50U penicillin/ml and 50ug streptomycin/ml were added. HepG2 cells were seeded at density of 5 x 10^5 in 12-well-plate (VWR, 29442-040), treated or not with 50ng/ml Recombinant Human VEGF 165 (PeproTech, 100-20-2UG) for 24 hours. Total cellular RNA was isolated using the Trizol reagent and extracted through Direct-zol RNA MicroPrep kit (Zymo Research, #R2060).

Acknowledgments

We thank Dr. John M. Asara at the Beth Israel Deaconess Medical Center for his help on lipidomic analysis. We thank the assistance provided by the Microscopy Resources on the North Quad (MicRoN) core and we thank Dr. Simon F. Norrelykke at the Image Analysis Collaboratory at Harvard Medical School for advice on AutoTube. We also thank Dr. Ah-Ram Kim, Dr. Sun Jin Moon at Harvard Medical School, Dr. Hua Bai at Iowa State University, Dr. Eileen White and Dr. Maria Gomez- Jenkins at Rutgers University for scientific discussion and insights. We thank Dr. Ju Xu at Chinese Academy of Sciences for the help generating Hnf4-127D01 fly strain. We also thank Christians Villalta for the help on fly dextran injection.

This article is subject to HHMI's Open Access to Publications policy. HHMI lab heads have previously granted a nonexclusive CC BY 4.0 license to the public and a sublicensable license to HHMI in their research articles. Pursuant to those licenses, the author-accepted manuscript of this article can be made freely available under a CC BY 4.0 license immediately upon publication.

Funding

This work is funded by NIH/NCI Grant #5P01CA120964-15 and is delivered as part of the CANCAN team supported by the Cancer Grand Challenges partnership funded by Cancer Research UK (CGCATF-2021/100022) and the National Cancer Institute (1 OT2 CA278685-01). N.P. is an investigator of the Howard Hughes Medical Institute.

Author contributions

Conceptualization: KH, TM, MG, NP; Methods, Data Acquisition, Resources, Visualization and Imaging: KH, TM; Mice sample preparation: ED, JS; Data Acquisition: KW, Bioinformatic Analysis: MH, YH; Independent replication of experiments at Harvard Medical School: KH; Writing – original draft: KH, Writing – review, editing & revising: KH, TM, NP.

Figure 1: Oenocyte Hnf4 regulates wax ester synthesis and supports Yki-induced tumor growth.

(A-B) Heatmap of the normalized total ion intensity of major lipid classes identified in whole bodies from control flies (*Esg-LexA*) or Yki flies (*Esg-LexA/LexAop-yki^{S3A}; PromE-GAL4, Tub-GAL80^{TS}/+*), with four biological replicates per condition. Scaled colors are presented as the values relative to the average of each row. (B) Untargeted lipidomic profiling using LC-MS/MS of whole-body extracts revealed increased wax ester (WE) levels in Yki flies. (C) WE level in oenocyte-specific knockdown of *Hnf4* using the RU486-inducible *PromE^{gs}* driver (+RU) compared to controls (-RU). (D) WE level in control flies, Yki flies, and *Hnf4* knockdown in oenocytes of Yki flies, N = 4 biological replicates. (E-F) Immunostaining of Nb127D01 for nuclear-localized, endogenously tagged Hnf4 (Hnf4-127D01) in oenocytes in controls and in Yki flies. DAPI staining marks nuclei, with dashed lines enclosing oenocytes. N = 8 independent experiments. (G) Quantification of intestinal stem cell (ISC) proliferation, as indicated by phospho-histone H3 (pH3+) cells in control. (H-I) Ovary sizes and quantification. (J) Abdominal bloating. (K) UMAP from whole body (without head) snRNAseq, *mElo* expression is enriched in adult oenocytes. (L) *mElo* expression in oenocytes, normalized to housekeeping gene, in Yki flies and in Yki flies with *Hnf4* knockdown. N = 2 biological replicates from 2 independent experiments. (M) Quantification of pH3+ cell number, N =8 flies. (N) Oil Red O staining in oenocytes of *Hnf4* knockdown or

mElo knockdown. N = 8 flies. (O-P) Lifespan analysis showing that oenocyte-specific

knockdown of *Hnf4* (O) or *mElo* (P) in Yki flies, compared to only Yki or control flies, N = 40.

Scale bar shows 20μm. Data are presented as mean ± s.e.m. Statistical significance was

determined using Student's t-test or one-way ANOVA with appropriate post hoc tests, or

Log-rank (Mantel-Cox) test. *$P < 0.05$, **$P < 0.01$, ****$P < 0.0001$.

Figure 1S: Hnf4 and mElo regulate metabolic and tumor-related phenotypes in Yki

flies.

(A) Confirmation of *Hnf4* knockdown efficiency via qPCR, N = 2 biological replicates from 2

independent experiments. (B) Immunostaining of Nb127D01 for nuclear-localized,

endogenously tagged Hnf4 (Hnf4-127D01) in oenocytes with *Hnf4* knockdown or control

flies, with dashed lines enclosing oenocytes. (C-D) Oenocyte-specific *Hnf4* knockdown

reduces ovary size compared to control flies, quantification in (D), n = 9 - 15. (E) Wet weight

measurement of fly whole body, N = 3 biological replicates. (F) Expression of VLCFA

biosynthesis and β-oxidation genes in control and Yki flies. (G-H) *mElo* expression

measured by qPCR in oenocyte specific *Hnf4* knockdown or *Hnf4* overexpression, N = 2

biological replicates from 2 independent experiments. (I) Confirmation of *mElo* knockdown

efficiency via qPCR. (J-K) Ovary size in oenocyte *mElo* knockdown flies compared to

control, quantification in (K). (L) Ovary size measurement in control and Yki flies, N = 8-10

flies. (M) Lifespan analysis in Yki-expressing flies upon oenocyte-specific *mElo* knockdown

in male flies. Data represent mean ± s.e.m.; significance assessed using appropriate

statistical tests.

Figure 2: TORC1 regulates Hnf4 activity and lipid metabolism in oenocytes.

(A) snRNAseq analysis showing *Hnf4* expression in fat body, muscle and oenocytes upon

overexpression of *TSC1,2* in oenocytes. (B-C) Immunostaining of Hnf4-127D01 in

oenocytes when *TSC1,2* is overexpressed in oenocytes compared to control. DAPI marks

nuclei, with dashed lines enclosing oenocytes, quantification of Hnf4-127D01 signal

intensity in the nucleus, N = 8-10 flies. (D) ModuleScore from snRNAseq analysis visualizes

VLCFA synthesis gene expression across muscle, fat body and oenocytes, in control and upon oenocyte *TSC1,2* overexpression. (E) ORO staining indicates steatosis in oenocytes induced by *TSC1,2* overexpression, and in *TSC1,2* with *Hnf4* overexpression, N = 8. (F-G) Expression of Hnf4 targets *mElo* and *FASN2* in *TSC1,2* overexpression and in *TSC1,2* with *Hnf4* overexpression. N = 2 biological replicates from 2 independent experiments. (H) Whole body WE level in ISC Yki flies and Yki with oenocyte *TSC1,2* overexpression. (I-J) P-4EBP immunostaining indicates TORC1 activity in oenocytes of Yki flies, quantification of P-4EBP intensity is shown on the right (J), N = 8-10 flies. (K) *mElo* expression in whole body mRNA extracts in Yki flies and Yki flies with *TSC1,2* overexpression in oenocytes, N = 6 biological samples. (L) Quantification of pH3+ cell number, N = 8 – 10 flies. (M) Ovary size measurements are rescued by *TSC1,2* overexpression in Yki flies, N = 15-30 flies.

Figure 2S: The role of oenocyte TORC1 in regulating lifespan, steatosis and Yki wasting phenotype.

(A) Sensitivity to dry starvation in flies overexpressing *TSC1,2* in oenocytes compared to controls. Adult oenocyte specific driver (*mFas-Gal4*) was used [66]. (B-C) ORO staining indicates steatosis in oenocytes with *PRAS40* overexpression, or in rapamycin treatment. (D) Host bloating quantification in fly abdomen in Yki flies and in Yki flies with *TSC1,2* overexpression, compared to control. Dashed lines enclose oenocytes in all immunostaining images. Data represent mean ± s.e.m.; statistical significance determined using Student's t-test or ANOVA as appropriate.

Figure 3: ISCs secreted Pvf1 activates Hnf4 through PvR signaling in oenocytes and affects tumor associated phenotypes.

(A-B) Immunostaining of Hnf4-127D01 nuclear localization in oenocytes upon *Pvf1* overexpression in ISCs (*Esg>Pvf1*). Dashed lines enclose oenocytes. N = 10 biological replicates. (C) *mElo* expression in oenocytes following ISC-specific *Pvf1* overexpression. N = 4 biological replicates from 2 independent experiments. (D-F) Immunostaining and qPCR analysis reveal Hnf4-127D01 nuclear localization and *mElo* expression in oenocytes

overexpressing active *PvR*. (G-H) Immunostaining of Hnf4-127D01 in oenocytes of Yki flies

(*Esg>yki³SA;Luc-i*) and flies with knockdown of *Pvf1* in Yki flies (*Esg> yki³SA;Pvf1-i*). N = 10

biological replicates. (I) *mElo* expression in oenocytes of *Esg>yki³SA;Pvf1-i* flies, N = 4

biological replicates from 2 independent experiments. (J-K) Inhibition of *Pvf1* in the gut

affects bloating and ovary size in Yki flies, n = 15-20 flies. (L) ISC mitosis (pH3+ cells) in

Esg>yki³SA;Pvf1-i flies, n = 8-10 biological replicates. (M) Yki-induced *mElo* expression. Dual

LexA and Gal4 systems to express *yki³SA* in ISCs and mediate knockdown of *PvR* in

oenocytes (*Esg-LexA> yki³SA; PromE>PvR-i*). N = 2 biological replicates from 2 independent

experiments. (N-P) *PvR* knockdown in oenocytes and its effect on tumor-associated

bloating (N), ovary wasting (O), and pH3+ cell number (P), N = 10-20 flies. (Q) Lifespan

extension in Yki flies with oenocyte-specific *PvR* knockdown, N = 40 flies. Dashed lines

enclose oenocytes in all immunostaining images. Data are presented as mean ± s.e.m.;

statistical significance was assessed using Student's t-test or ANOVA.

Figure 3S: Expression of *Pvf1* in fly whole body and oenocytes in Yki flies.

UMAP from whole body (without head) snRNAseq, shows *Pvf1* expression level, oenocyte

cluster is marked as "oe". (B) snRNAseq analysis on oenocyte *Pvf1* expression in control

(*Esg>+)* and Yki flies (*Esg>yki³SA*) at 5 days. (C) snRNAseq analysis of oenocyte *Pvf1*

expression at 8 days of *yki³SA* induction.

Figure 4: Circulating lipids target trachea but not ISCs to promote tracheal growth.

Untargeted lipidomic analysis of fly hemolymph showing lipid species in control, Yki flies,

and Yki flies with oenocyte-specific *Hnf4* knockdown (*Esg>yki³SA;PromE>Hnf4-i*). N = 4

biological replicates. (B-C) Levels of very-long-acylcarnitines (AcCa 24:1 and AcCa 24:2) in

Yki flies with or without *Hnf4* knockdown in oenocytes. (D-E) Levels of circulating

phospholipids, including PC 36:4 PE 32:5, across experimental conditions. (F-G) Levels of

circulating TG 18:3_16:0_16:1 and TG 18:2_16:0_18:0 in the fly hemolymph. (H) Counts of

pH3+ cells in the midgut of flies with oenocyte-specific *TSC1*,2 overexpression or *Hnf4*

knockdown or overexpression. N = 8-20 flies. (I-K) Number of tracheal branches, tube area,

and skeleton length in Yki flies with ISC-specific knockdown of *LpR1* and *LpR2*. (L) pH3+

cell counts in ISCs of Yki flies with ISC-specific knockdown of *LpR1* and *LpR2*. (M-O) Effects

of trachea-specific knockdown of *LpR1* and *LpR2* using the *Btl-Gal4* driver on trachea

parameters, including number of branches, tube area, and skeleton length, compared to

control flies (*Btl>ctrl*). N = 8-12 flies. Dashed lines enclose oenocytes in all relevant images.

Data are presented as mean ± s.e.m.; statistical analysis was performed using Student's t-

test or ANOVA as appropriate.

Figure 4S: Effects of ISC-specific *LpR1* and *LpR2* knockdown on tracheal morphology

and ISC proliferation.

(A-C) Trachea parameters, including number of branches, tube area, and skeleton length,

in flies with ISC-specific knockdown of *LpR1* and *LpR2* using *Esg* driver, compared to

control (*Esg>ctrl*). N = 7-11 flies. (D) Counts of pH3+ cells in the midgut of flies with ISC-

specific knockdown of *LpR1* and *LpR2*, compared to controls. (E) Counts of pH3+ cells in

the midgut of flies with trachea-specific knockdown of *LpR1* and *LpR2*, compared to

controls. N=8-10 flies. Data are presented as mean ± s.e.m.; statistical analysis was

performed using Student's t-test or ANOVA as appropriate.

Figure 5: Hnf4-mElo axis regulates tracheogenesis in the midgut.

(A-B) Whole-body wax ester (WE) levels in flies with oenocyte-specific *Hnf4* knockdown

(*PromE^{gs}>Hnf4-i*), *mElo* knockdown (*PromE^{gs}>mElo-i*), or *TSC1,2* overexpression

(*PromE^{gs}>TSC1,2*). N = 3-4 biological replicates (C) Transepithelial barrier function of the

trachea on the midgut, assessed using red dextran injection, in flies with oenocyte-specific

Hnf4 knockdown. N = 8 flies. (D-G) Trachea morphology on the midgut (R2-R4), including

branch number and skeleton length normalized to gut surface area, in flies with *Hnf4*

knockdown in oenocytes. N = 8 flies. (H-J) Branch number, tube area, and skeleton length

of the midgut trachea in flies with oenocyte-specific *mElo* knockdown. N = 6-8 flies (K-P)

Number of branches, tracheal tube area and skeleton length of the midgut in flies

overexpressing *Hnf4* or *mElo* in oenocytes (*PromE^{gs}>Hnf4* or *PromE^{gs}>mElo*). N = 8-15 flies.

Figure 5S: Effect of oenocyte Hnf4 on ovary trachea.

(A-D) Tracheal branch number, tube area and total skeleton length of the ovary in flies with *Hnf4* knockdown in oenocytes. N = 8 flies. (E-G) Comparison of gut trachea morphology in wild type flies (*PromE^{gs}>+*) with and without RU feeding. N = 12 flies. Dashed lines enclose oenocytes in all relevant images. Data are presented as mean ± s.e.m.; statistical analysis was performed using Student's t-test or ANOVA as appropriate.

Figure 6: Hnf4-mElo axis in oenocytes mediates lipid metabolism and tracheogenesis induced by Yki tumors.

(A-C) Number of tracheal branches, total tube area, and total tube length in the midgut of Yki flies with oenocyte-specific *TSC1,2* overexpression, *Hnf4* knockdown, or *mElo* knockdown. N = 9-17 flies. (D-F) Tracheal morphology measurements, including number of branches and skeleton length, in flies with ISC-specific knockdown of *Pvf1* combined with *yki* overexpression (*Esg>yki^{3SA};Pvf1-i*). N = 9-11 flies. Dashed lines enclose oenocytes in all relevant images. Data are presented as mean ± s.e.m.; statistical analysis was performed using Student's t-test or ANOVA as appropriate.

Figure 6S: Hnf4-mElo axis in oenocytes mediates lipid metabolism and tracheogenesis induced by Yki tumors.

(A-D) Tracheal morphology parameters, including number of branches, total tube area, total tube length, and skeleton length, in Yki midgut (*Esg>yki*), compared to wild-type controls (*Esg>ctrl*). N = 16-17. (E) Whole-body wax ester (WE) levels in Yki flies with oenocyte-specific *Hnf4* knockdown (Esg-LexA>LexAop-yki^{3SA}; *PromE>Hnf4-i*) or *TSC1,2* overexpression (Esg-LexA>LexAop-yki^{3SA}; *PromE>TSC1,2*), measured to investigate the role of lipid metabolism in trachea growth. N = 4 biological replicates.

Figure 7: Regulation of hepatocyte lipid metabolism by pro-angiogenic factors in tumor models.

(A-E) Relative expression levels of *ELOVL* fatty acid elongase 1–7 in HepG2 cells incubated

with vascular endothelial growth factor A (VEGF-A) for 24 hours, compared to controls. N =

4 biological replicates, from 2 independent experiments. (F-G) Relative expression levels of

Hnf4a and *Elovl7* in liver samples from *KrasG12D/+;Lkb1f/f* (KL) tumor-bearing mice

compared to control mice. N = 4 biological replicates, from 2 independent experiments. (H)

Schematic representation of the proposed mechanism, depicting tumor-secreted

PDGF/VEGF-like factors activating the mTOR-Hnf4 axis in hepatocyte-like oenocytes to

promote lipid synthesis, which are then transported to trachea. Dashed lines indicate

hepatocyte regions or cells analyzed. Data are presented as mean ± s.e.m.; statistical

analysis was performed using Student's t-test.

References

1.   Anderson, N.M. and M.C. Simon, *The tumor microenvironment.* Curr Biol, 2020.
30(16): p. R921-r925.

2.   Chen, X., et al., *Neural Circuitries between the Brain and Peripheral Solid Tumors.*
Cancer Res, 2024. **84**(21): p. 3509-3521.

3.   Goncalves, M.D., et al., *Fenofibrate prevents skeletal muscle loss in mice with lung*
cancer. Proceedings of the National Academy of Sciences, 2018. **115**(4): p. E743-
E752.

4.   Schutz, Y., *Protein turnover, ureagenesis and gluconeogenesis.* Int J Vitam Nutr Res,
2011. **81**(2-3): p. 101-7.

5.   Lee, J.S., et al., *Urea Cycle Dysregulation Generates Clinically Relevant Genomic*
and Biochemical Signatures. Cell, 2018. **174**(6): p. 1559-1570 e22.

6.   Al-Zoughbi, W. and G. Hoefler, *Tumor Macroenvironment: An Update.* Pathobiology,
2020. **87**(2): p. 58-60.

7.   Jin, H.-R., et al., *Lipid metabolic reprogramming in tumor microenvironment: from*
mechanisms to therapeutics. Journal of Hematology & Oncology, 2023. **16**(1): p.
103.

8.   Wang, H., et al., *CD36-mediated metabolic adaptation supports regulatory T cell*
survival and function in tumors. Nat Immunol, 2020. **21**(3): p. 298-308.

9.   Wu, L., et al., *RIPK3 Orchestrates Fatty Acid Metabolism in Tumor-Associated*
Macrophages and Hepatocarcinogenesis. Cancer Immunol Res, 2020. **8**(5): p. 710-
721.

10.   Veglia, F., et al., *Fatty acid transport protein 2 reprograms neutrophils in cancer.*
Nature, 2019. **569**(7754): p. 73-78.

11.   Bilder, D., et al., *Tumour-host interactions through the lens of Drosophila.* Nat Rev
Cancer, 2021. **21**(11): p. 687-700.

12.   Droujinine, I.A. and N. Perrimon, *Interorgan Communication Pathways in Physiology:*
912   *Focus on Drosophila.* Annu Rev Genet, 2016. **50**: p. 539-570.

913   13.   Gutierrez, E., et al., *Specialized hepatocyte-like cells regulate Drosophila lipid*
914   *metabolism.* Nature, 2007. **445**(7125): p. 275-80.

915   14.   Chatterjee, D., et al., *Control of metabolic adaptation to fasting by dILP6-induced*
916   *insulin signaling in Drosophila oenocytes.* Proc Natl Acad Sci U S A, 2014. **111**(50):
917   p. 17959-64.

918   15.   Billeter, J.C., et al., *Specialized cells tag sexual and species identity in Drosophila*
919   *melanogaster.* Nature, 2009. **461**(7266): p. 987-91.

920   16.   Ferveur, J.F., et al., *Genetic feminization of pheromones and its behavioral*
921   *consequences in Drosophila males.* Science, 1997. **276**(5318): p. 1555-8.

922   17.   Blomquist, G.J., A.J. Chu, and S. Remaley, *Biosynthesis of wax in the honeybee, Apis*
923   *mellifera L.* Insect Biochemistry, 1980. **10**(3): p. 313-321.

924   18.   Qiu, Y., et al., *An insect-specific P450 oxidative decarbonylase for cuticular*
925   *hydrocarbon biosynthesis.* Proc Natl Acad Sci U S A, 2012. **109**(37): p. 14858-63.

926   19.   Reed, J.R., et al., *Unusual mechanism of hydrocarbon formation in the housefly:*
927   *cytochrome P450 converts aldehyde to the sex pheromone component (Z)-9-*
928   *tricosene and CO2.* Proceedings of the National Academy of Sciences, 1994. **91**(21):
929   p. 10000-10004.

930   20.   Wicker-Thomas, C., et al., *Flexible origin of hydrocarbon/pheromone precursors in*
931   *Drosophila melanogaster.* J Lipid Res, 2015. **56**(11): p. 2094-101.

932   21.   Wang, Z., et al., *Evolution of a fatty acyl-CoA elongase underlies desert adaptation in*
933   *Drosophila.* Sci Adv, 2023. **9**(35): p. eadg0328.

934   22.   Labeur, C., R. Dallerac, and C. Wicker-Thomas, *Involvement of desat1 gene in the*
935   *control of Drosophila melanogaster pheromone biosynthesis.* Genetica, 2002.
936   **114**(3): p. 269-74.

937   23.   Storelli, G., et al., *Drosophila HNF4 Directs a Switch in Lipid Metabolism that*
938   *Supports the Transition to Adulthood.* Dev Cell, 2019. **48**(2): p. 200-214 e6.

939   24.   Ghabrial, A., et al., *Branching morphogenesis of the Drosophila tracheal system.*
940   Annu Rev Cell Dev Biol, 2003. **19**: p. 623-47.

941   25.   Li, Z., et al., *Trachea-derived dpp controls adult midgut homeostasis in Drosophila.*
942   Dev Cell, 2013. **24**(2): p. 133-43.

943   26.   Tamamouna, V., et al., *Remodelling of oxygen-transporting tracheoles drives*
944   *intestinal regeneration and tumorigenesis in Drosophila.* Nat Cell Biol, 2021. **23**(5):
945   p. 497-510.

946   27.   Perochon, J., et al., *Dynamic adult tracheal plasticity drives stem cell adaptation to*
947   *changes in intestinal homeostasis in Drosophila.* Nat Cell Biol, 2021. **23**(5): p. 485-
948   496.

949   28.   Kwon, Y., et al., *Systemic organ wasting induced by localized expression of the*
950   *secreted insulin/IGF antagonist ImpL2.* Dev Cell, 2015. **33**(1): p. 36-46.

951   29.   Kwon, Y., et al., *Systemic Organ Wasting Induced by Localized Expression of the*
952   *Secreted Insulin/IGF Antagonist ImpL2.* Developmental Cell, 2015. **33**(1): p. 36-46.

953   30.   Patel, S., D.R. Nelson, and A.G. Gibbs, *Chemical and physical analyses of wax ester*
954   *properties.* J Insect Sci, 2001. **1**: p. 4.

31.   Palanker, L., et al., *Drosophila HNF4 Regulates Lipid Mobilization and β-Oxidation*.
Cell Metabolism, 2009. **9**(3): p. 228-239.
32.   Xu, J., et al., *Protein visualization and manipulation in Drosophila through the use of*
epitope tags recognized by nanobodies. eLife, 2022. **11**: p. e74326.
33.   Bradley, M.E., et al., *Potent and Efficacious Inhibition of CXCR2 Signaling by*
Biparatopic Nanobodies Combining Two Distinct Modes of Action. Molecular
Pharmacology, 2015. **87**(2): p. 251-262.
34.   Palanker, L., et al., *Drosophila HNF4 regulates lipid mobilization and beta-oxidation*.
Cell Metab, 2009. **9**(3): p. 228-39.
35.   Ghosh, A.C., et al., *Drosophila PDGF/VEGF signaling from muscles to hepatocyte-*
like cells protects against obesity. eLife, 2020. **9**: p. e56969.
36.   Zhang, Y., et al., *Rheb is a direct target of the tuberous sclerosis tumour suppressor*
proteins. Nature Cell Biology, 2003. **5**(6): p. 578-581.
37.   Sopko, R., et al., *A systems-level interrogation identifies regulators of Drosophila*
blood cell number and survival. PLoS Genet, 2015. **11**(3): p. e1005056.
38.   Tran, T.A., et al., *Platelet-derived growth factor/vascular endothelial growth factor*
receptor inactivation by sunitinib results in Tsc1/Tsc2-dependent inhibition of
TORC1. Mol Cell Biol, 2013. **33**(19): p. 3762-79.
39.   Song, W., et al., *Tumor-Derived Ligands Trigger Tumor Growth and Host Wasting via*
Differential MEK Activation. Dev Cell, 2019. **48**(2): p. 277-286 e6.
40.   Otsubo, C., et al., *Long-chain Acylcarnitines Reduce Lung Function by Inhibiting*
Pulmonary Surfactant*. Journal of Biological Chemistry, 2015. **290**(39): p. 23897-
23904.
41.   Wolrab, D., et al., *Oncolipidomics: Mass spectrometric quantitation of lipids in*
cancer research. TrAC Trends in Analytical Chemistry, 2019. **120**: p. 115480.
42.   Cruciani, G., et al., *Diagnostic application of lipidomics fingerprints to bladder*
carcinoma. Translational Cancer Research, 2017. **6**(6): p. 1197-1206.
43.   Palm, W., et al., *Lipoproteins in Drosophila melanogaster--assembly, function, and*
influence on tissue lipid composition. PLoS Genet, 2012. **8**(7): p. e1002828.
44.   Khaliullina, H., et al., *Patched regulates Smoothened trafficking using lipoprotein-*
derived lipids. Development, 2009. **136**(24): p. 4111-21.
45.   Locke, M., *Pore canals and related structures in insect cuticle*. J Biophys Biochem
Cytol, 1961. **10**(4): p. 589-618.
46.   Moussian, B., *Recent advances in understanding mechanisms of insect cuticle*
differentiation. Insect Biochem Mol Biol, 2010. **40**(5): p. 363-75.
47.   Moussian, B., et al., *Cuticle differentiation during Drosophila embryogenesis*.
Arthropod Struct Dev, 2006. **35**(3): p. 137-52.
48.   Jaspers, M.H., et al., *The fatty acyl-CoA reductase Waterproof mediates airway*
clearance in Drosophila. Dev Biol, 2014. **385**(1): p. 23-31.
49.   Ding, G., et al., *Coordination of tumor growth and host wasting by tumor-derived*
Upd3. Cell Rep, 2021. **36**(7): p. 109553.
50.   Xu, J., et al., *Mechanistic characterization of a Drosophila model of paraneoplastic*
nephrotic syndrome. Nature Communications, 2024. **15**(1): p. 1241.

51.   Duvel, K., et al., *Activation of a metabolic gene regulatory network downstream of*
999   *mTOR complex 1*. Mol Cell, 2010. **39**(2): p. 171-83.
52.   Tiebe, M., et al., *REPTOR and REPTOR-BP Regulate Organismal Metabolism and*
Transcription Downstream of TORC1. Dev Cell, 2015. **33**(3): p. 272-84.
53.   Jiang, H., et al., *Cytokine/Jak/Stat signaling mediates regeneration and homeostasis*
in the Drosophila midgut. Cell, 2009. **137**(7): p. 1343-55.
54.   Buchon, N., et al., *Drosophila intestinal response to bacterial infection: activation of*
host defense and stem cell proliferation. Cell Host Microbe, 2009. **5**(2): p. 200-11.
55.   Apidianakis, Y., et al., *Synergy between bacterial infection and genetic*
predisposition in intestinal dysplasia. Proc Natl Acad Sci U S A, 2009. **106**(49): p.
20883-8.
56.   Amcheslavsky, A., et al., *Enteroendocrine Cells Support Intestinal Stem-Cell-*
Mediated Homeostasis in Drosophila. Cell Reports, 2014. **9**(1): p. 32-39.
57.   Lin, G., N. Xu, and R. Xi, *Paracrine Wingless signalling controls self-renewal of*
Drosophila intestinal stem cells. Nature, 2008. **455**(7216): p. 1119-1123.
58.   Schönfeld, P. and G. Reiser, *Brain Lipotoxicity of Phytanic Acid and Very Long-chain*
Fatty Acids. Harmful Cellular/Mitochondrial Activities in Refsum Disease and X-
Linked Adrenoleukodystrophy. Aging Dis, 2016. **7**(2): p. 136-49.
59.   Baarine, M., et al., *Evidence of oxidative stress in very long chain fatty acid--treated*
oligodendrocytes and potentialization of ROS production using RNA interference-
directed knockdown of ABCD1 and ACOX1 peroxisomal proteins. Neuroscience,
2012. **213**: p. 1-18.
60.   Zarrouk, A., et al., *Induction of mitochondrial changes associated with oxidative*
stress on very long chain fatty acids (C22:0, C24:0, or C26:0)-treated human
neuronal cells (SK-NB-E). Oxid Med Cell Longev, 2012. **2012**: p. 623257.
61.   Tamura, K., et al., *Novel lipogenic enzyme ELOVL7 is involved in prostate cancer*
growth through saturated long-chain fatty acid metabolism. Cancer Res, 2009.
69(20): p. 8133-40.
62.   Karlsson, M., et al., *A single–cell type transcriptomics map of human tissues*.
Science Advances, 2021. **7**(31): p. eabh2169.
63.   Koyama, T. and C.K. Mirth, *Growth-Blocking Peptides As Nutrition-Sensitive Signals*
for Insulin Secretion and Body Size Regulation. PLoS Biol, 2016. **14**(2): p. e1002392.
64.   Pallares-Cartes, C., G. Cakan-Akdogan, and Aurelio A. Teleman, *Tissue-Specific*
Coupling between Insulin/IGF and TORC1 Signaling via PRAS40 in
Drosophila. Developmental Cell, 2012. **22**(1): p. 172-182.
65.   Chatterjee, D., et al., *Control of metabolic adaptation to fasting by dILP6-induced*
insulin signaling in <i>Drosophila</i> oenocytes. Proceedings of the National
Academy of Sciences, 2014. **111**(50): p. 17959-17964.
66.   Chung, H., et al., *A Single Gene Affects Both Ecological Divergence and Mate*
Choice in <i>Drosophila</i>. Science, 2014. **343**(6175): p. 1148-1151.
67.   Wang, Z., et al., *Evolution of a fatty acyl–CoA elongase underlies desert*
adaptation in <i>Drosophila</i>. Science Advances, 2023. **9**(35): p. eadg0328.

68.   Montoya-Zegarra, J.A., et al., *AutoTube: a novel software for the automated*
morphometric analysis of vascular networks in tissues. Angiogenesis, 2019. **22**(2):
p. 223-236.
69.   Ji, H., et al., *LKB1 modulates lung cancer differentiation and metastasis.* Nature,
2007. **448**(7155): p. 807-810.
1045

A

B

PromE^{gs}> PRAS40

-RU   +RU

C

DMSO   Rapamycin 200uM

D

A RU

MtdTomato

PromE^{gs}>Hnf4-i;
QF6>MtdTomato

−

+

B Ovary
# of branches

ns

Arbitary unit / area

1500

1000

RU   −   +

C Ovary
Tube area

*

Arbitary unit / area

$2.5×10^5$

$2.0×10^5$

$1.5×10^5$

$1.0×10^5$

$5.0×10^4$

0.0

−   +

D Ovary
Skeleton length

ns

Arbitary unit / area

$4.0×10^4$

$3.5×10^4$

$3.0×10^4$

$2.5×10^4$

$2.0×10^4$

$1.5×10^4$

−   +

PromE^{gs}>Hnf4-i;
QF6>MtdTomato

E # of branches

ns

Arbitrary unit / area

4000

3000

2000

1000

RU   −   +

F Tube area

ns

Arbitrary unit / area

$6.0×10^5$

$5.5×10^5$

$5.0×10^5$

$4.5×10^5$

$4.0×10^5$

$3.5×10^5$

$3.0×10^5$

−   +

G Skeleton length

ns

Arbitrary unit / area

$1×10^5$

$9×10^4$

$8×10^4$

$7×10^4$

$6×10^4$

$5×10^4$

−   +

PromE^{gs}>ctrl