1 **Lipid metabolism of hepatocyte-like cells supports intestinal tumor growth in**
2 ***Drosophila***
3

4 Huang, K.*[1], Miao, T.*[1], Dantas, E.[3], Sanford, J.[3], Chen, Y. [1], Hu, Y.[1,2], Moon, SJ. [1], Han, M.[1,2],
5 Wang, K.[1], Goncalves, M.[3], and Perrimon, N[1,2]
6
7 1 Department of Genetics, Blavatnik Institute, Harvard Medical School, Harvard University,
8  Boston, MA 02115, USA
9 2 Howard Hughes Medical Institute, Boston, MA, USA
10 3 NYU Langone Health, New York, NY, USA.
11
12 ^Correspondence:
13 Kerui Huang: Kerui_Huang@hms.harvard.edu
14 Norbert Perrimon: perrimon@genetics.med.harvard.edu
15
*Authors contribute equally

**Abstract**

Tumors require metabolic adaptations to support their rapid growth, but how they

influence lipid metabolism in distant tissues remains poorly understood. Here, we uncover

a novel mechanism by which gut tumors in adult flies reprogram lipid metabolism in distal

hepatocyte-like cells, known as oenocytes, to promote tracheal development and tumor

growth. We show that tumors secrete a PDGF/VEGF-like factor, Pvf1, that activates the

TORC1-Hnf4 signaling pathway in oenocytes. This activation enhances the production of

specific lipids, including very long chain fatty acids and wax esters, that are required for

tracheal growth surrounding the gut tumor. Importantly, reducing expression in oenocytes

of either the transcription factor *Hnf4*, or the elongase *mElo* that generates very long chain

fatty acid robustly extends lifespan, suppresses tumor growth, tracheogenesis, and

associated organ wasting/cachexia-like phenotypes. Further, altering the expression level

of oenocyte *Hnf4* or *mElo* in control flies affects intestinal tracheogenesis. Lipoprotein-

mediated lipid transport and altered oenocyte lipid metabolism drive the observed

changes in tracheogenesis. Finally, we demonstrate that this regulatory pathway is

conserved in mammals, as VEGF-A stimulates lipid metabolism gene expression in human

hepatocytes, and lung tumor-bearing mice show increased hepatic expression of *Hnf4* and

the lipid elongation gene *Elovl7*. Our findings reveal a previously unrecognized tumor-host

interaction where tumors non-autonomously reprogram distal lipid metabolism to support

their growth. This study not only identifies a novel non-autonomous role of the TORC1-Hnf4

axis in lipid-mediated tumor progression but also highlights potential targets for

therapeutic intervention in cancer-associated metabolic disorders.

**Introduction**

Tumor growth demands a growing supply of nutrients and oxygen. The growing

tumor constantly shapes its microenvironment to favor fast growth and expansion, through

secreted factors and extracellular matrix [1-4]. The composition of the tumor

microenvironment (TME) varies between tumor types, but the hallmarks include increased

blood vessels, immune cells, stromal cells, and extracellular matrix [5]. To overcome the

increasingly hypoxic and acidic environment, TME promotes angiogenesis to restore

oxygen and nutrient supply. In addition to modulating the local TME, malignant cells can

exert systemic effects on distant organs—including the liver, skeletal muscle, and brain—to

reshape the host's metabolic pathways, ultimately promoting further tumor progression [6-

10]. Liver is a metabolic center for amino acids, carbohydrates, and lipid. A key liver-

specific metabolic function is the complete urea cycle (UC), which eliminates excess

ammonia by converting into urea, a less toxic waste. Previous studies have identified

evidence of decreased UC activity in the livers of 4T1 breast cancer bearing mice, and the

plasma of children with cancer [9], demonstrating an existing interaction between tumors

and the liver [11-15]. Delineating molecular pathways involving tumor-host interaction can

provide novel therapeutic targets.

Lipids play a critical role in cancer progression, by serving as building blocks, energy

sources and signaling molecules. It has been observed that tumor cells reprogram lipid

metabolism to enhance growth, characterized by increased lipid synthesis, lipid uptake,

fatty acid oxidation (FAO) and lipid storage [16]. Tumor cells also actively reshape lipid

metabolism in cells from the TME through secreted signaling molecules and drive TME to

favor tumor growth. For example, increased lipid uptake and FAO in regulatory T cells

(Tregs), tumor-associated macrophages (TAMs) and myeloid-derived suppressor cells

(MDSCs), promote their immunosuppressive function [17-19]. However, whether tumor

cells also reshape lipid metabolism in distal organs, such as liver, which is a metabolic

center for lipid transport, is not known.

The adult *Drosophila* midgut is a powerful model for investigating tumorigenesis and

tumor-driven metabolic remodeling. Many genetic alterations associated with human

cancers induce hyperproliferation of fly intestinal stem cells (ISCs) and lead to tumor

formation in the midgut [20].  In particular, expression of constitutively active Yorkie

(yki$^{3SA}$)—the *Drosophila* homolog of human YAP1—drives gut tumor development and

triggers cachexia-like phenotypes, including lipid depletion, hyperglycemia, and wasting of

muscle and ovaries [21, 22]. Mechanistically, yki$^{3SA}$ gut tumors secrete multiple systemic

factors, such as the IGF-antagonizing peptide *ImpL2* [21], *PDGF- and VEGF-related factor 1*

*(Pvf1)* [22] and unpaired 3 *(Upd3)* , a functional ortholog of mammalian interleukin-6 (IL-6)

[22]. Notably, IL-6- and PDGF/VEGF-associated inflammatory programs are also broadly

elevated across diverse cancer models [23-26]. Together, these observations highlight

tumor–host organ communication as an evolutionarily conserved process and establish

*Drosophila* as an excellent system to dissect tumor–host interactions *in vivo*.

As *Drosophila* has emerged as a simple yet powerful model to study tumor-host

interactions [27], we use this model to better understand the reciprocal communication

between liver and tumor cells. The fly midgut, oenocytes, fat body and Malpighian tubules

are functionally equivalent to human digestive tract, hepatocytes, adipose tissue and

kidney, respectively [28]. Oenocytes, which are hepatocyte-like cells, are important for

mobilizing stored lipids during starvation and interact closely with metabolic tissues, such

as the fat body to regulate lipid uptake and reprocessing [29, 30]. In addition to regulating

lipid metabolism under nutritional deprivation, oenocytes are the major site for synthesis

of cuticular hydrocarbons in pterygote insects [31-33]. Adult oenocytes synthesize

hydrocarbons using fatty acyl-CoA precursors via reduction to fatty aldehyde and oxidative

decarbonylation [34, 35]. Many fatty acid synthases, elongases and desaturases expressed

in adult oenocytes produce various types of fatty acyl-CoA substrates that are processed to

form a complex blend of cuticular hydrocarbons [36-39]. Hepatocyte nuclear factor 4

(Hnf4), an oenocyte enriched transcription factor, plays an important role in regulating the conversion of lipids to very long chain fatty acid, which are important substrates for hydrocarbon synthesis [39]. Oenocytes are the major site for lipid transport and processing, reminiscent of human hepatocytes. However, whether and how oenocytes regulate tumor growth is unknown.

The *Drosophila* trachea is a network of oxygen transporting tubes, the functional equivalent of mammalian blood vessels [40]. Trachea is composed of terminal tracheal cells (TTCs), which are highly plastic cells, and the terminal branches (TBs) or tracheoles which deliver oxygen to the tissues. TTCs are oxygen sensing cells, and hypoxic conditions promote the formation of TBs through hypoxia-inducible factor-1$\alpha$ (HIF-1$\alpha$; Similar (Sima) in *Drosophila*), similar to the role of HIF-1$\alpha$ in mammalian angiogenesis. Trachea resides in proximity of many tissues, including the midgut. Trachea penetrate the visceral muscles to access the intestinal epithelium to supply it with oxygen [41]. The intestinal epithelium is apicobasally polarized, composed of ISCs and their progeny. Damage induced by enteric infection, oxidative agents and tumor growth can increase tracheogenesis on the midgut, which is necessary for efficient damage-induced ISC-mediated regeneration [42, 43]. Additionally, tumor-induced tracheogenesis in *Drosophila* is reminiscent of cancer-induced angiogenesis in mammals [42].

Using a Drosophila gut tumor model, we uncover a novel mechanism through which gut tumors reprogram lipid metabolism in distal oenocytes to promote both tracheal development and tumor growth. We find that tumors secrete a PDGF/VEGF-like factor, Pvf1, which activates the TORC1-Hnf4 signaling pathway in oenocytes. This activation boosts the production of specific lipids, including very long-chain fatty acids and wax esters, essential for tracheal growth around the tumor. Notably, reducing the activity of Hnf4 and mElo, an elongase involved in generating very long-chain fatty acids, in oenocytes suppress tumor growth, tracheal development, and cachexia-like phenotypes, while also extending lifespan. Additionally, we demonstrate that this regulatory pathway is conserved in mammals, as VEGF-A stimulates lipid metabolism gene expression in human hepatocytes, and tumor-bearing mice exhibit increased hepatic expression of lipid

123    elongation genes. Our findings highlight a previously unrecognized tumor-host interaction,

124    where tumors non-autonomously reprogram distal lipid metabolism to facilitate their

125    growth.

126    **Results**

127    ***Oenocyte lipid metabolism regulated by Hnf4 controls tumor proliferation***

128           To investigate how tumor growth rewires lipid metabolism of the whole fly, we

expressed an active form of *Yorkie ( yki³ˢᴬ),* in adult gut stem cells, using the intestinal stem

cells (ISCs)/enteroblast driver (*Esg-Gal4* or *Esg-LexA*), hereafter referred to as Yki flies [21].

*Esg-Gal4* or *Esg-LexA* was combined with the *tubulin* promoter-driven temperature-

sensitive *GAL80 (tubP-GAL80ᵀˢ)* to control the temporal expression of Gal4 or LexA. Next,

we conducted untargeted lipidomic profiling using an LC-MS/MS-based platform of whole-

body samples of adult female flies (*Esg-LexA > LexAop-yki³ˢᴬ*) expressing *yki³ˢᴬ* for 3 days,

together with control flies (*Esg > +*). The complete *Drosophila* genotype information can be

found in Supplementary Table 1. Female flies are used in the analysis unless otherwise

noted because they show more significant and consistent tumor-associated cachexia

phenotypes. We detected 2471 lipid species in total representing 45 lipid classes

(Supplementary Table 2). The lipidomic analysis showed an increase in wax ester (WE),

monogalactosyldiacylglycerol (MGDG), phosphatidylglycerol (PG), phosphatidylcholine

(PC), ceramide (Cer) and acylcarnitine (AcCa) levels in Yki flies (Figure 1A). We also

observed a slight reduction of triglycerides (TG) in Yki flies at early stage, in agreement with

previous findings [21, 44]. WEs are major components of cuticular lipids, which cover

nearly all parts of terrestrial arthropods [45] to restrict water loss, and WE levels are

significantly increased in Yki flies (Figure 1B), yet the significance of this regulation is

unknown. By profiling the fatty acid carbon chain length of WEs, we found that Yki flies

exhibit elevated levels of WEs containing very-long-chain fatty acids (carbon length ≥22),

as well as WEs with shorter chains (<22) (Figure 1C). As the biosynthesis of cuticular lipids

mainly occurs in *Drosophila* oenocytes and Hnf4 is a main regulator of this process [39],

we tested whether oenocyte Hnf4 regulates the level of WEs in control flies. WEs were

slightly reduced when *Hnf4* was knocked down in control adult oenocytes using the

*PromE(800)-Gal4* temperature sensitive driver (*PromE-GAL4,Tub-GAL80$^{TS}$*, referred to as

*PromE*). PromE(800)-Gal4 is specifically expressed in adult female oenocytes, although it is

also expressed in the reproductive organs of adult males [31]. To ensure tissue specificity,

female flies were used in all experiments unless otherwise noted. Next, to test whether

Hnf4 regulates WE synthesis in Yki flies, we used the LexA-LexAop system to induce Yki

tumors in the gut and the Gal4/UAS system in oenocytes (*PromE-GAL4, Tub-GAL80$^{TS}$*,

referred to as *PromE*) to knockdown *Hnf4*. Knockdown of *Hnf4* in oenocytes decreased Yki-

induced WE levels (Figure 1D), indicating that *Hnf4* expression in oenocytes is required for

the increase of WE.

As a nuclear receptor, the localization of Hnf4 is highly enriched in oenocyte nuclei

compared to the fat body [39, 46]. To visualize the localization of Hnf4, we tagged the

endogenous *Hnf4* gene with 127D01, a small epitope tag that can be detected with a

specific nanobody [47, 48], using an homology directed repair pathway CRISPR-Cas9

targeted insertion method [47].  To validate the Hnf4-127D01 line, we knocked down *Hnf4* in

oenocytes, which significantly reduced its nuclear localization (Figure S1A). Hnf4-127D01

in control oenocytes showed high level of nuclear localization, consistent with previous

findings (Figure 1E)  [46]. Interestingly, increased nuclear localization of Hnf4-127D01 was

observed in the oenocytes of Yki flies (Figure 1E-1F).  Altogether, these data suggest that

oenocytes in Yki flies have increased Hnf4 activity. To test whether oenocyte Hnf4 signaling

is required for tumor growth, we measured ISC-mitosis using phospho-histone 3 (pH3+)

staining.  Downregulation of *Hnf4* in oenocytes of Yki flies *(Esg-LexA>yki$^{3SA}$-GFP;*

*PromE>Hnf4-i)* for 3 days showed a decrease in pH3+ staining compared to controls (*Esg>*

*UAS-yki$^{3SA}$-GFP*), and a slower growth rate for tumor (Figure 1G). Importantly, *Hnf4*

knockdown in wild-type animals did not alter pH3+ cell numbers relative to controls (Figure

1G), indicating that the requirement for Hnf4 in Yki-driven tumors reflects its tumor-

associated mis-regulation rather than a general role in proliferation. To quantify tumor

burden, we measured the percentage of GFP-positive area in the gut and found that

oenocyte-specific depletion of *Hnf4* in Yki flies markedly reduced GFP-positive area,

consistent with tumor regression (Figure S1B-C). We next assessed tumor-induced

cachexia phenotypes, including ovary wasting and abdominal bloating caused by fluid

accumulation (Figure S1D). Short-term oenocyte *Hnf4* knockdown (3 days) alleviated ovary

wasting (Figure S1E), reduced bloating (Figure 1H) and decreased whole-body wet weight

(Figure S1F), supporting a role for aberrant oenocyte Hnf4 activity in promoting tumor

growth and associated cachexia.

Our results suggest that Hnf4 transcriptional activity is induced in oenocytes of Yki

flies. To validate this, we measured whether the expression of Hnf4 targets in Yki oenocytes

also increased. Hnf4 target genes are involved in beta-oxidation and very long chain fatty

acid (VLCFA) biosynthesis in oenocytes [39, 49]. Thus, we tested whether VLCFA

biosynthesis genes (*Hnf4, mElo, CG30008, CG16904, FarO, ACC, HDAC, TER, Cyp4g1,*

*Cpr*)*,* as well as genes involved in beta-oxidation (*Whd, Scu, Yip2* and *CG7461)* were

dysregulated in Yki flies. *CG30008, CG16904, FarO, ACC, HDAC, TER, Cyp4g1, Cpr, Whd,*

*Scu, Yip2* showed no difference between control and Yki flies (Figure S1G). However, we

observed a significant increase in *mElo* (Figure 1I)*,* which encodes a methyl-branched

cuticular hydrocarbon elongase [37], and a minor increase for *Hnf4 and CG7461* (Figure

S1G).

*mElo* has recently been shown to produce long chain methyl-branched cuticular

hydrocarbons in oenocytes that help prevent desiccation [37], a key function of Hnf4 in

oenocytes [39]. Additionally, *mElo* expression is highly enriched in control adult oenocytes

according to snRNAseq data (Figure 1J). Inhibition of *Hnf4* reduced *mElo* expression, and

overexpression of *Hnf4* led to increased levels of *mElo* (Figure S1H-I), consistent with a

previous report that *mElo* is a transcriptional target of Hnf4 [39]. Further*, mElo* is highly

induced in the oenocytes of Yki flies and this induction is blocked when *Hnf4* is knocked

down (Figure 1I).

In control flies, reduction of *mElo* in oenocytes did not affect pH3+ number in the

gut (Figure 1K). To test the function of oenocyte *mElo* during tumor growth, we reduced the

level of oenocyte *mElo* in Yki flies and observed fewer pH3+ in the gut, suggesting that *mElo*

contributes to Yki tumor growth (Figure 1K). Similarly, we observed a reduced tumor burden

with *mElo* knockdown in Yki flies (Figure S1J). We validated the effectiveness of *mElo*

knockdown in the oenocytes (Figure S1K). Unlike *Hnf4,* reducing *mElo* expression in control

oenocytes using the gene-switch driver (hereafter referred to as PromE[gs]) [30], where the

driver is activated upon mifepristone feeding (RU), did not cause steatosis and affect ovary

size, suggesting that other targets of Hnf4 regulate oenocytes steatosis and ovary size

(Figure S1L and Figure S1M).  Consistently, *mElo* knockdown did not rescue ovary wasting

in Yki flies (Figure S1N).

To assess whether oenocyte lipid metabolism affects the viability of Yki flies, we

inhibited *Hnf4* in oenocytes and observed a significant lifespan extension (Figure

1L).Inhibition of mElo in oenocytes of Yki flies also significantly increased lifespan (Figure

1M). Finally, *mElo* has a sex specific function in regulating different types of hydrocarbons

[37]. Therefore, we tested whether *mElo* also regulates Yki viability in male flies. Strikingly,

inhibition of *mElo* also significantly increased the lifespan in Yki males (Figure S1O).

***Oenocyte Hnf4 activity is regulated by TORC1***

We next sought to determine the factors regulating Hnf4 level in oenocytes. Previous

studies have shown that inhibiting the TORC1 pathway in oenocytes leads to steatosis [50],

a phenotype that mimics reduced *Hnf4* expression in these cells [39], suggesting that the

TORC1 pathway positively regulates *Hnf4* transcription in oenocytes. The Tuberous

Sclerosis Complex (TSC) is known to negatively regulate TORC1 by inhibiting RAS homolog

enriched in brain (Rheb) activity [51]. Next, we also tested the effects of oenocyte *TSC1,2* in

tumor growth. Consistent with *Hnf4* and *mElo* knockdown, *TSC1,2* overexpression in

oenocytes reduced tumor induced WE levels (Figure 2A). We also observed an increased

level of TORC1 activity, as measured by phospho-4EBP (p4EBP) staining in oenocytes of Yki

flies (Figure 2B-C). Additionally, overexpressing *TSC1,2* blocked *mElo* expression in

oenocytes of Yki flies (Figure 2D), Yki-induced pH3[+] cells in the gut (Figure 2E), as well as

Yki-induced ovary wasting and bloating (Figure S2A and S2B). Together, these data identify

oenocyte TORC1 activation as a key driver of tumor growth and implicate Hnf4 as a

downstream effector.

To test whether TORC1 regulates *Hnf4* expression level, we analyzed a previously

published snRNAseq dataset [50]. Overexpression of *Tsc1* together with *Tsc2* (*TSC1*,2) in

oenocytes, which inhibits *TORC1*, resulted in reduced *Hnf4* expression specifically in

oenocytes (Figure 2F), but not in muscle nor fat body tissues, suggesting that TORC1

positively regulates Hnf4 levels in oenocytes. Next, we examined the protein level of

endogenously tagged Hnf4-127D01 following *TSC1,2* over-expression and observed a

significant reduction in Hnf4-127D01 levels in oenocytes, particularly in the nucleus as

Hnf4 is a transcription factor (Figure 2G-H). Given that Hnf4 regulates hydrocarbon

synthesis in oenocytes and its knockdown increases the susceptibility to dry starvation

[39], we overexpressed *TSC1,2* in oenocytes and found that it led to an increased sensitivity

of flies to dry starvation (Figure S2C). Additionally, as Hnf4 regulates the expression of

genes involved in very-long-chain fatty acid (VLCFA) synthesis [39], we analyzed the

expression of VLCFA synthesis genes in snRNAseq data from control (*PromE>EGFP*) versus

TSC1,2-overexpressing oenocytes (*PromE>TSC1,2*). Using uniform Manifold Approximation

and Projection (UMAP), which identifies cell types in mixed populations based on

snRNAseq data, we found that VLCFA synthesis genes were highly enriched in oenocytes

(Figure 2I). The ModuleScore, calculated by averaging the expression of the VLCFA

synthesis pathway genes and subtracting the expression of a randomly selected gene set

expression, decreased in oenocytes upon *TSC1,2* overexpression (Figure 2I).

Inhibition of TORC1 through overexpressing *TSC1,2,* or overexpressing Proline-rich

Akt substrate 40 kDa (*PRAS40)*, or by feeding flies with rapamycin, all induced steatosis in

oenocytes (Figure 2J and S2D-E), as observed following *Hnf4* knockdown in oenocytes

(Figure S1L). However, simultaneous overexpression of *Hnf4* with *TSC1*,2 prevented the

development of steatosis (Figure 2J), indicating that TORC1 regulates oenocyte lipid

metabolism via Hnf4. Lastly, *TSC1,2* overexpression also reduced the expression of *mElo*

and *FASN2,* which are Hnf4 target genes, and overexpressing *Hnf4* simultaneously blocked

this decrease (Figure 2K-L).  Together, these findings suggest that Hnf4 acts downstream of

TORC1 to regulate lipid metabolism in oenocytes.

**267**     ***ISCs secreted Pvf1 regulates Hnf4 through PvR signaling in oenocytes and affects***

**268**     ***tumor associated phenotypes***

**269**          TORC1 activity is regulated through various pathways, one of which includes

**270**     PDGF/VEGF signaling. Studies have shown that PvR activation can stimulate the TOR

**271**     pathway [50, 52, 53]. Consistent with this, muscle-secreted Pvf1 can regulate TORC1

**272**     activity in oenocytes, suggesting that the activation of the TORC1-Hnf4 axis in oenocytes of

**273**     Yki flies may occur through PvR activation. Yki-expressing ISCs can secrete higher levels of

**274**     the Pvf1 ligand, which activates PvR signaling in surrounding tissues (Figure S3A) [22].

**275**     However, *Pvf1* is expressed at low levels in oenocytes and is not significantly induced in

**276**     oenocytes and muscles of Yki flies (Figure S3A-B). To determine whether tumor secreted

**277**     Pvf1 non-autonomously activates Hnf4 level in oenocytes, we knocked down *Pvf1*

**278**     expression in the ISCs of Yki flies (*Esg>yki^3SA; Pvf1-i*) and observed its effect on Hnf4 nuclear

**279**     level in oenocytes. *Esg>yki^3SA; Pvf1-i* significantly reduced the level of Hnf4 nuclear

**280**     localization compared to *Yki* flies (*Esg>yki^3SA; Luc-i)* (Figure 3A-B). Consistently, *mElo*

**281**     expression was reduced in *Esg>yki^3SA; Pvf1-i* flies compared to Yki flies (Figure 3C). We also

**282**     tested whether *Pvf1* reduction in Yki flies affects tumor development, like *Hnf4* inhibition in

**283**     oenocytes. As reported previously [22], inhibition of *Pvf1* in the tumor gut rescues both

**284**     bloating and ovary wasting (Figure 3D and Figure S3C). Gut tumors in *Esg>yki^3SA;Pvf1-i* flies

**285**     exhibited slower growth at early stage, as shown by reduced number of pH3+ cells and less

**286**     GFP positive area (Figure 3E and Figure S3D).

**287**          To demonstrate that Pvf1 secreted from Yki tumors directly controls *mElo*

**288**     expression through PvR, we used the dual Gal4 and LexA systems to reduce *PvR* level in

**289**     oenocytes while inducing Yki in ISCs. *Esg-LexA>yki^3SA;PromE>PvR-i* animals showed

**290**     reduced *mElo* expression (Figure 3F), decreased the level of bloating (Figure 3G) and ovary

**291**     wasting (Figure S3E), reduced the number of pH3+ cells in gut tumors (Figure 3H) and

**292**     reduced tumor burden (Figure S3F), and extended lifespan compared to Yki flies (Figure 3I).

**293**     Altogether, these data suggest that PDGF/VEGF induced by Yki tumor non-autonomously

**294**     increases Hnf4 activity in oenocytes through PvR.

To test whether ISC secreted Pvf1 can induce oenocyte Hnf4 levels under normal

physiology, we overexpressed *Pvf1* specifically in adult ISCs (*Esg>Pvf1*) and measured Hnf4

nuclear level in oenocytes. *Pvf1* overexpression in ISCs increased nuclear Hnf4 levels

(Figure 3J-K), suggesting an increased transcriptional activity as measured by increased

level of *mElo* expression (Figure 3L). To test whether ISC-derived Pvf1 activates PvR in

oenocytes, we overexpressed an active form of PvR in oenocytes which was associated

with increased nuclear Hnf4 level, as well as increased level of *mElo* expression (Figure

3O).

***Circulating lipids target trachea but not ISCs to promote trachea growth***

To identify the lipid species induced by oenocytes in Yki flies*,* we performed

lipidomic analysis on the fly hemolymph of control flies, *Esg-LexA> yki³ˢᴬ* and *Esg-LexA>*

*yki³ˢᴬ; PromE>Hnf4-i* flies (Figure 4A, Supplementary Table 3). Among the lipid species that

are induced by Yki and decreased by *Esg-LexA > yki³ˢᴬ; PromE>Hnf4-i*, the majority are

phospholipids including phosphotidyl choline (PC), phosphatidylglycerol (PG) and

phosphatidylethanolamine (PE), triglycerides (TGs) and acylcarnitines (AcCa). More

specifically, AcCa 24:1 and AcCa 24:2, which are very-long-chain acylcarnitines ($C \geq 22$)

secreted by mitochondria when metabolism is inhibited, were elevated in Yki flies while

reduced by *Hnf4* knockdown (Figure 4B-C). Circulating PC 36:4, and PE 32:5 phospholipids

and TG(18:3_16:0_16:1)*,* TG(18:2_16:0_18:0) triglycerides were induced in Yki flies and

rescued by *Hnf4* knockdown (Figure 4D-G). Interestingly, PC 36:4 is elevated in the serum

and plasma samples from breast cancer patients  and TG(18:2_16:0_18:0)  is induced in

bladder cancer patients [54].

PC and PE are major membrane phospholipids and contribute to lipoprotein

assembly and lipid transport. Thus, elevated PC/PE could promote tracheogenesis by

supporting both membrane biogenesis and lipid delivery needed for tracheal membrane

expansion. Increased AcCas and WEs could contribute to tracheogenesis by supporting

local energetic demand or increasing the production of reactive oxygen species (ROS). For

example, hepatocyte-derived circulating AcCas are required for systemic thermogenesis

during cold exposure in mice; in this setting, peripheral tissues—including brown adipose

tissue, skeletal muscle, and heart—take up plasma acylcarnitines and oxidize them to

support energy metabolism [55].  In insects, lipid transport in circulation primarily relies on

lipophorins (Lpps), which are the major lipoproteins. It is produced predominantly by the

fat body and delivers neutral lipids to peripheral tissues via the lipophorin receptors

LpR1/LpR2, which mediate lipoprotein binding and internalization through endocytic

pathways [56, 57]. The increased circulation of PC 36:4 and PE 32:5 in Yki flies suggests an

increase in lipoproteins that transport lipids from oenocytes towards trachea or the midgut.

332           The transport of lipids, such as phospholipids, from oenocytes to the trachea or

ISCs to promote tumor growth has not been reported [42]. To test whether changes in

oenocyte metabolism regulates ISCs growth,  we inhibited Hnf4 activity in oenocytes by

overexpressing *TSC1,2*, knocking down or overexpressing *Hnf4*, and observed no significant

effect on the number of pH3+ cells in the midgut (Figure S4A). This suggests that changes

in oenocyte lipid metabolism does not promote ISC proliferation under normal condition.

Since 95% of the hemolymph lipids are transported through lipophorin (Lpp) in flies [56],

and that low-density lipoprotein (LDL) receptor homologues LpR1 and LpR2 promote Lpp

uptake [58], we reduced LpR1 or LpR2 level in the ISCs. However, this had no impact on the

trachea area, skeleton length, and only a minor increase in the number of branches in the

*LpR1* knockdown condition (Figure S4C-D). Reducing LpR1 and LpR2 activities in the ISCs

also did not affect the proliferation (Figure S4E). To validate RNAi knockdown efficiency, we

quantified *LpR1* expression in thoraces from flies expressing *LpR1* RNAi under the muscle-

specific *Mhc-Gal4* driver (Myosin heavy chain), and *LpR2* expression in fat bodies from flies

expressing *LpR2* RNAi under the fat body–specific *Lpp-Gal4* driver (Apolipophorin). Both

RNAi lines efficiently reduced their respective target transcripts (Figure S4F-G ). To further

examine the role of LpR1 or LpR2 in ISC tumors, we reduced LpR1 and LpR2 levels while

overexpressing Yki (*Esg> yki$^{3SA}$;LpR1-i and  Esg> yki$^{3SA}$;LpR2-i*), and found that these

manipulations did not impair Yki-induced tracheal growth (Figure 4H-J) or prevent Yki-

induced ISC proliferation (Figure 4K). These results suggest that ISC proliferation does not

rely on circulating lipids in the hemolymph and that circulating lipids control

tracheaogenesis instead.

We further tested whether disrupting *LpR1* and *LpR2* activities in the trachea could

affect tumor growth. It has been reported that restricting tracheal growth delays tumor

development [42]. Consistently, knocking down *LpR2*, but not *LpR1*, using a temperature

sensitive trachea-specific driver (*btl-GAL4, UAS-srcGFP, Tub-GAL80^{TS}*, referred to as *Btl*) in

Yki flies prevented Yki-induced ISC proliferation (Figure 4L) and the tumor burden (Figure

4M). Consistently, it also reduced the level of bloating (Figure 4N), tracheal number of

branches, tube area and skeleton length compared to control flies (*Btl*>+) (Figure 4O-Q).

Knocking down *LpR2* in the normal condition also reduced tracheal branch number

skeleton length and total tube area (Figure 4O-Q). Reducing *LpR1* and *LpR2* function in the

trachea did not alter ISC proliferation under basal conditions (Figure S4F).

Interestingly, oenocyte-specific depletion of *LpR2* also altered gut tracheal

morphology under basal conditions (Figure S4H-J), reflecting that the trafficking of

phospholipids, WEs and AcCas from oenocytes relies on the uptake of LPPs. Altogether,

these findings are consistent with a model in which tracheal morphology, rather than ISC

proliferation, depends on lipids generated or processed in oenocytes and delivered via

circulating LPP particles.

***Hnf4-mElo axis in oenocytes mediates lipid metabolism and tracheogenesis***

***induced by Yki tumors***

Tumor growth in the adult *Drosophila* midgut has been previously shown to promote

tracheole density [42]. Similarly, we observed an increase in the number of trachea

branches, total tube area, and total tube length in the surrounding tissue of the Yki midgut

(Figure S5A-D). This led us to investigate whether oenocyte lipid metabolism mediated by

Hnf4-mElo axis is crucial for trachea growth in the Yki midgut.

To explore whether perturbations of lipid metabolism in oenocytes could block Yki-

induced tracheogenesis in the midgut, we manipulated lipid metabolism by overexpressing

*TSC1,2* or knocking down *Hnf4* or *mElo*. Using the LexA/LexAop and Gal4/UAS systems,

overexpressing *TSC1,2* or reducing *mElo* significantly decreased the number of tracheal

branches, total tube area, and total tube length in the Yki midgut. Reducing *Hnf4* levels, on

the other hand, only decreased the total tube area (Figure 5A-C).

In addition, to verify whether midgut secreted Pvf1 regulates tracheal growth, we

knocked down *Pvf1* in ISC in addition to Yki overexpression. Reduction of *Pvf1* blocked Yki

induced number of branches, tube area and skeleton length (Figure 5D-F), indicating that

oenocyte lipid metabolism is important for tracheal growth induced by Yki.

***Oenocyte Hnf4 and mElo regulate tracheogenesis in the midgut***

Next, we aimed to understand whether oenocyte lipid metabolism controls trachea

growth in control guts. Consistent with *Hnf4* knockdown, oenocyte-specific *mElo* reduction

and *TSC1,2* overexpression showed reduced level of whole body WE in non-Yki flies (Figure

6A-B). WEs are essential parts of the outer layer of the trachea [59-62], and *waterproof*

(*wat*), which is involved in biosynthesis of wax, has been implicated in hydrophobic

tracheal coating in *Drosophila* embryos [62]. Knocking down *Hnf4* did not affect the

transepithelial barrier mediated by septate junctions in the midgut trachea on the midgut,

as revealed by red dextran injection experiment (Figure 6C). However, reduction of

oenocyte *Hnf4* altered the morphology of trachea associated with the midgut (R2-R4

regions) (Figure 6D). Specifically, we observed a reduction in the number of branches and

total skeleton length, normalized to the area of the gut surface (Figure 6E-G). Consistently,

we also observed a reduction in branch numbers, tube area and skeleton length when

*mElo* was knocked down in oenocytes of control flies (Figure 6H-J). The effect of oenocyte

Hnf4 on trachea number is not specific to the midgut, as we also observed a reduction of

trachea total tube area in the ovary of control flies (Figure S6A-D). We also examined

whether Hnf4 and mElo in the fat body contribute to tracheal growth, since the fat body is a

major metabolic organ involved in lipid homeostasis and, like oenocytes, shares

hepatocyte-like functions in flies [63, 64]. However, knockdown of *Hnf4* and *mElo* using the

fat body– and gut-specific GeneSwitch driver *S106-Gal4* [65] did not affect tracheal growth

(Figure S6E-J ), in contrast to the marked reduction observed upon oenocyte-specific

knockdown. It suggests that oenocytes play a more prominent role than the fat body in

regulating tracheal growth. Overexpressing *Hnf4* or *mElo* is sufficient to induce total

tracheal tube area and skeleton length in the midguts of control flies (Figure 6K-P). Finally,

to eliminate the effect of RU feeding on trachea morphology, we fed control (*PromE>ctrl*)

flies and observed no difference in trachea morphology (Figure S6K-M).

***VEGF-A reprograms lipid metabolism in hepatoma cells, while hepatic lipid***

***metabolism is altered in tumor-bearing mice***

Using the *Drosophila* Yki tumor model, we found that the tumor secreted

PDGF/VEGF-like factor Pvf1 can non-autonomously regulate Hnf4 activity in hepatocyte-

like cells to promote the production of VLCFA or related lipids. To test whether this

mechanism is conserved in mammals, we incubated HepG2 cells with vascular

endothelial growth factor A (VEGF-A), which is orthologous to Pvf1 in flies and a potent pro-

angiogenic growth factor expressed in many types of tumors. We measured the expression

level of ELOVL fatty acid elongase 1 – 7 (*ELOVL1-7*). Incubating HepG2 with VEGF-A for 24

hours can significantly increase the level of *ELOVL1, 2, 4, 6* and *7*, suggesting a conserved

regulation of PDGF/VEGF on lipid metabolism in hepatocytes (Figure 7A-E). HNF4$\alpha$ plays

an important role in hepatocyte lipid metabolism. Consistent with this, *APOB*, an early

transcriptional target of HNF4$\alpha$ [66, 67], showed increased expression in HepG2 cells

following VEGF-A treatment (Figure 7F), suggesting an increased HNF4$\alpha$ activity.

Consistently, we observed an increased level of HNF4$\alpha$ localization in HepG2 cells treated

with VEGF-A (Figure S7A-B). We performed lipidomic profiling of conditioned medium

collected from HepG2 cultures following VEGF-A treatment and observed a robust

increase in WE (Figure 7G), as seen in Yki flies (Figure 1A-C). Together, these results suggest

a conserved VEGF-A–driven regulatory mechanism in hepatic lipid metabolism mediated

by HNF4α.

In addition, we collected liver samples from *KrasG12D/+;Lkb1f/f* (KL) mice, in which

tumors were specifically induced in the lungs, and analyzed the expression levels of *Hnf4$\alpha$*,

*Elovl1-7* and HNF4$\alpha$ target genes *acetyl-Coenzyme A carboxylase alpha* (*Acaca*), *fatty acid*

*synthase* (*Fasn*) in the liver [68-70]. *Hnf4α, Elovl7, Acc1* and *Fasn* showed higher expression

level in the tumor bearing mice, compared to control, suggesting that tumor growth in the

lung increased HNA4$\alpha$ activity and induced genes involved in lipid metabolism in liver

tissues in the KL mouse model (Figure 7H-L). Consistent with the expression data,

lipidomic profiling of serum samples with tumor bearing mice and control mice revealed

higher level of WE in the tumor mice serum (Figure 7M), which are mostly comprised of very

long chain fatty acids (Figure S7C). The lipidomic analysis also showed that out of 147

significantly changed lipid species, most of them (118) are induced and only 29 are

reduced (Figure S7C). From the summed lipid species, we found that although

phosphatidylcholine (PC) and triglycerides (TG) are reduced in tumor bearing mice serum,

the majority of other lipid species, such as Acylcarnitine (AcCa), lysophosphatidylcholine

(LPC) are highly induced in tumor bearing mice (Figure 7N). Altogether, these data suggest

a conserved mechanism of lipid metabolism regulation in hepatocytes by a PDGF/VEGF-

like factor and Hnf4 (Figure 7O).

**Discussion**

We reveal a novel mechanism by which gut tumors in *Drosophila* reprogram lipid

metabolism in oenocytes to promote tracheal development and tumor growth. Our

findings demonstrate that tumors secrete a PDGF/VEGF-like factor, Pvf1, which activates

TORC1 activity, triggering the expression of *mElo* through Hnf4 in oenocytes. This TORC1-

Hnf4 signaling pathway enhances the production of specific lipids, such VLCFAs and wax

esters (WEs), which stimulate tracheal growth, a crucial process for tumor expansion. We

also show that reducing Hnf4 or mElo activity in oenocytes suppresses tumor growth,

tracheogenesis, and tumor-induced cachexia, while extending lifespan. Additionally, we

highlight the conservation of this regulatory pathway in mammals, where VEGF-A

stimulates lipid metabolism gene expression in human hepatocytes. Furthermore, lung

tumor-bearing mice exhibit increased *Hnf4α* and *Elovl7* expression in peripheral

hepatocytes.

Pvf1, a tumor-secreted factor, has previously been linked to tumor-induced muscle

and fat body wasting, as well as renal dysfunction in *Drosophila* [22, 71, 72]. In this study,

we uncover a novel role for tumor-secreted Pvf1 in modulating oenocyte lipid metabolism

through TORC1 and Hnf4, which regulates the expression of the oenocyte-enriched gene

*mElo* (Figure 3). While Hnf4 activity is known to be induced by starvation and during the

developmental transition from pupae to larvae [39, 49], its upstream regulators remain

poorly understood. In MEF cells, the HNF4$\alpha$ binding motif is highly enriched in the

promoters of TORC1-induced genes [73], suggesting HNF4$\alpha$ mediates the transcriptional

activities of TORC1. Here, we demonstrate that TORC1 positively regulates Hnf4 protein

levels in the oenocyte nucleus through transcriptional regulation (Figure 2G). However, the

precise mechanisms controlling *Hnf4* transcription downstream of TORC1 remain unclear.

REPTOR and REPTOR-BP are likely key players in this process, as they are the major

effectors of the transcriptional response induced by TORC1 inhibition [74].

Previous studies showed that muscle-derived Pvf1 suppresses lipid synthesis in

*Drosophila* oenocytes and limits fat accumulation in the fat body through PI3K/Akt/TOR

signaling [50]. In contrast, our results indicate that Yki tumor–derived Pvf1 promotes very-

long-chain fatty acid synthesis in oenocytes, thereby supporting tumor growth. This

difference may reflect tissue-specific effects of Pvf1. Notably, muscle-derived Pvf1 was

reported to inhibit oenocyte lipid synthesis, whereas Pvf1 from the fat body and midgut had

no detectable effect on oenocyte lipid metabolism or steatosis [50]. What a gut tumor–

derived Pvf1 regulates oenocyte lipid metabolism has not been previously examined. One

possible explanation is that tissue-specific post-translational modification of Pvf1 alters its

activity, as PDGF/VEGF ligands function as disulfide-linked dimers [75, 76]. In addition,

Pvf2 and Pvf3 are strongly induced in gut tumors [22], raising the possibility that distinct

PVF homo- or heterodimers may generate different downstream signaling outputs [53].

Our study identified a novel mode of communication between oenocytes and the

gut. Oenocytes produce lipid substrates that support gut tracheal growth, a process crucial

for tumor expansion and potentially ISC growth. Like tumors, bacterial infections and

epithelial damage also trigger ISC proliferation [77-79], suggesting that oenocyte-secreted

lipids may also support stress-induced ISC expansion. Lipids from oenocytes are possibly

acquired by the tracheal cells instead of ISCs, as knockdowns of *LpR1* and *LpR2* in ISCs

have no effects in tumor growth (Figure 4H-J). In contrast, reducing *LpR2* in tracheal cells

was sufficient to decrease tracheal development (Figure 5O-Q). This may be due to the

trachea's direct contact with hemolymph, allowing it to actively acquire lipids, while the

midgut is covered by visceral muscle and basement membrane [80, 81].

The mechanism by which VLCFA metabolism regulates tracheogenesis remains

unclear, though it may involve the induction of Reactive Oxygen Species (ROS) in the

trachea. ROS has been implicated in the activation of HIF-1$\alpha$/Sima, and VLCFAs is known

to induce mitochondrial dysfunction and ROS production [82-84]. Increased ROS levels,

through treatments such as paraquat, $H_2O_2$ or bacterial infection, have been shown to

induce tracheogenesis in the midgut [42]. Furthermore, manipulating ROS level in the

trachea alone can trigger tracheal branching and ISC proliferation [42]. Interestingly,

overexpressing *Hnf4* alone in oenocytes could promote tracheogenesis, but does not

significantly induce ISC proliferation (Figure S4A). This difference may be attributed to

variations in driver strength, with tracheal development occurring prior to ISC proliferation.

Our data indicate that wax esters (WEs) and acylcarnitines (AcCas) are elevated in

tumor-bearing flies and mice (Figure 1A and 7N). Notably, in both models we observe an

increase in WEs containing very-long-chain fatty acids (VLCFAs), consistent with the

elevated expression of Hnf4 and fatty acid elongases in the KL mouse liver as well as in

oenocytes from Yki flies. Functionally, increased AcCas and WEs could contribute to

tracheogenesis by supporting local energetic demand. For example, hepatocyte-derived

plasma acylcarnitines have been shown to be required for systemic thermogenesis under

cold exposure in mice, where peripheral tissues such as brown adipose tissue, skeletal

muscle, and heart take up circulating acylcarnitines and oxidize them to fuel metabolism

[55]. In addition, wax esters can serve as high-density neutral lipid reserves that are

mobilized to meet energetic demands in diverse physiological contexts (e.g., diapause and

reproduction in marine organisms). Alternatively, lipid elevation could promote

tracheogenesis by increasing reactive oxygen species (ROS), which has been implicated as

a driver of tracheal growth . WEs can be hydrolyzed into fatty acids and fatty alcohols [42, 85]. VLCFAs released from WEs may undergo peroxisomal oxidation, generating $H_2O_2$ as a byproduct, which can stimulate tracheogenesis [42]. In parallel, fatty acids can be transported to mitochondria for β-oxidation, and high fatty-acid flux through mitochondrial respiration can increase electron leak from the ETC and elevate ROS production, potentially contributing to tracheogenesis [86]. These studies also leave open the possibility that metabolites derived from these lipid species act as signaling molecules that promote tracheogenesis.

Additionally, increased AcCas and WEs could contribute to tracheogenesis by supporting local energetic demand. For example, hepatocyte-derived plasma acylcarnitines have been shown to be required for systemic thermogenesis under cold exposure in mice, where peripheral tissues such as brown adipose tissue, skeletal muscle, and heart take up circulating acylcarnitines and oxidize them to fuel metabolism [55]. In addition, WEs can serve as high-density neutral lipid reserves that are mobilized to meet energetic demands in diverse physiological contexts (e.g., diapause and reproduction in marine organisms). These studies also leave open the possibility that metabolites derived from these lipid species act as signaling molecules that promote tracheagenesis.

We observed that VEGF-A stimulates lipid metabolism gene expression in human hepatocytes, and lung tumor-bearing mice showed increased expression of *Hnf4α* and *Elovl7* in peripheral hepatocytes. This work provides new insights into tumor-host interactions, challenging the conventional view that tumor-associated metabolic changes are merely passive processes and instead demonstrating their role in promoting tumor growth. Specifically, inhibiting *Hnf4* and *mElo* in oenocytes slows tumor growth, halts tumor-induced cachexia, and extends the lifespan of flies (Figure 1 and Figure S1). Yki activation increased levels of AcCas, PCs, and TAGs in the hemolymph, and inhibiting *Hnf4* in oenocytes reversed these lipid changes induced by Yki (Figure 4A). This study identifies the TORC1-Hnf4-Elovl7 axis and tumor-induced lipid metabolism as potential therapeutic targets for managing tumor growth or as biomarkers for early tumor detection.

Hnf4 is a master regulator of multiple metabolic pathways in *Drosophila* and

inhibiting it could have broad effects on the host. For instance, knocking down *Hnf4* in non-

tumor-bearing flies led to increased steatosis (Figures S1L). These wide-ranging effects

may limit the efficacy of Hnf4 inhibition as an anti-tumor strategy. In contrast, mElo, which

is more specifically involved in lipid elongation, presents a more targeted inhibition with

lower toxicity and a greater extension of lifespan (Figures 1M, S1O). Our study also

suggests a potential role for elongases in tumor-induced angiogenesis, as overexpression

of ELOVL7 has been linked to prostate cancer and is required for cancer growth [87]. While

mElo is exclusively expressed in oenocytes in flies (Figure 1J), its ortholog, ELOVL7, is

widely expressed in multiple tissues, including the prostate, kidney, brain, small intestine,

and thyroid [88]. It is plausible that ELOVL7 is upregulated in various tissues in cancer

patients, indicating additional tissue targets for anti-angiogenic therapies.

The regulation of *mElo* by Hnf4 seems to be prominent and specific compared to

other transcriptional targets under normal physiological conditions (Figure 1I and S1G). It

further suggests that the tumor-associated metabolic milieu may reshape Hnf4-driven

transcription, potentially by altering (i) cofactor interactions and/or (ii) ligand availability,

thereby promoting selective activation of downstream targets such as mElo. Supporting

this concept, previous studies demonstrate that HNF4$\alpha$ transcriptional output is context-

dependent and influenced by partner proteins: for instance, HNF4$\alpha$ can cooperate with

ChREBP (carbohydrate response element-binding protein) to stimulate lipogenic gene *Fasn*

during feeding [68, 70], whereas under fasting conditions it can interact with *Foxo1* to

enhance expression of gluconeogenic genes such as *G6Pase* [89]. Moreover, HNF4$\alpha$

activity can be modulated by long-chain fatty acids [46], raising the possibility that changes

in lipid availability in tumor-bearing flies (e.g., due to TAG mobilization) may further

influence Hnf4 activity and target selection. Collectively, these findings support a model in

which tumor-induced shifts in Hnf4 cofactors and/or lipid ligands redirect Hnf4

transcriptional program in Yki oenocytes, leading to robust *mElo* induction.

In summary, our data reveal a novel paradigm in which growing tumors actively

acquire substrates for tracheogenesis by secreting a PDGF/VEGF-like factor that activates

the TORC1-Hnf4 axis in distal oenocytes, inducing the expression of mElo. This promotes

the synthesis of VLCFAs, which are transported to the trachea to further support tumor

growth (Figure 7O).

**Materials and Methods**

***Drosophila stocks and husbandry***

All experiments were conducted using female flies due to their more consistent

bloating phenotype and larger oenocyte, which facilitated dissection. For temperature-

sensitive experiments, fly crosses were set up and maintained under a 12:12 hour

light:dark cycle at 18°C. Crosses were discarded after 5 days to maintain consistent

population density. Following eclosion, adult flies were collected and kept at 18°C for 2–3

days. GAL80$^{TS}$ driver activation was achieved by incubating the flies at 29°C for the

durations specified in each experiment. For experiments involving the GeneSwitch driver,

fly crosses were maintained at 25°C under a 12:12 hour light:dark cycle and discarded after

3 days. Adult progenies were collected and fed on food containing 200 μM RU486

(Mifepristone, Cayman Chemicals, 10006317), prepared by dissolving the compound in

95% ethanol before adding it to standard fly food. Flies were fed on RU486-containing food

for 7 days. Esg$^{ts}$ (*esg-Gal4, UAS-GFP, tub-GAL80$^{TS}$*), *Esg-LexA* and *LexAop-yki$^{3SA}$-GFP, Attp40*

*UAS-Luciferase* are from the Perrimon lab stock collection.

The following strains were obtained from the Bloomington *Drosophila* Stock Center

(BL), Vienna Drosophila Resource Center (V) and flyORF (F): *UAS-Hnf4 RNAi* (BL# 29375),

*UAS-mElo RNAi* (BL# 44510), *UAS-yki$^{3SA}$* (BL#28817), *UAS-Pvf1-3xHA* (F002862), *UAS-*

*PvR[A]* (BL#58496), *UAS-Pvf1 RNAi* (V#102699), *UAS-PvR RNAi* (V#43459), *UAS-Luciferase*

*RNAi* (BL#31603), Oe$^{ts}$ (*Desat1-Gal4, tubP-Gal80*) (BL# 65406), *UAS-LpR1 RNAi* (BL#

50737), *UAS-LpR2 RNAi* (BL# 54461).

*UAS-Hnf4* is a gift from Dr. Carl Thummel [46]; *UAS-TSC1, TSC2* is a gift from Dr.

Christen K Mirth [90] and UAS-PRAS40 was a gift from Dr. Aurelio Teleman [91], Btl$^{TS}$ (*Btl-*

*Gal4, UAS-srcGFP*) and *QF6 >QUAS-mtdTomato* was a kind gift from Dr. Chrysoula Pitsouli

[42]. PromE$^{GS}$(*PromE-GeneSwitch-Gal4*) was a gift from Dr. Heinrich Jasper [92]. *UAS-*

*CG18609* and *mFas-Gal4* were gifts from Dr. Henry Chung [93, 94]. *UAS-PRAS40* is a gift

from Dr. Aurelio A Teleman [74].

Hnf4-127D01 was generated through CRISPR-mediated tagging of endogenous Hnf4

with 127D01-tag as described previously [47]. Briefly, we cloned the sequence 1 kb

upstream and 1 kb downstream of the stop codon into the donor vector pScarlessHD-C-

3x127D01-DsRed (Addgene#171578). A sgRNA plasmid pCFD3-Hnf4-sgRNA that target the

seed sequence (CCAGAGACTGGTTACTAGAAGAA) near the stop codon of Hnf4 was injected

into yw;nos-cas9/CyO embryos together with the donor plasmid. Positive transformants

with red eye fluorescent eyes were outcrossed and successful Kis were confirmed by PCR

(F: GTACAACCGGAGCGAGGGTA; R: CAAAATGGTTCAAGGCATAACA) and sequencing.

Subsequently, 3xP3dsRed was excised using piggyBac transposase (BL#8285).

***Untargeted lipidomic analysis***

Lipidomics and metabolomics of fly samples was performed following the Folch

method (Folch et al., 1957). 15 adult flies, 20ul of mice serum and hemolymph from 200-

300 adult flies were used for lipidomics and metabolomics. Solid samples were

homogenized in a 1 mL Dounce homogenizer (TIGHT) on ice with 0.6 mL chloroform and 0.3

mL methanol (2:1 chloroform-methanol mixture). Homogenized samples were transferred

to a 15 mL glass tube with a Teflon cap, and an additional 0.6 mL chloroform and 0.3 mL

methanol were added to make the final volume approximately 1.8 mL. Hemolymph

samples or conditioned medium was transferred to a 15 mL glass tube containing 2:1

chloroform-methanol mixture. Samples were vortexed briefly and incubated at room

temperature on a rotator for 30–45 minutes. Following incubation, 0.2 volumes (0.36 mL) of

HPLC-grade deionized water (HPLC dH2O) were added. The mixture was vortexed three

times, followed by centrifugation at 1000 × g for 10 minutes at 4°.

The upper aqueous phase, containing polar metabolites, was carefully collected

using a glass pipette, while avoiding the middle phase. Metabolites were transferred to

microcentrifuge tubes, evaporated under vacuum using a SpeedVac rotary evaporator

overnight, and stored at -80°C. Polar metabolites were resuspended in 20 μL of HPLC-grade

water and then injected and analyzed using a hybrid 6500 QTRAP triple quadrupole mass

spectrometer (AB/SCIEX) connected to a Prominence UFLC HPLC system (Shimadzu). The

lower phase, containing non-polar lipids, was transferred to a clean 1.85 mL glass test tube

using a glass pipette. The phase was dried under nitrogen gas for approximately 30

minutes, and dried lipids were stored at -80°C to prevent oxidation. Non-polar lipids were

resuspended in 35 µL of HPLC-grade 50% methanol/50% isopropyl alcohol (IPA) and then

injected and analyzed using untargeted high-resolution LC-MS/MS on a Thermo QExactive

Plus Orbitrap mass spectrometer. LipidSearch version 4.2 (Thermo Scientific) was used to

identify lipid molecules, with the quantification of ion intensity by measuring the area size

of identified peaks.

The full names of the lipid species can be found in Supplementary Table 4. For

lipidomic quantification using whole fly bodies, the same number of flies was used for lipid

extraction. Lipid species were normalized to the total reads in each sample. For

hemolymph lipidomic analysis, lipid species were normalized to both the volume of

hemolymph extracted and the total reads in each sample.

***RNA extraction Quantitative RT-PCR***

Adult female fly oenocytes were dissected in cold 1xPBS before RNA extraction. For

oenocyte dissection, the fat body was removed through liposuction and then oenocytes

were removed from the cuticle using a fine-tip glass needle. Tissue lysis, RNA extraction,

and cDNA synthesis were performed using Cells-to-CT Kit (Thermo Scientific, #44-029-53).

For female adult whole body mRNA extraction, four flies per biological replicate were

collected. Total RNA was extracted using Direct-zol RNA MicroPrep kit (Zymo Research,

#R2060). mRNA was extracted from HepG2 cells using Directo-zol RNA MicroPrep kit

(Zymo Research, #R2060). 30-50mg of mice livers were dissected and total RNA was

extracted using Qiagen RNeasy Mini kit (Qiagen #74104). All the cDNAs were synthesized

using the iScript cDNA synthesis kit (Bio-Rad, #1708896) in a 40ul reaction mixture using

1000ng total RNA. Quantitative real-time RT-PCR (RT-qPCR) assays were performed using

iQ SYBR Green Supermix (Bio-Rad, #1708880) on a CFX96 Real-Time PCR Detection

system (Bio-Rad). Two to four independent biological replicates were performed with two

technical replicates. The mRNA abundance of each gene was normalized to the expression

of *RpL32* or *αTub84B* for fly samples, Gapdh for mouse samples and ACTB for HepG2

samples. Primer sequences are listed in the Supplementary Table 5.

***Fly imaging***

*Drosophila* ovaries were dissected in cold PBS. ovaries were imaged immediately in

cold PBS using a ZEISS Axiozoom V16 fluorescence microscope. Flies were anesthetized

on $CO_2$ pad and imaged using ZEISS Axiozoom V16 fluorescence microscope. The size of

the ovary and abdomen is measured using Fiji (version: 2.14.0/1.54f).

***Oil Red O staining***

The method was as previously described [29]. Flies were dissected in PBS and fat

bodies were removed with liposuction. Specimens were fixed in 4% paraformaldehyde for

20 minutes. Specimens were rinsed twice with distilled water. Oil Red O (O0625-, Sigma-

Aldrich) was dissolved in isopropanol to make 0.1% Oil Red O stock. A working solution

was prepared by combining 3 mL of 0.1% Oil Red O in isopropanol with 2 mL of distilled

water. The solution was freshly prepared and filtered through a 0.2-μm syringe filter before

use. Specimens were incubated for 25 minutes in Oil Red O stain, rinsed briefly with PBS

and mounted in Vector Shield for imaging. Oil Red O images were taken using Olympus

VS200 Slide Scanner, equipped with color camera (2448x2048 pixels, 3.45μm).

***Immunostaining and image analysis***

For immunostaining using the nanobodies, fly abdomens were dissected in PBS at

room temperature with fat bodies removed and fixed in 4% formaldehyde in PBS for 30

minutes. Specimens were washed three times in PBST (0.1% Triton-X-100), 5 minutes each.

The specimens were then incubated in 5% NGS (normal goat serum in PBST) for 1 hour.

Nb127D01-ALFA, 0.2mg/ml was used at a dilution of 1:500, incubating the samples for 1

hour and a half with mild shaking. Followed by 0.1% PBST washes, 5 minutes each wash.

AlexaFluor 594 anti-Alpaca (Jackson Immuno Research, 128-585-232) was diluted 1:400 in

PBS-TX + 5% NGS and incubate with the specimens at room temperature with mild shaking

for 1 hours. After washes and DAPI staining, tissues were mounted in Vectashield and

imaged with Olympus VS200 Slide Scanner or confocal microscope Zeiss Axio Observer Z1.

For regular immunostainings, fly tissues were dissected in PBS at room temperature.

Tissues were fixed in 4% paraformaldehyde in PBS for 30 minutes. Specimens were washed

three times in 0.1% PBST and incubated in 5% NDS solution for 1 hour. Tissues were then

incubated with primary antibodies diluted in PBST and blocking solution overnight at 4°.

Then the tissues were washed three times with PBST at room temperature. Secondary

antibodies were applied for 1 hour. Excess antibodies were washed in PBST, and tissues

were incubated with DAPI (Invitrogen, D1306) dilute in PBST for 5 minutes. Tissues were

then mounted in Vectashield (Vector Laboratories Inc, H-1000). The following primary

antibodies were used: rabbit-anti-pH3 (1:500, Cell Signaling, 9701), rabbit-anti-p4E-BP

(1:800, Cell Signaling, 2855S). Secondary antibodies against mouse, rabbit, chicken or

alpaca conjugated to Alexa Fluor 488 and 594 (Jackson Immuno Research) were used at

1:800 for anti-p-4EBP, 1:1000 for anti-pH3 and 1:400 for Nb127D01-ALFA.For

immunostaining in HepG2 cells, cells were washed twice with PBS and fixed in 4%

paraformaldehyde for 25 min, followed by a PBS wash. Samples were permeabilized with

0.5% Triton X-100 in PBS for 10 min and washed with PBS. Non-specific binding was

blocked in 1% BSA in PBS for 1 h at room temperature. Cells were then incubated with anti–

HNF4α rabbit monoclonal antibody (C11F12, Cell Signaling Technology, #3113) diluted

1:1000 in PBS at 4 °C overnight. After washing twice with 0.05% Triton X-100 in PBS,

samples were incubated with Alexa Fluor 594–conjugated secondary antibody (1:500 in

PBS) at room temperature. Cells were washed three times with 0.05% Triton X-100 in PBS

and once with PBS, then mounted using SlowFade® Gold antifade reagent with DAPI

(Invitrogen, S36939) and sealed with a coverslip.

Quantification of nucleus Hnf4 intensity was conducted using ImageJ. Background

subtraction was performed using the rolling ball radius at 50 pixels. We apply threshold to

the DAPI channel and using analyze particles to select nucleus as Region Of Interest (ROI),

size: 10-2000; circularity: 0.1-1. Mean gray value was measured for each ROI.

***Lifespan analysis***

For survival analysis, flies were collected within 24 hours post-eclosion (20 females

and 3 males per vial) and maintained on standard laboratory fly food at 18°C, with 60%

humidity, and a 12-hour light/dark cycle. After two days, mature and mated females were

transferred to 29°C to induce gene expression. To maintain vial cleanliness, flies were

transferred to fresh vials every two days, and the number of dead flies was recorded daily.

***snRNAseq analysis***

Data and plots were generated from the Fly Cell Atlas, generated using the 10X

platform, "Stringent All" dataset. Other snRNAseq dataset was analyzed as described in

[50], where a resolution of 0.4 was chosen as the clustering parameter.

For the Seurat Module Score analysis, Seurat AddModuleScore function was used to

calculate the average expression of a group of genes in Very Long Chain Fatty Acid

Synthesis pathway in each cell (for details of the list, see Supplementary Table 6), then

subtracted by a randomly picked gene set expression.

***Rapamycin feeding***

Rapamycin was dissolved in DMSO to make 2mM rapamycin stock. Rapamycin

stock was further diluted to 200$\mu$M in ethanol and evenly distributed on to food surface. A

small piece of Kimwipe was used to remove excess liquids on the food. DMSO dissolved in

ethanol was used as control. Flies were transferred to new food every two days. Flies were

fed for 6-7 days before dissection.

***Dextran injection***

Using a fine glass needle, ~100nl of 50 mg/ml 10,000 MW TR-dextran (Thermo Fisher

Scientific, D-1817) diluted in water was injected in the abdomen of adult females. 6 hours

after injection, abdomen and thoraces were separated and whole abdomen were fixed in

4% paraformaldehyde for 30 minutes with mild shaking. After fixation, midguts were

dissected and washed briefly with PBS and mounted in Vectashield. Midguts were imaged

using a single point laser scanning confocal, Olympus IX83 to measure the trachea

leakage. A Z-stack was taken, and maximum projections were used to generate final image

for analysis.

***Trachea imaging***

Generating fly strains that incorporate both a trachea reporter and other genetic

modifications is challenging. Therefore, the following method, as described in [42], was

used to quantify the trachea. Flies were dissected in cold PBS, and intestines were

carefully extracted to maintain their original structure. Tissues were then fixed in 4%

paraformaldehyde for 20 minutes, mounted in Vectashield, and imaged within 3 days.

Whole intestine images were captured using an Olympus VS200 Slide Scanner equipped

with a color camera. Only the upper layer of the tissue, in contact with the coverslip, was

analyzed.

***Trachea analysis***

To automate trachea measurement, we used the previously developed software

AutoTube [95]. Only the topmost Z-stack, which provided the clearest image of the trachea,

was analyzed. Images corresponding to the R2-R4 regions of the gut were cropped,

avoiding areas with the major trunk of the trachea to prevent skewing the results. Images

were thresholded on a white background and analyzed with AutoTube using the following

settings:

- Input Color Channel: Green

- Intensity adjustment: Autocontrast

- Illumination correction: Illumination size set to 51

- Noise correction: BM3D

- Tube detection: Finer tubes with a tube size of 3

- Small regions removed: 0.01% of Ima

- Hole filling: Hole size set to 1

- Threshold type: MultiOtsu

- Short ramifications removed: Spur length set to 15

- Spatial branch merging: Spatial distance set to 10

The analysis provided results for "Number of Branches (µm)," "Total Tube Area (µm)," and

"Total Skeleton Length (µm)," which were normalized to the corresponding intestinal area

and were multiplied by an arbitrary number 1000000. Results from the R2-R4 regions were

averaged for each gut.

***Animal care and tumor induction***

$Kras^{G12D}/+;Lkb1^{f/f}$ mice, previously described [7, 96], were backcrossed to the FVB

strain. Male mice were maintained on a 12-hour light/dark cycle at an ambient temperature

of 22°C, with ad libitum access to rodent chow (PicoLab Rodent 5053; Lab Diet, 3.43

kcal/g) and drinking water. Tumors were induced in adult male and female mice (12 to 20

weeks old) via intranasal administration of 75 µL PBS containing $2.5 \times 10^7$ pfu of adenovirus

SMV-CRE (Ad5CMV-Cre) purchased from the University of Iowa Gene Transfer Vector Core

(Iowa City, IA). All animal experiments were conducted with the approval of the Institutional

Animal Care and Use Committee (IACUC) at Weill Cornell Medicine (WCM) under protocol

number 2013-0116.

***Mice tissue collection***

Male mice were euthanized with $CO_2$, followed by blood collection via cardiac

puncture. Liver, gonadal adipose tissue, and skeletal muscles were dissected and flash-

frozen in liquid nitrogen and stored at -80°. The control mice were a mix of fasted and fed

mice. "Fasted" mice were fasted for 12 hours and "fed" mice were fasted for 12 hours and

then allowed free access to food for 4 hours before tissue collection.

***VEGFA incubation***

HepG2 cells were cultured in HyClone Dulbecco's Modified Eagle Medium (DMEM)

with high glucose (SH30022.01): with L-glutamine; without sodium pyruvate. Additional

50U penicillin/ml and 50ug streptomycin/ml were added. HepG2 cells were seeded at

density of $5 \times 10^5$ in 12-well-plate (VWR, 29442-040), treated or not with 50ng/ml

Recombinant Human VEGF 165 (PeproTech, 100-20-2UG) for 24 hours. Total cellular RNA

was isolated using the Trizol reagent and extracted through Direct-zol RNA MicroPrep kit

(Zymo Research, #R2060).

**Acknowledgments**

We thank Dr. John M. Asara at the Beth Israel Deaconess Medical Center for his help

on lipidomic analysis. We thank the assistance provided by the Microscopy Resources on

the North Quad (MicRoN) core and we thank Dr. Simon F. Norrelykke at the Image Analysis

Collaboratory at Harvard Medical School for advice on AutoTube. We also thank Dr. Ah-

Ram Kim, Dr. Sun Jin Moon at Harvard Medical School, Dr. Hua Bai at Iowa State University,

Dr. Eileen White and Dr. Maria Gomez- Jenkins at Rutgers University for scientific

discussion and insights. We thank Dr. Ju Xu at Chinese Academy of Sciences for the help

generating Hnf4-127D01 fly strain. We thank Dr. Ying Liu at Harvard Medical School for

sharing fly lines used in this study. We also thank Christians Villalta for the help on fly

dextran injection.

This article is subject to HHMI's Open Access to Publications policy. HHMI lab

heads have previously granted a nonexclusive CC BY 4.0 license to the public and a

sublicensable license to HHMI in their research articles. Pursuant to those licenses, the

author-accepted manuscript of this article can be made freely available under a CC BY 4.0

license immediately upon publication.

**Funding**

This work is funded by NIH/NCI Grant #5P01CA120964-15 and is delivered as part of

the CANCAN team supported by the Cancer Grand Challenges partnership funded by

Cancer Research UK (CGCATF-2021/100022) and the National Cancer Institute (1 OT2

CA278685-01). This work is funded by the Hypothesis Development Award from DOD

CDMRP TSCRP (GRANT14223060). N.P. is an investigator of the Howard Hughes Medical

Institute.

**Author contributions**

Conceptualization: KH, TM, MG, NP; Methods, Data Acquisition, Resources,

Visualization and Imaging: KH, TM, YC; Mice sample preparation: ED, JS; Data Acquisition:

KW, Bioinformatic Analysis: MH, YH, SJM; Independent replication of experiments at

Harvard Medical School: KH; Writing – original draft: KH, Writing – review, editing & revising:

KH, TM, NP.

**Figure 1: Oenocyte Hnf4 regulates wax ester synthesis and supports Yki-induced**

**tumor growth.**

(A-B) Heatmap of the normalized total ion intensity of major lipid classes identified in

whole bodies from control flies (*Esg-LexA*) or Yki flies (*Esg-LexA/LexAop-yki$^{S3A}$; PromE-*

*GAL4, Tub-GAL80$^{TS}$/+*), with four biological replicates per condition. Scaled colors are

presented as the values relative to the average of each row. (B) Untargeted lipidomic

profiling using LC-MS/MS of whole-body extracts revealed increased wax ester (WE) levels

in Yki flies. Normalized intensity was calculated as the integrated LC–MS peak area for

each lipid species, normalized to the sample's total signal (total reads) and to the analyzed

sample volume. (C) Fatty acid chain-length distribution (C< 22 and C≥22) in detected WE

level from control and Yki flies. (D) WE level in control flies, Yki flies, and *Hnf4* knockdown

in oenocytes of Yki flies, N = 4 biological replicates. (E-F) Immunostaining of Nb127D01 for

nuclear-localized, endogenously tagged Hnf4 (Hnf4-127D01) in oenocytes in controls and

in Yki flies. DAPI staining marks nuclei, with dashed lines enclosing oenocytes. N = 8

independent experiments. (G) Quantification of intestinal stem cell (ISC) proliferation, as

indicated by phospho-histone H3 (pH3+) cells in control. (H) Bloating as measured by

abdominal area of the fly in control, oenocyte specific Hnf4-i, Yki flies and Yki flies with

oenocyte specific Hnf4-i. (I) *mElo* expression in oenocytes, normalized to housekeeping

gene, in control, oenocyte specific Hnf4-I, Yki flies and in Yki flies with *Hnf4* knockdown. N

= 2 biological replicates from 2 independent experiments. (J) UMAP from whole body

(without head) snRNAseq, *mElo* expression is enriched in adult oenocytes. (K)

Quantification of pH3+ cell number, N =8 flies. (L-M) Lifespan analysis showing that

oenocyte-specific knockdown of *Hnf4* (L) or *mElo* (M) in Yki flies, compared to only Yki or

control flies, N = 40. Scale bar shows 20μm. Data are presented as mean ± s.e.m.

Statistical significance was determined using Student's t-test or one-way ANOVA with

appropriate post hoc tests, or Log-rank (Mantel-Cox) test. *P < 0.05, **P < 0.01, ****P <

0.0001.

**Figure 1S: Hnf4 and mElo regulate metabolic and tumor-related phenotypes in Yki**

**flies.**

(A) Immunostaining of Nb127D01 for nuclear-localized, endogenously tagged Hnf4 (Hnf4-

127D01) in oenocytes with *Hnf4* knockdown or control flies, with dashed lines enclosing

oenocytes. (B-C) Level of GFP positive region in the whole fly midgut in control, oenocyte

specific *Hnf4-i* and *Yki* flies with *Hnf4-i*. (D) Abdominal area of control and Yki flies for

bloating measurement. (E) Ovary size in control flies, oenocyte specific *Hnf4-i* flies, *Yki* flies

and *Yki* flies with oenocyte specific *Hnf4-i*, n = 9 - 15. (F) Wet weight measurement of fly

whole body, where flies were weighed first and were dried at 65°C for 5 hours, the second

value was subtracted from the first to obtain the wet weight of the fly. N = 3 biological

replicates. (G) Expression of VLCFA biosynthesis and β-oxidation genes in control and Yki

flies. (H-I) *mElo* expression measured by qPCR in oenocyte specific *Hnf4* knockdown or

*Hnf4* overexpression, N = 2 biological replicates from 2 independent experiments. (J) Level

of GFP positive region in the whole fly midgut in control, Yki flies, oenocyte specific *Hnf4*-i

and *Yki* flies.  (K) Confirmation of *mElo* knockdown efficiency via qPCR, N = 2 biological

replicates from 2 independent experiments. (L) Oil Red O staining in oenocytes of *Hnf4*

knockdown or *mElo* knockdown. N = 8 flies. (M) Ovary size measurement in control and

*mElo-i* flies, N = 13-16 flies. (N) Ovary size in control, Yki and Yki with *mElo-i* flies. (O)

Lifespan analysis in Yki-expressing flies upon oenocyte-specific *mElo* knockdown in male

flies. Data represent mean ± s.e.m.; significance assessed using appropriate statistical

tests.

**Figure 2: TORC1 regulates Hnf4 activity and lipid metabolism in oenocytes.**

(A) Whole body WEs level in ISC Yki flies and Yki with oenocyte *TSC1,2* overexpression. (B-

C) P-4EBP immunostaining indicates TORC1 activity in oenocytes of Yki flies, quantification

of P-4EBP intensity is shown on the right (C), N = 8-10 flies. (D) *mElo* expression in whole

body mRNA extracts in Yki flies and Yki flies with *TSC1,2* overexpression in oenocytes, N = 6

biological samples. (E) Quantification of pH3+ cell number, N = 8 – 10 flies. (F) Violin plot of

*Hnf4* expression in fat body, muscle and oenocytes upon overexpression of *TSC1,2* in

oenocytes [12]. (G-H) Immunostaining of Hnf4-127D01 in oenocytes when *TSC1,2* is

overexpressed in oenocytes compared to control. DAPI marks nuclei, with dashed lines

enclosing oenocytes, quantification of Hnf4-127D01 signal intensity in the nucleus, N = 8-

10 flies. (I) ModuleScore from snRNAseq analysis visualizes VLCFA synthesis gene

expression across muscle, fat body and oenocytes, in control and upon oenocyte *TSC1,2*

overexpression. (J) ORO staining indicates steatosis in oenocytes induced by *TSC1,2*

overexpression, and in *TSC1,2* with *Hnf4* overexpression, N = 8. (K-L) Expression of Hnf4

targets *mElo* and *FASN2* in *TSC1,2* overexpression and in *TSC1,2* with *Hnf4* overexpression.

N = 2 biological replicates from 2 independent experiments.

**Figure 2S: The role of oenocyte TORC1 in regulating lifespan, steatosis and Yki wasting**

**phenotype.**

(A-B) Ovary size measurements and bloating are rescued by *TSC1,2* overexpression in Yki

flies, N = 15-30 flies. (C) Sensitivity to dry starvation in flies overexpressing *TSC1,2* in

oenocytes compared to controls. Adult oenocyte specific driver (*mFas-Gal4*) was used

[93]. (D-E) ORO staining indicates steatosis in oenocytes with *PRAS40* overexpression, or in

rapamycin treatment. Dashed lines enclose oenocytes in all images. Data represent mean

± s.e.m.; statistical significance determined using Student's t-test or ANOVA as

appropriate.

**Figure 3: ISCs secreted Pvf1 activates Hnf4 through PvR signaling in oenocytes and affects tumor associated phenotypes.**

(A-B) Hnf4 immunostaining in oenocytes of Yki flies with ISC-specific Pvf1 knockdown. Dashed lines enclose oenocytes. N = 10 biological replicates. (C) *mElo* expression in oenocytes of Yki flies with ISC-specific Pvf1 knockdown, N = 4 biological replicates from 2 independent experiments. (D) Inhibition of *Pvf1* in the gut affects bloating in Yki flies, n = 15-20 flies. (E) ISC proliferation in Yki flies with *Pvf1* knockdown, n = 8-10 biological replicates. (F) *mElo* expression in oenocytes of Yki flies with *PvR* knockdown. N = 2 biological replicates from 2 independent experiments. (G-H) *PvR* knockdown in oenocytes of Yki flies and its effect on tumor-associated bloating (G) and pH3+ cell number (H). (I) Lifespan extension in Yki flies with oenocyte-specific *PvR* knockdown, N = 40 flies. N = 10 biological replicates. (J) Hnf4 immunostaining in oenocytes of flies with ISC-specific *Pvf1* overexpression. Dashed lines enclose oenocytes. N = 10 biological replicates. (L) *mElo* expression in oenocytes following ISC-specific *Pvf1* overexpression. N = 4 biological replicates from 2 independent experiments. (M-O) Immunostaining and qPCR analysis reveal Hnf4 nuclear localization and *mElo* expression in oenocytes overexpressing active *PvR*. Dashed lines enclose oenocytes in all immunostaining images. Data are presented as mean ± s.e.m.; statistical significance was assessed using Student's t-test or ANOVA.

**Figure 3S: Expression of *Pvf1* in fly whole body and oenocytes in Yki flies.**

(A) Heatmap showing average gene expression changes (log2 fold change; Yki vs WT at day 8) across multiple tissues. (B) UMAP from whole body (without head) snRNAseq, shows *Pvf1* expression level, oenocyte cluster is marked as "oe". (C) Ovary wasting of Yki flies with ISC-specific *Pvf1* knockdown. Dashed lines enclose oenocytes. N = 10-20 biological replicates. (D) Tumor mass (GFP-positive region of the gut) in Yki flies with ISC-specific *Pvf1* knockdown. N = 8-10 biological replicates. (E) Ovary wasting of Yki flies with ISC-specific *Pvf1* knockdown. Dashed lines enclose oenocytes. N = 10-20 biological replicates. (F)

Tumor mass (GFP-positive region of the gut) in Yki flies with ISC-specific *Pvf1* knockdown.

N = 8-10 biological replicates.

**Figure 4: Circulating lipids target trachea but not ISCs to promote tracheal growth.**

(A) Untargeted lipidomic analysis of fly hemolymph showing lipid species in control, Yki

flies, and Yki flies with oenocyte-specific *Hnf4* knockdown (*Esg>yki$^{3SA}$;PromE>Hnf4-i*). N = 4

biological replicates. (B-C) Levels of very-long-acylcarnitines (AcCa 24:1 and AcCa 24:2) in

Yki flies with or without *Hnf4* knockdown in oenocytes. (D-E) Levels of circulating

phospholipids, including PC 36:4 PE 32:5, across experimental conditions. (F-G) Levels of

circulating TG 18:3_16:0_16:1 and TG 18:2_16:0_18:0 in the fly hemolymph. (H-J) Number

of tracheal branches, tube area, and skeleton length in Yki flies with ISC-specific

knockdown of *LpR1* and *LpR2*. (K) PH3+ cell counts in ISCs of Yki flies with ISC-specific

knockdown of *LpR1* and *LpR2*. (L) PH3+ cell counts in ISCs of Yki flies with trachea-specific

(*Btl-Gal4* driver) knockdown of *LpR1* and *LpR2*. (M) GFP positive region in the gut of tracheal

specific *LpR2* knockdown in Yki flies. N = 8-12 flies. (N) Abdominal bloating area in Yki flies

with trachea-specific LpR2 knockdown. (O-Q) Number of tracheal branches, tube area,

and skeleton length in Yki flies with trache-specific knockdown of *LpR2*. N = 8-12. Data are

presented as mean ± s.e.m.; statistical analysis was performed using Student's t-test or

ANOVA as appropriate.

**Figure 4S: Effects of ISC-specific *LpR1* and *LpR2* knockdown on tracheal morphology**

**and ISC proliferation.**

(A) Counts of pH3+ cells in the midgut of flies with oenocyte-specific *TSC1*,2

overexpression or *Hnf4* knockdown or overexpression. N = 8-20 flies. (B-D) Trachea

parameters, including number of branches, tube area, and skeleton length, in flies with

ISC-specific knockdown of *LpR1* and *LpR2* using *Esg* driver, compared to control (*Esg>ctrl*).

N = 7-11 flies. (E) Counts of pH3+ cells in the midgut of flies with ISC-specific knockdown of

*LpR1* and *LpR2*, compared to controls. (F) *LpR1* expression level in fly thoraces with *LpR1*

knockdown using temperature sensitive muscle specific driver (Mhc-Gal4) activated at

adult stage for 7 days. (G) *LpR2* expression level in fly fat bodies with *LpR2* knockdown

using temperature sensitive fat body specific driver (Lpp-Gal4) activated at adult stage for 7

days.  (H-J) Gut tracheal parameters in flies with oenocyte-specific *LpR2* knockdown.

N=20-25 flies. Data are presented as mean ± s.e.m.; statistical analysis was performed

using Student's t-test or ANOVA as appropriate.

**Figure 5: Hnf4-mElo axis in oenocytes mediates lipid metabolism and tracheogenesis**

**induced by Yki tumors.**

(A-C) Number of tracheal branches, total tube area, and total tube length in the midgut of

Yki flies with oenocyte-specific *TSC1,2* overexpression, *Hnf4* knockdown, or *mElo*

knockdown. N = 9-17 flies. (D-F) Tracheal morphology measurements, including number of

branches and skeleton length, in flies with ISC-specific knockdown of *Pvf1* combined with

*yki* overexpression (*Esg>yki³ˢᴬ;Pvf1-i*). N = 9-11 flies. Dashed lines enclose oenocytes in all

relevant images. Data are presented as mean ± s.e.m.; statistical analysis was performed

using Student's t-test or ANOVA as appropriate.

1001

1002    **Figure 5S: Hnf4-mElo axis in oenocytes mediates lipid metabolism and tracheogenesis**

1003    **induced by Yki tumors.**

1004    (A-D) Tracheal morphology parameters, including number of branches, total tube area,

1005    total tube length, and skeleton length, in Yki midgut (*Esg>yki*), compared to wild-type

1006    controls (*Esg>ctrl*). N = 16-17.

1007

**Figure 6: Hnf4-mElo axis regulates tracheogenesis in the midgut.**

(A-B) Whole-body wax ester (WE) levels in flies with oenocyte-specific *Hnf4* knockdown

(*PromEᵍˢ>Hnf4-i*), *mElo* knockdown (*PromEᵍˢ>mElo-i*), or *TSC1,2* overexpression

(*PromEᵍˢ>TSC1,2*). N = 3-4 biological replicates (C) Transepithelial barrier function of the

trachea on the midgut, assessed using red dextran injection, in flies with oenocyte-specific

*Hnf4* knockdown. N = 8 flies. (D-G) Trachea morphology on the midgut (R2-R4), including

branch number and skeleton length normalized to gut surface area, in flies with *Hnf4*

knockdown in oenocytes. N = 8 flies. (H-J) Branch number, tube area, and skeleton length

of the midgut trachea in flies with oenocyte-specific *mElo* knockdown. N = 6-8 flies (K-P)

Number of branches, tracheal tube area and skeleton length of the midgut in flies

overexpressing *Hnf4* or *mElo* in oenocytes (*PromE^{gs}>Hnf4* or *PromE^{gs}>mElo*). N = 8-15 flies.

1019

**Figure 6S: Effect of oenocyte Hnf4 on ovary trachea.**

(A-D) Tracheal branch number, tube area and total skeleton length of the ovary in flies with

*Hnf4* knockdown in oenocytes. Red (mtdTomato) marks the trachea (QF6>QUAS-

mtdTomato). N = 8 flies. (E-G) Tracheal parameters of mid gut in flies with *Hnf4* knockdown

in the fat body and gut. (N =18). (H-J) Tracheal parameters of mid gut in flies with *mElo*

knockdown in the fat body and gut. (N =18). (K-M) Comparison of gut trachea morphology in

control flies (*PromE^{gs}>+*) with and without RU feeding. N = 12 flies. Data are presented as

mean ± s.e.m.; statistical analysis was performed using Student's t-test or ANOVA as

appropriate.

1029

**Figure 7: Regulation of hepatocyte lipid metabolism by pro-angiogenic factor in KL**

**mice tumor model.**

(A-F) Relative expression levels of *ELOVL* fatty acid elongase 1,2,4, 6 and 7 and *APOB* in

HepG2 cells incubated with vascular endothelial growth factor A (VEGF-A) for 24 hours,

compared to controls. N = 4 biological replicates, from 2 independent experiments. (G) The

level of WE from conditioned medium collected from HepG2 cultures following VEGF-A

treatment. (H-L) Relative expression levels of *Hnf4α, Elovl7, Acc1* and *Fasn* in liver samples

from *KrasG12D/+;Lkb1f/f* (KL) tumor-bearing mice compared to control mice. N = 3

biological replicates, from 2 independent experiments. (M) The level of WEs from serum

samples of KL mice and control mice. (N) Heatmap showing the significantly changed lipid

species from serum samples of tumor-bearing and control mice. (H) Schematic

representation of the proposed mechanism, depicting tumor-secreted PDGF/VEGF-like

factors activating the mTOR-Hnf4 axis in hepatocyte-like oenocytes to promote lipid

synthesis, which are then transported to trachea. Data are presented as mean ± s.e.m.;

statistical analysis was performed using Student's t-test and significance analysis of

microarrays was used for lipid species.

1046

**Figure 7S: VEGF-A regulates lipid metabolism in hepatocytes.**

(A-B) Immunostaining of HepG2 cells treated with or without VEGF-A for 24 hours with

Hnf4$\alpha$ and quantification of Hnf4$\alpha$ intensity in the nucleus, blue color indicates DAPI

signal, scale bar represents 20$\mu$m. (C) The level of WEs with various fatty acid chain length

in the serum of tumor bearing or control mice. (D) Heatmap showing the significantly

altered lipid species from the serum of control and tumor bearing mice. Data are presented

as mean ± s.e.m.; statistical analysis was performed using Student's t-test, and

significance analysis of microarrays was used for (D).

1055

1056

**References**

1.  Bakopoulos, D., et al., *Convergent insulin and TGF‑$\beta$ signalling drives cancer*
*cachexia by promoting aberrant fat body ECM accumulation in a Drosophila tumour*
*model.* The EMBO Reports, 2023. **24**(12): p. EMBR202357695.
2.  Fukumura, D., et al., *Tumor induction of VEGF promoter activity in stromal cells.*
Cell, 1998. **94**(6): p. 715-25.
3.  Quail, D.F. and J.A. Joyce, *Microenvironmental regulation of tumor progression and*
*metastasis.* Nat Med, 2013. **19**(11): p. 1423-37.
4.  Winkler, J., et al., *Concepts of extracellular matrix remodelling in tumour*
*progression and metastasis.* Nat Commun, 2020. **11**(1): p. 5120.
5.  Anderson, N.M. and M.C. Simon, *The tumor microenvironment.* Curr Biol, 2020.
**30**(16): p. R921-r925.
6.  Chen, X., et al., *Neural Circuitries between the Brain and Peripheral Solid Tumors.*
Cancer Res, 2024. **84**(21): p. 3509-3521.
7.  Goncalves, M.D., et al., *Fenofibrate prevents skeletal muscle loss in mice with lung*
*cancer.* Proceedings of the National Academy of Sciences, 2018. **115**(4): p. E743-
E752.
8.  Schutz, Y., *Protein turnover, ureagenesis and gluconeogenesis.* Int J Vitam Nutr Res,
2011. **81**(2-3): p. 101-7.
9.  Lee, J.S., et al., *Urea Cycle Dysregulation Generates Clinically Relevant Genomic*
*and Biochemical Signatures.* Cell, 2018. **174**(6): p. 1559-1570 e22.
10. Al-Zoughbi, W. and G. Hoefler, *Tumor Macroenvironment: An Update.* Pathobiology,
2020. **87**(2): p. 58-60.

11. Goldman, O., et al., *Early Infiltration of Innate Immune Cells to the Liver Depletes*
*HNF4alpha and Promotes Extrahepatic Carcinogenesis.* Cancer Discov, 2023. **13**(7):
p. 1616-1635.
12. Liu, Y., et al., *Hepatic gluconeogenesis and PDK3 upregulation drive cancer*
*cachexia in flies and mice.* Nat Metab, 2025. **7**(4): p. 823-841.
13. Grossberg, A.J., et al., *Interleukin 6-independent metabolic reprogramming as a*
*driver of cancer-related fatigue.* Brain, Behavior, and Immunity, 2020. **88**: p. 230-241.
14. Fearon, K.C., et al., *Elevated circulating interleukin-6 is associated with an acute-*
*phase response but reduced fixed hepatic protein synthesis in patients with cancer.*
Ann Surg, 1991. **213**(1): p. 26-31.
15. Wang, G., et al., *Tumour extracellular vesicles and particles induce liver metabolic*
*dysfunction.* Nature, 2023. **618**(7964): p. 374-382.
16. Jin, H.-R., et al., *Lipid metabolic reprogramming in tumor microenvironment: from*
*mechanisms to therapeutics.* Journal of Hematology & Oncology, 2023. **16**(1): p.
103.
17. Wang, H., et al., *CD36-mediated metabolic adaptation supports regulatory T cell*
*survival and function in tumors.* Nat Immunol, 2020. **21**(3): p. 298-308.
18. Wu, L., et al., *RIPK3 Orchestrates Fatty Acid Metabolism in Tumor-Associated*
*Macrophages and Hepatocarcinogenesis.* Cancer Immunol Res, 2020. **8**(5): p. 710-
721.
19. Veglia, F., et al., *Fatty acid transport protein 2 reprograms neutrophils in cancer.*
Nature, 2019. **569**(7754): p. 73-78.
20. Patel, P.H. and B.A. Edgar, *Tissue design: How Drosophila tumors remodel their*
*neighborhood.* Seminars in Cell & Developmental Biology, 2014. **28**: p. 86-95.
21. Kwon, Y., et al., *Systemic organ wasting induced by localized expression of the*
*secreted insulin/IGF antagonist ImpL2.* Dev Cell, 2015. **33**(1): p. 36-46.
22. Song, W., et al., *Tumor-Derived Ligands Trigger Tumor Growth and Host Wasting via*
*Differential MEK Activation.* Dev Cell, 2019. **48**(2): p. 277-286 e6.
23. Konishi, T., et al., *The K-ras gene regulates vascular endothelial growth factor gene*
*expression in non-small cell lung cancers.* Int J Oncol, 2000. **16**(3): p. 501-11.
24. Sunaga, N., et al., *Targeting Oncogenic KRAS in Non-Small-Cell Lung Cancer.*
Cancers (Basel), 2021. **13**(23).
25. Brooks, G.D., et al., *IL6 Trans-signaling Promotes KRAS-Driven Lung Carcinogenesis.*
Cancer Res, 2016. **76**(4): p. 866-76.
26. Olivares-Hernández, A., et al., *Current and future perspectives in clinical practice in*
*KRAS-mutated non-small cell lung cancer: a literature review.* Precision Cancer
Medicine, 2024. **7**.
27. Bilder, D., et al., *Tumour-host interactions through the lens of Drosophila.* Nat Rev
Cancer, 2021. **21**(11): p. 687-700.
28. Droujinine, I.A. and N. Perrimon, *Interorgan Communication Pathways in Physiology:*
*Focus on Drosophila.* Annu Rev Genet, 2016. **50**: p. 539-570.
29. Gutierrez, E., et al., *Specialized hepatocyte-like cells regulate Drosophila lipid*
*metabolism.* Nature, 2007. **445**(7125): p. 275-80.

30. Chatterjee, D., et al., *Control of metabolic adaptation to fasting by dILP6-induced insulin signaling in Drosophila oenocytes.* Proc Natl Acad Sci U S A, 2014. **111**(50): p. 17959-64.

31. Billeter, J.C., et al., *Specialized cells tag sexual and species identity in Drosophila melanogaster.* Nature, 2009. **461**(7266): p. 987-91.

32. Ferveur, J.F., et al., *Genetic feminization of pheromones and its behavioral consequences in Drosophila males.* Science, 1997. **276**(5318): p. 1555-8.

33. Blomquist, G.J., A.J. Chu, and S. Remaley, *Biosynthesis of wax in the honeybee, Apis mellifera L.* Insect Biochemistry, 1980. **10**(3): p. 313-321.

34. Qiu, Y., et al., *An insect-specific P450 oxidative decarbonylase for cuticular hydrocarbon biosynthesis.* Proc Natl Acad Sci U S A, 2012. **109**(37): p. 14858-63.

35. Reed, J.R., et al., *Unusual mechanism of hydrocarbon formation in the housefly: cytochrome P450 converts aldehyde to the sex pheromone component (Z)-9-tricosene and CO2.* Proceedings of the National Academy of Sciences, 1994. **91**(21): p. 10000-10004.

36. Wicker-Thomas, C., et al., *Flexible origin of hydrocarbon/pheromone precursors in Drosophila melanogaster.* J Lipid Res, 2015. **56**(11): p. 2094-101.

37. Wang, Z., et al., *Evolution of a fatty acyl-CoA elongase underlies desert adaptation in Drosophila.* Sci Adv, 2023. **9**(35): p. eadg0328.

38. Labeur, C., R. Dallerac, and C. Wicker-Thomas, *Involvement of desat1 gene in the control of Drosophila melanogaster pheromone biosynthesis.* Genetica, 2002. **114**(3): p. 269-74.

39. Storelli, G., et al., *Drosophila HNF4 Directs a Switch in Lipid Metabolism that Supports the Transition to Adulthood.* Dev Cell, 2019. **48**(2): p. 200-214 e6.

40. Ghabrial, A., et al., *Branching morphogenesis of the Drosophila tracheal system.* Annu Rev Cell Dev Biol, 2003. **19**: p. 623-47.

41. Li, Z., et al., *Trachea-derived dpp controls adult midgut homeostasis in Drosophila.* Dev Cell, 2013. **24**(2): p. 133-43.

42. Tamamouna, V., et al., *Remodelling of oxygen-transporting tracheoles drives intestinal regeneration and tumorigenesis in Drosophila.* Nat Cell Biol, 2021. **23**(5): p. 497-510.

43. Perochon, J., et al., *Dynamic adult tracheal plasticity drives stem cell adaptation to changes in intestinal homeostasis in Drosophila.* Nat Cell Biol, 2021. **23**(5): p. 485-496.

44. Kwon, Y., et al., *Systemic Organ Wasting Induced by Localized Expression of the Secreted Insulin/IGF Antagonist ImpL2.* Developmental Cell, 2015. **33**(1): p. 36-46.

45. Patel, S., D.R. Nelson, and A.G. Gibbs, *Chemical and physical analyses of wax ester properties.* J Insect Sci, 2001. **1**: p. 4.

46. Palanker, L., et al., *Drosophila HNF4 Regulates Lipid Mobilization and β-Oxidation.* Cell Metabolism, 2009. **9**(3): p. 228-239.

47. Xu, J., et al., *Protein visualization and manipulation in Drosophila through the use of epitope tags recognized by nanobodies.* eLife, 2022. **11**: p. e74326.

48. Bradley, M.E., et al., *Potent and Efficacious Inhibition of CXCR2 Signaling by*
*Biparatopic Nanobodies Combining Two Distinct Modes of Action*. Molecular
Pharmacology, 2015. **87**(2): p. 251-262.

49. Palanker, L., et al., *Drosophila HNF4 regulates lipid mobilization and beta-oxidation*.
Cell Metab, 2009. **9**(3): p. 228-39.

50. Ghosh, A.C., et al., *Drosophila PDGF/VEGF signaling from muscles to hepatocyte-*
*like cells protects against obesity*. eLife, 2020. **9**: p. e56969.

51. Zhang, Y., et al., *Rheb is a direct target of the tuberous sclerosis tumour suppressor*
*proteins*. Nature Cell Biology, 2003. **5**(6): p. 578-581.

52. Sopko, R., et al., *A systems-level interrogation identifies regulators of Drosophila*
*blood cell number and survival*. PLoS Genet, 2015. **11**(3): p. e1005056.

53. Tran, T.A., et al., *Platelet-derived growth factor/vascular endothelial growth factor*
*receptor inactivation by sunitinib results in Tsc1/Tsc2-dependent inhibition of*
*TORC1*. Mol Cell Biol, 2013. **33**(19): p. 3762-79.

54. Cruciani, G., et al., *Diagnostic application of lipidomics fingerprints to bladder*
*carcinoma*. Translational Cancer Research, 2017. **6**(6): p. 1197-1206.

55. Simcox, J., et al., *Global Analysis of Plasma Lipids Identifies Liver-Derived*
*Acylcarnitines as a Fuel Source for Brown Fat Thermogenesis*. Cell Metabolism,
2017. **26**(3): p. 509-522.e6.

56. Palm, W., et al., *Lipoproteins in Drosophila melanogaster--assembly, function, and*
*influence on tissue lipid composition*. PLoS Genet, 2012. **8**(7): p. e1002828.

57. Vonolfen, M.C., et al., *Drosophila HNF4 acts in distinct tissues to direct a switch*
*between lipid storage and export in the gut*. Cell Rep, 2024. **43**(9): p. 114693.

58. Khaliullina, H., et al., *Patched regulates Smoothened trafficking using lipoprotein-*
*derived lipids*. Development, 2009. **136**(24): p. 4111-21.

59. Locke, M., *Pore canals and related structures in insect cuticle*. J Biophys Biochem
Cytol, 1961. **10**(4): p. 589-618.

60. Moussian, B., *Recent advances in understanding mechanisms of insect cuticle*
*differentiation*. Insect Biochem Mol Biol, 2010. **40**(5): p. 363-75.

61. Moussian, B., et al., *Cuticle differentiation during Drosophila embryogenesis*.
Arthropod Struct Dev, 2006. **35**(3): p. 137-52.

62. Jaspers, M.H., et al., *The fatty acyl-CoA reductase Waterproof mediates airway*
*clearance in Drosophila*. Dev Biol, 2014. **385**(1): p. 23-31.

63. Chatterjee, N. and N. Perrimon, *What fuels the fly: Energy metabolism in*
*<i>Drosophila</i> and its application to the study of obesity and diabetes*. Science
Advances, 2021. **7**(24): p. eabg4336.

64. Huang, K., et al., *RiboTag translatomic profiling of Drosophila oenocytes under aging*
*and induced oxidative stress*. BMC Genomics, 2019. **20**(1): p. 50.

65. Rera, M., et al., *Modulation of longevity and tissue homeostasis by the Drosophila*
*PGC-1 homolog*. Cell Metab, 2011. **14**(5): p. 623-34.

66. Radi, S.H., et al., *HNF4alpha isoforms: the fraternal twin master regulators of liver*
*function*. Front Endocrinol (Lausanne), 2023. **14**: p. 1226173.

67. Metzger, S., et al., *Orphan receptor HNF-4 and bZip protein C/EBP alpha bind to*
*overlapping regions of the apolipoprotein B gene promoter and synergistically*
*activate transcription.* J Biol Chem, 1993. **268**(22): p. 16831-8.
68. Adamson, A.W., et al., *Hepatocyte nuclear factor-4alpha contributes to*
*carbohydrate-induced transcriptional activation of hepatic fatty acid synthase.*
Biochem J, 2006. **399**(2): p. 285-95.
69. Meng, J., et al., *Identification of HNF-4α as a key transcription factor to promote*
*ChREBP expression in response to glucose.* Scientific Reports, 2016. **6**(1): p. 23944.
70. Bideyan, L., R. Nagari, and P. Tontonoz, *Hepatic transcriptional responses to fasting*
*and feeding.* Genes Dev, 2021. **35**(9-10): p. 635-657.
71. Ding, G., et al., *Coordination of tumor growth and host wasting by tumor-derived*
*Upd3.* Cell Rep, 2021. **36**(7): p. 109553.
72. Xu, J., et al., *Mechanistic characterization of a Drosophila model of paraneoplastic*
*nephrotic syndrome.* Nature Communications, 2024. **15**(1): p. 1241.
73. Duvel, K., et al., *Activation of a metabolic gene regulatory network downstream of*
*mTOR complex 1.* Mol Cell, 2010. **39**(2): p. 171-83.
74. Tiebe, M., et al., *REPTOR and REPTOR-BP Regulate Organismal Metabolism and*
*Transcription Downstream of TORC1.* Dev Cell, 2015. **33**(3): p. 272-84.
75. Hoch, R.V. and P. Soriano, *Roles of PDGF in animal development.* Development,
2003. **130**(20): p. 4769-84.
76. Holmes, D.I. and I. Zachary, *The vascular endothelial growth factor (VEGF) family:*
*angiogenic factors in health and disease.* Genome Biol, 2005. **6**(2): p. 209.
77. Jiang, H., et al., *Cytokine/Jak/Stat signaling mediates regeneration and homeostasis*
*in the Drosophila midgut.* Cell, 2009. **137**(7): p. 1343-55.
78. Buchon, N., et al., *Drosophila intestinal response to bacterial infection: activation of*
*host defense and stem cell proliferation.* Cell Host Microbe, 2009. **5**(2): p. 200-11.
79. Apidianakis, Y., et al., *Synergy between bacterial infection and genetic*
*predisposition in intestinal dysplasia.* Proc Natl Acad Sci U S A, 2009. **106**(49): p.
20883-8.
80. Amcheslavsky, A., et al., *Enteroendocrine Cells Support Intestinal Stem-Cell-*
*Mediated Homeostasis in <em>Drosophila</em>.* Cell Reports, 2014. **9**(1): p. 32-39.
81. Lin, G., N. Xu, and R. Xi, *Paracrine Wingless signalling controls self-renewal of*
*Drosophila intestinal stem cells.* Nature, 2008. **455**(7216): p. 1119-1123.
82. Schönfeld, P. and G. Reiser, *Brain Lipotoxicity of Phytanic Acid and Very Long-chain*
*Fatty Acids. Harmful Cellular/Mitochondrial Activities in Refsum Disease and X-*
*Linked Adrenoleukodystrophy.* Aging Dis, 2016. **7**(2): p. 136-49.
83. Baarine, M., et al., *Evidence of oxidative stress in very long chain fatty acid--treated*
*oligodendrocytes and potentialization of ROS production using RNA interference-*
*directed knockdown of ABCD1 and ACOX1 peroxisomal proteins.* Neuroscience,
2012. **213**: p. 1-18.
84. Zarrouk, A., et al., *Induction of mitochondrial changes associated with oxidative*
*stress on very long chain fatty acids (C22:0, C24:0, or C26:0)-treated human*
*neuronal cells (SK-NB-E).* Oxid Med Cell Longev, 2012. **2012**: p. 623257.

85.    Sargent, J.R., R.R. Gatten, and R. McIntosh, *Wax esters in the marine environment —*
1251            *their occurrence, formation, transformation and ultimate fates*. Marine Chemistry,
1252            1977. **5**(4): p. 573-584.
1253    86.    Rosca, M.G., et al., *Oxidation of Fatty Acids Is the Source of Increased Mitochondrial*
1254            *Reactive Oxygen Species Production in Kidney Cortical Tubules in Early Diabetes*.
1255            Diabetes, 2012. **61**(8): p. 2074-2083.
1256    87.    Tamura, K., et al., *Novel lipogenic enzyme ELOVL7 is involved in prostate cancer*
1257            *growth through saturated long-chain fatty acid metabolism*. Cancer Res, 2009.
1258            **69**(20): p. 8133-40.
1259    88.    Karlsson, M., et al., *A single–cell type transcriptomics map of human tissues*.
1260            Science Advances, 2021. **7**(31): p. eabh2169.
1261    89.    Hirota, K., et al., *A Combination of HNF-4 and Foxo1 Is Required for Reciprocal*
1262            *Transcriptional Regulation of Glucokinase and Glucose-6-phosphatase Genes in*
1263            *Response to Fasting and Feeding *<sup></sup>*. Journal of Biological Chemistry,
1264            2008. **283**(47): p. 32432-32441.
1265    90.    Koyama, T. and C.K. Mirth, *Growth-Blocking Peptides As Nutrition-Sensitive Signals*
1266            *for Insulin Secretion and Body Size Regulation*. PLoS Biol, 2016. **14**(2): p. e1002392.
1267    91.    Pallares-Cartes, C., G. Cakan-Akdogan, and Aurelio A. Teleman, *Tissue-Specific*
1268            *Coupling between Insulin/IGF and TORC1 Signaling via PRAS40 in*
1269            *<em>Drosophila</em>*. Developmental Cell, 2012. **22**(1): p. 172-182.
1270    92.    Chatterjee, D., et al., *Control of metabolic adaptation to fasting by dILP6-induced*
1271            *insulin signaling in <i>Drosophila</i> oenocytes*. Proceedings of the National
1272            Academy of Sciences, 2014. **111**(50): p. 17959-17964.
1273    93.    Chung, H., et al., *A Single Gene Affects Both Ecological Divergence and Mate*
1274            *Choice in <i>Drosophila</i>*. Science, 2014. **343**(6175): p. 1148-1151.
1275    94.    Wang, Z., et al., *Evolution of a fatty acyl–CoA elongase underlies desert*
1276            *adaptation in <i>Drosophila</i>*. Science Advances, 2023. **9**(35): p. eadg0328.
1277    95.    Montoya-Zegarra, J.A., et al., *AutoTube: a novel software for the automated*
1278            *morphometric analysis of vascular networks in tissues*. Angiogenesis, 2019. **22**(2):
1279            p. 223-236.
1280    96.    Ji, H., et al., *LKB1 modulates lung cancer differentiation and metastasis*. Nature,
1281            2007. **448**(7155): p. 807-810.
1282

A

Ovary size

*Esg-LexA>yki*     -    +    +
*PromE>TSC1,2*     -    -    +

B

Abdomen area

*Esg-LexA>yki*     -    +    +
*PromE>TSC1,2*     -    -    +

C

Dry starvation survival

****

*mFas>Luc*
*mFas>TSC1,2*

Hours

D

*PromE$^{gs}$> PRAS40*

-RU          +RU

E

DMSO          Rapamycin 200uM

**A** adult fly (female,body,no head), yki tumor
DEG Values :8D_Yki_vs_Wt

**B** *Pvf1*

**C**

Ovary size

*Esg>*
Luc-i   Pvf1-i   yki;Luc-i   yki;Pvf1-i

**D**

% of GFP+ region

*Esg>*
Luc-i   yki;Luc-i   yki;Pvf1-i

**E**

Ovary size

*Esg-LexA>yki*    −   −   +   +
*PromE>PvR-i*     −   +   −   +

**F**

% of GFP+ region

*Esg-LexA>yki*    −   +   +
*PromE>PvR-i*     −   −   +

**A**

*Esg-LexA>* *ctrl*

*yki*

**B** # of branches

*Esg>* ctrl    yki

**C** Tube area

ctrl    yki

**D** Tube length

ctrl    yki