1   **Title:** A mechanism linking dopamine's roles in reinforcement, movement and motivation.
2

3   **Authors**:
4   Allison E Hamilos*[a,b], Isabella C Wijsman[b], Qinxin Ding[a,b], Pichamon Assawaphadungsit[a,b],
5   Zeynep Ozcan[a] and John A Assad*[b,c]
6

7   **Affiliations:**
8   [a]Whitehead Institute, Cambridge, Massachusetts, 02142, USA
9   [b]Department of Neurobiology, Harvard Medical School, Boston, 02115, USA
10   [c]Istituto Italiano di Tecnologia, Genova, Italy
11   *For correspondence: A.E.H. (ahamilos@wi.mit.edu) or J.A.A. (jassad@hms.harvard.edu)
12

Summary:
Dopamine neurons (DANs) play seemingly distinct roles in reinforcement,[1-3] motivation,[4,5] and
movement,[6,7] and DA-modulating therapies relieve symptoms across a puzzling spectrum of
neurologic and psychiatric symptoms.[8] Yet, the mechanistic relationship among these roles is
unknown. Here, we show DA's tripartite roles are causally linked by a process in which phasic
striatal DA rapidly and persistently recalibrates the propensity to move, a measure of vigor.
Using a self-timed movement task, we found that single exposures to reward-related DA
transients (both endogenous and exogenously-induced) exerted one-shot updates to
movement timing—but in a surprising fashion. Rather than reinforce specific movement times,
DA transients quantitatively *changed* movement timing on the next trial, with larger transients
leading to earlier movements (and smaller to later), consistent with a stochastic search process
that calibrates the frequency of movement. Both abrupt and gradual changes in external and
internal contingencies—such as timing criterion, reward content, and satiety state—caused
changes to the amplitude of DA transients that causally altered movement timing. The rapidity
and bidirectionality of the one-shot effects are difficult to reconcile with gradual synaptic
plasticity, and instead point to more flexible cellular mechanisms, such as DA-dependent
modulation of neuronal excitability. Our findings shed light on how natural reinforcement, as
well as DA-related disorders such as Parkinson's disease, could affect behavioral vigor.
31

MAIN TEXT:
Both electrical[9] and optogenetic activation[10-12] of dopamine neurons (DANs) can gradually
reinforce actions.[10-15] It is widely believed DANs drive gradual learning via phasic dopamine (DA)
transients, with ongoing debate over what these transients represent in reinforcement learning
frameworks (e.g., causal association,[16] learning rate,[3] value,[4,5] reward/performance-prediction
errors[1,2,17-22], etc.). Yet, multiple lines of evidence suggest DA also plays "real-time" roles in motor
control[6,7,23-25] and motivation,[5,25] which are thought to be distinct from DA's role in gradual
learning.[6,22,26] For example, tonic DA levels are associated with movement vigor, with both tonic
DA and vigor increasing in reward-rich environments.[6,26] Exogenous DA/DAN manipulations also
causally modulate the probability,[7,23] number,[23] acceleration[23] and variability[11] of movements,
and DA-modulating drugs are used ubiquitously in clinical settings as a "tuning dial" to adjust
patients' propensity to move.[8,27] DA repletion/agonists increase movement propensity (e.g., to

treat Parkinson's disease), whereas DA antagonists have the opposite effect (e.g., to treat mania,
Huntington's disease, delirium, etc.). Yet, it remains a mystery whether *endogenous*, reward-
related DA transients are related to DA's role in movement.[23,26,28-30]
48
49 Because DA drugs bidirectionally influence the propensity to move, we suspected DA's roles in
50 reinforcement and movement might be mechanistically linked by a tuning process, whereby
51 endogenous DA transients calibrate the propensity to move based on feedback from recent
52 reward history. Individual DANs encode both reward- and movement-related information,[30,31]
53 suggesting these functions may be intertwined or somehow co-dependent.
54
55 We recently showed DANs play a causal, real-time role in movement initiation while mice
56 perform a self-timed movement task.[7] Given this task also shows outcome-related DA signaling
57 and quantitative behavioral changes related to motivational state, we realized this paradigm
58 would allow us to study DA's tripartite roles in the same context, allowing us to test the
59 calibration hypothesis.
60
Task for quantifying propensity to move
We trained head-fixed, water-deprived mice to perform a self-timed movement task[7] (**Fig 1a**). On
each trial, mice had to estimate elapsed time to decide when to lick after a brief cue (tone/LED).
Mice received a juice reward if (and only if) the first-lick occurred during a hidden reward window
(3.3-7s after the cue), which had to be discovered through trial-and-error. Only the first lick
determined the trial outcome (unrewarded/rewarded). Mice learned to self-time licks, with the
timing distribution stabilizing within 10-20 training sessions (**Fig 1b**, **Extended Data Fig 1**).
68
Timing behavior is naturally variable,[32,33] with both humans and well-trained animals making
many relatively early and late movements during the same session.[7,34-36] We exploited this
variability to ask how DAN signaling related to trial-by-trial updates to movement timing. DA has
long been implicated in timing behavior. For example, Parkinsonism/DA antagonists cause
systematically late timing,[33,37-39] whereas DA agonists cause early timing.[40,41] These findings
indicate that movement timing can serve as a quantitative, operational definition of what we
mean by "propensity to move" (earlier timing=higher propensity; later timing=lower). We thus
hypothesized that reward-related DAN signaling might adjust timing based on task reward/error
feedback.
78
Outcome-related DA signals
We used fiber photometry to monitor intracellular Ca^{++} in genetically-defined DANs expressing
GCaMP6f, recording from both cell bodies (substantia nigra, SNc: n=4 DAT-cre mice; ventral
tegmental area, VTA: n=3 mice) and axon terminals (dorsolateral striatum, DLS, n=4 mice). DANs
exhibited RPE-like transients in the timing task: unrewarded first-licks were followed by small dips
in intracellular Ca^{++}; rewarded trials were followed by large increases (**Fig 1c**). These signals were
surprisingly persistent, decaying for >10s after the reward, much longer than GCaMP6f's off-
kinetics (t½ <300 ms at 37°).[42]
87

88   The same basic RPE-like features were present across DAN recording sites (SNc, VTA, DLS) and in
89   striatal DA release (**Fig 1d**), which we monitored with the fluorescent DA sensors dLight or DA2m,
90   expressed in either dorsolateral (DLS) or ventrolateral dorsal striatum (VLS[43,44]; **Extended Data**
91   **Fig 2**).
92
93   These persistent RPE-like signals bled into the next trial (**Fig 1e**), becoming continuous with
94   differences in baseline DAN calcium signals that were correlated with timing on the next trial
95   (**Fig 1f**; Pearson's r: –0.79, 95% CI: [-0.84, -0.73]), suggesting outcome-related DA might influence
96   timing based on task feedback from the previous trial.
97
"Sloshing" strategy for timing updates
There were three possible trial outcomes: 1) early (no reward), 2) rewarded, or 3) no lick. Because
mice only rarely did not lick, we focused analyses on early versus rewarded outcomes. Intuitively,
an early trial should prompt slower timing on the next trial. Unrewarded DA dips persisted for
seconds and were associated with lower pre-cue DAN signals—which were indeed systematically
correlated with later timing on the next trial. Conversely, rewarded trials were associated with
disproportionate DA increases. Because exogenous DA can gradually reinforce actions,[10-15] we
expected animals would move at about the same time on the next trial. Surprisingly, this was not
the case. After rewards, DA signals remained elevated for >10s, and elevated baseline was
systematically correlated with *earlier* timing on the next trial. Rather than repeating the rewarded
time, mice *sped up*. This "sloshing" strategy may seem unintuitive, but is not necessarily irrational:
A thirsty mouse may want to move at the earliest possible rewarded time, but this time is
unknown—mice had to discover it through experience. By slowing down after early trials and
speeding up after rewarded trials, this sloshing effect can cause the timing distribution to build
up around the earliest possible rewarded time.
113
A potential confound is that, on average, relatively late trials might—by chance—tend to be
followed by earlier trials (and relatively early followed by later). This "regression to the median"
phenomenon could occur even in the absence of goal-directed updates to timing. We thus
compared the change in the movement time between consecutive trials, $\Delta T_{(n+1)-n}$, to the change
in movement time expected by chance from regression to the median (n=17 mice, 155 sessions;
Supplemental Methods). $\Delta T_{(n+1)-n}$ was in excess of chance: as suggested by the DAN signals,
animals indeed moved significantly earlier after rewarded trials and significantly later after
unrewarded trials (update after rewards: –317ms, 95% CI: [–406ms, –223ms]; update after early
trials: +74ms, 95% CI: [+41ms, +106ms]; 10,000x bootstrap procedure; **Fig 1g**).
123
DA transients explain *changes* in timing
To determine if DA explained sloshing updates, we derived a generalized linear model (GLM)
predicting the *change* in the movement time on the next trial, $\Delta T_{(n+1)-n}$, from the amplitude of
outcome-related DA transients (**Fig 2a**). This amplitude was encoded as the DA signal averaged
over 0-500ms following the previous trial's first-lick ("DA", **Fig 2b**). We controlled for optical
artifacts and ongoing body movements by including the tdTomato ("tdt") photometry channel,
accelerometer and/or neck EMG signals as predictors.
131

Across all recording sites (and for both GCaMP and DA sensors), DA was independently predictive
of $\Delta T_{(n+1)-n}$ (**Fig 2c**; **Extended Data Fig 3**), in excess of optical artifacts and ongoing body
movements. The larger the DA transient amplitude, the larger the timing update on the next trial
(highest DA: largest early-shifts; lowest DA: largest late-shifts).

The amplitude of the DA transient was strongly correlated with trial outcome (which was also a
strong predictor of $\Delta T_{(n+1)-n}$). Although this would not preclude a causal role for DA, we tested if
additional information was present in DA by fitting separate models for early and rewarded trials.
Both early and rewarded DA transient amplitudes showed considerable trial-by-trial variability,
which had additional predictive power beyond correlation with trial outcome (**Fig 2d**, **Extended
Data Fig 3**). These effects were also not explained away by a tendency to revert toward the
median movement time (DA coefficient significantly larger than expected by chance in shuffled
trial order model (p=7.2e-9, Wilcoxon rank sum test; **Fig 2e**; **Extended Data Fig 4**).

Outcome-related DA was thus positioned to make "one-shot" learning updates to timing. We use
the machine learning term "one-shot learning" here because a *single exposure* to an endogenous
DA transient was associated with a quantitative update to behavior. To our knowledge, DA
learning effects based on variability in endogenous DA transients have not previously been
observed.

DA causes one-shot timing updates
To test if outcome-related DA could cause one-shot timing updates, we virally expressed
channelrhodopsin (ChR2) in genetically defined DANs for optogenetic manipulations (n=4 DAT-
cre mice, **Fig 3a**). To modulate outcome-related DAN signaling, we triggered stimulation in the
SNc at the time of the first-lick on 30% of randomly interleaved trials (for inhibition: see
Supplemental Note 1). Stimulation was calibrated[7,30] outside the context of the task to
recapitulate endogenous rewarded DA transients (**Fig 3b**), which matched the amplitude and
width of unstimulated rewarded transients during the task (**Fig 3c; Extended Data Fig 5a**).

Stimulation on rewarded trials slightly increased DA transient amplitude, but on average, this
remained within the limits of unstimulated reward transients on single trials measured during the
same session (**Extended Data Fig 5c**). Control stimulations in mice not expressing opsin indicated
that rdLight1 recordings were not contaminated by blue light-induced photoswitching[45] artifacts
("No Opsin" mice, **Fig 3d; Extended Data Fig 5d**).

DAN activation caused mice to move significantly earlier than expected on the next trial, with 4/4
mice showing early-shifting (**Fig 3e-f**; see **Methods**; mean $\Delta\Delta T_{(n+1)-n}$: -88ms, p=0.0019, 10,000x
bootstrap test, n=4 mice, 40 sessions). By contrast, stimulation in No Opsin mice had no significant
effect on $\Delta T_{(n+1)-n}$ (mean $\Delta\Delta T_{(n+1)-n}$: -6ms, p=0.40, 10,000x bootstrap test, n=4 mice, 40 sessions).
These results were consistent across 3 additional complementary statistical analyses (**Extended
Data Fig 6**).

Activation also caused subtle early-shifting of the *entire* timing distribution across the session
compared to interleaved sham sessions (**Fig 3g**; p<0.025, 1000x bootstrap test; **Extended Data**

Fig 5b,e). This raised the possibility that outcome-related DA may normally control the "set-point"
of the movement timing distribution, which we operationally define as the median first-lick time.
178
DA explains the timing set-point
Surprisingly, across timing paradigms, both rodents and non-human primates frequently tend to
move too early, at the expense of reward[7,34-36,46] (**Extended Data Fig 7a**). In our mice, this early-
bias developed over days of experience (**Extended Data Fig 1**) and persisted even in expert mice
with >70 sessions' experience.[7] Outcome-related DA also appeared highly skewed in this task,
with disproportionately large reward-related transients compared to small unrewarded dips
(**Extended Data Fig 7b**). We observed that high amplitude (rewarded) DA transients caused much
larger early-shifts compared to relatively small late-shifts after DA dips, and overall, mice made
larger updates to timing after rewarded trials compared to unrewarded trials (median –317ms vs
74ms; **Fig 1g**). Together, this suggests bias in DA signaling may explain early-biased timing behavior.
189
190 But behavior was not early skewed for the entire session—all mice showed marked timing
191 changes *within* each session, initially moving too early, but progressively shifting to later times
192 (**Fig 4a-b; Extended Data Fig 7c**). If allowed to work *ad libitum*, the timing set-point ultimately
193 shifted into the reward window—suggestive of timing calibration. (Importantly, the sloshing
194 model GLMs were robust to this non-stationarity, **Extended Data Fig 4c**).
195
The Sloshing Model suggested that changes in outcome-related DA could explain changes in the
timing set-point. Across each session, as the set-point shifted later, 1) rewarded trials were
followed by progressively smaller timing updates (median timing change (in excess of chance),
beginning half of session: -400ms, 95% CI: [-519ms, -270ms]; **ending: -271ms**, 95% CI: [-403ms,
-143ms]; n=17 mice, 155 sessions; **Extended Data Fig 7d**); and, 2) rewarded DA transient
amplitudes progressively declined (**Fig 4c; Extended Data Fig 7e-f, Extended Data Fig 8**; *for*
*unrewarded transient effects: see **Supplemental Note 2***). Importantly, this DA decline was not
explained-away by bleaching or optical artifacts (**Extended Data Fig 7g**, **Supplemental Methods**).
204
Rewarded DA amplitude was tightly correlated with the timing set-point (median Pearson's r:
–0.81; **Fig 4d**). This was not the case for movement-related/optical artifacts (median Pearson's r:
+0.19, **Extended Data Fig 7h**). Strikingly, shifts were not always gradual—rewarded DA amplitude
decreased suddenly during abrupt shifts of the set-point into the reward window (**Fig 4e**; R^2=0.65,
p=0.0003), which was also not explained away by optical/movement related artifacts (R^2=0.19,
p=0.10; **Extended Data Fig 7i**).
211
DA calibrates the propensity to move
Altogether, across sessions and recording sites/modalities, rewarded DA transients decreased in
amplitude as timing behavior became more accurate. DA transients also controlled trial-by-trial
timing updates, and these updates became progressively smaller as behavior became more
accurate (**Extended Data Fig 7d**). Together, these findings suggest a calibration process for motor
timing. This adaptive decrease in stochastic learning is reminiscent of adaptive stochastic gradient
descent algorithms.[47] However, because rewarded DA and set-point tended to change
monotonically across the session, we tested several complementary manipulations to "break the

monotonicity" between these co-variates. Each manipulation produced marked changes in the
animals' timing strategy, including late-to-early timing shifts. Importantly, in all settings, rewarded
DA transients explained bidirectional changes in the timing set-point, as follows:
223
Different reward identities. First, we serendipitously discovered that well-trained mice continued
to participate even when juice rewards were replaced by the usual "correct" tone (**"No Juice"**
Task; solenoid valve disconnected)—but set-point dynamics were strikingly different (**Fig 4f-g**;
Extended Data Fig 7j). Mice still showed early bias for the first ~150 trials, but unexpectedly, the
set-point suddenly and dramatically shifted into the reward window—significantly earlier in the
session than in the +Juice task (p=0.001, Wilcoxon rank sum test; 5 mice, 18 paired sessions,
Fig 4h). Mice continued to work for as many as 1000 trials under these conditions with correct
timing on most trials, exhibiting much more "accurate" behavior than the same animals
performing the +Juice task on interleaved sessions (same mice as **Figs 1b, 4b-d**).
233
Surprisingly, mice still showed "rewarded" DA transients in the No Juice task (in response to the
correct tone, **Fig 4i**), but these were narrower and smaller in amplitude than reward transients in
the +Juice task (**Fig 4j**). Moreover, reward-transient amplitude declined significantly earlier in the
session during No Juice sessions (n=5 mice, 18 paired sessions; p=0.0073, Wilcoxon rank sum test,
Fig 4k, Extended Data Fig 7k-m), and the timing of this abrupt, early decline explained the rapid
shift in timing set-point (median Pearson's r: –0.78; **Fig 4l; Extended Data Fig 7o**). These effects
were not explained by changes in ongoing body movements or optical artifacts (**Extended Data**
Fig 7l,n).
242
243 Interestingly, the persistence of high amplitude rewarded transients in the +Juice task suggests
244 that juice skewed normal timing updates: despite apparent motivation, mice received many fewer
245 rewards/time compared to the No Juice task. The resulting early timing bias is reminiscent of the
246 effects of DA-modulating drugs of abuse on timing behavior, such as amphetamines.[40]
247
Abrupt changes in reward identity. Second, we trained mice to alternate between blocks of trials
with either No Juice rewards (correct tone only) or +Juice. During No Juice blocks, the timing set-
point rapidly late-shifted into the reward window. This was rapidly reversed in +Juice blocks, in
which the timing set-point rapidly shifted significantly earlier (n=3 mice, 5 block changes; p=0.032,
Wilcoxon rank sum test; **Fig 5a**). Again, the amplitude of rewarded DA transients explained these
set-point shifts (median Pearson's r: –0.89; **Fig 5b-c**).
254
The rapid changes in DA-transient amplitude might be correlated—but not causal—to abrupt set-
point shifts. Therefore, we also had mice alternate between blocks of trials with either No Juice
or No Juice+Stimulation, in which DANs were optogenetically activated at the time of first-licks
that occurred either within the reward window or on all trials (n=3 mice, 20 block changes;
Extended Data Fig 9a-e). Stimulation was calibrated to recapitulate usual juice-related DA
transients without delivering liquid rewards (as in **Fig 3b**). The set-point rapidly late-shifted during
No Juice blocks, but then rapidly and significantly early-shifted during +Stimulation blocks (off-on:
p=0.001, on-off: p=0.007; Wilcoxon rank sum test; **Extended Data Fig 9b**), even when stimulation
was only delivered on rewarded trials (off-on: p=0.009, on-off: p=0.002; n=2 mice, 12 block

changes, **Fig 5d**). Stimulation did not change the set-point in control mice expressing No Opsin
(n=4 mice, 20 block changes; **Fig 5d; Extended Data Fig 9b-e**). Again, the amplitude of rewarded
DA transients explained the set-point shifts (median stim: −0.66, no opsin: −0.56; **Fig 5e-f**,
Extended Data Fig 9d-f). These results emphasize that rewarded DA transients induced *shifts* to
earlier times—rather than reinforcing the rewarded movement time.

Rapid satiation. Third, in addition to sudden shifts, the timing set-point also showed gradual late-
drift across sessions, potentially related to fatigue or satiation. The sloshing model suggested DA
could also explain state/motivation-dependent shifts. We thus tested whether *rapid* satiation
could change rewarded DA and set-point. After completing >=100 trials of the +Juice task and
receiving >=10 rewards, the task was paused, and mice were allowed to drink *ad libitum* until
they stopped licking available juice (n=3 mice, 15 sessions; range: 0.75-1.8mL (daily water
consumption for a water-deprived mouse is ~1mL); **Fig 5g-h**). The task was then resumed.
Rewarded DA and timing set-point remained highly anticorrelated throughout the session
(median Pearson's r: −0.70, **Fig 5i**), but rewarded DA decreased rapidly after the pause (p=8.5e-5,
Wilcoxon rank sum test; n=11 photometry sessions; **Extended Data Fig 9g**), and the timing set-
point showed significant late-shifting (p=1.7e-5, Wilcoxon rank sum test; **Extended Data Fig 9h**).
In contrast, in interleaved control sessions (empty syringe; no juice given during the pause),
rewarded DA did not show this precipitous decrease (p=0.0026; **Fig 5h**), and there was no
significant change in timing set-point (p=0.18; **Fig 5g**). Altogether, rapid juice consumption caused
subsequent rewarded DA transients to rapidly decrease in amplitude, which was associated with
late-shifting.

Changing timing constraints. Finally, fourth, to test if endogenous changes in rewarded DA
transients could re-calibrate the timing set-point in response to different environmental timing
contingencies, we changed the rewarded time window in blocks of trials ("Timeshift Task;" e.g.,
from 7-12s to 2-3.3s; n=7 mice; **Fig 5j**). The new reward window was not signaled and had to be
discovered through trial-and-error. We hypothesized increases in rewarded DA would cause early-
shifting to match the constraints of the task (and decreases, late-shifting). Because fatigue/satiety
should increase monotonically over the course of the session, we reasoned early-shifts would
disentangle goal-directed processes from fatigue/satiation.

The Timeshift Task produced pronounced shifts in the animals' timing behavior, including directed
changes in timing in relation to the new reward window (**Fig 5j, Extended Data Fig 9i**) and
exploratory behavior that appeared to anticipate block changes (**Extended Data Fig 9j**). The
amplitude of rewarded DA transients continued to explain the timing set-point (median Pearson's
r: −0.64; 7 mice, 54 sessions; **Fig 5k-l**), and intriguingly, *unrewarded* DA transients also became
more correlated with the set-point (median r: −0.36; **Fig 5l**). Early-shifts were accompanied by
increases in DA transient amplitude (and late-shifts by decreases), and these effects were not
explained away by optical artifacts (tdt: Rewarded trials median r: −0.20; unrewarded r: 0.10;
Extended Data Fig 9k).

Together, these results show that changes in reward-related DA cause shifts in the behavioral
timing set-point, both in relation to changing environmental conditions (such as changes in

environmental timing contingencies for reward or the presence/absence of juice rewards) and
internal state (such as gradual or rapid satiation). These findings demonstrate that DA
quantitatively and adaptively calibrates the propensity to move in response to both external and
internal constraints.

DISCUSSION
A cornerstone hypothesis of biological reinforcement learning is that reward-related DA
transients (whatever their nature[3,5,16,19,48,49]) lead to changes in future behavior. However, to our
knowledge, variability in *endogenous* reward-related DA transients has not been shown to relate
to variability in subsequent behavior, despite serious effort to detect such effects.[3,30,50,51] Here,
we show for the first time that the amplitude of native DA reward transients is quantitatively
related to one-shot, goal-directed updates to movement timing on the *next* trial, >10s later. This
suggests a functional link between reward and the calibration of the propensity to move (an
aspect of vigor), unifying two seemingly distinct aspects of DA function.[6] In addition, we found
that asymmetry in the relative amplitude of rewarded/unrewarded DA transients explained
asymmetry in one-shot updates, and that changes in the relative amplitude of these transients
could causally shift the timing set-point. This process, reminiscent of adaptive stochastic gradient
descent,[52] allowed the animals to adjust their propensity to move in response to changing
experimental contingencies (e.g., timing constraints). Changes in DA transients and behavior
could also be caused by changes in internal/motivational state (reward identity, satiety/fatigue),
demonstrating that flexibility in reward-related DA signaling mediates both strategic and
motivationally driven changes to movement timing. Together, DA's tripartite roles in
reinforcement, movement and motivation are mechanistically linked via this calibration process.

Relationship to reported rapid DA effects. Others have observed rapid, "real-time" effects of DA
on movement vigor—but at much shorter latency than the reward-related effects we report here
(within 100s of milliseconds before movement), and those studies specifically excluded reward-
related DA/DAN transients from analyses.[23,25] We previously found that ongoing ramping
DAN/DA dynamics *during* the self-timed interval (between the cue and the first-lick) predicted
self-timed movement initiation (**Fig 1f, right**).[7] Moreover, physiologically calibrated optogenetic
DAN activation throughout the self-timed interval could increase the moment-by-moment
probability of initiating the self-timed movement (and inhibition, decrease it)[7]—a real-time DAN
effect on movement timing. However, even before the trial started, differences in baseline DAN
calcium signals were systematically related to the timing of the upcoming movement,[7] suggesting
that baseline might reflect a putative "stochastic set-point" that modulates timing by controlling
the start-point and slope of the ramping process.[53-55]

Here, we show that this variability in DAN baseline can be traced back *tens of seconds* to variability
in the amplitude of the previous trial's outcome-related DA transient—and that variability in this
transient can causally update timing on the next trial, 10-25s in the future. The effects of the
outcome-related transient were in the same direction as prior real-time stimulation effects
(higher DA, earlier movement); thus, parsimony would suggest both outcome-related and
baseline/ramping DA exert the same mechanistic effect on timing. But, importantly, the results

presented here demonstrate that, in addition to real-time effects (sub-second to a few seconds),
even transient DAN signaling causes effects on movement timing that persist over tens of seconds.
This time-course has important mechanistic implications. In particular, the rapid, one-shot nature
is difficult to square with conventional views of incremental DA-dependent synaptic plasticity.[56]
356
Relationship to reported gradual DA effects. Whereas we found one-shot effects of endogenous
reward-related DA signals on the next-trial, other studies have found that *exogenous* DA
manipulations can reinforce actions, but typically only *gradually*, over many exposures.[10,12,14,57]
Such gradual effects could be viewed as consistent with incremental synaptic plasticity. In some
cases, DA-mediated reinforcement has been specifically reported as imperceptible on the order
of tens of consecutive trials.[2] One study found that optogenetic DAN inhibition throughout the
previous trial of a reversal-learning task could affect choice probability on the subsequent trial,
but this inhibition had no impact when limited to the rewarded period and did not affect the next
trial's movement latency.[31] Other recent work showed an increased propensity to repeat
behavioral syllables associated with spontaneous DA transients in freely behaving mice,[11] but this
was outside a goal-directed context and did not explore the effects of variation in DA transients.
Endogenous DA transient amplitude can also gradually decrease over the course of learning in a
variety of other settings.[2,58,59] For example, in birds, endogenous DA signals related to song
performance prediction errors have been associated with slow changes in syllable pitch over
juvenile learning.[21,22] But again, these effects were gradual—and not linked to changes in
movement vigor.[22]
373
374 Others have proposed that real-time DA effects on movement are instead the domain of tonic or
375 "quasi-phasic[60]" DA signals rather than reward-related DA transients.[6,22,26,28] Here, we
376 demonstrate that these two modes of DA signaling (phasic and tonic/quasi-phasic) are actually
377 linked, potentially via a novel mechanism.
378
Potential mechanisms. Why could we detect one-shot learning effects of endogenous DA
transients when previous studies did not? One possibility is that our time-to-move metric
provided a more sensitive "analog" behavioral readout. Behavioral updates to binary behavioral
metrics or in discrete reinforcement-learning tasks, such as bandit tasks, may be difficult to
detect on a trial-by-trial basis,[3,50,57] instead requiring multiple repetitions before becoming
manifest.
385
Alternatively, we suspect that one-shot effects on vigor arise via a fundamentally different
biological mechanism than synaptic plasticity.[56] First, a common assumption in reinforcement-
learning (RL) models is that specific stimuli or actions (or other states of an agent) are reinforced
due to temporal association with reward-related DA transients.[56] Importantly, we did not find
that reward reinforced a *specific* time-to-move; rather, DA transients induced a bidirectional
update in timing on the next trial. For example, if an animal was rewarded for moving at 4s, it did
not repeat that 4s movement on the next trial—it tended to move *earlier*. Second, RL models
often assume incremental learning is mediated by gradual DA-dependent structural synaptic
plasticity in cortico-striatal synapses.[56] In contrast, we found that reward-related DA transients
changed timing on a trial-to-trial basis, and (in our previous study) DAN stimulation during the

timing interval affected movement time within only 100s of milliseconds to a few seconds.[7] This
dynamism seems unlikely to arise via incremental DA-dependent synaptic plasticity.
398
What is the cellular/circuit mechanism for reward-related DA transient updates to vigor?
Theoretical work has suggested frameworks for understanding DA's effects on vigor, but the
biological implementations were not clear.[6,61] However, any proposed mechanism should have
the following properties: 1) it should support *bidirectional* effects on timing; 2) it should have a
rapid onset (sub-second to a few seconds); and 3) it should *persist* for at least tens of seconds
(the duration of the ITI in our experiments). DA-dependent synaptic plasticity is likely too slow
and incremental to meet these constraints—and would also seem metabolically inefficient to
support back-and-forth, trial-by-trial updates.
407
By contrast, DA also modulates the excitability of striatal projection neurons (SPNs) via
metabotropic receptors.[62,63] A salient example is a careful striatal slice study, which found that
calibrated, transient optogenetic stimulation of DAN axons caused rapid (sub-second) yet long-
lasting (minutes) increases in the firing rate of direct pathway dSPNs in response to depolarizing
current injections.[63] This excitability effect was blocked by a specific inhibitor of Protein Kinase A
(PKA). PKA is an intriguing candidate effector, because DA transients *in vivo* have been shown to
drive increased PKA activity in dSPNs, whereas DA dips cause increased PKA activity in indirect
pathway iSPNs, with a latency of a few seconds and persistence for tens of seconds,[64] similar to
the dynamics of the behavioral effects we observe. It is possible that the relative balance of
dSPN:iSPN excitability could scale dynamics of the cortico-basal ganglia-thalamic circuit[65,66] in a
manner that allows for bidirectional changes in timing—and modulation of vigor more broadly.
Outcome-related DA transients could drive these PKA changes. (The persistence of outcome-
related DAN signals into the baseline of the next trial are suggestive of changes in tonic DA, which
could also drive and/or contribute to these effects, although we are cautious to make arguments
about tonic DA due to our dF/F-limited measurement window. This idea merits future
investigation with techniques more suitable to measuring tonic dopamine, such as with
quantitative fluorescence lifetime photometry measurements, which are still in development.[67])
This scaled-dynamics model provides a framework for understanding vigor-related symptoms of
DA-related disorders of movement, such as bradykinesia and akinesia in Parkinson's disease.
427
As a final note, we are not arguing against DA's role in synaptic plasticity. On the contrary, rapid
DA-mediated effects on striatal excitability could operate in parallel—or even be synergistic—with
slower changes in synaptic strength. For example, DA-induced changes in excitability could
modulate the probability of spike-timing-dependent synaptic plasticity.[56] Increased vigor could
also enhance reinforcement learning simply by allowing an agent to explore its environment more
briskly, resulting in more exposures to the reinforced action or association. When an animal
encounters a reward-rich environment, the ensuing increase in reward-related DA transients
would increase the propensity to move, which in turn could allow the animal more opportunity
to discover specific causal structure in its environment, potentially through synaptic plasticity
mechanisms. This interpretation could explain DA's association with exploratory behavior,[6] as well
as the longstanding observation that animals increase the frequency and speed of all
movements—not just instrumental, reward-associated movements—in reward-rich

environments.[6] Our findings here reveal a mechanistic link between reward and vigor that

provides a platform for further dissecting the relationship between reinforcement, motivation

and movement.

Supplemental Notes

Supplemental Note 1: Optogenetic inhibition

We also attempted to inhibit DANs (stGtACR2 or JAWS), but despite serious effort (5 mice), we could not identify stimulation parameters that would prevent reward-related DA transients, despite finding parameters that caused DA dips outside the context of the task. Furthermore, more aggressive inhibition at higher light powers (e.g., 5-30 mW) in the context of the task—but not outside the context of the task—caused *seconds-long* rebounds in baseline DA after the offset of inhibition (i.e., this was not the typical transient "rebound spike" sometimes seen after inhibition, but a more sustained increase in DA). Thus—paradoxically—aggressive inhibition tended to *raise* DA overall on stimulated trials. We suspect inhibition was ill-posed in this preparation: we could not detect the lick and initiate stimulation fast enough to prevent reward-related action potentials from triggering DA release, and more powerful inhibition caused paradoxical rebounds. Others have also been unable to effectively block reward-triggered DA with inhibition and have proposed similar rationales (e.g., Ref [31]).

Because we could not prevent first-lick related transients, we did not proceed with the Inhibition cohort. Instead, in lieu of optogenetic inhibition experiments, we caused reductions to outcome-related DA through non-optogenetic manipulations (**Fig 4f-l, Fig 5**), whose effects were consistent with the sloshing model.

Supplemental Note 2: Unrewarded DA transient changes across the session (+Juice Task)

By contrast to rewarded DA transients, unrewarded transients showed only a slight decrease in amplitude in the usual +Juice Task (**Extended Data Fig 7e**), and unrewarded timing updates showed relatively little change (**beginning: +64ms**, 95% CI: [+22ms, +103ms]; **end: +94ms**, 95% CI: [+35ms, +153ms]). Unrewarded transients did however show set-point related changes in the Timeshift Task (**Fig 5k**).

470 **METHODS**
471
Animals
We used adult male and female hemizygous DAT-cre mice[68] (B6.SJL-*Slc6a3*$^{tm1.1(cre)Bkmm}$/J,
RRID:IMSR_JAX:020080; The Jackson Laboratory, ME) or wild-type C57BL/6 mice (>2mo old).
Mice were housed in a temperature and humidity-controlled facility on a reversed night/day cycle
(12h dark/12h light). Behavioral sessions occurred during the dark cycle. All protocols were
approved by the Harvard Institutional Animal Care and Use Committee and MIT Committee on
Animal Care (Harvard IACUC protocol #1343, Animal Welfare Assurance Number #A3431-01; MIT
CAC protocol #2306000543). All work was in accordance with the NIH Guide for the Care and Use
of Laboratory Animals. **Surgeries** were conducted under aseptic conditions with every effort to
minimize suffering, as we described previously.[7] Mice were **water deprived** as we described
previously[7] (≥80% initial body weight) with *ad libitum* feeding. **Histology** to confirm viral
expression and fiber placement was conducted as we reported previously.[7]
484
Stereotaxic coordinates (from bregma and brain surface)
Viral Injection:
SNc[7]: 3.16mm posterior, +/-1.4mm lateral, 4.2mm ventral
VTA[7]: 3.1mm posterior, +/-0.6mm lateral, 4.2mm ventral
DLS[7]: 0mm anterior, +/-2.6mm lateral, 2.5mm ventral.
VLS[43]: 0.34mm anterior, +/- 2.75mm lateral, 3.7mm ventral (rdLight1/dLight1.3), 3.2mm
ventral (D1/A2A-cre; GCaMP)
Fiber Optic Tips (ventral position): 200µm 0.53NA or 0.55NA blunt fiber optic cannulae (Doric
Lenses, Quebec, Canada) or tapered fiber optic cannulae (200 µm, 0.60 NA, 1.5, 2 or
2.5mm tapered shank, OptogeniX, Lecce, Italy) were positioned at SNc, VTA, DLS or VLS:
SNc/VTA[7]: 4.0mm (photometry) or 3.9mm (optogenetics).
DLS[7]: 2.311mm (blunt) or 4.0mm (tapered)
VLS (rdLight1/dLight1.3): 3.4mm (blunt)
VLS[43] (D1/A2A-cre; GCaMP): 3.0mm (blunt or midpoint of 2.5mm tapered)
499
Viruses
Photometry:
tdTomato ("tdt"): AAV1-CAG-FLEX-tdT (5.3*10^{10} to 5.3*10^{12}gc/mL; UNC Vector Core, NC),
500nL alone or mixed with other fluorophores (below).
mCherry (D1/A2A-cre experiments): AAV8-Syn-DIO-mCherry (2*10^{12}gc/mL; Datta Lab),
mixed with GCaMP6f.
gCaMP6f (at SNc or VTA): 300nL AAV1.Syn.Flex.GCaMP6f.WPRE.SV40 (2.5*10^{13}gc/mL;
Penn Vector Core, PA) mixed 3:1 with tdt or mCherry.
DA$_{2m}$ (at DLS): 200-300nL AAV9-hSyn-DA4.4(DA2m) (3*10^{12}gc/mL; Vigene, MD) mixed
2:1 or 3:1 with tdt
dLight1.1 (at DLS): 300nL AAV9.hSyn.dLight1.1.wPRE (9.6*10^{12}gc/mL, Children's Hospital
Boston, MA) mixed 3:1 with AAV1.CB7.CI.TurboRFP.WPRE.rBG (10^{12}gc/mL, Penn
Vector Core).

rdLight1 (at VLS): 500nL AAV2/9-Syn-RdLight1.0 ($2*10^{12}$gc/mL; #construct-1646-aav2-9,
Canadian Neurophotonics Platform (CNP), Quebec, Canada).
dLight1.3 (at VLS): 500nL AAV2/DJ-CAG-dLight1.3b ($2*10^{12}$gc/mL, CNP)
Optogenetic stimulation/inhibition (bilateral at SNc):
ChR2: 1000nL AAV5-EF1a-DIO-hChR2(H134R)-EYFP-WPRE-pA ($3.2*10^{13}$gc/mL, UNC
Vector Core, NC)
stGtACR2: 300nL AAV2/8-hSyn1-SIO-stGtACR2-FusionRed ($4.7*10^{11}$gc/mL,
Addgene/Janelia Viral Core, VA)
521
522
523 **Behavior**
524 Neurophysiology and behavioral data was acquired as we described previously.[7]
525
Self-timed movement task (original version): Mice were head-fixed with a juice tube positioned
in front of the tongue. The spout was placed ~1-1.5mm ventral and ~1-1.5mm anterior to the
mouth. During periods when rewards were not available, a houselamp was illuminated. At trial
start, the houselamp turned off, and a random delay ensued (0.4-1.5s) before a cue (simultaneous
LED flash and 3300Hz tone, 100ms) indicated start of the timing interval. The timing interval was
divided into two windows, early (0-3.333s) and reward (3.333-7s), followed by the intertrial
interval (ITI, 7-17s). The window in which the mouse first licked determined the trial outcome
(early, reward, or no-lick). An early first-lick caused an error tone (440Hz, 200ms) and houselamp
illumination, and the mouse had to wait until the full trial interval had elapsed before beginning
the ITI. Thus, there was no advantage of licking early. A first-lick during the reward window caused
a reward tone (5050Hz, 200ms) and juice delivery (~4µL apple juice or Gatorade; **+Juice Task**),
and the houselamp remained off until the end of the trial. If the timing interval elapsed with no
lick, a time-out error tone played (131Hz, 2s), the houselamp turned on, and ITI commenced.
During the ITI and pre-cue delay ("lamp-off interval"), there was no penalty or feedback for licking.
Mice learned the task in 3 stages, as described previously.[7] All mice learned the task and worked
for ~400-1,800 trials/session.
542
543 **Pre-training/post-training.** To encourage participation, each session (except for SNc/VTA/DLS
544 GCaMP mice) commenced with 10 trials of "pre-training" during which juice was delivered
545 triggered on the first-lick within the reward window, but during which early licks were not
546 penalized. Pretraining was always conducted with juice rewards.
547
One-shot Stimulation (+Juice Task, Fig 3). Closed-loop stimulation was delivered at SNc and
triggered on first-lick on 30% of randomly interleaved trials in the context of the +Juice Task. Mice
were trained without stimulation for 13-16 sessions before starting stimulation sessions. Each
mouse completed 10 pairs (20 sessions) of interleaved stimulation and sham sessions, where
sham sessions were identical to stim sessions (patch cables attached, etc.), except the stimulation
LED was turned off and no light was delivered. Photometry was usually not collected during sham
sessions to reduce bleaching from recording on consecutive days. See **Optogenetics** section for
stimulation parameters, calibration and analysis.
556

No Juice Task (Fig 4f-l). In the usual (+Juice) task, feedback was either an error tone (440Hz,
200ms) or correct tone (5050 Hz, 200 ms) plus juice reward. The No Juice Task was identical to
the Juice+ Task and used the same feedback tones, but no juice was delivered (dispensing
solenoid valve disconnected). In a subset of animals trained on an open rig with a loud juice
solenoid, a decoy solenoid was triggered, but no juice was dispensed. After >=35 days of +Juice
Task, expert mice successfully performed the No Juice Task, some for >1000 trials/session (one
mouse tested on days 23 and 25 showed less engagement). Anecdotally, we observed that
consecutive No Juice sessions could reduce participation in subsequent sessions in some animals,
and so No Juice sessions were typically interleaved with +Juice sessions to promote participation.
Together, these factors limited the number of No Juice sessions that we could collect.
567
568 **Alternating Juice/No Juice Task (Fig 5a-c).** Mice alternated between unsignaled blocks of usual
569 +Juice or No Juice Task.
570
571 **Alternating Stimulation Blocks (Fig 5d-f).** Mice were trained in a No Juice Task variant in which
572 closed-loop SNc DAN stimulation was delivered in blocks of trials. During stimulation blocks,
573 stimulation was delivered on either 30%, 60% or 100% of first-licks on all trials (separate blocks,
574 **Extended Data Fig 9a**) *or* only on 100% of rewarded first-licks (**Fig 5d**). See **Optogenetics** section
575 for stimulation parameters, calibration and analysis.
576
Rapid Satiation Task (Fig 5g-i). At >20d of expert task training, Rapid Satiation sessions were
collected, where the ongoing task was paused once behavior met two criteria: ≥100 trials
completed and ≥10 juice rewards received. Video and photometry recording were continued
during the pause (mice remained headfixed on the behavior rig). The spout was retracted, and
the mice were hand-fed undiluted Cool Blue Gatorade from a 1mL syringe until they stopped
initiating licks to the offered syringe. The spout was then moved back into position and the task
resumed. In Control sessions, the same procedure was followed, except an empty syringe was
offered.
585
Both Rapid Satiation and Control sessions were collected on a background of the +Juice task with
undiluted Cool Blue Gatorade rewards. We used a newer version of the +Juice task, in which
there was also an unsignaled late window after the reward window (7-14s). In this version of the
task, the trial duration was always 14s, with the late window occurring between the unsignaled
end of the reward window and the start of a 3s-long ITI (total trial+ITI duration 17s, as in the
original +Juice Task). First-licks during the late window triggered a distinct, late error tone (132Hz,
500ms) and illumination of the houselamp (similar to early trials). Mice still had to wait until 14s
to start the ITI.
594
595 In analyses, to account for missing time while the task was paused, the total duration of the pause
596 was divided by 17 (the trial length was 17s) and rounded to recover the number of missing trials.
597 Rewarded DA transient amplitudes ("rewDA") were then interpolated linearly and smoothed with
598 a 20-trial gaussian kernel. Edges (trials before the first reward and after the last reward in the
599 session) were omitted.
600

Rapid Satiation sessions were collected in pairs with interleaved control sessions. However, some
sessions had to be excluded either because the mouse received no rewards before the pause (1
session), the mouse stopped participating after the pause (1 session), or there were issues with
lick detection due to a broken spout (3 sessions). Although including these sessions did not
change the result of analyses, they were omitted from the final rewDA analysis. Statistical
comparisons of rewDA before and after the pause were performed on the average rewDA signal
across 50 trials before and after the maximum duration pause (36 trials) and compared by
Wilcoxon rank sum test.
609
Timeshift Task (Fig 5j-l). In the usual +Juice and No Juice tasks, there was only one rewarded time
window (3.333-7s). In the Timeshift Task, this window changed dynamically in blocks of trials. The
Timeshift Task was introduced after >28 days of training on the usual +Juice Task. Each block used
one of five reward windows: a) 1-2s, b) 2-3.333s, c) 3.333-7s, d) 7-12s, e) 9-14s. There was no set
schedule for changing between blocks. Block changes were signaled by a ~90s timeout (task
paused). The new reward window was not indicated and had to be discovered through trial-and-
error. In a subset of sessions, a "late" window was added (as in the Rapid Satiation Task) such that
mice were not immediately alerted when the reward window had closed. Behavior was grossly
similar on both versions of the Timeshift Task, and thus, for all analyses, data from the standard
Timeshift Task and late window Timeshift Task was pooled. Mice were also trained on a No Juice
version in some Timeshift sessions (7 mice, 9 sessions), and these data were pooled with +Juice
Timeshift sessions.
622
623
Photometry
Signal collection. Fiber optics were illuminated with 475nm blue LED light (Plexon, TX)
(SNc/VTA: 50μW, DLS: 35μW, VLS: 30μW) measured at patch cable tip with a light-power meter
(Thorlabs, NJ). Green fluorescence was collected via a custom dichroic mirror (Doric Lenses,
Quebec, Canada) and detected with a Newport 1401 Photodiode (Newport Corporation, CA) or
integrated Doric cube. For red florescence, 550nm lime LEDs were used (Plexon, TX). Fluorescence
was allowed to recover ≥1d between recording sessions. To avoid optical crosstalk, the red control
channel was recorded at one of the other implanted sites (SNc, VTA, DLS or VLS, 550nm lime LED,
Plexon, TX) while GCaMP6f, dLight1.1/1.3/rdLight1 or DA$_{2m}$ were recorded simultaneously only
at the other implanted sites.
634
635 **dF/F.** Raw fluorescence for each session was pre-processed by removing rare singularities (single
636 points >15 STD from the mean) by interpolation to obtain F(t). To correct photometry signals for
637 bleaching, dF/F was calculated as:

$$\frac{dF}{F}(t) = \frac{F(t) - F_0(t)}{F_0(t)}$$

638
639 where $F_0(t)$ is the mean fluorescence during the 5s preceding the current trial, as well as 5 trials
640 before the current trial and 5 trials after the current trial (as we characterized and validated
641 previously[7]).
642

Correlation of baseline photometry signals with first-lick time. Baseline signals were de-noised
by pooling based on first-lick time into 0.25s pools from 0.75 to 4s after the cue, as in **Fig 1d**. Pools
with first-lick times after 4s were not included due to insufficient numbers of trials in each pool
to de-noise the measurement (inclusion criteria >=175 trials). Pearson's r was calculated from
each pool's average dF/F signal calculated within windows of time relative to lamp-off (left of
Fig 1d, r: –0.63; –10.5 to 10s before lamp-off) or the cue (right of **Fig 1d**, r: –0.79; –0.4 to 0s before
the cue (excludes lamp-off event)). These signals were related to the earliest first-lick time in each
pool.
651
652 **Sloshing Model.** To test the independent contribution of each task-related input to the
653 photometry signal and select the best model, we employed nested fitting, in which each dataset
654 was fit multiple times (in "nests"), with models becoming progressively more complex in
655 subsequent nests. The nests fit to the photometry data employed the inputs $X^{(j)}$ at each j^{th} nest:
656
657   Null Model: $X^{(0)} = x_0$
658   Nest 1: $X^{(1)} = X^{(0)} +$ EMG and/or accelerometer
659   Nest 2: $X^{(2)} = X^{(1)} +$ red control fluorophore ("tdt")
660   Nest 3: $X^{(3)} = X^{(2)} +$ outcome-related photometry signal
661
662 The outcome-related photometry signal was taken as the mean dF/F signal within the first 500ms
663 after the first-lick (peak or median dF/F produced similar results). All d_j design matrix predictors
664 were 0,1 normalized.
665
666 The GLM for each nest took the form:
667
668 $$Y = \Theta X^{(j)}$$
669
Where Y is the *1xN* vector of the change in the movement time between the current trial, n, and
the next trial, n+1 ($\Delta T_{n-(n+1)}$), where *N* is the total number of trials in the session; $X^{(j)}$ is the *dxN*
design matrix for nest *j*, where the rows correspond to the d_j predictors for nest *j* and the columns
correspond to each of the *N* trials of Y; and Θ is the *dx1* vector of fit weights. Because of day-to-
day/mouse-to-mouse variation (ascribable to many possible sources, *e.g.*, different neural
subpopulations, expression levels, behavioral states, *etc.*), each session was fit separately.
To guard against a trivial relationship between the trial outcome and DA signal, separate models
were fit for early or rewarded trials.
678
679 Among the nests, the best model for each session was selected by BIC (AIC and AICc were typically
680 in agreement with BIC), which was typically the model including the photometry signal. All
681 reported data is for the full model.
682
Residual sloshing models (Extended Data Fig 4a). To determine whether outcome-related DA
was predictive of ΔT in excess of chance (regression to the median effects), the next trial's
movement time was shuffled, and the sloshing model was refit on the shuffled data (10,000
shuffled datasets). The coefficients from the 50th percentile shuffled model (50th percentile

shuffled DA coefficient) were used to predict the true data (y-fit shuffle) and calculate shuffled-
R^2. Y-fit shuffle was subtracted from the true dataset to obtain a residual dataset, and the residual
model was derived by refitting the sloshing model on the residual dataset.
690
Shuffled models/regression to the median (Fig 2e, Extended Data Fig 4b-c). To guard against
spurious relationships between DA and ΔT arising from behavioral non-stationarity, each session
was divided into four quartiles of trials (same number of trials with participation in each quartile)
to isolate more stationary periods of behavior. The sloshing model was fit on each quartile
(including movement/optical artifact control signals and mean DA photometry signal during the
first 500ms after the first-lick). The trial order (for each quartile) was then shuffled 10,000x and
the sloshing model fit on each shuffled dataset to produce the distribution of DA coefficients for
each quartile assuming no trial-order dependence. To determine if the true fit coefficient was in
excess of the shuffle, a Wilcoxon rank sum test was applied to compare the set of true coefficients
across quartiles (and sessions) to the set of mean shuffled DA coefficients.
701
702
703   **Optogenetics**
704   **Opsin expression.** 3 cohorts of mice were tested: Activation (ChR2, 4 mice), Inhibition (stGtACR2,
705   4 mice or JAWS, 1 mouse), and No Opsin (no SNc injection, 4 mice). All mice co-expressed rdLight1
706   at VLS for co-monitoring of stimulation effects on DA release. All stimulations were delivered via
707   fiber optic cannulae implanted over SNc.
708
Co-monitored photometry. We selected ChR2 (blue-activated opsin)/rdLight1 (green-excited
fluorophore) as our DA co-monitoring strategy because we had previously observed issues with
off-target excitation of red light activated opsins (e.g., ChrimsonR) during prolonged co-recording
of dLight1.1 (a blue light-excited fluorophore) in DLS (red opsins are also activated by blue light,
albeit with lower efficiency, which becomes significant during hours-long timecourses of
photometry illumination in our behavioral sessions).[7] This off-target excitation would have
created a confound, because we had previously shown that DAN activation limited to the self-
timing interval increases the probability of early first-licks.[7] ChR2, by contrast, is not appreciably
activated by green light (used for rdLight1 excitation), and was thus a cleaner preparation. One
challenge of blue stimulation/green-excited photometry is that stimulation light causes a DC-
offset artifact in the red photometry channel while the stimulation LED is turned on. To avoid
crosstalk between the stimulation LED and the photometry recording, two complimentary
correction methods were used, with similar results: 1) the brief stimulation up-times were
omitted from the photometry signal (missing points filled by interpolation), or 2) the signal was
decontaminated by measuring the instantaneous increase in rdLight1 signal at the beginning of
each pulse of blue light, which was subtracted from the voltage trace during each pulse. Both
corrections were taken before dF/F correction and yielded similar photometry signals, however
the second method had the advantage of compatibility with continuous illuminations used during
stGtACR2 inhibition experiments (see **Stimulation Parameter Calibration**, below).
728
Stimulation parameter calibration. Activation parameters were calibrated outside the context of
the task. Mice sat on the typical behavioral platform and received 10 unexpected juice rewards

(ca. 4μL each), spaced at least 15s apart. A grid screen of stimulation parameters and light powers
identified conditions that best recapitulated both the amplitude and width of the typical reward-
related transient. Mice received 10 stimulations with each stimulation protocol, and DA release
in VLS was monitored throughout the calibration procedure with rdLight1. Light intensity and
pulse width were varied until the resulting rdLight transient matched the amplitude and width
typically seen with unexpected juice rewards.

The final parameters used in all experiments were as follows: Optogenetic stimulation was
delivered bilaterally via blunt fiber optics at SNc (5mW 465nm LED, 10Hz, 250ms, 35% duty cycle).
One ChR2 animal had partial response to 5mW blue light, and thus received 10mW blue light,
with a larger but still partial effect (Mouse X4-3-1M). We could not identify a set of inhibition
parameters that prevented first-lick related transients in the task and did not proceed with the
Inhibition cohort (**Supplemental Note 1**).

One-shot stimulation analyses. Initial power simulations indicated that ~5,000 trials (10x 500
trial sessions) would be necessary to achieve 80% power to detect a 20% change in first-lick time.
Thus, at least 10 sessions of stimulation (~500 trials each) were collected from each animal.

Three complementary analyses examined single-trial stimulation effects (**Extended Data Fig 6**):

1. **"deldel,"** difference in the timing update (del1) between stimulated vs unstimulated
trials (del2):
$$\Delta(\Delta T_{(n+1)-n})_{\text{stim-unstim}} = \Delta T_{(n+1)-n \mid \text{trial n stimulated}} - \Delta T_{(n+1)-n \mid \text{trial n unstimulated}}$$
2. **"dAUC,"** difference in the area under the cdf of the next trial (n+1) timing distribution for
stimulated vs unstimulated trials (as described previously[7]).
3. **"cdf-rank,"** percentage of points in the n+1th trial cdf after stimulated trials that were
less than matching points in the n+1th trial cdf after unstimulated trials. cdfs were
interpolated such that each point was matched between the stimulated and unstimulated
curves.

Analyses were performed at 3 levels, with consistent results across analyses and levels:

<u>By Session</u>
Within single session analyses: deldel, dAUC and cdf-rank analyses compared the distributions
or first-lick times on the next trial after stimulated vs unstimulated trials. As expected from
preliminary power calculations, these statistics were noisy (**Extended Data Fig 6 top row,**
single dots).

Between groups single session comparisons: The test statistic for each single session was
bootstrapped 10,000x, and the mean of the single session test statistics for each group were
subtracted to derive a between groups comparison test statistic (activation – no opsin),
which was then compared by Wilcoxon rank sum test (**Extended Data Fig 6 top row,**
between group statistic).

 By Mouse

 Within "Mega" session comparisons: Because initial power simulations suggested 5000 trials
 were needed for 80% power to detect a 20% change in first-lick time, data were pooled across
 stimulated sessions from individual animals to generate "mega" sessions, which had ~5000
 trials each. The test statistic for each mega session was reported as dots in **Extended Data**
 Fig 6, **second row**.

 Between groups mega session comparisons: The test statistics for each group of animals'
 mega sessions were compared by Wilcoxon rank sum test. 2-sided unpaired t-test found
 similar results.

 All Mice

 Within "super mega session" comparisons: All data from all sessions in each cohort were
 pooled to generate a "super mega session" to compare all stimulated and unstimulated trials
 by 10,000x bootstrap procedure (same as used on within-single session analyses). Test
 statistic and p-value reported in the Main Text and **Extended Data Fig 6, third row**.

 Whole session behavioral analyses
 "Set point" calculations: Running median first-lick time. For visualization, the timing set-point
 was defined as the 50-trial running median centered at the current trial, excluding missing trials
 (e.g., no-lick) and reactions (first-lick <0.5s). This captured the peak of the timing distribution
 (whereas including reactions tended to underestimate the peak). For quantitative comparison to
 rewarded DA transients (which were less abundant) and in tasks with rapidly changing set-point
 dynamics, the number of trials included in the running median was adaptively reduced to avoid
 over-smoothing (10-50 trials used, depending on analysis: e.g., Timeshift No Juice task used 15
 trials). Data provenance text files are included with each figure and specify the smoothing used
 in each analysis.

 +Juice/No Juice Task Comparisons. For animals with No Juice Task sessions (n=5 VLS rdLight mice,
 11 sessions), the nearest +Juice session with photometry was matched for comparisons of
 behavior (timing set-point) and outcome-related DA signals. For set-point analysis, 15 sessions
 were available for comparison (e.g., **Figs 1b, 4b-d**). Some flanking sessions were collected without
 photometry (we allow signals to recover from bleaching for at least 1d between recording
 sessions)—in these situations, the most recent +Juice photometry session was used for DA
 analyses. Together, this produced 18 paired photometry sessions for No Juice vs Juice+
 comparisons (**Fig 4i-k**; n=5 mice, 9 sessions each task). Except for one session (Randy_VLS_47),
 paired +Juice photometry sessions happened before the paired No Juice session. Because the
 +Juice sessions mostly had 500 trials total, for consistency, we used only the first 500 trials from
 each No Juice session for comparisons, although some No Juice sessions were recorded for >1000
 trials. The Q1 and Q4 lick-triggered averages shown in **Fig 4** were thus taken on trials 1-125 and
 375-500, respectively, irrespective of session duration.

Set-point cross-over analysis (Fig 4e). Sudden shifts of the set-point into the reward window were
eventually observed in all animals when allowed to work ad libitum, typically between trials 400-
1100. However, because many sessions were only recorded for 500 trials, only a subset of sessions
were recorded long enough to observe cross-overs (we validated this in a large behavioral dataset
from animals allowed to work until disengagement, typically between trial 1100-1800, 3 mice,
120 sessions). Sessions with set-point cross-over were identified in the SNc GCaMP or VLS rdlight1
datasets as those sessions with set-point non-stationarity of >=+1.5s (16 sessions total). If the set-
point started within the reward window at the beginning of the session, the set-point had to first
exit and then re-enter the reward window (1/16 sessions). Evolution in first-lick time and
rewarded DA transient amplitude ("rewDA") was taken as a running median over 75 trials (37-38
either side of each trial), where missing data was interpolated linearly (edges excluded as above
to avoid artifacts). To de-noise the derivative of the rewDA signal, rewDA was first downsampled
(every 15 trials taken). The downsampled trial with maximum derivative of the DA signal
($-1*$rewDA') was compared to the "set-point cross-over trial," defined as the trial on which the
median of the timing distribution first crossed into the reward window (>3.333s). The cross-over
trial number and $-$rewDA' were 0,1-normalized.
834
835
836　**Code Availability.** All custom behavioral software and analysis tools are available with sample
837　datasets at https://github.com/harvardschoolofmouse.
838
839　**Data Availability.** All datasets supporting the findings of this study will be made publicly available
840　at the time of publication via DANDI, with source data files provided for all figures.

Acknowledgements

We thank L. Hou, E. Sayed, J.G. Mikhael, and S. Hrvatin for discussions; M. Yang, S.L. Jacobs-Skolik, N. Hunter, V. Berezovskii, J. LeBlanc, T. LaFratta, O. Mazor, and P. Gorelik for technical assistance; and B. Sabatini, C. Harvey, S.R. Datta, M. Andermann, A. Mohebi and W. Regehr for reagents. The work was supported by NIH grants UF-NS108177 and U19 NS113201; NIH core grant EY-12196; the Valhalla Foundation and the Whitehead Fellows Program. The funders had no role in study design, data collection and interpretation, or the decision to submit the work for publication.

Author contributions

A.E.H. and J.A.A. conceived the study, designed the experiments, acquired funding, supervised the project, and wrote and edited the manuscript. **A.E.H.** Performed experiments, wrote the code for analysis, analyzed the data, and wrote the first draft. **I.C.W., Q.D., Z.O. and P.A.** assisted with experimental design, performed experiments, processed data, and assisted with analysis. All authors reviewed the final manuscript.

Author ORCIDs

Allison E Hamilos   https://orcid.org/0000-0001-9486-0017

Isabella C Wijsman   https://orcid.org/0009-0000-5107-7823

Qinxin Ding   https://orcid.org/0009-0009-6571-2678

Pichamon Assawaphadungsit https://orcid.org/0009-0003-0343-5368

Zeynep Ozcan   https://orcid.org/0000-0002-8311-1958

John A Assad   https://orcid.org/0000-0002-7689-5336

Competing Interests Declaration

J.A.A. is a co-founder of OptogeniX, which produces the tapered optical fibers used in some experiments.

Additional Information

Corresponding Author: Allison E. Hamilos and John A. Assad

Supplementary information:

    Extended Data Figures

    Supplemental Notes 1-2

    Supplemental Methods

a

House lamp ●
Lamp-off

LED + Cue tone (100 ms) 🔴 + 🔊
Cue

House lamp + Miss tone ⊙ + 🔊
Lamp-on

Random delay

Early window

First-lick → Error tone 🔊

0s

Reward window

First-lick → Reward tone + juice 🔊 + 🟡

3.3s

Intertrial (ITI)

First-lick → Nothing happens

7s   17s

b

0.3

0.05

Reward Window

5 mice
15 sessions

0   2   4   6   8   10s

c SNc GCaMP6f

First lick   early
              rewarded

|10% dF/F

SNc
4 mice

-2 0   6s

d Striatal Dopamine

First lick   First lick

DA2m (DLS)   rdlight1 (VLS)
4 mice   8 mice

|2% dF/F |1% dF/F

-2 0   6s   -2 0   6s

e

Trial n   Trial n+1

single trial
SNc GCaMP6f
1 session

first-lick trial n (early)   first-lick trial n (reward)   Cue trial n+1   first-lick trial n+1 (early)   first-lick trial n+1 (reward)

10% dF/F

-20   -15   -10   -5   0   5   10s

f

Trial n   Trial n+1

Lamp on (ITI start)   Lamp off (trial start)   Cue   self-timing interval

r= -0.63

SNc DAN GCaMP6f
n=4 mice, 20 sessions

r= -0.79

First-lick time (Trial n+1)
0   4s

0.5% dF/F

-10 -9 -8 -7 -6 -5 -4 -3 -2 -1 0   0   1   2   3   4s

g

Reward Window

n=17 mice
155 sessions

● p<0.05

0.6

-0.4

median ΔT in excess of chance (s)

Slow down
ΔT+
ΔT-
Speed up

0   first-lick time, trial n   9s

Fig 1. Outcome-related DA is related to timing on the next trial. a, Task. **b,** Whole-session timing distribution, same 5 mice as in **Fig 4**. **c-d,** Outcome-related signals; shading: std. **e-f,** Timing-related DAN signals: **e,** Single-trial. **f,** DAN signals pooled by movement time and plotted up to 150ms before spout contact. X-axis breaks indicate change of alignment. **g,** Change in timing set-point on next trial (ΔT); shading: 95% CI (10,000x bootstrap).

922
923
924 **Fig 2. Outcome-related DAN signals predict changes in timing on the next trial. a,** Sloshing GLM. **b,** Trial-by-trial
925 variability in outcome-related DAN signals. Left: average. Right: single-trial mean DAN signals within 0.5s post-first-
926 lick, jittered for clarity. **c-d,** Sloshing Model fit. Coefficient error bars: 95% CI. **e,** Model DA coefficients, true data vs
927 10,000x shuffle fit, all sessions. Standard box plot (red line: median). *p: Wilcoxon rank sum test. Models for all
928 signals: **Extended Data Figs 3-4**.

929
930
Fig 3. Outcome-related DA causes one-shot timing updates. a, Optogenetics strategy. **b,** Stimulation parameter
calibration, outside context of task. Colors: single trials, black: ave. Left: 10x stimulations; Right: 10x unexpected juice
rewards (~4μL each). **c-d,** Task DA. **e-f,** Single trial ("one-shot") stimulation effects. **e:** Standard box plot, red bar:
median. n=4 Activation mice; n=4 No Opsin mice, 10 sessions/mouse. Wilcoxon rank sum test, *p<0.05. **f:** First-lick
distribution on next trial (n+1). Arrow: direction of effect. Shading: 95% CI; Bar: *p<0.025 (1000x bootstrap). **g,** Whole
session effects (stimulated vs interleaved sham sessions). Shading: 95% CI; Bar: *p<0.025 (1000x bootstrap).

937
938
Figure 4. Timing changes across behavioral sessions. a-b, Whole-session behavior, +Juice Task. "Set-point": running
median first-lick time. **b,** Sessions shown for comparison to matched No Juice Task sessions (**Fig 4f-l**). Q1-Q4: quartiles
of trials in session. **c,** Outcome-related DA. **d,** Normalized rewarded DA transient amplitude ("rewDA") vs timing set-

point. **e,** max(-rewDA'): trial at maximum negative derivative of rewDA. Cross-over: trial at which set-point crossed

into reward window. Right: Circles—single sessions. Black circles—No Juice sessions. **f-g:** No Juice Task whole session

behavior, as in panels a-b. **h,** +Juice vs No Juice behavior. Boxplot, yellow line: median; Wilcoxon rank sum test. **i,** No

Juice Task outcome-related DA. **j-k:** rewDA, +Juice vs. No Juice Task. **j** shows Q1 signals. **l,** No Juice rewDA vs timing

set-point.

947
Figure 5. DA calibrates propensity to move. a-c, Alternating Juice/No Juice Task. Standard box plot, Wilxocon rank
sum, p*=0.032. **b-c,** Correlation of reward-related DA ("rewDA") and set-point across block changes. **d-f,** Alternating
Stimulation Blocks, Rewarded Trials only. *p<0.05, **p<0.01, ***p<0.001. **g-i,** Rapid Satiation task: causes abrupt
decline in rewDA amplitude accompanied by rapid late-shifting. **j-l,** Timeshift Task. **k,** Set-point remains strongly
correlated with rewDA; unrewarded DA peak amplitude (within 500ms after first-lick) shows stronger correlation
with set-point than in the standard +Juice/No Juice tasks.

Extended Data Fig 1. Timing behavior over the course of learning. Animals trained with Juice rewards (*but not No-*
*Juice rewards, see **Fig 4***) show early-biased timing behavior—we will address this in detail later in the paper. All mice
start by learning the +Juice Task, shown here. Timing peaks emerge over the course of learning, stabilize within the
first 20 sessions of training, and early bias in the timing peak persists throughout +Juice Task training. We will come
back to this early bias later in the paper. **a)** Histograms show the distribution of first-licks after the cue on each day
of +Juice Task training for one representative mouse. This mouse was trained for 33 consecutive sessions with 500
trials/session, with no training days on weekends. The mouse was transitioned from the Beginner task (blue traces,
Pavlovian reward at 5s, 0.5s reaction window) to the Intermediate task (purple traces; no Pavlovian rewards, 0.5s
reaction window) on day 4. Transition to the Expert task (red, no Pavlovian rewards and no reaction window)
occurred on day 7. Y-range of histograms was maximized to clearly show the timing distribution peak; reaction peak
(<0.5s) is truncated in some plots and indicated by open trace. +Juice Task distributions become stable by day 20 of
training, and early timing bias persists throughout training when juice rewards are used. **b)** Mean first-lick time for
each day of training. Early timing bias (mean first-lick time between 2-3s) is firmly established by day 10 and persists
throughout +Juice training. **c)** Standard deviation of first-lick time for each day of training declines as timing
distribution becomes more stereotyped over the course of training.

971
972
973 **Extended Data Fig 2.** Outcome related dopamine and striatal projection neuron (SPN) signals. Shading: standard
974 deviation.

1017
1018 **Extended Data Fig 3.** Sloshing model coefficients, all signals, all sessions. Small colored dots indicate coefficients from
1019 individual mice.

Extended Data Fig 4. Outcome-related DA explains ΔT in excess of chance. **a-b,** Shuffle models. The next trial's movement time was shuffled, and the sloshing model (**Fig 2a**) was refit on the shuffled data (10,000 shuffled datasets total). The coefficients from the 50th percentile shuffled model (50th percentile shuffled DA coefficient) were used to predict the true data (y-fit shuffle) and calculate R^2 (**a, middle panel**). Y-fit shuffle was subtracted from the true dataset to obtain a residual dataset, and the residual model was derived by refitting the sloshing model on the residual dataset (**a, right panel**). Outcome-related DA remained a significant predictor to the residual model, both in single sessions (**a, right-bottom panel**) and across sessions for SNc GCaMP, VLS rdlight1, and DLS DA2m. DLS dlight1.1 DA signals did not remain significant after the shuffle procedure, likely due to relatively poor S:N quality in the dlight1.1 dataset. **c.** Comparison of true vs shuffled model DA coefficients fit on quartiles of trials in the session. To guard against spuriously significant DA coefficients arising from non-stationarity in timing behavior across the session, true and shuffle models were fit on quartiles of trials in each session (e.g., first quarter of trials in the session, second quarter, etc.). Consistent with the residual models in **b**, the population of true model DA coefficients was significantly larger in magnitude (more negative) than the population of mean shuffled model DA coefficients (except for DLS dlight1.1; Wilcoxon rank sum test, *p<0.05). Outcome-related DA is thus predictive of ΔT in excess of regression to the median effects (see **Supplemental Methods**) and unlikely to be explained by behavioral non-stationarity.

1066
Extended Data Fig 5. One-shot optogenetic manipulations—all mice. LED triggered on 30% of randomly interleaved first-licks. **a,d.** VLS rdlight1, all stimulation sessions, 10 sessions/mouse. Each row is a different mouse. **b,e.** Comparison of trial n+1 first-lick distributions (all trials) from stimulation sessions (blue trace) vs interleaved sham sessions (black trace). Blue shading: early-shift effect of LED; red shading: late-shift effect. Arrow: indicates overall direction of effect. Top bars (blue/red): denotes significant difference between eCDF curves (*p < 0.025, 95% CI, 1000x bootstrap test). K-S D: signed Kolmolgorov-Smirnov distance metric (negative indicates early shift effect). dAUC: difference in area under eCDF curves, p: 10,000x permutation test. **c,** Peak rewarded DA transient amplitude ("rewDA"), 1 session. Purple arrow and dashed line indicate the first rewarded (unstimulated) trial, whose amplitude is ~0.065 dF/F—corresponding to the purple bar in the bar plot. Bar plot also shows mean peak rewDA amplitude for each stimulation condition and trial outcome (reward/no reward). Stimulated trials remained below the naturally occurring peak rewDA amplitude throughout the session and on average.

1078
1079
1080 **Extended Data Fig 6.** One-shot optogenetic manipulations – complementary analysis methods, all sessions and mice.
1081 Comparisons: stimulated vs unstimulated trials from the same (stimulated) session. Dot colors indicate whether the
1082 test statistic was significant at the individual session level (p<0.05). Note these metrics can also be used on whole
1083 session comparisons (stim sessions vs interleaved sham sessions), as in **Fig 3g**.

Extended Data Fig 7. Whole session timing behavior. **a,** Timing distributions from monkeys, rats, and mice show early-bias relative to the reward window. Figure panels/data adapted and used with permission. **b,** Skew in outcome-related DA responses appears similar to skew in the timing distribution with respect to the reward window. **c&j,** Heatmap view of first-lick distribution. Same mice shown in panels c and j, interleaved sessions. "Q:" quartile of trials in session with a first-lick. **d,** The effect of trial outcome on timing updates becomes less skewed toward the end of the session (rewarded trials followed by smaller updates). **e&k,** Single trial peak amplitude DA signal within 500ms after the first-lick. **f&l,** smoothed single trial data plotted for each session. Each line is one session. "rewDA:" reward-related DA peak amplitude; "no-rewDA:" unrewarded peak. **g,** Peak outcome-related tdTomato, same sessions as panel f. **h,** Correlation of all single-session outcome-related DA (top) and tdt (bottom) transients with timing set-point. Thin traces: single sessions; thick: average. Left: SNc GCaMP6f; Right: VLS rdLight1 signals. Control rew-tdt signals are

not consistently correlated with the timing set-point, arguing against a trivial correlation with ongoing body
movements/optical artifacts. **i,** max(-rew-tdt'): trial at maximum negative derivative of rewarded tdt signal amplitude.
Cross-over: trial at which set-point crossed into reward window. Circles: single sessions. Black circles indicate No Juice
sessions. The strong correlation of set-point cross-over with rewarded DA amplitude (**Fig 4e**) is not present in control
tdt signals, arguing against a trivial correlation with ongoing body movements/optical artifacts. **m-n,** Comparison of
Juice+ and No Juice Task rewDA shows that rewDA declines significantly earlier in the session in the No Juice Task
compared to the Juice+ Task, but there was no difference in the 50% max point for tdt data collected on the same
sessions. Standard box plot with Wilcoxon rank sum test. The scatterplot points are the trial at which rewDA (panel
m) or rew-tdt (panel n) reached 50% max peak amplitude for paired Juice+/No Juice session. Green/red dots: the
data for either rdLight1 (green) or tdt (red); light grey dots: control data obtained from randomly pairing Juice+
sessions from the same animal, which shows 50%-max tends to occur around the same trial on sessions of the same
task-variant within the same mouse. **o,** Additional single-session examples of cross-over analysis in No Juice task
(analogous to **Fig 4e**).

1109
1110
Extended Data Fig 8. Reward-related photometry across the session. Q1: Beginning of session (first quartile of
consecutive trials). Q4: End of session (fourth quartile). All DA/DAN recordings show decline in rewarded DA transient
amplitude across the session, with comparably little change in unrewarded (early) transient amplitude. Optical
artifacts (tdTomato) show the opposite direction of effect on unrewarded trials, whereas rewarded trial amplitude is
similar across the session. Recordings from direct pathway striatal projection neurons (dSPNs) show similar patterns
to DA/DAN recordings. Indirect pathway SPNs (iSPNs) show the *opposite* effect on average: both early- and reward-
related iSPN transient amplitude *increase* toward the end of the session. This difference is consistent with DA's effect
on iSPN excitability and provides supporting evidence against bleaching-related artifacts explaining the reward-
related DA downtrend (**Supplemental Methods**).

1161
1162
Extended Data Fig 9. Set-point manipulations: more detail. **a,** Alternating Stimulation Blocks task, 100% of trials. *Left*:
Example single sessions with same block schedule (1 ChR2 mouse, 1 No Opsin mouse, 1 session each). Exposure to

blue light initially caused a small early-shift in the No Opsin mouse, but this was much smaller than in the ChR2
mouse and did not recur during the second stimulation block. The No Opsin set-point trace is greyed after the arrow
because the mouse did not participate on sufficient trials during blocks 4-5 to meaningfully calculate the running
median first-lick time. *Right*: DA transient peak amplitude ("rewDA") in the example ChR2 session. Interestingly, a
slow late-shift dynamic was present even with ChR2 stimulation, and stimulated DA transients declined progressively
over the course of the session. **b,d,f,** Combined Alternating Stimulation Blocks paradigm results (30%, 60%, 100%
and rewarded-trial only versions combined). ChR2: n=3 mice, 20 block changes; No Opsin: n=4 mice, 20 block changes.
b, Same layout as **Fig 5d**, Wilcoxon rank sum, *p<0.05, **p<0.01, ***p<0.001. **c,** Comparison of ChR2 and No Opsin
set-point dynamics, single sessions (2 mice, 1 session each). Despite exposure to blue LED, No Opsin mouse does not
show early-shift during stimulation blocks. **d-e,** Single session block change set-point and dynamics. Thick trace: mean,
thin traces: individual sessions. Blue: ChR2 mice, Black: No Opsin Mice. **f,** *Top:* Overlay of mean set-point and rewDA
dynamics at block change. *Bottom:* Pearson's correlation of set-point and rewDA across all Alternating Stimulation
Block sessions. **g-h,** Rapid Satiation Task: comparison of set-point and rewDA dynamics for single sessions and across
all sessions. **g**: Left: Single session examples shown for paired, interleaved sessions from the same mouse. Black
line/dagger(‡) in the top single-session plot indicates that the interval without red dots had no rewarded trials during
this particular session (the set-point was highly late-shifted with reduced participation during this time, see panel *h*).
Right: All photometry sessions. Thin traces: single sessions; thick: average. **h**: Left: Behavior from the same paired
single sessions as in panel *g*, Right: all behavior sessions. **i-j,** Timeshift Task: example No Juice session set-point
dynamics. **k,** Unlike DA, the co-monitored tdt control channel was not consistently correlated with set-point changes
and did not explain away the correlation between outcome-related DA and set-point (compare to **Fig 5l**).

Supplemental Methods

Regression to the median analyses.

Timing behavior may arise from a stochastic generative process, in which timing on any trial can be modeled as stochastic "draws" from a latent distribution, T. This presents potential confounds to detecting trial-by-trial learning. In this framework, the expected median of the next trial's movement time would be the median of the latent distribution on the next trial, $T^{(n+1)}$. If the latent distribution is stationary trial-by-trial (i.e., $T^{(n)} = T^{(n+1)}$), we would still see changes in movement timing on the next trial (Δt_n) that arise by chance, but which are not necessarily in response to learning. Thus, even though some deterministic process (say, outcome-related DA signaling) might control this tendency to "regress to the median" of $T^{(n)}$, we could not rule out the possibility that regression to the median of $T^{(n)}$ happens by chance (i.e., due to a different process). Thus, to detect learning effects in excess of chance would require that Δt_n be in excess of the regression to the median effect. In a stochastic generative model of timing, we would interpret this kind of learning as an update to the latent distribution on the next trial, such that $T^{(n)} \neq T^{(n+1)}$. We define regression to the median on each trial, n, as:

$$R2M_n = \text{median}(T_n) - t_n$$

Where t_n is the movement time on trial n and T_n is the vector of all movement times in consideration. We say "in consideration" because we cannot simply use all movement times in the session due to slow, non-stationarity in the timing distribution across the session, which might arise from processes other than learning (e.g., satiation/fatigue) and which necessitates a more local calculation of median(T_n) to isolate "micro-nonstationarity" arising from trial-by-trial updates. Thus, the median(T_n) is an uncertain quantity. To address this, we calculated median(T_n) in 3 complementary ways:

    A. We broke each session into deciles of trials and calculated the median of the distribution within each decile.
    B. We broke the session into deciles as in A, but we bootstrapped the next trial's movement time by resampling the distribution of movement times within the decile 10,000 times.
    C. We calculated a local, running median movement time for each trial (100 trials, centered at current trial).

$R2M_n$ for each method was then compared to the true Δt_n on each trial, defined as in the rest of the paper ($\Delta t_n = t_{n+1} - t_n$; where late-shifting is positive by convention). Because of the high level of trial-by-trial variability, we resolved Δt in excess of R2M as a significant difference between the distributions of $R2M_n$ vs Δt_n across all trials (R2M and ΔT), calculated by Wilcoxon rank sum test. The median of each distribution was then used to calculate "ΔT in excess of chance" (**Fig 1g**) defined as:

$$\Delta T \text{ in excess of chance} = \text{median}(\Delta T - R2M)$$

1228 Assuming timing arises by a stochastic generative process, this excess term is interpreted as the
1229 change in the median of $T^{(n+1)}$ versus $T^{(n)}$.
1230
For the analysis dataset, we used all +Juice photometry sessions that recorded DAN/DA signals
from mice expressing no opsin (17 mice, 155 sessions). To ascribe excess effects to trials with
different outcomes, we pooled trials either within 0.5s bins of time with respect to the cue on
trial n (as in **Fig 1g**). We also tried more broadly pooling early and rewarded trials (statistics
reported in-line in the text). "Early trials" included all trials with a first-lick time falling within a
"notch" of the timing distribution (between the local median and opening of the reward window
at 3.333s). We looked at notch trials because we expected learning updates (late-shift) and R2M
effects (early-shift) for these trials to be in opposite directions, presenting the most challenging
conditions to detect learning effects. To put pooled early and rewarded trial analyses on the same
footing, the rewarded trial "notch" included trials pooled within an equivalent interval in the
reward window (e.g., if the notch were between 3 to 3.333s for a local measurement, the
rewarded notch interval was taken as 3.333-3.666s).
1243
All three R2M approximation methods (A, B and C) returned similar results in all analyses. R2M
effects were present in the dataset, indicating the mice did indeed have a tendency to "regress
to the median" movement time, consistent with a stochastic generative process underlying timing.
However, ΔT was in excess of R2M by all metrics and depended on the trial outcome: early trials
(>0.5 to 3.333s) showed late-shifting in excess of chance, whereas rewarded trials showed early-
shifting—consistent with learning effects. Interestingly, "reaction" trials, within 500ms of the cue,
also showed small but significant early-shifting, which is likely a multi-trial effect (i.e., rewarded
trials are often followed by reactions, and the subsequent trial (n+2) also appears to show some
residual early-shift). Although there was no opportunity for mice to be "late" in this version of the
timing task, rare trials in which the first-lick fell within the ITI (unrewarded) showed subsequent
late-shifting, consistent with DA dips observed on these trials. It is important to note that these
late-shifts represent shifts in the *median* of the putative latent distribution on trial n+1, which
were small (100s of milliseconds) compared to the first-lick time (>7s)—which means that the
first-lick time on trial n+1 was still more likely to be much earlier than that on trial n. Finally, we
note that the size of the excess ΔT effects was on the same order of magnitude as one-shot
optogenetic effects (10s-100s of milliseconds, **Fig 3**), suggesting that outcome-related DA controls
updates to the stochastic "set-point" of the latent distribution on trial n+1.
1261
1262
1263 **Bleaching analyses**
1264 As is standard for photometry measurements, we corrected for bleaching using dF/F.
1265 Nevertheless, there remains some possibility that bleaching is either not fully corrected by this
1266 processing *or* that dF/F itself introduces some insidious artifact that could explain the downtrend
1267 in rewarded DA transients ("rewDA") across the session. *A priori*, this is unlikely:
1268
1. The amplitude of unrewarded transients showed relatively little change across the session
(**Extended Data Fig 7e,f,k**)—bleaching artifacts would likely affect signals similarly for all
trial outcomes.

2. Indirect pathway striatal projection neuron (iSPN) GCaMP signals showed the *opposite*
trend across the session on average: rewarded iSPN transient amplitude *increased*, on
average (n=3 A2A-cre mice, 30 sessions; **Extended Data Fig 8 bottom row**). iSPNs receive
negative input from DANs, thus these findings are consistent with decreased DAN
signaling.
3. The timing set-point showed abrupt shifts that occurred at the moments of the greatest
decline in rewDA (**Fig 4e**), these changes occurred at different times in the session
(**Extended Data Fig 7o**) and occurred significantly earlier in the No Juice Task (**Fig 4k**).
Bleaching would likely cause similar decay rates across all sessions from the same mouse.
4. Reward-related tdTomato transients (rew-tdt) did not show the same pattern of decline
across the session (**Extended Data Fig 7g,l, Extended Data Fig 8 top row**) and did not
explain abrupt shifts in set-point (**Extended Data Fig 7i**).
5. Causal effects of rewDA on set-point cannot be explained away by bleaching (**Fig 5**,
Extended Data Fig 9).
6. Rapid Satiation (**Fig 5g-i**) caused a sudden, precipitous decline in rewDA that was
significantly different from that observed in control sessions in the same mice, which
cannot be explained by bleaching.
1289

1290 Nevertheless, to directly assess the contribution of bleaching to the decline in rewDA, we derived
1291 a GLM predicting the peak amplitude of rewDA from bleaching-related and behavioral predictors.
1292 To derive a bleaching predictor, we applied the raw fluorescence, F0, from the dF/F correction. F0
1293 was measured for each trial as the average fluorescence signal during the 5s preceding the Lamp-
1294 off event, averaged across the 10 nearest neighboring trials.[7]
1295

1296 We first fit the model with F0 alone. As expected, because F0 decreases monotonically, F0 was
1297 positively correlated with set-point and the decline in rewDA, and thus was a significant positive
1298 coefficient in the "bleaching-only model." This verified that bleaching was encoded properly.
1299

1300 Subsequently, we introduced behavioral predictors to derive nested models containing
1301 permutations of the following predictors:
1302

1303
1304
Across DAN/DA recordings, predictors related to ongoing body movements, optical artifacts, and
first-lick time were relatively poor predictors of rewDA, as evaluated by variance explained and 4
complementary model criterion metrics (AIC, BIC, AICc, and CAIC). For the +Juice Task, the best
models always included the number of rewards received (nRR) as a significant predictor, which
weakened F0's coefficient, indicating that variance in rewDA was independently explained by

variables related to satiation and/or behavioral accuracy, in excess of bleaching. Supporting this,
when the models were fit on data from the Timeshift Task, in which the set-point no longer
trivially correlated with nRR, set-point became a stronger predictor in the model, and all 4
information criterion metrics agreed that the most parsimonious model of rewDA required both
set-point and nRR as significant predictors, in excess of bleaching. Together, these data indicate
that rewDA signals contained information related to propensity to move +/- satiation, in excess
of bleaching. Thus, bleaching did not explain away the relationship of rewDA to changes in the
timing set-point.

References

| 1318 | **References** |
| 1320 | 1 | Schultz, W., Dayan, P. & Montague, P. R. A neural substrate of prediction and reward. *Science* **275**, 1593-1599 (1997). https://doi.org/10.1126/science.275.5306.1593 |

1318   **References**
1319

1320   1   Schultz, W., Dayan, P. & Montague, P. R. A neural substrate of prediction and reward.
1321   *Science* **275**, 1593-1599 (1997). https://doi.org/10.1126/science.275.5306.1593

1322   2   Schultz, W., Apicella, P. & Ljungberg, T. Responses of monkey dopamine neurons to
1323   reward and conditioned stimuli during successive steps of learning a delayed response
1324   task. *J Neurosci* **13**, 900-913 (1993). https://doi.org/10.1523/JNEUROSCI.13-03-
1325   00900.1993

1326   3   Coddington, L. T., Lindo, S. E. & Dudman, J. T. Mesolimbic dopamine adapts the rate of
1327   learning from action. *Nature* **614**, 294-302 (2023). https://doi.org/10.1038/s41586-022-
1328   05614-z

1329   4   Mohebi, A. *et al.* Dissociable dopamine dynamics for learning and motivation. *Nature*
1330   **570**, 65-70 (2019). https://doi.org/10.1038/s41586-019-1235-y

1331   5   Hamid, A. A. *et al.* Mesolimbic dopamine signals the value of work. *Nat Neurosci* **19**,
1332   117-126 (2016). https://doi.org/10.1038/nn.4173

1333   6   Niv, Y., Daw, N. D., Joel, D. & Dayan, P. Tonic dopamine: opportunity costs and the control
1334   of response vigor. *Psychopharmacology (Berl)* **191**, 507-520 (2007).
1335   https://doi.org/10.1007/s00213-006-0502-4

1336   7   Hamilos, A. E. *et al.* Slowly evolving dopaminergic activity modulates the moment-to-
1337   moment probability of reward-related self-timed movements. *Elife* **10** (2021).
1338   https://doi.org/10.7554/eLife.62583

1339   8   Albin, R. L., Young, A. B. & Penney, J. B. The functional anatomy of basal ganglia
1340   disorders. *Trends Neurosci* **12**, 366-375 (1989). https://doi.org/10.1016/0166-
1341   2236(89)90074-x

1342   9   Olds, J. & Milner, P. Positive reinforcement produced by electrical stimulation of septal
1343   area and other regions of rat brain. *J Comp Physiol Psychol* **47**, 419-427 (1954).
1344   https://doi.org/10.1037/h0058775

1345   10   Steinberg, E. E. *et al.* A causal link between prediction errors, dopamine neurons and
1346   learning. *Nat Neurosci* **16**, 966-973 (2013). https://doi.org/10.1038/nn.3413

1347   11   Markowitz, J. E. *et al.* Spontaneous behaviour is structured by reinforcement without
1348   explicit reward. *Nature* **614**, 108-117 (2023). https://doi.org/10.1038/s41586-022-
1349   05611-2

1350   12   Xiao, L. *et al.* A Basal Ganglia Circuit Sufficient to Guide Birdsong Learning. *Neuron* **98**,
1351   208-221 e205 (2018). https://doi.org/10.1016/j.neuron.2018.02.020

1352   13   Adamantidis, A. R. *et al.* Optogenetic interrogation of dopaminergic modulation of the
1353   multiple phases of reward-seeking behavior. *J Neurosci* **31**, 10829-10835 (2011).
1354   https://doi.org/10.1523/JNEUROSCI.2246-11.2011

1355   14   Kim, K. M. *et al.* Optogenetic mimicry of the transient activation of dopamine neurons by
1356   natural reward is sufficient for operant reinforcement. *PLoS One* **7**, e33612 (2012).
1357   https://doi.org/10.1371/journal.pone.0033612

1358   15   Tsai, H. C. *et al.* Phasic firing in dopaminergic neurons is sufficient for behavioral
1359   conditioning. *Science* **324**, 1080-1084 (2009). https://doi.org/10.1126/science.1168878

1360   16   Jeong, H. *et al.* Mesolimbic dopamine release conveys causal associations. *Science* **378**,
1361   eabq6740 (2022). https://doi.org/10.1126/science.abq6740

17 Starkweather, C. K., Babayan, B. M., Uchida, N. & Gershman, S. J. Dopamine reward
prediction errors reflect hidden-state inference across time. *Nat Neurosci* **20**, 581-589
(2017). https://doi.org/10.1038/nn.4520
18 Mikhael, J. G., Kim, H. R., Uchida, N. & Gershman, S. J. Ramping and State Uncertainty in
the Dopamine Signal. *bioRxiv*, 805366 (2019). https://doi.org/10.1101/805366
19 Kim, H. R. *et al.* A unified framework for dopamine signals across timescales. *bioRxiv*,
803437 (2019). https://doi.org/10.1101/803437
20 Qian, L. *et al.* Prospective contingency explains behavior and dopamine signals during
associative learning. *Nat Neurosci* (2025). https://doi.org/10.1038/s41593-025-01915-4
21 Gadagkar, V. *et al.* Dopamine neurons encode performance error in singing birds. *Science*
354, 1278-1282 (2016). https://doi.org/10.1126/science.aah6837
22 Kasdin, J. *et al.* Natural behaviour is learned through dopamine-mediated reinforcement.
Nature (2025). https://doi.org/10.1038/s41586-025-08729-1
23 da Silva, J. A., Tecuapetla, F., Paixão, V. & Costa, R. M. Dopamine neuron activity before
action initiation gates and invigorates future movements. *Nature* **554**, 244-248 (2018).
https://doi.org/10.1038/nature25457
24 Howe, M. W. & Dombeck, D. A. Rapid signalling in distinct dopaminergic axons during
locomotion and reward. *Nature* **535**, 505-510 (2016).
https://doi.org/10.1038/nature18942
25 Ko, D. & Wanat, M. J. Phasic Dopamine Transmission Reflects Initiation Vigor and Exerted
Effort in an Action- and Region-Specific Manner. *J Neurosci* **36**, 2202-2211 (2016).
https://doi.org/10.1523/JNEUROSCI.1279-15.2016
26 Cai, X. *et al.* Dopamine dynamics are dispensable for movement but promote reward
responses. *Nature* **635**, 406-414 (2024). https://doi.org/10.1038/s41586-024-08038-z
27 Cepeda, C., Murphy, K. P., Parent, M. & Levine, M. S. The role of dopamine in
Huntington's disease. *Prog Brain Res* **211**, 235-254 (2014).
https://doi.org/10.1016/B978-0-444-63425-2.00010-6
28 Niv, Y. Neuroscience: Dopamine ramps up. *Nature* **500**, 533-535 (2013).
https://doi.org/10.1038/500533a
29 Salamone, J. D. & Correa, M. The Neurobiology of Activational Aspects of Motivation:
Exertion of Effort, Effort-Based Decision Making, and the Role of Dopamine. *Annu Rev*
Psychol **75**, 1-32 (2024). https://doi.org/10.1146/annurev-psych-020223-012208
30 Coddington, L. T. & Dudman, J. T. The timing of action determines reward prediction
signals in identified midbrain dopamine neurons. *Nat Neurosci* **21**, 1563-1573 (2018).
https://doi.org/10.1038/s41593-018-0245-7
31 Parker, N. F. *et al.* Reward and choice encoding in terminals of midbrain dopamine
neurons depends on striatal target. *Nat Neurosci* **19**, 845-854 (2016).
https://doi.org/10.1038/nn.4287
32 Gallistel, C. R. & Gibbon, J. Time, rate, and conditioning. *Psychol Rev* **107**, 289-344
(2000). https://doi.org/10.1037/0033-295x.107.2.289
33 Merchant, H., Harrington, D. L. & Meck, W. H. Neural basis of the perception and
estimation of time. *Annu Rev Neurosci* **36**, 313-336 (2013).
https://doi.org/10.1146/annurev-neuro-062012-170349

34 Lee, I. H. & Assad, J. A. Putaminal activity for simple reactions or self-timed movements.
J Neurophysiol **89**, 2528-2537 (2003). https://doi.org/10.1152/jn.01055.2002

35 Kirshenbaum, A. P., Brown, S. J., Hughes, D. M. & Doughty, A. H. Differential-
reinforcement-of-low-rate-schedule performance and nicotine administration: a
systematic investigation of dose, dose-regimen, and schedule requirement. *Behav*
Pharmacol **19**, 683-697 (2008). https://doi.org/10.1097/FBP.0b013e328315ecbb

36 Eckard, M. L. & Kyonka, E. G. E. Differential reinforcement of low rates differentially
decreased timing precision. *Behav Processes* **151**, 111-118 (2018).
https://doi.org/10.1016/j.beproc.2018.02.022

37 Malapani, C. *et al.* Coupled temporal memories in Parkinson's disease: a dopamine-
related dysfunction. *J Cogn Neurosci* **10**, 316-331 (1998).
https://doi.org/10.1162/089892998562762

38 Meck, W. H. Affinity for the dopamine D2 receptor predicts neuroleptic potency in
decreasing the speed of an internal clock. *Pharmacol Biochem Behav* **25**, 1185-1189
(1986). https://doi.org/10.1016/0091-3057(86)90109-7

39 Meck, W. H. Neuroanatomical localization of an internal clock: a functional link between
mesolimbic, nigrostriatal, and mesocortical dopaminergic systems. *Brain Res* **1109**, 93-
107 (2006). https://doi.org/10.1016/j.brainres.2006.06.031

40 Dews, P. B. & Morse, W. H. Some observations on an operant in human subjects and its
modification by dextro amphetamine. *J Exp Anal Behav* **1**, 359-364 (1958).
https://doi.org/10.1901/jeab.1958.1-359

41 Schuster, C. R. & Zimmerman, J. Timing behavior during prolonged treatment with dl-
amphetamine. *J Exp Anal Behav* **4**, 327-330 (1961). https://doi.org/10.1901/jeab.1961.4-
327

42 Helassa, N., Podor, B., Fine, A. & Torok, K. Design and mechanistic insight into ultrafast
calcium indicators for monitoring intracellular calcium dynamics. *Sci Rep* **6**, 38276
(2016). https://doi.org/10.1038/srep38276

43 Chen, Z. *et al.* Direct and indirect pathway neurons in ventrolateral striatum differentially
regulate licking movement and nigral responses. *Cell Rep* **37**, 109847 (2021).
https://doi.org/10.1016/j.celrep.2021.109847

44 Lee, J., Wang, W. & Sabatini, B. L. Anatomically segregated basal ganglia pathways allow
parallel behavioral modulation. *Nat Neurosci* **23**, 1388-1398 (2020).
https://doi.org/10.1038/s41593-020-00712-5

45 Taniguchi, J. *et al.* Comment on 'Accumbens cholinergic interneurons dynamically
promote dopamine release and enable motivation'. *bioRxiv*, 2023.2012.2027.573485
(2024). https://doi.org/10.1101/2023.12.27.573485

46 Yang, Z., Inagaki, M., Gerfen, C. R., Fontolan, L. & Inagaki, H. K. Integrator dynamics in
the cortico-basal ganglia loop underlie flexible motor timing. *bioRxiv* (2024).
https://doi.org/10.1101/2024.06.29.601348

47 Ward, R., Wu, X. & Bottou, L. AdaGrad stepsizes: Sharp convergence over nonconvex
landscapes. *J Mach Learn Res* **21**, 1-30 (2020).

48 Berke, J. D. What does dopamine mean? *Nat Neurosci* **21**, 787-793 (2018).
https://doi.org/10.1038/s41593-018-0152-y

49 Gardner, M. P. H., Schoenbaum, G. & Gershman, S. J. Rethinking dopamine as
generalized prediction error. *Proc Biol Sci* **285** (2018).
https://doi.org/10.1098/rspb.2018.1645
50 Beron, C. C., Neufeld, S. Q., Linderman, S. W. & Sabatini, B. L. Mice exhibit stochastic and
efficient action switching during probabilistic decision making. *Proc Natl Acad Sci U S A*
119, e2113961119 (2022). https://doi.org/10.1073/pnas.2113961119
51 Long, C. *et al.* Physiological constraints on the rapid dopaminergic modulation of striatal
reward activity. *bioRxiv* (2022). https://doi.org/10.1101/2022.09.16.508310
52 Duchi, J., Hazan, E. & Singer, Y. Adaptive subgradient methods for online learning and
stochastic optimization. *J Mach Learn Res* **12(Jul)**, 2121-2159 (2011).
53 Remington, E. D., Narain, D., Hosseini, E. A. & Jazayeri, M. Flexible Sensorimotor
Computations through Rapid Reconfiguration of Cortical Dynamics. *Neuron* **98**, 1005-
1019 e1005 (2018). https://doi.org/10.1016/j.neuron.2018.05.020
54 Sohn, H., Narain, D., Meirhaeghe, N. & Jazayeri, M. Bayesian Computation through
Cortical Latent Dynamics. *Neuron* **103**, 934-947 e935 (2019).
https://doi.org/10.1016/j.neuron.2019.06.012
55 Maimon, G. & Assad, J. A. A cognitive signal for the proactive timing of action in
macaque LIP. *Nat Neurosci* **9**, 948-955 (2006). https://doi.org/10.1038/nn1716
56 Yagishita, S. *et al.* A critical time window for dopamine actions on the structural plasticity
of dendritic spines. *Science* **345**, 1616-1620 (2014).
https://doi.org/10.1126/science.1255514
57 Lee, K. *et al.* Temporally restricted dopaminergic control of reward-conditioned
movements. *Nat Neurosci* **23**, 209-216 (2020). https://doi.org/10.1038/s41593-019-
0567-0
58 Salinas-Hernandez, X. I., Zafiri, D., Sigurdsson, T. & Duvarci, S. Functional architecture of
dopamine neurons driving fear extinction learning. *Neuron* **111**, 3854-3870 e3855
(2023). https://doi.org/10.1016/j.neuron.2023.08.025
59 Salinas-Hernandez, X. I. *et al.* Dopamine neurons drive fear extinction learning by
signaling the omission of expected aversive outcomes. *Elife* **7** (2018).
https://doi.org/10.7554/eLife.38818
60 Floeder, J. R., Jeong, H., Mohebi, A. & Namboodiri, V. M. K. Mesolimbic dopamine ramps
reflect environmental timescales. *bioRxiv* (2024).
https://doi.org/10.1101/2024.03.27.587103
61 McClure, S. M., Daw, N. D. & Montague, P. R. A computational substrate for incentive
salience. *Trends Neurosci* **26**, 423-428 (2003). https://doi.org/10.1016/s0166-
2236(03)00177-2
62 Tritsch, N. X. & Sabatini, B. L. Dopaminergic modulation of synaptic transmission in
cortex and striatum. *Neuron* **76**, 33-50 (2012).
https://doi.org/10.1016/j.neuron.2012.09.023
63 Lahiri, A. K. & Bevan, M. D. Dopaminergic Transmission Rapidly and Persistently
Enhances Excitability of D1 Receptor-Expressing Striatal Projection Neurons. *Neuron* **106**,
277-290 e276 (2020). https://doi.org/10.1016/j.neuron.2020.01.028
64 Lee, S. J. *et al.* Cell-type-specific asynchronous modulation of PKA by dopamine in
learning. *Nature* **590**, 451-456 (2021). https://doi.org/10.1038/s41586-020-03050-5

65   Mello, G. B., Soares, S. & Paton, J. J. A scalable population code for time in the striatum.
Curr Biol **25**, 1113-1122 (2015). https://doi.org/10.1016/j.cub.2015.02.036

66   Costacurta, J. C., Bhandarkar, S., Zoltowski, D. M. & Linderman, S. W. Structured
flexibility in recurrent neural networks via neuromodulation. *bioRxiv* (2024).
https://doi.org/10.1101/2024.07.26.605315

67   Lodder, B. *et al.* Absolute measurement of fast and slow neuronal signals with
fluorescence lifetime photometry at high temporal resolution. *bioRxiv* (2025).
https://doi.org/10.1101/2025.01.10.632162

68   Backman, C. M. *et al.* Characterization of a mouse strain expressing Cre recombinase
from the 3' untranslated region of the dopamine transporter locus. *Genesis* **44**, 383-390
(2006). https://doi.org/10.1002/dvg.20228

1503